

COMPARING COHOMOLOGY VIA EXACT SPLIT PAIRS IN DIAGRAM ALGEBRAS

SULAKHANA CHOWDHURY AND GEETHA THANGAVELU

ABSTRACT. In this article, we compare the cohomology between the categories of modules of the diagram algebras and the categories of modules of its input algebras. Our main result establishes a sufficient condition for exact split pairs between these two categories, analogous to a work by Diracca and Koenig [7]. To be precise, we prove the existence of the exact split pairs in A -Brauer algebras, cyclotomic Brauer algebras, and walled Brauer algebras with their respective input algebras.

1. INTRODUCTION

Representation theory stands as one of the cornerstones among the disciplines in mathematics, with its primary objectives revolving around the classification and description of irreducible (also known as simple) representations. In the context of ring homomorphisms, directly comparing the cohomology of two rings is often a challenging task. However, in [7], a method is introduced by Diracca and Koenig to overcome this difficulty. This approach combines the aspects of subring and quotient ring situations, allowing for cohomology comparison without the need for strong assumptions typically used in either of these scenarios. This technique can be used for the non-vanishing of cohomology in specific situations. The concept of exact split pair of functors provides an interesting tool to compare the cohomology between rings under particular conditions.

Diagram algebras play a significant role in the Schur-Weyl duality, a connection between representations of diagram algebras and algebraic groups. The classical Schur-Weyl duality, establishes a relationship between the polynomial representations of general linear groups GL_n over the complex numbers \mathbb{C} and representations of the symmetric groups \mathfrak{S}_r , for details see [10, 20]. Later on, Brauer examined an analogous problem for the symplectic and the orthogonal groups. The results on the Brauer algebras have been extended to walled Brauer algebras, Brauer algebras of type C, cyclotomic Brauer algebras, and A -Brauer algebras. The walled Brauer algebras were introduced as centralizer algebras of GL_n acting on the mixed tensor space in [1], and are subalgebras of the Brauer algebras. The cyclotomic Brauer algebras were studied by Rui, Yu in [18], Bowman, Cox, and Visscher in [3], and A -Brauer algebras were studied in [8]. The primary focus of this article is to study an exact split pair for these algebras.

In this article, we use the framework of cellular algebras introduced by Koenig and Xi in [15, 17], where every cellular algebra can be written as an iterated inflations involving smaller cellular algebras. Using this framework Hartmann, Henke, Koenig, and Paget studied the cellularly stratified algebras and their split pairs in [[12], Section 3]. A -Brauer algebras are known to be cyclic cellular but the cellularly stratified structure is yet to be known. Therefore, we used results from [7] to obtain an exact split pair for A -Brauer algebras, cyclotomic Brauer algebras and the walled Brauer algebras.

2020 *Mathematics Subject Classification.* 18E30,20J05,20C30,16D40.

Key words and phrases. Cellular algebras, cohomological comparison, exact split pair, A -Brauer algebras, cyclotomic Brauer algebras, walled Brauer algebras.

The primary objective is to determine the sufficient condition for the existence of an exact split pair for the above mentioned class of diagram algebras and we use these to compare Hom, Ext and study the associated numerical invariants, including the global dimensions.

This article is organized as follows. In Section 2, we review definitions and properties of an exact split pair. In Section 3, we show an iterated inflation for A -Brauer algebras, which allows us to prove the existence of an exact split pair for the A -Brauer algebras. In Section 4 and Section 5, we prove our main result for the cyclotomic Brauer algebras and the walled Brauer algebras.

2. PRELIMINARIES

In this section, we recall some definitions from [7], which will be used in this article.

Let \mathcal{A} and \mathcal{B} be two additive categories. A pair (F, G) of two additive functors $F : \mathcal{A} \rightarrow \mathcal{B}$ and $G : \mathcal{B} \rightarrow \mathcal{A}$ is a *split pair of functors* if the composition $F \circ G$ is an auto-equivalence of \mathcal{B} . If \mathcal{A} and \mathcal{B} are equipped with exact structures, and F and G are exact functors with respect to these exact structures, then the split pair is called an *exact split pair of functors*.

Let A and B be two rings. We call B a *split quotient* of A if there is an embedding ϵ sending the unit of B to the unit of A and there exists a surjective homomorphism $\pi : A \rightarrow B$ such that $\pi \circ \epsilon = 1_B$. Let $\mathbf{mod}\text{-}A$ and $\mathbf{mod}\text{-}B$ be the categories of finitely generated right A -modules and right B -modules, respectively. The homomorphism π and ϵ induce two exact functors F and G between the exact categories $\mathbf{mod}\text{-}A$ and $\mathbf{mod}\text{-}B$ such that $F \circ G$ is the identity on $\mathbf{mod}\text{-}B$, where $F = - \otimes_A A_B$ and $G = - \otimes_B B_A$.

Let e be an idempotent of a ring A . Let B be the centralizer subring eAe . Consider eA and Ae to be the B - A and A - B bimodules, respectively. Assume Ae to be eAe -projective. Then the functors

$$\begin{aligned} F : \mathbf{mod}\text{-}A &\longrightarrow \mathbf{mod}\text{-}B & G : \mathbf{mod}\text{-}B &\longrightarrow \mathbf{mod}\text{-}A \\ N &\longmapsto N \otimes_A Ae & M &\longmapsto M \otimes_B eA \end{aligned}$$

form an exact split pair of functors. We call the centralizer subrings eAe as *corner rings*.

Let e be an idempotent in A , and B a split quotient of eAe , viewed as a subring of eAe . Then we call B a *corner split quotient* of A with respect to e , if there is an eAe - A bimodule S , which is projective as left B -module via the embedding of B into eAe , and also satisfies the condition $Se \cong B$ as right B -modules.

To prove the sufficient condition for the existence of an exact split pair, we will frequently use the following result.

Theorem 2.1 ([7], Lemma 3.2). *Let A and B be any arbitrary rings and S a left B -module. If B is a corner split quotient of A . Then the functors $F = - \otimes_A Ae$ and $G = - \otimes_B B \otimes_{eAe} S$ form an exact split pair.*

3. THE A -BRAUER ALGEBRA

Let R be a commutative unital ring, with $\delta \in R$. Let A be a unitary associative algebra over R , not necessarily commutative. Assume that A possesses an R -linear algebra involution denoted by $*$, along with a $*$ -invariant R -valued trace, denoted as tr , where $\text{tr}(1) = \delta$. An A -Brauer diagram is a Brauer diagram with A -labeled edges and has an orientation of edges. We denote the A -Brauer algebra by $D_n(A)$. Note that the algebra $D_n(A)$ is generated by s_i , e_i , and h_j^a where $a \in A$, $1 \leq i \leq n-1$, and $1 \leq j \leq n$ (see Figure 1).




FIGURE 1. Generators of the A -Brauer algebra.

Let X and Y be two A -Brauer diagrams. The product $X \cdot Y$ is defined by placing X on top of Y and concatenating the bottom row of X with the top row of Y to form a new diagram Z . The product rule is similar to that of Brauer diagrams. Fix an orientation of labels in the concatenated diagram and apply $*$ on the label to reverse the orientation if needed. Each edge in the product is labeled by the product of the labels of the component edges. For each loop, multiply the diagram by the R -valued trace of its label. Then, $X \cdot Y = \alpha Z$, where the scalar α is the product of R -valued trace of all closed loops. For example, see [[8], Figure 5.3]. For more details, see [[8], Section 5].

Let $A \wr \mathfrak{S}_n$ be the wreath product algebra, a subalgebra of $D_n(A)$ generated by s_i and h_j^a for $1 \leq i \leq n - 1, a \in A$, and $1 \leq j \leq n$. A cellular algebra is said to be *cyclic cellular* if every cell module of A is cyclic [[8], Section 2.4.3]. Let (Γ, \geq) be a finite poset. Let A be a cyclic cellular algebra with cell datum $(A, *, \Gamma, \geq, \mathcal{T}, \mathcal{C})$, described as in [[8], Lemma 2.5]. If the algebra A is cyclic cellular, then $A \wr \mathfrak{S}_n$ is cyclic cellular with cell datum $(A \wr \mathfrak{S}_n, *, \Lambda_n^\Gamma, \supseteq_\Gamma, \mathcal{M}, \mathcal{B})$, see [[8], Theorem 4.1].

When δ is invertible in R , the idempotent is given by Figure 2. When $\delta = 0$ and n is odd, the idempotent is defined as in Figure 3. When $\delta = 0$ and n is even, using the same idempotent as in Figure 2 results in the square of the two-sided ideal J being zero, where J is the two-sided ideal of $D_n(A)$ containing all A -Brauer diagrams with only horizontal edges. Hence, we shall not consider the case $\delta = 0$ and n is even.




FIGURE 2. Idempotent of the A -Brauer algebra for an invertible δ .




FIGURE 3. Idempotent of the A -Brauer algebra when $\delta = 0$, and n is odd.

The cellular structure of A -Brauer algebra is known from [8], we will prove that the algebra $D_n(A)$ can be written as an iterated inflation of the wreath product algebra $A \wr \mathfrak{S}_n$. The “iterated inflation” is an inductive construction of cellular algebra from a smaller cellular algebra, see [[17], Section 3] for more details. Let $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$, where $\lfloor \frac{n}{2} \rfloor$ is the integer part of $\frac{n}{2}$ and J_l be the free R -module generated by all A -Brauer diagrams with at least l horizontal edges. Throughout the paper, we assume that the diagram having l horizontal edges means it has l horizontal edges in each row.

Lemma 3.1. *If $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$, then J_l is a two-sided ideal of $D_n(A)$.*

Proof. For any $d_1, d_2 \in D_n(A)$ with m_1 and m_2 horizontal edges, the product $d_1 \cdot d_2$ have at least $\max\{m_1, m_2\}$ horizontal edges. Thus, J_l is a two-sided ideal, where $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$. \square

Let $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$. Consider the chain of two-sided ideals of $D_n(A)$

$$0 \subseteq J_{\lfloor \frac{n}{2} \rfloor} \subseteq \cdots \subseteq J_1 \subseteq J_0 = D_n(A),$$

where each J_l is generated by the idempotent e_l , which we defined already. An (n, l) A -labeled partial diagram is a one-row diagram on n vertices with l A -labeled horizontal edges, and $n - 2l$ free vertices. Let $V(n, l, A)$ be the free R -module generated by all (n, l) A -labeled partial diagrams. From now on, let us assume that R is a commutative Noetherian domain.

Consider the R -algebra $V(n, l, A) \otimes_R V(n, l, A) \otimes_R A \wr \mathfrak{S}_{n-2l}$ having a basis of the form $e \otimes f \otimes \pi_d$, where e and f are respectively the top and bottom configuration of an A -Brauer diagram d , and $\pi_d \in A \wr \mathfrak{S}_{n-2l}$. The multiplication in $V(n, l, A) \otimes V(n, l, A) \otimes A \wr \mathfrak{S}_{n-2l}$ is defined by

$$(e \otimes f \otimes \pi_d) \cdot (e' \otimes f' \otimes \pi_{d'}) = e \otimes f' \otimes \pi_d \phi(f, e') \pi_{d'},$$

where $\phi : V(n, l, A) \otimes_R V(n, l, A) \rightarrow A \wr \mathfrak{S}_{n-2l}$ is a bilinear form. Let the R -algebra J_l/J_{l+1} be denoted by B .

Lemma 3.2. *For $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$, B is isomorphic to the inflation $V(n, l, A) \otimes V(n, l, A) \otimes A \wr \mathfrak{S}_{n-2l}$ of the wreath product algebra along $V(n, l, A)$.*

Proof. The R -algebra B has a basis consisting of all diagrams with exactly l horizontal edges in the top row, another l horizontal edges in bottom row and $n - 2l$ vertical edges. Define an R -linear map

$$\begin{aligned} \Psi : B &\longrightarrow V(n, l, A) \otimes V(n, l, A) \otimes A \wr \mathfrak{S}_{n-2l} \\ d &\mapsto e \otimes f \otimes \pi_d \end{aligned}$$

as follows: Fix $d \in B$ and denote by e (resp. f) the (n, l) A -labeled partial diagram in top (resp. bottom) configuration of d . Renumber the free vertices of e and f as $1, 2, \dots, n - 2l$ from left to right. Then the vertical edges of d define a permutation $\pi_d \in A \wr \mathfrak{S}_{n-2l}$ sending the number assigned to top vertex of an edge to the number in its bottom vertex. We need to define a bilinear form ϕ to give a multiplicative structure on $V(n, l, A) \otimes V(n, l, A) \otimes A \wr \mathfrak{S}_{n-2l}$. This construction will use the map Ψ , thereby confirming that Ψ is indeed an R -algebra isomorphism. Clearly, Ψ is R -module isomorphism and we need to check that Ψ preserves the multiplication of two diagrams in $V(n, l, A) \otimes V(n, l, A) \otimes A \wr \mathfrak{S}_{n-2l}$. Let $d, d' \in B$ with $\Psi(d) = X$, and $\Psi(d') = Y$, where X and Y are of the form $e \otimes f \otimes \pi_d$ and $e' \otimes f' \otimes \pi_{d'}$, respectively. Then the product $X \cdot Y$ has to be of the form $e \otimes f' \otimes \pi_d \phi(f, e') \pi_{d'}$. The product $X \cdot Y$ is obtained by identifying the vertices in f with those in e . Now, we have two cases:

- (i) In the concatenated diagram if each horizontal edge of f and e' shares at least one common vertex, then we define the bilinear form $\phi : V(n, l, A) \times V(n, l, A) \rightarrow A \wr \mathfrak{S}_{n-2l}$ by $\phi(f, e') = \prod_i \text{tr}(a_i)^{l_i} \cdot \sigma$, where l_i is the number of connected components that lies entirely in the middle row of the concatenated diagram and σ is the permutation defined by the vertical edges of the product $\Psi^{-1}(f \otimes f \otimes \text{id}) \cdot \Psi^{-1}(e' \otimes e' \otimes \text{id})$. Then the product $X \cdot Y$ coincides with $\Psi(\Psi^{-1}(X) \Psi^{-1}(Y))$.
- (ii) If at least one horizontal edges in f which does not share a vertex with any horizontal edges in e' , then we define $\phi(f', e) = 0$. The number of horizontal edges of $\Psi^{-1}(X) \cdot \Psi^{-1}(Y)$ in $D_n(A)$ is strictly greater than l which implies that the product is zero inside B .

Hence, Ψ is an R -algebra isomorphism. \square

Remark 3.3. The R -rank of $V(n, l, A)$ is $(\dim A)^n \left(\frac{n!}{l!(n-2l)!2^l} \right)$.

Lemma 3.4. *Let $0 \leq k, l \leq \lfloor \frac{n}{2} \rfloor$. Let $d \in J_k/J_{k+1}$ and $d' \in J_l/J_{l+1}$ be two diagrams in $D_n(A)$, which maps under Ψ to $e \otimes f \otimes \pi_d$ and $e' \otimes f' \otimes \pi_{d'}$, respectively. Then the product $d \cdot d'$ is either an element of J_s/J_{s+1} or lies in J_{s+1} , where $s = \max\{k, l\}$. Moreover, if $\max\{k, l\} = l$, then the product is of the form $c(e'' \otimes e' \otimes \sigma \pi_{d'})$, where c is a scalar, $e'' \in V(n, l, A)$, and $\sigma \in A \wr \mathfrak{S}_{n-2l}$.*

Proof. The proof is similar to that of the Lemma 3.2. \square

In the following lemma, an R -linear anti-automorphism f of an R -algebra with $f^2 = \text{id}$ will be called an *involution*.

Lemma 3.5. *If $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$, then the involution $*$: $A \wr \mathfrak{S}_{n-2l} \rightarrow A \wr \mathfrak{S}_{n-2l}$ extends to an involution i on $D_n(A)$ given by $i(e \otimes f \otimes \pi_d) = (f \otimes e \otimes \pi_d^*)$.*

Proof. Straightforward. \square

As an immediate consequence, we have the following:

Theorem 3.6. *The algebra $D_n(A)$ is an iterated inflation of wreath product algebras $A \wr \mathfrak{S}_{n-2l}$ with the following decomposition*

$$D_n(A) \cong \bigoplus_{l=0}^{\lfloor \frac{n}{2} \rfloor} V(n, l, A) \otimes V(n, l, A) \otimes A \wr \mathfrak{S}_{n-2l}.$$

As a consequence of this theorem and [[17], Proposition 3.5], we recover the following result from [[8], Theorem 6.1].

Corollary 3.7. *Let A be a cyclic cellular algebra. Then, for $n \geq 1$, $D_n(A)$ is a cyclic cellular algebra.*

Proof. The assumption on A ensures that the wreath product algebra $A \wr \mathfrak{S}_n$ is cyclic cellular. Therefore, the A -Brauer algebra $D_n(A)$ is cyclic cellular by [[17], Proposition 3.5]. \square

3.1. Exact split pair $(D_n(A), A \wr \mathfrak{S}_n)$. The main objective of this article is to establish the sufficient condition for the existence of an exact split pair for the case of A -Brauer algebras using the method from [7]. Constructing an exact split pair involves identifying a corner split quotient, as outlined in the Theorem 2.1.

Proposition 3.8. *The algebra $A \wr \mathfrak{S}_n$ is a split quotient of $D_n(A)$.*

Proof. The subalgebra $A \wr \mathfrak{S}_n$ of $D_n(A)$ is a free R -module generated by s_i, h_j^a for $1 \leq i \leq n-1, 1 \leq j \leq n$, and $a \in A$. Let ϵ be a canonical inclusion of $A \wr \mathfrak{S}_n$ into $D_n(A)$, where the unit element of $A \wr \mathfrak{S}_n$ is mapped to the unit element of $D_n(A)$. Let us define a map $\pi : D_n(A) \rightarrow A \wr \mathfrak{S}_n$ in such a way that π takes each $d \in D_n(A)$ to a diagram that has no horizontal edges. Since π is surjective, the kernel of π contains all A -Brauer diagrams with at least one horizontal edge. Therefore, $\ker \pi$ coincides with J_1 , and $D_n(A)/J_1 \cong A \wr \mathfrak{S}_n$. Thus, $\pi \circ \epsilon$ is the identity on $A \wr \mathfrak{S}_n$. \square

From now on, assume δ is invertible in R , where R is a commutative Noetherian domain.

Proposition 3.9. *If $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$, then $D_{n-2l}(A)$ is isomorphic to the corner ring $e_l D_n(A) e_l$.*

Proof. The algebra $e_l D_n(A) e_l$ has a basis consisting of diagrams in $D_n(A)$ with at least l horizontal edges in the top (resp. bottom) row placed at the right side of the diagram. Consider the map ϕ defined by

$$\begin{aligned} \phi : D_{n-2l}(A) &\longrightarrow e_l D_n(A) e_l \\ d &\longmapsto e_l d e_l, \end{aligned}$$

where we identify d inside $D_n(A)$ by adding vertical edges from the vertices j in top row to the vertices j in bottom row, where $n-2l+1 \leq j \leq n$. Clearly, ϕ is a surjective homomorphism, and $\ker(\phi)$ contains only the zero element. This implies ϕ is injective, and hence, $D_{n-2l}(A) \cong e_l D_n(A) e_l$. \square

Theorem 3.10. *If $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$, then the algebra $A \wr \mathfrak{S}_{n-2l}$ is a corner split quotient of $D_n(A)$.*

Proof. By Proposition 3.8, $A \wr \mathfrak{S}_{n-2l}$ is a split quotient of $D_{n-2l}(A)$ and is isomorphic to the corner ring $e_l D_n(A) e_l$ by Proposition 3.9. Therefore, we only need to show that there is an $e_l D_n(A) e_l$ - $D_n(A)$ -bimodule S_l , which is projective as a left $A \wr \mathfrak{S}_{n-2l}$ -module and satisfies $S_l e_l \cong A \wr \mathfrak{S}_{n-2l}$ as a right $A \wr \mathfrak{S}_{n-2l}$ -modules.

As the right $D_n(A)$ -module, $e_l D_n(A)$ is a projective module generated by A -Brauer diagrams with at least l horizontal edges, where the top row contains the partial diagram of e_l . Recall that $e_l D_n(A) e_l$ consists of A -Brauer diagrams having at least l horizontal edges with the partial diagram of e_l on its top and bottom configurations. Therefore, $e_l D_n(A)$ is a left $e_l D_n(A) e_l$ -module. Now, the right action of $e_l D_n(A) e_l$ on $A \wr \mathfrak{S}_{n-2l}$ is given as follows: let $d \in D_n(A)$, and $\sigma \in A \wr \mathfrak{S}_{n-2l}$

$$\sigma \cdot d = \begin{cases} \sigma d & ; \text{ if } d \text{ has exactly } l \text{ horizontal edges} \\ 0 & ; \text{ otherwise.} \end{cases}$$

If d has more than l horizontal edges, then the quotient map $\alpha : e_l D_n(A) e_l \rightarrow e_l(D_n(A)/J_{l+1})e_l$ ($\cong \frac{e_l D_n(A) e_l}{e_l J_{l+1} e_l}$) is defined by $\alpha(e_l d e_l) = e_l d e_l \pmod{(J_{l+1})}$. Clearly, $\ker \alpha$ contains all diagrams having at least $l+1$ horizontal edges. The map α induces an action of $e_l D_n(A) e_l$ on $A \wr \mathfrak{S}_{n-2l}$. Define the $e_l D_n(A) e_l$ - $D_n(A)$ -bimodule S_l to be $A \wr \mathfrak{S}_{n-2l} \otimes_{e_l D_n(A) e_l} e_l D_n(A)$.

Next, we claim that S_l is a free left $A \wr \mathfrak{S}_{n-2l}$ -module. As an R -module, S_l is generated by elements of the form $\sigma \otimes e_l d$, where $d \in D_n(A)$, and $\sigma \in A \wr \mathfrak{S}_{n-2l}$. If d has more than l horizontal edges, say m , then we have

$$\begin{aligned} \sigma \otimes e_l d &= \sigma \otimes e_m d' \\ &= \sigma e_m \otimes d' \\ &= 0 \otimes d' = 0, \end{aligned}$$

where $d' \in D_n(A)$ contains the bottom configuration of d . The last equality holds because the idempotent e_m is in $\ker \alpha$. In particular, $A \wr \mathfrak{S}_{n-2l} \otimes_{e_l D_n(A) e_l} J_{l+1} e_l$ vanishes. Consequently, all the generators of S_l are of the form $\sigma \otimes d$, where d has precisely l horizontal edges containing the partial diagram of e_l at its top, and l horizontal edges at the bottom can be arranged arbitrarily. That is, the partial diagram of its bottom configuration lies in $V(n, l, A)$. Therefore, the generators must be of the form $1_{A \wr \mathfrak{S}_{n-2l}} \otimes \sigma d$. Hence, $S_l = (A \wr \mathfrak{S}_{n-2l})^{\oplus n_l}$ where n_l is the rank of $V(n, l, A)$. Thus, S_l is a free $A \wr \mathfrak{S}_{n-2l}$ - $D_n(A)$ -bimodule. Thus, S_l is projective as a left $A \wr \mathfrak{S}_{n-2l}$ -module.

On the other hand, the right action of $e_l D_n(A) e_l$ on $A \wr \mathfrak{S}_{n-2l}$ coincides with the action of $e_l(D_n(A)/J_{l+1})e_l$ on $A \wr \mathfrak{S}_{n-2l}$. Hence, we have

$$\begin{aligned} (3.1) \quad S_l &= A \wr \mathfrak{S}_{n-2l} \otimes_{e_l D_n(A) e_l} e_l D_n(A) \\ &\cong A \wr \mathfrak{S}_{n-2l} \otimes_{e_l D_n(A) e_l} e_l(D_n(A)/J_{l+1}) \\ &\cong A \wr \mathfrak{S}_{n-2l} \otimes_{e_l(D_n(A)/J_{l+1})e_l} e_l(D_n(A)/J_{l+1}) \\ &\cong A \wr \mathfrak{S}_{n-2l} \otimes_{A \wr \mathfrak{S}_{n-2l}} e_l(D_n(A)/J_{l+1}) \\ &\cong e_l(D_n(A)/J_{l+1}), \end{aligned}$$

where (3.1) follows the fact that $e_l(D_n(A)/J_{l+1})e_l$ has exactly l horizontal edges so that it will be isomorphic to $A \wr \mathfrak{S}_{n-2l}$. One can easily obtain the following right $A \wr \mathfrak{S}_{n-2l}$ -module isomorphism:

$$\begin{aligned} S_l e_l &\cong (e_l(D_n(A)/J_{l+1}))e_l \\ &\cong A \wr \mathfrak{S}_{n-2l}. \end{aligned}$$

This ends the proof. \square

Remark 3.11. Consider the categories $\mathbf{mod}\text{-}A \wr \mathfrak{S}_{n-2l}$, and $\mathbf{mod}\text{-}D_n(A)$. Using the bimodule structure of S_l , we define the layer induction functor $\text{ind}_l(M) = M \otimes_{e_l D_n(A) e_l} S_l$, where $M \in \mathbf{mod}\text{-}A \wr \mathfrak{S}_{n-2l}$ and $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$. One can define the restriction functor by $\text{Res}_l(N) = N \otimes_{D_n(A)} D_n(A) e_l$ for all $N \in \mathbf{mod}\text{-}D_n(A)$ and Res_l keeps all the layers.

Corollary 3.12. *For $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$, the pair $(\text{ind}_l, \text{Res}_l)$ forms an exact split pair between $\mathbf{mod}\text{-}A \wr \mathfrak{S}_{n-2l}$ and $\mathbf{mod}\text{-}D_n(A)$.*

Proof. If $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$, then $A \wr \mathfrak{S}_{n-2l}$ is a corner split quotient of $D_n(A)$ by Theorem 3.10. By applying Theorem 2.1, $(\text{ind}_l, \text{Res}_l)$ form an exact split pair of functors between $\mathbf{mod}\text{-}D_n(A)$ and $\mathbf{mod}\text{-}A \wr \mathfrak{S}_{n-2l}$. \square

Note that in the case of $\delta = 0$ and n is odd, the statements in Proposition 3.8, Proposition 3.9 and Corollary 3.12 are still valid.

4. THE CYCLOTOMIC BRAUER ALGEBRA

For fixed natural numbers r and n , let R be a commutative ring containing a primitive r -th root of unity and parameters $\delta_0, \delta_1, \dots, \delta_{r-1}$. Let $\boldsymbol{\delta} = (\delta_0, \dots, \delta_{r-1}) \in R^r$ and A be the group algebra of the cyclic group $\mathbb{Z}/r\mathbb{Z}$. The associated A -Brauer algebra is known as the cyclotomic Brauer algebra $\mathcal{B}_n^r(\boldsymbol{\delta})$. These algebras were studied in [3, 19, 18]. Similar to the A -Brauer diagrams, all the edges are equipped with an orientation. Reversing the orientation results in relabeling the edges using the inverse element in $\mathbb{Z}/r\mathbb{Z}$. It is generated by s_i, e_i and h_j^m subject to the relations [[19], Section 2], for $1 \leq i \leq n-1, 1 \leq j \leq n$ and $m \in \mathbb{Z}/r\mathbb{Z}$.

Let $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$. If $\boldsymbol{\delta}$ is invertible, then we consider the idempotent as in Figure 2. If $\boldsymbol{\delta} = 0$ and n is odd, then the idempotent is given as in Figure 3.

4.1. The Generalized symmetric group. Let \mathcal{H}_n^r be the group algebra of the generalized symmetric group $\mathbb{Z}/r\mathbb{Z} \wr \mathfrak{S}_n$. It is a subalgebra of $\mathcal{B}_n^r(\boldsymbol{\delta})$ generated by s_i , and h_j^m for $1 \leq i \leq n-1, 1 \leq j \leq n$, and $m \in \mathbb{Z}/r\mathbb{Z}$. The group algebra \mathcal{H}_n^r is cyclic cellular with cell datum $(\mathcal{H}_n^r, *, \Lambda_n^r, \triangleright, \mathcal{M}, \mathcal{B})$, see [[8], Theorem 4.1] where Λ_n^r is the set of all multipartitions of n with r parts which is a poset under the dominance order for the multipartitions.

From now on, assume R to be a commutative Noetherian domain with an invertible $\boldsymbol{\delta}$. Let V_l be the free R -module generated by all (n, l) - $\mathbb{Z}/r\mathbb{Z}$ labeled partial diagrams. The algebra $\mathcal{B}_n^r(\boldsymbol{\delta})$ can be decomposed as an iterated inflation of \mathcal{H}_{n-2l}^r along V_l ,

$$\mathcal{B}_n^r(\boldsymbol{\delta}) \cong \bigoplus_{l=0}^{\lfloor \frac{n}{2} \rfloor} V_l \otimes V_l \otimes \mathcal{H}_{n-2l}^r,$$

where \mathcal{H}_{n-2l}^r is a cellular algebra, see [[3], Theorem 3.1.3]. The involution is given by reflecting the diagram along its horizontal axis.

Define the index set Λ by $\Lambda := \{(l, \boldsymbol{\lambda}) : 0 \leq l \leq \lfloor \frac{n}{2} \rfloor, \boldsymbol{\lambda} \in \Lambda_n^r\}$. We extend the dominance order of \mathcal{H}_n^r to $\mathcal{B}_n^r(\boldsymbol{\delta})$ to get the dominance order “ \geq ” on Λ , which is defined as follows:

$$(l, \boldsymbol{\lambda}) \geq (m, \boldsymbol{\mu}) \text{ if either } l < m, \text{ or } l = m \text{ and } \boldsymbol{\mu} \triangleright \boldsymbol{\lambda}.$$

The cell modules for $\mathcal{B}_n^r(\boldsymbol{\delta})$ are indexed by $(l, \boldsymbol{\lambda}) \in \Lambda$ by [[18], Theorem 5.11].

4.2. Exact split pair $(\mathcal{B}_n^r(\boldsymbol{\delta}), \mathcal{H}_n^r)$. In this subsection, we show that there exists an exact split pair of functors between $\mathbf{mod}\text{-}\mathcal{B}_n^r(\boldsymbol{\delta})$ and $\mathbf{mod}\text{-}\mathcal{H}_n^r$. As an application of this, we obtain the global dimension of $\mathcal{B}_n^r(\boldsymbol{\delta})$.

Let us define the induction and restriction functor as follows:

$$\begin{aligned} \text{ind}_l : \mathbf{mod}\text{-}\mathcal{H}_{n-2l}^r &\longrightarrow \mathbf{mod}\text{-}\mathcal{B}_n^r(\boldsymbol{\delta}) & \text{Res}_l : \mathbf{mod}\text{-}\mathcal{B}_n^r(\boldsymbol{\delta}) &\longrightarrow \mathbf{mod}\text{-}\mathcal{H}_{n-2l}^r \\ M &\longmapsto M \otimes_{e_l \mathcal{B}_n^r(\boldsymbol{\delta}) e_l} e_l \mathcal{B}_n^r(\boldsymbol{\delta}) & N &\longmapsto N \otimes_{\mathcal{B}_n^r(\boldsymbol{\delta})} \mathcal{B}_n^r(\boldsymbol{\delta}) e_l. \end{aligned}$$

Corollary 4.1. *The pair $(\text{ind}_l, \text{Res}_l)$ forms an exact split pair of functors between $\mathbf{mod}\text{-}\mathcal{B}_n^r(\boldsymbol{\delta})$ and $\mathbf{mod}\text{-}\mathcal{H}_{n-2l}^r$, where $0 \leq l \leq \lfloor \frac{n}{2} \rfloor$.*

Proof. By Theorem 3.10, \mathcal{H}_{n-2l}^r is a corner split quotient of $\mathcal{B}_n^r(\delta)$. Therefore, by Theorem 2.1, the pair of functors $(\text{ind}_l, \text{Res}_l)$ form an exact split pair. \square

The global dimension of an algebra A , denoted by $\text{gl.dim } A$, is the maximum of the projective dimensions of the A -modules. The computation of the global dimension of $\mathcal{B}_n^r(\delta)$ is challenging due to its non-self-symmetric nature, the existence of an exact split pair simplifies the calculation.

Proposition 4.2. *If R is a field of characteristic p , then $\mathcal{B}_n^r(\delta)$ has finite global dimension if and only if $p > n$ and $p \nmid r$, or $p = 0$.*

Proof. Assume that $\mathcal{B}_n^r(\delta)$ has finite global dimension. It follows that \mathcal{H}_n^r also has finite global dimension because $\text{gl.dim } \mathcal{B}_n^r(\delta) \geq \text{gl.dim } \mathcal{H}_n^r$ by [[7], Corollary 1.6] and Corollary 4.1. This implies that $p > n$ and $p \nmid r$, or $p = 0$. Conversely, if $p > n$ and $p \nmid r$ or $p = 0$, then $\mathcal{B}_n^r(\delta)$ is quasi-hereditary by [[3], Theorem 3.3.1], and hence has finite global dimension by [[16], Theorem 1.1]. \square

Note that in the case of $\delta = 0$ and n is odd, the statements in Corollary 4.1 and Proposition 4.2 are still valid.

Remark 4.3. The Brauer algebra of type C , denoted by $\mathcal{B}(C_n, \delta)$, is an associative R -algebra over a field R with δ a non-zero element in R . It was first introduced in [5], and can be written as an iterated inflation of group algebras of hyper-octahedral groups, for more details see [2]. The hyper-octahedral group is a special case of the generalized symmetric group when we take $r = 2$. The diagram in $\mathcal{B}(C_n, \delta)$ can be viewed as “unfolding” the diagram vertically along the wall. Similarly one can prove that $R[\mathfrak{S}_2 \wr \mathfrak{S}_{n-l}]$ is a corner split quotient of $\mathcal{B}(C_n, \delta)$, which gives an exact split pair $(\text{ind}_l, \text{Res}_l)$ of functors between the categories $\mathbf{mod}\text{-}R[\mathfrak{S}_2 \wr \mathfrak{S}_{n-l}]$ and $\mathbf{mod}\text{-}\mathcal{B}(C_n, \delta)$, where $0 \leq l \leq n$. Note that $\mathcal{B}(C_n, \delta)$ also has finite global dimension if and only if $\delta \neq 0$ and $p > n$.

5. THE WALLED BRAUER ALGEBRA

Let R be a commutative ring with 1, and δ a distinguished element in R . Fix two positive integers r and t . The walled Brauer algebra $\mathcal{B}_{r,t}(\delta)$ is a subalgebra of the ordinary Brauer algebra $\mathcal{B}_{r+t}(\delta)$, and if either r or t is 0, then $\mathcal{B}_{r,t}(\delta)$ coincides with the ordinary Brauer algebra. An (r, t) -walled Brauer diagram is an ordinary Brauer diagram with $r + t$ vertices arranged in two rows, separated by a wall between the r th and $(r + 1)$ th vertices. All horizontal edges cross the wall, but vertical edges do not. The walled Brauer algebra $\mathcal{B}_{r,t}(\delta)$ is an associative R -algebra spanned by all (r, t) -walled Brauer diagrams, with multiplication similar to that of ordinary Brauer algebras. It contains the group algebra of the direct product of two symmetric groups \mathfrak{S}_r and \mathfrak{S}_t as a subalgebra and is denoted by $R\mathfrak{S}_{r,t}$. The generators are s_i , s_{r+j} , and $e_{k,l}$, where $1 \leq i \leq r - 1$, $1 \leq j \leq t - 1$, and $1 \leq k \leq r$, $r + 1 \leq l \leq r + t$, see Figure 4.






FIGURE 4. Generators of the walled Brauer algebra.

Let $s = \min\{r, t\}$ and $0 \leq l \leq s$. When δ is invertible in R , the idempotent is defined as shown in Figure 5. However, when $\delta = 0$ and either r or t is at least 2, the idempotent is defined as in Figure 6.


 FIGURE 5. Idempotent of $\mathcal{B}_{r,t}(\delta)$ for invertible δ .

 FIGURE 6. Idempotent of $\mathcal{B}_{r,t}(\delta)$ when $\delta = 0$, and one of r or $t \geq 2$.

5.1. Cellular structure. Let $s := \min\{r, t\}$, and $0 \leq l \leq s$. Assume that R is a commutative Noetherian domain with invertible δ . Let V_l be the free R -module generated by all (r, t) -partial diagrams with l horizontal edges, $r - l$ free vertices on the left side, and $t - l$ free vertices on the right side of the wall. The involution i is defined by flipping the diagram along the horizontal axis. By [[6], Proposition 2.6], the algebra $\mathcal{B}_{r,t}(\delta)$ can be expressed as an iterated inflation of $R\mathfrak{S}_{r-l,t-l}$ along V_l :

$$\mathcal{B}_{r,t}(\delta) = \bigoplus_{l=0}^s V_l \otimes V_l \otimes R\mathfrak{S}_{r-l,t-l}.$$

The group algebra $R\mathfrak{S}_{r-l,t-l}$ being a direct product of two cellular algebras is cellular by [[11], Example 3.1.6]. Therefore, by [[17], Proposition 3.5], $\mathcal{B}_{r,t}(\delta)$ is also cellular.

5.2. Exact split pair $(\mathcal{B}_{r,t}(\delta), R\mathfrak{S}_{r,t})$. We now show the existence of an exact split pair for $\mathcal{B}_{r,t}(\delta)$. This will allow us to determine the criteria for the non-vanishing of Hom and Ext spaces between the cell modules of $\mathcal{B}_{r,t}(\delta)$.

Let $0 \leq l \leq s$. Define the induction functor ind_l , and restriction functor Res_l as follows: $\text{ind}_l(M) = M \otimes_{e_l \mathcal{B}_{r,t}(\delta) e_l} e_l \mathcal{B}_{r,t}(\delta)$ for $M \in \mathbf{mod}\text{-}R\mathfrak{S}_{r-l,t-l}$ and $\text{Res}_l(N) = N \otimes_{\mathcal{B}_{r,t}(\delta)} \mathcal{B}_{r,t}(\delta) e_l$ for $N \in \mathbf{mod}\text{-}\mathcal{B}_{r,t}(\delta)$.

Theorem 5.1. *If δ is invertible, then for each $l \in \{1, \dots, s\}$, there exists an exact split pair of functors $(\text{ind}_l, \text{Res}_l)$ between the categories $\mathbf{mod}\text{-}R\mathfrak{S}_{r-l,t-l}$, and $\mathbf{mod}\text{-}\mathcal{B}_{r,t}(\delta)$.*

Proof. The proof uses a similar argument as in Corollary 4.1. □

Next, we will show some applications of these exact split pairs. The poset Λ is the set of all tuples $(l, (\lambda, \mu))$ where $0 \leq l \leq s$ and λ, μ are partitions of $r - l$ and $t - l$, respectively. Let $(l, (\lambda, \mu)), (m, (\lambda', \mu')) \in \Lambda$, the partial order " \leq " on Λ is defined as follows:

$$(l, (\lambda, \mu)) \leq (m, (\lambda', \mu')) \text{ if } m < l, \text{ or } m = l \text{ and } \lambda \supseteq \lambda', \mu \supseteq \mu'.$$

Therefore, by using the induction functor, we write the cell modules of $\mathcal{B}_{r,t}(\delta)$ as $\text{ind}_l S^{\lambda, \mu}$ where $(l, (\lambda, \mu)) \in \Lambda$ and $S^{\lambda, \mu}$ is the Specht module of $R\mathfrak{S}_{r-l,t-l}$. From now on, we assume R to be a field.

Corollary 5.2. *If $\text{char } R \neq 2$, then $\text{Hom}_{\mathcal{B}_{r,t}(\delta)}(\text{ind}_l S^{\lambda, \mu}, \text{ind}_l S^{\lambda', \mu'}) \neq 0$ implies that $\lambda \triangleright \lambda'$, and $\mu \triangleright \mu'$.*

Proof. We have the following isomorphism using [[7], Proposition 1.3], and Theorem 5.1

$$\begin{aligned} \text{Hom}_{\mathcal{B}_{r,t}(\delta)}(\text{ind}_l S^{\lambda, \mu}, \text{ind}_l S^{\lambda', \mu'}) &\cong \text{Hom}_{R\mathfrak{S}_{r-l,t-l}}(S^{\lambda, \mu}, S^{\lambda', \mu'}) \\ &\cong \text{Hom}_{R\mathfrak{S}_{r-l}}(S^\lambda, S^{\lambda'}) \boxtimes_R \text{Hom}_{R\mathfrak{S}_{t-l}}(S^\mu, S^{\mu'}), \end{aligned}$$

and the last isomorphism follows from [[21], Lemma 3.2]. If $\lambda \not\triangleright \lambda'$ or $\mu \not\triangleright \mu'$, then the last isomorphism is zero by [[14], Corollary 13.17], provided $\text{char } R \neq 2$. Hence, $\text{Hom}_{\mathcal{B}_{r,t}(\delta)}(\text{ind}_l S^{\lambda,\mu}, \text{ind}_l S^{\lambda',\mu'}) \neq 0$ implies $\lambda \triangleright \lambda'$ and $\mu \triangleright \mu'$. \square

Corollary 5.3. *If $\text{char } R \neq 2, 3$, then $\text{Ext}_{\mathcal{B}_{r,t}(\delta)}^1(\text{ind}_l S^{\lambda,\mu}, \text{ind}_l S^{\lambda',\mu'}) \neq 0$ implies $\lambda \triangleright \lambda'$ and $\mu \triangleright \mu'$.*

Proof. We have the following isomorphism from [[7], Proposition 1.6], and Theorem 5.1

$$\begin{aligned} \text{Ext}_{\mathcal{B}_{r,t}(\delta)}^1(\text{ind}_l S^{\lambda,\mu}, \text{ind}_l S^{\lambda',\mu'}) &\cong \text{Ext}_{R\mathfrak{S}_{r-l,t-l}}^1(S^{\lambda,\mu}, S^{\lambda',\mu'}) \\ &\cong (\text{Ext}_{R\mathfrak{S}_{r-l}}^1(S^\lambda, S^{\lambda'}) \boxtimes_R \text{Hom}_{R\mathfrak{S}_{t-l}}(S^\mu, S^{\mu'})) \bigoplus (\text{Hom}_{R\mathfrak{S}_{r-l}}(S^\lambda, S^{\lambda'}) \boxtimes_R \text{Ext}_{R\mathfrak{S}_{t-l}}^1(S^\mu, S^{\mu'})), \end{aligned}$$

and the last isomorphism follows from [[4], Chapter XI, Theorem 3.1]. This first direct summand is zero whenever $\lambda \not\triangleright \lambda'$ or $\mu \not\triangleright \mu'$ by [[13], Theorem 4.2.1] and [[14], Corollary 13.17], provided $\text{char } R \neq 2, 3$. Similarly, if $\text{char } R \neq 2, 3$ the other direct summand is zero whenever $\lambda \not\triangleright \lambda'$ or $\mu \not\triangleright \mu'$. Hence, the proof follows. \square

Remark 5.4. We can determine the criteria for non-vanishing cohomology in the case of $\mathcal{B}_{r,t}(\delta)$ because the existence of an exact split pair ensures that the Ext between two cell modules of $\mathcal{B}_{r,t}(\delta)$ is isomorphic to the Ext between two cell modules of $R\mathfrak{S}_{r-l,t-l}$. While a similar isomorphism holds for other algebras, the criteria for non-vanishing cohomology for the case of wreath product algebras $A \wr \mathfrak{S}_n$ remain unknown, which is needed to determine the non-vanishing cohomology for $D_n(A)$, and $\mathcal{B}_n^r(\delta)$.

Let $\text{char } R = p$. If $p \leq r$ (resp. $p \leq t$) then $R\mathfrak{S}_r$ (resp. $R\mathfrak{S}_t$) has infinite global dimension. This implies that $R\mathfrak{S}_{r,t}$ has infinite global dimension if $p \leq \min\{r, t\}$.

Corollary 5.5. *$\mathcal{B}_{r,t}(\delta)$ has finite global dimension if and only if $p > \max\{r, t\}$, or $p = 0$.*

Proof. If the R -algebra $\mathcal{B}_{r,t}(\delta)$ has finite global dimension, then $R\mathfrak{S}_{r,t}$ has finite global dimension by [[7], Corollary 1.6], and Theorem 5.1. This gives $p > \max\{r, t\}$, or $p = 0$. Conversely, if the given conditions hold, then $\mathcal{B}_{r,t}(\delta)$ is quasi-hereditary by [[6], Corollary 2.8], which ensures $\mathcal{B}_{r,t}(\delta)$ has finite global dimension by [[16], Theorem 1.1]. \square

The statements in Theorem 5.1, Corollary 5.2 and Corollary 5.3 holds true for the case $\delta = 0$, and one of r and t greater than or equal to 2. The Corollary 5.5 is still valid if $\delta = 0$ with one of r and t greater than or equal to 2, and $r \neq t$.

Acknowledgment. The authors express their gratitude to Prof. Steffen Koenig for his time and valuable discussions. We would like to sincerely thank the referees for their thoughtful and constructive comments, which have significantly improved the quality of this paper. The first author's research is supported by Indian Institute of Science Education and Research Thiruvananthapuram PhD fellowship. The second author's research was partially supported by IISER-Thiruvananthapuram, SERB-Power Grant SPG/2021/004200 and Prof. Steffen Koenig's research grant. The second author also would like to acknowledge the Alexander Von Humboldt Foundation for their support.

REFERENCES

- [1] Benkart, G., Chakrabarti, M., Halverson, T., Leduc, R., Lee, C., Stroome, J., Tensor product representations of general linear groups and their connections with Brauer algebras *J. Algebra* **166(3)** (1994) 596–567.
- [2] Bowman, C., Brauer algebra of type C are cellularly stratified, *Math. Proc. Cambridge Philos. Soc.* **153(1)** (2012) 1–7.
- [3] Bowman, C., Cox, A.G., De, Visscher, M., Decomposition number for the cyclotomic Brauer algebras in characteristic zero, *J. Algebra* **378** (2013) 80–102.
- [4] Cartan, H., Eilenberg, S. Homological algebra (PMS-19), Volume 19. Princeton University Press, Princeton (1956).
- [5] Cohen, A.M., Liu, S., Yu, S. Brauer algebras of type C, *J. Pure Appl. Algebra* **216(2)** (2012) 407–426.
- [6] Cox, A., De Visscher, M., Doty, S., Martin, P., On blocks of the walled Brauer algebra, *J. Algebra* **320(1)** (2008) 169–212.
- [7] Diracca, L., Koenig, S., Cohomological reduction by split pairs, *J. Pure Appl. Algebra*, **212** (2008) 471–485.
- [8] Geetha, T., Goodman, F. M., Cellularity of the Wreath products and A-Brauer algebras, *J. Algebra* **389** (2013) 151–190.

- [9] Graham, J.J., Lehrer, G.I., Cellular algebras, *Invent. Math.* **123(1)** (1996) 1–34.
- [10] Green, J.A., Polynomial Representations of GL_n , *Lecture Notes in Mathematics* **Vol. 830** Springer-Verlag, Berlin Heidelberg, 1980. [3]
- [11] Green, R.M., Martin, P.P., Constructing cell data for diagram algebras, *J. Pure Appl. Algebra* **209(2)** (2007) 551–569.
- [12] Hartmann, R., Henke, A., Koenig, S., Paget, R., Cohomological stratification of diagram algebras, *Math. Ann.* **347(4)** (2010) 765–804.
- [13] Hemmer, D.J., Nakano, D.K. Specht filtrations for Hecke algebras of type A, *J. London Math. Soc.(2)* **69(3)** (2004) 623–638 .
- [14] James, G.D., The Representation Theory of the Symmetric Groups, *Lecture Notes in Mathematics* **682** (1978) 156 Springer.
- [15] Koenig, S., Xi, C, On the structure of cellular algebras, *Amer. Math. Soc., Providence, RI* **24** (Algebras and modules, II (Geiranger, 1996)) 365–386.
- [16] Koenig, S., Xi, C., When is a cellular algebra quasi-hereditary?, *Math. Ann.* **315(2)** (1999) 281–293.
- [17] Koenig, S., Xi, C., Cellular algebras: Inflation and Morita equivalences, *J. London Math. Soc.* **60(3)** (1999) 700–722.
- [18] Rui, H., Yu, W., On the semisimplicity of the cyclotomic Brauer algebras *J. Algebra*, **277(1)** (2004) 187–221.
- [19] Rui, H., Xu, J., On the semisimplicity of the cyclotomic Brauer algebras, II, *J. Algebra* **312(2)** (2007) 995–1010.
- [20] Schur, I., Über die rationalen darstellungen der allgemeinen linearen gruppe. *Sitzungsberichte der Preussischen Akademie der Wissenschaften zu Berlin*, pages 58–75, 1927.
- [21] Xi, C., On the representation dimension of finite dimensional algebras, *J. Algebra* **226(1)** (2000) 332–346.

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH THIRUVANANTHAPURAM, THIRUVANANTHAPURAM,
EMAIL: SULAKHANA17@IISERTVM.AC.IN

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH THIRUVANANTHAPURAM, THIRUVANANTHAPURAM,
EMAIL: TGEETHA@IISERTVM.AC.IN