Some consequences of Frobenius descent for torsors

Niels Borne^{*} Mohamed Rafik Mammeri[†]

February 20, 2025

Abstract

We show how the formalism of Frobenius descent for torsors enables to study torsors under Frobenius kernels in terms of non-commutative, Lie-valued differential forms. We pay particular attention to affine line bundles trivialized by the Frobenius.

1 Introduction

Let k be a field of positive characteristic p, that in this introduction we will assume perfect. Let X/k be a smooth scheme, and denote by $F_X : X \to X$ the Frobenius morphism.

1.1 The prototype

It is well known that the set of isomorphism classes of μ_p -torsors on X for the fppf topology is in one to one correspondence with the set of global 1-differentials forms on X that are closed and locally logarithmic.

It is fairly easy to give a rough idea of the correspondence: given such a form ω , verifying locally $\omega = df/f$, where f is a unit, one defines a μ_p -torsor locally by the Kummer equation $x^p = f$. Since two such f differ by the p-th power of a *unique* unit, the global form ω enables to glue the local torsors.

The actual proof relies on a delicate comparison of cohomologies on the étale and on the flat site, see [Mil80, Corollary 4.14].

There is a parallel characterization of the set of isomorphism classes of α_p torsors on X in terms of 1-differentials forms on X that are closed and locally exact. Moreover, these two cases have been generalized by Artin and Milne (see [AM76]) who give a description of $H^1(X, A)$ in terms of suitable Lie-valued differential forms if A/k is an arbitrary abelian group scheme of height 1.

^{*}Niels.Borne@univ-lille.fr

 $^{^{\}dagger}$ mammeri.mohamed@gmail.com

1.2 Frobenius descent enters the scene

Our main point is that one can avoid the use of cohomology altogether and reinterpret the correspondence between μ_p -torsors and closed and locally logarithmic differential forms in terms of descent along the Frobenius. Namely, Cartier's Frobenius descent for vector bundles ensures that if \mathcal{E} is a locally free sheaf on X, then $F_X^*\mathcal{E}$ is endowed with a canonical connection ∇_{can} , whose moreover both curvature C and p-curvature ψ vanish. And conversely, it is possible, in an essentially unique way, to descend such a vector bundle equipped with a suitable connection along $F_X: X \to X$.

If now ω is a closed 1-form on X, and \mathscr{C} is the Cartier operator, the condition $\mathscr{C} \omega = \omega$ not only characterizes the fact that ω is locally logarithmic, but also the fact that the connection $d + \omega$ on the structure sheaf \mathcal{O}_X is of vanishing p-curvature. By Frobenius descent, $(\mathcal{O}_X, d + \omega)$ descends to an invertible sheaf \mathcal{L} such that $F_X^* \mathcal{L} \simeq \mathcal{O}_X$. Since there is a natural isomorphism $F_X^* \mathcal{L} \simeq \mathcal{L}^{\otimes p}$, this gives rise to a μ_p -torsor on X, as expected.

As Frobenius descent has been generalized from $G = \operatorname{GL}_n$ to *G*-torsors, where G/k is a smooth affine group scheme (see [CZ15; Mam16] and Theorem 3.2) the merits of this geometric approach are clear.

1.3 Main results

To illustrate the usefulness of Frobenius descent for *G*-torsors, we work out completely the simplest non-commutative case, that is when the structure group is $G = \mathbb{G}_a \ltimes \mathbb{G}_m$:

Corollary A (Corollary 4.8). Assume that X/k is smooth, and that k is perfect. Then the set of isomorphism classes of affine line bundles on X trivialized along the Frobenius is in one to one correspondence with the set of pairs (ω, ω') of 1-forms on X such that :

- 1. $d\omega = 0 = d\omega' + \omega \wedge \omega'$,
- 2. $\mathscr{C}\omega = \omega$,
- 3. for each Zariski open $U \subset X$ and each $f \in \mathbb{G}_m(U)$ such that $\omega_{|U} = df/f$, the equality $\mathscr{C}(f\omega'_{|U}) = 0$ holds.

The first condition illustrates a new feature of the non-commutative situation, the appearance of non-closed forms.

Moreover, one gets a complete description of torsors under Frobenius kernels $G^F = \ker(F_G)$ in terms of suitable non-commutative Lie-valued differential forms:

Theorem B (Theorem 5.4). Assume that X/k is smooth, and k is perfect. There is a natural one to one map $\mathrm{H}^1(X, G^F) \to \mathrm{H}^0(X, \Omega^1_{X/k} \otimes_k \mathrm{Lie}(G))^{C=\psi=0}$ that for each $g \in G(X)$ sends the image $\partial(g)$ of g via the cobord map $\partial : G(X) \to \mathrm{H}^1(X, G^F)$ to $\mathrm{dlog}(g)$, commutes with étale base change, and is characterized by these properties.

1.4 Some context

The second result (Theorem B) appeared initially in the second author's phd ([Mam16]). The exposition of its proof has been vastly improved.

The first result (Corollary A) also appeared in [Mam16], although in a more hidden form. The exposition chosen here is motivated by the recent surge of interest in affine bundles (see [Ple19a; Ple19b; Bou24]). Affine line bundles are in fact a venerable subject, initiated by their classification on Riemann surfaces by André Weil (see [Wei55]).

Our study leads to the following conclusions. In the case where the structure group G is abelian, Theorem B completely solves the issue of classifying G-torsors trivialized by the Frobenius, because in this situation there exists an explicit formula for the *p*-curvature (see Appendix C). However, Corollary A shows that even for the simplest non-abelian structure group, this classification is much more difficult. The case of a general non-abelian structure group remains mysterious.

1.5 Structure of the text

In §2 we recall Atiyah's definition of a connection on a G-torsor, when G/k is a smooth affine group scheme, and use it to define the Maurer-Cartan map d log. We also recall the definition of the associated curvatures.

In the next paragraph, \$3, we give an account of Frobenius descent for G-torsors, and show how it enables to describe the local structure of connections. We pay particular attention to the case of special groups.

Paragraph 4 is devoted to the study of affine line bundles. We apply the machinery of Frobenius descent to prove Corollary A.

Finally, in §5 we gather the previous results and prove Theorem B.

1.6 Notations and conventions

We work over an arbitrary field k of positive characteristic p and denote by $S = \operatorname{spec} k$. Throughout X is a smooth k-scheme. The tangent sheaf of X is denoted by \mathcal{T}_X . By definition, this is the sheaf $\operatorname{Der}_k(\mathcal{O}_X)$ of derivations of \mathcal{O}_X with respect to k.

We use G for an affine k-group scheme. We consider mainly the fppf topology on X, and G-torsors $p: P \to X$ for this topology. We stick to EGA's convention and consider *right* G-torsors. We denote as usual by BG(X) the category of G-torsors on X. We sometimes (especially when G is smooth) use the étale topology on X; we then stick to the small étale site.

If \mathcal{F} is a quasi-coherent *G*-sheaf on *P*, we denote as usual by $p_*^G \mathcal{F}$ the quasicoherent sheaf on *X* obtained by pushing forward to *X* and then taking fixed sections under the action of *G*. The adjoint action of *G* on Lie(*G*) give rises to a *G*-sheaf $\mathcal{O}_P \otimes_k \text{Lie}(G)$ on *P*, and we denote by $\operatorname{ad}(P) = p_*^G(\mathcal{O}_P \otimes_k \text{Lie}(G))$ the associated locally free sheaf on *X* (so $\operatorname{ad}(P)$ is the sheaf of sections of the adjoint vector bundle). Finally, we denote by $F_{X/S}: X \to X^{(p)}$ the relative Frobenius of X (see Appendix A for details).

1.7 Acknowledgements

The first author wants to thank Cédric Pépin and Angelo Vistoli for many stimulating conversations. This work was supported in part by the Labex CEMPI (ANR-11-LABX-0007-01).

2 Atiyah's exact sequence

Throughout this section, G will denote a smooth affine group scheme over k.

2.1 Invariant vector fields on a torsor

Atiyah's definition of a connection on a *G*-torsor relies on the following elementary fact:

Lemma 2.1. Let G/k be a smooth affine group scheme and $p: P \to X$ a *G*-torsor. Then there is a natural exact sequence :

$$0 \to \mathrm{ad}(P) \to p^G_*(\mathcal{T}_P) \to \mathcal{T}_X \to 0 \tag{1}$$

Proof. [Ati57, Theorem 1].

Definition 2.2 ([Ati57]). An Atiyah-connection on a *G*-torsor $p : P \to X$ is an \mathcal{O}_X -linear splitting $s : \mathcal{T}_X \to p^G_*(\mathcal{T}_P)$ of the exact sequence (1).

We will use the notation At(G, X) for the category of G-torsors on X endowed with a connection.

When $G = \operatorname{GL}_n$, one recovers the familiar notion of Koszul-connection, that is, a locally free sheaf \mathcal{E} endowed with a k-linear map $\nabla : \mathcal{E} \to \mathcal{E} \otimes_{\mathcal{O}_X} \Omega^1_{X/k}$, satisfying Leibniz rule.

It is clear that any section $\sigma : X \to P$ of a *G*-torsor $p : P \to X$ induces an Atiyah-connection $s_{\sigma} = p_*^G(\mathcal{T}_{\sigma}) : \mathcal{T}_X \to p_*^G(\mathcal{T}_P)$. In particular, the unit section $e : X \to X \times G$ of the trivial torsor $\operatorname{pr}_1 : X \times G \to X$ induces an Atiyah-connection s_e on $X \times G$.

Definition 2.3 (Maurer-Cartan map). The Maurer-Cartan map

$$d \log : G(X) \to H^0(X, \Omega^1_{X/k} \otimes_k \operatorname{Lie}(G))$$

denoted by $g \mapsto \Omega_g$, is the map given by, for all $g \in G(X)$: $\Omega_g = s_{g^{-1}} - s_e$.

When $G = GL_n$, there is an explicit and well-known formula for the Maurer-Cartan map.

Lemma 2.4. The map $d\log : \operatorname{GL}_n(X) \to \operatorname{H}^0(X, \operatorname{Lie}(\operatorname{GL}_n) \otimes_k \Omega^1_{X/k})$ is given by: $d\log(g) = g^{-1}dg$. Proof. Given $g \in G$, the element $d\log(g) = \Omega_g$ of $\mathrm{H}^0(X, \mathrm{Lie}(\mathrm{GL}_n) \otimes_k \Omega^1_{X/k})$ is characterized by the fact that left multiplication $L_g : (X \times G, s_e + \Omega_g) \to (X \times G, s_e)$ is an isomorphism of torsors endowed with Atiyah-connections. This translates into the fact that $g : (\mathcal{O}_X^{\otimes n}, \mathrm{d} + \Omega_g) \to (\mathcal{O}_X^{\otimes n}, \mathrm{d})$ is a morphism of free sheaves endowed with Koszul-connections, which is equivalent to the displayed formula for $d\log(g) = \Omega_g$.

By functoriality in G, we get the same formula for any smooth closed subgroup of GL_n . For instance, by using the standard embedding $\mathbb{G}_a \to \operatorname{GL}_2$, $a \mapsto \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$, we see that the Maurer-Cartan map for \mathbb{G}_a is given by $a \mapsto da$.

For later use, we record the following lemma describing the image of d log.

Lemma 2.5. Let $\Omega \in \mathrm{H}^0(X, \Omega^1_{X/k} \otimes_k \mathrm{Lie}(G))$. Then there exists $g \in G(X)$ such that $\Omega = \Omega_g$ if and only if $(X \times G, s_e + \Omega) \simeq (X \times G, s_e)$.

Proof. This is clear, since both assertions are equivalent to : there exists $g \in G(X)$ such that $s_e + \Omega = s_q$.

2.2 Curvatures

2.2.1 Usual curvature

Definition 2.6. Let $s : \mathcal{T}_X \to p^G_*(\mathcal{T}_P)$ be an Atiyah-connection on a *G*-torsor $p : P \to X$. Its curvature is the morphism $C_s : \mathcal{T}_X \times \mathcal{T}_X \to \mathrm{ad}(P)$ given for all D, D' in \mathcal{O}_X by :

$$C_s(D, D') = [s(D), s(D')] - s([D, D'])$$
.

The curvature C_s of an Atiyah-connection is a skew-symmetric \mathcal{O}_X -bilinear morphism (see [Lan14, §3.2]; in this language, the morphism $p_s^G \mathcal{T}_p : p_s^G (\mathcal{T}_P) \to \mathcal{T}_X$ can be seen as the anchor of a natural restricted Lie algebroid over X/k with kernel $(p_*^G(\mathcal{T}_P))^{(0)} = \operatorname{ad}(P)$). We will also denote by $C_s : \bigwedge^2 \mathcal{T}_X \to \operatorname{ad}(P)$ the corresponding \mathcal{O}_X -linear morphism. It can be seen as a global section of the sheaf $\Omega_{X/S}^2 \otimes_{\mathcal{O}_X} \operatorname{ad}(P)$.

When $G = GL_n$, it is easy to compute the curvature of connections on the trivial bundle:

Lemma 2.7. Let $\Omega \in \mathrm{H}^{0}(X, \mathrm{Lie}(\mathrm{GL}_{n}) \otimes_{k} \Omega^{1}_{X/k})$. Then there is equality in $\mathrm{H}^{0}(X, \mathrm{Lie}(\mathrm{GL}_{n}) \otimes_{k} \Omega^{2}_{X/k}) : C_{s_{e}+\Omega} = \mathrm{d}\Omega + \Omega \wedge \Omega.$

Proof. [Del70, I (3.2.2)].

2.2.2 *p*-curvature

Definition 2.8. Let $s : \mathcal{T}_X \to p^G_*(\mathcal{T}_P)$ be an Atiyah-connection on a *G*-torsor $p : P \to X$. The *p*-curvature $\psi_s : \mathcal{T}_X \to \operatorname{ad} P$ is the morphism given for $D \in \mathcal{T}_X$ by

$$\psi_s(D) = s(D)^p - s(D^p)$$

If the Atiyah connection s is integrable, then the morphism ψ_s is p-linear (see [Lan14, Lemma 4.9]).

If F_X denotes the absolute Frobenius of X, we will also write $\psi_s : \mathcal{T}_X \to F_{X*}(\operatorname{ad} P)$ for the associated \mathcal{O}_X -linear morphism, or any of the equivalent following pieces of data : a \mathcal{O}_X -linear morphism $F_{X/S}^*\mathcal{T}_{X^{(p)}} \simeq F_X^*\mathcal{T}_X \to \operatorname{ad} P$ (where $F_{X/S} : X \to X^{(p)}$ is the relative Frobenius), or an element of $\operatorname{H}^0(X, F_{X/S}^*\Omega^1_{X^{(p)}/k} \otimes_{\mathcal{O}_X} \operatorname{ad}(P))$.

In contrast to the usual curvature, when G is non-abelian, there is no explicit formula (known to us) for the *p*-curvature of a connection on a trivial *G*-torsor, even when $G = \operatorname{GL}_n$. Nevertheless, we will be able to characterize, thanks to Cartier descent, the vanishing of *p*-curvature, see forthcoming Corollary 3.4. The first step is the following.

Lemma 2.9. Let $s : \mathcal{T}_X \to p^G_*(\mathcal{T}_P)$ be an Atiyah-connection on a G-torsor $p : P \to X$. The cancellation of C_s is a local property with respect to the étale topology on X. If the connection s is integrable, the same holds for the cancellation of ψ_s .

Proof. Let $\mathcal{F} = \Omega^2_{X/S} \otimes_{\mathcal{O}_X} \operatorname{ad}(P)$ (resp. $\mathcal{F} = F^*_X(\Omega^1_{X/S}) \otimes_{\mathcal{O}_X} \operatorname{ad}(P)$). Since \mathcal{F} is quasi-coherent on X, by flat descent, the pre-sheaf $(f : X' \to X) \mapsto \operatorname{H}^0(X', f^*\mathcal{F})$ is a sheaf on the small étale site of X. Besides, since formation of $\Omega^1_{X/S}$ and of $\operatorname{ad}(P)$ commutes with étale base change, we have that $f^*\mathcal{F} = \Omega^2_{X'/S} \otimes_{\mathcal{O}_{X'}}$ $\operatorname{ad}(P')$ (resp. $f^*\mathcal{F} = F^*_{X'}(\Omega^1_{X'/S}) \otimes_{\mathcal{O}_{X'}} \operatorname{ad}(P')$), and the restriction morphism $\operatorname{H}^0(X,\mathcal{F}) \to \operatorname{H}^0(X', f^*\mathcal{F})$ sends C_s to $C_{s'}$ (resp. ψ_s to $\psi_{s'}$), where s' is the induced Atiyah-connection on the induced G-torsor $P' \to X'$.

3 Frobenius descent for torsors

3.1 Frobenius descent for torsors under a smooth group scheme

Let G/k be a smooth affine group scheme. We first generalize the fact that if \mathcal{F} is a locally free sheaf on $X^{(p)}$, then $F^*_{X/S}\mathcal{F}$ is endowed with a canonical connection ∇_{can} (see Appendix A).

Lemma 3.1. The functor $F^*_{X/S}$: $BG(X^{(p)}) \to BG(X)$ factors canonically through $At(G, X) \to BG(X)$.

Proof. Let $p: P \to X^{(p)}$ be a *G*-torsor on $X^{(p)}$ and $q: X \times_{X^{(p)}} P \to X$ its pullback along $F_{X/S}: X \to X^{(p)}$. There is a natural morphism of exact sequences:

Since the Frobenius morphism induces 0 on tangent spaces, the morphism $q^G_*(\mathcal{T}_{X\times_{X^{(p)}}P}) \to F^*_{X/S}p^G_*(\mathcal{T}_P)$ factors through $\operatorname{ad}(X\times_{X^{(p)}}P) \to F^*_{X/S}p^G_*(\mathcal{T}_P)$, and this gives a retraction of the top exact sequence, or equivalently a section $s_{can}: \mathcal{T}_X \to q^G_*(\mathcal{T}_{X\times_{X^{(p)}}P})$. Thus we get a canonical Atiyah-connection (Definition 2.2) s_{can} on $X\times_{X^{(p)}}P$.

We use the symbol $(F^*_{X/S}, s_{can}) : B G(X^{(p)}) \to At(G, X)$ (or simply $F^*_{X/S}$ when there is no ambiguity) to denote the functor induced by Lemma 3.1.

Theorem 3.2 (Frobenius descent for G-torsors). Let X/k be a smooth scheme and G/k a smooth affine group scheme. The functor $(F^*_{X/S}, s_{can})$ induces an equivalence of categories:

$$(F^*_{X/S}\cdot,s_{can}):\operatorname{B} G(X^{(p)})\to\operatorname{At}(G,X)^{C=\psi=0}$$

between the category of G-torsors on $X^{(p)}$ and the category of G-torsors on X endowed with an Atiyah-connection with vanishing curvature and vanishing p-curvature.

The aim of this note is not to discuss a proof of this theorem, but to give some applications. For a tannakian proof, see [Mam16]. For a sketch of a more direct proof, using connexions on the algebras of the torsors, see [CZ15]. Both proofs ultimately rely on the $G = GL_n$ case.

3.2 Torsors with vanishing curvatures

We give a first application of Theorem 3.2 to the local structure of connections on torsors with vanishing curvature and vanishing p-curvature.

Proposition 3.3. With the hypothesis of Theorem 3.2, an Atiyah-connection s on a G-torsor $p: P \to X$ verifies $C_s = \psi_s = 0$ if and only if étale locally on X, there exists an isomorphism $(P, s) \simeq (X \times G, s_e)$.

Proof. The "if" part is clear, since the cancellation of the (p-)curvature is a property that is étale-local (see Lemma 2.9). For the "only if" direction: if we assume that $C_s = \psi_s = 0$, then by Theorem 3.2, $P \to X$ descends along $F_{X/S} : X \to X^{(p)}$ into a *G*-torsor $\tilde{P} \to X^{(p)}$. Since *G* is smooth, there exists an étale covering $(X_i^{(p)} \to X^{(p)})_{i \in I}$ such that the *G*-torsor $\tilde{P} \to X^{(p)}$ is trivial when restricted along $X_i^{(p)} \to X^{(p)}$, for each $i \in I$. By applying the functor $(F_{X/S}^*, s_{can})$, we see that $(P, s) \simeq (G_X, s_e)$ when restricted along each arrow of the pull-back étale cover $(X_i \to X)_{i \in I}$ of X.

Corollary 3.4. Let $\Omega \in \mathrm{H}^0(X, \Omega^1_{X/k} \otimes_k \mathrm{Lie}(G))$. Then the Atiyah-connection $s_e + \Omega$ on $X \times G \to G$ has vanishing curvature and p-curvature if and only if étale locally on X, Ω is in the image of the Maurer-Cartan map.

Proof. This follows from Proposition 3.3 and Lemma 2.5.

3.3 Special groups


Let G/k be a group scheme, and X a scheme. We recall that a fpqc G-torsor $P \to X$ is called *locally isotrivial* if for any point $x \in X$, there exists a Zariski open $U \subset X$ containing x, and a finite étale cover $U' \to U$, such that $U' \times_X P \to U'$ is trivial as a G-torsor. In fact, when G/k is a smooth affine group scheme, any fpqc torsor is locally isotrivial (see [Ray70, Lemme XIV.1.4]).

Serre in [Ser58, §4.4] (see also [Gro58, §3]) defines a special group as a group scheme G/k such that any fpqc G-torsor that is locally isotrivial is already locally trivial in the Zariski topology (the definition is given for an algebraically closed field but makes sense over an arbitrary field). For a modern treatment, see [RT20]. Classical examples include \mathbb{G}_m (more generally GL_n) and \mathbb{G}_a .

If G/k is a smooth affine group scheme, that is special in the above sense, then Proposition 3.3 and Corollary 3.4 hold verbatim if we replace the word 'étale' by the word 'Zariski'.

3.4 Torsors trivialized along the relative Frobenius


Definition 3.5. We define the category $B G(X^{(p)})^{F_{X/S}-triv}$ as the 2-product fitting in the following diagram :



where * denotes the discrete category with a single object.

3.5 Atiyah connections on trivialized bundles

Definition 3.6. Similarly, we define the category $\operatorname{At}^{triv}(G, X)$ as the 2-product fitting in the following diagram :



It is clear that $\operatorname{At}^{triv}(G, X)$ is a discrete category, more precisely: the functor

$$s_e + \cdot : \mathrm{H}^0(X, \Omega^1_{X/k} \otimes_k \mathrm{Lie}(G)) \to \mathrm{At}^{triv}(G, X)$$
 (2)

sending Ω to $((X \times G \to X, s_e + \Omega), id)$ is an equivalence of discrete categories, where we see the set on the left as a category with only identities as morphisms. There is even an explicit inverse equivalence given by transfer of structure.


We denote by $\operatorname{At}^{triv}(G,X)^{C=\psi=0}$ the full subcategory of $\operatorname{At}^{triv}(G,X)$ consisting of connections with vanishing curvatures, and similarly $\operatorname{H}^0(X, \Omega^1_{X/k} \otimes_k$

 $\operatorname{Lie}(G)^{C=\psi=0}$ for the corresponding Lie-valued 1-forms. It follows from Theorem 3.2 that the functor $(F^*_{X/S}, s_{can})$ induces an equivalence of categories:

$$(F_{X/S}^*, s_{can}) : \operatorname{B} G(X^{(p)})^{F_{X/S} - triv} \to \operatorname{At}^{triv}(G, X)^{C = \psi = 0}$$

3.6 Torsors trivialized along the absolute Frobenius

Definition 3.7. We define the category $B G(X)^{F_X - triv}$ as the 2-product fitting in the following diagram :



We will see a natural interpretation of this definition in section 5 but first use it in section 4. When the base field is perfect, it is clear that the functor $\mathrm{B}\,G(X)^{F_X-triv} \to \mathrm{B}\,G^{(p)}(X^{(p)})^{F_{X/S}-triv}$ is an equivalence, so that we get a natural equivalence $\mathrm{B}\,G(X)^{F_X-triv} \to \mathrm{H}^0(X, \Omega^1_{X/k} \otimes_k \mathrm{Lie}(G^{(p)}))^{C=\psi=0}$.

4 Affine line bundles trivialized along the Frobenius

4.1 Affine line bundles

Definition 4.1. An affine line bundle over a scheme S is a couple (L, A) where L is a line bundle and A is a torsor under L.

The morphisms are the pairs $(\phi, \psi) : (L, A) \to (L', A')$ so that $\phi : L \to L'$ is linear and $\psi : A \to A'$ is ϕ -equivariant.

Since each object is Zariski locally isomorphic to the trivial object $(\mathbb{G}_a, \mathbb{A}^1)$, affine line bundles form a neutral gerbe banded by $\operatorname{Aut}(\mathbb{G}_a, \mathbb{A}^1) = \mathbb{G}_a \ltimes \mathbb{G}_m$. This affine group will be denoted by aff(1).

Remark 4.2. V. Plechinger gives another description of this category as the category of epimorphisms $\mathcal{E} \twoheadrightarrow \mathcal{O}_S$ where \mathcal{E} is a rank 2 vector bundle on S, see [Ple19b; Ple19a]. It is easy to work out an explicit equivalence.

4.2 de Rham interpretation

We are now back to our main setup where X/k is a smooth variety. After a brief reminder of the cases $G = \mathbb{G}_m$ and $G = \mathbb{G}_a$, we give an interpretation of affine line bundles (trivialized by the Frobenius) in terms of differential forms.

4.2.1 Cartier operator

We denote by $\mathscr{C} : \mathrm{H}^{0}(X, \mathrm{Z}(\Omega^{1}_{X/k})) \to \mathrm{H}^{0}(X^{(p)}, \Omega^{1}_{X^{(p)}/k})$ the Cartier operator. For its definition, we refer for instance to L.Illusie's article in [Ber+96], in particular, to §3 therein. More generally, the Cartier operator defines an isomorphism

$$\mathscr{C}: \oplus_{i \ge 0} \mathcal{H}^i\left((F_{X/k})_* \Omega^{\bullet}_{X/k} \right) \to \oplus_{i \ge 0} \Omega^i_{X^{(p)}/k}$$

4.2.2 Line bundles and 1-forms

Proposition 4.3 ([Kat72, Corollary 7.1.3]). Let $\omega \in \mathrm{H}^0(X, \mathrm{Z}(\Omega^1_{X/k}))$ be a closed 1-form. The following statements are equivalent:

- 1. $\psi(\mathcal{O}_X, \mathbf{d} + \omega) = 0,$
- 2. $\mathscr{C}\omega = \pi^*_{X/k}\omega$,
- 3. ω is locally logarithmic.

Remark 4.4. Here $\pi_{X/k} : X^{(p)} \to X$ denotes the canonical morphism (see Appendix A for details). When X is perfect, we identify the isomorphism $\pi_{X/k}$ with id_X and write the above equality as $\mathscr{C} \omega = \omega$.

Proof. The assertions 1 and 3 are equivalent by Corollary 3.4. The equivalence of 1 and 2 follows from a direct computation, see [Kat72, Proposition 7.1.2] \Box

4.2.3 \mathbb{G}_a -torsors and 1-forms

Since $\text{Lie}(\mathbb{G}_a) = k$, any 1-form ω also defines a connection on the trivial \mathbb{G}_a -torsor. The analogue of Proposition 4.3 is well known:

Proposition 4.5. Let $\omega \in \mathrm{H}^0(X, \mathrm{Z}(\Omega^1_{X/k}))$ be a closed 1-form. The following statements are equivalent:

- 1. $\psi(\mathbb{G}_a, \mathbf{d} + \omega) = 0,$ 2. $\mathscr{C} \omega = 0,$
- 3. ω is locally exact.

Proof. Again, the assertions 1 and 3 are equivalent by Corollary 3.4. The equivalence of 2 and 3 follows from Cartier's theorem ([Ber+96, Théorème 3.5, p.130]). \Box

4.2.4 de Rham interpretation of affine line bundles

For the rest of this section we set G = aff(1). This is a special group, so according to §3.3 we can work with the Zariski topology rather than the étale topology.

We use the standard embedding $G \to \operatorname{GL}_2$ given by $(a,b) \mapsto \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$. Thus Lie *G* identifies with $\left\{ \begin{pmatrix} a & b \\ 0 & 0 \end{pmatrix}, a, b \in k \right\}$, hence connections on the trivial *G*-torsor are given by matrices

$$\Omega = \begin{pmatrix} \omega & \omega' \\ 0 & 0 \end{pmatrix}$$

where ω, ω' are two 1-forms on X. Lemma 2.7 gives immediately :

$$C_{s_e+\Omega} = \begin{pmatrix} \mathrm{d}\omega & \mathrm{d}\omega' + \omega \wedge \omega' \\ 0 & 0 \end{pmatrix}$$

Proposition 4.6. With notations as above, assume that $d\omega = 0 = d\omega' + \omega \wedge \omega'$. Then $\psi_{s_e+\Omega} = 0$ if and only if $\mathscr{C} \omega = \pi^*_{X/k} \omega$ and for each Zariski open $U \subset X$ and each $f \in \mathbb{G}_m(U)$ such that $\omega_{|U} = df/f$, the equality $\mathscr{C}(f\omega'_{|U}) = 0$ holds.

Remark 4.7. The condition $\mathscr{C}(f\omega'_{|U}) = 0$ makes sense because $d\omega' + \omega \wedge \omega' = 0$ ensures that $f\omega'_{|U}$ is closed.

Proof of Proposition 4.6. Frobenius descent, and in particular Corollary 3.4, implies that $\psi_{s_e+\Omega} = 0$ if and only if Ω is Zariski locally in the image of the Maurer-Cartan map. If $g = \begin{pmatrix} f & f' \\ 0 & 1 \end{pmatrix}$; then $g^{-1}dg = \begin{pmatrix} f^{-1}df & -f^{-1}df' \\ 0 \end{pmatrix}$, so $\psi_{s_e+\Omega} = 0$ if and only if Zariski locally there exists $f \in \mathbb{G}_m$ and $f' \in \mathbb{G}_a$ such that $\omega = f^{-1}df$ and $\omega' = -f^{-1}df'$. Hence the 'if' direction follows from the cases $G = \mathbb{G}_m$ and $G = \mathbb{G}_a$ (Propositions 4.3 and 4.5). For the 'only if' direction : it follows from the $G = \mathbb{G}_m$ case that $\mathscr{C}\omega = \pi^*_{X/k}\omega$. We now fix a Zariski open $U \subset X$ and $f \in \mathbb{G}_m(U)$ such that $\omega_{|U|} = df/f$. By hypothesis, there exists a Zariski cover $(U_i)_{i\in I}$ of X, and for each $i \in I$, a pair (f_i, f_i') in $G(U_i)$ such that $\omega_{|U_i} = f_i^{-1}df_i$ and $\omega'_{|U_i} = -f_i^{-1}df_i'$. On each $U \cap U_i$, since we assume that X/k is smooth, there exists f_i'' in $\mathbb{G}_m((U \cap U_i)^{(p)})$ such that $f_{|U \cap U_i} = F^*_{U \cap U_i/S}(f_i'')f_{i|U \cap U_i}$ Hence by p^{-1} -linearity of the Cartier operator, for each $i \in I$ we have : $\mathscr{C}(f_{|U \cap U_i}\omega'_{|U \cap U_i}) = f_i'' \mathscr{C}(f_i|_{U \cap U_i}\omega'_{|U \cap U_i}) = 0$, which implies $\mathscr{C}(f\omega'_{|U}) = 0$.

We get the following quite complete description of the set of affine line bundles on X trivialized along the absolute Frobenius :

Corollary 4.8. Assume that X/k is smooth, and that k is perfect. Then the set of isomorphism classes of affine line bundles on X trivialized along the Frobenius is in one to one correspondence with the set of pairs (ω, ω') of 1-forms on X such that :

1.
$$d\omega = 0 = d\omega' + \omega \wedge \omega'$$
,

- 2. $\mathscr{C} \omega = \omega$,
- 3. for each Zariski open $U \subset X$ and each $f \in \mathbb{G}_m(U)$ such that $\omega_{|U} = df/f$, the equality $\mathscr{C}(f\omega'_{|U}) = 0$ holds.

Proof. This follows from section 3.6 and Proposition 4.6.

5 Torsors under Frobenius kernels

5.1 Frobenius kernels

Let k be a field and G/k be a smooth affine group scheme. We define its Frobenius kernel as $G^F := \ker(G \to G^{(p)})$. In this §5.1, we will give a complete description of G^F -torsors on a smooth scheme X/k, when k is perfect.

Lemma B.1 implies that the functor $F_X^* : BG(X) \to BG^{(p)}(X)$ is isomorphic with restriction of the structure group along $G \to G^{(p)}$. It follows from Definition 3.7 that there is a natural equivalence $BG^F(X) \to BG(X)^{F_X-triv}$. Now from §3.5 and §3.6 one get a natural functor $BG^F(X) \to At^{triv}(G^{(p)}, X)$.

5.2 Torsors under Frobenius kernels and Maurer-Cartan map

This functor $\mathcal{B} G^F(X) \to \operatorname{At}^{triv}(G^{(p)}, X)$ is strongly related to the Maurer-Cartan map in the following sense. For $g \in G^{(p)}(X)$, we define $\partial(g) = X \times_{g,G^{(p)}} G$, seen as a G^F -torsor on X.

Lemma 5.1. The following diagram is 2-commutative :

$$\begin{array}{ccc} \operatorname{B} G^{F}(X) & \longrightarrow & \operatorname{At}^{triv}(G^{(p)}, X) \\ & & & & \uparrow & & \uparrow^{s_{e}+\cdot} \\ & & & & G^{(p)}(X) & \xrightarrow{\operatorname{d} \log} & \operatorname{H}^{0}(X, \Omega^{1}_{X/k} \otimes_{k} \operatorname{Lie}(G^{(p)})) \end{array}$$

Proof. It is enough to show that the following diagram is commutative :

$$\begin{array}{ccc} \operatorname{B} G^{F}(X) & \longrightarrow \operatorname{At}^{triv}(G^{(p)}, X) \\ & & & \downarrow \\ & & & & \\ G^{(p)}(X) \xrightarrow{\operatorname{d} \log} \operatorname{H}^{0}(X, \Omega^{1}_{X/k} \otimes_{k} \operatorname{Lie}(G^{(p)})) \end{array}$$

where the right vertical functor is the inverse equivalence of (2). As this is just a matter of unraveling the definitions, we skip the proof.

By construction, the functor $\mathcal{B}G^F(X) \to \operatorname{At}^{triv}(G^{(p)}, X)$ factors through $\operatorname{At}^{triv}(G^{(p)}, X)^{C=\psi=0}$. Moreover:

Lemma 5.2. If k is perfect, the functor $B G^F(X) \to At^{triv}(G^{(p)}, X)^{C=\psi=0}$ is an equivalence of categories.

Proof. This has already been observed in §3.6.

Remark 5.3. In particular, $B G^F(X)$ is rigid, which is also straightforward to show directly, since X/k is smooth and G^F is an infinitesimal group scheme over k.

Theorem 5.4. Assume that k is perfect. There is a natural one to one map $\mathrm{H}^1(X, G^F) \to \mathrm{H}^0(X, \Omega^1_{X/k} \otimes_k \mathrm{Lie}(G))^{C=\psi=0}$ that for each $g \in G(X)$ sends the image $\partial(g)$ of g via the cobord map $\partial : G^{(p)}(X) \to \mathrm{H}^1(X, G^F)$ to $\mathrm{dlog}(g)$, commutes with étale base change, and is characterized by these properties.

Proof. According to equation (2), Lemmas 5.1 and 5.2 there is a natural 2-commutative diagram:

$$\begin{array}{ccc} \operatorname{B} G^{F}(X) & \longrightarrow & \operatorname{At}^{triv}(G^{(p)}, X)^{C=\psi=0} \\ & & & & & \\ \partial \uparrow & & & & & \\ G^{(p)}(X) & \xrightarrow{\operatorname{d} \log} & \operatorname{H}^{0}(X, \Omega^{1}_{X/k} \otimes_{k} \operatorname{Lie}(G^{(p)}))^{C=\psi=0} \end{array}$$

All categories are rigid, so we can identify $BG^F(X)$ with $H^1(X, G^F)$ and because the functor $s_e + \cdot$ admits an explicit inverse, we get a (1-)commutative diagram :

$$\begin{array}{c} \operatorname{B} G^{F}(X) \\ \stackrel{}{ \partial \uparrow} \\ G^{(p)}(X) \xrightarrow{\operatorname{d} \log} \operatorname{H}^{0}(X, \Omega^{1}_{X/k} \otimes_{k} \operatorname{Lie}(G^{(p)}))^{C=\psi=0} \end{array}$$


where the diagonal map α is one to one. The fact that the formation of α commutes with étale base change on X is clear, and the unicity follows, since any element of B $G^F(X)$ is étale locally in the image of ∂ (by Corollary 3.4, or an easy direct proof).

- **Remark 5.5.** 1. The rigidity of $B G^F(X)$ is the key fact that enables, starting from $\Omega \in H^0(X, \Omega^1_{X/k} \otimes_k \operatorname{Lie}(G^{(p)}))^{C=\psi=0}$, corresponding on an étale cover $(X_i \to X)_{i \in I}$ to elements $g_i \in G^{(p)}(X_i)$, to glue together the local torsors $T_i = \partial(g_i)$. Since the T_i 's are given by explicit equations, this gives a very concrete description of the map α^{-1} .
 - 2. The assumption that k is perfect is essential. Namely, in the simple case $X = \mathbb{A}^1, G = \mathbb{G}_m$, it is easy to see that $\mathrm{H}^1(X, G^F) \simeq k^*/k^{*p}$ whereas $\mathrm{H}^0(X, \Omega^1_{X/k} \otimes_k \mathrm{Lie}(G))^{C=\psi=0}$ is trivial.

3. To describe explicitly the set $\mathrm{H}^0(X, \Omega^1_{X/k} \otimes_k \mathrm{Lie}(G))^{C=\psi=0}$ is probably a difficult question, as there is for a general, non-commutative group G, no explicit formula (at least known to us) for the *p*-curvature. Interestingly, when G is commutative, one can express the *p*-curvature ψ using only the Cartier operator \mathscr{C} and the *p*-th power operation on $\mathrm{Lie}(G)$, see Appendix C.

A Classical Frobenius descent

In this paragraph, we put $S = \operatorname{spec} k$. We recall Cartier's celebrated result on Frobenius descent. This is nowadays seen as fppf descent along the relative Frobenius morphism $F_{X/S}$ of X/S, that is the unique morphism making the following diagram commute :



where F_S (respectively F_X) is the absolute Frobenius of S (respectively of X).

Let \mathcal{F} be a quasi-coherent sheaf on $X^{(p)}$. Then $F_{X/S}^*\mathcal{F} = \mathcal{O}_X \otimes_{\mathcal{O}_{X^{(p)}}} \mathcal{F}$ is endowed with a canonical connection ∇^{can} given by $f \otimes s \mapsto (1 \otimes s) \otimes df$. This connection is integrable with vanishing *p*-curvature.

In the other direction, if (\mathcal{E}, ∇) is a quasi-coherent sheaf endowed with an integrable connection X, of vanishing p-curvature, then the sub-sheaf of horizontal sections $\mathcal{E}^{\nabla} = \{s \in \mathcal{E}/\nabla(s) = 0\}$ is a quasi-coherent sheaf on $X^{(p)}$.

Theorem A.1 (Cartier). Assume X/S is smooth.

The functors $\mathcal{F} \mapsto (F_{X/S}^*\mathcal{F}, \nabla_{can})$ and $(\mathcal{E}, \nabla) \mapsto \mathcal{E}^{\nabla}$ are inverse equivalences between the category of quasi-coherent sheaves on $X^{(p)}$ and the category of quasicoherent sheaves on X endowed with an integrable connection of vanishing pcurvature.

Proof. See [Kat70, Theorem (5.1)].

B Frobenius pull-back of torsors

We use notations of diagram (3).

Lemma B.1. Let G/S be an affine group scheme and $P \to X$ be a G-torsor. There exists a natural isomorphism of $G^{(p)}$ -torsors:

$$P \wedge^G G^{(p)} \simeq F_X^* P$$
 .

Proof. Notice that $F_X^*P = X \times_{F_X,X} P = P^{(p/X)}$, moreover the relative Frobenius morphism $F_{P/X} : P \to P^{(p/X)} = F_X^*P$ is equivariant with respect to the morphism $F_G : G \to G^{(p)}$, hence give rise to an (iso)morphism of $G^{(p)}$ -torsors $P \wedge^G G^{(p)} \simeq F_X^*P$.

C *p*-curvature for abelian structure groups

In this section, we assume that $\operatorname{car} k \neq 2$.

Let G/k be an affine, smooth, abelian group scheme. In this situation, there is an explicit description of the *p*-curvature map in terms of the Cartier operator $\mathscr{C}: F_{X/S}_*(\mathbb{Z}(\Omega^1_{X/k})) \to \Omega^1_{X^{(p)}/k}.$

Let us first remark that if $\Omega \in \mathrm{H}^0(X, \Omega^1_{X/k} \otimes_k \mathrm{Lie}(G))$, since G is abelian, according to Lemma 2.7, we have that $C_{s+\Omega} = \mathrm{d}\Omega$. Hence the source of the *p*-curvature map can be written as $\mathrm{H}^0(X, \mathrm{Z}(\Omega^1_{X/k})) \otimes_k \mathrm{Lie}(G)$.

Proposition C.1. If G/k be an affine, smooth, abelian group scheme, the pcurvature morphism :

$$\psi : \mathrm{H}^{0}(X, \mathrm{Z}(\Omega^{1}_{X/k})) \otimes_{k} \mathrm{Lie}(G) \to \mathrm{H}^{0}(X^{(p)}, \Omega^{1}_{X^{(p)}/k}) \otimes_{k} \mathrm{Lie}(G)$$

equals $\pi^*_{X/k} \otimes P - \mathscr{C} \otimes \mathrm{id}$, where $\pi_{X/k} : X^{(p)} \to X$ is the canonical morphism, P is the p-th power map on $\mathrm{Lie}(G)$, and \mathscr{C} is the Cartier operator.

Proof. This is [CZ15, (A.8)] (but see also [AM76, Lemma (2.2)]).

References

- [AM76] M. Artin and J. S. Milne. "Duality in the flat cohomology of curves". In: *Invent. Math.* 35 (1976), pp. 111–129. ISSN: 0020-9910.
- [Ati57] M. F. Atiyah. "Complex analytic connections in fibre bundles". In: Trans. Amer. Math. Soc. 85 (1957), pp. 181–207. ISSN: 0002-9947.
- [Ber+96] José Bertin et al. Introduction à La Théorie de Hodge. Vol. 3. Panoramas et Synthèses [Panoramas and Syntheses]. Société Mathématique de France, Paris, 1996. vi+273. ISBN: 978-2-85629-049-1.
- [Bou24] Naoufal Bouchareb. Affine Holomorphic Bundles over P¹_ℂ and Apolar Ideals. Oct. 24, 2024. arXiv: 2410.18706. URL: http://arxiv.org/abs/2410.18706. Pre-published.
- [CZ15] Tsao-Hsien Chen and Xinwen Zhu. "Non-abelian Hodge theory for algebraic curves in characteristic p". In: Geom. Funct. Anal. 25.6 (2015), pp. 1706–1733. ISSN: 1016-443X. URL: http://dx.doi.org/10.1007/s00039-015-0343-6.
- [Del70] Pierre Deligne. Équations différentielles à points singuliers réguliers. Lecture Notes in Mathematics, Vol. 163. Springer-Verlag, Berlin-New York, 1970, pp. iii+133.

- [Gro58] A. Grothendieck. "Torsion homologique et sections rationnelles". fre.
 In: Séminaire Claude Chevalley 3 (1958), pp. 1–29. URL: http://eudml.org/doc/110353.
- [Kat70] Nicholas M. Katz. "Nilpotent connections and the monodromy theorem: Applications of a result of Turrittin". In: Inst. Hautes Études Sci. Publ. Math. 39 (1970), pp. 175–232. ISSN: 0073-8301.
- [Kat72] Nicholas M. Katz. "Algebraic Solutions of Differential Equations (\$p\$-Curvature and the Hodge Filtration)". In: Inventiones Mathematicae 18 (1972), pp. 1–118. ISSN: 0020-9910.
- [Lan14] Adrian Langer. "Semistable modules over Lie algebroids in positive characteristic". In: Doc. Math. 19 (2014), pp. 509–540. ISSN: 1431-0635.
- [Mam16] Mohamed Rafik Mammeri. "Descente de Cartier et Torseurs Sous Le Noyau de Frobenius". These de doctorat. Lille 1, Dec. 14, 2016. URL: https://theses.fr/2016LIL10170.
- [Mil80] James S. Milne. Étale Cohomology. Vol. 33. Princeton Mathematical Series. Princeton, N.J.: Princeton University Press, 1980. xiii+323. ISBN: 0-691-08238-3.
- [Ple19a] Valentin Plechinger. "Classifying Affine Line Bundles on a Compact Complex Space". In: *Complex Manifolds* 6.1 (2019), pp. 103–117.
- [Ple19b] Valentin Plechinger. "Espaces de Modules de Fibrés En Droites Affines". These de doctorat. Aix-Marseille, Oct. 4, 2019. URL: http://www.theses.fr/2019AIXM0367.
- [Ray70] Michel Raynaud. Faisceaux amples sur les schémas en groupes et les espaces homogènes. Lecture Notes in Mathematics, Vol. 119. Springer-Verlag, Berlin-New York, 1970, pp. ii+218.
- [RT20] Zinovy Reichstein and Dajano Tossici. "Special Groups, Versality and the Grothendieck-Serre Conjecture". In: Documenta Mathematica 25 (Jan. 1, 2020), pp. 171–188. ISSN: 1431-0635. URL: https://ems.press/journals/dm/articl
- [Ser58] J.-P. Serre. "Espaces fibrés algébriques". fre. In: Séminaire Claude Chevalley 3 (1958), pp. 1–37. URL: http://eudml.org/doc/108928.
- [Wei55] André Weil. Fibre spaces in algebraic geometry. English. Notes by A. Wallace. Department of Mathematics, University of Chicago 48 p. 1955.