

ESSENTIAL p -CAPACITY-VOLUME ESTIMATES FOR ROTATIONALLY SYMMETRIC MANIFOLDS

XIAOSHANG JIN AND JIE XIAO

ABSTRACT. Given $p \in [1, \infty]$, this article presents the novel basic volumetric estimates for the relative p -capacities with their applications to finding not only the sharp weak (p, q) -imbeddings but also the precise lower bounds of the principal p -frequencies, which principally live in the rotationally symmetric manifolds.

CONTENTS

Introduction	1
1. Vanishing relative p -capacities under $\alpha < 1 - \frac{p}{n}$	4
2. Nonvanishing relative p -capacities under $\alpha \geq 1 - \frac{p}{n}$	6
3. Applying $\alpha = 1 - \frac{p}{n}$ to sharp weak (p, q) -imbeddings	12
4. Applying $\alpha = 1 > 1 - \frac{p}{n}$ to principal p -frequencies	16
References	18

INTRODUCTION

Given $p \in (0, \infty]$, if

$$v_n = |\mathbb{B}^n| \quad \& \quad nv_n = |\mathbb{S}^{n-1}|$$

are respectively the volume of the unit open ball \mathbb{B}^n & the surface area of the unit closed sphere $\mathbb{S}^{n-1} = \partial\mathbb{B}^n$ within the $1 \leq n$ -dimensional Euclidean space \mathbb{R}^n , then it is well-known that for any compact subset K of a bounded open subdomain Ω of an n -dimensional complete Riemannian manifold (\mathbb{M}^n, g) with volume element dv_g , the relative functional/variational p -capacity

$$\text{cap}_p(K, \Omega) = \begin{cases} \inf \left\{ \int_{\Omega} |\nabla f|^p dv_g : 1_K \leq f \in C_0^1(\Omega) \right\} & \text{as } p \in (0, \infty); \\ \inf \left\{ \sup_{x \in \Omega} |\nabla f(x)| : 1_K \leq f \in C_0^1(\Omega) \right\} & \text{as } p = \infty, \end{cases}$$


defines the minimum energy of a condenser (K, Ω) (cf. [14] for $p = 2 = n - 1$) whose importance in mathematical physics can be three-dimensionally pictured as:

2010 *Mathematics Subject Classification.* 31B15, 49Q10, 53C21, 74G65.

Key words and phrases. Rotationally symmetric manifold, surface area, volume, relative p -capacity, principal p -frequency, weak (p, q) -imbedding,

The 1st-named & 2nd-named authors were supported by NNSF of China # 12201225 and NSERC of Canada # 202979 respectively.

Nature always prefer minimum energy state.



Also nature:



made with mematic

https://www.reddit.com/r/physicsmemes/comments/tcbosm/principle_of_minimum_energy/?rdt=55327

Referring to not only [16] for the Euclidean space \mathbb{R}^n but also [8] for the hyperbolic space \mathbb{H}^{n+1} , we have

$$\text{cap}_p(K, \Omega) = 0 \quad \forall p \in (0, 1).$$

Consequently, our main interest in these capacities then lies with using the volume pair $\{|K|, |\Omega|\}$ to find the essential lower bound of each relative capacity

$$\text{cap}_p(K, \Omega) \quad \forall p \in [1, \infty].$$

Note that if

$$\begin{cases} dv_g(x) = dx = \text{the Lebesgue volume element } \forall x \in \mathbb{M}^n = \mathbb{R}^n; \\ rK = \{rx \in \mathbb{R}^n : x \in K\}; \\ \emptyset \neq K^\circ = \text{the interior of } K; \\ f \in C^1(K^\circ); \\ f_r(\cdot) = f(\frac{\cdot}{r}) : rK \rightarrow \mathbb{R}, \end{cases}$$

then

$$\begin{cases} |K|^{\frac{p}{n}-1} \int_K |\nabla f(x)|^p dx = |rK|^{\frac{p}{n}-1} \int_{rK} |\nabla f_r(y)|^p dy & \text{as } p \in [1, \infty); \\ |K|^{\frac{1}{n}} \sup_{x \in K} |\nabla f(x)| = |rK|^{\frac{1}{n}} \sup_{y \in rK} |\nabla f_r(y)| & \text{as } p = \infty. \end{cases}$$

In other words,

$$\int_K \left(\frac{|\nabla f(x)|^p}{|K|^{1-\frac{p}{n}}} \right) dx \quad \& \quad \sup_{x \in K} \left(\frac{|\nabla f(x)|}{|K|^{-\frac{1}{n}}} \right)$$

are scaling invariant. Thus, within the 1st two sections of this paper we are naturally led to utilize the powers

$$1 - \frac{p}{n} \leq 1 - \frac{1}{n}$$

for a brand-new investigation of the precise-to-optimal lower bound of $\text{cap}_p(K, \Omega)$ based on such a power of the volume-rate as $(|K|/|\Omega|)^\alpha$, see also:

$$\begin{cases} \text{Theorem 1.1 when } \alpha < 1 - \frac{p}{n} \text{ lives in a complete Riemannian manifold;} \\ \text{Theorem 2.1 when } \alpha \geq 1 - \frac{p}{n} \text{ lives in a rotationally symmetric manifold.} \end{cases}$$

Furthermore, within the 2nd two sections of this paper, we employ Theorem 2.1 to not only sharply embed the Sobolev p -space

$$W_0^{1,p}(\Omega) = \overline{\left\{ f \in C_0^1(\Omega) : \|\nabla f\|_p = \left(\int_\Omega |\nabla f|^p dv_g \right)^{\frac{1}{p}} < \infty \right\}}$$

into the weak Lebesgue q -space

$$L^{q,\infty}(\Omega) = \begin{cases} \left\{ f : \|f\|_{q,\infty} = \sup_{\lambda>0} \lambda |\{x \in \Omega : |f(x)| \geq \lambda\}|^{\frac{1}{q}} < \infty \right\} & \text{as } q \in (0, \infty); \\ \left\{ f : \|f\|_\infty = \text{esssup}_{x \in \Omega} |f(x)| < \infty \right\} & \text{as } q = \infty, \end{cases}$$

but also precisely handle the volumetric behavior of the principal p -frequency (cf. [15, p. 24] for $p = 2 = n$)

$$\lambda_{1,p}(\Omega) = \begin{cases} \inf \left\{ \frac{\int_\Omega |\nabla f|^p dv_g}{\int_\Omega |f|^p dv_g} : f \in C_0^1(\Omega) \right\} & \text{as } p \in [1, \infty); \\ \inf \left\{ \frac{\sup_\Omega |\nabla f|}{\sup_\Omega |f|} : f \in C_0^1(\Omega) \right\} & \text{as } p = \infty, \end{cases}$$

see also:

$$\begin{cases} \text{Theorem 3.1 when } \alpha = 1 - \frac{p}{n} \text{ lives in a rotationally symmetric manifold;} \\ \text{Theorem 4.1 when } \alpha = 1 > 1 - \frac{p}{n} \text{ lives in a rotationally symmetric manifold.} \end{cases}$$

1. VANISHING RELATIVE p -CAPACITIES UNDER $\alpha < 1 - \frac{p}{n}$

Theorem 1.1. *Given*

$$\alpha < 1 - \frac{p}{n} \leq 1 - \frac{1}{n},$$

suppose that (\mathbb{M}^n, g) is an n -dimensional complete Riemannian manifold. There hold two vanishing results for $\text{cap}_p(K, \Omega)$.

(i) *If $p \neq n$, then*

$$(1.1) \quad 0 = \inf_{K \subset \subset \Omega} \begin{cases} |K|^{-\alpha} \text{cap}_p(K, \Omega) & \text{as } p \neq \infty; \\ |K|^{\frac{1}{n}} \text{cap}_\infty(K, \Omega) & \text{as } p = \infty. \end{cases}$$

(ii) *If $p = n$, then*

$$(1.2) \quad 0 = \inf_{K \subset \subset \Omega} |K|^{-1} \exp\left(-\beta(\text{cap}_n(K, \Omega))^{\frac{1}{1-n}}\right) \quad \forall \beta > n^{\frac{n}{n-1}} v_n^{\frac{1}{n-1}}.$$

Proof. (i) For $p \neq n$, we select a point $x \in \Omega$ such that

$$B(x, r) \subseteq B(x, R) \subseteq \Omega \quad \text{for some small } r < R,$$

where $B(x, r)$ is x -centered geodesic open ball with radius r , thereby estimating

$$\frac{\text{cap}_p(\overline{B(x, r)}, \Omega)}{|B(x, r)|^\alpha} \leq \frac{\text{cap}_p(\overline{B(x, r)}, B(x, R))}{|B(x, r)|^\alpha} \leq \frac{\left(\int_r^R S_t^{\frac{1}{1-p}} dt\right)^{1-p}}{|B(x, r)|^\alpha},$$

where

$$\begin{cases} S_t = |\{y \in \Omega : \text{dist}(y, x) = t\}| \sim n v_n t^{n-1} & \text{as } t \rightarrow 0; \\ |B(x, t)| \sim v_n t^n & \text{as } t \rightarrow 0; \\ |\partial B(x, t)| \sim n v_n t^{n-1} & \text{as } t \rightarrow 0. \end{cases}$$

In the above & below, $U \sim V$ means that either U & V or U^{-1} & V^{-1} are equivalent infinitesimals.

If $p = 1$, then [12, p.107, Lemma] yields a constant C_1 such that

$$\begin{aligned} \lim_{r \rightarrow 0} \frac{\text{cap}_1(\overline{B(x, r)}, \Omega)}{|B(x, r)|^\alpha} &\leq \lim_{r \rightarrow 0} \frac{|\partial B(x, r)|}{|B(x, r)|^\alpha} \\ &\leq C_1 \lim_{r \rightarrow 0} r^{n-1-\alpha n} \\ &= 0 \quad \text{since } \alpha < 1 - n^{-1}. \end{aligned}$$

Thus, the argument for (1.1) is divided into three situations.

If $1 < p < n$, then there is a constant $C_p > 0$ such that

$$\lim_{r \rightarrow 0} \frac{\left(\int_r^R S_t^{\frac{1}{1-p}} dt\right)^{1-p}}{|B(x, r)|^\alpha} = C_p \lim_{r \rightarrow 0} \frac{r^{n-1}}{r^{n\alpha}} = 0 \quad \text{since } n\alpha < n - p < n - 1.$$

However, if $n < p < \infty$, then $\alpha < 1 - \frac{p}{n} < 0$. We can still derive that the above limit is 0 because not only the numerator is bounded but also the denominator tends on infinity. Therefore, we always have

$$\lim_{r \rightarrow 0} \frac{\text{cap}_p(\overline{B(x, r)}, \Omega)}{|\overline{B(x, r)}|^\alpha} = 0.$$

If $p \rightarrow \infty$, then (cf. [6, Theorem 1])

$$\begin{cases} \frac{\alpha}{p} < \frac{1}{p} - \frac{1}{n} \rightarrow -\frac{1}{n} < 0; \\ (\text{cap}_p(\overline{B(x, r)}, \Omega))^{\frac{1}{p}} \rightarrow \text{cap}_\infty(\overline{B(x, r)}, \Omega), \end{cases}$$

and hence

$$\begin{aligned} \limsup_{p \rightarrow \infty} \left(\frac{\text{cap}_p(\overline{B(x, r)}, \Omega)}{|\overline{B(x, r)}|^\alpha} \right)^{\frac{1}{p}} &= \limsup_{p \rightarrow \infty} \frac{(\text{cap}_p(\overline{B(x, r)}, \Omega))^{\frac{1}{p}}}{|\overline{B(x, r)}|^{\frac{\alpha}{p}}} \\ &\leq |\overline{B(x, r)}|^{\frac{1}{n}} (\text{cap}_\infty(\overline{B(x, r)}, \Omega)) \\ &\rightarrow 0 \text{ as } r \rightarrow 0. \end{aligned}$$

(ii) If $p = n$, then

$$\begin{aligned} \frac{\exp\left(-\beta(\text{cap}_n(\overline{B(x, r)}, \Omega))^{\frac{1}{1-n}}\right)}{|\overline{B(x, r)}|} &\leq \frac{\exp\left(-\beta(\text{cap}_n(\overline{B(x, r)}, B(x, R)))^{\frac{1}{1-n}}\right)}{|\overline{B(x, r)}|} \\ &\leq \frac{\exp\left(-\beta \int_r^R S_t^{\frac{1}{1-n}} dt\right)}{|\overline{B(x, r)}|} \end{aligned}$$

For

$$\beta > n^{\frac{n}{n-1}} \nu_n^{\frac{1}{n-1}},$$

there are sufficiently small ε & R such that

$$\frac{S_t}{t^{n-1}} < n\nu_n + \varepsilon < \left(\frac{\beta}{n}\right)^{n-1} \quad \forall t < R.$$

Accordingly, we get

$$\begin{aligned} \frac{\exp\left(-\beta \int_r^R S_t^{\frac{1}{1-n}} dt\right)}{|\overline{B(x, r)}|} &\leq \frac{\exp\left(-\beta(n\nu_n + \varepsilon)^{\frac{1}{1-n}} \int_r^R \frac{1}{t} dt\right)}{|\overline{B(x, r)}|} \\ &= \left(\frac{r}{R}\right)^{\beta(n\nu_n + \varepsilon)^{\frac{1}{1-n}}} \frac{1}{|\overline{B(x, r)}|}. \end{aligned}$$

Notice that

$$\beta(n\nu_n + \varepsilon)^{\frac{1}{1-n}} > n.$$

So, via setting $r \rightarrow 0$, we derive

$$\lim_{r \rightarrow 0} \frac{\exp\left(-\beta(\text{cap}_n(\overline{B(x, r)}, \Omega))^{\frac{1}{1-n}}\right)}{|\overline{B(x, r)}|} = 0,$$

as desired in (1.2). □

2. NONVANISHING RELATIVE p -CAPACITIES UNDER $\alpha \geq 1 - \frac{p}{n}$

Theorem 2.1. *Given*

$$\alpha \geq 1 - \frac{p}{n} \leq 1 - \frac{1}{n},$$

suppose that not only

$$(\mathbb{M}^n, g) = ([0, +\infty) \times \mathbb{S}^{n-1}, dt^2 + \varphi^2(t)g_{\mathbb{S}^{n-1}}) \text{ with } \varphi'' \geq 0.$$

but also Ω obeys the isoperimetry of $o = \{t = 0\}$ -centered balls - namely - if the surface area $|\partial\tilde{K}|$ of any compact subset \tilde{K} of Ω with the same volume as a geodesic ball $B(o, r)$ is not less than the surface area of $B(o, r)$ -i.e.-

$$|\partial\tilde{K}| \geq |\partial B(o, r)|.$$

There hold four nonvanishing results for $\text{cap}_p(K, \Omega)$.

(i) *If $1 \leq p < n$, then*

$$(2.1) \quad \inf_{K \subset \subset \Omega} \frac{\text{cap}_p(K, \Omega)}{|K|^\alpha |\Omega|^{1-\frac{p}{n}-\alpha}} \geq \begin{cases} nv_n^{\frac{p}{n}} \left(\frac{n-p}{p-1}\right)^{p-1} & \text{as } p \neq 1; \\ nv_n^{\frac{1}{n}} & \text{as } p = 1. \end{cases}$$

with equality being approachable whenever $\alpha = 1 - \frac{p}{n}$.

(ii) *If $p = n$, then*

$$(2.2) \quad \inf_{K \subset \subset \Omega} \frac{\exp\left(-\beta(\text{cap}_n(K, \Omega))^{\frac{1}{1-n}}\right)}{|K||\Omega|^{-1}} \geq 1 \quad \forall \beta \in (0, n^{\frac{n}{n-1}} v_n^{\frac{1}{n-1}}],$$

with equality being approachable whenever $\beta = n^{\frac{n}{n-1}} v_n^{\frac{1}{n-1}}$.

(iii) *If $n < p < \infty$, then*

$$(2.3) \quad \inf_{K \subset \subset \Omega} \frac{\text{cap}_p(K, \Omega)}{|\Omega|^{1-\frac{p}{n}}} \geq nv_n^{\frac{p}{n}} \left(\frac{p-n}{p-1}\right)^{p-1},$$

with equality being approachable.

(iv) *If $p = \infty$, then*

$$(2.4) \quad \inf_{K \subset \subset \Omega} \frac{\text{cap}_\infty(K, \Omega)}{|\Omega|^{-\frac{1}{n}}} \geq v_n^{\frac{1}{n}},$$

with equality being approachable.

Proof. (i) We only need to consider

$$(2.1) \text{ under } \alpha = 1 - \frac{p}{n} < 1 - \frac{1}{n},$$

thanks to

$$\frac{\text{cap}_p(K, \Omega)}{|K|^\alpha |\Omega|^{1-\frac{p}{n}-\alpha}} = \left(\frac{\text{cap}_p(K, \Omega)}{|K|^{1-\frac{p}{n}}} \right) \left(\frac{|\Omega|^{\alpha-(1-\frac{p}{n})}}{|K|^{\alpha-(1-\frac{p}{n})}} \right) \geq \frac{\text{cap}_p(K, \Omega)}{|K|^{1-\frac{p}{n}}}.$$

For any compact set K in Ω , assume that

$$|K| = |\overline{B(o, r)}| \quad \& \quad \Omega = |B(o, R)| \quad \text{for some } r < R,$$

where $B(o, r)$ & $B(o, R)$ are geodesic balls centered at $o = \{t = 0\}$.

Since (\mathbb{M}^n, g) satisfies the isoperimetry of $o = \{t = 0\}$ -centered balls, there is the isoperimetric function:

$$\begin{cases} I(\tau) = \inf \{|\partial K| : K \text{ is precompact open in } \Omega \ \& \ |K| \geq \tau\} = |\partial B(o, t)|; \\ t \text{ is determined by } |B(o, t)| = \tau. \end{cases}$$

Consequently, Maz'ya's isocapacity inequalities within either [12, (2.2.8)] or [5] indicates

$$\begin{aligned} \text{cap}_p(K, \Omega) &\geq \left(\int_{|K|}^{|\Omega|} (I(\tau))^{\frac{p}{1-p}} d\tau \right)^{1-p} \\ &= \left(\int_r^R (I(|B(o, t)|))^{\frac{p}{1-p}} \left(\frac{d\tau}{dt} \right) dt \right)^{1-p} \\ &= \left(\int_r^R |\partial B(o, t)|^{\frac{p}{1-p}} |\partial B(o, t)| dt \right)^{1-p} \\ &= \text{cap}_p(\overline{B(o, r)}, B(o, R)) \end{aligned}$$

This in turn implies

$$\begin{aligned} \frac{\text{cap}_p(K, \Omega)}{|K|^\alpha} &\geq \frac{\text{cap}_p(\overline{B(o, r)}, B(o, R))}{|\overline{B(o, r)}|^\alpha} \\ &= \frac{\left(\int_r^R S_t^{\frac{1}{1-p}} dt \right)^{1-p}}{|\overline{B(o, r)}|^\alpha} \\ &= \left(\frac{\int_r^R S_t^{\frac{1}{1-p}} dt}{\left(\int_0^r S_t dt \right)^{\alpha(1-p)^{-1}}} \right)^{1-p} \\ &= \left(\frac{\int_r^R (nv_n \varphi^{n-1}(t))^{\frac{1}{1-p}} dt}{\left(\int_0^r (nv_n \varphi^{n-1}(t)) dt \right)^{\alpha(1-p)^{-1}}} \right)^{1-p} \\ &= (F(r))^{1-p} (nv_n)^{1-\alpha}, \end{aligned}$$

where

$$F(r) = \left(\int_0^r \varphi^{n-1}(t) dt \right)^{\frac{\alpha}{p-1}} \int_r^R \varphi^{\frac{n-1}{1-p}}(t) dt.$$

- On the one hand, upon noticing that

$$\varphi(r) \sim r \quad \text{as } r \rightarrow 0,$$

we get

$$\begin{aligned}
F(0) &= \lim_{r \rightarrow 0} F(r) \\
&= \lim_{r \rightarrow 0} \frac{-\varphi^{\frac{n-1}{1-p}}(r)}{\alpha(1-p)^{-1} \left(\int_0^r \varphi^{n-1}(t) dt \right)^{\alpha(1-p)^{-1}-1} \varphi^{n-1}(r)} \\
&= \lim_{r \rightarrow 0} \frac{-r^{\frac{n-1}{1-p}}}{\alpha(1-p)^{-1} \left(\int_0^r \varphi^{n-1}(t) dt \right)^{\alpha(1-p)^{-1}-1} r^{n-1}} \\
&= \alpha^{-1}(p-1) \lim_{r \rightarrow 0} \frac{r^{\frac{(n-1)p}{1-p}}}{\left(\int_0^r \varphi^{n-1}(t) dt \right)^{\alpha(1-p)^{-1}-1}} \\
&= \left(\frac{n(p-1)}{n-1} \right) \left(\lim_{r \rightarrow 0} \frac{r^n}{\int_0^r \varphi^{n-1}(t) dt} \right)^{\frac{(n-1)p}{n(1-p)}} \\
&= \left(\frac{n(p-1)}{n-1} \right) n^{\frac{(n-1)p}{n(1-p)}}.
\end{aligned}$$

- On the other hand, we can prove that $F(r)$ obtains its maximum at $r = 0$. In order to achieve this, we compute

$$F'(r) = \left(\int_0^r \varphi^{n-1}(t) dt \right)^{\frac{\alpha}{p-1}-1} (\varphi(r))^{n-1} G(r),$$

where

$$G(r) = \alpha(p-1)^{-1} \int_r^R \varphi^{\frac{n-1}{1-p}}(t) dt - \varphi^{\frac{(n-1)p}{1-p}}(r) \int_0^r \varphi^{n-1}(t) dt$$

satisfies

$$\begin{cases} G(R) < 0; \\ G'(r) = \left(\frac{(n-1)p}{p-1} \right) (\varphi(r))^{\frac{n-1}{1-p}-n} H(r); \\ H(r) = \varphi'(r) \int_0^r \varphi^{n-1}(t) dt - n^{-1} \varphi^n(r). \end{cases}$$

That

$$H(0) = 0 \quad \& \quad H'(r) = \varphi''(r) \int_0^r \varphi^{n-1}(t) dt \geq 0$$

would imply that

$$H(r) \geq 0 \quad \& \quad G(r) \leq G(R) < 0.$$

As a consequence, we achieve that not only $F'(r) \leq 0$ but also $F(r)$ obtains its maximum at $r = 0$. Accordingly, there holds

$$\inf_{K \subset \subset \Omega} \frac{\text{cap}_p(K, \Omega)}{|K|^{1-\frac{p}{n}}} \geq \inf_{0 < r < R} (F(r))^{1-p} (n\nu_n)^{1-(1-\frac{p}{n})} = (F(0))^{1-p} (n\nu_n)^{\frac{p}{n}}.$$

In the end, from the above argument we know that the above-verified estimation is optimal in the sense that if

$$(K, \Omega) = (\overline{B(o, r)}, B(o, R)) \quad \& \quad r \rightarrow 0.$$

then (2.1)'s inequality approaches an equality whenever $\alpha = 1 - \frac{p}{n}$.

(ii) Assume that

$$|K| = |\overline{B(o, r)}| \quad \& \quad \Omega = |B(o, R)| \quad \text{for some } r < R.$$

Then

$$\begin{aligned} \frac{\exp\left(-\beta(\text{cap}_n(K, \Omega))^{\frac{1}{1-n}}\right)}{|K|} &\geq \frac{\exp\left(-n^{\frac{n}{n-1}} v_n^{\frac{1}{n-1}} \int_r^R (nv_n \varphi^{n-1}(t))^{\frac{1}{1-n}} dt\right)}{|\overline{B(o, r)}|} \\ &= \frac{\exp\left(-n \int_r^R \frac{dt}{\varphi(t)}\right)}{\int_0^r nv_n \varphi^{n-1}(t) dt} \\ &= \frac{1}{nv_n \bar{F}(r)}, \end{aligned}$$

where

$$\bar{F}(r) = \exp\left(n \int_r^R \frac{dt}{\varphi(t)}\right) \int_0^r \varphi^{n-1}(t) dt.$$

We are going to show that $\bar{F}(r)$ is an increasing function for $r \in [0, R]$. Notice that not only

$$\bar{F}'(r) = \exp\left(n \int_r^R \frac{dt}{\varphi(t)}\right) \frac{1}{\varphi(r)} \left(\varphi^n(r) - n \int_0^r \varphi^{n-1}(t) dt\right)$$

but also

$$\bar{F}'(r) \geq 0 \Leftrightarrow \left(\varphi^n(r) - n \int_0^r \varphi^{n-1}(t) dt\right) \geq 0,$$

whose right-hand inequality holds provided $\varphi' \geq 1$ which holds as $\varphi'' \geq 0$. Hence, we obtain

$$\inf_{K \subset \subset \Omega} |K|^{-1} \exp\left(-\beta(\text{cap}_n(K, \Omega))^{\frac{1}{1-n}}\right) \geq \inf_{0 < r < R} \frac{1}{nv_n \bar{F}(r)} \geq \frac{1}{nv_n \bar{F}(R)} = |\Omega|^{-1}.$$

From the proof we know that the above-verified inequality is optimal in the sense that if

$$\begin{cases} \beta = n^{\frac{n}{n-1}} v_n^{\frac{1}{n-1}}; \\ (K, \Omega) = (\overline{B(o, r)}, B(o, R)); \\ r \rightarrow R, \end{cases}$$

then (2.2)'s inequality approaches an equality.

(iii) Assume that

$$|K| = |\overline{B(o, r)}| \quad \& \quad \Omega = |B(o, R)| \quad \text{for some } r < R.$$

Then

$$\begin{aligned}
\text{cap}_p(K, \Omega) |\Omega|^{\frac{p}{n}-1} &\geq \left(\int_r^R (nv_n \varphi^{n-1}(t))^{\frac{1}{1-p}} dt \right)^{1-p} |\Omega|^{\frac{p}{n}-1} \\
&\geq \left(\int_0^R (nv_n \varphi^{n-1}(t))^{\frac{1}{1-p}} dt \right)^{1-p} \left(\int_0^R nv_n \varphi^{n-1}(t) dt \right)^{\frac{p}{n}-1} \\
&= (nv_n)^{\frac{p}{n}} (\tilde{F}(R))^{1-p},
\end{aligned}$$

where

$$\tilde{F}(R) = \left(\int_0^R \varphi^{n-1}(t) dt \right)^{-\frac{p-n}{n(p-1)}} \int_0^R \varphi^{\frac{n-1}{1-p}}(t) dt.$$

- On the one hand, there holds

$$\begin{aligned}
\tilde{F}(0) &= \lim_{R \rightarrow 0} \left(\frac{\varphi^{\frac{n-1}{1-p}}(R)}{\left(\frac{p-n}{n(p-1)} \left(\int_0^R \varphi^{n-1}(t) dt \right)^{\frac{p-n}{n(p-1)}-1} \varphi^{n-1}(R) \right)} \right) \\
&= \left(\frac{n(p-1)}{p-n} \right) \lim_{R \rightarrow 0} \left(\frac{R^{\frac{n-1}{1-p}}}{\left(\left(\frac{R^n}{n} \right)^{\frac{p-n}{n(p-1)}-1} R^{n-1} \right)} \right) \\
&= \left(\frac{p-1}{p-n} \right) n^{\frac{p-n}{n(p-1)}}.
\end{aligned}$$

- On the other hand, by a similar analysis to (i), we can get that $\tilde{F}' \leq 0$ provided $\varphi'' \geq 0$. Then

$$\inf_{K \subset \subset \Omega} \text{cap}_p(K, \Omega) |\Omega|^{\frac{p}{n}-1} \geq (nv_n)^{\frac{p}{n}} (\tilde{F}(0))^{1-p} = nv_n^{\frac{p}{n}} \left(\frac{p-n}{p-1} \right)^{p-1}.$$

Of course, (2.3)'s equality is approachable in the sense that if

$$(K, \Omega) = (\overline{B(o, r)}, B(o, R)) \quad \& \quad r < R \rightarrow 0,$$

then (2.3)'s inequality tends to an equality.

(iv) (2.4) follows from [6, Theorem 1] & (2.3) via

$$\inf_{K \subset \subset \Omega} \left(\frac{\text{cap}_p(K, \Omega)}{|\Omega|^{1-\frac{p}{n}}} \right)^{\frac{1}{p}} \geq n^{\frac{1}{p}} v_n^{\frac{1}{n}} \left(\frac{p-n}{p-1} \right)^{\frac{p-1}{p}} \rightarrow v_n^{\frac{1}{n}} \text{ as } p \rightarrow \infty,$$

Similarily, (2.4)'s equality is approachable in the sense that if

$$(K, \Omega) = (\overline{B(o, r)}, B(o, R)) \quad \& \quad r < R \rightarrow 0,$$

then (2.4)'s inequality tends to an equality. □

Remark 2.2. Below are six comments.

- Under the hypothesis that Ω enjoys the isoperimetry of $o = \{t = 0\}$ -centered balls, we can obtain the following stronger isocapacitary $(1, \infty) \ni p$ -inequality in a similar way to establish [8, Theorem 2.1]:

$$\exists \tau_0 \in (0, 1) \text{ such that } \left(\frac{\text{cap}_p(K, \Omega)}{\text{cap}_p(\overline{B(o, r)}, B(o, R))} \right)^{\frac{1}{p}} \geq \frac{|\partial D_{\tau_0}|}{|\partial B(o, t_0)|} \geq 1,$$

where

$$\begin{cases} D_\tau = \{x \in \Omega \setminus K : u(x) \geq \tau\} \cup K; \\ u \text{ is the } p\text{-capacitary potential of } (K, \Omega); \\ t_0 \text{ is determined by } |B(o, t_0)| = |D_{\tau_0}|. \end{cases}$$

- In (2.2), the condition “ $\varphi'' \geq 0$ ” can be weakened to “ $\varphi' \geq 1$.”
- From the geometric point of view, if we let not only $\text{Sec}[g]$ but also $\text{Ric}[g]$ denote not only the sectional curvature of (\mathbb{M}^n, g) but also the Ricci curvature of (\mathbb{M}^n, g) , then

$$\varphi'' \geq 0 \Leftrightarrow \text{Sec}[g] \leq 0 \Leftrightarrow \text{Ric}[g] \leq 0.$$

- If

$$\varphi(t) = t \text{ or } \varphi(t) = \sinh t,$$

then \mathbb{M}^n is the Euclidean space \mathbb{R}^n or the hyperbolic space \mathbb{H}^n . Thus, Theorem 2.1 holds for these two special manifolds; see also [1, 8] for certain of the related results.

- According to [4, Lemma 2.1] (cf. [11]), if

$$\begin{cases} \mathbb{M}^n = \mathbb{R}^n; \\ n - 1 < p \leq n; \\ \text{diam}(K) = \text{the diameter of } K, \end{cases}$$

then there is a constant $c_{p,n} > 0$ depending on $\{p, n\}$ such that

$$\frac{\text{cap}_p(K, \Omega)}{(\text{diam}(K))^{\frac{p}{n-1}} |\Omega|^{\frac{n-1-p}{n-1}}} \geq c_{p,n}.$$

Obviously, this last inequality can be treated as a rough variant of the special case $\alpha = \frac{p}{n(n-1)}$ of (2.1) with $p \in (n-1, n)$. While for $p = n$, (2.2) is stronger than the last rough inequality.

- Theorem 2.1 is a local version of the capacity-volume estimates. So, it is natural to consider the global version by setting not only

$$\Omega \rightarrow \mathbb{M}^n = [0, +\infty) \times \mathbb{S}^{n-1},$$

but also

$$\left\{ \begin{array}{ll} \overline{\text{cap}}_p(K) = \inf_{\Omega \subset \mathbb{M}^n} \text{cap}_p(K, \Omega) & \text{as } p \in [1, n); \\ \overline{\text{cap}}_n(K) = \inf_{\Omega \subset \mathbb{M}^n} |\Omega| \exp\left(-n^{\frac{n}{n-1}} \nu_n^{\frac{1}{n-1}} (\text{cap}_n(K, \Omega))^{\frac{1}{1-n}}\right) & \text{as } p = n; \\ \overline{\text{cap}}_p(K) = \inf_{\Omega \subset \mathbb{M}^n} |\Omega|^{\frac{p}{n-1}} \text{cap}_p(K, \Omega) & \text{as } p \in (n, \infty); \\ \overline{\text{cap}}_\infty(K) = \inf_{\Omega \subset \mathbb{M}^n} |\Omega|^{\frac{1}{n}} \text{cap}_\infty(K, \Omega) & \text{as } p = \infty. \end{array} \right.$$

Accordingly, we have the global p -isocapacitary inequalities below:

$$\left\{ \begin{array}{ll} \inf_{K \subset \subset \mathbb{M}^n} \frac{\overline{\text{cap}}_p(K)}{|K|^{1-\frac{1}{n}}} \geq n \nu_n^{\frac{p}{n}} \left(\frac{n-p}{p-1}\right)^{p-1} & \text{as } p \in [1, n); \\ \inf_{K \subset \subset \mathbb{M}^n} \frac{\overline{\text{cap}}_n(K)}{|K|} \geq 1 & \text{as } p = n; \\ \inf_{K \subset \subset \mathbb{M}^n} \overline{\text{cap}}_p(K) \geq n \nu_n^{\frac{p}{n}} \left(\frac{p-n}{p-1}\right)^{p-1} & \text{as } p \in (n, \infty); \\ \inf_{K \subset \subset \mathbb{M}^n} \overline{\text{cap}}_\infty(K) \geq \nu_n^{\frac{1}{n}} & \text{as } p = \infty; \end{array} \right.$$

thereby obtaining the generalized isoperimetric inequality (cf. [12, Lemma 2.2.5])

$$\overline{\text{cap}}_1(K) = \inf \left\{ |\Sigma| : \text{all admissible surfaces } \Sigma \text{ enclosing } \partial K \right\} \geq n \nu_n^{\frac{1}{n}} |K|^{1-\frac{1}{n}}.$$

3. APPLYING $\alpha = 1 - \frac{p}{n}$ TO SHARP WEAK (p, q) -IMBEDDINGS

Theorem 3.1. *Let*

$$\left\{ \begin{array}{l} (\mathbb{M}^n, g) = ([0, +\infty) \times \mathbb{S}^{n-1}, dt^2 + \varphi^2(t)g_{\mathbb{S}^{n-1}}) \text{ with } \varphi'' \geq 0; \\ \Omega = \text{a bounded domain in } \mathbb{M}^n \text{ with the isoperimetry of } o = \{t = 0\}\text{-centered balls;} \\ c_{n,p} = \begin{cases} n^{-1} \nu_n^{-\frac{1}{n}} & \text{as } p = 1; \\ n^{-\frac{1}{p}} \nu_n^{-\frac{1}{n}} \left(\frac{n-p}{p-1}\right)^{\frac{1-p}{p}} & \text{as } p \in (1, n); \\ n^{-1} \nu_n^{-\frac{1}{n}} & \text{as } p = n; \\ n^{-\frac{1}{p}} \nu_n^{-\frac{1}{n}} \left(\frac{p-n}{p-1}\right)^{\frac{1-p}{p}} & \text{as } p \in (n, \infty); \\ \nu_n^{-\frac{1}{n}} & \text{as } p = \infty. \end{cases} \end{array} \right.$$

Then there hold four sharp weak (p, q) -imbeddings.

(i) *If $1 \leq p < n$, then*

$$(3.1) \quad \|u\|_{\frac{np}{n-p}, \infty} \leq c_{n,p} \|\nabla u\|_p \quad \forall u \in W_0^{1,p}(\mathbb{M}^n).$$

with equality being approachable.

(ii) *If $p = n$, then*

$$(3.2) \quad \sup_{\lambda > 0} \lambda \left(\ln \left(\frac{|\Omega|}{|\{x \in \Omega : |u(x)| \geq \lambda\}|} \right) \right)^{\frac{1}{n}-1} \leq c_{n,n} \|\nabla u\|_n \quad \forall u \in W_0^{1,n}(\Omega).$$

with equality being approachable.

(iii) If $p \in (n, \infty)$, then

$$(3.3) \quad \|u\|_\infty \leq c_{n,p} |\Omega|^{\frac{p-n}{np}} \|\nabla u\|_p \quad \forall u \in W_0^{1,p}(\Omega).$$

with equality being approachable.

(iv) If $p = \infty$, then

$$(3.4) \quad \|u\|_\infty \leq c_{n,\infty} |\Omega|^{\frac{1}{n}} \|\nabla u\|_\infty \quad \forall u \in W_0^{1,\infty}(\Omega).$$

with equality being approachable.

Proof. In the sequel, we always assume that not only

$$\begin{cases} u \in W_0^{1,p}(\mathbb{M}^n) & \text{as } p \in [1, n) \text{ via letting } \Omega \rightarrow \mathbb{M}^n; \\ u \in W_0^{1,p}(\Omega) & \text{as } p \in [n, \infty), \end{cases}$$

but also

$$[u]_\lambda = \{x \in \mathbb{M}^n \text{ or } \Omega : |u(x)| \geq \lambda\} \text{ with } \lambda^{-1}|u(\cdot)| \geq 1_{[u]_\lambda} \quad \forall \lambda \in (0, \infty).$$

(i) Via choosing $\alpha = 1 - \frac{p}{n}$ within (2.1), we can obtain

$$\begin{aligned} c_{n,p}^{-p} |[u]_\lambda|^{1-\frac{p}{n}} &\leq \inf_{\Omega \subseteq \mathbb{M}^n} \text{cap}_p([u]_\lambda, \Omega) \\ &\leq \int_{\mathbb{M}^n} |\nabla(\lambda^{-1}|u|)|^p dv_g \\ &= \lambda^{-p} \int_{\mathbb{M}^n} |\nabla u|^p dv_g \end{aligned}$$

Hence (3.1) follows from

$$\lambda |[u]_\lambda|^{\frac{n-p}{np}} \leq c_{n,p} \|\nabla u\|_p \quad \forall \lambda > 0.$$

However, regarding the (3.1)'s sharpness in the sense that $c_{n,p}$ is approachable, we deal with two cases.

- If $1 < p < n$, then we construct the $(1, n) \ni p$ -capacity potential of a geodesic ball $\overline{B(o, r)}$ according to the formula

$$u_{p,r}(x) = \begin{cases} 1 & \text{as } x \in \overline{B(o, r)}; \\ 1 - \frac{\int_r^{d(o,x)} \varphi^{\frac{n-1}{1-p}}(s) ds}{\int_r^\infty \varphi^{\frac{n-1}{1-p}}(s) ds} & \text{as } x \in \mathbb{M}^n \setminus \overline{B(o, r)}, \end{cases}$$

where not only $d(o, \cdot)$ denotes the distance function of o but also the infinite integral is convergent due to $\varphi'' \geq 0$, thereby getting that not only

$$\|u_{p,r}\|_{\frac{np}{n-p}, \infty} \geq |[u_{p,r}]_1|^{\frac{1}{p}-\frac{1}{n}} = |\overline{B(o, r)}|^{\frac{1}{p}-\frac{1}{n}}$$

but also

$$\begin{aligned}
(3.5) \quad \|\nabla u_{p,r}\|_p^p &= \left(\int_r^\infty \varphi^{\frac{n-1}{1-p}}(s) ds \right)^{-p} \int_{\mathbb{M}^n} |\varphi^{\frac{n-1}{1-p}}(d(o,x)) \nabla d(o,x)|^p dv_g(x) \\
&= \left(\int_r^\infty \varphi^{\frac{n-1}{1-p}}(s) ds \right)^{-p} \int_r^\infty |\varphi^{\frac{n-1}{1-p}}(s)|^p n v_n \varphi^{n-1}(s) ds \\
&= \widehat{\text{cap}}_p(\overline{B(o,r)}).
\end{aligned}$$

Consequently, we achieve

$$\frac{\|u_{p,r}\|_{\frac{np}{n-p},\infty}}{\|\nabla u_{p,r}\|_p} \geq \frac{|\overline{B(o,r)}|^{\frac{n-p}{np}}}{(\widehat{\text{cap}}_p(\overline{B(o,r)}))^{\frac{1}{p}}} \rightarrow \left(n v_n^{\frac{p}{n}} \left(\frac{n-p}{p-1} \right)^{p-1} \right)^{-\frac{1}{p}} = c_{n,p} \text{ as } r \rightarrow 0$$

thereby finding that (3.1)'s equality for $p \in (1, n)$ is approachable.

- If $p = 1$, then we choose

$$u_{1,r}(x) = \begin{cases} 1 & \text{as } x \in \overline{B(o,r)}; \\ 1 + r^{-2}(r - d(o,x)) & \text{as } x \in \overline{B(o,r+r^2)} \setminus \overline{B(o,r)}; \\ 0 & \text{as } x \in \mathbb{M}^n \setminus \overline{B(o,r+r^2)}, \end{cases}$$

to calculate

$$\begin{aligned}
\liminf_{r \rightarrow 0} \frac{\|u_{1,r}\|_{\frac{n}{n-1},\infty}}{\|\nabla u_{1,r}\|_1} &\geq \lim_{r \rightarrow 0} \frac{[u_{1,r}]_1^{1-\frac{1}{n}}}{\int_{\mathbb{M}^n} |\nabla u_{1,r}| dv_g} \\
&= \lim_{r \rightarrow 0} \frac{|\overline{B(o,r)}|^{1-\frac{1}{n}}}{\int_r^{r+r^2} r^{-2} n v_n \varphi^{n-1}(t) dt} \\
&= \lim_{r \rightarrow 0} \frac{(v_n r^n)^{1-\frac{1}{n}}}{\int_r^{r+r^2} r^{-2} n v_n t^{n-1} dt} \\
&= n^{-1} v_n^{-\frac{1}{n}}.
\end{aligned}$$

This just says that (3.1)'s equality for $p = 1$ is approachable.

- (ii) According to (2.2), for any $\lambda > 0$ there holds

$$\begin{aligned}
\frac{|[u]_\lambda|}{|\Omega|} &\leq \exp\left(-n^{\frac{n}{n-1}} v_n^{\frac{1}{n-1}} (\text{cap}_n([u]_\lambda, \Omega))^{\frac{1}{1-n}}\right) \\
&\leq \exp\left(-n^{\frac{n}{n-1}} v_n^{\frac{1}{n-1}} \left(\int_\Omega |\nabla(\lambda^{-1}|u|)|^n dv_g\right)^{\frac{1}{1-n}}\right) \\
&= \exp\left(-n^{\frac{n}{n-1}} v_n^{\frac{1}{n-1}} \lambda^{-\frac{n}{1-n}} \left(\int_\Omega |\nabla u|^n dv_g\right)^{\frac{1}{1-n}}\right),
\end{aligned}$$

which is actually equivalent to (3.2).

In order to study the sharpness of (3.2), given a geodesic open ball $\Omega = B(o, R)$, we consider the n -capacitary potential $u_{n,r}$ of $(\overline{B(o, r)}, B(o, R))$ for some $r \in (0, R)$ given as below:

$$u_{n,r}(x) = \begin{cases} 1 & \text{as } x \in \overline{B(o, r)}; \\ 1 - \frac{\int_r^{d(o,x)} \varphi^{-1}(s) ds}{\int_r^R \varphi^{-1}(s) ds} & \text{as } x \in B(o, R) \setminus \overline{B(o, r)}. \end{cases}$$

In a similar way to establish (3.5), we have

$$\|\nabla u_{n,r}\|_n^n = \text{cap}_p(\overline{B(o, r)}, B(o, R)) = n\nu_n \left(\int_r^R \varphi^{-1}(s) ds \right)^{1-n},$$

whence estimating

$$\begin{aligned} \frac{\sup_{\lambda > 0} \lambda \left(\ln \frac{|B(o, R)|}{|[u_{n,r}]_\lambda|} \right)^{\frac{1}{n}-1}}{\|\nabla u_{n,r}\|_n} &\geq \frac{\left(\ln |B(o, R)| - \ln |\overline{B(o, r)}| \right)^{\frac{1}{n}-1}}{\left(\text{cap}_p(\overline{B(o, r)}, B(o, R)) \right)^{\frac{1}{n}}} \\ &= (n\nu_n)^{-\frac{1}{n}} \left(\frac{\ln |B(o, R)| - \ln |\overline{B(o, r)}|}{\int_r^R \varphi^{-1}(s) ds} \right)^{\frac{1}{n}-1} \\ &\rightarrow (n\nu_n)^{-\frac{1}{n}} \left(\frac{n\nu_n \varphi^n(R)}{|B(o, R)|} \right)^{\frac{1}{n}-1} \text{ as } r \rightarrow R \\ &\rightarrow n^{-1} \nu_n^{-\frac{1}{n}} \text{ as } R \rightarrow 0. \end{aligned}$$

Of course, this tells that (3.2)'s equality is approachable.

(iii) In accordance with (2.3), for

$$\begin{cases} 0 < \lambda < \|u\|_\infty; \\ [u]_\lambda \neq \emptyset; \\ |[u]_\lambda| > 0, \end{cases}$$

we can evaluate

$$\begin{aligned} c_{n,p}^{-p} |\Omega|^{1-\frac{p}{n}} &\leq \text{cap}_p([u]_\lambda, \Omega) \\ &\leq \int_\Omega |\nabla(\lambda^{-1}|u|)|^p dv_g \\ &= \lambda^{-p} \int_\Omega |\nabla u|^p dv_g. \end{aligned}$$

So, sending λ to $\|u\|_\infty$ in the above estimation gives

$$\|u\|_\infty \leq c_{n,p} |\Omega|^{\frac{p-n}{np}} \|\nabla u\|_p,$$

as desired in (3.3).

In order to reach (3.3)'s equality, let

$$\begin{cases} \Omega = B(o, R); \\ u_{p,r} = \text{the } p\text{-capacitary potential of } (\{o\}, B(o, R)) \text{ - i.e. -} \\ u_{p,r}(x) = 1 - \frac{\int_0^{d(o,x)} \varphi^{\frac{n-1}{1-p}}(s) ds}{\int_0^R \varphi^{\frac{n-1}{1-p}}(s) ds} \quad \forall x \in B(o, R). \end{cases}$$

Then

$$\begin{cases} \|u\|_\infty = 1; \\ \|\nabla u\|_p^p = nv_n \left(\int_0^R \varphi^{\frac{n-1}{1-p}}(s) ds \right)^{1-p}. \end{cases}$$

Needless to say, the last two formulas derive

$$\begin{aligned} |\Omega|^{\frac{p-n}{np}} \|\nabla u\|_p &= |B(o, R)|^{\frac{p-n}{np}} (nv_n)^{\frac{1}{p}} \left(\int_0^R \varphi^{\frac{n-1}{1-p}}(s) ds \right)^{\frac{1-p}{p}} \\ &\rightarrow (v_n R^n)^{\frac{p-n}{np}} (nv_n)^{\frac{1}{p}} \left((p-1)(p-n)^{-1} R^{\frac{p-n}{p-1}} \right)^{\frac{1-p}{p}} \\ &= c_{n,p}^{-1} \text{ as } R \rightarrow 0. \end{aligned}$$

Therefore, (3.3)'s equality is approachable.

(iv) Upon taking into account of (2.4), we see that the proof is analogous to that of verifying (iii) so it is omitted here. \square

Remark 3.2. *By tracing the proof process of Theorem 3.1, we find that if*

$$\begin{cases} \varphi(t) = t; \\ \mathbb{M}^n = \mathbb{R}^n; \\ 1 < p \leq \infty; \\ 0 < r < R < \infty, \end{cases}$$

the equalities within (3.1)-(3.2)-(3.3)-(3.4) can be achieved via the next functions

$$u_p(x) = \begin{cases} \left(\frac{\max\{|x|, r\}}{r} \right)^{\frac{p-n}{p-1}} \quad \forall x \in \mathbb{R}^n & \text{as } 1 < p < n; \\ \frac{\ln R - \ln \max\{|x|, r\}}{\ln R - \ln r} \quad \forall x \in B(o, R) & \text{as } p = n; \\ 1 - \left(\frac{|x|}{R} \right)^{\frac{p-n}{p-1}} \quad \forall x \in B(o, R) & \text{as } n < p < \infty; \\ 1 - \frac{|x|}{R} \quad \forall x \in B(o, R) & \text{as } p = \infty. \end{cases}$$

For certain of the closely-related results over \mathbb{R}^n , see also [3, 2, 1, 18].

4. APPLYING $\alpha = 1 > 1 - \frac{p}{n}$ TO PRINCIPAL p -FREQUENCIES

Theorem 4.1. *Let*

$$\begin{cases} (\mathbb{M}^n, g) = ([0, +\infty) \times \mathbb{S}^{n-1}, dt^2 + \varphi^2(t)g_{\mathbb{S}^{n-1}}) \text{ with } \varphi'' \geq 0; \\ \Omega \text{ obey the isoperimetry of } o = \{t = 0\}\text{-centered balls.} \end{cases}$$

Then

$$(4.1) \quad \lambda_{1,p}(\Omega) \geq \begin{cases} nv_n^{\frac{1}{n}} |\Omega|^{-\frac{1}{n}} & \text{as } p = 1; \\ nv_n^{\frac{p}{n}} p^{-p} (\max\{n, p-n\})^{p-1} |\Omega|^{-\frac{p}{n}} & \text{as } p \in (1, \infty); \\ v_n^{\frac{1}{n}} |\Omega|^{-\frac{1}{n}} & \text{as } p = \infty. \end{cases}$$

In particular, (4.1)'s case at $p = 1$ or $p = \infty$ is sharp in the sense that

$$\mathbb{M}^n = \mathbb{R}^n \text{ \& } \Omega = \mathbb{B}^n \Rightarrow \lambda_{1,1}(\Omega) = n \text{ \& } \lambda_{1,\infty}(\Omega) = 1.$$

Proof. The sharpness of (4.1) at $p \in \{1, \infty\}$ follows from [17, Proposition 3.1 & Remark 3.4]. Thus, it remains to verify (4.1).

First of all, recall such an equivalent property of the Maz'ya constant $\gamma_p(\Omega)$ that (cf. [13, 5])

$$(4.2) \quad \begin{aligned} \lambda_{1,p}(\Omega) &\leq \gamma_p(\Omega) \\ &\equiv \inf_{K \subset \subset \Omega} |K|^{-1} \text{cap}_p(K, \Omega) \\ &\leq \begin{cases} \lambda_{1,1}(\Omega) & \text{as } p = 1; \\ p^p (p-1)^{1-p} \lambda_{1,p}(\Omega) & \text{as } p \in (1, \infty). \end{cases} \end{aligned}$$

Next is a three-fold consideration.

- If $p \in [1, n)$, then using Theorem 2.1's case $\alpha = 1$, along with (4.2), derives

$$(4.3) \quad \lambda_{1,p}(\Omega) \geq \begin{cases} nv_n^{\frac{1}{n}} |\Omega|^{-\frac{1}{n}} & \text{as } p = 1; \\ (p-1)^{p-1} p^{-p} \gamma_p(\Omega) \geq nv_n^{\frac{p}{n}} (n-p)^{p-1} p^{-p} |\Omega|^{-\frac{p}{n}} & \text{as } p \in (1, n). \end{cases}$$

- However, thanks to [7, Theorem 1.2], we know that

$$[1, \infty) \ni p \mapsto p(\lambda_{1,p}(\Omega))^{\frac{1}{p}}$$

is an increasing function, whence getting

$$(4.4) \quad \lambda_{1,p}(\Omega) \geq \left(p^{-1} \lambda_{1,1}(\Omega)\right)^p \geq n^p v_n^{\frac{p}{n}} p^{-p} |\Omega|^{-\frac{p}{n}} \quad \forall p \in [1, \infty).$$

It is easy to see that (4.4) not only is stronger than (4.3)'s case $p \in (1, n)$ but also ensures (4.1) when $p \in (1, 2n]$.

- If $n < p < \infty$, then (2.3) tells us that not only

$$\gamma_p(\Omega) \geq \inf_{K \subset \subset \Omega} |\Omega|^{-1} \text{cap}_p(K, \Omega) \geq nv_n^{\frac{p}{n}} \left(\frac{p-n}{p-1}\right)^{p-1} |\Omega|^{-\frac{p}{n}},$$

but also

$$(4.5) \quad \lambda_{1,p}(\Omega) \geq (p-1)^{p-1} p^{-p} \gamma_p(\Omega) \geq nv_n^{\frac{p}{n}} (p-n)^{p-1} p^{-p} |\Omega|^{-\frac{p}{n}},$$

which ensures (4.1)'s case $p \in [2n, \infty)$.

Finally, a combination of (4.5) & the limiting formula (cf. [17, Remark 3.4] or [9, 10] for $\mathbb{M}^n = \mathbb{R}^n$)

$$\lambda_{1,\infty}(\Omega) = \lim_{p \rightarrow \infty} (\lambda_{1,p}(\Omega))^{\frac{1}{p}} = \lim_{p \rightarrow \infty} (\gamma_p(\Omega))^{\frac{1}{p}}$$

derives

$$\lambda_{1,\infty}(\Omega) \geq \lim_{p \rightarrow \infty} n^{\frac{1}{p}} v_n^{\frac{1}{n}} (1 - p^{-1}n)(p - n)^{-\frac{1}{p}} |\Omega|^{-\frac{1}{n}} = v_n^{\frac{1}{n}} |\Omega|^{-\frac{1}{n}},$$

as desired within (4.1)'s case $p = \infty$. □

REFERENCES

- [1] D.R. Adams & J. Xiao, Full capacity-volumetry of sharp exp-integrability law. Preprint (submitted) 23pages. [11, 16]
- [2] D. Cassani, B. Ruf & C. Tarsi, Group invariance and Pohozaev identity in Moser type inequalities. *Comm. Contemp. Math.* (2012).doi:10.1142/S021919971250054X. [16]
- [3] D. Cassani, B. Ruf & C. Tarsi, Optimal Sobolev type inequalities in Lorentz spaces. *Potential Anal.* 39(2013)265-285. [16]
- [4] V. Gol'dshtein & A. Ukhlov, Sobolev homeomorphisms and composition operators. Around the Research of Vladimir Maz'ya I: Function Spaces. *International Math. Series 11*, DOI 10.1007/978-1-4419-1341-8_8. pp. 207-220. Springer Science + Business Media, LLC2010. [11]
- [5] A. Grigor'yan, Isoperimetric inequalities and capacities on riemannian manifolds. In *The Maz'ya Anniversary Collection: Volume 1: On Maz'ya's work in functional analysis, partial differential equations and applications*, pp. 139-153. Springer, 1999. [7, 17]
- [6] R. Jiang, J. Xiao & D. Yang, Relative ∞ -capacity and its affinization. *Adv. Calc. Var.* 11(2018)95-110. [5, 10]
- [7] X. Jin, The lower bound of first dirichlet frequency of p -Laplacian in Riemannian manifolds. arXiv preprint arXiv:2405.02669, 2024. [17]
- [8] X. Jin & J. Xiao, Sharply estimating hyperbolic capacities. Preprint (submitted) 22 pages. [2, 11]
- [9] P. Jutila, P. Lindqvist & J. Manfredi, The ∞ -frequency problem. *Arch. Ration. Meh. Anal.* 148(1999)89-105. [17]
- [10] B. Kawohl & V. Fridman, Isoperimetric estimates for the first frequency of the p -Laplace operator and the Cheeger constant. *Commun. Math. Univ. Carol.* 44(2003)659-667. [17]
- [11] V.I. Kruglikov, Capacities of condensers and spatial mappings quasiconformal in the mean (Russian). *Mat. Sb.* 130(1986)185-206. [11]
- [12] V. Maz'ya, *Sobolev Spaces*. Springer-Verlag, Berline, 1985. [4, 7, 12]
- [13] V. Maz'ya. Lectures on isoperimetric and isocapacitary inequalities in the theory of sobolev spaces. *Contemp. Math.* 338(2003)307-340. [17]
- [14] G. Pólya & G. Szegő, Inequalities for the capacity of a condenser. *Amer. J. Math.* 67(1945)1-32. [1]
- [15] G. Pólya & G. Szegő, *Isoperimetric Inequalities in Mathematical Physics*. *Annals of Math. Studies*, No. 27, Princeton, Princeton Univ. Press, 1951. [3]
- [16] J. Sarvas, Symmetrization of condensers in n -space. *Ann. Acad. Sci. Fenn. Ser. AI* 522, 1-44. (1972) [2]
- [17] J. Xiao, The p -Faber-Krahn inequality noted. A. Laptev (ed.), Around the Research of Vladimir Maz'ya I: Function Spaces. *International Math. Series 11*, DOI 10.1007/978-1-4419-1341-8_17. pp. 373-390. Springer Science + Business Media, LLC2010. [17]
- [18] J. Xiao, Optimal weak-type estimates for Riesz potentials within geometric physics. Preprint (submitted) 24 pages. [16]

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND
TECHNOLOGY, WUHAN, HUBEI 430074, CHINA

Email address: jinxs@hust.edu.cn

DEPARTMENT OF MATHEMATICS & STATISTICS, MEMORIAL UNIVERSITY, ST. JOHN'S, NL
A1C 5S7, CANADA

Email address: jxiao@math.mun.ca