Subgroup Conjugacy Separability in Residually Free Groups

S. C. Chagas^{*1} and I. Kazachkov^{**2}

¹UnB - University of Brasília, Campus Universitário Darcy Ribeiro, Brasília-DF | CEP 70910-900, Brasil; email: sheilachagas@gmail.com ²Ikerbasque - Basque Foundation for Science and Matematika Saila, UPV/EHU, Sarriena s/n, 48940, Leioa - Bizkaia, Spain; email: ilya.kazachkov@gmail.com

Abstract

We prove that finitely presented residually free groups are subgroup conjugacy separable. Furthermore, if they are of type FP_{∞} , then they are also subgroup conjugacy distinguished.

Using a connection between conjugacy separability and residual finiteness of outer automorphism group established by Grossman in [Gr-75], we show that finitely presented residually free groups have residually finite outer automorphism groups.

1 Introduction

The three fundamental decision problems for groups: the word, conjugacy and the isomorphism problems, were formulated by Max Dehn in the early 1900's and since then have shaped entire areas of group theory.

One of the first important decidability results is due to A. I. Mal'cev, who observed a relation between decidability of the word problem and its residual properties, see [M-58]. He proved that if a finitely presented group is residually finite, then it has decidable word problem.

^{*}partially supported by Capes

^{**} partially supported by Basque Government grant IT1483-22

In order to apply Mal'cev's argument and solve the conjugacy problem a stronger residual property is required. A group G is said to be conjugacy separable if one can distinguish its conjugacy classes by looking at finite quotients of G. If a finitely presented group is conjugacy separable, then it has decidable conjugacy problem.

Another natural generalisation of residually finiteness that extends from elements to subgroups, is subgroup separability. A group G is subgroup separable if one can distinguish its finitely generated subgroups by looking at its finite quotients. This property is more general and assures the decidability of the membership problem in G.

Just as conjugacy separability generalises residual finiteness, the notion of subgroup conjugacy separability introduced by O. Bogopolski and F. Grunewald in [BG-10] generalises the notion of conjugacy separability. A group G is said to be subgroup conjugacy separable if for every pair of non-conjugate finitely generated subgroups H and K of G there exists a finite quotient of G where the images of these subgroups are not conjugate.

Finitely presented subgroup conjugacy separable groups have solvable conjugacy problem for finitely generated subgroups. The latter means, that there is an algorithm which given a finitely presented subgroup conjugacy separable group $G = \langle X | R \rangle$ and two finite sets of elements Y and Z decides whether or not the subgroups $\langle Y \rangle$ and $\langle Z \rangle$ are conjugate in G.

In this paper we study subgroup conjugacy separability in residually free groups. Residually free groups is a prominent class of groups that provides a context for a rich and powerful interplay among group theory, topology and logic.

By definition, a residually free group embeds in a direct product of (perhaps, infinitely many) free groups. For example, the fundamental group of a closed orientable surface is residually free, but it cannot be embedded in a finite direct product if it has negative Euler characteristic. However, it is shown in [BMR-99], see also [BHMS-07] that one can embed finitely generated residually free groups into a finite product of fully residually free groups.

In turn, fully residually free groups have been extensively studied in connection with Tarski?s problems on the first order logic of free groups and they appear under different guises: as groups having the same existential theory as free groups, as coordinate groups of irreducible varieties as well as so-called limit groups.

A large body of work on the structure of residually free groups culminated in an important paper of Bridson-Howie-Miller-Short [BHMS-07]. One of the outcomes of their work is a stark contrast between finitely generated residually free groups and residually free groups that satisfy higher finiteness properties. For instance, while the membership and conjugacy problem is undecidable for finitely generated residually free groups, they are decidble for finitely presented ones, see [BHMS-07].

The results of this paper constitute a further step in this direction and can be regarded as a generalisation of the aforementioned result from [BHMS-07] as we prove

Theorem. Let G be a finitely presented residually free group. Then G is finitely presented subgroup conjugacy separable.

Moreover, we show that residually free group G of type FP_{∞} are also subgroup conjugacy distinguished, see Definition 2.2 and Theorem 2.4 for the precise statement.

Note that if G is residually finite and finitely presented, then for any finitely generated conjugacy distinguished subgroup H of G there exists an algorithm that decides if a given element g of G is conjugate to some element of H.

Finally, we establish residual finiteness of the outer automorphism group of a finitely presented residually free group, see Corollary 4.3. In order to do so we use a result of Grossman, see [Gr-75], who proved that if G is a finitely generated conjugacy separable group such that all its pointwise inner automorphisms are inner, then Out(G) is residually finite; we establish this for finitely generated residually free groups.

2 Residually free groups of type FP_{∞}

In this section we study conjugacy distinguished subgroups of residually free groups. We first prove that virtual retracts of hereditarily conjugacy separable groups are conjugacy distguished and then show that subgroups of type FP_m (for large enough m) over \mathbb{Q} of a finitely presented residually free group are conjugacy distinguished.

We say that a group G is hereditarily conjugacy separable if every finite index subgroup of G is conjugacy separable.

Recall that a subgroup H of a group K is called a retract if there is a homomorphism $\rho : K \to H$ which restricted to H is the identity map. This is equivalent to K splitting as a semidirect product $N \rtimes H$, where $N = ker \rho$. In this case the map ρ is called a retraction of K onto H. **Definition 2.1.** Let G be a group and let H be a subgroup of G. We say that H is a virtual retract of G if there exists a subgroup $K \leq G$ such that $|G:K| < \infty$, $H \subseteq K$ and H is a retract of K.

Given a group G, the profinite topology on G has as a basis all the cosets of finite index subgroups of G. Each such coset is both open and closed in the profinite topology. While the separability properties of G can be described in terms of the finite quotients of G, it is sometimes more convenient to talk about the profinite topology instead. For instance, G is called residually finite if for any $1 \neq g \in G$, there exists a finite quotient of G in which g does not map to the identity element. In other words, there is a finite index normal subgroup N of G such that $g \notin N$. Equivalently, this means that $\{1\}$ is closed in the profinite topology of G. In fact, this is the same as saying that any one element subset of G is closed or, the seemingly stronger statement, that G is Hausdorff.

The group G is called subgroup separable if for every finitely generated subgroup H of G and every $g \in G \setminus H$ there exists a normal subgroup of finite index, N of G such that $g \notin HN$. Thus, there is a finite quotient of G in which the images of g and H are disjoint. As before, this is the same as saying that every finitely generated subgroup is closed in the profinite topology of G. More generally, a subset $S \subseteq G$ is called separable if for every $g \notin S$, there is a finite quotient of G in which g and S have disjoint images. Equivalently, S is separable if it is closed.

A group G is called conjugacy separable if the conjugacy classes of elements are separable. If we have a subgroup H of G, then there are two possible topologies one can put on H. Namely, the subspace topology and the profinite topology of H itself. In general the subspace topology may be more coarse, but not if H has finite index.

In the above notation, denote by \widehat{G} the profinite completion of G and by \overline{H} the closure of H in \widehat{G} .

Definition 2.2. A group G is called subgroup conjugacy distinguished if for any finitely generated subgroup H of G and any element $x \in G$ such that x is not conjugate into H there exists a finite quotient of G where the image of x is not conjugate into the image of H.

A subgroup H of a group G is said to be conjugacy distinguished if $\cup_{g \in G} H^g$ is closed in the profinite topology of G. Equivalently, H is conjugacy distinguished if whenever y is not conjugate to an element of H, there exists a finite quotient of G where the image of y is not conjugate to an element of the image of H.

Theorem 2.3. Let G be a hereditarily conjugacy separable group, and H a virtual retract of G. Then H is conjugacy distinguished.

Proof. To prove the theorem, we shall replace H and g by their conjugates in G and change γ accordingly until we achieve the statement of the theorem holding for them.

Since H is a virtual retract of G, there exist a finite index subgroup U of G containing H and a homomorphism $f: U \to H$ such that $f_{|_H}$ is the identity map. Moreover, $G\overline{U} = \widehat{G}$.

Hence replacing g by some conjugate in G we may assume that γ belongs to \widehat{U} . Since $g^{\gamma} \in \overline{H}, \overline{H} \leq \overline{U}$ and U is closed in the profinite topology of G we obtain $g \in \widehat{U} \cap G = U$.

Let $\widehat{f} : \widehat{U} \to \overline{H}$ be the continuous extension f. We have, $g^{\gamma} \in \overline{H}$, so $g^{\gamma} = \widehat{f}(g^{\gamma}) = f(g)^{\widehat{f}(\gamma)} \in \widehat{f}(\overline{H}) = \overline{H}$. Then, g and f(g) are conjugate in \overline{U} . Since U is conjugacy separable by hypothesis, g and f(g) are conjugate in U and lemma is proved.

Theorem 2.4. Let G be a finitely presented residually free group. Then there exists m such that all subgroups of G of type FP_m over \mathbb{Q} are conjugacy distinguished.

Proof. Let G be a finitely presented residually free group. By [BHMS-07, Corollary 19], G embeds in a direct product of finitely many limit groups $L = \prod_{i=1}^{n} L_i$. Choose minimal such n. Without loss of generality we may assume that G is a subdirect product of the L_i 's.

Let H be a FP_n subgroup of G. By [BW-08, Lemma 7], H is a virtual retract of L. Then H also is a virtual retract of G. By [CZ-09, Theorem 3.5], G is conjugacy separable. Hence, by Theorem 2.3, H is conjugacy distinguished in G.

3 Finitely presented residually free groups

The goal of this section is to prove that finitely presented residually free groups are finitely presented subgroup conjugacy separable.

Definition 3.1. Let G_1, \ldots, G_n be groups. A subgroup $H \le G_1 \times \cdots \times G_n$ is called full if H intersects each factor non-trivially, i.e. $H \cap G_i \ne 1$ for all $i \in \{1, 2, \ldots, n\}$.

Notice that if H is full, in particular it is nontrivial and if H intersects trivially one of the factors, say $H \cap G_1 = 1$, then H is isomorphic to a subgroup of $G_2 \times \cdots \times G_n$.

Proposition 3.2. Let G_1, G_2 be finitely presented subgroups of the direct product $L = \prod_{i=1}^{n} L_i$, where L_i is a limit group. If G_1 and G_2 are conjugate in \hat{L} , then they are conjugate in L.

Proof. Suppose that G_1 is not full in L, say $G_1 \cap L_1 = 1$ and assume that G_2 is full. Consider $L' = L/L_1$ and note that it is a subgroup of L. So $G_1 \cong G_1 L_1/L_1 \leq L'$ and since any finitely presented subgroup of L' is separable, $\widehat{G}_1 \cong \widehat{G_1 L_1/L_1} \leq \widehat{L'}$. Hence $\widehat{L}_1 \cap \widehat{G}_1 = 1$ by hopfian property of a finitely generated profinite group.

By assumption, there exists $\gamma \in \hat{L}$ so that $\hat{G}_1^{\gamma} = \hat{G}_2$. It follows that

$$\widehat{G_1L_1/L_1}^{\gamma} = \widehat{G_2L_1/L_1} \cong \widehat{G}_2/\widehat{G}_2 \cap \widehat{L}_1$$

are conjugate, and so isomorphic. Therefore $\widehat{G}_1 \cong \widehat{G}_2/\widehat{G}_2 \cap \widehat{L}_1$. Since a finitely generated profinite group is hopfian, it follows that $\widehat{L}_1 \cap \widehat{G}_2 = 1$. In particular we have $L_1 \cap G_2 = 1$. Thus, G_1 is full if and only if so is G_2 . Hence, factoring out L_i that intersect G_1 and G_2 trivially we may assume that G_1 and G_2 intersect all L_i non-trivially.

Let π_i be the natural projection from L onto L_i . Observe that the images $\pi_i(\hat{G}_1)$ and $\pi_i(\hat{G}_2)$ are conjugate in \hat{L}_i , since L_i is subgroup conjugacy separable by [CZ-16]; $\pi_i(G_1)$ and $\pi_i(G_2)$ are conjugate in L_i , i.e. $\pi_i(G_1)^{l_i} = \pi_i(G_2)$ for some $l_i \in L_i$. Thus, conjugating G_1 by the element (l_1, \ldots, l_n) if necessary, we may assume that $\pi_i(G_1) = \pi_i(G_2) = N_i$ for all i.

Now observe that $\pi_i(G_j)$ is a finitely generated subgroup of L_i . By the main result of [W-08], L_i has a virtual retract, so there exist a finite index K_i that contained G_i and a homomorphism $\psi : K_i \to \pi_i(G_j)$ such that $\psi_{|_{\pi_i(G_j)}}$ is the identity map. Thus, substituting L_i by K_i we may assume that $L_i = K_i$, and then N_i is a retract of L_i for every i.

By [BW-08, Theorem 8], the finitely presented subgroups G_1 and G_2 contain some term of the lower central series $\gamma_m(L)$, then $\gamma_m(L) \subseteq G_1 \cap G_2$. Hence $\gamma_m(\hat{L}) \subseteq \hat{G}_1 \cap \hat{G}_2$.

Now in the quotient $\widehat{L}/\gamma_m(\widehat{L})$ the images of $\widehat{G}_1/\gamma_m(\widehat{L})$ and $\widehat{G}_2/\gamma_m(\widehat{L})$ are conjugate. By Theorem 7 in Chapter 4 of [S-83], $G_1/\gamma_m(L)$ and $G_2/\gamma_m(L)$ are conjugate in $L/\gamma_m(L)$. It follows that G_1 and G_2 are conjugate in L.

We denote by $N_L(G)$ the normalizer of G in L, that is

$$N_L(G) = \{ l \in L \mid lGl^{-1} = G \}.$$

Lemma 3.3. Let $L = \prod_{i=1}^{n} L_i$ be the direct product of limit groups and G be a full finitely presented subgroup of L. Then the normalizer $N_L(G)$ is dense in $N_{\widehat{L}}(\widehat{G})$.

Proof. By [CZ-16, Theorem 2.5], L_i is hereditarily subgroup conjugacy separable and hence, by [BZ-24, Lemma 2.3], $N_{L_i}(\pi_i(G))$ is dense in $N_{\widehat{L}_i}(\pi_i(\widehat{G}))$. Therefore replacing L_i by $N_{L_i}(\pi_i(G))$ we may assume that $\pi_i(G)$ is normal in L_i . Then, by [BH-07, Theorem 1], either the index of $\pi_i(G)$ in L_i is finite or L_i is abelian. Put $K = \prod_{i=1}^n \pi_i(G)$. By [BW-08, Thereom 8], $G \ge \gamma_m(K)$. Since $\gamma_m(\overline{K}) \le \widehat{G} \le \overline{N_L(G)}$, it suffices to show that $N_{L/\gamma_m(K)}(G/\gamma_m(K))$ is dense in $N_{\overline{L/\gamma_m(K)}}(\overline{G/\gamma_m(K)})$. But this is the subject of [RSZ-98, Proposition 3.3]. The lemma is proved.

Theorem 3.4. Let G be a finitely presented residually free group. Then G is finitely presented subgroup conjugacy separable.

Proof. By [BHMS-07, Corollary 19], G embeds in a direct product of finitely many limit groups $L = \prod_{i=1}^{n} L_i$. Choose minimal such n. As we argued in the proof of Proposition 3.2, without loss of generality we may assume that G is a subdirect product of the L_i 's.

Let G_1, G_2 be finitely presented subgroups of G such that $\overline{G}_1^{\gamma} = \overline{G}_2$, for some $\gamma \in \widehat{G}$. By Theorem 3.2, G_1 and G_2 are conjugate in L, i.e. there exist $l \in L$ such that $G_1^l = G_2$. Hence, $\delta = \gamma l^{-1} \in N_{\widehat{G}}(\widehat{G}_1)$. Then, $l = \delta^{-1}\gamma \in N_{\widehat{G}}(\widehat{G}_1)\widehat{G}$.

Now by [BW-08, Theorem 8], $G \ge \gamma_m(L)$ for some m. Hence $N_G(G_1)G \ge \gamma_m(L)$. Observe that $N_G(G_1)G/\gamma_m(L)$ is closed in the profinite topology of $L/\gamma_m(L)$ (see [LW-77]). Therefore $N_G(G_1)G$ which is the preimage of $N_G(G_1)G/\gamma_m(L)$ in L is closed in the profinite topology of L.

By Lemma 3.3, $N_G(G_1)$ is dense in $N_{\widehat{G}}(\widehat{G}_1)$. Therefore $\overline{N_G(G_1)G} = N_{\widehat{G}}(\widehat{G}_1)\widehat{G}$. Thus $N_{\widehat{G}}(\widehat{G}_1)\widehat{G} \cap L = N_G(G_1)G$, so $l \in N_{\widehat{G}}(\widehat{G}_1)\widehat{G} \cap L$ and also $l \in N_G(G_1)G$. Then we may write $l = ng_0$, where $n \in N_G(G_1)$ and $g_0 \in G$. Therefore $G_1^{g_0} = G_2$ as desired.

Corollary 3.5 (cf. [BW-08]). *Finitely presented residually free groups are conjugacy separable.*

Proof. Let G be a finitely presented residually free group. From the definition of being residually free, it follows that any 2-generated subgroup of G is either free or free abelian. Hence, in particular G does not contain non-abelian Baumslag-Solitar groups.

It follows that $\langle x \rangle$ and $\langle y \rangle$ are conjugate if and only if either so are x and y or x and y^{-1} and the statement follows.

4 Outer automorphisms of residually free groups

In this section we first show that every commensurating endomorphism of a finitely generated residually free group is an inner automorphism. We combine this fact with a result of Grossman's to conclude that the outer automorphism group of a finitely presented residually free group is residually finite. Our arguments are adjustments of [AMS16, Section 9 and Theorem 1.6].

Let H be a subgroup of a group G and let $\varphi \colon H \to G$ be a homomorphism. Then φ is *commensurating* if for all $h \in H$ there are $z \in G$ and $n, m \in \mathbb{Z} \setminus \{0\}$ such that $h^n = z\varphi(h)^m z^{-1}$.

We record the following observation.

Remark 4.1. If *H* is a free abelian group then the only commensurating endomorphisms of *H* are endomorphisms of the form $h \mapsto h^s$ for some $s \in \mathbb{Z} \setminus \{0\}$ and for all $h \in H$.

Theorem 4.2. Let H be a finitely generated non-abelian residually free group. Then every commensurating endomorphism $\varphi \colon H \to H$ is an inner automorphism of H.

Proof. Let H be a residually free group and let H < G, where G is the direct product of limit groups. Since subgroups of limit groups are again limit groups, without loss of generality, we further assume that H is a subdirect product of G, see [BHMS-07].

Let $G = L_0 \times L_1 \times \cdots \times L_l$ be the standard factorization of G, where L_0 is the center of G and L_1, \ldots, L_l are all non-abelian limit groups, see [BMR-99, BHMS-07]. Observe that L_0 is a finitely generated free abelian group and $l \ge 1$ as H is non-abelian. Let $\pi_i \colon G \to L_i$ denote the canonical retraction, $i = 0, 1, \ldots, l$. We have $\pi_i(H) = L_i$.

Observe that $N_i := H \cap L_i \triangleleft H$ is non-trivial whenever $i = 1, \ldots, l$, because otherwise H would embed into the direct product of $L_0 \times L_1 \times L_{i-1} \times L_{i+1} \times \cdots \times L_l$, see [BHMS-07]. Moreover, N_i is non-abelian and hence there is an element $h_i \in N_i \setminus \{1\}$ such that $E_{L_i}(h_i) = \langle h_i \rangle \subseteq N_i$, $i = 1, \ldots, l$, where

$$E_G(h) = \left\{ x \in G \mid xh^k x^{-1} = h^l \text{ for some } k, l \in \mathbb{Z} \setminus \{0\} \right\}.$$

Suppose that $xh^kx^{-1} = h^l$ in a residually free group H. Let ψ be a homomorphism to a free group F so that $\psi(xh^kx^{-1}) = \psi(h^l) \neq 1$ (and

so $\psi(h) \neq 1$). In a free group $yg^k y^{-1} = g^l$ implies k = l, so the same is true in any residually free group. It follows that if H is residually free, then $E_H(h) = C_H(h)$.

Now consider any commensurating endomorphism $\varphi \colon H \to H$. For each $i \in \{0, 1, \ldots, l\}$ let $B_i \triangleleft G$ denote the product of all L_j , $j \neq i$; thus $G = L_i B_i \cong L_i \times B_i$ and $B_i = \ker \pi_i$. By the hypothesis, for any $g \in H \cap B_i, \varphi(g) \in H$ and $\varphi(g)^m = ug^n u^{-1} \in B_i$ for some $m, n \in \mathbb{Z} \setminus \{0\}$ and $u \in G$. And since $G/B_i \cong L_i$ is torsion-free, we can conclude that $\varphi(g) \in B_i$. The latter shows that φ preserves the kernel of the restriction of π_i to $H, i = 0, 1, \ldots, l$. Therefore φ naturally induces an endomorphism $\varphi_i \colon \pi_i(H) \to \pi_i(H)$ for $i = 0, 1, \ldots, l$, defined by the formula $\varphi_i(\pi_i(g)) = \pi_i(\varphi(g))$ for all $g \in H$.

Evidently, φ_i is a commensurating endomorphism of $\pi_i(H)$ for each i = 0, 1, ..., l. Therefore, by Remark 4.1, there exists $s \in \mathbb{Z} \setminus \{0\}$ such that $\varphi_0(a) = a^s$ for all $a \in \pi_0(H)$. On the other hand, if $i \neq 0$, then L_i is a non-abelian limit group and $\pi_i(H) = L_i$. Therefore there exists $w_i \in L_i$ such that $\varphi_i(a) = w_i a w_i^{-1}$ for all $a \in \pi_i(H)$ (here we used the fact that $E_{L_i}(\pi_i(H)) = \bigcap_{h \in \pi_i(H)} E_{L_i}(h) = \bigcap_{h \in \pi_i(H)} C_{L_i}(h) = \{1\}$), i = 1, ..., l.

Let $\psi \in Inn(G)$ be the inner automorphism defined by $\psi(g) = wgw^{-1}$ for all $g \in G$, where $w = w_1 \cdots w_l \in G$. Let us show that the endomorphism φ is actually the restriction of ψ to H. The preceding paragraph implies that this is true if the abelian factor L_0 is trivial, because in this case for every $g \in H$ one would have $g = \pi_1(g) \cdots \pi_l(g)$, and so

$$\varphi(g) = \pi_1(\varphi(g)) \cdots \pi_l(\varphi(g)) = \varphi_1(\pi_1(g)) \cdots \varphi_l(\pi_l(g)) = \\ = \pi_1(g)^{w_1} \cdots \pi_l(g)^{w_l} = g^w.$$

On the other hand, if L_0 is non-trivial, then $N_0 = H \cap L_0$ is also non-trivial (by the minimality of G). So, pick any $h_0 \in N_0 \setminus \{1\}$. Let $h_1 \in N_1 = H \cap L_1$ be the element constructed above. Since φ is commensurating and $h_0h_1 \in H$, there must exist $m, n \in \mathbb{Z} \setminus \{0\}$ and $u \in H$ such that

$$\varphi(h_0h_1)^m = u(h_0h_1)^n u^{-1} = h_0^n u_1 h_1^n u_1^{-1}$$
, where $u_1 = \pi_1(u) \in L_1$.

But we also have $\varphi(h_0h_1) = \varphi_0(h_0)\varphi_1(h_1) = h_0^s w_1 h_1 w_1^{-1}$. Therefore

$$h_0^{sm}w_1h_1^mw_1^{-1} = h_0^nu_1h_1^nu_1^{-1}.$$

Applying π_0 and π_1 to the above equation we obtain $h_0^{sm} = h_0^n$ and $u_1^{-1}w_1h_1^mw_1^{-1}u_1 = h^n$. The former yields that n = sm; and the latter shows that $u_1^{-1}w_1 \in E_{L_1}(h_1) = \langle h_1 \rangle$, in particular this element commutes with h_1 . Thus $h_1^m = h_1^n$, and so m = n. Consequently, s = 1, which

implies that $\varphi(g) = wgw^{-1} = \psi(g)$ for all $g \in H$. If $w \in H$ then the proof would have been finished. However, this may not be the case, so one more step is needed.

Let $h_i \in N_i = H \cap L_i$, i = 1, ..., l, be the elements constructed above so that $E_{L_i}(h_i) = \langle h_i \rangle \subseteq H$, and set $h = h_1 \cdots h_l \in H$. By the assumption, there exist $m, n \in \mathbb{Z} \setminus \{0\}$ and $u \in H$ such that $\varphi(h)^m = uh^n u^{-1}$. On the other hand, we know that $\varphi(h) = whw^{-1}$. Combining these two equalities one gets $wh^m w^{-1} = uh^n u^{-1}$ in *G*. Applying π_i yields that $u_i^{-1}w_i \in E_{L_i}(h_i) = \langle h_i \rangle$, where $u_i = \pi_i(u) \in L_i$, for i = 1, ..., l. It follows that for every i = 1, ..., l, there exists $t_i \in \mathbb{Z}$ such that $w_i = u_i h_i^{t_i}$ in L_i . Thus, denoting $u_0 = \pi_0(u) \in L_0$, we achieve

$$w = w_1 \cdots w_l = u_1 h_1^{t_1} \cdots u_l h_l^{t_l} = u_0^{-1} u h_1^{t_1} \cdots h_l^{t_l} = u_0^{-1} v,$$

where the element $v = uh_1^{t_1} \cdots h_l^{t_l}$ belongs to H by construction. Since $u_0 \in L_0$ is central in G, we see that $\varphi(g) = wgw^{-1} = vgv^{-1}$ for all $g \in H$, thus φ is indeed an inner automorphism of H.

An automorphism $\alpha \in Aut(G)$ is said to be *pointwise inner* if $\alpha(g)$ is conjugate to g for each $g \in G$. The set of all pointwise inner automorphisms, $Aut_{pi}(G)$ is a normal subgroup of Aut(G).

In [Gr-75] Grossman proved that if G is a finitely generated conjugacy separable group such that $Aut_{pi}(G) = Inn(G)$ then Out(G) is residually finite. We thus arrive at the following

Corollary 4.3. Let G be a finitely presented residually free group. Then Out(G) is residually finite.

References

- [AMS16] Antolín, Y., Minasyan, A., Sisto, A., Commensurating endomorphisms of acylindrically hyperbolic groups and applications. Groups Geom. Dyn. 10 (2016), no. 4, pp. 1149-1210.
- [BG-10] Bogopolski, O., Grunewald, F. On subgroup conjugacy separability in the class of virtually free groups, Max-Planck-Institute of Mathematics Preprint Series, n. 110 (2010), 18 pages.
- [BMR-99] Baumslag, B., Myasnikov, A., and Remeslennikov, V., Algebraic geometry over groups. I. Algebraic sets and ideal theory, J. Algebra, 219 (1999) 402-418.

- [BZ-24] M. Boggi, P. Zalesskii, *Finite subgroups of the profinite completion of good groups*, Bulletin of the London Mathematical society 57 (2025) 236-255.
- [BH-07] Bridson, M.R., Howie, J. Normalisers in limit groups. Math. Ann. 337, (2007) 385-394.
- [BHMS-07] Bridson, M. R., Howie J., Miller S.F., Short, H., Subgroups of direct products of limit groups, Ann. Math. v.170 (2009) 1447-1467
- [BW-08] Bridson, M. R., Wilton, H., Subgroup separability in residually free groups, Math. Z., **260** (2008) 25-30.
- [CZ-09] Chagas, S. C., Zalesskii, P. A., *Finite index subgroups of conjugacy separable groups*, Forum Math., **21** (2009) 347-353.
- [CZ-16] Chagas, S. C., Zalesskii, P. A., Limit groups are subgroup conjugacy separable. J. Algebra 461 (2016) 121-128.
- [Gr-75] E. Grossman, On the residual finiteness of certain mapping class groups. J. London Math. Soc. 9 (1975/75), No. 2, pp. 160-164.
- [M-58] Mal'cev, A. I., On Homomorphisms onto Finite Groups, Uchen. Zap. Ivanovskogo Gos. Ped. Inst., 18 (1958) 40-60.
- [RSZ-98] Ribes, L., Segal, D., Zalesskii, P. A., Conjugacy Separability and Free Products of Groups with Cyclic Amalgamation, Journal of the London Mathematical Society, 57, Issue3 (1998) 609-628
- [S-83] Segal, D., Polycyclic groups, Cambridge University Press, 1983.
- [LW-77] Lennox, J. C., Wilson, J. S., A note on permutable groups, Arch. Math. (Basel) 28 (1977) 113-116.
- [W-08] Wilton, H. Hall's Theorem for Limit Groups, GAFA Geom. funct. anal. 18, 271-303 (2008).