

FOOTBALLS OF CONSTANT CURVATURE ONE IN \mathbb{E}^3

ZHIQIANG WEI

ABSTRACT. In this paper, we construct a family of surfaces with constant curvature one, each featuring two conical singularities, and map them from football metrics into \mathbb{E}^3 . Additionally, we provide visualizations of these surfaces, revealing a novel geometric representation of a football in \mathbb{E}^3 .

Key words Constant curvature one, Conical singularities, Isometric immersion, Euclidean space.

2020 MR Subject Classification: 53A05, 53C42.

1. INTRODUCTION

It is a classical problem in differential geometry to determine when a Riemannian surface (M^2, g) can be isometrically immersed or embedded into 3-dimensional Euclidean space \mathbb{E}^3 , and whether such an isometric realization is unique up to rigid motions in \mathbb{E}^3 . On the other hand, the class of conformal metrics of constant Gaussian curvature 1 with conical singularities on a Riemann surface has been extensively studied, both from a local and global perspectives. For a comprehensive survey on this topic, we refer the reader to Eremenko [6]. For further details on the existence of constant curvature one metrics with conical singularities, readers may consult [1, 2, 3, 4, 5, 7, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20] and references cited therein. It is hence natural to investigate the isometric immersion problem for such metrics of constant curvature with conical singularities.

A metric of constant curvature one with two conical singularities on S^2 referred to as a football metric or simply a football if the distance between the two singularities equals π . The existence of constant curvature one metric with two conical singularities on S^2 was established by Troyanov in [17]. Specifically, a constant curvature one metric with two conical singularities on S^2 exists if and only if the two conical angles are identical. The shape of a football with conical angles in $(0, 2\pi)$ embedded in \mathbb{E}^3 is a rotationally symmetric surface, as described by Gálvez, Hauswirth and Mira in [8]. However, when the conical angles exceed 2π , it remains unknown whether other shapes of the football exist. In this paper, we demonstrate the existence of a family of isometric immersions from each football into \mathbb{E}^3 , revealing a novel geometric representation of a football in \mathbb{E}^3 . Our main result is as follows.

Theorem 1.1. *Let $\alpha > 0$ be a real number. Define the following coordinate functions:*

$$\begin{cases} x^1(u, \theta) = B \sin u \cos(\lambda\theta), \\ x^2(u, \theta) = B \sin u \sin(\lambda\theta), \\ x^3(u, \theta) = \int_0^u \sqrt{1 - B^2 \cos^2 t} dt, \end{cases}$$

where $(u, \theta) \in (0, \pi) \times [0, 2\pi)$, $0 < B < 1$ and $\lambda = \frac{\alpha}{B} \in \mathbb{Z}^+$. Regard S^2 as $\mathbb{C} \cup \{\infty\}$ and consider the map $F : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{E}^3$ defined by $z = re^{\sqrt{-1}\theta} \mapsto F(r, \theta)$, where

$$F(r, \theta) = (x^1(2 \arctan r, \theta), x^2(2 \arctan r, \theta), x^3(2 \arctan r, \theta)).$$

This map is an isometric immersion, and the pull back $F^*ds_E^2$ of the Euclidean metric ds_E^2 induces a football metric on S^2 with conical angles $2\pi\alpha$ both at points 0 and ∞ .

Remark 1.1. *When $\alpha \in (0, 1)$ and $\lambda = 1$, F is an embedding and the resulting surface is rotationally symmetric. However, when $\lambda \neq 1$, the surface loses its rotational symmetry. When $\alpha > 1$, since $2 \leq \lambda = \frac{\alpha}{B} \in \mathbb{Z}^+$, F fails to be embedding.*

It is well-known that solving the Gauss and Codazzi equations is essential for addressing isometric immersion problems in \mathbb{E}^3 . If the underlying surface is simply connected and the Gauss and Codazzi equations admit a solution, the existence of an isometric immersion can be established. However, obtaining an explicit expression for the immersion is generally challenging. Motivated by the form of rotationally symmetric surfaces, we successfully derive the explicit expression presented in **Theorem 1.1**.

This paper is organized as follows. In Sect.2, we review the classification of constant curvature one with two conical singularities on S^2 and provide the explicit expressions for football metrics. In Sect.3, we present the proof of **Theorem 1.1**. In Sect.4, we elaborate on the derivation of the immersion expressions stated in Sect.3. Finally, in Sect.5, we illustrate several visualizations of footballs in \mathbb{E}^3 for various parameter values.

2. PRELIMINARY

Definition 2.1. ([18]) *Let M be a Riemann surface, and let $p \in M$. A conformal metric ds^2 on M is said to have a conical singularity at p with the singular angle $2\pi\alpha$ ($\alpha > 0, \alpha \neq 1$) if, in a neighborhood of p ,*

$$ds^2 = e^{2\varphi} |dz|^2,$$

where z is a local complex coordinate defined in the neighborhood of p with $z(p) = 0$ and

$$\varphi - (\alpha - 1) \ln |z|$$

is continuous at 0.

In [17], Troyanov classified the constant curvature one metric with two conical singularities on S^2 . Recently, Wei, Wu and Xu [20] provided a new proof of this classification.

Theorem 2.1. ([17],[20]) *Regard S^2 as $\mathbb{C} \cup \{\infty\}$. Let ds^2 be a spherical conical metric on $\mathbb{C} \cup \{\infty\}$ with conical singularities at $z = 0$ and $z = \infty$, each with conical angles $2\pi\alpha$ and $2\pi\alpha$ respectively, where $0 < \alpha \neq 1$. Then, up to a change of coordinate $z \mapsto pz$, where $p \in \mathbb{C} \setminus \{0\}$ is a constant, the metric can be expressed as follows.*

- (1) *If $\alpha \notin \mathbb{Z}^+$, the metric is given by $ds^2 = \frac{4\alpha^2|z|^{2(\alpha-1)}}{(1+|z|^2)^2}|dz|^2$.*
- (2) *If $\alpha \in \mathbb{Z}^+$, the metric is given by $ds^2 = \frac{4\alpha^2|z|^{2(\alpha-1)}}{(1+|z^\alpha+b|^2)^2}|dz|^2$, where $b \in \mathbb{R}$ is a constant.*

In [20], Wei, Wu and Xu introduced the definition of a football metric, which was used to construct constant curvature one metric with finitely many conical singularities on compact Riemann surfaces.

Definition 2.2. ([20]) *Let ds^2 be a constant curvature one metric with two conical singularities on a 2-sphere. If the distance between these two singularities is π , we define ds^2 as a football metric. Furthermore, a 2-sphere equipped with a football metric is referred to as a football.*

By direct calculation, one can easily obtain the following lemma:

Lemma 2.1. *Let ds^2 be a football metric on S^2 with conical angles $2\pi\alpha$ and $2\pi\alpha$. Then, under the geodesic coordinate, ds^2 can be expressed as*

$$ds^2 = dr^2 + (\alpha \sin r)^2 d\theta^2,$$

where $(r, \theta) \in (0, \pi) \times [0, 2\pi)$.

3. PROOF OF **Theorem 1.1**

First, consider the map $\sigma : (0, \pi) \times [0, 2\pi) \rightarrow \mathbb{C} \setminus \{0\} = \mathbb{R}^2 \setminus \{(0, 0)\}$, defined by

$$\sigma(u, \theta) = \left(\tan \frac{u}{2} \cos \theta, \tan \frac{u}{2} \sin \theta \right).$$

Clearly, σ is smooth and bijective, meaning it serves as a parameter transformation. Therefore, to prove **Theorem 1.1**, it suffices to show that the map

$$\vec{r} : (0, \pi) \times [0, 2\pi) \rightarrow \mathbb{E}^3, (u, \theta) \mapsto \vec{r}(u, \theta) = (x^1(u, \theta), x^2(u, \theta), x^3(u, \theta))$$

is an immersion and $\vec{r}^* ds_E^2$ induces a football metric.

Since

$$\begin{aligned} \vec{r}_u &= (B \cos u \cos(\lambda\theta), B \cos u \sin(\lambda\theta), \sqrt{1 - B^2 \cos^2 u}), \\ \vec{r}_\theta &= (-B\lambda \sin u \sin(\lambda\theta), B\lambda \sin u \cos(\lambda\theta), 0), \end{aligned}$$

and \vec{r}_u and \vec{r}'_θ are linearly independent, \vec{r} is indeed an immersion. Furthermore, a unit normal vector \vec{n} is given by

$$\vec{n} = (-\sqrt{1 - B^2 \cos^2 u} \cos(\lambda\theta), -\sqrt{1 - B^2 \cos^2 u} \sin(\lambda\theta), B \cos u).$$

Next, we compute the second derivatives:

$$\vec{r}_{uu} = (-B \sin u \cos(\lambda\theta), -B \sin u \sin(\lambda\theta), \frac{B^2 \cos u \sin u}{\sqrt{1 - B^2 \cos^2 u}}),$$

$$\vec{r}_{u\theta} = (-B\lambda \sin u \sin(\lambda\theta), B\lambda \sin u \cos(\lambda\theta), 0),$$

$$\vec{r}_{\theta\theta} = (-B\lambda^2 \sin u \cos(\lambda\theta), -B\lambda^2 \sin u \sin(\lambda\theta), 0).$$

The coefficients of the first fundamental form are:

$$E = \vec{r}_u \cdot \vec{r}_u = 1, \quad F = \vec{r}'_u \cdot \vec{r}_\theta = 0, \quad G = \vec{r}_\theta \cdot \vec{r}_\theta = (B\lambda)^2 \sin^2 u.$$

The coefficients of the second fundamental form are:

$$L = \vec{n} \cdot \vec{r}_{uu} = \frac{B \sin u}{\sqrt{1 - B^2 \cos^2 u}}, \quad M = \vec{n} \cdot \vec{r}_{u\theta} = 0,$$

$$N = \vec{n} \cdot \vec{r}_{\theta\theta} = \lambda^2 B \sin u \sqrt{1 - B^2 \cos^2 u}.$$

Thus, the first and second fundamental forms are:

$$I = \vec{r}^* ds_E^2 = du^2 + (B\lambda)^2 \sin^2 u d\theta^2 = du^2 + \alpha^2 \sin^2 u d\theta^2,$$

$$II = \frac{B \sin u}{\sqrt{1 - B^2 \cos^2 u}} du^2 + \lambda^2 B \sin u \sqrt{1 - B^2 \cos^2 u} d\theta^2.$$

Moreover, the Gauss curvature is computed to be 1.

From the above proof, we derive the following proposition.

Proposition 3.1. *The mean curvature H of the immersion F in Theorem 1.1 is given by*

$$H = \frac{1}{2} \left(\frac{B \sin u}{\sqrt{1 - B^2 \cos^2 u}} + \frac{\sqrt{1 - B^2 \cos^2 u}}{B \sin u} \right) > 1,$$

where $u = 2 \arctan r$.

Proof. First, we observe that $\frac{1}{2} \left(\frac{B \sin u}{\sqrt{1 - B^2 \cos^2 u}} + \frac{\sqrt{1 - B^2 \cos^2 u}}{B \sin u} \right) \geq 1$. Equality holds if and only if

$$\frac{B \sin u}{\sqrt{1 - B^2 \cos^2 u}} = \frac{\sqrt{1 - B^2 \cos^2 u}}{B \sin u}$$

which implies $B = 1$. However, this leads to a contradiction. \square

4. HOW TO OBTAIN THE MAP \vec{r} IN SECTION 2

We derive the expression for \vec{r} in Section 2 using the method of rotationally symmetric surfaces, with slight modifications, as follows.

Let $f, g : [0, \pi] \rightarrow \mathbb{R}$ be two smooth functions satisfying

$$\begin{cases} f(0) = f(\pi) = g(0) = 0, \\ (f'(u))^2 + (g'(u))^2 = 1, \\ g'(u) \neq 0, \\ f(u), g(u) > 0, \forall u \in (0, \pi). \end{cases}$$

Suppose $\vec{r} : (0, \pi) \times [0, 2\pi) \rightarrow \mathbb{E}^3$, defined by

$$(u, \theta) \mapsto \vec{r}(u, \theta) = (x^1(u, \theta), x^2(u, \theta), x^3(u, \theta))$$

is a football with conical angles $2\pi\alpha$ and $2\pi\alpha$ in \mathbb{E}^3 , where

$$\begin{cases} x^1(u, \theta) = f(u) \cos(\lambda\theta), \\ x^2(u, \theta) = f(u) \sin(\lambda\theta), \\ x^3(u, \theta) = g(u), \end{cases}$$

and $\lambda \in \mathbb{Z}^+$ is a parameter.

Since

$$\begin{aligned} \vec{r}_u &= (f'(u) \cos(\lambda\theta), f'(u) \sin(\lambda\theta), g'(u)), \\ \vec{r}_\theta &= (-\lambda f(u) \sin(\lambda\theta), \lambda f(u) \cos(\lambda\theta), 0), \end{aligned}$$

and \vec{r}_u and \vec{r}_θ are linearly independent, \vec{r} is indeed an immersion. Furthermore, a unit normal vector \vec{n} is given by

$$\vec{n} = (-g'(u) \cos(\lambda\theta), -g'(u) \sin(\lambda\theta), f'(u)).$$

Next, we compute the second derivatives:

$$\begin{aligned} \vec{r}_{uu} &= (f''(u) \cos(\lambda\theta), f''(u) \sin(\lambda\theta), g''(u)), \\ \vec{r}_{u\theta} &= (-\lambda f'(u) \sin(\lambda\theta), \lambda f'(u) \cos(\lambda\theta), 0), \\ \vec{r}_{\theta\theta} &= (-\lambda^2 f(u) \cos(\lambda\theta), -\lambda^2 f(u) \sin(\lambda\theta), 0). \end{aligned}$$

The coefficients of the first fundamental form are:

$$E = \vec{r}_u \cdot \vec{r}_u = 1, \quad F = \vec{r}_u \cdot \vec{r}_\theta = 0, \quad G = \vec{r}_\theta \cdot \vec{r}_\theta = (\lambda f(u))^2.$$

The coefficients of the second fundamental form are:

$$\begin{aligned} L = \vec{n} \cdot \vec{r}_{uu} &= f'(u)g''(u) - f''(u)g'(u), \quad M = \vec{n} \cdot \vec{r}_{u\theta} = 0, \\ N = \vec{n} \cdot \vec{r}_{\theta\theta} &= \lambda^2 f(u)g'(u). \end{aligned}$$

Since $(f'(u))^2 + (g'(u))^2 = 1$, we have

$$f'(u)f''(u) + g'(u)g''(u) = 0,$$

and

$$(f'(u)g''(u) - f''(u)g'(u))g'(u) = -f''(u).$$

Moreover,

$$L = -\frac{f''(u)}{g'(u)}.$$

Thus, the first and second fundamental forms are:

$$\begin{aligned} I &= du^2 + (\lambda f(u))^2 d\theta^2, \\ II &= -\frac{f''(u)}{g'(u)} du^2 + \lambda^2 f(u) g'(u) d\theta^2. \end{aligned}$$

Since the Gauss curvature equals 1, we obtain the following equation

$$f''(u) + f(u) = 0.$$

Thus, f can be expressed as

$$f(u) = A \cos u + B \sin u.$$

Noting that $f(0) = f(\pi) = 0$, we obtain

$$f(u) = B \sin u,$$

and

$$g(u) = \int_0^u \sqrt{1 - B^2 \cos^2 t} dt.$$

Furthermore, the first fundamental form is

$$I = du^2 + (B\lambda)^2 \sin^2 u d\theta^2.$$

Thus, by **lemma 2.1**, $\lambda B = \alpha$, i.e., $\lambda = \frac{\alpha}{B} \in \mathbb{Z}^+$.

Consequently,

$$\begin{cases} x^1(u, \theta) = B \sin u \cos(\lambda\theta), \\ x^2(u, \theta) = B \sin u \sin(\lambda\theta), \\ x^3(u, \theta) = \int_0^u \sqrt{1 - B^2 \cos^2 t} dt, \end{cases}$$

where $\lambda B = \alpha$.

5. SOME EXPLICIT EXAMPLES


In this section, we will present graphical representations of these immersions based on Theorem 1.1.

Example 5.1. Let $\alpha = \frac{1}{2}$ and $B = \frac{1}{2}$, which implies $\lambda = 1$. The isometric immersion F can then be expressed as

$$F(r, \theta) = (x^1(2 \arctan r, \theta), x^2(2 \arctan r, \theta), x^3(2 \arctan r, \theta)),$$

where

$$\begin{cases} x^1(u, \theta) = \frac{1}{2} \sin u \cos \theta, \\ x^2(u, \theta) = \frac{1}{2} \sin u \sin \theta, \\ x^3(u, \theta) = \int_0^u \sqrt{1 - \frac{1}{4} \sin^2 t} dt. \end{cases}$$


TABLE 1. Football of angles π and π .

Example 5.2. Let $\alpha = 1$ and $B = \frac{1}{2}$, which implies $\lambda = 2$. The isometric immersion F can then be expressed as

$$F(r, \theta) = (x^1(2 \arctan r, \theta), x^2(2 \arctan r, \theta), x^3(2 \arctan r, \theta)),$$

where

$$\begin{cases} x^1(u, \theta) = \frac{1}{2} \sin u \cos 2\theta, \\ x^2(u, \theta) = \frac{1}{2} \sin u \sin 2\theta, \\ x^3(u, \theta) = \int_0^u \sqrt{1 - \frac{1}{4} \sin^2 t} dt. \end{cases}$$


TABLE 2. Football of angles 2π and 2π .

Example 5.3. Let $\alpha = 2$ and $B = \frac{1}{2}$, which implies $\lambda = 4$. The isometric immersion F can then be expressed as

$$F(r, \theta) = (x^1(2 \arctan r, \theta), x^2(2 \arctan r, \theta), x^3(2 \arctan r, \theta)),$$

where

$$\begin{cases} x^1(u, \theta) = \frac{1}{2} \sin u \cos 4\theta, \\ x^2(u, \theta) = \frac{1}{2} \sin u \sin 4\theta, \\ x^3(u, \theta) = \int_0^u \sqrt{1 - \frac{1}{4} \sin^2 t} dt. \end{cases}$$


TABLE 3. Football of angles 4π and 4π .

Example 5.4. Let $\alpha = 2$ and $B = \frac{1}{4}$, which implies $\lambda = 8$. The isometric immersion F can then be expressed as

$$F(r, \theta) = (x^1(2 \arctan r, \theta), x^2(2 \arctan r, \theta), x^3(2 \arctan r, \theta)),$$

where

$$\begin{cases} x^1(u, \theta) = \frac{1}{4} \sin u \cos 4\theta, \\ x^2(u, \theta) = \frac{1}{4} \sin u \sin 4\theta, \\ x^3(u, \theta) = \int_0^u \sqrt{1 - \frac{1}{16} \sin^2 t} dt. \end{cases}$$


TABLE 4. Football of angles 4π and 4π .

Example 5.5. Let $\alpha = 2$ and $B = \frac{1}{8}$, which implies $\lambda = 16$. The isometric immersion F can then be expressed as

$$F(r, \theta) = (x^1(2 \arctan r, \theta), x^2(2 \arctan r, \theta), x^3(2 \arctan r, \theta)),$$

where

$$\begin{cases} x^1(u, \theta) = \frac{1}{8} \sin u \cos 16\theta, \\ x^2(u, \theta) = \frac{1}{8} \sin u \sin 16\theta, \\ x^3(u, \theta) = \int_0^u \sqrt{1 - \frac{1}{64} \sin^2 t} dt. \end{cases}$$

TABLE 5. Football of angles 4π and 4π .

Declarations

Data Availability Statement This manuscript has no associated data.

Competing interests On behalf of all authors, the corresponding author states that there is no conflict of interest.

REFERENCES

- [1] Bartolucci D., De Marchis F., Malchiodi A. : Supercritical conformal metrics on surfaces with conical singularities, *Int. Math. Res. Not.*, 24 (2011), 5625-5643
- [2] Bryant R. L. : Surfaces of mean curvature one in hyperbolic space, *Astérisque*, 154-155 (1988), 321-347
- [3] Chen Q., Wang, W., Wu, Y. Y., Xu, B. : Conformal metrics with constant curvature one and finitely many conical singularities on compact Riemann surfaces, *Pacific J. Math.*, 273 (2015), no. 1, 75-100
- [4] Chen W. X., Li, C. : Classification of solutions of some nonlinear elliptic equations, *Duke Math. J.*, 63 (3) (1991), 615-622
- [5] Eremenko A. : Metrics of positive curvature with conic singularities on the sphere, *Proc. Amer. Math. Soc.*, 132(11) (2004), 3349-3355
- [6] Eremenko A. : Metrics of constant positive curvature with conic singularities. A survey. [arXiv:2103.13364](https://arxiv.org/abs/2103.13364)
- [7] Eremenko A., Mondello M. , Panov D. : Moduli of spherical tori with one conical point, *Geom. Topol.*, 27 (9) (2023), 3619-3698
- [8] José A. G., Laurent H., Pablo M. : Surfaces of constant curvature in \mathbb{R}^3 with isolated singularities, *Adv. Math.*, 241 (1) (2013), 103-126
- [9] Li L. G., Song J. J., Xu B. : Irreducible cone spherical metrics and stable extensions of two line bundles, *Adv. Math.*, 388 (2021), 36 pp
- [10] Luo F., Tian G. : Liouville equation and spherical convex polytopes. *Proc. Amer. Math. Soc.*, 116 (4) (1992), 1119-1129
- [11] Mazzeo R., Zhu X. W. : Conical metrics on Riemann surfaces, I: The compactified configuration space and regularity, *Geom. Topol.*, 24 (1) (2020), 309-372
- [12] Mazzeo R., Zhu X. W. : Conical metrics on Riemann surfaces, II: Spherical metrics, *Int. Math. Res. Not.*, (12) (2022), 9044-9113
- [13] McOwen R. C. : Point singularities and conformal metrics on Riemann surfaces, *Proc. Amer. Math. Soc.*, 103 (1) (1988), 222-224
- [14] Mondello D., Panov D. : Spherical metrics with conical singularities on a 2-sphere: angle constraints, *Int. Math. Res. Not.*, (16) (2016), 4937-4995

- [15] Mondello D., Panov D. : Spherical surfaces with conical points: systole inequality and moduli spaces with many connected component, *Geom. Funct. Anal.*, 29 (4) (2019), 1110-1193
- [16] Tahar, G. : Geometric decompositions of surfaces with spherical metric and conical singularities, *Q. J. Math.*, 73 (2) (2022), 657-678
- [17] Troyanov, M. : Metrics of constant curvature on a sphere with two conical singularities, *Differential geometry*, 296-306, *Lecture Notes in Math.*, 1410, Springer, Berlin, 1989
- [18] Troyanov, M. : Prescribing curvature on compact surfaces with conical singularities, *Trans. Amer. Math. Soc.*, 324 (2) (1991), 793-821
- [19] Umehara, M., Yamada, K. : Metrics of constant curvature 1 with three conical singularities on the 2-sphere, *Illinois J. Math.*, 44 (1) (2000), 72-94
- [20] Wei, Z. Q., Wu, Y. Y., Xu, B. : Geometric structure and the existence of reducible spherical conical metrics, submitted

Zhiqiang Wei

School of Mathematics and Statistics, Henan University, Kaifeng 475004 P. R. China

Center for Applied Mathematics of Henan Province, Henan University, Zhengzhou 450046 P. R. China

Email: weizhiqiang15@mails.ucas.edu.cn. or 10100123@vip.henu.edu.cn.