A SHARP BOUND ON LARGE PLANAR SIGNED VECTOR SUMS

FLORIAN GRUNDBACHER

ABSTRACT. We give a sharp lower bound to the largest possible Euclidean norm of signed sums of n vectors in the plane. This is achieved by connecting the signed vector sum problem to the isoperimetric problem for the circumradius of polygons. In turn, we apply the sharp bound for the signed vector sum problem to establish a sharp lower bound to the circumradius of the Minkowski sum of n planar symmetric convex bodies. We also determine a tight lower bound to the circumradius of the Minkowski sum of general convex bodies in any dimension independent of their number.

1. INTRODUCTION

Vector sum problems have a long history. Especially problems concerned with finding small signed sums or small sums of subsets of prescribed cardinality have gained attention in the past literature since they provide a unified framework to deal with various problems from other areas. A general overview of the many variants of these vector sum problems and related results is provided in [2, 4].

Recently, a new variant of these vector sum problems that is in spirit dual to the ones mentioned above has been considered by Ambrus and González Merino in [2]. Instead of considering how small appropriate choices of (signed) vector (subset) sums can be made, they asked for best possible lower bounds to the *largest* of these sums. Formally, for integers $n \ge k \ge 1$, they asked to determine

$$c(d,n,k) \coloneqq \min_{\substack{u^1,\dots,u^n \in \mathbb{S}^{d-1} \\ \varepsilon_{i_1},\dots,\varepsilon_{i_k} \in \{-1,1\}}} \max_{\substack{u^1,\dots,u^n \in \mathbb{S}^{d-1} \\ \varepsilon_{i_1},\dots,\varepsilon_{i_k} \in \{-1,1\}}} \left\| \sum_{j=1}^k \varepsilon_{i_j} u^{i_j} \right\|,$$

where \mathbb{S}^{d-1} is the *Euclidean unit sphere* in \mathbb{R}^d and $\|\cdot\|$ denotes the *Euclidean norm*. Already in [2], multiple sharp results on the asymptotic behavior of c(d, n, k) in its three parameters are obtained by again establishing connections to many other mathematical areas. Furthermore, in the planar case, improved lower bounds to c(2, n, k) and its precise value when k - 1 divides n are provided.

The results in [2] leave open the very natural problem of computing c(d, n, n). This appears to be a difficult task, as even c(d, d + 1, d + 1) is unknown for $d \ge 3$ (see [2, Conjecture 1]). Given that the precise value of c(2, n, k) has been obtained in special cases, one may hope that at least c(2, n, n) can be found for general n. The main problem in this direction with the lower bounds to c(2, n, k) in [2, Theorem 5] is that their strength comparatively declines as k becomes larger. They eventually even decrease despite c(2, n, k) clearly increasing in k.

The main goal of this note is to remedy the above problem in the planar case by computing c(2, n, n) precisely for all n. As a corollary, we also improve the lower bound to c(2, n, k) for large k.

Our approach is based on the connection between the problem of computing c(d, n, n) and the isoperimetric problem for the circumradius of polytopes established by Joós and Lángi in [18]. Although using different terminology, they point out that c(d, n, n) is equal to the minimal circumradius of a zonotope in \mathbb{R}^d generated from n segments of length 2. Using a Dowker-type result for the circumradius-perimeter-ratio of polygons, which is discussed in Section 2, this leads us to a straightforward proof for the precise value of c(2, n, n) in Section 3.

Date: February 19, 2025.

²⁰²⁰ Mathematics Subject Classification. Primary 52A40; Secondary 05A99, 52A10, 52A37, 52B60.

Key words and phrases. Vector sums, Minkowski addition, Circumradius, Isoperimetric problem.

F. GRUNDBACHER

In Section 4, we show that the connection between signed vector sum problems and problems related to the circumradius is mutually beneficial. As already discussed, c(2, n, n) can be computed via the isoperimetric problem for the circumradius of polygons. In turn, this approach allows us to find a sharp lower bound to the circumradius of the Minkowski sum of n planar symmetric convex bodies. Problems of the latter type were first studied in [16] and have since been generalized to various other settings (cf. [1, 10, 15, 17, 19]), typically only for sums of two convex bodies. See also [9], where sums of n convex bodies are considered in connection to other vector sum problems.

2. Preliminaries

For $X, Y \subset \mathbb{R}^d$, their *Minkowski sum* is given by $X + Y \coloneqq \{x + y : x \in X, y \in Y\}$. The *t*-translation and ρ -dilatation of X for $t \in \mathbb{R}^d$ and $\rho \in \mathbb{R}$ are defined as $t + X \coloneqq \{t\} + X$ and $\rho X \coloneqq \{\rho x : x \in X\}$. We abbreviate $-X \coloneqq (-1)X$. The closed *segment* connecting $x, y \in \mathbb{R}^d$ is denoted by [x, y]. A convex body $K \subset \mathbb{R}^d$ is a non-empty compact convex set. It is (t-)symmetric if there exists $t \in \mathbb{R}^d$ with -t + K = t - K, or equivalently $K = t + \frac{K - K}{2}$.

The *circumradius* of K is given by $R(K) := \min\{\rho \ge 0 : K \subset t + \rho \mathbb{B}^d, t \in \mathbb{R}^d\}$, where \mathbb{B}^d is the *Euclidean unit ball* in \mathbb{R}^d . This functional is by definition translation invariant. It is well-known (see, e.g., [7, Section 35]) that K possesses a unique circumball. The center of this ball is not an extreme point of K (unless K is a singleton) and coincides with the center of K if K is symmetric. For d = 2, we denote the *perimeter* of K by L(K). Let us point out that the perimeter is Minkowski additive and strictly increasing (see, e.g., [7, Paragraph 7]).

The key ingredient for our proof of Theorem 3.1 in the next section is the following solution to the isoperimetric problem for the circumradius of polygons.

Proposition 2.1. Let $P \subset \mathbb{R}^2$ be a convex m-gon for integers $1 \le m \le n$. Then $2n \sin\left(\frac{\pi}{n}\right) R(P) \ge L(P),$

with equality if and only if P is a regular n-gon or a singleton.

The above proposition falls into the category of *Dowker-type results* (in reference to the fundamental paper [11] by Dowker; see [12, 13, 20] for the corresponding results involving the perimeter). Despite the elementary nature of the problem underlying the proposition, a clear proof of the complete result appears difficult to establish in the literature (see [5, 14] for the inequality without the characterization of the equality case, and [3, Open Problems (3a)] for mention of the entire result but without proof). For the sake of completeness, we provide a short proof below.

Proof. Let $t \in \mathbb{R}^2$ and $\rho \ge 0$ be such that $t + \rho \mathbb{B}^n$ is the circumcircle of P. All claims are clear if P is a singleton, so we may assume $\rho > 0$. It is well-known that t must be a non-extreme point of P in this case. If P' denotes the convex m-gon whose vertices are the intersection points of $t + \rho \mathbb{S}^1$ with the rays emanating from t through the vertices of P, then $t \in P \subset t + \rho \mathbb{B}^2$ implies $P \subset P'$. We may further enlargen P' to a convex n-gon P^* whose vertices all lie in $t + \rho \mathbb{S}^1$. The strict monotonicity of the perimeter shows $L(P) \le L(P^*)$, with equality if and only if P is an n-gon with all of its vertices in $t + \rho \mathbb{S}^1$. Since $R(P^*) = R(P)$, it suffices to prove the proposition for P^* .

The assumption $\rho > 0$ implies by $R(P^*) = \rho$ that $n \ge 2$. Let v^1, \ldots, v^n be the distinct vertices of P^* and write $v^{n+1} \coloneqq v^1$. We may further assume that the vertices are indexed such that v^i and $v^{i+1}, i \in \{1, \ldots, n\}$, are connected by an edge of P^* . For $i \in \{1, \ldots, n\}$, let $\alpha_i \in [0, \pi]$ be the angle enclosed between $v^i - t$ and $v^{i+1} - t$. Since these two vectors have norm ρ , the distance between them, and consequently the distance between v^i and v^{i+1} , equals $2\rho \sin\left(\frac{\alpha_i}{2}\right)$. Note that $t \in P^*$ implies $\sum_{i=1}^n \alpha_i = 2\pi$. Hence, the concavity of sine on the interval $[0, \pi]$ yields

$$L(P^*) = \sum_{i=1}^n 2\rho \sin\left(\frac{\alpha_i}{2}\right) = 2n\rho \sum_{i=1}^n \frac{\sin(\frac{\alpha_i}{2})}{n} \le 2n\rho \sin\left(\sum_{i=1}^n \frac{\alpha_i}{2n}\right) = 2n \sin\left(\frac{\pi}{n}\right) R(P^*).$$

Equality holds if and only if every α_i equals $\frac{2\pi}{n}$, or equivalently if P^* is a regular *n*-gon as claimed. \Box

3. Signed Sums of Planar Vectors

Our main result on signed vector sums concerns a generalization of the problem underlying c(2, n, n). Instead of dealing only with unit vectors, we can allow arbitrary vectors.

Theorem 3.1. Let $u^1, \ldots, u^n \in \mathbb{R}^2$. Then

$$\max_{\varepsilon_1,\ldots,\varepsilon_n\in\{-1,1\}} \left\|\sum_{i=1}^n \varepsilon_i u^i\right\| \ge \frac{1}{n\sin\left(\frac{\pi}{2n}\right)} \sum_{i=1}^n \left\|u^i\right\|,$$

with equality if and only if $\{\pm u^1, \ldots, \pm u^n\}$ forms the vertex set of a regular 2n-gon or equals $\{0\}$.

Our main proof idea is to relate the expressions in the above inequality to the circumradius and perimeter of a certain zonotope. A direct application of Proposition 2.1 then yields the claimed inequality. As initially mentioned, this approach (up to the last step) is also pointed out in [18, Section 2]. While the identities (1) and (2) below are also derived in [18, Corollary 1 and (5)], as well as the inequality in Proposition 2.1 being mentioned in the proof of [18, Lemma 6], they were not combined to obtain the above inequality.

Proof. Let $P \coloneqq \sum_{i=1}^{n} [-u^{i}, u^{i}]$ be a zonotope. Since the perimeter is Minkowski additive, we have

(1)
$$L(P) = 4 \sum_{i=1}^{n} ||u^{i}||$$

The zonotope P is 0-symmetric, so it is further clear that

(2)
$$R(P) = \max_{v \in P} \|v\| = \max_{\varepsilon_1, \dots, \varepsilon_n \in \{-1, 1\}} \left\| \sum_{i=1}^n \varepsilon_i u^i \right\|.$$

It is easy to see that any edge of P must be parallel to one of the segments $[-u^i, u^i]$. Therefore, P has at most 2n edges (less if some u^i are scalar multiples of each other), which means P is an m-gon for some $m \leq 2n$. Proposition 2.1 now shows

$$\max_{\varepsilon_1,\dots,\varepsilon_n\in\{-1,1\}} \left\|\sum_{i=1}^n \varepsilon_i u^i\right\| = R(P) \ge \frac{L(P)}{4n\sin\left(\frac{\pi}{2n}\right)} = \frac{1}{n\sin\left(\frac{\pi}{2n}\right)} \sum_{i=1}^n \left\|u^i\right\|,$$

which, read from left to right, is the claimed inequality.

Equality holds if and only if P is a regular 2n-gon or a singleton. The latter is equivalent to all u^i being zero, which is the second claimed equality case. For the other equality case, we first note that P is a 2n-gon if and only if all u^i are non-zero and no two of them are scalar multiples of each other. In this case, we may assume that u^1, \ldots, u^n are ordered and oriented in sign such that they induce consecutive edges of P, i.e., P has a vertex v^0 such that $v^i \coloneqq v^{i-1} + 2u^i$ is a vertex of P for all $i \in \{1, \ldots, n\}$. Now, P is additionally regular if and only if all of its edges are of equal length and all interior angles at its vertices coincide. The first condition is equivalent to all u^i having the same positive norm. The second condition is equivalent to the angles enclosed between u^i and u^{i+1} for all $i \in \{1, \ldots, n-1\}$, as well as the angle enclosed between u^n and $-u^1$, all being the same. Altogether, P is a regular 2n-gon if and only if $\{\pm u^1, \ldots, \pm u^n\}$ itself forms the vertex set of a regular 2n-gon, which is the first claimed equality case.

The special case of the above theorem where all u^i are unit vectors immediately establishes the precise value of c(2, n, n). We also obtain the desired improvement of the lower bound to c(2, n, k) for large k compared to [2, Theorems 2 and 5].

Corollary 3.2. Let $n \ge k \ge 1$ be integers. Then

$$c(2, n, k) \ge \frac{1}{\sin\left(\frac{\pi}{2k}\right)},$$

with equality if and only if $k \in \{1, n\}$.

Proof. The inequality is immediately obtained from the above theorem. Equality holds if and only if there exist n unit vectors in \mathbb{R}^2 such that for every choice of k of them, the chosen vectors and their negatives together form the vertex set of a regular 2k-gon. This is easily seen to be the case if and only if $k \in \{1, n\}$.

Our result also impacts the upper bound to c(d, n, k) for general dimensions $d \ge 2$. The monotonicity of c(d, n, k) in its three parameters (see the discussion below [2, Definition 1]) yields

$$c(d, n, k) \le c(2, n, n) = \frac{1}{\sin\left(\frac{\pi}{2n}\right)}$$

By the above corollary and [2, Proposition 2], equality holds if k = n and $\min\{d, n\} \leq 2$. A closer analysis of the monotonicity of c(d, n, k) shows that these are the only equality cases, though we omit the details. While the above constitutes for $d \geq 2$ the best possible upper bound that is independent of d and k, the asymptotically much stronger bounds in [2, Theorem 2] show that the new estimate is rather weak if d is large or k is small.

4. CIRCUMRADIUS OF MINKOWSKI SUMS OF CONVEX BODIES

As initially mentioned, the theory of circumradii also benefits from the link to vector sum problems. First results derived from this connection appeared in [9]. Our focus lies on the setting of [9, Corollary 1.1], where a sharp lower bound to the circumradius of the Minkowski sum of three planar convex bodies with unit circumradii is established. The theorem below generalizes this result to an arbitrary number of planar convex bodies with any circumradii, though under the additional assumption that the convex bodies involved are symmetric. Afterward, we use a simplified (but also coarser) method to determine the best possible absolute constants (independent of the number of convex bodies) that can be used in such a lower bound for sums of general convex bodies in arbitrary dimensions.

Theorem 4.1. Let $K^1, \ldots, K^n \subset \mathbb{R}^2$ be symmetric convex bodies. Then

$$R(K^1 + \ldots + K^n) \ge \frac{1}{n \sin\left(\frac{\pi}{2n}\right)} \left(R(K^1) + \ldots + R(K^n) \right)$$

For $n \geq 2$, equality holds only if there exist $\rho \geq 0$ and $\varphi \in [0, \pi)$ such that each of the segments

$$\left[-\rho\left(\cos\left(\frac{j\pi}{n}+\varphi\right),\sin\left(\frac{j\pi}{n}+\varphi\right)\right),\rho\left(\cos\left(\frac{j\pi}{n}+\varphi\right),\sin\left(\frac{j\pi}{n}+\varphi\right)\right)\right], j \in \{1,\dots,n\},$$

is the unique longest segment in one of the convex bodies $\frac{K^i-K^i}{2}$. Moreover, the inequality is sharp.




FIGURE 1. Two examples for Theorem 4.1: K^1 (dashed, blue), K^2 (dotted, red), $K^1 + K^2$ (dash-dotted, purple), $R(K^1)\mathbb{B}^2 = R(K^2)\mathbb{B}^2$ and $R(K^1 + K^2)\mathbb{B}^2$ (solid, black). The constant in Theorem 4.1 for n = 2 equals $\frac{1}{\sqrt{2}}$. The left-hand example shows that the necessary equality condition in Theorem 4.1 cannot be reduced to K^1 and K^2 consisting of only the described segments. The right-hand example shows that the necessary condition is, in general, not sufficient for equality.

Proof. Let $C^1 := \frac{K^1 - K^1}{2}, \ldots, C^n := \frac{K^n - K^n}{2}$ be the 0-symmetric translates of K^1, \ldots, K^n . By the translation invariance of the circumradius, it suffices to prove the theorem for the C^i .

It is well-known that the circumcircle of any C^i has its center at the origin. We may therefore choose some $u^i \in C^i$ with $||u^i|| = R(C^i)$ for every $i \in \{1, \ldots, n\}$. Since the circumcircle of the set $C^1 + \ldots + C^n$ is also centered at the origin, we obtain from Theorem 3.1 that

(3)

$$R(C^{1} + \ldots + C^{n}) = \max_{v \in C^{1} + \ldots + C^{n}} \|v\| \ge \max_{\varepsilon_{1}, \ldots, \varepsilon_{n} \in \{-1, 1\}} \left\| \sum_{i=1}^{n} \varepsilon_{i} u^{i} \right\|$$

$$\ge \frac{1}{n \sin\left(\frac{\pi}{2n}\right)} \sum_{i=1}^{n} \|u^{i}\| = \frac{1}{n \sin\left(\frac{\pi}{2n}\right)} \left(R(C^{1}) + \ldots + R(C^{n}) \right).$$

If equality holds and excluding the trivial instance $\{\pm u^1, \ldots, \pm u^n\} = \{0\}$, Theorem 3.1 implies that $\{\pm u^1, \ldots, \pm u^n\}$ forms the vertex set of a regular 2*n*-gon. In this case, it is easy to see that the segments $[-u^i, u^i]$ can be written in the way outlined in the present theorem. Since $C^i \subset R(C^i)\mathbb{B}^2$, it is further clear that these segments are always longest segments in the C^i . For $n \geq 2$, they must indeed be the unique longest segments in the C^i : Again by $C^i \subset R(C^i)\mathbb{B}^2$, any other longest segment in C^i could be written in the form $[-\tilde{u}^i, \tilde{u}^i]$ for some $\tilde{u}^i \in C^i$ with $\|\tilde{u}^i\| = R(C^i)$. However, replacing u^i with \tilde{u}^i in (3) would now lead to a strict inequality since $(\{\pm u^1, \ldots, \pm u^n\} \setminus \{\pm u^i\}) \cup \{\pm \tilde{u}^i\}$ cannot also form the vertex set of a regular 2n-gon for $n \geq 2$.

Finally, it is clear that (3) is satisfied with equality from left to right if $C^i = [-u^i, u^i]$ for every $i \in \{1, ..., n\}$ and $\{\pm u^1, ..., \pm u^n\}$ forms the vertex set of a regular 2*n*-gon.

With $1 < \frac{x}{\sin(x)} \to 1$ for $0 < x \to 0$ and $0 < \frac{\pi}{2n} \to 0$ for $n \to \infty$, the above theorem yields the inequality $R(K^1 + \ldots + K^n) \geq \frac{2}{\pi} \left(R(K^1) + \ldots + R(K^n) \right)$ for any planar symmetric convex bodies $K^1, \ldots, K^n \subset \mathbb{R}^2$. This result is generalized with a more direct proof in the remark below. We write κ_d for the volume of \mathbb{B}^d and $W_{d-1}(K)$ for the (d-1)-th quermassintegral of a convex body $K \subset \mathbb{R}^d$. Recall that W_{d-1} is translation invariant, Minkowski additive, positive homogeneous of degree 1, and strictly increasing (see, e.g., [7, Paragraph 7]). Moreover, we have $W_{d-1}(\mathbb{B}^d) = \kappa_d$ and

$$\kappa_d R(K) \ge W_{d-1}(K) \ge \frac{2\kappa_{d-1}}{d} R(K).$$

For non-singleton K, equality holds in the first inequality if and only if K is a Euclidean ball, and in the second inequality if and only if K is a segment. The first inequality is immediate from the properties of W_{d-1} and the fact that K is contained in a Euclidean ball of radius R(K). The second inequality is shown in [8, Theorem 1.4] (using the *first intrinsic volume* $V_1(K) := \frac{d}{K_{d-1}} W_{d-1}(K)$).

Remark 4.2. For convex bodies $K^1, \ldots, K^n \subset \mathbb{R}^d$, we have the inequality

$$R(K^{1} + \ldots + K^{n}) \geq \frac{1}{\kappa_{d}} W_{d-1}(K^{1} + \ldots + K^{n})$$

= $\frac{1}{\kappa_{d}} (W_{d-1}(K^{1}) + \ldots + W_{d-1}(K^{n})) \geq \frac{2\kappa_{d-1}}{d\kappa_{d}} (R(K^{1}) + \ldots + R(K^{n})).$

Equality holds in the second inequality if and only if all K^i are singletons or segments. In this case, $K^1 + \ldots + K^n$ is a polytope, which implies that the first inequality is strict unless d = 1 or all K^i are singletons. Thus, the inequality from left to right is strict unless d = 1 or all K^i are singletons.

To see that $\frac{2\kappa_{d-1}}{d\kappa_d}$ is nevertheless the best possible absolute constant for the inequality from left to right, we use the well-known fact that \mathbb{B}^d is a zonoid [6]. This means that for any $\varepsilon > 0$, there exist $n \in \mathbb{N}$ and segments $S^1, \ldots, S^n \subset \mathbb{R}^d$ such that $S^1 + \ldots + S^n \subset \mathbb{B}^d \subset (1 + \varepsilon)(S^1 + \ldots + S^n)$. Now,

$$R(S^{1} + \ldots + S^{n}) \leq R(\mathbb{B}^{d}) = \frac{1}{\kappa_{d}} W_{d-1}(\mathbb{B}^{n}) \leq \frac{1}{\kappa_{d}} W_{d-1}\left((1+\varepsilon)(S^{1} + \ldots + S^{n})\right)$$
$$= \frac{1+\varepsilon}{\kappa_{d}} \left(W_{d-1}(S^{1}) + \ldots + W_{d-1}(S^{n})\right) = (1+\varepsilon)\frac{2\kappa_{d-1}}{d\kappa_{d}} \left(R(S^{1}) + \ldots + R(S^{n})\right).$$

We end this note with some final remarks. It would be interesting to know if Theorem 4.1 remains true if the convex bodies are allowed to be non-symmetric. The results in [9] already partially verify this for $n \leq 3$. Moreover, Remark 4.2 shows that an only slightly weaker inequality is, in fact, true. One approach to extend Theorem 4.1 to general planar convex bodies might be to generalize Theorem 3.1 in the sense of the vector sum problems considered in [9].

Another interesting direction for future research would be to generalize Theorem 4.1 to dimensions $d \geq 3$. As the proof of Theorem 4.1 indicates, this problem is naturally connected to computing c(d, n, n). The latter appears to be a difficult task for n > d as outlined in the introduction.

References

- J. Abardia-Evéquoz, M. A. Hernández Cifre, and E. Saorín Gómez, Mean projection and section radii of convex bodies, Acta Math. Hungar. 155 (2018), no. 1, 89–103.
- [2] G. Ambrus and B. González Merino, Large signed subset sums, Mathematika 67 (2021), no. 3, 579–595.
- [3] C. Audet, P. Hansen, and F. Messine, Extremal problems for convex polygons, J. Glob. Optim. 38 (2007), no. 2, 163–179.
- [4] I. Bárány, On the power of linear dependencies, In: M. Grötschel, G. O. H. Katona, and G. Sági (eds.), Building Bridges: Between Mathematics and Computer Science, Bolyai Society Mathematical Studies 19 (2008), Springer Berlin, Heidelberg, 31–45.
- [5] B. Basit and Z. Lángi, Dowker-type theorems for disk-polygons in normed planes, Discrete Math. 347 (2024), no. 6, 114019.
- [6] E. D. Bolker, A class of convex bodies, Trans. Amer. Math. Soc. 145 (1969), 323-345.
- [7] T. Bonnesen and W. Fenchel, *Theory of convex bodies*, BCS Associates, Moscow, Idaho (1987) (transl. from German and ed. by L. F. Boron, C. O. Christenson, and B. A. Smith, original 1934).
- [8] K. J. Böröczky and D. Hug, A reverse Minkowski-type inequality, Proc. Amer. Math. Soc. 148 (2020), no. 11, 4907–4922.
- M. Brugger, M. Fiedler, B. González Merino, and A. Kirschbaum, Additive colourful Carathéodory type results with an application to radii, Linear Algebra Appl. 554 (2018), 342–357.
- [10] F. Chen, C. Yang, and M. Luo, Successive radii and Orlicz Minkowski sum, Monatsh. Math. 179 (2016), no. 2, 201–219.
- [11] C. H. Dowker, On minimum circumscribed polygons, Bull. Amer. Math. Soc. 50 (1944), no. 2, 120–122.
- [12] H. G. Eggleston, Approximation to plane convex curves. (I) Dowker-type theorems, Proc. London. Math. Soc. (3) 7 (1957), no. 1, 351–377.
- [13] L. Fejes Tóth, Remarks on polygon theorems of Dowker, Mat. Lapok 6 (1955), 176–179 (in Hungarian).
- [14] G. Fejes Tóth and L. Fejes Tóth, Remark on a paper of C. H. Dowker, Period. Math. Hungar. 3 (1973), no. 3–4, 271–274.
- [15] B. González Merino, On the families of successive radii and the sum of convex sets, Beitr. Algebra Geom. 55 (2014), no. 2, 433–440.
- [16] B. González Merino and M. A. Hernández Cifre, Successive radii and Minkowski addition, Monatsh. Math. 166 (2012), no. 3–4, 395–409.
- [17] B. González Merino and M. A. Hernández Cifre, On successive radii and p-sums of convex bodies, Adv. Geom. 14 (2014), no. 1, 117–128.
- [18] A. Joós and Z. Lángi, Isoperimetric problems for zonotopes, Mathematika 69 (2023), no. 2, 508–534.
- [19] N. Lombardi and E. Saorín Gómez, Mean radii of symmetrizations of a convex body, Beitr. Algebra Geom. 65 (2024), no. 2, 415–440.
- [20] J. Molnár, On inscribed and circumscribed polygons of convex regions, Mat. Lapok 6 (1955), 210–218 (in Hungarian).

Florian Grundbacher – Technical University of Munich, Department of Mathematics, Germany. florian.grundbacher@tum.de