A REMARK ON THE WEYL QUANTISATION OF PALEY-WIENER FUNCTIONS

HELGE J. SAMUELSEN

ABSTRACT. We present a short proof of the fact that the Weyl quantisation of a tempered distribution with compactly supported Fourier transform is in the Schatten *p*-class if and only if the symbol is L^p -integrable. The proof is based on Werner-Young's inequality from quantum harmonic analysis and a quantum version of Wiener's division lemma.

1. INTRODUCTION

The theory of quantum harmonic analysis was introduced by Werner, and allows for Fourier analysis of operators [12]. In this setting, the appropriate replacement of the classical L^p spaces are the Schatten *p*-classes of compact operators T on $L^2(\mathbb{R}^d)$ such that

$$||T||_{\mathcal{S}^p} = \left(\operatorname{tr}(|T|^p)\right)^{\frac{1}{p}} < \infty.$$

Several classical results have been extended to the setting of quantum harmonic analysis, such as Wiener's Tauberian theorem [1, 7], Fourier restriction [5, 9] and decoupling [11].

From the inherent symplectic structure on \mathbb{R}^{2d} , it is convenient to introduce the symplectic Fourier transform when working with quantum harmonic analysis. Let σ denote the standard symplectic form on \mathbb{R}^{2d} , and define the symplectic Fourier transform by

$$\mathcal{F}_{\sigma}(\Psi)(\zeta) = \int_{\mathbb{R}^{2d}} e^{-2\pi i \sigma(\zeta, z)} \Psi(z) \, dz,$$

for $\Psi \in \mathscr{S}(\mathbb{R}^{2d})$. The symplectic Fourier transform extends to an isomorphism on $\mathscr{S}'(\mathbb{R}^{2d})$ through duality by

$$\langle \mathcal{F}_{\sigma}(\tau), \Psi \rangle_{\mathscr{S}',\mathscr{S}} = \langle \tau, \mathcal{F}_{\sigma}\Psi \rangle_{\mathscr{S}',\mathscr{S}},$$

for $\tau \in \mathscr{S}'(\mathbb{R}^{2d})$ and all $\Psi \in \mathscr{S}(\mathbb{R}^{2d})$.

The theory of quantum harmonic analysis is based on the isometric isomorphism between $L^2(\mathbb{R}^{2d})$ and the class of Hilbert-Schmidt operators \mathcal{S}^2 for the Weyl quantisation [10]. For a tempered distribution $\tau \in \mathscr{S}'(\mathbb{R}^{2d})$, the Weyl quantisation is defined through a sesquilinear dual pairing as the bounded linear operator $L_{\tau} : \mathscr{S}(\mathbb{R}^d) \to \mathscr{S}'(\mathbb{R}^d)$ such that

$$\langle L_{\tau}\varphi,\psi\rangle_{\mathscr{S}',\mathscr{S}} = \langle \tau,\mathcal{W}(\psi,\varphi)\rangle_{\mathscr{S}',\mathscr{S}},$$

for every $\varphi, \psi \in \mathscr{S}(\mathbb{R}^d)$. The tempered distribution τ is called the symbol of the operator L_{τ} , and $\mathcal{W}(\psi, \varphi)$ is the cross-Wigner distribution defined as

$$\mathcal{W}(\psi,\varphi)(x,\xi) = \int_{\mathbb{R}^d} \varphi\left(x + \frac{t}{2}\right) \overline{\psi\left(x - \frac{t}{2}\right)} e^{-2\pi i \xi \cdot t} dt.$$

Date: February 20, 2025.

²⁰²⁰ Mathematics Subject Classification. 42B10, 47G30, 47B10.

Unlike the case p = 2, there is generally no equivalence between L^{p} -integrability of the symbol and Schatten properties of the Weyl quantisation. Nevertheless, there is an equivalence whenever the symbol has compact Fourier support.

Theorem 1. Let $\tau \in \mathscr{S}'(\mathbb{R}^{2d})$, and assume that $\mathcal{F}_{\sigma}(\tau)$ is a compactly supported distribution on \mathbb{R}^{2d} . Then L_{τ} is a compact operator on $L^2(\mathbb{R}^{2d})$ if and only if $\tau \in C_0(\mathbb{R}^{2d})$, and $L_{\tau} \in \mathcal{S}^p$ if and only if $\tau \in L^p(\mathbb{R}^{2d})$ for $1 \leq p \leq \infty$.

Distributions with compact Fourier support are necessarily real analytic functions on \mathbb{R}^{2d} by [3, Thm. 7.3.1], and thus Theorem 1 says that the Weyl quantisation is compact if and only if the symbol vanishes at infinity.

A version of Theorem 1 first surfaced in a paper on Fourier restriction in quantum harmonic analysis [5]. Here it was proved under the assumption that $\mathcal{F}_{\sigma}(\tau)$ is a compactly supported Radon measure. The theorem was later extended to tempered distributions with compact Fourier support by Mishra and Vemuri [8], and independently by Müller [9].

The aim of this paper is to present a short proof of Theorem 1 using techniques from quantum harmonic analysis. For a detailed exposition of quantum harmonic analysis, we refer to [5-7, 11, 12].

2. Proof of Theorem 1

The main tool from quantum harmonic analysis needed for the proof of Theorem 1 is Werner's operator convolution given by

(1)
$$T \star S(z) = \operatorname{tr}(T\alpha_z(PSP)), \quad z \in \mathbb{R}^{2d}.$$

Here Pf(t) = f(-t) denotes the parity operator and α_z the operator translation

$$\alpha_z(T) = \rho(z)T\rho(-z).$$

The operator $\rho : \mathbb{R}^{2d} \to \mathcal{L}(L^2(\mathbb{R}^d))$ is the symmetric time-frequency shift given by

$$\rho(x,\xi)f(t) = e^{-\pi i x \cdot \xi} e^{2\pi i \xi \cdot t} f(t-x).$$

An important property of the operator convolution is that it coincides with the convolution of the Weyl symbols. More specifically, for $\tau \in \mathscr{S}'(\mathbb{R}^{2d})$ and $\Phi \in \mathscr{S}(\mathbb{R}^{2d})$ we have

(2)
$$L_{\tau} \star L_{\Phi} = \tau * \Phi,$$

as they both have the same symplectic Fourier transform [6].

The proof of Theorem 1 relies on two theorems related to the operator convolution defined by (1). The first is a version of Young's inequality for the operator convolution [12, Thm 3.3].

Proposition 2.1 (Werner-Young's inequality). Suppose $S \in S^p$ and $T \in S^q$ with $1 + r^{-1} = p^{-1} + q^{-1}$. Then $S \star T \in L^r(\mathbb{R}^{2d})$ and

$$\|S \star T\|_{L^r} \le \|S\|_{\mathcal{S}^p} \|T\|_{\mathcal{S}^q}.$$

The final ingredient of the proof is the following result, first used to derive an equivalence between classical and quantum decoupling.

Theorem 2.2 ([11], Cor. 3.2.1). Let $\Omega \subseteq \mathbb{R}^{2d}$ be a bounded set, and let $1 \leq p \leq \infty$. Then there exist L^2 -normalised $g, h \in \mathscr{S}(\mathbb{R}^{2d})$ and $C = C(\Omega) > 0$ such that if $\tau \in \mathscr{S}'(\mathbb{R}^{2d})$ and $\mathcal{F}_{\sigma}(\tau)$ is supported on Ω , then

$$||L_{\tau}||_{\mathcal{S}^p} \le C(\Omega) ||L_{\tau} \star (g \otimes h)||_{L^p}.$$

REFERENCES

Proof of Theorem 1. Let $\tau \in \mathscr{S}'(\mathbb{R}^{2d})$. If $\mathcal{F}_{\sigma}(\tau)$ is compactly supported, then there exists a bounded subset $\Omega \subseteq \mathbb{R}^{2d}$ such that supp $(\mathcal{F}_{\sigma}(\tau)) \subseteq \Omega$. Assume first that $L_{\tau} \in \mathcal{S}^p$. If $\Psi \in C_c^{\infty}(\mathbb{R}^{2d})$ is a smooth cut-off function equal to 1 on Ω ,

then

$$\mathcal{F}_{\sigma}(\tau) = \mathcal{F}_{\sigma}(\tau)\Psi,$$

and thus $\tau = \tau * \mathcal{F}_{\sigma}(\Psi)$. By (2), and Theorem 2.1, it follows that

(3)
$$\|\tau\|_{L^p} = \|\tau * \mathcal{F}_{\sigma}(\Psi)\|_{L^p} = \|L_{\tau} \star L_{\mathcal{F}_{\sigma}(\Psi)}\|_{L^p} \le \|L_{\mathcal{F}_{\sigma}(\Psi)}\|_{\mathcal{S}^1}\|L_{\tau}\|_{\mathcal{S}^p}$$

Since the Weyl quantisation of a Schwartz function belongs to S^1 by [13, Thm. C.16], it follows that $\tau \in L^p(\mathbb{R}^{2d})$.

Let $\tau \in L^p(\mathbb{R}^{2d})$. Then by (2) and Theorem 2.2, it follows that

(4)
$$||L_{\tau}||_{\mathcal{S}^p} \leq C(\Omega) ||L_{\tau} \star (g \otimes h)||_{L^p} = C(\Omega) ||\tau \star \mathcal{W}(g,h)||_{L^p} \leq C(\Omega) ||\mathcal{W}(g,h)||_{L^1} ||\tau||_{L^p},$$

from the classical Young's inequality for convolutions. Since the cross-Wigner distribution of two Schwartz functions is again a Schwartz function [2, Thm. 11.2.5], it follows that $L_{\tau} \in \mathcal{S}^p$.

Assume now that $\tau \in C_0(\mathbb{R}^{2d})$. Since $C_0(\mathbb{R}^{2d})$ is the L^{∞} -closure of $\mathscr{S}(\mathbb{R}^{2d})$, there exists a sequence $\{\varphi_n\} \subseteq \mathscr{S}(\mathbb{R}^{2d})$ converging uniformly to τ . By (4), the sequence of compact operators $\{L_{\varphi_n}\}$ converges to L_{τ} in the norm topology on $\mathcal{L}(L^2(\mathbb{R}^d))$ which implies that L_{τ} is a compact operator.

If L_{τ} is a compact operator, then there exists a sequence of Schwartz operators $\{L_{\varphi_n}\}$ converging to L_{τ} in the norm topology of $\mathcal{L}(L^2(\mathbb{R}^d))$ by the density of Schwartz operators in the finite rank operators [4]. By (3), it follows that φ_n converges uniformly to τ . This implies that $\tau \in C_0(\mathbb{R}^{2d})$ as it is the uniform closure of $\mathscr{S}(\mathbb{R}^{2d})$. \square

Acknowledgement

The author would like to thank Sigrid Grepstad and Franz Luef for valuable feedback on earlier versions of the manuscript.

References

- R. Fulsche, F. Luef, and R. F. Werner, Wiener's Tauberian theorem in classical and [1]quantum harmonic analysis, 2024. arXiv: 2405.08678 [math.FA].
- K. Gröchenig, Foundations of Time-Frequency Analysis (Applied and Numerical Har-|2|monic Analysis). Boston, MA: Birkhäuser Boston, 2013.
- L. Hörmander, The analysis of linear partial differential operators. I (Classics in Math-3 ematics). Springer-Verlag, Berlin, 2003.
- M. Keyl, J. Kiukas, and R. F. Werner, "Schwartz operators," Rev. Math. Phys., vol. 28, |4|no. 3, pp. 1630001, 60, 2016.
- F. Luef and H. J. Samuelsen, "Fourier Restriction for Schatten Class Operators and $\left[5\right]$ Functions on Phase Space," Int. Math. Res. Not. IMRN, no. 2, Paper No. 291, 22, 2025.
- [6] F. Luef and E. Skrettingland, "Mixed-state localization operators: Cohen's class and trace class operators," J. Fourier Anal. Appl., vol. 25, no. 4, pp. 2064–2108, 2019.
- [7]F. Luef and E. Skrettingland, "A Wiener Tauberian theorem for operators and functions," J. Funct. Anal., vol. 280, no. 6, p. 108 883, 2021.

REFERENCES

- [8] M. Mishra and M. K. Vemuri, *The Weyl transform of a compactly supported distribution*, 2025. arXiv: 2409.16835 [math.CA].
- [9] D. Müller, Bounds on pseudodifferential operators and Fourier restriction for Schatten classes, 2024. arXiv: 2412.05879 [math.CA].
- [10] J. C. T. Pool, "Mathematical aspects of the Weyl correspondence," J. Math. Phys., vol. 7, pp. 66–76, 1966.
- [11] H. J. Samuelsen, "Decoupling for Schatten class operators in the setting of quantum harmonic analysis," *Bull. Lond. Math. Soc.*, vol. 57, no. 1, pp. 23–37, 2025.
- [12] R. Werner, "Quantum harmonic analysis on phase space," J. Math. Phys., vol. 25, no. 5, pp. 1404–1411, 1984.
- [13] M. Zworski, Semiclassical analysis. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2012, vol. 138.

DEPARTMENT OF MATHEMATICAL SCIENCES, NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, TRONDHEIM, NORWAY

Email address: helge.j.samuelsen@ntnu.no