NÉRON MODELS, MINIMAL MODELS, AND BIRATIONAL GROUP ACTIONS

JÁNOS KOLLÁR

ABSTRACT. Let A_K be an Abelian variety over the quotient field of a Dedekind domain R. We show that the identity component of the Néron model of A_K acts regularly on any minimal model of A_K over Spec R, and discuss when the Néron model is an open subset of a minimal model.

The main technical result is the rigidity of small modifications, leading to a regularity criterion for birational group actions.

Let S be a 1-dimensional, irreducible, regular scheme with function field K, and A_K an Abelian variety over K.

Finding a model of A_K over S attracted considerable attention. Kodaira described minimal, elliptic surfaces over \mathbb{C} [Kod60, Kod63]; this was extended to arbitrary Dedekind domains in [Nér64, Chap.III]. For higher dimensional Abelian varieties, the optimal smooth group scheme Nér (A_K) —called the *Néron model*—is constructed in [Nér64, Chap.II]; see Section 2 for a summary.

Minimal model theory—also known as Mori's program—aims to provide a (nonunique) projective, minimal model of any variety or scheme; see (6.4) and (16). For Abelian generic fibers, minimal models give higher dimensional analogs of the minimal, elliptic surfaces of Kodaira and Néron.

The aim of this note is to study actions of group schemes on minimal models, and then derive connections between the Néron model and the minimal models. The main result is Theorem 7, which gives the following for the Néron model.

Theorem 1. Let S be a 1-dimensional, irreducible, regular scheme with function field K, and A_K an Abelian variety over K. Assume that A_K has a projective, minimal model $\pi : A_S \to S$. Let $N\acute{er}^{\circ}(A_K) \subset N\acute{er}(A_K)$ be the identity component of the Néron model (14).

Then the A_K -action on itself extends to a regular Nér^o (A_K) -action on A_S .

Let $p: Y_S \to S$ be a morphism with generic fiber A_K , and $Y_S^{\mathrm{sm}} \subset Y_S$ the *p*-smooth locus. By definition, there is a birational morphism $Y_S^{\mathrm{sm}} \to \operatorname{N\acute{e}r}(A_K)$, which is an isomorphism if Y_S is a minimal, elliptic surface by [Nér64, Chap.III]. In higher dimensions the induced $A_S^{\mathrm{sm}} \to \operatorname{N\acute{e}r}(A_K)$ is an isomorphism in some cases.

Corollary 2. Let $\pi : A_S \to S$ be a projective, minimal model. Then the natural morphism $A_S^{sm} \to N\acute{e}r(A_K)$ is an

(2.1) open embedding, whose image is $Nér(A_K) \setminus (finitely many fibers)$, and an

- (2.2) isomorphism in either of the following cases:
- (2.2.a) A_S is regular, or
- (2.2.b) A_S is factorial and the residue fields of S are perfect.

Previous results 3. There is a vast literature on degenerations of Abelian varieties, focusing on the case of semiabelian reduction, which can be always be

achieved after a suitable finite base change $S' \to S$; see [Mum72, FC90, Kün98, AN99, Ale02, MN24, Nak24] and the references there. However, the situation seems quite different without a base change.

The case (2.2.a) is treated in [HN18, BHPT18, BKV24]; see (23) for details and comments.

The group of birational automorphisms of a projective minimal model is studied in [Han87, 3.3]. A detailed description of Néron models of Lagrangian fibrations over higher dimensional bases is given in [Kim24]. In particular, [Kim24, Thm.1.3] proves that the identity component of the Néron model acts regularly on the total space of a Lagrangian fibration.

In Theorem 1 and Corollary 2 we assume that a minimal model exists, but after that the proofs do not use any minimal model techniques. It is sufficient to know that something called a minimal model with generic fiber A_K exists, it is normal, and (16.4) holds.

One can get further results relying on the construction of additional minimal models involving boundary divisors, and using more information about the occurring singularities. Currently these are known in characteristic 0 (6.4). The main result of this type is the extension of Theorems 1 and 7 to log canonical pairs, given in (51) and (47).

Following [Lan83], a good completion of a Néron model is an open embedding Nér $(A_K) \hookrightarrow Y$ such that Y is normal, projective over S, and the induced rational Nér (A_K) -action on Y is regular; equivariant completion may be a better name. Their existence is stated as conjecture GCNM on [Lan83, p.318], and attributed to Mumford. We prove it in Section 5 in characteristic 0.

Corollary 4. If S is over a field of characteristic 0, then

(4.1) there is a minimal model A_S^* on which Nér (A_K) acts regularly, and

(4.2) Nér $(A_K) \rightarrow S$ has an equivariant completion.

The papers [MN24, Nak24] construct an equivariant completion of the Néron model for the semiabelian reduction cases, but do not check that its singularities are terminal. Using minimal model theory, we get the following.

Corollary 5. Assume that all fibers of $Nér(A_K) \to S$ are semiabelian, and S is over a field of characteristic 0. Then

(5.1) all fibers of $\pi: A_S \to S$ are reduced, semi-log-canonical,

(5.2) Nér (A_K) is isomorphic to the π -smooth locus,

- (5.3) for every $s \in S$, $N\acute{er}(A_K)_s$ acts with a dense, open orbit on A_s , and
- (5.4) A_s is semiabelic in the terminology of [Ale02, 1.1.5].

Remarks 6.

(6.1) Minimal models are not unique, and our methods establish properties that hold for *all* of them. Fixing a suitable polarization on A_K may lead to a unique model A_S . The semiabelian cases are worked out in [Mum72, FC90, Kün98, AN99, Ale02, MN24, Nak24].

(6.2) Usually there is no regular Nér (A_K) -action on a given minimal model; see (21) for some examples.

(6.3) Let $\sum_{i \in I} m_i E_i$ be a fiber of π . If dim $A_K = 1$, then either $m_i = 1$ for some i, or $gcd(m_i : i \in I) > 1$. I do not know whether this holds in higher dimensions.

 $\mathbf{2}$

(6.4) Roughly speaking, minimal models are known to exist in characteristic 0, and are conjectured to exist in general. In characteristic 0 the main references are [HM07, BCHM10, LM22], while [BMP⁺23] settles the case dim $A_K \leq 2$, S excellent, and all residue characteristics are > 5. For Abelian generic fibers, see also [Fuj11, Lai11] for key steps of the proof.

(6.5) By Nagata's theorem, every separated S-scheme X of finite type is isomorphic to an open subscheme of a proper S-scheme \bar{X} ; see [Nag63]. Over a field, equivariant completions are established in [Sum74, Bri22]. The case of smooth, affine group schemes is considered in [Sum75], but it is not clear whether all the assumptions there are necessary. Very little seems to be known about equivariant completions of actions of non-affine group schemes.

(6.6) A stronger form of the existence of equivariant completions is conjectured in [Nak24, 12.4].

(6.7) Y.-J. Kim pointed out that Nér[°](A_K) is isomorphic to the main component of $\mathbf{Aut}_S(A_S)$ using [Kim24, 2.42].

Acknowledgments. I thank V. Alexeev, Y.-J. Kim, R. Laza, S. Molcho and I. Nakamura for comments and references. Partial financial support was provided by the NSF (grant number DMS-1901855) and by the Simons Foundation (grant number SFI-MPS-MOV-00006719-02).

1. Main technical theorems

Theorem 1 is a special case of the following regularity criterion, whose proof is given in (38). It can be viewed as a generalization of [Han87, 3.3] and of [Kim24, Thm.1.3].

Theorem 7. Let S be a normal scheme with function field K, $G_S \to S$ a smooth group scheme with connected fibers, and $\pi : X_S \to S$ a dominant, projective morphism. Assume that X_S has terminal singularities and $K_{X_S/S}$ is π -nef.

Then any rational G_K -action $\mu_K : G_K \times_K X_K \dashrightarrow X_K$ on the generic fiber X_K extends to a regular G_S -action $\mu_S : G_S \times_S X_S \to X_S$.

Note that the statement tacitly assumes that $K_{X_S/S}$ exists. This holds if S is excellent and has a dualizing complex.

Theorem 7 is derived from a more general regularity criterion on rational group actions (37). The full scheme theoretic version is a little technical, the following variant for varieties, to be proved in (32), gives the right idea.

Theorem 8. Let X be a normal, projective variety over an algebraically closed field, and G a connected, smooth algebraic group. Let $\mu : G \times X \dashrightarrow X$ be a rational action. Assume that, for general $g \in G$, the map $\mu_g : X \dashrightarrow X$ is an isomorphism in codimension 1. Then $\mu : G \times X \to X$ is a regular action.

Note that we do not assume that there is an open subset $U \subset X$ where all the μ_q are defined.

9 (Small modifications). Let X_i be normal, proper schemes over S. A birational map $\phi : X_1 \dashrightarrow X_2$ is a *small modification*, if there are closed subsets $Z_i \subset X_i$ of codimension ≥ 2 such that ϕ restricts to an isomorphism $\phi : (X_1 \setminus Z_1) \cong (X_2 \setminus Z_2)$.

The best known examples are flips and flops, much studied when the X_i have log terminal singularities [KM98]. The case of arbitrary normal varieties attracted much less attention [Kol91, Sec.2].

The key technical result behind Theorem 8 is the rigidity of small modifications, see (30) and (36) for the precise statements.

Very little seems to be known about the existence and structure of small modifications, beyond the log terminal case. As the next example shows, there are some unexpected flops of 3-folds with log canonical singularities.

Example 10 (Flopping a fiber of a product). We construct an example of a pointed surface $v \in S$ with simple elliptic (hence log canonical) singularity at v, and an elliptic curve E such that the fiber $E \times \{v\} \subset E \times S$ has infinitely many different flops.

Let $E \subset \mathbb{P}^2$ be a smooth cubic, $S \subset \mathbb{A}^3$ the cone over E with vertex v and projection $\pi: S \dashrightarrow E$. For $m \in \mathbb{Z}$ consider the map

$$\phi_m : E \times S \dashrightarrow E \times S$$
 given by $(p, s) \mapsto (p + m\pi(s), s)$

Then ϕ_m is an isomorphism over $S \setminus \{v\}$, but it is not defined along $E \times \{v\}$ for $m \neq 0$. Note that $E \times S$ has log canonical singularities, and the ϕ_m give infinitely many different flops of the curve $E \times \{v\}$.

2. Néron models

We summarize the properties of Néron models that we use; see [Art86, BLR90] for complete treatments, and [Lor17] for further results.

Notation 11. As explained in [BLR90, p.13], the theory of Néron models is local on the base. Thus, from now on, R denotes a DVR with quotient field K, residue field k and $T := \operatorname{Spec} R$. For a morphism $\pi : X_T \to T$, we use X_K (resp. X_k) to denote the generic (resp. closed or central) fiber.

Definition 12. A smooth morphism $X_T \to T$ satisfies the Néron mapping property iff the following holds.

(12.1) Let $Y_T \to T$ be any smooth morphism. Then every morphism $\phi_K : Y_K \to X_K$ extends to a morphism $\phi_T : Y_T \to X_T$.

A morphism $X_T \to T$ is a *Néron model* of its generic fiber X_K , if it is smooth, of finite type, and satisfies the Néron mapping property. We use Nér (X_K) to denote the Néron model of X_K .

It is clear that X_K uniquely determines its Néron model Nér (X_K) . Néron models commute with étale base changes $T' \to T$ [BLR90, p.13]. Thus many questions about Néron models can be treaded over the strict Henselisation of T.

However, ramification usually changes the Néron model drastically.

[Nér62, Nér64] construct Néron models when X_K is a *torsor* (or *principal homo*geneous space) under an Abelian variety over K. Note that [Art86] and [BLR90] discuss mainly (13.1).

Theorem 13 (Néron models). Let A_K be an Abelian variety over K, and X_K an A_K -torsor. Then the following hold.

(13.1) Nér (A_K) exists, and it is a smooth, finite type group scheme over T.

(13.2) Nér (X_K) exists.

(13.3) Nér
$$(X_K)$$
 is either X_K or a Nér (A_K) -torsor.

Definition 14. The union of A_K and the identity component of $N\acute{e}r(A_K)_k$ is the *identity component* of $N\acute{e}r(A_K)$, denoted by $N\acute{e}r^{\circ}(A_K)$. Note that $N\acute{e}r^{\circ}(A_K)$ is open

4

and dense in Nér (A_K) , and Nér $^{\circ}(A_K) \to T$ is smooth group scheme with connected fibers.

The central fiber $N\acute{e}r(X_K)_k$ is usually disconnected. Its group of connected components is a finite abelian group.

15 (Pseudo-Abelian varieties). When K is not perfact, it is natural to also consider pseudo-Abelian varieties P_K as defined in [Tot13]. These also have Néron models Nér (P_K) by [BLR90, Sec.10.2]; see also [Ove24] for further results. Theorem 7 also applies to actions of Nér (P_K) .

3. MINIMAL MODELS AND NÉRON MODELS

An introduction to minimal models is in [KM98], and a summary of later results is in [Kol23, Secs.11.1–2]. For the purposes of Theorem 1 we need only the following special setting.

16 (Minimal models of Abelian torsors). Let S be a 1-dimensional, irreducible, regular scheme with function field K. Let X_K be a proper K-variety with terminal singularities, whose canonical class is numerically trivial.

A proper morphism $\pi: X_S \to S$ is a minimal model of X_K if

(16.1) X_S has terminal singularities [KM98, 2.34],

(16.2) K_{X_S} is numerically π -trivial, and

(16.3) the generic fiber is isomorphic to X_K .

Numerically π -trivial means that $(K_{X_S} \cdot C) = 0$ for every curve contracted by π .

Being terminal is preserved by étale base change [Kol13, 2.15], so being a minimal model is also preserved by étale base change. Note that minimal model theory frequently works with \mathbb{Q} -factorial schemes, that is, some multiple of any Weil divisor is required to be Cartier. However, being \mathbb{Q} -factorial is not an étale-local property, so, as in [Kol21], we work without it.

Minimal models of X_K need not be unique, but the following is easy to see directly; see [KM98, 3.52].

Claim 16.4. Let B be a normal scheme and $\pi_i : X_i \to B$ proper morphisms. Assume that X_i is irreducible, has terminal singularities, and $K_{X_i/B}$ is π_i -nef for i = 1, 2. Let $\phi : X_1 \dashrightarrow X_2$ be a birational map.

Then there are closed $Z_i \subset X_i$ of codimension ≥ 2 , such that ϕ induces an isomorphism $X_1 \setminus Z_1 \cong X_2 \setminus Z_2$.

For the existence of minimal models, the main references are [HM07, BCHM10], with additional steps in [Fuj11, Lai11, LM22].

Theorem 16.5. Assume that S is over a field of characteristic 0. Let X_K be a projective K-variety with terminal singularities, whose canonical class is numerically trivial. Then X_K has a \mathbb{Q} -factorial, projective, minimal model $\pi : X_S \to S$.

In particular, every Abelian torsor over K has a minimal model.

The following generalization of Theorem 1 connects minimal models and Néron models. We follow Notation 11, thus T is the spectrum of a DVR R with quotient field K and residue field k.

Theorem 17. Let X_K be a torsor under an Abelian variety A_K , and $\pi : X_T \to T$ a minimal model of X_K . Let $N\acute{er}^{\circ}(A_K) \subset N\acute{er}(A_K)$ be the identity component of the

Néron model. Then the A_K -action on X_K extends to a regular Nér° (A_K) -action on X_T .

Proof. The A_K -action on X_K extends to a rational Nér^o (A_K) -action on X_T which we write as

$$\mu: \operatorname{N\acute{e}r}^{\circ}(A_K) \times_T X_T \dashrightarrow X_T.$$
(17.1)

Being an morphism can be checked after a surjective, étale base change. We may thus assume that T is strictly Henselian.

Given any section $\tau : T \to \operatorname{N\acute{e}r}^{\circ}(A_K)$, the induced map $\mu_{\tau} : X_T \dashrightarrow X_T$ is a small modification by (16.4). Thus μ is a morphism by (37).

The Nér^{\circ}(A_K) action gives strong restrictions on the smooth locus.

Lemma 18. Let $U_T \to T$ be a smooth morphism with a regular $\operatorname{N\acute{e}r}^\circ(A_K)$ action, whose generic fiber U_K is an A_K -torsor. Then $u : U_T \to \operatorname{N\acute{e}r}(U_K)$ is an open immersion, and the complement of the image is a union of $\operatorname{N\acute{e}r}^\circ(A_K)_k$ -orbits.

Proof. By the universal property of Néron models there is a birational morphism $u : U_T \to \text{Nér}(U_K)$, which is $\text{Nér}^{\circ}(A_K)$ -equivariant. The restriction of u gives a $\text{Nér}^{\circ}(A_K)_k$ -equivariant morphism of U_k to $\text{Nér}(U_K)_k$, which is a union of $\text{Nér}^{\circ}(A_K)_k$ -orbits. Thus u_k is an isomorphism onto an union of connected components of $\text{Nér}(X_K)_k$.

The following is a generalization of Corollary 2.

Corollary 19. Continuing with the notation of (17), let $X_T^{sm} \subset X_T$ denote the π -smooth locus, and $u: X_T^{sm} \to \operatorname{N\acute{e}r}(X_K)$ the induced morphism. Then

(19.1) either u is an isomorphism,

(19.2) or $X_T^{\mathrm{sm}} \cong X_K$.

Furthermore, u is an isomorphism in either of the following cases.

(19.3) X_T is regular.

(19.4) X_T is factorial and k is perfect.

Proof. We may assume that T is strictly Henselian. If X_k^{sm} is empty, we have (19.2). Otherwise there is at least one irreducible component $E_1 \subset X_k$, such that u maps $E_1 \cap X_k^{\text{sm}}$ onto an irreducible component $F_1 \subset \text{Nér}(X_K)_k$.

Let $F_2 \subset \operatorname{Nér}(X_K)_k$ be any other irreducible component. There is a section $\tau: T \to \operatorname{Nér}(A_K)$ that maps F_1 onto F_2 . This gives a rational map $\mu_{\tau}: X_T \dashrightarrow X_T$ that sends E_1 to another irreducible component $E_2 \subset X_k$ by (16.4). Then u maps $E_2 \cap X_k^{\mathrm{sm}}$ onto F_2 . Thus (19.1) holds by (18).

It remains to show that X_k^{sm} is not empty in cases (19.3–4). Let $C \subset X_T$ be a section. Then $(X_k \cdot C) = 1$, and $(\operatorname{red} X_k \cdot C) \geq 1$ if $\operatorname{red} X_k$ is a Cartier divisor. Thus X_k is reduced near $X_k \cap C$, hence geometrically reduced if k is perfect.

If X_T is regular, then $(X_k \cdot C) = 1$ implies that X_k is regular at the k-point $X_k \cap C$, hence smooth. These show (19.3–4).

The next examples show that in general Nér^{\circ}(A_K) is not isomorphic to an open subset of any minimal model, and that minimal models are not unique.

Example 20. Let *B* be an Abelian variety and σ an automorphism of order *m* fixing the identity. Set $X := X(B, \sigma) := B \times \mathbb{A}^1/(\sigma, \epsilon)$, where ϵ is a primitive *m*th root of unity. The central fiber has multiplicity *m*, and the reduced central fiber is the Kummer-type variety $B/\langle \sigma \rangle$. The singularities of *X* are analytically quotient

 $\mathbf{6}$

singularities. The Reid-Thai criterion determines whether they are terminal [Rei80]. If they are, then $X \to \mathbb{A}^1$ is a minimal model which does not contain the Néron model. Let us see some concrete examples.

(20.1) Let $\sigma := -1$ and $n = \dim B$. The singularities of X are analytically of the form $\mathbb{A}^{n+1}/\frac{1}{2}(1, \cdots, 1)$. These are terminal for $n \ge 2$, and $K_X \sim 0$ iff n is odd. The central fiber of the Néron model is $\mathbb{G}^n_a \times (\mathbb{Z}/2)^{2n}$.

(20.2) Set $E = (x^3 + y^3 + z^3 = 0)$ with $\mathbb{Z}/3$ -action $(x:y:z) \mapsto (\epsilon x:y:z)$. Set $B = E \times E$ and let $\mathbb{Z}/3$ -act as $(\epsilon, \epsilon, \epsilon^2)$. There are 9 fixed points, all are analytically of the form $\mathbb{A}^3/\frac{1}{3}(1, 1, 2)$. These are terminal.

(20.3) With $B = E \times E$ as above, let $\mathbb{Z}/3$ -act as $(\epsilon, \epsilon, \epsilon)$. There are 9 fixed points, all are analytically of the form $\mathbb{A}^3/\frac{1}{3}(1,1,1)$. These are canonical but not terminal. A minimal model is obtained by blowing them up. We have now 9 reduced divisors in the certal fiber, which cointains the Néron model. The central fiber of the Néron model is $\mathbb{G}_a^2 \times (\mathbb{Z}/3)^2$.

(20.4) Let B be a simple abelian variety of dimension ≥ 4 . Then $B/\langle \sigma \rangle$ has canonical singularities by [KL09, Thm.9], hence $X(B, \sigma)$ has terminal singularities, and it is a minimal model.

Example 21. Let $g_i(x, y, z)$ be 2 general cubics and $\pi : X \to \mathbb{A}^1$ the fiber product of the 2 elliptic surfaces $(xyz + tg_i(x, y, z) = 0) \subset \mathbb{P}^2 \times \mathbb{A}^1$.

 K_X is trivial and the general fiber is a product of 2 elliptic curves. The central fiber is a union of 9 copies $L_{ij} \cong \mathbb{P}^1 \times \mathbb{P}^1$. X has 9 singular points of the form (xy = uv) where 4 of the L_{ij} meet. The central fiber of the Néron model is $\mathbb{G}_m^2 \times (\mathbb{Z}/3)^2$. It acts regularly on X, transitively on both the L_{ij} and the singular points.

Each singular point has 2 small resolutions, giving 3⁹ different minimal models, among them 2⁹ are smooth. The projective ones can be obtained by blowing up a divisor $\sum_{ij} m_{ij} L_{ij}$. For example, $\sum_i L_{ii}$ gives a projective, smooth minimal model.

There are 2 smooth minimal models on which the whole Néron model acts, but neither is projective.

Example 22. As a variant of (21), work over the base field \mathbb{Q} and start with the elliptic surfaces

$$S_a := (x^3 + ay^3 + a^2z^3 - 3axyz = tg(x, y, z)),$$

whose central fiber is a union of the 3 lines $(x + \alpha y + \alpha^2 z = 0)$ for $\alpha^3 = a$.

Now consider the fiber product of S_a and S_c , say for a = 2, c = 3. The central fiber is the union of 9 divisors, which are conjugate to each other. It has 9 singular points, and no projective, smooth, small modifications.

23 (Kulikov models). Working over \mathbb{C} , the papers [HN18, BHPT18, BKV24] state that the π -smooth locus of a *Kulikov model* π : Kul $(A_K) \to T$ of an Abelian variety A_K is isomorphic to its Néron model.

Note however, that [BHPT18, BKV24] and [HN18] use non-equivalent definitions of a Kulikov model.

In all 3 papers, a Kulikov model is a flat, proper morphism $\pi : Y \to T$, where Y is regular, and the general fiber has trivial canonical class (in our case, an Abelian variety). The additional assumptions are, however, quite different.

In [HN18, 5.1.2] the central fiber Y_0 is required to be a normal crossing divisor, and the log canonical class $K_{Y/T}$ (red Y_0) is assumed to be numerically π -trivial. [BKV24, 2.26] refers to [BHPT18, 3.3]. In the latter the central fiber of a Kulikov model is allowed to have arbitrary singularities, and $K_{Y/T}$ is assumed to be numerically π -trivial; see [BHPT18, 3.2].

The arguments in [BHPT18, BKV24] rely on [HN18], sliding over the incompatibility of the definitions.

Note also that [BHPT18, 3.3] refers to [HN18, 6.7]. There does not seem to be any entry numbered (6.7) in [HN18]; neither in the published version nor in any of the three ArXiV versions.

4. Small modifications

We work over a Noetherian base scheme. To avoid some possible problems with dimension, let us assume that it is universally catenary.

Definition 24 (Families of birational maps). A birational map $\phi : X \dashrightarrow Y$ between (reduced, separated) schemes X, Y is an (equivalence class of) isomorphisms between dense, open subsets $X \supset U_X \cong U_Y \subset Y$. The largest such is called the *domain of isomorphy* of ϕ , denoted by domi(ϕ), and usually viewed as a subset of X.

Let $p_i : X_i \to S$ be equidimenisonal, proper morphisms. A birational map $\phi : X_1 \dashrightarrow X_2$ over S is a *family of birational maps* if domi(ϕ) contains the generic points of each fiber. Thus the restrictions $\phi_s : \operatorname{red}(X_1)_s \dashrightarrow \operatorname{red}(X_2)_s$ are birational for every $s \in S$.

Warning 24.1. If each ϕ_s is an isomorphism, then ϕ is an isomorphism over a dense, open subset of S by (25.2), but it can happen that ϕ is not an isomorphism, see (27.2) for such an example.

25 (Restriction of birational maps). Let $\phi : X \dashrightarrow Y$ be a birational map, and $W_X \subset X$ and $W_Y \subset Y$ closed subsets such that ϕ restricts to a birational map $\phi_W : W_X \dashrightarrow W_Y$. Then $\operatorname{domi}(\phi) \cap W_X \subset \operatorname{domi}(\phi_W)$, but they need not be equal. That is, it can happen that ϕ_W is an isomorphism at a point $x \in W_X$, but ϕ is not defined at x.

It is useful to know that this does not happen if W_X is in general position.

Let S be a normal scheme and $\pi: X \to S$ an irreducible, normal scheme, proper over S, with geometrically irreducible generic fiber. There is a dense, open $S^{\circ} \subset S$ over which the fibers are geometrically irreducible. Set $S_1 := S \setminus S^{\circ}$. For $i \geq 2$ let $S_i \subset S$ be the closed subset over which the fiber dimension is $\geq i-1+\dim X - \dim S$. By a dimension count we see the following.

Claim 25.1. Let $W \subset S$ be an irreducible, local complete intersection of codimension m. If $\operatorname{codim}(W \cap S_i, S_i) \geq m$ for every i, then $\pi^{-1}(W)$ is irreducible. \Box

Next let $\pi_i : X_i \to S$ be irreducible, normal schemes, proper over S, and $\phi : X_1 \dashrightarrow X_2$ a birational map with closed graph $\Gamma(\phi)$. Applying (25.1) to the $X_i \to S$ and to $\Gamma(\phi) \to S$, we get the following.

Claim 25.2. There are closed subsets $S_i \subset S$ with the following property.

Let $Z \subset S$ be an irreducible, local complete intersection of codimension m such that $\operatorname{codim}(Z \cap S_i, S_i) \geq m$ for every i. Then ϕ restricts to a birational map $\phi_Z : \operatorname{red} \pi_1^{-1}(Z) \dashrightarrow \operatorname{red} \pi_2^{-1}(Z)$, and $\operatorname{domi}(\phi) \cap \pi_1^{-1}(Z) = \operatorname{domi}(\phi_Z)$. \Box

We are especially interested in cases when $\operatorname{domi}(\phi)$ is large.

Definition 26 (Small modification). An open subset $U \subset X$ is called *large* if $X \setminus U$ has codimension ≥ 2 in X. A birational map $\phi : X_1 \dashrightarrow X_2$ is a *small modification* if domi(ϕ) is large in both X_1 and X_2 .

The composite of small modifications is a small modification.

A family of birational maps $\phi : X_1 \dashrightarrow X_2$ over S is a family of small modifications if $\phi_s : (X_1)_s \dashrightarrow (X_2)_s$ is a small modification for every $s \in S$.

Remark 27. Given $X_i \to S$, let $\phi : X_1 \dashrightarrow X_2$ be a small modification. By definition there are large open subsets $U_i \subset X_i$ such that ϕ restricts to an isomorphism $\phi_U : U_1 \cong U_2$. There is a dense, open subset $S^\circ \subset S$ such that $(U_i)_s \subset (X_i)_s$ are large open subsets for $s \in S^\circ$.

Thus ϕ is a family of small modifications over S° , but not always over S. The following example illustrates some of the possibilities.

Start with the deformation of the quadric cone

$$X:=(x_1^2-x_2^2+x_3^2=t^2)\subset \mathbb{P}^3_x\times \mathbb{A}^1_t$$

It has 2 small simultaneous resolutions. We get $\pi_1 : X_1 \to X$ by blowing up $(x_1 - x_2 = x_3 - t = 0)$, and $\pi_2 : X_2 \to X$ by blowing up $(x_1 + x_2 = x_3 - t = 0)$.

- (27.1) The $\pi_i: X_i \to X$ are small modifications, but not families of small modifications over \mathbb{A}^1 .
- (27.2) The composite $\phi := \pi_2^{-1} \circ \pi_1 : X_1 \dashrightarrow X_2$ is a family of small modifications, where each ϕ_t is an isomorphism. However ϕ is not an isomorphism.

Recall that a divisor $D \subset X$ is π -exceptional for $\pi : X \to S$ if $\pi(D) \subset S$ has codimension ≥ 2 .

Lemma 28. Let S be a normal scheme, $\pi_i : X_i \to S$ irreducible, normal schemes, proper over X, and $\phi : X_1 \dashrightarrow X_2$ a birational map. Assume that there are no π_i -exceptional divisors. Then ϕ is a small modification iff $\phi_s : (X_1)_s \dashrightarrow (X_2)_s$ is (28.1) a small modification for the generic point $s \in S$, and

(28.2) birational for each codimension 1 point $s \in S$.

The following isomorphism criterion is proved in [MM64]; see [Kol23, 11.39] for the relative version.

Theorem 29 (Matsusaka-Mumford criterion). Let $\pi_i : X_i \to S$ be normal schemes, projective over S, and $\phi : X_1 \dashrightarrow X_2$ a small modification. Let H_1 be a π_1 -ample \mathbb{Q} -Cartier divisor. Assume that $H_2 := \phi_*(H_1)$ is \mathbb{Q} -Cartier and π_2 -ample.

Then ϕ is an isomorphism.

Next we show that small, projective, normal modifications are rigid. It is likely that the same holds for proper modifications, but the proof uses projectivity.

Theorem 30. Let X, S be geometrically irreducible and geometrically normal kvarieties, $Y \to S$ a projective morphism with geometrically normal generic fiber, and $\phi: X \times S \dashrightarrow Y$ a small modification. Assume that

(30.1) either ϕ is a family of birational maps,

(30.2) or S has rational singularities.

Then there is a unique, small, projective modification $\sigma: X \dashrightarrow X'$ such that the composite

 $\phi \circ (\sigma, 1_S)^{-1} : X' \times S \dashrightarrow X \times S \dashrightarrow Y$ is an isomorphism.

Example 10 shows that assumption (30.2) is necessary.

Proof. Assume first that the base field is algebraically closed. As we noted in (25.2), there is a dense, open subset $S^{\circ} \subset S$ such that $Y \to S$ is flat with geometrically normal fibers over S° , and $\operatorname{domi}(\phi^{-1}) \cap Y_s = \operatorname{domi}(\phi_s^{-1})$ for $s \in S^{\circ}$, and ϕ is a family of small modifications over S° .

Pick $s \in S^{\circ}$ and take $X' := Y_s$. The composite map

$$\psi := \phi \circ (\phi_s, 1_S)^{-1} : X' \times S \dashrightarrow X \times S \dashrightarrow Y$$

is then a small modification such that $\operatorname{domi}(\psi) \cap X' \times \{s\} \subset X' \times \{s\} \cong Y_s$ is large, and $\psi_s : Y_s = X' \times \{s\} \dashrightarrow Y_s$ extends to the identity map.

Let $H'_s \subset X'$ be a normal, ample divisor (33.1). Set $H' := H'_s \times S$ and $H := \psi_*(H')$ its birational transform. Note that $H_s = H'_s$. Next (34) says that H is Cartier along Y_s . Since $H|_{X_s} = H'_s$ is ample, H is relatively ample over S° , after shrinking S° as needed. Then (29) shows that ψ is an isomorphism over S° .

Thus $\psi : X' \times S \dashrightarrow Y$ is a small modification, which is an isomorphism over S° . Let A be an ample divisor on Y and $A' := \psi_*^{-1}(A)$ its birational transform.

Since ψ is an isomorphism over S° , A' is Cartier and ample over S° .

If ϕ is a family of birational maps, then suppA' does not contain any fiber X'_s , so A' is a family of Weil divisors. Thus A' is Cartier and ample over S by (33.2). We can now use (29) again to get that ψ is an isomorphism.

Note that A' gives a morphism $S^{\circ} \to \operatorname{Pic}(X')$, which extends to a morphism $S \to \operatorname{Pic}(X')$ if S has rational singularities by (35). Then A' is Cartier and ample over S. As before, ψ is an isomorphism by (29).

For an arbitrary base field k, we can choose the above $s \in S(\bar{k})$ to be Galois over k. Then $Y_{\bar{k}} \cong Y_s \times S_{\bar{k}}$, thus the conjugate fibers of $Y_{\sigma(s)}$ are isomorphic to Y_s . Therefore Y_s is defined over k.

Remark 31. Note that there may be several ways of writing Y as $X' \times S$. For example, if G is an algebraic group and X is a nontrivial G-torsor, then $G \times X \cong X \times X$.

The claim of (30) is that the rational product structure ϕ can be promoted to a regular product structure in a unique way.

32 (Proof of Theorem 8). Take 2 copies $Y_i := X \times G$ and consider the rational map

 $\phi: Y_1 \dashrightarrow Y_2$ given by $(x,g) \mapsto (\mu(g,x),g)$.

Let $\Gamma \subset Y_1 \times_G Y_2$ be its closed graph. By (25.2) there is a dense, open subset $U \subset G$ such that, for $g \in U$, the fiber Γ_g equals the closed graph of $\mu_g : X \dashrightarrow X$. Thus ϕ is a family of small modifications over U by (25.2).

By (30) there is a small, projective modification $\sigma: X \dashrightarrow X'$ such that the composite

 $\psi: (\sigma, 1_U) \circ \phi: X \times U \dashrightarrow X \times U \dashrightarrow X' \times U$ is an isomorphism.

Thus, we get that the composite

$$\mu_h^{-1} \circ \mu_g = \psi_h^{-1} \circ \psi_g : X \dashrightarrow X$$

is an isomorphism for $g, h \in U$. Since G is connected, $(h, g) \mapsto h^{-1}g$ is a surjective map $U \times U \to G$. Hence μ_g is an isomorphism for all $g \in G$.

As we noted in (24.1), this implies that ϕ is an isomorphism over a dense, open subset of G. The G-action now shows that it is an isomorphism everywhere.

10

33 (Divisors on normal varieties). We used two results about normal, projective varieties X.

(33.1) Let |H| be a very ample linear system. Then a general $H \in |H|$ is normal; see [Gro60, IV.12.1.6].

(33.2) $\operatorname{Pic}^{\circ}(X)$ is proper. Thus if $D \subset X \times S$ is a family of Weil divisors that is Cartier along the generic fiber, then it is Cartier. Over \mathbb{C} this goes back to Picard and Severi, the general case is in [Mat52]; see [KLOS23, Sec.11.3] for a detailed discussion.

34 (Lifting normal, Cartier divisors). Let Y be a normal scheme, $X \subset Y$ a Cartier divisor and D an effective Weil divisor on Y. Assume that the *divisorial restriction* $D|_X^{\text{div}}$ is normal and Cartier. Then D is Cartier along X.

Recall that the divisorial restriction is obtained by first taking the schemetheoretic restriction $\mathcal{O}_D|_X$, and then the quotient by its torsion subsheaf; see [Kol23, p.106] for details.

Indeed, since $D|_X^{\text{div}}$ is normal, there is a closed $Z \subset X$ of codimension ≥ 3 such that $D|_X^{\text{div}}$ is regular on $X \setminus Z$. Since $D|_X^{\text{div}}$ is Cartier, X and hence Y are regular along $(D \cap X) \setminus Z$. Thus D is Cartier along $X \setminus Z$.

Next note that depth_Z $(D|_X^{\text{div}}) \ge 2$ since $D|_X^{\text{div}}$ is normal, hence depth_Z $(X) \ge 3$ since $D|_X^{\text{div}}$ is Cartier. Thus D is Cartier along Z by [Gro68, XI.2.2]; see also [Kol23, 10.73].

35 (Rational maps to Abelian varieties). Let X be a normal variety, A an Abelian variety and $\phi : X \dashrightarrow A$ a rational map. We would like to know that it is necessarily a morphism.

If X is smooth, this is [Wei48, II.15.Prop.1]. For the general case, let $Y \to X \times A$ be the normalization of the closed graph of ϕ . Let L be the pull-back of the universal bundle from $A \times \operatorname{Pic}^{\circ}(A)$ to $Y \times \operatorname{Pic}^{\circ}(A)$.

If $p: Y \to X$ is not an isomorphism, then there is a curve $C \subset Y$ whose second projection to A is not constant. Thus the restriction of L to $C \times \operatorname{Pic}^{\circ}(A)$ is nonconstant.

On the other hand, the restriction of L to $Y \times [\mathcal{O}_A]$ is \mathcal{O}_Y . Since $R^1 p_* \mathcal{O}_Y = 0$, the constant 1 section of \mathcal{O}_Y lifts to a section of L in a neighborhood of $[\mathcal{O}_A] \in \mathbf{Pic}^{\circ}(A)$. This is a contradiction.

We also need a scheme theoretic version of (30). The proof is essentially the same, but seems to need more assumptions and different references. Also, it is not clear to me how to descend projectivity to arbitrary base schemes from the strictly Henselian case, so we state only the latter.

Theorem 36. Let S be a normal, local, strictly Henselian scheme, X a normal S-scheme, and $W \to S$ a smooth morphism with geometrically connected fibers. Let $Y \to W$ be a projective morphism and $\phi : X \times_S W \dashrightarrow Y$ a small modification. Assume that

(36.1) $Y_{\tau} := S \times_{\tau, W} Y$ is normal for every section $\tau : S \to W$, and

(36.2) the induced map $\phi_{\tau}: X \dashrightarrow Y_{\tau}$ is a small modification.

Then there is a small, projective modification $\sigma: X \dashrightarrow X'$ such that the composite

 $\phi \circ (\sigma, 1_W)^{-1} : X' \times_S W \dashrightarrow X \times_S W \dashrightarrow Y$ is an isomorphism.

Proof. Note that section of $W \to S$ are local complete intersections. Thus by (25.2) there is section $\tau : S \to W$ such that $\operatorname{domi}(\phi^{-1}) \cap Y_{\tau} = \operatorname{domi}(\phi_{\tau}^{-1})$. Set

 $X' := Y_{\tau}$ and consider the composite map

$$\psi := \phi \circ (\phi_{\tau}, 1_W)^{-1} : X' \times_S W \dashrightarrow X \times_S W \dashrightarrow Y.$$

Then domi(ψ) contains a large open subset of $X' \times_S \tau(S)$, and

$$\psi_{\tau}: X' \times_S \tau(S) \cong X' = Y_{\tau} \dashrightarrow Y_{\tau}$$

extends to the identity map.

Let $H'_{\tau} \subset X'$ be a normal, relatively ample divisor (33.1). Set $H' := H'_{\tau} \times_S W$ and $H := \psi_*(H')$ its birational transform on Y. Note that $H_{\tau} = H'_{\tau}$. As before, (34) implies that H is Cartier along Y_{τ} .

Since $H|_{X_{\tau}} = H'_{\tau}$ is relatively ample, H is relatively ample over a suitable open subset $\tau(S) \subset W^{\circ} \subset W$. Then (29) shows that ψ is an isomorphism over W° , and W° dominates S since it contains a section.

Thus $\psi : X' \times_S W \dashrightarrow Y$ is a small modification, which is an isomorphism over W° . Let A be an ample divisor on Y and $A' := \psi_*^{-1}(A)$ its birational transform.

Since ψ is an isomorphism over W° , A' is Cartier and ample over W° . Thus A' is Cartier by (39), and ample over W. We can now use (29) again to get that ψ is an isomorphism.

Proposition 37. Let S be a normal scheme, X a normal, projective S-scheme, and $G \to S$ a smooth group scheme with irreducible fibers. Let $\mu_S : G \times_S X \dashrightarrow X$ be a rational action. Assume that for every étale section $\tau : S' \to G$ (that is, $S' \to G \to S$ is étale), the induced map

$$\tau_X : X_{S'} = (S' \times_S X) \xrightarrow{(1_{S'}, \tau, 1_X)} S' \times_S G \times_S X \xrightarrow{(1_{S'}, \mu_S)} (S' \times_S X) = X_{S'}$$
(37.1)

is a small modification. Then $\mu_S : G \times_S X \dashrightarrow X$ is a regular action.

Proof. Being a morphism is an étale-local property, we may thus assume that S is local and strictly Henselian.

Take 2 copies $Y_i := X \times_S G$ and consider the rational map

 $\phi: Y_1 \dashrightarrow Y_2$ given by $(x,g) \mapsto (\mu(g,x),g)$.

By (37.1) the assumption (36.2) is satisfied. Thus by (36) there is a small, projective modification $\sigma: X \dashrightarrow X'$ such that the composite

$$\psi: (\sigma, 1_G) \circ \phi: X \times_S G \dashrightarrow X \times_S G \dashrightarrow X' \times_S G \tag{37.2}$$

is an isomorphism. Let $\tau_i : S \to G$ be general sections. Restricting (37.2) to $\tau_i(S)$ and composing, we get that

$$\mu_{\tau_1}^{-1} \circ \mu_{\tau_2} = \psi_{\tau_1}^{-1} \circ \psi_{\tau_2} : X \dashrightarrow X$$

is an isomorphism. Since G has connected fibers, these $\mu_{\tau_1}^{-1} \circ \mu_{\tau_2}$ generate it. Thus, by (25.2), μ_S is an isomorphism over an open dense set of G that contains a section. By the G-action, it is an isomorphism everywhere.

38 (Proof of Theorem 7). Let $\tau : S' \to G$ be an étale section. As in (37.1) we get a birational map $\tau_X : X_{S'} \dashrightarrow X_{S'}$. Also, $X_{S'}$ has terminal singularities and $K_{X_{S'}}/S$ is reletively nef. Thus τ_X is a small modification by (16.4). Thus $\mu_S : G \times_S X \dashrightarrow X$ is regular by (37).

The following can be viewed as a generalization of (33.2); for proofs see [Gro60, IV.21.14.1] or [Kol23, 10.65].

Theorem 39 (Ramanujam-Samuel). Let Z be a normal scheme and $g: X \to Z$ a smooth morphism. Let D be a Weil divisor on X that is Cartier on a dense open subset of every fiber of g. Then D is Cartier.

5. Equivariant completions

We prove Corollary 4 in 2 steps.

40 (Proof of 4.1). This part uses characteristic 0.

Let $\pi : X_T \to T$ be a Q-factorial minimal model of A_K , and H_0 a π -ample divisor on X_T .

By (16.4) Nér (A_K) acts on the Néron-Severi group of X_T , and Nér (A_K) acts trivially. So the Nér (A_K) -orbit of $[H_0] \in NS(X_T)$ is finite. Choose representatives $\{H_i : i \in I\}$ from each divisor class in the orbit, and set $H := \sum_{i \in I} H_i$. Let $X_T^c \to T$ be the canonical model of (X_T, H) . That is

$$X_T^{\mathrm{c}} := \operatorname{Proj}_T \oplus_m \pi_* \mathcal{O}_{X_T}(mH),$$

provided this sheaf of algebras is finitely generated. A technical problem is that minimal model theory is usually formulated for effective divisors, and it is not clear that the H_i can be chosen effective. However, finite generation can be checked after an étale base change by [KM98, 6.6], so it is enough to show that X_T^c exists when T is strictly Henselian.

In this case there are sections $\sigma_i : T \to \operatorname{N\acute{e}r}(A_K)$ passing through each irreducible component of the central fiber. We can then take $H_i := \sigma_i(H_0)$ and choose $0 < \epsilon \ll$ 1 such that $(X_T, \epsilon H)$ is log terminal. Then X_T^c is its canonical model by [KM98, 3.52], and $X_T \dashrightarrow X_T^c$ is a small modification.

Then H descends to a relatively ample divisor H^c on X_T^c , and its class $[H^c] \in NS(X_T^c)$ is Nér (A_K) -invariant. Thus the Nér (A_K) -action is regular on X_T^c by (29).

41 (Proof of $4.1 \Rightarrow 4.2$). The arguments work over any 1-dimensional, regular base scheme.

We have $Y_0 := X_T^c \to T$ with a Nér (A_K) -action and a section $e_0 : T \to Y_0$, giving $p_0 : \text{Nér}(A_K) \to Y_0$ that is an isomorphism on the generic fiber.

If Y_i is aready defined with a Nér (A_K) -action and a section $e_i : T \to Y_i$, let $p_i : \text{Nér}(A_K) \to Y_0$ be the corresponding morphism.

Let $Z_i \subset Y_i$ be the closure of the image of the central fiber $N\acute{e}r(A_K)_k$. If Z_i is a divisor and X_i is regular at the generic points of Z_i , then p_i restricts to an open embedding $N\acute{e}r(A_K)_k \hookrightarrow Z_i$, and so $p_i : N\acute{e}r(A_K) \to Y_0$ is an open embedding. Thus the normalization of Y_i is an equivariant completion of $N\acute{e}r(A_K)$.

Otherwise let $Y_{i+1} := B_{Z_i}Y_i$ be the blow-up of Z_i . Since Z_i is Nér (A_K) -equivariant, the Nér (A_K) -action on Y_i lifts to a Nér (A_K) -action on Y_{i+1} , and the section e_i also lifts to $e_{i+1}: T \to Y_{i+1}$.

By a lemma of Abhyankar and Zariski (cf. [KM98, 2.45]), the process eventually stops, and we get the required equivariant completion of Nér (A_K) .

6. Semiabelian reduction

The correspondence between the minimal and Néron models is especially strong if A_K has semiabelian reduction, that is, the central fiber of Nér^o (A_K) is an extension of an Abelian variety by a torus.

One can work with log terminal minimal models. That is, (16.2-3) hold, but X_T is allowed to have log terminal singularities. Using minimal model theory, we get the following in characteristic 0.

Theorem 42. Let A_K be an Abelian variety, and $\pi: X_T \to T$ a log terminal minimal model of A_K with central fiber X_k . Let A_k denote the identity component of the central fiber of $N\acute{e}r(A_K)$. The following are equivalent.

(42.1) A_k is a semiabelian variety.

(42.2) X_k is reduced, and the pair (X_T, X_k) is log canonical.

If (42.2) holds then $\pi: X_T \to T$ is *locally stable* in the terminology of [Kol23]. We start the proof with some auxiliary lemmas.

Lemma 43. Let $\pi : Y_T \to T$ be a proper morphism such that Y_T is normal and K_{Y_T} is Q-Cartier. Let $E \subset Y_k$ be an irreducible component of multiplicity 1. Let $E^* \subset E$ be the open set where Y_k is smooth, and $D \subset E$ its complement. Let $\tau: \overline{E} \to E$ be normalization and $\overline{D} \subset \overline{E}$ the divisorial part of the preimage of D. Then there is an effective \mathbb{Q} -divisor D^* suported on \overline{D} such that

$$K_{\bar{E}} + \bar{D} + D^* \sim_{\mathbb{Q}} \tau^* (K_{Y_T} + Y_k) \sim_{\mathbb{Q}} \tau^* K_{Y_T}.$$

$$(43.1)$$

Proof. Write $\Delta := Y_k - E$, as a divisor on Y_T . By [Kol13, 4.2], there is an effective divisor $\operatorname{Diff}_{\bar{E}}\Delta$ such that

$$K_{\bar{E}} + \operatorname{Diff}_{\bar{E}} \Delta \sim_{\mathbb{Q}} \tau^* (K_{Y_T} + Y_k) \sim_{\mathbb{Q}} \tau^* K_{Y_T}, \qquad (43.2)$$

Let $D_i \subset D$ be an irreducible component, and $\overline{D}_i \subset \overline{D}$ its preimage.

If Y_k is singular along D_i , then (Y_T, Y_k) is not klt along D_i , hence \overline{D}_i appears in Diff_E Δ with coefficient ≥ 1 by [Kol13, 2.35.5–6]. We can thus set $D^* :=$ $\operatorname{Diff}_{\bar{E}}\Delta - \bar{D}.$

The following is a special case of [BZ17].

Lemma 44. Let A be a connected, commutative algebraic group with unipotent radical $U \subset A$. Let $\overline{A} \supset A$ be a normal, A-equivariant compactification, and $\overline{D} \subset \overline{A}$ the divisorial part of $\overline{A} \setminus A$.

(44.1) If $U \neq \{0\}$, then $K_{\bar{A}} + \bar{D}$ is not pseudo-effective.

(44.2) If $U = \{0\}$, then $K_{\bar{A}} + \bar{D} \sim 0$ and (\bar{A}, \bar{D}) is log canonical.

45 (Proof of (42.2) \Rightarrow (42.1)). We check in (51) that there is a regular Nér[°](A_K)action on X_T .

Apply (43) to $Y_T := X_T$ and any irreducible component $E \subset X_k$. Note that $E^* \subset E$ is the dense A_k -orbit by (18).

Then (43.2) gives that $K_{\bar{E}} + \text{Diff}_{\bar{E}} \Delta \sim_{\mathbb{Q}} 0$. Since (X_T, X_k) is log canonical, $\operatorname{Diff}_{\bar{E}}\Delta = \bar{D}$ by [Kol13, 4.9]. Thus $K_{\bar{E}} + \bar{D} \sim_{\mathbb{Q}} 0$, hence A_k is semiabelian by (44.1). \square

46 (Proof of (42.1) \Rightarrow (42.2)). As in (41), let $Y_T \rightarrow T$ be a Nér^o(A_K)-equivariant resolution of X_T that contains Nér (A_K) as an open subset. Running a minimal model program over X_T gives $Y_T = Y_T^1 \dashrightarrow \cdots \dashrightarrow Y_T^m =: Z_T$.

Let $E \subset Y_k$ be any divisor corresponding to an irreducible component of $Nér(A_K)_k$. We claim that E is not contracted by $Y_T \dashrightarrow Y_T^m$.

14

Assume by induction that E appears on Y_T^i . If A_k is semiabelian then $K_{\bar{E}} + \bar{D} \sim 0$ by (44.2), hence $\tau^* K_{Y_T^i} \sim_{\mathbb{Q}} K_{\bar{E}} + \bar{D} + D^* \sim_{\mathbb{Q}} D^*$ is effective. Thus $Y_T^i \to Y_T^{i+1}$ does not contract E.

Therefore Z_k contains at least 1 divisor E with multiplicity 1. Here $K_{Z_T} \sim_{\mathbb{Q}} 0$, hence $K_{\bar{E}} + \text{Diff}_{\bar{E}} \Delta \sim_{\mathbb{Q}} 0$, and so $\text{Diff}_{\bar{E}} \Delta = \bar{D}$. Inversion of adjunction [Kol13, 4.9] implies that (Z_T, Z_k) is log canonical along E. In particular, every irreducible component of Z_k meeting E also has multiplicity 1. Repeating the argument for the other irreducible components we get that Z_k is reduced and (Z_T, Z_k) is log canonical. Then the same holds for (X_T, X_k) .

7. Log canonical models

In this section S denotes a normal, integral base scheme with function field K. The proof uses several results of the minimal model program, so we need to assume that S is excellent, has a dualizing complex, and is over a field of characteristic 0. With these restrictions, regularity of actions also holds for almost all pairs over higher dimensional bases.

Theorem 47. Let $p: (X_S, \Delta_S) \to S$ be a dominant, projective morphism, and $G \to S$ a smooth group scheme with irreducible fibers. Let $\mu_K: G_K \times X_K \to X_K$ be a regular action such that Δ_K is G_K -invariant. Assume that

(47.1) (X_S, Δ_S) is log canonical and (X_K, Δ_K) is terminal,

(47.2) $K_{X_S} + \Delta_S$ is p-semiample, and

(47.3) $p^{-1}(p(Z)) \subset \operatorname{Supp} \Delta_S$ for every log canonical center Z of (X_S, Δ_S) .

Then the induced rational action $\mu_S : G \times_S X \dashrightarrow X$ is regular.

Remarks 47.4. See [Kol13, 4.28] for log canonical centers.

Assumptions (47.1–2) generalize the notion of a minimal model to pairs (allowing worse singularities than usual).

The abundance conjecture says that if $K_{X_S} + \Delta_S$ is *p*-nef, then it is also *p*-semiample, This is known if (X_S, Δ_S) is klt and $K_{X_K} + \Delta_K \sim_{\mathbb{Q}} 0$ [Fuj11, Lai11].

Let E be a smooth, elliptic curve and X the blow-up of a point $(e, 0) \in E \times \mathbb{A}^1$. Let $\Delta := E'$ be the birational transform of E_0 with coefficient 1. Then $K_X + \Delta$ is numerically trivial but the E-action does not lift to X. Note that Δ has a component with coefficient 1 (namely E') and also a component with coefficient 0 (the exceptional \mathbb{P}^1). Assumption (47.3) is there to exclude such extreme cases.

Proof. Let $g: X'_S \to X_S$ be a Q-factorial, minimal modification. That is, X'_S has terminal singularities, and $K_{X'_S}$ is g-nef. Thus there is a unique effective Q-divisor Δ'_S such that $K_{X'_S} + \Delta'_S$ is numerically g-trivial and $g_* \Delta'_S = \Delta_S$. These imply that (X'_S, Δ'_S) is log canonical and $\operatorname{Supp} \Delta'_S \supset g^{-1}(\operatorname{Supp} \Delta_S)$ by [KM98, 3.39].

 (X'_S, Δ'_S) is log canonical and $\operatorname{Supp} \Delta'_S \supset g^{-1}(\operatorname{Supp} \Delta_S)$ by [KM98, 3.39]. By (47.2) there is a factorization $p: X \to S' \to S$ such that $S' \to S$ is birational and $K_{X_S} + \Delta_S$ is numerically trivial on $X_S \to S'$. MMP for $X'_S \to S'$ gives a minimal model $X^m \to S'$. Let $\tau: X'_S \dashrightarrow X^m$ be the corresponding birational map.

At each step of the MMP the image of $K_{X'_S} + \Delta'_S$ is numerically trivial, and we end with a log canonical pair (X^m, Δ^m) such that $K_{X^m} + \Delta^m$ is also numerically trivial over S'. For any irreducible divisor $E \subset X'_S$,

$$a(E, X^{\mathrm{m}}, \Delta^{\mathrm{m}}) = a(E, X'_{S}, \Delta'_{S}) = a(E, X_{S}, \Delta_{S})$$

is the negative of its coefficient in Δ'_S . Thus, using the notation in (49),

 $a(E, X^{\mathrm{m}}, \Delta^{\mathrm{m}}) \leq 0 \leq 1 + \text{totaldiscrep}(\tau(E), X^{\mathrm{m}}, \Delta^{\mathrm{m}}),$

and equality holds iff

$$a(E, X^{\mathrm{m}}, \Delta^{\mathrm{m}}) = 0$$
 and totaldiscrep $(\tau(E), X^{\mathrm{m}}, \Delta^{\mathrm{m}}) = -1$.

The first says that E is not contained in the support of Δ'_S , the second that g(E) is contained in a log canonical center of (X_S, Δ_S) . This is excluded by (47.3).

Note that X^{m} is a minimal model over S, so the induced G-action on X^{m} is regular by Theorem 7. It then lifts to a regular G-action on X'_S by (50), which pushes forward to a regular G-action on X_S by (48).

Lemma 48. Let S be a normal scheme, X, Y normal, projective S-schemes, and $G \to S$ a smooth group scheme with irreducible fibers. Let $\mu_X : G \times_S X \to X$ be a regular action and $g : X \dashrightarrow Y$ a birarational map. Assume that

(48.1) g^{-1} has no exceptional divisors, and

(48.2) every g-exceptional divisor is μ_X -invariant.

Then μ_X pushes forward to a regular action $\mu_Y : G \times_S Y \to Y$.

Proof. We have a rational action $\mu_Y : G \times_S Y \dashrightarrow Y$. For every étale section $\tau : S' \to G$ the induced map $\tau_Y := g \circ \tau_X \circ g^{-1}$ is a small modification by the assumptions. Thus μ_Y is a regular action by (37).

Definition 49. Let (X, Δ) be a pair and $W \subset X$ an irreducible subset. Let totaldiscrep (W, X, Δ) denote the infimum of the discrepancies $a(E, X, \Delta)$, where E runs through all divisors for which $W \subset \text{center}_X E$.

Lemma 50. Let S be a normal scheme, $p: (X, \Delta) \to S$ a projective morphism, and $G \to S$ a smooth group scheme with connected fibers acting on X.

Let $g: Y \to X$ be a projective, birational morphism such that Y is normal, and $a(E, X, \Delta) < 1 + \text{totaldiscrep}(g(E), X, \Delta)$ for every g-exceptional divisor E.

Then the G-action on X lifts to a G-action on Y.

Proof. Let $h: X' \to X$ be a *G*-equivariant log resolution of (X, Δ) . There is a unique snc dvisor Δ' such that $K_{X'} + \Delta' \sim_{\mathbb{Q}} h^*(K_X + \Delta)$ and $h_*\Delta' = \Delta$. Then $a(E, X', \Delta') = a(E, X, \Delta)$ by [KM98, 2.30]. Since $a(E, X, \Delta) < 1 +$ totaldiscrep $(g(E), X, \Delta)$, the center of *E* on *X'* is a stratum of (X', Δ') by [KM98, 2.31]. All strata are *G*-equivariant. Thus, after further *G*-equivariant blow-ups $X'' \to X'$, we may assume that every *g* exceptional divisor *E* is a divisor on *X''* by [KM98, 2.45]. Then the *G*-action on *X''* pushes forward to a *G*-action on *Y* by (48).

Specializing (47) to the case when S = T is the spectrum of a DVR and X_K is an Abelian torsor gives the following.

Corollary 51. Let X_K be a torsor under an Abelian variety A_K , and $\pi : (X_T, \Delta) \to T$ a projective morphism. Assume that

- (51.1) either (X_T, Δ) is log canonical and $\operatorname{Supp} \Delta = X_k$,
- (51.2) or (X_T, Δ) is klt and Supp $\Delta \subset X_k$.

Then the A_K -action on X_K extends to a regular Nér^o (A_K) -action on X_T .

Corollary 52. Continuing with the notation of (51), assume in addition that $\Delta = \operatorname{red}(X_k)$, (X_T, Δ) is divisorial log terminal, and X_k has an irreducible component of multiplicity 1. Then

(52.1) Nér (X_K) is isomorphic to the π -smooth locus of X_T , and (52.2) Nér $(A_K)_k$ is semiabelian.

Proof. We follow the proof of (47). For $g: X'_T \to X_T$ we choose a *thrifty* modification as in [Kol13, 2.79]. That is, all g-exceptional divisors have coefficient < 1 in Δ' . Note that

$$K_{X'_{\mathcal{T}}} \sim_{\mathbb{Q}} -\Delta' \sim_{\mathbb{Q}} X'_k - \Delta'.$$

If $E \subset X'_k$ is an irreducible component that dominates a reduced component of X_k , then E appears in $X'_k - \Delta'$ with coefficient 0, all other irreducible components appear with positve coefficient. Therefore the $K_{X'_T}$ -MMP contracts precisely the support of $X'_k - \Delta'$.

By (19.5) the smooth locus of $X_T^m \to T$ is isomorphic to Nér (X_K) , hence the same holds for $X_T \to T$ by (18). The second claim follow from (42).

Examples 53. These examples show that there are many log minimal models, and the Supp $\Delta = Y_k$ assumption in (51) is needed.

(53.1) Let E be an elliptic curve. Blowing up a point in the central fiber of $E \times \mathbb{A}^1 \to \mathbb{A}^1$, and then contracting the birational transform of E gives a log minimal model $\pi: S \to \mathbb{A}^1$. A pair (S, Δ) is log canonical only for $\Delta = 0$. There is no regular Nér° (A_K) -action on S.

(53.2) Start with $(xyz = t(x^3 + y^3 + z^3)) \subset \mathbb{P}^2_{xyz} \times \mathbb{A}^1_t$. All log minimal models with $\Delta = X_k$ are constructed as follows.

Repeatedly blow up singular points of the central fiber, then contract any subset of the central fiber (except we can not contract the whole central fiber).

(53.3) If in (53.2) we contract all but 1 or 2 of the irreducible components of the central fiber, we get a log minimal model that does not have a Nér(A_K)-action.

References

- [Ale02] Valery Alexeev, Complete moduli in the presence of semiabelian group action, Ann. of Math. (2) 155 (2002), no. 3, 611–708.
- [AN99] Valery Alexeev and Iku Nakamura, On Mumford's construction of degenerating abelian varieties, Tohoku Math. J. (2) 51 (1999), no. 3, 399–420. MR MR1707764 (2001g:14013)
- [Art86] Michael Artin, Néron models, Arithmetic geometry (Storrs, Conn., 1984), Springer, New York, 1986, pp. 213–230. MR MR861977
- [BCHM10] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James M^cKernan, Existence of minimal models for varieties of log general type, J. Amer. Math. Soc. 23 (2010), no. 2, 405–468.
- [BHPT18] Fedor Bogomolov, Lars Halvard Halle, Fabien Pazuki, and Sho Tanimoto, Abelian Calabi-Yau threefolds: Néron models and rational points, Math. Res. Lett. 25 (2018), no. 2, 367–392. MR 3826826
- [BKV24] Fedor Bogomolov, Ljudmila Kamenova, and Misha Verbitsky, Sections of Lagrangian fibrations on holomorphic symplectic manifolds, https://arxiv.org/abs/2407.07877, 2024.
- [BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud, Néron models, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), vol. 21, Springer-Verlag, Berlin, 1990. MR 1045822 (91i:14034)

- [BMP+23] Bhargav Bhatt, Linquan Ma, Zsolt Patakfalvi, Karl Schwede, Kevin Tucker, Joe Waldron, and Jakub Witaszek, Globally +-regular varieties and the minimal model program for threefolds in mixed characteristic, Publ. Math. Inst. Hautes Études Sci. 138 (2023), 69–227. MR 4666931
- [Bri22] Michel Brion, On models of algebraic group actions, Proc. Indian Acad. Sci. Math. Sci. 132 (2022), no. 2, Paper No. 61, 17. MR 4504264
- [BZ17] Michel Brion and De-Qi Zhang, Log Kodaira dimension of homogeneous varieties, Algebraic varieties and automorphism groups, Adv. Stud. Pure Math., vol. 75, Math. Soc. Japan, Tokyo, 2017, pp. 1–6. MR 3793359
- [FC90] Gerd Faltings and Ching-Li Chai, Degeneration of abelian varieties, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), vol. 22, Springer-Verlag, Berlin, 1990, With an appendix by David Mumford. MR MR1083353 (92d:14036)
- [Fuj11] Osamu Fujino, Semi-stable minimal model program for varieties with trivial canonical divisor, Proc. Japan Acad. Ser. A Math. Sci. 87 (2011), no. 3, 25–30. MR 2802603 (2012j:14023)
- [Gro60] Alexander Grothendieck, Éléments de géométrie algébrique. I–IV., Inst. Hautes Études Sci. Publ. Math. (1960), no. 4,8,11,17,20,24,28,32.
- [Gro68] _____, Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2), North-Holland Publishing Co., Amsterdam, 1968, Augmenté d'un exposé par Michèle Raynaud, Séminaire de Géométrie Algébrique du Bois-Marie, 1962, Advanced Studies in Pure Mathematics, Vol. 2. MR 0476737
- [Han87] Masaki Hanamura, On the birational automorphism groups of algebraic varieties, Compositio Math. 63 (1987), no. 1, 123–142. MR MR906382 (88h:14018)
- [HM07] Christopher D. Hacon and James McKernan, Extension theorems and the existence of flips, Flips for 3-folds and 4-folds, Oxford Lecture Ser. Math. Appl., vol. 35, Oxford Univ. Press, Oxford, 2007, pp. 76–110.
- [HN18] Lars Halvard Halle and Johannes Nicaise, Motivic zeta functions of degenerating Calabi-Yau varieties, Math. Ann. 370 (2018), no. 3-4, 1277–1320. MR 3770167
- [Kim24] Yoon-Joo Kim, The Néron model of a higher-dimensional Lagrangian fibration, =https://arxiv.org/abs/2410.21193, 2024.
- [KL09] János Kollár and Michael Larsen, Quotients of Calabi-Yau varieties, Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, Progr. Math., vol. 270, Birkhäuser Boston Inc., Boston, MA, 2009, pp. 179–211. MR 2641190 (2011c:14117)
- [KLOS23] János Kollár, Max Lieblich, Martin Olsson, and Will Sawin, What determines an algebraic variety?, Princeton University Press, Princeton, NJ, 2023.
- [KM98] János Kollár and Shigefumi Mori, Birational geometry of algebraic varieties, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [Kod60] Kunihiko Kodaira, On compact complex analytic surfaces. I, Ann. of Math. (2) 71 (1960), 111–152. MR 132556
- [Kod63] _____, On compact analytic surfaces. II, III, Ann. of Math. (2) **77** (1963), 563–626; 78 (1963), 1–40. MR 184257
- [Kol91] János Kollár, Flips, flops, minimal models, etc, Surveys in differential geometry (Cambridge, MA, 1990), Lehigh Univ., Bethlehem, PA, 1991, pp. 113–199. MR 1144527
- [Kol13] _____, Singularities of the minimal model program, Cambridge Tracts in Mathematics, vol. 200, Cambridge University Press, Cambridge, 2013, With the collaboration of Sándor Kovács.
- [Kol21] _____, *Relative MMP without* Q-factoriality, Electron. Res. Arch. **29** (2021), no. 5, 3193–3203. MR 4342251
- [Kol23] _____, Families of varieties of general type, Cambridge Tracts in Mathematics, vol. 231, Cambridge University Press, Cambridge, 2023, With the collaboration of Klaus Altmann and Sándor Kovács. MR 4566297
- [Kün98] Klaus Künnemann, Projective regular models for abelian varieties, semistable reduction, and the height pairing, Duke Math. J. 95 (1998), no. 1, 161–212. MR 1646554
- [Lai11] Ching-Jui Lai, Varieties fibered by good minimal models, Math. Ann. 350 (2011), no. 3, 533–547. MR 2805635 (2012e:14032)

- [Lan83] Serge Lang, Fundamentals of Diophantine geometry, Springer-Verlag, New York, 1983. MR 715605
- [LM22] Shiji Lyu and Takumi Murayama, The relative minimal model program for excellent algebraic spaces and analytic spaces in equal characteristic zero, =https://arxiv.org/abs/2209.08732, 2022.
- [Lor17] Dino Lorenzini, Néron models, Eur. J. Math. 3 (2017), no. 2, 171–198. MR 3695053
- [Mat52] Teruhisa Matsusaka, On the algebraic construction of the Picard variety. II, Jap. J. Math. 22 (1952), 51–62. MR 0062471
- [MM64] Teruhisa Matsusaka and David Mumford, Two fundamental theorems on deformations of polarized varieties, Amer. J. Math. 86 (1964), 668–684. MR MR0171778 (30 #2005)
- [MN24] Kentaro Mitsui and Iku Nakamura, Relative compactifications of semiabelian Néron models, I, =https://arxiv.org/abs/2201.08113, 2024.
- [Mum72] David Mumford, An analytic construction of degenerating abelian varieties over complete rings, Compositio Math. 24 (1972), 239–272. MR MR0352106 (50 #4593)
- [Nag63] Masayoshi Nagata, A generalization of the imbedding problem of an abstract variety in a complete variety, J. Math. Kyoto Univ. 3 (1963), 89–102. MR MR0158892 (28 #2114)
- [Nak24] Iku Nakamura, Relative compactification of semiabelian Néron models, II, =https://arxiv.org/abs/2405.09172, 2024.
- [Nér62] André Néron, Modèles minimaux des variétés abéliennes, Séminaire Bourbaki, 14ième année: 1961/62. Fasc. 1, 2 et 3: Textes des Conférences, Exp. 223 à 240, Secrétariat mathématique, Paris, 1962, 2ième édition, corrigée. MR 146035
- [Nér64] _____, Modèles minimaux des variétés abéliennes sur les corps locaux et globaux, Inst. Hautes Études Sci. Publ. Math. (1964), no. 21, 1–128. MR 179172
- [Ove24] Otto Overkamp, Néron models of pseudo-Abelian varieties, Rend. Semin. Mat. Univ. Padova 152 (2024), 83–116. MR 4784847
- [Rei80] Miles Reid, Canonical 3-folds, Journées de Géometrie Algébrique d'Angers, Juillet 1979/Algebraic Geometry, Angers, 1979, Sijthoff & Noordhoff, Alphen aan den Rijn, 1980, pp. 273–310. MR 605348 (82i:14025)
- [Sum74] Hideyasu Sumihiro, Equivariant completion, J. Math. Kyoto Univ. 14 (1974), 1–28. MR 337963
- [Sum75] _____, Equivariant completion. II, J. Math. Kyoto Univ. 15 (1975), no. 3, 573–605. MR 387294
- [Tot13] Burt Totaro, Pseudo-abelian varieties, Ann. Sci. Éc. Norm. Supér. (4) 46 (2013), no. 5, 693–721. MR 3185350
- [Wei48] André Weil, Variétés abéliennes et courbes algébriques, Publications de l'Institut de Mathématiques de l'Université de Strasbourg, vol. 8, Hermann & Cie, Paris, 1948, Actualités Scientifiques et Industrielles, No. 1064. MR 29522

Princeton University, Princeton NJ 08544-1000, kollar@math.princeton.edu