Optimal Overlap Detection of Shotgun Reads

Nir Luria and Nir Weinberger

The Viterbi Faculty of Electrical and Computer Engineering Technion - Israel Institute of Technology Technion City, Haifa 3200004, Israel nir.luria@campus.technion.ac.il, nirwein@technion.ac.il.

Abstract

We consider the problem of detecting the overlap between a pair of short fragments sampled in random locations from an exponentially longer sequence, via their possibly noisy reads. We consider a noiseless setting, in which the reads are noiseless, and the sequence is only assumed to be stationary and ergodic. Under mild conditions on the mixing property of the process generating the sequence, we characterize exactly the asymptotic error probability of the optimal Bayesian detector. Similarly, we consider a noisy setting, in which the reads are noisy versions of the sampled fragments obtained via a memoryless channel. We further assume that the sequence is stationary and memoryless, and similarly characterize exactly the asymptotic error probability of the optimal Bayesian detector for this case.

Index Terms

Bayesian error probability, DNA sequencing, hypothesis testing, overlap detection, shotgun sequencing.

I. INTRODUCTION

Consider an archaeologist who discovers two short torn fragments from an ancient long text corpus. When it can be reliably determined that these two fragments come from the same location in the text, and thus should be stitched together to form a longer contiguous fragment? Alternatively, when it can be determined that these two fragments are parts of different locations in the text? A correct answer to this question would affect the conclusions that can be made from this discovery. Notwithstanding the above example, the most well-known instance of this problem arises in *DNA shotgun sequencing* [27], in which the goal is to assembly a long genomic sequence based on *shotgun reads*. In this setup, short fragments are sampled from random starting locations on

Nir Luria and Nir Weinberger are with the Viterbi Faculty of Electrical and Computer Engineering, Technion-Israel Institute of Technology, Haifa 3200004, Israel (Emails: nir.luria@campus.technion.ac.il, nirwein@technion.ac.il.). Their work is supported by the Israel Science Foundation (ISF), grant no. 1782/22. A short version of this paper was submitted to the 2025 IEEE International Symposium on Information Theory. This work has been submitted to the IEEE for possible publication. Copyright may be transferred without notice, after which this version may no longer be accessible.

the sequence, and then each of them is independently sequenced to obtain a *read* – a possibly corrupted version of this fragment. The set of noisy reads is then input to an assembly algorithm, which outputs a reconstruction of the sequence. From a more general perspective, alignment of short signals is a widely used pre-processing technique. In image processing, one can crate a panoramic view from smaller images, or connect localized medical scans to form one whole medical image. In hardware security, many power traces are recorded and should be aligned in order to understand the power behavior of the design. In this paper, we address a distilled version of this setting: Given a *single pair* of reads from a long sequence, what are the fundamental limits of the detection of the overlap length, or the relative alignment, between these reads? To address this question, we develop the optimal Bayesian detector for this problem, and derive exact asymptotic expressions for its optimal Bayesian error probability.

A. Motivation and Background: Fundamental Limits of DNA Shotgun Sequencing

As said, a prominent example of the alignment problem is DNA sequencing, and so we next focus on this setting. Our motivation for addressing the alignment problem in isolation arises, because, as we shall next justify, an analysis of the fundamental limits of the reconstruction of the full sequence from multiple reads masks this basic question. To this end, we briefly review the information-theoretic study of fundamental limits of DNA shotgun sequencing, pioneered in [22]. The genome \mathbf{X} is modeled as a length n sequence over an alphabet $\mathcal{X} = \{A, C, G, T\}$. Each of the *m* reads has length ℓ , and a starting point that is chosen uniformly over the sequence,¹ independently of the starting points of all other reads. The interesting analytical regime is when the reads are exponentially shorter than the sequence, concretely $\ell = \beta \log n$, for some fixed scaling constant $\beta > 0$. Trivially, if $m \leq \frac{n}{\ell}$ then the sequence is not even fully covered by the reads, and thus cannot be fully reconstructed from the m reads. In their classic paper [16], Lander and Waterman used a coupon-collection argument to show that if the coverage condition $m \gtrsim \frac{n}{\ell} \log(\frac{n}{\ell\epsilon}) \sim \frac{n}{\beta}$ holds, then m reads will fully cover the entire sequence with probability $1 - \epsilon$. As argued later on in [1], this does not suffice to assure a reliable reconstruction of the sequence, due possible repetitions in the sequence. Indeed, it was shown in [33] that if there is either a substring of length $\ell - 1$ that repeats itself *three* times, or there are two interleaved pairs of repeats of length $\ell - 1$, then perfect reconstruction is impossible, even given the ℓ -spectrum – the set of all reads of length ℓ in the sequence (see [4] for refined conditions on the required coverage depth). It was proved in [1] that for a sequence generated from a memoryless process with a symbol probability mass function (p.m.f.) P_X , such interleaved repeats will occur with high probability whenever $\beta < \frac{2}{H_2(P_X)}$, where $H_2(P_X)$ is the second-order Rényi entropy $H_2(P_X) := -\log \sum_{x \in \mathcal{X}} P_X^2(x)$. It was then proved in [22] that if the sequence is

¹That is, uniformly over $[n] := \{1, 2, ..., n\}$, with some adjustment at the boundaries; [22] analyzed a cyclic sequence model, and we will consider here a slightly different model at the boundaries. Both choice are made mainly for simplicity, and do not have any impact on the asymptotic analytical results.

a memoryless random process, and the *repeat condition* $\beta \ge \frac{2}{H_2(P_X)}$ holds on top of the coverage condition, it is possible to perfectly reconstruct the sequence from *m* reads.

Importantly, the requirement of the covering condition for reliable reconstruction greatly influenced the reconstruction algorithm proposed and analyzed in [22]. This algorithm is greedy in its nature, and keeps track of a set of *contigs* – a contiguous fragment of overlapping reads. At first, the set of contigs is initialized to the set of sequenced reads. Then, at each of m-1 steps, the algorithm finds a pair of contigs with maximal overlap and merges them into a single contig. The analysis of the probability that the algorithm fails to perfectly reconstruct the sequence leads to the necessary repeat-condition $\beta \geq \frac{2}{H_2(P_X)}$. Intuitively speaking, it is based on the following arguments: (i) Long overlaps between reads from distant locations are unlikely. Indeed, consider overlaps of length $\alpha \ell$ for some $\alpha \in (0, 1)$. For a memoryless process over $|\mathcal{X}|$, the probability that two reads from distant locations in the sequence will share the exact same symbols is $|\mathcal{X}|^{-H_2(P_X)\alpha\ell}$, and since $\ell = \beta \log n$, this probability polynomially decays with n (with degree $\alpha\beta H_2(P_X)$). This probability decays fast enough if α is large enough. (ii) It is highly unlikely for a specific symbol in the sequence *not* to be covered by at least one pair of reads that has a *long* overlap. Indeed, given the minimum number of reads to satisfy the coverage condition, on the average, each symbol is covered by $\frac{m\ell}{n} \sim \log n$ reads. Since the relative alignment between the reads is uniformly, it is highly unlikely that neither pair from these reads does not have a long overlap. Therefore, reliable reconstruction is essentially obtained by almost exclusively merging reads with *long* overlaps.

So, since the algorithm is based on merging reads with sufficiently long overlaps, the analysis of this algorithm masks the basic question regarding the ability to detect an overlap, without any prior guarantee that this overlap is long enough. In fact, detection of short overlaps is more efficient, because they result longer contigs. Interestingly, [16], dated to the early days of the genome sequencing projects, considered the case in which the genome is not fully covered (without a rigorous analysis). In the parlance of [16], an ideal alignment of the available reads in such cases will result in a set of contigs, possibly separated by uncovered symbols. The reconstruction algorithm results *islands*, whose correct order cannot be determined from the available reads. In this under-coverage regime, $m \leq \frac{n}{\ell}$, it is possible for a read to have a relatively short maximal overlap with all other reads. Such shorts overlaps are not used in the reconstruction algorithm of [22] (with high probability), as it is unclear when a detection of such an overlap is actually reliable, and, as said, it is also necessary to use such overlaps whenever the coverage condition holds. This question remains a timely challenge, even with contemporary high-throughput DNA sequencing technologies: In *metagenomics analysis* such as that of the Human Microbiome Project [17], multiple sequences from different genomes are analyzed at once, and typically the total number of reads does not suffice for a full coverage of all the genomes [10], [14].²

The reconstruction algorithm and analysis were extended in [22] in two directions. First, the assumption on the input process generating the sequence was generalized from a memoryless process to a Markov process

²In metagenomics, even the identifiability of the sequences from the reads is a challenging question [11].

(assumed for simplicity, to be a first-order Markov process), in which the second-order Rényi entropy rate³ $\mathcal{H}_2(\mathbf{X})$ replaces $H_2(P_X)$ used in the memoryless case, and was shown to be optimal [22, Th. 6] (ii) Noisy sequencing case of a memoryless sequence, in which the clean fragments are corrupted by a memoryless channel. In this case, $H_2(P_X)$ is replaced by a information measure denoted I^* [22, Th. 10], without a claim for optimality. Evidently, while the *scaling* in all cases is the same, it is the exact minimal value of β required for perfect reconstruction that is affected by the statistics of the process generating the sequence or the reads $(H_2(P_X), \mathcal{H}_2(\mathbf{X}) \text{ or } I^*)$. As we show, a similar property holds in our setting as well.

In general, a central part of the operation of an assembly algorithm of a sequence form reads is merging of pairs of fragments based on their overlap (either reads, or shorter fragments called *k*-mers). Various papers have shown that the ability to do so successfully fundamentally depends on the statistics of repeats in the long sequence. For example, in [4], a multi-bridging algorithm based on *k*-mers was proposed, and conditions on β were derived based on the appearance of repeats in the sequence. This idea was extended to noisy reads in [15], and to an assembly algorithm based on the reads themselves in [31]. Information-theoretic limits on assembly for the noisy case were developed in [23] for independent and identically distributed (i.i.d.) sequences and in [29] for arbitrary sequences under an adversarial noise model, showing the effect of noise on the assembly algorithm performance. In [30], the computational problem of overlap detection was addressed based on sketches, which are compressed versions of the pair of reads. The optimal number of bits required for reliable approximate overlap was tightly bounded, where the approximation is under the mean squared error loss function. In [30], it was *assumed* that the alignment parameter is larger than some $\theta_0 \ell$ for some constant $\theta_0 \in (0, 1)$. Here, we show that sufficiently short overlaps cannot be reliably detected. See also [3], [12], [28], [30] and references therein for practical overlap detection algorithms.

B. Our Results

We focus on the distilled problem of overlap detection between a *single* pair of reads, and analyze a detector that is required to decide on the length of the overlap between them. We emphasize here the Bayesian setting, because each of the possibilities have a substantially different prior. Given a length n sequence X_1^n and two reads of length $\ell = \beta \log n$ starting at random locations, the probability that they overlap at a given length $T \in \{-[\ell - 1]\} \cup [\ell] \text{ is } \frac{1}{n},^4$ whereas the probability that they do not overlap at all is $\frac{2\ell - 1}{n} \approx \frac{2\beta \log n}{n}$, an event which we denote by T = 0. Therefore, for a detector \hat{T} , we let the type-I error be the *false-overlap-detection* error event, whose probability is $p_{type-I}(\hat{T}) := \mathbb{P}[\hat{T} \neq 0 \mid T = 0]$, and the type-II error be the *misdetection or*

³To be formally defined later on.

⁴We distinguish between a positive and negative overlaps, to wit, between the case that the first read appears before the second one and vice-versa.

erroneous-overlap error event, whose probability is $p_{\text{type-II}}(\hat{T}) := \mathbb{P}[\hat{T} \neq T \mid T \neq 0]$. The total error probability is thus tightly upper bounded as

$$P_{\text{error}}(\hat{T}) \lessapprox p_{\text{type-I}}(\hat{T}) + \frac{2\beta \log n}{n} p_{\text{type-II}}(\hat{T}), \tag{1}$$

which highlights the non-equal contribution of the two types of error to the total error probability.

Similarly to the reconstruction problem, we characterize the scaling as well as the exact *constant* of the error probability (see [22, Section I.C]). It is clear here that the scaling of the optimal error probability is $P_{\text{error}}(\hat{T}) = \Theta(\frac{\log n}{n})$, because a trivial detector $\hat{T}_0 \equiv 0$, which always decides "no-overlap", has an error probability of $P_{\text{error}}(\hat{T}_0) = \mathbb{P}[T \neq 0] \sim 2\beta \cdot \frac{\log n}{n}$. Therefore, the main problem is to characterize the optimal constant

$$\phi := \min_{\hat{T}} \lim_{n \to \infty} \frac{P_{\text{error}}(\hat{T})}{(\log n)/n},\tag{2}$$

for different probabilistic models of the process generating the sequence. That is, taking into account memory in the process generating the sequence, which might lead to various fragment repeat properties, as well as the noise in the reading mechanism of the fragments. For tractability, we focus on two settings – the *noiseless setting* and the *noisy setting* – for which the optimal Bayesian detector can be easily derived and the error probability analysis is tractable.

In the first setting, referred to for brevity as the *noiseless setting*, we assume that the reading process is noiseless, but the sequence is generated by a stationary and ergodic process. We do not assume that the process is memoryless or a Markov process of a finite order. We show that under a mild condition on the strong mixing coefficient of the process, as well as a mild condition on the recurrence probability of the process (to be formally defined later on), it holds that

$$\phi = 2 \cdot \left[\beta \wedge \frac{1}{\mathcal{H}_1(\mathbf{X})} \right],\tag{3}$$

where $\mathcal{H}_1(\mathbf{X})$ is the Shannon entropy rate of the process.

Interestingly, the constant is *not* determined by the second-order Rényi entropy rate $\mathcal{H}_2(\mathbf{X})$, as might be hypothesized from the analysis of sequence reconstruction discussed above. As we show and discuss in length, the root of this is directly related to the imbalance in the prior probability. Indeed, a simplified, yet very useful, way of thinking on the alignment detection problem is as a collection of $2\ell - 1$ binary detection problems between T = 0 and T = t, for some $t \in \{-[\ell - 1]\} \cup [\ell]$, with a prior probability on which detection problem is active. The testing problem is more difficult for short alignments, and so we upper bound the error probability by 1 if the active detection problem is in one of the most $2t^*$ difficult problems (those of shortest overlaps). The error probability upper bound is thus at least the probability that one of these difficult detection problems is active, which is $\sim 2t^* \cdot \frac{\log n}{n}$. The challenge in the proof is to accurately identify that t^* for which both the probability of both false-overlap-detection, as well as the probability to err from one detection problem to another (that is, a wrong alignment value), is negligible compared to this probability. While this intuition may appear to be just a bounding technique, it is actually also used to obtain a matching lower bound.

In a nutshell, the optimal Bayesian detector will detect an overlap of some specific $T = t \neq 0$ only if the likelihood for T = t compared to the likelihood for T = 0 (no overlap) is larger than the ratio between their inverse prior probabilities $\frac{\mathbb{P}[T=0]}{\mathbb{P}[T=t]} \sim \frac{1}{1/n} = n$. However, for overlaps shorter than $t \sim \frac{\log n}{\mathcal{H}_1(\mathbf{X})}$, it holds that the expectation of this log-likelihood-ratio is less than n, and so, such overlaps are not likely to be detected by the optimal detector. Consequently, whenever $1 \leq |T| \leq \frac{\log n}{\mathcal{H}_1(\mathbf{X})}$ the conditional error probability is 1. Evidently, when $\beta > \frac{1}{\mathcal{H}_1(\mathbf{X})}$ this is the dominating error event. The technical challenge in upper bounding the error probability is to show that all other error events, either type-I error event or other sub-events of the type-II error event, have negligible $o(\frac{\log n}{n})$ probability compared to the probability of this event, without making strong assumptions on the process generating the sequence.

An easy generalization of our upper bounds leads to a different scaling of the contributions of the two type of errors. If we replace the prior probability ratio $\frac{\mathbb{P}[T=0]}{\mathbb{P}[T=t]} \sim n$ by n^{μ} for $\mu > 1$ then we decrease the type-I error probability to

$$p_{\text{type-I}}(\hat{T}) = o\left(\frac{\log n}{n^{\mu}}\right),\tag{4}$$

a faster decay rate, with power almost μ , whereas overlaps $|T| \leq \frac{\mu \log n}{\mathcal{H}_1(\mathbf{X})}$ are not detected, and in turn, the contribution of the type-II error, and so also the total error probability is increased by a factor of μ , it is

$$P_{\text{error}}(\hat{T}) \sim 2\mu \cdot \left[\beta \wedge \frac{1}{\mathcal{H}_1(\mathbf{X})}\right] \cdot \frac{\log n}{n}.$$
(5)

In the second setting, referred to for brevity as the *noisy setting*, we assume that the process is generated by a stationary memoryless process with p.m.f. P_X , and that the reads are obtained by observing the fragments through a discrete memoryless channel $P_{Y|X}$ (independently for each read). Let Y, \tilde{Y} be two independent observations of the same $X \sim P_X$ through $P_{Y|X}$ so that the Markov relation $Y - X - \tilde{Y}$ holds. Here we show that

$$\phi = 2 \cdot \left[\beta \wedge \frac{1}{I(Y; \tilde{Y})} \right],\tag{6}$$

where $I(Y; \tilde{Y})$ is the mutual information. Similarly to the noiseless setting, this is related to the log-likelihoodratio between an overlap, characterized by the joint distribution $P_{Y\tilde{Y}}$ and no-overlap, characterized by the product distribution $P_Y \otimes P_{\tilde{Y}}$. The expectation of this log-likelihood ratio under $P_{Y\tilde{Y}}$ is exactly $I(Y; \tilde{Y})$, which explains why it appears in the bound.

There are various technical challenges associated with the error probability analysis. First, it is challenging to show that the type-I error probability is upper bounded as $o(\frac{\log n}{n})$ (thus negligible). In the noiseless setting,

this requires a tight analysis of a term of the form

$$\sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}^2 \left(x_1^t \right) \mathbb{1} \left[\frac{1}{P_{X_1^t}(x_1^t)} \ge n \right]$$

$$\tag{7}$$

which is a partial sum of the sum involved in the second-order Rényi entropy of $P_{X_1^t}$, the p.m.f. of $X_1^t := (X_1, \ldots, X_t)$. Interestingly, an analysis that is based on trivially upper bounding the indicator function by 1, results an sub-optimal upper bound on the constant of total error probability of $\phi \leq \beta \wedge \frac{1}{\mathcal{H}_2(\mathbf{X})}$. In the noisy setting, the type-I error probability pertains to the probability that a sum of independent random variables (r.v.'s) crosses a threshold. Here, an application of the Chernoff bound does not suffice, despite the fact that it is exponentially tight. We thus turn to stronger versions, similar to that of Bahadur–Rao [2] [7, Th. 3.7.4.] (see also [20], [32]), utilizing their accurate pre-exponent. Second, there are challenge in establishing that the probability of an erroneous overlap is negligible. This is because conditioned on the event that T = t > 0, the analysis of the probability of the event that the detector decides $\hat{T} \neq t$ involves analyzing sum of possibly *dependent* r.v.'s. The proofs propose methods to circumvent the need to analyze these sums of dependent r.v.'s.

C. Paper Outline

The outline of the rest of the paper is as follows. In Sec. II we state notation conventions and formulate the problem. In Sec. III we address noiseless reading kernels, and state a tight asymptotic expression for the Bayesian error probability. In Sec. IV we similarly address noisy reading kernels, yet restrict to memoryless processes. In Sec. V we conclude the paper. In the appendices, we provide the proofs for the statements made throughout the paper.

II. PROBLEM FORMULATION

A. Notation conventions

We denote r.v.'s by capital letters, specific values they may take by the corresponding lower case letters, and their alphabets by calligraphic letters. We superscript random sequences and their realizations by their dimension. For example, the random sequence $X_1^n = (X_1, \ldots, X_n)$, where *n* is a positive integer, may take a specific vector value $x^n = (x_1, \ldots, x_n) \in \mathcal{X}^{\otimes n}$, the *n*th order Cartesian power of \mathcal{X} , which is the alphabet of each component of this vector. We denote a two-sided random process by $\mathbf{X} := X_{-\infty}^{\infty} = \{X_i\}_{i \in \mathbb{Z}}$. We denote the Cartesian product of \mathcal{X} and \mathcal{Y} (finite alphabets) by $\mathcal{X} \otimes \mathcal{Y}$, the p.m.f. of X by P_X , the p.m.f. of X_i^j by $P_{X_i^j}$, and by the joint p.m.f. of (X, Y) by P_{XY} , and by $P_{Y|X}$ the conditional p.m.f. of Y conditioned on X. We denote the complement of an event \mathcal{A} (or a set) by \mathcal{A}^c . We denote the order- α Rényi entropy of a p.m.f. P_X by

$$H_{\alpha}(P_X) := \begin{cases} \frac{1}{1-\alpha} \log \sum_{x \in \mathcal{X}} P_X^{\alpha}(x), & \alpha \neq 1\\ -\sum_{x \in \mathcal{X}} P_X(x) \log P_X(x), & \alpha = 1 \end{cases},$$
(8)

where we notice that $H_1(P_X)$ is the Shannon entropy. We will also use the minus-infinity-order Rényi entropy, defined as

$$H_{-\infty} := \lim_{\alpha \downarrow -\infty} H_{\alpha}(P_X) = -\log \min_{x \in \operatorname{supp}(P_X)} P_X(x), \tag{9}$$

which is determined by the minimal probability in the support of P_X . For a given random process **X**, with marginal p.m.f.'s $X_{-t}^t \sim P_{X_{-t}^t}$, we let

$$\overline{\mathcal{H}}_{\alpha}(\mathbf{X}) := \limsup_{t \to \infty} \frac{1}{2t+1} H_{\alpha}(P_{X_{-t}^{t}})$$
(10)

and

$$\underline{\mathcal{H}}_{\alpha}(\mathbf{X}) := \liminf_{t \to \infty} \frac{1}{2t+1} H_{\alpha}(P_{X_{-t}^{t}})$$
(11)

be the supremum (resp. infimum) order- α Rényi entropy rate. If the limit exists and $\underline{\mathcal{H}}_{\alpha}(\mathbf{X}) = \overline{\mathcal{H}}_{\alpha}(\mathbf{X})$ then the common limit $\mathcal{H}_{\alpha}(\mathbf{X})$ is the order α Rényi entropy rate. For the sake of simplicity, we assume henceforth that all considered entropy rates exists. Otherwise, one can replace the entropy rate $\mathcal{H}_{\alpha}(\mathbf{X})$ with either $\underline{\mathcal{H}}_{\alpha}(\mathbf{X})$ or $\overline{\mathcal{H}}_{\alpha}(\mathbf{X})$ (depending on the context).⁵

We denote the Rényi divergence pair of probability distributions $P_X \ll Q_X$ over \mathcal{X} by

$$D_{\alpha}(P_X \mid\mid Q_X) := \begin{cases} \frac{1}{\alpha - 1} \log\left(\sum_{x \in \mathcal{X}} P_X^{\alpha}(x) Q_X^{1 - \alpha}(x)\right), & \alpha \neq 1\\ \sum_{x \in \mathcal{X}} P_X(x) \log \frac{P_X(x)}{Q_X(x)}, & \alpha = 1 \end{cases}$$
(12)

where we notice that $D_1(P_X || Q_X)$ is the Kullback–Leibler divergence. We denote the total variation distance between P_X and Q_X by

$$||P - Q||_{\text{TV}} := \frac{1}{2} \sum_{x \in \mathcal{X}} |P(x) - Q(x)|.$$
(13)

For an integer $n \in \mathbb{N}_+$, we denote $[n] := \{1, 2, ..., n\}$ and for brevity, we also denote $-[n] := \{-1, -2, ..., -n\}$. We denote $a \lor b := \max\{a, b\}$ and $a \land b := \min\{a, b\}$ for $a, b \in \mathbb{R}$. Unless otherwise stated, we take logarithms to the base $|\mathcal{X}|$, and we also denote $\exp_a(t) := a^t$. We will use the standard Bachmann–Landau asymptotic notation, such as $O(\frac{\log n}{n})$, $o_n(1)$ (where here the subscript indicates that this term is vanishing when $n \to \infty$), and $\omega(\log n)$, and also use $a_n \sim b_n$ to denote that $\frac{a_n}{b_n} \to 1$ as $n \to \infty$. We will use \equiv to denote equivalence, mainly to simplify notation. For example, we will denote an element from a sequence $\{\tau(n)\}_{n\in\mathbb{N}}$ by τ , and indicate this by $\tau \equiv \tau(n)$.

⁵An exception is Theorem 9, in which the existence of $\mathcal{H}_1(\mathbf{X})$ is required to be assumed.




Figure 1. Reads of length $\ell = 5$. X(1) and X(2) have a positive overlap of T = 2, whereas $\tilde{X}(1)$ and X(2) have a negative overlap T = -2.

B. Problem Setting

We next model the process of sampling a pair of short substrings (sub-sequences of consecutive symbols) called *reads*, from a sequence, and the possible overlap between them. Let \mathcal{X} be a finite alphabet, and let $\mathbf{X} := \{X_i\}_{i \in \mathbb{Z}}$ be a random process defined over \mathcal{X} . Let $\ell \equiv \ell(n)$ denote the length of each read, where n determines the length of the available sequence X_1^n . We will assume that $\ell \ll n$, and concretely, we will focus on the regime $\ell = \Theta(\log n)$. For a reason that will be clarified next, we will slightly extend the available sequence to $X_{2-\ell}^{n+2\ell-2}$, whose length is $n + 3\ell - 3 = [1 + o_n(1)] \cdot n$ in this regime, and thus is essentially n.

Let $\{I(j)\}_{j \in [2]}$ be two starting indices, and let the *noiseless reads* be the substrings

$$X_1^{\ell}(j) = X_{I(j)}^{I(j)+\ell-1} \tag{14}$$

for $j \in [2]$. The pair of reads *overlap* if the indices of the reads have at least a single common index, that is, if $|I(1) - I(2)| < \ell$. We will be interested in the *overlap* between the reads. To this end, we define the *(signed) overlap* as

$$T := \begin{cases} 0 \lor [\ell - (I(2) - I(1))], & I(2) \ge I(1) \\ -\{0 \lor [\ell - (I(1) - I(2))]\}, & I(2) < I(1) \end{cases}.$$
(15)

Thus, T = 0 if the reads do not overlap, the overlap is positive $T \in [\ell]$ if the reads overlap and the last T indices of $X_1^{\ell}(1)$ in the sequence are the same as the first T indices of $X_1^{\ell}(2)$, and the overlap is negative $T \in -[\ell - 1]$ in the opposite case.⁶ See Fig. 1 for an illustration.

The direct way of modeling the random sampling process of the reads is to independently choose two random starting indices $I(j) \sim \text{Uniform}[n]$. As is, this read-sampling process results in an edge effect in the distribution of the amount of overlap, occurring when I(1) is sampled close to the edges, i.e., $I(1) \in [\ell]$ or $I(1) \in [n] \setminus [n - \ell - 1]$ (or, similarly, I(2)). Since we are interested in $n \gg \ell$, these edge effects are negligible,

⁶Note that when $T = \ell$, there is no distinction between the read with smaller index and the read with larger index. We arbitrarily attribute this case to a positive overlap of ℓ .

and so we slightly modify the sampling process in order to make later derivations less cumbersome. Concretely, we assume that $I(1) \sim \text{Uniform}[n]$ (or even arbitrary chosen in [n]), and that

$$I(2) \mid I(1) \sim \begin{cases} I(1) + \text{Uniform}\{-\ell + 1, -\ell + 2, \dots, n - \ell\}, & 1 \le I(1) \le \ell - 1\\ \text{Uniform}[n], & \ell \le I(1) \le n - \ell + 1\\ I(1) + \text{Uniform}\{\ell - n, \ell - n + 1, \dots, \ell - 1\}, & n - \ell + 2 \le I(1) \le n \end{cases}$$
(16)

So, the distribution of I(2) is always uniform over a window of length n regardless of I(1), the reads are always sampled from the extended available sequence $X_{2-\ell}^{n+2\ell-2}$, and the sampling window location of I(2) only slightly changes at the edges, whose total size is $2\ell \ll n$. With this slight simplification, it holds that T is an r.v. whose p.m.f. is

$$P_{T}(t) = \begin{cases} 1 - \frac{(2\ell - 1)}{n}, & t = 0\\ \frac{1}{n}, & t \in -[\ell - 1] \cup [\ell] \end{cases}$$
$$= \begin{cases} \frac{n_{\ell}}{n}, & t = 0\\ \frac{1}{n}, & t \in -[\ell - 1] \cup [\ell] \end{cases}$$
(17)

where, for brevity, we denoted $n_{\ell} := n - (2\ell - 1)$ and note that $n_{\ell} \sim n$ whenever $\ell = o(n)$.

Let \mathcal{Y} be another finite alphabet, and let $P_{Y_1^{\ell}|X_1^{\ell}}$ be a Markov kernel from $\mathcal{X}^{\otimes \ell}$ to $\mathcal{Y}^{\otimes \ell}$. For $j \in [2]$, let

$$Y_1^{\ell}(j) \sim P_{Y_1^{\ell} | X_1^{\ell}} \left(\cdot \mid x_1^{\ell}(j) \right)$$
(18)

be the noisy read of $x_1^{\ell}(j)$, where conditioned on $\{X_1^{\ell}(j) = x_1^{\ell}(j)\}_{j \in [2]}$ it holds that $\{Y_1^{\ell}(j)\}_{j \in [2]}$ are pairwise independent. The goal is to detect the signed overlap T based on these noisy reads. Letting $\mathcal{T} := \{-[\ell - 1]\} \cup \{0\} \cup [\ell]$, an overlap detector based on a pair of noisy reads is a mapping $\hat{T}: \mathcal{Y}^{\otimes \ell} \times \mathcal{Y}^{\otimes \ell} \to \mathcal{T}$. The Bayesian error probability of a detector \hat{T} is given by

$$P_{\text{error}}(\hat{T}) := \mathbb{P}\left[\hat{T}\left(Y_1^{\ell}(1), Y_1^{\ell}(2)\right) \neq T\right].$$
(19)

In many cases, there is a conceptual distinction between detection of $\hat{T} = 0$ – no overlap, and $\hat{T} \neq 0$ – a discovery of an overlap. Thus, with a slight deviation from the standard statistical terminology, we define the type-I error of a detector \hat{T} as the event that T = 0, but \hat{T} is detected as a discovery overlap, and denote its probability by

$$p_{\text{type-I}}(\hat{T}) := \mathbb{P}\left[\hat{T} \neq 0 \mid T = 0\right].$$
(20)

We then also define the type-II error as the event that T = t for some non-zero overlap $t \neq 0$, and it is either

misdetected or erroneously detected, and denote its probability by

$$p_{\text{type-II}}(\hat{T};t) := \mathbb{P}\left[\hat{T} \neq t \mid T = t\right],\tag{21}$$

where we also denote, again with a slight abuse of notation,

$$p_{\text{type-II}}(\hat{T}) = \sum_{t \in \{-[\ell-1]\} \cup [\ell]} \mathbb{P}[T = t \mid T \neq 0] \cdot p_{\text{type-II}}(\hat{T}; t).$$
(22)

The Bayesian error probability is then given by

$$P_{\text{error}}(\hat{T}) = P_{T}(0) \cdot p_{\text{type-I}}(\hat{T}) + \mathbb{P}[T \neq 0] \cdot \sum_{t \in \{-[\ell-1]\} \cup [\ell]} \mathbb{P}[T = t \mid T \neq 0] \cdot p_{\text{type-II}}(\hat{T}; t).$$
(23)

For the considered overlap p.m.f. P_T in (17), the Bayesian error probability is (tightly) upper bounded as

$$P_{\text{error}}(\hat{T}) \sim p_{\text{type-I}}(\hat{T}) + \frac{2\ell}{n} \cdot p_{\text{type-II}}(\hat{T}).$$
(24)

We will be interested in the minimal error probability

$$P_{\text{error}}^* \equiv P_{\text{error}}^* \left(\beta, P_{\boldsymbol{X}}, P_{Y_1^{\ell} | X_1^{\ell}} \right) := \min_{\hat{T}} P_{\text{error}}(\hat{T}).$$
(25)

As the error probability (19) is averaged over the prior distribution, the optimal detector is the maximum *a* posteriori (MAP) detector. For a given realization of a pair of noisy reads $y_1^{\ell}(1), y_1^{\ell}(2)$, it is given by the Bayes rule as

$$\hat{T}_{\text{MAP}}\left(y_1^{\ell}(1), y_1^{\ell}(2)\right) = \operatorname*{arg\,max}_{t \in \mathcal{T}} P_{T|Y_1^{\ell}(1)Y_1^{\ell}(2)}\left(t \mid y_1^{\ell}(1), y_1^{\ell}(2)\right)$$
(26)

$$= \underset{t \in \mathcal{T}}{\arg\max} P_T(t) \cdot P_{Y_1^{\ell}(1)Y_1^{\ell}(2)|T}\left(y_1^{\ell}(1), y_1^{\ell}(2) \mid T = t\right),$$
(27)

where we denote the likelihood by

$$f\left(y_1^{\ell}(1), y_1^{\ell}(2); t\right) := P_{Y_1^{\ell}(1)Y_1^{\ell}(2)|T}\left(y_1^{\ell}(1), y_1^{\ell}(2) \mid T = t\right).$$

$$(28)$$

For any given overlap $t \in [\ell]$, we may partition $Y_1^{\ell}(1)$ into two segments, given by $Y_1^{\ell-t}(1)$, which does not overlap with $Y_1^{\ell}(2)$, and $Y_{\ell-t+1}^{\ell}(1)$, which overlaps with $Y_1^{t}(2)$. Similarly, we also partition $Y_1^{\ell}(2)$ into $Y_1^{t}(2)$ which overlaps with $Y_{\ell-t+1}^{\ell}(1)$, and $Y_{t+1}^{\ell}(2)$, which does not overlap with $Y_1^{\ell}(1)$. The two reads may be analogously partitioned when $t \in -[\ell-1]$.

We will assume throughout that the read length scales as $\ell = \beta \log n$, where $\beta > 0$ is a fixed parameter (in fact, the interesting regime is in which $\beta > \log |\mathcal{X}| = 1$). We will focus on two specific settings:

Definition 1 (Noiseless setting). **X** is a stationary and ergodic process, and the reading kernel $P_{Y_1^{\ell}|X_1^{\ell}}$ is a noiseless channel, that is $Y_1^{\ell} = X_1^{\ell}$, with probability 1.

For simplicity, when analyzing this setting, we will use $\{X_1^{\ell}(j)\}_{j \in [2]}$ to denote the reads.

Definition 2 (Noisy setting). X is stationary and memoryless, that is, $P_{X^n}(x^n) = \prod_{i=1}^n P_X(x_i)$ for any $n \in \mathbb{N}_+$, and the reading channel is also stationary and memoryless, that is, $P_{Y_1^\ell|X_1^\ell} = \prod_{i=1}^\ell P_{Y|X}(y_i \mid x_i)$ for all $\ell \in \mathbb{N}_+$.

Finally, we abbreviate $P_{\text{error}}^* \equiv P_{\text{error}}(\hat{T}_{\text{MAP}})$. Our goal is to determine

$$\phi\left(\beta, P_{\boldsymbol{X}}, P_{Y_1^{\ell}|X_1^{\ell}}\right) := \lim_{n \to \infty} \frac{n}{\log n} \cdot P_{\text{error}}^*$$
(29)

for both the noiseless and the noisy settings.

III. OVERLAP DETECTION FOR NOISELESS READING KERNELS

In this section, we address the noiseless setting of Definition 1.

A. The MAP Detection Rule

Our first step in deriving the MAP detection rule is to obtain an expression for the likelihood function.

Lemma 3. Consider the noiseless setting (Def. 1). Let

$$\tilde{\Gamma}_{+}(t; x_{1}^{\ell-t}) := \frac{1}{P_{X_{1}^{t} \mid X_{-(\ell-t-1)}^{0}}(x_{1}^{t}(2) \mid x_{1}^{\ell-t})}$$
(30)

and let

$$\tilde{\Gamma}_{-}(t; x_{1}^{\ell-t}) := \frac{1}{P_{X_{1}^{t} \mid X_{-(\ell-t-1)}^{0}}(x_{1}^{t}(1) \mid x_{1}^{\ell-t})}$$
(31)

Then, the likelihood of overlap $t \in [\ell]$ given reads $x_1^{\ell}(1)$ and $x_1^{\ell}(2)$ is

$$f\left(x_{1}^{\ell}(1), x_{1}^{\ell}(2); t\right) \propto \mathbb{1}\left[x_{\ell-t+1}^{\ell}(1) = x_{1}^{t}(2)\right] \cdot P_{X_{1}^{\ell} \mid X_{-(\ell-t-1)}^{0}}\left(x_{1}^{\ell}(2) \mid x_{1}^{\ell-t}(1)\right) \cdot \tilde{\Gamma}_{+}(t; x_{1}^{\ell-t}(1)),$$
(32)

and for $t \in -[\ell - 1]$

$$f\left(x_{1}^{\ell}(1), x_{1}^{\ell}(2); t\right) \propto \mathbb{1}\left[x_{\ell-t+1}^{\ell}(2) = x_{1}^{t}(1)\right] \cdot P_{X_{1}^{\ell}|X_{-(\ell-t-1)}^{0}}\left(x_{1}^{\ell}(1) \mid x_{1}^{\ell-t}(2)\right) \cdot \tilde{\Gamma}_{-}(t; x_{1}^{\ell-t}(2)) \cdot,$$
(33)

and where we set $\Gamma_{\pm}(0) = 1$.

Proof: It holds that

$$f(x_{1}^{\ell}(1), x_{1}^{\ell}(2); t) = P_{X_{1}^{\ell}}\left(x_{1}^{\ell}(1)\right) \cdot \mathbb{1}\left[x_{\ell-t+1}^{\ell}(1) = x_{1}^{t}(2)\right] P_{X_{t+1}^{\ell}|X_{-(\ell-t-1)}^{t}}\left(x_{t+1}^{\ell}(2) \mid (x_{1}^{t}(2), x_{1}^{\ell-t}(1))\right)$$

$$P_{X_{1}^{\ell}}\left(x_{1}^{\ell}(2) \mid x_{1}^{\ell-t}(1)\right)$$

$$(34)$$

$$\stackrel{(a)}{=} P_{X_1^{\ell}}\left(x_1^{\ell}(1)\right) \mathbb{1}\left[x_{\ell-t+1}^{\ell}(1) = x_1^{t}(2)\right] \cdot \frac{P_{X_1^{\ell}|X_{-(\ell-t-1)}^{0}}\left(x_1^{t}(2) \mid x_1^{t-1}(1)\right)}{P_{X_1^{t}|X_{-(\ell-t-1)}^{0}}\left(x_1^{t}(2) \mid x_1^{\ell-t}(1)\right)}$$
(35)

$$\stackrel{(b)}{=} \mathbb{1} \left[x_{\ell-t+1}^{\ell}(1) = x_1^t(2) \right] \cdot P_{X_1^{\ell}} \left(x_1^{\ell}(1) \right) \cdot P_{X_1^{\ell} \mid X_{-(\ell-t-1)}^0} \left(x_1^{\ell}(2) \mid x_1^{\ell-t}(1) \right) \cdot \tilde{\Gamma}_+(t; x_1^{\ell-t}(1)), \tag{36}$$

where (a) follows from Bayes rule, and in (b) we have used the notation for $\tilde{\Gamma}_+(t; x_1^{\ell-t})$. This shows (32), and (33) follows by analogous arguments.

While the likelihood function appears to be rather cumbersome, and leads to a complicated MAP rule, it is greatly simplified when \mathbf{X} is a memoryless process. Moreover, it turns out that the MAP rule in the memoryless case (in the form we state it) is a good approximation in the case that the process \mathbf{X} has memory, yet ergodic. Specifically, in the memoryless case, the MAP detector takes the following simple form:

Proposition 4. Consider the noiseless setting (Def. 1), and further assume that X is a memoryless process. Let the maximum overlap with strictly positive (resp., negative) posterior be

$$t_{+} := \max_{t \in [\ell]} \mathbb{1} \left[x_{\ell-t+1}^{\ell}(1) = x_{1}^{t}(2) \right]$$
(37)

$$t_{-} := \max_{t \in [\ell-1]} \mathbb{1} \left[x_{\ell-t+1}^{\ell}(2) = x_{1}^{t}(1) \right]$$
(38)

and let

$$\Gamma_{+}(t) := \frac{1}{P_{X_{1}^{t}}(x_{1}^{t}(2))}$$
(39)

$$\Gamma_{-}(t) := \frac{1}{P_{X_{1}^{t}}(x_{1}^{t}(1))}.$$
(40)

Then,

$$\hat{T}_{\text{MAP}}\left(x_{1}^{\ell}(1), x_{1}^{\ell}(2)\right) = \begin{cases} t_{+}, & \Gamma_{+}(t_{+}) \ge [\Gamma_{-}(t_{-}) \lor n_{\ell}] \\ -t_{-}, & \Gamma_{-}(t_{-}) \ge [\Gamma_{+}(t_{+}) \lor n_{\ell}] \\ 0, & \text{otherwise} \end{cases}$$
(41)

Proof: The Bayesian detector compares the posterior probabilities among the different possibilities $t \in -[\ell - 1], t = 0$ and $t \in [\ell]$. First, let us compare the posterior probabilities only among overlaps in $t \in [\ell]$. Using (32) from Lemma 3, it holds for $t \in [\ell]$ under the assumption that X is a memoryless process that

$$\arg\max_{t\in[\ell]} f(x_1^{\ell}(1), x_1^{\ell}(2); t) = \arg\max_{t\in[\ell]} \mathbb{1} \left[x_{\ell-t+1}^{\ell}(1) = x_1^{t}(2) \right] \cdot P_{X_1^{\ell}} \left(x_1^{\ell}(1) \right) \cdot P_{X_1^{\ell}} \left(x_1^{\ell}(2) \right) \cdot \Gamma_+(t),$$
(42)

where $\Gamma_{+}(t)$ is as defined in (39). Since the prior distribution $P_{T}(t) = \frac{1}{n}$ for all $t \in [\ell]$, and as $\Gamma_{+}(t)$ is a monotonic non-decreasing function of $t \in [\ell]$ we deduce that if $\hat{T}_{MAP} \in [\ell]$ then $\hat{T}_{MAP} = t_{+}$. Analogously, if $\hat{T}_{MAP} \in -[\ell-1]$ then $\hat{T}_{MAP} = t_{-}$. The decision between $\hat{T}_{MAP} = 0$, $\hat{T}_{MAP} = t_{+}$ and $\hat{T}_{MAP} = t_{-}$ is again according to the maximal posterior probability, which is proportional to $P_{T}(t) \cdot f(x^{\ell}(1), x^{\ell}(2); t)$. As the likelihood for t = 0 is

$$f\left(x_1^{\ell}(1), x_1^{\ell}(2); 0\right) = P_{X_1^{\ell}}\left(x_1^{\ell}(1)\right) \cdot P_{X_1^{\ell}}\left(x_1^{\ell}(2)\right), \tag{43}$$

and as the priors are $P_T(0) = \frac{n_\ell}{n}$ and $P_T(t) = \frac{1}{n}$ for $t \in -[\ell-1] \cup [\ell]$, utilizing (42) and (43) implies the stated claim.

In words, the MAP detector in the memoryless case finds the maximal positive overlap and the maximal negative overlap, and chooses between them based on the maximal value of the corresponding inverse sequence probability, computed at the potential overlapping segments. If this value crosses the threshold n_{ℓ} then an overlap is detected. Otherwise, the detector declares that there is no overlap.

Comparing the likelihood in the memoryless case (42) and the case with memory (32), we see that $P_{X_1^\ell}(x_1^\ell(2))$ is replaced by $P_{X_1^\ell|X_{-(\ell-t-1)}^0}(x_1^\ell(2) | x_1^{\ell-t}(1))$, and that $\Gamma_+(t)$ is replaced by $\tilde{\Gamma}_+(t; x_1^{\ell-t}(1))$. To wit, both terms now have a dependency on past symbols $X_{-(\ell-t-1)}^0 = x_1^{\ell-t}(1)$, assuming I(1) = 1 without loss of generality (w.l.o.g.). Moreover, the time span of this past symbols depends on t, and so it cannot be omitted from the likelihood, unlike the first term in the memoryless case $P_{X_1^\ell}(x_1^\ell(2))$. An analogous expression can be derived for $t \in -[\ell-1]$, which introduces an analogous situation. Similarly, the expression for the likelihood condition on t = 0 is invalid, in general, since due to the memory in the process, the reads $X_1^\ell(1)$ and $X_1^l(2)$ may be dependent even if they do not overlap. Nonetheless, we argue that under mild conditions, the MAP detector (41) derived in Prop. 4 for the memoryless case is a viable option for the case with memory, as it is a sufficiently good approximation to it. We will fully justify this argument in what follows by showing that the detector (41) of Prop. 4 achieves asymptotically optimal error probability, as the exact MAP detector. To make this notion precise, let us recall the following definition mixing of a random process:

Definition 5 (Strong mixing coefficient). Let $\mathbf{X} := \{X_i\}_{i \in \mathbb{Z}}$ be a stationary and ergodic random process. For a given delay $s \in \mathbb{N}_+$, the *strong mixing coefficient* is

$$d(s) \equiv d(s; \mathbf{X}) := \sup\left\{ |\mathbb{P}[\mathcal{F} \mid \mathcal{E}] - \mathbb{P}[\mathcal{F}]| : i \in \mathbb{Z}, \ \mathcal{E} \in \sigma\left(X_{-\infty}^{i}\right), \ \mathcal{F} \in \sigma\left(X_{i+s}^{\infty}\right) \right\},$$
(44)

where $\sigma(X_i^j)$ is the sigma-algebra generated by X_i^j . The process **X** is *strongly mixing* if $d(s) \to 0$ as $s \to \infty$. In this case, since we can always replace d(s) with $\tilde{d}(s) := \sup_{s' \ge s} d(s)$, we may assume, w.l.o.g., that d(s) is monotonically non-increasing.

Hence, for strongly mixing processes, we expect that the exact likelihood for the case with memory is well approximated by the likelihood in Prop. 4. Indeed, for $T = t \in [\ell]$ the conditioning on past symbols $X_{-(\ell-t-1)}^0 = x_1^{\ell-t}(1)$ will only affect a relatively small number of symbols at the beginning of the read, and so

$$f\left(x_{1}^{\ell}(1), x_{1}^{\ell}(2); t\right) \approx \mathbb{1}\left[x_{\ell-t+1}^{\ell}(1) = x_{1}^{t}(2)\right] \cdot \frac{P_{X_{1}^{\ell}}\left(x_{1}^{t}(2)\right)}{P_{X_{1}^{t}}(x_{1}^{t}(2))},\tag{45}$$

as the likelihood in Prop. 4. Similarly, for T = 0, since the reads length is $\ell = \Theta(\log n)$ and the length of the entire sequence is n, under mild assumption on the mixing of process they will be approximately independent

for most instances of I(1) and I(2). Then, we may expect that

$$f\left(x_1^{\ell}(1), x_1^{\ell}(2); 0\right) \approx P_{X_1^{\ell}}\left(x_1^{\ell}(1)\right) \cdot P_{X_1^{\ell}}\left(x_1^{\ell}(2)\right), \tag{46}$$

which is again, similar to the likelihood in Prop. 4. At this point, we do not get into precise conditions on d(s) for sufficient approximate independence to holds,⁷ but only say that following this line of thought, it makes sense to use the detector (41) of Prop. 4 even when the source has memory (note that $\Gamma_{\pm}(t)$ are well defined for this case too). In this case, the error probability of this sub-optimal detector is an upper bound on the Bayesian error probability.

For the detector (41) of Prop. 4, there exists a minimal absolute overlap value that can be detected. Specifically, let the *minimal detectable overlap* be defined as

$$t_{\rm MDO} \equiv t_{\rm MDO}(n) := \max\left\{ t \in [\ell]: \max_{x_1^t \in \mathrm{supp}(P_{X_1^t})} \frac{1}{P_{X_1^t}(x_1^t)} \le n_\ell \right\}$$
(47)

$$= \max\left\{t \in [\ell]: \log \frac{1}{\min_{x_{1}^{t} \in \operatorname{supp}(P_{X_{1}^{t}})} P_{X_{1}^{t}}(x_{1}^{t})} \le \log n_{\ell}\right\}$$
(48)

$$= \max\left\{t \in [\ell]: H_{-\infty}(P_{X_1^t}) \le \log n_{\ell}\right\}.$$
(49)

Prop. 4 implies that $|t| \le t_{\text{MDO}}$ will never be detected by the detector (41). It is readily verified that $t_{\text{MDO}}(n)$ is a monotonic non-decreasing sequence, which also satisfies $\lim_{n\to\infty} t_{\text{MDO}}(n) = \infty$ (assuming $\mathcal{H}_{-\infty}(\mathbf{X}) < \infty$). Furthermore, if the minus-infinity-order Rényi entropy rate exists, then

$$t_{\text{MDO}}(n) \sim \frac{\log n_{\ell}}{\frac{1}{t_{\text{MDO}}(n)} H_{-\infty}(P_{X_1^{t_{\text{MDO}}(n)}})} \sim \frac{\log n}{\mathcal{H}_{-\infty}(\mathbf{X})},\tag{50}$$

which scales linearly with the read length whenever $\ell = \Theta(\log n)$. Nonetheless, in the *typical* case, even longer overlaps are not expected to be reliably detected. The scaling of $t_{\text{MDO}}(n) = \Theta(\log n)$ shows that we need to consider the typical value of $\frac{1}{t} \log \frac{1}{P_{X_1^t}(x_1^t)}$, and indeed, if we take the *expected* value of $\frac{1}{t} \log \frac{1}{P_{X_1^t}(x_1^t)}$ instead of its maximal value in (49), it will tend to $\mathcal{H}_1(\mathbf{X})$ (whenever this entropy rate exists). Following this intuition, we may expect that

$$t^* \equiv t^*(n) \sim \frac{\log n}{\mathcal{H}_1(\mathbf{X})} \tag{51}$$

is a minimal reliably detectable overlap.⁸ As we shall see in the next section, this intuition is correct.

B. Error Probability Analysis

We next separately state an upper bound on the Bayesian error probability, and then a matching lower bound.

We begin with the upper bound:

⁷See Lemma 19 and its proof in what follows for the exact likelihood for t = 0, and a rigorous quantitative statement of the approximation (46).

⁸This is indeed a longer overlap, since the standard entropy rate $\mathcal{H}_1(\mathbf{X})$ (assuming it exists) is smaller or equal to $\mathcal{H}_{-\infty}(\mathbf{X})$.

Theorem 6. Consider the noiseless setting (Def. 1), and assume that the following conditions hold:

- 1) Finite minus-infinity-order Rényi entropy rate: $\mathcal{H}_{-\infty}(\mathbf{X}) < \infty$.
- 2) The process is moderately-fast strongly mixing: $\sum_{s=0}^{\infty} d(s; \mathbf{X}) < \infty$.
- 3) The process is aperiodic:

$$\mathcal{R}(\mathbf{X}) := \sup_{s \in \mathbb{N}_+} \mathbb{P}\left[X_1 = X_{1+s}\right] < 1.$$
(52)

Then,

$$P_{\text{error}}^* \le 2\left[1 + o_n(1)\right] \cdot \left\{\beta \wedge \frac{1}{\mathcal{H}_1(\mathbf{X})}\right\} \cdot \frac{\log n}{n}.$$
(53)

The proof of Prop. 4, is deferred to Appendix A, as all the other proofs in this section. We next discuss the result of Theorem 6.

The dependence of the error probability on β : The root of this dependence is simple to explain: A trivial detector, which always decides that there is no overlap, will err only if $T \neq 0$. The probability of this event is $\frac{2\ell}{n} \sim \frac{2\beta \log n}{n}$, and thus the error probability of the optimal MAP detector is less than $\frac{2\beta \log n}{n}$ (asymptotically). This explains the upper bound in (53) when $\beta \leq \frac{1}{\mathcal{H}_1(\mathbf{X})}$, and so we assume in the rest of the discussion that $\beta > \frac{1}{\mathcal{H}_1(\mathbf{X})}$.

The dependence of the error on the entropy rate $\mathcal{H}_1(\mathbf{X})$: This dependence is much more subtle. Let us assume for the sake of this discussion that the source is memoryless. In this case, the MAP detector is \hat{T} given in (41) (Prop. 4). On the face of it, one might expect that the error probability is determined by the second-order Rényi entropy, also known as the *collision entropy*, which is given by

$$H_2(P_X) = -\log \sum_{x \in \mathcal{X}} P_X^2(x).$$
(54)

This quantity indeed appears in various previous works on reads alignment, e.g., [22]. This is because whenever T = 0, the two reads do not overlap, and the probability that two independent substrings of length t exactly match and a false overlap is detected is $|\mathcal{X}|^{-tH_2(P_{X_1})}$. Following this line of thought, one might expect, that the Bayesian error probability is upper bounded as

$$P_{\text{error}}^* \le [1 + o_n(1)] \cdot \frac{2}{H_2(P_X)} \cdot \frac{\log n}{n}.$$
 (55)

The fallacy of such an argument is that it ignores that an overlap is detected only if the probability of the matching substring is sufficiently small, as expressed in the condition

$$\Gamma_{+}(t) = \frac{1}{P_{X_{1}^{t}}(x_{1}^{t}(2))} \ge n_{\ell}$$
(56)

in the MAP detector (41). The root of this condition is that the likelihood of the two reads at the given (false) overlap should overcome the strong prior probability that the two reads do not overlap at all. As we have

discussed above, the typical length of strings which satisfy this condition is the minimal reliably detectable overlap given by $t^* \sim \frac{\log n}{H_1(P_X)}$, and this introduces dependence on $H_1(P_{X_1})$. Nonetheless, from the discussion at this level it is not obvious which of the two entropy values is the relevant one. Next, we will outline the proof of Theorem 6, and this will clarify this issue.

The main ideas of the proof: The proof of Theorem 6 is fairly long and involved, so we next outline its steps. The proof carefully analyzes the different types of error probability in the decomposition (24) for the detector (41) of Prop. 4. In the case of memory, this detector is sub-optimal, and its error probability is thus an upper bound on the Bayesian error probability of the MAP detector.

Let us begin by considering $p_{\text{type-1}}(\hat{T})$, that is, the error conditioned on T = 0. Recall that the detector \hat{T} never decides $0 < |\hat{T}| < t_{\text{MDO}}$, where $t_{\text{MDO}} \sim \frac{\log n}{\mathcal{H}_{-\infty}(\mathbf{X})}$, because for $0 < |T| < t_{\text{MDO}}$, the posterior evidence of any substring common to both reads is not strong enough compared to the prior knowledge that there is no overlap between the reads. This has two implications. First, the error probability is at least the probability that $0 < |T| < t_{\text{MDO}}$, which is $\frac{t_{\text{MDO}}(n)}{n} = \Omega(\frac{\log n}{n})$. Second, the detector will never err from T = 0 to $0 < |\hat{T}| < t_{\text{MDO}}$, only to longer overlaps (either positive or negative). As it turns out, however, this property alone does not suffice to tightly bound $p_{\text{type-I}}(\hat{T})$. So, let us generalize this observation and consider, instead of $t_{\text{MDO}} \sim \frac{\log n}{\mathcal{H}_{-\infty}(\mathbf{X})}$, a general effective minimal overlap value, given by $t(h) \sim \frac{\log n}{h}$ where $h < \mathcal{H}_{-\infty}(\mathbf{X})$, and aim to find the minimal h which leads to the tightest upper bound on $p_{\text{type-I}}(\hat{T})$. Conditioned on T = 0, according to the detector of Prop. 4, an error will occur whenever two non-overlapping reads of length longer than t(h) match, and their probability is smaller than $1/n_{\ell}$. We then take a union bound over all possible lengths $t(h) \leq t \leq \ell = \beta \log n$, so there are $\Theta(\log n)$ such possibilities. To discuss the analysis of one of these possibilities, that is, an error from T = 0 to some $\hat{T} = t > t(h)$, let us assume for simplicity that \mathbf{X} is memoryless. By ignoring here several subtle technical details, simple algebraic manipulations show that

$$\mathbb{P}\left[\hat{T}=t \mid T=0\right] \lesssim \sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}^2\left(x_1^t\right) \cdot \mathbb{1}\left[\frac{1}{P_{X_1^t}(x_1^t)} \ge n\right].$$
(57)

If we can guarantee that this probability is $o(\frac{1}{n})$, then after the union bound over all the candidates $t(h) \le t \le \ell$ ($\Theta(\log n)$ terms) we obtain that

$$p_{\text{type-I}}(\hat{T}) = o\left(\frac{\log n}{n}\right).$$
 (58)

Since, as explained above, the error probability is $\Omega(\frac{\log n}{n})$, this means that $p_{\text{type-I}}(\hat{T})$ has a negligible effect on the error probability. Thus, let us explore what is the minimal value of h required to make the right-hand side (r.h.s.) of (57) $o(\frac{1}{n})$. Evidently, if we trivially upper bound the indicator by 1 we get

$$\sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}^2(x_1^t) = \exp_{|\mathcal{X}|} \left[-tH_2(P_{X_1}) \right] \le \exp_{|\mathcal{X}|} \left[-\frac{H_2(P_{X_1})}{h} \log n \right] = n^{-\frac{H_2(P_{X_1})}{h}}$$
(59)

and so $h < H_2(P_{X_1})$ suffices. However, the indicator in the sum (57) is $\mathbb{1}[P_{X_1^t}(x_1^t) \le 1/n]$, and retains only small probabilities in the summation. Thus, one may conjecture that taking this indicator into account may substantially reduce the value of the sum below $n^{-\frac{H_2(P_{X_1})}{\hbar}}$ (as obtained when the indicator is trivially bounded by 1). The proof of Theorem 6 begins with Lemma 14 (Appendix A), which uses a Chernoff-style bound on the indicator function to show that this is indeed the case, and that the sum in (57) is essentially bounded by

$$\exp_{|\mathcal{X}|}\left[-\frac{H_1(P_{X_1})}{h}\log n\right] = n^{-\frac{H_1(P_{X_1})}{h}}.$$
(60)

This allows to show that (57) is $o(\frac{1}{n})$, for all $h < H_1(P_{X_1})$, which improves the previous sufficient condition $h < H_2(P_{X_1})$ (recall that the monotonicity property of the Rényi entropy assures that $H_2(P_{X_1}) \le H_1(P_{X_1})$). Lemma 14 shows this property in greater generality, for any strongly mixing process **X**, with condition given by the entropy rate $\mathcal{H}_1(\mathbf{X})$. This justifies that $t^* \sim \frac{\log n}{\mathcal{H}_1(\mathbf{X})}$ is the *effective, or reliable*, minimal detectable overlap.

Given that the type-I error probability is a negligible $o(\frac{\log n}{n})$, we now turn to consider the type-II error probability. Due to the symmetry between positive and negative overlaps, let us upper bound the contribution of T > 0, given by

$$\sum_{t \in [\ell]} \mathbb{P}[T=t] \cdot p_{\text{type-II}}(\hat{T};t) = \frac{1}{n} \sum_{t \in [\ell]} p_{\text{type-II}}(\hat{T};t).$$
(61)

We may then double it to include also the contribution of T < 0. Now, if $0 < T < t_{\text{MDO}}$, then the conditional error probability is 1. However, for the type-I error probability to be negligible it is required to avoid deciding $T < t^*$, we upper bound the conditional error probability by 1 for all $0 < T < t^*$. The total contribution of such events to the error probability is thus $\frac{t^*}{n} \sim \frac{1}{\mathcal{H}_1(\mathbf{X})} \frac{\log n}{n}$. This is the dominating term in the bound, and multiplying it by 2, to account for negative overlaps, leads to the bound of Theorem 6. Our previous refinement of the upper bound in (57) becomes crucial here, because had our necessary condition was $h < \mathcal{H}_2(\mathbf{X})$, then this term was larger $\sim \frac{1}{\mathcal{H}_2(\mathbf{X})} \frac{\log n}{n}$.

It is thus remain to bound $p_{type-II}(\hat{T};t)$ for $T = t > t^*$. Specifically, there are $\ell - t^* = \Theta(\log n)$ such terms, and so if we show that each of them is $o_n(1)$, then after multiplying by the prior $\mathbb{P}[T = t] = \frac{1}{n}$, their overall contribution to the error probability will be negligible $o(\frac{\log n}{n})$. An error from $T = t > t^*$ may be to one of three possibilities: First, to $\hat{T} = 0$, second, to a longer overlap $\hat{T} > T$, and third, to a longer negative overlap. Let us begin with the first case, which is somewhat dual to the type-I error probability discussed above. In the memoryless case, it can be shown to be upper bounded as

$$\mathbb{P}\left[\hat{T}=0 \mid T=t\right] \lesssim \sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}(x_1^t) \cdot \mathbb{1}\left[\frac{1}{P_{X_1^t}(x^t)} \le n\right],\tag{62}$$

which is similar to the upper bound in (57), albeit the summed probability is not squared here, and the inequality in the indicator is reversed, that is, the sum retains large probabilities. An additional result of Lemma 14, proved similarly, is that the same condition $t > t^* \sim \frac{\log n}{\mathcal{H}_1(\mathbf{X})}$ suffices to make this upper bound $o_n(1)$ (which




Figure 2. A case in which X(1) and X(2) have both overlap of length $\tilde{t} = 2$ and $\bar{t} = 3$ for $\ell = 5$. We show X(1) and the two possible alignments of X(2) to X(1). By relating the two possible alignments we obtain relations between the symbols of X(1). For example, the overlap of length \tilde{t} (top) implies that the sub-sequence marked in red (square) in X(1) equals the sub-sequence marked in blue (x) in X(2). The overlap of length \bar{t} (bottom) implies that the sub-sequence marked in grey (disk) in X(1) also equals the sub-sequence marked in grey (disk) in X(2). Both together imply that the sub-sequence marked in red (square) in X(1) equals the sub-sequence marked in grey (disk) in X(1). Using similar arguments, it stems that $X_{I(1)+2}^{I(1)+4} = X_{I(1)+3}^{I(1)+5}$, a repetition of length $\bar{t} = 3$ (here it is "tail-biting", but this is not the case for larger differences $\bar{t} - \tilde{t}$).

is less demanding than the $o(\frac{1}{n})$ derived above for (57)). As for (57), Lemma 14 shows this result for general processes, possibly with memory. The second and third cases, to wit, an error to $\hat{T} > T$ or to $\hat{T} < -T$ can be similarly bounded, and so we only discuss the second case. Specifically, let us focus on an error from $T = \tilde{t} > t^*$ to $\hat{T} = \tilde{t} > \tilde{t}$. According to the detection rule of \hat{T} ((41) in Prop. 4), such an error occurs when both $X_{\ell-\tilde{t}+1}^{\ell}(1) = X_1^{\tilde{t}}(2)$ and $1/P_{X_1^{\tilde{t}}}(X_1^{\tilde{t}}(2)) \ge n_{\ell}$ occur. For an upper bound, we may ignore the second condition, and only upper bound the probability of the first one. This event implies that while the true overlap is \tilde{t} , there is also a matching for a longer overlap \tilde{t} . It is not difficult to verify that for this to occur, it must be that the process X_1^n contains a *repetition* of length \tilde{t} , that is, an event of the form $\{X_1^{\tilde{t}} = X_{1+s}^{\tilde{t}+s}\}$ for some $s \in \mathbb{N}_+$. See Fig. 2 for an illustration. Lemma 15 establishes that if the process is aperiodic and strongly mixing then the probability of any such repetition is $o(\frac{1}{\log n})$. After taking a union bound over all possible $\tilde{t} > \tilde{t}$ (there are $\Theta(\log n)$ such terms), the resulting contribution to the conditional error probability is $\mathbb{P}[\hat{T} > T \mid T = \tilde{t}] = o(\frac{1}{n})$. Taking a union bound over all possible $T = \tilde{t}$ (again, there are $\Theta(\log n)$ such terms) results a total contribution to the type-II error probability, and thus to the Bayesian error probability of $o(\frac{\log n}{n})$, which is negligible compared to the $\frac{1}{\mathcal{H}_1(\mathbf{X})} \frac{\log n}{n}$ contribution from $T < t^*$. This completes the upper bounds of the various types of error probability.

To conclude the description of the proof's main ideas, we next shortly outline the proof of Lemma 15, which upper bounds the probability of the repetition $\{X_1^{\overline{t}} = X_{1+s}^{\overline{t}+s}\}$. The motivation for the proof technique is that the process **X** is strongly mixing, and so events which are at time points that are sufficiently far apart from one another are approximately independent, even if the process has memory. Thus, we may consider repetition at jumps of $\tau \equiv \tau(n)$ time points, for (slowly) increasing sequence $\tau(n)$, that is, the event

$$\{X_1 = X_{1+s}\} \cap \{X_{1+\tau} = X_{1+\tau+s}\} \cap \cdots.$$
(63)

Nonetheless, the probability of this event cannot be upper bounded as if all the symbols $\{X_i\}$ involved in this event are approximately independent, since for a general value of s, the index of the end symbol in the matching of a pair might be closer than τ to the start index of a different pair. For example, 1 + s might be close by less than τ time points from some $1 + j\tau$, $j \in \mathbb{N}_+$. In other words, the time indices of the symbols in the events defining (63) are intertwined, and not trivially separated. One technical challenge of the proof is to adequately dilute pair matching events from (63), in order to assure a separation of τ time indices between any two symbols.

The trade-off between the two types of error: Evidently, the optimal Bayesian trade-off is as follows: To make the type-I error probability a negligible $o(\frac{\log n}{n})$, the MAP detector rarely decides on overlaps shorter than $\sim \frac{\log n}{\mathcal{H}_1(\mathbf{X})}$, and in turn, when such short overlaps actually occur the detector err. This is the dominating error event. Now, suppose that we would like to further reduce the contribution of the type-I error probability, to make it decay faster than $o(\frac{\log n}{n})$. This can be achieved by modifying the detection rule (41) of Prop. 4 by replacing $n_{\ell} \sim n$ with n^{μ} for some $\mu > 1$. It can be shown that this reduces the decay rate of the type-I error probability to $o(\frac{\log n}{n^{\mu}})$, and increases the Bayesian error probability to $\sim \frac{\mu}{\mathcal{H}_1(\mathbf{X})} \frac{\log n}{n}$, that is, by a factor of μ . A short outline of the details appears in Appendix A. By contrast, the effect of reducing μ below 1 will make the type-I error probability dominating, by detecting short, unreliable, overlaps. When $\mu < 1$, a characterization of the dependence of the type-II contribution to the error probability as a function of the increased type-I error probability is more delicate, since we still need to assure that $p_{type-I}(\hat{T}) \sim \frac{\varphi_{\mu}}{\mathcal{H}_1(\mathbf{X})} \frac{\log n}{n}$ for some $\varphi_{\mu} < 1$ to be in the interesting regime of $P_{error}^* = \Theta(\frac{\log n}{n})$. ⁹ We thus refrain from pursuing the analysis of this case.

Interestingly, the trade-off involved in our upper bound on the Bayesian error probability does *not* balance between the contributions of the type-I and type-II error probabilities to the total error probability (while taking into account the prior probability). Indeed, the contribution of the type-I error probability is $o(\frac{\log n}{n})$ whereas that of the type-II error probability is $\Theta(\frac{\log n}{n})$. As explained in the introduction, this is because the setting here is, loosely speaking, should be described as a collection of $2\ell - 1$ binary detection problems.

We next discuss the various assumptions of Theorem 6.

The finite minus-infinity-order Rényi entropy rate assumption: This assumption implies a lower bound on the minimal error probability of the process. Specifically, for any $\epsilon > 0$, there exists $s_0(\epsilon)$ such that for all $s > s_0$ it holds that

$$\min_{\tilde{x}_1^s \in \text{supp}(P_{X_1^s})} P_{X_1^s}(\tilde{x}_1^s) \ge \exp_{|\mathcal{X}|} \left[-s[\mathcal{H}_{-\infty}(\mathbf{X}) - \epsilon] \right].$$
(64)

⁹Bounds on the type-I error probability of the form $o(\frac{\log n}{n^q})$ for q < 1 are most likely easier to derive, but are already rather large, and less interesting given that the prior probability of $T \neq 0$ is $\Theta(\frac{\log n}{n})$.

Thus, the minimal probability of a sequence of length s from the process decays exponentially, but not faster. Since the Rényi entropy is monotonically decreasing in its order, this guarantees that $\mathcal{H}_{\alpha}(\mathbf{X}) < \infty$ for any order $\alpha \in \mathbb{R}$. This assumption is rather mild, but of technical nature and so it is not obvious that it cannot be relaxed. Its origin is in the analysis of the type-I error probability, and can be explained as follows: In this analysis, T = 0 is assumed, and so the reads do not overlap. However, it is possible that they are close, or even adjacent, i.e., have start indices I(1) = 1 and $I(2) + \ell + 1$. When the process X has memory, such reads, e.g., $X(1) = X_1^{\ell}$ and $X(2) = X_{\ell+1}^{2\ell}$ may be highly correlated, even though T = 0. In those cases, the analysis of the type-I error, which is based on upper bounding the probability of matching events such as $X_{\ell-t+1}^{\ell}(1) = X_1^{\ell}(2)$, should be done under the true joint probability of the suffix/prefix of the reads. We circumvent this challenging aspect by upper bound the probability of the event $\{X_{\ell-t+1}^{\ell}(1) = X_1^t(2)\}$, with the probability of a slightly shorter match $\{X_{\ell-t+1+\tau}^{\ell}(1) = X_{\tau}^{t}(2)\}$ for some $\tau \equiv \tau(n)$. This results a distance of τ time points between the symbols defining this event. Now, if we choose $\tau(n)$ to increase slowly with n (it turns out that $\tau(n) = \left[\sqrt{\log n}\right]$ is a valid choice), then the strongly mixing property of X implies that the symbols in $X_{\ell-t+1+\tau}^{\ell}(1)$ are approximately independent from the symbols in $X_{\tau}^{t}(2)$, and this leads to tractable analysis. Nonetheless, the error event also includes the condition $\frac{1}{P_{X_{\ell}^t}(X_1^t(2))} \ge n_{\ell}$ but in order to agree with the reduced match length, we would like to modify this condition to $1/P_{X_{\tau}^{t}}(X_{\tau}^{t}(2)) \geq n_{\ell}$, that is, for a read shorter by τ symbols. The assumption on $\mathcal{H}_{-\infty}(\mathbf{X})$ guarantees that this modification has a mild effect on the probability, and thus this modification is inconsequential.

The moderately strongly mixing assumption: It is obvious that some type of mixing property is required for reliable overlap detection. Indeed, if the process may output long repetitions of the same symbol, of typical length $\Omega(\log n)$, then it is impossible to reliably detect the overlap in case the two reads belong to that segment in the process. Our assumption is fairly mild, because for a process to be strongly mixing, it is only required that $d(s) \to 0$ as $s \to \infty$. One common class of processes mixes with an *exponential* decay rate, that is $d(s) = \gamma^s$ for some $\gamma < 1$ (otherwise it is only sub-exponential), for which $\sum \gamma^s = \frac{1}{1-\gamma} < \infty$. However, our assumption is much more lenient because it suffices that $d(s) = o(\frac{1}{s})$ for $\sum_{s=0}^{\infty} d(s) < \infty$ to converge.

The aperiodicity assumption: The probability $\mathcal{R}(\mathbf{X}) = \mathbb{P}[X_1 = X_{1+s}]$ is the recurrence probability at time delay s. Now, if there exists an $s^* > 0$ such that $\mathbb{P}[X_1 = X_{1+s^*}] = 1$, then the process is periodic with a period s^* . An aperiodicity property is clearly also necessary for reliable overlap detection, because if the process has period s^* then it is impossible to distinguish between overlaps T and $T + ms^*$, for any $m \in \mathbb{N}_+$. We assume that the supremum over all s is bounded away from 1, but under the strong mixing assumption, it is satisfied naturally for all s sufficiently large. Indeed, let $p_{\min} := \min_{x \in \mathcal{X}} P_{X_1}(x)$, for which we may assume, w.l.o.g., that $p_{\min} > 0$ (otherwise we may remove the irrelevant symbol from the alphabet), and anyhow holds under the assumption $\mathcal{H}_{-\infty}(\mathbf{X}) < \infty$. It holds that $d(s) \to 0$ as $s \to \infty$ and so

$$\mathbb{P}[X_1 = X_{1+s}] = \sum_{x \in \mathcal{X}} \mathbb{P}[X_1 = x] \mathbb{P}[X_{1+s} = x \mid X_1 = x]$$
(65)

$$\leq \sum_{x \in \mathcal{X}} \mathbb{P}[X_1 = x] \left(\mathbb{P}[X_{1+s} = x] + d(s) \right) \tag{66}$$

$$\leq \left(1 + \frac{d(s)}{p_{\min}}\right) \sum_{x \in \mathcal{X}} P_{X_1}^2[X_1 = x]$$
(67)

$$= \left(1 + \frac{d(s)}{p_{\min}}\right) |\mathcal{X}|^{-H_2(P_{X_1})}$$
(68)

for all $s > s_0$, where s_0 is determined by $(\{d(s)\}, p_{\min}, H_2(P_{X_1}))$.

The parameters determining the convergence to asymptotic limits: The convergence of the upper bound to its asymptotic limit depends on the problem parameters: The normalized length of the read β , the minus-infinityorder Rényi entropy $\mathcal{H}_{-\infty}(\mathbf{X})$, and the related quantity p_{\min} , the decay rate of the strong mixing coefficient d(s) and on the recurrence probability $\mathcal{R}(\mathbf{X})$.

We conclude with two simple examples that illustrate the result of Theorem 6.

Example 7 (Memoryless sources). If **X** is memoryless process, then the assumptions of Theorem 6 trivially hold $(\mathcal{H}_{-\infty}(\mathbf{X}) = -\log p_{\min} < \infty, d(s) \equiv 0$ for all $s \in \mathbb{N}$, and $\mathcal{R}(\mathbf{X}) = |\mathcal{X}|^{-H_2(P_{X_1})} < 1$). Also, the entropy rate exists, and so

$$P_{\text{error}}^* \le 2\left[1 + o_n(1)\right] \cdot \left\{\beta \wedge \frac{1}{H_1(P_{X_1})}\right\} \cdot \frac{\log n}{n}.$$
(70)

Example 8 (First-order Markov Process (Markov chains)). If **X** is a first-order Markov process, which forms an irreducible and aperiodic Markov chain, then it has a unique stationary distribution (e.g., [18, Sec. 1.5]). Let us denote its transition matrix (Markov kernel) by

$$K(x, x') := \mathbb{P}\left[X_i = x' \mid X_{i-1} = x\right]$$
(71)

for any $x, x' \in \mathcal{X}$ and $i \in \mathbb{Z}$. As a simple and concrete example, let us focus on the case

$$K(x,x') = \begin{cases} 1 - (|\mathcal{X}| - 1)\epsilon, & x = x'\\ \epsilon, & x \neq x' \end{cases}$$
(72)

which has a uniform stationary distribution $P_{X_1}(x) = \frac{1}{|\mathcal{X}|}$ for all $x \in \mathcal{X}$.

We next discuss the various assumptions of Theorem 6. First, from the Markov property

$$\mathcal{H}_{-\infty}(\mathbf{X}) = \lim_{t \to \infty} \frac{1}{t} \max_{x_1^t \in \operatorname{supp}(P_{X_1^t})} \log \frac{1}{P_{X_1^t}(x_1^t)}$$
(73)

$$= \lim_{t \to \infty} \frac{1}{t} \max_{x_1^t \in \text{supp}(P_{X_1^t})} \log \frac{1}{P_{X_1}(x_1) \prod_{i=2}^t K(x_{i-1}, x_i)}$$
(74)

$$= \lim_{t \to \infty} \frac{1}{t - 1} \max_{x_1^t \in \text{supp}(P_{X_1^t})} \sum_{i=2}^t \log \frac{1}{K(x_{i-1}, x_i)}.$$
(75)

This value is upper bounded by the maximal possible sum of $\log \frac{1}{K(x_{i-1},x_i)}$ along a cycle (i.e., a path that starts and end at the same symbol) on the state space graph of the chain (for which each node is $x \in \mathcal{X}$). In our concrete example, it holds that

$$\mathcal{H}_{-\infty}(\mathbf{X}) = \begin{cases} -\log\left[\left(1 - (|\mathcal{X}| - 1)\epsilon\right)\right], & \epsilon \ge \frac{1}{|\mathcal{X}|} \\ -\log(\epsilon), & \epsilon < \frac{1}{|\mathcal{X}|} \end{cases}$$
(76)

which indeed satisfies the condition $\mathcal{H}_{-\infty}(\mathbf{X}) < \infty$. Second, for a given delay $s \in \mathbb{N}_+$ and length t, Markovity implies that

$$\begin{aligned} \left| \mathbb{P}[X_{i+s+1}^{i+s+t} \mid X_{-\infty}^{i+1}] - \mathbb{P}[X_{i+s+1}^{i+s+t}] \right| \\ &= \left| \mathbb{P}[X_{i+s+1}^{i+s+t} \mid X_{i+1}] - \mathbb{P}[X_{i+s+1}^{i+s+t}] \right| \end{aligned}$$
(77)

$$= |\mathbb{P}[X_{i+s+1} \mid X_{i+1}] - \mathbb{P}[X_{i+s+1}]| \cdot \prod_{j=2}^{t} \mathbb{P}[X_{i+s+j} \mid X_{i+s+j-1}]$$
(78)

$$\leq |\mathbb{P}[X_{i+s+1} \mid X_{i+1}] - \mathbb{P}[X_{i+s+1}]|$$
(79)

$$\leq \max_{x \in \mathcal{X}} \|K^s(x, \cdot) - P_{X_1}\|_{\mathrm{TV}},\tag{80}$$

where the latter is commonly take as the definition of the strong mixing coefficients of a Markov chain. In our concrete example, it is not difficult to verify (See details on Appendix A) that $d(s) \leq e^{-|\mathcal{X}| \epsilon \cdot s}$, which clearly implies that the chain is strongly mixing. Third, aperiodicity holds by assumption. Finally, the error probability is determined by the entropy rate, easily verified to be given by

$$\mathcal{H}_1(\mathbf{X}) = \frac{1}{|\mathcal{X}|} \log \frac{\epsilon}{1 - (|\mathcal{X}| - 1)\epsilon} + \log \frac{1}{\epsilon}.$$
(81)

We next turn to derive a lower bound, and prove that the bound of Theorem 6 is tight (under proper conditions).

Theorem 9. Consider the noiseless setting (Def. 1). Assume that for some $\eta > 0$, it holds that

$$d(s) \le \frac{1}{s^{\beta \mathcal{H}_{-\infty}(\mathbf{X}) + \eta}},\tag{82}$$

for all $s \in \mathbb{N}_+$, and that the entropy rate $\mathcal{H}_1(\mathbf{X})$ exists. Then

$$P_{\text{error}}^* \ge 2\left[1 + o_n(1)\right] \cdot \left\{\beta \wedge \frac{1}{\mathcal{H}_1(\mathbf{X})}\right\} \cdot \frac{\log n}{n}.$$
(83)

Theorem 9 thus establishes that the bound in Theorem 6 is tight.

The main ideas of the proof: In its core, the proof relies on the fact that the dominant error event is when $|T| \leq t^* \sim \frac{\log n}{\mathcal{H}_1(\mathbf{X})}$, in which the probability of the overlapping part of the read $X_1^T(2)$ is not sufficiently small to assure that this overlap will actually be detected, i.e., $1/P_{X_1^T(2)}(X_1^T(2)) \leq n_\ell$, and so $\hat{T} = 0$ is erroneously decided. To prove this rigorously, we lower bound the error probability of the MAP detector by the error probability of a *genie-aided detector*, whose operation can be described as follows, with some simplifications of technical details: If $T > t^* \sim \frac{\log n}{\mathcal{H}_1(\mathbf{X})}$, then the genie reveals the detector the true value of T, and so there is no error. If $0 < |T| \leq t^*$, the genie only reveals to the detector that the overlap is either T = 0 or $T = t^*$. If the overlap is T = 0, then the genie reveals to the detector that the true value is either T = 0 or some random T = t, chosen uniformly in $[t^*]$. This reduces the operation of the genie-aided detector to that of $[t^*]$ binary detectors. The average error probability of each of these binary detectors can be lower bounded similarly, so we may focus on one of them. As usual, this lower bound is based on comparing the likelihood ratio to the ratio of the prior probabilities. In turn, this requires evaluating both $f(y_1^\ell(1), y_1^\ell(2); t)$ and $f(y_1^\ell(1), y_1^\ell(2); 0)$. As we have seen in the proof of Prop. 4, both likelihoods are challenging to handle in case of memory. For $T = t \neq 0$, we utilize the asymptotic equipartition property (AEP) theorem of Shannon–McMillan–Breiman (e.g., [6, Th. 16.8.1]) to show that, with high probability, the likelihood

$$f(y_1^{\ell}(1), y_1^{\ell}(2); t) \le e^{o_n(1) \cdot \log n} \cdot \mathbb{1} \left[x_{\ell-t+1}^{\ell}(1) = x_1^t(2) \right] \cdot P_{X_1^{\ell}} \left(x_1^{\ell}(1) \right) \cdot P_{X_1^{\ell}} \left(x_1^{\ell}(2) \right) \cdot \Gamma_+(t), \tag{84}$$

where the r.h.s. is the likelihood given for a memoryless processes, or more generally, for any finite-order Markov chain, as developed in (42) (proof of Prop. 4), multiplied by a $e^{o_n(1) \cdot \log n}$ factor. For T = 0, we argued that for ergodic processes (46) the reads are approximately independent, and the likelihood is a multiplication of the likelihood of each read. Lemma 19 shows that under the stronger mixing conditions imposed by Theorem 9, the approximate relation (46) is actually an accurate asymptotic lower bound

$$f\left(y_{1}^{\ell}(1), y_{1}^{\ell}(2); 0\right) \ge \left[1 - o_{n}(1)\right] \cdot P_{X_{1}^{\ell}}\left(x_{1}^{\ell}(1)\right) \cdot P_{X_{1}^{\ell}}\left(x_{1}^{\ell}(2)\right).$$
(85)

With these asymptotic bounds, we can upper bound the likelihood ratio, and prove a lower bound on the error probability of this binary detector. It should be mentioned that this lower bound requires the almost sure convergence of $-\frac{1}{t} \log P_{X_1^t}(x_1^t)$ to the entropy rate $\mathcal{H}_1(\mathbf{X})$, as assured by the Shannon–McMillan–Breiman AEP, and thus also the existence of the entropy rate $\mathcal{H}_1(\mathbf{X})$.

The strongly mixing assumption: Compared to Theorem 6, which only assumed $d(s) = o(\frac{1}{s})$, the lower bound makes a slightly stronger assumption that d(s) is polynomially decaying with power strictly larger than $\beta \mathcal{H}_{-\infty}(\mathbf{X})$. Since for many processes the strong mixing coefficient decays much faster, this difference is rather minor. Nonetheless, pinpointing the exact dependence on the strong mixing coefficient remains an elusive open problem. We mention that the source of this assumption is Lemma 19, and there it is used to show that $d(\tau)$ is negligible compared to $P_{X_1^{\ell}}(x_1^{\ell}(1)) \cdot P_{X_1^{\ell}}(x_1^{\ell}(2))$ whenever $\tau \equiv \tau(n) = \frac{n}{\log n}$.

IV. OVERLAP DETECTION FOR MEMORYLESS PROCESSES AND NOISY MEMORYLESS READING KERNELS

In this section, we address the noisy setting of Definition 2. For this analysis, we make the following additional notation conventions. We denote by P_Y the Y-marginal of $P_X \otimes P_{Y|X}$, and let \tilde{Y} be such that $P_{\tilde{Y}|X} = P_{Y|X}$ and the Markov chain relation $Y - X - \tilde{Y}$ holds. We then denote by $P_{Y\tilde{Y}}$ the (Y, \tilde{Y}) -marginal, that is,

$$P_{Y\tilde{Y}}(y,\tilde{y}) = \sum_{x \in \mathcal{X}} P_X(x) \cdot P_{Y|X}(y \mid x) P_{Y|X}(\tilde{y} \mid x)$$
(86)

for any $(y, \tilde{y}) \in \mathcal{Y}^{\otimes 2}$ (hence $P_Y = P_{\tilde{Y}}$ holds). Furthermore, we denote the *likelihood ratio* between $y \in \mathcal{Y}$ and $\tilde{y} \in \mathcal{Y}$ as

$$\lambda(y,\tilde{y}) := \frac{P_{Y\tilde{Y}}(y,\tilde{y})}{P_{Y}(y)P_{\tilde{Y}}(\tilde{y})}.$$
(87)

We may note that

$$\lambda_{\max} := \max_{(y,\tilde{y})\in\mathcal{Y}^{\otimes 2}} \lambda(y,\tilde{y}) \in [1,\infty), \tag{88}$$

and, in addition, since $P_{Y\tilde{Y}} \ll P_Y \otimes P_Y$ then

$$\lambda(Y, \tilde{Y}) \ge \lambda_{\min} := \min\left\{\lambda(y, \tilde{y}) \colon (y, \tilde{y}) \in \operatorname{supp}(P_{Y\tilde{Y}})\right\} > 0$$
(89)

with probability 1.

A. The MAP Detection Rule

Our first step in deriving the MAP detection rule is to obtain an expression for the likelihood function.

Lemma 10. Consider the noisy setting (Def. 2). Then, the likelihood of overlap $t \in [\ell]$ given reads $y_1^{\ell}(1)$ and $y_1^{\ell}(2)$ is

$$f\left(y_{1}^{\ell}(1), y_{1}^{\ell}(2); t\right) \propto \Gamma_{+}(t) := \prod_{i=1}^{t} \lambda\left(y_{\ell-t+i}(1), y_{i}(2)\right),$$
(90)

and for $t \in -[\ell - 1]$

$$f\left(y_{1}^{\ell}(1), y_{1}^{\ell}(2); t\right) \propto \Gamma_{-}(t) := \prod_{i=1}^{t} \lambda\left(\left(y_{\ell-t+i}(2), y_{i}(1)\right),$$
(91)

and where we set $\Gamma_{\pm}(0) = 1$.

Inspecting the likelihood in (90), the function $t \mapsto \Gamma_+(t)$ is not necessarily monotonic *increasing* with t. Thus, unlike the noiseless case, any potential overlap $t \in [\ell]$ should be considered, as each has possibly nonzero likelihood, and an overlap is possible even if the overlapping noisy parts do not match. Nonetheless, the detector still has a rather simple form, as follows: **Proposition 11.** Consider the noisy setting (Def. 2). Let $\Gamma_{\pm}(t)$ be as defined in Lemma 10, and let the most likely positive and negative overlap be

$$t_{+} := \underset{t \in [\ell]}{\arg \max} \Gamma_{+}(t) \tag{92}$$

$$t_{-} := \arg\max_{t \in -[\ell-1]} \Gamma_{-}(t).$$
(93)

Then, the MAP detector is as in (41). Furthermore, let the minimum detectable overlap be denoted as

$$t_{\rm MDO} := \frac{\log(n_\ell)}{\log \lambda_{\rm max}}.$$
(94)

Then, $0 < |t| \le t_{MDO}$ will never be detected by the MAP detector.

The proof of Prop. 11 is similar to the proof of Prop. 4, and thus omitted.

B. Error Probability Analysis

Theorem 12. For the noisy setting (Def. 2)

$$P_{\text{error}}^* \le 2\left[1 + o_n(1)\right] \cdot \left\{\beta \land \frac{1}{I(Y;\tilde{Y})}\right\} \cdot \frac{\log n}{n}.$$
(95)

The dependence of the error on the mutual information: When the reading kernel is noisy, the mutual information $I(Y; \tilde{Y})$ replaces the role of the Shannon entropy rate (and indeed reduces to it when the reading kernel is noiseless). It emerges as the key measure because it is the expected log-likelihood ratio between the likelihood that the overlapping part of the reads are from separated locations in the sequence and the likelihood that this part are from the same location.

The main ideas of the proof: As in the proof of Theorem 6, we carefully analyze the probabilities of the various types of error. To simplify the proof and the description here, we only focus on errors from and to positive overlaps. Let us begin by considering the type-I error. Since $\hat{T} = t < t_{\text{MDO}}$ is never detected, we may take a union bound of the $\Theta(\log n)$ possibilities to err from T = 0 to $t_{\text{MDO}} < \hat{T} = t < \ell$. For each one of these possibilities, it can be shown that according to the detection rule of Prop. 11, the error probability is upper bounded by

$$\mathbb{P}\left[\sum_{i=1}^{t} \log \lambda\left(Y_{\ell-t+i}(1), Y_{i}(2)\right) \ge \log n_{\ell} \middle| T = 0\right].$$
(96)

This is a sum of t independent r.v.'s, where the expected value of each (conditioned on T = 0) is $I(Y; \tilde{Y})$. Hence, we expect that such probability will vanish only if $t > t^* \sim \frac{\log n}{I(Y; \tilde{Y})}$. Following the same arguments as in the proof of Theorem 6, the contribution of the type-II error probability is $\frac{1}{I(Y; \tilde{Y})} \frac{\log n}{n}$. However, to establish that the type-I error probability is negligible, we need to prove that it is $o(\frac{\log n}{n})$, and so taking into account the union bound over $\Theta(\log n)$ terms, we need to prove that the probability in (96) is $o(\frac{1}{n})$. To this end, if we deploy the standard Chernoff bound, then we only obtain $O(\frac{1}{n})$, which does not suffice for our purpose. Therefore, we deploy a stronger large-deviations bound, similar to that of Bahadur–Rao [2] [7, Th. 3.7.4.], which bounds this probability with the correct pre-exponent. In our case, this pre-exponent contributes a factor of $\frac{1}{\sqrt{t}} = \Theta(\frac{1}{\sqrt{\log n}})$ to the upper bound, which shows that the overall type-I error probability is $O(\frac{\sqrt{\log n}}{n}) = o(\frac{\log n}{n})$. The type-II error is similarly bounded using Chernoff's bounding technique. The upper bound on the error probability from T > 0 to T = 0 involves the analysis of a probability similar to (96), albeit with a reverse sign, and under the condition T = t > 0. Such probability is analyzed in Lemma 22. The error probability for a wrong detection, that is from T = t to a different $\hat{T} \neq 0$, is more complicated compared to the noiseless setting because there is also a possibility to underestimate the true overlap. As in the noiseless setting, the conditioning on T > 0 results dependencies between the terms involved in the sum of log-likelihoods. This is addressed in Lemma 23.

The trade-off between the two types of error: As in the noiseless setting, it can be verified from the proof of Theorem 12, that if n_{ℓ} is replaced in the detector by n^{μ} for some $\mu > 0$, then the type-I error would be $p_{\text{type-I}}(\hat{T}) = o(\frac{1}{n^{\mu}})$ whereas the contribution of the type-II error probability would increase by a factor of μ , that is, given by $[1 + o_n(1)] \frac{\mu}{I(Y;\tilde{Y})} \cdot \frac{\log n}{n}$.

The parameters determining the convergence to asymptotic limits: Naturally, here the convergence to the asymptotic limit depends on the source distribution P_X , the noise kernel $P_{Y|X}$, and the scaled read length β .

We conclude with a matching lower bound.

Theorem 13. For the noisy setting (Def. 2),

$$P_{\text{error}}^* \ge 2\left[1 - o_n(1)\right] \cdot \left\{\beta \wedge \frac{1}{I(Y;\tilde{Y})}\right\} \cdot \frac{\log n}{n}.$$
(97)

The proof of Theorem 13 follows similar lines to that of the lower bound in the noiseless reading kernel case.

V. CONCLUSION

We considered the problem of detecting the overlap between two short reads of length $\ell = \beta \log n$ drawn from two random locations in a long sequence of length *n*, taken from a process **X**. We considered the optimal Bayesian MAP detector, and provided tight asymptotic expressions for its error probability, both in the case the reading kernel is noiseless, allowing **X** to be stationary and ergodic, as well as in the case the reading kernel is noisy and memoryless, while restricting **X** to be memoryless. We next discuss a few possible future research directions.

First, it is plausible to conjecture that in the general case, in which both the process has memory and the reading kernel is noisy (perhaps also with memory), the error probability is given by $[1+o_n(1)] \cdot [\beta \wedge (1/\mathcal{I}(\mathbf{Y}, \tilde{\mathbf{Y}}))] \cdot \frac{\log n}{n}$, where, here, $\mathcal{I}(\mathbf{Y}, \tilde{\mathbf{Y}})$ is the mutual-information rate between two reads satisfying $\mathbf{Y} - \mathbf{X} - \tilde{\mathbf{Y}}$ (that is, two independent noisy observations $\mathbf{Y}, \tilde{\mathbf{Y}}$ of the same process \mathbf{X}). One of the challenges in proving such a result, beyond the additional anticipated technical effort, is that even if \mathbf{X} is a first-order Markov process and the

reading kernel is memoryless, each read is a *hidden* Markov process, for which the likelihood more complicated than the cases we considered. In a similar spirit, it would be interesting to consider noisy reading channels which introduce synchronization errors, such as deletions and insertions, and to characterize the error probability of the optimal detector.

Second, it would be interesting to utilize techniques similar to the ones used here in order to derive the fundamental limits of the perfect reconstruction of the set of contigs (islands in the parlance of [16]) from reads, whenever the reads do not fully cover the sequence. This is challenging, because without any prior assumption on the coverage, there are $\Theta(m^2 \cdot \frac{\log n}{n})$ pairs of reads on the average that only have a short overlap, and the detection of this overlap is likely to either not to be detected, or erroneously detected. The problem is even more challenging to analyze in the noisy setting, because in this case multiple reads of the same symbol may improve the de-noising of that symbol, and this must be taken into account in the analysis of the optimal detector.

Third, following [30], another open problem is related to detection of the alignment based on *sketches* of the read. In this setting, each of the reads $Y_1^{\ell}(1)$ and $Y_1^{\ell}(2)$ is separately compressed, or sketched, to $Z_1^B(1)$ and $Z_1^B(2)$. The detector is then required to detect the correct overlap based on the sketches $(Z_1^B(1), Z_1^B(2))$. The question is what is the optimal error probability, where here the sketching operation can also be optimized. Clearly, for B = 0 the error probability is $\beta \frac{\log n}{n}$, whereas for $B \ge H(Y_1^{\ell})$ the error probability, is as in our setting. The exact dependence of the Bayesian error probability on B is interesting to characterize.

Fourth, a variation of our problem may address soft-detection, in which the detector outputs the posterior probability of the overlap, and the performance is measure by the logarithmic loss functions, thus given by $H(T \mid Y_1^{\ell}(1), Y_1^{\ell}(2))$. It is then of interest to tightly characterize the asymptotic value of this equivocation. It would also be interesting to embed the posterior of the overlap into assembly algorithms that merge reads that approximately overlap [13], [31].

Fifth, following [21], [36], we may consider the overlap detection problem in case that there is a *reference sequence* that is correlated with the main sequence, and aids the detection task. It is then of interest to characterize how the asymptotic error probability improves as a function of the amount of dependence between the sequence and the reference sequence.

APPENDIX A

PROOFS AND DETAILS FOR SECTION III

A. Proof of Theorem 6

We begin with the following lemma, which states bounds that will be instrumental in the analysis of the probability of error events from T = 0 to $\hat{T} \neq 0$ (the bound (A.1)), and from $T \neq 0$ to $\hat{T} = 0$ (the bound (A.2)).

Lemma 14. Let $\mathbf{X} := \{X_i\}_{i \in \mathbb{Z}}$ be a stationary ergodic process for which $\mathcal{H}_1(\mathbf{X}) < \infty$. Let $\rho \in (0, 1)$ be given and $\{\tau(n)\}_{n \in \mathbb{N}}$ be a sequence of non-negative integers such that $\tau(n) = \omega_n(1)$ and $\tau(n) = o(\log n)$. Then, for any $\eta > 0$

$$\max_{t > \frac{(1+\eta)}{\mathcal{H}_1(\mathbf{X})} \log n} \sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}^2(x_1^t) \cdot \mathbb{1}\left[\frac{1}{P_{X_1^t}(x^t)} \ge \rho^{\tau - 1} \cdot n_\ell\right] = o\left(\frac{1}{n}\right),\tag{A.1}$$

and

$$\max_{t > \frac{(1+\eta)}{\mathcal{H}_1(\mathbf{X})} \log n} \sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}(x_1^t) \cdot \mathbb{1}\left[\frac{1}{P_{X_1^t}(x^t)} \le n_\ell\right] = o_n(1).$$
(A.2)

The constants involved in the asymptotic expressions depend on η .

Proof: Let any $t \ge \frac{(1+\eta)\log n}{\mathcal{H}_1(\mathbf{X})}$ be given. We first prove (A.1). As $n_\ell = n - (2\ell - 1)$ and $\tau \equiv \tau(n) = \omega_n(1)$, it holds that for all n sufficiently large (depending only on (ρ, β)) that

$$\rho^{\tau-1} \cdot n_{\ell} = \rho^{\tau-1} \cdot (n - (2\ell - 1)) \tag{A.3}$$

$$\geq \rho^{\tau-1} \left(1 - \frac{2\beta \log n}{n} \right) \cdot n \tag{A.4}$$

$$\geq \left(\frac{\rho}{2}\right)^{\tau} \cdot n. \tag{A.5}$$

In addition, since $\mathcal{H}_{2-r}(\mathbf{X}) \uparrow \mathcal{H}_1(\mathbf{X})$ as $r \uparrow 1$ (monotonicity of Rényi entropy rate [26]), there exists $r_0(\eta) \in (0,1)$ and $\epsilon_0(\eta) > 0$ for which

$$\frac{\left(1+\eta\right)\left(\mathcal{H}_{2-r_{0}}(\mathbf{X})-\epsilon_{0}\right)}{\mathcal{H}_{1}(\mathbf{X})}>1,\tag{A.6}$$

such that $r_0(\eta) \to 1$ and $\epsilon_0(\eta) \to 0$ as $\eta \to 0$. So, for all n sufficiently large

$$\sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}^2(x_1^t) \cdot \mathbb{1}\left[\frac{1}{P_{X_1^t}(x^t)} \ge \rho^{\tau - 1} \cdot n_\ell\right]$$
(A.7)

$$\stackrel{(a)}{\leq} \sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}^2(x^t) \cdot \mathbb{1}\left[\frac{1}{P_{X_1^t}(x^t)} \ge \left(\frac{\rho}{2}\right)^{\tau} \cdot n\right]$$
(A.8)

$$\stackrel{(b)}{=} \sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}^2(x^t) \cdot \mathbb{1}\left[\log P_{X_1^t}(x^t) \le -(1+\kappa)\log n\right]$$
(A.9)

$$\leq \sum_{x_{1}^{t} \in \mathcal{X}^{\otimes t}} P_{X_{1}^{t}}^{2}(x^{t}) \cdot \min_{r \geq 0} e^{r \left[-(1+\kappa) \log n - \log P_{X_{1}^{t}}(x^{t}) \right]}$$
(A.10)

$$\leq \frac{1}{n^{(1+\kappa)r_0}} \sum_{x^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}^{2-r_0}(x^t)$$
(A.11)

$$=\frac{1}{n^{(1+\kappa)r_0}}|\mathcal{X}|^{-(1-r_0)H_{2-r_0}(P_{X_1^t})}$$
(A.12)

$$= \exp_{n} \left(-(1+\kappa)r_{0} - \frac{(1-r_{0})H_{2-r_{0}}(P_{X_{1}^{t}})}{\log n} \right)$$
(A.13)

$$\stackrel{(d)}{\leq} \exp_n\left(-(1+\kappa)r_0 - (1-r_0)\frac{(1+\eta)}{\mathcal{H}_1(\mathbf{X})}\left(\mathcal{H}_{2-r_0}(\mathbf{X}) - \epsilon_0\right)\right) \tag{A.14}$$

$$\stackrel{(e)}{=} o\left(\frac{1}{n}\right),\tag{A.15}$$

where (a) follows from (A.5), (b) follows by denoting

$$\kappa \equiv \kappa(n) := \frac{\tau(n)}{\log n} \log \frac{\rho}{2},\tag{A.16}$$

(c) follows by $\mathbb{1}[a > 0] \le \min_{r \ge 0} e^{ra}$, and (d) holds for all n sufficiently large, since for $\epsilon_0 > 0$ there exists $n_0(\epsilon_0)$ such that for all $n \ge n_0$

$$H_{2-r_0}(P_{X_1^t}) \ge t(\mathcal{H}_{2-r_0}(\mathbf{X}) - \epsilon) \ge \frac{(1+\eta)}{\mathcal{H}_1(\mathbf{X})} \left(\mathcal{H}_{2-r_0}(\mathbf{X}) - \epsilon_0\right) \log(n).$$
(A.17)

Finally, (e) follows since (A.6) and $\kappa \equiv \kappa(n) = o_n(1)$ (which holds by the assumption $\tau_n = o(\log n)$) imply that

$$(1+\kappa(n)) r_0 + (1-r_0) \frac{(1+\eta)}{\mathcal{H}_1(\mathbf{X})} \left(\underline{\mathcal{H}}_{2-r_0}(\mathbf{X}) - \epsilon_0 \right) > 1$$
(A.18)

for all n sufficiently large (which depends on $(\rho, \{\tau(n)\})$). Since these constants do not depend on t, the result (A.1) is obtained.

We turn to prove (A.2). For any $r \in (0,1)$ and $\epsilon > 0$, there exists $n_0(\epsilon)$ such that for all $n \ge n_0(\epsilon)$

$$\sum_{\substack{x_1^t \in \mathcal{X}^{\otimes t} \\ x_1^t \in \mathcal{X}^{\otimes t}}} P_{X_1^t}(x_1^t) \cdot \mathbb{1}\left[\frac{1}{P_{X_1^t}(x^t)} \le n_\ell\right]$$
$$= \mathbb{P}\left[P_{X_1^t}^{1-r}(X_1^t) \ge \frac{1}{n_\ell^{1-r}}\right]$$
(A.19)

$$\stackrel{(a)}{\leq} n^{1-r} \cdot \mathbb{E}\left[P_{X_1^t}^{1-r}\left(X_1^t(1)\right)\right] \tag{A.20}$$

$$= n^{1-r} \sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}^{2-r}(x_1^t)$$
(A.21)

$$\leq n_{\ell}^{1-r} \cdot \exp_{|\mathcal{X}|} \left[-(1-r)t(\mathcal{H}_{2-r}(\mathbf{X}) - \epsilon) \right]$$
(A.22)

$$= \exp_{|\mathcal{X}|} \left[-(1-r) \left((1+\eta) \left(\frac{\mathcal{H}_{2-r}(\mathbf{X}) - \epsilon}{\mathcal{H}_1(\mathbf{X})} \right) - 1 \right) \log n \right]$$
(A.23)

$$\stackrel{(b)}{=} o_n(1), \tag{A.24}$$

where (a) follows from Markov's inequality and $n_{\ell} \leq n$, and (b) follows from (A.6), by setting $r = r_0(\eta)$ and $\epsilon = \epsilon_0(\eta)$.

We continue with the following lemma, which will be used to analyze error events from a non-zero overlap T = t to a longer overlap $\hat{T} > t$.

Lemma 15. Let $\mathbf{X} := \{X_i\}_{i \in \mathbb{Z}}$ be a stationary strongly mixing process with mixing coefficients $\{d(s)\}_{s \in \mathbb{N}}$,

which satisfies $\sum_{s=0}^{\infty} d(s) < \infty$, and is also aperiodic in the sense that

$$\mathcal{R}(\mathbf{X}) := \sup_{s \in \mathbb{N}_+} \mathbb{P}\left[X_1 = X_{1+s}\right] < 1.$$
(A.25)

Then,

$$\sup_{1 \le s \le \ell} \max_{\substack{(1+\eta)\\ \mathcal{H}_1(\mathbf{x})}} \max_{\log n \le t \le \beta \log n} \mathbb{P}\left[X_1^t = X_{1+s}^{t+s}\right] = o\left(\frac{1}{\log n}\right).$$
(A.26)

The constants depend on $p_{\min} := \min_{x \in \text{supp}(\mathcal{X})} P_{X_1}(x)$.

To prove Lemma 15, we will be assisted by the following lemma. Recall first Definition 5 of the strong mixing coefficient, which compares between the probability of an event and its probability conditioned on a distant past event. The next lemma upper bounds a probability conditioned on both past and *future* events, which are sufficiently distant apart.

Lemma 16. Let $\mathbf{X} := \{X_i\}_{i \in \mathbb{Z}}$ be a stationary strongly mixing process with strong mixing coefficients $\{d(s)\}_{s \in \mathbb{N}}$. Let $\underline{r}, \overline{r} \in \mathbb{N}_+$ and let

$$t_{-\underline{r}} < t_{-\underline{r}+1} < \dots t_{-1} < t_0 < t_1 < \dots < t_{\overline{r}}$$
(A.27)

be a given sequence of $\underline{r} + \overline{r} + 1$ times points. Also let $p_{\min} := \min_{x \in \text{supp}(\mathcal{X})} P_{X_1}(x) > 0$ and

$$d_{\min} := \min_{0 \le i \le \overline{r} - 1} |t_{i+1} - t_i| \wedge \min_{0 \le i \le \underline{r} - 1} |t_{-i} - t_{i-1}|, \qquad (A.28)$$

as well as

$$\overline{d} := \frac{d_{\min}}{p_{\min}}.$$
(A.29)

If $\overline{d} \leq \frac{1}{2}$ then

$$\mathbb{P}\left[X_{t_0} \mid X_{-t_1}, \dots, X_{-t_{\underline{r}}}, X_{t_1}, \dots, X_{t_{\overline{r}}}\right] \le \mathbb{P}[X_{t_0}] \cdot e^{2\overline{d}(\overline{r}+1)}.$$
(A.30)

Proof: To ease the notation, let us denote $V_i := X_{t_i}$. Then,

$$\mathbb{P}\left[V_0 \mid V_{-\underline{r}}, \dots, V_{-1}, V_1, \dots, V_{\overline{r}}\right] \\
\stackrel{(a)}{=} \frac{\mathbb{P}\left[V_{-\underline{r}}, \dots, V_{-1}, V_0, V_1, \dots, V_{\overline{r}}\right]}{\mathbb{P}\left[V_{-\underline{r}}, \dots, V_{-1}, V_1, \dots, V_{\overline{r}}\right]}$$
(A.31)

$$\stackrel{(b)}{=} \frac{\mathbb{P}\left[V_{-\underline{r}}, \dots, V_{-1}\right] \mathbb{P}\left[V_0 \mid V_{-\underline{r}}, \dots, V_{-1}\right] \mathbb{P}\left[V_1, \dots, V_{\overline{r}} \mid V_0, V_{-\underline{r}}, \dots, V_{-1}\right]}{\mathbb{P}\left[V_{-r}, \dots, V_{-1}\right] \mathbb{P}\left[V_1, \dots, V_{\overline{r}} \mid V_{-r}, \dots, V_{-1}\right]}$$
(A.32)

$$=\frac{\mathbb{P}\left[V_{0}\mid V_{-\underline{r}},\dots,V_{-1}\right]\mathbb{P}\left[V_{1},\dots,V_{\overline{r}}\mid V_{0},V_{-\underline{r}},\dots,V_{-1}\right]}{\mathbb{P}\left[V_{1},\dots,V_{\overline{r}}\mid V_{-\underline{r}},\dots,V_{-1}\right]}$$
(A.33)

$$= \mathbb{P}\left[V_{0} \mid V_{-\underline{r}}, \dots, V_{-1}\right] \cdot \prod_{i=1}^{\overline{r}} \frac{\mathbb{P}\left[V_{i} \mid V_{i-1}, \dots, V_{1}, V_{0}, V_{-\underline{r}}, \dots, V_{-1}\right]}{\mathbb{P}\left[V_{i} \mid V_{i-1}, \dots, V_{1}, V_{-\underline{r}}, \dots, V_{-1}\right]}$$
(A.34)

$$\stackrel{(c)}{\leq} \left(\mathbb{P}[V_0] + d_{\min}\right) \cdot \prod_{i=1}^{\overline{r}} \left(\frac{\mathbb{P}[V_i] + d_{\min}}{\mathbb{P}[V_i] - d_{\min}}\right) \tag{A.35}$$

$$\stackrel{(d)}{\leq} \mathbb{P}[V_0]\left(1+\overline{d}\right) \cdot \prod_{i=1}^r \left(1+2\overline{d}\right) \tag{A.36}$$

$$\leq \mathbb{P}[V_0] \left(1 + 2\overline{d}\right)^{\overline{r}+1} \tag{A.37}$$

$$\stackrel{(e)}{\leq} \mathbb{P}[V_0] \cdot e^{2\overline{d}(\overline{r}+1)},\tag{A.38}$$

where (a) and (b) follow from Bayes rule, (c) follows from the definition of the strong mixing coefficient d(s)and its assumed monotonicity, as well as the assumption $\mathbb{P}[V_i] \ge p_{\min} \ge 2d_{\min}$, (d) follows since for $0 < a \le \frac{1}{2}$ it holds that $\frac{1}{1-a} \le 1+2a$ and using this for $a = \frac{d_{\min}}{\mathbb{P}[V_1]} \le \frac{1}{2}$, and (e) from $1 + a \le e^a$ for $a \in \mathbb{R}$.

Using Lemma 16 we may now prove Lemma 15.

Proof of Lemma 15: Let arbitrary $t \ge t^*(n) = \frac{(1+\eta)}{\mathcal{H}_1(\mathbf{X})} \log n$ and $1 \le s \le \ell$ be given. We let $p_{\min} := \min_{x \in \text{supp}(\mathcal{X})} P_{X_1}(x)$. Since $\sum_{s=0}^{\infty} d(s) < \infty$ it holds that $d(s) = O(\frac{1}{s \cdot \log s})$. Thus, there must exists $\tau \equiv \tau(n) \in \mathbb{N}_+$ which satisfies that

$$k \equiv k(n) = \left\lfloor \frac{t^*(n) + 1}{\tau(n)} \right\rfloor = \omega(\log \log n), \tag{A.39}$$

and

$$d(\tau(n)) \cdot k(n) = o_n(1) \tag{A.40}$$

(as an extreme illustrative example, if $d(s) = \frac{1}{s \cdot \log s}$ then we can take $\tau(n) = \frac{\log n}{(\log \log n)^2}$ so that

$$d(\tau(n)) \cdot k(n) = O\left(\frac{1}{\tau(n) \cdot \log \tau(n)} \cdot \frac{\log n}{\tau(n)}\right)$$
(A.41)

$$= O\left(\frac{(\log \log n)^3}{\log n}\right) \tag{A.42}$$

and $k(n) = \Omega((\log \log n))^2$). Assume further for simplicity that k is even, otherwise we may substitute k by k-1.

We consider two cases. First, we consider the case $1 \le s \le 4\tau$. The idea is to focus on a subset of matching pairs from $X_1^t = X_{1+s}^{t+s}$, which are at sufficiently distant time indices. We have

$$\mathbb{P}\left[X_{1}^{t} = X_{1+s}^{t+s}\right] \overset{(a)}{\leq} \mathbb{P}\left[\{X_{1} = X_{1+s}\} \cap \{X_{1+8\tau} = X_{1+8\tau+s}\} \cap \{X_{1+16\tau} = X_{1+16\tau+s}\} \cap \dots \cap \{X_{1+k\tau} = X_{1+k\tau+s}\}\right] \quad (A.43)$$

$$\mathbb{P}\left[\{X_{1} = X_{1+s}\} \cap \{X_{1+8\tau} = X_{1+8\tau+s}\} \cap \{X_{1+16\tau} = X_{1+16\tau+s}\} \cap \dots \cap \{X_{1+k\tau} = X_{1+k\tau+s}\}\right] \quad (A.43)$$

$$= \mathbb{P}\left[X_1 = X_{1+s}\right] \cdot \prod_{i=1}^{n/s} \mathbb{P}\left[X_{1+8i\tau} = X_{1+8i\tau+s} \middle| \bigcap_{j=0}^{r-1} \{X_{1+8j\tau} = X_{1+8j\tau+s}\}\right]$$
(A.44)

$$\leq \prod_{i=1}^{k/8} \mathbb{P} \left[X_{1+8i\tau} = X_{1+8i\tau+s} \left| \bigcap_{j=0}^{i-1} \{ X_{1+8j\tau} = X_{1+8j\tau+s} \} \right]$$
(A.45)

$$\stackrel{(b)}{\leq} \prod_{i=1}^{k/8} \left(\mathbb{P}\left[X_{1+8i\tau} = X_{1+8i\tau+s} \right] + d(8\tau - s) \right) \tag{A.46}$$

$$\stackrel{(c)}{\leq} \prod^{k/8} \left(\mathbb{P} \left[X_{1+8i\tau} = X_{1+8i\tau+s} \right] + d(4\tau) \right) \tag{A.47}$$

$$\stackrel{i=1}{\stackrel{(d)}{=}} \left(\mathbb{P} \left[X_1 = X_{1+s} \right] + d(4\tau) \right)^{k/8} \tag{A.48}$$

$$\stackrel{(e)}{\leq} \left(\mathbb{P}\left[X_1 = X_{1+s} \right] + o_n(1) \right)^{k/8} \tag{A.49}$$

$$\stackrel{(f)}{=} o\left(\frac{1}{\log n}\right),\tag{A.50}$$

where (a) follows by only taking into account symbol pair matching events at distant 8τ apart, (b) follows from the definition of the strong mixing coefficient, and since the events $\{X_{1+2i\tau} = X_{1+2i\tau+s}\}$ and $\bigcap_{j=0}^{i-1} \{X_{1+2j\tau} = X_{1+2j\tau+s}\}$ are $8\tau - s$ time points apart (obtained at the symbols and $X_{1+2(i-1)\tau+s}$ and $X_{1+2i\tau}$, (c) follows from the assumed monotonicity of d(s) w.l.o.g., and since $s \leq 4\tau$, (d) follows from stationarity, (e) follows since $d(\tau(n)) = o_n(1)$, and (f) follows since by assumption $\sup_{s \in \mathbb{N}_+} \mathbb{P}[X_1 = X_{1+s}] < 1$ and as $k \equiv k(n) = \omega(\log \log n)$.

Second, we consider the case $4\tau \le s \le \ell$. Let $m = \lfloor \frac{s}{\tau} \rfloor$, where $m \ge 4$. The idea is again to focus on a subset of matching pairs from the event $\{X_1^t = X_{1+s}^{t+s}\}$. In principle, as in the previous case, we would like to choose them to be τ apart (or a multiple of τ), to allow for approximate independence between these events. A natural candidate for a loosening of the event $\overline{\mathcal{E}} := \{X_1^t = X_{1+s}^{t+s}\}$ is the subset

$$\mathcal{E} := \bigcap_{i=0}^{k} \mathcal{E}_i \tag{A.51}$$

where

$$\mathcal{E}_i := \{ X_{1+i\tau} = X_{1+i\tau+s} \} \,. \tag{A.52}$$

However, for the case $4\tau \le s \le \ell$, this does not guarantee that the symbols involved in the definition of \mathcal{E} are τ apart. Indeed, suppose, for example, that $m_0 = \frac{s}{\tau}$ is integer. Then, the end symbol in the matching of \mathcal{E}_0 , i.e., X_{1+s} is equal to the start symbol in the matching of \mathcal{E}_{m_0} , to wit, $X_{1+m_0\tau}$, and thus they are definitely not τ symbols apart. To circumvent this, we will modify the event (A.51) by diluting some of the events \mathcal{E}_i , and consider the event

$$\tilde{\mathcal{E}} := \bigcap_{i \in \mathcal{I}} \{ X_{1+i\tau} = X_{1+i\tau+s} \} \subset \mathcal{E},$$
(A.53)

where $\mathcal{I} \subset \{0\} \cup [k]$ is a retained set of indices. Specifically, we would like to guarantee *any* two symbols that are a part of the definition the even $\tilde{\mathcal{E}}$ to wit, $\{X_{1+i\tau}, X_{1+i\tau+s}\}_{i\in\mathcal{I}}$ are at least τ time points apart. This can be seen to be achieved by the following choice. First, we include the events $\{\mathcal{E}_j\}_{j\in\{0,1,\dots,m-1\}}$, but exclude \mathcal{E}_m and \mathcal{E}_{m+1} , as their start symbol indices, i.e., $1 + m\tau$, $1 + (m+1)\tau$, resp., may be too close to the final




Figure 3. Time indices of pair matching of the modified event (A.53). Here m = 4, and the symbols $X_{1+4\tau}$ and $X_{1+5\tau}$ are less than τ time points from X_{1+s} , thus their corresponding events are removed from the event in (A.51) (appearing in dash points in the figure). Symbols with time indices that are less than τ time points are marked by a circle of a similar color.

symbol index 1 + s of \mathcal{E}_0 . For a similar reason, we exclude \mathcal{E}_{m+2} (closeness to \mathcal{E}_1), and so on. The index of the smallest event \mathcal{E}_i that can be included while satisfying the requirements, to wit, whose start symbol index is assured to be more than τ apart from the final symbol index of \mathcal{E}_{m-1} , is \mathcal{E}_{2m+1} . Similarly to the inclusion of $\{\mathcal{E}_i\}_{\{0,1,\dots,m-1\}}$, we may also include all the m-2 following events, that is, $\{\mathcal{E}_j\}_{j\in\{2m+2,\dots,3m-1\}}$. See Fig. 3 for an illustration. Generalizing the above considerations, traversing from i = 0 to i = k, we allow inclusion of a sequence of at least m-1 events, followed by exclusion of the following m+1 events, inclusion of at least m-1 events, and so on. The cardinality of the retained set of indices is thus at least $|\mathcal{I}| \ge \frac{k}{3}$. With this in mind, we bound

$$\mathbb{P}\left[X_{1}^{t} = X_{1+s}^{t+s}\right]$$

$$\stackrel{(a)}{\leq} \mathbb{P}\left[\bigcap_{i\in\mathcal{I}} \left\{X_{1+i\tau} = X_{1+i\tau+s}\right\}\right]$$
(A.54)

$$\stackrel{(b)}{=} \prod_{i \in \mathcal{I}} \mathbb{P} \left[X_{1+i\tau} = X_{1+i\tau+s} \middle| \bigcap_{j \in \mathcal{I}: j \le i-1} \{ X_{1+j\tau} = X_{1+j\tau+s} \} \right]$$

$$\stackrel{(c)}{=} \prod \sum_{i \in \mathcal{I}} \mathbb{P} \left[X_{1+i\tau+s} = x_i \middle| \bigcap_{j \in \mathcal{I}: j \le i-1} \{ X_{1+j\tau} = X_{1+j\tau+s} \} \right] \times$$

$$(A.55)$$

$$\stackrel{c}{=} \prod_{i \in \mathcal{I}} \sum_{x \in \mathcal{X}} \mathbb{P} \left[X_{1+i\tau+s} = x \left| \bigcap_{j \in \mathcal{I}: j \leq i-1} \{ X_{1+j\tau} = X_{1+j\tau+s} \} \right] \times \mathbb{P} \left[X_{1+i\tau} = x \left| \bigcap_{j \in \mathcal{I}: j \leq i-1} \{ X_{1+j\tau} = X_{1+j\tau+s} \} \cap \{ X_{1+i\tau+s} = x \} \right]$$
(A.56)

$$\stackrel{(d)}{\leq} \prod_{i \in \mathcal{I}} \sum_{x \in \mathcal{X}} \mathbb{P}\left[X_{1+i\tau+s} = x\right] \mathbb{P}\left[X_{1+i\tau} = x\right] e^{\frac{d(\tau)}{p_{\min}}(2k+1)}$$
(A.57)

$$\stackrel{(e)}{=} \prod_{i \in \mathcal{I}} \exp_{|\mathcal{X}|} \left[\log(e) \cdot \frac{d(\tau)}{p_{\min}} (2k+1) - H_2(P_{X_1}) \right]$$
(A.58)

$$\leq \exp_{|\mathcal{X}|} \left[-\frac{k}{4} \left(H_2(P_{X_1}) - \log(e) \cdot \frac{d(\tau)}{p_{\min}} (2k+1) \right) \right]$$
(A.59)

$$\stackrel{(g)}{\leq} \exp_{|\mathcal{X}|} \left[-\frac{k}{8} \cdot H_2(P_{X_1}) \right] \tag{A.60}$$

$$\stackrel{(g)}{=} o\left(\frac{1}{\log n}\right),\tag{A.61}$$

where (a) follows by considering the probability $\mathcal{E} \subset \overline{\mathcal{E}}$ in (A.53), (b) follows from Bayes rule, (c) follows again from Bayes rule, (d) follows due to the following bounds on the two probabilities: The first probability is bounded as

$$\mathbb{P}\left[X_{1+i\tau+s} = x \left| \bigcap_{j \in \mathcal{I}: j \le i-1} \{X_{1+j\tau} = X_{1+j\tau+s}\}\right] \le \mathbb{P}\left[X_{1+i\tau+s} = x\right] + d(\tau)$$
(A.62)

$$\leq \mathbb{P}\left[X_{1+i\tau+s} = x\right] \left[1 + \frac{d(\tau)}{p_{\min}}\right] \tag{A.63}$$

$$\leq \mathbb{P}\left[X_{1+i\tau+s} = x\right] \cdot e^{\frac{a(\tau)}{p_{\min}}},\tag{A.64}$$

since the intersection $\bigcap_{j \in \mathcal{I}: j \leq i-1} \{X_{1+j\tau} = X_{1+j\tau+s}\} \in \sigma(X_{-\infty}^{1+(i-1)\tau+s})$, that is, it is at distance of τ times points from $X_{1+i\tau+s}$. The second probability is bounded as

$$\mathbb{P}\left[X_{1+i\tau} = x \left| \bigcap_{j \in \mathcal{I}: j \leq i-1} \{X_{1+j\tau} = X_{1+j\tau+s}\} \cap \{X_{1+i\tau+s} = x\} \right] \\ \leq \mathbb{P}\left[X_{1+i\tau} = x\right] e^{2\frac{d(\tau)}{p_{\min}}|\mathcal{I}|}$$
(A.65)

$$\leq \mathbb{P}\left[X_{1+i\tau} = x\right] e^{2\frac{d(\tau)}{p_{\min}}k},\tag{A.66}$$

using Lemma 16, as the conditioning event involve symbols which are at least τ time points apart from $X_{1+i\tau}$ (either to the past or the future), (e) follows from stationarity

$$\sum_{x \in \mathcal{X}} \mathbb{P}\left[X_{1+i\tau+s} = x\right] \mathbb{P}\left[X_{1+i\tau} = x\right] = \sum_{x \in \mathcal{X}} P_{X_1}^2(x) = |\mathcal{X}|^{-H_2(P_{X_1})},\tag{A.67}$$

and $1 + a \le e^a = |\mathcal{X}|^{\log(e) \cdot a}$, (f) follows since there are at least $|\mathcal{I}| \ge \frac{k}{3}$ terms in the product, and (g) follows since $d(\tau(n)) \cdot k(n) = o_n(1)$ while $k(n) = \omega(\log \log n)$.

Combining the two cases (A.50) and (A.61), and noting that the bounds do not depend on t and s, proves the claim of the lemma.

With Lemma 14 and Lemma 15 at hand, may now prove Theorem 6.

Proof of Theorem 6: We first consider the case that $\beta \leq \frac{1}{\mathcal{H}_1(\mathbf{X})}$, which is simple to analyze. Indeed,

consider a trivial detector that always outputs T = 0. This detector will err if $T \neq 0$, which happens with probability $\frac{2\ell-1}{n} = \frac{2\beta \log n - 1}{n}$. Hence,

$$P_{\text{error}}^* \le \frac{2\ell}{n} = 2\beta \cdot \frac{\log n}{n}.$$
(A.68)

The rest of the proof is devoted to the case that $\beta > \frac{1}{\mathcal{H}_1(\mathbf{X})}$ and to prove the bound

$$P_{\text{error}}^* \le [1 + o_n(1)] \cdot \frac{2}{\mathcal{H}_1(\mathbf{X})} \cdot \frac{\log n}{n}, \tag{A.69}$$

under the assumptions of the theorem, thus establishing the upper bound on the error probability in (53). To this end, let

$$t^* := \frac{\log n}{\mathcal{H}_1(\mathbf{X})},\tag{A.70}$$

let $\eta > 0$ be given and set

$$t_{\eta}^* := \left\lceil (1+2\eta)t^* \right\rceil = \left\lceil (1+2\eta)\frac{\log n}{\mathcal{H}_1(\mathbf{X})} \right\rceil,\tag{A.71}$$

where η is also small enough so that $t_{\eta}^* \leq \ell$ (we will eventually take $\eta \downarrow 0$). The idea is to use the detector of Prop. 4, which is possibly sub-optimal when **X** has memory. To slightly simplify the analysis, we modify this detector further, and analyze the detector

$$\overline{T} := \begin{cases} \hat{T}, & |\hat{T}| > t_{\eta}^{*} \\ 0, & |\hat{T}| \le t_{\eta}^{*} \end{cases}$$
(A.72)

The error probability of this sub-optimal detector is clearly an upper bound on the error probability of the MAP detector.

Let us decompose the error probability \overline{T} as

$$P_{\text{error}}(\overline{T}) \le \sum_{t \in \mathcal{T}} P_T(t) \cdot \mathbb{P}\left[\overline{T} \neq T \mid T = t\right],$$
(A.73)

so the summation over $t \in \mathcal{T}$ can be partitioned into the following five index sets: $-\{[\ell - 1] \setminus [t_{\eta}^*]\}, \{-[t_{\eta}^*]\}, \{0\}, [t_{\eta}^*], \text{ and } [\ell] \setminus [t_{\eta}^*]$. We next separately analyze each of them.

Analysis of the type-I error probability:

We begin with the set $\{0\}$, that is, analyzing the type-I error probability – the probability of a wrong detection in case T = 0 to $\{\overline{T} \neq 0\} = \{\overline{T} > 0\} \cup \{\overline{T} < 0\}$. For $\overline{T} > 0$, we bound for all n sufficiently large,

$$\mathbb{P}\left[\overline{T} > 0 \mid T = 0\right]$$

$$\stackrel{(a)}{=} \mathbb{P}\left[\bigcup_{t \in [\ell] \setminus [t_{\eta}^*]} \left\{ X_{\ell-t+1}^{\ell}(1) = X_1^t(2) \right\} \cap \left\{ \frac{1}{P_{X_1^t}(X_1^t(2))} \ge n_{\ell} \right\} \middle| T = 0 \right]$$
(A.74)

$$\overset{(b)}{\leq} \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \mathbb{P}\left[\left\{X_{\ell-t+1}^{\ell}(1) = X_{1}^{t}(2)\right\} \cap \left\{\frac{1}{P_{X_{1}^{t}}\left(X_{1}^{t}(2)\right)} \ge n_{\ell}\right\} \middle| T = 0\right]$$

$$\leq \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \sum_{k=-(n-1)}^{n-1} \mathbb{P}\left[I(1) - I(2) = k \mid T = 0\right] \times$$

$$\mathbb{P}\left[\left\{X_{\ell-t+I(1)}^{\ell+I(1)-1} = X_{I(2)}^{t+I(2)-1}\right\} \cap \left\{\frac{1}{P_{X_{I(2)}^{t+I(2)-1}}\left(X_{I(2)}^{t+I(2)-1}\right)} \ge n_{\ell}\right\} \middle| I(1) - I(2) = k, T = 0\right]$$

$$(A.76)$$

$$\stackrel{(c)}{=} \sum_{t \in [\ell] \setminus [t_{\eta}^*]} \sum_{k=-n}^n \mathbb{P}\left[I(1) - I(2) = k \mid T=0\right] \mathbb{P}\left[\left\{X_{\ell-t+k+1}^{\ell+k} = X_1^t\right\} \cap \left\{\frac{1}{P_{X_1^t}\left(X_1^t\right)} \ge n_\ell\right\}\right]$$
(A.77)

$$\stackrel{(d)}{\leq} \frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \sum_{k=-n}^{-\ell} \mathbb{P}\left[\left\{X_{\ell-t+k+1}^{\ell+k} = X_{1}^{t}\right\} \cap \left\{\frac{1}{P_{X_{1}^{t}}\left(X_{1}^{t}\right)} \ge n_{\ell}\right\}\right] + \frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \sum_{k=\ell}^{n} \mathbb{P}\left[\left\{X_{\ell-t+k+1}^{\ell+k} = X_{1}^{t}\right\} \cap \left\{\frac{1}{P_{X_{1}^{t}}\left(X_{1}^{t}\right)} \ge n_{\ell}\right\}\right],$$

$$(A.78)$$

where (a) follows from the detection rule of Prop. 4 which states that an error occurs if there is a matching overlapping fragment of length larger than t_{MDO} , and it has small enough probability, as well as the modification in (A.72), which results that an error to $0 < \overline{T} \le t_{\eta}^*$ is impossible, (b) follows from the union bound, (c) follows from stationarity, and (d) follows since given T = 0, $\mathbb{P}[I(1) - I(2) = k | T = 0] > 0$ only if either $\ell \le k \le n$ or $-n \le k \le -\ell$, and otherwise it follows from the conditional distribution of I(2) given I(1) in (16) that

$$\mathbb{P}[I(1) - I(2) = k \mid T = 0] \le \frac{1}{n}.$$
(A.79)

Let us continue by upper bounding the first double summation in (A.78), and the second double summation can be similarly upper bounded. To this end, let $\{\tau(n)\}_{n\in\mathbb{N}}$ be a monotonic non-decreasing sequence of integers such that $\tau(n) = \omega_n(1)$ and $\tau(n) = o(\log n)$. For example, $\tau(n) = \lceil \sqrt{\log n} \rceil$ is a suitable choice. For brevity, let us next denote $\tau \equiv \tau(n)$. We now bound the first double sum in (A.78) as

$$\frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \sum_{k=-n}^{-\ell} \mathbb{P}\left[\left\{X_{\ell-t+k+1}^{\ell+k} = X_{1}^{t}\right\} \cap \left\{\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})} \ge n_{\ell}\right\}\right]$$

$$\stackrel{(a)}{\leq} \frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \sum_{k=-n}^{-\ell} \mathbb{P}\left[\left\{X_{\ell-t+k+\tau}^{\ell+k} = X_{\tau}^{t}\right\} \cap \left\{\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})} \ge n_{\ell}\right\}\right]$$
(A.80)

$$\stackrel{(b)}{\leq} \frac{1}{n} \sum_{t \in [\ell] \setminus [t^*_{\tau}]} \sum_{k=-n}^{-\ell} \mathbb{P}\left[\left\{ X^{\ell+k}_{\ell-t+k+\tau} = X^t_{\tau} \right\} \cap \left\{ \frac{1}{P_{X^t_{\tau}}\left(x^t_{\tau}\right)} \ge \rho^{\tau-1} \cdot n_{\ell} \right\} \right]$$
(A.81)

$$= \frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \sum_{k=-n}^{-\ell} \sum_{x_{\tau}^{t} \in \mathcal{X}^{t-\tau+1}} P_{X_{\ell-t+k+\tau}^{\ell+k}} \left(x_{\tau}^{t} \right) \mathbb{P} \left[X_{\tau}^{t} = x_{\tau}^{t} \mid X_{\ell-t+k+\tau}^{\ell+k} = x_{\tau}^{t} \right] \cdot \mathbb{1} \left[\frac{1}{P_{X_{\tau}^{t}} \left(x_{\tau}^{t} \right)} \ge \rho^{\tau-1} \cdot n_{\ell} \right]$$
(A.82)

$$\stackrel{(c)}{\leq} \frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \sum_{k=-n}^{-\ell} \sum_{x_{1}^{t-\tau+1} \in \mathcal{X}^{t-\tau+1}} P_{X_{1}^{t-\tau+1}}^{2} \left(x_{1}^{t-\tau+1}\right) \mathbb{1} \left[\frac{1}{P_{X_{1}^{t-\tau+1}}(x_{1}^{t-\tau+1})} \ge \rho^{\tau-1} \cdot n_{\ell} \right]$$

$$+ \frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \sum_{k=-n}^{-\ell} \sum_{x_{1}^{t-\tau+1} \in \mathcal{X}^{t-\tau+1}} P_{X_{1}^{t-\tau+1}} \left(x_{1}^{t-\tau+1}\right) d \left(\tau-\ell-k\right) \cdot \mathbb{1} \left[\frac{1}{P_{X_{1}^{t-\tau+1}}(x_{1}^{t-\tau+1})} \ge \rho^{\tau-1} \cdot n_{\ell} \right],$$

$$(A.83)$$

where (a) follows by slightly shortening the required match length by $\tau + 1$ symbols, (b) follows from the assumption $\mathcal{H}_{-\infty}(\mathbf{X}) < \infty$, which assures that for any $\rho < |\mathcal{X}|^{-\mathcal{H}_{-\infty}(\mathbf{X})} < 1$ there exists n_0 such that for all $n \ge n_0$,

$$\frac{1}{P_{X_{1}^{t}}(x_{1}^{t})} \leq \frac{1}{P_{X_{\tau}^{t}}(x_{\tau}^{t}) \cdot \min_{\tilde{x}_{1}^{\tau-1} \in \operatorname{supp}\left[P_{X_{1}^{\tau-1}}\right]} P_{X_{1}^{\tau-1}}(\tilde{x}_{1}^{\tau-1})}$$
(A.84)

$$\leq \frac{1}{P_{X_{\tau}^t}\left(x_{\tau}^t\right)} \left(\frac{1}{\rho}\right)^{\tau-1},\tag{A.85}$$

(c) follows from the definition of the strong mixing coefficient (Definition 5), which implies that

$$\mathbb{P}\left[X_{\tau}^{t} = x_{\tau}^{t} \mid X_{\ell-t+k+\tau}^{\ell+k} = x_{\tau}^{t}\right] \leq \mathbb{P}\left[X_{\tau}^{t} = x_{\tau}^{t}\right] + d\left(\tau - \ell - k\right)$$
(A.86)

and utilizing stationarity to replace $P_{X_{\ell-t+k+1+\tau}^{\ell+k}} = P_{X_{\tau}^{t}} = P_{X_{1}^{t-\tau+1}}$. We now consider each of the triple summations in (A.83). The first triple summation is upper bounded as

$$\frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \sum_{k=-n}^{-\ell} \sum_{x_{1}^{t-\tau+1} \in \mathcal{X}^{t-\tau+1}} P_{X_{1}^{t-\tau+1}}^{2} \left(x_{1}^{t-\tau+1}\right) \mathbb{1} \left[\frac{1}{P_{X_{1}^{t-\tau+1}}(x_{1}^{t-\tau+1})} \ge \rho^{\tau-1} \cdot n_{\ell} \right] \\
\leq \ell \cdot \max_{t_{\eta}^{*} \le t \le \ell} \sum_{x_{1}^{t-\tau+1} \in \mathcal{X}^{t-\tau+1}} P_{X_{1}^{t-\tau+1}}^{2} \left(x_{1}^{t-\tau+1}\right) \mathbb{1} \left[\frac{1}{P_{X_{1}^{t-\tau+1}}(x_{1}^{t-\tau+1})} \ge \rho^{\tau-1} \cdot n_{\ell} \right]$$
(A.87)

$$\stackrel{(a)}{=} \ell \cdot o\left(\frac{1}{n}\right) \tag{A.88}$$

$$\stackrel{(b)}{=} o\left(\frac{\log n}{n}\right),\tag{A.89}$$

where (a) follows from (A.1) in Lemma 14, by first replacing $t \leftrightarrow t - \tau + 1$ and noting that for all $t \ge t_{\eta}^* = (1+2\eta)t^*$ the assumption $\tau = o(\log n) = o(t_{\eta}^*)$ implies that

$$t - \tau + 1 \ge (1 + 2\eta)t_{\eta}^* - \tau + 1 \tag{A.90}$$

$$\geq (1+\eta)t_{\eta}^{*} \tag{A.91}$$

$$=\frac{(1+\eta)}{\mathcal{H}_1(\mathbf{X})}\log n,\tag{A.92}$$

and (b) follows since $\ell = \beta \log n$. The second triple summation in (A.83) is upper bounded as

$$\frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \sum_{k=-n}^{-\ell} \sum_{x_{1}^{t-\tau+1} \in \mathcal{X}^{t-\tau+1}} P_{X_{1}^{t-\tau+1}} \left(x_{1}^{t-\tau+1}\right) d\left(\tau-\ell-k\right) \cdot \mathbb{1}\left[\frac{1}{P_{X_{1}^{t-\tau+1}}(x_{1}^{t-\tau+1})} \ge \rho^{\tau-1} \cdot n_{\ell}\right]$$

$$\leq \frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^{*}]} \sum_{k=-n}^{r} d\left(\tau - \ell - k\right) \sum_{x_{1}^{t-\tau+1} \in \mathcal{X}^{t-\tau+1}} P_{X_{1}^{t-\tau+1}}\left(x_{1}^{t-\tau+1}\right)$$
(A.93)

$$= \frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^*]} \sum_{k=-n}^{\infty} d(\tau - \ell - k)$$
(A.94)

$$\leq \frac{\ell}{n} \cdot \sum_{s=\tau}^{\infty} d(s) \tag{A.95}$$

$$\stackrel{(a)}{=} \frac{\ell}{n} \cdot o_n(1) \tag{A.96}$$

$$\stackrel{(b)}{=} o\left(\frac{\log n}{n}\right),\tag{A.97}$$

where (a) follows since the assumption $\sum_{s=0}^{\infty} d(s) < \infty$ of the theorem, combined with $\tau \equiv \tau(n) = \omega_n(1)$ implies that $\sum_{s=\tau(n)}^{\infty} d(s) \to 0$ as $n \to \infty$, and (b) follows again since $\ell = \beta \log n$. Substituting (A.89) and (A.97) into (A.83) results that the first double summation in (A.78) is

$$\frac{1}{n} \sum_{t \in [\ell] \setminus [t_{\eta}^*]} \sum_{k=-n}^{-\ell} \mathbb{P}\left[\left\{X_{\ell-t+k+1}^{\ell+k} = X_1^t\right\} \cap \left\{\frac{1}{P_{X_1^t}(X_1^t)} \ge n_\ell\right\}\right] = o\left(\frac{\log n}{n}\right).$$
(A.98)

Similarly, the second double summation in (A.78) is also $o(\frac{\log n}{n})$, and so (A.78) implies that

$$\mathbb{P}\left[\overline{T} > 0 \mid T = 0\right] = o\left(\frac{\log n}{n}\right).$$
(A.99)

It can be proved using analogous arguments that

$$\mathbb{P}\left[\overline{T} < 0 \mid T = 0\right] = o\left(\frac{\log n}{n}\right) \tag{A.100}$$

and so the type-I error probability is upper bounded as

$$\mathbb{P}\left[\overline{T} \neq 0 \mid T=0\right] = \mathbb{P}\left[\overline{T} > 0 \mid T=0\right] + \mathbb{P}\left[\overline{T} < 0 \mid T=0\right] = o\left(\frac{\log n}{n}\right).$$
(A.101)

<u>Analysis of the type-II error probability from</u> $T \in \{-[t^*_{\eta}]\} \cup [t^*_{\eta}]$:

In this case, we trivially bound the conditional error probability by 1 and use $P_T(t) = \frac{1}{n}$ to obtain that the contribution of these sets to the error probability in (A.73) is upper bounded by

$$\sum_{t \in \{-[t_{\eta}^*]\} \cup [t_{\eta}^*]} P_T(t) \cdot \mathbb{P}\left[\overline{T} \neq T \mid T = t\right] \le \frac{2t_{\eta}^*}{n}.$$
(A.102)

<u>Analysis of the type-II error probability from</u> $T \in [\ell] \setminus [t_{\eta}^*]$:

In this case, an error may occur in one of three events: (1) An error to $\overline{T} = 0$, that is, deciding that the reads do not overlap. (2) An error to a longer positive overlap $\overline{T} \in [\ell] \setminus [\tilde{t}]$. (3) An error to a longer negative overlap $\overline{T} \in -\{[\ell-1] \setminus [t_{\eta}^*]\}$. We next upper bound the probability of each of these error events, and then use a union bound to upper bound $\mathbb{P}[\overline{T} \neq T \mid T = \tilde{t}]$.

First, the error probability of an error from $T = \tilde{t}$ to $\overline{T} = 0$, is upper bounded as

$$\mathbb{P}\left[\overline{T}=0 \mid T=\tilde{t}\right] \stackrel{(a)}{\leq} \mathbb{P}\left[\frac{1}{P_{X_{1}^{\tilde{t}}}\left(X_{1}^{\tilde{t}}(1)\right)} \leq n_{\ell}\right]$$
(A.103)

$$=\sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}(x_1^t) \cdot \mathbb{1} \left[\frac{1}{P_{X_1^{\tilde{t}}}\left(X_1^{\tilde{t}}(1)\right)} \le n_\ell \right]$$
(A.104)

$$=o_n(1), \tag{A.105}$$

where (a) follows since an error will occur only if $P_{X_1^{\tilde{t}}}^{-1}(X_1^{\tilde{t}}(1)) \leq n_{\ell}$ because the reads are noiseless and so $X_{\ell-\tilde{t}+1}^{\ell}(1) = X^{\tilde{t}}(2)$ occurs with probability 1, and (b) follows from (A.2) in Lemma 14 since $\tilde{t} \geq t_{\eta}^* \geq \frac{(1+\eta)}{\mathcal{H}_1(\mathbf{X})} \log n$. Second, the probability of an error from $T = \tilde{t}$ to $\overline{T} \in [\ell] \setminus [\tilde{t}]$ is upper bounded from the union bound as

$$\mathbb{P}\left[\overline{T} \in [\ell] \setminus [\tilde{t}] \mid T = \tilde{t}\right] \le \sum_{\overline{t} = \tilde{t} + 1}^{\ell} \mathbb{P}\left[\overline{T} = \overline{t} \mid T = \tilde{t}\right]$$
(A.106)

$$\leq \ell \cdot \max_{\tilde{t}+1 \leq \bar{t} \leq \ell} \mathbb{P}\left[\overline{T} = \bar{t} \mid T = \tilde{t}\right].$$
(A.107)

Now, for such \overline{t} which satisfies $\ell \geq \overline{t} \geq \tilde{t} + 1$,

$$\mathbb{P}\left[\overline{T} = \overline{t} \mid T = \widetilde{t}\right]$$

$$\stackrel{(a)}{=} \mathbb{P}\left[\left\{X_{\ell-\overline{t}+1}^{\ell}(1) = X_{1}^{\overline{t}}(2)\right\} \cap \left\{\frac{1}{P_{X_{1}^{\overline{t}}}\left(X_{1}^{\overline{t}}(2)\right)} \ge n_{\ell}\right\} \middle| T = \widetilde{t}\right]$$
(A.108)

$$\stackrel{(b)}{\leq} \mathbb{P}\left[X_{\ell-\tilde{t}+1}^{\ell}(1) = X_1^{\tilde{t}}(2) \mid T = \tilde{t}\right]$$
(A.109)

$$= \mathbb{P}\left[X_{\ell-\bar{t}+1+I(1)}^{\ell+I(1)} = X_{1+I(2)}^{\bar{t}+I(2)} \mid T = \tilde{t}\right]$$
(A.110)

$$\stackrel{(c)}{=} \mathbb{P}\left[X_{\ell-\tilde{t}+1+I(1)}^{\ell+I(1)} = X_{1+I(1)+\ell-\tilde{t}}^{\tilde{t}+I(1)+\ell-\tilde{t}} \mid T = \tilde{t}\right]$$
(A.111)

$$\stackrel{(d)}{=} \mathbb{P}\left[X_1^{\overline{t}} = X_{1+\overline{t}-\widetilde{t}}^{2\overline{t}-\widetilde{t}} \mid T = \widetilde{t}\right]$$
(A.112)

$$\stackrel{(e)}{=} \mathbb{P}\left[X_1^{\overline{t}} = X_{1+\overline{t}-\overline{t}}^{2\overline{t}-\overline{t}}\right] \tag{A.113}$$

$$\stackrel{(f)}{=} o\left(\frac{1}{\log n}\right),\tag{A.114}$$

where (a) follows from the definition of the detection rule \overline{T} , (b) follows by relaxing the condition $\frac{1}{P_{X_{1}^{\overline{t}}}(X_{1}^{\overline{t}}(2))} \ge n_{\ell}$, (c) follows since for $T = \tilde{t} > 0$ it holds that $I(2) = \ell - \tilde{t} - I(1)$, (d) follows from stationarity, (e) follows

from independence of T and $\{X_i\}_{i\in\mathbb{Z}}$, and (f) follows from Lemma 15, with the notation change $t \leftarrow \overline{t}$ and $s \leftarrow \overline{t} - \tilde{t}$. Substituting (A.114) into (A.107) implies that

$$\mathbb{P}\left[\overline{T} \in [\ell] \setminus [\tilde{t}] \mid T = \tilde{t}\right] = o_n(1).$$
(A.115)

Third, similarly to (A.107), the probability of an error from $T = \tilde{t}$ to $\overline{T} \in -\{[\ell - 1] \setminus [\tilde{t}]\}$ is upper bounded from the union bound as

$$\mathbb{P}\left[\overline{T} \in -\{[\ell-1] \setminus [\tilde{t}]\} \mid T = \tilde{t}\right] \le \sum_{\overline{t} = \tilde{t}+1}^{\ell} \mathbb{P}\left[\overline{T} = -\overline{t} \mid T = \tilde{t}\right]$$
(A.116)

$$\leq \ell \cdot \max_{\tilde{t}+1 \leq \bar{t} \leq \ell} \mathbb{P}\left[\overline{T} = -\overline{t} \mid T = \tilde{t}\right].$$
(A.117)

Now, for such $\ell \ge -\overline{t} \ge \tilde{t} + 1$ we may analyze similarly to (A.114), by replacing the role of X(1) and X(2), as:

$$\mathbb{P}\left[\overline{T} = -\overline{t} \mid T = \widetilde{t}\right]$$

$$= \mathbb{P}\left[\left\{X_{\ell-\overline{t}+1}^{\ell}(2) = X_{1}^{\overline{t}}(1)\right\} \cap \left\{\frac{1}{P_{X_{1}^{\overline{t}}}\left(X_{1}^{\overline{t}}(1)\right)} \ge n_{\ell}\right\} \middle| T = \widetilde{t}\right]$$
(A.118)

$$\leq \mathbb{P}\left[X_{\ell-\overline{t}+1}^{\ell}(2) = X_1^{\overline{t}}(1) \mid T = \tilde{t}\right]$$
(A.119)

$$= \mathbb{P}\left[X_{\ell-\bar{t}+1+I(2)}^{\ell+I(2)} = X_{1+I(1)}^{\bar{t}+I(1)} \mid T = \tilde{t}\right]$$
(A.120)

$$= \mathbb{P}\left[X_{\ell-\bar{t}+1+I(2)}^{\ell+I(2)} = X_{1+\ell-\bar{t}-I(2)}^{\bar{t}+\ell-\bar{t}-I(2)} \mid T = \tilde{t}\right]$$
(A.121)

$$= \mathbb{P}\left[X_{1+\tilde{t}-\tilde{t}+2I(2)}^{\tilde{t}+2I(2)} = X_1^{\tilde{t}} \mid T = \tilde{t}\right]$$
(A.122)

$$= \mathbb{P}\left[X_{1+\tilde{t}-\tilde{t}+2I(2)}^{\tilde{t}+2I(2)} = X_{1}^{\tilde{t}}\right]$$
(A.123)

$$= o\left(\frac{1}{\log n}\right),\tag{A.124}$$

where the transitions follow from the same arguments as in (A.114), by replacing the role of X(1) and X(2)(while $I(2) = \ell - \tilde{t} - I(1)$ still holds). Substituting (A.124) into (A.117) implies that

$$\mathbb{P}\left[t_{\eta}^* \in -\{[\ell-1] \setminus [\tilde{t}]\} \mid T = \tilde{t}\right] = o_n(1).$$
(A.125)

Combining all three conditional error probabilities (A.105) (A.115) and (A.125) we obtain

$$\mathbb{P}\left[\overline{T} \neq T \mid T = \tilde{t}\right] = o_n(1). \tag{A.126}$$

Hence, also

$$\sum_{\tilde{t} \in [\ell] \setminus [t_{\eta}^*]} P_T(t) \cdot \mathbb{P}\left[\overline{T} \neq T \mid T = t\right] = o\left(\frac{\ell}{n}\right) = o\left(\frac{\log n}{n}\right).$$
(A.127)

This completes the analysis of the contribution of $\tilde{t} \in [\ell] \setminus [t_{\eta}]$ to the error probability in the decomposition (A.73).

<u>Analysis of the type-II error probability from</u> $T = \tilde{t} \in -\{[\ell - 1] \setminus [t_{\eta}^*]\}$:

This case can be analyzed in a complete analogous manner to the previous case, and results, similarly to (A.127), the upper bound

$$\sum_{\tilde{t} \in -\{[\ell-1] \setminus [t_{\eta}^*]\}} P_T(t) \cdot \mathbb{P}\left[\overline{T} \neq T \mid T = t\right] = o\left(\frac{\log n}{n}\right).$$
(A.128)

Combining the different error probabilities:

Substituting (A.101), (A.102), (A.127) and (A.128) into (A.73) results

$$P_{\text{error}}^* \le P_{\text{error}}(\overline{T}) \tag{A.129}$$

$$\leq \frac{2t_{\eta}^{*}}{n} + o\left(\frac{1}{n}\right) + o\left(\frac{\log n}{n}\right) \tag{A.130}$$

$$\leq 2(1+2\eta) \cdot [1+o_n(1)] \cdot \frac{1}{\mathcal{H}_1(\mathbf{X})} \frac{\log n}{n}.$$
(A.131)

The proof of the upper bound (53) is then completed by taking $\eta \downarrow 0$. This completes the proof of Theorem 6.

B. Details for analysis of trade-off between two types of error 8

Suppose that we modify the detection rule (41) of Prop. 4 and make the condition of overlap detection more stringent, with the goal of reducing the type-I error probability to $o(\frac{\log n}{n^q})$ for some degree q > 1. We can achieve this by replacing $n_\ell \sim n$ in the detection rule (41) of Prop. 4 by n^μ for some $\mu > 1$. In turn, the minimal reliably detectable overlap $t^* \sim \frac{\log n}{\mathcal{H}_1(\mathbf{X})}$ will now be replaced by a modified parameter $t^{\#} \sim \frac{\log n}{h}$ for some $h < \mathcal{H}_1(\mathbf{X})$ which in turn will increase the dominating contribution of the type-II error probability (misdetection of overlaps) to $\frac{1}{h} \cdot \frac{\log n}{n}$. Our goal is thus to characterize the trade-off between q and h (and how they can be tuned by controlling μ). The analysis of the error probability of this modified detector, follows the same events as in the analysis of the proof of Theorem 6. As the technical details can be easily discerned from that proof, we only provide here a sketch of required modifications. Since the contribution of error in detecting a wrong overlap to the type-II error, that is, an error from T = t to $\hat{T} = \tilde{t}$, is negligible compared to the error probability from T = 0 to $\hat{T} = t$ for some $t > t^{\#}$ and vice-versa.

First, the probability of an error from T = 0 to $\hat{T} = t > 0$, previously bounded in (57), will now be bounded as

$$\mathbb{P}\left[\hat{T}=t \mid T=0\right] \lesssim \sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}^2\left(x_1^t\right) \mathbb{1}\left[\frac{1}{P_{X_1^t}(x_1^t)} \ge n^{\mu}\right].$$
(A.132)

$$q = \mu r - (r-1)\frac{\mathcal{H}_{2-r}(\mathbf{X})}{h}$$
(A.133)

for some choice of r > 0. Second, the probability of an error from T = t to $\hat{T} = 0$, which was previously bounded in (62), will now be bounded as

$$\mathbb{P}\left[\hat{T}=0 \mid T=t\right] \lessapprox \sum_{x_1^t \in \mathcal{X}^{\otimes t}} P_{X_1^t}\left(x_1^t\right) \mathbb{1}\left[\frac{1}{P_{X_1^t}(x_1^t)} \le n^{\mu}\right],\tag{A.134}$$

and this is $o_n(1)$ only if

$$(1-r)\left[\frac{\mathcal{H}_{2-r}(\mathbf{X})}{h} - \mu\right] > 0 \tag{A.135}$$

for some choice of $r \in (0,1)$. This last condition means that $\mu \leq \frac{\mathcal{H}_{2-r}(\mathbf{X})}{h}$. Let us assume that the same parameter r is used for both types of error. Then, by choosing μ to its maximal value $\frac{\mathcal{H}_{2-r}(\mathbf{X})}{h}$, the maximal degree of the type-I error upper bound is

$$q = \frac{\mathcal{H}_{2-r}(\mathbf{X})}{h}r - (r-1)\frac{\mathcal{H}_{2-r}(\mathbf{X})}{h} = \frac{\mathcal{H}_{2-r}(\mathbf{X})}{h},$$
(A.136)

which could be at most μ for an optimized choice of r. Consequently, we deduce that for any given choice of μ in the modified detector, it holds that the contribution of the type-I error to the error probability is

$$P_T(0) \cdot p_{\text{type-I}}(\hat{T}) = o\left(\frac{\log n}{n^{\mu}}\right) \tag{A.137}$$

whereas the contribution of the type-II error probability is

$$\sum_{t \in \{-[\ell-1]\} \cup [\ell]} \mathbb{P}[T=t] \cdot p_{\text{type-II}}(\hat{T};t) \sim \frac{\mu}{\mathcal{H}_1(\mathbf{X})} \frac{\log n}{n}.$$
(A.138)

In words, the cost of improving the decay rate of the type-I error probability to $o(\frac{\log n}{n^{\mu}})$ results an increase of a factor of μ in the total error probability.

C. Details for Example 8

We next provide details on the computation of the strong mixing coefficient for the Markov example (Example 8). We may write the Markov kernel in a matrix form

$$K = \epsilon \cdot \mathbf{1}\mathbf{1}^{\top} + (1 - |\mathcal{X}|\epsilon) \cdot \mathbf{I}, \tag{A.139}$$

where $\mathbf{1}^{\top} := (1, 1, \dots, 1) \in \mathbb{R}^{|\mathcal{X}|}$ is the all ones vector (here, of dimension $|\mathcal{X}|$), and $\mathbf{I} \in \mathbb{R}^{|\mathcal{X}| \times |\mathcal{X}|}$ is the identity matrix. Then, for an integer s > 1, it is easy to verify that

$$K^s = a_s \cdot \mathbf{1}\mathbf{1}^\top + b_s \cdot \boldsymbol{I},\tag{A.140}$$

where $a_1 = \epsilon$, $b_1 = (1 - |\mathcal{X}|\epsilon)$, and the recursive formulas hold:

$$a_{s+1} = a_s + b_s \epsilon \tag{A.141}$$

$$b_{s+1} = b_s \cdot (1 - |\mathcal{X}|\epsilon) . \tag{A.142}$$

Thus, $b_s = (1 - |\mathcal{X}|\epsilon)^s$ and

$$a_s = \epsilon \cdot \sum_{i=0}^{s-1} \left(1 - |\mathcal{X}|\epsilon\right)^i = \epsilon \cdot \frac{1 - \left(1 - |\mathcal{X}|\epsilon\right)^s}{|\mathcal{X}|\epsilon} = \frac{1}{|\mathcal{X}|} \left[1 - \left(1 - |\mathcal{X}|\epsilon\right)^s\right].$$
 (A.143)

Due to the symmetry between the letters, the mixing coefficient is determined as the total variation distance between Π_X and any of the rows of K^s , say the first, given by $(a_s + b_s, a_s, a_s, \dots, a_s)$. So, since $a_s \leq \frac{1}{|\mathcal{X}|}$ it must hold that $a_s + b_s \geq \frac{1}{|\mathcal{X}|}$. Hence,

$$d(s) = \max_{x \in \mathcal{X}} \|K^{s}(x, \cdot) - P_{X_{1}}\|_{\mathrm{TV}}$$
(A.144)

$$= \frac{1}{2} \left(a_s + b_s - \frac{1}{|\mathcal{X}|} \right) + \frac{1}{2} (|\mathcal{X}| - 1) \left(\frac{1}{|\mathcal{X}|} - a_s \right)$$
(A.145)

$$=\frac{|\mathcal{X}|-1}{|\mathcal{X}|}\left(1-|\mathcal{X}|\epsilon\right)^{s}\tag{A.146}$$

$$\leq \frac{|\mathcal{X}| - 1}{|\mathcal{X}|} \cdot e^{-|\mathcal{X}|\epsilon \cdot s}.$$
(A.147)

D. Proof of Theorem 9

We next turn to the proof of Theorem 9. To this end, recall that in case the process has memory, the likelihoods for both $T = t \neq 0$ and T = 0 do not take the simple form of the memoryless case (Prop. 4). Nonetheless, we show that these likelihoods are sufficiently tight approximations. We show this separately for the case $T = t \neq 0$ and then for T = 0. To this end, will use the following known result from ergodic theory, which is an easy consequence of the AEP (see, for example, [9, Lemma 3.4.1.]). We provide for completeness the simple proof.

Lemma 17. Let X be a stationary and ergodic process. Let $m \in \mathbb{N}_+$ and $k \equiv k(m)$ be such that $k(m) = \omega_m(1)$. Then,

$$\frac{1}{m+k}\log\frac{P_{X_1^{m+k}}(X_1^{m+k})}{P_{X_1^k}(X_1^k) \cdot P_{X_{k+1}^{m+k}}(X_{k+1}^{m+k})} \to 0$$
(A.148)

as $m \to \infty$, with probability 1.

Proof: The AEP for ergodic sources [6, Th. 16.8.1] implies that

$$\frac{1}{m+k}\log P_{X_1^{m+k}}(X_1^{m+k}) \to -\mathcal{H}_1(\mathbf{X})$$
(A.149)

and

$$\frac{1}{m+k}\log\frac{1}{P_{X_{1}^{k}}(X_{1}^{k})\cdot P_{X_{k+1}^{m+k}}(X_{k+1}^{m+k})} = \frac{k}{m+k}\cdot\frac{1}{k}\log\frac{1}{P_{X_{1}^{k}}(X_{1}^{k})} + \frac{m}{m+k}\cdot\frac{1}{m}\log\frac{1}{P_{X_{k+1}^{m+k}}(X_{k+1}^{m+k})}$$
(A.150)

$$\rightarrow \frac{k}{m+k}\mathcal{H}_1(\mathbf{X}) + \frac{m}{m+k}\mathcal{H}_1(\mathbf{X})$$
(A.151)

$$=\mathcal{H}_1(\mathbf{X}),\tag{A.152}$$

as $m \to \infty$, with probability 1. The result then follows by summing both limits.

Lemma 18. Consider the noiseless setting (Def. 1). Let $0 < \theta_0 < \theta_1 < \beta$ be given. For any $q \in (0,1)$ and $\nu > 0$ there exists $n_0(\nu, q)$ such that for all $n \ge n_0$ it holds with probability 1 - q that

$$\max_{\theta_0 \log n \le t \le \theta_1 \log n} \log \frac{f\left(X_1^{\ell}(1), X_1^{\ell}(2); t\right)}{\mathbbm{1}\left[X_{\ell-t+1}^{\ell}(1) = X_1^{t}(2)\right] \cdot P_{X_1^{\ell}}\left(X_1^{\ell}(1)\right) \cdot P_{X_1^{\ell}}\left(X_1^{\ell}(2)\right) \cdot \Gamma_+(t)} \le \nu \cdot \log n.$$
(A.153)

Proof: For T = t, the likelihood was derived in Lemma 3, and is given by

$$f\left(x_{1}^{\ell}(1), x_{1}^{\ell}(2); t\right) = \mathbb{1}\left[x_{\ell-t+1}^{\ell}(1) = x_{1}^{t}(2)\right] \cdot P_{X_{1}^{\ell}}\left(x_{1}^{\ell}(1)\right) \cdot \frac{P_{X_{1}^{\ell}|X_{-(\ell-t-1)}^{0}}\left(x_{1}^{\ell}(2) \mid x_{1}^{\ell-t}(1)\right)}{P_{X_{1}^{t}|X_{-(\ell-t-1)}^{0}}\left(x_{1}^{t}(2) \mid x_{1}^{\ell-t}(1)\right)}.$$
 (A.154)

For clarity, we will next omit the arguments and use the abbreviated notation $P_{X_1^{\ell}|X_{-(\ell-t-1)}^0} \equiv \mathbb{P}[X_1^{\ell} \mid X_{-(\ell-t-1)}^0]$. Our goal is to show that the ratio in (A.154), given by

$$\frac{\mathbb{P}[X_1^{\ell} \mid X_{-(\ell-t-1)}^0]}{\mathbb{P}[X_1^{t} \mid X_{-(\ell-t-1)}^0]},\tag{A.155}$$

is tightly upper bounded by $\mathbb{P}[X_1^{\ell}]/\mathbb{P}[X_1^t]$, up to a sufficiently small multiplicative term. We begin with the probability on the numerator. Let us write

$$\mathbb{P}[X_1^{\ell} \mid X_{-(\ell-t-1)}^0] = \mathbb{P}[X_1^{\ell}] \cdot \frac{\mathbb{P}[X_{-(\ell-t-1)}^{\ell}]}{\mathbb{P}[X_{-(\ell-t-1)}^0] \cdot \mathbb{P}[X_1^{\ell}]}.$$
(A.156)

To bound the ratio appearing in this term, we use Lemma 17 that states that if $m \equiv m(n) \rightarrow \infty$ and $k \equiv k(n) \rightarrow \infty$ as $n \rightarrow \infty$ then

$$\frac{1}{m+k} \log \frac{\mathbb{P}[X_1^{m+k}]}{\mathbb{P}[X_1^k] \cdot \mathbb{P}[X_{k+1}^{m+k}]} \to 0,$$
(A.157)

as $n \to \infty$ with probability 1. Let (Ω, \mathcal{F}, P) be the probability space underlying the process $X \equiv X(\omega)$, where $\omega \in \Omega$. In principle, the rate of the convergence to the limit 0 in (A.157) may depend on $\omega \in \Omega$. However, by

Egorov's theorem (e.g., [5, p. 79, Ex. 5]), for any $q \in (0, 1)$ there exists a set $\mathcal{B}_1 \in \mathcal{F}$ such that $\mathbb{P}[\mathcal{B}_1] \leq q/2$ and the convergence is *uniform* on \mathcal{B}_1^c . Thus, for any $\nu > 0$ there exists $n_0(\nu, q)$ and such that

$$\sup_{\omega \in \mathcal{B}^c} \frac{1}{m+k} \log \frac{\mathbb{P}[X_1^{m+k}(\omega)]}{\mathbb{P}[X_1^k(\omega)] \cdot \mathbb{P}[X_{k+1}^{m+k}(\omega)]} \le \nu$$
(A.158)

for all $n \ge n_0$ sufficiently large so that $m(n) \ge m(n_0)$ and $k(n) \ge k(n_0)$. Using $m = \ell - t$ and $k = \ell$ and noting that for any $\theta_0 \log n \le t \le \theta_1$ it holds that $m \equiv m(n) \to \infty$ as $n \to \infty$, we deduce that

$$\sup_{\omega \in \mathcal{B}^c} \max_{\theta_0 \log n \le t \le \theta_1 \log n} \frac{1}{2\ell - t} \log \frac{\mathbb{P}[X^{\ell}_{-(\ell - t - 1)}(\omega)]}{\mathbb{P}[X^0_{-(\ell - t - 1)}(\omega)] \cdot \mathbb{P}[X^{\ell}_1(\omega)]} \le \nu.$$
(A.159)

Thus, also

$$\sup_{\omega \in \mathcal{B}^c} \max_{\theta_0 \log n \le t \le \theta_1 \log n} \log \frac{\mathbb{P}[X_{-(\ell-t-1)}^\ell]}{\mathbb{P}[X_{-(\ell-t-1)}^0] \cdot \mathbb{P}[X_1^\ell]} \le 2\nu\beta \cdot \log n$$
(A.160)

In a completely similar fashion, we can lower bound the probability on the denominator, that is, $\mathbb{P}[X_1^t \mid X_{-(\ell-t-1)}^0]$, showing uniform convergence, except on a set $\mathcal{B}_2 \in \mathcal{F}$ which satisfy $\mathbb{P}[\mathcal{B}_2] \leq q/2$. The result then follows by taking $\mathcal{B} = \mathcal{B}_1 \cup \mathcal{B}_2$ for which $\mathbb{P}[\mathcal{B}] \leq q$, and combining the bounds for both the numerator and denominator.

Next, we consider the case T = 0, and state the approximation bound for this case.

Lemma 19. Consider the noiseless setting (Def. 1). Assume that there exists $\eta > 0$ so that

$$d(s) \le \frac{1}{s^{\beta \mathcal{H}_{-\infty}(\mathbf{X}) + \eta}},\tag{A.161}$$

for all $s \in \mathbb{N}_+$. Then, the likelihood for T = 0 is lower bounded as

$$f\left(x_{1}^{\ell}(1), x_{1}^{\ell}(2); 0\right) \ge \xi(n) \cdot P_{X_{1}^{\ell}}\left(x_{1}^{\ell}(1)\right) P_{X_{1}^{\ell}}\left(x_{1}^{\ell}(2)\right),$$
(A.162)

where $\xi(n) \to 1$ and does not depend on $(x_1^{\ell}(1), x_1^{\ell}(2))$.

Proof: Under the conditions of the lemma, the likelihood is bounded as

$$f\left(x_{1}^{\ell}(1), x_{1}^{\ell}(2); 0\right)$$

$$= \mathbb{P}\left[\left\{X_{1}^{\ell}(1) = x_{1}^{\ell}(1)\right\} \cap \left\{X_{1}^{\ell}(2) = x_{1}^{\ell}(2)\right\} \middle| T = 0\right]$$

$$= \sum_{i_{1}=1}^{n} \sum_{i_{2}=1}^{n} \mathbb{P}\left[\left\{I(1) = i_{1}\right\} \cap \left\{I(2) = i_{2}\right\} \middle| T = 0\right] \times$$

$$\mathbb{P}\left[\left\{X_{1}^{\ell}(1) = x_{1}^{\ell}(1)\right\} \cap \left\{X_{1}^{\ell}(2) = x_{1}^{\ell}(2)\right\} \middle| \left\{I(1) = i_{1}\right\} \cap \left\{I(2) = i_{2}\right\}\right]$$
(A.164)

$$= \frac{1}{n} \sum_{i_1=1}^n \sum_{i_2=1}^n \mathbb{P}\left[I(2) = i_2 \mid T = 0, I(1) = i_1\right] \cdot \mathbb{P}\left[\left\{X_{i_1}^{i_1+\ell-1} = x_1^{\ell}(1)\right\} \cap \left\{X_{i_2}^{i_2+\ell-1} = x_1^{\ell}(2)\right\}\right]$$
(A.165)

$$\stackrel{(a)}{=} \frac{1}{n} \sum_{i_1=1}^n \sum_{i_2=1}^n \mathbb{P}\left[I(2) = i_2 \mid T = 0, I(1) = i_1\right] \cdot \mathbb{P}\left[\left\{X_1^{\ell} = x_1^{\ell}(1)\right\} \cap \left\{X_{i_2-i_1+1}^{i_2-i_1+\ell} = x_1^{\ell}(2)\right\}\right]$$
(A.166)

$$\stackrel{(b)}{\geq} \frac{1}{n} \sum_{i_1=\tau(n)}^{n-\tau(n)} \sum_{i_2=1}^{n} \mathbb{P}\left[I(2) = i_2 \mid T = 0, I(1) = i_1\right] \cdot \mathbb{P}\left[\left\{X_1^{\ell} = x_1^{\ell}(1)\right\} \cap \left\{X_{i_2-i_1+\ell}^{i_2-i_1+\ell} = x_1^{\ell}(2)\right\}\right], \quad (A.167)$$

where (a) follows from stationarity, and in (b) we choose $\tau \equiv \tau(n) = 2\lfloor \frac{n}{2\log n} \rfloor$, which is an even integer. It holds that $\frac{\tau(n)}{\ell} = \frac{\tau(n)}{\beta\log n} = \omega_n(1)$, and so according to the way that I(1) and I(2) are drawn, as described in (16), whenever $\ell \leq I(1) \leq n - \ell + 1$ it holds that $I(2) \sim \text{Uniform}[n]$. So, conditioned on $I(1) = i_1$ and T = 0 for some $i_1 \in \{\tau(n) + 1, \dots, n - \tau(n)\}$, it holds that

$$\mathbb{P}\left[I(2) = i_2 \mid T = 0, I(1) = i_1\right] \sim \text{Uniform}\left[[n] \setminus \{i_1 - \ell + 1, \dots, i_1 + \ell - 1\}\right], \quad (A.168)$$

which is supported over $n - (2\ell - 1) = n_\ell$ possible time indices. So, the inner summation in (A.167) is given by

$$\sum_{i_{2}=1}^{n} \mathbb{P}\left[I(2) = i_{2} \mid T = 0, I(1) = i_{1}\right] \cdot \mathbb{P}\left[\left\{X_{1}^{\ell} = x_{1}^{\ell}(1)\right\} \cap \left\{X_{i_{2}-i_{1}+1}^{i_{2}-i_{1}+\ell} = x_{1}^{\ell}(2)\right\}\right]$$

$$= \frac{1}{n_{\ell}} \sum_{i_{2}=1}^{i_{1}-\ell} \mathbb{P}\left[\left\{X_{1}^{\ell} = x_{1}^{\ell}(1)\right\} \cap \left\{X_{i_{2}-i_{1}+1}^{i_{2}-i_{1}+\ell} = x_{1}^{\ell}(2)\right\}\right] + \frac{1}{n_{\ell}} \sum_{i_{2}=i_{1}+\ell}^{n} \mathbb{P}\left[\left\{X_{1}^{\ell} = x_{1}^{\ell}(1)\right\} \cap \left\{X_{i_{2}-i_{1}+1}^{i_{2}-i_{1}+\ell} = x_{1}^{\ell}(2)\right\}\right].$$
(A.169)

We may lower bound the first sum in the r.h.s. of (A.169) as

$$\frac{1}{n_{\ell}} \sum_{i_{2}=1}^{i_{1}-\ell} \mathbb{P}\left[\left\{X_{1}^{\ell} = x_{1}^{\ell}(1)\right\} \cap \left\{X_{i_{2}-i_{1}+1}^{i_{2}-i_{1}+\ell} = x_{1}^{\ell}(2)\right\}\right]$$

$$\stackrel{(a)}{=} \frac{1}{n_{\ell}} \sum_{s=\ell}^{i_{1}-1} \mathbb{P}\left[\left\{X_{1}^{\ell} = x_{1}^{\ell}(1)\right\} \cap \left\{X_{-s+1}^{-s+\ell} = x_{1}^{\ell}(2)\right\}\right]$$
(A.170)

$$\geq \frac{1}{n_{\ell}} \sum_{s=\ell+\tau/2}^{i_1-1} \mathbb{P}\left[\left\{X_1^{\ell} = x_1^{\ell}(1)\right\} \cap \left\{X_{-s+1}^{-s+\ell} = x_1^{\ell}(2)\right\}\right]$$
(A.171)

$$\stackrel{(b)}{\geq} \frac{1}{n_{\ell}} \sum_{s=\ell+\tau/2}^{i_1-1} \mathbb{P}\left[X_1^{\ell} = x_1^{\ell}(1)\right] \left(\mathbb{P}\left[X_{-s+1}^{-s+\ell} = x_1^{\ell}(2)\right] - d(s-\ell)\right)$$
(A.172)

$$\stackrel{(c)}{\geq} \frac{1}{n_{\ell}} \sum_{s=\ell+\tau/2}^{i_1-1} \mathbb{P}\left[X_1^{\ell} = x_1^{\ell}(1)\right] \left(\mathbb{P}\left[X_{-s+1}^{-s+\ell} = x_1^{\ell}(2)\right] - d(\tau/2)\right)$$
(A.173)

$$\stackrel{(d)}{=} \frac{1}{n_{\ell}} \sum_{s=\ell+\tau/2}^{i_1-1} \mathbb{P}\left[X_1^{\ell} = x_1^{\ell}(1)\right] \left(\mathbb{P}\left[X_1^{\ell} = x_1^{\ell}(2)\right] - d(\tau/2)\right)$$
(A.174)

$$= \left(\frac{i_1 - \ell - \tau/2}{n_\ell}\right) \mathbb{P}\left[X_1^\ell = x_1^\ell(1)\right] \left(\mathbb{P}\left[X_1^\ell = x_1^\ell(2)\right] - d(\tau/2)\right),\tag{A.175}$$

where (a) follows from the change of variables $s = i_1 - i_2$, (b) follows from the strong mixing property, (c)

follows from the assumed monotonicity of d(s), and (d) follows from stationarity.

Similarly, we may lower bound the second sum in the r.h.s. of (A.169) as

$$\frac{1}{n_{\ell}} \sum_{i_{2}=i_{1}+\ell}^{n} \mathbb{P}\left[\left\{X_{1}^{\ell} = x_{1}^{\ell}(1)\right\} \cap \left\{X_{i_{2}-i_{1}+1}^{i_{2}-i_{1}+\ell} = x_{1}^{\ell}(2)\right\}\right]$$

$$\stackrel{(a)}{=} \frac{1}{n_{\ell}} \sum_{s=\ell}^{n-i_{1}} \mathbb{P}\left[\left\{X_{1}^{\ell} = x_{1}^{\ell}(1)\right\} \cap \left\{X_{s+1}^{s+\ell} = x_{1}^{\ell}(2)\right\}\right]$$
(A.176)

$$\stackrel{(b)}{\geq} \frac{1}{n_{\ell}} \sum_{s=\ell+\tau/2}^{n-i_1} \mathbb{P}\left[\left\{X_1^{\ell} = x_1^{\ell}(1)\right\} \cap \left\{X_{s+1}^{s+\ell} = x_1^{\ell}(2)\right\}\right]$$
(A.177)

$$\geq \frac{1}{n_{\ell}} \sum_{s=\ell+\tau/2}^{n-i_1} \mathbb{P}\left[X_1^{\ell} = x_1^{\ell}(1)\right] \left(\mathbb{P}\left[X_{-s+1}^{-s+\ell} = x_1^{\ell}(2)\right] - d(s-\ell)\right)$$
(A.178)

$$\geq \frac{1}{n_{\ell}} \sum_{s=\ell+\tau/2}^{n-i_1} \mathbb{P}\left[X_1^{\ell} = x_1^{\ell}(1)\right] \left(\mathbb{P}\left[X_{-s+1}^{-s+\ell} = x_1^{\ell}(2)\right] - d(\tau/2)\right)$$
(A.179)

$$\geq \left(\frac{n-i_1-\ell-\tau/2+1}{n_\ell}\right) \mathbb{P}\left[X_1^\ell = x_1^\ell(1)\right] \left(\mathbb{P}\left[X_1^\ell = x_1^\ell(2)\right] - d(\tau)\right),\tag{A.180}$$

where (a) follows from the change of variables $s = i_2 - i_1$, and (b) is a non-empty summation (for all n sufficiently large) since $i_1 \le n - \tau$.

Summing the lower bounds on the first and second terms in the r.h.s. of (A.169) leads to the lower bound

$$\sum_{i_{2}=1}^{n} \mathbb{P}\left[I(2) = i_{2} \mid T = 0, I(1) = i_{1}\right] \cdot \mathbb{P}\left[\left\{X_{1}^{\ell} = x_{1}^{\ell}(1)\right\} \cap \left\{X_{i_{2}-i_{1}+1}^{i_{2}-i_{1}+\ell} = x_{1}^{\ell}(2)\right\}\right]$$

$$\geq \left(\frac{i_{1}-\ell-\tau/2}{n_{\ell}} + \frac{n-i_{1}-\ell-\tau/2+1}{n_{\ell}}\right) \cdot \mathbb{P}\left[X_{1}^{\ell} = x_{1}^{\ell}(1)\right] \left(\mathbb{P}\left[X_{1}^{\ell} = x_{1}^{\ell}(2)\right] - d(\tau/2)\right)$$
(A.181)

$$= \left(\frac{n-2\ell-\tau+1}{n_\ell}\right) \mathbb{P}\left[X_1^\ell = x_1^\ell(1)\right] \cdot \left(\mathbb{P}\left[X_1^\ell = x_1^\ell(2)\right] - d(\tau/2)\right)$$
(A.182)

$$= \left(\frac{n-2\ell-2\tau-1}{n_\ell}\right) \mathbb{P}\left[X_1^\ell = x_1^\ell(1)\right] \mathbb{P}\left[X_1^\ell = x_1^\ell(2)\right] \left(1 - \frac{d(\tau/2)}{\mathbb{P}\left[X_1^\ell = x_1^\ell(2)\right]}\right)$$
(A.183)

$$\sim \mathbb{P}\left[X_1^\ell = x_1^\ell(1)\right] \mathbb{P}\left[X_1^\ell = x_1^\ell(2)\right],\tag{A.184}$$

where last step holds since

$$\frac{n - 2\ell - 2\tau - 1}{n_{\ell}} = 1 - O\left(\frac{1}{\log n}\right) \sim 1,$$
(A.185)

and since

$$\frac{d(\tau/2)}{\mathbb{P}\left[X_1^{\ell} = x_1^{\ell}(2)\right]} \leq \frac{d\left(\frac{n}{2\log n}\right)}{\mathbb{P}\left[X_1^{\ell} = x_1^{\ell}(2)\right]}$$
(A.186)

$$\stackrel{(i)}{\leq} \frac{\frac{(2\log n)^{\beta\mathcal{H}_{-\infty}(\mathbf{X})+\eta}}{n^{\beta\mathcal{H}_{-\infty}(\mathbf{X})+\eta}}}{\mathbb{P}\left[X_1^{\ell} = x_1^{\ell}(2)\right]}$$
(A.187)

$$\stackrel{(ii)}{\leq} \frac{\frac{(2\log n)^{\beta\mathcal{H}_{-\infty}(\mathbf{X})+\eta}}{n^{\beta\mathcal{H}_{-\infty}(\mathbf{X})+\epsilon(n)]}}$$
(A.188)

$$=\frac{(2\log n)^{\beta\mathcal{H}_{-\infty}(\mathbf{X})+\eta}}{n^{\beta[\eta-\epsilon(n)]}}$$
(A.189)

$$=o_n(1),\tag{A.190}$$

where, here, (i) follows from the assumption of the lemma, and (ii) holds for some $\epsilon(n) = o_n(1)$.

Before getting into the proof of Theorem 9, we remind the reader the following lower bound on the error probability in binary hypothesis testing. This result will also be used in the proof of Theorem 9.

Lemma 20 (Strong converse on errors in binary hypothesis testing [25, Th. 14.9]). Consider a binary hypothesis testing problem between $H_0: X \sim P$ and $H_1: X \sim Q$. Let $Z \in \{0, 1\}$ be a test based on X. Then, for any $\eta > 0$ it holds that the two types of error must satisfy

$$P[Z=1] + \eta \cdot Q[Z=0] \ge 1 - \mathbb{P}_{X \sim P}\left[\log \frac{\mathrm{d}P}{\mathrm{d}Q}(X) > \log \eta\right].$$
(A.191)

We may now prove Theorem 9.

Proof of Theorem 9: Let us first assume that $\beta > \frac{1}{\mathcal{H}_1(\mathbf{X})}$. As we have seen in the proof of Theorem 6, the possibility of both positive and negative overlaps simply doubles the error probability, and only involves repeating similar arguments for both the positive and negative overlaps. Therefore, in this proof, as well as the rest of the proofs to follow, we slightly simplify the setting, and assume that either no overlap or a positive overlap are possible. That is, we assume that either T = 0 or $T \in [\ell]$, and that

$$P_T(t) = \begin{cases} \frac{1-\ell}{n}, & t = 0\\ \frac{1}{n}, & t \in [\ell] \end{cases}$$
(A.192)

(c.f. (17)).

We consider the following genie-aided detector. Let $t_0 := \lceil \theta_0 \log n \rceil$ and $t_1 := \lfloor \theta_1 \log n \rfloor$ so that $0 < \theta_0 < \theta_1 < \beta$, where

$$\theta_1 = \frac{1 - \theta_0}{\mathcal{H}_1(\mathbf{X})} \tag{A.193}$$

(we take θ_0 as a small constant). The genie provides the detector with side-information S, so that the optimal Bayesian detector is $\hat{T}_{ga}(X_1^{\ell}(1), X_1^{\ell}(2), S)$ (with a slight abuse of notation). Let $\tilde{T}_0 \sim \text{Uniform}([t_1] \setminus [t_0])$. The

side-information is given by

$$S = \begin{cases} \{0, \tilde{T}_0\}, & T = 0\\ \{0, T\}, & T \in [t_1] \setminus [t_0] \\ T, & \text{otherwise} \end{cases}$$
(A.194)

That is, if $T \in \{1, 2, ..., t_0\}$ or $T \in \{t_1 + 1, ..., \ell\}$ then the genie reveals to the detector the exact value of T. Otherwise, it reveals to the detector that the overlap is either zero or a positive overlap in $[t_1] \setminus [t_0]$, where one of these options is the true overlap. We next lower bound the Bayesian error probability of *any* genie-aided detector. First, conditioned on the event $T \in \{1, ..., t_0\} \cup \{t_1 + 1, ..., \ell\}$ an optimal genie-aided detector will decide $\hat{T}_{ga} = S$, and obtain zero error. Second, conditioned on the event $\mathcal{E}_t := \{S = \{0, t\}\}$ for some $t \in [t_1] \setminus [t_0]$, an optimal genie-aided detector will output either 0 or t. Consequently, any optimal genie-aided detector defines a set of $t_1 - t_0$ binary hypothesis testers. Let us focus on a single index $t \in [t_1] \setminus [t_0]$. The probability of the event \mathcal{E}_t is given by

$$\mathbb{P}[\mathcal{E}_t] = \frac{P_T(0)}{t_1 - t_0} + P_T(t) = \frac{1 - \frac{\ell}{n}}{t_1 - t_0} + \frac{1}{n} = \frac{1}{n} \left(\frac{n - \ell + t_1 - t_0}{t_1 - t_0}\right) \sim \frac{1}{t_1 - t_0}.$$
(A.195)

Furthermore, the binary hypothesis testing problem between T = 0 and T = t has priors

$$\pi_t := \mathbb{P}\left[T = t \mid S = \{0, t\}\right] = \frac{t_1 - t_0}{n - \ell + t_1 - t_0} \sim (\theta_1 - \theta_0) \cdot \frac{\log n}{n},\tag{A.196}$$

and $\pi_0 = 1 - \pi_t$. Let \hat{D}_t be any binary test between T = 0 and T = t under the prior $\{\pi_0, \pi_t\}$. So, for the MAP detector

$$P_{\text{error}}^* \stackrel{(a)}{\geq} P_{\text{error}}(\hat{T}_{\text{ga}}) \tag{A.197}$$

$$=\sum_{t=t_0}^{t_1} \mathbb{P}[\mathcal{E}_t] \cdot \mathbb{P}[\hat{T}_{ga} \neq T \mid \mathcal{E}_t] + \mathbb{P}\left[\bigcap_{t=t_0}^{t_1} \mathcal{E}_t^c\right] \cdot \mathbb{P}\left[\hat{T}_{ga} \neq T \mid \bigcap_{t=t_0}^{t_1} \mathcal{E}_t^c\right]$$
(A.198)

$$\stackrel{(b)}{=} \sum_{t=t_0}^{t_1} \mathbb{P}[\mathcal{E}_t] \cdot \mathbb{P}[\hat{T}_{ga} \neq T \mid \mathcal{E}_t]$$
(A.199)

$$\stackrel{(c)}{=} \sum_{t=t_0}^{t_1} \mathbb{P}[\mathcal{E}_t] \cdot \left(\pi_t \mathbb{P}\left[\hat{D}_t = 0 \mid T = t \right] + \pi_0 \cdot \mathbb{P}\left[\hat{D}_t = t \mid T = 0 \right] \right)$$
(A.200)

$$\overset{(d)}{\sim} \frac{1}{t_1 - t_0} \cdot \sum_{t = t_0}^{t_1} \pi_t \mathbb{P}\left[\hat{D}_t = 0 \mid T = t\right] + \pi_0 \cdot \mathbb{P}\left[\hat{D}_t = t \mid T = 0\right]$$
(A.201)

$$\geq \min_{t \in [t_1] \setminus [t_0]} \pi_t \mathbb{P}\left[\hat{D}_t = 0 \mid T = t\right] + \pi_0 \cdot \mathbb{P}\left[\hat{D}_t = t \mid T = 0\right]$$
(A.202)

$$\geq \min_{t \in [t_1] \setminus [t_0]} \pi_t \cdot \left(\mathbb{P}\left[\hat{D}_t = 0 \mid T = t \right] + \frac{\pi_0}{\pi_t} \cdot \mathbb{P}\left[\hat{D}_t = t \mid T = 0 \right] \right), \tag{A.203}$$

where (a) follows since the Bayesian error probability of an optimal genie-aided detector is lower than that of

the MAP detector, (b) holds since conditioned on $\bigcap_{t=t_0}^{t_1} \mathcal{E}_t^c$ it holds that T is known exactly to the genie-aided detector, (c) holds since, conditioned on \mathcal{E}_t , the genie-aided detector is reduced to a binary hypothesis test, and (d) holds since $\mathbb{P}[\mathcal{E}_t] \sim \frac{1}{t_1-t_0}$.

We now invoke Lemma 20 (or [25, Th. 14.9]), by setting therein $\eta = \frac{\pi_0}{\pi_t}$. Let $q \in (0, 1)$ be a small given probability, and let $\nu = \theta_0/8$. We will use Lemma 18 which assures that there exists an event \mathcal{B} such that $\mathbb{P}[\mathcal{B}] \leq q$ and on \mathcal{B}^c it holds that

$$f\left(X_{1}^{\ell}(1), X_{1}^{\ell}(2); t\right) \leq e^{\nu \log n} \cdot \mathbb{1}\left[X_{\ell-t+1}^{\ell}(1) = X_{1}^{t}(2)\right] \cdot P_{X_{1}^{\ell}}\left(X_{1}^{\ell}(1)\right) \cdot P_{X_{1}^{\ell}}\left(X_{1}^{\ell}(2)\right) \cdot \Gamma_{+}(t).$$
(A.204)

We then have that

$$\mathbb{P}\left[\hat{D}_{t}=0 \mid T=t\right] + \frac{\pi_{0}}{\pi_{t}} \cdot \mathbb{P}\left[\hat{D}_{t}=t \mid T=0\right] \\
\stackrel{(a)}{\geq} 1 - \mathbb{P}\left[\log\frac{f\left(X_{1}^{\ell}(1), X_{1}^{\ell}(2); t\right)}{f\left(X_{1}^{\ell}(1), X_{1}^{\ell}(2); 0\right)} > \log\frac{\pi_{0}}{\pi_{t}} \mid T=t\right]$$
(A.205)

$$\stackrel{(b)}{\geq} 1 - \mathbb{P}\left[\log\frac{f\left(X_{1}^{\ell}(1), X_{1}^{\ell}(2); t\right)}{\xi(n)P_{X_{1}^{\ell}}\left(X_{1}^{\ell}(1)\right) \cdot P_{X_{1}^{\ell}}\left(X_{1}^{\ell}(2)\right)} > \log\frac{\pi_{0}}{\pi_{t}} \middle| T = t \right]$$
(A.206)

$$\stackrel{(c)}{=} 1 - \mathbb{P}\left[\{ \mathcal{B}^{c} \} \cap \left\{ \log \frac{f\left(X_{1}^{\ell}(1), X_{1}^{\ell}(2); t\right)}{\xi(n) P_{X_{1}^{\ell}}\left(X_{1}^{\ell}(1)\right) \cdot P_{X_{1}^{\ell}}\left(X_{1}^{\ell}(2)\right)} > \log \frac{\pi_{0}}{\pi_{t}} \right\} \middle| T = t \right]$$

$$- \mathbb{P}\left[\{ \mathcal{B} \} \cap \left\{ \log \frac{f\left(X_{1}^{\ell}(1), X_{1}^{\ell}(2); t\right)}{\xi(n) P_{X_{1}^{\ell}}\left(X_{1}^{\ell}(1)\right) \cdot P_{X_{1}^{\ell}}\left(X_{1}^{\ell}(2)\right)} > \log \frac{\pi_{0}}{\pi_{t}} \right\} \middle| T = t \right]$$

$$(A.207)$$

$$\stackrel{(d)}{\geq} 1 - q - \mathbb{P}\left[\log\frac{\mathbb{1}\left[X_{\ell-t+1}^{\ell}(1) = X_{1}^{t}(2)\right] \cdot \Gamma_{+}(t)}{\xi(n)} + \nu \log n > \log\frac{\pi_{0}}{\pi_{t}} \,\middle| \, T = t\right]$$
(A.208)

$$\stackrel{(e)}{=} 1 - q - \mathbb{P}\left[\log\frac{\Gamma_{+}(t)}{\xi(n)} + \nu\log n > \log\frac{\pi_{0}}{\pi_{t}} \middle| T = t\right]$$
(A.209)

$$\stackrel{(f)}{=} 1 - q - \mathbb{P}\left[\frac{1}{P_{X_1^t}(X_1^t)} > \log \xi(n) - \nu \log n + \log \frac{\pi_0}{\pi_t} \,\middle| \, T = t\right]$$
(A.210)

$$\stackrel{(g)}{\sim} 1 - q,\tag{A.211}$$

where (a) follows from Lemma 20, (b) follows since under the alternative hypothesis T = 0,¹⁰ Lemma 19 implies that

$$f\left(x_{1}^{\ell}(1), x_{1}^{\ell}(2); 0\right) \ge \xi(n) P_{X_{1}^{\ell}}\left(x_{1}^{\ell}(1)\right) \cdot P_{X_{1}^{\ell}}\left(x_{1}^{\ell}(2)\right),$$
(A.212)

(c) follows by using the definition of the event \mathcal{B} , (d) follows by utilizing the bound (A.204) on \mathcal{B} , and $\mathbb{P}[\mathcal{B}] \leq q$, (e) holds since the reading kernel is noiseless, and so under T = t it holds with probability 1 that $X_{\ell-t+1}^{\ell}(1) = X_1^t(2)$, (f) holds from the definition of $\Gamma_+(t)$ and stationarity. Finally, (g) follows from the

¹⁰We name the two hypothesis this way in order to comply with the next use of [25, Th. 14.9].

following derivation, which holds for all n sufficiently large:

$$\mathbb{P}\left[\log\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})} > \log\xi(n) - \nu\log n + \log\frac{\pi_{0}}{\pi_{t}} \middle| T = t\right] \\
\stackrel{(i)}{\leq} \mathbb{P}\left[\log\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})} > \log\xi(n) + (1 - \zeta(n) - \nu)\log n\right]$$
(A.213)

$$= \mathbb{P}\left[\log\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})} - \mathbb{E}\left[\log\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})}\right] > \log\xi(n) + (1 - \zeta(n) - \nu)\log n - \mathbb{E}\left[\log\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})}\right]\right] \quad (A.214)$$

$$\stackrel{(ii)}{\leq} \mathbb{P}\left[\log\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})} - \mathbb{E}\left[\log\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})}\right] > (1 - 2\zeta(n) - \nu)\log n - \mathbb{E}\left[\log\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})}\right]\right]$$
(A.215)

$$\stackrel{(iii)}{\leq} \mathbb{P}\left[\log\frac{1}{P_{X_1^t}\left(X_1^t\right)} - \mathbb{E}\left[\log\frac{1}{P_{X_1^t}\left(X_1^t\right)}\right] > \frac{\theta_0}{8}\log n\right]$$
(A.216)

$$= \mathbb{P}\left[\frac{1}{t}\log\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})} - \frac{1}{t}\mathbb{E}\left[\log\frac{1}{P_{X_{1}^{t}}(X_{1}^{t})}\right] > \frac{\theta_{0}}{8\theta_{1}}\right]$$
(A.217)

$$\stackrel{(iv)}{=} o_n(1), \tag{A.218}$$

where here (i) follows by setting

$$\log \frac{\pi_0}{\pi_t} = \log \frac{1 - \pi_t}{\pi_t} \tag{A.219}$$

$$=\log\frac{n-\ell}{t_1-t_0}\tag{A.220}$$

$$\geq \log \frac{n-\ell}{(\theta_1 - \theta_0)\log n} \tag{A.221}$$

$$= \log n + \log \left(1 - \frac{\ell}{n}\right) - \log(\theta_1 - \theta_0) - \log\log(n)$$
(A.222)

$$\geq \log n + \log \left(1 - \frac{\beta \log n}{n}\right) - \log \beta - \log \log(n)$$
(A.223)

$$=: (1 - \zeta(n)) \log n, \tag{A.224}$$

where $\zeta(n) := \Theta(\frac{\log \log(n)}{\log n}) = o_n(1), (ii)$ follows since $\log \xi(n) \to 0$ whereas $(1 - \zeta(n)) \log n \to \infty$ as $n \to \infty$, (*iii*) follows since for all n sufficiently large

$$\mathbb{E}\left[\log\frac{1}{P_{X_{1}^{t}}\left(X_{1}^{t}\right)}\right] \leq \mathbb{E}\left[\log\frac{1}{P_{X_{1}^{t_{1}}}\left(X_{1}^{t_{1}}\right)}\right]$$
(A.225)

$$\leq t_1 \cdot [\mathcal{H}_1(\mathbf{X}) + \epsilon(n)] \tag{A.226}$$

$$\leq (1 - \theta_0/2) \log n, \tag{A.227}$$

where $\epsilon(n) = o_n(1)$ and $\nu = \frac{\theta_0}{8}$, (iv) follows from the AEP for ergodic sources [6, Th. 16.8.1], which implies that

$$-\frac{1}{t}\log P_{X_1^t}\left(X_1^t\right) \to \mathcal{H}_1(\mathbf{X}) \tag{A.228}$$

with probability 1, as $t \to \infty$, and since any $t \in [t_1] \setminus [t_0]$ satisfies $t > \theta_0 \log n \to \infty$ as $n \to \infty$ and $\frac{\theta_0}{8\theta_1}$ is a

constant.

Overall, we obtain that

$$P_{\text{error}}(\hat{T}) \ge [1 - q - o_n(1)]\pi_t$$
 (A.229)

$$\sim (1-q)(\theta_1 - \theta_0) \frac{\log n}{n} \tag{A.230}$$

$$= (1-q)\left(\frac{1-\theta_0}{\mathcal{H}_1(\mathbf{X})} - \theta_0\right) \cdot \frac{\log n}{n}.$$
 (A.231)

The proof is then completed by noting that both θ_0 and q can be made arbitrarily small, and taking $\theta_0 \downarrow 0$ and $q \downarrow 0$. Finally, in case that $\beta \leq \frac{1}{\mathcal{H}_1(\mathbf{X})}$ a similar proof holds with t_1 chosen as $t_1 \sim \frac{1-\theta_0}{\beta} \cdot \log n$.

APPENDIX B

PROOFS FOR SECTION IV

We begin by proving Lemma 10, which provides an exact expression for the likelihood function in the noisy setting.

Proof of Lemma 10: We only prove the expression for $\Gamma_+(t)$, as the expression for $\Gamma_-(t)$ can be obtained by replacing the role of the two reads. Recall that the pair of sequences may be partitioned into 3 parts. The three parts – the left side of the left fragment, the common fragment, and the right side of the right fragment – are statistically independent conditioned on T. Using this, we may write the likelihood as

$$f\left(y_{1}^{\ell}(1), y_{1}^{\ell}(2); t\right) := P_{Y_{1}^{\ell}(1)Y_{1}^{\ell}(2)|T}\left(y_{1}^{\ell}(1), y_{1}^{\ell}(2) \mid T = t\right)$$

$$= P_{Y^{\ell-t}(1)}\left(y_{1}^{\ell-t}(1)\right) \cdot P_{Y_{t+1}^{\ell}(2)}\left(y_{t+1}^{\ell}(2)\right)$$

$$\times \mathbb{P}\left[Y_{\ell-t+1}^{\ell}(1) = y_{\ell-t+1}^{\ell}(1), Y_{1}^{t}(2) = y^{t}(2) \mid X_{\ell-t+1}^{\ell}(1) = X_{1}^{t}(2)\right].$$
(B.2)

Now, recalling that P_Y is the Y-marginal of $P_X \otimes P_{Y|X}$, since both the source and the channel are memoryless and stationary, the product of the first two terms in (B.2) is

$$P_{Y^{\ell-t}(1)}\left(y_1^{\ell-t}(1)\right) \cdot P_{Y_{t+1}^{\ell}(2)}\left(y_{t+1}^{\ell}(2)\right) = \prod_{i=1}^{\ell-t} P_Y(y_i(1)) \prod_{i=t+1}^{\ell} P_Y(y_i(2)).$$
(B.3)

The third term in (B.2) is given by

$$\mathbb{P}\left[Y_{\ell-t+1}^{\ell}(1) = y_{\ell-t+1}^{\ell}(1), Y_{1}^{t}(2) = y_{1}^{t}(2) \mid X_{\ell-t+1}^{\ell}(1) = X_{1}^{t}(2)\right]$$

$$= \prod_{i=1}^{t} P_{Y\tilde{Y}}(y_{\ell-t+i}(1), y_{i}(2)),$$
(B.4)

due to the independence of the noisy channels generating the reads. Combining (B.3) and (B.4) results the

likelihood

$$\begin{split} f\left(y_{1}^{\ell}(1), y_{1}^{\ell}(2); t\right) \\ &= \prod_{i=1}^{\ell-t} P_{Y}(y_{i}(1)) \prod_{i=1}^{t} P_{Y\tilde{Y}}(y_{\ell-t+i}(1), y_{i}(2)) \prod_{i=t+1}^{\ell} P_{Y}(y_{i}(2)) \\ &= \prod_{i=1}^{\ell-t} P_{Y}(y_{i}(1)) \prod_{i=1}^{t} P_{Y}(y_{\ell-t+i}(1)) \frac{\prod_{i=1}^{t} P_{Y\tilde{Y}}(y_{\ell-t+i}(1), y_{i}(2))}{\prod_{i=1}^{t} P_{Y}(y_{\ell-t+i}(1)) P_{Y}(y_{i}(2))} \prod_{i=1}^{t} P_{Y}(y_{i}(2)) \prod_{i=t+1}^{\ell} P_{Y}(y_{i}(2)) \quad (B.6) \end{split}$$

$$=\prod_{i=1}^{\ell} P_Y(y_i(1)) \times \prod_{i=1}^{t} \lambda(y_{\ell-t+i}(1), y_i(2)) \times \prod_{i=1}^{\ell} P_Y(y_i(2)).$$
(B.7)

Finally, since the first term $\prod_{i=1}^{\ell} P_Y(y_i(1))$ and the last term $\prod_{i=1}^{\ell} P_Y(y_i(2))$ do not depend on t, they can be omitted from the likelihood (without affecting the maximizer), and (90) thus follows.

A. Proof of Theorem 12

Before we prove Theorem 12, we state and prove a few useful lemmas. The first lemma will be used in the analysis of the type-I error probability. It is a reminder from [24, Lemma 47], and is commonly used in *finite blocklength* information theory. This result is based on a change-of-measure argument combined with Berry–Esseen central limit theorem (e.g. [8, Th. 2, Chapter XVI.5]) to obtain strong large-deviations bounds, similar to the one of Bahadur and Rao [2] [7, Th. 3.7.4.] (which has the exact pre-exponent). Compared to Bahadur–Rao bound, this lemma does not have the exact pre-exponent constant, which is anyway immaterial for our analysis. However, it is non-asymptotic, and its applicable to both lattice and non-lattice variables. See [25, Th. 14.5] and [32, Th. 1.3] for a discussion, and how it is used for the higher order terms in capacity expansions. See also [20, Ch. 3] on its relation saddle-point methods and [19], [35] for two additional applications of those techniques. This lemma will be used in the analysis of the type-I error probability.

Lemma 21 (From [24, Lemma 47] (simplified for i.i.d. r.v.'s)). Let $\{Z_i\}_{i \in [n]}$ be i.i.d. r.v's with $\sigma^2 = \mathbb{V}[Z_i] > 0$ and $m_3^3 := \mathbb{E}[|Z_i - \mathbb{E}[Z_i]|^3] < \infty$. Then, for any $a \in \mathbb{R}$

$$\mathbb{E}\left[\exp\left\{-\sum_{i=1}^{n} Z_i\right\} \cdot \mathbb{1}\left(\sum_{i=1}^{n} Z_i > a\right)\right] \le 2 \cdot \left(\frac{\log 2}{\sqrt{2\pi}} + \frac{12m_3^3}{\sigma^2}\right) \frac{1}{\sigma}e^{-a}.$$
(B.8)

The second lemma will be used in the analysis of the type-II error probability, and specifically, to an error event from $T \neq 0$ to T = 0 (misdetection of an overlap). This is a large-deviations result, which is based in Chernoff's bound.

Lemma 22. Let $\{(Y_i, \tilde{Y}_i, \overline{Y}_i)\}_{i \in \mathbb{Z}}$ be a sequence of i.i.d. r.v.'s, distributed as $P_{Y\tilde{Y}} \otimes P_Y$, and assume that

 $I(Y; \tilde{Y}) > 0$. Let a > 0 be given. Then,

$$\mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) \le a \log n_\ell\right] \le \exp\left[-t \cdot E_-^{(t)}(a, P_{XY})\right],\tag{B.9}$$

for

$$E_{-}^{(t)}(a, P_{XY}) := \sup_{\nu \le 1} \frac{(\nu - 1)a \log n_{\ell}}{t} - (\nu - 1) \cdot D_{\nu}(P_{Y\tilde{Y}} \mid\mid P_Y \otimes P_Y),$$
(B.10)

and there exists $E_{-}(0, P_{XY}) > 0$ such that

$$\mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) \le 0\right] \le \exp\left[-t \cdot E_-(0, P_{XY})\right].$$
(B.11)

Finally, there exists $E_+(P_{XY}) > 0$ such that

$$\mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \overline{Y}_i) > 0\right] \le \exp\left[-t \cdot E_+(P_{XY})\right].$$
(B.12)

Proof: It follows from Chernoff's bound that

$$\log \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) \le a \log n_{\ell}\right]$$

$$\leq \log \inf_{\nu \ge 0} \mathbb{E}\left[\exp\left\{\nu a \log n_{\ell} - \nu \sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i)\right\}\right]$$
(B.13)

$$\stackrel{(a)}{=} \log \inf_{\nu \ge 0} n_{\ell}^{\nu a} \prod_{i=1}^{t} \mathbb{E} \left[\frac{P_{Y}^{\nu}(Y_{i}) P_{Y}^{\nu}(\tilde{Y}_{i})}{P_{Y\tilde{Y}}^{\nu}(Y_{i}, \tilde{Y}_{i})} \right]$$
(B.14)

$$= \inf_{\nu \ge 0} \nu a \log n_{\ell} + t \log \sum_{(y,\tilde{y}) \in \mathcal{Y}^{\otimes 2}} P_{Y\tilde{Y}}^{1-\nu}(y,\tilde{y}) P_{Y}^{\nu}(y) P_{Y}^{\nu}(\tilde{y})$$
(B.15)

$$= t \cdot \left(\inf_{\nu \le 1} \frac{(1-\nu)a \log n_{\ell}}{t} + \log \sum_{(y,\tilde{y}) \in \mathcal{Y}^{\otimes 2}} P_{Y\tilde{Y}}^{\nu}(y,\tilde{y}) P_{Y}^{1-\nu}(y) P_{Y}^{1-\nu}(\tilde{y}) \right)$$
(B.16)

$$\stackrel{(b)}{=:} -t \cdot E_{-}^{(t)}(a, P_{XY}), \tag{B.17}$$

where (a) holds since $\{(Y_i, \tilde{Y}_i)\}_{i=1}^t$ are i.i.d., and (b) utilizes the definition of the Rényi divergence in (12). This shows (B.9). Next, we focus on the case a = 0, and show that the infimum, which we denote by $E_-(0, P_{XY})$ is *strictly* negative. To this end, let

$$f(\nu) := \log \sum_{(y,\tilde{y})\in\mathcal{Y}^{\otimes 2}} P^{\nu}_{Y\tilde{Y}}(y,\tilde{y})P^{1-\nu}_{Y}(y)P^{1-\nu}_{Y}(\tilde{y}).$$
(B.18)

Then, f(1) = 0 and

$$f(0) = \log P_Y \otimes P_Y(\operatorname{supp}(P_{Y\tilde{Y}})) \le 0.$$
(B.19)

If f(0) < 0 then (B.9) holds with $E_{-}(0, P_{XY}) \geq -f(0)$. Otherwise, f(0) = f(1) = 0, but we next show

that $f(\nu)$ is strictly convex, and so there exists $\nu_* \in (0,1)$ so that $f(\nu_*) < 0$, and then (B.9) holds with $E_-(0, P_{XY}) \ge -f(\nu_*)$. Indeed, let $P^{(\nu)} \propto P^{\nu}_{Y\tilde{Y}}(y, \tilde{y})P^{1-\nu}_Y(y)P^{1-\nu}_Y(\tilde{y})$ denote a tilted probability measure, and let $\mathbb{E}_{\nu}[\cdot]$ (resp. $\mathbb{V}_{\nu}[\cdot]$) denote expectation (resp. variance) with respect to this measure. Then,

$$\frac{\mathrm{d}}{\mathrm{d}\nu}f(\nu) := \frac{\sum_{(y,\tilde{y})\in\mathcal{Y}^{\otimes 2}} P_Y(y) P_Y(\tilde{y}) \frac{P_{Y\tilde{Y}}(y,\tilde{y})}{P_Y^{\nu}(y) P_Y^{\nu}(\tilde{y})} \log \frac{P_{Y\tilde{Y}}(y,\tilde{y})}{P_Y(y) P_Y(\tilde{y})}}{\sum_{(y,\tilde{y})\in\mathcal{Y}^{\otimes 2}} P_{Y\tilde{Y}}^{\nu}(y,\tilde{y}) P_Y^{1-\nu}(y) P_Y^{1-\nu}(\tilde{y})}$$
(B.20)

$$= \mathbb{E}_{\nu} \left[\log \frac{P_{Y\tilde{Y}}(y,\tilde{y})}{P_{Y}(y)P_{Y}(\tilde{y})} \right], \tag{B.21}$$

and

$$\frac{d^{2}}{d\nu^{2}}f(\nu) := \frac{d}{d\nu} \frac{\sum_{(y,\tilde{y})\in\mathcal{Y}^{\otimes 2}} P_{Y}(y)P_{Y}(\tilde{y})\frac{P_{Y\tilde{Y}}^{\nu}(y,\tilde{y})}{P_{Y}^{\nu}(y)P_{Y}^{\nu}(\tilde{y})}\log\frac{P_{Y\tilde{Y}}(y,\tilde{y})P_{Y}^{\nu}(y,\tilde{y})}{P_{Y}(y)P_{Y}(\tilde{y})P_{Y}^{1-\nu}(y)P_{Y}^{1-\nu}(\tilde{y})} \qquad (B.22)$$

$$= \frac{\sum_{(y,\tilde{y})\in\mathcal{Y}^{\otimes 2}} P_{Y}(y)P_{Y}(\tilde{y})\frac{P_{Y\tilde{Y}}^{\nu}(y,\tilde{y})}{P_{Y}^{\nu}(y)P_{Y}^{\nu}(\tilde{y})}\log^{2}\frac{P_{Y\tilde{Y}}(y,\tilde{y})}{P_{Y}(y)P_{Y}(\tilde{y})}}{\sum_{(y,\tilde{y})\in\mathcal{Y}^{\otimes 2}} P_{Y\tilde{Y}}^{\nu}(y,\tilde{y})P_{Y}^{1-\nu}(y)P_{Y}^{1-\nu}(\tilde{y})} \\
- \frac{\left(\sum_{(y,\tilde{y})\in\mathcal{Y}^{\otimes 2}} P_{Y}(y)P_{Y}(\tilde{y})\frac{P_{Y\tilde{Y}}^{\nu}(y,\tilde{y})}{P_{Y}^{\nu}(y)P_{Y}^{\nu}(\tilde{y})}\log\frac{P_{Y\tilde{Y}}(y,\tilde{y})}{P_{Y}(y)P_{Y}(\tilde{y})}\right)^{2}}{\left(\sum_{(y,\tilde{y})\in\mathcal{Y}^{\otimes 2}} P_{Y\tilde{Y}}^{\nu}(y,\tilde{y})P_{Y}^{1-\nu}(y)P_{Y}^{1-\nu}(\tilde{y})\right)^{2}} \qquad (B.23)$$

$$= \mathbb{E}_{\nu} \left[\log^2 \frac{P_{Y\tilde{Y}}(y,y)}{P_Y(y)P_Y(\tilde{y})} \right] - \mathbb{E}_{\nu}^2 \left[\log \frac{P_{Y\tilde{Y}}(y,y)}{P_Y(y)P_Y(\tilde{y})} \right]$$
(B.24)
$$= \left[\sum_{V \in \tilde{V}} \frac{P_{V\tilde{Y}}(y,\tilde{y})}{P_Y(y)P_Y(\tilde{y})} \right]$$

$$= \mathbb{V}_{\nu} \left[\log \frac{P_{YY}(y, y)}{P_Y(y) P_Y(\tilde{y})} \right]$$
(B.25)

$$> 0,$$
 (B.26)

where the last inequality holds since due to the assumption that $I(Y; \tilde{Y}) > 0$ it holds that Y and \tilde{Y} are not statistically independent, and so $\log[P_{Y\tilde{Y}}(y,\tilde{y})/P_Y(y)P_Y(\tilde{y})]$ is not a constant under the tilted measure (note that since $P_Y \otimes P_Y \gg P_{Y\tilde{Y}}$, the tilted measure $P^{(\nu)}$ has the same support as $P_{Y\tilde{Y}}$).

Similarly, it follows from Chernoff's bound that

$$\log \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \overline{Y}_i) > 0\right] \le \log \inf_{\nu \ge 0} \mathbb{E}\left[\exp\left\{\nu \sum_{i=1}^{t} \log \lambda(Y_i, \overline{Y}_i)\right\}\right]$$
(B.27)

$$= \log \inf_{\nu \ge 0} \prod_{i=1}^{t} \mathbb{E} \left[\frac{P_{Y\tilde{Y}}^{\nu}(Y_i, \overline{Y}_i)}{P_Y^{\nu}(Y_i) P_Y^{\nu}(\overline{Y}_i)} \right]$$
(B.28)

$$= t \cdot \inf_{\nu \ge 0} \log \sum_{(y,\overline{y}) \in \mathcal{Y}^{\otimes 2}} P^{\nu}_{Y\tilde{Y}}(y,\overline{y}) P^{1-\nu}_{Y}(y) P^{1-\nu}_{Y}(\overline{y}).$$
(B.29)

$$= t \cdot \inf_{\nu \le 1} \log \sum_{(y,\overline{y}) \in \mathcal{Y}^{\otimes 2}} P_Y^{\nu}(y) P_Y^{\nu}(\overline{y}) P_{Y\tilde{Y}}^{1-\nu}(y,\overline{y}), \tag{B.30}$$

which is similar to the previous Chernoff-based bound, albeit with the roles of $P_{Y\tilde{Y}}$ and $P_Y \otimes P_Y$ interchanged.

Defining similarly

$$g(\nu) := \log \sum_{(y,\overline{y})\in\mathcal{Y}^{\otimes 2}} P_Y^{\nu}(y) P_Y^{\nu}(\overline{y}) P_{Y\tilde{Y}}^{1-\nu}(y,\overline{y}), \tag{B.31}$$

it holds that g(1) = 0 and

$$g(0) = \log P_{Y\tilde{Y}}(\operatorname{supp}(P_Y \otimes P_Y)) = 0$$
(B.32)

since $P_Y \otimes P_Y \gg P_{Y\tilde{Y}}$. As before $g(\nu)$ is strictly convex, and so there exists $\nu_* \in (0,1)$ so that $g(\nu_*) < 0$, and then (B.12) holds with $E_+(P_{XY}) \ge -g(\nu_*)$.

The next lemma will also be used to analyze the type-II error probability, however, it will be used for the case that the true overlap is either overestimated or underestimated.

Lemma 23. It holds that

$$\mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_{k+i}, \tilde{Y}_i) > 0\right] \le 2 \exp\left[-(t-1) \cdot \frac{E_+(P_{XY})}{2}\right].$$
(B.33)

Proof: In general, the r.v.'s $\{\lambda(Y_{k+i}, \tilde{Y}_i)\}_{i=1}^t$ are possibly statistically dependent, due to the dependence between Y_i and \tilde{Y}_i for any given *i*. Following [22, Lemma 16], we note that [t] can be partitioned into two sets I_1, I_2 of almost equal size $(I_1 \cup I_2 = [t], I_1 \cap I_2 = \emptyset)$, for which the Y_i 's and \tilde{Y}_i 's involved in each sum are all statistically independent. Specifically, iteratively construct I_1 as follows: Initializing $I_1^{(0)} = \emptyset$, at the *r*th iteration, set

$$I_1^{(r)} \leftarrow I_1^{(r-1)} \cup \{\{r, 2k+r, \ldots\} \cap [t]\}$$
(B.34)

and

$$I_2^{(r)} \leftarrow I_2^{(r-1)} \cup \{\{k+r, 3k+r, \ldots\} \cap [t]\}.$$
 (B.35)

The construction halts at the first iteration r_* for which $|I_1^{(r_*)}| \in \{t/2 - 1, t/2\}$.

For simplicity, let us assume that t is odd, and so $|I_1| = (t-1)/2$ and $|I_2| = (t+1)/2$. The case of even t is similar. Then,

$$\mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_{k+i}, \tilde{Y}_i) > 0\right] = \mathbb{P}\left[\sum_{i \in I_1} \log \lambda(Y_{k+i}, \tilde{Y}_i) + \sum_{i \in I_2} \log \lambda(Y_{k+i}, \tilde{Y}_i) > 0\right]$$
(B.36)

$$\stackrel{(a)}{\leq} \mathbb{P}\left[\sum_{i \in I_1} \log \lambda(Y_{k+i}, \tilde{Y}_i) > 0\right] + \mathbb{P}\left[\sum_{i \in I_2} \log \lambda(Y_{k+i}, \tilde{Y}_i) > 0\right]$$
(B.37)

$$\stackrel{(b)}{=} \mathbb{P}\left[\sum_{i \in I_1} \log \lambda(Y_i, \overline{Y}_i) > 0\right] + \mathbb{P}\left[\sum_{i \in I_2} \log \lambda(Y_i, \overline{Y}_i) > 0\right]$$
(B.38)

$$\stackrel{(c)}{\leq} 2 \exp\left[-(t-1) \cdot \frac{E_+(P_{XY})}{2}\right] \tag{B.39}$$

where (a) follows from the union bound (at least one of the sums must be positive), (b) follows since within I_1 ,

 $\{Y_{k+i}, \tilde{Y}_i\}_{i \in I_1}$ is a collection of t statistically independent r.v.'s, all distributed according to P_Y , and similarly for I_2 , and (c) follows from Lemma 22.

Using the above lemmas, we may now prove Theorem 12,

Proof of Theorem 12: As can be discerned from the proof of Theorem 6, and as discussed in the beginning of the proof of Theorem 9, the analysis of positive and negative overlaps is analogous. Thus, in order to simplify the analysis, we only consider positive overlaps, and then double the error to account for negative overlaps, at the end of the proof. As also discussed in the proof of Theorem 6, the case $\beta < \frac{1}{I(Y;\tilde{Y})}$ is trivial, since this is the error of a detector that never decides on an overlap. We thus focus on the opposite case. With a slight abuse of notation, we let $\{(Y_i, \tilde{Y}_i, \overline{Y}_i)\}_{i \in \mathbb{Z}}$ be a sequence of i.i.d. r.v.'s, distributed as $P_{Y\tilde{Y}} \otimes P_Y$. We also let $\hat{T} \equiv \hat{T}_{MAP}$ be the optimal MAP detector.

Analysis of the type-I error probability:

We begin by analyzing $p_{type-I}(\hat{T})$. By the union bound and Prop. 10

$$p_{\text{type-I}}(\hat{T}) = \mathbb{P}\left[\hat{T} > 0 \mid T = 0\right]$$
(B.40)

$$= \mathbb{P}\left[\bigcup_{t=t_{\mathsf{MDO}}}^{\ell} \left\{\sum_{i=1}^{t} \log \lambda\left(Y_{\ell-t+i}(1), Y_{i}(2)\right) \ge \log n_{\ell}\right\} \middle| T = 0\right]$$
(B.41)

$$\leq \sum_{t=t_{\text{MDO}}}^{\ell} \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda\left(Y_{\ell-t+i}(1), Y_{i}(2)\right) \geq \log n_{\ell} \middle| T=0\right].$$
(B.42)

Fix any $t \ge t_{MDO}$. To bound each of the probabilities in (B.42), we will use the truncated MGF bound of Lemma 21. To this end, let

$$m_3^3 := \mathbb{E}\left[\left|\log\lambda(Y_i, \tilde{Y}_i) - \mathbb{E}[\log\lambda(Y_i, \tilde{Y}_i)]\right|^3\right] < \log\lambda_{\max}^3 < \infty,$$
(B.43)

and let $\sigma^2 := \mathbb{V}[\log \lambda(Y_i, \tilde{Y}_i)]$, where $\sigma^2 > 0$ since the channel $P_{Y|X}$ is not completely noiseless and so Yand \tilde{Y} are not independent (as $\frac{1}{I(Y,\tilde{Y})} \leq \beta < \infty$). Hence, it does not hold that $P_{Y\tilde{Y}}(y, \tilde{y}) = P_Y(y)P_Y(\tilde{y})$ with probability 1. In addition, conditioned on T = 0, it holds that $\{(Y_{\ell-t+i}(1), Y_i(2))\}_{i \in [t]} \stackrel{d}{=} \{(Y_i, \overline{Y}_i)\}_{i \in [t]}$. So, for any fixed $t \geq t_{\text{MDO}}$, any of the probabilities in the sum (B.42) is bounded as

$$\mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \overline{Y}_i) \ge \log n_{\ell}\right]$$
$$= \sum_{(y^t, \overline{y}^t) \in [\mathcal{Y}^{\otimes 2}]^t} \prod_{i=1}^{t} P_Y(y_i) P_Y(\overline{y}_i) \cdot \mathbb{1}\left[\sum_{i=1}^{t} \log \lambda(y_i, \overline{y}_i) \ge \log n_{\ell}\right]$$
(B.44)

$$\stackrel{(a)}{=} \sum_{(y^t, \overline{y}^t) \in [\mathcal{Y}^{\otimes 2}]^t} P_{Y_1^t \tilde{Y}_1^t}(y^t, \overline{y}^t) \cdot \exp\left[-\sum_{i=1}^t \log \lambda(y_i, \overline{y}_i)\right] \mathbb{1}\left[\sum_{i=1}^t \log \lambda(y_i, \overline{y}_i) \ge \log n_\ell\right]$$
(B.45)

$$= \mathbb{E}\left[\exp\left[-\sum_{i=1}^{t}\log\lambda(Y_i, \tilde{Y}_i)\right] \mathbb{1}\left[\sum_{i=1}^{t}\log\lambda(Y_i, \tilde{Y}_i) \ge \log n_\ell\right]\right]$$
(B.46)

$$\stackrel{(b)}{\leq} 2\left(\frac{\log 2}{2} + \frac{12m_3^3}{2}\right) \frac{\exp(-\log n_\ell)}{2} \tag{B.47}$$

$$= \tilde{a} \left(\sqrt{2\pi\sigma^2} + \sigma^3 \right) \qquad \sqrt{t}$$

$$\leq \tilde{a}(P_{XY}, \beta) \cdot \frac{1}{\sqrt{\log n}} \frac{1}{n_\ell},$$
(B.48)

for some constant $\tilde{a}(P_{XY},\beta)$ that does not depend on n, where (a) follows since

$$\exp\left[-\sum_{i=1}^{t}\log\lambda(y_i,\overline{y}_i)\right] = \prod_{i=1}^{t}\frac{P_Y(y_i)P_Y(\overline{y}_i)}{P_{Y\tilde{Y}}(y_i,\overline{y}_i)},\tag{B.49}$$

and (b) follows from Lemma 21. Plugging this back to (B.42) results for $\ell = \beta \log n$

$$p_{\text{type-I}}(\hat{T}) \le (\ell - t_{\text{MDO}})\tilde{a}(P_{XY}, \beta) \cdot \frac{1}{\sqrt{\log n}} \frac{1}{n}$$
(B.50)

$$\leq \ell \tilde{a}(P_{XY},\beta) \cdot \frac{1}{\sqrt{\log n}} \frac{1}{n}$$
(B.51)

$$=a(P_{XY},\beta)\frac{\sqrt{\log n}}{n},\tag{B.52}$$

with $a(P_{XY},\beta) = \beta \cdot \tilde{a}(P_{XY},\beta)$, as claimed.

Analysis of the type-II error probability:

We next turn to analyze $p_{\text{type-II}}(\hat{T};t)$. Let $\epsilon > 0$ be fixed. To this end, we will use the Chernoff-bound based analysis carried in Lemma 22. It is shown therein that for any $t = \theta \log n_{\ell}$ with $\frac{1}{\log \lambda_{\text{max}}} \le \theta \le \beta$ (this domain is not empty since $\frac{1}{\log \lambda_{\text{max}}} \le \frac{1}{I(Y;\tilde{Y})} \le \beta$) it holds that

$$\mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) \le \log n_\ell\right] \le \exp\left[-t \cdot E_-^{(t)}(1, P_{XY})\right],\tag{B.53}$$

where

$$E_{-}^{(t)}(1, P_{XY}) := \sup_{\nu \le 1} (\nu - 1) \cdot \left[\frac{\log n_{\ell}}{t} - D_{\nu}(P_{Y\tilde{Y}} || P_{Y} \otimes P_{Y}) \right].$$
(B.54)

So, the negative log-probability in (B.53) is lower bounded by $\log n_{\ell} \cdot A(\theta)$ where

$$A(\theta) := \sup_{\nu \le 1} f(\nu) := \sup_{\nu \le 1} (\nu - 1) - (\nu - 1)\theta \cdot D_{\nu}(P_{Y\tilde{Y}} || P_Y \otimes P_Y),$$
(B.55)

and where $f(\nu)$ was implicitly defined. In addition, $\theta \to A(\theta)$ is monotonically non-decreasing. Indeed, this will follow if we show that the supremum in $A(\theta)$ is always achieved at $\nu \in [0,1]$ since¹¹

$$(\nu - 1)D_{\nu}(P_{Y\tilde{Y}} \parallel P_Y \otimes P_Y) \le 0 \tag{B.56}$$

for all $\nu \in [0,1]$ (which holds since $D_{\nu}(P_{Y\tilde{Y}} || P_Y \otimes P_Y) \ge 0$ for $\nu \ge 0$ [34, Th. 8]). To show this property,

¹¹Anyway, we may always restrict $\nu \in [0, 1]$ in the Chernoff bound (B.17) of Lemma 22. We show here that this does not compromises the bound.

note that f(1) = 0 and, as in the proof of Lemma 22,

$$\left. \frac{\partial}{\partial \nu} f(\nu) \right|_{\nu=0} = \theta \cdot D_1(P_Y \otimes P_Y \mid\mid P_{Y\tilde{Y}}) > 0, \tag{B.57}$$

so that $\nu \mapsto f(\nu)$ is increasing on $\nu = 0$ and

$$\frac{\partial^2}{\partial\nu^2} f(\nu) = -\theta \cdot \mathbb{V}_{\nu} \left[\log \frac{P_{Y\tilde{Y}}(y,\tilde{y})}{P_Y(y)P_Y(\tilde{y})} \right] < 0, \tag{B.58}$$

so that $\nu \mapsto f(\nu)$ is strictly concave. From the above three properties, its supremum over $\nu \leq 1$ occurs in [0, 1]. Accordingly, we may define θ_{ϵ}^* as the minimal value of θ such that $A(\theta) \geq \epsilon$, that is,

$$\theta_{\epsilon}^* \equiv \theta_{\epsilon}^*(P_{XY}) := \min\left\{\theta \in \left(\frac{1}{\log \lambda_{\max}}, \beta\right) : A(\theta) \ge \epsilon\right\}$$

$$(B.59)$$

$$1 - \nu - \epsilon$$

$$= \min_{0 \le \nu \le 1} \frac{1 - \nu - \epsilon}{(1 - \nu) \cdot D_{\nu}(P_{Y\tilde{Y}} \parallel P_Y \otimes P_Y)},$$
(B.60)

where the last equality follows from the following consideration: If $\theta \ge \theta_{\epsilon}^*$, then there exists $0 \le \nu \le 1$ such that

$$(\nu - 1) - (\nu - 1)\theta \cdot D_{\nu}(P_{Y\tilde{Y}} \mid\mid P_Y \otimes P_Y) \ge \epsilon$$
(B.61)

which is equivalent to

$$\theta \ge \frac{1 - \nu - \epsilon}{(1 - \nu)D_{\nu}(P_{Y\tilde{Y}} \mid\mid P_Y \otimes P_Y)}$$
(B.62)

as $D_{\nu}(P_{Y\tilde{Y}} || P_Y \otimes P_Y) > 0$ for $\nu \ge 0$ [34, Th. 8], and so θ_{ϵ}^* is as claimed.

Let $t_{\epsilon}^* := \lceil \theta_{\epsilon}^* \log n_{\ell} \rceil$. Then, if $t \leq t_{\epsilon}^*$ we trivially bound the error by 1. For $t > t_{\epsilon}^*$ we decompose the error probability into three terms, as

$$p_{\text{type-II}}(\hat{T}_{\text{MAP}};t) = \mathbb{P}\left[\hat{T} \neq t \mid T = t\right]$$
(B.63)

$$= \mathbb{P}\left[\hat{T} = 0 \mid T = t\right] + \sum_{k=1}^{t} \mathbb{P}\left[\hat{T} = t - k \mid T = t\right] + \sum_{k=1}^{\ell-t} \mathbb{P}\left[\hat{T} = t + k \mid T = t\right], \quad (B.64)$$

that is, an error to $\hat{T} = 0$, an error that underestimate T (but not zero), and an error that overestimate T. We next analyze each of these terms separately.

First, for an error to $\hat{T} = 0$, it holds that

$$\mathbb{P}\left[\hat{T}=0 \mid T=t\right]$$

$$\stackrel{(a)}{=} \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_{\ell-t+i}(1), Y_i(2)) \le \log n_{\ell} \mid T=t\right]$$
(B.65)

$$\stackrel{(b)}{=} \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) \le \log n_\ell\right]$$
(B.66)

$$\stackrel{(c)}{\leq} \exp\left[-A(\theta) \cdot \log n_{\ell}\right] \tag{B.67}$$

$$\stackrel{(d)}{\leq} \exp\left[-\epsilon \log n_{\ell}\right] \tag{B.68}$$

$$= \exp\left[-\epsilon \log(n-\ell)\right] \tag{B.69}$$

$$=\frac{1}{(n-\ell)^{\epsilon}} \tag{B.70}$$

$$=\frac{1}{n^{\epsilon}} \cdot \frac{1}{\left[1 - \frac{\beta \log n}{n}\right]^{\epsilon}}$$
(B.71)

$$\sim \frac{1}{n^{\epsilon}},$$
 (B.72)

where (a) follows from Prop. 10, (b) follows since conditioned on T = t, it holds that $Y_{\ell-t+i}(1) - X_{\ell-t+i} - Y_i(2)$ for $i \in [t]$ and that $\{Y_{\ell-t+i}(1) - X_{\ell-t+i} - Y_i(2)\}_{i \in [t]}$ are i.i.d., hence $\{Y_{\ell-t+i}(1), Y_i(2)\}_{i \in [t]} \stackrel{d}{=} \{(Y_i, \tilde{Y}_i)\}_{i \in [t]}, (c)$ follows from the statements made in the discussion after (B.53), and (d) follows since $\theta \to A(\theta)$ is monotonic non-decreasing, and since $\theta \ge \theta_{\epsilon}^*$.

Second, for an error to $\hat{T} = t - k$ for some $k \in \{1, \dots, t - t_{\text{MDO}}\}$, it holds that

$$\mathbb{P}\left[\hat{T} = t - k \mid T = t\right]
\stackrel{(a)}{\leq} \mathbb{P}\left[\log \frac{\prod_{i=1}^{t-k} \lambda(Y_{\ell-t+k+i}(1), Y_i(2))}{\prod_{i=1}^{t} \lambda(Y_{\ell-t+i}(1), Y_i(2))} \ge 0 \mid T = t\right]$$
(B.73)

$$\stackrel{(b)}{=} \mathbb{P}\left[\log\frac{\prod_{i=1}^{t-k}\lambda(Y_{k+i},\tilde{Y}_i)}{\prod_{i=1}^{t}\lambda(Y_i,\tilde{Y}_i)} \ge 0\right]$$
(B.74)

$$= \mathbb{P}\left[\sum_{i=1}^{t-k} \log \lambda(Y_{k+i}, \tilde{Y}_i) \ge \sum_{i=1}^t \log \lambda(Y_i, \tilde{Y}_i)\right]$$
(B.75)

$$\stackrel{(c)}{\leq} \mathbb{P}\left[\left\{\sum_{i=1}^{t-k} \log \lambda(Y_{k+i}, \tilde{Y}_i) > 0\right\} \cup \left\{\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) < 0\right\}\right]$$
(B.76)

$$\overset{(d)}{\leq} \mathbb{P}\left[\sum_{i=1}^{t-k} \log \lambda(Y_{k+i}, \tilde{Y}_i) > 0\right] + \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) < 0\right]$$
(B.77)

$$\stackrel{(e)}{=} \mathbb{P}\left[\sum_{i=1}^{t-k} \log \lambda(Y_i, \overline{Y}_i) > 0\right] + \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) < 0\right]$$
(B.78)

$$\stackrel{(f)}{\leq} \exp\left[-\frac{(t-k-1)}{2} \cdot E_+\right] + \exp\left[-t \cdot E_0\right] \tag{B.79}$$

$$\stackrel{(g)}{\leq} \exp\left[-\frac{E_{+}}{2\log\lambda_{\max}}\log n_{\ell}\right] + \exp\left[-\frac{E_{0}}{\log\lambda_{\max}}\log n_{\ell}\right]$$
(B.80)

$$\stackrel{(n)}{\leq} \frac{1}{n^{\eta(P_{XY},\beta)}},\tag{B.81}$$

where (a) follows from Prop. 10, (b) follows since, conditioned on T = t, it holds that

$$\{Y_{\ell-t+k+i}(1), Y_i(2), Y_{\ell-t+i}(1)\}_{i \in [t]} \stackrel{d}{=} \{(Y_{i+k}, \tilde{Y}_i, Y_i)\}_{i \in [t]},$$
(B.82)

(c) follows since for any pair or r.v.'s (U, V) it holds for any $r \in \mathbb{R}$ that

$$\mathbb{P}[U \ge V] \le \mathbb{P}\left[\{U \ge r\} \cup \{V \le r\}\right] \tag{B.83}$$

and by choosing r = 0, (d) follows from the union bound, (e) follows since $\{(Y_{k+i}, \tilde{Y}_i)\}_{i \in [t-k]} \stackrel{d}{=} \{(Y_i, \overline{Y}_i)\}_{i \in [t-k]}$, (f) follows from the Chernoff-bound analysis in Lemma 22, for some strictly positive constants $E_0 \equiv E_-(0, P_{XY}) > 0$ and $E_+ \equiv E_+(P_{XY}) > 0$, and from Lemma 23, (g) follows since $t - k \ge t_{\text{MDO}} = \frac{\log n_\ell}{\log \lambda_{\text{max}}}$ and $t > t_{\epsilon}^* \ge t_{\text{MDO}}$ both hold, and (h) holds for some constant $\eta(P_{XY}, \beta) > 0$.

Third, and similarly to the previous case, for an error to $\hat{T} = t + k$ for some $k \in \{1, \dots, \ell - t\}$, it holds that

$$\mathbb{P}\left[\hat{T} = t + k \mid T = t\right]$$

$$\stackrel{(a)}{=} \mathbb{P}\left[\log \frac{\prod_{i=1}^{t+k} \lambda(Y_{\ell-t-k+i}(1), Y_i(2))}{\prod_{i=1}^{t} \lambda(Y_{\ell-t+i}(1), Y_i(2))} \ge 0 \mid T = t\right]$$
(B.84)

$$\stackrel{(b)}{=} \mathbb{P}\left[\log\frac{\prod_{i=1}^{t+k}\lambda(Y_{i-k},\tilde{Y}_i)}{\prod_{i=1}^{t}\lambda(Y_i,\tilde{Y}_i)} \ge 0\right]$$
(B.85)

$$= \mathbb{P}\left[\sum_{i=1}^{t+k} \log \lambda(Y_{i-k}, \tilde{Y}_i) \ge \sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i)\right]$$
(B.86)

$$\leq \mathbb{P}\left[\sum_{i=1}^{t+k} \log \lambda(Y_{i-k}, \tilde{Y}_i) \ge 0\right] + \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) < 0\right]$$
(B.87)

$$= \mathbb{P}\left[\sum_{i=1}^{t+k} \log \lambda(Y_i, \overline{Y}_i) \ge 0\right] + \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) < 0\right]$$
(B.88)

$$\leq \exp\left[-\frac{(t+k-1)}{2} \cdot E_{+}\right] + \exp\left[-t \cdot E_{-}\right]$$
(B.89)

$$\leq \exp\left[-\frac{E_{+}}{2\log\lambda_{\max}}\log n_{\ell}\right] + \exp\left[-\frac{E_{-}}{\log\lambda_{\max}}\log n_{\ell}\right]$$
(B.90)

$$\leq \frac{1}{n^{\eta(P_{XY},\beta)}},\tag{B.91}$$

using similar reasoning as in the previous case.

Thus, $p_{\text{type-II}}(\hat{T};t) \leq 1$ if $t \leq t_{\epsilon}^*$ and the decomposition (B.64) and the bounds (B.72) (B.81) and (B.91) result in that for $t \geq t_{\epsilon}^*$

$$p_{\text{type-II}}(\hat{T};t) \le \frac{1}{n^{\epsilon/2}} + \frac{\beta \log n}{n^{\eta(P_{XY},\beta)}} \le \frac{1}{n^{\epsilon/4}}$$
(B.92)

for all $n \ge n_0(P_{XY}, \beta, \epsilon)$. This completes the analysis of $p_{\text{type-II}}(\hat{T}; t)$.

To complete the proof, we bound the Bayesian error probability as

$$P_{\text{error}}^* = \sum_{t=0}^{\ell} P_T(t) \cdot \mathbb{P}\left[\hat{T} \neq T \mid T = t\right]$$
(B.93)

$$=P_T(0) \cdot \mathbb{P}\left[\hat{T} > 0 \mid T=0\right] + \sum_{t=1}^{t_{\epsilon}^*} P_T(t) \cdot \mathbb{P}\left[\hat{T} \neq T \mid T=t\right] + \sum_{t=t_{\epsilon}^*+1}^{\ell} P_T(t) \cdot \mathbb{P}\left[\hat{T} \neq T \mid T=t\right]$$
(B.94)

$$\leq \mathbb{P}\left[\hat{T} > 0 \mid T = 0\right] + \frac{t_{\epsilon}^*}{n} + \frac{\ell - t_{\epsilon}^*}{n} \max_{t \in \{t_{\epsilon}^* + 1, \dots, \ell\}} \mathbb{P}\left[\hat{T} \neq T \mid T = t\right]$$
(B.95)

$$\leq a(P_{XY},\beta)\frac{\sqrt{\log n_{\ell}}}{n} + \theta_{\epsilon}^* \frac{\log n_{\ell}}{n} + \frac{\beta \log n_{\ell}}{n^{1+\epsilon/4}}$$
(B.96)

for all $n \ge n_0(P_{XY}, \beta, \epsilon)$. The result is then completed since $\epsilon > 0$ is arbitrary, and since $\alpha \to D_\alpha(P_{Y\tilde{Y}} || P_Y \otimes P_Y)$ is monotonic non-decreasing [34, Summary table], it holds that

$$\theta_{\epsilon}^* \to \frac{1}{D(P_{Y\tilde{Y}} \parallel P_Y \otimes P_Y)} = \frac{1}{I(Y; \tilde{Y})}$$
(B.97)

as $\epsilon \downarrow 0$.

B. Proof of Theorem 13

The proof of Theorem 13 follows similar lines to the proof of Theorem 9, and so we just highlight the differences.

Proof of Theorem 13: We use the same argument as in the proof of Theorem 9, and focus on the case that $\frac{1}{I(Y;\tilde{Y})} \leq \beta$ (the opposite case is proved similarly, and even simpler). Using the notation therein, we obtain

$$P_{\text{error}}(\hat{T}) \stackrel{(a)}{\geq} \min_{t \in [t_1] \setminus [t_0]} \pi_t \cdot \left(\mathbb{P}\left[\hat{D}_t = 0 \mid T = t \right] + \frac{\pi_0}{\pi_t} \cdot \mathbb{P}\left[\hat{D}_t = t \mid T = 0 \right] \right) \tag{B.98}$$

$$\stackrel{(b)}{\geq} \min_{t \in [t_1] \setminus [t_0]} \pi_t \cdot \left(1 - \mathbb{P}\left[\sum_{i=1}^t \log \lambda(Y_i, \tilde{Y}_i) > \log \frac{\pi_0}{\pi_t} \right] \right)$$
(B.99)

$$\stackrel{(c)}{\sim} \pi_t, \tag{B.100}$$

where (a) follows using the arguments that lead to (A.203) (proof of Theorem 9), (b) follows by setting $\eta = \frac{\pi_0}{\pi_t}$ in Lemma 20 (or [25, Theorem 14.9]), and as the hypothesis depends on $Y_1^{\ell}(1)$ and $Y_1^{\ell}(2)$ only through their possibly overlapping part. Under the null hypothesis T = t, and each distribution of joint letters is $P_{Y\tilde{Y}}$. Under the alternative the joint distribution of every pair of letters is $P_Y \otimes P_Y$. Finally, (c) holds for any $t \in [t_1] \setminus [t_0]$ due to the following derivation:

$$\mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) > \log \frac{\pi_0}{\pi_t}\right] \\
\stackrel{(i)}{\leq} \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) > (1 - \zeta(n)) \log n\right]$$
(B.101)

$$\leq \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) - \mathbb{E}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i)\right] > (1 - \zeta(n)) \log n - \mathbb{E}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i)\right]\right]$$
(B.102)

64

$$\stackrel{(ii)}{\leq} \mathbb{P}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i) - \mathbb{E}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i)\right] > \frac{\theta_0}{2} \log n\right]$$
(B.103)

$$= \mathbb{P}\left[\frac{1}{t}\sum_{i=1}^{t}\log\lambda(Y_i, \tilde{Y}_i) - \mathbb{E}\left[\frac{1}{t}\sum_{i=1}^{t}\log\lambda(Y_i, \tilde{Y}_i)\right] > \frac{\theta_0}{2\theta_1}\right]$$
(B.104)

$$\stackrel{(iii)}{\leq} \frac{4\theta_1^2 \mathbb{V}[\log \lambda(Y_i, Y_i)]}{\theta_0^2} \cdot \frac{1}{t}$$
(B.105)

$$=o_n(1), (B.106)$$

where here (i) follows by setting, exactly as in (A.224),

$$\log \frac{\pi_0}{\pi_t} = (1 - \zeta(n)) \log n,$$
(B.107)

(ii) follows by choosing

$$t_1 = \frac{(1-\theta_0)}{I(Y;\tilde{Y})} \log n \tag{B.108}$$

so that for any $t \in [t_0]$

$$\mathbb{E}\left[\sum_{i=1}^{t} \log \lambda(Y_i, \tilde{Y}_i)\right] = t \cdot I(Y; \tilde{Y}) \le (1 - \theta_0) \log n,$$
(B.109)

and $\theta_0 - \zeta(n) \ge \theta_0/2$, (*iii*) follows from Chebyshev's inequality, as $\mathbb{V}[\log \lambda(Y_i, \tilde{Y}_i)] < \infty$ (where the latter holds since $0 < \lambda_{\min} \le \lambda(Y, \tilde{Y}) \le \lambda_{\max} < \infty$ with probability 1). Overall, we obtain that

$$P_{\text{error}}(\hat{T}) \ge [1 - o_n(1)]\pi_t$$
 (B.110)

$$\sim \frac{(1-\theta_0/2)}{I(Y;\tilde{Y})} \frac{\log n}{n},\tag{B.111}$$

and the claim of the proof follows by taking $\theta_0 \downarrow 0$.

REFERENCES

- R. Arratia, D. Martin, G. Reinert, and M. S. Waterman. Poisson process approximation for sequence repeats, and sequencing by hybridization. *Journal of Computational Biology*, 3(3):425–463, 1996. 2
- [2] R. R. Bahadur and R. R. Rao. On deviations of the sample mean. Ann. Math. Statist., 31(4):1015–1027, December 1960. 7, 27, 54
- [3] T. Z. Baharav, G. M. Kamath, D. N. C. Tse, and I. Shomorony. Spectral Jaccard similarity: A new approach to estimating pairwise sequence alignments. *Patterns*, 1(6), 2020. 4
- [4] G. Bresler, M. Bresler, and D. N. C. Tse. Optimal assembly for high throughput shotgun sequencing. *BMC bioinformatics*, 14(Suppl 5):S18, 2013. 2, 4
- [5] K. L. Chung. A course in probability theory. Elsevier, 2000. 46
- [6] T. M. Cover and J. A. Thomas. Elements of Information Theory. Wiley-Interscience, Hoboken, NJ, USA, 2006. 24, 45, 52
- [7] A. Dembo. Large deviations techniques and applications. Springer, 2009. 7, 27, 54
- [8] W. Feller. An introduction to probability theory and its applications, Volume 2, volume 81. John Wiley & Sons, 1991. 54
- [9] R. M. Gray. Entropy and information theory. Springer Science & Business Media, 2011. 44

- [10] G. Greenberg and I. Shomorony. The metagenomic binning problem: Clustering Markov sequences. In 2019 IEEE Information Theory Workshop (ITW), pages 1–5. IEEE, 2019. 3
- [11] M. Herring. A probabilistic analysis of shotgun sequencing for metagenomics. arXiv preprint arXiv:2201.05029, 2022. 3
- [12] G. Kamath, T. Baharav, and I. Shomorony. Adaptive learning of rank-one models for efficient pairwise sequence alignment. Advances in Neural Information Processing Systems, 33:7513–7525, 2020. 4
- [13] G. M. Kamath, I. Shomorony, F. Xia, T. A. Courtade, and D. N. C. Tse. HINGE: long-read assembly achieves optimal repeat resolution. *Genome research*, 27(5):747–756, 2017. 28
- [14] D. D. Kang, J. Froula, R. Egan, and Z. Wang. MetaBAT, an efficient tool for accurately reconstructing single genomes from complex microbial communities. *PeerJ*, 3:e1165, 2015. 3
- [15] K-K. Lam, A. Khalak, and D. N. C. Tse. Near-optimal assembly for shotgun sequencing with noisy reads. BMC bioinformatics, 15(Suppl 9):S4. 4
- [16] E. S. Lander and M. S. Waterman. Genomic mapping by fingerprinting random clones: A mathematical analysis. *Genomics*, 2(3):231–239, 1988. 2, 3, 28
- [17] E. Le Chatelier, T. Nielsen, J. Qin, E. Prifti, F. Hildebrand, G. Falony, M. Almeida, M. Arumugam, J-M Batto, and S. Kennedy. Richness of human gut microbiome correlates with metabolic markers. *Nature*, 500(7464):541–546, 2013. 3
- [18] D. A. Levin and Y. Peres. Markov chains and mixing times, volume 107. American Mathematical Soc., 2017. 22
- [19] N. Merhav. d-semifaithful codes that are universal over both memoryless sources and distortion measures. IEEE Transactions on Information Theory, 69(7):4746–4757, 2023. 54
- [20] N. Merhav and N. Weinberger. A toolbox for refined information-theoretic analyses with applications. *arXiv preprint arXiv:2406.00744*, 2024. 7, 54
- [21] S. Mohajer, A. S. Motahari, and D. N. C. Tse. Reference-based DNA shotgun sequencing: Information theoretic limits. In 2013 IEEE International Symposium on Information Theory, pages 1635–1639. IEEE, 2013. 28
- [22] A. S. Motahari, G. Bresler, and D. N. C. Tse. Information theory of DNA shotgun sequencing. *IEEE Transactions on Information Theory*, 59(10):6273–6289, 2013. 2, 3, 4, 5, 16, 57
- [23] A. S. Motahari, K. Ramchandran, D. N. C. Tse, and N. Ma. Optimal DNA shotgun sequencing: Noisy reads are as good as noiseless reads. In 2013 IEEE International Symposium on Information Theory, pages 1640–1644. IEEE, 2013. 4
- [24] Y. Polyanskiy, H. V. Poor, and S. Verdú. Channel coding rate in the finite blocklength regime. *IEEE Transactions on Information Theory*, 56(5):2307–2359, 2010. 54
- [25] Y. Polyanskiy and Y. Wu. Information Theory: From Coding to Learning. Cambridge University Press, 2023+. 49, 51, 54, 63
- [26] Z. Rached, F. Alajaji, and L. L. Campbell. Rényi's divergence and entropy rates for finite alphabet Markov sources. *IEEE Transactions on Information theory*, 47(4):1553–1561, 2001. 29
- [27] J. A. Reuter, D. V. Spacek, and M. P. Snyder. High-throughput sequencing technologies. Molecular cell, 58(4):586-597, 2015. 1
- [28] K. Sahlin. Strobealign: Flexible seed size enables ultra-fast and accurate read alignment. Genome Biology, 23(1):260, 2022. 4
- [29] I. Shomorony, T. A. Courtade, and D. N. C. Tse. Fundamental limits of genome assembly under an adversarial erasure model. IEEE Transactions on Molecular, Biological and Multi-Scale Communications, 2(2):199–208, 2016. 4
- [30] I. Shomorony and G. M. Kamath. Sketching and sequence alignment: A rate-distortion perspective. In 2021 IEEE International Symposium on Information Theory (ISIT), pages 3308–3313. IEEE, 2021. 4, 28
- [31] I. Shomorony, S. H. Kim, T. A. Courtade, and D. N. C. Tse. Information-optimal genome assembly via sparse read-overlap graphs. *Bioinformatics*, 32(17):i494–i502, 2016. 4, 28
- [32] V. Y. F. Tan. Asymptotic estimates in information theory with non-vanishing error probabilities. *Foundations and Trends*® *in Communications and Information Theory*, 11(1-2):1–184, 2014. 7, 54
- [33] E. Ukkonen. Approximate string-matching with q-grams and maximal matches. *Theoretical computer science*, 92(1):191–211, 1992. 2

- [34] T. Van Erven and P. Harremos. Rényi divergence and Kullback-Leibler divergence. *IEEE Transactions on Information Theory*, 60(7):3797–3820, 2014. 59, 60, 63
- [35] G. Vázquez-Vilar, A. Albert, Guillén i Fàbregas, T. Koch, and A. Lancho. Saddlepoint approximation of the error probability of binary hypothesis testing. In 2018 IEEE International Symposium on Information Theory (ISIT), pages 2306–2310. IEEE, 2018. 54
- [36] N. Weinberger and I. Shomorony. Fundamental limits of reference-based sequence reordering. *IEEE Transactions on Information Theory*, 2024. 28