A UNIFYING FRAMEWORK FOR COMPLEX-VALUED EIGENFUNCTIONS VIA THE CARTAN EMBEDDING

SIGMUNDUR GUDMUNDSSON AND ADAM LINDSTRÖM

ABSTRACT. In this work we find a unifying scheme for the known explicit complex-valued eigenfunctions on the classical compact Riemannian symmetric spaces. For this we employ the well-known Cartan embedding for those spaces. This also leads to the construction of new eigenfunctions on the quaternionic Grassmannians.

1. INTRODUCTION

A complex-valued function $\phi : (M, g) \to \mathbb{C}$ on a Riemannian manifold is called an *eigenfunction* if it is eigen both with respect to the classical *Laplace-Beltrami* operator and the so called *conformality* operator. These objects have turned out be very useful for the construction of explicit complexvalued harmonic morphisms, p-harmonic functions and even minimal submanifolds of (M, g) of codimension two. For the original references to these constructions see Theorem 2.6 of [12], Theorem 3.1 in [14] and Theorem 1.3 from [11]. For the general theory of harmonic morphisms between Riemannian manifolds we refer to the excellent book [2] and the regularly updated online bibliography [9].

Explicit complex-valued eigenfunctions have been constructed on all the classical compact irreducible *Riemannian symmetric spaces* G/K, see Table 1. The main aim of this paper is to provide a unified framework for the construction for the seven last examples in the table. For this we employ the well-known *Cartan embedding* mapping such a symmetric space G/K into its isometry group G as a totally geodesic submanifold. Our main results can be summarised as follows.

Theorem 1.1. Let G/K be one of the classical compact irreducible symmetric spaces 4.-10. in Table 1. Denote by $\hat{\Phi} : G \to G$ the corresponding Cartan map. Then there exists a complex matrix $A \in \mathbb{C}^{n \times n}$ such that the function $\hat{\phi}_A : G \to \mathbb{C}$ with

$$\phi_A(p) = \operatorname{trace}(\Phi(p) \cdot A)$$

²⁰²⁰ Mathematics Subject Classification. 53C35, 53C43, 58E20.

Key words and phrases. symmetric spaces, complex-valued eigenfunctions, Cartan embedding.

is a K-invariant (λ, μ) -eigenfunction on G, for suitable $\lambda, \mu \in \mathbb{C}$. This induces a (λ, μ) -eigenfunction ϕ_A on the quotient G/K with the same eigenvalues.

As an important example we explicitly investigate the special case of the quaternionic Grassmannians $\mathbf{Sp}(m+n)/\mathbf{Sp}(m) \times \mathbf{Sp}(n)$. As a by-product we yield new eigenfunctions on these spaces. We then conclude with a new general construction method for eigenfunctions from Riemannian manifolds. The reader interested in further details is referred to [17].

No.	G/K	λ	μ	Eigenfunctions
1.	$\mathbf{SO}(n)$	$-\frac{(n-1)}{2}$	$-\frac{1}{2}$	see [12]
2.	$\mathbf{SU}(n)$	$-\frac{n^2-1}{n}$	$-\frac{n-1}{n}$	see [14]
3.	$\mathbf{Sp}(n)$	$-\frac{2n+1}{2}$	$-\frac{1}{2}$	see [12]
4.	$\mathbf{SU}(n)/\mathbf{SO}(n)$	$-rac{2(n^2+n-2)}{n}$	$-\frac{4(n-1)}{n}$	see [13]
5.	$\mathbf{Sp}(n)/\mathbf{U}(n)$	-2(n+1)	-2	see [13]
6.	$\mathbf{SO}(2n)/\mathbf{U}(n)$	-2(n-1)	-1	see [13]
7.	$\mathbf{SU}(2n)/\mathbf{Sp}(n)$	$-\frac{2(2n^2-n-1)}{n}$	$-\frac{2(n-1)}{n}$	see [13]
8.	$\mathbf{SO}(m+n)/\mathbf{SO}(m) \times \mathbf{SO}(n)$	-(m+n)	-2	see $[7]$
9.	$\mathbf{U}(m+n)/\mathbf{U}(m)\times\mathbf{U}(n)$	-2(m+n)	-2	see [8]
10.	$\mathbf{Sp}(m+n)/\mathbf{Sp}(m) \times \mathbf{Sp}(n)$	-2(m+n)	-1	see [8]

TABLE 1. Eigenfamilies on the classical compact irreducible Riemannian symmetric spaces.

2. Eigenfunctions and Eigenfamilies

Let (M, g) be an *m*-dimensional Riemannian manifold and $T^{\mathbb{C}}M$ be the complexification of the tangent bundle TM of M. We extend the metric g to a complex bilinear form on $T^{\mathbb{C}}M$. Then the gradient $\nabla \phi$ of a complexvalued function $\phi : (M, g) \to \mathbb{C}$ is a section of $T^{\mathbb{C}}M$. In this situation, the well-known complex linear Laplace-Beltrami operator (alt. tension field) τ on (M, g) acts locally on ϕ as follows

$$\tau(\phi) = \operatorname{div}(\nabla\phi) = \sum_{i,j=1}^{m} \frac{1}{\sqrt{|g|}} \frac{\partial}{\partial x_j} \left(g^{ij} \sqrt{|g|} \frac{\partial\phi}{\partial x_i} \right).$$

For two complex-valued functions $\phi,\psi:(M,g)\to\mathbb{C}$ we have the following well-known fundamental relation

$$\tau(\phi \cdot \psi) = \tau(\phi) \cdot \psi + 2\kappa(\phi, \psi) + \phi \cdot \tau(\psi),$$

where the complex bilinear *conformality operator* κ is given by

$$\kappa(\phi, \psi) = g(\nabla \phi, \nabla \psi).$$

In local coordinates this satisfies

$$\kappa(\phi,\psi) = \sum_{i,j=1}^{m} g^{ij} \cdot \frac{\partial \phi}{\partial x_i} \frac{\partial \psi}{\partial x_j}.$$

Definition 2.1. [12] Let (M, g) be a Riemannian manifold. Then a complexvalued function $\phi : M \to \mathbb{C}$ is said to be a (λ, μ) -eigenfunction if it is eigen both with respect to the Laplace-Beltrami operator τ and the conformality operator κ i.e. there exist complex numbers $\lambda, \mu \in \mathbb{C}$ such that

$$\tau(\phi) = \lambda \cdot \phi \text{ and } \kappa(\phi, \phi) = \mu \cdot \phi^2.$$

A set $\mathcal{E} = \{\phi_i : M \to \mathbb{C} \mid i \in I\}$ of complex-valued functions is said to be a (λ, μ) -eigenfamily on M if there exist complex numbers $\lambda, \mu \in \mathbb{C}$ such that for all $\phi, \psi \in \mathcal{E}$ we have

$$\tau(\phi) = \lambda \cdot \phi \text{ and } \kappa(\phi, \psi) = \mu \cdot \phi \psi.$$

For the standard odd-dimensional round spheres we have the following eigenfamilies based on the classical real-valued spherical harmonics.

Example 2.2. [11] Let S^{2n-1} be the odd-dimensional unit sphere in the standard Euclidean space $\mathbb{C}^n \cong \mathbb{R}^{2n}$ and define $\phi_1, \ldots, \phi_n : S^{2n-1} \to \mathbb{C}$ by

$$\phi_j: (z_1, \dots, z_n) \mapsto \frac{z_j}{\sqrt{|z_1|^2 + \dots + |z_n|^2}}.$$

Then the tension field τ and the conformality operator κ on S^{2n-1} satisfy

$$\tau(\phi_j) = -(2n-1) \cdot \phi_j$$
 and $\kappa(\phi_j, \phi_k) = -1 \cdot \phi_j \cdot \phi_k.$

For the standard complex projective space $\mathbb{C}P^n$ we have a complex multidimensional eigenfamily, obtained in a similar way.

Example 2.3. [11] Let $\mathbb{C}P^n$ be the standard *n*-dimensional complex projective space. For a fixed integer $1 \leq \alpha < n+1$ and some $1 \leq j \leq \alpha < k \leq n+1$ define the function $\phi_{jk} : \mathbb{C}P^n \to \mathbb{C}$ by

$$\phi_{jk}: [z_1, \dots, z_{n+1}] \mapsto \frac{z_j \cdot z_k}{z_1 \cdot \overline{z}_1 + \dots + z_{n+1} \cdot \overline{z}_{n+1}}.$$

Then the tension field τ and the conformality operator κ on $\mathbb{C}P^n$ satisfy

 $\tau(\phi_{jk}) = -4(n+1) \cdot \phi_{jk}$ and $\kappa(\phi_{jk}, \phi_{lm}) = -4 \cdot \phi_{jk} \cdot \phi_{lm}$.

The next result shows that a given eigenfamily \mathcal{E} on (M, g) induces a large collection $\mathcal{P}_d(\mathcal{E})$ of such objects. This is particularly interesting in light of Table 1.

Theorem 2.4. [8] Let (M,g) be a Riemannian manifold and the set of complex-valued functions

$$\mathcal{E} = \{\phi_i : M \to \mathbb{C} \mid i = 1, 2, \dots, n\}$$

be a finite eigenfamily i.e. there exist complex numbers $\lambda, \mu \in \mathbb{C}$ such that for all $\phi, \psi \in \mathcal{E}$

$$\tau(\phi) = \lambda \cdot \phi \text{ and } \kappa(\phi, \psi) = \mu \cdot \phi \psi.$$

Then the set of complex homogeneous polynomials of degree d

$$\mathcal{P}_d(\mathcal{E}) = \{ P : M \to \mathbb{C} \mid P \in \mathbb{C}[\phi_1, \phi_2, \dots, \phi_n], \ P(\alpha \cdot \phi) = \alpha^d \cdot P(\phi), \ \alpha \in \mathbb{C} \}$$

is an eigenfamily on M such that for all $P, Q \in \mathcal{P}_d(\mathcal{E})$ we have

$$\tau(P) = (d\lambda + d(d-1)\mu) \cdot P \quad and \quad \kappa(P,Q) = d^2\mu \cdot PQ.$$

3. The Cartan Embedding

In this section we describe the Cartan embedding which is our most important tool for the unifying scheme that we are introducing in this work. For the general theory of Riemannian symmetric spaces we refer to the much celebrated standard text [15] of Helgason.

Definition 3.1. Let (G, K, σ) be a symmetric triple of compact type. Then the *Cartan map* $\hat{\Phi} : G \to G$ is defined by

$$\hat{\Phi}: p \mapsto p \cdot \sigma(p^{-1}).$$

Since $\hat{\Phi}(p \cdot k) = \hat{\Phi}(p)$, for each $k \in K$, $\hat{\Phi}$ induces a map $\Phi : G/K \to G$ on the symmetric space G/K with

$$\Phi: pK \mapsto \hat{\Phi}(p) = p \cdot \sigma(p^{-1}),$$

known as the *Cartan embedding*.

The following result plays an important role in this work. It can be found in [3] as Proposition 3.42.

Theorem 3.2. [3] Let (G, K, σ) be a symmetric triple of compact type and let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the standard decomposition of the Lie algebra \mathfrak{g} of G. Then the Cartan embedding $\Phi : G/K \to G$ satisfies

- (i) Φ is an immersion,
- (ii) the image of Φ is totally geodesic in G,
- (*iii*) $\Phi(G/K) = \exp(\mathfrak{p}),$
- (iv) Φ is conformal with constant conformal factor 2.

Remark 3.3. What usually is called the "Cartan embedding" in the literature, is actually in some cases only an immersion and not an embedding. The best known case is $\Phi: S^n = \mathbf{SO}(n+1)/\mathbf{SO}(n) \to \mathbf{SO}(n+1)$ mapping the *n*-dimensional unit sphere S^n in \mathbb{R}^{n+1} onto the totally geodesic image in $\mathbf{SO}(n+1)$. This is diffeomorphic to the real projective space $\mathbb{R}P^n$. For this see Example 5.26 of [10].

Proof. (Theorem 3.2) The proofs of parts (i)-(iii) can be found in [3]. We have not been able to find the arguments for (iv) in the literature. For this reason they are provided here.

In order to calculate the differential of the Cartan map at a point $p \in G$ we rewrite $\hat{\Phi}$ as a composition $\hat{\Phi} = \mu \circ \xi$, where $\xi : G \to G \times G$ is the map

$$\xi: p \mapsto (p, \sigma(p^{-1}))$$

and $\mu: G \times G \to G$ is the group multiplication. Let $X_p \in T_pG$ be a tangent vector at p and $X \in \mathfrak{g}$ be the corresponding left-invariant vector field with $X_p = (dL_p)_e(X)$. Then we have

$$d\phi_p(X_p) = \frac{d}{dt} (\xi(p \cdot \exp(tX))) \Big|_{t=0}$$

=
$$\frac{d}{dt} ((p \cdot \exp(tX), \sigma(\exp(-tX)\sigma(p^{-1})))) \Big|_{t=0}$$

=
$$((dL_p)_e(X), -(dR_{\sigma(p^{-1})})_e(d\sigma(X))).$$

Employing the chain rule we see that the differential $d\hat{\Phi}$ of $\hat{\Phi}$ satisfies

$$\begin{aligned} d\hat{\Phi}_p(X_p) &= d\mu_{(p,\sigma(p^{-1}))}((dL_p)_e(X), -(dR_{\sigma(p^{-1})})_e(d\sigma(X))) \\ &= (dR_{\sigma(p^{-1})})_p(dL_p)_e(X) + (dL_p)_{\sigma(p^{-1})}(dR_{\sigma(p^{-1})})_e(-d\sigma(X)) \\ &= d(R_{\sigma(p^{-1})} \circ L_p)_e(X) + d(L_p \circ R_{\sigma(p^{-1})})_e(-d\sigma(X)) \\ &= d(R_{\sigma(p^{-1})} \circ L_p)_e(X - d\sigma(X)). \end{aligned}$$

In the last step we have used the fact that right and left translations commute with each other. For $X \in \mathfrak{k}$, we have $d\sigma(X) = X$ and the inner term vanishes as expected. On the other hand for $X \in \mathfrak{p}$, we have $d\sigma(X) = -X$ and we yield

$$d\Phi_p(X_p) = d(R_{\sigma(p^{-1})} \circ L_p)_e(X - d\sigma(X))$$

= $d(R_{\sigma(p^{-1})} \circ L_p)_e(2X)$
= $2(dR_{\sigma(p^{-1})})_p(X_p).$

Since the natural projection $\pi : G \to G/K$ is a Riemannian submersion and the Cartan map $\hat{\Phi}$ satisfies $\hat{\Phi} = \Phi \circ \pi$, we only need to show that the Cartan map is conformal when restricted to the horizontal distribution on G induced by the submersion π . In this case, it is easy to check that the horizontal distribution is given by left-translations of the subspace \mathfrak{p} of \mathfrak{g} . If $p \in G, \, X_p, Y_p$ are horizontal vectors at p and g is the bi-invariant metric on G we then have

$$g_{\hat{\Phi}(p)}(d\hat{\Phi}_p(X_p), d\hat{\Phi}_p(Y_p)) = g_{\Phi(p)}(2dR_{\sigma(p^{-1})}(X_p), 2dR_{\sigma(p^{-1})}(Y_p)) = 4 \cdot g_p(X_p, Y_p),$$

where the last equality follows from the right-invariance of the metric. \Box

Remark 3.4. The Cartan embedding $\Phi : G/K \to G$ is a conformal immersion with constant conformal factor and the image $\Phi(G/K)$ is totally geodesic in G. It then follows from the Proposition on page 119, of the seminal work [5] of Eells and Sampson, that Φ is a harmonic map.

4. EIGENFAMILIES VIA THE CARTAN EMBEDDING

In this section we show how the Cartan embedding can be used to construct eigenfunctions and eigenfamilies. For this we employ the following well-known result.

Theorem 4.1. Let (G, K, σ) be a symmetric triple of compact type with and $\dim(G/K) \geq 2$. Then the Cartan map $\hat{\Phi} : G \mapsto G$ is harmonic i.e. it satisfies

$$\tau(\hat{\Phi}) = 0$$

Proof. The projection $\pi : G \to G/K$ is a Riemannian submersion with totally geodesic fibres, and hence a harmonic morphism (see [6, 16] and [1]). It then follows from Proposition 7.13 in [4] that the composition $\hat{\Phi} = \Phi \circ \pi$ is harmonic as well.

When combined with the composition law, this has the following consequences given by equations (4.2) and (4.3). We let $\phi : G \to \mathbb{C}$ be a smooth complex-valued function and also denote by ϕ the restriction of ϕ to the image $\hat{\Phi}(G)$. Applying the composition law to the function $\phi \circ \hat{\Phi} : G \to \mathbb{C}$ then gives

$$\tau(\phi \circ \hat{\Phi}) = d\phi(\tau(\hat{\Phi})) \circ \hat{\Phi} + \text{trace} \,\nabla d\phi(d\hat{\Phi}, d\hat{\Phi}) \circ \hat{\Phi}$$

= $\text{trace} \,\nabla d\phi(d\hat{\Phi}, d\hat{\Phi}) \circ \hat{\Phi},$ (4.1)

since $\hat{\Phi}$ is harmonic. We now choose an orthonormal basis

$$\mathcal{B}_{\mathfrak{g}} = \{X_1, \dots, X_k, Y_1, \dots, Y_l\}$$

for \mathfrak{g} with $\mathcal{B}_{\mathfrak{k}} = \{X_1, \ldots, X_k\}$ and $\mathcal{B}_{\mathfrak{p}} = \{Y_1, \ldots, Y_l\}$ orthonormal bases for \mathfrak{k} and \mathfrak{p} , respectively. Expanding the right hand side of equation (4.1), we obtain

trace
$$\nabla d\phi(d\hat{\Phi}, d\hat{\Phi}) \circ \hat{\Phi} = \left\{ \sum_{X \in \mathcal{B}_{\mathfrak{g}}} d\hat{\Phi}(X)^2(\phi) - \left(\nabla_{\!\!\!d} \hat{\Phi}(X) d\hat{\Phi}(X) \right)(\phi) \right\} \circ \hat{\Phi}.$$

.

It can be seen in Section 5, that the second term of the sum vanishes, and for the first term, only the vector fields from \mathcal{B}_{p} contribute. This implies that

$$\tau(\phi \circ \hat{\Phi}) = \left\{ \sum_{X \in \mathcal{B}_{\mathfrak{p}}} d\hat{\Phi}(X)^2(\phi) \right\} \circ \hat{\Phi}.$$
 (4.2)

We get a similar formula for the conformality operator. Let $\phi, \psi: G \mapsto \mathbb{C}$ be smooth. Then

$$\kappa(\phi \circ \hat{\Phi}, \psi \circ \hat{\Phi}) = \left\{ \sum_{X \in \mathcal{B}_{\mathfrak{p}}} d\hat{\Phi}(X)(\phi) \cdot d\hat{\Phi}(X)(\psi) \right\} \circ \hat{\Phi}.$$
(4.3)

Remark 4.2. At a point $p \in G$ and for $X \in \mathfrak{p}$ we have the expression

$$d\hat{\Phi}_p(X_p) = 2(dR_{\sigma(p^{-1})})_p(X_p).$$

Since we assume G to be compact and equipped with a bi-invariant metric then the right translations are isometries. Hence if \mathcal{B}_p is an orthonormal basis for \mathfrak{p}_p , the set

$$\{(dR_{\sigma(p^{-1})})_p(X_p) \mid X_p \in \mathcal{B}_p\}$$

is an orthonormal basis for $T_{\hat{\Phi}(p)}N$, where N is the image $\hat{\Phi}(G)$ of $\hat{\Phi}$.

The next result then follows by Remark 4.2

Proposition 4.3. Let (G, K, σ) be a symmetric triple of compact type and let $\hat{\Phi} : G \to G$ be the associated Cartan map. Let N be the image $\hat{\Phi}(G)$ of $\hat{\Phi}, \tau, \kappa$ denote the tension field and conformality operator on the Lie group G and τ_N , κ_N denote the same on N. If $\phi, \psi : G \to \mathbb{C}$ are complex-valued C^2 -functions on G, then the compositions

$$\phi \circ \hat{\Phi}, \psi \circ \hat{\Phi} : G \to \mathbb{C}$$

are K-invariant C^2 -functions on G. The tension fields and conformality operators then satisfy the relations

$$\tau(\phi \circ \hat{\Phi}) = 4 \cdot \tau_N(\phi) \circ \hat{\Phi}$$

and

$$\kappa(\phi \circ \hat{\Phi}, \psi \circ \hat{\Phi}) = 4 \cdot \kappa_N(\phi, \psi) \circ \hat{\Phi}.$$

The following is useful when explicitly calculating the Laplace-Beltrami and conformality operators on the image of the Cartan embedding.

Remark 4.4. Let (G, K, σ) be a symmetric triple of compact type and $\hat{\Phi}: G \to G$ be the associated Cartan map. Let N be the image $\hat{\Phi}(G)$ of $\hat{\Phi}$. By Remark 4.2, given a point $p \in G$ and an orthonormal basis \mathcal{B}_p for \mathfrak{p}_p , an orthonormal basis for $T_{\Phi(p)}N$ is given by

$$\{(dR_{\sigma(p^{-1})})_p(X_p) \mid X_p \in \mathcal{B}_p\}.$$

If we now fix $X_p \in \mathcal{B}_p$ and take $X \in \mathfrak{p}$ such that $(dL_p)_e(X) = X_p$ we have

$$(dR_{\sigma(p^{-1})})_p(X_p) = (dR_{\sigma(p^{-1})})_p(dL_p)_e(dL_{\sigma(p^{-1})})_{\sigma(p)}(dL_{\sigma(p)})_e(X)$$

$$= (dL_{\hat{\Phi}(p)})_e \operatorname{Ad}_{\sigma(p)}(X).$$

Hence given an orthonormal basis $\mathcal{B}_{\mathfrak{p}}$ for \mathfrak{p} we can construct an orthonormal basis for $T_{\hat{\Phi}(p)}N$ as

$$\mathcal{B}_{\hat{\Phi}(p)} = \{ \mathrm{Ad}_{\sigma(p)}(X)_{\hat{\Phi}(p)} \mid X \in \mathcal{B}_{\mathfrak{p}} \}.$$

5. The General Linear Group $\mathbf{GL}_n(\mathbb{C})$

In this section we now turn our attention to the concrete Riemannian matrix Lie groups embedded as subgroups of the complex general linear group.

The group of linear automorphism of \mathbb{C}^n is the complex general linear group $\mathbf{GL}_n(\mathbb{C}) = \{z \in \mathbb{C}^{n \times n} \mid \det z \neq 0\}$ of invertible $n \times n$ matrices with its standard representation

$$z \mapsto \begin{bmatrix} z_{11} & \cdots & z_{1n} \\ \vdots & \ddots & \vdots \\ z_{n1} & \cdots & z_{nn} \end{bmatrix}.$$

Its Lie algebra $\mathfrak{gl}_n(\mathbb{C})$ of left-invariant vector fields on $\mathbf{GL}_n(\mathbb{C})$ can be identified with $\mathbb{C}^{n \times n}$ i.e. the complex linear space of $n \times n$ matrices. We equip $\mathbf{GL}_n(\mathbb{C})$ with its natural left-invariant Riemannian metric g induced by the standard Euclidean inner product $\mathfrak{gl}_n(\mathbb{C}) \times \mathfrak{gl}_n(\mathbb{C}) \to \mathbb{R}$ on its Lie algebra $\mathfrak{gl}_n(\mathbb{C})$ satisfying

$$g(Z, W) \mapsto \mathfrak{Re} \operatorname{trace} (Z \cdot \overline{W}^t).$$

For $1 \leq i, j \leq n$, we shall by E_{ij} denote the element of $\mathbb{R}^{n \times n}$ satisfying

$$(E_{ij})_{kl} = \delta_{ik}\delta_{jl}$$

and by D_t the diagonal matrices $D_t = E_{tt}$. For $1 \le r < s \le n$, let X_{rs} and Y_{rs} be the matrices satisfying

$$X_{rs} = \frac{1}{\sqrt{2}}(E_{rs} + E_{sr}), \quad Y_{rs} = \frac{1}{\sqrt{2}}(E_{rs} - E_{sr}).$$

For the real vector space $\mathfrak{gl}_n(\mathbb{C})$ we then have the canonical orthonormal basis $\mathcal{B}^{\mathbb{C}} = \mathcal{B} \cup i\mathcal{B}$, where

$$\mathcal{B} = \{Y_{rs}, X_{rs} \mid 1 \le r < s \le n\} \cup \{D_t \mid t = 1, 2, \dots, n\}.$$

The following relations shall prove useful.

Lemma 5.1. Let n > 1 be an integer. Then we have

$$\sum_{1 \le r < s \le n} Y_{rs}^2 = -\frac{n-1}{2} \cdot I_n, \sum_{1 \le r < s \le n} X_{rs}^2 = \frac{n-1}{2} \cdot I_n, \quad \sum_{t=1}^n D_t^2 = I_n.$$

Let G be a classical Lie subgroup of $\mathbf{GL}_n(\mathbb{C})$ with Lie algebra \mathfrak{g} inheriting the induced left-invariant Riemannian metric, which we shall also denote by g. In the cases considered in this paper, $\mathcal{B}_{\mathfrak{g}} = \mathcal{B}^{\mathbb{C}} \cap \mathfrak{g}$ will be an orthonormal basis for the subalgebra \mathfrak{g} of $\mathfrak{gl}_n(\mathbb{C})$. By employing the Koszul formula for the Levi-Civita connection ∇ on (G,g), we see that for all $Z, W \in \mathcal{B}_{\mathfrak{g}}$ we have

$$g(\nabla_{Z}Z, W) = g([W, Z], Z)$$

= $\Re \mathfrak{e} \operatorname{trace} ((WZ - ZW)\overline{Z}^{t})$
= $\Re \mathfrak{e} \operatorname{trace} (W(Z\overline{Z}^{t} - \overline{Z}^{t}Z))$
= 0.

If $Z \in \mathfrak{g}$ is a left-invariant vector field on G and $\phi : U \to \mathbb{C}$ is a local complex-valued function on G then the k-th order derivatives $Z^k(\phi)$ satisfy

$$Z^{k}(\phi)(p) = \frac{d^{k}}{ds^{k}} \left(\phi(p \cdot \exp(sZ)) \right) \Big|_{s=0},$$

This implies that the tension field τ and the conformality operator κ on G fulfill

where $\mathcal{B}_{\mathfrak{g}}$ is the orthonormal basis $\mathcal{B}^{\mathbb{C}} \cap \mathfrak{g}$ for the Lie algebra \mathfrak{g} .

6. The Quaternionic Unitary Group

For a positive integer $n \in \mathbb{Z}^+$, the quaternionic unitary group $\mathbf{Sp}(n)$ is the intersection of the unitary group $\mathbf{U}(2n)$ and the standard representation of the quaternionic general linear group $\mathbf{GL}_n(\mathbb{H})$ in $\mathbb{C}^{2n \times 2n}$ given by

$$(z+jw) \mapsto q = \begin{bmatrix} z & w \\ -\bar{w} & \bar{z} \end{bmatrix}.$$
 (6.1)

The Lie algebra $\mathfrak{sp}(n)$ of $\mathbf{Sp}(n)$ satisfies

$$\mathfrak{sp}(n) = \left\{ \begin{bmatrix} Z & W \\ -\bar{W} & \bar{Z} \end{bmatrix} \in \mathbb{C}^{2n \times 2n} \mid Z^* + Z = 0, \ W^t - W = 0 \right\}.$$

The following elements form the orthonormal basis $\mathcal{B}_{\mathfrak{sp}(n)}$ for the Lie algebra $\mathfrak{sp}(n)$ of $\mathbf{Sp}(n)$

$$\begin{split} Y_{rs}^{a} &= \frac{1}{\sqrt{2}} \begin{bmatrix} Y_{rs} & 0\\ 0 & Y_{rs} \end{bmatrix}, X_{rs}^{a} &= \frac{1}{\sqrt{2}} \begin{bmatrix} iX_{rs} & 0\\ 0 & -iX_{rs} \end{bmatrix}, \\ X_{rs}^{b} &= \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & iX_{rs}\\ iX_{rs} & 0 \end{bmatrix}, X_{rs}^{c} &= \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & X_{rs}\\ -X_{rs} & 0 \end{bmatrix}, \end{split}$$

$$D_t^a = \frac{1}{\sqrt{2}} \begin{bmatrix} iD_t & 0\\ 0 & -iD_t \end{bmatrix}, D_t^b = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & iD_t\\ iD_t & 0 \end{bmatrix}, D_t^c = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & D_t\\ -D_t & 0 \end{bmatrix}$$

Here $1 \le r < s \le n$ and $1 \le t \le n$.

For those interested in the construction of eigenfamilies on the quaternionic unitary group we recommend the paper [12].

7. The Quaternionic Grassmannians

In this section we describe the compact quaternionic Grassmannians which are classical Riemannian symmetric spaces.

For positive integers $m, n \in \mathbb{Z}^+$, let G be the quaternionic unitary group $\mathbf{Sp}(m+n)$ with the subgroup $K = \mathbf{Sp}(m) \times \mathbf{Sp}(n)$ given in terms of quaternionic matrices by

$$\mathbf{Sp}(n) \times \mathbf{Sp}(m) = \left\{ \begin{bmatrix} q_1 & 0\\ 0 & q_2 \end{bmatrix} \in \mathbf{Sp}(m+n) \mid q_1 \in \mathbf{Sp}(m), \ q_2 \in \mathbf{Sp}(n) \right\}.$$

Let $I_m \in \mathbb{R}^{m \times m}$ denote the identity matrix and define $I_{m,n} \in \mathbb{R}^{(m+n) \times (m+n)}$ and $\hat{I}_{m,n} \in \mathbb{R}^{2(m+n) \times 2(m+n)}$ by

$$I_{m,n} = \begin{bmatrix} I_m & 0\\ 0 & -I_n \end{bmatrix} \text{ and } \hat{I}_{m,n} = \begin{bmatrix} I_{m,n} & 0\\ 0 & I_{m,n} \end{bmatrix}$$

respectively. Let $\sigma : G \to G$ be the involutive automorphism with $\sigma(q) = I_{m,n} \cdot q \cdot I_{m,n}$. Then it is easily seen that the subgroup K is the fixed-point set of σ . Thus the compact quotient

$$\mathbf{Sp}(m+n)/\mathbf{Sp}(m) \times \mathbf{Sp}(n)$$

is a Riemannian symmetric space, called the quaternionic Grassmannian. In terms of the standard complex representation of $\mathbf{Sp}(m+n)$ one easily checks that $\sigma(q) = \hat{I}_{m,n} \cdot q \cdot \hat{I}_{m,n}$. The Cartan map $\hat{\Phi}: G \to G$ is given by

$$\hat{\Phi}: q \mapsto q \cdot \hat{I}_{m,n} \cdot \bar{q}^t \cdot \hat{I}_{m,n}$$

and is clearly K-invariant. This induces the Cartan embedding $\Phi: G/K \to G$ with $\Phi(qK) = \hat{\Phi}(q)$.

Let $\mathfrak{sp}(m+n) = \mathfrak{k} \oplus \mathfrak{p}$ be the Cartan decomposition induced by σ . Then $\mathfrak{k} \cong \mathfrak{sp}(m) \oplus \mathfrak{sp}(n)$ and the set

$$\left\{Y^{a}_{rs}, X^{a}_{rs}, X^{b}_{rs}, X^{c}_{rs} \; \middle| \; m+1 \le r \le m+n, 1 \le s \le m\right\}$$

is an orthonormal basis for \mathfrak{p} .

8. EIGENFAMILIES ON THE QUATERNIONIC GRASSMANNIANS

The aim of this section is to construct eigenfamilies on the quaternionic Grassmannians. Here we employ the Cartan embedding to obtain known examples but also new ones, see Theorem 8.2.

For $1 \leq j, \alpha \leq 2(m+n)$, we define the maps $\hat{\psi}_{j\alpha} : \mathbf{Sp}(m+n) \to \mathbb{C}$ on the quaternionic group $\mathbf{Sp}(m+n)$ by

$$\hat{\psi}_{j\alpha}: q \mapsto \frac{1}{2} \cdot (q\hat{I}_{m,n}\bar{q}^t + I_{2(m+n)})_{j\alpha}.$$

It is easy to verify that $\hat{\psi}_{j\alpha}$ then satisfies

$$\hat{\psi}_{j\alpha}(q) = \sum_{r=1}^{m} q_{jr} \bar{q}_{\alpha r} + \sum_{r=m+n+1}^{2m+n} q_{jr} \bar{q}_{\alpha r}.$$
(8.1)

Further we see that we may write $\hat{\psi}_{j\alpha}$ as a composition with the Cartan map $\hat{\psi}_{j\alpha} = \eta_{j\alpha} \circ \hat{\Phi}$, where $\eta_{j\alpha} : \mathbf{Sp}(m+n) \to \mathbb{C}$ is given by

$$\eta_{j\alpha}: q \mapsto \frac{1}{2} \cdot (q \cdot \hat{I}_{m,n} + I_{2(m+n)})_{j\alpha}$$

Hence $\hat{\psi}_{j\alpha}$ is $\mathbf{Sp}(m) \times \mathbf{Sp}(n)$ -invariant.

Proposition 8.1. Let $1 \leq j, k, \alpha \leq 2(m+n)$ with $j, k \neq \alpha$. Then the tension field τ and the conformality operator κ on the quaternionic unitary group $\mathbf{Sp}(m+n)$ satisfy

$$\tau(\hat{\psi}_{j\alpha}) = -2(m+n) \cdot \hat{\psi}_{j\alpha} \quad and \quad \kappa(\hat{\psi}_{j\alpha}, \hat{\psi}_{k\alpha}) = -\hat{\psi}_{j\alpha}\hat{\psi}_{k\alpha}.$$

Proof. Denote by N the totally geodesic image of the Cartan embedding Φ in $\mathbf{Sp}(m+n)$. It follows from Proposition 4.3 that in order to compute $\tau(\hat{\psi}_{j\alpha})$ and $\kappa(\hat{\psi}_{j\alpha},\hat{\psi}_{k\beta})$ at a point $q \in \mathbf{Sp}(m+n)$ it is sufficient to compute

$$\tau_N(\eta_{j\alpha})(\Phi(q))$$
 and $\kappa_N(\eta_{j\alpha},\eta_{k\beta})(\Phi(q))$

at $\hat{\Phi}(q) \in N$. By Remark 4.4, for each $q \in \mathbf{Sp}(m+n)$ an orthonormal basis for $T_{\hat{\Phi}(q)}N$ is given by

$$\mathcal{B}_{\hat{\Phi}(q)} = \{ \mathrm{Ad}_{\sigma(q)}(Q)_{\hat{\Phi}(q)} \mid Q \in \mathcal{B}_{\mathfrak{p}} \}.$$

We then have

$$\begin{aligned} \operatorname{Ad}_{\sigma(q)}(Q)(\eta_{j\alpha})(\hat{\Phi}(q)) \\ &= \frac{d}{dt} \Big|_{t=0} \eta_{j\alpha}(\hat{\Phi}(q) \cdot \exp(t \operatorname{Ad}_{\sigma(q)} Q)) \\ &= \frac{1}{2} \frac{d}{dt} \Big|_{t=0} e_j \cdot (\hat{\Phi}(q)\sigma(q) \exp(tQ)\sigma(q)^{-1}\hat{I}_{m,n} + I_{2(m+n)}) \cdot e_{\alpha}^t \\ &= \frac{1}{2} \frac{d}{dt} \Big|_{t=0} e_j \cdot (q \exp(tQ)\hat{I}_{m,n}\bar{q}^t + I_{2(m+n)}) \cdot e_{\alpha}^t \\ &= \frac{1}{2} e_j \cdot q Q \hat{I}_{m,n} \bar{q}^t \cdot e_{\alpha}^t. \end{aligned}$$

Similarly we compute

$$\operatorname{Ad}_{\sigma(q)}(Q)^{2}(\eta_{j\alpha})(\hat{\Phi}(q)) = \frac{d^{2}}{dt^{2}}\Big|_{t=0} \eta_{j\alpha}(\hat{\Phi}(q) \cdot \exp(t \operatorname{Ad}_{\sigma(q)} Q))$$

$$= \frac{1}{2} e_j \cdot q \, Q^2 \hat{I}_{m,n} \, \bar{q}^t \cdot e^t_\alpha$$

For the tension field τ_N we then have

$$\begin{aligned} &\tau_{N}(\eta_{j\alpha})(\Phi(q)) \\ &= \frac{1}{2} \sum_{Q \in \mathcal{B}_{p}} e_{j} \cdot q Q^{2} \tilde{I}_{m,n} \bar{q}^{t} \cdot e_{\alpha}^{t} \\ &= \frac{1}{2} \sum_{r=1}^{m} \sum_{s=m+1}^{m+n} e_{j} \cdot q \left((Y_{rs}^{a})^{2} + (X_{rs}^{a})^{2} + (X_{rs}^{b})^{2} + (X_{rs}^{c})^{2} \right) \tilde{I}_{m,n} \bar{q}^{t} \cdot e_{\alpha}^{t} \\ &= \frac{1}{4} e_{j} \cdot q \left[\sum_{r=1}^{m} \sum_{s=m+1}^{m+n} {Y_{rs}^{2} \ 0 \ Y_{rs}^{2}} \right) - 3 {X_{rs}^{2} \ 0 \ X_{rs}^{2}} \right] \tilde{I}_{m,n} \bar{q}^{t} \cdot e_{\alpha}^{t} \\ &= e_{j} \cdot q \left[\sum_{r=1}^{m} \sum_{s=m+1}^{m+n} {Y_{rs}^{2} \ 0 \ Y_{rs}^{2}} \right] \tilde{I}_{m,n} \bar{q}^{t} \cdot e_{\alpha}^{t}. \end{aligned}$$
(8.2)

From Lemma 5.1, we can deduce

$$\sum_{r=1}^{m} \sum_{s=m+1}^{m+n} \begin{pmatrix} Y_{rs}^2 & 0\\ 0 & Y_{rs}^2 \end{pmatrix} = -\frac{m+n}{4} \cdot I_{2(m+n)} + \frac{m-n}{4} \cdot \tilde{I}_{m,n},$$

which yields

$$\begin{aligned} &\tau_N(\eta_{j\alpha})(\Phi(q)) \\ &= e_j \cdot q \left[-\frac{m+n}{4} \cdot I_{2(m+n)} + \frac{m-n}{4} \cdot \tilde{I}_{m,n} \right] \tilde{I}_{m,n} \bar{q}^t \cdot e_\alpha^t \\ &= -\frac{m+n}{2} \cdot \frac{1}{2} e_j \cdot (q \tilde{I}_{m,n} \bar{q}^t + I_{2(m+n)}) \cdot e_\alpha^t + \frac{m+n}{4} \cdot \delta_{j\alpha} + \frac{m-n}{4} \cdot \delta_{j\alpha} \\ &= -\frac{m+n}{2} \cdot \psi_{j\alpha}(q) + 2m \cdot \delta_{j\alpha}. \end{aligned}$$

Under our assumption that $j\neq \alpha$ this simplifies further to

$$\tau_N(\eta_{j\alpha})(\hat{\Phi}(q)) = -\frac{m+n}{2} \cdot \hat{\psi}_{j\alpha}(q).$$
(8.3)

We now turn to the conformality operator. For $1 \leq j,k,\alpha,\beta \leq$ we have

$$\begin{split} \kappa_N(\eta_{j\alpha},\eta_{k\beta})(\Phi(q)) \\ &= \frac{1}{4} \sum_{r=1}^m \sum_{s=m+1}^{m+n} \left[e_j \cdot q Y_{rs}^a \tilde{I}_{m,n} \bar{q}^t \cdot e_\alpha^t \cdot e_k \cdot q Y_{rs}^a \tilde{I}_{m,n} \bar{q}^t \cdot e_\beta^t \right. \\ &\quad + e_j \cdot q X_{rs}^a \tilde{I}_{m,n} \bar{q}^t \cdot e_\alpha^t \cdot e_k \cdot q X_{rs}^a \tilde{I}_{m,n} \bar{q}^t \cdot e_\beta^t \\ &\quad + e_j \cdot q x_{rs}^b \tilde{I}_{m,n} \bar{q}^t \cdot e_\alpha^t \cdot e_k \cdot q X_{rs}^b \tilde{I}_{m,n} \bar{q}^t \cdot e_\beta^t \\ &\quad + e_j \cdot q X_{rs}^c \tilde{I}_{m,n} \bar{q}^t \cdot e_\alpha^t \cdot e_k \cdot q X_{rs}^c \tilde{I}_{m,n} \bar{q}^t \cdot e_\beta^t \right]. \end{split}$$

12

Expanding this as a polynomial in the matrix components of q and \bar{q} and making use of the notation $\tilde{r} = r + m + n$ and $\tilde{s} = s + m + n$ we arrive, after a lengthy calculation, at

$$\begin{aligned} \kappa_N(\eta_{j\alpha},\eta_{k\beta})(\Phi(q)) \\ &= \frac{1}{8} \cdot \left(\delta_{j\beta} \cdot \hat{\psi}_{k\alpha}(q) + \delta_{j\alpha} \cdot \hat{\psi}_{j\beta}(q) - 2 \cdot \hat{\psi}_{j\beta}(q) \cdot \hat{\psi}_{k\alpha}(q) \right) \\ &+ \frac{1}{8} \sum_{r=1}^m \sum_{s=m+1}^{m+n} \left[(q_{jr}q_{k\tilde{r}} - q_{j\tilde{r}}q_{kr})(\bar{q}_{\alpha s}\bar{q}_{\beta\tilde{s}} - \bar{q}_{\alpha\tilde{s}}\bar{q}_{\beta s}) \right. \\ &+ (q_{j\tilde{s}}q_{ks} - q_{js}q_{k\tilde{s}})(\bar{q}_{\alpha\tilde{r}}\bar{q}_{\beta r} - \bar{q}_{\alpha r}\bar{q}_{\beta\tilde{r}}) \right]. \end{aligned}$$

Recalling our assumption that $\alpha = \beta \neq j, k$, we see that the terms involving sums and Kronecker deltas cancel and we conclude

$$\kappa_N(\eta_{j\alpha},\eta_{k\alpha})(\hat{\Phi}(q)) = -\frac{1}{4} \cdot \hat{\psi}_{j\alpha}(q) \cdot \hat{\psi}_{k\alpha}(q).$$
(8.4)

The result now follows by applying Proposition 4.3 to the equations (8.3) and (8.4).

The $\mathbf{Sp}(m) \times \mathbf{Sp}(n)$ -invariant functions $\hat{\psi}_{j\alpha} : \mathbf{Sp}(m+n) \to \mathbb{C}$ induce functions $\psi_{j\alpha} : \mathbf{Sp}(m+n)/\mathbf{Sp}(m) \times \mathbf{Sp}(n) \to \mathbb{C}$ on the quotient space. The eigenfunctions obtained by taking $1 \leq j, \alpha \leq m+n$ coincide with those found by Gudmundsson and Ghandour in [8], while the remaining cases are new. Hence, by the following result, we yield the eigenfamilies found in [8] and obtain new ones.

Theorem 8.2. For a fixed natural number $1 \le \alpha \le 2(m+n)$ the set

$$\mathcal{E}_{\alpha} = \{\psi_{j\alpha} : \mathbf{Sp}(m+n) / \mathbf{Sp}(n) \times \mathbf{Sp}(m) \to \mathbb{C} \mid 1 \le j \le 2(m+n), \ j \ne \alpha\}$$

is an eigenfamily on the quaternionic Grassmannian such that the tension field τ and conformality operator κ satisfy

$$\tau(\phi) = -2(m+n) \cdot \phi \text{ and } \kappa(\phi,\psi) = -\phi \cdot \psi,$$

for all $\phi, \psi \in \mathcal{E}_{\alpha}$.

9. The Complex Grassmannians

As a homogenous space, the complex Grassmannians are obtained as $G_n(\mathbb{C}^{m+n}) = \mathbf{U}(m+n)/\mathbf{U}(m) \times \mathbf{U}(n)$. The Cartan involution of $\mathbf{U}(m+n)$ is given by conjugation with the block matrix $I_{m,n}$. So explicitly

$$\sigma(z) = I_{m,n} \cdot z \cdot I_{m,n}.$$

The Cartan map is thus $\Phi(z) = z \cdot I_{m,n} \cdot \overline{z}^t \cdot I_{m,n}$. For $1 \leq j, \alpha \leq m+n$ set $\eta_{j\alpha}(z) = (z \cdot I_{m,n} + I_{m+n})_{j\alpha}$

and let $\hat{\psi}_{j\alpha} = \eta_{j\alpha} \circ \Phi$. Then one easily checks that these functions coincide with the $\mathbf{U}(m) \times \mathbf{U}(n)$ -invariant functions described in [8]. Denoting by $\psi_{j\alpha}$ the induced functions on the Grassmannians, we thus have the following. **Theorem 9.1.** [8] Let M be the complex Grassmannian manifold

$$\mathbf{U}(m+n)/\mathbf{U}(m) \times \mathbf{U}(n).$$

For a fixed natural number $1 \leq \alpha \leq m+n$, the set

$$\mathcal{E}_{\alpha} = \{\psi_{j\alpha} : M \to \mathbb{C} \mid 1 \le j \le m + n, j \ne \alpha\}$$

is an eigenfamily on the complex Grassmannian M such that the tension field τ and conformality operator κ satisfy

$$\tau(\phi) = -2(m+n) \cdot \phi \text{ and } \kappa(\phi,\psi) = -2 \cdot \phi \psi,$$

for all $\phi, \psi \in \mathcal{E}_{\alpha}$.

10. The Remaining Classical Compact Symmetric Spaces

A detailed description of the Riemannian symmetric spaces of this section can be found in Helgason's book [15]. There the Cartan involutions are also listed, from which the Cartan embeddings are easily derived.

The eigenfunctions on the real Grassmannians found in [7] fall into our unifying scheme as follows.

Theorem 10.1. Let M be the real Grassmannian of oriented n-planes in \mathbb{R}^{m+n} i.e.

$$\mathbf{SO}(m+n)/\mathbf{SO}(m) \times \mathbf{SO}(n)$$

Let A be an $(m+n) \times (m+n)$ complex symmetric matrix with rank A = 1, $A^2 = 0$ and trace A = 0. Define $\eta_A : \mathbf{SO}(m+n) \to \mathbb{C}$ by

$$\eta_A : x \mapsto \frac{1}{2} \operatorname{trace}((x \cdot I_{m,n} + I_{m+n}) \cdot A).$$

Then the function $\psi_A : M \to \mathbb{C}$ induced by the composition $\hat{\psi}_A = \eta_A \circ \hat{\Phi}$ of η_A with the Cartan map is an eigenfunction on M satisfying

$$au(\psi_A) = -(m+n) \cdot \psi_A$$
 and $\kappa(\psi_A, \psi_A) = -2 \cdot \psi_A^2$.

The same applies for the solutions on the Riemannian symmetric space $\mathbf{SU}(n)/\mathbf{SO}(n)$ constructed in [13] with the following.

Theorem 10.2. Let M be the Riemannian symmetric space $\mathbf{SU}(n)/\mathbf{SO}(n)$. Let $A = a^t a \in \mathbb{C}^{n \times n}$ for some non-zero complex vector $a \in \mathbb{C}^n$ and define $\eta_A : \mathbf{SU}(n) \to \mathbb{C}$ with

$$\eta_A: z \mapsto \operatorname{trace}(z \cdot A).$$

Then the function $\phi_A : M \to \mathbb{C}$ induced by the composition $\hat{\phi}_A = \eta_A \circ \Phi$ of η_A with the Cartan map is an eigenfunction on M satisfying

$$\tau(\phi_A) = -\frac{2(n^2 + n - 2)}{n} \cdot \phi_A, \quad \kappa(\phi_A, \phi_A) = -\frac{4(n - 1)}{n} \cdot \phi_A^2.$$

In their work [13] the authors provide complex-valued eigenfunctions on $\mathbf{SO}(2n)/\mathbf{U}(n)$. In our unifying scheme the are described as follows.

Theorem 10.3. Let M be the Riemannian symmetric space $\mathbf{SO}(2n)/\mathbf{U}(n)$. Let V be an isotropic subspace of \mathbb{C}^{2n} , $a, b \in V$ be linearly independent and A the skew-symmetric matrix

$$A = \sum_{r,s=1}^{2n} a_r b_s Y_{rs}.$$

Define the complex-valued function $\eta_A : \mathbf{SO}(2n) \mapsto \mathbb{C}$ by

$$\eta_A : x \mapsto \operatorname{trace}(x \cdot JA).$$

Then the function $\psi_A : M \to \mathbb{C}$ induced by the composition $\hat{\psi}_A = \eta_A \circ \Phi$ of η_A with the Cartan map is an eigenfunction on M satisfying

$$au(\psi_A) = -2(n-1)\cdot\psi_A, \quad \kappa(\psi_A,\psi_A) = -\psi_A^2.$$

The same applies to the eigenfunctions on $\mathbf{Sp}(n)/\mathbf{U}(n)$ found again in [13].

Theorem 10.4. Let M be the Riemannian symmetric space $\mathbf{Sp}(n)/\mathbf{U}(n)$. Further let $A = a^t a$ for some non-zero vector $a \in \mathbb{C}^{2n}$ and define the complex-valued function $\eta_A : \mathbf{Sp}(n) \to \mathbb{C}$ with

$$\eta_A: q \mapsto \operatorname{trace}(q \cdot A).$$

Then the function $\phi_A : M \to \mathbb{C}$ induced by the composition $\phi_A = \eta_A \circ \Phi$ of η_A with the Cartan map is an eigenfunction on M satisfying

$$\tau(\phi_A) = -2(n+1) \cdot \phi_A$$
 and $\kappa(\phi_A, \phi_A) = -2 \cdot \phi_A^2$.

In the last case of $\mathbf{SU}(2n)/\mathbf{Sp}(n)$ the eigenfunctions produced in [13] can be described as follows.

Theorem 10.5. Let M be the Riemannian symmetric space $\mathbf{SU}(2n)/\mathbf{Sp}(n)$. For two non-zero linearly independent vectors $a, b \in \mathbb{C}^{2n}$, let $A \in \mathbb{C}^{2n \times 2n}$ be the skew-symmetric matrix

$$A = \sum_{r,s=1}^{2n} a_r b_s Y_{rs}.$$

Define the complex-valued function $\eta_A : \mathbf{SU}(2n) \to \mathbb{C}$ by

$$\eta_A: z \mapsto \operatorname{trace}(z \cdot A).$$

Then the function $\phi_A : M \to \mathbb{C}$ induced by the composition $\phi_A = \eta_A \circ \Phi$ of η_A with the Cartan map is an eigenfunction on M satisfying

$$au(\phi_A) = \frac{2(2n^2 - n - 1)}{n} \cdot \phi_A \text{ and } \kappa(\phi_A, \phi_A) = -\frac{2(n - 1)}{n} \cdot \phi_A^2.$$

11. A New Constructions

In this section we introduce a new method for the construction complexvalued harmonic morphisms on the product of Riemannian manifolds. The main result is given by the following statement.

Theorem 11.1. Let (M_1, g_1) and (M_2, g_2) be two Riemannian manifolds and let $N = M_1 \times M_2$ be their Riemannian product, with product metric g. Let τ_1, τ_2, τ and $\kappa_1, \kappa_2, \kappa$ be the tension fields and conformality operators on M_1 , M_2 and N, respectively. Suppose that \mathcal{E}_1 and \mathcal{E}_2 are eigenfamilies on M_1 and M_2 with eigenvalues λ_1, μ_1 and λ_2, μ_2 , respectively. Then the set

$$\mathcal{E} = \{ \phi : N \to \mathbb{C} \mid \phi : (p_1, p_2) \mapsto \phi_1(p_1) \cdot \phi_2(p_2), \quad \phi_1 \in \mathcal{E}_1, \phi_2 \in \mathcal{E}_2 \}$$

is an eigenfamily on the product space N, with

$$\tau(\phi) = (\lambda_1 + \lambda_2) \cdot \phi$$
 and $\kappa(\phi, \psi) = (\mu_1 + \mu_2) \cdot \phi \psi$

for all $\phi, \psi \in \mathcal{E}$.

Proof. Recall that at each point (p_1, p_2) of M we have the orthogonal decomposition

$$T_{(p_1,p_2)}N = T_{p_1}M_1 \oplus T_{p_2}M_2.$$

Take $p = (p_1, p_2) \in M$ and let $\{X_1, \ldots, X_{m_1}, Y_1, \ldots, Y_{m_2}\}$ be a local orthonormal frame for TM on an open subset $U = U_1 \times U_2$ containing (p_1, p_2) such that $\{X_1, \ldots, X_{m_1}\}$ and $\{Y_1, \ldots, Y_{m_2}\}$ are local orthonormal frames for TM_1 and TM_2 , respectively. Employing the above formulae we can now determine the tension field

$$\begin{aligned} \tau(\phi)(p) &= \sum_{k=1}^{m_1} \left(X_k^2(\phi)(p) - \left(\nabla_{X_k} X_k \right)(\phi)(p) \right) \\ &+ \sum_{l=1}^{m_2} \left(Y_l^2(\phi)(p) - \left(\nabla_{Y_l} Y_l \right)(\phi)(p) \right) \\ &= \sum_{k=1}^{m_1} \left(X_k^2(\phi_1)(p_1) \cdot \phi_2(p_2) - \left(\nabla_{X_k} X_k \right)(\phi_1)(p_1) \cdot \phi_2(p_2) \right) \\ &+ \sum_{l=1}^{m_2} \left(\phi_1(p_1) \cdot Y_l^2(\phi_2)(p_2) - \phi_1(p_1) \cdot \left(\nabla_{Y_l} Y_l \right)(\phi_2)(p_2) \right) \\ &= \tau_1(\phi_1)(p_1) \cdot \phi_2(p_2) + \phi_1(p_1) \cdot \tau_2(\phi_2)(p_2) \\ &= \lambda_1 \cdot \phi_1(p_1)\phi_2(p_2) + \lambda_2 \cdot \phi_1(p_1)\phi_2(p_2) \\ &= (\lambda_1 + \lambda_2) \cdot \psi(p). \end{aligned}$$

For the conformality operator κ we now have

$$\kappa(\phi,\phi)(p) = \sum_{k=1}^{m_1} X_k(\phi)(p)^2 + \sum_{l=1}^{m_2} Y_l(\phi)(p)^2$$

16

A UNIFYING FRAMEWORK

$$= \phi_2(p_2)^2 \cdot \sum_{k=1}^{m_1} X_k(\phi_1)(p_1)^2 + \phi_1(p_1)^2 \cdot \sum_{l=1}^{m_2} Y_l(\phi_2)(p_2)^2$$

$$= \cdot \phi_2(p_2)^2 \cdot \kappa_1(\phi_1, \phi_1)(p_1) + \phi_1(p_1)^2 \cdot \kappa_2(\phi_2, \phi_2)(p_2)$$

$$= (\mu_1 + \mu_2) \cdot \phi(p)^2.$$

The general formula for the conformality operator κ follows from its symmetry. Since the point p was chosen arbitrarily we have completed the proof.

12. Acknowledgements

The authors would like to thank Thomas Jack Munn for useful discussions on this work. The second author gratefully acknowledges the support of the Austrian Science Fund (FWF) through the project "The Standard Model as a Geometric Variational Problem" (DOI: 10.55776/P36862).

References

- P. Baird, J. Eells, A conservation law for harmonic maps, Geometry Symposium Utrecht 1980, Lecture Notes in Mathematics 894, 1-25, Springer (1981).
- [2] P. Baird, J.C. Wood, *Harmonic Morphisms Between Riemannian Manifolds*, The London Mathematical Society Monographs 29, Oxford University Press (2003).
- [3] J. Cheeger, D. G. Ebin, Comparison Theorems in Riemannian Geometry, North-Holland Mathematical Library 9 (1975).
- [4] J.Eells, L. Lemaire, Selected topics in harmonic maps CBMS Regional Conf. Ser. in Math. 50, AMS (1983).
- [5] J.Eells, J.H.Sampson, Harmonic mappings of Riemannian manifolds, Amer. J. Math. 86 (1964), 109-160.
- [6] B. Fuglede, Harmonic morphisms between Riemannian manifolds, Ann. Inst. Fourier 28 (1978), 107-144
- [7] E. Ghandour, S. Gudmundsson, Explicit p-harmonic functions on the real Grassmannians, Adv. Geom. 23 (2023), 315–321.
- [8] E. Ghandour, S. Gudmundsson, Explicit harmonic morphisms and p-harmonic functions from the complex and quaternionic Grassmannians, Ann. Global Anal. Geom. 64 (2023), 15.
- [9] S. Gudmundsson, The Bibliography of Harmonic Morphisms, www.matematik.lu.se/ matematiklu/personal/sigma/harmonic/bibliography.html
- [10] S. Gudmundsson, An Introduction to Riemannian Geometry, www.matematik.lu.se/matematiklu/personal/sigma/Riemann.pdf
- [11] S. Gudmundsson, T. J. Munn, Minimal submanifolds via complex-valued eigenfunctions, J. Geom. Anal. 34 (2024), 190.
- [12] S. Gudmundsson, A. Sakovich, Harmonic morphisms from the classical compact semisimple Lie groups, Ann. Global. Anal. Geom. 33 (2008) 343-356.
- [13] S. Gudmundsson, A. Siffert, M. Sobak, Explicit proper p-harmonic functions on the Riemannian symmetric spaces $\mathbf{SU}(n)/\mathbf{SO}(n)$, $\mathbf{Sp}(n)/\mathbf{U}(n)$, $\mathbf{SO}(2n)/\mathbf{U}(n)$, $\mathbf{SU}(2n)/\mathbf{Sp}(n)$, J. Geom. Anal. **32** (2022), 147.
- [14] S. Gudmundsson, M. Sobak, Proper r-harmonic functions from Riemannian manifolds, Ann. Global Anal. Geom. 57 (2020), 217-223.
- [15] S. Helgason, Differential Geometry, Lie Groups, and Symmetric Spaces, Graduate Studies in Mathematics 34, AMS (2001).

- T. Ishihara, A mapping of Riemannian manifolds which preserves harmonic functions, J. Math. Soc. Japan 7 (1979), 345-370
- [17] A. Lindström, Harmonic Morphisms and p-Harmonic Functions on the Classical Compact Symmetric Spaces via the Cartan Embedding, Master's dissertation, University of Lund (2023).

Mathematics, Faculty of Science, Lund University, Box 118, Lund 221 00, Sweden

Email address: Sigmundur.Gudmundsson@math.lu.se

Institut für Mathematik, Universität Wien, Oskar-Morgenstern-Platz 1, 1090 Wien, Austria

 $Email \ address: {\tt Adam.Lindstroem@univie.ac.at}$

18