

Ribbon blocks for centraliser algebras of symmetric groups

Matthew Fayers

Queen Mary University of London
m.fayers@qmul.ac.uk

Lorenzo Putignano

Università degli Studi di Firenze
lorenzo.putignano@unifi.it

2020 Mathematics subject classification: 20C08, 20C30, 05A17, 05E10

Abstract

Suppose l, m are natural numbers with $l \leq m$, and \mathbb{F} a field of characteristic p , and let $\mathcal{C}_{l,m}^{\mathbb{F}}$ denote the centraliser of the group algebra $\mathbb{F}S_l$ inside $\mathbb{F}S_m$. Ellers and Murray give a conjectured classification of the blocks of $\mathcal{C}_{l,m}^{\mathbb{F}}$ in terms of the p -blocks of S_l and S_m . We prove this conjecture for a family of blocks that we call *ribbon blocks* and *belt blocks*. These are the blocks containing Specht modules labelled by skew-partitions having no repeated entries in their p -content.

Contents

1	Introduction	1
2	Background and notation	2
2.1	Partitions and Specht modules	3
2.2	Young diagrams and standard tableaux	3
2.3	Skew-partitions and skew Specht modules	4
2.4	The abacus	5
2.5	p -cores and p -blocks	6
2.6	The degenerate affine Hecke algebra	7
2.7	The centraliser algebra for symmetric groups	8
2.8	Combinatorial blocks	9
3	Ribbon blocks and belt blocks	11
3.1	Generalised ribbon blocks	12
3.2	Belt blocks	20

1 Introduction

Let G be a finite group, H a subgroup of G and \mathbb{F} an algebraically closed field. The *centraliser algebra* $\mathbb{F}G^H$ is defined by

$$\mathbb{F}G^H = \{a \in \mathbb{F}G \mid ah = ha \forall h \in H\}.$$

This algebra appears in a series of papers by Ellers [E1, E2, E3, E4] mainly motivated by the attempt to extend some *local-global* conjectures and theorems whenever \mathbb{F} has positive characteristic. Such statements relate representation-theoretic information on $\mathbb{F}G^H$ to local information. In particular, Ellers' main motivation lies in constructing a theory in which Alperin's Weight Conjecture and Brauer's First Main Theorem are just special cases of a more general setting.

With this in mind, the centraliser algebra has become an important object of study. If \mathbb{F} has characteristic zero, then a more general argument by Curtis and Reiner [CR, Section 11D] shows that $\mathbb{F}G^H$ is a semisimple algebra, with a straightforward construction of its simple modules. So the main focus is on the case where \mathbb{F} has positive characteristic.

In this paper we focus on the case where both G and H are symmetric groups. Let l, m be non-negative integers with $l \leq m$. The symmetric group S_l is naturally a subgroup of S_m and hence it makes sense to consider the centraliser algebra $\mathbb{F}S_m^{S_l} := \mathcal{C}_{l,m}^{\mathbb{F}}$. The main goal is to find a suitable labelling for the simple $\mathcal{C}_{l,m}^{\mathbb{F}}$ -modules and a complete description of its blocks, i.e. minimal two-sided ideals whose direct sum gives the whole algebra. A powerful result would be to give answers to these issues in terms of the combinatorics of integer partitions, somehow extending the symmetric group case pioneered by James [J] which we recall in Section 2. Following [K], a complete set of pairwise non-isomorphic simple $\mathcal{C}_{l,m}^{\mathbb{F}}$ -modules when $\text{char}(\mathbb{F}) = 0$ is given by modules of the form

$$S^{\mu \setminus \lambda} := \text{Hom}_{\mathbb{F}S_l}(S^{\mu \downarrow S_l}, S^{\lambda})$$

where μ and λ denote partitions of m and l , respectively, and $\mu \setminus \lambda$ is a *skew-partition* in the sense of Section 2.3. In view of the similarity with the symmetric group case, we name these modules *Specht modules*. The Specht modules are also defined in positive characteristic, where they are generally reducible, and (unlike in the symmetric group case) there does not seem to be a straightforward way to obtain the simple modules from the Specht modules. In positive characteristic, a full description of the simple modules of $\mathcal{C}_{l,m}^{\mathbb{F}}$ is far out of reach.

The goal of the present paper is to characterise certain blocks of $\mathcal{C}_{l,m}^{\mathbb{F}}$. As in the symmetric group case, describing the blocks amounts to finding the appropriate subdivision of the set of Specht modules (or equivalently, of the set of skew-partitions). In [EM2], Ellers and Murray suggested a classification of the blocks in positive characteristic and checked it for $m - l \leq 3$. Their conjecture (Conjecture 2.5) is inspired by the Nakayama Conjecture (Theorem 2.1) and hence is in terms of cores of partitions. We define a *combinatorial block* to be a set of skew-partitions which is predicted to comprise a single block by Conjecture 2.5. The work of Ellers and Murray shows that every combinatorial block is a union of blocks, so the remaining task to prove their conjecture is to show that the Specht modules labelled by two skew-partitions in the same combinatorial block really do lie in the same block.

In this paper we prove Ellers and Murray's conjecture for a class of combinatorial blocks of $\mathcal{C}_{l,m}^{\mathbb{F}}$ we call *ribbon blocks* and *belt blocks*. These are precisely the combinatorial blocks containin skew-partitions having no repetitions in their content (see Section 2.3). In Section 3 we show that (a proper approximation of) the decomposition matrix of a ribbon or belt block is a connected matrix; this implies that combinatorial blocks of this kind are blocks of the centraliser algebra.

Acknowledgements. The second author is supported by GNSAGA (INdAM) and has been funded by the European Union - Next Generation EU, Missione 4 Componente 1, PRIN 2022-2022PSTWLB - Group Theory and Applications, CUP B53D23009410006.

2 Background and notation

In this chapter we introduce all the background we need in Section 3. We begin with some basic notations. Let R be a commutative ring with field of fractions \mathbb{F} . Let \mathfrak{p} be a maximal ideal of R and denote by k the residue field R/\mathfrak{p} of positive characteristic p . We suppose that both \mathbb{F} and k are algebraically closed.

If $e \geq 2$ is an integer, then we write \bar{a} to mean $a + e\mathbb{Z}$, for $a \in \mathbb{Z}$, so that $\mathbb{Z}/e\mathbb{Z} = \{\bar{a} \mid a \in \mathbb{Z}\} = \{\bar{0}, \bar{1}, \dots, \overline{e-1}\}$.

In this paper, \mathbb{N} denotes the set of positive integers.

2.1 Partitions and Specht modules

In this section we introduce the main combinatorial objects in this paper. A *partition* is a non-decreasing sequence $\nu = (\nu_1, \nu_2, \dots)$ of non-negative integers such that $\nu_N = 0$ for $N \gg 0$. The integer ν_k is called the k -th *part* of ν and the number of non-zero parts is referred as the *length* of ν and denoted by $l(\nu)$. The (finite) sum $\nu_1 + \dots + \nu_{l(\nu)}$ is referred as the *size* of ν and denoted by $|\nu|$. If $n \geq 0$ and ν is a partition of size n , we say that ν is a *partition of n* and we write $\nu \vdash n$. We denote by \emptyset the unique partition of 0. When writing partitions we omit the trailing zeroes and group together equal parts with a superscript, e.g. $(3, 2^3, 1)$ stands for the length 5 partition $(3, 2, 2, 2, 1, 0, \dots) \vdash 10$. We denote by \mathcal{P} the set of all partitions and by \mathcal{P}_n the set of partitions of a fixed non-negative integer n .


The set of partitions \mathcal{P}_n plays the role of a powerful labelling set for a relevant class of $RS_n, \mathbb{F}S_n$ and kS_n -modules known in the literature as *Specht modules*. The set of Specht modules $\{S^\nu \mid \nu \in \mathcal{P}_n\}$ is a crucial object in the study of the representation theory of the symmetric group S_n because it provides a complete set of pairwise non-isomorphic simple $\mathbb{F}S_n$ -modules (in fact, more generally, over any characteristic zero field). If \mathbb{F} is replaced by the residue field k , an analogous statement holds if $n < p$. When $n \geq p$, a complete family of pairwise non-isomorphic simple kS_n -modules can be constructed from the Specht modules: if μ is a *p -regular* partition (meaning that it does not have p equal positive parts), then the Specht module S^μ over k has a simple head D^μ , and the simple modules obtained in this way give a complete set of non-isomorphic simple kS_n -modules. The main outstanding problem in the modular representation theory of the symmetric group is the determination of the *decomposition numbers* $[S^\lambda : D^\mu]$ for all partitions λ, μ of n with μ p -regular.

2.2 Young diagrams and standard tableaux

We now introduce a good method for visualising partitions. The *Young diagram* of a partition ν is the set

$$[\nu] = \{(r, c) \in \mathbb{N}^2 \mid c \leq \nu_r\}.$$

The elements of $[\nu]$, and more generally of \mathbb{N}^2 , are called *nodes*. We draw a Young diagram as an array of boxes in the plane such that the horizontal axis is oriented left-to-right and the vertical axis top-to-bottom (commonly referred as the *English* notation). For example, $[(3, 2^3, 1)]$ appears as follows.



Motivated by the drawing, we say that the node (r, c) is *above* the node (s, d) , or that (s, d) is *below* (r, c) , if $r < s$. Analogously, we say that (r, c) lies *to the left* of (s, d) , or that (s, d) lies *to the right* of (r, c) , whenever $c < d$.

Fix e is a positive integer. The *content* of a node (r, c) is the integer $c - r$, and the *e -residue* of (r, c) is $\overline{c - r}$. Note that nodes lying in the same diagonal of \mathbb{N}^2 have the same content and therefore the same e -residue for all e . If $\nu \in \mathcal{P}$, we define the *e -content* and we write $\text{cont}_e(\nu)$, for the multiset of e -residues of the nodes in $[\nu]$. For example,

$$\text{cont}_3(3, 2^3, 1) = \{\overline{-4}, \overline{-3}, \overline{-2}, \overline{-2}, \overline{-1}, \overline{-1}, \overline{0}, \overline{0}, \overline{1}, \overline{2}\} = \{\overline{0}, \overline{0}, \overline{1}, \overline{1}, \overline{2}, \overline{2}, \overline{2}, \overline{2}\}.$$

The *rim* of a partition ν is the set of nodes $(r, c) \in [\nu]$ such that $(r + 1, c + 1) \notin [\nu]$. We regard this set as ordered from from SW to NE saying that a node (r, c) comes after (s, d) if (r, c) lies above or to the right of (s, d) . For $h \geq 1$, we define a *removable h -hook* of ν to be a set H of h consecutive nodes $(r_1, c_1), \dots, (r_h, c_h)$ in the rim of ν such that $(r_1, c_1 + 1) \notin [\nu]$ and $(r_h + 1, c_h) \notin [\nu]$. We call (r_1, c_1) and (r_h, c_h) the *hand node* and the *foot node* of H , respectively. By definition the nodes of a

removable h -hook can be removed from $[\nu]$ to leave the Young diagram of a partition of $|\nu| - h$. In particular, if $h = 1$, we call a removable 1-hook simply a *removable node*. Dually, we define the *neighbours* of $[\nu]$ to be the nodes $(s, d) \notin [\nu]$ such that either $s = 1$ or $d = 1$, or $(s - 1, d - 1) \in [\nu]$. The neighbours of ν are naturally ordered in increasing order of content. For $h \geq 1$, a set H of h consecutive neighbours $(s_1, d_1), \dots, (s_h, d_h)$ is called an *addable h -hook* if $s_1 = 1$ or $(s_1 - 1, d_1) \in [\nu]$, and $s_h = 1$ or $(s_h, d_h - 1) \in [\nu]$. As in the dual case, we call (s_1, d_1) the *hand node* and (s_h, d_h) the *foot node* of H . Again we can see that, pictorially, an addable h -hook is a series of nodes which can be added to $[\nu]$ to obtain the Young diagram of a partition of $|\nu| + h$. An addable 1-hook is called an *addable node*.

Fix $n \geq 0$ and $\nu \in \mathcal{P}_n$. We define a ν -*tableau* as a bijection $T : [\nu] \rightarrow \{1, \dots, n\}$ represented by filling the nodes of the Young diagram $[\nu]$ with their images under T . A ν -tableau is called *standard* if its entries increase along rows and down columns.

Example. Let $\nu = (5, 3, 2, 1) \vdash 11$. The following are two ν -tableaux, the one on the left being standard.

1	3	7	8	10
2	5	9		
4	6			
11				


4	9	2	8	10
11	7	3		
1	6			
5				

2.3 Skew-partitions and skew Specht modules

This section gets the reader in touch with the combinatorial protagonist of the paper. Since the algebra of interest – as we shall see in Section 2.7 – relies on two natural numbers, it seems natural to consider a relation of *containment* between pairs of partitions.

Let $\lambda, \mu \in \mathcal{P}$. We say that λ lies inside μ if $[\lambda] \subseteq [\mu]$. In this case we say that $\mu \setminus \lambda$ is a *skew-partition*. We immediately see that, if $\mu \setminus \lambda$ is a skew-partition, then $|\mu| \geq |\lambda|$ with equality holding if and only if $\mu = \lambda$. We call the non-negative integer $|\mu| - |\lambda|$ the *size* of $\mu \setminus \lambda$. Given two non-negative integers $l \leq m$, we set $\mathcal{P}_{l,m} := \{\mu \setminus \lambda \mid \lambda \in \mathcal{P}_l, \mu \in \mathcal{P}_m\}$.

It is useful to visualise skew-partitions as Young diagrams. For a skew-partition $\mu \setminus \lambda$, we define the *skew-diagram* $[\mu \setminus \lambda]$ as the set subset of \mathbb{N}^2 comprising nodes $[\mu] \setminus [\lambda]$. For example, if $\mu = (6, 4^2, 3, 2) \vdash 19$ and $\lambda = (3, 2^3, 1) \vdash 10$, the skew-diagram $[\mu \setminus \lambda]$ comprises the coloured nodes in the picture below.



Note that two different skew-partitions can have the same Young diagram; for example, $[(3, 2) \setminus (3, 1)] = [(2^2, 1) \setminus (2, 1^2)]$.

We remark that if λ lies inside μ , the Young diagram $[\lambda]$ can be recovered by repeatedly deleting removable hooks from $[\mu]$. The union of these hooks comprises the skew-diagram $[\mu \setminus \lambda]$. Note that this series of removals is not unique.

Similarly to the case of partitions, for e a positive integer, we define the *e -content* of a skew-partition $\mu \setminus \lambda$, and we write $\text{cont}_e(\mu \setminus \lambda)$, as the multiset of e -residues of the nodes in $[\mu \setminus \lambda]$, e.g.

$$\text{cont}_4((6, 4^2, 3, 2) \setminus (3, 2^3, 1)) = \{\overline{-3}, \overline{-1}, \overline{0}, \overline{1}, \overline{1}, \overline{2}, \overline{3}, \overline{4}, \overline{5}\} = \{\overline{0}, \overline{0}, \overline{1}, \overline{1}, \overline{1}, \overline{2}, \overline{3}, \overline{3}\}.$$

In conclusion to this section, we generalise the ideas of tableau and standard tableau to the case of skew-partitions. If $\mu \setminus \lambda$ is a skew-partition of size n , a $\mu \setminus \lambda$ -*tableau* is a bijection $T : [\mu \setminus \lambda] \rightarrow \{1, \dots, n\}$ represented by filling the nodes of $[\mu \setminus \lambda]$ with their images under T . A tableau is *standard*

if its entries increase along rows and down columns. Standard *skew-tableaux* play a role in the representation of symmetric groups: if $\mu \setminus \lambda \in \mathcal{P}_{l,m}$, the set of standard $\mu \setminus \lambda$ -tableaux is in bijection with the paths between the Specht modules S^λ and S^μ in the branching graph of symmetric groups (see for example [S, Theorem 2.8.3]). The branching rule itself also ensures that if A is a commutative ring A , then the A -module

$$S^{\mu \setminus \lambda} := \text{Hom}_{AS_l}(S^\lambda, S^\mu \downarrow_{S_l})$$

is non-trivial if and only if $\mu \setminus \lambda$ is a skew-partition, in which the dimension of $S^{\mu \setminus \lambda}$ equals the number of standard $\mu \setminus \lambda$ -tableaux. Unlike in the case of symmetric groups, no closed formula for this number is known. Modules of this kind will come back in Section 2.7 and will be the heart of our treatment.


From now on throughout this paper we will abuse notation by omitting square brackets and not distinguishing between partitions and Young diagrams and between skew-partitions and skew-diagrams.

2.4 The abacus

In this section we introduce an idea of James and Kerber [JK] that gives another combinatorial perspective to the objects mentioned in Sections 2.1 and 2.3. Fix e a positive integer. We consider an abacus display with e vertical runners labelled from left to right by the symbols $0, \dots, e - 1$. We index the *positions* in the i -th runner by the integers $i, i + e, i + 2e, \dots$ from the top down. Then we align runners so that position x is immediately to the right of position $x - 1$ whenever $e \nmid x$. Each position in the abacus is either a *bead* \bullet or a *space* \dagger .

Now consider a partition ν and an integer $C \geq l(\nu)$. We put a bead in the abacus display just constructed at position $\nu_j - j + C$ for $1 \leq j \leq C$. We refer to this drawing as the *abacus configuration* of ν (of charge C) and we denote it by $\text{Ab}_e(\nu)$, or simply $\text{Ab}(\nu)$ if e is clear from the context. For convenience we always choose $C \equiv 0 \pmod{e}$ (this means that the total number of beads in an abacus display is a multiple of e). Observe that increasing C by e entails adding a row of beads at the top of the abacus configuration.

Note that, by construction, every runner of $\text{Ab}(\nu)$ has infinitely many consecutive spaces downwards. When we draw an abacus configuration, we will use the convention that all positions below those shown are spaces. The picture below exhibits the abacus configuration of $(7, 3, 2^2, 1^2) \vdash 16$ in an abacus display with 4 runners and 12 beads.



We remark that given the abacus configuration $\text{Ab}(\nu)$, we can read off the j -th part of ν counting the number of spaces which precede the j -th largest bead position in $\text{Ab}(\nu)$. Note that, by definition, there is a correspondence between the rim of ν and the set of positions $-l(\nu) + 1 + K \leq x \leq \nu_1 - 1 + K$ in $\text{Ab}(\nu)$, where a node (r, c) corresponds to the position marked by $c - r + K$; this position is a bead if (r, c) lies below the next node in the rim of ν , or (r, c) is the last node of the rim, i.e. $r = 1$ and $c = \lambda_1$, and otherwise this position is a space.

This correspondence gives another way to recover the shape of ν – and especially of its rim – from the abacus configuration $\text{Ab}(\nu)$, and will be useful in the next section. We remark that for $h \geq 1$, a removable h -hook arises from a position x such that x is a bead and $x - h$ is a space in $\text{Ab}(\nu)$. The removal of this h -hook from ν corresponds to moving the bead at x into the space at $x - h$. Similarly, adding an addable h -hook corresponds to moving a bead at position x to position $x + h$, for some x . We will say that a removable or addable hook is *at runner i* if it corresponds to a position


$x \equiv i \pmod{e}$. In the abacus configuration above, for example, $(7, 3, 2^2, 1^2)$ has a removable 3-hook at runner 2 since position 6 is a bead and position 3 is a space. Switching these two positions gives the abacus configuration of the partition $(4, 3, 2^2, 1^2) \vdash 13$ obtained by removing nodes $(1, 5), (1, 6)$ and $(1, 7)$ from $(7, 3, 2^2, 1^2)$.

2.5 p -cores and p -blocks

Our main interest in this paper is understanding blocks, and to prepare the ground we recall the combinatorics underlying the p -block theory of the symmetric groups.

Fix an integer $e \geq 2$, we say that a partition is an e -core if it has no removable e -hooks. Given $\nu \in \mathcal{P}$, we define the e -core of ν as the partition obtained by successively deleting removable e -hooks from ν until an e -core is reached. We denote this partition by $\text{core}_e(\nu)$. In terms of the abacus, evaluating $\text{core}_e(\nu)$ coincides with sliding all beads up as far as possible in the runners of $\text{Ab}(\nu)$. Since this process clearly produces a well-defined abacus configuration, we deduce that the e -core of a partition is well-defined. We define the e -weight of ν , written $\text{weight}_e(\nu)$, to be the number of e -hooks removed in evaluating $\text{core}_e(\nu)$. Note that $|\nu| = |\text{core}_e(\nu)| + e \text{weight}_e(\nu)$. For $i = 0, \dots, e - 1$, we write $\nu^{(i)}$ for the partition which corresponds to the i -th runner of $\text{Ab}(\nu)$ considered as an abacus configuration with one single runner (with positions properly numbered). The e -tuple of partitions $(\nu^{(0)}, \dots, \nu^{(e-1)})$ is called the e -quotient of ν . From the abacus interpretation of the weight, we immediately deduce that $\text{weight}_e(\nu) = |\nu^{(0)}| + \dots + |\nu^{(e-1)}|$.

Example. Let $\nu = (12, 7^2, 5, 4, 2, 1^2) \vdash 39$ and $e = 4$. Then $\text{core}_4(\nu) = (2, 1) \vdash 3$, as we see from the following abacus configurations.



We find that $\text{weight}_4(\nu) = 9$, and the 4-quotient of ν is $((2, 1), (1^2), \emptyset, (3, 1))$.

Now let n be a positive integer, and consider the p -blocks of S_n ; that is, the indecomposable summands of the algebra kS_n . It is a standard fact that each Specht module is a module for a single p -block. Conversely, every block contains at least one Specht module (because every simple module is a composition factor of a Specht module), so in order to describe the p -block structure of S_n it suffices to say when two Specht modules lie the same block. The following famous result, known as the *Nakayama conjecture*, gives a combinatorial condition to understand when two Specht modules belong to the same p -block of S_n .

Theorem 2.1. *Let $\alpha, \beta \in \mathcal{P}_n$. The Specht modules S^α, S^β belong to the same p -block of S_n if and only if $\text{core}_p(\alpha) = \text{core}_p(\beta)$.*

Theorem 2.1 was stated by Nakayama in 1940 and proved independently by Brauer and Robinson. Thorough the paper we will abuse notation regarding a p -block B of S_n as the subset of partitions of n which labels the Specht kS_n -modules belonging to B .

Clearly if two partitions of n have the same p -core then they have the same p -weight. Given a p -block B of S_n , we denote by γ_B and w_B , respectively, the p -core and the p -weight of *any* partition in B . It is clear that (the abacus configuration of) all partitions in B can be recovered starting from γ_B and then sliding w_B times beads down in $\text{Ab}(\gamma_B)$ in all possible ways.

We now give some more conditions equivalent to those in Theorem 2.1. Note that nothing of what follows requires p to be a prime number. We need all the abacus configurations to contain the

same number of beads. Since the length of a partition of n is at least n , to ensure this is enough to fix the (common) charge C of the abaci to be at least n , maintaining the convention that $C \equiv 0 \pmod{p}$. For every $\nu \in \mathcal{P}_n$ and every $i = 0, \dots, p-1$, let $b_i(\nu)$ denote the number of beads in the i -th runner of the abacus configuration $\text{Ab}_p(\nu)$. It turns out that the p -tuple $(b_0(\nu), \dots, b_{p-1}(\nu))$ is another invariant of the p -block to which ν belongs. Actually, by definition, the p -core of a partition is obtained by only sliding beads up in its abacus configuration. It follows that the number of beads in every runner remains unchanged at the end of the process. In particular, if α and β are in the same p -block B , then $b_i(\alpha) = b_i(\gamma_B) = b_i(\beta)$ for all $i = 0, \dots, p-1$. (Note that the numbers b_i depend on the choice of the charge; in fact C can be recovered as $b_0 + \dots + b_{p-1}$. If C is increased by p , then each b_i is increased by 1.)

Analysing the integers just defined, we can prove a result we will need later.

Proposition 2.2. *Let B be a p -block of S_n and let $\alpha, \beta \in B$. Suppose that for some $h \geq 1$ and some $i \in \{0, \dots, p-1\}$, α has a removable h -hook at runner i . Then at least one of the following occurs:*

- ◇ α has an addable h -hook at runner $j \equiv i - h \pmod{p}$;
- ◇ β has a removable h -hook at runner i .

The proof of this result is given in [EM2, Proposition 6] for $h = 1$. Our slight generation follows with the same arguments.

Another reinterpretation of the combinatorial condition for two Specht modules to be in the same block concerns p -contents. Recall that the p -content of a partition λ is the multiset of p -residues of the nodes of λ . The following result goes back to Littlewood [L].

Theorem 2.3. *Suppose $\lambda, \mu \in \mathcal{P}_n$. Then λ and μ have the same p -core if and only if they have the same p -content.*

So we can label a p -block B of S_n by a multiset of cardinality n of elements of $\mathbb{Z}/p\mathbb{Z}$, corresponding to the p -content of any partition in B . Using this interpretation, we immediately gain a useful fact which will come in handy in Section 3.2.

Proposition 2.4. *Let m be an integer with $m \geq p$. If $\mu \setminus \lambda \in \mathcal{P}_{m-p, m}$ and $\text{cont}_p(\mu \setminus \lambda) = \{\bar{0}, \dots, \overline{p-1}\}$, then $\text{core}_p(\lambda) = \text{core}_p(\mu)$ and $\text{weight}_p(\lambda) = \text{weight}_p(\mu) - 1$.*

Proof. Let $\nu = \text{core}_p(\lambda)$, let B be the block of S_{m-p} containing λ , and let C be the block of S_m with p -core ν . Observe that the nodes of a rim p -hook have contents $a, a+1, \dots, a+p-1$ for some integer a , and therefore have p -residues $\bar{0}, \bar{1}, \dots, \overline{p-1}$ in some order. Therefore the p -content of C is obtained from the p -content of B by adding one copy of each element $\bar{0}, \bar{1}, \dots, \overline{p-1}$. From Theorem 2.3, this means that μ lies in C . \square

2.6 The degenerate affine Hecke algebra

The next two sections are devoted to introducing the main algebraic objects in the paper. Before meeting the algebra which is at the heart of the treatment, we introduce an auxiliary structure and some interesting related concepts. Let A be a field and n be a positive integer.

Definition. The *degenerate affine Hecke algebra of degree n over A* , denoted by \mathcal{H}_n^A , is the unital, associative A -algebra with generators $z_1, \dots, z_n, s_1, \dots, s_{n-1}$ subject to the following relations:

1. $z_i z_j = z_j z_i$ for all i, j ;
2. $(s_i s_j)^{m_{ij}} = 1$, where $m_{ii} = 1$, $m_{i(i+1)} = 3$ and $m_{ij} = 2$ for $|i - j| > 1$;
3. $s_i z_j = z_j s_i$, for $j \neq i, i+1$;

$$4. \quad s_i z_i = z_{i+1} s_i - 1.$$

Looking at the relation (1), we see that the *polynomial* generators z_1, \dots, z_n generate a subalgebra $A[z_1, \dots, z_n]$ of \mathcal{H}_n^A isomorphic to the polynomial algebra in n indeterminates over A . Moreover, this is a maximal commutative subalgebra of \mathcal{H}_n^A . Considering relations in (2), we also observe that the Coxeter generators s_1, \dots, s_{n-1} generate a subalgebra of \mathcal{H}_n^A isomorphic to AS_n . In fact, as A -modules, $\mathcal{H}_n^A \cong A[z_1, \dots, z_n] \otimes_A AS_n$. We remark that the centre $Z(\mathcal{H}_n^A)$ of \mathcal{H}_n^A is given by the ring of symmetric polynomials in the indeterminates z_1, \dots, z_n [K, Theorem 3.3.1].

We now focus on the polynomial subalgebra in the case where A is an algebraically closed field. Since $A[z_1, \dots, z_n]$ is commutative, every irreducible $A[z_1, \dots, z_n]$ -module M is one-dimensional and corresponds (via a 1:1 correspondence) to the n -tuple $(a_1, \dots, a_n) \in A^n$ such that z_i acts as the scalar a_i on M . We define the *formal character* of a $A[z_1, \dots, z_n]$ -module as the formal \mathbb{Z} -linear combination of the elements of A^n corresponding to its composition factors. The *formal character* of a finite dimensional \mathcal{H}_n^A -module M is then defined as the formal character of its restriction $M \downarrow_{A[z_1, \dots, z_n]}$. More formally, following [K], one identifies $A[z_1, \dots, z_n]$ with the *parabolic subalgebra*

$$\mathcal{H}_{(1, \dots, 1)}^A \cong \mathcal{H}_1^A \otimes_A \cdots \otimes_A \mathcal{H}_1^A$$

of \mathcal{H}_n^A , and defines the formal character $\text{ch}(M)$ of M as the image of the class $[M]$ in the Grothendieck group $K(\mathcal{H}_n^A\text{-mod})$, under the homomorphism

$$\text{ch} : K(\mathcal{H}_n^A\text{-mod}) \longrightarrow K(\mathcal{H}_{(1, \dots, 1)}^A\text{-mod})$$

induced by the restriction from \mathcal{H}_n^A to $\mathcal{H}_{(1, \dots, 1)}^A$. By [K, Theorem 5.3.1], ch is injective, therefore the composition factors of an \mathcal{H}_n^A -module are determined by its formal character. This fact will be particularly useful in what follows due to the strong relationship between the degenerate affine Hecke algebra and the algebra we are going to meet in the next section.

Before going on we associate another important object to any indecomposable \mathcal{H}_n^A -module M . Let (a_1, \dots, a_n) be a summand in the formal character of M . The *central character* of M is defined as the map $Z(\mathcal{H}_n^A) \rightarrow A$ such that $f \mapsto f(a_1, \dots, a_n)$ for all $f \in Z(\mathcal{H}_n^A)$. Since M is indecomposable, all summands of $\text{ch}(M)$ lie in the same S_n -orbit [K, Lemma 4.2.2], so the previous definition does not depend on the choice of the summand (a_1, \dots, a_n) . So we say that $\{a_1, \dots, a_n\}$ is the central character of M and we write $\mathcal{C}(M)$ for it. Then two indecomposable \mathcal{H}_n^A -modules lie in the same block if and only if they have the same central character.

2.7 The centraliser algebra for symmetric groups

We are ready to meet the algebraic protagonist of the present work. Most of the material in this section is taken from [EM2]. Let A be a commutative ring and take m, l non-negative integers with $l \leq m$ and set $n := m - l$. Then AS_l is an A -subalgebra of the group ring AS_m . We define the *centraliser algebra* $\mathcal{C}_{l,m}^A$ to be the centraliser of AS_l in AS_m ; that is, the algebra

$$\mathcal{C}_{l,m}^A := \{a \in AS_m \mid ab = ba \text{ for every } b \in AS_l\}.$$

This algebra has been studied extensively by Ellers and Murray [EM1, EM2], following earlier more general work on centraliser algebras for subgroups of finite groups by Ellers. However, $\mathcal{C}_{l,m}^A$ remains poorly understood, and in particular its blocks are not yet known. This is the focus of the present paper.

By the very effective description in [K, Section 2.1] we know a straightforward generating set for $\mathcal{C}_{l,m}^A$. This is the union of three subsets:

- ◊ the centre $Z(AS_m)$;

- ◇ the symmetric group (isomorphic to S_n) on the symbols $\{l+1, \dots, m\}$;
- ◇ the set of *Jucys–Murphy elements* $L_j = (1\ j) + (2\ j) + \dots + (j-1\ j)$ for $j = l+1, \dots, m$.

We introduce a family of $\mathcal{C}_{l,m}^A$ -modules which will be at the heart of our treatment. Suppose that $\lambda \vdash l$ and $\mu \vdash m$. Then the A -module $S^{\mu \setminus \lambda} = \text{Hom}_{AS_l}(S^\lambda, S^\mu \downarrow_{S_l})$ introduced at the end of Section 2.3 is naturally a module for $\mathcal{C}_{l,m}^A$, via the action

$$(c\phi)(v) = c\phi(v) \quad \text{for } c \in \mathcal{C}_{l,m}^A, \phi \in S^{\mu \setminus \lambda}, v \in S^\lambda.$$

As already remarked, $S^{\mu \setminus \lambda}$ is non-trivial if and only if $\mu \setminus \lambda \in \mathcal{P}_{l,m}$. To emphasise the similarity with the symmetric group case, we refer to these $\mathcal{C}_{l,m}^A$ -modules as *Specht modules*.

We now recall our p -modular system (R, \mathbb{F}, k) , where p is the prime characteristic of the residue field k . Since $\text{char}(\mathbb{F}) = 0$ the algebras $\mathbb{F}S_m$ and $\mathbb{F}S_l$ are both semisimple by Maschke’s Theorem. The general theory of centraliser algebras then implies that $\mathcal{C}_{l,m}^{\mathbb{F}}$ is semisimple as well, with a complete irredundant set of simple modules given by the set

$$\{S^{\mu \setminus \lambda} \mid \mu \setminus \lambda \in \mathcal{P}_{l,m}\}$$

of Specht modules.

Things change if the fraction field \mathbb{F} is replaced by the residue field k : the centraliser algebra $\mathcal{C}_{l,m}^k$ is in general not semisimple and its Specht modules generally fail to be simple. The decomposition number problem for $\mathcal{C}_{l,m}^k$ asks for the composition factors of the Specht modules; unfortunately, at present we do not have a good labelling of simple module for $\mathcal{C}_{l,m}^k$. This paper is concerned with determining the p -blocks of $\mathcal{C}_{l,m}^k$. The Specht module $S^{\mu \setminus \lambda}$ over k is a p -modular reduction of the corresponding (simple) Specht module over \mathbb{F} . So (analogously to the case of symmetric groups) describing the p -block structure of $\mathcal{C}_{l,m}^k$ reduces to finding the appropriate partition of the set of Specht $\mathcal{C}_{l,m}^k$ -modules. For this, we use an approach pioneered by Ellers and Murray, which we describe in the next section.

2.8 Combinatorial blocks

Let l, m, n be as in Section 2.7. We have pointed out that in order to find the decomposition of $\mathcal{C}_{l,m}^k$ into p -blocks, we need to look for a way to partition the set of Specht $\mathcal{C}_{l,m}^k$ -modules. We know by Theorem 2.1 how the Specht modules for the symmetric group split into p -blocks, this is a characterisation based on the concept of p -core. Our main conjecture (which is implicit in the work of Ellers and Murray [EM2]) predicts that the case of the centraliser algebra works in the same way.

Conjecture 2.5. *Let $\mu_1 \setminus \lambda_1, \mu_2 \setminus \lambda_2 \in \mathcal{P}_{l,m}$. The Specht $\mathcal{C}_{l,m}^k$ -modules $S^{\mu_1 \setminus \lambda_1}, S^{\mu_2 \setminus \lambda_2}$ belong to the same p -block of $\mathcal{C}_{l,m}^k$ if and only if $\text{core}_p(\mu_1) = \text{core}_p(\mu_2)$ and $\text{core}_p(\lambda_1) = \text{core}_p(\lambda_2)$.*

We can consider Conjecture 2.5 in terms of idempotents. We say that a non-zero central idempotent of $\mathcal{C}_{l,m}^k$ is a *block idempotent* if it is primitive, i.e. it cannot be written as the sum of two non-zero central idempotents. The blocks of $\mathcal{C}_{l,m}^k$ are then precisely the algebras $e\mathcal{C}_{l,m}^k$ for e a primitive central idempotent.

In the case of the symmetric group S_n , Theorem 2.1 tells that there is a bijection between the set of p -block idempotents of kS_n and the set of p -core partitions which are the p -core of at least one partition in \mathcal{P}_n . Given p -blocks B and B' of kS_m and kS_l respectively, let e_B and $e_{B'}$ the corresponding p -block idempotents. Then it is straightforward to see that $e_B e_{B'}$ is a central idempotent of $\mathcal{C}_{l,m}^k$, so is a sum of block idempotents, and $e_B e_{B'} \mathcal{C}_{l,m}^k$ is a (possibly zero) sum of p -blocks of $\mathcal{C}_{l,m}^k$. Moreover, $e_B e_{B'}$ acts as the identity on $S^{\mu \setminus \lambda}$ if S^μ lies in B and S^λ lies in B' (that is, if $\text{core}_p(\mu) = \gamma_B$ and $\text{core}_p(\lambda) =$

$\gamma_{B'}),$ and otherwise $e_B e_{B'} S^{\mu \setminus \lambda} = 0$. If $e_B e_{B'} \neq 0$, then we refer to $e_B e_{B'} C_{l,m}^k$ as a *combinatorial block* of $C_{l,m}^k$. Conjecture 2.5 then predicts that every combinatorial block is a p -block. Since the sum of the elements $e_B e_{B'}$ over all choices of B and B' is 1, this would then mean that every p -block of $C_{l,m}^k$ is a combinatorial block. We will abuse notation by identifying a combinatorial block with the set of Specht modules in that combinatorial block, or with the set of labelling skew-partitions.

In [EM2], Ellers and Murray prove Conjecture 2.5 in the case $n \leq 3$. The aim of the present paper is to prove Conjecture 2.5 for a large family of combinatorial blocks of $C_{l,m}^k$. We follow the idea of Ellers and Murray [EM2, Section 1.2] to exploit the relationship between the centraliser algebra $C_{l,m}^k$ and the degenerate affine Hecke algebra \mathcal{H}_n^k . Their work shows that every $C_{l,m}^k$ -module naturally admits the structure of an \mathcal{H}_n^k -module: the symmetric group on $\{l+1, \dots, m\}$ and the Jucys–Murphy elements L_{l+1}, \dots, L_m generate a subalgebra of $C_{l,m}^k$ which is a quotient of \mathcal{H}_n^k ; restricting to this subalgebra and then inflating yields an \mathcal{H}_n^k -module. With respect to this structure, every Specht module is equipped with a formal character and a central character and it is an easy task to combinatorially evaluate these. Crucially, two $C_{l,m}^k$ -modules in the same combinatorial block have the same composition factors if and only if the corresponding \mathcal{H}_n^k -modules do. This enables us to work with formal characters of Specht modules to prove the p -block structure.

Let $\mu \setminus \lambda \in \mathcal{P}_{l,m}$ and consider a standard $\mu \setminus \lambda$ -tableau T . We associate to T the permutation $a_T = (a_1, \dots, a_n)$ of $\text{cont}_p(\mu \setminus \lambda)$ such that a_j is the p -residue of the node containing with number j in T , for $j = 1, \dots, n$. The formal character of the Specht module $S^{\mu \setminus \lambda}$, we denote by $\text{ch}(\mu \setminus \lambda)$, is the formal sum of the n -tuples a_T with T varying in the set of standard $\mu \setminus \lambda$ -tableaux.

Example. Let $p = 3$ and $\mu \setminus \lambda = (4^2, 2) \setminus (2^2, 1) \in \mathcal{P}_{5,10}$. The formal character of the Specht $C_{5,10}^k$ -module $S^{\mu \setminus \lambda}$ is as follows (we draw the skew-diagram $\mu \setminus \lambda$ filling every node with its 3-residue, omitting overlines):

		2	0
		1	2
	2		

$$2 \cdot (2, 2, 0, 1, 2) + 2 \cdot (2, 2, 1, 0, 2) + 2 \cdot (2, 0, 1, 2, 2) + 2 \cdot (2, 1, 0, 2, 2) + (2, 0, 2, 1, 2) + (2, 1, 2, 0, 2).$$

By construction, every term of the formal character $\text{ch}(\mu \setminus \lambda)$ is a permutation of the p -content of $\mu \setminus \lambda$. The central character of the Specht module $S^{\mu \setminus \lambda}$, as an \mathcal{H}_n^k -module, then exactly coincides with the multiset $\text{cont}_p(\mu \setminus \lambda)$. We observe the following fact.

Proposition 2.6. *Let $\mu_1 \setminus \lambda_1$ and $\mu_2 \setminus \lambda_2$ be skew-partitions in the same combinatorial block of $C_{l,m}^k$. Then $\text{cont}_p(\mu_1 \setminus \lambda_1) = \text{cont}_p(\mu_2 \setminus \lambda_2)$.*

Proof. This follows from Theorem 2.3. □

Motivated by the above result, we say that a combinatorial block B has central character $\mathcal{C}(B)$ equal to the p -content of *any* skew-partition in B .

One of the main tools in the paper is the *decomposition matrix*. Let B be a combinatorial block and let D_B be the matrix whose rows are indexed by B and columns by those simple \mathcal{H}_n^k -modules appearing as composition factors of some Specht module in B . To detect the entries of the row of D_B labelled by the skew-partition $\mu \setminus \lambda$, write $\text{ch}(\mu \setminus \lambda)$ as a \mathbb{Z} -linear combination of formal characters of simple \mathcal{H}_n^k -modules. The coefficients in this expression give the desired decomposition numbers. B is then a p -block of $C_{l,m}^k$ if and only if the decomposition matrix D_B is a connected matrix (i.e. if it cannot be put in block-diagonal form by permuting rows and columns).

We end this section and this chapter with a crucial remark.

Remark. Given a skew-partition $\mu \setminus \lambda$, fill the boxes of the skew-diagram $\mu \setminus \lambda$ with their p -residues, and define the p -shape of $\mu \setminus \lambda$ to be the multiset of connected components of the resulting diagram. It is easy to see that if two skew-partitions in B have the same p -shape, then the corresponding Specht modules have the same formal character, and therefore the same composition factors (with multiplicity) as \mathcal{H}_n^k -modules. This means that for our purposes, we can largely shorten the number of rows of D_B relabelling them with the set \mathcal{B} that comes out quotienting B by the equivalence relation that identifies skew-partitions with the same p -shape. We say that \mathcal{B} is the set of p -shapes of B . We regard the elements of \mathcal{B} as \mathcal{H}_n^k -modules and when talking about the formal character and the composition factors of a p -shape X we mean those of any Specht module in B whose p -shape is X . Also, saying that a p -shape belongs to a combinatorial block we mean that the combinatorial block contains a skew-partition with that p -shape.

When drawing p -shapes, we follow [EM2] by drawing the connected components in a diagonal line in arbitrary order.

For example, take $p = 3$, and consider the skew-partitions $(4, 1) \setminus (2)$ and $(3, 2) \setminus (2)$:




Both skew-partitions have the same 3-shape and the corresponding Specht modules both have formal character $(2, 0, 2) + 2 \cdot (2, 2, 0)$.

3 Ribbon blocks and belt blocks

This chapter is devoted to establishing Conjecture 2.5 for a special family of combinatorial blocks of the centraliser algebra. Again let m, l be non-negative integers with $m \geq l$ and let $n = m - l$. Recall that (R, \mathbb{F}, k) is a p -modular system where p denotes the (prime) characteristic of the residue field k . Let B be a combinatorial block of $\mathcal{C}_{l,m}^k$. The central character of B is a multiset of elements of $\mathbb{Z}/p\mathbb{Z}$. We will be concerned with the case where there are no repeated entries in the central character, so that the central character is a subset of $\mathbb{Z}/p\mathbb{Z}$. In this case clearly $n \leq p$ and we say that B is a generalised ribbon block if $n < p$ (so that the central character is a proper subset of $\mathbb{Z}/p\mathbb{Z}$), or a belt block if $n = p$. We will assume for the rest of Section 3 that B is either a generalised ribbon block or a belt block.

In the last section we explained that when working modulo p we can identify skew-partitions with the same formal character (since the rows of the decomposition matrix D_B labelled by them are identical), so from now on we will be concerned with just the set \mathcal{B} of p -shapes of B . Firstly it is a good idea to give a visualisation of the elements of \mathcal{B} . If $\mu \setminus \lambda$ is a skew-partition in B , then its skew-diagram does not contain a 2×2 -subdiagram (because otherwise it would have boxes with equal p -residue, contrarily to our assumption on B) and hence it is a disjoint union of rim-hooks of μ . If X is the p -shape of $\mu \setminus \lambda$, we refer to a connected component of X for any of these rim-hooks and we say that X is a ribbon if $\mu \setminus \lambda$ is connected, i.e. it is a single rim-hook of μ . For example, if $p > n = 3$, and B has central character $\{\bar{0}, \bar{1}, \bar{2}\}$, then the p -shapes in \mathcal{B} are among the nine drawings below (the first four on the left being the possible ribbons).




We close this long preface noting that, since there are no repetitions in $\mathcal{C}(B)$, if X is a p -shape in B , there is no risk of confusion writing \boxed{i} for the unique node labelled by i in X , for every $i \in \mathcal{C}(B)$.

3.1 Generalised ribbon blocks

In this section we fix a generalised ribbon block B , with $\mathcal{C} = \mathcal{C}(B) \subset \mathbb{Z}/p\mathbb{Z}$ its central character. Our aim is to prove Conjecture 2.5 for B .

Before starting we give another combinatorial perspective to the p -shapes in a generalised ribbon block. The reason why we do this will be clarified in the next section. We define an *arrow graph* to be a directed graph with vertex set \mathcal{C} such that every arrow has the form $i \rightarrow i + 1$ or $i + 1 \rightarrow i$. Given a p -shape X , we define an associated arrow graph Γ_X by drawing an arrow $i \rightarrow i + 1$ [resp. $i + 1 \rightarrow i$] whenever \overline{i} lies on the left of [resp. below] $\overline{i+1}$ in X , or no arrow between i and $i + 1$ if these nodes lie in different connected components of X . It is clear that the set of all possible p -shapes is in bijection with the set of all possible arrow graphs. We define a p -shape to be *maximal* if its arrow graph is maximal, i.e. contains an arrow between i and $i + 1$ whenever $i, i + 1 \in \mathcal{C}$. Note that starting from an arrow graph we can recover the corresponding p -shape by mirroring the above procedure. Moreover, we can evaluate its formal character by taking the formal sum of all permutations (a_1, \dots, a_n) of \mathcal{C} with the property that i appears before j for every arrow $i \rightarrow j$.


Example. Suppose $p = 7$ and $\mathcal{C} = \{\overline{0}, \overline{1}, \overline{2}, \overline{3}, \overline{6}\}$. We give an example of a p -shape and the corresponding arrow graph (omitting overlines).



The corresponding formal character is then (omitting bars)

$$\begin{aligned} & (0, 1, 3, 2, 6) + (0, 1, 3, 6, 2) + (0, 1, 6, 3, 2) + (0, 3, 1, 2, 6) + (0, 3, 1, 6, 2) \\ & + (0, 3, 6, 1, 2) + (0, 6, 1, 3, 2) + (0, 6, 3, 1, 2) + (1, 0, 3, 2, 6) + (1, 0, 3, 6, 2) \\ & + (1, 0, 6, 3, 2) + (1, 3, 0, 6, 2) + (1, 3, 0, 2, 6) + (1, 3, 2, 0, 6) + (3, 0, 1, 2, 6) \\ & + (3, 0, 1, 6, 2) + (3, 0, 6, 1, 2) + (3, 1, 0, 6, 2) + (3, 1, 0, 2, 6) + (3, 1, 2, 0, 6). \end{aligned}$$


Now we introduce a fundamental tool in our treatment. Let X, Y be two p -shapes and let $D(X, Y)$ the subset of \mathcal{C} containing those integers i such that $i + 1 \in \mathcal{C}$ and \overline{i} lies on the left of $\overline{i+1}$ in X and below $\overline{i+1}$ in Y (see the picture below), or vice versa.



Define the *distance* $d(X, Y)$ between X and Y as the cardinality of $D(X, Y)$. This concept can be rephrased also in terms of arrow graphs: take Γ_X and Γ_Y and denote by $\Gamma_X \cup \Gamma_Y$ the picture obtained by overlapping these two arrow graphs. Then $d(X, Y)$ coincides with the number of edges where two differently oriented arrows overlap in $\Gamma_X \cup \Gamma_Y$.

Note that d is *not* a metric: $d(X, Y)$ can be zero even when $X \neq Y$.

Example. Suppose $p \geq 11$ and $\mathcal{C} = \{\overline{0}, \overline{1}, \overline{2}, \overline{3}, \overline{4}, \overline{5}, \overline{6}, \overline{7}, \overline{8}, \overline{9}\}$ and consider the two following p -shapes:



Then $D(X, Y) = \{\bar{1}, \bar{3}, \bar{7}\}$ and $d(X, Y) = 3$. The union of the two arrow graphs $\Gamma_X \cup \Gamma_Y$ is given as follows, with Γ_X in red and Γ_Y in blue.



We start our analysis of generalised ribbon blocks by observing that the formal characters of two different maximal p -shapes have no summands in common.

Lemma 3.1. Consider $\mathbf{a} = (a_1, \dots, a_n)$ where a_1, \dots, a_n are the elements of \mathcal{C} in some order. Then \mathbf{a} appears in the formal character of exactly one maximal p -shape.

Proof. We construct the unique maximal p -shape X such that \mathbf{a} is a term of the formal character $\text{ch}(X)$. Start drawing \boxed{i} , for some $i \in \mathcal{C}$ such that $i - 1 \notin \mathcal{C}$. Then consider the order of i and $i + 1$ in \mathbf{a} : if $i + 1$ occurs before [resp. after] i , then draw $\boxed{i+1}$ just above [resp. to the right of] the node \boxed{i} . Now do the same for $i + 1$ and $i + 2$, and continue until an element $k \in \mathcal{C}$ is reached for which $k + 1 \notin \mathcal{C}$. This determines one component of X . Now repeat with the other elements $i \in \mathcal{C}$ for which $i - 1 \notin \mathcal{C}$ to obtain the other components.


By construction, \mathbf{a} is a summand of $\text{ch}(X)$. We now see that X is unique. If Y is different maximal p -shape, then the set $D(X, Y)$ is non-empty, because Γ_X and Γ_Y both have arrows joining i and $i + 1$ whenever $i, i + 1 \in \mathcal{C}$. Take $i \in D(X, Y)$, and suppose that i appears before $i + 1$ in \mathbf{a} (the other case being similar). This means that we have an arrow $i \rightarrow i + 1$ in Γ_X , and therefore an arrow $i \leftarrow i + 1$ in Γ_Y , so that \mathbf{a} does not appear in $\text{ch}(Y)$. \square

Corollary 3.2. Different maximal p -shapes have no composition factors in common.


The next three results will finally give us a labelling set for the columns of the decomposition matrix of a ribbon block. In Lemma 3.3, if X is a p -shape, a standard X -tableau indicates a standard tableau of any skew-partition with p -shape X . We begin with a definition.

Definition. Let X be a p -shape. A refinement of X is a maximal p -shape Y that can be obtained joining the connected components of X together.

Example. Suppose $p = 7$, and let X be the p -shape from an earlier example:



Then X has exactly two refinements, as follows.



Lemma 3.3. Let X be a p -shape. The formal character of X is the sum of the formal characters of the refinements of X .

Proof. This follows from a correspondence between standard tableaux: given a refinement Y of X , each standard Y -tableau gives rise to a standard X -tableau (just by breaking into individual connected components). Conversely, each standard X -tableau arises in this way for a unique refinement Y : for each i such that the nodes \boxed{i} and $\boxed{i+1}$ lie in different connected components of X , the relative order of the entries $i, i + 1$ tells us how they need to be joined together if the resulting tableau is to be standard. \square

Corollary 3.4. *If X is a p -shape, then the composition factors of X are the composition factors of the refinements of X .*

Proof. This follows from the fact that the formal characters of simple modules of the degenerate affine Hecke algebra are linearly independent [K, Theorem 5.3.1]. \square

Theorem 3.5. *Let X and Y be two p -shapes. The following are equivalent:*

1. X and Y have a composition factor in common;
2. The formal characters of X and Y have a term in common;
3. $d(X, Y) = 0$;
4. There is a maximal p -shape which is a refinement of both X and Y .

Proof.

(1 \Rightarrow 2) This follows from the fact that the formal character of a \mathcal{H}_n^k -module is the sum of the formal characters of its composition factors.

(2 \Rightarrow 3) If there exists an integer $i \in D(X, Y)$, then $i + 1$ occurs before i in every summand of $\text{ch}(X)$ and after i in every summand of $\text{ch}(Y)$, or vice versa. So $\text{ch}(X)$ and $\text{ch}(Y)$ have no terms in common.

(3 \Rightarrow 4) Consider the graph $\Gamma_X \cup \Gamma_Y$. Since $d(X, Y) = 0$, this is an admissible arrow graph. Then every arrow graph obtained by filling the vacant edges in $\Gamma_X \cup \Gamma_Y$ corresponds to a refinement of both X and Y .

(4 \Rightarrow 1) This follows from Corollary 3.4. \square

Theorem 3.5 tells us that the set of maximal p -shapes serves as a labelling set for an approximation to the decomposition matrix of B : by Corollary 3.4 any simple module S occurs as a composition factor of a unique maximal p -shape Y (say with multiplicity c). So by Corollary 3.4, if X is any p -shape then S occurs as a composition factor of X with multiplicity c if Y is a refinement of X , and 0 otherwise.

So we can define a matrix D_B with rows indexed by the set of p -shapes, and columns indexed by the set of maximal p -shapes, with entry $d_{XY} = 1$ if Y is a refinement of X and 0 otherwise. Then the actual decomposition matrix of B is obtained from D_B by possibly duplicating rows (because a given p -shape may correspond to more than one Specht module) and possibly duplicating and rescaling columns (because a Specht module labelled by a maximal p -shape might be reducible). As a consequence, to show that B is a single p -block of $\mathcal{C}_{l,m}^k$, we just need to show that D_B is a connected matrix, and our problem becomes purely combinatorial.

(In fact we suspect that every Specht module labelled by a maximal p -shape is irreducible, but we could not find a proof of this statement, and we do not need it.)

Example. Suppose $p = 5$, $l = 5$ and $m = 8$, and let B be the block consisting of Specht modules $S^{\mu \setminus \lambda}$ where $\text{core}_5(\mu) = (2, 1)$ and $\text{core}_5(\lambda) = \emptyset$. Then $\mathcal{C} = \{\bar{0}, \bar{1}, \bar{4}\}$, and the p -shapes in B are as follows:

$$X_1 = \begin{array}{|c|c|c|} \hline 4 & 0 & 1 \\ \hline \end{array}, \quad X_2 = \begin{array}{|c|c|} \hline 0 & 1 \\ \hline 4 & \\ \hline \end{array}, \quad X_3 = \begin{array}{|c|c|} \hline 0 & 1 \\ \hline 4 & \\ \hline \end{array}, \quad X_4 = \begin{array}{|c|} \hline 1 \\ \hline 0 \\ \hline 4 \\ \hline \end{array}, \quad X_5 = \begin{array}{|c|} \hline 1 \\ \hline 0 \\ \hline 4 \\ \hline \end{array}.$$

(For example, the p -shape X_2 arises from the skew-partitions $(7, 1) \setminus (5)$ and $(5, 3) \setminus (4, 1)$.)


The matrix D_B is then given as follows.

	X_1	X_3	X_5
X_1	1	0	0
X_2	1	1	0
X_3	0	1	0
X_4	0	1	1
X_5	0	0	1


Definition. Let X, Y be two p -shapes in \mathcal{B} . We say that X and Y are p -linked if there exists a series of p -shapes $X = X_1, X_2, \dots, X_{t-1}, X_t = Y$ in \mathcal{B} such that $d(X_j, X_{j+1}) = 0$ for every $1 \leq j \leq t - 1$.

Our aim in this section is to show that any two p -shapes in B are p -linked. To do this, we will mainly work with abacus configurations. As remarked in Section 2.3, a skew-partition $\mu \setminus \lambda$ in B (that contains no 2×2 -subdiagrams) can be thought as the union of a family of rim-hooks of μ whose removal gives λ (recall that each of these rim-hooks corresponds to a connected component of the p -shape of $\mu \setminus \lambda$). If H is one of them and $h, f \in \mathbb{Z}$ are the positions of $\text{Ab}(\mu)$ corresponding to the hand node and foot node of H , respectively, then h is a bead and $f - 1$ is a space (and vice versa in $\text{Ab}(\lambda)$). We represent the bead at position f in $\text{Ab}(\mu)$ by a new symbol \star called a *star bead*. (So the star beads are precisely the beads occurring in the abacus configuration of μ but not of λ .)

Example. Take $p = 5$ and $\mu \setminus \lambda = (7, 3, 1^2) \setminus (7, 2)$, giving the following p -shape.



This p -shape has two connected components, so that $\mu \setminus \lambda$ comprises two rim-hooks. We draw abacus configurations of μ and λ as follows, with star beads in μ .



Now we can give our main result.

Theorem 3.6. Let $X, Y \in \mathcal{B}$ such that $d(X, Y) > 0$. Then there is $Z \in \mathcal{B}$ such that $d(X, Z) < d(X, Y)$ and $d(Y, Z) < d(X, Y)$.

Proof. For every $j \in \mathcal{C}$, consider the node \boxed{j} in $\mu_X \setminus \lambda_X$ and $\mu_Y \setminus \lambda_Y$ and denote by x_j and y_j the positions corresponding to these nodes in the abacus display, respectively. (Note that by the choice of the charge, positions x_j and y_j are both in the j -th runner.) Fix $i \in D(X, Y)$, and assume without loss of generality that the node \boxed{i} lies on the left of $\boxed{i+1}$ in X and below it in Y . Then x_i is a space in $\text{Ab}(\mu_X)$ and y_i is a bead (and *not* a star bead) in $\text{Ab}(\mu_Y)$. We let r be maximal such that $i, i + 1, \dots, i + r \in \mathcal{C}$ (where we add modulo p); note that there must be such an r because by assumption $\mathcal{C} \neq \mathbb{Z}/p\mathbb{Z}$. We prove that we can find the required p -shape Z by contradiction, via the following claim.

Claim 1. Suppose that the required p -shape Z does not exist. Then for every $k \in \{i + 1, \dots, i + r\}$ with $k \notin D(X, Y)$, the following statements hold.

- ◇ If x_k is a bead or a star bead in $\text{Ab}(\mu_X)$, then
 - μ_X has no addable $(k - i)$ -hooks at runner i , and
 - λ_Y has a removable $(k - i)$ -hook at runner k .
- ◇ If y_k is a space or a star bead in $\text{Ab}(\mu_Y)$, then
 - μ_X has an addable $(k - i)$ -hook at runner i , and
 - λ_Y has no removable $(k - i)$ -hooks at runner k .

Note that the condition that x_k is a bead or star bead in $\text{Ab}(\mu_X)$ is the same as saying that $\boxed{k+1}$ does not lie immediately to the right of \boxed{k} in X . Similarly, the condition that y_k is a space or star bead in $\text{Ab}(\mu_Y)$ means that $\boxed{k+1}$ does not lie immediately above \boxed{k} in Y .

Claim 1 then gives the desired contradiction to complete the proof: if we let $k = i + r$, then x_k and y_k are star beads in $\text{Ab}(\mu_X)$ and $\text{Ab}(\mu_Y)$ respectively because $i + r + 1 \notin \mathcal{C}$. So Claim 1 cannot be true in the case $k = i + r$.

We prove Claim 1 by induction on k . So take $k \in \{i + 1, \dots, i + r\} \setminus D(X, Y)$, and assume that Claim 1 holds for all smaller values of k . The following statement is crucial.

Claim 2. Suppose k is as in Claim 1, and that the inductive hypothesis holds. Then position $x_k + i - k$ is a space in $\text{Ab}(\mu_X)$.

Proof. Consider the node \boxed{k} in X . We have that $x_k = x_i - i + k + tp$, for some $t \in \mathbb{Z}$. If $t = 0$, then the claim is trivially true since we are supposing that x_i is a space. If $t \neq 0$, then the nodes \boxed{i} and \boxed{k} lie in different connected components of X . Let $j \in \{i + 2, \dots, k\}$ be such that \boxed{j} is the foot node of the connected component of X containing \boxed{k} . It follows that $\boxed{j-1}$ is the hand node of a connected component of X , so that x_{j-1} is a star bead in $\text{Ab}(\mu_X)$ and $j - 1 \notin D(X, Y)$. By the inductive hypothesis, μ_X has no addable $(j - 1 - i)$ -hooks at runner i , so position $x_j + i - j$ must be a space since $x_j - 1$ is a space in $\text{Ab}(\mu_X)$. Finally, since \boxed{j} and \boxed{k} lie in the same connected component of X , we see that $x_j = x_k + j - k$. It follows that $x_j + i - j = x_k + i - k$, proving what we have claimed.

With a perfectly dual argument it can be shown that the inductive hypothesis forces the position $y_k + i - k$ to be a bead in $\text{Ab}(\lambda_Y)$.

With Claim 2 established, we deal with the inductive step of Claim 1. We consider only the case when x_k is a bead or a star bead in $\text{Ab}(\mu_X)$, the other case being similar. By Claim 2, $x_k + i - k$ is a space in $\text{Ab}(\mu_X)$, and so μ_X has a removable $(k - i)$ -hook at position x_k . Assume that there exists $a_1 \equiv i \pmod{p}$ such that μ_X has an addable $(k - i)$ -hook at a_1 . Let $1 \leq k_1 \leq k - i$ be the least integer such that $k - k_1 \notin D(X, Y)$, position x_{k-k_1} is a space and $a_1 + k - i - k_1$ is a bead in $\text{Ab}(\mu_X)$. The assumptions that $k - k_1 \notin D(X, Y)$ and x_{k-k_1} is a space mean that y_{k-k_1} is a space or a star bead in $\text{Ab}(\mu_Y)$. So if $k_1 < k - i$, then the inductive hypothesis says that there exists an integer $a_2 \equiv i$ such that μ_X has an addable $(k - i - k_1)$ -hook at position a_2 . Then we repeat the above process, letting $1 \leq k_2 \leq k - i - k_1$ be minimal such that $k - k_1 - k_2 \notin D(X, Y)$, $x_{k-k_1-k_2}$ is a space and $a_2 + k - i - k_1 - k_2$ is a bead in $\text{Ab}(\mu_X)$. It is clear that this procedure always terminates at a step s such that $i = k - \sum_{1 \leq u \leq s} k_u$. Summarising, we have found integers $a_1, \dots, a_s \equiv i \pmod{p}$ and k_1, \dots, k_s such that, for every $1 \leq v \leq s$, in $\text{Ab}(\mu_X)$, there is a bead at position $c_v := a_v + k - i - \sum_{1 \leq u \leq v} k_u$ and a space at position $d_v := a_v + k - i - \sum_{1 \leq u \leq v-1} k_u$. Let $i + 1 \leq j_1 < \dots < j_t \leq k$ be those residues such that $\boxed{j_1}, \dots, \boxed{j_t}$ are hand nodes of connected components of X .


We now construct a new abacus configuration. We start from $\text{Ab}(\mu_X)$ and make the following moves:

- ◇ move the bead (or star bead) at x_k to the space at $x_{j_t+1} - 1$,

- ◇ for every $2 \leq v \leq t$, move the star bead at x_{j_v} to the space at $x_{j_{v-1}+1} - 1$,
- ◇ move the star bead at x_{j_1} to the space at x_i ,
- ◇ for every $1 \leq v \leq s$, move the bead at c_v to the space at d_v .

Note that in terms of the Young diagram, the first three types of moves simply entail removing all the nodes $\boxed{i+1}, \dots, \boxed{k}$ from μ_X , and the moves of the last type entail putting these nodes back in new positions. So if we let μ_Z be the partition defined by this new abacus configuration, then $\lambda_X \subset \mu_Z$, and we have a skew-partition $\mu_Z \setminus \lambda_X$ in B . We let Z be the p -shape of this skew-partition. We will show that in most cases, Z is the p -shape desired in the theorem.

The diagram below illustrates the changes made in the abacus. Note that the rows of the diagram in which the given positions appear are illustrative only; these rows could easily appear in a different order in practice.



For every $j \in \mathcal{C}$, let z_j be the position of the node \boxed{j} in $\text{Ab}(\mu_Z)$. Now take $j \in \{i+1, \dots, k-1\}$ with $j \notin D(X, Y)$. Then we claim that $j \notin D(X, Z)$ and $j \notin D(Y, Z)$. If x_j is a bead or a star bead then by the inductive hypothesis μ_X has no addable $(j-i)$ -hooks at runner i , so that z_j is a bead in $\text{Ab}(\mu_Z)$. If x_j is a space, then the choice of k_1, \dots, k_s means that z_j is a space or a star bead. This gives $j \notin D(X, Z)$. We deduce similarly that $j \notin D(Y, Z)$ (note that the inductive hypothesis means that it cannot be the case that x_j is a star bead and y_j is a space).

So our claim is true. Additionally, it is easy to see that $i \notin D(X, Z)$ and $i \notin D(Y, Z)$. Furthermore, $k \notin D(X, Z)$ because x_k and z_k are both beads. It follows that $d(X, Z) < d(X, Y)$, and that $d(Y, Z) < d(X, Y)$ unless y_k is a space in $\text{Ab}(\lambda_Y)$ and, in $\text{Ab}(\mu_X)$, x_k is a star bead and $a_1 = x_{k+1} + k - i - 1$ is the only addable $(k-i)$ -hook at runner i . If this is not the case, then Z is the desired p -shape and we conclude. Hence assume the opposite from now on.

We are under the assumption that y_k is a space in $\text{Ab}(\lambda_Y)$. Then λ_Y has an addable $(k-i)$ -hook at runner i because position $y_k + i - k$ is a bead by the dual version of Claim 2. The fact that μ_X has a removable $(k-i)$ -hook at runner k and exactly an addable $(k-i)$ -hook at runner i , together with Proposition 2.2, implies that μ_Y has a removable $(k-i)$ -hook at runner k . Say that this removable hook occurs at position $r_1 \equiv k \pmod{p}$. Note that $r_1 \neq y_k$ because y_k is a space, so r_1 is a removable $(k-i)$ -hook at runner k of λ_Y as well. Let $1 \leq k_1 \leq k-i$ with $k-k_1 \notin D(X, Y)$ be the least integer such that y_{k-k_1} is a bead and $r_1 - k_1$ is a space in $\text{Ab}(\lambda_Y)$. If $k_1 < k-i$, the inductive hypothesis ensures that λ_Y has a removable $(k-i-k_1)$ -hook at some position $r_2 \equiv r_1 - k_1 \pmod{p}$. Then we define $1 \leq k_2 \leq$

$k - i - k_1$ with $k - k_1 - k_2 \notin D(X, Y)$ to be minimal such that $y_{k-k_1-k_2}$ is a bead and $r_2 - k_2$ is a space in $\text{Ab}(\lambda_Y)$. We continue in this way; the process terminates at a step s such that $i = k - \sum_{1 \leq u \leq s} k_u$. Summarising we have collected integers k_1, \dots, k_s and r_1, \dots, r_s with $r_v \equiv k - \sum_{1 \leq u \leq v-1} k_u \pmod{p}$, for $1 \leq v \leq s$, such that, in $\text{Ab}(\lambda_Y)$, there is a bead at position r_v and a space at position $r_v - k_v$ for every $1 \leq v \leq s$. Let $i + 1 \leq j_1 < \dots < j_t \leq k$ be the residues of the hand nodes of the connected components of Y .


We modify $\text{Ab}(\lambda_Y)$ as follows:

- ◇ move the bead at y_i to the space at y_{j_1} ,
- ◇ for every $1 \leq v \leq t - 1$, move the star bead at $y_{j_{v+1}} - 1$ to the space at $y_{j_{v+1}} - 1$,
- ◇ move the star bead at $y_{j_t} - 1$ to the space at x_k ,
- ◇ for every $1 \leq v \leq s$, move the bead at r_v to the space at $r_v - k_v$.




Let λ_W be the resulting partition. Then $\mu_Y \setminus \lambda_W \in B$. We let W be the p -shape of $\mu_Y \setminus \lambda_W$. Then we can deduce that $d(X, W) < d(X, Y)$ and $d(Y, W) < d(X, Y)$; this is done as for Z above, but now note that k cannot belong to $D(X, W)$, e.g. because we are supposing that x_k is a star bead in $\text{Ab}(\mu_X)$. The p -shape W is finally the desired one, contradicting the hypothesis of Claim 1.

We therefore assume that μ_X has no addable $(k - i)$ -hooks at runner i and, hence, by Proposition 2.2, that μ_Y has a removable $(k - i)$ -hook at runner k . Finally we note that this removable hook does not occur at position y_k since by the previous remark position $y_k + i - k$ is a bead in $\text{Ab}(\mu_Y)$. So we conclude that λ_Y has a removable $(k - i)$ -hook at runner i ending both the inductive step and the proof. \square

Example. Let $m = 42$, $l = 37$ and $p = 7$. The skew-partitions $\mu_X \setminus \lambda_X = (14^2, 7, 5, 2) \setminus (14^2, 4, 3, 2)$ and $\mu_Y \setminus \lambda_Y = (9, 8, 7, 5^2, 4^2) \setminus (9, 8, 4^5)$ have size $n = m - l = 5$ and belong to the same combinatorial block of $\mathcal{C}_{37,42}^k$ having 7-content $\mathcal{C} = \{\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}\}$. Indeed, $\text{core}_7(\mu_X) = \text{core}_7(\mu_Y) = (5, 3, 2^3)$ and $\text{core}_7(\lambda_X) = \text{core}_7(\lambda_Y) = (7, 3, 2^3)$. Let X and Y be the p -shapes of $\mu_X \setminus \lambda_X$ and $\mu_Y \setminus \lambda_Y$, respectively. The Young diagrams and the abacus configurations (with $K = 14$ beads) are shown below.



We see that $D(X, Y) = \{\bar{0}\}$, so that $d(X, Y) = 1$. We use the procedure explained in the proof of Theorem 3.6, faithfully retracing the notation, to construct a 7-shape that has distance 0 from both X and Y . Let $i = \bar{0}$: position $x_0 = y_0 = 14$ is a space in $\text{Ab}(\mu_X)$ and a bead in $\text{Ab}(\mu_Y)$. In this case the inductive procedure stops at $k = 4$ because $y_4 = 18$ is an addable 4-hook at runner 0 of $\text{Ab}(\lambda_Y)$ and $r_1 = 11$ is a removable 4-hook at runner 4. It follows that $k_1 = 3$ since $r_1 - k_1 = 8$ is a space and $y_1 = 15$ is a bead in $\text{Ab}(\mu_Y)$. The procedure ensures that λ_Y has a removable node at runner 1. Actually, this is at position $r_2 = 22$. Here we stop at the first step because $r_2 - 1 \equiv 0 \pmod{7}$. This gives $k_2 = 1$ and $s = 2$ ending the ‘removing’ part. For the ‘adding’ one we have that $t = 1$ and $j_1 = \bar{4}$ is the only hand node of a connected component of X – that is actually a ribbon. We then move the nodes as explained in the proof of Corollary 3.7 and shown in the figure above. We find a partition $\lambda_Z = (8^2, 7, 5, 4^2, 1) \vdash 37$ such that $\mu_Y \setminus \lambda_Z$ has 7-shape Z such that $d(X, Z) = d(Y, Z) = 0$ as can be easily deduced from the diagrams.



We immediately deduce the following corollary, using induction on $d(X, Y)$.

Corollary 3.7. *Suppose $X, Y \in \mathcal{B}$. Then X and Y are p -linked.*

We deduce as a corollary the main result of this section.

Corollary 3.8. *Conjecture 2.5 holds for generalised ribbon blocks.*

Proof. Let B be a generalised ribbon block as above. It follows from Corollary 3.7 and the implication $(3 \Rightarrow 4)$ in Theorem 3.5 that the matrix D_B is connected. As explained in the discussion following Theorem 3.5, this implies that the decomposition matrix of B is connected, so that B is a single p -block of $\mathcal{C}_{l,m}^k$. \square


3.2 Belt blocks

For the rest of this section suppose that $p = n$. Our aim is to establish Conjecture 2.5 for a *belt block* B , i.e. a combinatorial block of $\mathcal{C}_{l,m}^k$ whose central character equals $\mathbb{Z}/p\mathbb{Z} = \{\overline{0}, \dots, \overline{p-1}\}$.

As in Section 3.1, we first replace B (again regarded as a subset of $\mathcal{P}_{l,m}$) with its set \mathcal{B} of p -shapes. We keep going with the same notation established in the preface of the chapter talking about the connected components of a p -shape. Now the maximal p -shapes are ribbons. Unfortunately in this situation we cannot mimic at all the arguments we have used for ribbon blocks. Indeed, even if it is still true that the formal character of a disconnected p -shape is the sum of the formal characters of its refinements, a basic step of the reasoning in Section 3.1, namely Lemma 3.1, is no longer true in this setting. This means that, in general, different ribbons may share composition factors.

Example. Let $p = 5$. The formal characters of the following two ribbons share two summands:


$(1, 2, 0, 3, 4)$ and $(1, 2, 3, 0, 4)$.




To fix this inconvenience, we need to introduce a functional ‘refinement’ of the set of ribbons. To do this, we begin with the introduction of a graph analogue to that in Section 3.1. We define a *(circular) arrow graph* to be a directed graph on the set $\mathbb{Z}/p\mathbb{Z}$ such that every arrow has the form $i \rightarrow i + 1$ or $i + 1 \rightarrow i$ for some i and the graph is not a directed cycle. An arrow graph is a *belt* if for every i there is either an arrow $i \rightarrow i + 1$ or an arrow $i + 1 \rightarrow i$.

We will draw arrow graphs with $\overline{0}, \overline{1}, \dots, \overline{p-1}$ in clockwise order, omitting overlines. So we may say that the arrows of an arrow graph are all oriented clockwise to mean that they all have the form $i \rightarrow i + 1$.

Example. Let $p = 7$. The following are two arrow graphs, the one on the right being a belt.



We observe that every p -shape X in a belt block is uniquely associated to an arrow graph Γ_X through the same procedure explained in Section 3.1. For example, the arrow graph on the left of the above example corresponds to the p -shape below.



Note that in this case the procedure does not give rise to a bijection between all possible p -shapes and all possible arrow graphs. In particular, no p -shape corresponds to a belt. If Γ is an arrow graph, we define its formal character $\text{ch}(\Gamma)$ to be the formal sum of all permutations $\mathbf{b} = (b_0, \dots, b_{p-1})$ of $\mathbb{Z}/p\mathbb{Z}$ with the property that i appears before [resp. after] $i + 1$ for every arrow $i \rightarrow i + 1$ [resp. $i + 1 \rightarrow i$]. Note that if X is a p -shape, then $\text{ch}(\Gamma_X) = \text{ch}(X)$.

Our next task is to build for every belt Γ a \mathcal{H}_p^k -module whose formal character is $\text{ch}(\Gamma)$. We take a vector space M_Γ with basis $\{e_{\mathbf{b}}\}$ labelled by the different summands of $\text{ch}(\Gamma)$. If $\mathbf{b} = (b_0, \dots, b_{p-1})$ is a summand of $\text{ch}(\Gamma)$, then for each $k \in \{1, \dots, p-1\}$ we define $\sigma_k \mathbf{b} = (b_0, \dots, b_{k+1}, b_k, \dots, b_{p-1})$. We define an action of \mathcal{H}_p^k on M_Γ as follows. Fix a summand \mathbf{a} of $\text{ch}(\Gamma)$. We set

$$z_k e_{\mathbf{b}} = b_k e_{\mathbf{b}}$$

and

$$s_k e_{\mathbf{b}} = \begin{cases} \frac{1}{b_{k+1}-b_k} e_{\mathbf{b}} + e_{\sigma_k \mathbf{b}} & \text{if } b_k \text{ appears after } b_{k+1} \text{ in } \mathbf{a}, \\ \frac{1}{b_{k+1}-b_k} e_{\mathbf{b}} + \left(1 - \frac{1}{(b_{k+1}-b_k)^2}\right) e_{\sigma_k \mathbf{b}} & \text{if } b_k \text{ appears before } b_{k+1} \text{ in } \mathbf{a}. \end{cases}$$

Theorem 3.9. *With the above rule M_Γ is a \mathcal{H}_p^k -module with formal character $\text{ch}(\Gamma)$.*

Proof. We just need to prove the first statement; the second part is then obvious from the definition of the formal character of a \mathcal{H}_p^k -module.

We prove that M_Γ is an \mathcal{H}_p^k -module by showing that the above defined rules respect the defining relations of \mathcal{H}_p^k shown in Section 2.6. Relations (1) and (3) are trivially satisfied and need no further investigations. Therefore we only consider the *braid* relations in (2) and the relations between the polynomial and the Coxeter generators in (4).

We start from the braid relations. These divide into three types depending on the integers $m_{i,j}$ such that $(s_i s_j)^{m_{i,j}} = 1$. For each relation several cases arise depending on the positions of the involved entries of \mathbf{b} in the fixed summand \mathbf{a} of $\text{ch}(\Gamma)$. We only analyse one case for each relation, the others being similar.

◇ $s_k^2 = 1$. Suppose that b_k appears before b_{k+1} in \mathbf{a} . We have that

$$\begin{aligned} s_k \mathbf{e}_{\mathbf{b}} &= \frac{1}{b_{k+1} - b_k} \mathbf{e}_{\mathbf{b}} + \left(1 - \frac{1}{(b_{k+1} - b_k)^2}\right) \mathbf{e}_{\sigma_k \mathbf{b}} \Rightarrow \\ s_k^2 \mathbf{e}_{\mathbf{b}} &= \frac{1}{(b_{k+1} - b_k)^2} \mathbf{e}_{\mathbf{b}} + \frac{1}{b_{k+1} - b_k} \left(1 - \frac{1}{(b_{k+1} - b_k)^2}\right) \mathbf{e}_{\sigma_k \mathbf{b}} \\ &\quad - \frac{1}{b_{k+1} - b_k} \left(1 - \frac{1}{(b_{k+1} - b_k)^2}\right) \mathbf{e}_{\sigma_k \mathbf{b}} + \left(1 - \frac{1}{(b_{k+1} - b_k)^2}\right) \mathbf{e}_{\mathbf{b}} \\ &= \mathbf{e}_{\mathbf{b}}. \end{aligned}$$

◇ $s_j s_k = s_k s_j$ with $|k - j| > 1$. Suppose that b_k appears after b_{k+1} and that b_j appears before b_{j+1} in \mathbf{a} . For the left-hand side we have

$$\begin{aligned} s_k \mathbf{e}_{\mathbf{b}} &= \frac{1}{b_{k+1} - b_k} \mathbf{e}_{\mathbf{b}} + \mathbf{e}_{\sigma_k \mathbf{b}} \Rightarrow \\ s_j s_k \mathbf{e}_{\mathbf{b}} &= \frac{1}{(b_{k+1} - b_k)(b_{j+1} - b_j)} \mathbf{e}_{\mathbf{b}} + \frac{1}{b_{k+1} - b_k} \left(1 - \frac{1}{(b_{j+1} - b_j)^2}\right) \mathbf{e}_{\sigma_j \mathbf{b}} \\ &\quad + \frac{1}{b_{j+1} - b_j} \mathbf{e}_{\sigma_k \mathbf{b}} + \left(1 - \frac{1}{(b_{j+1} - b_j)^2}\right) \mathbf{e}_{\sigma_j \sigma_k \mathbf{b}}, \end{aligned}$$

while for the right-hand side we have

$$\begin{aligned} s_j \mathbf{e}_{\mathbf{b}} &= \frac{1}{b_{j+1} - b_j} \mathbf{e}_{\mathbf{b}} + \left(1 - \frac{1}{(b_{j+1} - b_j)^2}\right) \mathbf{e}_{\sigma_j \mathbf{b}} \Rightarrow \\ s_k s_j \mathbf{e}_{\mathbf{b}} &= \frac{1}{(b_{k+1} - b_k)(b_{j+1} - b_j)} \mathbf{e}_{\mathbf{b}} + \frac{1}{b_{j+1} - b_j} \mathbf{e}_{\sigma_k \mathbf{b}} \\ &\quad + \frac{1}{b_{k+1} - b_k} \left(1 - \frac{1}{(b_{j+1} - b_j)^2}\right) \mathbf{e}_{\sigma_j \mathbf{b}} + \left(1 - \frac{1}{(b_{j+1} - b_j)^2}\right) \mathbf{e}_{\sigma_k \sigma_j \mathbf{b}}. \end{aligned}$$

Comparing the coefficients of the different basis elements we easily gain the proof.

◇ $s_k s_{k+1} s_k = s_{k+1} s_k s_{k+1}$. Suppose that b_{k+1} precedes b_{k+2} which precedes b_k in \mathbf{a} . The left-hand

and right-hand sides are as follows

$$\begin{aligned}
 s_k e_{\mathbf{b}} &= \frac{1}{b_{k+1} - b_k} e_{\mathbf{b}} + e_{\sigma_k \mathbf{b}} \Rightarrow \\
 s_{k+1} s_k e_{\mathbf{b}} &= \frac{1}{(b_{k+1} - b_k)(b_{k+2} - b_{k+1})} e_{\mathbf{b}} + \frac{1}{b_{k+1} - b_k} \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\sigma_{k+1} \mathbf{b}} \\
 &\quad + \frac{1}{b_{k+2} - b_k} e_{\sigma_k \mathbf{b}} + e_{\sigma_{k+1} \sigma_k \mathbf{b}} \Rightarrow \\
 s_k s_{k+1} s_k e_{\mathbf{b}} &= \frac{1}{(b_{k+1} - b_k)^2 (b_{k+2} - b_{k+1})} e_{\mathbf{b}} + \frac{1}{(b_{k+1} - b_k)(b_{k+2} - b_{k+1})} e_{\sigma_k \mathbf{b}} \\
 &\quad + \frac{1}{(b_{k+1} - b_k)(b_{k+2} - b_k)} \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\sigma_{k+1} \mathbf{b}} \\
 &\quad + \frac{1}{b_{k+1} - b_k} \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\sigma_k \sigma_{k+1} \mathbf{b}} \\
 &\quad - \frac{1}{(b_{k+2} - b_k)(b_{k+1} - b_k)} e_{\sigma_k \mathbf{b}} + \frac{1}{b_{k+2} - b_k} \left(1 - \frac{1}{(b_{k+1} - b_k)^2} \right) e_{\mathbf{b}} \\
 &\quad + \frac{1}{b_{k+2} - b_{k+1}} e_{\sigma_{k+1} \sigma_k \mathbf{b}} + \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\sigma_k \sigma_{k+1} \sigma_k \mathbf{b}}
 \end{aligned}$$

and

$$\begin{aligned}
 s_{k+1} e_{\mathbf{b}} &= \frac{1}{b_{k+2} - b_{k+1}} e_{\mathbf{b}} + \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\sigma_{k+1} \mathbf{b}} \Rightarrow \\
 s_k s_{k+1} e_{\mathbf{b}} &= \frac{1}{(b_{k+2} - b_{k+1})(b_{k+1} - b_k)} e_{\mathbf{b}} + \frac{1}{b_{k+2} - b_{k+1}} e_{\sigma_k \mathbf{b}} \\
 &\quad + \frac{1}{b_{k+2} - b_k} \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\sigma_{k+1} \mathbf{b}} + \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\sigma_k \sigma_{k+1} \mathbf{b}} \Rightarrow \\
 s_{k+1} s_k s_{k+1} e_{\mathbf{b}} &= \frac{1}{(b_{k+2} - b_{k+1})^2 (b_{k+1} - b_k)} e_{\mathbf{b}} \\
 &\quad + \frac{1}{(b_{k+2} - b_{k+1})(b_{k+1} - b_k)} \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\sigma_{k+1} \mathbf{b}} \\
 &\quad + \frac{1}{(b_{k+2} - b_k)(b_{k+2} - b_{k+1})} e_{\sigma_k \mathbf{b}} + \frac{1}{b_{k+2} - b_{k+1}} e_{\sigma_{k+1} \sigma_k \mathbf{b}} \\
 &\quad - \frac{1}{(b_{k+2} - b_k)(b_{k+2} - b_{k+1})} \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\sigma_{k+1} \mathbf{b}} \\
 &\quad + \frac{1}{b_{k+2} - b_k} \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\mathbf{b}} \\
 &\quad + \frac{1}{b_{k+1} - b_k} \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\sigma_k \sigma_{k+1} \mathbf{b}} + \left(1 - \frac{1}{(b_{k+2} - b_{k+1})^2} \right) e_{\sigma_{k+1} \sigma_k \sigma_{k+1} \mathbf{b}}.
 \end{aligned}$$

The coefficients of $e_{\sigma_k \sigma_{k+1} \mathbf{b}}$, $e_{\sigma_{k+1} \sigma_k \mathbf{b}}$ and $e_{\sigma_k \sigma_{k+1} \sigma_k \mathbf{b}} = e_{\sigma_{k+1} \sigma_k \sigma_{k+1} \mathbf{b}}$ are the same on both sides. The coefficients of $e_{\mathbf{b}}$, $e_{\sigma_k \mathbf{b}}$ and $e_{\sigma_{k+1} \mathbf{b}}$ turn out to be equal from easy calculations comparing the two above formulas.

We finally focus on the relation (4): $s_k z_k = z_{k+1} s_k - 1$. Suppose that b_k precedes b_{k+1} in \mathbf{a} . We evaluate the action of both sides of (4) on $e_{\mathbf{b}}$. On the left-hand we have

$$\begin{aligned}
 z_k e_{\mathbf{b}} &= b_k e_{\mathbf{b}} \Rightarrow \\
 s_k z_k e_{\mathbf{b}} &= \frac{b_k}{b_{k+1} - b_k} e_{\mathbf{b}} + b_k \left(1 - \frac{1}{(b_{k+1} - b_k)^2} \right) e_{\sigma_k \mathbf{b}},
 \end{aligned}$$

while on the right we have

$$s_k e_{\mathbf{b}} = \frac{1}{b_{k+1} - b_k} e_{\mathbf{b}} + \left(1 - \frac{1}{(b_{k+1} - b_k)^2}\right) e_{\sigma_k \mathbf{b}} \Rightarrow$$


$$(z_{k+1} s_k - 1) e_{\mathbf{b}} = \frac{b_{k+1}}{b_{k+1} - b_k} e_{\mathbf{b}} + b_k \left(1 - \frac{1}{(b_{k+1} - b_k)^2}\right) e_{\sigma_k \mathbf{b}} - e_{\mathbf{b}}.$$

It is now straightforward to check that the coefficients of $e_{\mathbf{b}}$ and $e_{\sigma_k \mathbf{b}}$ are equal on both sides. This ends the proof. \square

Theorem 3.9 strengthens the above discussion and shows that every arrow graph corresponds to a \mathcal{H}_p^k -module. In particular, we can talk about the composition factors of a belt, meaning the composition factors of the corresponding \mathcal{H}_p^k -module. Later on we may look at the set of p -shapes \mathcal{B} as a subset of the set of all possible arrow graphs.

As in Section 3.1 we rely on the notion of distance between p -shapes. Given p -shapes X and Y , we define $D(X, Y)$ and $d(X, Y)$ as in Section 3.1.

Example. Let $p = 7$, and let X and Y be the p -shapes below. From the corresponding arrow graphs, we see that $D(X, Y) = \{\bar{0}, \bar{4}\}$, and therefore $d(X, Y) = 2$.



We can now state a variant of Lemma 3.1.

Lemma 3.10. Consider $\mathbf{a} = (a_0, \dots, a_{p-1})$ where a_0, \dots, a_{p-1} equal $\bar{0}, \dots, \overline{p-1}$ in some order. Then \mathbf{a} appears in the formal character of exactly one belt.

Proof. This result follows similarly to Lemma 3.1. Take an empty arrow graph and fill its edges according to the relative position between the entries of \mathbf{a} until a belt X is reached. By construction \mathbf{a} is a term of the formal character $\text{ch}(X)$. If a different belt Y is considered, then $D(X, Y) \neq \emptyset$ and repeating the last argument in the proof of Lemma 3.1, we see that $\text{ch}(X)$ and $\text{ch}(Y)$ have no terms in common. \square

Corollary 3.11. Different belts have no composition factors in common.

At this point we can mimic the arguments in Section 3.1 using belts in place of maximal p -shapes.

Definition. Let X be a p -shape. A belt Y is a refinement of X if Y can be obtained by adding edges to X .

Corollary 3.12. Let X be a p -shape. The formal character of X is the sum of the formal characters of the refinements of X .

Proof. Given Y a refinement of X , every term of $\text{ch}(Y)$ is also a term of $\text{ch}(X)$ (just because X is obtained forgetting some arrows of Y). Conversely, each term of $\text{ch}(X)$ is a term of the formal character of a unique refinement of X : for every j and $j + 1 \pmod{p}$ that are not linked by an arrow in X , the relative order of these entries in the chosen term of $\text{ch}(X)$ indicate how they need to be linked to find the desired belt. \square

As in Corollary 3.4, [K, Theorem 5.3.1] ensures the following.

Corollary 3.13. *If X is a p -shape, then the composition factors of X are the composition factors of its refinements.*


We are now ready to establish a result which gives us a way to recognise when two Specht $\mathcal{C}_{l,m}^k$ -modules share a composition factor.

Theorem 3.14. *Let X and Y be p -shapes. The following are equivalent:*

1. X and Y have a composition factor in common;
2. The formal characters of X and Y have a term in common;
3. $d(X, Y) = 0$ and $\Gamma_X \cup \Gamma_Y$ is not a directed cycle;
4. There is a belt which is a refinement of both X and Y .

Remark. Before proving Theorem 3.14 we dwell a moment on the meaning of condition (3). Since $d(X, Y) = 0$, the overlapping $\Gamma_X \cup \Gamma_Y$ is an admissible arrow graph. We have that $\Gamma_X \cup \Gamma_Y$ is a directed cycle if and only if all arrows in Γ_X and Γ_Y are anticlockwise (or all clockwise) and there is no $k \in \mathbb{Z}/p\mathbb{Z}$ such that \boxed{k} is simultaneously the foot node of a connected component of X and a connected component of Y .

Example. Let $p = 7$, and let X and Y be the p -shapes given below. All arrows in Γ_X and Γ_Y are oriented clockwise (because neither X nor Y has a node immediately above another). Moreover, X and Y do not have foot nodes of the same residue. So $\Gamma_X \cup \Gamma_Y$ is a directed cycle. As a consequence, the formal characters of X and Y have no term in common, so X and Y do not share a composition factor.



Proof of Theorem 3.14.

(1 \Rightarrow 2) This is trivial because the formal character of a \mathcal{H}_p^k -module is the sum of the formal characters of its composition factors.

(2 \Rightarrow 3) Suppose that $\text{ch}(X)$ and $\text{ch}(Y)$ have a term in common. Then it immediately follows that $d(X, Y) = 0$. Suppose for a contradiction that $\Gamma_X \cup \Gamma_Y$ is a clockwise directed cycle and set

$$F_X := \{k \in \mathcal{C}(B) \mid \boxed{k} \text{ is the foot node of a connected component of } X\}$$

and

$$F_Y := \{k \in \mathcal{C}(B) \mid \boxed{k} \text{ is the foot node of a connected component of } Y\}.$$

(For example, for the 7-shapes in the example above, $F_X = \{\bar{0}, \bar{4}\}$ and $F_Y = \{\bar{5}\}$.) The remark above shows that $F_X \cap F_Y = \emptyset$. Now since every term of $\text{ch}(X)$ has an element of F_X as first component and every term of $\text{ch}(Y)$ has an element of F_Y as first component, it follows that $\text{ch}(X)$ and $\text{ch}(Y)$ have no terms in common and this contradicts the hypothesis. (If the arrows in $\Gamma_X \cup \Gamma_Y$ are all anticlockwise, the proof follows similarly replacing "foot" with "hand" in the definitions of F_X and F_Y above.)

(3 \Rightarrow 4) By the hypothesis the union $\Gamma_X \cup \Gamma_Y$ is an arrow graph. This can be refined to a belt by adding in arrows if necessary (ensuring that these arrows are not all oriented the same way as all the existing arrows). This belt is a refinement of both X and Y .

(4 \Rightarrow 1) This follows from Corollary 3.13. \square

It seems natural that the set of belts gives a family of simple \mathcal{H}_p^k -modules. Despite this, we will not focus on this question in the present paper. We write our suspect as a conjecture such that it can inspire further research.

Conjecture 3.15. *Let Γ be a belt. Then the \mathcal{H}_p^k -module M_Γ is simple.*

Now we construct an approximation to the decomposition matrix as in Section 3.1: we construct the matrix

$$D_B = (d_{X,Y})_{X \in \mathcal{B}, Y \in \text{Belt}}$$

where $d_{X,Y} = 1$ if Y is a refinement of X , and 0 otherwise. As in Section 3.1, showing that D_B is a connected matrix is sufficient to show that B is a block.

Definition. Let X, Y be p -shapes in \mathcal{B} . We say that X and Y are p -linked if there exists a series of p -shapes $X = X_1, X_2, \dots, X_{t-1}, X_t = Y$ in \mathcal{B} such that X_j and X_{j+1} satisfy *any* of the equivalent conditions in Theorem 3.14, for every $1 \leq j \leq t - 1$.

We now move to the heart of this section. First of all we show that we can reduce to the case where two p -shapes in a belt block have distance zero.


Proposition 3.16. *Let X and Y be p -shapes in \mathcal{B} with $d(X, Y) > 0$. There exists a p -shape $Z \in \mathcal{B}$ such that $d(X, Z) < d(X, Y)$ and $d(Y, Z) < d(X, Y)$.*

Proof. Let $\mu_X \setminus \lambda_X$ and $\mu_Y \setminus \lambda_Y$ be skew-partitions in B having p -shapes X and Y , respectively. We want to show that there is a p -shape $Z \in \mathcal{B}$ that satisfies the conditions in the statement. The proof of this result is very similar to that of Theorem 3.6. Let $i \in D(X, Y)$. Here we can state and prove an intermediate claim, i.e. a perfect analogue of Claim 1 in Theorem 3.6, running for $k \in \{i + 1, \dots, i + p + 1\}$ with $k \notin D(X, Y)$. If for any of these steps, a modifications of μ_X or λ_Y is allowed, then we are done performing the appropriate modification exploited in Theorem 3.6. Alternatively, if for all k no modifications have been allowed, we first find integers k_1, \dots, k_s with $\sum_{1 \leq u \leq s} k_u = p$, and positions $c_1, \dots, c_s, d_1, \dots, d_s$ of the abacus display such that c_1, \dots, c_s are beads and d_1, \dots, d_s are spaces in $\text{Ab}(\mu_X)$. The proof then follows observing that μ_X has a addable p -hook at runner i (realised by the biggest bead position at runner i of $\text{Ab}(\mu_X)$): the desired partition $\mu_Z \vdash m$ is obtained from μ_X deleting *all* the nodes in $\mu_X \setminus \lambda_X$ and adding them in different positions (moving the bead at c_v to the space at d_v for all v). The choice of k_1, \dots, k_s implies that $\mu_Z \setminus \lambda_X$ has the desired p -shape Z . \square


Example. Let $m = 119, l = 112$ and $p = n = 7$. Consider the skew-partitions

$$\begin{aligned} \mu_X \setminus \lambda_X &= (20, 18, 17, 16, 12^4) \setminus (20, 18, 17, 16, 12^2, 11, 6), \\ \mu_Y \setminus \lambda_Y &= (21^2, 17, 15, 14, 13, 12, 6) \setminus (20, 16^2, 15, 14, 13, 12, 6) \end{aligned}$$

in $\mathcal{P}_{112,119}$ having 7-shapes X and Y , respectively. These belong to the same combinatorial block of $C_{112,119}^k$ as can be easily checked (all the partitions involved have 7-core $(13,7,3,2^2,1) \vdash 28$). Below we show the abacus configurations with charge 21 and the arrow graphs of X and Y .



We see that $D(X, Y) = \{\bar{0}, \bar{4}\}$ and hence $d(X, Y) = 2$. We follow the notation of Theorem 3.6. Let $i = \bar{0}$: position $x_0 = 21$ is a space in $\text{Ab}(\mu_X)$ and position $y_0 = 35$ is a bead in $\text{Ab}(\mu_Y)$. Here, for every $1 \leq k \leq 6$, no modifications of μ_X or λ_Y are allowed by the procedure. Consider the addable 7-hook at runner 0 of $\text{Ab}(\mu_X)$ at position $a_1 = 35$. This gives $d_1 = 42$ from which follows that $k_1 = 5$ and $c_1 = 37$. The partition μ_X has an addable 2-hook at runner 0, this is at position $a_2 = 28$. Hence, $d_2 = 30$ and consequently $k_2 = 2$ and $c_2 = a_2 = 28$. Then $s = 2$ and we stop. For the 'removing' part, since X is a ribbon, $t = 1$ and $j_1 = \bar{5}$. We modify $\text{Ab}(\mu_X)$ according to the rule given in Theorem 3.6 (as shown above), obtaining the partition $\mu_Z = (22, 21, 17, 16, 14, 12, 11, 6) \vdash 119$. Denoting by Z the 7-shape of $\mu_Z \setminus \lambda_X$, we see that $d(X, Z) = 1$ and $d(Y, Z) = 0$.



Now we use Proposition 3.16 to show that the combinatorial block B is actually a p -block of $\mathcal{C}_{l,m}^k$. This is slightly more complicated than in Section 3.1, because it is no longer the case that two p -shapes at distance 0 are p -linked. Take two p -shapes $X, Y \in \mathcal{B}$ such that $d(X, Y) = 0$. By Theorem 3.14, X and Y are p -linked if $\Gamma_X \cup \Gamma_Y$ is not a directed cycle. Hence from now on we suppose that $\Gamma_X \cup \Gamma_Y$ is a directed cycle and we assume without loss of generality that all its arrows are oriented clockwise. The aim of the rest of this section is to prove that X and Y are p -linked.

First of all we establish some notation. Every skew-partition $\mu \setminus \lambda$ in B satisfies the hypothesis of Proposition 2.4 and hence $\text{core}_p(\mu) = \text{core}_p(\lambda)$. So there is a p -core γ and an integer w such that

$$\text{core}_p(\mu) = \text{core}_p(\lambda) = \gamma, \quad \text{weight}_p(\lambda) = w, \quad \text{weight}_p(\mu) = w + 1$$

for every skew-partition $\mu \setminus \lambda$ in B . Let $c \in \mathbb{Z}/p\mathbb{Z}$ be the p -residue of the right-most node in the first part of γ , i.e. $c = \overline{\gamma_1 - 1}$. If γ is the empty partition, we set $c := \overline{p - 1}$.

Let L be the p -block of S_l whose partitions all have p -core γ and p -weight w . For each $i \in \mathbb{Z}/p\mathbb{Z}$, let

$$L_i = \{\lambda \in L \mid \overline{\lambda_1 - 1} = i\}.$$

We remark that some L_i can be empty and that

$$L = \bigsqcup_{i \in \mathbb{Z}/p\mathbb{Z}} L_i.$$

For every $\lambda \in L$, let O_λ be the set of partitions $\mu \vdash m$ such that $\mu \setminus \lambda \in B$ and all the arrows of the p -shape of $\mu \setminus \lambda$ are clockwise (i.e. $\mu \setminus \lambda$ does not contain two nodes in the same column).

Remark. Suppose $\lambda_X, \lambda_Y \in L$, $\mu_X \in O_{\lambda_X}$ and $\mu_Y \in O_{\lambda_Y}$, and let X and Y be the p -shapes of $\mu_X \setminus \lambda_X$ and $\mu_Y \setminus \lambda_Y$ respectively. Then $d(X, Y) = 0$, because all edges are oriented clockwise. Moreover, every pair X, Y such that $\Gamma_X \cup \Gamma_Y$ is a clockwise directed cycle arises in this way.


For every $i \in \mathbb{Z}/p\mathbb{Z}$, let X_i be the (ribbon) p -shape

$$\boxed{i+1 \mid \cdots \cdots \mid i-1 \mid i}.$$

For example, if $p = 7$ and $i = 4$, X_4 is the p -shape Y in the example in Theorem 3.14.

Let $\lambda \in L_i$ and let $\mu_\lambda = (\lambda_1 + p, \lambda_2, \dots, \lambda_{l(\lambda)})$. Then the abacus configuration for μ_λ is obtained from the abacus configuration for λ by moving the last bead down one position. It is easy to see that $\mu_\lambda \in O_\lambda$ and $\mu_\lambda \setminus \lambda$ has p -shape X_i .

Example. Let $m = 37$, $l = 30$ and $p = n = 7$. Consider the partition $\lambda = (12, 8, 5, 4, 1) \vdash 30$. Then $\mu_\lambda = (19, 8, 5, 4, 1)$ and $\mu_\lambda \setminus \lambda$ has 7-shape X_4 .




Now take $\mu \in O_\lambda$ with $\mu \neq \mu_\lambda$, and call X the p -shape of $\mu \setminus \lambda$.

Proposition 3.17. X and X_i are p -linked.


Proof. If a connected component of X has hand node \boxed{i} then we are done because $\text{ch}(X_i)$ and $\text{ch}(X)$ share the term $(i + 1, \dots, i)$. Suppose that this is not the case, and let $j \in \mathbb{Z}/p\mathbb{Z}$ be the p -residue of the hand node of the connected component of X containing i . We firstly observe that $\mu_1 = \lambda_1$ because otherwise the nodes $(1, \lambda_1 + 1), \dots, (1, \mu_1)$ of μ would form a connected component of X with $\boxed{i+1}$ as foot node and this would imply that X has a connected component whose hand node is \boxed{i} .

Now we can modify μ by moving the nodes of residue $i + 1, i + 2, \dots, j$ in $\mu \setminus \lambda$ to the end of the first row. Let $\tilde{\mu}$ be the resulting partition. Then $\tilde{\mu} \setminus \lambda$ is in B and its p -shape \tilde{X} is p -linked with both X_i and X : $\text{ch}(X)$ and $\text{ch}(\tilde{X})$ share the term $(i + 1, i + 2, \dots, i)$ while $\text{ch}(X_i)$ and $\text{ch}(\tilde{X})$ share $(j + 1, j + 2, \dots, j)$. So X and X_i are p -linked. □

Example. Let m, l, p, n and λ be as in the previous example. Let $\mu = (12^2, 5, 4^2) \in O_\lambda$. The skew-partition $\mu \setminus \lambda$ belongs to the combinatorial block B of $C_{30,37}^k$ characterised by the 7-core $\gamma := \text{core}_7(B) = (1^2) \vdash 2$. Call X the 7-shape of $\mu \setminus \lambda$.



Retracing the notation of Proposition 3.17, we have that $i = \bar{4}$ (because $\lambda_1 - 1 = 11 \equiv 4 \pmod{7}$) and then $\lambda \in L_4$ and $j = \bar{6}$ because the node $\boxed{4}$ belongs to the connected component of X having hand node $\boxed{6}$. So we find $\tilde{\mu}$ by moving the nodes $\boxed{5} \boxed{6}$ to the end of the first row, giving $\tilde{\mu} = (14, 12, 5, 4, 2) \vdash 37$.



Now the 7-shape \tilde{X} of $\tilde{\mu} \setminus \lambda$, is 7-linked with both X and X_4 .

Remark. If $\lambda_1, \lambda_2 \in L_i$, then $\mu_{\lambda_1} \setminus \lambda_1$ and $\mu_{\lambda_2} \setminus \lambda_2$ have the same p -shape X_i . By the previous proposition we deduce that, for every $\mu_1 \in O_{\lambda_1}$ and $\mu_2 \in O_{\lambda_2}$, the p -shapes of $\mu_1 \setminus \lambda_1$ and $\mu_2 \setminus \lambda_2$ are p -linked.

We are now very close to the end of this section. The next result will link p -shapes X_i, X_j for all pairs of $i, j \in \mathbb{Z}/p\mathbb{Z}$. Recall that $c = \overline{\gamma_1 - 1}$, and observe that L_c is non-empty: for example, it contains the partition $\overline{\gamma} = (\gamma_1 + wp, \gamma_2, \gamma_3, \dots)$. In particular this shows that B contains at least one skew-partition with p -shape X_c .

Proposition 3.18. *Let $i \in \mathbb{Z}/p\mathbb{Z}$ such that $L_i \neq \emptyset$. Then X_i and X_c are p -linked.*

Proof. Let $\lambda \in L_i$ and consider the skew-partitions $\mu_\lambda \setminus \lambda$ and $\mu_{\overline{\gamma}} \setminus \overline{\gamma}$ in B having p -shapes X_i and X_c , respectively. For convenience we suppose that $c = \overline{0}$, though the proof can easily be modified for other values of c .

If $i = \overline{0}$, the result follows by the previous remark.

Suppose that $i \neq \overline{0}$. First of all we observe that in the abacus configuration of γ , the leftmost runner, i.e. runner 0, has more beads than any other runner: this is because the last bead on the abacus occurs on this runner and (because γ is a p -core) the beads are at the topmost positions of their runners. So, following the notation of Section 2.5, we have that for every $j \in \mathbb{Z}/p\mathbb{Z} \setminus \{\overline{0}\}$,

$$b_0(\gamma) > b_j(\gamma). \quad (1)$$

The abacus configuration of λ is obtained by lowering down beads in $\text{Ab}(\gamma)$ a total amount of w times. Now recall the p -quotient $(\lambda^{(0)}, \dots, \lambda^{(p-1)})$ of λ . The fact that $\lambda \in L_i$ implies that

$$\lambda^{(i)}_1 \geq \lambda^{(0)}_1 + b_0(\lambda) - b_i(\lambda), \quad (2)$$

where $\lambda^{(i)}_1$ and $\lambda^{(0)}_1$ denote the number of spaces above the lowest beads in the i -th and 0-th runners of $\text{Ab}(\lambda)$, respectively. Recalling that $b_j(\lambda) = b_j(\gamma)$ for all $j \in \mathbb{Z}/p\mathbb{Z}$, we combine (1) and (2) getting

$$\lambda^{(i)}_1 > \lambda^{(0)}_1. \quad (3)$$

By construction, if $(\mu_\lambda^{(0)}, \dots, \mu_\lambda^{(p-1)})$ denotes the p -quotient of μ_λ , then $(\mu_\lambda^{(j)})_1 = \lambda^{(j)}_1$ for every $j \in \mathbb{Z}/p\mathbb{Z} \setminus \{i\}$ and

$$(\mu_\lambda^{(i)})_1 = \lambda^{(i)}_1 + 1. \quad (4)$$

Comparing (3) and (4) we find that


$$(\mu_\lambda^{(i)})_1 > (\mu_\lambda^{(0)})_1 + 1.$$

This inequality implies that μ_λ has at least two addable i -hooks at runner 0. We consider one of them, say at position $a_1 \equiv 0 \pmod{p}$, avoiding the one that affects the space immediately above the lowest bead in the i -runner of $\text{Ab}(\mu_\lambda)$, if necessary. Now let $1 \leq k_1 \leq i$ be the least integer such that $a_1 + i - k_1$ is a bead. If $k_1 < i$, by (1) and the fact that $b_j(\mu_\lambda) = b_j(\gamma)$ for all $j \in \mathbb{Z}/p\mathbb{Z}$, μ_λ has an addable $(i - k_1)$ -hook at runner 0, say at position a_2 . Then we start again the process from position a_2 : let k_2 be minimal such that position $a_2 + i - k_1 - k_2$ is a bead. It is clear that this procedure terminates at a step s such that $i = \sum_{1 \leq u \leq s} k_u$. Then for $1 \leq v \leq s$ we have positions $c_v := a_v + i - \sum_{1 \leq u \leq v} k_u$ and $d_v := a_v + i - \sum_{1 \leq u \leq v-1} k_u$ such that c_v is a bead and $c_v + 1, \dots, d_v$ are all spaces in $\text{Ab}(\mu_\lambda)$. Then we modify $\text{Ab}(\mu_\lambda)$ as follows (remind that C denotes the charge of the abacus displays):

- ◇ move the star bead at $\lambda_1 + C + p - 1$ to the space at $\lambda_1 + C + p - i - 1$,
- ◇ for every $1 \leq v \leq s$, move the bead at c_v to the space d_v .

The resulting partition $\hat{\mu} \vdash m$ is such that $\hat{\mu} \setminus \lambda \in B$ and has p -shape \hat{X} that is p -linked with both X_i and X_c : the formal characters $\text{ch}(X_i)$ and $\text{ch}(\hat{X})$ share the term $(i + 1, \dots, i)$ while $\text{ch}(X_c)$ and $\text{ch}(\hat{X})$ share the term $(\overline{1}, \overline{2}, \dots, \overline{p-1}, \overline{0})$. This ends the proof. \square


Example. Let B be the combinatorial block of $C_{30,37}^k$ from the previous examples. As already pointed out, following the notation, $\gamma = (1^2)$ and then $c = \gamma_1 - 1 \equiv 0 \pmod{7}$. We see that $\bar{\gamma} = (29,1)$ and $\mu_{\bar{\gamma}} = (36,1)$.



$\mu_{\bar{\gamma}} \setminus \bar{\gamma}$



We show how to modify the skew-partition μ_λ from the previous example, to find a 7-shape that is 7-linked with both X_4 and X_0 .

We first draw the abacus configuration of γ (all abacus configurations in the example have charge $C = 14$).



γ

Note that as remarked in the proof of Proposition 3.18, the number of beads in runner 0 of γ , $b_0(\gamma) = 3$, is bigger than the number of beads in any other runner of $\text{Ab}(\gamma)$. Our proof points out that μ_λ has at least an addable 4-hook at runner 0 (in fact it has two). Take $a_1 = 14$ and $d_1 = 18$. We have that $k_1 = 2$ and so $c_1 = 16$. Now we consider the addable 2-hook of μ_λ at position $a_2 = 7$. This gives $d_2 = 9$ and consequently $k_2 = 1$ and $c_2 = 8$. Finally we take the addable node at runner 0 of μ_λ at position $a_3 = 14$. We have that $d_3 = 15$, giving $k_3 = 1$ and $c_3 = 14$ (then $s = 3$). We modify $\text{Ab}(\mu_\lambda)$ following the rule given in the proof of Proposition 3.18 (shown in figure). The resulting partition $\hat{\mu} = (15, 8, 7, 5, 1^2) \vdash 37$ is such that $\hat{\mu} \setminus \lambda$ has the desired 7-shape \hat{X} .



$\hat{\mu} \setminus \lambda$

We have finally gained the key result of this section.

Corollary 3.19. Suppose $X, Y \in \mathcal{B}$. Then X and Y are p -linked.

Proof. By Proposition 3.16, we can reduce to the case when $d(X, Y) = 0$. Moreover, Theorem 3.14 tells us that we are done if $\Gamma_X \cup \Gamma_Y$ is not an oriented cycle. In the opposite case, the description of the remaining cases together with Propositions 3.17 and 3.18 ends the proof. \square

Now, Corollary 3.19 and Theorem 3.14 implies that the decomposition matrix D_B of the belt block B is connected. As remarked along the treatment, this is enough to state our main result.

Corollary 3.20. *Conjecture 2.5 holds for belt blocks.*

References

- [CR] C. W. Curtis & I. Reiner, *Methods of representation theory Vol 1*, John Wiley & Sons, New York, 1981. [cited on p. 2]
- [E1] H. Ellers, ‘Cliques of irreducible representations of p-solvable groups and a theorem analogous to Alperin’s conjecture’, *Math. Z.* **217** (4) (1994), 607–634. [1]
- [E2] H. Ellers, ‘Cliques of irreducible representations, quotient groups, and Brauer’s theorems on blocks’, *Canad. J. Math.* **47** (5) (1995), 929–945. [1]
- [E3] H. Ellers, ‘The defect groups of a clique, p-solvable groups, and Alperin’s conjecture’, *J. Reine Angew. Math.* **468** (1995), 1–48. [1]
- [E4] H. Ellers, ‘Searching for more general weight conjectures, using the symmetric group as an example’, *J. Algebra* **225** (2000), 602–629. [1]
- [EM1] H. Ellers & J. Murray, ‘Block theory, branching rules, and centraliser algebras’, *J. Algebra* **276** (2004), 236–258. [8]
- [EM2] H. Ellers & J. Murray, ‘Blocks of centraliser algebras and degenerate affine Hecke algebras’, *Commun. Algebra* **42** (2014), 1074–1094. [2, 7, 8, 9, 10, 11]
- [J] G. James, *The representation theory of symmetric groups*, Lecture Notes in Mathematics **682**, Springer-Verlag, Berlin-Heidelberg, 1978. [2]
- [JK] G. James & A. Kerber, *The representation theory of the symmetric group*, Encyclopædia of Mathematics and its Applications **16**, Addison–Wesley, 1981. [5]
- [K] A. Kleshchev, *Linear and projective representations of symmetric groups*, Cambridge Tracts in Mathematics **163**, Cambridge University Press, 2005. [2, 8, 14, 25]
- [L] D. Littlewood, ‘Modular representations of the symmetric group’, *Proc. Roy. Soc. London (A)* **209** (1951), 333–352. [7]
- [S] B. E. Sagan, *The symmetric group: Representations, combinatorial algorithms, and symmetric functions (2nd ed.)*, Graduate Texts in Mathematics **203**, Springer, 2001. [5]