THE GALOIS ACTION ON CONSISTENT MAPS

CHARLES L. SAMUELS

ABSTRACT. A 2009 article of Allcock and Vaaler explored of the \mathbb{Q} -vector space $\mathcal{G} := \overline{\mathbb{Q}}^{\times} / \overline{\mathbb{Q}}_{\text{tors}}^{\times}$, showing how to represent it as part of a function space on the places of $\overline{\mathbb{Q}}$. Several years later, the author began attempts to examine dual spaces related to \mathcal{G} in an effort to obtain Riesz-type representation theorems. Those results required the construction of an object called a *consistent map*. We study a natural Galois action on consistent maps and establish when consistent maps are invariant under this action. Our results generalize earlier work of the author regarding rational valued consistent maps over non-Archimedean places of \mathbb{Q} .

1. INTRODUCTION

Let K be a number field and L/K a finite extension. Each place w of L restricts to a unique place v of K, and in this case, we say that w divides v. Writing L_w and K_v to denote the respective completions, we note the important identity

(1.1)
$$[L:K] = \sum_{w|v} [L_w:K_v]$$

for each place v of K (see [5, Sec. 4, Eq. (2)] or [10, Cor. 8.4], for example). If v divides the place p of \mathbb{Q} , then we write $\|\cdot\|_v$ to denote the unique extension of the p-adic absolute value on \mathbb{Q}_p , and define

(1.2)
$$\|\alpha\|_{\mathcal{G}} = \sum_{v \in M_K} \frac{[K_v : \mathbb{Q}_v]}{[K : \mathbb{Q}]} \cdot \left|\log\|\alpha\|_v\right|,$$

where M_K denotes the set of all places of K. The right hand side of (1.2) does not depend on K, and moreover, its value is unchanged when α is multiplied by a root of unity. Now fixing an algebraic closure $\overline{\mathbb{Q}}$ of \mathbb{Q} and setting $\mathcal{G} = \overline{\mathbb{Q}}^{\times}/\overline{\mathbb{Q}}_{\text{tors}}^{\times}$, the properties of absolute values ensure that $\|\cdot\|_{\mathcal{G}}$ defines a norm on \mathcal{G} with respect to the usual absolute value on \mathbb{Q} . The value $\frac{1}{2} \|\alpha\|_{\mathcal{G}}$, often called the *Weil height of* α , has an extensive history dating back to a famous 1933 problem of D.H. Lehmer [8].

As part of a 2009 article [1], Allcock and Vaaler provided a deep exploration of the normed space \mathcal{G} . Let Y denote the set of all places of $\overline{\mathbb{Q}}$, and for each number field K and each place v of K, we let $Y(K, v) = \{y \in Y : y \mid v\}$. The collection of sets of the form Y(K, v) forms a basis for a totally disconnected, Hausdorff topology on Y, and moreover, there exists a regular measure λ on the Borel sets \mathcal{B} of Y with the property that

$$\lambda(Y(K,v)) = \frac{[K_v : \mathbb{Q}_v]}{[K : \mathbb{Q}]}$$

for each number field K and each place v of K. Among other results, Allcock and Vaaler [1, Theorem 1.1] identified an isometric isomorphism from the completion \mathcal{G} to a subspace of $L^1(Y, \mathcal{B}, \lambda)$. Their methodology opened a pathway for other authors to study the Weil height using techniques of functional analysis (see [2–4, 6, 7, 13], for example).

The author [11] later began the study of various dual spaces related to \mathcal{G} . Specifically, let $\mathcal{V} = \overline{\mathbb{Q}}^{\times}/\overline{\mathbb{Z}}^{\times}$, where $\overline{\mathbb{Z}}$ is the ring of algebraic integers and $\overline{\mathbb{Z}}^{\times}$ is its group of units. The main results of [11] established

²⁰²⁰ Mathematics Subject Classification. 11R32, 11G50, 46E15, 46E27.

Key words and phrases. Linear Functionals, Galois Action, Places, Consistent Maps.

This work was funded in part by the AMS-Simons Research Enhancement Grant for PUI Faculty.

a representation theorem for the algebraic dual of \mathcal{V} roughly in the following way. Defining the set $\mathcal{I} = \{(K, v) : [K : \mathbb{Q}] < \infty, v \in M_K^0\}$, a map $c : \mathcal{I} \to \mathbb{Q}$ is called *consistent* if we have

(1.3)
$$c(K,v) = \sum_{w|v} c(L,w)$$

for all $(K, v) \in \mathcal{I}$ and all finite extensions L/K. If $c, d: \mathcal{I} \to \mathbb{Q}$ are consistent maps and $r \in \mathbb{Q}$ then we let

(1.4)
$$(c+d)(K,v) = c(K,v) + d(K,v)$$
 and $(rc)(K,v) = rc(K,v)$.

These operations cause the set of consistent maps $c : \mathcal{I} \to \mathbb{Q}$ to be a vector space over \mathbb{Q} , which we shall denote by \mathcal{I}^* . If K is a number field and $\alpha \in K$, then we write

(1.5)
$$\Phi_c(\alpha) = \sum_{v \in M_K^0} \frac{c(K,v)}{\log p_v} \log \|\alpha\|_v.$$

The right hand side of (1.5) is unchanged whenever K is replaced by a different number field containing α or when α is multiplied by a unit. Therefore, we have constructed a well-defined element Φ_c belonging to the algebraic dual \mathcal{V}^* of \mathcal{V} which happens to satisfy the following property.

Theorem 1.1 (Theorem 1.1 of [11]). The map $c \mapsto \Phi_c$ is a \mathbb{Q} -vector space isomorphism from \mathcal{I}^* to \mathcal{V}^* .

Our primary interest in [11] arises from a family of theorems about an action of the absolute Galois group $G := \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ on \mathcal{I}^* . If $\sigma \in G$ and $\alpha \in \mathcal{V}$, then $\sigma(\alpha)$ is a well-defined element of \mathcal{V} , and in this way, we obtain a left action of G on \mathcal{V} . This left action on \mathcal{V} induces a corresponding right action of G on \mathcal{V}^* given by $(\Phi, \sigma) \mapsto \Phi \circ \sigma$. Now if $\rho : \mathcal{I}^* \to \mathcal{V}^*$ denotes the isomorphism of Theorem 1.1, then

(1.6)
$$(c,\sigma) \mapsto \rho^{-1}(\Phi_c \circ \sigma)$$

is a right action of G on \mathcal{I}^* .

The author [11] explored this Galois action extensively with an emphasis on the subgroups $\operatorname{Stab}_G(c)$. When G is equipped with the Krull Topology as in [9, Ch. 7], these stabilizers are always closed but not necessarily open. In fact, we established the following (Theorems 2.4 and 2.2 of [11], respectively):

- (A) For every open subgroup H of G, there exists $c \in \mathcal{I}^*$ such that $\operatorname{Stab}_G(c) = H$.
- (B) There exists a consistent map $c \in \mathcal{I}^*$ such that $\operatorname{Stab}_G(c)$ is not open.

Although we presented (A) and (B) as main theorems, they should really be interpreted as corollaries to the same result, a theorem on Galois Invariance [11, Lemma 4.1]. For each number field E, we say that $c \in \mathcal{I}^*$ is *E*-Galois-Invariant if $c = c\sigma$ for all $\sigma \in \operatorname{Gal}(\overline{\mathbb{Q}}/E)$, or equivalently, c is *E*-Galois-Invariant if $\operatorname{Gal}(\overline{\mathbb{Q}}/E) \subseteq \operatorname{Stab}_G(c)$.

Theorem 1.2 (Lemma 4.1 of [11]). Suppose E is a number field and $c : \mathcal{I} \to \mathbb{Q}$ is a consistent map. Then c is E-Galois-Invariant if and only if

$$c(K, v) = \frac{[K_v : E_u]}{[K : E]} c(E, u)$$

for all non-Archimedean places u of E, all finite extensions K/E, and all places v of K dividing u.

This theorem is a powerful tool to construct consistent maps whose stabilizers have properties such as (A) or (B), however, it suffers from an unfortunate limitation. Specifically, the proof of Theorem 1.2 is indirect, relying entirely on the isomorphism given by Theorem 1.1 rather than on any direct Galois action on consistent maps. This is important because the consistent maps used in [11] are only a special case of an object that is needed for other applications as in [12]. Indeed, that article selects a set S of places of \mathbb{Q} and defines

$$\mathcal{J}_S = \{ (K, v) : [K : \mathbb{Q}] < \infty \text{ and } v \in M_{K,S} \},\$$

where $M_{K,S}$ is the set of places of K dividing a place in S. Then a map $c : \mathcal{J}_S \to \mathbb{R}$ is called *consistent* over S if (1.3) is satisfied for all $(K, v) \in \mathcal{J}_S$ and all finite extensions L/K. While [12] presents versions of Theorem 1.1 for these generalized consistent maps, there is no obvious Galois action as in [11]. Therefore, it remains open to provide an analog of Theorem 1.2 for these objects. The present article accomplishes four goals, including a complete resolution to this problem.

- (I) We construct a further generalization of consistent maps beyond that of [12] which replaces S with an object called a *decomposable set*.
- (II) We define a corresponding Galois action on these objects.
- (III) We prove that this action agrees with that of [11] in the special case of (1.6).
- (IV) We establish an analog of Theorem 1.2 for our generalized consistent maps.

All of these results are presented in Section 2 with their proofs appearing in the subsequent sections. As we shall find, our Galois action does not rely on any isomorphism from the space of consistent maps to another \mathbb{R} -vector space. Therefore, not only is our main result more general than Theorem 1.2, but our proof is more direct.

2. Main Results

Let K be a number field. A set $X \subseteq Y$ is called *decomposable over* K if there exists a set S of places of K such that

(2.1)
$$X = \bigcup_{v \in S} Y(K, v)$$

If $v_1, v_2 \in S$ then $Y(K, v_1)$ and $Y(K, v_2)$ are either equal or disjoint. As a result, we may assume without loss of generality that the right had side of (2.1) is a pairwise disjoint union. This observation also implies that the set S is unique, depending only on X and K. Our results require the following preliminary facts about decomposable sets over number fields.

Proposition 2.1. Suppose X is a set of places of $\overline{\mathbb{Q}}$ and let K and L be number fields.

(i) If X is decomposable over K and L, then X is also decomposable over $K \cap L$.

(ii) If $K \subseteq L$ and X is decomposable over K, then X is also decomposable over L.

If a set $X \subseteq Y$ is decomposable over some number field K, then we shall simply say that X is *decomposable*. As a result of Proposition 2.1(i), every decomposable set X has a unique minimal number field F_X over which it is decomposable. Given a number field K, Proposition 2.1(ii) implies that X is decomposable over K if and only if $F_X \subseteq K$. We further write S_X denote the unique set of places of F_X such that

$$X = \bigcup_{p \in S_X} Y(F_X, p).$$

If K is a finite extension of F_X , we easily verify that

$$\{v \in M_K : v \mid p \text{ for some } p \in S_X\} = \{v \in M_K : x \mid v \text{ for some } x \in X\}$$

We shall let $M_{K,X}$ denote the set given by (2.2) and define

$$\mathcal{J}_X = \{ (K, v) : F_X \subseteq K, \ [K : F_X] < \infty, \ v \in M_{K, X} \}$$

Clearly our definition of \mathcal{J}_X generalizes the definitions of \mathcal{I} and \mathcal{J}_S from [11] and [12], respectively, both of which required X to be decomposable over \mathbb{Q} . In the case of [11], we further required X to be the set of all non-Archimedean places of $\overline{\mathbb{Q}}$.

We say that a map $c: \mathcal{J}_X \to \mathbb{R}$ is consistent over X if

(2.3)
$$c(K,v) = \sum_{w|v} c(L,w)$$

for all for all $(K, v) \in \mathcal{J}_X$ and all finite extensions L/K. We write \mathcal{J}_X^* to denote the space of all consistent maps over X, and observe that \mathcal{J}_X^* is an \mathbb{R} -vector space with addition and scalar multiplication defined as in (1.4).

Now let G_X denote the absolute Galois group $G_X = \operatorname{Gal}(\overline{\mathbb{Q}}/F_X)$, where G_X is equipped with the Krull Topology as in [9, Ch. 7]. If K is any field for which $F_X \subseteq K \subseteq \overline{\mathbb{Q}}$ and $\sigma \in G_X$, then $F_X \subseteq \sigma(K) \subseteq \overline{\mathbb{Q}}$, and moreover, K/F_X is Galois if and only if $\sigma(K) = K$ for all $\sigma \in G_X$. If $\{\alpha_1, \alpha_2, \ldots, \alpha_n\}$ is a basis for K over F_X , then $\{\sigma(\alpha_1), \sigma(\alpha_2), \ldots, \sigma(\alpha_n)\}$ is a basis for $\sigma(K)$ over F_X . In particular, K/F_X is finite if and only if $\sigma(K)/F_X$ is finite, and in this case $[K:F_X] = [\sigma(K):F_X]$. Given a place v of K, we let $|\cdot|_v$ be an absolute value from v, and we define the absolute value $|\cdot|_{\sigma(v)}$ on $\sigma(K)$ according to the formula

(2.4)
$$|\alpha|_{\sigma(v)} = |\sigma^{-1}(\alpha)|_v.$$

We shall later prove in Lemma 4.1 that $\sigma(v)$ defines a unique place of $\sigma(K)$ which is independent of the particular absolute value $|\cdot|_v$ from v. Since σ^{-1} fixes the elements of F_X , $\sigma(v)$ and v must divide the same place of F_X . These observations yield a left action of G_X on \mathcal{J}_X given by the formula

(2.5)
$$(\sigma, (K, v)) \mapsto (\sigma(K), \sigma(v)).$$

For simplicity, we shall often write $\sigma(K, v) = (\sigma(K), \sigma(v))$. Based on this notation, we obtain a right action of G_X on \mathcal{J}_X^* given by

$$(c,\sigma)\mapsto c\circ\sigma,$$

and write $c\sigma = c \circ \sigma$. As promised, this Galois action agrees with that of [11] in the special case where X is the set of non-Archimedean places of $\overline{\mathbb{Q}}$.

Theorem 2.2. If $c \in \mathcal{I}^*$ and $\sigma \in G$ then $\Phi_{c\sigma} = \Phi_c \circ \sigma$.

We now proceed with our main result, a generalization of Theorem 1.2 for consistent maps of the type described in this section. Given a finite extension E/F_X , a consistent map $c: \mathcal{J}_X \to \mathbb{R}$ is called *E-Galois-Invariant over* X if $c = c\sigma$ for all $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/E)$.

Theorem 2.3. Suppose that X is a decomposable subset of Y, $c \in \mathcal{J}_X^*$, and E is a finite extension of F_X . Then c is E-Galois-Invariant over X if and only if

(2.6)
$$c(K,v) = \frac{[K_v : E_u]}{[K : E]} c(E,u)$$

for all $u \in M_{E,X}$, all finite extensions K/E, and all places v of K dividing u.

Theorem 2.3 shows that each *E*-Galois-Invariant consistent map over X is uniquely determined by its values at (E, u), for $u \in M_{E,X}$. To make this assertion more precise, we let $\mathcal{J}_{E,X}^*$ denote the set of *E*-Galois-Invariant consistent maps over X, i.e.,

$$\mathcal{J}_{E,X}^* = \left\{ c \in \mathcal{J}_X^* : c = c\sigma \text{ for all } \sigma \in \operatorname{Gal}(\overline{\mathbb{Q}}/E) \right\}.$$

It is straightforward to verify that $\mathcal{J}_{E,X}^*$ is a subspace of \mathcal{J}_X^* for every finite extension E/F_X . Now also define

$$\mathbb{R}^{E,X} = \left\{ (x_u)_{u \in M_{E,X}} : x_u \in \mathbb{R} \right\}.$$

If addition and scalar multiplication are defined coordinate-wise, then clearly $\mathbb{R}^{E,X}$ is a vector space over \mathbb{R} . If $M_{E,X}$ is finite, then $\mathbb{R}^{E,X}$ is isomorphic to \mathbb{R}^n , where $n = \#M_{E,X}$. Otherwise, $M_{E,X}$ is countably infinite and $\mathbb{R}^{E,X}$ is isomorphic to $\mathbb{R}^{\infty} := \{(x_i)_{i \in \mathbb{N}} : x_i \in \mathbb{R}\}$. Regardless, Theorem 2.3 has the following consequence.

Corollary 2.4. Suppose that X is a decomposable subset of Y and E is a finite extension of F_X . Then the map $c \mapsto (c(E, u))_{u \in M_{E,X}}$ defines an \mathbb{R} -vector space isomorphism from $\mathcal{J}_{E,X}^*$ to $\mathbb{R}^{E,X}$.

This observation establishes our earlier remark that E-Galois-Invariant consistent maps over X are uniquely determined by their values at (E, u). For the reader's reference, Corollary 2.4 generalizes another known result [11, Lemma 5.1].

3. Decomposable Sets

This section is dedicated only to the proof of our Proposition 2.1.

Proof of Proposition 2.1. We begin with the proof of (i) and let $F = K \cap L$. Since we have assumed that X is decomposable over K and L, we may write

(3.1)
$$X = \bigcup_{v \in S_K} Y(K, v) = \bigcup_{w \in S_L} Y(L, w),$$

where S_K and S_L are sets of places of K and L, respectively. Each place v of K divides a unique place p of F, so we may let

$$S_F = \{ p \in M_F : v \mid p \text{ for some } v \in S_K \}.$$

To complete the proof, it is enough to establish that

(3.2)
$$X = \bigcup_{p \in S_F} Y(F, p).$$

If $x \in X$ then certainly $x \in Y(K, v)$ for some $v \in S_K$. Then by definition of S_F , there exists $q \in S_F$ such that $v \mid q$. This means that $x \in Y(F, q)$, and hence

$$X \subseteq \bigcup_{p \in S_F} Y(F, p).$$

For the reverse containment, we assume y belongs to the right hand side of (3.2), and let $p_0 \in S_F$ be such that $y \in Y(F, p_0)$. By definition of S_F , there must exist $v_0 \in S_K$ such that $v_0 \mid p_0$, meaning that $Y(K, v_0) \subseteq Y(F, p_0)$. Additionally, since $Y(K, v_0)$ is non-empty, we may select a point $x \in Y(K, v_0)$.

Now let $G = \text{Gal}(\overline{\mathbb{Q}}/F)$. According to the remarks immediately prior to [1, Lemma 3], G acts transitively on $Y(F, p_0)$. Therefore, since $x, y \in Y(F, p_0)$, there exists $\sigma \in G$ such that $\sigma(x) = y$. If we also define

$$H_K = \operatorname{Gal}(\overline{\mathbb{Q}}/K) \text{ and } H_L = \operatorname{Gal}(\overline{\mathbb{Q}}/L),$$

then in the usual Galois correspondence, the field $F = K \cap L$ corresponds to the subgroup of G generated by H_K and H_L . In other words, G is itself generated by H_K and H_L . Therefore, there exist $k_1, k_2, \ldots, k_n \in H_K$ and $\ell_1, \ell_2, \ldots, \ell_n \in H_L$ such that

$$\sigma = \ell_n k_n \cdots \ell_2 k_2 \ell_1 k_1.$$

We now claim that, for all $0 \le m \le n$, we have

$$(3.3) \qquad \qquad (\ell_m k_m \cdots \ell_2 k_2 \ell_1 k_1)(x) \in X$$

We shall proceed by induction on m. When m = 0, the product on the left hand side of (3.3) is empty, so the base case is simply the result of our assumption that $x \in Y(K, v_0) \subseteq X$.

For the inductive step, we let $z = (\ell_{m-1}k_{m-1}\cdots\ell_2k_2\ell_1k_1)(x)$ and assume that $z \in X$. The left hand equality of (3.1) yields a place $v_1 \in S_K$ such that $z \in Y(K, v_1)$. Using the fact that H_K acts on $Y(K, v_1)$, we obtain that $k_m(z) \in Y(K, v_1) \subseteq X$. Now applying the right hand equality of (3.1), we see that $k_m(z) \in$ $Y(L, w_1)$ for some place $w_1 \in S_L$. Of course, H_L acts on $Y(L, w_1)$, implying that $(\ell_m k_m)(z) \in Y(L, w_1) \subseteq X$, and (3.3) follows immediately.

If we apply (3.3) with m = n, we see that

$$y = \sigma(x) = (\ell_n k_n \cdots \ell_2 k_2 \ell_1 k_1)(x) \in X,$$

as required.

Now to prove (ii), we assume only the left hand equality of (3.1). For each $v \in S_K$, we let T_v be the set of places of L that divide v so that

$$Y(K, v) = \bigcup_{w \in T_v} Y(L, w)$$
 for all $v \in S$.

Let $S_L = \bigcup_{v \in S_K} T_v$ and combine this observation with the left hand equality of (3.1) to obtain

. . . .

$$X = \bigcup_{v \in S_K} \bigcup_{w \in T_v} Y(L, w) = \bigcup_{v \in S_L} Y(L, w)$$

so that X satisfies the required property.

4. Galois Actions

Now that we have provided the necessary background on decomposable sets, we proceed with the formal definition of our right group action of G_X on \mathcal{J}_X^* . This action arises from a left action of G_X on \mathcal{J}_X which we vaguely described in Section 2 using (2.4) and (2.5). Nevertheless, it remains necessary to show that (2.4) defines a unique place $\sigma(v)$ of $\sigma(K)$ for all finite extensions K/F_X . Specifically, we must show that the right hand side of (2.4) is an absolute value on $\sigma(K)$ and that the place containing $|\cdot|_{\sigma(v)}$ depends only on v, not on the specific absolute value $|\cdot|_v$ from v. These goals are accomplished in the following lemma.

Lemma 4.1. Let X be a decomposable subset of Y. Suppose p is a place of F_X and K is any field such that $F_X \subseteq K \subseteq \overline{\mathbb{Q}}$. For each $\sigma \in G_X$, the following properties hold:

(i) If $|\cdot|$ is an absolute value on K dividing p then

$$\beta \mapsto |\sigma^{-1}(\beta)|$$

defines an absolute value on $\sigma(K)$ dividing p.

(ii) If $\|\cdot\|$ is an absolute value on $\sigma(K)$ then there exists a unique absolute value $|\cdot|$ on K such that

$$\|\beta\| = |\sigma^{-1}(\beta)| \quad for \ all \ \beta \in \sigma(K).$$

In this case, $\|\cdot\|$ and $|\cdot|$ extend the same absolute value on F_X .

(iii) Suppose $|\cdot|_1, |\cdot|_2$ are absolute values on K and $\|\cdot\|_1, \|\cdot\|_2$ are absolute values on $\sigma(K)$ for which

$$\|\beta\|_1 = |\sigma^{-1}(\beta)|_1 \quad and \quad \|\beta\|_2 = |\sigma^{-1}(\beta)|_2$$

for all $\beta \in \sigma(K)$. Then $|\cdot|_1 \sim |\cdot|_2$ if and only if $\|\cdot\|_1 \sim \|\cdot\|_2$.

Proof. To establish (i), we observe clearly that $|\sigma^{-1}(0)| = 0$. If $|\sigma^{-1}(\alpha)| = 0$, then since $|\cdot|$ is already assumed to be an absolute value on K, we have $\sigma^{-1}(\alpha) = 0$. It now follows that $\alpha = 0$. The remaining two absolute value axioms follow from the fact that $|\cdot|$ is an absolute value on K and σ^{-1} is an automorphism. Now assume that $|\cdot|$ divides the place p of F_X and $\alpha \in F_X$. We clearly have that σ^{-1} fixes the elements of F_X , and therefore

$$|\sigma^{-1}(\alpha)| = |\alpha|,$$

implying that the given map yields an absolute value dividing p.

For (ii), we assume that $\|\cdot\|$ is an absolute value on $\sigma(K)$ and define $|\cdot|$ by

$$|\alpha| = \|\sigma(\alpha)\| \quad \text{for all } \alpha \in K.$$

By the same argument as in our proof of (i), $|\cdot|$ defines an absolute value on K. Moreover, if $\beta \in \sigma(K)$ then

$$\|\beta\| = \|\sigma(\sigma^{-1}(\beta))\| = |\sigma^{-1}(\beta)|,$$

establishing the required equality. To establish the uniqueness property, we assume $|\cdot|'$ is an absolute value on K such that $\|\beta\| = |\sigma^{-1}(\beta)|'$ for all $\beta \in \sigma(K)$. Then if $\alpha \in K$, we have

$$|\alpha|' = |\sigma^{-1}(\sigma(\alpha))|' = ||\sigma(\alpha)|| = |\sigma^{-1}(\sigma(\alpha))| = |\alpha|$$

completing the proof of (ii).

Finally for (iii), we assume that $|\cdot|_1 \sim |\cdot|_2$ so there exists $\theta > 0$ such that $|\alpha|_1 = |\alpha|_2^{\theta}$ for all $\alpha \in K$. Now if $\beta \in \sigma(K)$, we obtain

$$\|\beta\|_1 = |\sigma^{-1}(\beta)|_1 = |\sigma^{-1}(\beta)|_2^{\theta} = \|\beta\|_2^{\theta}$$

as required. The reverse direction has a similar proof.

The combination of Lemma 4.1 (ii) and (iii) shows that (2.4) defines a unique place of $\sigma(K)$. We are now prepared to formalize the corresponding action of G_X on \mathcal{J}_X that was described in (2.5).

Lemma 4.2. The map $(\sigma, (K, v)) \mapsto (\sigma(K), \sigma(v))$ defines a left group action of G_X on \mathcal{J}_X .

Proof. If 1 denotes the identity element of G_X then obviously (1(K), 1(v)) = (K, v) for all $(K, v) \in \mathcal{J}_X$. Now we assume that $(K, v) \in \mathcal{J}_X$ and $\sigma, \tau \in G_X$. By definition we have that $\sigma(\tau(K)) = (\sigma\tau)(K)$. Suppose $|\cdot|_v$ is an absolute value on K from the place v and $\gamma \in \sigma(\tau(K))$. Then by (2.4), we have that

$$\begin{aligned} \gamma|_{(\sigma\tau)(v)} &= |(\sigma\tau)^{-1}(\gamma)|_v \\ &= |\tau^{-1}(\sigma^{-1}(\gamma))|_u \\ &= |\sigma^{-1}(\gamma)|_{\tau(v)} \\ &= |\gamma|_{\sigma(\tau(v))}, \end{aligned}$$

proving that $(\sigma \tau)(v)$ and $\sigma(\tau(v))$ are equal as places of $\sigma(\tau(K))$. We have now established that

$$((\sigma\tau)(K), (\sigma\tau)(v)) = (\sigma(\tau(K)), \sigma(\tau(v)))$$

as is required.

For simplicity, we shall often write $\sigma(K, v) = (\sigma(K), \sigma(v))$ for the action described in Lemma 4.2. At times in our work, it will be useful to restrict this action to various subsets of \mathcal{J}_X . If E/F_X is a finite extension, we define

(4.1)
$$\mathcal{J}_X(E) = \{ (K, v) : E \subseteq K, \ [K : E] < \infty, \ v \in M_{K,X} \}.$$

Certainly whenever $[K : E] < \infty$ we also have $[K : F_X] < \infty$ so that X is decomposable over K for all such fields K. Therefore, the expression $M_{K,X}$ appearing on the right hand side of (4.1) is well-defined. If E'/E is any finite extension, then $\mathcal{J}_X(E') \subseteq \mathcal{J}_X(E)$ with equality if and only if E = E'. Now letting $G_X(E) = \operatorname{Gal}(\overline{\mathbb{Q}}/E)$, we obtain the following additional facts about the group action of Lemma 4.2.

Lemma 4.3. Suppose that X is a decomposable set.

- (i) If E/F_X is a finite extension and E'/E is a finite Galois extension, then $(K, v) \mapsto \sigma(K, v)$ defines an action of $G_X(E)$ on $\mathcal{J}_X(E')$.
- (ii) If E/F_X is a finite extension, then $(K, v) \mapsto \sigma(K, v)$ defines an action of $G_X(E)$ on $\mathcal{J}_X(E)$.
- (iii) If E' is a finite Galois extension of \mathbb{Q} for which X is decomposable over E', then $(K, v) \mapsto \sigma(K, v)$ defines an action of $G_X(E')$ on $\mathcal{J}_X(E')$.

Proof. For (i), it is sufficient to prove that if $(K, v) \in \mathcal{J}_X(E')$ and $\sigma \in G_X(E)$

(4.2)
$$\sigma(K,v) \in \mathcal{J}_X(E').$$

Indeed, since $G_X(E)$ is a subgroup of $G_X = G_X(F_X)$, the remaining group action properties follow from Lemma 4.2.

Therefore, we assume $(K, v) \in \mathcal{J}_X(E')$ and $\sigma \in G_X(E)$. From the fact that $(K, v) \in \mathcal{J}_X(E')$, we know that K is a finite extension of E' and $v \in M_{K,X}$. By definition (2.4) along with Lemma 4.1(ii), $\sigma(v)$ is a place of $\sigma(K)$ dividing the same place of F_X as v. In particular, we have that $\sigma(v) \in M_{\sigma(K),X}$. Now of course, $\sigma(K)$ is a finite extension of $\sigma(E')$. However, $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/E)$ and E'/E is Galois, which implies that $\sigma(E') = E'$. As a result, $\sigma(K)$ is a finite extension of E' and $(\sigma(K), \sigma(v)) \in \mathcal{J}_X(E')$, establishing (4.2).

We obtain (ii) immediately by applying (i) with E = E'. For (iii), the definition of F_X ensures that E' is a finite Galois extension of F_X , and the statement follows by applying (i) with $E = F_X$.

Following the process described in Section 2, the left action of G_X on \mathcal{J}_X given by Lemma 4.2 yields a corresponding right action of G_X on \mathcal{J}_X^* . Specifically, $\sigma \in G_X$ and $c \in \mathcal{J}_X^*$, then clearly $c\sigma : \mathcal{J}_X \to \mathbb{R}$ given by

$$(c\sigma)(K,v) = c(\sigma(K),\sigma(v))$$

(or equivalently, $c\sigma = c \circ \sigma$) is a well defined map. In fact, that map is also consistent over X.

Lemma 4.4. If $c: \mathcal{J}_X \to \mathbb{R}$ is a consistent map over X and $\sigma \in G_X$ then $c\sigma$ is also consistent over X.

Proof. Let $(K, v) \in \mathcal{J}_X$ and let L/K be a finite extension. We need to show that

$$c(\sigma(K), \sigma(v)) = \sum_{\substack{w|v\\7}} c(\sigma(L), \sigma(w)).$$

Clearly $\sigma(L)$ is a finite extension of $\sigma(K)$ and $\sigma(K)$ is a finite extension of F_X . Now let $|\cdot|_v$ be an absolute value on K from the place v, and for each place w of L dividing v, let $|\cdot|_w$ be an absolute value from w that extends $|\cdot|_v$. For each $\alpha \in \sigma(K)$, we have that

$$|\alpha|_{\sigma(w)} = |\sigma^{-1}(\alpha)|_w = |\sigma^{-1}(\alpha)|_v = |\alpha|_{\sigma(v)}$$

which shows that $\sigma(w)$ is a place of $\sigma(L)$ dividing $\sigma(v)$.

Now let $W_v(L/K)$ be the set of places of L that divide v. Our above observations ensure that σ : $W_v(L/K) \to W_{\sigma(v)}(\sigma(L)/\sigma(K))$ is a well-defined map. However, if $x \in W_{\sigma(v)}(\sigma(L)/\sigma(K))$ then clearly $\sigma^{-1}(x) \in W_v(L/K)$ and $\sigma(\sigma^{-1}(x)) = x$, showing that σ is surjective. Moreover, if $\sigma(w_1) = \sigma(w_2)$ then

$$|\alpha|_{\sigma(w_1)} = |\alpha|_{\sigma(w_2)}$$
 for all $\alpha \in \sigma(L)$.

By the definition (2.4), we obtain that

$$\sigma^{-1}(\alpha)|_{w_1} = |\sigma^{-1}(\alpha)|_{w_2}$$
 for all $\alpha \in \sigma(L)$.

From the fact that σ is a bijection from L to $\sigma(L)$, we conclude that $w_1 = w_2$. We have now shown that $\sigma: W_v(L/K) \to W_{\sigma(v)}(\sigma(L)/\sigma(K))$ is a bijection. Finally, we obtain

$$\sum_{w|v} c(\sigma(L), \sigma(w)) = \sum_{w|\sigma(v)} c(\sigma(L), w) = c(\sigma(K), \sigma(v)),$$

where the last equality follows from the fact that c is consistent.

As a result of Lemma 4.4, we obtain easily that $c \mapsto c\sigma$ defines a right action of G_X on \mathcal{J}_X^* . With this information in mind, we are now able to prove Theorem 2.2.

Proof of Theorem 2.2. Let $\alpha \in \overline{\mathbb{Q}}^{\times}$ and suppose that K is a number field containing α . Then clearly $\sigma(K)$ contains $\sigma(\alpha)$ and we obtain

$$\Phi_c(\sigma(\alpha)) = \sum_{w \in M^0_{\sigma(K)}} \frac{c(\sigma(K), w)}{\log p_w} \log \|\sigma(\alpha)\|_w.$$

Now using definition (2.4), we have that

$$\Phi_c(\sigma(\alpha)) = \sum_{w \in M^0_{\sigma(K)}} \frac{c(\sigma(K), w)}{\log p_w} \log \|\alpha\|_{\sigma^{-1}(w)}$$
$$= \sum_{w \in M^0_{\sigma(K)}} \frac{c(\sigma(K), \sigma(\sigma^{-1}(w)))}{\log p_w} \log \|\alpha\|_{\sigma^{-1}(w)}$$

For each $w \in M^0_{\sigma(K)}$, there exists a unique place v of K such that $v = \sigma^{-1}(w)$, and in this case, v and w divide the same place of \mathbb{Q} . Therefore, we have that

$$\Phi_c(\sigma(\alpha)) = \sum_{v \in M_K^0} \frac{c(\sigma(K), \sigma(v))}{\log p_v} \log \|\alpha\|_v = \Phi_{c\sigma}(\alpha)$$

as required.

5. Proof of Theorem 2.3

Our proof requires a preliminary theorem and corollary that generalizes [12, Theorem 5.2] and heavily borrows from its proof. Assuming that E is a finite extension of F_X , we may also define a version of consistent for the more specialized set $\mathcal{J}_X(E)$. Specifically, we say that $c : \mathcal{J}_X(E) \to \mathbb{R}$ is consistent over X with respect to E if

$$c(K,v) = \sum_{w|v} c(L,w)$$

for all $(K, v) \in \mathcal{J}_X(E)$ and all finite extensions L/K. Extending our earlier notation, we write $\mathcal{J}_X^*(E)$ to denote the space of all consistent maps over X with respect to E. Clearly, $\mathcal{J}_X^*(E)$ forms a vector space

over \mathbb{R} with operations defined as in (1.4). Although $\mathcal{J}_X^*(E)$ and \mathcal{J}_X^* are technically distinct spaces as their elements have different domains, there is a canonical isomorphism between them.

Theorem 5.1. Suppose X is a decomposable set and E is a finite extension of F_X . Then the map $c \mapsto c \mid_{\mathcal{J}_K(E)}$ defines an \mathbb{R} -vector space isomorphism from \mathcal{J}_X^* to $\mathcal{J}_X^*(E)$.

Proof. let $\rho: \mathcal{J}_X^* \to \mathcal{J}_X^*(E)$ be the map given in the statement of the theorem, i.e.,

$$\rho(c) = c \mid_{\mathcal{J}_K(E)}$$

To see that ρ is a well-defined map, we let $c \in \mathcal{J}_X^*$ and $(K, v) \in \mathcal{J}_X(E)$. Certainly $(K, v) \in \mathcal{J}_X$, so we apply (2.3) to obtain

$$[\rho(c)](K,v) = c(K,v) = \sum_{w|v} c(L,w)$$

showing that $\rho(c)$ is, in fact, a consistent map over X with respect to E. Now assuming that $c, d \in \mathcal{J}_X^*$ and $(K, v) \in \mathcal{J}_K(E)$, we have

$$\begin{aligned} [\rho(c+d)](K,v) &= (c+d)(K,v) \\ &= c(K,v) + d(K,v) \\ &= [\rho(c)](K,v) + [\rho(d)](K,v) \\ &= [\rho(c) + \rho(d)](K,v). \end{aligned}$$

Further supposing that $r \in \mathbb{R}$, we get that

$$[\rho(rc)](K,v) = (rc)(K,v) = rc(K,v) = r[\rho(c)](K,v)$$

proving ρ is a linear transformation.

To prove that ρ is surjective, we will show that each consistent map $c : \mathcal{J}_X(E) \to \mathbb{R}$ extends to a consistent map $d : \mathcal{J}_X \to \mathbb{R}$. To accomplish this goal, we let $(K, v) \in \mathcal{J}_X$ and let L be a finite extension of both K and E. If w is a place of L dividing v, then w must also divide a place in S_X . This means that $(L, w) \in \mathcal{J}_X(E)$ and we may define

$$d_L(K,v) = \sum_{w|v} c(L,w).$$

We now claim that d_L is independent of L. To see this, we assume that M is a finite extension of L. Now using that fact that $c \in \mathcal{J}_X^*(E)$, we obtain

$$d_L(K,v) = \sum_{w|v} c(L,w) = \sum_{w|v} \sum_{u|w} c(M,u) = \sum_{u|v} c(M,u) = d_M(K,v).$$

Since there exists a smallest number field containing both E and K, we conclude that $d_L(K, v)$ does not depend on K and we may define

(5.1)
$$d(K, v) = \sum_{w|v} c(L, w),$$

where L is any finite extension of K containing E. In this way, $d : \mathcal{J}_X \to real$ is a well-defined map. Furthermore, if $(K, v) \in \mathcal{J}_X(E)$, then we may select L = K and it follows that d(K, v) = c(K, v). In other words, $\rho(d) = c$.

To see that d is consistent over X, we let $(K, v) \in \mathcal{J}_X$ and let L/K be a finite extension. If M is a finite extension of L and E we apply definition (5.1) twice to obtain

$$\sum_{w|v} d(L, w) = \sum_{w|v} \sum_{u|w} c(M, u) = \sum_{u|v} c(M, u) = d(K, v).$$

These observations prove that d is consistent and ρ is surjective.

Finally, to show that ρ is injective, we let $d \in \ker(\rho)$ so that d(L, w) = 0 for all $(L, w) \in \mathcal{J}_X(E)$. Now let $(K, v) \in \mathcal{J}_X$ and choose L to be a finite extension of both K and E. Therefore, $(L, w) \in \mathcal{J}_X(E)$ for all places w of L dividing v. Now using the consistency of d over X, we conclude that

$$d(K,v) = \sum_{\substack{w|v\\9}} d(L,w) = 0,$$

as required.

Theorem 5.1 has the following simple but important consequence, showing essentially that checking *E*-Galois-Invariance requires only considering points in $\mathcal{J}_X(E)$.

Corollary 5.2. Suppose X is a decomposable subset of Y and $c \in \mathcal{J}_X^*$. Further assume that E is a finite extension of F_X and $\sigma \in G_X(E)$. Then $c = c\sigma$ if and only if

(5.2)
$$c(K,v) = c(\sigma(K), \sigma(v)) \quad for \ all \ (K,v) \in \mathcal{J}_X(E).$$

Proof. If $c = c\sigma$ as elements of \mathcal{J}_X^* , this means that $c(K, v) = (c\sigma)(K, v) = c(\sigma(K), \sigma(v))$ for all $(K, v) \in \mathcal{J}_X$. Since $\mathcal{J}_X(E) \subseteq \mathcal{J}_X$, we immediately obtain (5.2).

Now assume that (5.2) holds. This means that $c(K, v) = (c\sigma)(K, v)$ for all $(K, v) \in \mathcal{J}_X(E)$, and hence, we may define $d: \mathcal{J}_X(E) \to \mathbb{R}$ by

$$d = c \mid_{\mathcal{J}_X(E)} = (c\sigma) \mid_{\mathcal{J}_X(E)}.$$

Clearly $d \in \mathcal{J}_X^*(E)$, so according to Theorem 5.1, d must have a unique extension to an element of \mathcal{J}_X^* . As c and $c\sigma$ are both extensions of d to an element of \mathcal{J}_X^* , it follows that $c = c\sigma$ as elements of \mathcal{J}_X^* .

Now that we may work in $\mathcal{J}_X(E)$ rather than needing to work in \mathcal{J}_X , we are prepared to prove Theorem 2.3.

Proof of Theorem 2.3. We begin by assuming that c is E-Galois-Invariant over X. Suppose that $u \in M_{E,X}$ and K is a finite extension of E. In this case, there exists a finite Galois extension L of E such that $K \subseteq L$. We shall first prove that

(5.3)
$$c(L,w) = \frac{[L_w : E_u]}{[L : E]} c(E,u)$$

for all places w of L dividing u. Now let $H = \operatorname{Gal}(L/E)$ and let $W_u(L/E)$ be the set of places of L that divide u as in the notation of [1].

It is well-known that $(\bar{\sigma}, v) \mapsto \bar{\sigma}(v)$ defines a transitive action of H on $W_u(L/E)$. Therefore, if $w_1, w_2 \in W_u(L/E)$, then there exists $\bar{\sigma} \in H$ such that $\bar{\sigma}(v_1) = v_2$. Since $\sigma \mapsto \sigma|_L$ is a surjective group homomorphism from $G_X(E)$ to H, there exists $\sigma \in G_X(E)$ such that $\sigma(\alpha) = \bar{\sigma}(\alpha)$ for all $\alpha \in L$. Since $\bar{\sigma}^{-1}$ is certainly a bijection, we may replace α by $\bar{\sigma}^{-1}(\alpha)$ to obtain

(5.4)
$$\sigma^{-1}(\alpha) = \bar{\sigma}^{-1}(\alpha) \quad \text{for all } \alpha \in L.$$

Additionally, $\sigma(w_1)$ defines a place of $\sigma(L)$ dividing u. In this situation, however, L/E is Galois and $\sigma \in G_X(E)$, meaning that $\sigma(L) = L$ and $\sigma(w_1)$ is a place of L. Combining this observation with (5.4), we conclude that $\sigma(w_1) = \bar{\sigma}(w_1) = w_2$. Now since $(L, w_1) \in J_X(E)$ and $\sigma \in G_X(E)$ we obtain

$$\sigma(L, w_1) = (\sigma(L), \sigma(w_1)) = (L, w_2).$$

Applying c to both sides of this equality yields $(c\sigma)(L, w_1) = c(L, w_2)$. We have assumed that c is E-Galois-Invariant over X and $\sigma \in G_X(E)$, and hence, we obtain

$$c(L, w_1) = c(L, w_2).$$

Since w_1 and w_2 were arbitrary elements of $W_u(L/E)$, we have shown that c(L, w) has the same value for all $w \in W_u(L/E)$.

In order to prove (5.3), we now fix a place $w \in W_u(L/E)$ and note that

$$c(L, w) = c(L, w')$$
 for all $w' \in W_u(L/E)$.

Using the fact that c is consistent over X, we obtain that

(5.5)
$$c(E,u) = \sum_{w'|u} c(L,w') = c(L,w) \cdot \#W_u(L/E).$$

From (1.1), we also have that

$$[L:E] = \sum_{\substack{w' \mid u \\ 10}} [L_{w'}:E_u],$$

but since L/E is Galois, all of the local degrees on the right hand side are equal. Consequently, we have $[L:E] = [L_w:E_u] \cdot \#W_u(L/E)$. Combining this formula with (5.5), we obtain

$$\frac{c(E,u)}{c(L,w)} = \frac{[L:E]}{[L_w:E_u]}$$

and (5.3) follows.

Now assume that v is a place of K dividing u. Since $v \in M_{K,X}$ we may apply the fact that c is consistent over X to obtain

$$c(K, v) = \sum_{w|v} c(L, w).$$

Then (5.3) yields

$$c(K,v) = \sum_{w|v} \frac{[L_w:E_u]}{[L:E]} c(E,u)$$
$$= c(E,u) \cdot \sum_{w|v} \frac{[L_w:E_u]}{[L:E]}$$
$$= c(E,u) \cdot \sum_{w|v} \frac{[L_w:K_v] \cdot [K_v:E_u]}{[L:K] \cdot [K:E]}$$
$$= \frac{[K_v:E_u]}{[K:E]} \cdot c(E,u) \cdot \sum_{w|v} \frac{[L_w:K_v]}{[L:K]}.$$

Applying (1.1), the sum on the right hand side is equal to 1 and (2.6) follows.

We now assume (2.6), let $(K, v) \in \mathcal{J}_X(E)$, and let $\sigma \in G_X(E)$. Because of Lemma 5.2, to show that c is E-Galois-Invariant over X, it is enough to prove that

(5.6)
$$c(K,v) = c(\sigma(K), \sigma(v)).$$

Since K is a finite extension of E, v must divide a unique place u of E. Since v is assumed to divide a place in S_X , it follows that $u \in M_{E,X}$. As a result, (2.6) applies to yield

(5.7)
$$c(K,v) = \frac{[K_v : E_u]}{[K : E]} c(E,u).$$

By Lemma 4.3(ii), $G_X(E)$ acts on $J_X(E)$ so we have $(\sigma(K), \sigma(v)) \in \mathcal{J}_X(E)$. Additionally, since σ^{-1} fixes the elements of E, both v and $\sigma(v)$ must divide the same place of E. Therefore, $\sigma(v)$ divides u and we may apply (2.6) with $(\sigma(K), \sigma(v))$ in place of (K, v). To simplify the notation we write $L = \sigma(K)$ and obtain

(5.8)
$$c(L, \sigma(v)) = \frac{[L_{\sigma(v)} : E_u]}{[L : E]} c(E, u).$$

The map $\sigma: K \to L$ is certainly a vector space isomorphism over E which implies that [K:E] = [L:E]. Also, if $\alpha, \beta \in K$ then

$$|\alpha - \beta|_v = |\sigma(\alpha - \beta)|_{\sigma(v)} = |\sigma(\alpha) - \sigma(\beta)|_{\sigma(v)}$$

meaning that σ is also an isometry that fixes the elements of E_u . Consequently, we also have $[K_v : E_u] = [L_{\sigma(v)} : E_u]$. We have now shown that the right hand side of (5.7) and (5.8) are equal so that (5.6) follows. \Box

Proof of Corollary 2.4. Let ρ be the map described in the statement of the theorem and note that ρ is easily verified to be a linear transformation. To see that ρ is injective, assume that $c \in \mathcal{J}_{E,X}^*$ is such that $\rho(c) = 0$. By definition of ρ , we obtain that c(E, u) = 0 for all $u \in M_{E,X}$. Now since c is E-Galois-Invariant, we apply Theorem 2.3 to conclude that c(K, v) = 0 for all $(K, v) \in \mathcal{J}_X(E)$. In other words, we have shown that $c \mid_{\mathcal{J}_X(E)} = 0$ as an element of $\mathcal{J}_X^*(E)$. Now it follows from Theorem 5.1 that c = 0 in \mathcal{J}_X^* , proving that ρ is injective. To establish surjectivity, we assume that $(x_u)_{u \in M_{E,X}} \in \mathbb{R}^{E,X}$. For each finite extension K/E and $v \in M_{K,X}$, there exists a unique place $u \in M_{E,X}$ such that $v \mid u$. Then we may define

$$c(K,v) = \frac{[K_v : E_u]}{[K : E]} \cdot x_v$$

and easily verify using (1.1) that $c \in \mathcal{J}_X^*(E)$. Clearly we also have that $\rho(c) = (x_u)_{u \in M_{E,X}}$ Because of Theorem 5.1, c has a unique extension to an element $c \in \mathcal{J}_X^*$ which certainly satisfies (2.6), and as a result of Theorem 2.3, $c \in \mathcal{J}_{E,X}^*$.

References

- [1] D. Allcock and J. D. Vaaler. A Banach space determined by the Weil height. Acta Arith., 136(3):279–298, 2009.
- [2] P. Fili and Z. Miner. Norms extremal with respect to the Mahler measure. J. Number Theory, 132(1):275–300, 2012.
- [3] P. Fili and Z. Miner. Orthogonal decomposition of the space of algebraic numbers and Lehmer's problem. J. Number Theory, 133(11):3941–3981, 2013.
- [4] P. Fili and Z. Miner. A generalization of Dirichlet's unit theorem. Acta Arith., 162(4):355-368, 2014.
- [5] A. Fröhlich. Local fields. In Algebraic Number Theory (Proc. Instructional Conf., Brighton, 1965), pages 1–41. Academic Press, London, 1967.
- [6] R. Grizzard and J. D. Vaaler. Multiplicative approximation by the Weil height. Trans. Amer. Math. Soc., 373(5):3235–3259, 2020.
- [7] A. M. Hughes. A Weil Banach algebra for multiplicative algebraic numbers. J. Number Theory, 196:244-271, 2019.
- [8] D. H. Lehmer. Factorization of certain cyclotomic functions. Ann. of Math. (2), 34(3):461-479, 1933.
- [9] J. S. Milne. Fields and Galois theory. Kea Books, Ann Arbor, MI, [2022] ©2022.
- [10] J. Neukirch. Algebraic number theory, volume 322 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1999. Translated from the 1992 German original and with a note by Norbert Schappacher, With a foreword by G. Harder.
- [11] C. L. Samuels. A classification of \mathbb{Q} -valued linear functionals on $\overline{\mathbb{Q}}^{\times}$ modulo units. Acta Arith., 205(4):341–370, 2022.
- [12] C. L. Samuels. Consistent maps and their associated dual representation theorems. *Preprint*, 2023. https://drive.google.com/file/d/17ogsVpfzucjGcf307UDA8CAbMOcocodc/view?usp=drive_link.
- [13] J. D. Vaaler. Heights on groups and small multiplicative dependencies. Trans. Amer. Math. Soc., 366(6):3295–3323, 2014.

Christopher Newport University, Department of Mathematics, 1 Avenue of the Arts, Newport News, VA 23606, USA

Email address: charles.samuels@cnu.edu