INTERIOR CONTROL FOR SURFACES WITH POSITIVE SCALAR CURVATURE AND ITS APPLICATION

SHULI CHEN, JIANCHUN CHU, AND JINTIAN ZHU

ABSTRACT. Let M^n , $n \in \{3, 4, 5\}$, be a closed aspherical *n*-manifold and $S \subset M$ a subset consisting of disjoint incompressible embedded closed aspherical submanifolds (possibly with different dimensions). When n = 3, 4, we show that $M \setminus S$ cannot admit any complete metric with positive scalar curvature. When n = 5, we obtain the same result either when S contains a submanifold of codimension 1 or 2, or when S itself is a connected submaifold of codimension ≥ 3 . The key ingredient is a new interior control for the extrinsic diameter of surfaces with positive scalar curvature.

1. INTRODUCTION

Scalar curvature is the weakest curvature invariant of a Riemannian metric, but the existence of metrics with positive scalar curvature still places strong topological constraints on the underlying manifolds. The well-known aspherical conjecture asserts that closed aspherical manifolds cannot admit any smooth metric with positive scalar curvature (see [17, 19, 8]). Recall that a manifold is called *aspherical* if it has contractible universal cover (or equivalently, if the homotopy group π_i vanishes for all $i \geq 2$). Aspherical manifolds are also known as $K(\pi, 1)$ -spaces or Eilenberg–MacLane spaces in topology.

So far, there have been many progresses on the aspherical conjecture in low dimensions. The two dimensional case follows directly from the Gauss-Bonnet formula combined with the classification of closed surfaces. In 1983, Gromov and Lawson [11] verified the three dimensional case with the help of some filling radius estimate. Closely related, it is worth mentioning that Schoen and Yau [18] previously obtained a fundamental group obstruction for 3-manifolds with positive scalar curvature, which in fact can be also used to confirm the three dimensional aspherical conjecture, combined with the final resolution of the virtually Haken conjecture [1]. In dimension four, the special case with non-zero first Betti number was first confirmed by Wang [21]. We note that there exist infinitely many closed aspherical 4-manifolds that are homology 4-spheres [16], which have zero first Betti number. The general case in dimension four was recently established by Chodosh and Li [6] based on a previous outline from Schoen and Yau [19]. In dimension five, the aspherical conjecture was recently confirmed independently by Chodosh and Li in the same paper [6] and also Gromov in his paper [10] based on the recent development of Gromov's μ -bubble method [9]. Shortly after that, Chodosh, Li and Liokumovich [7] improved the aspherical conjecture into a mapping version in dimensions four and five. Despite of these progresses up to dimension five, the aspherical conjecture is still widely open in dimensions greater than five.

With the help of the μ -bubble method, there have been various efforts to generalize known topological obstructions for positive scalar curvature on closed manifolds to those on non-compact complete manifolds, where the situations are more complicated since the scalar curvature may decay to zero at infinity. For results on non-compact manifolds obtained from (generalized) connected sum operations from *n*-torus (or even closed Scheon-Yau-Schick manifolds), the readers can refer to [6, 13, 3, 4]. When the underlying manifold is only aspherical, due to the lack of nonzero homology classes, the μ -bubble method cannot be applied in a direct way to overcome the issue of non-compactness. As a result, not many results are known for non-compact manifolds constructed from closed aspherical ones. In particular, Gromov [9, p. 151] asked the following start-up question:

Question 1.1. Are there complete metrics with positive scalar curvature on closed aspherical manifolds with punctures of dimension 4 and 5?

In the previous work [5], we are able to give a negative answer to Gromov's question. More generally, we considered the question whether there is any complete metric with positive scalar curvature on connected sums of non-compact manifolds with closed aspherical ones, and we proved the following:

Theorem 1.2 (Theorem 1.2 of [5]). Let N^n , $n \in \{3, 4, 5\}$, be a closed aspherical manifold and let X^n be an arbitrary n-manifold. Then the connected sum Y = N # X admits no complete metric with positive scalar curvature.

In this paper, we want to generalize Gromov's start-up question by removing other submanifolds beyond punctures. Namely, we consider the following question:

Question 1.3. Let M^n be a closed aspherical n-manifold and $S \subset M$ a closed subset. For which S can we guarantee that $M \setminus S$ cannot admit any complete metric with positive scalar curvature?

We note that there are several previous results related to this question. In [11], Gromov and Lawson showed that manifolds of the form $\mathbb{T}^n \setminus \mathbb{T}^k$ for $0 \leq k < n$ and $n \geq 2$ do not carry a complete metric with positive scalar curvature. In [20], Shi, Wang, Wu, and the third-named author showed that manifolds of the form $M \setminus \Gamma$ do not carry a complete metric with positive scalar curvature, where M^n , $3 \leq n \leq 7$ is a closed Schoen–Yau–Schick manifold and Γ is a submanifold such that either dim $\Gamma \leq 1$ or the first Betti number $b_1(\Gamma) \leq n-3$. We also note that Question 1.4 is also closely related to the Schoen conjecture regarding uniformly Euclidean (L^{∞}) metrics up to a blow-up trick [14].

The work in this paper can be considered as a generalization of the abovementioned result by Gromov and Lawson in [11]. In our view, the key feature of the pair $(\mathbb{T}^n, \mathbb{T}^k)$ is that the inclusion map $\pi_1(\mathbb{T}^k) \to \pi_1(\mathbb{T}^n)$ is injective. Recall that if a submanifold N in M has injective map $\pi_1(N) \to \pi_1(M)$, it is called incompressible in M. Under such consideration, we restrict our attention to the case when S consists of incompressible submanifolds, and we propose the following

Question 1.4. Let M^n be a closed aspherical n-manifold and $S \subset M$ a subset consisting of disjoint incompressible embedded closed aspherical submanifolds (possibly with different dimensions). Can $M \setminus S$ admit any complete metric with positive scalar curvature?

In this paper, we fully answer Question 1.4 for n = 3 and 4. Namely, we are able to show

Theorem 1.5. Let M^n , n = 3, 4, be a closed aspherical n-manifold and $S \subset M$ a subset consisting of disjoint incompressible embedded closed aspherical submanifolds (possibly with different dimensions). Then $M \setminus S$ cannot admit any complete metric with positive scalar curvature. Moreover, if g is a complete smooth metric on $M \setminus S$ with nonnegative scalar curvature, then g is flat and there is a finite cover (\tilde{M}, \tilde{S}) of the pair (M, S) such that $\tilde{M} \setminus \tilde{S}$ consists of cylinders $\mathbb{T}^{n-1} \times \mathbb{R}$.

Throughout this paper, we call (\tilde{M}, \tilde{S}) a cover of the pair (M, S) if \tilde{M} is a cover of M associated with the covering map $\pi : \tilde{M} \to M$ and $\tilde{S} = \pi^{-1}(S)$. If \tilde{M} is a finite cover of M, then we call (\tilde{M}, \tilde{S}) a finite cover of (M, S).

For n = 5, we can only give a partial answer to Question 1.4 either when S contains a submanifold with codimension 1 or 2, or when S itself is a connected submanifold with codimension ≥ 3 .

Theorem 1.6. Let M be a closed aspherical 5-manifold and $S \subset M$ a subset consisting of disjoint incompressible closed aspherical embedded submanifolds (possibly with different dimensions). If either S contains a submanifold of codimension 1 or 2, or S is a connected submanifold of codimension ≥ 3 , then $M \setminus S$ cannot admit any complete metric with positive scalar curvature. Moreover, if g is a complete smooth metric on $M \setminus S$ with nonnegative scalar curvature, then g is flat and there is a finite cover (\tilde{M}, \tilde{S}) of the pair (M, S)such that $\tilde{M} \setminus \tilde{S}$ consists of cylinders $\mathbb{T}^{n-1} \times \mathbb{R}$.

The strategy for the proof of the above theorems essentially follows from the one developed in [5], except that some necessary modifications will be made to obtain extrinsic diameter estimates. In particular, the following interior control of the extrinsic diameter plays a key role when n = 3 and 4.

Proposition 1.7. Given a positive and continuous function $L: [0, +\infty) \rightarrow (0, +\infty)$ there is a positive function $K: (0, +\infty) \rightarrow (0, +\infty)$ such that the following property holds: for any $s_0 > 0$, if (Σ, g, ρ) is a triple satisfying

- (Σ, g) is a connected Riemannian surface with non-empty boundary and $\rho: \Sigma \to [0, +\infty)$ is a smooth and proper function with Lip $\rho < 1$;
- $\rho > K(s_0)$ on $\partial \Sigma$;

4

• (Σ, g) has \mathbb{T}^* -stabilized scalar curvature $R \ge L \circ \rho$ in $\Sigma_{K(s_0)}$, where we use the notation $\Sigma_s := \rho^{-1}([0, s]),$

then we can find finitely many pairwise disjoint disks $\{D_j\}$ in $\Sigma_{K(s_0)}$ such that

$$\Sigma_{s_0} \subset \bigcup_j D_j$$

and for each connected component $\Sigma_{s_0}^o$ of Σ_{s_0} , the extrinsic diameter of $\Sigma_{s_0}^o$ in $(\Sigma_{K(s_0)}, g)$ is bounded by

$$4\pi \left(\min_{[0,K(s_0)]} L\right)^{-\frac{1}{2}}.$$

1.1. Outline of the proof. In this subsection, we shall outline the proof for Theorem 1.5 and Theorem 1.6. For readers' convenience, we first recall the previous strategy for Theorem 1.2 from [5], and then introduce necessary modifications to be made for Theorem 1.5 and Theorem 1.6.

1.1.1. The previous strategy for Theorem 1.2. First let us briefly recall the strategy developed in [5] for the proof of Theorem 1.2, which is essentially a relative version of the chain-closing program first proposed in [19] and further developed in [6].

For $n \in \{3, 4, 5\}$, we consider the non-compact connected sum N # X, where N is a closed aspherical n-manifold, and X denotes an arbitrary noncompact n-manifold. For simplicity, we denote N # X by Y. Throughout this paper, all homology groups are assumed to have Z coefficients.

Suppose that g is a complete metric of positive scalar curvature on Y = N # X, then we aim to derive a contradiction. Since N has a contractible universal cover \tilde{N} , we can consider the corresponding cover $\tilde{Y} = \tilde{N} \#_{\pi_1(N)} X$ of Y. By definition the connected sum $\tilde{Y} = \tilde{N} \#_{\pi_1(N)} X$ induces a natural decomposition

$$X = \tilde{N}_{\varepsilon} \cup \tilde{X}_{\varepsilon}$$

in the following way: for each X we take away a small ball and label it by $\tilde{X}_{\varepsilon,i}$ with index $i \in \pi_1(N)$, then we can take the union

$$\tilde{X}_{\varepsilon} = \left(\bigcup_{i} \tilde{X}_{\varepsilon,i}\right)$$

such that the complement \tilde{N}_{ε} of \tilde{X}_{ε} in \tilde{Y} is diffeomorphic to

$$\tilde{N} - \left(\bigcup_i B_i\right).$$

It is clear that we can guarantee both \tilde{N}_{ε} and \tilde{X}_{ε} to be $\pi_1(N)$ -invariant under the Deck transformation of the covering $\tilde{Y} \to Y$.

The whole proof of Theorem 1.2 relies on the following two topological properties of \tilde{Y} :

- (Homological property) From the contractibility of \tilde{N} , one can show that the relative homology group $H_k(\tilde{Y}, \tilde{X}_{\varepsilon})$ is zero for k = n - 1and the map $H_k(\tilde{Y}) \to H_k(\tilde{Y}, \tilde{X}_{\varepsilon})$ is zero for all k.
- (Quantitative filling property) Since \tilde{N}_{ε} is the cover of a compact manifold, for any $k \in \mathbb{N}_+$ there is a function $F: (0, +\infty) \to (0, +\infty)$ depending only on the triple (Y, N_{ε}, g) such that if C is a relative k-boundary in $\mathcal{B}_k(\tilde{Y}, \tilde{X}_{\varepsilon})$ contained in some geodesic ball $B_r^{\tilde{g}}(\tilde{q})$ of (\tilde{Y}, \tilde{g}) with $\tilde{q} \in \tilde{N}_{\varepsilon}$, then one can find a relative (k + 1)-chain $\Gamma \in \mathcal{Z}_{k+1}(\tilde{Y}, \tilde{X}_{\varepsilon})$ contained in the geodesic ball $B_{F(r)}^{\tilde{g}}(\tilde{q})$ of (\tilde{Y}, \tilde{g}) with $\partial \Gamma = C \mod \tilde{X}_{\varepsilon}$.

The basic idea is to complete a chain-closing program in \tilde{Y} . Namely, we fix a proper line $\tilde{\sigma}$ in \tilde{N}_{ε} and take the boundary of a tubular neighborhood \mathcal{N} of $\tilde{\sigma}|_{[0,+\infty)}$. From our construction $\partial \mathcal{N}$ is a locally-finite relative chain having non-zero intersection number with $\tilde{\sigma}$. Due to the homology property of \tilde{Y} , if one can close this chain without creating further intersections, then the newly obtained relative cycle represents a non-trivial relative homology class in $H_{n-1}(\tilde{Y}, \tilde{X}_{\varepsilon})$, which leads to a desired contradiction.

To be more precise, through cutting $\partial \mathcal{N}$ at a finite length we can obtain a hypersurface \tilde{M}_{n-1} with boundary in \tilde{Y} , which has non-zero algebraic intersection with $\tilde{\sigma}$. In this step, we use the fact $H_{n-1}(\tilde{Y}, \tilde{X}_{\varepsilon}) = 0$. Let M_{n-1} be the minimizing hypersurface obtained from solving the Plateau problem with the prescribed boundary $\partial \tilde{M}_{n-1}$. After using μ -bubble method we can further construct a codimension-two closed submanifold M_{n-2} in \tilde{Y} . Through adjusting the size of the tubular neighborhood as well as the cutting length we can guarantee

- M_{n-2} and ∂M_{n-1} enclose a bounded region Ω_{n-1} in M_{n-1} which is disjoint from $\tilde{\sigma}$;
- dist $(M_{n-2}, \tilde{\sigma}) > L$ for arbitrarily large L > 0;
- M_{n-2} has \mathbb{T}^* -stabilized scalar curvature no less than $R(g) \mu_{\text{loss}}$ for arbitrarily small $\mu_{\text{loss}} > 0$.

The last property above guarantees that the \mathbb{T}^* -stabilized scalar curvature on $M_{n-2} \cap \tilde{N}_{\varepsilon}$ has a positive lower bound <u>R</u>. The next step is to fill M_{n-2} with chain without creating further intersections with $\tilde{\sigma}$ and we have to make discussion according to the dimension n.

In dimensions 3 and 4, we have the following two properties:

(1) Note that M_{n-2} is either a curve or a surface, so the scalar curvature lower bound <u>R</u> guarantees that the surface $M_{n-2} \cap \tilde{N}_{\varepsilon}$ is within a distance $C(\underline{R})$ from its boundary.

S. CHEN, J. CHU, AND J. ZHU

(2) After enlarging \tilde{N}_{ε} we can guarantee that the ends $\tilde{X}_{\varepsilon,i}$ are at a far enough distance away from each other. Then the surface M_{n-2} can only enter one $\tilde{X}_{\varepsilon,i}$. Since $\partial \tilde{X}_{\varepsilon,i}$ is closed, we can bound the *extrinsic* diameter of the boundary of $M_{n-2} \cap \tilde{N}_{\varepsilon}$ by the extrinsic diameter of $\partial \tilde{X}_{\varepsilon,i}$, which is a constant independent of *i*.

Combining these two points, we can bound the extrinsic diameter of $M_{n-2} \cap \tilde{N}_{\varepsilon}$ by $C(\underline{R}) + \operatorname{diam}(\partial \tilde{X}_{\varepsilon,i})$.

Once we obtain the above extrinsic diameter estimate, we can then apply the quantitative filling property and fill in M_{n-2} by some relative (n-1)chain Γ within some fixed distance independent of L. When L is large this chain will have no intersection with $\tilde{\sigma}$. Then we can form the relative (n-1)-cycle

$$\tilde{\Gamma}_{n-1} := \tilde{M}_{n-1} + \Omega_{n-1} + \Gamma.$$

This relative cycle has non-zero algebraic intersection number with the line $\tilde{\sigma}$, which contradicts $H_{n-1}(\tilde{Y}, \tilde{X}_{\varepsilon}) = 0$.

In dimension 5, since M_{n-2} is a 3-manifold, the property (1) no longer holds for M_3 . In order to deal with this issue we have to make a refinement of Chodosh-Li's slice-and-dice argument, to which the most effort in [5] is devoted. The idea for the original slice-and-dice argument of Chodosh-Li in the closed case is as follows: Given the closed 3-manifold M_3 with \mathbb{T}^* stabilized positive scalar curvature, one can first use closed minimal surfaces S_k to slice M_3 into a tree-like manifold with simpler second homology group, and then one can use free boundary μ -bubbles D_l to further dice the tree-like manifold into blocks. The positive lower bound of the \mathbb{T}^* -stabilized positive scalar curvature of M_3 can be used to bound the intrinsic diameter of the slicing surfaces S_k and the dicing surfaces D_l , and therefore one can bound the diameter of the blocks. In our case, however, we can only obtain an inradius bound for $S_k \cap N_{\varepsilon}$ and $D_l \cap N_{\varepsilon}$ from the positive lower bound of the \mathbb{T}^* -stabilized positive scalar curvature of $M_3 \cap N_{\varepsilon}$ due to the existence of possibly mean-concave boundary. Roughly speaking, the key observation is that we can establish analogies of properties (1) and (2) for surfaces S_k and D_l , which gives an *extrinsic diameter* bound of the surfaces $S_k \cap \tilde{N}_{\varepsilon}$ and $D_l \cap N_{\varepsilon}$, and thereby we can still bound the extrinsic diameter of the blocks. A similar filling argument based on the quantitative filling property gives the desired contradiction again.

1.1.2. The modification. In the following discussion, let M^n , $n \in \{3, 4, 5\}$, be a closed aspherical *n*-manifold and $S \subset M$ a subset consisting of disjoint incompressible embedded closed aspherical submanifolds (possibly with different dimensions). We assume that $(M \setminus S, g)$ is a complete manifold with positive scalar curvature and aim to derive a contradiction.

We follow the same strategy as above. Let $\pi : M \to M$ be the universal covering of M. Take $\tilde{S} = \pi^{-1}(S)$ and $\tilde{g} = \pi^* g$. Since S consists of incompressible aspherical submanifolds in M, its lift \tilde{S} consists of contractible

6

submanifolds in \tilde{M} . Take S_{ε} to be a tubular neighborhood of S in M and denote $\tilde{S}_{\varepsilon} = \pi^{-1}(S_{\varepsilon})$. We consider the manifold $(\tilde{M} \setminus \tilde{S}, \tilde{g})$, and the relative homology groups $H_k(\tilde{M} \setminus \tilde{S}, \tilde{S}_{\varepsilon})$. Since the components of \tilde{S} are contractible, $\tilde{M} \setminus \tilde{S}$ still satisfy the same topological properties as \tilde{Y} . Then we can take a proper line $\tilde{\sigma}$ in $\tilde{M} \setminus \tilde{S}_{\varepsilon}$ and find M_{n-2} the same way. If we can fill in M_{n-2} by some relative (n-1)-chain within some fixed distance then we are done by the same arguments as above.

The difference here is that we don't have the property (2) anymore because now each component of $\partial \tilde{S}_{\varepsilon}$ is no longer compact. So we need to find another way to control the extrinsic diameter of $M_{n-2} \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon})$ such that the filling argument can run smoothly. In dimensions 3 and 4, recall that we have the inradius estimate for $M_{n-2} \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon})$. When n = 3, the submanifold M_{n-2} consists of curves and the inradius estimate gives the diameter estimate for each component of $M_{n-2} \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon})$. Then we can fill $M_{n-2} \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon})$ with relative chains of bounded size, component by component. When n = 4, we take the same idea and try to bound the extrinsic diameter of each component of $M_{n-2} \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon})$. Since the \mathbb{T}^* -stabilized scalar curvature on $M_{n-2} \cap \pi^{-1}(K)$ has a positive lower bound for any compact subset Kof $M \setminus S$, the worrying case is that the surface M_{n-2} stretches to infinity. Fortunately, since M_{n-2} is a surface, this issue can be handled with the help of Proposition 1.7, which bounds the extrinsic diameter of every interior region of a complete surface with positive \mathbb{T}^* -stabilized scalar curvature.

However, this type of estimate no longer holds for 3-manifolds. Moreover, if we run the slice-and-dice argument in dimension 5, for the slicing surfaces S_k and dicing surfaces D_l , we can only bound the extrinsic diameter of each component of $S_k \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon})$ and $D_l \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon})$, which is not enough to obtain the extrinsic diameter estimate for blocks. As a result, we can only partially answer Question 1.4 in dimension 5 either when S contains a submanifold with codimension 1 or 2, or when S itself is a connected submanifold with codimension ≥ 3 .

In order to obtain the partial result — Theorem 1.6, we make the following two observations. First, if S contains one submanifold of codimension 1 or 2, then this essentially implies $M \setminus S$ has a closed incompressible aspherical hypersurface. The presence of such a hypersurface already forbids any metric with positive scalar curvature on $M \setminus S$ from [3, Theorem 1.1].

Second, if S is a connected submanifold with codimension at least 3, we can pass to a cover \tilde{M} such that $\pi_1(\tilde{M}) = \pi_1(S)$ and repeat the strategy from [5] on $\tilde{M} \setminus \tilde{S}$. The strategy is still valid from the facts that each component of $\partial \tilde{S}_{\varepsilon}$ is compact in $\tilde{M} \setminus \tilde{S}$, and that $\tilde{M} \setminus \tilde{S}$ still satisfies the homological property since the codimension of S is high enough. Notice in this case, if S contains more than one submanifolds $\{S_j\}$, then it could happen that these $\pi_1(S_j)$ together generate the whole $\pi_1(M)$, and it is unclear what cover we can pass to.

The organization of this paper is as follows. In Section 2, we prove the key extrinsic diameter estimate, Proposition 1.7. In Section 3, we prove our main result in dimension 3 and 4, Theorem 1.5. In Section 4, we prove our result in dimension 5, Theorem 1.6.

Acknowledgments. We thank Professors Otis Chodosh, Antoine Song, Ailana Fraser, and Rudolf Zeidler for their interest and helpful suggestions. Part of this research was performed while the first-named author was in residence at the Simons Laufer Mathematical Sciences Institute (formerly MSRI), which is supported by the National Science Foundation (Grant No. DMS-1928930). The second-named author was partially supported by National Key R&D Program of China 2024YFA1014800 and 2023YFA1009900, and NSFC grants 12471052 and 12271008. The third-named author was partially supported by National Key R&D Program of China 2023YFA1009900 and NSFC grant 12401072 as well as the start-up fund from Westlake University.

2. Extrinsic diameter estimate

We begin with some useful definitions and results.

Definition 2.1. Let (M, g) be a Riemannian manifold possibly with nonempty boundary and R a smooth function on M. We say that (M, g) has \mathbb{T}^l -stabilized scalar curvature R if there are l positive smooth functions v_1, \ldots, v_l on M such that the scalar curvature \tilde{R} of $(M \times \mathbb{T}^l, g + \sum_i v_i^2 d\theta_i^2)$ is the \mathbb{T}^l -invariant extension of R. We say that (M, g) has \mathbb{T}^* -stabilized scalar curvature R if (M, g) has \mathbb{T}^l -stabilized scalar curvature R for some positive integer l.

Definition 2.2. We say that (M, ∂_{\pm}, g) is a Riemannian band if (M, g) is a compact Riemannian manifold with boundary and ∂_+ and ∂_- are two disjoint non-empty smooth compact portions of ∂M such that $\overline{\partial M} - (\partial_+ \cup \partial_-)$ is also a compact smooth portion of ∂M .

Lemma 2.3 (Lemma 2.12 of [20]). Given any positive constant c_0 , there is a positive constant $D_0 = D_0(c_0)$ such that the following holds. Let (M^n, ∂_{\pm}, g) be a Riemannian band with $n \leq 7$ and $\overline{\partial M} - (\partial_{+} \cup \partial_{-}) = \emptyset$. If

$$\rho: (M, \partial_{\pm}) \to ([-D, D], \pm D)$$

is a smooth function satisfying

- Lip $\rho < 1$;
- (M,g) has \mathbb{T}^* -stabilized scalar curvature R such that $R \ge c_0$ in $\rho^{-1}([-1,1])$ and $R \ge 0$ in M;
- Any closed hypersurface Σ in M separating ∂_− from ∂₊ does not admit metric with positive T^{*}-stabilized scalar curvature,

then $D \leq D_0$.

Lemma 2.4 (Lemma 2.14 of [5]). Let (M^n, ∂_{\pm}, g) be a Riemannian band with $2 \leq n \leq 7$ and $\operatorname{dist}(\partial_+, \partial_-) > 2d_0$. If $\Gamma := \overline{\partial M - (\partial_+ \cup \partial_-)}$ does not intersect ∂_{\pm} and (M, g) has \mathbb{T}^l -stabilized scalar curvature R, then there exists a hypersurface Σ (possibly with non-empty boundary $\partial \Sigma$) intersecting Γ orthogonally such that

- Σ and ∂_{-} bound a relative region Ω relative to Γ , i.e. $\partial \Omega (\Sigma \cup \partial_{-}) \subset \Gamma$;
- (Σ, g) has \mathbb{T}^{l+1} -stabilized scalar curvature R' where $R' \geq R \pi^2/d_0^2$.

We can then give a proof of Proposition 1.7.

Proof of Proposition 1.7. It suffices to determine the value of $K(s_0)$ for s_0 . Set

$$c_0 = \min_{[0,s_0+2]} L, \quad s_1 = s_0 + D_0(c_0) + D_0(3c_0/4) + 2,$$

$$c_1 = \min_{[0,s_1+3]} L, \quad K(s_0) = s_1 + D_0(c_1) + 3,$$

where $D_0(c_0)$, $D_0(3c_0/4)$ and $D_0(c_1)$ are the constants defined in Lemma 2.3. Let s' be a regular value of ρ in $(K(s_0) - 1, K(s_0))$. We deal with interior topology and interior geometry separately.

I. Interior topology. The argument is broken into three steps.

Step 1. Let s_1^* be a regular value of ρ in $(s_1, s_1 + 1)$. Then each component $\Sigma_{s_1^*}^o$ of $\Sigma_{s_1^*}$ is homeomorphic to a 2-sphere with finitely many disks removed.

Otherwise, some component $\Sigma_{s_1^*}^o$ must be $m\mathbb{T}^2$ or $m\mathbb{RP}^2$ after filling with disks. In particular, $\Sigma_{s_1^*}^o$ has the form of $\mathbb{T}^2 \# S$ or $\mathbb{RP}^2 \# S$ for some surface S, then we can find $\hat{\Sigma}_{s_1^*}^o \subset \Sigma_{s_1^*}^o$ homeomorphic to $\mathbb{T}^2 - B$ or $\mathbb{RP}^2 - B$. Let $\Sigma_{s'}^o$ denote the component of $\Sigma_{s'}$ containing $\Sigma_{s_1^*}^o$, and let S_+ and S_- denote copies of $\Sigma_{s'}^o \setminus \hat{\Sigma}_{s_1^*}^o$. Consider a double cover $\tilde{\Sigma}_{s_1^*}^o$ of $\hat{\Sigma}_{s_1^*}^o$ and clearly $\tilde{\Sigma}_{s_1^*}^o$ has the form of $\mathbb{T}^2 - (B_+ \sqcup B_-)$ or $\mathbb{S}^2 - (B_+ \sqcup B_-)$, where B_+ and B_- are lifts of B. Correspondingly, the surface $\Sigma_{s'}^o$ is lifted to some double cover

$$\tilde{\Sigma}_{s'}^o = S_+ \cup \tilde{\Sigma}_{s_1^*}^o \cup S_-.$$

Denote $\pi: \tilde{\Sigma}^o_{s'} \to \Sigma^o_{s'}$ to be the covering map. Then the map

$$\tilde{\rho}(x) = \begin{cases} (\rho \circ \pi - s_1^*)_+, & x \in S_+; \\ -(\rho \circ \pi - s_1^*)_+, & x \in S_-; \\ 0, & \text{otherwise.} \end{cases}$$

is a Lipschitz function on $\tilde{\Sigma}_{s'}^o$ with $\operatorname{Lip} \tilde{\rho} < 1$. Also we see that the function $\tilde{\rho}$ has range $[s_1^* - s', s' - s_1^*]$, and that $\tilde{\Sigma}_{s'}^o$ has \mathbb{T}^* -stabilized scalar curvature $R \geq c_1 \chi_{[-2,2]} \circ \tilde{\rho}$. By smooth approximation and Lemma 2.3 we obtain

$$D_0(c_1) + 1 = K(s_0) - 1 - (s_1 + 1) < s' - s_1^* \le D_0(c_1),$$

which leads to a contradiction.

Step 2. Let s_0^* be a regular value of ρ in $(s_0, s_0 + 1)$. Then each circle γ in $\rho^{-1}(s_0^*)$ bounds a disk in $\Sigma_{s_1^*}$.

Let $\Sigma_{s_1^*}^o$ be the component of $\Sigma_{s_1^*}$ containing γ . Suppose that γ does not bound a disk in $\Sigma_{s_1^*}$. Recall that $\Sigma_{s_1^*}^o$ is homeomorphic to a 2-sphere with finitely disks removed. From this we conclude that γ separates the boundary components of $\partial \Sigma_{s_1^*}^o$ into two non-empty collections. Denote

$$\Sigma_{s_1^*}^o-\gamma=\Omega_+\sqcup\Omega_-$$

Define

$$\rho_{s_1^*}(x) = \begin{cases} \min\{\operatorname{dist}(x,\gamma), s_1^* - s_0^*\}, & x \in \Omega_+; \\ -\min\{\operatorname{dist}(x,\gamma), s_1^* - s_0^*\}, & x \in \Omega_-; \\ 0, & \text{otherwise.} \end{cases}$$

Then $\rho_{s_1^*}$ is a Lipschitz function on $\Sigma_{s_1^*}^o$ with Lip $\rho_{s_1^*} \leq 1$. Also $\rho_{s_1^*}$ has range $[s_0^* - s_1^*, s_1^* - s_0^*]$, and that $\Sigma_{s_1^*}^o$ has T*-stabilized scalar curvature

$$R \ge c_0 \chi_{[-s_0 - 2 + s_0^*, s_0 + 2 - s_0^*]} \circ \rho_{s_1^*}$$

By smooth approximation and Lemma 2.3 we obtain

$$D_0(c_0) + D_0(3c_0/4) + 1 = s_1 - s_0 - 1 < s_1^* - s_0^* \le D_0(c_0),$$

which leads to a contradiction.

Step 3. Σ_{s_0} is contained in $\sqcup_j D_j$, where $\{D_j\}$ are finitely many disjoint disks in $\Sigma_{s_1^*}$.

By passing to each component we can assume $\Sigma_{s_1^*}$ to be connected. Due to the fact $\Sigma_{s_0} \subset \Sigma_{s_0^*}$ it suffices to show that $\Sigma_{s_0^*}$ is contained in $\sqcup_j D_j$, where $\{D_j\}$ are finitely many disjoint disks in $\Sigma_{s_1^*}$. From our assumption we know that Σ is connected and so $\Sigma_{s'}$ has non-empty boundary. In particular, $\Sigma_{s_1^*}$ has non-empty boundary as well. Denote $\rho^{-1}(s_0^*) = \sqcup_k \gamma_k$. From the previous steps we conclude that each γ_k bounds a unique disk D_k in $\Sigma_{s_1^*}$. We alaim that for $k \neq l$ one of the following happens:

We claim that for $k \neq l$ one of the following happens:

- D_k and D_l are disjoint;
- $D_k \subset D_l$ or $D_l \subset D_k$.

Otherwise, we can find a point $p_{kl} \in D_k \cap D_l$ and a pair of points $p_k \in D_k \setminus D_l$ and $p_l \in D_l \setminus D_k$. Connecting p_{kl} and p_k with path in D_k we see $\gamma_l \cap D_k \neq \emptyset$. Since γ_k and γ_l are disjoint, we conclude $\gamma_l \subset D_k$ and similarly $\gamma_k \subset D_l$. In particular, the union $D_k \cup D_l$ is a closed surface in $\Sigma_{s_1^*}$, which is impossible since $\Sigma_{s_1^*}$ is connected with non-empty boundary.

Now we can take the collection of maximal disks $\{D_{j,\max}\}$ with respect to the inclusion. From previous claim we conclude that $D_{j,\max}$ are pairwise disjoint.

It remains to show $\Sigma_{s_0^*} \subset \sqcup D_{j,\max}$. Take any component $\Sigma_{s_0^*}^o$ of $\Sigma_{s_0^*}$. Since $\Sigma_{s_0^*}^o$ does not cross $\sqcup_k \gamma_k$, either $\Sigma_{s_0^*}^o \subset \sqcup D_{j,\max}$ or $\Sigma_{s_0^*}^o$ is disjoint with all disks D_k . Denote $\partial \Sigma_{s_0^*}^o = \sqcup_l \gamma_{k_l}$. Then the union

$$\Sigma_{s_0^*}^o \cup \left(\bigsqcup_l D_{k_l}\right)$$

is a closed surface in $\Sigma_{s_1^*}$, which leads to the same contradiction as before.

II. Interior geometry. To establish the extrinsic diameter estimate, we argue by contradiction. Set

$$c_2 = \min_{[0,K(s_0)]} L.$$

If

diam
$$(\Sigma_{s_0}^o \subset (\Sigma_{s_1^*}, g)) > 4\pi \left(\min_{[0, K(s_0)]} L\right)^{-\frac{1}{2}} = \frac{4\pi}{\sqrt{c_2}}$$

then there exist two points $p_+, p_- \in \Sigma_{s_0}^o$ such that $\operatorname{dist}(p_+, p_-) > 4\pi/\sqrt{c_2}$. Recall that s' is a regular value of ρ in $(K(s_0) - 1, K(s_0))$. For sufficiently small $\varepsilon > 0$, set

 $M = \Sigma_{s'} \setminus (B^g_{\varepsilon}(p_-) \cup B^g_{\varepsilon}(p_+)), \quad \partial_- = \partial B^g_{\varepsilon}(p_-), \quad \partial_+ = \partial B^g_{\varepsilon}(p_+), \quad \Gamma = \partial \Sigma_{s'},$ and then (M, ∂_{\pm}, g) is a Riemannian band with dist $(\partial_+, \partial_-) > 4\pi/\sqrt{c_2}$ and $\Gamma \cap (\partial_+ \cup \partial_-) = \emptyset$. Recall that (Σ, g) has \mathbb{T}^l -stabilized scalar curvature $R \geq L \circ \rho$ in $\Sigma_{K(s_0)}$ and so in $\Sigma_{s'}$. By Lemma 2.4 (with the choice $d_0 = 2\pi/\sqrt{c_2}$), there exists a curve γ satisfying

- (a) γ and ∂_{-} bound a relative region Ω relative to Γ ;
- (b) γ has the positive \mathbb{T}^{l+1} -stabilized scalar curvature

$$R' \ge R - \frac{c_2}{4} \ge L \circ \rho - \frac{c_2}{4} \ge c_2 - \frac{c_2}{4} = \frac{3}{4}c_2 > 0.$$

We split the argument into two cases:

Case 1. $\gamma \cap \rho^{-1}(s') = \emptyset$. In this case, $\partial \gamma = \emptyset$ and each component of γ is \mathbb{S}^1 . Then (b) shows that \mathbb{S}^1 has positive \mathbb{T}^{l+1} -stabilized scalar curvature. This contradicts the fact that \mathbb{T}^n cannot admit any smooth metric with positive scalar curvature.

Case 2. $\gamma \cap \rho^{-1}(s') \neq \emptyset$. Let σ be a path in $\Sigma_{s_0}^o$ such that

- σ connects p_- and p_+ ;
- σ has non-zero algebraic intersection with ∂_{-} .

Using (a), γ is relative homologous to ∂_{-} relative to Γ and then γ has nonzero algebraic intersection with σ . In particular, we obtain $\gamma \cap \sigma \neq \emptyset$ and so $\gamma \cap \Sigma_{s_0} \neq \emptyset$. Fix a point $x_0 \in \gamma \cap \Sigma_{s_0}$ and the component of γ containing x_0 . For convenience, we still denote this component by γ . Choose an arc length parametrization of γ such that

$$\gamma: [-D', D''] \to M, \ \gamma(0) = x_0, \ \gamma(-D'), \ \gamma(D'') \in \rho^{-1}(s'),$$

where D' and D'' are two positive constants. We next assume that $D' \ge D''$, as the case D'' > D' can be proved similarly. Since Lip $\rho < 1$, then

$$|\rho(\gamma(t)) - \rho(\gamma(0))| < |t| \text{ for } t \in [-D', D''].$$
(2.1)

Choosing t = D'' and recalling the facts $\gamma(D'') \in \rho^{-1}(s')$ and $\rho(\gamma(0)) \leq s_0$, we can derive from (2.1) that

$$D'' > s' - s_0 > K(s_0) - 2 - s_0 = D_0(c_0) + D_0(3c_0/4) + D_0(c_1) + 3.$$
(2.2)

Using (2.1) again, we obtain

$$\rho(\gamma(t)) < s_0 + 1 \text{ for } t \in [-1, 1],$$

which implies $\gamma([-1,1]) \subset \Sigma_{s_0+1}$. Together with (b) and $c_2 \leq c_0$, we know that γ has positive \mathbb{T}^{l+1} -stabilized scalar curvature R' with

$$R' \ge R - \frac{c_2}{4} \ge L \circ \rho - \frac{c_2}{4} \ge c_0 - \frac{c_2}{4} \ge \frac{3c_0}{4} > 0 \text{ in } \gamma([-1,1]).$$

Applying Lemma 2.3 to $\gamma([-D'', D''])$ and $\rho|_{\gamma([-D'', D''])}$, we obtain $D'' \leq D_0(3c_0/4)$, which contradicts with (2.2).

3. Proof of Theorem 1.5

In the rest of this paper, we define the orientable cover of a manifold N to be

- N when N is orientable;
- the orientation double cover of N when N is non-orientable.

Proof of Theorem 1.5. By passing to the orientable cover of M, we can always assume that M is orientable. We just need to deal with the case when n = 4 since we can make Riemannian product with \mathbb{S}^1 in dimension three. First we take an embedded closed curve σ disjoint from S such that $[\sigma] \neq 0 \in \pi_1(M)$, which can be done as follows.

Case 1. S contains some hypersurface S_0 . Then we can know $\pi_1(S_0) \neq 0$ from its aspherical property. Take an embedded circle σ_0 in S_0 with $[\sigma_0] \neq 0 \in \pi_1(S_0)$. If $\sigma_0^*(NS_0)$ is a trivial bundle, we pick a nowhere-zero normal vector field along σ_0 and push σ_0 away from S_0 to obtain an embedded closed curve σ . Otherwise, we do the same thing for the double cover of σ_0 to obtain an embedded closed curve σ . From the construction we know that σ is homotopic to σ_0 or σ_0^2 . Since M is aspherical, $\pi_1(M)$ has no torsion. Since the map $\pi_1(S_0) \to \pi_1(M)$ is injective, we see $[\sigma] \neq 0 \in \pi_1(M)$.

Case 2. S contains no hypersurface. Then we just take an embedded closed curve σ in M with $[\sigma] \neq 0 \in \pi_1(M)$. Since S has codimension greater than one, we can perturb σ such that σ is disjoint from S.

In the following, we are going to deduce a contradiction assuming that there is a complete metric g on $M \setminus S$ with positive scalar curvature. We begin with the basic set-up. Let $\pi : \tilde{M} \to M$ denote the universal covering of M. Take $\tilde{S} = \pi^{-1}(S)$ and $\tilde{g} = \pi^* g$. Since S consists of incompressible aspherical submanifolds in M, its lift \tilde{S} consists of contractible submanifolds in \tilde{M} . On the other hand, we take a line $\tilde{\sigma}$ in \tilde{M} which is a lift of σ .

Take S_{ε} to be a tubular neighborhood of S in M disjoint from σ . Denote $\tilde{S}_{\varepsilon} = \pi^{-1}(S_{\varepsilon})$. Since \tilde{S} is contractible, we have $H_i(\tilde{S}_{\varepsilon}) = 0$ for all $i \geq 1$. Recall that \tilde{M} is contractible as well. The composed map

$$H_1(\tilde{M} \setminus \tilde{S}) \to H_1(\tilde{M}) \to H_1(\tilde{M}, \tilde{S}_{\varepsilon})$$

is the zero map. Moreover, from the exact sequence

$$H_i(M) \to H_i(M, S_{\varepsilon}) \to H_{i-1}(S_{\varepsilon})$$

we conclude $H_i(\tilde{M}, \tilde{S}_{\varepsilon}) = 0$ for all $i \geq 2$.

Let $\rho: M \setminus S \to [0, +\infty)$ denote a proper smooth function such that $\rho \equiv 0$ in $M \setminus S_{\varepsilon}$ and Lip $\rho < 1$. We define

$$L(s) = \frac{1}{2} \min_{\{\rho \le s\}} R(g).$$

Fix $s_0 = 1$ and let $K(s_0)$ be the constant from Proposition 1.7.

Similar as in [5] for any large constant \tilde{L} we can construct a piecewisely smooth hypersurface M_3 in $\tilde{M} \setminus \tilde{S}$ with non-empty boundary such that

- M_3 has non-zero intersection number with $\tilde{\sigma}$;
- dist $(\partial M_3, \tilde{\sigma}) \geq \tilde{L};$
- ∂M_3 has \mathbb{T}^* -stabilized scalar curvature $R_3 \geq R(\tilde{g})|_{\partial M_3} c\tilde{L}^{-2}$, where c is an absolute positive constant independent of \tilde{L} .

Take L large enough such that

$$c\tilde{L}^{-2} < \min_{[0,K(s_0)]} L = L(K(s_0)).$$

Let Σ be any component of ∂M_3 . It is clear that

(

$$\Sigma = \Sigma \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon})$$
 modulo \tilde{S}_{ε} .

Next we show how to establish the diameter estimate for each component of $\Sigma \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon})$. There are three cases:

Case 1. $\Sigma \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon}) = \emptyset$. This case is trivial.

Case 2. $\Sigma \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon}) \neq \emptyset$ and $\Sigma \subset \{\rho \circ \pi \leq K(s_0)\}$. Then we know that Σ has \mathbb{T}^* -stabilized scalar curvature

$$R_3 \ge R(\tilde{g})|_{\partial M_3} - c\tilde{L}^{-2} \ge L(K(s_0)).$$

and so it follows from [10] that its diameter is no greater than

$$D_1 := 2\pi L(K(s_0))^{-\frac{1}{2}}$$

Case 3. $\Sigma \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon}) \neq \emptyset$ and $\Sigma \cap \{\rho \circ \pi > K(s_0)\} \neq \emptyset$. We can apply Proposition 1.7 to Σ associated with the map $(\rho \circ \pi)|_{\Sigma}$ to obtain that each component of $\Sigma \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon})$ has diameter no greater than

$$D_2 := 4\pi L(K(s_0))^{-\frac{1}{2}}$$

From $H_2(M, S_{\varepsilon}) = 0$ and the above discussion, we conclude that each component of $\Sigma \cap (\tilde{M} \setminus \tilde{S}_{\varepsilon})$ is a relative boundary and has diameter no greater than max $\{D_1, D_2\}$. Using the quantitative filling lemma [5, Lemma 2.8] we conclude that there is a 3-chain Γ lying in D_3 -neighborhood of ∂M_3 such that

$$\partial \Gamma = \Sigma \mod \tilde{S}_{\varepsilon}$$
.

where D_3 is an absolute constant depending only on D_1 , D_2 and (M,g)but independent of \tilde{L} . When $\tilde{L} > D_3$ the relative cycle $M_3 + \Gamma$ has nonzero intersection number with $\tilde{\sigma}$ and so it contributes to some non-trivial element in $H_3(\tilde{M}, \tilde{S}_{\varepsilon})$, which contradicts the fact $H_3(\tilde{M}, \tilde{S}_{\varepsilon}) = 0$. This contradiction shows that $M \setminus S$ cannot admit a complete metric with positive scalar curvature.

Lastly, we prove the rigidity part. Suppose that $M \setminus S$ admits a complete metric g of nonnegative scalar curvature. Again by passing to the orientable cover we can assume that M is orientable. By a result of Kazdan [12], if $(M \setminus S, g)$ is not Ricci-flat, then there exists a complete metric with positive scalar curvature. Thus $(M \setminus S, g)$ is Ricci-flat.

If S contains any component S_0^k of dimension $k \leq n-2$ that is not a hypersurface, then we claim that S contains no hypersurface. Otherwise, we can find a component U_0 of $M \setminus S$ such that U_0 has at least two ends, for which one is compactified by S_0 in M and the other is compactified by some hypersurface S_1 in M. Then it follows from the Cheeger–Gromoll splitting theorem [2, Theorem 4] that U_0 is isometric to some Riemannian product $N \times \mathbb{R}$, where N is a closed Riemannian manifold. Denote the orientable cover of S_1 by \check{S}_1 and the unit tangent sphere bundle of S_0 in M by T_1S_0 . Realize T_1S_0 to be the boundary of a tubular neighborhood of S_0 in M. Then \check{S}_1 and T_1S_0 are homotopy equivalent since they are both homotopy equivalent to N. However, by considering the long exact sequence of homotopy groups for fibration, we have

$$\pi_{n-k+1}(S_0) \to \pi_{n-k}(\mathbb{S}^{n-k}) \to \pi_{n-k}(T_1S_0).$$

Since $n - k + 1 \ge 3$, and S_0 is aspherical, we have $\pi_{n-k+1}(S_0) = 0$, so $\mathbb{Z} \cong \pi_{n-k}(\mathbb{S}^{n-k}) \to \pi_{n-k}(T_1S_0)$ is injective. But this shows $0 \ne \pi_{n-k}(T_1S_0) = \pi_{n-k}(\check{S}_1)$, which contradicts that \check{S}_1 is aspherical.

By the homotopy classification of aspherical spaces [15, Theorem 1.1], since S_0 is not homotopy equivalent to any finite cover of M, $\pi_1(S_0)$ must be a proper subgroup of $\pi_1(M)$ with infinite index. We can consider the covering $\hat{\pi} : \hat{M} \to M$ such that $\pi_1(\hat{M}) = \pi_1(S_0)$. Let $\hat{S} = \hat{\pi}^{-1}(S)$. Then we can lift the Ricci-flat metric g to $\hat{M} \setminus \hat{S}$ as \hat{g} , and $\pi^{-1}(S_0)$ consists of infinitely many components diffeomorphic to S_0 , and each of them gives one end in $(\hat{M} \setminus \hat{S}, \hat{g})$. Recall that S contains no hypersurface and so does \hat{S} , which means that $\hat{M} \setminus \hat{S}$ is connected. However, by the Cheeger–Gromoll splitting theorem, a complete connected Ricci-flat manifold has at most two ends, which leads to a contradiction.

We have shown that S consists of hypersurfaces only, whose collection will be denoted by $\{S_j\}$. For each connected component U of $M \setminus S$, we know that it can have at most two ends. Fix an arbitrary metric on M and denote the metric completion of U by \overline{U} .

If U has two ends, then U is isometric to a cylinder $N \times \mathbb{R}$ for some closed Ricci-flat manifold N by the Cheeger–Gromoll splitting theorem, and each component of the boundary $\partial \overline{U}$ is the orientable cover of some S_j .

If U has only one end, we claim that we can find a two-sheeted cover \overline{U}^* of \overline{U} such that the interior of \overline{U}^* is a cylinder $N \times \mathbb{R}$ for a closed manifold and the boundary $\partial \overline{U}^*$ consists two copies of the orientable cover of some S_j . To see this, we start with a careful analysis of the topology of \overline{U} . Since U has only one end, the boundary $\partial \overline{U}$ is connected and it is the orientable cover \check{S}_j of some S_j . Recall that S_j is incompressible in M, then we can conclude that $\partial \overline{U}$ is incompressible in \overline{U} , which yields that the map $\pi_1(\partial \overline{U}) \to \pi_1(\overline{U})$ is injective. Let us show that

$$\pi_1(\partial \overline{U}) \neq \pi_1(\overline{U}).$$

Otherwise, $i_* : \pi_1(\partial \overline{U}) \to \pi_1(\overline{U})$ is an isomorphism. Recall that $\partial \overline{U} = \check{S}_j$ is an Eilenberg–MacLane space, then we can find a continuous map

$$f:\overline{U}\to\partial\overline{U}$$

such that $f_*: \pi_1(\overline{U}) \to \pi_1(\partial \overline{U})$ is the inverse map of $i_*: \pi_1(\partial \overline{U}) \to \pi_1(\overline{U})$. Since $f \circ i$ induces an identity map on $\pi_1(\partial \overline{U})$, we have the homology relation

$$f_*i_*([\partial \overline{U}]) = [\partial \overline{U}].$$

However, this is impossible since $\partial \overline{U}$ is null-homologous in \overline{U} . Therefore, We have shown that $\pi_1(\partial \overline{U})$ is a proper subgroup of $\pi_1(\overline{U})$. In particular, we can find a cover \overline{U}^* of \overline{U} with $\pi_1(\overline{U}^*) = \pi_1(\partial \overline{U})$ such that $\partial \overline{U}^*$ consists of multiple copies of $\partial \overline{U}$. Correspondingly, the interior of \overline{U}^* has multiple ends, and it admits a complete Ricci-flat metric as a cover of U. Then the Cheeger–Gromoll splitting theorem yields that the interior of \overline{U}^* is isometric to a cylinder $N \times \mathbb{R}$ for some closed Ricci-flat manifold N and so $\partial \overline{U}^*$ consists of two copies of \check{S}_j .

Thus by lifting to a finite cover of the pair (M, S), we can assume that each connected component U of $M \setminus S$ is isometric to a cylinder $N \times \mathbb{R}$ for some closed Ricci-flat manifold N and that all S_j are orientable. Let us label the components of $M \setminus S$ by U_k and denote $U_k = N_k \times \mathbb{R}$. From the cylindrical structure of U_k we see that each N_k admits a non-zero degree map to some closed aspherical manifold S_j . It follows from [5, Proposition 4.4] that N_k is a closed flat manifold and so the metric g is flat. From the cylindrical structure of U_k we also have the isomorphisms

$$\pi_1(S_j) = \pi_1(U_k) = \pi_1(N_k)$$

through homotopies in M. By lifting further to a finite cover of the pair (M, S), we can assume $\pi_1(N_k) = \mathbb{Z}^{n-1}$ for all k. From the second Bieberbach

rigidity theorem we know that each N_k is diffeomorphic to \mathbb{T}^{n-1} , and so all U_k are cylinders $\mathbb{T}^{n-1} \times \mathbb{R}$. We complete the proof.

4. Proof of Theorem 1.6

Proof of Theorem 1.6. Let n be the dimension of M. Here n = 5 but the same proof also works if n equals 3 or 4. We note that the rigidity part is exactly the same as in the proof of Theorem 1.5. Thus we only need to show that $M \setminus S$ cannot admit any complete metric with positive scalar curvature. According to our assumption, there are three cases.

Case 1. S contains a hypersurface Σ^{n-1} of codimension 1. By passing to the orientable cover of M, we can assume that M is orientable. Denote the orientable cover of Σ by $\check{\Sigma}$ and we can realize it as a boundary component of some small tubular neighborhood of Σ in M. Clearly, $\check{\Sigma}$ is orientable and incompressible in M and so in $M \setminus S$. Since $\check{\Sigma}$ is aspherical, by [3, Theorem 1.1] $M \setminus S$ cannot admit any complete metric with positive scalar curvature.

Case 2. S contains a submanifold Σ^{n-2} of codimension 2. Again we can assume that M is orientable. We make a discussion depending on whether Σ is orientable or not.

Suppose that Σ is orientable. Take Σ_{ε} to be a small tubular neighborhood of Σ in M. Then $\partial \Sigma_{\varepsilon}$ is a circle bundle over Σ . By considering the long exact sequence of homotopy groups for fibration, we see that $\partial \Sigma_{\varepsilon}$ is again an aspherical manifold, and $\pi_1(\partial \Sigma_{\varepsilon})$ is generated by curves in $\pi_1(\partial D)$ and $\pi_1(\Sigma)$, where D is the disk fiber of Σ_{ε} over a point $p \in \Sigma$. Our goal is to show that $\pi_1(\partial \Sigma_{\varepsilon}) \to \pi_1(M \setminus S)$ is injective. Then it follows from [3, Theorem 1.1] that $M \setminus S$ cannot admit any complete metric with positive scalar curvature.

Orient D such that the intersection number of D and Σ in M is 1. Let us consider the map $\pi_1(\partial D) \to \pi_1(M \setminus S)$ induced by the inclusion. We claim that the map $\pi_1(\partial D) \to \pi_1(M \setminus S)$ is injective. Otherwise, an element $k[\partial D]$ for some $k \in \mathbb{Z}$ is mapped to zero in $\pi_1(M \setminus S)$. In other words, it bounds a disk D' in $M \setminus S$. On the other hand, $k[\partial D]$ bounds a disk D'' in Σ_{ε} having intersection number k with Σ . Then $D' \cup D''$ is a sphere in Mwhich has intersection number k with Σ . However, since M is aspherical, we have $\pi_2(M) = 0$. As a consequence, the sphere $D' \cup D''$ must be trivial in $H_2(M)$ and so k = 0. Thus the map $\pi_1(\partial D) \to \pi_1(M \setminus S)$ is injective.

Now we consider the inclusion maps

$$i_1: \partial \Sigma_{\varepsilon} \to M \setminus S \text{ and } i_2: M \setminus S \to M.$$

To see the injectivity of the map $\pi_1(\partial \Sigma_{\varepsilon}) \to \pi_1(M \setminus S)$, we take an element γ in $\pi_1(\partial \Sigma_{\varepsilon})$ such that $(i_1)_*\gamma = 0$ in $\pi_1(M \setminus S)$, and try to show $\gamma = 0$ in $\pi_1(\partial \Sigma_{\varepsilon})$. It is clear that we have $(i_2 \circ i_1)_*\gamma = 0$ in $\pi_1(M)$. Since Σ is incompressible, we conclude that $\pi_*\gamma = 0$ in $\pi_1(\Sigma)$, where $\pi : \partial \Sigma_{\varepsilon} \to \Sigma$ is the projection map of the circle bundle. From the long exact sequence of

homotopy groups for fibration, the sequence

$$\pi_1(\partial D) \to \pi_1(\partial \Sigma_{\varepsilon}) \to \pi_1(\Sigma)$$

is exact. As a result, we can find an element ξ in $\pi_1(\partial D)$ such that $i_*\xi = \gamma$ in $\pi_1(\partial \Sigma_{\varepsilon})$, where $i : \partial D \to \partial \Sigma_{\varepsilon}$ is the inclusion map of the circle bundle. In particular, we have $(i_1)_*i_*\xi = 0$ in $\pi_1(M \setminus S)$. Then we can conclude from the injectivity of the map $\pi_1(\partial D) \to \pi_1(M \setminus S)$ that $\xi = 0$ in $\pi_1(\partial D)$ and so $\gamma = 0$ in $\pi_1(\partial \Sigma_{\varepsilon})$.

Suppose that Σ is not orientable. Denote the orientable cover of Σ by $\check{\Sigma}$. Then we can find a cover (\check{M}, \check{S}) of the pair (M, S) such that $\pi_1(\check{M}) = \pi_1(\check{\Sigma})$. In particular, the subset \check{S} contains $\check{\Sigma}$ as a component. Similarly, we can show $\pi_1(\partial \check{\Sigma}_{\varepsilon}) \to \pi_1(\check{M} \setminus \check{S})$ is injective. This implies that $\check{M} \setminus \check{S}$ admits no complete metric with positive scalar curvature and so does $M \setminus S$.

Case 3. S is a connected submanifold of codimension ≥ 3 . Set

$$d = \dim S \le n - 3.$$

By the homotopy classification of aspherical spaces, since S is not homotopy equivalent to any finite cover of M, $\pi_1(S)$ must be a proper subgroup of $\pi_1(M)$ with infinite index.

Denote the orientable cover of S by \check{S} and take the cover (\tilde{M}, \tilde{S}) of the pair (M, S) such that $\pi_1(\tilde{M}) = \pi_1(\check{S})$. It is easy to see that \tilde{S} consists of disjoint copies of \check{S} , and so each component of \tilde{S} is homotopy equivalent to \tilde{M} through inclusion. By passing to the orientable cover of \tilde{M} , we can further assume that \tilde{M} is orientable. Take S_{ε} to be a small tubular neighborhood of S in M and denote $\tilde{S}_{\varepsilon} = \pi^{-1}(S_{\varepsilon})$, where $\pi : \tilde{M} \to M$ is the covering map.

We have the following topological lemma.

Lemma 4.1. We have

$$H_k(\tilde{M} \setminus \tilde{S}, \tilde{S}_{\varepsilon}) = 0$$

for all integers $k \ge d+2$. Moreover, the map

$$H_k(\tilde{M} \setminus \tilde{S}) \to H_k(\tilde{M} \setminus \tilde{S}, \tilde{S}_{\varepsilon})$$

is the zero map for all integers $k \ge 0$.

Proof. Let $i : \tilde{M} \setminus \tilde{S} \to \tilde{M}$ denote the inclusion map. From excision we conclude that the map

$$i_*: H_k(M \setminus S, S_{\varepsilon}) \to H_k(M, S_{\varepsilon})$$

is an isomorphism. From the exact sequence of the pair $(\tilde{M}, \tilde{S}_{\varepsilon})$ we have the long exact sequence

$$\cdots \to H_k(\tilde{S}_{\varepsilon}) \to H_k(\tilde{M}) \to H_k(\tilde{M}, \tilde{S}_{\varepsilon}) \to H_{k-1}(\tilde{S}_{\varepsilon}) \to \cdots$$

Notice that for $k \ge d+1$ we have $H_k(\tilde{S}_{\varepsilon}) = H_k(\tilde{M}) = 0$ since each component of \tilde{S} is homotopy equivalent to \tilde{M} . This gives

$$H_k(\tilde{M}\setminus\tilde{S},\tilde{S}_\varepsilon)=0$$

for all integers $k \ge d+2$.

Again by the homotopy equivalence of \tilde{M} and each component of \tilde{S} , we have that the map $H_k(\tilde{S}_{\varepsilon}) \to H_k(\tilde{M})$ is surjective for all k. Exactness of the sequence then implies that the map $H_k(\tilde{M}) \to H_k(\tilde{M}, \tilde{S}_{\varepsilon})$ is the zero map. Thus the composed map

$$H_k(\tilde{M} \setminus \tilde{S}) \to H_k(\tilde{M}) \to H_k(\tilde{M}, \tilde{S}_{\varepsilon}) \xrightarrow{i_*^{-1}} H_k(\tilde{M} \setminus \tilde{S}, \tilde{S}_{\varepsilon})$$

is the zero map.

We are ready to complete the proof of Case 3. Since $\pi_1(S)$ is a proper subgroup of $\pi_1(M)$ with infinite index, there exists a simple closed noncontractible loop σ in M such that $[\sigma]$ is in $\pi_1(M) \setminus \pi_1(S)$ and $\sigma \cap S_{\varepsilon} = \emptyset$.

Suppose that $M \setminus S$ admits a complete metric of positive scalar curvature and so does the cover $\tilde{M} \setminus \tilde{S}$ by lifting. In the following, we are going to run the chain-closing program of [5] to derive a contradiction, with Lemma 4.1 playing the role of [5, Lemma 2.7].

To be more precise, we construct a proper line

$$\tilde{\sigma}: (-\infty, +\infty) \to M \setminus S_{\varepsilon}$$

by lifting σ and then take the boundary $\partial \mathcal{N}$ of a tubular neighborhood \mathcal{N} of $\tilde{\sigma}([0, +\infty))$. Through cutting $\partial \mathcal{N}$ at a finite length we can obtain a hypersurface \tilde{M}_{n-1} with boundary which has non-zero algebraic intersection with $\tilde{\sigma}$. We recall that in this step we need $H_{n-1}(\tilde{M} \setminus \tilde{S}, \tilde{S}_{\varepsilon}) = 0$ (see [5, Lemma 3.1, Step 3]), which is guaranteed by Lemma 4.1 and the fact $d \leq n-3$. Let M_{n-1} be the area-minimizing hypersurface obtained from solving the Plateau problem with the prescribed boundary $\partial \tilde{M}_{n-1}$. Then using the μ -bubble method we can construct a codimension-two closed submanifold $M_{n-2} \subset M_{n-1}$. By adjusting the size of the tubular neighborhood as well as the cutting length we can guarantee

- M_{n-2} and ∂M_{n-1} enclose a bounded region Ω_{n-1} in M_{n-1} which is disjoint from $\tilde{\sigma}$;
- dist $(M_{n-2}, \tilde{\sigma}) > L$ for arbitrarily large L > 0;
- M_{n-2} has \mathbb{T}^* -stabilized scalar curvature no less than $R(g) \mu_{\text{loss}}$ for arbitrarily small $\mu_{\text{loss}} > 0$.

Since \tilde{S} consists of disjoint copies of some finite cover of S, each component of \tilde{S} is compact and so $\partial \tilde{S}_{\varepsilon}$ is a disjoint union of closed manifolds. Then we can find a chain Γ disjoint from $\tilde{\sigma}$ such that $\partial \Gamma = M_{n-2}$ modulo \tilde{S}_{ε} by repeating the proof of [5, Proposition 3.10]. We point out that the proof of [5, Proposition 3.10] uses the property that closed k-manifolds in $\tilde{M} \setminus \tilde{S}$ represent the zero class in $H_k(\tilde{M} \setminus \tilde{S}, \tilde{S}_{\varepsilon})$ for all $k \leq n-2$, which is again guaranteed by Lemma 4.1. Now we consider the chain

$$\tilde{\Gamma}_{n-1} := \tilde{M}_{n-1} + \Omega_{n-1} + \Gamma_{\cdot}$$

18

This is a relative (n-1)-cycle having non-zero algebraic intersection number with the line $\tilde{\sigma}$, contradicting the fact $H_{n-1}(\tilde{M} \setminus \tilde{S}, \tilde{S}_{\varepsilon}) = 0$ from Lemma 4.1.

References

- [1] Agol, I. The virtual Haken conjecture, Doc. Math. 18 (2013), 1045–1087.
- [2] Cheeger, J., Gromoll, D. The splitting theorem for manifolds of nonnegative Ricci curvature, J. Differential Geom. 6 (1971), 119–128.
- [3] Chen, J., Liu, P., Shi, Y., Zhu, J. Incompressible hypersurface, positive scalar curvature and positive mass theorem, preprint, arXiv: 2112.14442.
- [4] Chen, S. A generalization of the Geroch conjecture with arbitrary ends, Math. Ann. 389 (2024), 489–513.
- [5] Chen, S., Chu, J., Zhu, J. Positive scalar curvature metrics and aspherical summands, preprint, arXiv: 2312.04698.
- [6] Chodosh, O., Li, C. Generalized soap bubbles and the topology of manifolds with positive scalar curvature, Ann. of Math. (2) 199 (2024), no. 2, 707–740.
- [7] Chodosh, O., Li, C., Liokumovich, Y. Classifying sufficiently connected PSC manifolds in 4 and 5 dimensions, Geom. Topol. 27 (2023), no. 4, 1635–1655.
- [8] Gromov, M. Large Riemannian manifolds, Curvature and topology of Riemannian manifolds (Katata, 1985), 108–121. Lecture Notes in Math., 1201 Springer-Verlag, Berlin, 1986.
- [9] Gromov, M. Four Lectures on Scalar Curvature, preprint, arXiv: 1908.10612.
- [10] Gromov, M. No metrics with positive scalar curvatures on aspherical 5-manifolds, preprint, arXiv: 2009.05332.
- [11] Gromov, M., Lawson, H. B. Jr. Positive scalar curvature and the Dirac operator on complete Riemannian manifolds, Inst. Hautes Études Sci. Publ. Math. (1983), no. 58, 83–196.
- [12] Kazdan, J. L. Deformation to positive scalar curvature on complete manifolds, Math. Ann. 261 (1982), no. 2, 227–234.
- [13] Lesourd, M., Unger, R., Yau, S.-T. Positive scalar curvature on noncompact manifolds and the Liouville theorem, Comm. Anal. Geom. 32 (2024), no. 5, 1311–1337.
- [14] Li, C., Mantoulidis, C. Positive scalar curvature with skeleton singularities, Math. Ann. 374 (2019), no. 1-2, 99–131.
- [15] Lück, W. Survey on aspherical manifolds, European Congress of Mathematics, 53–82, Eur. Math. Soc., Zürich, 2010.
- [16] Ratcliffe, J. G., Tschantz, S. T. Some examples of aspherical 4-manifolds that are homology 4-spheres, Topology 44 (2005), no. 2, 341–350.
- [17] Rosenberg, J. C^{*}-algebras, positive scalar curvature, and the Novikov conjecture, Inst. Hautes Études Sci. Publ. Math. 58 (1983), 197–212.
- Schoen, R., Yau, S.-T. Existence of incompressible minimal surfaces and the topology of three-dimensional manifolds with nonnegative scalar curvature, Ann. of Math. (2) 110 (1979), no. 1, 127–142.
- [19] Schoen, R., Yau, S.-T. The structure of manifolds with positive scalar curvature, Directions in partial differential equations (Madison, WI, 1985), 235–242. Publ. Math. Res. Center Univ. Wisconsin, 54 Academic Press, Inc., Boston, MA, 1987.
- [20] Shi, Y., Wang, J., Wu, R., Zhu, J. On open manifolds admitting no complete metric with positive scalar curvature, preprint, arXiv: 2404.01660.
- [21] Wang, J. Contractible 3-manifolds and positive scalar curvature, Ph.D. thesis, Université Grenoble Alpes, 2019.

(Shuli Chen) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 S UNI-VERSITY AVE, CHICAGO IL, 60637, UNITED STATES *Email address*: shulichen@uchicago.edu

(Jianchun Chu) SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, YI-HEYUAN ROAD 5, BEIJING 100871, PEOPLE'S REPUBLIC OF CHINA *Email address*: jianchunchu@math.pku.edu.cn

(Jintian Zhu) INSTITUTE FOR THEORETICAL SCIENCES, WESTLAKE UNIVERSITY, 600 DUNYU ROAD, HANGZHOU, ZHEJIANG 310030, PEOPLE'S REPUBLIC OF CHINA *Email address*: zhujintian@westlake.edu.cn

20