Divisibility Relations Between Ring Homomorphisms and Surjective Group Homomorphisms in Finite Cyclic Structures

Sonu Kumar^a, Priyabrata Mandal^a

^aDepartment of Mathematics, Manipal Institute of Technology, Manipal Academy of Higher Education, Manipal, 576104, India

Abstract

In this article, we delve into the intricate relationship between the number of ring homomorphisms and surjective group homomorphisms between two finite cyclic structures, specifically \mathbb{Z}_m and \mathbb{Z}_n . We demonstrate that the number of ring homomorphisms from \mathbb{Z}_m to \mathbb{Z}_n is a divisor of the number of surjective group homomorphisms from \mathbb{Z}_m to \mathbb{Z}_n , provided that n is not of the form $2 \cdot \alpha$, where each prime factor p of α satisfies $p \equiv 3 \pmod{4}$.

Keywords: Group homomorphism, Ring homomorphism, Prime factorization 2000 MSC: 20K01, 20K30, 11A41

Introduction

Homomorphisms are essential tools in algebra for understanding the structural connections between different algebraic objects, such as groups and rings. These mappings not only preserve the fundamental operations of these structures (like addition and multiplication) but also capture the inherent symmetries and compatibilities between them. By studying homomorphisms, we gain valuable insights that have numerous applications in both theoretical and practical mathematics.

Cyclic groups and rings, denoted as \mathbb{Z}_m and \mathbb{Z}_n , respectively, are fundamental algebraic structures. A cyclic group \mathbb{Z}_m consists of the integers modulo m equipped with the operation of addition, while a cyclic ring \mathbb{Z}_m comprises the same elements but with both addition and multiplication modulo m. Despite sharing a common notation, the nature of homomorphisms

Preprint submitted to Communications in Algebra

February 21, 2025

between these structures differs substantially due to the additional ring structure of \mathbb{Z}_m .

This paper investigates the relationship between the number of ring homomorphisms from \mathbb{Z}_m to \mathbb{Z}_n and the number of surjective group homomorphisms between the corresponding groups. Unlike previous studies that examined these homomorphisms separately, our main finding reveals a specific divisibility relationship between them. This relationship holds unless n is not of the form $2 \cdot \alpha$, where each prime factor p of α satisfies $p \equiv 3$ (mod 4). This condition underscores certain algebraic constraints that influence the count of homomorphisms, indicating a complex interaction between the ring and group structures.

1. Preliminaries

In this section, we provide essential definitions, proof outlines, and notation used throughout the paper. For a more in-depth discussion, we refer the reader to [1].

Definition 1.1. Let G and G' be groups. A function $\psi : G \to G'$ is called a group homomorphism if, for all $a, b \in G$, the following condition holds:

$$\psi(ab) = \psi(a) \cdot \psi(b),$$

where \cdot represents the group operation in G'.

The homomorphism ψ is surjective (or onto) if, for every $y \in G'$, there exists an element $x \in G$ such that $\psi(x) = y$.

Definition 1.2. A ring homomorphism is a function between two rings that preserves the fundamental operations of addition and multiplication.

More formally, let R and S be rings. A function $\psi : R \to S$ is called a ring homomorphism if it meets the following conditions:

1. Additive Preservation: For all $a, b \in R$,

$$\psi(a+b) = \psi(a) + \psi(b),$$

where + denotes addition in both R and S.

2. Multiplicative Preservation: For all $a, b \in R$,

$$\psi(ab) = \psi(a) \cdot \psi(b)$$

where \cdot denotes multiplication in both R and S.

Theorem 1.3. Let G be a cyclic group of order m and H be a cyclic group of order n. The number of surjective group homomorphisms from G to H is $\varphi(n)$ if n divides m, where φ denotes Euler's totient function.

Proof. Let $G = \langle a \rangle$ be a cyclic group, and let $\psi : G \to H$ be a surjective group homomorphism such that $\psi(a) = x$, where $x \in H$.

Since $a^m = e_G$, it follows that $\psi(a^m) = (\psi(a))^m = \psi(e_G) = e_H$. Thus, n divides m.

Next, we show that $\varphi(n)$ is the number of surjective group homomorphisms.

Consider the subgroup $\text{Im}(\psi) = \langle x \rangle$ in *H*. Since ψ is surjective, we have $\text{Im}(\psi) = H$, meaning $H = \langle x \rangle$.

As there are $\varphi(n)$ elements in H with order n, the number of surjective homomorphisms from G to H is therefore $\varphi(n)$.

Theorem 1.4. The number of idempotent elements in any integral domain R is exactly two.

Proof. Let a be an idempotent element in R. Then we have

$$a^{2} = a,$$

$$\implies a^{2} - a = 0,$$

$$\implies a(a - 1) = 0.$$

Since R is an integral domain, we must have either a = 0 or a - 1 = 0, i.e., a = 1 or a = 0.

Hence, the only idempotent elements in R are 0 and 1.

Remark 1.5. Note that, the above Theorem 1.4 does not necessarily holds for a ring which is not an integral domain. For example, consider the ring \mathbb{Z}_{10} , which is not an integral domain. It is easy to see that 0, 1, 5, 6 are the idempotent elements in \mathbb{Z}_{10} .

2. Main results

Lemma 2.1. There exists exactly one surjective group homomorphism and two distinct ring homomorphisms from \mathbb{Z}_m to \mathbb{Z}_2 when m is even.

_____ .

Proof. Let $\psi : (\mathbb{Z}_m, +) \to (\mathbb{Z}_2, +)$ be a group homomorphism. For any $x \in \mathbb{Z}_m$, we have

$$\psi(x) = \psi\left(\underbrace{1+1+\dots+1}_{x \text{ times}}\right) = x \cdot \psi(1).$$

Thus, the homomorphism is fully determined by the value of $\psi(1)$, and since $\psi(1)$ can take only two possible values in \mathbb{Z}_2 , we have exactly two choices for ψ . If $\psi(1) = 0$, then ψ is the zero map, which serves as a trivial homomorphism. The other homomorphism is defined by setting $\psi(1) = 1$. Consequently, there is exactly one surjective group homomorphism.

Next, consider the ring homomorphism $\theta : (\mathbb{Z}_m, +, \cdot) \to (\mathbb{Z}_2, +, \cdot)$. By definition, a ring homomorphism satisfies

$$\theta(a+b) = \theta(a) + \theta(b)$$
 and $\theta(a \cdot b) = \theta(a) \cdot \theta(b)$ for all $a, b \in \mathbb{Z}_m$.

We observe that $\theta(1) = \theta(1 \cdot 1) = \theta(1)^2$. Thus, $\theta(1)$ must be an idempotent element within \mathbb{Z}_2 . Since \mathbb{Z}_2 contains only two idempotent elements, there are two possible cases: either $\theta(1) = 0$ or $\theta(1) = 1$. Therefore, there exist precisely two distinct ring homomorphisms from \mathbb{Z}_m to \mathbb{Z}_2 .

In Lemma 2.1, we observe that the number of ring homomorphisms from the ring \mathbb{Z}_m to \mathbb{Z}_2 does not divide the number of surjective group homomorphisms from the group \mathbb{Z}_m to \mathbb{Z}_2 . This observation leads us to establish the following theorem in a more general context.

Theorem 2.2. Let n > 2. The number of ring homomorphisms from the ring \mathbb{Z}_m to \mathbb{Z}_n divides the number of surjective group homomorphisms from the group \mathbb{Z}_m to \mathbb{Z}_n , except in the case when $n = 2 \cdot \alpha$, where each prime factor p of α satisfies $p \equiv 3 \pmod{4}$.

Proof. As shown in [1], the number of ring homomorphisms from \mathbb{Z}_m to \mathbb{Z}_n is given by $2^{w(n)}$, where w(n) denotes the number of distinct prime divisors of n. Additionally, by Theorem 1.3, the number of surjective group homomorphisms from \mathbb{Z}_m to \mathbb{Z}_n is $\varphi(n)$. We need to show that $2^{w(n)}$ divides $\varphi(n)$. Consider the following cases based on the structure of n.

Case I: Suppose *n* is odd. Let $n = p_1^{r_1} p_2^{r_2} \cdots p_k^{r_k}$, where the p_i 's are distinct odd primes. In this case, we have w(n) = k. Now,

$$\varphi(n) = \varphi(p_1^{r_1} p_2^{r_2} \cdots p_k^{r_k})$$
$$= \prod_{i=1}^k \varphi(p_i^{r_i})$$
$$= \prod_{i=1}^k (p_i^{r_i} - p_i^{r_i-1})$$
$$= \prod_{i=1}^k p_i^{r_i-1} (p_i - 1)$$

Since the p_i 's are odd primes, it follows that 2 divides $(p_i - 1)$ for each $1 \leq i \leq k$. Therefore, 2^k divides $\varphi(n)$. Consequently, we conclude that $2^{w(n)} | \varphi(n)$.

Case II: Suppose *n* is even and is of the form $n = 2^s \cdot p_1^{r_1} p_2^{r_2} \cdots p_k^{r_k}$, where the p_i 's are distinct primes and s > 1. In this case, the number of distinct prime divisors of *n* is given by w(n) = k + 1. We have

$$\varphi(n) = \varphi(2^{s} \cdot p_{1}^{r_{1}} p_{2}^{r_{2}} \cdots p_{k}^{r_{k}})$$

= $\varphi(2^{s}) \cdot \prod_{i=1}^{k} \varphi(p_{i}^{r_{i}})$
= $\varphi(2^{s}) \cdot \prod_{i=1}^{k} (p_{i}^{r_{i}} - p_{i}^{r_{i}-1})$
= $(2^{s} - 2^{s-1}) \cdot \prod_{i=1}^{k} p_{i}^{r_{i}-1} (p_{i} - 1).$

Since s > 1, we deduce that 2 divides $(2^s - 2^{s-1})$. Additionally, by Case I, we know that 2^k divides the remaining product $\prod_{i=1}^k p_i^{r_i-1}(p_i-1)$. Therefore, we conclude that 2^{k+1} divides $\varphi(n)$, which implies that $2^{w(n)} \mid \varphi(n)$.

Case III: Let $n = 2 \cdot \alpha$ where α is odd. We can express α as $\alpha = p_1^{r_1} p_2^{r_2} \cdots p_k^{r_k}$. The number of distinct prime divisors is given by w(n) = k + 1. As shown above in Case II, we have

$$\varphi(n) = \varphi(2) \cdot \prod_{i=1}^{k} p_i^{r_i - 1}(p_i - 1) = \prod_{i=1}^{k} p_i^{r_i - 1}(p_i - 1).$$

Since each p_i is an odd prime, we deduce that 2^k divides $\varphi(n)$. However, we want to determine whether 2^{k+1} also divides $\varphi(n)$. To investigate this, we will consider the following two subcases.

Subcase I: If $p_i \equiv 1 \pmod{4}$ for some *i*, then $4 \mid (p_i - 1)$, which implies that $(p_i - 1) = 4t_i$ for some $t_i \in \mathbb{Z}$. Therefore, we can express $\varphi(n)$ as follows:

$$\varphi(n) = \prod_{i=1}^{k} p_i^{r_i - 1}(p_i - 1) = \prod_{i=1}^{k} p_i^{r_i - 1} \cdot 4t_i = 2^{2k} \cdot \prod_{i=1}^{k} p_i^{r_i - 1} \cdot t_i.$$

From this, it follows that 2^{k+1} divides $\varphi(n)$.

Subcase II: Let $p_i \equiv 3 \pmod{4}$ for all $1 \leq i \leq n$, which implies that $(p_i - 1) \equiv 2 \mod{4}$. Consequently, we can express $(p_i - 1)$ as $4t_i + 2$ for some $t_i \in \mathbb{Z}$. Thus, we have:

$$\varphi(n) = \prod_{i=1}^{k} p_i^{r_i - 1}(p_i - 1) = \prod_{i=1}^{k} p_i^{r_i - 1} \cdot 2(2t_i + 1) = 2^k \cdot \underbrace{\prod_{i=1}^{k} p_i^{r_i - 1}(2t_i + 1)}_{\text{odd integer}}.$$

In this case, it is evident that 2^k divides $\varphi(n)$, but 2^{k+1} does not divide $\varphi(n)$.

Thus, we conclude that if n divides m, then the number of ring homomorphisms from the ring \mathbb{Z}_m to \mathbb{Z}_n divides the number of surjective group homomorphisms from the group \mathbb{Z}_m to \mathbb{Z}_n , except when $n = 2 \cdot \alpha$, where each prime factor p of α satisfies $p \equiv 3 \pmod{4}$.

Acknowledgement

We extend our sincere gratitude to Prof. Jaikumar Radhakrishnan of ICTS-TIFR, Bengaluru, for his invaluable comments on an earlier version of this paper.

Disclosure statement

No potential conflict of interest was reported by the author(s).

References

 Joseph A. Gallian, James Van Buskirk, The Number of Homomorphisms From Z_m Into Z_n, The American Mathematical Monthly, 91:3, 196-197, 1984.