ORIENTATION-REVERSING CRYSTALLOGRAPHIC RIGIDITY

J. ESSON, E. KASTIS, AND B. SCHULZE

ABSTRACT. This paper provides a combinatorial characterisation for generic forced symmetric rigidity of bar-joint frameworks in the Euclidean plane that are symmetric with respect to the orientation-reversing wallpaper group $\mathbb{Z}^2 \rtimes C_s$, also known as pm in crystallography, under a fixed lattice representation. Corresponding results for the wallpaper groups cm and pg follow directly from this. The method used also provides an inductive construction for the corresponding gain graphs, in terms of Henneberg-type graph operations.

Keywords: infinitesimal rigidity; crystallographic framework; wallpaper group; gain graph; sparsity counts.

1. INTRODUCTION

A d-dimensional (bar-joint) framework is a pair (G, p) consisting of a simple graph G = (V, E)and a map $p: V \to \mathbb{R}^d$ assigning positions to the vertices of G, with $p(v_i) \neq p(v_j)$ for $v_i v_j \in E$. The map p is also called a *configuration* of G in \mathbb{R}^d . Frameworks provide suitable mathematical models for a variety of real-world structures whose rigidity and flexibility properties are essential for their behaviour and functioning. Loosely speaking, a framework is rigid if the vertices cannot be moved continuously into another non-congruent framework while maintaining the edge lengths, and flexible otherwise. Since determining whether a framework is rigid is computationally challenging [1], it is common to linearise the problem by differentiating the length constraints, leading to the classical notion of infinitesimal rigidity [30, 34]. An *infinitesimal motion* of a framework (G, p) is a map $u: V \to \mathbb{R}^d$ such that, for all $v_i v_j \in E$,

$$(p(v_i) - p(v_j)) \cdot (u(v_i) - u(v_j)) = 0.$$

The infinitesimal motion u is *trivial* if it corresponds to a rigid body motion, and *non-trivial* otherwise. The matrix corresponding to the linear system above is called the *rigidity matrix*, denoted R(G, p), and (G, p) is *infinitesimally rigid* if G is complete on at most d + 1 vertices or R(G, p) has maximum rank (equal to $d|V| - \binom{d+1}{2}$). An infinitesimally rigid framework which is also *independent* (i.e. the rigidity matrix has linearly independent rows) is also called *isostatic*. For generic configurations, infinitesimal rigidity is equivalent to rigidity [2], and both depend only on the underlying graph G. Generic rigidity is well understood in the plane, and there are significant partial results in higher dimensions (see e.g. [30, 6, 34]).

In applications, one often encounters non-generic configurations, because both natural and man-made structures typically exhibit symmetry. This has led to a surge of interest in the rigidity and flexibility of finite symmetric and infinite periodic frameworks. Crucially, if we consider configurations that are as generic as possible with the given symmetry constraints, then (infinitesimal) rigidity may still be characterised combinatorially in terms of the corresponding group-labelled quotient graph. For a summary of combinatorial results regarding the rigidity of finite symmetric frameworks, we refer the reader to [31, 28]. Let us briefly summarise what is known regarding the rigidity of generic infinite periodic frameworks. Here the allowed continuous motions are those that preserve the bar lengths and the periodicity. The forced periodicity is a key feature of this model: there are frameworks that only have trivial periodicity-preserving motions, but can be deformed non-trivially if larger classes of motions are allowed. Such frameworks are still rigid in the forced periodic model. Note that the lattice representation may either be fixed or allowed to change as the framework moves. See [22] and [4] for the respective mathematical models.

Generic rigid periodic frameworks with a fixed lattice representation in the plane have been characterised by E. Ross [23]. The analogous result for the fully flexible lattice representation in the plane was obtained by J. Malestein and L. Theran [18]. Similar results have also been established for partially flexible lattice representations [21, 20]. For frameworks with crystallographic symmetry, specifically periodic frameworks with added symmetry in the fundamental cells, necessary conditions for forced symmetric rigidity for various types of lattice flexibility were given in [24]. In [19], J. Malestein and L. Theran established complete combinatorial characterisations of forced symmetric generic rigid frameworks with crystallographic symmetry in the plane, where the group is generated by translations and rotations, and the lattice representation is fully flexible. The analogous result for the fixed lattice representation was recently obtained by D. Bernstein in [3] using a new approach based on tropical geometry.

A key gap in this theory is to deal with orientation-reversing wallpaper groups. We will address this gap here for the fixed lattice representation in the plane by giving a characterisation of generic forced symmetric rigidity for the group $\mathbb{Z}^2 \rtimes C_s$, also denoted by pm in the Hermann-Maguin notation used in crystallography [26, 25] or ** in the orbifold notation advocated by J.H. Conway [7]. Characterisations for forced symmetric rigidity with respect to the subgroups cm and pg of pm (called $*\times$ and $\times\times$ in the orbifold notation, respectively) follow directly from this, as will be discussed at the end of Section 5.

The paper is organised as follows. In Section 2 we review the necessary notions from the rigidity theory of symmetric frameworks. Section 3 introduces the relevant sparsity conditions for characterising $\mathbb{Z}^2 \rtimes C_s$ -symmetric rigidity, states the main result, and proves the necessity of the conditions. In Section 4 we introduce the Henneberg-type graph operations we will use for the induction proof of the main result and show that these operations preserve rigidity. These mostly follow standard methods, but a different approach is required for the case of a 1-extension that adds a triple of parallel edges to the gain graph. Section 5 is then dedicated to the main combinatorial part of the proof. We conclude the paper in Section 6 with a discussion on future research directions.

2. Symmetric frameworks

In this paper, Γ will always be a subgroup of the Euclidean group $\operatorname{Isom}(\mathbb{R}^d)$. A simple graph \tilde{G} is Γ -symmetric if there is a group action $\Gamma \to \operatorname{Aut}(\tilde{G})$. We will only consider actions that are free, meaning that no non-identity elements of the group Γ fix any vertices. A useful tool for studying symmetric graphs is the (group-labelled) gain graph (see e.g. [10, 22, 8]), which is defined as follows.

Let G = (V, E) be a multigraph and let Γ be a group. Choose an orientation for each edge and label the oriented edge set \vec{E} . A Γ -gain assignment is a function $m : \vec{E} \to \Gamma$, which assigns a gain to each edge such that parallel edges with the same orientation have different gains and every loop has a non-identity gain. A Γ -gain graph is a pair (G, m) consisting of a multigraph G = (V, E) and a Γ -gain assignment $m : \vec{E} \to \Gamma$. An edge e from vertex v_i to vertex v_j with gain m(e) is denoted by $e = (v_i, v_j; m(e))$.

For a Γ -gain graph (G, m), the derived graph \tilde{G} is the Γ -symmetric graph that is represented by (G, m). If G = (V, E), then $\tilde{G} = (\tilde{V}, \tilde{E})$, where $\tilde{V} = \{(v, \gamma) : v \in V, \gamma \in \Gamma\}$ and $((v_i, \gamma_i), (v_j, \gamma_j))$ is an edge of \tilde{E} if and only if there exists $(v_i, v_j; \gamma_i^{-1}\gamma_j) \in \tilde{E}$. Conversely, given a Γ -symmetric graph \tilde{G} , we may construct the quotient Γ -gain graph G that has \tilde{G} as its derived graph by choosing a representative for each vertex orbit and adding an edge for each edge orbit with the corresponding gain. An edge $((v_i, \gamma_i), (v_j, \gamma_j)) \in \tilde{E}$ is represented by $(v_i, v_j; \gamma_i^{-1}\gamma_j) \in \tilde{E}$ in the gain graph. The choice of orientation for each edge in the gain graph is unimportant, as it is equivalent to choose the opposite orientation and then assign the inverse gain to that edge.

For some $\Gamma \leq \text{Isom}(\mathbb{R}^d)$, let (G, m) be a Γ -gain graph with derived graph \tilde{G} . A configuration $\tilde{p}: \tilde{V} \to \mathbb{R}^d$ of \tilde{G} is Γ -symmetric if, for all $v \in V$ and $\gamma \in \Gamma$, we have

$$\tilde{p}(v,\gamma) = \gamma(\tilde{p}(v,\mathrm{id})).$$

A framework (\tilde{G}, \tilde{p}) is Γ -symmetric if \tilde{G} is a Γ -symmetric graph and $\tilde{p} : \tilde{V} \to \mathbb{R}^d$ is a corresponding Γ -symmetric configuration. Note that a Γ -gain framework (G, m, p) consisting of a Γ -gain graph (G, m) and $p : V \to \mathbb{R}^d$ uniquely determines the derived Γ -symmetric framework (\tilde{G}, \tilde{p}) .

For $\gamma \in \text{Isom}(\mathbb{R}^2)$, let γ_l denote the linear part of the isometry γ . An infinitesimal motion $\tilde{u}: \tilde{V} \to \mathbb{R}^d$ of a Γ -symmetric framework (\tilde{G}, \tilde{p}) is Γ -symmetric if, for all $(v, \gamma) \in \tilde{V}$, we have

$$\tilde{u}((v,\gamma)) = \gamma_l(\tilde{u}((v,\mathrm{id}))).$$

A Γ -symmetric framework is Γ -symmetrically infinitesimally rigid if all of its Γ -symmetric infinitesimal motions are trivial. Otherwise, it is Γ -symmetrically infinitesimally flexible. Note that if Γ contains translations, then the infinite framework (\tilde{G}, \tilde{p}) has periodic symmetry, and the lattice representation is fixed under any motion, since Γ is fixed.

A now classical object to check Γ -symmetric infinitesimal rigidity is the orbit rigidity matrix [29, 33].

Let (G, m, p) be a Γ -gain framework with derived Γ -symmetric framework (\tilde{G}, \tilde{p}) . Then the orbit rigidity matrix $\mathbf{O}(\tilde{G}, \tilde{p}, \Gamma)$ of (\tilde{G}, \tilde{p}) is the $|E| \times d|V|$ matrix with the following row for each edge $e = (v_i, v_j; m(e)) \in E$:

(1) If e is not a loop $(v_i \neq v_j)$, then the row for e is:

$$e \begin{bmatrix} 0 & \dots & 0 & (p(v_i) - m(e)p(v_j))^T & 0 & \dots & 0 & (p(v_j) - (m(e))^{-1}p(v_i))^T & 0 & \dots & 0 \end{bmatrix}$$

(2) If e is a loop $(v_i = v_j)$, then the row for e is:

$$e \begin{bmatrix} 0 & \dots & 0 & (2p(v_i) - m(e)p(v_i) - (m(e))^{-1}p(v_i))^T & 0 & \dots & 0 \end{bmatrix}.$$

Note that $u \in \mathbb{R}^{d|V|}$ extends to a Γ -symmetric infinitesimal motion $\tilde{u} \in \mathbb{R}^{d|\tilde{V}|}$ of (\tilde{G}, \tilde{p}) if and only if $\mathbf{O}(\tilde{G}, \tilde{p}, \Gamma)u = 0$. In this case, u acts as a restriction of \tilde{u} to the set of vertex orbit representatives used for the gain graph. As such, ker $(\mathbf{O}(\tilde{G}, \tilde{p}, \Gamma))$ is isomorphic to the space of Γ -symmetric infinitesimal motions. Since the dimension of the space of trivial Γ -symmetric infinitesimal motions can easily be determined (see e.g. [27, 24]) Γ -symmetric infinitesimal rigidity may be checked by computing the rank of $\mathbf{O}(\tilde{G}, \tilde{p}, \Gamma)$.

A configuration $p: V \to \mathbb{R}^d$ of a Γ -gain graph (G, m) is called *generic* if the corresponding orbit rigidity matrix $\mathbf{O}(\tilde{G}, \tilde{p}, \Gamma)$ achieves the maximum possible rank among configurations of (G, m). The set of generic configurations is open and dense in the set of all such configurations. Like for non-symmetric rigidity, assuming genericity allows rigidity to be treated as a property of the gain graph. A Γ -gain graph (G, m) is said to be *rigid* in \mathbb{R}^d if, for every (equivalently, for some) generic configuration $p: V \to \mathbb{R}^d$, the derived framework (\tilde{G}, \tilde{p}) is Γ -symmetrically infinitesimally rigid. Otherwise, it is called *flexible*. A Γ -gain graph (G, m) is *independent* in \mathbb{R}^d if $\mathbf{O}(\tilde{G}, \tilde{p}, \Gamma)$ is row-independent, and *dependent* otherwise. A gain graph is *minimally rigid* in \mathbb{R}^d if it is rigid and independent in \mathbb{R}^d .

A walk in a gain graph (G, m) is denoted by $e_1^{\alpha_1} e_2^{\alpha_2} \dots e_k^{\alpha_k}$, where $e_1, e_2, \dots, e_k \in E$ and the values $\alpha_1, \alpha_2, \dots, \alpha_k$ are +1 for edges that are traversed forwards, and -1 for edges that are traversed backwards. The *net gain* for this walk is the element $\prod_{i=1}^k m(e_i)^{\alpha_i}$. For a given vertex $v \in V$, the *gain space* $\langle (G, m) \rangle_v$ of (G, m) is the subgroup of Γ that is generated by the net gains on all closed walks in (G, m) that start and end at v [10, 22, 31].

A gain graph (G, m) is balanced if, for every $v \in V$, the gain space $\langle (G, m) \rangle_v$ is trivial. Otherwise, it is unbalanced.

Let (G, m) be a Γ -gain graph. For some $v \in V$ and $\gamma \in \Gamma$, a switching operation at v by γ defines a new gain assignment $m' : \vec{E} \to \Gamma$ as follows:

$$m'(e) = \begin{cases} \gamma m(e)\gamma^{-1} & \text{if } e \text{ is a loop incident to } v;\\ \gamma m(e) & \text{if } e \text{ is a non-loop edge directed from } v;\\ m(e)\gamma^{-1} & \text{if } e \text{ is a non-loop edge directed to } v;\\ m(e) & \text{otherwise.} \end{cases}$$

Gain assignments are *equivalent* if one can be reached from the other by a sequence of switching operations. In effect, a switching operation at v changes the choice of vertex orbit representative for v that is used to obtain the gain graph from the derived graph. Since the derived graph does not change, switching operations preserve rigidity and independence of gain graphs. For point groups, this was shown in part of the proof of [10, Lemma 5.2]. When, for a given starting vertex, the gain space is a normal subgroup of the overall gain group, switching operations will preserve the gain space. In particular, performing switching operations on a balanced gain graph.

Using switching operations, the edges of any spanning tree in a Γ -gain graph can be assigned identity labels. Based on this simple observation, we have the following result, which is an extension of [10, Lemma 2.3].

Lemma 2.1. Let (G,m) be a Γ -gain graph such that for every vertex v of G, the gain space $\langle (G,m) \rangle_v$ is contained in $\Gamma' \leq \Gamma$. Then there is an equivalent gain graph (G,m') in which every gain is an element of Γ' .

3. The conditions for reflectional periodic rigidity

From now on we will work in the Euclidean plane. In the following we will let Γ be the group $\mathbb{Z}^2 \rtimes \mathcal{C}_s$, which is the wallpaper group formed by taking the semi-direct product of the group \mathbb{Z}^2 of translations (w.l.o.g. generated by the vectors (1,0) and (0,1)) with the reflectional group \mathcal{C}_s , generated by the reflection s, in the plane (where w.l.o.g. the mirror line of s is the x-axis) [32, Section 3.2]. In the Hermann–Mauguin notation used in crystallography, this group is denoted by pm.

Our main goal is to characterise the $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -gain graphs that are rigid in \mathbb{R}^2 . To state the necessary and sufficient conditions, we need the following definitions.

Let $k \in \mathbb{N}$ and $l \in \mathbb{N}_0$. A multigraph G = (V, E) is (k, l)-sparse if all subgraphs $G' = (V', E') \subseteq G$ with $|E'| \ge 1$ satisfy $|E'| \le k|V'| - l$. The multigraph G is (k, l)-tight if it is (k, l)-sparse and it satisfies |E| = k|V| - l.

A Γ -gain graph that, for every choice of starting vertex, has a gain space consisting only of translations is said to be *purely periodic*.

Definition 3.1. A $\mathbb{Z}^2 \rtimes C_s$ -gain graph (G, m) is said to be $\mathbb{Z}^2 \rtimes C_s$ -tight if it satisfies the following conditions:

- (1) (2,1)-tight Condition: G is (2,1)-tight;
- (2) Purely Periodic Condition: Every purely periodic subgraph of G is (2,2)-sparse;
- (3) Balanced Condition: Every balanced subgraph of G is (2,3)-sparse.

We will prove the following main theorem.

Theorem 3.2. A $\mathbb{Z}^2 \rtimes C_s$ -gain graph (G,m) is minimally rigid if and only if it is $\mathbb{Z}^2 \rtimes C_s$ -tight.

The necessity of the conditions is easy to see and follows from the lemma below. The sufficiency will be proved in Sections 4 and 5.

Lemma 3.3. Every minimally rigid $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -gain graph is $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight.

Proof. Suppose that (G, m) is a minimally rigid $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -gain graph and let $p: V \to \mathbb{R}^2$ be a generic configuration of (G, m) that derives the $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -symmetric framework (\tilde{G}, \tilde{p}) . We begin by proving that (G, m) is (2, 1)-tight. It is easy to see that the only trivial $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -symmetric infinitesimal motions in the plane are those induced by translations parallel to the reflection axis, giving a space of dimension 1. Thus, we must have |E| = 2|V| - 1. Similarly, for any subgraph $G' = (V', E') \subseteq G$ that derives a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -symmetric subgraph $\tilde{G}' = (\tilde{V}', \tilde{E}') \subseteq \tilde{G}$, we have $\operatorname{rank}(\mathbf{O}(\tilde{G}', \tilde{p}|_{\tilde{V}'}, \mathbb{Z}^2 \rtimes \mathcal{C}_s)) \leq 2|V'| - 1$. Since (G, m) is independent, it follows that $|E'| \leq 2|V'| - 1$. This shows that G is (2, 1)-tight.

Suppose next that G' = (V', E') is a purely periodic subgraph of G with derived $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ symmetric graph $\tilde{G}' = (\tilde{V}', \tilde{E}')$. By Lemma 2.1 and the argument in [10, Lemma 5.2], switching
operations can be used to consider (G', m) as a \mathbb{Z}^2 -gain graph without changing the rank of the
orbit rigidity matrix. Since \mathbb{Z}^2 admits a 2-dimensional space of trivial periodic infinitesimal motions
(corresponding to the space of translations), it follows that $\operatorname{rank}(\mathbf{O}(\tilde{G}', \tilde{p}|_{\tilde{V}'}, \mathbb{Z}^2 \rtimes \mathcal{C}_s)) \leq 2|V'| - 2$.
Since (G, m) is independent, we have $|E'| \leq 2|V'| - 2$, proving the purely periodic condition.

Finally, to prove necessity of the balanced condition, suppose that G' = (V', E'), where $E' \neq \emptyset$, is a balanced subgraph of G with derived $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -symmetric graph $\tilde{G}' = (\tilde{V}', \tilde{E}')$. By Lemma 2.1 and the argument in [10, Lemma 5.2], switching operations can be used to assign trivial gain to every edge of G', while preserving the rank of the orbit rigidity matrix. As such, G' can be thought of as a simple non-symmetric finite graph. Thus, $\operatorname{rank}(\mathbf{O}(\tilde{G}', \tilde{p}|_{\tilde{V}'}, \mathbb{Z}^2 \rtimes \mathcal{C}_s)) \leq 2|V'| - 3$, and since (G, m) is independent, it follows that $|E'| \le 2|V'| - 3$, proving the balanced condition. This gives the result.

4. Reflectional Periodic Symmetry: Extensions

To prove that $\mathbb{Z}^2 \rtimes C_s$ -tightness is sufficient for minimal rigidity, we will use an inductive approach, which adapts the one in [23, Theorem 5.1] for purely periodic graphs.

The base case of our induction is a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -gain graph on K_1^1 , which consists of a single vertex with a single loop. Clearly, a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -gain graph on K_1^1 is $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight if and only if the loop is assigned a gain with a non-trivial \mathcal{C}_s -component. When this is the case, the orbit rigidity matrix for any generic $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -symmetric framework derived from this gain graph consists of a single non-zero row vector of size 2. Since the space of trivial $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -symmetric infinitesimal motions in the plane is of dimension 1, any such gain graph is minimally rigid.

We will require the following types of graph extension moves, which are variations of some well established moves in symmetric rigidity theory (see e.g. [8] and [10]).

Definition 4.1. Let (G, m) be a $\mathbb{Z}^2 \rtimes C_s$ -gain graph. A gained 0-extension forms a new gain graph (G', m) by adding a new vertex v_0 with incident edges $e_1 = (v_0, v_1; m(e_1))$ and $e_2 = (v_0, v_2; m(e_2))$, for some (not necessarily distinct) $v_1, v_2 \in V$. This move is illustrated in Figure 4.1.




FIGURE 4.1. The two variations of the 0-extension.

A gained 1-extension forms a new gain graph (G', m) by removing an edge $e = (v_1, v_2; m(e)) \in E$ and adding a new vertex v_0 with incident edges $e_1 = (v_0, v_1; m(e_1)), e_2 = (v_0, v_2; m(e_2))$ and $e_3 = (v_0, v_3; m(e_3))$, for some $v_3 \in V$, with the additional requirement that $(m(e_1))^{-1}m(e_2) = m(e)$. Note that the vertices $v_1, v_2, v_3 \in V$ need not be distinct. This move is illustrated in Figure 4.2.




FIGURE 4.2. The four variations of the 1-extension.

A gained loop-1-extension forms a new gain graph (G', m) by adding a new vertex v_0 with incident edges $l = (v_0, v_0; m(l))$ and $e = (v_0, v_1; m(e))$, for some $v_1 \in V$, with the additional requirement that m(l) has a non-trivial C_s -component. This move is illustrated in Figure 4.3.




FIGURE 4.3. The loop-1-extension. Note that the gain m(l) must have a non-trivial C_s -component.

Gained 0-reductions, 1-reductions and loop-1-reductions are the inverse moves of gained 0extensions, 1-extensions and loop-1-extensions respectively.

We now show that each of these extensions preserves minimal rigidity of $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -gain graphs.

Proposition 4.2. Let (G, m) be a minimally rigid $\mathbb{Z}^2 \rtimes C_s$ -gain graph and let (G', m) be formed by a gained 0-extension of (G, m). Then (G', m) is minimally rigid.

Proof. This follows from a standard argument, which involves placing the new vertex so that it is not collinear with its neighbours. See sources such as [10, Lemma 6.1] for details. \Box

Proposition 4.3. Let (G, m) be a minimally rigid $\mathbb{Z}^2 \rtimes C_s$ -gain graph and let (G', m) be formed by a gained 1-extension of (G, m). Then (G', m) is minimally rigid.

Proof. We consider each of the cases seen in subfigures of Figure 4.2. The proofs for the cases illustrated in Figures 4.2a, 4.2b and 4.2c follow by a standard argument, such as that seen in [10, Lemma 6.1]. This argument uses genericity to assume that the derived neighbours of the new vertex are not collinear.

This leaves the case illustrated in Figure 4.2d, where a 1-extension is performed on a loop and gives a triple of parallel edges. If there exists a configuration of the derived graph in which the derived neighbours of the new vertex are not collinear, then it is again possible to apply the standard method from [10, Lemma 6.1]. The only case where this is not possible is where the edges incident to the new vertex all have the same horizontal-translation-gain component (this being parallel to the axis of reflection). A different method is needed to complete the proof in this case, which we now describe.

Suppose that (G', m) is formed from (G, m) by a gained 1-extension that adds a new vertex v_0 with incident edges $e_1 = (v_0, v_1; m(e_1))$, $e_2 = (v_0, v_1; m(e_2))$ and $e_3 = (v_0, v_1; m(e_3))$ such that these edges all have the same horizontal-translation-gain component. By switching operations, we may assume w.l.o.g. that $m(e_1) = (0, 0, s)$, $m(e_2) = (0, d_2, 0)$ and $m(e_3) = (0, d_3, 0)$ for some $d_2 \neq d_3 \in \mathbb{Z}$. (To see this, note that the gain of the loop at v_0 in G must have a non-trivial C_s -component, for otherwise the loop is dependent. By switching by the reflection s at v_0 , we may make e_3 have a trivial C_s -component. By definition of 1-extension, exactly one of e_1 and e_2 will also have trivial C_s -component. Since the horizontal-translation gain component must be the same for all three edges, we may then switch by translations to obtain the desired form). Let $p: V \to \mathbb{R}^2$ be a generic configuration of G with $p(v_1) = (a_1, b)$, for some $a_1, b \in \mathbb{R}$ with $b \neq 0$. Let (\tilde{G}, \tilde{p}) be the $\mathbb{Z}^2 \rtimes C_s$ -symmetric framework derived by p. Extend p to a configuration $p': V' \to \mathbb{R}^2$ of (G', m) with $p'|_V = p$ and $p'(v_0) = (a_0, b)$, for some $a_0 \in \mathbb{R}$ with $a_0 \neq a_1$. Let (\tilde{G}', \tilde{p}') be the $\mathbb{Z}^2 \rtimes C_s$ -symmetric framework derived by p' and suppose that this has a $\mathbb{Z}^2 \rtimes C_s$ -symmetric framework derived by p' and suppose that this has a $\mathbb{Z}^2 \rtimes C_s$ -symmetric infinitesimal motion $\tilde{u}: \tilde{V}' \to \mathbb{R}^2$. We aim to show that \tilde{u} must be trivial.

Let $u: V' \to \mathbb{R}^2$ be the restriction of \tilde{u} to the set of vertex orbit representatives used to form the gain graph (G', m). By adding a trivial infinitesimal motion induced by a horizontal translation, it can be assumed that $u(v_1)$ is a vertical vector. For some $x_0, y_0, y_1 \in \mathbb{R}$, let $u(v_1) = (0, y_1)$ and let $u(v_0) = (x_0, y_0)$. We will show that $u(v_0) = u(v_1) = (0, 0)$, which will imply that \tilde{u} is trivial, since (\tilde{G}, \tilde{p}) is $\mathbb{Z}^2 \rtimes C_s$ -symmetrically infinitesimally rigid. The constraint imposed by e_2 requires that

$$\begin{pmatrix} a_0 - a_1 \\ -d_2 \end{pmatrix} \cdot \begin{pmatrix} x_0 \\ y_0 - y_1 \end{pmatrix} = 0.$$

Hence, $x_0(a_0 - a_1) - d_2(y_0 - y_1) = 0$. Likewise, the constraint imposed by e_3 is

$$\begin{pmatrix} a_0 - a_1 \\ -d_3 \end{pmatrix} \cdot \begin{pmatrix} x_0 \\ y_0 - y_1 \end{pmatrix} = 0.$$

Hence, $x_0(a_0 - a_1) - d_3(y_0 - y_1) = 0$. Combining these constraints shows that $d_2(y_0 - y_1) = d_3(y_0 - y_1)$. Since $d_2 \neq d_3$, this implies that $y_0 = y_1$. Applying this to the constraint imposed by e_2 shows that $x_0(a_0 - a_1) = 0$. Since $a_0 \neq a_1$, it follows that $x_0 = 0$.

Since $x_0 = 0$ and $y_0 = y_1$, the constraint imposed by e_1 is

$$\begin{pmatrix} a_0 - a_1 \\ 2b \end{pmatrix} \cdot \begin{pmatrix} 0 \\ 2y_1 \end{pmatrix} = 0$$

Hence, $4by_1 = 0$. By the choice of configuration, $b \neq 0$. This means that $y_1 = 0$ and therefore $u(v_0) = u(v_1) = (0, 0)$, as desired.

Proposition 4.4. Let (G, m) be a minimally rigid $\mathbb{Z}^2 \rtimes C_s$ -gain graph and let (G', m) be formed by a gained loop-1-extension of (G, m). Then (G', m) is minimally rigid.

Proof. Again, this can be proved using the standard approach from [10, Lemma 6.1]

5. Reflectional Periodic Symmetry: Reductions

The main aim of this section is to complete the proof of Theorem 3.2 by proving the following result.

Theorem 5.1. A $\mathbb{Z}^2 \rtimes C_s$ -gain graph is $\mathbb{Z}^2 \rtimes C_s$ -tight if and only if it can be obtained from a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph on K_1^1 using a sequence of gained 0-extensions, 1-extensions and loop-1-extensions.

As previously mentioned, any $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph on K_1^1 is minimally rigid. Propositions 4.2, 4.3 and 4.4 show that each of the extensions listed in Theorem 5.1 preserves minimal rigidity. Hence, Theorem 5.1 will show that every $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph is minimally rigid, completing the proof of Theorem 3.2.

We split the proof of Theorem 5.1 into several lemmas. To begin the proof, it is easy to see that each of the extensions preserves $\mathbb{Z}^2 \rtimes C_s$ -tightness. It therefore remains only to show that every $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph can be obtained from a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph on K_1^1 by a sequence of extensions. This can be proved by induction on the number of vertices. Basic counting arguments show that every (2, 1)-tight multigraph has a vertex of degree 2 or 3. It is therefore enough to show that every vertex of degree 2 or 3 in a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph admits a 0-reduction, 1-reduction or loop-1-reduction that preserves $\mathbb{Z}^2 \rtimes C_s$ -tightness. Since K_1^1 is the only (2, 1)-tight multigraph on a single vertex, such a sequence of reductions will eventually terminate at a gain graph on K_1^1 . This will prove Theorem 5.1 by induction.

Different methods will be required depending on the degree and the number of neighbours of the vertex. In the case of a degree 3 vertex with only one neighbour, there are two separate cases, corresponding to either a loop and another edge, or a triple of parallel edges (recall Figures 4.2(d) and 4.3).

The following lemma will be used repeatedly throughout this section. Similar calculations have been used to prove related results in various sources, such as [17, Theorem 5] and [23, Lemma 4.4].

Lemma 5.2. Let $l_A, l_B \in \mathbb{N}_0$. Let G be a multigraph with subgraphs $G_A = (V_A, E_A)$ and $G_B = (V_B, E_B)$ such that $|E_A| = 2|V_A| - l_A$ and $|E_B| = 2|V_B| - l_B$. Then

$$|E_A \cup E_B| + |E_A \cap E_B| = 2|V_A \cup V_B| + 2|V_A \cap V_B| - l_A - l_B.$$

Proof. Calculate

$$|E_A \cup E_B| + |E_A \cap E_B| = |E_A| + |E_B|$$

= $(2|V_A| - l_A) + (2|V_B| - l_B)$
= $2|V_A| + 2|V_B| - l_A - l_B$
= $2|V_A \cup V_B| + 2|V_A \cap V_B| - l_A - l_B.$

Hence, the lemma holds.

Here is another result that will be useful throughout this section.

Lemma 5.3. [17, Theorem 5] Let k, l > 0 be integers. Any (k, l)-tight multigraph with at least one edge is connected.

5.1. Straightforward Reductions. We begin the investigation of reductions by considering the case where the gain graph has a vertex of degree 2.

Proposition 5.4. Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 2. Form the gain graph $G-v_0$ by a gained 0-reduction on G at v_0 . Then $G-v_0$ is also $\mathbb{Z}^2 \rtimes C_s$ -tight.

Proof. Removing v_0 from G removes 1 vertex and 2 edges, so $G - v_0$ satisfies the overall (2, 1)edge-count. Since $G - v_0$ is a subgraph of G, every subgraph of $G - v_0$ is also a subgraph of G and therefore satisfies the other conditions of $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tightness. Hence, $G - v_0$ is $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight. \Box

Now consider a vertex of degree 3, where one of the incident edges is a loop. The same argument that was used to prove Proposition 5.4 can be used to prove that a loop-1-reduction is always admissible here.

Proposition 5.5. Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 3, which is incident to a loop. Form the gain graph $G - v_0$ by a gained loop-1-reduction on G at v_0 . Then $G - v_0$ is also $\mathbb{Z}^2 \rtimes C_s$ -tight.

Next, consider a vertex v_0 of degree 3 that has 3 incident non-loop edges that all join v_0 to the same neighbour.

Proposition 5.6. Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 3 with a triple of parallel incident edges. Then it is possible to form a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph by a gained 1-reduction at v_0 .

Proof. Suppose that v_0 has a triple of parallel incident edges: $e_1 = (v_0, v_1; m(e_1)), e_2 = (v_0, v_1; m(e_2))$ and $e_3 = (v_0, v_1; m(e_3))$ for some $v_1 \in V$. A 1-reduction at v_0 would involve removing v_0 and adding one of the candidate loop edges: $e_{12} = (v_1, v_1; (m(e_1))^{-1}m(e_2)), e_{23} = (v_1, v_1; (m(e_2))^{-1}m(e_3))$ or $e_{31} = (v_1, v_1; (m(e_3))^{-1}m(e_1))$. Hence, the aim is to show that one of $G - v_0 + e_{12}, G - v_0 + e_{23}$ or $G - v_0 + e_{31}$ is $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight.

Since $\{v_0, v_1\}$ induces a (2, 1)-tight subgraph of G, this subgraph cannot be purely periodic. Thus, among $\{e_1, e_2, e_3\}$, there is a pair (e_i, e_j) such that $(m(e_i))^{-1}m(e_j)$ has a non-trivial \mathcal{C}_s component.

Note that $G - v_0 + e_{ij}$ clearly satisfies the overall (2, 1)-edge-count. It is easy to see that any subgraph of $G - v_0 + e_{ij}$ containing e_{ij} is (2, 1)-sparse, as performing the corresponding 1-extension on an overcounted subgraph of $G - v_0 + e_{ij}$ would give an overcounted subgraph of G. Since e_{ij} is a loop and $m(e_{ij})$ has a non-trivial \mathcal{C}_s -component, any subgraph of $G - v_0 + e_{ij}$ containing e_{ij} will be unbalanced and not purely periodic. Hence, any subgraph containing e_{ij} satisfies all of the conditions on subgraphs that are required for the gain graph to be $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight. Any subgraph of $G - v_0 + e_{ij}$ that does not contain e_{ij} is itself a subgraph of G and therefore satisfies the required conditions on subgraphs. Hence, $G - v_0 + e_{ij}$ is $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight.

5.2. **1-reductions on Vertices with Two Neighbours.** The next consideration is the case where a vertex of degree 3 has 2 neighbours. Proving that there is an admissible reduction in this case is considerably more involved than in any of the previous cases.

Proposition 5.7. Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 3 with 2 distinct neighbours. Then it is possible to form a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph by a gained 1-reduction at v_0 .

Suppose that $v_0 \in V$ is a vertex of degree 3 with incident edges $e_1 = (v_0, v_1; m(e_1)), e_2 = (v_0, v_2; m(e_2))$ and $e_3 = (v_0, v_2; m(e_3))$, for some distinct $v_1, v_2 \in V$. After deleting v_0 , the possible options for edges to add for a 1-reduction are $e_{12} = (v_1, v_2; (m(e_1))^{-1}m(e_2)), e_{13} = (v_1, v_2; (m(e_1))^{-1}m(e_3))$ or $e_{23} = (v_2, v_2; (m(e_2))^{-1}m(e_3))$. To prove Proposition 5.7, the aim is to show that one of $G - v_0 + e_{12}, G - v_0 + e_{13}$ or $G - v_0 + e_{23}$ is $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight. Figure 5.1 illustrates the neighbourhood of v_0 , with the candidate edges e_{12}, e_{13} and e_{23} represented by dashed lines.




FIGURE 5.1. A vertex v_0 of degree 3 with two neighbours.

The proof of Proposition 5.7 is split into lemmas covering different cases. The general approach used is similar to that seen for \mathbb{Z}^2 -gain graphs in the corresponding part of the inductive proof of the Periodic Laman Theorem [23, Proposition 5.3], although the version for $\mathbb{Z}^2 \rtimes C_s$ -gain graphs is more complicated.

Note that if e_2 and e_3 have the same C_s -gain component, then the loop e_{23} in $G - v_0 + e_{23}$ will have a purely periodic gain, violating the (2, 2)-sparsity count for the purely periodic condition. So we will treat the cases where e_2 and e_3 have the same and different C_s -components separately.

We begin with the case where the parallel edges e_2 and e_3 have different C_s -gain components and $G - v_0 + e_{23}$ is (2, 1)-tight.

Lemma 5.8. Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 3, with incident edges $e_1 = (v_0, v_1; m(e_1)), e_2 = (v_0, v_2; m(e_2))$ and $e_3 = (v_0, v_2; m(e_3))$ such that $m(e_2)$ and $m(e_3)$ have different C_s -components. If $G - v_0 + e_{23}$ is (2, 1)-tight, then it is $\mathbb{Z}^2 \rtimes C_s$ -tight.

Proof. Since $G - v_0 + e_{23}$ is (2, 1)-tight, it remains only to check that it satisfies the purely periodic condition and the balanced condition. Since $m(e_2)$ and $m(e_3)$ have different C_s -components, $(m(e_2))^{-1}m(e_3)$ has a non-trivial C_s -component. Since e_{23} is a loop, it can be seen that any subgraph of $G - v_0 + e_{23}$ that contains e_{23} will be unbalanced and not purely periodic. Any subgraph of $G - v_0 + e_{23}$ that does not contain e_{23} is itself a subgraph of G and therefore satisfies the required conditions on subgraphs. Hence, $G - v_0 + e_{23}$ is $\mathbb{Z}^2 \rtimes C_s$ -tight. \Box

For other cases, the aim is to perform a reduction to either $G - v_0 + e_{12}$ or $G - v_0 + e_{13}$. Any subgraph of either of these that does not contain the new edge is also a subgraph of G and therefore satisfies all of the conditions required on subgraphs for $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tightness. It is therefore only necessary to consider subgraphs that contain the new edge. The cases where $G - v_0 + e_{12}$ or $G - v_0 + e_{13}$ fails each condition can be characterised in terms of subgraphs of G, known as blockers. These are described in the following definition.

Definition 5.9. Let (G, m) be a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph. Let $v_0 \in V$ be a vertex of degree 3, with two of its incident edges being $e_i = (v_0, v_i; m(e_i))$ and $e_j = (v_0, v_j; m(e_j))$, for some distinct $v_i, v_j \in V$. Let $e_{ij} = (v_i, v_j; (m(e_i))^{-1}m(e_j))$.

A (2,1)-tight blocker for the 1-reduction to $G - v_0 + e_{ij}$ is a (2,1)-tight subgraph $G_{ij} = (V_{ij}, E_{ij}) \subset G$ such that $v_i, v_j \in V_{ij}$ and $v_0 \notin V_{ij}$.

A purely periodic blocker for the 1-reduction to $G - v_0 + e_{ij}$ is a (2,2)-tight purely periodic subgraph $G_{ij} = (V_{ij}, E_{ij}) \subset G$ such that $v_i, v_j \in V_{ij}, v_0 \notin V_{ij}$ and every path in G_{ij} from v_i to v_j has a net gain with the same \mathcal{C}_s -component as $(m(e_i))^{-1}m(e_j)$.

A balanced blocker for the 1-reduction to $G - v_0 + e_{ij}$ is a (2,3)-tight balanced subgraph $G_{ij} = (V_{ij}, E_{ij}) \subset G$ such that $v_i, v_j \in V_{ij}$, $v_0 \notin V_{ij}$ and every path in G_{ij} from v_i to v_j has net gain $(m(e_i))^{-1}m(e_j)$.

Each type of blocker corresponds to a failure of one of the conditions. A 1-reduction that adds a non-loop edge gives a graph that fails the (2, 1)-tight condition if and only if it has a (2, 1)-tight blocker. It gives a graph that fails the purely periodic condition if and only if it has a purely periodic blocker. It gives a graph that fails the balanced condition if and only if it has a balanced blocker.

Now consider the case where e_2 and e_3 have different C_s -gain components but $G - v_0 + e_{23}$ is not (2, 1)-tight.

Lemma 5.10. Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph that has a vertex v_0 of degree 3, with incident edges $e_1 = (v_0, v_1; m(e_1)), e_2 = (v_0, v_2; m(e_2))$ and $e_3 = (v_0, v_2; m(e_3))$ such that $m(e_2)$ and $m(e_3)$ have different \mathcal{C}_s -components, but $G - v_0 + e_{23}$ is not (2, 1)-tight. Then one of $G - v_0 + e_{12}$ or $G - v_0 + e_{13}$ is $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight.

Proof. Since $G-v_0+e_{23}$ is not (2, 1)-tight, there exists a (2, 1)-tight subgraph $G_{23} = (V_{23}, E_{23}) \subset G$ with $v_2 \in V_{23}$ and $v_0 \notin V_{23}$. Note that $v_1 \notin V_{23}$, as otherwise adding v_0 with its incident edges to G_{23} would break the (2, 1)-sparsity of G.

If the reduction to either $G - v_0 + e_{12}$ or $G - v_0 + e_{13}$ has a (2, 1)-tight blocker, then adding v_0 with its incident edges to this blocker gives an overcounted subgraph of G, contradicting the (2, 1)-sparsity of G. Hence, both of these graphs are (2, 1)-tight.

Suppose that $G - v_0 + e_{12}$ fails the purely periodic condition, so it has a purely periodic blocker G_{12} . By Lemma 5.2,

$$E_{12} \cup E_{23}| + |E_{12} \cap E_{23}| = 2|V_{12} \cup V_{23}| + 2|V_{12} \cap V_{23}| - 3.$$

$$(5.1)$$

Since G_{12} is (2, 2)-tight, $|E_{12} \cap E_{23}| \leq 2|V_{12} \cap V_{23}| - 2$. Combining this with Equation (5.1), it follows that $|E_{12} \cup E_{23}| \geq 2|V_{12} \cup V_{23}| - 1$. Adding v_0 with its incident edges to $G_{12} \cup G_{23}$ breaks the (2, 1)-sparsity of G, giving a contradiction. Hence, $G - v_0 + e_{12}$ satisfies the purely periodic condition. Exactly the same method shows that $G - v_0 + e_{13}$ also satisfies the purely periodic condition.

Suppose that $G - v_0 + e_{12}$ and $G - v_0 + e_{13}$ both fail the balanced condition, so they have balanced blockers G_{12} and G_{13} respectively. Consider $G_{12} \cup G_{23}$. By Lemma 5.2,

$$|E_{12} \cup E_{23}| + |E_{12} \cap E_{23}| = 2|V_{12} \cup V_{23}| + 2|V_{12} \cap V_{23}| - 4.$$
(5.2)

Suppose that $|V_{12} \cap V_{23}| > 1$. Since G_{12} is (2, 3)-tight, $|E_{12} \cap E_{23}| \le 2|V_{12} \cap V_{23}| - 3$. Combining this with Equation (5.2), it follows that $|E_{12} \cup E_{23}| \ge 2|V_{12} \cup V_{23}| - 1$. Adding v_0 and its incident edges to $G_{12} \cup G_{23}$ forms a subgraph that breaks the (2, 1)-sparsity of G, a contradiction. The same method gives a contradiction for the case where $|V_{13} \cap V_{23}| > 1$.

This leaves the case where $|V_{12} \cap V_{23}| = |V_{13} \cap V_{23}| = 1$. In this case, note that $|E_{12} \cap E_{23}| = |E_{13} \cap E_{23}| = 0$. Equation (5.2) then shows that $|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| - 2$. Let $G^* = G_{12} \cup G_{23} = (V^*, E^*)$ and consider $G^* \cup G_{13}$. By Lemma 5.2,

$$|E^* \cup E_{13}| + |E^* \cap E_{13}| = 2|V^* \cup V_{13}| + 2|V^* \cap V_{13}| - 5.$$

Note that G_{13} is (2, 3)-tight and that $|V^* \cap V_{13}| > 1$, as $v_1, v_2 \in V^* \cap V_{13}$. Hence, $|E^* \cap E_{13}| \le 2|V^* \cap V_{13}| - 3$ and thus $|E^* \cup E_{13}| \ge 2|V^* \cup V_{13}| - 2$. Since $v_1, v_2 \in V^* \cup V_{13}$ we also have $|E^* \cup E_{13}| \le 2|V^* \cup V_{13}| - 2$, as otherwise adding v_0 and its three incident edges would violate (2, 1)-sparsity. Thus, $|E^* \cup E_{13}| = 2|V^* \cup V_{13}| - 2$. This shows that $G^* \cap G_{13}$ is (2, 3)-tight and hence connected by Lemma 5.3. It therefore contains a path from v_1 to v_2 that is in both G_{12} and G_{13} (the path does not use any edges from G_{23} , as $|E_{13} \cap E_{23}| = 0$). By the definition of a balanced blocker, such a path is required to have gain equal to both $(m(e_1))^{-1}m(e_2)$ and $(m(e_1))^{-1}m(e_3)$, so clearly $(m(e_1))^{-1}m(e_2) = (m(e_1))^{-1}m(e_3)$. However, this implies that $m(e_2) = m(e_3)$, which is a contradiction to the requirement that parallel edges must have different gains. Every case leads to a contradiction, so one of $G - v_0 + e_{12}$ or $G - v_0 + e_{13}$ satisfies the balanced condition. Hence, one of $G - v_0 + e_{12}$ or $G - v_0 + e_{13}$ must be $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight.

Lemmas 5.8 and 5.10 cover the case where $m(e_2)$ and $m(e_3)$ have different C_s -components, leaving the case where they have the same C_s -component. Recall that in this case it is not feasible to perform the 1-reduction to $G - v_0 + e_{23}$, as the loop e_{23} would violate the purely periodic condition. Consequently, we no longer have the (2, 1)-tight blocker G_{23} available to create subgraphs violating the sparsity counts as in the proof of Lemma 5.10. We will show that we can still perform a 1reduction to either $G - v_0 + e_{12}$ or $G - v_0 + e_{13}$. This will be covered by Lemma 5.12, for which the following result regarding connectivity will be needed.

Lemma 5.11. Let G = (V, E) be a (2,3)-sparse multigraph with |E| = 2|V| - 4. Then either G is connected or G consists only of 2 vertices and no edges.

Proof. Suppose that G is disconnected and has a connected component $G_A = (V_A, E_A)$, where $|E_A| \ge 1$. Let $G_B = G \setminus G_A = (V_B, E_B)$, where G_B may or may not contain an edge. Since G

is (2,3)-sparse, we have $|E_A| \leq 2|V_A| - 3$. If $|E_B| \geq 1$, then the (2,3)-sparsity of G implies that $|E_B| \le 2|V_B| - 3$. If $|E_B| = 0$, then $|E_B| = 0 \le 2|V_B| - 2$. In either case, $|E_B| \le 2|V_B| - 2$. Hence, $|E| = |E_A| + |E_B| \le (2|V_A| - 3) + (2|V_B| - 2) = 2|V| - 5.$

This contradicts that fact that |E| = 2|V| - 4, showing that it is not possible for G to contain a proper connected component with an edge. If G is disconnected, then this shows that it has no edges. The only graph with no edges that satisfies |E| = 2|V| - 4 is that on 2 vertices. This gives the result.

Lemma 5.12. Suppose that (G,m) is a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph that has a vertex v_0 of degree 3, with incident edges $e_1 = (v_0, v_1; m(e_1)), e_2 = (v_0, v_2; m(e_2))$ and $e_3 = (v_0, v_2; m(e_3))$ such that $m(e_2)$ and $m(e_3)$ have the same C_s -component. Then one of $G - v_0 + e_{12}$ or $G - v_0 + e_{13}$ is $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight.

Proof. By the same argument that was used in the proof of Lemma 5.10, both $G - v_0 + e_{12}$ and $G - v_0 + e_{13}$ are (2, 1)-tight. Suppose that $G - v_0 + e_{12}$ fails the purely periodic condition, so it has a purely periodic blocker G_{12} . Adding v_0 with its incident edges to G_{12} gives a (2,1)-tight purely periodic subgraph of G, contradicting the fact that G satisfies the purely periodic condition. Hence, $G - v_0 + e_{12}$ satisfies the purely periodic condition. The same method shows that $G - v_0 + e_{13}$ also satisfies the purely periodic condition.

Now suppose that $G - v_0 + e_{12}$ and $G - v_0 + e_{13}$ both fail the balanced condition. This means that they have balanced blockers G_{12} and G_{13} respectively. By Lemma 5.2,

$$|E_{12} \cup E_{13}| + |E_{12} \cap E_{13}| = 2|V_{12} \cup V_{13}| + 2|V_{12} \cap V_{13}| - 6.$$
(5.3)

Since $v_1, v_2 \in V_{12} \cap V_{13}$, we have $|V_{12} \cap V_{13}| > 1$. Note that G_{12} is (2,3)-tight, so $|E_{12} \cap E_{13}| \leq 1$ $2|V_{12} \cap V_{13}| - 3$. Also, note that $|E_{12} \cup E_{13}| \le 2|V_{12} \cup V_{13}| - 2$, as otherwise adding v_0 with its incident edges would break the (2, 1)-sparsity of G. Combining these bounds with Equation (5.3)gives the following possible cases:

- (1) $|E_{12} \cup E_{13}| = 2|V_{12} \cup V_{13}| 3$ and $|E_{12} \cap E_{13}| = 2|V_{12} \cap V_{13}| 3$; (2) $|E_{12} \cup E_{13}| = 2|V_{12} \cup V_{13}| 2$ and $|E_{12} \cap E_{13}| = 2|V_{12} \cap V_{13}| 4$.

In case 1, $G_{12} \cap G_{13}$ is (2,3)-tight and thus connected by Lemma 5.3. Hence, $G_{12} \cap G_{13}$ contains at least one path from v_1 to v_2 that is contained in both G_{12} and G_{13} . It can therefore be seen that $(m(e_1))^{-1}m(e_2) = (m(e_1))^{-1}m(e_3)$ and thus $m(e_2) = m(e_3)$. This is a contradiction, as parallel edges are required to have different gains.

Now consider case 2. By Lemma 5.11, $G_{12} \cap G_{13}$ is either connected or consists of just 2 vertices and no edges. If $G_{12} \cap G_{13}$ is connected, then the same contradiction as for case 1 can be reached. It therefore remains only to consider the case where $G_{12} \cap G_{13}$ is the graph on 2 vertices and no edges. In this case, note that $V_{12} \cap V_{13} = \{v_1, v_2\}$. We now show that $G_{12} \cup G_{13}$ is purely periodic.




FIGURE 5.2. Illustration for the proof of Lemma 5.12, showing the case where the balanced blockers G_{12} and G_{13} intersect in exactly 2 vertices.

To see this, note that any closed walk that is fully contained in one of G_{12} or G_{13} has trivial net gain. Any other closed walk in $G_{12} \cup G_{13}$ can be expressed as a concatenation of walks in G_{12} and walks in G_{13} . Any closed walks in either of these subgraphs will have trivial net gain, so, w.l.o.g., the only other case to consider is a concatenation of a walk from v_1 to v_2 in G_{12} followed by a walk from v_2 to v_1 in G_{13} . This has net gain $(m(e_1))^{-1}m(e_2)(m(e_3))^{-1}m(e_1)$. Figure 5.2 illustrates the paths through G_{12} and G_{13} .

The C_s -components of $(m(e_1))^{-1}$ and $m(e_1)$ cancel out, so $(m(e_1))^{-1}m(e_2)(m(e_3))^{-1}m(e_1)$ has the same C_s -component as $m(e_2)(m(e_3))^{-1}$. Since $m(e_2)$ and $m(e_3)$ have equal C_s -components, a closed walk of this form has a net gain in \mathbb{Z}^2 . Hence, $G_{12} \cup G_{13}$ is purely periodic. Adding v_0 with its incident edges to $G_{12} \cup G_{13}$ gives a (2, 1)-tight subgraph of G. This subgraph is purely periodic, since e_2 and e_3 have equal C_s -gain components, and hence every path from v_1 to v_2 has the same C_s -gain component. This is a contradiction to G satisfying the purely periodic condition.

This completes the proof, showing that one of $G - v_0 + e_{12}$ or $G - v_0 + e_{13}$ is $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight. \Box

Combining Lemmas 5.8, 5.10 and 5.12 shows that any vertex of degree 3 with exactly 2 neighbours admits a 1-reduction that preserves $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tightness. This completes the proof of Proposition 5.7.

5.3. 1-reductions on Vertices with Three Neighbours. For a degree 3 vertex, the last type of neighbourhood to consider is that where the vertex has 3 distinct neighbours.

Proposition 5.13. Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 3 with 3 distinct neighbours. Then it is possible to form a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph by a gained 1-reduction at v_0 .

Suppose that $v_0 \in V$ is a vertex of degree 3 with incident edges $e_1 = (v_0, v_1; m(e_1)), e_2 = (v_0, v_2; m(e_2))$ and $e_3 = (v_0, v_3; m(e_3))$, for some distinct $v_1, v_2, v_3 \in V$. After deleting v_0 , the possible options for edges to add for a 1-reduction are $e_{12} = (v_1, v_2; (m(e_1))^{-1}m(e_2)), e_{23} = (v_2, v_3; (m(e_2))^{-1}m(e_3))$ or $e_{31} = (v_3, v_1; (m(e_3))^{-1}m(e_1))$. Figure 5.3 illustrates the neighbourhood of v_0 , with the candidate edges e_{12}, e_{23} and e_{31} represented by dashed lines.




FIGURE 5.3. A vertex v_0 of degree 3 with three neighbours.

To prove Proposition 5.13, the aim is to show that one of $G-v_0+e_{12}$, $G-v_0+e_{23}$ or $G-v_0+e_{31}$ is $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight. Like for a vertex with 2 neighbours, only subgraphs containing the new edges need to be considered when investigating subgraphs that fail each condition. Again, the cases where each reduction fails each condition can be characterised in terms of the blockers described in Definition 5.9.

Lemmas 5.21 to 5.28 will show that any vertex of this type admits a 1-reduction to a $\mathbb{Z}^2 \rtimes C_s$ -tight graph, proving Proposition 5.13. As steps towards this, Lemmas 5.14 to 5.18 are combinatorial results that involve taking the unions and intersections of subgraphs with specific edge counts. Lemmas 5.19 and 5.20 discuss the gain spaces of unions of subgraphs with connected intersections. These results will be used later to examine the unions and intersections of blockers.

Lemma 5.14. [17, Theorem 5] Let $k, l \in \mathbb{N}$ with $0 < l \le k$. Let G be a (k, l)-tight multigraph with (k, l)-tight subgraphs $G_A = (V_A, E_A)$ and $G_B = (V_B, E_B)$ such that $|V_A \cap V_B| \ge 1$. Then $G_A \cup G_B$ and $G_A \cap G_B$ are both (k, l)-tight.

The following lemma is based on a result from [23, Lemma 4.4] regarding (2, 2)-tight multigraphs. It has been adapted here for (2, 1)-tight multigraphs. **Lemma 5.15.** Let G be a (2,1)-tight multigraph with a vertex $v_0 \in V$ of degree 3, with distinct neighbours $v_1, v_2, v_3 \in V$. Let $G - v_0$ be obtained from G by deleting v_0 with its incident edges. Then the following all hold:

- (1) There is no (2,1)-tight subgraph of $G v_0$ containing v_1, v_2 and v_3 .
- (2) If v_1 and v_2 are in a (2,1)-tight subgraph of $G v_0$, then neither the pair $\{v_1, v_3\}$ nor the pair $\{v_2, v_3\}$ is in a (2,1)-tight subgraph of $G v_0$.
- (3) If v_1 and v_2 are in a (2, 1)-tight subgraph of $G v_0$, then neither the pair $\{v_1, v_3\}$ nor the pair $\{v_2, v_3\}$ is in a (2, 2)-tight subgraph of $G v_0$.

Proof. If $G - v_0$ has a (2, 1)-tight subgraph containing all neighbours of v_0 , then adding v_0 with its incident edges to this subgraph would give an overcounted subgraph of G, breaking (2, 1)-sparsity. Hence, point 1 holds.

Suppose that v_1 and v_2 are in a (2, 1)-tight subgraph $G_{12} = (V_{12}, E_{12})$ of $G - v_0$ and that v_2 and v_3 are in a (2, 1)-tight subgraph $G_{23} = (V_{23}, E_{23})$ of $G - v_0$. By applying Lemma 5.14 on subgraphs of G, it follows that $G_{12} \cup G_{23}$ is a (2, 1)-tight subgraph of $G - v_0$ containing v_1, v_2 and v_3 . This contradicts point 1, so point 2 holds.

Suppose that v_1 and v_2 are in a (2, 1)-tight subgraph $G_{12} = (V_{12}, E_{12})$ of $G - v_0$ and that v_2 and v_3 are in a (2, 2)-tight subgraph $G_{23} = (V_{23}, E_{23})$ of $G - v_0$. By Lemma 5.2,

$$|E_{12} \cup E_{23}| + |E_{12} \cap E_{23}| = 2|V_{12} \cup V_{23}| + 2|V_{12} \cap V_{23}| - 3.$$
(5.4)

Since G_{23} is (2, 2)-tight, $|E_{12} \cap E_{23}| \le 2|V_{12} \cap V_{23}| - 2$. Combining this with Equation (5.4) shows that $|E_{12} \cup E_{23}| \ge 2|V_{12} \cup V_{23}| - 1$. This implies that $G_{12} \cup G_{23}$ is a (2, 1)-tight subgraph of $G - v_0$ containing v_1, v_2 and v_3 . This contradicts point 1, so point 3 holds.

The proofs of the next three lemmas are inspired by the version of [23, Lemma 4.2] given in the preprint of that article on ResearchGate. See also a similar method in [17, Theorem 5(1)].

Lemma 5.16. Let G be a (2,1)-tight multigraph that has a vertex v_0 of degree 3 with distinct neighbours $v_1, v_2, v_3 \in V$. Let $G_{12} = (V_{12}, E_{12})$ be a (2,3)-tight subgraph of G with $v_1, v_2 \in V_{12}$ and $v_0 \notin V_{12}$. Let $G_{23} = (V_{23}, E_{23})$ be a (2,3)-tight subgraph of G with $v_2, v_3 \in V_{23}$ and $v_0 \notin V_{23}$. Then one of the following holds:

- (1) $|V_{12} \cap V_{23}| > 1$ and $|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| 3$ and $|E_{12} \cap E_{23}| = 2|V_{12} \cap V_{23}| 3$,
- (2) $|V_{12} \cap V_{23}| > 1$ and $|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| 2$ and $|E_{12} \cap E_{23}| = 2|V_{12} \cap V_{23}| 4$,
- (3) $|V_{12} \cap V_{23}| = 1$ and $|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| 4$.

Proof. By Lemma 5.2,

$$|E_{12} \cup E_{23}| + |E_{12} \cap E_{23}| = 2|V_{12} \cup V_{23}| + 2|V_{12} \cap V_{23}| - 6.$$

$$(5.5)$$

If $|V_{12} \cap V_{23}| = 1$, then $|E_{12} \cap E_{23}| = 0$ and it clearly follows that $|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| - 4$.

Now suppose that $|V_{12} \cap V_{23}| > 1$. Since G_{12} and G_{23} are (2, 3)-tight, $|E_{12} \cap E_{23}| \le 2|V_{12} \cap V_{23}| - 3$. Using this with Equation (5.5) shows that $|E_{12} \cup E_{23}| \ge 2|V_{12} \cup V_{23}| - 3$. Also, Lemma 5.15(1) shows that $|E_{12} \cup E_{23}| \le 2|V_{12} \cup V_{23}| - 3$. Also, Lemma 5.15(1) shows that $|E_{12} \cup E_{23}| \le 2|V_{12} \cup V_{23}| - 2$. This gives the two possible edge counts for $G_{12} \cup G_{23}$. Substituting these into Equation (5.5) gives the corresponding edge counts for $G_{12} \cap G_{23}$. Hence, the lemma holds.

Lemma 5.17. Let G be a (2,1)-tight multigraph that has a vertex v_0 of degree 3 with distinct neighbours $v_1, v_2, v_3 \in V$. Let $G_{12} = (V_{12}, E_{12})$ be a subgraph of G that satisfies $|E_{12}| = 2|V_{12}| - 2$ with $v_1, v_2 \in V_{12}$ and $v_0 \notin V_{12}$. Let $G_{23} = (V_{23}, E_{23})$ be a (2, 2)-tight subgraph of G with $v_2, v_3 \in V_{23}$ and $v_0 \notin V_{23}$. Then

$$E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| - 2$$
 and $|E_{12} \cap E_{23}| = 2|V_{12} \cap V_{23}| - 2$.

Proof. By Lemma 5.2,

$$|E_{12} \cup E_{23}| + |E_{12} \cap E_{23}| = 2|V_{12} \cup V_{23}| + 2|V_{12} \cap V_{23}| - 4.$$
(5.6)

Since G_{23} is (2, 2)-tight, $|E_{12} \cap E_{23}| \le 2|V_{12} \cap V_{23}| - 2$. Also, Lemma 5.15(1) shows that $|E_{12} \cup E_{23}| \le 2|V_{12} \cup V_{23}| - 2$. Combining these bounds with Equation (5.6) shows that $|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| - 2$ and $|E_{12} \cap E_{23}| = 2|V_{12} \cap V_{23}| - 2$.

Lemma 5.18. Let G be a (2,1)-tight multigraph that has a vertex v_0 of degree 3 with distinct neighbours $v_1, v_2, v_3 \in V$. Let $G_{12} = (V_{12}, E_{12})$ be a subgraph of G that satisfies $|E_{12}| = 2|V_{12}| - 2$ with $v_1, v_2 \in V_{12}$ and $v_0 \notin V_{12}$. Let $G_{23} = (V_{23}, E_{23})$ be a (2,3)-tight subgraph of G with $v_2, v_3 \in V_{23}$ and $v_0 \notin V_{23}$. Suppose that $|V_{12} \cap V_{23}| > 1$. Then

$$|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| - 2$$
 and $|E_{12} \cap E_{23}| = 2|V_{12} \cap V_{23}| - 3$.

Proof. By Lemma 5.2,

$$|E_{12} \cup E_{23}| + |E_{12} \cap E_{23}| = 2|V_{12} \cup V_{23}| + 2|V_{12} \cap V_{23}| - 5.$$

$$(5.7)$$

Since G_{23} is (2,3)-tight and $|V_{12} \cap V_{23}| > 1$, we have $|E_{12} \cap E_{23}| \le 2|V_{12} \cap V_{23}| - 3$. Also, Lemma 5.15(1) shows that $|E_{12} \cup E_{23}| \le 2|V_{12} \cup V_{23}| - 2$. Combining these bounds with Equation (5.7) shows that $|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| - 2$ and $|E_{12} \cap E_{23}| = 2|V_{12} \cap V_{23}| - 3$.

The following lemma will be used later when considering pairs of balanced blockers.

Lemma 5.19. [10, Lemma 2.4] Let Γ be a group and let (G, m) be a Γ -gain graph with balanced subgraphs G_A and G_B . If $G_A \cap G_B$ is connected, then $G_A \cup G_B$ is balanced.

The following is a variation of Lemma 5.19 for purely periodic subgraphs of $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -gain graphs. The proof here is based on part of the proof of [23, Proposition 5.4], where an argument is described that proves a variant of Lemma 5.19 for \mathbb{Z}^2 -gain graphs.

Lemma 5.20. Let (G,m) be a $\mathbb{Z}^2 \rtimes C_s$ -gain graph with purely periodic subgraphs G_A and G_B . If $G_A \cap G_B$ is connected, then $G_A \cup G_B$ is purely periodic.

Proof. Suppose that $G_A \cap G_B$ is connected and consider a closed walk C in $G_A \cup G_B$. If C is entirely contained within one of G_A or G_B , then it is obvious that its net gain is in \mathbb{Z}^2 . Otherwise, suppose that C passes through both $G_A \setminus G_B$ and $G_B \setminus G_A$. Any such closed walk is formed of a composition of walks in G_A and walks in G_B , which meet at vertices in $G_A \cap G_B$. First, suppose that C starts at $x \in V_A \cap V_B$, follows a path P_1 through G_A with gain $m_1 \in \mathbb{Z}^2 \rtimes C_s$ to $y \in V_A \cap V_B$ and then follows a path P_2 through G_B with gain $m_2 \in \mathbb{Z}^2 \rtimes C_s$. See Figure 5.4.




FIGURE 5.4. An illustration of the paths through G_A and G_B .

Note that this closed walk has gain m_1m_2 . Since $G_A \cap G_B$ is connected, there is a path P_3 in $G_A \cap G_B$ from x to y. Suppose that this path has gain m_3 and insert it with its reverse into the middle of the closed walk C to obtain the following walk:

$$x \xrightarrow{m_1} y \xrightarrow{m_3^{-1}} x \xrightarrow{m_3} y \xrightarrow{m_2} x$$

The net gain of this augmented walk is $m_1m_3^{-1}m_3m_2 = m_1m_2$, which is the same as that of C. This walk can be split into two smaller closed walks: $P_1P_3^{-1}$ (with gain $m_1m_3^{-1}$) in G_A and P_3P_2 (with gain m_3m_2) in G_B . Since G_A and G_B are purely periodic, $m_1m_3^{-1} \in \mathbb{Z}^2$ and $m_3m_2 \in \mathbb{Z}^2$. Composing these shows that $m_1m_2 \in \mathbb{Z}^2$. Hence, the net gain of the closed walk C is in \mathbb{Z}^2 .

Inductively, this method can be used for any closed walk that moves between $G_A \setminus G_B$ and $G_B \setminus G_A$ multiple times. Each time the closed walk passes through a vertex in $G_A \cap G_B$, augment a path back to the starting vertex in $G_A \cap G_B$, along with its reverse. This shows that each closed walk in $G_A \cup G_B$ has the same gain as the composition of some sequence of closed walks that are each in one of G_A or G_B . Since G_A and G_B are purely periodic, each of these closed walks will

have a net gain in \mathbb{Z}^2 . The net gain on the full closed walk is the composition of the gains on these smaller closed walks, all of which are in \mathbb{Z}^2 . Hence, $G_A \cup G_B$ is purely periodic.

Now attention can be turned back to Proposition 5.13, aiming to show that, in a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph, any vertex of degree 3 with 3 distinct neighbours admits a 1-reduction to a smaller $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph. The overall approach is similar to that used to prove the corresponding result for \mathbb{Z}^2 -gain graphs [23, Proposition 5.4]. First, consider the (2, 1)-tight condition.

Lemma 5.21. Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 3 with edges e_1 , e_2 and e_3 from v_0 to distinct vertices v_1 , v_2 and v_3 respectively. Then at least two of $G - v_0 + e_{12}$, $G - v_0 + e_{23}$ or $G - v_0 + e_{31}$ are (2, 1)-tight.

Proof. By Lemma 5.15(2), at most one of the possible 1-reductions can have a (2, 1)-tight blocker. Hence, at least two of these graphs are (2, 1)-tight.

There are now 2 cases to consider: the case where all reductions give (2, 1)-tight graphs and the case where exactly one of the reductions gives a graph that is not (2, 1)-tight. We begin with the former case, where it remains to show that at least one of the reductions gives a graph that satisfies both the purely periodic condition and the balanced condition.

Lemma 5.22. Suppose that (G,m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 3 with edges e_1 , e_2 and e_3 from v_0 to distinct vertices v_1 , v_2 and v_3 respectively. Suppose that $G - v_0 + e_{12}$, $G - v_0 + e_{23}$ and $G - v_0 + e_{31}$ are all (2, 1)-tight. Then at least two of them satisfy the purely periodic condition.

Proof. Suppose that $G - v_0 + e_{12}$ and $G - v_0 + e_{23}$ both fail the purely periodic condition. This means that they have purely periodic blockers $G_{12} = (V_{12}, E_{12})$ and $G_{23} = (V_{23}, E_{23})$ respectively. Since purely periodic blockers are (2, 2)-tight, Lemma 5.17 shows that $|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| - 2$ and $|E_{12} \cap E_{23}| = 2|V_{12} \cap V_{23}| - 2$. Note that $G_{12} \cap G_{23}$ is (2, 2)-tight and thus connected by Lemma 5.3. Hence, Lemma 5.20 shows that $G_{12} \cup G_{23}$ is purely periodic. Moreover, adding v_0 with its incident edges to $G_{12} \cup G_{23}$ will give a purely periodic subgraph of G. To see this, observe that the C_s -gain components of paths between neighbours of v_0 that pass through v_0 will match those of other paths between the neighbours. This shows that any closed walk that passes through v_0 has a net gain in \mathbb{Z}^2 . Any closed walk that does not pass through v_0 is contained in $G_{12} \cup G_{23}$, which is purely periodic. Hence, $(G_{12} \cup G_{23}) + v_0$ is purely periodic. Since $(G_{12} \cup G_{23}) + v_0$ is (2, 1)-tight, this contradicts the fact that G satisfies the purely periodic condition. This contradiction can be reached for any pair of reductions. Hence, no more than one of the reductions gives a graph that fails the purely periodic condition.

Lemma 5.23. Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 3 with edges e_1 , e_2 and e_3 from v_0 to distinct vertices v_1 , v_2 and v_3 respectively. Suppose that $G - v_0 + e_{12}$, $G - v_0 + e_{23}$ and $G - v_0 + e_{31}$ are all (2, 1)-tight. Then at least one of them satisfies the balanced condition.

Proof. For contradiction, suppose that all of these graphs fail the balanced condition. This means that the reductions to $G - v_0 + e_{12}$, $G - v_0 + e_{23}$ and $G - v_0 + e_{31}$ have balanced blockers $G_{12} = (V_{12}, E_{12}), G_{23} = (V_{23}, E_{23})$ and $G_{31} = (V_{31}, E_{31})$ respectively.

To begin, consider the case where $|V_{12} \cap V_{23}| = |V_{23} \cap V_{31}| = |V_{31} \cap V_{12}| = 1$, so $|E_{12} \cap E_{23}| = |E_{23} \cap E_{31}| = |E_{31} \cap E_{12}| = 0$ and $|V_{12} \cap V_{23} \cap V_{31}| = 0$. In this case, consider the graph $G_{12} \cup G_{23} \cup G_{31}$. A basic counting argument shows that $|E_{12} \cup E_{23} \cup E_{31}| = 2|V_{12} \cup V_{23} \cup V_{31}| - 3$. We now show that $G_{12} \cup G_{23} \cup G_{31}$ is balanced. Any closed walk that is entirely contained within one of G_{12} , G_{23} or G_{31} clearly has trivial net gain. Since each intersection of a pair of balanced blockers has just one vertex, the only other possibility that needs to be considered for a closed walk in $G_{12} \cup G_{23} \cup G_{31}$ is a closed walk that passes through each of v_1 , v_2 and v_3 . Such a closed walk takes the form illustrated in Figure 5.5.




FIGURE 5.5. An illustration of the case where all pairs of balanced blockers intersect in exactly one vertex in the proof of Lemma 5.23.

Note that the net gain on this closed walk is trivial. Hence, $G_{12} \cup G_{23} \cup G_{31}$ is balanced and has a (2,3)-edge-count. Adding v_0 with its incident edges to $G_{12} \cup G_{23} \cup G_{31}$ gives a balanced subgraph of G with a (2,2)-edge-count, which contradicts the fact that G satisfies the balanced condition.

Now consider the case where at least one of the pairwise intersections of the blockers has more than one vertex. W.l.o.g., suppose that $|V_{12} \cap V_{23}| > 1$. Since balanced blockers are (2, 3)-tight, Lemma 5.16 gives the following possibilities for the edge counts of $G_{12} \cup G_{23}$ and $G_{12} \cap G_{23}$:

(1)
$$|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| - 3$$
 and $|E_{12} \cap E_{23}| = 2|V_{12} \cap V_{23}| - 3$

(2) $|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| - 2$ and $|E_{12} \cap E_{23}| = 2|V_{12} \cap V_{23}| - 4$.

In case 1, observe that $G_{12} \cap G_{23}$ is (2,3)-tight and thus connected by Lemma 5.3. Lemma 5.19 therefore shows that $G_{12} \cup G_{23}$ is balanced. Adding v_0 with its incident edges to $G_{12} \cup G_{23}$ then gives a balanced subgraph of G with a (2,2)-edge-count. This contradicts the fact that G satisfies the balanced condition.

In case 2, Lemma 5.11 shows that $G_{12} \cap G_{23}$ is either connected or consists only of 2 vertices and no edges. If $G_{12} \cap G_{23}$ is connected, then $G_{12} \cup G_{23}$ is balanced by Lemma 5.19. Since $G_{12} \cup G_{23}$ has a (2, 2)-edge-count, this contradicts the fact that G satisfies the balanced condition.

This leaves the case where $G_{12} \cap G_{23}$ is the graph with 2 vertices and no edges. If either of $G_{12} \cap G_{31}$ or $G_{23} \cap G_{31}$ contains at least one edge, then previous arguments can be applied on that pair of blockers to reach a contradiction. Hence, suppose that all of the pairwise intersections of the balanced blockers contain no edges. For ease, let $G^* = G_{12} \cup G_{23} = (V^*, E^*)$, so $|E^*| = 2|V^*| - 2$. Note that $G^* \cap G_{31} = (G_{12} \cap G_{31}) \cup (G_{23} \cap G_{31})$. Since both $G_{12} \cap G_{31}$ and $G_{23} \cap G_{31}$ contain no edges, $|E^* \cap E_{31}| = 0$. Since $v_1, v_3 \in V^* \cap V_{31}$, it is clear that $|V^* \cap V_{31}| > 1$. Hence, $|E^* \cap E_{31}| \neq 2|V^* \cap V_{31}| - 3$, which is a contradiction to Lemma 5.18.

Every case leads to a contradiction. Hence, one of the candidate reductions gives a graph that satisfies the balanced condition. $\hfill \Box$

In the case where every reduction gives a (2, 1)-tight graph, Lemma 5.22 shows that at least two of the reductions give graphs that satisfy the purely periodic condition and Lemma 5.23 shows that at least one gives a graph that satisfies the balanced condition. If two of the reductions give graphs that satisfy the balanced condition, then it is clear that at least one of these graphs also satisfies the purely periodic condition. It remains to consider the case where every reduction gives a (2, 1)-tight graph and only one reduction gives a graph that satisfies the balanced condition. In this case, it must be shown that this graph also satisfies the purely periodic condition. The following lemma covers this. **Lemma 5.24.** Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 3 with edges e_1 , e_2 and e_3 from v_0 to distinct vertices v_1 , v_2 and v_3 respectively. Suppose that $G - v_0 + e_{12}$, $G - v_0 + e_{23}$ and $G - v_0 + e_{31}$ are all (2, 1)-tight. Also, suppose that $G - v_0 + e_{12}$ and $G - v_0 + e_{23}$ both fail the balanced condition. Then $G - v_0 + e_{31}$ is $\mathbb{Z}^2 \rtimes C_s$ -tight.

Proof. By Lemma 5.23, the fact that $G - v_0 + e_{12}$ and $G - v_0 + e_{23}$ both fail the balanced condition implies that $G - v_0 + e_{31}$ satisfies the balanced condition. It therefore remains only to show that $G - v_0 + e_{31}$ satisfies the purely periodic condition.

For contradiction, suppose that $G - v_0 + e_{31}$ fails the purely periodic condition due to a purely periodic blocker G_{31} . Also, suppose that G_{12} and G_{23} are balanced blockers that cause $G - v_0 + e_{12}$ and $G - v_0 + e_{23}$ respectively to fail the balanced condition.

First, consider the case where $|V_{12} \cap V_{23}| = |V_{23} \cap V_{31}| = |V_{31} \cap V_{12}| = 1$, so $|E_{12} \cap E_{23}| = |E_{23} \cap E_{31}| = |E_{31} \cap E_{12}| = 0$ and $|V_{12} \cap V_{23} \cap V_{31}| = 0$. A basic counting argument shows that $|E_{12} \cup E_{23} \cup E_{31}| = 2|V_{12} \cup V_{23} \cup V_{31}| - 2$. By a similar method to that which was used in the first part of the proof of Lemma 5.23, it can then be shown that $G_{12} \cup G_{23} \cup G_{31}$ is purely periodic. Adding v_0 with its incident edges to $G_{12} \cup G_{23} \cup G_{31}$ gives a (2, 1)-tight purely periodic subgraph of G, which is a contradiction to the purely periodic condition.

Now consider the case where $|V_{12} \cap V_{31}| > 1$. Since G_{12} is (2, 3)-tight and G_{31} is (2, 2)-tight, Lemma 5.18 shows that $|E_{12} \cup E_{31}| = 2|V_{12} \cup V_{31}| - 2$ and $|E_{12} \cap E_{31}| = 2|V_{12} \cap V_{31}| - 3$. Since $G_{12} \cap G_{31}$ is (2, 3)-tight, combining Lemmas 5.3 and 5.20 shows that $G_{12} \cup G_{31}$ is purely periodic. Adding v_0 with its incident edges to $G_{12} \cup G_{31}$ gives a (2, 1)-tight purely periodic subgraph of G, which is a contradiction to the purely periodic condition. The same method also gives a contradiction in the case where $|V_{23} \cap V_{31}| > 1$.

The only case left to consider for this proof is that where $|V_{12} \cap V_{31}| = |V_{23} \cap V_{31}| = 1$ and $|V_{12} \cap V_{23}| > 1$. Given this, consider the edge count of $G_{12} \cup G_{23}$. Lemma 5.16 shows that one of the following holds:

- (1) $|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| 3$ and $|E_{12} \cap E_{23}| = 2|V_{12} \cap V_{23}| 3$,
- (2) $|E_{12} \cup E_{23}| = 2|V_{12} \cup V_{23}| 2$ and $|E_{12} \cap E_{23}| = 2|V_{12} \cap V_{23}| 4$.

In case 1, $G_{12} \cap G_{23}$ is (2,3)-tight, so combining Lemmas 5.3 and 5.19 shows that $G_{12} \cup G_{23}$ is balanced. Adding v_0 with its incident edges to $G_{12} \cup G_{23}$ gives a balanced subgraph of G with a (2,2)-edge-count. This contradicts the fact that G satisfies the balanced condition.

Now consider case 2. For ease of notation, let $G^* = G_{12} \cup G_{23} = (V^*, E^*)$, so $|E^*| = 2|V^*| - 2$. Since $V_{12} \cap V_{31} = \{v_1\}$ and $V_{23} \cap V_{31} = \{v_3\}$, it can be seen that $|V^* \cap V_{31}| = 2$ and $|E^* \cap E_{31}| = 0$. This means that $|E^* \cap E_{31}| \neq 2|V^* \cap V_{31}| - 2$, which contradicts Lemma 5.17.

Every case leads to a contradiction. Hence, $G - v_0 + e_{31}$ satisfies the purely periodic condition and is thus $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight.

For the proof of Proposition 5.13, combining Lemmas 5.22, 5.23 and 5.24 completes the case where every reduction gives a (2, 1)-tight graph, by showing that one of the reductions gives a graph that also satisfies the balanced condition and the purely periodic condition.

We now consider the case where one of the reductions gives a graph that is not (2, 1)-tight. The aim is to show that one of the other reductions gives a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph. We begin by considering the purely periodic condition.

Lemma 5.25. Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 3 with edges e_1 , e_2 and e_3 from v_0 to distinct vertices v_1 , v_2 and v_3 respectively. Suppose that $G - v_0 + e_{12}$ is not (2, 1)-tight. Then both $G - v_0 + e_{23}$ and $G - v_0 + e_{31}$ satisfy the purely periodic condition.

Proof. Since $G - v_0 + e_{12}$ is not (2, 1)-tight, it has a (2, 1)-tight blocker G_{12} . By Lemma 5.15(3), there are no purely periodic blockers for the reductions to $G - v_0 + e_{23}$ or $G - v_0 + e_{31}$. Hence, both of these graphs satisfy the purely periodic condition.

To complete the case where $G - v_0 + e_{12}$ is not (2, 1)-tight, it now remains only to show that one of $G - v_0 + e_{23}$ or $G - v_0 + e_{31}$ satisfies the balanced condition. In order to prove this, the following combinatorial lemmas will be needed. **Lemma 5.26.** Let $G_A = (V_A, E_A)$ and $G_B = (V_B, E_B)$ be (2,3)-sparse multigraphs with $|V_A \cap V_B| = 1$. Then $G_A \cup G_B$ is (2,3)-sparse.

Proof. Since G_A and G_B are (2,3)-sparse, it is clear that any subgraph of $G_A \cup G_B$ that has its edge set contained entirely within one of G_A or G_B is (2,3)-sparse. Now consider a subgraph $G'_A \cup G'_B \subseteq G_A \cup G_B$, where $G'_A = (V'_A, E'_A) \subseteq G_A$ and $G'_B = (V'_B, E'_B) \subseteq G_B$, both of which have at least one edge. Since G_A and G_B are (2,3)-sparse, $|E'_A| \leq 2|V'_A| - 3$ and $|E'_B| \leq 2|V'_B| - 3$. Using this, consider the edge count of $G'_A \cup G'_B$:

$$\begin{aligned} |E'_A \cup E'_B| + |E'_A \cap E'_B| &= |E'_A| + |E'_B| \\ &\leq (2|V'_A| - 3) + (2|V'_B| - 3) \\ &= 2|V'_A| + 2|V'_B| - 6 \\ &= 2|V'_A \cup V'_B| + 2|V'_A \cap V'_B| - 6 \end{aligned}$$

Note that $|V'_A \cap V'_B| \leq |V_A \cap V_B| = 1$ and $|E'_A \cap E'_B| \leq |E_A \cap E_B| = 0$. Hence, $|E'_A \cup E'_B| \leq 2|V'_A \cup V'_B| - 4$. This shows that $G_A \cup G_B$ is (2,3)-sparse.

Lemma 5.27. Let $G_A = (V_A, E_A)$ and $G_B = (V_B, E_B)$ be (2,3)-sparse multigraphs with $|V_A \cap V_B| = 2$ and $|E_A \cap E_B| = 0$. Then $G_A \cup G_B$ is (2,2)-sparse.

Proof. Again, it is clear that any subgraph of $G_A \cup G_B$ that has its edge set contained entirely within one of G_A or G_B is (2,3)-sparse. As before, consider a subgraph $G'_A \cup G'_B \subseteq G_A \cup G_B$, where $G'_A = (V'_A, E'_A) \subseteq G_A$ and $G'_B = (V'_B, E'_B) \subseteq G_B$, both of which have at least one edge. The same calculation from the proof of Lemma 5.26 gives

$$|E'_A \cup E'_B| + |E'_A \cap E'_B| \le 2|V'_A \cup V'_B| + 2|V'_A \cap V'_B| - 6.$$

Note that $|V'_A \cap V'_B| \leq |V_A \cap V_B| = 2$ and $|E'_A \cap E'_B| \leq |E_A \cap E_B| = 0$. Hence, $|E'_A \cup E'_B| \leq 2|V'_A \cup V'_B| - 2$. This shows that $G_A \cup G_B$ is (2,2)-sparse.

Now it is possible to complete the last step in the proof of Proposition 5.13. This involves showing that, when one of the 1-reductions gives a graph that is not (2, 1)-tight, one of the other 1-reductions gives a graph that satisfies the balanced condition.

Lemma 5.28. Suppose that (G, m) is a $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph that has a vertex v_0 of degree 3 with edges e_1 , e_2 and e_3 from v_0 to distinct vertices v_1 , v_2 and v_3 respectively. Suppose that $G - v_0 + e_{12}$ is not (2, 1)-tight. Then at least one of $G - v_0 + e_{23}$ or $G - v_0 + e_{31}$ satisfies the balanced condition.

Proof. Since $G - v_0 + e_{12}$ is not (2, 1)-tight, it has a (2, 1)-tight blocker G_{12} . For contradiction, suppose that both $G - v_0 + e_{23}$ and $G - v_0 + e_{31}$ fail the balanced condition. This would mean that their reductions have balanced blockers G_{23} and G_{31} respectively. Lemma 5.16 shows that one of the following holds:

- (1) $|V_{23} \cap V_{31}| > 1$ and $|E_{23} \cup E_{31}| = 2|V_{23} \cup V_{31}| 3$ and $|E_{23} \cap E_{31}| = 2|V_{23} \cap V_{31}| 3$,
- (2) $|V_{23} \cap V_{31}| > 1$ and $|E_{23} \cup E_{31}| = 2|V_{23} \cup V_{31}| 2$ and $|E_{23} \cap E_{31}| = 2|V_{23} \cap V_{31}| 4$,
- (3) $|V_{23} \cap V_{31}| = 1$ and $|E_{23} \cup E_{31}| = 2|V_{23} \cup V_{31}| 4$.

In case 1, $G_{23} \cap G_{31}$ is (2, 3)-tight. Lemmas 5.3 and 5.19 therefore show that $G_{23} \cup G_{31}$ is balanced. Adding v_0 with its incident edges to $G_{23} \cup G_{31}$ gives a balanced subgraph of G with a (2, 2)-edgecount, which contradicts the fact that G satisfies the balanced condition. Hence, case 1 is not possible.

Now consider case 2. By Lemma 5.11, $G_{23} \cap G_{31}$ is either connected or is formed of two vertices and no edges. If $G_{23} \cap G_{31}$ is connected, then Lemma 5.19 shows that $G_{23} \cup G_{31}$ is balanced, which breaks the balanced condition on G. If $G_{23} \cap G_{31}$ consists of two vertices and no edges, then Lemma 5.27 shows that $G_{23} \cup G_{31}$ is (2,2)-tight. However, this, combined with the existence of the (2,1)-tight blocker G_{12} , contradicts Lemma 5.15(3). Hence, case 2 is not possible.

Finally, consider case 3. For ease, let $G^* = G_{23} \cup G_{31} = (V^*, E^*)$, so $|E^*| = 2|V^*| - 4$. By Lemma 5.2,

$$|E_{12} \cup E^*| + |E_{12} \cap E^*| = 2|V_{12} \cup V^*| + 2|V_{12} \cap V^*| - 5.$$
(5.8)

Note that $|V_{12} \cap V^*| > 1$, as $v_1, v_2 \in V_{12} \cap V^*$. By Lemma 5.26, G^* is (2, 3)-sparse, so $|E_{12} \cap E^*| \le 2|V_{12} \cap V^*| - 3$. By Lemma 5.15(1), we have that $|E_{12} \cup E^*| \le 2|V_{12} \cup V^*| - 2$. Applying these bounds to Equation (5.8) shows that $|E_{12} \cup E^*| = 2|V_{12} \cup V^*| - 2$ and $|E_{12} \cap E^*| = 2|V_{12} \cap V^*| - 3$.

Note that $G_{12} \cap G^*$ is (2, 3)-tight and thus connected by Lemma 5.3. This means that $G_{12} \cap G^*$ contains a path from v_1 to v_2 . Note that $v_1 \in V_{12} \cap V_{31}$ and $v_2 \in V_{12} \cap V_{23}$. Since $V_{23} \cap V_{31} = \{v_3\}$, the path from v_1 to v_2 must pass through v_3 . Hence, $v_3 \in G_{12} \cap G^*$. However, this means that G_{12} contains all neighbours of v_0 . Since G_{12} is (2, 1)-tight, this contradicts Lemma 5.15(1). Hence, case 3 is not possible.

Every case leads to a contradiction. Hence, one of the reductions will give a graph that satisfies the balanced condition. $\hfill \Box$

All cases have now been covered to complete the proof of Proposition 5.13, showing that, in a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph, every vertex of degree 3 with 3 neighbours admits a 1-reduction to a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph.

Proof of Proposition 5.13. Lemma 5.21 shows that no more than one of the possible 1-reductions can give a graph that is not (2, 1)-tight. In the case where all reductions give (2, 1)-tight graphs, combining Lemmas 5.22, 5.23 and 5.24 shows that one of the reductions gives a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph. In the case where exactly one of the reductions gives a graph that is not (2, 1)-tight, combining Lemmas 5.25 and 5.28 shows that one of the other reductions gives a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph. All cases have been covered, completing the proof of Proposition 5.13.

This then completes the proof of Theorem 5.1, showing that every $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph can be obtained from a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph on K_1^1 by a sequence of extensions.

Proof of Theorem 5.1. It is clear that gained 0-extensions, 1-extensions and loop-1-extensions always preserve $\mathbb{Z}^2 \rtimes C_s$ -tightness.

Let (G, m) be a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph. Basic counting arguments on (2, 1)-tight graphs show that G has a vertex of degree 2 or 3. If G has a vertex of degree 2, then Proposition 5.4 shows that (G, m) admits a 0-reduction to a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph. If G has a vertex of degree 3 with an incident loop, then Proposition 5.5 shows that (G, m) admits a loop-1-reduction to a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph. If G has a vertex of degree 3 without an incident loop, then, depending on the number of distinct neighbours, Proposition 5.6, 5.7 or 5.13 shows that (G, m) admits a 1-reduction to a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph. Hence, every $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph admits a reduction to a smaller $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph.

Since K_1^1 is the only (2, 1)-tight graph on a single vertex, repeatedly applying these reductions will eventually lead to a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph on K_1^1 . Reverse this sequence of reductions to get the required sequence of extensions to complete the proof.

Now the proof of Theorem 3.2 can be completed, showing that every $\mathbb{Z}^2 \rtimes C_s$ -tight gain graph is minimally rigid.

Proof of sufficiency for Theorem 3.2. Every $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph on K_1^1 is minimally rigid. Theorem 5.1 states that every $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph can be reached from a $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph on K_1^1 by a sequence of 0-extensions, 1-extensions and loop-1-extensions. Propositions 4.2, 4.3 and 4.4 show that each of these extensions preserves minimal rigidity, so any $\mathbb{Z}^2 \rtimes \mathcal{C}_s$ -tight gain graph is minimally rigid.

The wallpaper groups cm and pg can be viewed as subgroups of the group pm. Hence, the proof of Theorem 3.2 may be applied almost directly to characterise minimal symmetric rigidity with respect to these groups.

Theorem 5.29. Let $\Gamma \in \{pm, cm, pg\}$. A Γ -gain graph (G, m) is minimally rigid if and only if it satisfies all of the following conditions:

- (1) (2,1)-tight Condition: G is (2,1)-tight;
- (2) Purely Periodic Condition: Every purely periodic subgraph of G is (2, 2)-sparse;
- (3) Balanced Condition: Every balanced subgraph of G is (2,3)-sparse.

Proof. For pm, this result is Theorem 3.2.

The group pg can be thought of as the subgroup of pm generated by (0, 1, 0) and (1, 0, s). It also admits exactly a 1-dimensional space of trivial pg-symmetric infinitesimal motions. Hence, the proof of Lemma 3.3 also proves necessity for the group pg. For sufficiency, we can follow mostly the same method used for Theorem 3.2 in this paper. The only significant difference is in proving that 1-extensions preserve minimal rigidity. In Lemma 4.3, a special case was described where a 1-extension is performed on a loop to add a triple of parallel edges in such a way that the derived neighbours of the new vertex are collinear for all configurations, which occurs if and only if the new edges all have the same horizontal-translation-gain component. This does not occur for pg, as it is required that two of the new edges have different C_s -gain components. This prevents them from having the same horizontal-translation-gain component. Hence, the usual method can be used to show that all 1-extensions preserve minimal rigidity of pg-gain graphs.

The group cm can be thought of as the subgroup of pm generated by (1, 1, 0), (1, -1, 0) and (0, 0, s). Again, the proof of necessity follows by exactly the same argument used for Lemma 3.3. Likewise, sufficiency can be proved by exactly the same method used for Theorem 3.2.

6. Further work

6.1. Other wallpaper groups. There are a number of other wallpaper groups for which there is still no known characterisation of conditions for symmetric rigidity. In the Hermann-Maguin notation, these are pmm, cmm, p4m, p4g, p6m, pmg, pgg, p3m1 and p31m [26]. As the groups pmm, cmm, p4m, p4g and p6m all contain even dihedral subgroups, the Bottema's Mechanism gain graph [29] shows that the sparsity conditions analogous to Theorem 3.2 are not sufficient for these groups. It is not known whether the sparsity conditions are sufficient for pmg, pgg, p3m1 or p31m.

Although D. Bernstein used tropical geometry to characterise conditions for forced symmetric rigidity with respect to p2, p3, p4 and p6 [3], this approach does not provide an inductive construction of the minimally rigid gain graphs. Finding such an inductive construction is likely to be challenging, as minimally rigid gain graphs with these groups would need to be (2, 0)-tight and inductive constructions of (2, 0)-tight graphs are generally quite difficult, since they may not have any vertices of degree less than 4. However, an inductive construction of (2, 0)-tight graphs was used to characterise conditions for "odd" dihedral-symmetric rigidity in [10]. See also [12], which dealt with (2, 0)-tight graphs in the context of symmetric rigidity in non-Euclidean normed planes. It may be possible to modify these approaches so that they can be applied to wallpaper groups.

6.2. Flexible lattice representations. An immediate question is how to extend the results of this paper to the fully flexible lattice representation, or at least to partially flexible lattice representations. Conditions for minimal rigidity of a gain graph for groups p1, p2, p3 p4 and p6 on a flexible lattice in the Euclidean plane were characterised in [18, Theorem A] and [19, Theorem 1], although these methods do not involve an inductive construction. To do this for any wallpaper group using an inductive method is again likely to be challenging, as minimal rigidity on the flexible or partially flexible lattice requires an overall edge count of (2, 0) or even higher.

For a partially flexible lattice with the group p1, an inductive construction for minimally rigid gain graphs was found in [21, Theorem 2]. The counts would make it difficult to apply this to any other wallpaper group.

6.3. Higher dimensions. A combinatorial characterisation of generic rigid (finite) frameworks is not known for dimension 3 and higher. Thus, there are also no characterisations of forced symmetric rigidity for bar-joint frameworks in dimension $d \ge 3$, for any symmetry group. However, for the special class of body-bar frameworks in *d*-space, a complete characterisation of symmetry forced generic rigidity was obtained by S. Tanigawa for all crystallographic groups (for the fixed lattice representation). See Theorem 7.2 in [33]. Extensions of this result to flexible lattice representations, or to periodic body-hinge or molecular frameworks are not known.

6.4. Non-Euclidean spaces. There has recently been a surge of interest in the rigidity of frameworks in non-Euclidean normed spaces, see for example [13, 11, 14, 15, 16, 12, 5]. Combinatorial characterisations for forced symmetric rigidity in the plane have so far only been obtained for finite frameworks with reflection symmetry [16] and half-turn symmetry [12] for the ℓ^1 and ℓ^∞ norms.

Analogous results for finite frameworks with other symmetries or other normed planes, or for infinite periodic or crystallographic frameworks with forced symmetry (for fixed or flexible lattice representations) have not been established yet. We will address some of these problems in the companion paper [9].

References

- T. G. Abbott. Generalizations of Kempe's universality theorem, 2008. Master's thesis, Massachusetts Institute of Technology.
- [2] L. Asimow and B. Roth. The rigidity of graphs. II. J. Math. Anal. Appl., 68(1):171-190, 1979.
- [3] D. I. Bernstein. Generic symmetry-forced infinitesimal rigidity: translations and rotations. SIAM J. Appl. Algebra Geom., 6(2):190-215, 2022.
- [4] C. S. Borcea and I. Streinu. Periodic frameworks and flexibility. Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci., 466(2121):2633-2649, 2010.
- [5] K. Clinch and D. Kitson. Constructing isostatic frameworks for the ℓ¹ and ℓ[∞] plane. Electron. J. Combin., 27(2):Paper No. 2.49, 21, 2020.
- [6] R. Connelly and S. D. Guest. Frameworks, Tensegrities, and Symmetry. Cambridge University Press, 2022.
- [7] J. H. Conway, H. Burgiel, and C. Goodman-Strauss. The symmetries of things. A K Peters, Ltd., Wellesley, MA, 2008.
- [8] S. Dewar, G. Grasegger, E. Kastis, and A. Nixon. Quotient graphs of symmetrically rigid frameworks. Doc. Math., 29(3):561–595, 2024.
- [9] J. Esson, E. Kastis, and B. Schulze. Periodic and crystallographic rigidity in non-Euclidean normed planes. preprint. 2025.
- [10] T. Jordán, V. E. Kaszanitzky, and S. Tanigawa. Gain-sparsity and symmetry-forced rigidity in the plane. Discrete Comput. Geom., 55(2):314–372, 2016.
- [11] D. Kitson. Rigidity with polyhedral norms. In Handbook of geometric constraint systems principles, Discrete Math. Appl. (Boca Raton), pages 525–541. CRC Press, Boca Raton, FL, 2019.
- [12] D. Kitson, A. Nixon, and B. Schulze. Rigidity of symmetric frameworks in normed spaces. *Linear Algebra Appl.*, 607:231–285, 2020.
- [13] D. Kitson and S. C. Power. Infinitesimal rigidity for non-Euclidean bar-joint frameworks. Bull. Lond. Math. Soc., 46(4):685–697, 2014.
- [14] D. Kitson and B. Schulze. Maxwell-Laman counts for bar-joint frameworks in normed spaces. Linear Algebra Appl., 481:313–329, 2015.
- [15] D. Kitson and B. Schulze. Symmetric isostatic frameworks with ℓ^1 or ℓ^{∞} distance constraints. *Electron. J. Combin.*, 23(4):Paper 4.23, 23, 2016.
- [16] D. Kitson and B. Schulze. Motions of grid-like reflection frameworks. J. Symbolic Comput., 88:47-66, 2018.
- [17] A. Lee and I. Streinu. Pebble game algorithms and sparse graphs. Discrete Math., 308(8):1425–1437, 2008.
- [18] J. Malestein and L. Theran. Generic combinatorial rigidity of periodic frameworks. Adv. Math., 233:291–331, 2013.
- [19] J. Malestein and L. Theran. Frameworks with forced symmetry II: orientation-preserving crystallographic groups. Geom. Dedicata, 170:219–262, 2014.
- [20] J. Malestein and L. Theran. Generic rigidity with forced symmetry and sparse colored graphs. In Rigidity and symmetry, volume 70 of Fields Inst. Commun., pages 227–252. Springer, New York, 2014.
- [21] A. Nixon and E. Ross. Periodic rigidity on a variable torus using inductive constructions. Electron. J. Combin., 22(1):Paper 1.1, 30, 2015.
- [22] E. Ross. The rigidity of periodic frameworks as graphs on a fixed torus. Contrib. Discrete Math., 9(1):11–45, 2014.
- [23] E. Ross. Inductive constructions for frameworks on a two-dimensional fixed torus. Discrete Comput. Geom., 54(1):78–109, 2015.
- [24] E. Ross, B. Schulze, and W. Whiteley. Finite motions from periodic frameworks with added symmetry. International Journal of Solids and Structures, 48(11):1711–1729, 2011.
- [25] D. E. Sands. Introduction to Crystallography. Dover Publications Inc, 2003.
- [26] D. Schattschneider. The plane symmetry groups: Their recognition and notation. The American Mathematical Monthly, 85(6):439–450, 1978.
- [27] B. Schulze. Block-diagonalised rigidity matrices of symmetric frameworks and applications. Contributions to Algebra and Geometry, 51:427–466, 2010.
- [28] B. Schulze. Combinatorial rigidity of symmetric and periodic frameworks. In M. Sitharam, A. St. John, and J. Sidman, editors, *Handbook of Geometric Constraint Systems Principles*. CRC Press, 2019.
- [29] B. Schulze and W. Whiteley. The orbit rigidity matrix of a symmetric framework. Discrete Comput. Geom., 46(3):561–598, 2011.
- [30] B. Schulze and W. Whiteley. Rigidity and scene analysis. In Handbook of Discrete and Computational Geometry, Third Edition, C.D. Toth, J. O'Rourke, J.E. Goodman, editors. Chapman & Hall CRC, 2018.
- [31] B. Schulze and W. Whiteley. Rigidity of symmetric frameworks. In Handbook of Discrete and Computational Geometry, Third Edition, C.D. Toth, J. O'Rourke, J.E. Goodman, editors. Chapman & Hall CRC, 2018.
- [32] A. Szczepanski. Geometry of crystallographic groups, volume 4 of Algebra and discrete mathematics. World Scientific Publishing, 2012.

- [33] S. Tanigawa. Matroids of gain graphs in applied discrete geometry. Trans. Amer. Math. Soc., 367(12):8597– 8641, 2015.
- [34] W. Whiteley. Some matroids from discrete applied geometry. In Matroid theory (Seattle, WA, 1995), volume 197 of Contemp. Math., pages 171–311. Amer. Math. Soc., Providence, RI, 1996.

School of Mathematical Sciences, Lancaster University, Lancaster LA1 4YF, United Kingdom *Email address:* j.esson@lancaster.ac.uk

School of Mathematical Sciences, Lancaster University, Lancaster LA1 4YF, United Kingdom *Email address:* 1.kastis@lancaster.ac.uk

School of Mathematical Sciences, Lancaster University, Lancaster LA1 4YF, United Kingdom *Email address:* b.schulze@lancaster.ac.uk