# arXiv:2502.14530v1 [math.AG] 20 Feb 2025

# PROJECTIVITY OF THE MODULI OF HIGHER RANK PT-STABLE PAIRS ON A THREEFOLD

MIHAI PAVEL AND TUOMAS TAJAKKA

ABSTRACT. We construct a globally generated line bundle on the moduli stack of higher-rank PT-semistable objects over a smooth projective threefold and analyze the extent to which it separates points. Furthermore, when the rank and degree are coprime, we refine our construction to obtain an explicit ample line bundle on the corresponding coarse moduli space of PT-stable objects, thereby establishing its projectivity.

### 1. INTRODUCTION

The study of moduli spaces of stable pairs on projective threefolds traces back to the work of Pandharipande and Thomas [PT09], where they were introduced to define new enumerative invariants of curves on projective Calabi-Yau threefolds. By definition, stable pairs on a smooth projective threefold X are 2-term complexes of the form

$$\mathcal{O}_X \xrightarrow{s} F$$

where F is a pure sheaf of dimension 1 and the cokernel of s is 0-dimensional. As shown in [PT09], such objects are parametrized by a fine moduli space PT(X), representing a distinguished component of the moduli stack of complexes with trivial determinant in  $D^{b}(X)$  [Lie06] that enjoys particularly nice geometric properties. The moduli space PT(X) arises as a GIT quotient of coherent systems, constructed by Le Potier [LP93], which endows PT(X) with the structure of a projective scheme. Additionally, PT-pairs admit a well-behaved obstruction theory, which allows for the definition of curve-counting invariants, known as *Pandharipande-Thomas (PT) invariants*, when X is Calabi-Yau. This raised further questions about the relationship between PT invariants and other established enumerative theories, most notably Donaldson-Thomas (DT) theory, eventually leading to the formulation of the DT/PT correspondence.

The PT-stable pairs can be understood through certain stability conditions on the derived category of coherent sheaves on X, which arise as "large volume limits" of Bridgeland stability conditions. This connection was independently observed by Bayer [Bay09] and Toda [Tod09], who studied the DT/PT correspondence via wall-crossing within their family of stability conditions. We recall the notion of PTstability in Section 2, in a framework that recovers the stable pairs of Pandharipande and Thomas as stable objects of rank 1 and trivial determinant.

Moduli spaces of higher-rank PT-pairs provide alternative compactifications of the moduli of  $\mu$ -stable reflexive sheaves. Indeed, the moduli theory of PT-semistable objects was developed by J. Lo in [Lo11, Lo13], who showed that the moduli stack of PT-semistable objects with fixed Chern character is a universally closed algebraic stack of finite type, which moreover admits a proper coarse moduli space in the absence of strictly semistable objects. However, unlike the rank 1 and trivial determinant case considered by Pandharipande and Thomas, the moduli problem for higher-rank PT-pairs is not known to be associated with a GIT problem, and hence it is unknown whether the moduli spaces are projective. The primary goal of this paper is to answer this question affirmatively in the case of coprime rank and degree, thereby confirming a result expected in the field (see [Tod09, Remark 3.22]). We summarize our main results in the following.

**Theorem 1.1.** Let (X, H) be a polarized, smooth, projective threefold over  $\mathbb{C}$ , let  $v \in K_{num}(X)$  be a class of positive rank, and let Z be a PT-stability function on (X, H).

- (a) There is a globally generated line bundle on the moduli stack  $\mathcal{M}_Z^{\mathrm{PT}}(v)$  of PT-semistable objects of class v on X whose sections separate  $\mu$ -stable reflexive sheaves.
- (b) Furthermore, if rk(v) and  $H^2 ch_1(v)$  are coprime, and  $U = td_X$  in the definition of Z, then the coarse moduli space  $M_Z^{PT}(v)$  of PT-stable objects is projective.

In the absence of GIT, we employ stack-theoretic techniques to analyze the positivity of explicit determinantal line bundles on the stack  $\mathcal{M}_Z^{\mathrm{PT}}(v)$ . We prove in Theorem 5.8 the existence of a semiample line bundle  $\mathcal{L}_2$  on  $\mathcal{M}_Z^{\mathrm{PT}}(v)$ , and then describe in Theorem 5.9 the extent to which its sections separate points, showing part (a). We then proceed to improve the positivity of  $\mathcal{L}_2$  by twisting it with two additional determinantal line bundles  $\mathcal{L}_1$  and  $\mathcal{L}_0$ , in a manner analogous to a construction in the classical theory of Gieseker moduli spaces [HL10, Section 8]. Indeed, when the rank and degree are coprime, we show in Theorem 5.21 that a positive linear combination of  $\mathcal{L}_2$ ,  $\mathcal{L}_1$ , and  $\mathcal{L}_0$  descends to an *ample* line bundle on the coarse moduli space  $M_Z^{\mathrm{PT}}(v)$  of PT-stable objects, thus establishing part (b).

Our approach is based on two key ingredients: Mehta-Ramanathan type restriction theorems [MR82, MR84], and an embedding result due to Beentjes and Ricolfi [BR21, Proposition 5.5]. We use the first one to generate sections of our determinantal line bundles  $\mathcal{L}_2$  and  $\mathcal{L}_1$  through a technique that involves restricting PT-semistable objects to various curves and surfaces in X, respectively; see Section 5.1 and Section 5.3. This method originates from the works of Le Potier [Le 92] and J. Li [Li93], who employed it to construct an algebro-geometric analogue of the Donaldson-Uhlenbeck compactification. We refer to [Taj23] for an application of this idea in the context of Bridgeland stability conditions on surfaces.

Our proof of the projectivity result relies crucially on the coprime assumption, which implies three useful properties. First, it ensures that there are no strictly PT-semistable objects, which in particular guarantees the existence of the proper coarse moduli space  $M_Z^{PT}(v)$ . Second, it allows for a concrete characterization of PT-stable objects due to J. Lo, see Proposition 2.3, which is helpful for describing certain strata of  $M_Z^{PT}(v)$  [Lo21]. Third, under the coprime assumption, Beentjes and Ricolfi constructed a stratification of the moduli space of PT-stable objects by relative Quot schemes, see [BR21, Proposition 5.5 and Appendix B]. This last property becomes important in Section 5.4, where we use it to analyze the positivity of  $\mathcal{L}_0$  and finally establish our main ampleness result.

1.1. Acknowledgements. M.P. would like to thank Daniel Greb, Jochen Heinloth, Emanuele Macrì, Cameron Ruether, Saket Shah, Andrei Stoenică and Matei Toma for helpful conversations and insightful suggestions related to the project. M.P. was partly supported by the PNRR grant CF 44/14.11.2022 *Cohomological Hall algebras of smooth surfaces and applications*, and also partially supported by a grant of the Ministry of Research, Innovation and Digitalization, CNCS - UEFIS-CDI, project number PN-IV-P2-2.1-TE-2023-2040, within PNCDI IV.

T.T. would like to thank Jarod Alper, Aaron Bertram, Jochen Heinloth, Nicholas Kuhn, Jason Lo, and Siddharth Mathur for insightful conversations. T.T. was partially supported by the Knut and Alice Wallenberg foundation under grant no. KAW 2019.0493, as well as enjoyed the hospitality of Institut Mittag-Leffler during the program Moduli and Algebraic Cycles in Autumn 2021. Some of the material in this article appeared in T.T.'s doctoral dissertation at the University of Washington [Taj21].

### 2. PT-SEMISTABLE OBJECTS

In this section, we recall the definition and basic properties of PT-stability conditions. They are examples of polynomial stability conditions defined in [Bay09] as a generalization of Bridgeland stability conditions in order to understand the large volume limit of Bridgeland stability, as well as to study relations between various curve counting invariants. We largely follow [Lo11] and [Lo13] in our presentation, except that we use a slightly different convention for the category of perverse sheaves that appears as the heart of a PT-stability condition.

Let (X, H) be a smooth, projective, polarized threefold over  $\mathbb{C}$ , where  $H \subset X$  is a very ample divisor. We define the *heart of perverse sheaves*  $\mathcal{A}^p(X) \subseteq D^{\mathrm{b}}(X)$  as the full subcategory consisting of objects  $E \in D^{\mathrm{b}}(X)$  that fit in an exact triangle

$$F \to E \to T[-1],$$

where F and T are coherent sheaves on X such that  $\dim(T) \leq 1$  and  $\dim(F') \geq 2$ for every coherent subsheaf  $F' \subseteq F$ . The category  $\mathcal{A}^p(X)$  is the heart of a bounded t-structure on  $D^{\mathrm{b}}(X)$  obtained by tilting along a certain torsion pair.

In the following, we let  $\phi(z) \in (0, \pi]$  denote the phase of a complex number z in the extended upper half-plane

$$\mathbb{H} = \{ z \in \mathbb{C} \mid \operatorname{Im} z > 0 \text{ or } \operatorname{Re} z < 0 \}.$$

**Definition 2.1.** A **PT**-stability condition on X consists of the data of the heart  $\mathcal{A}^p(X)$  together with a group homomorphism  $Z : \mathrm{K}_{\mathrm{num}}(X) \to \mathbb{C}[m]$ , called the central charge, of the form

$$Z(E)(m) = \sum_{d=0}^{3} \rho_d \left( \int_X H^d \cdot \operatorname{ch}(E) \cdot U \right) m^d,$$

where

(a) the  $\rho_d \in \mathbb{C}^*$  are nonzero complex numbers such that  $-\rho_0, -\rho_1, \rho_2, \rho_3 \in \mathbb{H}$ , and whose phases satisfy

 $\phi(\rho_2) > \phi(-\rho_0) > \phi(\rho_3) > \phi(-\rho_1).$ 

(b)  $U = 1 + U_1 + U_2 + U_3 \in A^*(X)$  is a class with  $U_i \in A^i(X)$  for i = 1, 2, 3.

The configuration of the complex numbers  $\rho_i$  is compatible with the heart  $\mathcal{A}^p(X)$ in the sense that for any nonzero  $E \in \mathcal{A}^p(X)$ , we have  $Z(E)(m) \in \mathbb{H}$  for  $m \gg 0$ . This allows us to define a notion of stability on  $\mathcal{A}^p(X)$ : an object  $E \in \mathcal{A}^p(X)$  is called Z-stable (resp. Z-semistable) if for every proper nonzero subobject  $F \subseteq E$ , we have

 $\phi(Z(F)(m)) < \phi(Z(E)(m) \quad (\text{resp. } \phi(Z(F)(m)) \leq \phi(Z(E)(m)) \quad \text{for} \quad m \gg 0.$ 

Remark 2.2. Our definition of the heart  $\mathcal{A}^p(X)$  differs from that in [Lo11], [Lo13], and [Bay09] by a shift: the nonzero cohomology sheaves are in degrees 0 and 1 rather than -1 and 0. To account for this, our definition of the charge Z also differs in that  $-\rho_0, -\rho_1, \rho_2, \rho_3$ , rather than  $\rho_0, \rho_1, -\rho_2, -\rho_3$ , are in  $\mathbb{H}$ . The reason for this choice is purely psychological: if  $E \in \operatorname{Coh}(X)$  is a torsion-free sheaf, then E, rather than E[1], is contained in  $\mathcal{A}^p(X)$ . This will let us view the moduli of PT-semistable objects as an enlargement of the moduli of  $\mu$ -stable reflexive sheaves without having to perform a shift.

In [PT09], Pandharipande and Thomas define a *stable pair* on X to be a map of the form

$$\mathcal{O}_X \xrightarrow{s} F$$
,

where F is a sheaf of pure dimension 1 and the cokernel of s is 0-dimensional. In [Bay09, Proposition 6.1.1], Bayer shows that for any PT-stability condition, the stable objects in  $\mathcal{A}^{p}(X)$  with numerical invariants ch =  $(1, 0, -\beta, -n)$  and trivial determinant coincide precisely with these stable pairs. The following partial characterization of PT-semistable objects generalizes this fact to higher rank. We refer to [HL10, Section 1.6] for the classical notion of  $\mu$ -semistability and the notion of semistability on  $\operatorname{Coh}_{3,1}(X)$  appearing in the statement, both considered here with respect to the polarization H.

**Proposition 2.3** ([Lo11, Lemma 3.3], [Lo13, Proposition 2.24]). If  $v \in K_{num}(X)$  is a class of rank rk(v) > 0, then any PT-semistable object  $E \in \mathcal{A}^p(X)$  of class v satisfies the following conditions:

- (i)  $\mathcal{H}^0(E)$  is torsion-free and semistable in  $\operatorname{Coh}_{3,1}(X)$ ,
- (ii)  $\mathcal{H}^1(E)$  is 0-dimensional,

(iii)  $\operatorname{Hom}_{D^{b}(X)}(T[-1], E) = 0$  for any 0-dimensional sheaf T.

In particular  $\mathcal{H}^0(E)$  is also  $\mu$ -semistable. If moreover  $\operatorname{rk}(v)$  and  $H^2 \cdot \operatorname{ch}_1(v)$  are coprime, then any object E of class v in  $\mathcal{A}^p(X)$  satisfying these conditions is PT-stable,  $\mathcal{H}^0(E)$  is  $\mu$ -stable, and there are no strictly semistable objects.

We also note that for any PT-semistable object E of positive rank, the torsionfree sheaf  $\mathcal{H}^0(E)$  has homological dimension at most 1, cf. [Lo21, Example 5.5], a fact which will become important later in Proposition 5.19.

We will need the following observation.

**Lemma 2.4.** Let  $E \in \mathcal{A}^p(X)$  be a *PT*-semistable object with respect to a charge  $Z : \mathrm{K}_{\mathrm{num}}(X) \to \mathbb{C}[m]$  as in Definition 2.1, and assume  $\mathrm{rk}(E) > 0$ . If  $F \subseteq E$  is a subobject in  $\mathcal{A}^p(X)$  such that

$$\phi(Z(F)(m)) = \phi(Z(E)(m)) \quad for \ m \gg 0,$$

then  $\operatorname{rk}(F) > 0$ .

*Proof.* Let Q denote the cokernel of the inclusion  $F \subseteq E$  in  $\mathcal{A}^p(X)$ , so that we have a short exact sequence

$$0 \to F \to E \to Q \to 0$$

in  $\mathcal{A}^p(X)$ . This induces an exact sequence

$$0 \to \mathcal{H}^0(F) \to \mathcal{H}^0(E) \to \mathcal{H}^0(Q) \to \mathcal{H}^1(F) \to \mathcal{H}^1(E) \to \mathcal{H}^1(Q) \to 0$$

in Coh(X). Since by Proposition 2.3, the sheaf  $\mathcal{H}^0(E)$  is torsion-free, if  $\operatorname{rk}(F) = 0$ , then F = F'[-1], where  $F' = \mathcal{H}^1(F)$  is a coherent sheaf with  $\dim(\operatorname{Supp}(F')) \leq 1$ . If  $\dim(\operatorname{Supp}(F')) = 1$ , then

$$\lim_{m \to \infty} \phi(Z(F)(m)) = \phi(\rho_1) < \phi(-\rho_3) = \lim_{m \to \infty} \phi(Z(E)(m))$$

Similarly, if  $\dim(\operatorname{Supp}(F')) = 0$ , then

$$\lim_{m \to \infty} \phi(Z(F)(m)) = \phi(\rho_0) > \phi(-\rho_3) = \lim_{m \to \infty} \phi(Z(E)(m)).$$

In neither case can we have  $\phi(Z(F)(m)) = \phi(Z(E)(m))$  for  $m \gg 0$ .

2.1. Moduli spaces of PT-semistable objects. The theory of moduli of PT-semistable objects was developed by Lo in [Lo11] and [Lo13], culminating in [Lo13, Theorem 1.1], where the author constructs the moduli stack of PT-semistable objects of fixed Chern character as a universally closed algebraic stack of finite type, and, in the absence of strictly semistable objects, as a proper algebraic space.

Here we revisit these results through the framework of good moduli spaces, developed by Alper in [Alp08]. To recall the definition, let  $\mathcal{M}$  be an algebraic stack and  $\pi : \mathcal{M} \to M$  a quasi-compact, quasi-separated morphism to an algebraic space M. Then  $\pi : \mathcal{M} \to M$  is called a **good moduli space** if the pushforward functor  $\pi_* : \operatorname{Qcoh}(\mathcal{M}) \to \operatorname{Qcoh}(M)$  is exact, and the natural map  $\mathcal{O}_M \to \pi_*\mathcal{O}_M$  is an isomorphism.

Now let us establish our moduli setup, which will remain consistent throughout the subsequent sections. As before, we work over a smooth, projective, polarized threefold (X, H) over  $\mathbb{C}$ . Let  $v \in K_{num}(X)$  be a class of positive rank, and let  $Z : K_{num}(X) \to \mathbb{C}[m]$  define a PT-stability condition on the heart  $\mathcal{A}^p(X)$ . The moduli stack of PT-semistable objects of class v is defined to be the category fibered in groupoids  $\mathcal{M}_Z^{\text{PT}}(v)$  over the category of  $\mathbb{C}$ -schemes that to a scheme S of finite type over  $\mathbb{C}$  associates the groupoid of objects  $E \in D^{\mathrm{b}}(S \times X)$  such that

- (1) E is relatively perfect over S, see [Sta20, Tag 0DHZ], and
- (2) for all  $\mathbb{C}$ -points  $s \in S$ , the derived restriction  $E|_{\{s\}\times X}^{\mathbb{L}}$  to the fiber over s lies in  $\mathcal{A}^p(X)$ , is semistable with respect to Z, and has numerical class  $v \in \mathrm{K}_{\mathrm{num}}(X)$ .

By [Lo13, Theorem 1.1], the stack  $\mathcal{M}_Z^{\mathrm{PT}}(v)$  is universally closed and of finite type over  $\mathbb{C}$ , and moreover, in the case of coprime  $\mathrm{rk}(v)$  and  $H^2 \cdot \mathrm{ch}_1(v)$ , it is a  $\mathbb{G}_m$ -gerbe over a proper algebraic space  $\mathrm{M}_Z^{\mathrm{PT}}(v)$ . In particular,  $\mathrm{M}_Z^{\mathrm{PT}}(v)$  is both a coarse and a good moduli space for  $\mathcal{M}_Z^{\mathrm{PT}}(v)$  in this case.

However, in the presence of strictly semistable objects we do not know if  $\mathcal{M}_Z^{\mathrm{PT}}(v)$  admits a good moduli space – since the heart  $\mathcal{A}^p(X)$  is not noetherian, the machinery of [AHLH23, Chapter 7] does not readily apply.

### 3. Restriction of semistable objects to curves and surfaces

In this section we examine the behavior of PT-semistable objects when restricted to smooth curves and surfaces in X. The following results rely on the characterization of PT-semistable objects given in Proposition 2.3 and will play a key role in Section 5.

**Lemma 3.1.** Assume  $E \in \mathcal{A}^p(X)$  is PT-semistable and fits in the triangle

$$F \to E \to T[-1]$$

where  $F \in Coh(X)$  is torsion-free and  $T \in Coh(X)$  is 0-dimensional. The sheaf  $Q = F^{\vee \vee}/F$  is pure of dimension 1.

*Proof.* If Q is not pure, the maximal 0-dimensional subsheaf  $Q_0 \subseteq Q$  is nonzero. We have exact sequences

$$0 \to Q[-1] \to F \to F^{\vee \vee} \to 0$$

and

$$0 \rightarrow Q_0[-1] \rightarrow Q[-1] \rightarrow Q/Q_0[-1] \rightarrow 0$$

in  $\mathcal{A}^p(X)$ . Thus, we get a nonzero map

$$Q_0[-1] \to Q[-1] \to F \to E$$

as a sequence of inclusions in the heart of a bounded t-structure of perverse sheaves. But this is impossible since  $\operatorname{Hom}_{D^{b}(X)}(Q_{0}[-1], E) = 0$  by assumption.  $\Box$ 

The next result is an analogue of [Taj23, Lemma 6.3].

**Proposition 3.2.** Let  $E \in \mathcal{A}^p(X)$  be a PT-semistable object fitting in a triangle  $F \to E \to T[-1]$ 

where  $F \in \operatorname{Coh}(X)$  is torsion-free and  $T \in \operatorname{Coh}(X)$  is 0-dimensional. Let a > 0 and let  $D, D' \subset X$  be two smooth surfaces in  $|\mathcal{O}_X(a)|$  whose intersection  $C = D \cap D'$ is a smooth, connected curve that does not contain any of the components of the support of the pure 1-dimensional sheaf  $Q = F^{\vee\vee}/F$ . The derived restriction  $E|_C^{\mathbb{L}}$ fits in an exact triangle

$$\mathcal{H}^0(E|_C^{\mathbb{L}}) \to E|_C^{\mathbb{L}} \to \mathcal{H}^1(E|_C^{\mathbb{L}})[-1]$$

where  $\mathcal{H}^1(E|_C^{\mathbb{L}})$  is a 0-dimensional sheaf.

*Proof.* Let  $s, s' \in \mathrm{H}^0(X, \mathcal{O}_X(a))$  be sections corresponding to D and D' respectively. We begin with analyzing the restrictions  $F|_C^{\mathbb{L}}$  and  $T|_C^{\mathbb{L}}$ . First, the restriction  $Q|_D^{\mathbb{L}}$  fits in an exact triangle

$$Q(-a) \xrightarrow{s} Q \to Q|_D^{\mathbb{L}}$$

which induces an exact sequence on cohomology sheaves

$$0 \to \mathcal{H}^{-1}(Q|_D^{\mathbb{L}}) \to Q(-a) \to Q \to \mathcal{H}^0(Q|_D^{\mathbb{L}}) \to 0$$

and all other cohomology sheaves vanish. Thus,  $\mathcal{H}^{-1}(Q|_D^{\mathbb{L}})$  is a subsheaf of Q(-a), hence pure 1-dimensional, and moreover the associated points of  $\mathcal{H}^{-1}(Q|_D^{\mathbb{L}})$  are precisely those associated points of Q where s vanishes. The assumption that Ccontains no component of  $\operatorname{Supp}(Q)$  thus implies that s' does not vanish at any associated point of  $\mathcal{H}^{-1}(Q|_D^{\mathbb{L}})$ .

Next, the short exact sequence

$$0 \to F \to F^{\vee \vee} \to Q \to 0$$

induces a triangle

$$F|_D^{\mathbb{L}} \to F^{\vee \vee}|_D^{\mathbb{L}} \to Q|_D^{\mathbb{L}}$$

which further leads to an exact sequence of cohomology sheaves

$$0 \to \mathcal{H}^{-1}(Q|_D^{\mathbb{L}}) \to F|_D^{\mathbb{L}} = F|_D \to F^{\vee \vee}|_D \to \mathcal{H}^0(Q|_H^{\mathbb{L}}) \to 0$$

as all other cohomology sheaves vanish. Thus  $F|_D^{\mathbb{L}} = F|_D$  is a sheaf and s' vanishes in none of its associated points, so the restriction  $F|_C^{\mathbb{L}}$  fits in the triangle

$$F|_D(-a) \to F|_D \to F|_C^{\mathbb{L}}.$$

The exact sequence of cohomology sheaves

$$0 \to \mathcal{H}^{-1}(F|_C^{\mathbb{L}}) \to F|_D(-a) \to F|_D \to F|_C \to 0$$

shows that the associated points of  $\mathcal{H}^{-1}(F|_C^{\mathbb{L}})$  form a subset of the associated points of  $F|_D(-a)$ . But  $\mathcal{H}^{-1}(F|_C^{\mathbb{L}})$  is supported on C which contains none of the associated points of  $F|_D(-a)$ . Thus  $\mathcal{H}^{-1}(F|_C^{\mathbb{L}}) = 0$ , showing that  $F|_C^{\mathbb{L}} = F|_C$  is a sheaf.

Similarly, we have triangles

$$T(-a) = T \xrightarrow{s} T \to T|_D^{\mathbb{L}}$$

and

$$T|_D^{\mathbb{L}}(-a) \to T|_D^{\mathbb{L}} \to T|_C^{\mathbb{L}}.$$

Combining the associated long exact sequences of cohomology sheaves implies that  $\mathcal{H}^i(T|_C^{\mathbb{L}})$  is 0-dimensional for i = -2, -1, 0 and vanishes otherwise.

Now the long exact sequence of sheaf cohomology associated to the triangle

$$F|_C \to E|_C^{\mathbb{L}} \to T|_C^{\mathbb{L}}[-1]$$

gives

$$0 \to \mathcal{H}^{-1}(E|_C^{\mathbb{L}}) \to \mathcal{H}^{-2}(T|_C^{\mathbb{L}}) \to F|_C \to \mathcal{H}^0(E|_C^{\mathbb{L}}) \to \mathcal{H}^{-1}(T|_C^{\mathbb{L}}) \to 0$$

and  $\mathcal{H}^1(E|_C^{\mathbb{L}}) \cong \mathcal{H}^0(T|_C^{\mathbb{L}})$ . The latter implies that  $\mathcal{H}^1(E|_C^{\mathbb{L}})$  is a 0-dimensional sheaf, so to conclude, we have to show that  $\mathcal{H}^{-1}(E|_C^{\mathbb{L}}) = 0$ . If not, then as a subsheaf of  $\mathcal{H}^{-2}(T|_C^{\mathbb{L}})$ , it is 0-dimensional, so for some  $p \in C$ , we have  $\operatorname{Hom}(\mathcal{O}_p, \mathcal{H}^{-1}(E|_C^{\mathbb{L}})) \neq 0$ . Since

$$\operatorname{Hom}(\mathcal{O}_p, \mathcal{H}^{-1}(E|_C^{\mathbb{L}})) \hookrightarrow \operatorname{Hom}(\mathcal{O}_p, E|_C^{\mathbb{L}}[-1])$$

is injective as  $\mathcal{H}^i(E|_C^{\mathbb{L}}) = 0$  for i < -1, we also get  $\operatorname{Hom}(\mathcal{O}_p, E|_C^{\mathbb{L}}[-1]) \neq 0$ .

Let  $i : C \to X$  denote the closed embedding. Since C and X are smooth, they have dualizing line bundles  $\omega_C$  and  $\omega_X$  respectively. Moreover, the derived restriction and pushforward along i are adjoints. Thus, we have

$$\operatorname{Hom}_{\mathrm{D^{b}}(C)}(\mathcal{O}_{p}, E|_{C}^{\mathbb{L}}[-1]) \cong \operatorname{Hom}_{\mathrm{D^{b}}(C)}(E|_{C}^{\mathbb{L}}[-1], \mathcal{O}_{p} \otimes \omega_{C}[1])^{\vee}$$
$$\cong \operatorname{Hom}_{\mathrm{D^{b}}(C)}(E|_{C}^{\mathbb{L}}, \mathcal{O}_{p}[2])^{\vee}$$
$$\cong \operatorname{Hom}_{\mathrm{D^{b}}(X)}(E, \mathcal{O}_{p}[2])^{\vee}$$
$$\cong \operatorname{Hom}_{\mathrm{D^{b}}(X)}(\mathcal{O}_{p}[2], E \otimes \omega_{X}[3])$$
$$\cong \operatorname{Hom}_{\mathrm{D^{b}}(X)}(\mathcal{O}_{p} \otimes \omega_{X}^{\vee}[-1], E)$$
$$\cong \operatorname{Hom}_{\mathrm{D^{b}}(X)}(\mathcal{O}_{p}[-1], E).$$

But by assumption  $\operatorname{Hom}_{\operatorname{D^b}(X)}(\mathcal{O}_p[-1], E) = 0$ . This concludes the proof.

We will use the following result in Section 5.2, where we restrict families of PT-semistable objects to various smooth curves in X.

**Lemma 3.3.** Let  $E \in D^{b}(X)$  be a PT-semistable object fitting in a triangle

$$F \to E \to T[-1]$$

where  $F \in \operatorname{Coh}(X)$  is torsion-free and  $T \in \operatorname{Coh}(X)$  is 0-dimensional, and denote  $Q = F^{\vee\vee}/F$ . Let  $C \subseteq X$  be a smooth, proper curve that does not contain any components of  $\operatorname{Supp}(Q)$ , and let G be a nonzero vector bundle on C.

- (a) If C meets Supp(T), then  $\mathcal{H}^1(E|_C^{\mathbb{L}})$  is a nonzero torsion sheaf, and we have  $\mathbb{H}^1(C, E|_C^{\mathbb{L}} \otimes G) \neq 0.$
- (b) If C does not meet Supp(T) but does meet Supp(Q), then  $\mathcal{H}^0(E|_C^{\mathbb{L}})$  has a nonzero torsion subsheaf, and we have  $\mathbb{H}^0(C, E|_C^{\mathbb{L}} \otimes G) \neq 0$ .
- (c) If C meets neither Supp(T) nor Supp(Q), then  $E|_C^{\mathbb{L}} = F^{\vee\vee}|_C$  is a sheaf.
- *Proof.* (a) Since C passes through the support of T, the sheaf  $\mathcal{H}^0(T|_C^{\mathbb{L}})$  is nonzero and torsion. The long exact sequence of cohomology sheaves associated to the triangle

$$F|_C^{\mathbb{L}} \to E|_C^{\mathbb{L}} \to T|_C^{\mathbb{L}}[-1]$$

shows that  $\mathcal{H}^1(E|_C^{\mathbb{L}}) \cong \mathcal{H}^1(T|_C^{\mathbb{L}}[-1]) = \mathcal{H}^0(T|_C^{\mathbb{L}})$  is nonzero torsion, and hence so is  $\mathcal{H}^1(E|_C^{\mathbb{L}} \otimes G) = \mathcal{H}^1(E|_C^{\mathbb{L}}) \otimes G$ . The long exact sequence of hypercohomology groups associated to the triangle

$$\mathcal{H}^0(E|_C^{\mathbb{L}}) \otimes G \to E|_C^{\mathbb{L}} \otimes G \to \mathcal{H}^1(E|_C^{\mathbb{L}})[-1] \otimes G$$

now gives a surjection

$$\mathbb{H}^{1}(C, E|_{C}^{\mathbb{L}} \otimes G) \twoheadrightarrow \mathbb{H}^{1}(C, \mathcal{H}^{1}(E|_{C}^{\mathbb{L}})[-1] \otimes G) = \mathrm{H}^{0}(C, \mathcal{H}^{1}(E|_{C}^{\mathbb{L}}) \otimes G) \neq 0.$$

(b) The question is local on X, so in a neighborhood of a point in the intersection of C and Supp(Q), we can express C as the intersection of two smooth surfaces D and D' cut out by functions f and g respectively, neither of which vanishes at the associated points of Q. Thus, we have an exact triangle

$$Q \xrightarrow{J} Q \to Q|_D^{\mathbb{L}}$$

where the map induced by f is an injective map of sheaves, showing that  $Q|_D^{\mathbb{L}} = Q|_D$  is a 0-dimensional sheaf. Now since g vanishes at some point in the support of  $Q|_D$ , the first map in the triangle

$$Q|_D \xrightarrow{g} Q|_D \to Q|_C^{\mathbb{L}}$$

is not an injection, so the sheaf  $\mathcal{H}^{-1}(Q|_C^{\mathbb{L}})$  is a nonzero torsion sheaf on C.

Now since C does not meet  $\operatorname{Supp}(T)$  we have  $E|_C^{\mathbb{L}} = F|_C^{\mathbb{L}}$ . The short exact sequence

$$0 \to F \to F^{\vee \vee} \to Q \to 0$$

induces an exact triangle

$$F|_C^{\mathbb{L}} \to F^{\vee \vee}|_C^{\mathbb{L}} \to Q|_C^{\mathbb{L}}.$$

As above,  $F^{\vee\vee}|_D^{\mathbb{L}}$  is a torsion-free sheaf, hence  $\mathcal{H}^{-1}(F^{\vee\vee}|_C^{\mathbb{L}}) = 0$ , and so the long exact sequence of cohomology sheaves gives an injection

$$\mathcal{H}^{-1}(Q|_C^{\mathbb{L}}) \hookrightarrow F|_C = \mathcal{H}^0(E|_C^{\mathbb{L}}).$$

Finally, taking the long exact sequence in hypercohomology associated to the triangle

$$\mathcal{H}^0(E|_C^{\mathbb{L}}) \otimes G \to E|_C^{\mathbb{L}} \otimes G \to \mathcal{H}^1(E|_C^{\mathbb{L}})[-1] \otimes G$$

gives an inclusion

$$\mathbb{H}^{0}(C, \mathcal{H}^{0}(E|_{C}^{\mathbb{L}}) \otimes G)) \hookrightarrow \mathbb{H}^{0}(C, E|_{C}^{\mathbb{L}} \otimes G),$$

and the first group is nonzero since  $\mathcal{H}^0(E|_C^{\mathbb{L}}) \otimes G$  contains a nonzero torsion subsheaf.

(c) Since C does not meet  $\operatorname{Supp}(T)$ , we have  $E|_C^{\mathbb{L}} = F|_C^{\mathbb{L}}$ , and since C does not meet  $\operatorname{Supp}(Q)$ , we have  $F|_C^{\mathbb{L}} = F^{\vee\vee}|_C^{\mathbb{L}}$ . Working again locally, we saw above that  $F^{\vee\vee}|_D^{\mathbb{L}}$  is torsion-free, so g acts on it as a nonzero divisor. This further implies that  $F^{\vee\vee}|_C^{\mathbb{L}} = F^{\vee\vee}|_C$  is a sheaf.

We end this section with a result that describes the restriction of PT-semistable objects to surfaces.

**Lemma 3.4.** Let  $D \subset X$  be a smooth surface, and let  $E \in \mathcal{A}^p(X)$  be a PT-semistable object fitting in an exact triangle

$$F \to E \to T[-1]$$

where  $F \in Coh(X)$  is a torsion-free and  $T \in Coh(X)$  is 0-dimensional. The derived restriction  $E|_D^{\mathbb{L}}$  fits in an exact triangle

$$\mathcal{H}^0(E|_D^{\mathbb{L}}) \to E|_D^{\mathbb{L}} \to \mathcal{H}^1(E|_D^{\mathbb{L}})[-1]$$

in  $D^{b}(D)$ , and  $\mathcal{H}^{1}(E|_{D}^{\mathbb{L}})$  is isomorphic to  $T|_{D}$ . Moreover, if D does not meet  $\operatorname{Supp}(T)$ , then  $E|_{D}^{\mathbb{L}}$  is isomorphic to the restricted sheaf  $F|_{D}$ .

*Proof.* By using the short exact sequence

$$0 \to F(-D) \to F \to F|_D \to 0$$

one computes that  $\mathcal{H}^0(F|_D^{\mathbb{L}}) \cong F|_D$  and  $\mathcal{H}^i(F|_D^{\mathbb{L}}) = 0$  for  $i \neq 0$ . Similarly, one shows that  $\mathcal{H}^i(T|_D^{\mathbb{L}}) = 0$  for  $i \neq -1, 0$ . Then the exact triangle

$$F|_D^{\mathbb{L}} \to E|_D^{\mathbb{L}} \to T|_D^{\mathbb{L}}[-1]$$

leads to the following long exact sequence in sheaf cohomology

$$0 \to F|_D \to \mathcal{H}^0(E|_D^{\mathbb{L}}) \to \mathcal{H}^{-1}(T|_D^{\mathbb{L}}) \to 0 \to \mathcal{H}^1(E|_D^{\mathbb{L}}) \to \mathcal{H}^0(T|_D^{\mathbb{L}}) \to 0.$$

We see that  $\mathcal{H}^{i}(E|_{D}^{\mathbb{L}}) = 0$  for  $i \neq 0, 1$ , and so  $E|_{D}^{\mathbb{L}}$  is supported only in degrees 0 and 1. Moreover,

$$\mathcal{H}^1(E|_D^{\mathbb{L}}) \cong \mathcal{H}^0(T|_D^{\mathbb{L}}) \cong \operatorname{Coker}(T(-D) \to T),$$

hence  $\mathcal{H}^1(E|_D^{\mathbb{L}})$  is isomorphic to  $T|_D$ . In particular  $\mathcal{H}^1(E|_D^{\mathbb{L}}) = 0$  and  $\mathcal{H}^{-1}(T|_D^{\mathbb{L}}) = 0$ when D avoids the support of T, from which one gets the statement.  $\Box$ 

### 4. Determinantal line bundles

In this section we review the construction and properties of determinantal line bundles on algebraic stacks. Additionally, we construct certain determinantal line bundles on the moduli stack of PT-semistable objects that descend to the good moduli space (assuming one exists). 4.1. Preliminaries on determinantal line bundles. Let S be an algebraic stack of finite type over  $\mathbb{C}$ , and let  $\mathcal{E} \in D^{b}(S \times X)$  be a complex relatively perfect over S. Consider the diagram



We obtain a group homomorphism

$$\lambda_{\mathcal{E}} : \mathrm{K}(X) \to \mathrm{Pic}(S),$$

called the **Donaldson morphism**, defined by sending a vector bundle F on X to the line bundle

$$\lambda_{\mathcal{E}}(F) = \det(\mathrm{R}p_*(\mathcal{E} \otimes q^*F))$$

and extending linearly to K(X). Moreover, if the complex  $Rp_*(\mathcal{E} \otimes q^*F)$  can be locally on S expressed as a 2-term complex of locally free sheaves

$$\cdots \to 0 \to \mathcal{G}_0 \xrightarrow{f} \mathcal{G}_1 \to 0 \to \cdots$$

with  $\operatorname{rk}(\mathcal{G}_0) = \operatorname{rk}(\mathcal{G}_1)$ , then the local sections  $f : \mathcal{O}_S \to \operatorname{det}(\mathcal{G}_1) \otimes \operatorname{det}(\mathcal{G}_0)^{\vee}$  glue to a global section of  $\lambda_{\mathcal{E}}(F)^{\vee}$ . We recall here the following straightforward application of the cohomology and base change theorem [Sta20, Tag 0A1K], which gives a criterion for the existence and non-vanishing of such a section. The proof of [Taj23, Lemma 4.1] applies verbatim.

**Lemma 4.1.** Let X be a smooth, projective variety and S an algebraic stack of finite type over  $\mathbb{C}$ . Let  $\mathcal{E} \in D^{\mathrm{b}}(S \times X)$  be an S-perfect family of objects of class  $v \in \mathrm{K}_{\mathrm{num}}(X)$ , and let  $K \in D^{\mathrm{b}}(X)$ .

(a) If for all  $\mathbb{C}$ -points  $s \in S$ , we have  $\mathbb{H}^i(X, \mathcal{E}_s \otimes^{\mathbb{L}} K) = 0$  whenever  $i \neq 0, 1$ , and

$$\chi(X, \mathcal{E}_s \otimes^{\mathbb{L}} K) = \dim \mathbb{H}^0(X, \mathcal{E}_s \otimes^{\mathbb{L}} K) - \dim \mathbb{H}^1(X, \mathcal{E}_s \otimes^{\mathbb{L}} K) = 0,$$

then the line bundle  $\lambda_{\mathcal{E}}(K)^{\vee}$  on S has a canonical section  $\delta_K$ . (b) In addition, if for some  $s_0 \in S$  we have

$$\mathbb{H}^{0}(X, \mathcal{E}_{s_{0}} \otimes^{\mathbb{L}} K) = \mathbb{H}^{1}(X, \mathcal{E}_{s_{0}} \otimes^{\mathbb{L}} K) = 0,$$

then the section  $\delta_K$  is nonzero at  $s_0$ .

4.2. Determinantal line bundles on PT-moduli spaces. We now specialize to the situation of PT-semistability. Let  $v \in K_{num}(X)$  be a numerical class of positive rank and Z a PT-stability function on  $\mathcal{A}^p(X)$ . Consider the stack  $\mathcal{M}_Z^{PT}(v)$ of PT-semistable objects of class v in  $\mathcal{A}^p(X)$ , and let  $\mathcal{E}$  be the universal complex on  $\mathcal{M}_Z^{PT}(v) \times X$ , so that we have the diagram



As in [HL10, Example 8.1.8 (iii)], define

$$v_{2}(v) = -\chi(v \cdot h^{3})h^{2} + \chi(v \cdot h^{2})h^{3} \in \mathcal{K}(X),$$
  

$$v_{1}(v) = -\chi(v \cdot h^{3})h + \chi(v \cdot h)h^{3} \in \mathcal{K}(X),$$
  

$$v_{0}(v) = -\chi(v \cdot h^{3})[\mathcal{O}_{X}] + \chi(v)h^{3} \in \mathcal{K}(X),$$

where  $h = [\mathcal{O}_H] \in \mathcal{K}(X)$ , and define

$$\mathcal{L}_i = \lambda_{\mathcal{E}}(v_i(v)) \in \operatorname{Pic}(\mathcal{M}_Z^{\operatorname{PT}}(v)) \text{ for } i = 0, 1, 2.$$

The line bundles  $\mathcal{L}_i$  are our main objects of study. We will show that each  $\mathcal{L}_i$  descends to a line bundle on the good moduli space  $M_Z^{PT}(v)$  (assuming that this space exists) when U in the definition of Z is taken to be the Todd class of X.

We recall the following criterion [Alp08, Theorem 10.3] for a locally free sheaf  $\mathcal{F}$  on  $\mathcal{M}$  to descend to the good moduli space M.

**Proposition 4.2.** If  $\pi : \mathcal{M} \to M$  is a good moduli space and  $\mathcal{M}$  is locally Noetherian, then the pullback morphism  $\pi^* : \operatorname{Coh}(\mathcal{M}) \to \operatorname{Coh}(\mathcal{M})$  induces an equivalence of categories between locally free sheaves on  $\mathcal{M}$  and those locally free sheaves  $\mathcal{F}$  on  $\mathcal{M}$ such that for every geometric point  $x : \operatorname{Spec} k \to \mathcal{M}$  with closed image, the induced representation  $x^*\mathcal{F}$  of the stabilizer  $G_x$  is trivial.

By Proposition 4.2 we must control the action of the stabilizer group  $G_x$  of  $M_Z^{\text{PT}}(v)$  on the fiber  $\mathcal{L}_i|_x$  for closed points  $x \in M_Z^{\text{PT}}(v)$ . We first prove the following.

**Lemma 4.3.** If E is a PT-semistable of class v and  $F \subseteq E$  is a subobject in  $\mathcal{A}^p(X)$  such that

$$\phi(Z(F)(m)) = \phi(Z(E)(m)) \quad \text{for } m \gg 0,$$

then

(4.1) 
$$\int_X H^d \cdot \operatorname{ch}(F) \cdot U = \frac{\operatorname{rk}(F)}{\operatorname{rk}(E)} \int_X H^d \cdot \operatorname{ch}(E) \cdot U$$

for d = 0, 1, 2.

*Proof.* The assumption on F is equivalent to saying that  $p_v(F) = 0$ , where  $p_v$  is defined by

$$p_v : \mathbf{K}(X) \to \mathbb{R}[m], \quad p_v(F) = \operatorname{Im} Z(v) \operatorname{Re} Z(F) - \operatorname{Re} Z(v) \operatorname{Im} Z(F)$$

To lighten the notation, we set

$$I_d(G) = \int_X H^d \cdot \operatorname{ch}(G) \cdot U, \quad d = 0, \dots, 3$$

We note that since

$$I_3(G) = \int_X H^3 \cdot \operatorname{ch}(G) \cdot U = \operatorname{deg}(X) \operatorname{rk}(G),$$

and since rk(F) > 0 by Lemma 2.4, equation (4.1) is equivalent to

$$I_3(E)I_d(F) = I_3(F)I_d(E)$$

Moreover, we set

$$r_{ij} = \operatorname{Re}(\rho_i) \operatorname{Im}(\rho_j), \quad i, j = 0, \dots, 3.$$

and note that since none of the complex numbers  $\rho_i$  are collinear, the real numbers  $r_{ij} - r_{ji}$  are all nonzero for  $i \neq j$ .

The condition  $p_v(F) = 0$  can now be written as

(4.2) 
$$\sum_{d=0}^{\circ} \sum_{i+j=d} r_{ij} (I_i(E)I_j(F) - I_i(F)I_j(E))m^d = 0.$$

We compare coefficients on both sides of this equation. First, the  $m^5$  term in (4.2) gives

$$(r_{32} - r_{23})(I_2(E)I_3(F) - I_2(F)I_3(E)) = 0,$$

so dividing by deg(X) and  $r_{32} - r_{23}$  gives (4.1) for d = 2. Similarly from the  $m^4$  term, noting that the i = j = 2 term cancels out, we get

$$(r_{31} - r_{13})(I_1(E)I_3(F) - I_1(F)I_3(E)) = 0,$$

giving (4.1) for d = 1. Finally, the  $m^3$  term gives

$$(r_{30} - r_{03})(I_0(E)I_3(F) - I_0(F)I_3(E)) + (r_{21} - r_{12})(I_1(E)I_2(F) - I_1(F)I_2(E)) = 0.$$

The second term on the left cancels since  $I_3(E) \neq 0$  and by what we have already proven, we have

$$I_3(E)I_1(E)I_2(F) = I_3(F)I_1(E)I_2(E) = I_3(E)I_1(E)I_2(E).$$

Thus, we obtain (4.1) for d = 0, completing the proof.

**Proposition 4.4.** Let Z be a PT-stability function on  $\mathcal{A}^p(X)$  and assume that  $U = \operatorname{td}_X$  in the definition of Z is the Todd class of X. Assuming a good moduli space  $\operatorname{M}_Z^{\operatorname{PT}}(v)$  exists, the line bundle

$$\mathcal{L}_i = \lambda_{\mathcal{E}}(v_i(v)) \in \operatorname{Pic}(\mathcal{M}_Z^{\operatorname{PT}}(v))$$

descends to  $M_Z^{PT}(v)$  for i = 0, 1, 2.

*Proof.* Let  $x \in M_Z^{PT}(v)$  be a closed point corresponding to the Z-polystable object

$$E = \bigoplus_{j} F_j$$
, where  $p_v(F_j) = 0$  for all  $j$ .

By [Taj23, Proposition 4.2], the automorphism group of E acts trivially on the fiber of  $\mathcal{L}_i$  at x if and only if  $\chi([F_j] \cdot v_i(v)) = 0$  for each j. By the Hirzebruch-Riemann-Roch formula,

$$\chi([F_j] \cdot v_i(v)) = \int_X \operatorname{ch}(F_j) \operatorname{ch}(v_i(v)) \operatorname{td}_X.$$

Now

$$\operatorname{ch}(v_i(v)) = -\chi(v \cdot h^3) \operatorname{ch}(h)^i + \chi(v \cdot h^i) \operatorname{ch}(h)^3,$$

and

$$ch(h) = ch(\mathcal{O}_X) - ch(\mathcal{O}_X(-H)) = H - \frac{1}{2}H^2 + \frac{1}{6}H^3$$

Thus,  $ch(v_i(v))$  is a linear combination of powers of H. Hence, by Lemma 4.3 and linearity, we obtain

$$\int_X \operatorname{ch}(F_j) \operatorname{ch}(v_i(v)) \operatorname{td}_X = \frac{\operatorname{rk}(F_j)}{\operatorname{rk}(E)} \int_X \operatorname{ch}(E) \operatorname{ch}(v_i(v)) \operatorname{td}_X.$$

Since  $[E] = v \in K_{num}(X)$ , by the Hirzebruch-Riemann-Roch formula again,

$$\int_X \operatorname{ch}(E) \operatorname{ch}(v_i(v)) \operatorname{td}_X = \chi(v \cdot v_i(v)) = 0.$$

We also recall the following general descent result that will be useful in Section 5.

**Proposition 4.5** ([BDF<sup>+</sup>22, Lemma 5.1.5 (iii)]). Let  $\pi : \mathcal{M} \to \mathcal{M}$  be a good moduli space, and let  $\mathcal{L}$  be a globally generated line bundle on  $\mathcal{M}$ . Then  $\mathcal{L}$  descends to a globally generated line bundle L on  $\mathcal{M}$ .

### 5. PROJECTIVITY OF THE MODULI SPACE OF PT-SEMISTABLE OBJECTS

In this section we study the positivity of the line bundles  $\mathcal{L}_i$  that were previously constructed on  $\mathcal{M}_Z^{\mathrm{PT}}(v)$ , with the goal of establishing the projectivity of the good moduli space  $\mathcal{M}_Z^{\mathrm{PT}}(v)$  in the case of coprime rank and degree.

5.1. **Positivity of**  $\mathcal{L}_2$ . Following [Taj21], we start by showing that a power of  $\mathcal{L}_2$  is globally generated on  $\mathcal{M}_Z^{\mathrm{PT}}(v)$ . The main idea consists of relating  $\mathcal{L}_2$  to restrictions of the universal complex  $\mathcal{E}$  on  $\mathcal{M}_Z^{\mathrm{PT}}(v) \times X$  to various curves in X. Let a, b > 0 be integers and let  $H' \in |\mathcal{O}_X(a)|$  and  $H'' \in |\mathcal{O}_X(b)|$  be smooth

Let a, b > 0 be integers and let  $H' \in |\mathcal{O}_X(a)|$  and  $H'' \in |\mathcal{O}_X(b)|$  be smooth hyperplane sections whose intersection  $C = H' \cap H''$  is a smooth, connected curve. Consider the diagram



Here  $\mathcal{E}_C$  denotes the derived restrictions of  $\mathcal{E}$  to  $\mathcal{M}_Z^{\mathrm{PT}}(v) \times C$ . Consider the Donaldson morphisms

$$\lambda_{\mathcal{E}} : \mathrm{K}(X) \to \mathrm{Pic}(\mathcal{M}_Z^{\mathrm{PT}}(v)), \quad \lambda_{\mathcal{E}_C} : \mathrm{K}(C) \to \mathrm{Pic}(\mathcal{M}_Z^{\mathrm{PT}}(v)).$$

Lemma 5.1. We have a commutative diagram

$$\begin{array}{c} \mathrm{K}(X) & \stackrel{\cdot[\mathcal{O}_C]}{\longrightarrow} \mathrm{K}(X) \\ i^* \downarrow & \lambda_{\mathcal{E}} \downarrow \\ \mathrm{K}(C) & \stackrel{}{\longrightarrow} \mathrm{Pic}(\mathcal{M}_Z^{\mathrm{PT}}(v)) \end{array}$$

*Proof.* It suffices to show that  $\lambda_{\mathcal{E}}(F \otimes i_* \mathcal{O}_C) = \lambda_{\mathcal{E}_C}(F|_C)$  when F is a locally free sheaf on X. By flat base change and the projection formula, we have

$$j_*\mathcal{E}_C = j_*j^*\mathcal{E} = \mathcal{E} \otimes j_*q_C^*\mathcal{O}_C = \mathcal{E} \otimes q^*i_*\mathcal{O}_C$$

Thus,

$$\lambda_{\mathcal{E}_C}(F|_C) = \det \operatorname{R}_{p_{\mathcal{E}}}(\mathcal{E}_C \otimes q_C^* i^* F)$$
  
= det Rp\_\*j\_\*( $\mathcal{E}_C \otimes j^* q^* F$ )  
= det Rp\_\*(j\_\*  $\mathcal{E}_C \otimes q^* F$ )  
= det Rp\_\*( $\mathcal{E} \otimes q^* (i_* \mathcal{O}_C \otimes F)$ )  
=  $\lambda_{\mathcal{E}}(i_* \mathcal{O}_C \otimes F)$ .

**Proposition 5.2.** Consider the class

$$w \coloneqq -\chi(v \cdot h \cdot [\mathcal{O}_C]) \cdot 1 + \chi(v \cdot [\mathcal{O}_C]) \cdot h \in \mathcal{K}(X),$$

where  $h = [\mathcal{O}_H] \in \mathcal{K}(X)$ . Then

$$\lambda_{\mathcal{E}_C}(w|_C) = \mathcal{L}_2^{\otimes a^2 b^2}.$$

Moreover,  $-w|_C$  has positive rank and so is represented by a locally free sheaf on C.

*Proof.* For the first claim, by Lemma 5.1 it suffices to show that  $w \cdot [\mathcal{O}_C] = a^2 b^2 v_2(v)$ . Note that  $[\mathcal{O}_C] = [\mathcal{O}_{H'}][\mathcal{O}_{H''}]$ , and that  $\mathcal{O}_{H'}$  fits in the exact sequence

$$0 \to \mathcal{O}_X(-a) \to \mathcal{O}_X \to \mathcal{O}_{H'} \to 0,$$

so that

$$[\mathcal{O}_{H'}] = [\mathcal{O}_X] - [\mathcal{O}_X(-a)] = 1 - [\mathcal{O}_X(-1)]^a = 1 - (1-h)^a = ah - \binom{a}{2}h^2 + \binom{a}{3}h^3$$

and similarly  $[\mathcal{O}_{H''}] = bh - {b \choose 2}h^2 + {b \choose 3}h^3$ , so that

$$[\mathcal{O}_C] = (ah - \binom{a}{2}h^2 + \binom{a}{3}h^3)(bh - \binom{b}{2}h^2 + \binom{b}{3}h^3)$$
$$= abh^2 - \left(a\binom{b}{2} + b\binom{a}{2}\right)h^3.$$

Thus,  $h \cdot [\mathcal{O}_C] = abh^3$ , hence

$$w = -ab\chi(v \cdot h^3) \cdot 1 + ab\chi(v \cdot h^2) \cdot h - \left(a\binom{b}{2} + b\binom{a}{2}\right)\chi(v \cdot h^3) \cdot h,$$

and so

$$w \cdot [\mathcal{O}_C] = -a^2 b^2 \chi(v \cdot h^3) \cdot h^2 + a^2 b^2 \chi(v \cdot h^2) h^3 = a^2 b^2 v_2(v).$$

For the second claim, we note that  $\operatorname{rk}(w) = -a^4 \chi(v \cdot h^3) = -a^2 b^2 \operatorname{rk}(v) \operatorname{deg}(X) < 0$ . Since restriction to C preserves rank, we see that  $\operatorname{rk}(-w|_C) > 0$ , so  $-w|_C$  is represented for example by the sheaf

$$\mathcal{O}_C^{\oplus \operatorname{rk}(-w|_C)-1} \oplus \det(-w|_C).$$

Let a > 0 be an integer and let  $S_a = |\mathcal{O}_X(a)| \times |\mathcal{O}_X(a)|$ , where  $|\mathcal{O}_X(a)|$  is the complete linear system of  $\mathcal{O}_X(a)$ . Let  $Z^a \subseteq S_a \times X$  denote the incidence correspondence of complete intersections  $D_1 \cap D_2 \subseteq X$  with  $D_1, D_2 \in |\mathcal{O}_X(a)|$ . Consider the diagram



**Lemma 5.3.** If  $U \subseteq S_a$  is a nonempty open set, then  $\bigcup_{t \in U} Z_t^a = q(p^{-1}(U)) \subseteq X$  is an open subset whose complement has codimension at least 2 in X.

Proof. The map  $q: Z^a \to X$  is a product of projective bundles (see e.g. [HL10, Section 3.1]), hence flat and in particular open, and so  $q(p^{-1}(U))$  is nonempty and open. If  $\eta \in X$  is a point of codimension 1 with closure  $Y \subseteq X$ , then the intersection  $D_1 \cap D_2 \cap Y$  is nonempty for any  $D_1, D_2 \in |\mathcal{O}_X(a)|$  since  $\mathcal{O}_X(a)$  is ample. Thus,  $q(p^{-1}(U)) \cap Y$  is nonempty and open in Y, hence contains  $\eta$  since Y is irreducible.

**Lemma 5.4.** There exists  $a_0 > 0$  such that for any  $a \ge a_0$  there exists a nonempty open subset  $U \subseteq S_a$  with the following property: for every  $t \in U$ , the fiber  $Z_t^a$  is a smooth, connected curve, and if  $E \in \mathcal{A}^p(X)$  is any PT-semistable object of class v, then  $Z_t^a$  contains no component of  $\operatorname{Supp}(\mathcal{H}^0(E)^{\vee\vee}/\mathcal{H}^0(E))$ .

Proof. Since the set of isomorphism classes of PT-semistable objects E of class v is bounded, so is the set of isomorphism classes of the quotients  $Q = \mathcal{H}^0(E)^{\vee\vee}/\mathcal{H}^0(E)$ , and hence the degree of  $\operatorname{Supp}(Q)$  is bounded by some b > 0, where the scheme structure of  $\operatorname{Supp}(Q)$  is given by the annihilator ideal sheaf. Let  $a_0 = b + 1$ . By Bertini's theorem, there is a nonempty open subset  $U \subseteq S_a$  such that for every  $t \in$ U, the fiber  $Z_t^a$  is a smooth, connected curve, and since  $\operatorname{deg}(Z_t^a) > \operatorname{deg}(\operatorname{Supp}(Q))$ when  $a \ge a_0$ , the curve  $Z_t^a$  cannot contain components of  $\operatorname{Supp}(Q)$ .  $\Box$ 

**Lemma 5.5.** There exists  $a_0 > 0$  such that for any  $a \ge a_0$  and any PT-semistable object  $E \in \mathcal{A}^p(X)$  of class v, there exists a nonempty open subset  $U \subseteq S_a$  such that for every  $t \in U$ , the fiber  $Z_t^a$  is a smooth, connected curve and the restriction  $E|_{Z_t^a}^{\mathbb{L}}$  is a semistable sheaf on  $Z_t^a$ .

*Proof.* Recall that E fits in an exact triangle

$$F \to E \to T[-1]$$

where F is a  $\mu$ -semistable torsion-free sheaf and T is a 0-dimensional sheaf. Denote  $r = \operatorname{rk}(E) = \operatorname{rk}(F)$ . By Flenner's Theorem [HL10, Theorem 7.1.1], if  $a \in \mathbb{N}$  satisfies

$$\frac{\binom{a+3}{a} - 2a - 1}{a} > \deg(X) \cdot \max\left\{\frac{r^2 - 1}{4}, 1\right\},\$$

there exists a nonempty open subset  $U' \subseteq S_a$  such that for any  $t \in U'$ , the fiber  $Z_t^a$  is a smooth, connected curve, and the restriction  $F|_{Z_t^a}$  is a semistable sheaf. Now the set of those  $t \in U'$  such that  $Z_t^a$  intersects  $\operatorname{Supp}(T)$  is a proper, closed subset of U', and if we take U to be the complement of this subset in U', then for any  $t \in U$ , we have  $E|_{Z_t^a}^{\mathbb{L}} = F|_{Z_t^a}$ .

The following result, originally due to Faltings, is key to proving the semiampleness of the line bundle  $\mathcal{L}_2$  and to describing which points can be separated by it. We refer to the work of Seshadri [Ses93, Lemma 3.1 and Lemma 4.2] for a proof.

**Lemma 5.6.** Let C and S be two smooth, projective, connected curves, and let  $\mathcal{F} \in \operatorname{Coh}(S \times X)$  be a family of semistable locally free sheaves on C of rank r > 0 and degree d. Let  $E \in \operatorname{Coh}(X)$  be a locally free sheaf such that

 $r \deg E + (d + r(1 - g)) \operatorname{rk} E = 0.$ 

(i) If E is semistable and  $r \gg 0$  is sufficiently large, then for any line bundle L of degree d, there exists a locally free sheaf G with  $\operatorname{rk} G = r$  and  $\det G \cong L$  such that

$$\mathrm{H}^{0}(C, E \otimes G) = \mathrm{H}^{1}(C, E \otimes G) = 0.$$

(ii) If the determinantal line bundle  $\lambda_{\mathcal{F}}(E) \in \operatorname{Pic}(S)$  has degree 0, then the semistable sheaves  $\mathcal{F}_s$  are all S-equivalent.

Remark 5.7. The following converse of Lemma 5.6 (i) also holds: if E is a locally free sheaf on a smooth, projective, connected curve C and there exists a locally free sheaf G such that

$$\mathrm{H}^{0}(C, E \otimes G) = \mathrm{H}^{1}(C, E \otimes G) = 0,$$

then E is semistable. We refer to [MS17] for a proof.

**Theorem 5.8.** There is an integer  $a_0 > 0$  such that for any  $a \ge a_0$  and any *PT*-semistable object  $E_0 \in \mathcal{A}^p(X)$  of class v, there exists a nonempty open subset  $U \subseteq S_a$  satisfying the following property: for every  $t \in U$ , the fiber  $C = Z_t^a$  is a smooth, connected curve, and there exists a locally free sheaf G on C such that we have

$$\mathcal{L}_2^{\otimes ma^4} = \lambda_{\mathcal{E}_C}(G)^{\vee}$$

for some m > 0, and G induces a global section  $\delta_G \in \Gamma(\mathcal{M}_Z^{\mathrm{PT}}(v), \mathcal{L}_2^{\otimes ma^4})$  that is nonvanishing at  $[E_0] \in \mathcal{M}_Z^{\mathrm{PT}}(v)$ . In particular, for sufficiently large m > 0, the line bundle  $\mathcal{L}_2^{\otimes ma^4} \in \operatorname{Pic}(\mathcal{M}_Z^{\mathrm{PT}}(v))$  is globally generated.

*Proof.* Combining Lemmas 5.4 and 5.5, we find  $a_0 > 0$  such that for any  $a \ge a_0$  the following two conditions hold:

- (i) there exists a nonempty open set  $U_1 \subseteq S_a$  such that for every  $t \in U_1$ , the fiber  $Z_t^a$  is a smooth, connected curve, and if  $E \in \mathcal{A}^p(X)$  is any PT-semistable object of class v, then  $Z_t^a$  does not contain any components of  $\mathcal{H}^0(E)^{\vee\vee}/\mathcal{H}^0(E)$ ,
- (ii) there exists an open subset  $U_2 \subseteq S_a$  such that for every  $t \in U_2$ , the fiber  $Z_t^a$  is a smooth, connected curve, and the restriction  $E_0|_{Z_t^a}^{\mathbb{L}}$  is a  $\mu$ -semistable sheaf.

Set  $U = U_1 \cap U_2 \subseteq S_a$ . Let  $t \in U$  and denote  $C := Z_t^a$ . Recall from Proposition 5.2 that we defined

$$w \coloneqq -\chi(v \cdot h \cdot [\mathcal{O}_C]) \cdot 1 + \chi(v \cdot [\mathcal{O}_C]) \cdot h \in \mathbf{K}(X)$$

and observed that  $\mathcal{L}_2^{\otimes a^4} = \lambda_{\mathcal{E}_C}(w|_C)$ . Also, notice that for m > 0 we have

 $\operatorname{rk}(-mw|_{C}) = ma^{2} \operatorname{deg}(X) \operatorname{rk}(v) > 0,$ 

and that  $\chi(mw|_C \cdot v|_C) = 0$  since

$$w|_C = -\chi(v|_C \cdot h|_C) \cdot 1 + \chi(v|_C) \cdot h|_C.$$

Thus, by Lemma 5.6 (i), for m sufficiently large there exists a locally free sheaf G of class  $-mw|_C$  on C such that  $\mathbb{H}^i(C, E_0|_C^{\mathbb{L}} \otimes G) = 0$  for all  $i \in \mathbb{Z}$ .

Let now  $E \in \mathcal{A}^p(X)$  be any PT-semistable object. By Proposition 3.2, we have an exact triangle

$$F \to E|_C^{\mathbb{L}} \otimes G \to T[-1]$$

in  $D^{b}(C)$ , where F is a coherent sheaf and T is a 0-dimensional coherent sheaf. Thus, the long exact sequence in hypercohomology shows that  $\mathbb{H}^i(C, E|_C^{\mathbb{L}} \otimes G) = 0$ for  $i \neq 0, 1$ , and since  $\chi(C, E|_C^{\mathbb{L}} \otimes G) = \chi(-mw|_C \cdot v|_C) = 0$ , we have

$$\dim \mathbb{H}^0(C, E|_C^{\mathbb{L}} \otimes G) = \dim \mathbb{H}^1(C, E|_C^{\mathbb{L}} \otimes G).$$

Thus, by Lemma 4.1, the line bundle

$$\mathcal{L}_2^{\otimes ma^4} = \lambda_{\mathcal{E}_C}(mw|_C) = \lambda_{\mathcal{E}_C}(G)^{\vee}$$

has a global section  $\delta_G$  that does not vanish at the point  $[E_0] \in \mathcal{M}_Z^{\mathrm{PT}}(v)$ . Since  $\mathcal{M}_Z^{\mathrm{PT}}(v)$  is quasi-compact, we can choose a sufficiently large m > 0 such that  $\mathcal{L}_2^{\otimes ma^4}$  is globally generated. 

5.2. Description of the fibers. From now on we assume the existence of the (proper) good moduli space  $\mathcal{M}_Z^{\mathrm{PT}}(v) \to \mathrm{M}_Z^{\mathrm{PT}}(v)$ . By Theorem 5.8 we know that  $\mathcal{L}_{2}^{\otimes ma^{4}}$  is globally generated on  $\mathcal{M}_{Z}^{\mathrm{PT}}(v)$  for some integers m, a > 0. By using Proposition 4.5 we obtain that  $\mathcal{L}_{2}^{\otimes ma^{4}}$  descends to a globally generated line bundle L on  $M_Z^{PT}(v)$ . We choose a sufficiently large integer k > 0 such that the map

$$\phi: \mathbf{M}_Z^{\mathrm{PT}}(v) \to \mathbb{P} \coloneqq \mathbb{P}\Gamma(\mathbf{M}_Z^{\mathrm{PT}}(v), L^{\otimes k})$$

determined by the complete linear system  $|L^{\otimes k}|$  has connected fibers (see [Laz04, Lemma 2.1.28]). Notice that the vector space  $\Gamma(\mathbf{M}_{Z}^{\mathrm{PT}}(v), L^{\otimes k})$  is finite-dimensional since  $M_Z^{PT}(v)$  is proper.

Next we study the fibers of  $\phi$ . To state our result, we introduce the following notation. Given a  $\mu$ -semistable torsion-free sheaf F on X, let  $F^{**} := \operatorname{gr}(F)^{\vee \vee}$ denote the double dual of the polystable sheaf  $gr(F) = \bigoplus_i gr_i(F)$  associated to a Jordan-Hölder filtration of F with torsion-free factors  $gr_i(F)$ . Note that  $F^{**}$  is independent of the Jordan-Hölder filtration, cf. [HL10, Corollary 1.6.10]. We aim to prove the following result, which together with Theorem 5.8, establishes the first part of Theorem 1.1.

**Theorem 5.9.** Let  $E_1, E_2 \in \mathcal{M}_Z^{\mathrm{PT}}(v)(\mathbb{C})$  be two PT-semistable objects mapping to the same point under the map  $\mathcal{M}_Z^{\mathrm{PT}}(v) \to \mathrm{M}_Z^{\mathrm{PT}}(v) \xrightarrow{\phi} \mathbb{P}$ . Denote  $F_i = \mathcal{H}^0(E_i)$  for i = 1, 2.

- (i) The sheaves F<sub>1</sub><sup>\*\*</sup> and F<sub>2</sub><sup>\*\*</sup> are isomorphic.
  (ii) If gcd(rk(v), H<sup>2</sup> · ch<sub>1</sub>(v)) = 1, then F<sub>1</sub><sup>∨∨</sup> and F<sub>2</sub><sup>∨∨</sup> are isomorphic, and for every point η ∈ X of codimension 2, the stalks of the sheaves F<sub>1</sub><sup>∨∨</sup>/F<sub>1</sub> and  $F_2^{\vee\vee}/F_2$  at  $\eta$  have the same lengths as modules over the local ring  $\mathcal{O}_{X,\eta}$ .

Since the fiber  $M_p \subset \mathcal{M}_Z^{\mathrm{PT}}(v)$  of  $\phi$  over a point  $p \in \mathbb{P}$  is a proper and connected algebraic space, it can be covered by images of maps  $S \to M_p$  where S is a smooth, proper, connected curve. Moreover, after possibly taking a finite cover of S, we may assume that the map lifts to  $S \to \mathcal{M}_Z^{\mathrm{PT}}(v)$ ; see for example [Taj23, Lemma 5.4]. Notice that in particular  $\deg(\mathcal{L}_2^{\otimes k}|_S) = 0$ . So let S be a smooth, proper, connected curve and let  $\mathcal{E} \in D^b(S \times X)$  be a

family of PT-semistable objects corresponding to a map  $S \to \mathcal{M}_Z^{\mathrm{PT}}(v)$  such that  $\deg(\mathcal{L}_2^{\otimes k}|_S) = 0.$  For each  $s \in S$ , denote

$$\mathcal{E}_s = \mathcal{E}|_{\{s\} \times X}, \quad F_s = \mathcal{H}^0(\mathcal{E}_s), \quad T_s = \mathcal{H}^1(\mathcal{E}_s), \quad Q_s = F_s^{\vee \vee}/F_s.$$

The following in particular proves the first part of Theorem 5.9, since any two points in  $\mathcal{M}_p$  can be connected by a chain of such curves.

**Proposition 5.10.** Under the above notation, the sheaves  $F_s^{**}$  are isomorphic for all  $s \in S$ , and there exists a closed 1-dimensional subset  $Y \subset X$  such that for every  $s \in S$ , the support of  $Q_s$  is contained in Y.

Proof. Fix a point  $s_0 \in S$ , choose an integer a and an open subset  $U \subseteq |\mathcal{O}_X(a)| \times |\mathcal{O}_X(a)|$  as in Theorem 5.8 so that for each  $t \in U$  there exists an integer m > 0 and a sheaf G on the fiber  $C := Z_t^a$  such that the section  $\delta_G$  of  $\mathcal{L}_2^{\otimes ma^4}|_S = \lambda_{\mathcal{E}_C}(G)^{\vee}|_S$  is nonzero at  $s_0 \in S$ . By assumption, we can choose m appropriately so that  $\deg(\mathcal{L}_2^{\otimes ma^4}|_S) = 0$ , which further implies that the section  $\delta_G$  must be nonzero at every point  $s \in S$ , and so

$$\mathbb{H}^{0}(C, \mathcal{E}_{s}|_{C}^{\mathbb{L}} \otimes G) = \mathbb{H}^{1}(C, \mathcal{E}_{s}|_{C}^{\mathbb{L}} \otimes G) = 0.$$

By Lemma 3.3, the curve  $Z_t^a$  does not meet the supports of  $Q_s$  or  $T_s$  for any  $t \in U$ and  $s \in S$ . In particular, the supports of the sheaves  $Q_s$  must be contained in the complement Y of the union  $\bigcup_{t \in U} Z_t^a \subseteq X$  of all the  $Z_t^a$  for  $t \in U$ , which by Lemma 5.3 above is a closed 1-dimensional subset. This proves the second claim.

To prove the first claim, we give a variant of the restriction argument in the proof of Theorem 5.8. Fix  $s_1, s_2 \in S$ , and for j = 1, 2 fix a Jordan-Hölder filtration

$$0 \subset F_{s_j}^{(1)} \subset \cdots \subset F_{s_j}^{(k_j-1)} \subset F_{s_j}^{(k_j)} = F_{s_j}$$

with  $\mu$ -stable torsion-free factors  $G_{s_j}^{(i)} = F_{s_j}^{(i)} / F_{s_j}^{(i-1)}$ . Note that

$$F_{s_j}^{**} = \left(\bigoplus_i G_{s_j}^{(i)}\right)^{\vee \vee}$$

First, we can choose an integer  $a \gg 0$  and a smooth, connected surface  $D \in |\mathcal{O}_X(a)|$  such that D avoids the supports of  $T_{s_1}$  and  $T_{s_2}$ , the restrictions  $F_{s_1}|_D$  and  $F_{s_2}|_D$  are  $\mu$ -semistable and torsion-free [HL10, Theorem 7.1.1] and the restrictions  $G_{s_1}^{(i)}|_D$  and  $G_{s_2}^{(i)}|_D$  of all the Jordan-Hölder factors are  $\mu$ -stable and torsion-free [HL10, Theorem 7.2.8] on D. This implies that for j = 1, 2 the restricted filtration

$$0 \subset F_{s_j}^{(1)}|_D \subset \dots \subset F_{s_j}^{(k_j-1)}|_D \subset F_{s_j}^{(k_j)}|_D = F_{s_j}|_D$$

is a Jordan-Hölder filtration for  $F_{s_j}|_D$ . We can also assume that D avoids the finitely many (codimension 2 or 3) associated points of each of the sheaves  $(G_{s_1}^{(i)})^{\vee\vee}/G_{s_1}^{(i)}$  and  $(G_{s_2}^{(i)})^{\vee\vee}/G_{s_1}^{(i)}$ . We may also assume that D avoids the finitely many singular points of  $F_{s_1}^{**}$  and  $F_{s_2}^{**}$ , implying that  $F_{s_1}^{**}|_D$  and  $F_{s_2}^{**}|_D$  are locally free. Moreover, since  $F_{s_1}^{**}$  and  $F_{s_2}^{**}$  are reflexive, by increasing a if necessary, we obtain that

$$\operatorname{Ext}^{l}(F_{s_{1}}^{**}, F_{s_{2}}^{**}(-D)) = \operatorname{Ext}^{l}(F_{s_{2}}^{**}, F_{s_{1}}^{**}(-D)) = 0$$

for l = 0, 1. Hence  $\operatorname{Hom}(F_{s_1}^{**}, F_{s_2}^{**}) = \operatorname{Hom}(F_{s_1}^{**}|_D, F_{s_2}^{**}|_D)$  and similarly with  $F_{s_1}$  and  $F_{s_2}$  interchanged. This implies that  $F_{s_1}^{**} \cong F_{s_2}^{**}$  if and only if  $F_{s_1}^{**}|_D \cong F_{s_2}^{**}|_D$ . As above, using [HL10, Theorem 7.1.1] and [HL10, Theorem 7.2.8], we can choose

As above, using [HL10, Theorem 7.1.1] and [HL10, Theorem 7.2.8], we can choose an integer  $b \gg 0$  and a curve  $C \in |\mathcal{O}_D(b)|$  such that again the restrictions  $F_{s_1}|_C$ and  $F_{s_2}|_C$  are semistable and the restrictions  $G_{s_1}^{(i)}|_C$  and  $G_{s_2}^{(i)}|_C$  are stable. This implies that

$$\operatorname{gr}(F_{s_1})|_C \cong \operatorname{gr}(F_{s_1}|_C)$$
 and  $\operatorname{gr}(F_{s_2})|_C \cong \operatorname{gr}(F_{s_2}|_C).$ 

Since D avoids the associated points of each of  $(G_{s_1}^{(i)})^{\vee\vee}/G_{s_1}^{(i)}$  and  $(G_{s_2}^{(i)})^{\vee\vee}/G_{s_2}^{(i)}$ , we may further assume that C avoids the supports of  $(G_{s_1}^{(i)})^{\vee\vee}/G_{s_1}^{(i)}$  and  $(G_{s_2}^{(i)})^{\vee\vee}/G_{s_2}^{(i)}$  altogether, implying that

$$F_{s_1}^{**}|_C\cong \operatorname{gr}(F_{s_1})|_C\cong \operatorname{gr}(F_{s_1}|_C) \qquad \text{and} \qquad F_{s_2}^{**}|_C\cong \operatorname{gr}(F_{s_2})|_C\cong \operatorname{gr}(F_{s_2}|_C).$$

Moreover, since  $F_{s_1}^{**}|_D$  and  $F_{s_2}^{**}|_D$  are locally free, we may, by increasing *b* if necessary, assume that  $\operatorname{Hom}(F_{s_1}^{**}, F_{s_2}^{**}) = \operatorname{Hom}(F_{s_1}^{**}|_C, F_{s_2}^{**}|_C)$  and  $\operatorname{Hom}(F_{s_2}^{**}, F_{s_1}^{**}) = \operatorname{Hom}(F_{s_2}^{**}|_C, F_{s_1}^{**}|_C)$ . Finally, we can assume that  $\operatorname{H}^1(X, \mathcal{O}_X(b-a)) = 0$  so that  $\operatorname{H}^0(X, \mathcal{O}_X(b)) \to \operatorname{H}^0(X, \mathcal{O}_X(b))$  is surjective, implying that  $C = D \cap D'$  for a surface  $D' \in |\mathcal{O}_X(b)|$ . With these choices, it is sufficient to show that  $F_{s_1}|_C$  and  $F_{s_2}|_C$  are S-equivalent.

By Proposition 5.2, we have  $\lambda_{\mathcal{E}_C}(mw|_C) \cong \mathcal{L}_2^{\otimes ma^2b^2}$ , and the class  $-mw|_C \in K(X)$  is represented by a locally free sheaf M on C. Since  $F_{s_1}|_C$  is semistable on C, by Lemma 5.6 (i), for large enough m > 0 we can choose M such that

$$\mathrm{H}^{0}(C, F_{s_{1}}|_{C} \otimes M) = \mathrm{H}^{1}(C, F_{s_{1}}|_{C} \otimes M) = 0.$$

This implies that the section  $\delta_M$  of  $\lambda_{\mathcal{E}_C}(M)^{\vee} = \mathcal{L}_2^{\otimes ma^2b^2}$  is nonvanishing at  $s_1 \in S$ , hence nonvanishing everywhere. Then

$$\mathrm{H}^{0}(C, F_{s}|_{C} \otimes M) = \mathrm{H}^{1}(C, F_{s}|_{C} \otimes M) = 0$$

for every  $s \in S$ . This shows by Lemma 3.3 that C does not meet the support of  $T_s$  and  $Q_s$  for all  $s \in S$ , hence  $\mathcal{E}_C$  is a family of locally free sheaves on C parameterized by S. By Remark 5.7 the sheaves in this family are semistable, and by Lemma 5.6 (ii), they are all S-equivalent. This concludes the proof.

Assume now that  $gcd(rk(v), H^2 \cdot ch_1(v)) = 1$ , so there are no strictly PTsemistable objects and  $F_s$  is a  $\mu$ -stable sheaf for every  $s \in S$ . Applying Proposition 5.10 to this case, we obtain that the double duals  $F_s^{\vee\vee}$  are all isomorphic for  $s \in S$ , showing the first part of Theorem 5.9 (ii). Moreover, there exists a closed 1-dimensional subset  $Y \subseteq X$  with the property that  $\operatorname{Supp}(Q_s) \subseteq Y$  for all  $s \in S$ . Let  $\eta_1, \ldots, \eta_n \in X$  denote the generic points of the irreducible components of Y. The following proves the second part of Theorem 5.9 (ii).

# **Proposition 5.11.** The length $l_{\eta_i}(Q_s)$ is constant for i = 1, ..., n.

*Proof.* From the triangles

 $F_s \to \mathcal{E}_s \to T_s[-1]$  and  $F_s \to F_s^{\vee \vee} \to Q_s$ 

we get an equation for Hilbert polynomials

$$P(Q_s, m) + P(T_s, m) = P(F_s^{\vee \vee}, m) - P(\mathcal{E}_s, m),$$

where  $P(-,m) = \chi(X, (-) \otimes^{\mathbb{L}} \mathcal{O}_X(m))$ . The right hand side is independent of s since  $\mathcal{E}$  is S-perfect and  $F_s^{\vee\vee}$  is independent of s. Thus, the left hand side is independent of s. Moreover, the degrees of  $P(Q_s, m)$  and  $P(T_s, m)$  are respectively 1 and 0, so we see that the leading coefficient of  $P(Q_s, m)$  is independent of s. Now on the one hand, the leading coefficient is  $\sum_i l_{\eta_i}(Q_s)$  by the Riemann-Roch theorem, and on the other hand each quantity  $l_{\eta_i}(Q_s)$  is upper semicontinuous by Lemma 5.12 below. Since S is connected, this implies that each  $l_{\eta_i}(Q_s)$  must be constant.

**Lemma 5.12.** Let  $\eta \in X$  be a point of codimension 1. The length  $l_{\eta}(Q_s)$  is upper semicontinuous as a function of  $s \in S$ .

*Proof.* Let  $W = \operatorname{Spec} \mathcal{O}_{X,\eta}$  be the spectrum of the local ring of X at  $\eta$  and let  $\iota : W \to X$  denote the canonical monomorphism and let  $\iota_S : S \times W \to S \times X$  denote the induced map. We replace  $F_s$  and  $Q_s$  by their restrictions to W – this does not change  $l_\eta(Q_s)$ .

Define  $F := \iota_S^* \mathcal{E} \in D^b(S \times W)$ . We have  $F|_{\{s\} \times W} = \iota^* \mathcal{E}_s$ , and since localization is exact, pulling back along  $\iota$  commutes with taking cohomology sheaves, so that

$$\mathcal{H}^{i}(F|_{\{s\}\times W}) = \mathcal{H}^{i}(\iota^{*}\mathcal{E}_{s}) = \iota^{*}\mathcal{H}^{i}(\mathcal{E}_{s}), \quad i \in \mathbb{Z}.$$

In particular,  $\mathcal{H}^i(F|_{\{s\}\times W}) = 0$  for  $i \neq 0, 1$ , and also

$$\mathcal{H}^1(F|_{\{s\}\times W}) = \iota^* T_s = 0$$

since  $T_s$  is supported in codimension 3. Thus, F is a sheaf on  $S \times W$ , flat over S by [Huy06, Lemma 3.31]. Moreover, for each  $s \in S$  we have a short exact sequence

$$0 \to F_s \to F_s^{\vee \vee} \to Q_s \to 0.$$

Now  $\mathscr{E}xt^i(F_s^{\vee\vee}, \mathcal{O}_W) = 0$  for i > 0 since  $F_s^{\vee\vee}$  is reflexive on the regular 2dimensional scheme W, hence locally free. Thus, applying  $\mathscr{H}om(-, \mathcal{O}_W)$  to the above sequence and taking the long exact sequence gives isomorphisms

$$\mathscr{E}xt^1(F_s,\mathcal{O}_W)\cong \mathscr{E}xt^2(Q_s,\mathcal{O}_W), \quad \mathscr{E}xt^2(F_s,\mathcal{O}_W)\cong \mathscr{E}xt^3(Q_s,\mathcal{O}_W).$$

We claim that  $\mathscr{E}xt^3(Q_s, \mathcal{O}_W) = 0$  and  $l_\eta(Q_s) = l_\eta(\mathscr{E}xt^2(Q_s, \mathcal{O}_W))$ . To see this, we observe that  $Q_s$  has a filtration by copies of the residue field  $k(\eta)$  and length is additive in short exact sequences, so by induction it suffices to show

$$U_{\eta}(\mathscr{E}xt^{2}(k(\eta),\mathcal{O}_{W})=1, \quad \mathscr{E}xt^{3}(k(\eta),\mathcal{O}_{W})=0$$

Since  $\mathcal{O}_{X,\eta}$  is a regular local ring of dimension 2, these follow by applying  $\mathscr{H}om(-,\mathcal{O}_W)$  to the Koszul complex

$$0 \to \mathcal{O}_W \to \mathcal{O}_W^{\oplus 2} \to \mathcal{O}_W \to k(\eta) \to 0.$$

Thus, we must show that  $l_{\eta}(\mathscr{E}xt^{1}(F_{s}, \mathcal{O}_{W}))$  is upper semicontinuous as a function of s.

We temporarily spread out and replace W by a scheme finite type over  $\mathbb{C}$  and F by a coherent sheaf on  $S \times W$  flat over S in order to apply [AK80, Theorem 1.9] to the sheaves  $\mathscr{E}xt^i(F, \mathcal{O}_{S \times W})$ . First, for i = 2 and any  $s \in S$  the map

$$\mathscr{E}xt^2(F, \mathcal{O}_{S \times W})|_{\{s\} \times W} \to \mathscr{E}xt^2(F_s, \mathcal{O}_W) = 0$$

is clearly surjective, hence an isomorphism, so we get  $\mathscr{E}xt^2(F, \mathcal{O}_{S \times W}) = 0$ . Next, for i = 1 this implies that

$$\mathscr{E}xt^1(F, \mathcal{O}_{S \times W})|_{\{s\} \times W} \to \mathscr{E}xt^1(F_s, \mathcal{O}_W)$$

is an isomorphism. Thus, we have reduced to showing that  $l_{(s,\eta)}(\mathscr{E}xt^1(F, \mathcal{O}_{S\times W}))$  is upper semicontinuous as a function of s, which follows from Lemma 5.13 below.  $\Box$ 

**Lemma 5.13.** Let X and S be schemes over  $\mathbb{C}$  and let F be a quasicoherent sheaf of finite type on  $S \times X$ . Assume that the restriction  $F_s$  of F to the fiber  $\{s\} \times X$  is supported in codimension d for every  $s \in S$ , and let  $\eta \in X$  be a point of codimension d. The function that assigns to  $s \in S$  the length  $l_{\eta}((F_s)_{\eta})$  of the stalk of  $F_s$  at  $\eta$  as a module over the local ring  $\mathcal{O}_{X,\eta}$  is upper semicontinuous. *Proof.* Note that, by the assumption on dimensions, the length of  $F_s$  at  $\eta$  is indeed finite. We want to reduce the statement to the familiar fact that the fiber dimension of a quasicoherent sheaf of finite type is upper semicontinuous. We may first replace X by the Spec  $\mathcal{O}_{X,\eta}$ . Now F is set-theoretically supported on  $S \times \{\eta\}$ , so we may even replace X by Spec  $\mathcal{O}_{X,\eta}/\mathfrak{m}^n$  for sufficiently large n, where  $\mathfrak{m} \subseteq \mathcal{O}_{X,\eta}$  denotes the maximal ideal. Thus, we may assume that X is the spectrum of a local artinian ring A with maximal ideal  $\mathfrak{m}$  whose residue field  $L = A/\mathfrak{m}$  has transcendence degree  $c = \dim X - d$  over  $\mathbb{C}$ .

We can choose a set of elements  $y_1, \ldots, y_c \in A$  whose images  $\overline{y}_1, \ldots, \overline{y}_c$  in L form a transcendence basis over  $\mathbb{C}$ . These elements determine a ring homomorphism

$$\phi : \mathbb{C}[x_1, \dots, x_c] \xrightarrow{x_i \mapsto y_i} A.$$

If  $f \in \mathbb{C}[x_1, \ldots, x_c]$  is a nonzero polynomial, then the image of  $\phi(f)$  in L is nonzero since there are no algebraic relations among the  $\overline{y}_i$ 's. Thus,  $\phi(f)$  lies outside the maximal ideal  $\mathfrak{m}$ , hence is a unit. Thus, we obtain a map  $K := \mathbb{C}(x_1, \ldots, x_c) \to A$ . Since A has a filtration by copies of L and L is a finite extension of K, this map makes A into a finitely generated K-module. Thus, the induced map  $S \times \operatorname{Spec} A \to S \times \operatorname{Spec} K$  is finite, and we can view F as a quasicoherent sheaf of finite type on  $S \times \operatorname{Spec} K$ . Let  $\xi \in \operatorname{Spec} K$  be the unique point. Now on the one hand

$$l_{\xi}(F_s) = \deg(L/K) \, l_{\eta}(F_s),$$

and on the other hand  $l_{\xi}(F_s)$  is just the dimension of the fiber of  $F_s$  at  $\xi$  since Spec K is reduced, and this is an upper semicontinuous function of s.  $\Box$ 

5.3. Positivity of  $\mathcal{L}_1$ . Next we study the positivity of  $\mathcal{L}_1$  on the fibers of the morphism

$$\mathcal{M}_Z^{\mathrm{PT}}(v) \to \mathrm{M}_Z^{\mathrm{PT}}(v) \xrightarrow{\phi} \mathbb{P}$$

constructed in the previous section. Fix a  $\mathbb{C}$ -point  $p \in \mathbb{P}$ , and let  $\mathcal{M}_p$  and  $M_p$  denote the corresponding fibers over p, as described by the following cartesian diagram

(5.1) 
$$\begin{array}{c} \mathcal{M}_{Z}^{\mathrm{PT}}(v) \longrightarrow \mathrm{M}_{Z}^{\mathrm{PT}}(v) \xrightarrow{\phi} \mathbb{P} \\ \uparrow & \uparrow & \uparrow \\ \mathcal{M}_{p} \longrightarrow M_{p} \longrightarrow \mathrm{Spec}\,\mathbb{C} \end{array}$$

We aim to show the semiampleness of  $\mathcal{L}_1$  over  $\mathcal{M}_p$ . The method for producing sections involves restricting the universal complex on  $\mathcal{M}_p \times X$  to certain smooth surfaces in X. For the proof we need two auxiliary lemmas.

To lighten the notation, in the following we will denote by  $\mathcal{E}_p$  the restriction of the universal complex  $\mathcal{E}$  to  $\mathcal{M}_p \times X$ .

**Lemma 5.14.** Let a > 0 be an integer and  $D \in |\mathcal{O}_X(a)|$  a smooth surface. Consider the class

$$w = -\chi(v(m) \cdot [\mathcal{O}_D])[\mathcal{O}_X(n)] + \chi(v(n) \cdot [\mathcal{O}_D])[\mathcal{O}_X(m)] \in \mathcal{K}(X)$$

for some integers m, n. Then there exist sufficiently large choices of m, n and a such that

$$\lambda_{\mathcal{E}_p}(w \cdot [\mathcal{O}_D])^{\otimes e} = \mathcal{L}_1^{\otimes e'}$$

in  $\operatorname{Pic}(\mathcal{M}_p)$  for some integers e, e' > 0.

*Proof.* To simplify the computations, we introduce the following equivalence relation on K(X):

$$u \equiv_{\mathcal{M}_p} u'$$
 if and only if  $\lambda_{\mathcal{E}_p}(u) = \lambda_{\mathcal{E}_p}(u') \in \operatorname{Pic}(\mathcal{M}_p).$ 

Using this notation, we will show that  $ew \cdot [\mathcal{O}_D] \equiv_{\mathcal{M}_p} e'v_1(v)$  for some integers e, e' > 0. We may assume without loss of generality that m = am' and n = an' with m', n' integers. Setting  $\alpha := [\mathcal{O}_D] \in \mathcal{K}(X)$ , we can write

$$[\mathcal{O}_X(m)] = \sum_{i \ge 0} \binom{m'+i-1}{i} \alpha^i,$$

and similarly for  $[\mathcal{O}_X(n)]$ . Then a direct computation yields

$$w \cdot [\mathcal{O}_D] = \sum_{i>j} c(i,j)(-\chi(v \cdot \alpha^{i+1})\alpha^{j+1} + \chi(v \cdot \alpha^{j+1})\alpha^{i+1}),$$

where

$$c(i,j) \coloneqq \binom{m'+i-1}{i} \binom{n'+j-1}{j} - \binom{m'+j-1}{j} \binom{n'+i-1}{i} > 0$$

for  $m' \gg n' > 0$ .

Recall that by construction  $\mathcal{L}_2^{\otimes k} = \lambda_{\mathcal{E}}(v_2(v))^{\otimes k}$  is trivial on the fiber  $\mathcal{M}_p$  for some integer k > 0. Thus, we may choose a > 0 sufficiently large such that

$$-\chi(v\cdot\alpha^3)\alpha^2 + \chi(v\cdot\alpha^2)\alpha^3 = a^5v_2(v) \equiv_{\mathcal{M}_p} 0,$$

which further gives

$$\chi(v \cdot \alpha^3)(-\chi(v \cdot \alpha^2)\alpha + \chi(v \cdot \alpha)\alpha^2) \equiv_{\mathcal{M}_p} \chi(v \cdot \alpha^2)(-\chi(v \cdot \alpha^3)\alpha + \chi(v \cdot \alpha)\alpha^3)$$
$$\equiv_{\mathcal{M}_p} a^4 \chi(v \cdot \alpha^2)v_1(v).$$

Putting together the above identities modulo  $\mathcal{M}_p$ , we obtain

$$a^{3}\chi(v \cdot h^{3})w \cdot [\mathcal{O}_{D}] \equiv_{\mathcal{M}_{p}} (a^{3}\chi(v \cdot h^{3})c(2,0) + a^{4}\chi(v \cdot \alpha^{2})c(1,0))v_{1}(v).$$

We get the desired result by choosing  $m \gg n$  such that

$$(a^{3}\chi(v \cdot h^{3})c(2,0) + a^{4}\chi(v \cdot \alpha^{2})c(1,0)) > 0.$$

We will also need the following characterization of Gieseker-semistability on a given smooth surface  $D \subset X$ , which restates the results from [ÁCK07, Theorem 7.2 and Theorem 7.10]. Here, Gieseker-semistability is defined with respect to the polarization H.

**Theorem 5.15.** Let  $D \subset X$  be a smooth surface and  $P \in \mathbb{Q}[T]$  a fixed polynomial. For  $m \gg n \gg 0$ , any Castelnuovo-Mumford n-regular pure sheaf E of Hilbert polynomial P on D is Gieseker-semistable if and only if there is a map

$$U_1 \otimes \mathcal{O}_D(-m) \xrightarrow{\theta} U_0 \otimes \mathcal{O}_D(-n)$$

where  $U_1$  and  $U_0$  are vector spaces, such that the linear map

$$\operatorname{Hom}(\theta, E) : \operatorname{Hom}(U_0, \operatorname{H}^0(E(n))) \to \operatorname{Hom}(U_1, \operatorname{H}^0(E(m)))$$

is invertible, i.e.  $\delta_{\theta}(E) \coloneqq \det \operatorname{Hom}(\theta, E) \neq 0$ . Moreover, if E and F are Giesekersemistable sheaves of Hilbert polynomial P that are not S-equivalent on D, then there is a map  $\theta$  as above such that  $\delta_{\theta}(E) \neq 0$  and  $\delta_{\theta}(F) = 0$ . Remark 5.16. a) The statement in [ÁCK07, Theorem 7.2] is more exact, as it gives precise conditions on m and n under which the conclusion holds. Due to the boundedness of Gieseker-semistable sheaves of fixed Hilbert polynomial [HL10, Theorem 3.3.7], these conditions are always fulfilled for  $m \gg n \gg 0$ , i.e. there is an integer  $n_0 > 0$  such that we can take any  $n \ge n_0$ , and once n is fixed, there is some m(n) > n such that we can take any  $m \ge m(n)$ .

b) If E is Gieseker-semistable as in the statement, then there exists a dense open subset  $V \subset \operatorname{Hom}(U_1 \otimes \mathcal{O}_D(-m), U_0 \otimes \mathcal{O}_D(-n))$  such that  $\operatorname{Hom}(\theta, E)$  is invertible for every  $\theta \in V$ .

c) If we consider the complex

$$K: K_1 = U_1 \otimes \mathcal{O}_D(-m) \xrightarrow{\theta} U_0 \otimes \mathcal{O}_D(-n) = K_0,$$

then the condition that  $\operatorname{Hom}(\theta, E)$  is invertible is equivalent to  $\operatorname{R}\Gamma(E \otimes^{\mathbb{L}} K^{\vee}) = 0$ . To avoid any confusion, in the following we will view  $K^{\vee} = [K_0^{\vee} \to K_1^{\vee}]$  as a complex with  $K_0^{\vee}$  in degree 0 and  $K_1^{\vee}$  in degree 1.

In what follows, for every  $\mathbb{C}$ -point  $t \in \mathcal{M}_p$ , we denote by  $E_t \in \mathcal{A}^p(X)$  the corresponding PT-semistable object and denote

$$F_t = \mathcal{H}^0(E_t), \quad T_t = \mathcal{H}^1(E_t), \quad Q_t = F_t^{\vee \vee}/F_t, \quad F_t^{**} = \operatorname{gr}(F)^{\vee \vee},$$

where  $\operatorname{gr}(F)$  is the graded module corresponding to a Jordan-Hölder filtration of  $F_t$ . Since  $\mathcal{M}_p$  is quasi-compact, Theorem 5.9 and Proposition 5.10 ensure that we have the following description of the fiber  $\mathcal{M}_p$ :

(\*) There exists a closed 1-dimensional subset  $Y \subset X$  such that for every  $\mathbb{C}$ -point  $t \in \mathcal{M}_p$  the sheaf  $Q_t$  supported on Y. Moreover, the sheaves  $F_t^{**}$  for  $t \in \mathcal{M}_p$  are all isomorphic.

## **Theorem 5.17.** The restriction of the line bundle $\mathcal{L}_1$ to $\mathcal{M}_p$ is semiample.

*Proof.* Fix a  $\mathbb{C}$ -point  $t_0 \in \mathcal{M}_p$ . By Proposition 2.3 we know that  $F_{t_0}$  is torsion-free and semistable in  $\operatorname{Coh}_{3,1}(X)$ . Hence, for a general surface  $D \in |\mathcal{O}_X(a)|$  of sufficiently large degree and avoiding the support of  $T_{t_0}$ , we get by the Mehta-Ramanathan type restriction theorem in [Pav24] that  $E_{t_0}|_D^{\mathbb{L}} = F_{t_0}|_D$  is a Gieseker-semistable torsion-free sheaf on D. According to Theorem 5.15 and the remarks below it, for  $m \gg n \gg 0$  there is a general map

$$K: K_1 = U_1 \otimes \mathcal{O}_D(-m) \xrightarrow{\theta} U_0 \otimes \mathcal{O}_D(-n) = K_0,$$

with  $U_0, U_1$  vector spaces, such that  $\mathbb{H}^i(E_{t_0}|_D^{\mathbb{L}} \otimes^{\mathbb{L}} K^{\vee}) = 0$  for all *i*. Choosing *D* such that it meets *Y* transversely, we may further assume by  $(\star)$  that the cohomology sheaf  $\mathcal{H}^1(E_t|_D^{\mathbb{L}}) = T_t|_D$  is supported on the 0-dimensional subset  $Y \cap D$  for all  $t \in \mathcal{M}_p$ .

Let P denote the Hilbert polynomial of  $F_{t_0}|_D$ . Then the equality

$$P(n)\dim(U_0) = P(m)\dim(U_1)$$

implies that  $\dim(U_1) < \dim(U_0)$  (since we can assume that P(N) grows monotonically for  $N \ge n$ ). Thus, since for any point  $x \in D$ , a general map  $\mathcal{O}_D(n) \to \mathcal{O}_D(m)$ is surjective at x, the same is true for a general map  $U_0^{\vee} \otimes \mathcal{O}_D(n) \to U_1^{\vee} \otimes \mathcal{O}_D(m)$ . In particular, we may assume that  $\theta^{\vee}$  is surjective at the finitely many points of  $Y \cap D$ , and so the map

$$T_t|_D \otimes K_0^{\vee} \xrightarrow{\mathrm{id} \otimes \theta^{\vee}} T_t|_D \otimes K_1^{\vee}$$

is surjective for all  $t \in \mathcal{M}_p$ . Since  $\mathcal{M}_p$  is quasi-compact, by choosing *n* sufficiently large, we may also assume that  $F_t|_D$  is *n*-regular in the sense of Castelnuovo-Mumford for all  $t \in \mathcal{M}_p$ . Thus by Lemma 5.18 below we get

$$\mathbb{H}^{2}(E_{t}|_{D}^{\mathbb{L}}\otimes^{\mathbb{L}}K^{\vee}) = \mathrm{H}^{0}(\mathrm{Coker}(T_{t}|_{D}\otimes K_{0}^{\vee} \xrightarrow{\mathrm{id}\otimes\theta^{\vee}} T_{t}|_{D}\otimes K_{1}^{\vee})),$$

implying that  $\mathbb{H}^i(E_t|_D^{\mathbb{L}} \otimes^{\mathbb{L}} K^{\vee}) = 0$  for  $i \neq 0, 1$ . Then by Lemma 4.1 the line bundle

$$\lambda_{\mathcal{E}_D}([K^{\vee}])^{\vee} \cong \lambda_{\mathcal{E}_D}([K])$$

has a global section  $\delta_K$  on  $\mathcal{M}_p$  non-vanishing at  $t_0$ , where we denoted by  $\mathcal{E}_D$  the restriction of the universal complex  $\mathcal{E}_p$  to  $\mathcal{M}_p \times D$ . Using again the fact that  $\mathcal{M}_p$  is quasi-compact, we obtain that  $\lambda_{\mathcal{E}_D}([K])$  is semiample on  $\mathcal{M}_p$ .

Next we relate  $\lambda_{\mathcal{E}_D}([K])$  and  $\mathcal{L}_1|_{\mathcal{M}_p}$ . Since  $\mathbb{H}^i(E_t|_D^{\mathbb{L}} \otimes^{\mathbb{L}} K^{\vee}) = 0$  for  $i \neq 0, 1$  and for all  $t \in \mathcal{M}_p$ , we get by cohomology and base change (see [Sta20, Tag 0A1K]) that  $\mathbb{R}^i p_*(\mathcal{E}_D \otimes^{\mathbb{L}} p^* K^{\vee}) = 0$  for  $i \neq 0, 1$ , where  $p : \mathcal{M}_p \times D \to \mathcal{M}_p$  is the first projection. Therefore

$$\lambda_{\mathcal{E}_D}([K^{\vee}]) \cong \lambda_{\mathcal{E}_D}([K_0^{\vee}]) \otimes \lambda_{\mathcal{E}_D}([K_1^{\vee}])^{\vee} \cong \lambda_{\mathcal{E}_D}([K_0^{\vee}] - [K_1^{\vee}]),$$

and also

$$\lambda_{\mathcal{E}_D}([K]) \cong \lambda_{\mathcal{E}_D}([K_0] - [K_1]).$$

Now let  $d_1 = \dim(U_1)$  and  $d_0 = \dim(U_0)$ . Using  $d_1P(m) = d_0P(n)$ , we get

$$P(m)([K_0] - [K_1]) = d_0(P(n)[\mathcal{O}_D(m)] - P(m)[\mathcal{O}_D(n)]) = d_0w|_D$$

in K(D), where

$$w \coloneqq -\chi(v(m) \cdot [\mathcal{O}_D])[\mathcal{O}_X(n)] + \chi(v(n) \cdot [\mathcal{O}_D])[\mathcal{O}_X(m)] \in \mathcal{K}(X).$$

Hence

$$\lambda_{\mathcal{E}_p}(w \cdot [\mathcal{O}_D])^{d_0} \cong \lambda_{\mathcal{E}_D}(w|_D)^{d_0} \cong \lambda_{\mathcal{E}_D}([K_0] - [K_1])^{P(m)}$$

(the first isomorphism follows as in Lemma 5.1). By using Lemma 5.14 we finally obtain the semiampleness of  $\mathcal{L}_1$  over  $\mathcal{M}_p$ .

Lemma 5.18. Let E be a PT-semistable object fitting in

$$F \to E \to T[-1],$$

and let  $D \subset X$  be a smooth divisor. Let  $m \gg n \gg 0$  such that  $F|_D$  is n-regular, and consider a complex

$$K: K_1 = U_1 \otimes \mathcal{O}_D(-m) \xrightarrow{\theta} U_0 \otimes \mathcal{O}_D(-n) = K_0$$

as before. Then

$$\mathbb{H}^{2}(E|_{D}^{\mathbb{L}} \otimes^{\mathbb{L}} K^{\vee}) = \mathrm{H}^{0}(\mathrm{Coker}(T|_{D} \otimes K_{0}^{\vee} \xrightarrow{\mathrm{id} \otimes \theta^{\vee}} T|_{D} \otimes K_{1}^{\vee})).$$

In particular, if  $\theta^{\vee}$  is surjective on the support of  $T|_D$ , then  $\mathbb{H}^2(E|_D^{\mathbb{L}} \otimes^{\mathbb{L}} K^{\vee}) = 0$ .

*Proof.* By Lemma 3.4 there is an exact triangle

$$\mathcal{H}^0(E|_D^{\mathbb{L}}) \to E|_D^{\mathbb{L}} \to T|_D[-1],$$

leading to a long exact sequence in cohomology

$$\ldots \to \mathbb{H}^2(\mathcal{H}^0(E|_D^{\mathbb{L}}) \otimes^{\mathbb{L}} K^{\vee}) \to \mathbb{H}^2(E|_D^{\mathbb{L}} \otimes^{\mathbb{L}} K^{\vee}) \to \mathbb{H}^1(T|_D \otimes^{\mathbb{L}} K^{\vee}) \to 0.$$

Using the spectral sequence

$$E_2^{p,q} = \mathrm{H}^p(\mathcal{H}^q(T|_D \otimes^{\mathbb{L}} K^{\vee})) \Rightarrow \mathbb{H}^{p+q}(T|_D \otimes^{\mathbb{L}} K^{\vee})$$

and the fact that  $T|_D$  is a 0-dimensional sheaf, one obtains

$$\begin{split} \mathbb{H}^{1}(T|_{D} \otimes^{\mathbb{L}} K^{\vee}) &= \mathrm{H}^{0}(\mathcal{H}^{1}(T|_{D} \otimes^{\mathbb{L}} K^{\vee})) \\ &= \mathrm{H}^{0}(\mathrm{Coker}(T|_{D} \otimes K_{0}^{\vee} \xrightarrow{\mathrm{id} \otimes \theta^{\vee}} T|_{D} \otimes K_{1}^{\vee})). \end{split}$$

We show next that  $\mathbb{H}^2(\mathcal{H}^0(E|_D^{\mathbb{L}}) \otimes^{\mathbb{L}} K^{\vee}) = 0$ . As we saw in the proof of Lemma 3.4, there is a short exact sequence

$$0 \to F|_D \to \mathcal{H}^0(E|_D^{\mathbb{L}}) \to \mathcal{H}^{-1}(T|_D^{\mathbb{L}}) \to 0$$

with  $\mathcal{H}^{-1}(T|_D^{\mathbb{L}})$  a 0-dimensional sheaf, and so we obtain a surjective map

$$\mathbb{H}^{2}(F|_{D} \otimes^{\mathbb{L}} K^{\vee}) \to \mathbb{H}^{2}(\mathcal{H}^{0}(E_{t}|_{D}^{\mathbb{L}}) \otimes^{\mathbb{L}} K^{\vee}) \to 0.$$

To compute  $\mathbb{H}^2(F|_D \otimes^{\mathbb{L}} K^{\vee})$  we will use the exact triangle

$$F|_D \otimes^{\mathbb{L}} K_0^{\vee} \to F|_D \otimes^{\mathbb{L}} K^{\vee} \to F|_D \otimes^{\mathbb{L}} K_1^{\vee}[-1].$$

As  $K_0$  and  $K_1$  are locally free we have  $F|_D \otimes^{\mathbb{L}} K_0^{\vee} = F|_D \otimes K_0^{\vee}$  and  $F|_D \otimes^{\mathbb{L}} K_1^{\vee} = F|_D \otimes K_1^{\vee}$ . Moreover, since  $F|_D$  is *n*-regular, we know that

(5.2) 
$$\mathrm{H}^{i}(F|_{D} \otimes^{\mathbb{L}} K_{0}^{\vee}) = \mathrm{H}^{i}(F|_{D} \otimes^{\mathbb{L}} K_{1}^{\vee}) = 0 \quad \text{for } i \geq 1.$$

Hence the long exact sequences in cohomology induced by the above triangle shows that  $\mathbb{H}^2(F|_D \otimes^{\mathbb{L}} K^{\vee}) = 0$ , from which one gets the statement.  $\Box$ 

For the next result, we recall that any coherent sheaf F on X that is semistable in  $\operatorname{Coh}_{3,1}(X)$  admits Jordan-Hölder filtrations with respect to this notion of semistability, and moreover the graded modules of any two such filtrations are isomorphic in  $\operatorname{Coh}_{3,1}(X)$  [HL10, Theorem 1.6.7]. In this case, we say that two coherent sheaves are *S*-equivalent in  $\operatorname{Coh}_{3,1}(X)$  if they are semistable and have isomorphic graded modules in  $\operatorname{Coh}_{3,1}(X)$ .

**Proposition 5.19.** Let  $t_1, t_2 \in \mathcal{M}_p$  be two  $\mathbb{C}$ -points such that  $F_{t_1}$  and  $F_{t_2}$  are not S-equivalent in  $\operatorname{Coh}_{3,1}(X)$ . Then there is a positive power of the line bundle  $\mathcal{L}_1|_{\mathcal{M}_p}$  which separates  $t_1$  and  $t_2$ .

Proof. Since  $F_{t_1}$  and  $F_{t_2}$  are semistable in  $\operatorname{Coh}_{3,1}(X)$ , they admit some Jordan-Hölder filtrations  $F_{t_1}^{\bullet}$  and  $F_{t_2}^{\bullet}$ , respectively, whose factors are torsion-free and stable in  $\operatorname{Coh}_{3,1}(X)$ . By assumption the corresponding graded modules  $G_{t_1} \coloneqq \operatorname{gr}(F_{t_1}^{\bullet})$  and  $G_{t_2} \coloneqq \operatorname{gr}(F_{t_1}^{\bullet})$  are non-isomorphic. Recall that for j = 1, 2, we have  $\operatorname{hd}(F_{t_j}) \leq 1$  (as we remarked after Proposition 2.3). By using [Sta20, Tag 065S], which characterizes homological dimension in short exact sequences, we may choose the filtrations  $F_{t_j}^{\bullet}$  so that also  $\operatorname{hd}(G_{t_j}) \leq 1$ . Moreover, by the Mehta-Ramanathan type restriction theorems in [Pav24], for a general surface  $D \in |\mathcal{O}_X(a)|$  of sufficiently large degree we may also assume that the restrictions  $F_{t_1}|_D$  and  $F_{t_2}|_D$  are Gieseker-semistable, and that  $G_{t_1}|_D$  and  $G_{t_2}|_D$  are Gieseker-polystable. In particular the restrictions

 $F_{t_1}^{\bullet}|_D$  and  $F_{t_2}^{\bullet}|_D$  yield Jordan-Hölder filtrations of  $F_{t_1}|_D$  and  $F_{t_2}|_D$ , respectively, with respect to the notion of Gieseker-semistability on D, implying that

$$G_{t_1}|_D = \operatorname{gr}(F_{t_1}^{\bullet}|_D)$$
 and  $G_{t_2}|_D = \operatorname{gr}(F_{t_1}^{\bullet}|_D).$ 

Now consider the short exact sequence

$$0 \to G_{t_2}(-D) \to G_{t_2} \to G_{t_2}|_D \to 0$$

Applying  $Hom(G_{t_1}, -)$  to the above sequence, we get the long exact sequence

 $\operatorname{Hom}(G_{t_1}, G_{t_2}(-D)) \to \operatorname{Hom}(G_{t_1}, G_{t_2}) \to \operatorname{Hom}(G_{t_1}|_D, G_{t_2}|_D) \to \operatorname{Ext}^1(G_{t_1}, G_{t_2}(-D)).$ Since  $\operatorname{hd}(G_{t_1}) \leq 1$ , we have  $\mathscr{E}xt^i(G_{t_1}, G_{t_2}) = 0$  for  $i \geq 2$ . Therefore, by using Serre Duality and the local-to-global Ext spectral sequence

$$E_2^{p,q} = \mathrm{H}^p(X, \mathscr{E}xt^q(G_{t_1}, G_{t_2})) \Rightarrow \mathrm{Ext}^{p+q}(G_{t_1}, G_{t_2}),$$

one obtains

$$\operatorname{Ext}^{i}(G_{t_{1}},G_{t_{2}}(-D)) \cong \operatorname{H}^{0}(X,\mathscr{E}xt^{3-i}(G_{t_{1}},G_{t_{2}}) \otimes \omega_{X}(D)) = 0$$

for i = 0, 1 and  $a \gg 0$ , where  $\omega_X$  is the dualizing line bundle on X. Then

$$\operatorname{Hom}(G_{t_1}, G_{t_2}) \to \operatorname{Hom}(G_{t_1}|_D, G_{t_2}|_D)$$

is an isomorphism, and the same holds with  $G_{t_1}$  and  $G_{t_2}$  interchanged. We deduce that the restrictions  $G_{t_1}|_D$  and  $G_{t_2}|_D$  are still non-isomorphic. Thus, for  $D \in$  $|\mathcal{O}_X(a)|$  general of large enough degree, we get that  $E_{t_1}|_D^{\mathbb{L}} = F_{t_1}|_D$  and  $E_{t_2}|_D^{\mathbb{L}} =$  $F_{t_2}|_D$  are Gieseker-semistable sheaves of the same Hilbert polynomial that are not S-equivalent on D. By Theorem 5.15 there is a map

$$K: K_1 = U_1 \otimes \mathcal{O}_D(-m) \xrightarrow{\theta} U_0 \otimes \mathcal{O}_D(-n) = K_0$$

such that  $\operatorname{Hom}(\theta, F_{t_1}|_D)$  is invertible, but  $\operatorname{Hom}(\theta, F_{t_2}|_D)$  is not so. As in the proof of Theorem 5.17 we obtain a global section  $\delta_K$  of  $\lambda_{\mathcal{E}_D}([K])$  on  $\mathcal{M}_p$  such that  $\delta_K(t_1) \neq 0$ but  $\delta_K(t_2) = 0$ . As we saw before,  $\lambda_{\mathcal{E}_D}([K])^{\otimes e} \cong \mathcal{L}_1|_{\mathcal{M}_p}^{\otimes e'}$  for some integers e, e' > 0, which is enough to produce global sections for some power of  $\mathcal{L}_1|_{\mathcal{M}_p}$  separating  $t_1$ and  $t_2$ .  $\Box$ 

5.4. **Positivity of**  $\mathcal{L}_0$ . It remains to study the positivity of  $\mathcal{L}_0$ . Here we must assume that  $\operatorname{rk}(v)$  and  $H^2 \cdot \operatorname{ch}_1(v)$  are coprime since we use [BR21, Proposition 5.5], which crucially depends on the characterization of PT-stable objects from Proposition 2.3. We also assume that  $U = \operatorname{td}_X$  in the definition of the stability function Z to ensure that the line bundles  $\mathcal{L}_i$  for i = 0, 1, 2 descend to line bundles  $L_i$  on the good moduli space  $\operatorname{M}_Z^{\operatorname{PT}}(v)$ , cf. Proposition 4.4.

Fix a C-point  $p \in \mathbb{P}$ , and consider the good moduli space  $\mathcal{M}_p \to \mathcal{M}_p$  given by (5.1). By Theorem 5.17 we we can find an integer n > 0 such that  $L_1|_{\mathcal{M}_p}^{\otimes n}$  is globally generated on  $\mathcal{M}_p$ . Moreover, by choosing n sufficiently large, we may assume that the induced morphism

$$M_p \xrightarrow{\eta} W \coloneqq \mathbb{P}\Gamma(M_p, L_1|_{M_p}^{\otimes n})$$

has connected fibers. The vector space  $\Gamma(M_p, L_1|_{M_p}^{\otimes n})$  is finitely generated since  $M_p$  is proper.

Now fix a  $\mathbb{C}$ -point  $y \in W$ , and consider the following commutative diagram



where  $\mathcal{N}$ , resp. N, denotes the fiber of  $\mathcal{M}_p$ , resp.  $\mathcal{M}_p$ , over y. In particular the canonical map  $\pi : \mathcal{N} \to N$  is a proper good moduli space. As we are in the coprime case, by Proposition 5.19 and Proposition 2.3, the geometric points  $s \in \mathcal{N}$  correspond to exact triangles of the form

$$F \to E_s \to T_s[-1]$$

where F is a fixed  $\mu$ -stable torsion-free sheaf of homological dimension at most 1.

Set  $r := \operatorname{rk}(v)$  and  $\beta := \operatorname{ch}_1(v)$ . In the following we denote by  $\mathcal{M}_Z^{\operatorname{PT}}(r,\beta)$ , resp.  $\operatorname{M}_Z^{\operatorname{PT}}(r,\beta)$ , the disjoint union of all moduli stacks  $\mathcal{M}_Z^{\operatorname{PT}}(c)$ , resp. all moduli spaces  $\operatorname{M}_Z^{\operatorname{PT}}(c)$ , over all numerical classes  $c \in \operatorname{K}_{\operatorname{num}}(X)$  with  $\operatorname{rk}(c) = r$  and  $\operatorname{ch}_1(c) = \beta$ .

**Proposition 5.20.** If rk(v) and  $H^2 \cdot ch_1(v)$  are coprime, then the restriction of the line bundle  $L_0$  to N is ample.

*Proof.* Denote by  $\operatorname{Quot}(\mathscr{E}xt^1(F, \mathcal{O}_X))$  the Quot scheme of zero-dimensional quotients of  $\mathscr{E}xt^1(F, \mathcal{O}_X)$  on X. By [BR21, Proposition 5.5] there exists a closed embedding  $\psi_F$  described by the following diagram



sending a geometric quotient  $[q : \mathscr{E}xt^1(F, \mathcal{O}_X) \to Q]$  in  $\operatorname{Quot}(\mathscr{E}xt^1(F, \mathcal{O}_X))$  to a suitable PT-stable object

$$F \to E \to Q^D[-1]$$

with  $Q^D = \mathscr{E}xt^3(Q, \mathcal{O}_X)$  is the dual of Q on X (see below Remark 5.2 in [BR21] for the construction). We note that the characterization of PT-stable objects from Proposition 2.3 is necessary for the map  $\psi_F$  to be well-defined with codomain  $M_Z^{PT}(r,\beta)$ . Then the base change of  $\psi_F$  via the closed subspace  $N \subset M_Z^{PT}(r,\beta)$  gives a commutative diagram



where  $Q_1 \subset \text{Quot}(\mathscr{E}xt^1(F, \mathcal{O}_X))$  is the induced closed subscheme and  $\psi_{F,N}$  is a closed embedding. Moreover, note that  $\psi_{F,N}$  is surjective since any  $\mathbb{C}$ -point  $s \in N$  corresponding to a PT-stable object

$$F \to E_s \to T_s[-1]$$

gives rise to a surjection  $\mathscr{E}xt^1(F, \mathcal{O}_X) \to Q$ , with  $Q = \mathscr{E}xt^3(T_s, \mathcal{O}_X)$ . Indeed, this can be seen by applying  $\mathcal{H}om(-, \mathcal{O}_X)$  to the above exact triangle.

Let  $q: Q_1 \times X \to X$  denote the natural projection, and let  $\mathcal{Q}$  denote the universal quotient on  $Q_1 \times X$ . Then the pullback  $\mathcal{G}$  of the universal complex  $\mathcal{E}$  on  $\mathcal{N} \times X$  via  $\varphi_{F,N}$  fits in an extension

$$q^*F \to \mathcal{G} \to \mathcal{Q}^D[-1]$$

over  $Q_1 \times X$ , viewed as an element in  $\operatorname{Ext}^1_{Q_1 \times X}(\mathcal{Q}^D[-1], q^*F)$ , cf. [BR21, Proposition 5.5]. We obtain

$$\psi_{F,N}^*(L_0|_N) \cong \varphi_{F,N}^*(\mathcal{L}_0|_{\mathcal{N}}) \cong \lambda_{\varphi_{F,N}^*} \mathcal{E}(v_0(v)) \cong \lambda_{\mathcal{Q}^D}(v_0(v))^{\vee} \cong \lambda_{\mathcal{Q}}(v_0(v))^{\vee},$$

where the last isomorphism holds true since by duality we have

$$\operatorname{Ext}^{1}_{Q_{1} \times X}(\mathcal{Q}^{D}[-1], q^{*}F) \cong \operatorname{Ext}^{1}_{Q_{1} \times X}((q^{*}F)^{\vee}[1], \mathcal{Q}[-1]).$$

We saw above that  $\psi_{F,N}$  is a surjective closed embedding, hence the ampleness of  $L_0$  over N will follow once we show that  $\lambda_{\mathcal{Q}}(v_0(v))^{\vee}$  is ample over  $Q_1$ . Since  $\mathcal{Q}$  is a flat family of 0-dimensional sheaves, we get

$$\lambda_{\mathcal{Q}}(v_0(v))^{\vee} \cong \lambda_{\mathcal{Q}}([\mathcal{O}_X])^{\chi(v \cdot h^3)}$$

which is ample on  $Q_1$ , cf. [HL10, Proposition 2.2.5].

Putting everything together, we obtain our main ampleness result, which in turn implies the second part of Theorem 1.1.

**Theorem 5.21.** Assume that  $\operatorname{rk}(v)$  and  $H^2 \cdot \operatorname{ch}_1(v)$  are coprime, and that  $U = \operatorname{td}_X$  in the definition of the PT-stability function Z. Then there exists an ample line bundle obtained as a linear combination with positive coefficients of  $L_2$ ,  $L_1$  and  $L_0$  on  $\operatorname{M}_{Z}^{\operatorname{PT}}(v)$ .

*Proof.* We work under the same notation as above, and consider the proper map  $\eta: M_p \to W$  induced by the semiample line bundle  $L_1|_{M_p}$  for some  $p \in \mathbb{P}$ . By Proposition 5.20 the line bundle  $L_0|_{M_p}$  is ample on each fiber of  $\eta$ , which implies that  $L_0|_{M_p}$  is  $\eta$ -ample, cf. [Laz04, Theorem 1.7.8]. Moreover, by [Laz04, Proposition 1.7.10], the line bundle

$$L_0|_{M_p} \otimes \eta^* \mathcal{O}_W(m)$$

is ample on  $M_p$  for  $m \gg 0$ . By construction  $\eta^* \mathcal{O}_W(1)$  is isomorphic to a power of  $L_1|_{M_p}$ , from which we deduce that  $L_0 \otimes L_1^m$  is ample on  $M_p$  for some large integer m > 0. As we will see below, we can choose m independently of p.

Consider now the map  $\phi : M_Z^{PT}(v) \to \mathbb{P}$  induced by the semiample line bundle  $L_2$ , as defined in Section 5.2. In particular  $\phi^* \mathcal{O}_{\mathbb{P}}(1)$  is isomorphic to a power of  $L_2$ . Since  $\mathbb{P}$  is noetherian and the property of a line bundle of being ample is open in families, cf. [Laz04, Theorem 1.2.17], we can find a large enough integer m > 0 such that  $L_0 \otimes L_1^m$  is ample on every fiber of  $\phi$ . Applying the same argument as before, we obtain that the line bundle

$$L_0 \otimes L_1^m \otimes L_2^n$$

is ample on  $\mathcal{M}_Z^{\mathrm{PT}}(v)$  for  $m, n \gg 0$ .

$$\square$$

### References

- [ÁCK07] Luis Álvarez-Cónsul and Alastair King. A functorial construction of moduli of sheaves. Invent. Math., 168(3):613–666, 2007.
- [AHLH23] Jarod Alper, Daniel Halpern-Leistner, and Jochen Heinloth. Existence of moduli spaces for algebraic stacks. Invent. Math., 234(3):949–1038, 2023.
- [AK80] Allen B. Altman and Steven L. Kleiman. Compactifying the Picard scheme. Adv. in Math., 35(1):50–112, 1980.
- [Alp08] Jarod Alper. Good moduli spaces for Artin stacks. ProQuest LLC, Ann Arbor, MI, 2008. Thesis (Ph.D.)–Stanford University.
- [Bay09] Arend Bayer. Polynomial Bridgeland stability conditions and the large volume limit. Geom. Topol., 13(4):2389–2425, 2009.
- [BDF+22] Pieter Belmans, Chiara Damiolini, Hans Franzen, Victoria Hoskins, Svetlana Makarova, and Tuomas Tajakka. Projectivity and effective global generation of determinantal line bundles on quiver moduli. Preprint, arXiv:2210.00033 [math.AG] (2022), 2022.
- [BR21] Sjoerd V. Beentjes and Andrea T. Ricolfi. Virtual counts on Quot schemes and the higher rank local DT/PT correspondence. *Math. Res. Lett.*, 28(4):967–1032, 2021.
- [HL10] Daniel Huybrechts and Manfred Lehn. The geometry of moduli spaces of sheaves. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 2010.
- [Huy06] D. Huybrechts. Fourier-Mukai transforms in algebraic geometry. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2006.
- [Laz04] Robert Lazarsfeld. Positivity in algebraic geometry I: Classical setting: Line bundles and linear series. II: Positivity for vector bundles, and multiplier ideals. Berlin: Springer, 2004.
- [Le 92] Joseph Le Potier. Fibré déterminant et courbes de saut sur les surfaces algébriques. In Complex projective geometry, 1989 in Bergen, pages 213–240. Cambridge University Press, 1992.
- [Li93] Jun Li. Algebraic geometric interpretation of Donaldson's polynomial invariants. J. Differ. Geom., 37(2):417–466, 1993.
- [Lie06] Max Lieblich. Moduli of complexes on a proper morphism. J. Algebraic Geom., 15(1):175–206, 2006.
- [Lo11] Jason Lo. Moduli of PT-semistable objects I. J. Algebra, 339:203–222, 2011.
- [Lo13] Jason Lo. Moduli of PT-semistable objects II. Trans. Amer. Math. Soc., 365(9):4539– 4573, 2013.
- [Lo21] Jason Lo. A relation between higher-rank PT-stable objects and quotients of coherent sheaves. Kyoto J. Math., 61(4):815–842, 2021.
- [LP93] Joseph Le Potier. Systèmes cohérents et structures de niveau., volume 214 of Astérisque. Paris: Société Mathématique de France, 1993.
- [MR82] V. B. Mehta and A. Ramanathan. Semistable sheaves on projective varieties and their restriction to curves. *Math. Ann.*, 258:213–224, 1982.
- [MR84] V. B. Mehta and A. Ramanathan. Restriction of stable sheaves and representations of the fundamental group. *Invent. Math.*, 77:163–172, 1984.
- [MS17] Emanuele Macrì and Benjamin Schmidt. Lectures on Bridgeland stability. In Moduli of curves. CIMAT Guanajuato, Mexico 2016. Lecture notes of a CIMPA-ICTP school, Guanajuato, Mexico, February 22 – March 4, 2016, pages 139–211. Cham: Springer, 2017.
- [Pav24] Mihai Pavel. Restriction theorems for semistable sheaves. Doc. Math., 29(3):597–625, 2024.
- [PT09] R. Pandharipande and R. P. Thomas. Curve counting via stable pairs in the derived category. Invent. Math., 178(2):407–447, 2009.
- [Ses93] C. S. Seshadri. Vector bundles on curves. In *Linear algebraic groups and their repre*sentations (Los Angeles, CA, 1992), volume 153 of Contemp. Math., pages 163–200. Amer. Math. Soc., Providence, RI, 1993.
- [Sta20] The Stacks project authors. The stacks project. https://stacks.math.columbia.edu, 2020.
- [Taj21] Tuomas Tajakka. Projective moduli spaces of complexes of sheaves. 2021. PhD Thesis.

# MIHAI PAVEL AND TUOMAS TAJAKKA

[Taj23] Tuomas Tajakka. Uhlenbeck compactification as a Bridgeland moduli space. Int. Math. Res. Not., 2023(6):4952–4997, 2023.

[Tod09] Yukinobu Toda. Limit stable objects on Calabi-Yau 3-folds. Duke Math. J., 149(1):157–208, 2009.

MIHAI PAVEL SIMION STOILOW INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY 21 CALEA GRIVITEI STREET, 010702, BUCHAREST, ROMANIA Email address: cpavel@imar.ro

TUOMAS TAJAKKA Email address: tuomas.tajakka@gmail.com