UNIFORM ESTIMATES FOR ELLIPTIC EQUATIONS WITH CARATHEODORY NONLINEARITIES AT THE INTERIOR AND ON THE BOUNDARY

EDGAR ANTONIO, MARTÍN P. ÁRCIGA-ALEJANDRE, ROSA PARDO, AND JORGE SÁNCHEZ ORTIZ

ABSTRACT. We establish an explicit uniform a priori estimate for weak solutions to slightly subcritical elliptic problems with nonlinearities simultaneously at the interior and on the boundary. Our explicit $L^{\infty}(\Omega)$ a priori estimates are in terms of powers of their $H^1(\Omega)$ norms. To prove our result, we combine a De Giorgi-Nash-Moser's iteration scheme together with elliptic regularity and the Gagliardo-Nirenberg's interpolation inequality.

MSC 2020: Primary 35B45; Secondary 35B66, 35B33, 35J75, 35J25.

Keywords: A priori estimates, slightly subcritical non-linearities, L^{∞} a priori bounds, nonlinear boundary conditions.

1. INTRODUCTION

Let us consider the boundary value problem with nonlinear boundary conditions

(1.1) $\begin{cases} -\Delta u + u = f(x, u), & x \in \Omega, \\ \frac{\partial u}{\partial \eta} = f_B(x, u), & x \in \partial \Omega, \end{cases}$

where $\Omega \subset \mathbb{R}^N$, (N > 2), is an open, connected, bounded domain with C^2 boundary, $\partial/\partial \eta = \eta \cdot \nabla$ is the (unit) outer normal derivative, and the functions $f: \Omega \times \mathbb{R} \to \mathbb{R}$, and $f_B: \partial\Omega \times \mathbb{R} \to \mathbb{R}$, are both slightly subcritical Carathéodory functions. We give in (f1)-(f2) and (f_B1)-(f_B2) the precise statement of the hypothesis on the nonlinearities at the interior, and on the boundary respectively.

The third author is supported by grants PID2022-137074NB-I00, MICINN, Spain, and by UCM, Spain, Grupo 920894.

2 E. ANTONIO, M. ARCIGA-ALEJANDRE, R. PARDO, AND J. SANCHEZ-ORTIZ

Our goal is to establish explicit $L^{\infty}(\Omega)$ a priori estimates for weak solutions to (1.1), in terms of powers of their $H^1(\Omega)$ norms (see Theorem 2.2). Our estimates are valid to positive solutions and to changing sign solutions. Consequently, any sequence of solutions to (1.1), uniformly bounded in the $H^1(\Omega)$ norm, is also uniformly bounded in the $L^{\infty}(\Omega)$ norm.

Our techniques are based in an iterative process due to Moser, in the elliptic regularity theory, and in the Gagliardo–Nirenberg interpolation inequalities.

For the Homogeneous Dirichlet boundary conditions, by a Moser's type procedure, it is well known that weak solutions to a subcritical or even critical elliptic problem are in $L^q(\Omega)$ for all $1 < q < \infty$ (see [18, Lemma 1], see also [8, Section 2.2], [24, Lemma B.3]. Moreover, by elliptic regularity, the solutions are in $L^{\infty}(\Omega)$.

Moser's results can be extended to the case of nonlinear boundary conditions, and also to a general quasilinear problem, which includes in particular (1.1), see, for instance, [15, Theorem 3.1]. In [15] the authors state that weak solutions to some quasilinear problem are in $L^{\infty}(\Omega) \cap$ $L^{\infty}(\partial\Omega)$. By elliptic regularity, weak solutions to (1.1) are in fact more regular, and in particular, they are uniformly continuous functions. Indeed, the elliptic regularity theory, applied to weak solutions of a subcritical or even critical problem implies that they are in $C(\overline{\Omega})$, see estimate (B.2) in Theorem B.1. So, in that case,

(1.2)
$$\|u\|_{L^{\infty}(\partial\Omega)} \le \|u\|_{C(\overline{\Omega})} = \|u\|_{L^{\infty}(\Omega)}.$$

In addition to improving the regularity, uniform $L^{\infty}(\Omega)$ a priori bounds, joint with Leray-Schauder degree theory, allow us to investigate results on the existence of solutions. Leray-Schauder degree theory (see [13]) is a powerful tool to obtain results on the existence of solutions. To define the degree, uniform $L^{\infty}(\Omega)$ a priori bounds are needed. Results on uniform $L^{\infty}(\Omega)$ a priori bounds by Gidas and Spruck can be found in [9] and by de Figueiredo, Lions, and Nussbaum in [7], see also [3] for a slightly subcritical Dirichlet problem, [4] for a non-convex domain, [17] for an elliptic system,[6] for the *p*-laplacian case, and [20, 21] for a review.

The type of $L^{\infty}(\Omega)$ estimates given by (2.14) are known for slightly subcritical nonlinearities in the homogeneous Dirichlet problem with the laplacian operator, see [22, Theorem 1.5], with the *p*-Laplacian operator, see [23, Theorem 1.6], and also with a linear problem at the interior joint with nonlinear boundary conditions on the boundary of power type, see [5].

In this paper, we analyze the combined effect of both nonlinearities simultaneously. Our main result establishes the explicit estimates provided by Theorem 2.2, where both nonlinearities in the interior and on the boundary are slightly subcritical, not necessarily of power type.

This paper is organized in the following way. Section 2 contains the statement of our main result, Theorem 2.2. The proof of Theorem 2.2 is achieved in Section 3. By the sake of completeness, we include two appendix; in Appendix A, we recall the regularity of weak solution to the linear problem with non homogeneous data both at the interior and on the boundary, see Theorem A.1; Appendix B deals with further regularity of weak solutions to (1.1), see Theorem B.1.

2. Our main result

In this section we state our main result.

For p > 1, we define the trace operator:

$$\Gamma: W^{1,p}(\Omega) \to L^p(\partial\Omega),$$

in the following way

- (1) $\Gamma u = u|_{\partial\Omega}$ if $u \in W^{1,p}(\Omega) \cap C(\overline{\Omega})$,
- (2) $\|\Gamma u\|_{L^p(\partial\Omega)} \leq C \|u\|_{W^{1,p}(\Omega)},$

where $C = C(p, |\Omega|)$ is a constant and $\partial \Omega$ is C^1 . Since the surjectivity and the continuity of the trace operator, we get

$$\Gamma: W^{1,p}(\Omega) \to W^{1-\frac{1}{p},p}(\partial\Omega) \hookrightarrow L^q(\partial\Omega), \quad \text{for} \quad 1 \le q \le \frac{(N-1)p}{N-p},$$

and

$$\|\Gamma u\|_{L^q(\partial\Omega)} \le C \|u\|_{W^{1,p}(\Omega)}, \quad \text{for some} \quad C > 0,$$

this operator is continuous for $1 \leq q \leq \frac{(N-1)p}{N-p}$, and compact for $1 \leq q < \frac{(N-1)p}{N-p}$ (see [11, Theorem 6.4.1] and [2, Lemma 9.9]).

Throughout this paper, we use the Sobolev embedding

(2.3)
$$H^1(\Omega) \hookrightarrow L^{2^*}(\Omega),$$

and the continuity of the trace operator

$$H^1(\Omega) \hookrightarrow L^{2*}(\partial\Omega),$$

where

(2.4)
$$2^* := \frac{2N}{N-2}$$
 and $2_* := \frac{2(N-1)}{N-2}$,

are the critical Sobolev exponent and the critical exponent in the sense of the trace, respectively.

For $1 < p, p_B \leq \infty$, we will denote

(2.5)
$$2_{N/p}^* := \frac{2^*}{p'} = 2^* \left(1 - \frac{1}{p}\right)$$
 and $2_{*,N/p_B} := \frac{2_*}{p'_B} = 2_* \left(1 - \frac{1}{p_B}\right)$,

where p' is the conjugate exponent of p, that is $\frac{1}{p} + \frac{1}{p'} = 1$. And also

$$p^* := \frac{Np}{N-p}, \ p_* := \frac{(N-1)p}{N-p} = \frac{(N-1)p^*}{N}, \quad \text{for } 1 \le p < N,$$

will denote the critical Sobolev exponent and the critical exponent in the sense of the trace, respectively.

Given $f: \Omega \times \mathbb{R} \to \mathbb{R}$, we will assume the following hypothesis on the nonlinearity at the interior:

(f1) f is a Carathéodory function:
(a) f(·,t) is measurable for each t ∈ ℝ;
(b) f(x,·), is continuous for each x ∈ Ω.
(f2) f is slightly subcritical (at infinity), that is:

(2.6)
$$|f(x,t)| \le |a(x)|f(|t|)$$

with $a(x) \in L^r(\Omega)$ for $r > \frac{N}{2}$, $\tilde{f} : [0, +\infty) \to [0, +\infty)$ is continuous, and such that

(2.7)
$$\lim_{t \to +\infty} \frac{f(t)}{t^{2^*_{N/r} - 1}} = 0.$$

Likewise, given $f_B : \partial \Omega \times \mathbb{R} \to \mathbb{R}$, we will assume the following hypothesis for the nonlinearity on the boundary:

- $(\mathbf{f}_B \mathbf{1}) f_B$ is a Carathéodory function:
 - (a) $f_B(\cdot, t)$ is measurable for each $t \in \mathbb{R}$;
 - (b) $f_B(x, \cdot)$ is continuous for each $x \in \partial \Omega$.

5

 $(\mathbf{f}_B \mathbf{2})$ f_B is slightly subcritical (at infinity), that is:

(2.8)
$$|f_B(x,t)| \le |a_B(x)| \, \tilde{f}_B(|t|),$$

with $a_B(x) \in L^{r_B}(\partial\Omega)$ for $r_B > N-1$, and $\tilde{f}_B : [0, +\infty) \to [0, +\infty)$ is continuous, such that

(2.9)
$$\lim_{t \to +\infty} \frac{f_B(t)}{t^{2_{*,N/r_B}-1}} = 0.$$

We say that $u \in H^1(\Omega)$ is a *weak solution* to (1.1) if $f(\cdot, u) \in L^{(2^*)'}(\Omega)$, and $f_B(\cdot, u) \in L^{(2^*)'}(\partial\Omega)$ are such that for all $\psi \in H^1(\Omega)$,

$$\int_{\Omega} \nabla u \nabla \psi \, dx + \int_{\Omega} u \psi \, dx = \int_{\Omega} f(x, u) \psi \, dx + \int_{\partial \Omega} f_B(x, u) \psi \, dS \,,$$

being $(2^*)' = \frac{2N}{N+2}$ and $(2_*)' = \frac{2(N-1)}{N}$ the conjugate exponents of 2^* and 2_* , respectively.

Remark 2.1. (i) Let $u \in H^1(\Omega)$. By Sobolev embeddings, for f and f_B slightly subcritical (satisfying (f2) and (f_B2) respectively), we have

$$\tilde{f}(u) \in L^{\frac{2^*}{2^{N/r}-1}}(\Omega), \quad \text{where} \quad \frac{2^*_{N/r}-1}{2^*} = \frac{1}{2} + \frac{1}{N} - \frac{1}{r},$$
$$\tilde{f}_B(u) \in L^{\frac{2^*}{2^*,N/r_B}-1}(\partial\Omega), \quad \text{where} \quad \frac{2_{*,N/r_B}-1}{2_*} = \frac{N}{2(N-1)} - \frac{1}{r_B}.$$

Hence,

$$f(\cdot, u) \in L^{(2^*)'}(\Omega)$$
 and $f_B(\cdot, u) \in L^{(2_*)'}(\partial\Omega)$,

(ii) We can allways choose \tilde{f} and \tilde{f}_B such that $\tilde{f}(t) > 0$ and $\tilde{f}_B(t) > 0$ for s > 0. Note that redefining both functions, $\tilde{f}(t)$ and $\tilde{f}_B(t)$, as $\max_{[0,t]} \tilde{f}$ and $\max_{[0,t]} \tilde{f}_B$, respectively, we can always choose \tilde{f} and \tilde{f}_B as non decreasing functions for s > 0.

Now, we will define two new functions, h and h_B , which will be essential for the statement of our main result. Let us define

(2.10)
$$h(s) := \frac{s^{2^*_{N/r}-1}}{\tilde{f}(s)}$$
 and $h_B(s) := \frac{s^{2^*_{N/r_B}-1}}{\tilde{f}_B(s)}$ for $s > 0$,

Since the nonlinearities f and f_B are both slightly subcritical, in other words, satisfy conditions (2.6)-(2.7) and (2.8)-(2.9), respectively, then

(2.11)
$$h(s) \to \infty$$
 and $h_B(s) \to \infty$ as $s \to \infty$.

Our main result is given in the following theorem, which applies for weak solutions to (1.1) with slightly subcritical Carathéodory nonlinearities. Our estimates apply to positive or even changing sign solutions.

We will denote as a_M the maximum of the corresponding norms of $a \in L^r(\Omega)$ and of $a_B \in L^{r_B}(\partial\Omega)$, that is

(2.12)
$$a_M := \max\{\|a\|_{L^r(\Omega)}, \|a_B\|_{L^{r_B}(\partial\Omega)}\}$$

Let u be a solution to (1.1). Let h_m be defined as the minimum of h and a certain power of h_B , specifically

(2.13)
$$h_m(s) := \min\left\{h(s), h_B^{\frac{2^*_{N/r} - 1}{2^*_{*,N/r_B} - 1}}(s)\right\},$$

with h and h_B defined in (2.10).

The following Theorem contains our estimates of $h_m(||u||_{L^{\infty}(\Omega)})$ in terms of their $H^1(\Omega)$ norms.

Theorem 2.2. Let $f : \Omega \times \mathbb{R} \to \mathbb{R}$ and $f_B = \partial \Omega \times \mathbb{R} \to \mathbb{R}$ be Carathéodory functions, satisfying (f1)–(f2) and (f_B1)–(f_B2), respectively. Let $u \in H^1(\Omega)$ be an arbitrary weak solution to (1.1).

Then, for all $\varepsilon > 0$, there exist $C_{\varepsilon} > 0$ depending of ε , N, $|\Omega|$ and $|\partial \Omega|$, but independent of u, such that

(2.14)
$$h_m(\|u\|_{L^{\infty}(\Omega)}) \le C_{\varepsilon} a_M^{A+\varepsilon} \left(1 + \|u\|_{H^1(\Omega)}^{(2^*_{N/r}-2)(A+\varepsilon)}\right),$$

where h_m is defined by (2.13), a_M by (2.12), and

$$(2.15) \quad A := \begin{cases} \frac{1}{2} - \frac{N-r}{Nr} \\ \frac{1}{2} - \frac{N-1}{Nr_B} \\ 1 \end{cases} \quad if \begin{cases} either \ r \ge N, \\ or \ N/2 < r < N \ and \ r^* \ge \frac{Nr_B}{N-1}, \\ 1 \end{cases}$$
$$(2.15) \quad if \ N/2 < r < N \ and \ r^* \le \frac{Nr_B}{N-1}. \end{cases}$$

Remark 2.3. Since (1.2), in fact

$$h_m(\|u\|_{C(\overline{\Omega})}) \le C_{\varepsilon} a_M^{A+\varepsilon} \left(1 + \|u\|_{H^1(\Omega)}^{(2^*_{N/r}-2)(A+\varepsilon)}\right)$$

Remark 2.4. From the definitions of h and h_B given in (2.10), we note that

$$h(s) = \frac{s^{2^*_{N/r} - 1}}{\tilde{f}(s)} \qquad and \quad h_B^{\frac{2^*_{N/r} - 1}{2_{*,N/r_B} - 1}}(s) = \left(\frac{s^{2^*_{*,N/r_B} - 1}}{\tilde{f}_B(s)}\right)^{\frac{2^*_{N/r} - 1}{2_{*,N/r} - 1}}.$$

Thus,

$$h_m(s) = \min\left\{\frac{s^{2^*_{N/r}-1}}{\tilde{f}(s)}, \frac{s^{2^*_{N/r}-1}}{\tilde{f}_B^{\frac{2^*_{N/r}-1}{2^*_{N/r}-1}}(s)}\right\}.$$

3. $L^{\infty}(\Omega)$ a priori explicit estimates

In this section, assuming that f and f_B are Carathéodory functions satisfying slightly subcritical growth conditions, we prove our main result.

Our method combines elliptic regularity with the Gagliardo-Nirenberg interpolation inequality. Let u be an arbitrary solution to (1.1). First, we find estimates of the nonlinearities in terms of products of the $H^1(\Omega)$ -norm of u and their $L^{\infty}(\Omega)$ -norm. With it, using elliptic regularity (see Theorem (A.1)), we obtain estimates of the $W^{1,m}(\Omega)$ -norm, with m > N, of the solutions to (1.1). Finally, applying the Gagliardo-Nirenberg interpolation inequality, (see [19]), we obtain an explicit estimate of the $L^{\infty}(\Omega)$ -norm of u in terms of the $H^1(\Omega)$ norm of u.

Proof of Theorem 2.2. Let $u \in H^1(\Omega)$ be a weak solution to (1.1). By Theorem B.1, $u \in H^1(\Omega) \cap L^{\infty}(\Omega)$.

Firstly, we will estimate both nonlinearities (the interior and the boundary nonlinearities) in terms of the $H^1(\Omega)$ -norm and the $L^{\infty}(\Omega)$ -norm of u.

Step 1. $W^{1,m}(\Omega)$ estimates for m > N.

By hypothesis, \tilde{f} and \tilde{f}_B are both increasing. By (1.2)we denote

(3.16)
$$M := \tilde{f}(\|u\|_{L^{\infty}(\Omega)}) = \max_{\begin{bmatrix} 0, \|u\|_{L^{\infty}(\Omega)} \end{bmatrix}} \tilde{f},$$
$$M_B := \tilde{f}_B(\|u\|_{L^{\infty}(\Omega)}) = \max_{\begin{bmatrix} 0, \|u\|_{L^{\infty}(\Omega)} \end{bmatrix}} \tilde{f}_B.$$

8 E. ANTONIO, M. ARCIGA-ALEJANDRE, R. PARDO, AND J. SANCHEZ-ORTIZ

Along this proof, we will use the obvious fact that for any $\gamma > 0$, there exist two constants C_1 and C_2 , only dependent on γ , such that

(3.17)
$$C_1(1+x^{\gamma}) \le (1+x)^{\gamma} \le C_2(1+x^{\gamma}), \quad \text{for all} \quad x \ge 0.$$

From now on, all throughout this proof, C denotes several constants independent of u.

By the growth condition (2.6) and the definition of M (3.16), we have that

(3.18)
$$\begin{aligned} \int_{\Omega} |f(\cdot, u)|^q dx &\leq \int_{\Omega} |a(x)|^q \tilde{f}(u)^{q-t+t} dx \\ &\leq C M^{q-t} \int_{\Omega} |a(x)|^q \tilde{f}(u)^t dx, \end{aligned}$$

for any t < q, and any

(3.19)
$$q \in \left(\frac{N}{2}, \min\{r, N\}\right)$$

Using the Hölder's inequality, for all $1 < s < \infty$, we can write

$$(3.20) \qquad \int_{\Omega} |a(x)|^q \tilde{f}(u)^t \, dx \le \left(\int_{\Omega} |a(x)|^{qs} dx\right)^{\frac{1}{s}} \left(\int_{\Omega} \tilde{f}(u)^{ts'} dx\right)^{\frac{1}{s'}},$$

where s' is such that $\frac{1}{s} + \frac{1}{s'} = 1$. Choosing s and t < q, so that qs = r and $ts' = \frac{2^*}{2^*_{N/r} - 1}$, thus,

(3.21)
$$t := \frac{2^*}{2^*_{N/r} - 1} \left(1 - \frac{q}{r} \right) < q$$
$$\iff \frac{1}{q} - \frac{1}{r} < \frac{2^*_{N/r} - 1}{2^*} = 1 - \frac{1}{r} - \frac{1}{2} + \frac{1}{N}$$
$$\iff q > \frac{2N}{N+2}, \quad \checkmark$$

since $q > \frac{N}{2} > \frac{2N}{N+2}$.

(3.22)

On the other hand, by subcriticality, see (2.7), and Sobolev embeddings, see (2.3),

$$\int_{\Omega} |\tilde{f}(u)|^{\frac{2^*}{2^*_{N/r}-1}} dx \le C \int_{\Omega} \left(1 + |u|^{2^*}\right) dx$$
$$\le C \left(1 + \|u\|_{L^{2^*}(\Omega)}^{2^*}\right)$$

(3.23)
$$= C \left(1 + \|u\|_{H^{1}(\Omega)}^{2^{*}} \right).$$

Finally, substituting (3.23) in the second factor on the RHS of (3.20), this result in (3.18) and since 1/(qs') = 1/q - 1/r, we get

(3.24)
$$\left(\int_{\Omega} |f(\cdot, u)|^q dx \right)^{\frac{1}{q}} \le C M^{1 - \frac{t}{q}} \|a\|_{L^r(\Omega)} \left(1 + \|u\|_{H^1(\Omega)}^{2^* \left(\frac{1}{q} - \frac{1}{r}\right)} \right).$$

Likewise, by the condition (2.8) and the subcriticality (2.9), we get

(3.25)
$$\int_{\partial\Omega} |f_B(\cdot, u)|^{q_B} dS \leq \int_{\partial\Omega} |a_B(x)|^{q_B} \tilde{f}_B(u)^{q_B-t_B+t_B} dS$$
$$\leq C M_B^{q_B-t_B} \int_{\partial\Omega} |a_B(x)|^{q_B} \tilde{f}_B(u)^{t_B} dS,$$

for any $t_B < q_B$, and any

$$(3.26) q_B \in (N-1, r_B).$$

Using Hölder's inequality, for all $1 < s_B < \infty$, we obtain

(3.27)
$$\int_{\partial\Omega} |a_B(x)|^{q_B} \tilde{f}_B(u)^{t_B} dS$$
$$\leq \left(\int_{\partial\Omega} |a_B(x)|^{q_Bs_B} dS \right)^{\frac{1}{s_B}} \left(\int_{\partial\Omega} \tilde{f}_B(u)^{t_Bs'_B} dS \right)^{\frac{1}{s'_B}}$$

where s'_B is such that $\frac{1}{s_B} + \frac{1}{s'_B} = 1$. Choosing, as before, s_B , $t_B < q_B$, so that $q_B s_B = r_B$, and $t_B s'_B = \frac{2_*}{2_{*,N/r_B} - 1}$, thus,

(3.28)
$$t_{B} := \frac{2_{*}}{2_{*,N/r_{B}} - 1} \left(1 - \frac{q_{B}}{r_{B}} \right) < q_{B}$$
$$\iff \frac{1}{q_{B}} - \frac{1}{r_{B}} < \frac{2_{*,N/r_{B}} - 1}{2_{*}} = 1 - \frac{1}{r_{B}} - \frac{N - 2}{2(N - 1)}$$
$$\iff \frac{1}{q_{B}} < \frac{N}{2(N - 1)} \iff q_{B} > \frac{2(N - 1)}{N}, \quad \checkmark$$

and the last inequality is satisfied since $q_B > N - 1$ and N > 2. On the other hand, again by subcriticality, see (2.8) and (2.9),

$$\int_{\partial\Omega} |f_B(u)|^{\frac{2_*}{2_*,N/r_B}-1} dx \le C \int_{\partial\Omega} \left(1+|u|^{2_*}\right) dS$$

$$\le C \left(1+\|u\|^{2_*}\right) \le C \left(1+\|u\|^{2_*}\right)$$

(3.29)
$$\leq C \left(1 + \|u\|_{L^{2_*}(\partial\Omega)}^2 \right)$$

(3.30)
$$\leq C \left(1 + \|u\|_{H^1(\Omega)}^{2_*} \right),$$

10E. ANTONIO, M. ARCIGA-ALEJANDRE, R. PARDO, AND J. SANCHEZ-ORTIZ

then, substituting (3.30) in the second factor on the RHS of (3.27), this result in (3.25), and since $1/(q_B s'_B) = 1/q_B - 1/r_B$, we get

(3.31)
$$\left(\int_{\partial\Omega} |\tilde{f}_B(\cdot, u)|^{q_B} dx \right)^{\frac{1}{q_B}} \leq C M_B^{1 - \frac{t_B}{q_B}} \|a_B\|_{L^{r_B}(\partial\Omega)} \\ \times \left(1 + \|u\|_{H^1(\Omega)}^{2*\left(\frac{1}{q_B} - \frac{1}{r_B}\right)} \right)$$

Now, using elliptic regularity, we estimate the norm $||u||_{W^{1,m}(\Omega)}$ in terms of the corresponding norms of the nonlinearities, see TheoremA.1, equation (A.2). Specifically, using (3.24) and (3.31), we obtain that

$$\|u\|_{W^{1,m}(\Omega)} \leq C \left[M^{1-\frac{t}{q}} \|a\|_{L^{r}(\Omega)} \left(1 + \|u\|_{H^{1}(\Omega)}^{2^{*}\left(\frac{1}{q} - \frac{1}{r}\right)} \right) + M_{B}^{1-\frac{t_{B}}{q_{B}}} \|a_{B}\|_{L^{r_{B}}(\partial\Omega)} \left(1 + \|u\|_{H^{1}(\Omega)}^{2^{*}\left(\frac{1}{q_{B}} - \frac{1}{r_{B}}\right)} \right) \right],$$

where $m = \min\{q^*, \frac{Nq_B}{N-1}\}$ $(q^* := \frac{Nq}{N-q})$, whenever $1 \le q < N$, see Theorem A.1.

Fixing

(3.33)
$$q_B := \frac{(N-1)q^*}{N} \implies m = q^* = \frac{Nq_B}{N-1} > N_F$$

(in the forthcoming Remark 3.1, we explain the necessity of the election for q_B), moreover, we have the following equivalences

(3.34)
$$q_B := \frac{(N-1)q^*}{N} \iff \frac{2_*}{q_B} = \frac{2^*}{q^*} \iff 2_{*,N/q_B} = 2_{N/q^*}^*$$

Indeed, we only have to notice that, using the definitions (2.4), (2.5) and (3.33), we can conclude that

$$2_{*,N/q_B} = 2_* - \frac{2_*}{q_B} = 2_* + \frac{2^*}{N} - \frac{2^*}{q} = 2^* - \frac{2^*}{q} = 2_{N/q}^*.$$

With that election of q_B , we also need to restrict q in order to satisfy (3.26). Specifically

(3.35)
$$q \in \left(\frac{N}{2}, \min\left\{r, \frac{Nr_B}{N-1+r_B}\right\}\right).$$

Indeed, note that, because of the definition of q_B , see (3.33), and their restriction, (3.26), the following inequality have to be satisfied

$$N - 1 < \frac{(N - 1)q^*}{N} = q_B < r_B.$$

By (3.19), we obtain that $q^* > N$ so $\frac{(N-1)q^*}{N} > N-1$. Thus, we only need to check

$$\begin{aligned} q^* < & \frac{Nr_B}{N-1} \iff \frac{1}{q} - \frac{1}{N} > \frac{N-1}{Nr_B} \iff \frac{1}{q} > \frac{N-1}{Nr_B} + \frac{1}{N} \\ \iff q < \frac{Nr_B}{N-1+r_B}, \end{aligned}$$

from which, using (3.19), and that

$$(3.36) \qquad \qquad \frac{Nr_B}{N-1+r_B} < N,$$

we conclude (3.35).

Step 2. Gagliardo-Nirenberg interpolation inequality.

The Gagliardo-Nirengberg's interpolation inequalities (see [19]), implies that there exist a constant $C = C(N, q, |\Omega|)$, such that

(3.37)
$$\|u\|_{L^{\infty}(\Omega)} \leq C \|u\|_{W^{1,m}(\Omega)}^{\sigma} \|u\|_{L^{2^{*}}(\Omega)}^{1-\sigma},$$

where

(3.38)
$$\frac{1}{\sigma} = 1 + 2^* \left(\frac{2}{N} - \frac{1}{q}\right)$$

From (3.34), due to the definition of $2^*_{N/q}$, see (2.5), it is easy to check that

(3.39)
$$\frac{1}{\sigma} = 1 + 2^* \left(\frac{2}{N} \mp 1 - \frac{1}{q}\right) = 2^*_{N/q} - 1.$$

Substituting the estimate of $||u||_{W^{1,m}(\Omega)}$, see 3.32, and using (3.17) in the inequality (3.37), we get

$$\begin{aligned} \|u\|_{L^{\infty}(\Omega)} &\leq C \left[M^{1-\frac{t}{q}} \|a\|_{L^{r}(\Omega)} \left(1 + \|u\|_{H^{1}(\Omega)}^{2^{*}\left(\frac{1}{q} - \frac{1}{r}\right)} \right) \right. \\ &+ M_{B}^{1-\frac{t_{B}}{q_{B}}} \|a_{B}\|_{L^{r_{B}}(\partial\Omega)} \left(1 + \|u\|_{H^{1}(\Omega)}^{2^{*}\left(\frac{1}{q_{B}} - \frac{1}{r_{B}}\right)} \right) \right]^{\sigma} \|u\|_{L^{2^{*}}(\Omega)}^{(1-\sigma)} \\ &\leq C \left[M^{\left(1-\frac{t}{q}\right)\sigma} \|a\|_{L^{r}(\Omega)}^{\sigma} \left(1 + \|u\|_{H^{1}(\Omega)}^{2^{*}\left(\frac{1}{q_{B}} - \frac{1}{r_{B}}\right)\sigma} \right) \right. \\ &\left. (3.40) + M_{B}^{\left(1-\frac{t_{B}}{q_{B}}\right)\sigma} \|a_{B}\|_{L^{r_{B}}(\partial\Omega)}^{\sigma} \left(1 + \|u\|_{H^{1}(\Omega)}^{2^{*}\left(\frac{1}{q_{B}} - \frac{1}{r_{B}}\right)\sigma} \right) \right] \|u\|_{L^{2^{*}}(\Omega)}^{(1-\sigma)}, \end{aligned}$$

We now look closely at the exponents of $||u||_{L^{\infty}(\Omega)}$ in the RHS, in order to achieve our estimates.

Taking into account the definitions of M and M_B , see (3.16), that f and f_B are non decreasing, and the definition of the functions h and h_B , see (2.10), we can write the following relation between them,

(3.41)
$$M = \frac{\|u\|_{L^{\infty}(\Omega)}^{2^*_{N/r}-1}}{h(\|u\|_{L^{\infty}(\Omega)})} \quad \text{and} \quad M_B = \frac{\|u\|_{L^{\infty}(\Omega)}^{2_{*,N/r_B}-1}}{h_B(\|u\|_{L^{\infty}(\Omega)})}.$$

Moreover, using the definitions of t, see (3.21), and of $2^*_{N/p}$, see (2.5), we can get

(3.42)
$$1 - \frac{t}{q} = 1 - \frac{2^*}{2^*_{N/r} - 1} \left(\frac{1}{q} - \frac{1}{r}\right) = \frac{2^*_{N/q} - 1}{2^*_{N/r} - 1}.$$

Thus, because of the expression (3.42), we deduce

$$(2_{N/r}^* - 1)\left(1 - \frac{t}{q}\right) = (2_{*,N/q} - 1),$$

and because of the definition of σ , see (3.39)

(3.43)
$$\left(2_{N/r}^* - 1\right) \left(1 - \frac{t}{q}\right) \sigma = 1.$$

Similarly, from the definitions of t_B , see (3.28), and of $2_{*,N/q_B}$, see (2.5), we obtain

(3.44)
$$1 - \frac{t_B}{q_B} = 1 - \frac{2_*}{2_{*,N/r_B} - 1} \left(\frac{1}{q_B} - \frac{1}{r_B}\right) = \frac{2_{*,N/q_B} - 1}{2_{*,N/r_B} - 1}.$$

Likewise, since (3.44), the definition of σ , see (3.39), and the equivalences (3.34), we get

(3.45)
$$\left(2_{*,N/r_B} - 1\right) \left(1 - \frac{t_B}{q_B}\right) \sigma = \frac{2_{*,N/q_B} - 1}{2_{*,N/q} - 1} = 1.$$

Now, we divide both sides of the inequality (3.40) by $||u||_{L^{\infty}(\Omega)}$. Using the definitions of M and M_B , also the two expressions concerning σ ; (3.43), (3.45), and the definition of a_M , see (2.12), we obtain that

$$(3.46) \qquad 1 \le Ca_M^{\sigma} \left(\frac{\left(1 + \|u\|_{H^1(\Omega)}^{2^*\left(\frac{1}{q} - \frac{1}{r}\right)\sigma}\right)}{h^{\frac{1}{2^*/r^{-1}}}(\|u\|_{L^{\infty}(\Omega)})} + \frac{\left(1 + \|u\|_{H^1(\Omega)}^{2^*\left(\frac{1}{q_B} - \frac{1}{r_B}\right)\sigma}\right)}{h_B^{\frac{1}{2^*,N/r_B}^{-1}}(\|u\|_{L^{\infty}(\Omega)})} \right) \|u\|_{L^{2^*}(\Omega)}^{(1-\sigma)}.$$

Now, the definition of h_m (see (2.13)), implies that

$$\frac{1}{h_m^{\frac{1}{2^*_{N/r}-1}}(\|u\|_{L^{\infty}(\Omega)})} = \max\left\{\frac{1}{h^{\frac{1}{2^*_{N/r}-1}}(\|u\|_{L^{\infty}(\Omega)})}, \frac{1}{h_B^{\frac{1}{2_{*,N/r_B}-1}}(\|u\|_{L^{\infty}(\Omega)})}\right\}.$$

So, substituting this maximum in the inequality (3.46), we get

(3.47)
$$h_{m}^{\frac{1}{2_{N/r}^{*}-1}}(\|u\|_{L^{\infty}(\Omega)}) \leq Ca_{M}^{\sigma}\left(1+\|u\|_{H^{1}(\Omega)}^{2^{*}\left(\frac{1}{q}-\frac{1}{r}\right)\sigma} +\|u\|_{H^{1}(\Omega)}^{2_{*}\left(\frac{1}{q_{B}}-\frac{1}{r_{B}}\right)\sigma}\right)\|u\|_{L^{2^{*}}(\Omega)}^{(1-\sigma)}.$$

The RHS in the above inequality is upper bounded by a term with the largest exponent of both addends. Let us denote this maximum by

(3.48)
$$E_M := \max\left\{2^*\left(\frac{1}{q} - \frac{1}{r}\right), 2_*\left(\frac{1}{q_B} - \frac{1}{r_B}\right)\right\}.$$

Since the inequality (3.47) the definition (3.48) and Sobolev's embedding, we obtain

(3.49)
$$h_m(||u||_{L^{\infty}(\Omega)}) \le Ca_M^{\theta} \left(1 + ||u||_{H^1(\Omega)}^{\beta}\right),$$

where

(3.50)
$$\theta := \left(2_{N/r}^* - 1\right)\sigma = \frac{2_{N/r}^* - 1}{2_{N/q}^* - 1},$$

and

(3.51)
$$\beta := \left(E_M + \frac{1-\sigma}{\sigma}\right)\theta.$$

At this moment, we look closely at the definition of E_M .

Firstly by definition of 2^* and of 2_* , see (2.5), secondly by election of q_B , see (3.33), and finally rearranging terms, we observe that

$$(3.52) 2^* \left(\frac{1}{q} - \frac{1}{r}\right) \ge 2_* \left(\frac{1}{q_B} - \frac{1}{r_B}\right) \iff \frac{1}{q} \mp \frac{1}{N} - \frac{1}{r} \ge \frac{N-1}{N} \left(\frac{1}{q_B} - \frac{1}{r_B}\right) \iff \frac{1}{r} - \frac{1}{N} \le \frac{N-1}{Nr_B}.$$

If $r \ge N$, then $1/r - 1/N \le 0$, and the last inequality holds. Moreover, if N/2 < r < N, then the last inequality holds if and only if

$$(1/r - 1/N)^{-1} =: r^* \ge Nr_B/(N-1).$$

Observe that

$$r^* \ge Nr_B/(N-1) \iff 2^*_{N/r} \ge 2_{*,N/r_B}$$

On the contrary, the reverse inequality to (3.52), will be satisfied whenever N/2 < r < N, and $r^* \leq Nr_B/(N-1)$. Hence

(3.53)
$$E_M = \begin{cases} 2^* \left(\frac{1}{q} - \frac{1}{r}\right) & \{ \text{if } r \ge N, \\ \text{or } N/2 < r < N \text{ and } r^* \ge \frac{Nr_B}{N-1}, \\ 2_* \left(\frac{1}{q_B} - \frac{1}{r_B}\right) & \text{if } N/2 < r < N \text{ and } r^* \le \frac{Nr_B}{N-1}. \end{cases}$$

Consequently, we have two cases in the search for the optimum exponents θ and β varying q, see (3.35):

Case (I): Either $r \ge N$, or N/2 < r < N and $r^* \ge \frac{Nr_B}{N-1}$.

Using the definition of β , (3.51), the first equality for E_M in (3.53), and the expression for σ , see(3.38), and for $2^*_{N/r}$, see (2.5),

(3.54)
$$\beta = \left[2^* \left(\frac{1}{q} - \frac{1}{r}\right) + \frac{1 - \sigma}{\sigma}\right] \theta$$

(3.55)
$$= \left[2^* \left(\frac{1}{q} - \frac{1}{r}\right) + 2^* \left(\frac{2}{N} - \frac{1}{q}\right)\right] \theta = (2^*_{N/r} - 2)\theta.$$

The function $\theta: q \mapsto \theta(q)$, defined by (3.50) is decreasing. We look for their infimum for q in the interval (3.19).

Assume $r \ge N$. Since (3.35)–(3.36), we deduce that $q \in \left(\frac{N}{2}, \frac{Nr_B}{N-1+r_B}\right)$.

Assume N/2 < r < N and $r^* \ge \frac{Nr_B}{N-1}$. Note that

$$(3.56) \quad r^* \ge \frac{Nr_B}{N-1} \iff \frac{1}{r} - \frac{1}{N} \le \frac{N-1}{Nr_B} \iff r \ge \frac{Nr_B}{N-1+r_B},$$

and also $q \in \left(\frac{N}{2}, \frac{Nr_B}{N-1+r_B}\right)$. Hence, in case I,

(3.57)
$$\inf_{q \in \left(\frac{N}{2}, \frac{Nr_B}{N-1+r_B}\right)} \theta(q) = \theta\left(\frac{Nr_B}{N-1+r_B}\right) = \frac{\frac{1}{2} - \frac{N-r}{Nr}}{\frac{1}{2} - \frac{N-1}{Nr_B}}$$

Case (II): N/2 < r < N and $r^* \leq \frac{Nr_B}{N-1}$.

Likewise, using the definition of β , (3.51), the second equality for E_M in (3.53), and the expressions for σ (3.38), for $2^*_{N/p}$, $2_{*,N/p_B}$ (2.5), and the equivalence (3.34)

$$\beta = \left[2_* \left(\frac{1}{q_B} - \frac{1}{r_B}\right) + \frac{1 - \sigma}{\sigma}\right] \theta$$

= $\left[2_* \left(\frac{1}{q_B} \mp 1 - \frac{1}{r_B}\right) + 2^* \left(\frac{2}{N} \mp 1 - \frac{1}{q}\right)\right] \theta$
(3.58) = $\left[2_{*,N/r_B} - 2_{*,N/q_B} - 2 + 2_{N/q}^*\right] \theta = \left[2_{*,N/r_B} - 2\right] \theta.$

Now, for q satisfying (3.35), thanks to (3.56), we deduce that $q \in (\frac{N}{2}, r)$, hence

(3.59)
$$\inf_{q \in \left(\frac{N}{2}, r\right)} \theta(q) = \theta(r) = 1.$$

Finally, we introduce into the inequality (3.49), the infima of θ and β given by (3.57) and (3.55) respectively in case I, and by (3.59) and (3.58), in case II. Since these infima are not attained in the set where q belongs, for any $\varepsilon > 0$, there exists a constant $C_{\varepsilon} > 0$ such that,

$$h_m(\|u\|_{L^{\infty}(\Omega)}) \le C_{\varepsilon} a_M^{A+\varepsilon} \left(1 + \|u\|_{H^1(\Omega)}^{(2^*_{N/r}-2)(A+\varepsilon)}\right)$$

where A is defined in (2.15), and $C_{\varepsilon} = C_{\varepsilon}(\varepsilon, N, |\Omega|, |\partial \Omega|)$ and it is independent of u.

In the following Remark, we observe the necessity of the election for q_B , see (3.33).

Remark 3.1. Assume for a while that (3.33) do not hold and, to fix ideas, that

$$q_B < \frac{(N-1)q^*}{N} \implies m = \frac{Nq_B}{N-1} > N.$$

We also have the following equivalence

$$q_B < \frac{(N-1)q^*}{N} \iff \frac{2^*}{q^*} < \frac{2_*}{q_B} \iff 2_{*,N/q_B} < 2^*_{N/q_B}$$

Indeed, the first equivalence is obvious. With respect to the second one, notice that, due to the definitions of $2^*_{N/q}$ and of $2_{*,N/q_B}$, see (2.5), we can conclude that

$$2_{*,N/q_B} = 2_* - \frac{2_*}{q_B} < 2_* - \frac{2^*}{q^*} = 2^* - \frac{2^*}{q} = 2_{N/q}^*$$

Now, in the Gagliardo-Nirengberg's interpolation inequality, see (3.37), the parameter σ is given by

$$\frac{1}{\sigma} = 1 + 2^* \left(\frac{1}{N} - \frac{1}{m}\right) = 1 + 2^* \left(\frac{1}{N} \mp 1 - \frac{N-1}{Nq_B}\right)$$
$$= 2_{*,N/q_B} - 1 < 2^*_{N/q} - 1.$$

And the expression (3.43) becomes now

$$\left(2_{N/r}^*-1\right)\left(1-\frac{t}{q}\right)\sigma=\frac{2_{*,N/q}-1}{2_{*,N/q_B}-1}>1.$$

The above inequality means that the exponent of $||u||_{L^{\infty}(\Omega)}$ in the RHS will dominate 1, which is the exponent of $||u||_{L^{\infty}(\Omega)}$ in the LHS, and the bounds can not be reached.

Likewise, if $q_B > \frac{(N-1)q^*}{N}$, then $m = q^*$ and it can be proved that

$$\frac{1}{\sigma} = 1 + 2^* \left(\frac{2}{N} \mp 1 - \frac{1}{q}\right) = 2^*_{N/q} - 1 < 2_{*,N/q_B} - 1,$$

and so

$$\left(2_{*,N/r_B} - 1\right) \left(1 - \frac{t_B}{q_B}\right) \sigma = (2_{*,N/q_B} - 1)\sigma = \frac{2_{*,N/q_B} - 1}{2_{N/q}^* - 1} > 1,$$

concluding that necessarily, q_B have to be chosen as in (3.33).

Throughout that proof, we have explicit estimates of $h_m(||u||_{L^{\infty}(\Omega)})$ expressed in their $L^{2^*}(\Omega)$ norm and $L^{2_*}(\partial\Omega)$ norm (see (3.22) and (3.29)). Previously, we unify those estimates in their $H^1(\Omega)$ norm to simplify the expression.

In the next Corollary, we split those estimates in terms of the $L^{2^*}(\Omega)$ norm and the $L^{2_*}(\partial\Omega)$ norm.

Corollary 3.2. Assume that all the hypotheses of Theorem 2.2 hold. Then, for all $\varepsilon > 0$, there exist $C_{\varepsilon} > 0$ depending of ε , N, $|\Omega|$ and $|\partial \Omega|$, but independent of u, such that

$$h_m(\|u\|_{L^{\infty}(\Omega)}) \le Ca_M^{A+\varepsilon} \left(1 + \|u\|_{L^{2^*}(\Omega)}^{A_1+\varepsilon} + \|u\|_{L^{2^*}(\partial\Omega)}^{A_2+\varepsilon} \|u\|_{L^{2^*}(\Omega)}^{A_3+\varepsilon} \right),$$

where A is defined in (2.15),

$$\begin{aligned} A_1 &:= (2^*_{N/r} - 2)A \\ A_2 &:= 0 \\ A_3 &:= (2_{*,N/r_B} - 2)A \end{aligned} if \begin{cases} either \ r \ge N, \\ or \ N/2 < r < N \ and \ r^* \ge \frac{Nr_B}{N-1}, \end{aligned}$$

$$\begin{array}{rcl} A_1 := & 2^*_{N/r} - 2 \\ A_2 := & 2_{*,N/r_B} - 2^*_{N/r} & \mbox{if} & N/2 < r < N \mbox{ and } r^* \leq \frac{Nr_B}{N-1} \\ A_3 := & 2^*_{N/r} - 2 \end{array}$$

Proof. The proof is similar to the proof of the Theorem (2.2).

Step 1. $W^{1,m}(\Omega)$ estimates for m > N.

Substituting (3.22) in the second factor on the RHS of (3.20), and this is (3.18), we get

(3.60)
$$\left(\int_{\Omega} |f(\cdot, u)|^q dx\right)^{\frac{1}{q}} \le CM^{1-\frac{t}{q}} \|a\|_{L^r(\Omega)} \left(1 + \|u\|_{L^{2^*}(\Omega)}^{2^*\left(\frac{1}{q}-\frac{1}{r}\right)}\right),$$

with M defined in (3.16), t in (3.21) and q in (3.19), respectively. See the analogy with (3.24).

On the other hand, replacing (3.29) in the second factor on the RHS of (3.27), and this in (3.25), we get

(3.61)
$$\left(\int_{\partial\Omega} |\tilde{f}_B(\cdot, u)|^{q_B} dx \right)^{\frac{1}{q_B}} \leq C M_B^{1 - \frac{t_B}{q_B}} \|a_B\|_{L^{r_B}(\partial\Omega)} \\ \times \left(1 + \|u\|_{L^{2*}(\partial\Omega)}^{2*\left(\frac{1}{q_B} - \frac{1}{r_B}\right)} \right),$$

with M_B defined in (3.16), t_B in (3.28), and q_B in (3.26) respectively. See the analogy with(3.31).

By elliptic regularity, we estimate the norm $||u||_{W^{1,m}(\Omega)}$ in terms of (3.60) and (3.61), see Theorem A.1, obtaining

$$\|u\|_{W^{1,m}(\Omega)} \leq C \left[M^{1-\frac{t}{q}} \|a\|_{L^{r}(\Omega)} \left(1 + \|u\|_{L^{2^{*}}(\Omega)}^{2^{*}\left(\frac{1}{q}-\frac{1}{r}\right)} \right) + M_{B}^{1-\frac{t_{B}}{q_{B}}} \|a_{B}\|_{L^{r_{B}}(\partial\Omega)} \left(1 + \|u\|_{L^{2^{*}}(\partial\Omega)}^{2^{*}\left(\frac{1}{q_{B}}-\frac{1}{r_{B}}\right)} \right) \right]$$
(3.62)

with m > N. See also the analogy with (3.32).

Step 2. Gagliardo-Nirenberg interpolation inequality.

Substituting 3.62 in the Gagliardo-Nirengberg's inequality (3.37) and using the inequality (3.17), we get

$$\|u\|_{L^{\infty}(\Omega)} \leq C \left[M^{\left(1-\frac{t}{q}\right)\sigma} \|a\|_{L^{r}(\Omega)}^{\sigma} \left(1+\|u\|_{L^{2^{*}}(\Omega)}^{2^{*}\left(\frac{1}{q}-\frac{1}{r}\right)\sigma}\right) + M_{B}^{\left(1-\frac{t_{B}}{q_{B}}\right)\sigma} \|a_{B}\|_{L^{r_{B}}(\partial\Omega)}^{\sigma} \left(1+\|u\|_{L^{2^{*}}(\partial\Omega)}^{2^{*}\left(\frac{1}{q_{B}}-\frac{1}{r_{B}}\right)\sigma}\right) \right] \|u\|_{L^{2^{*}}(\Omega)}^{(1-\sigma)}$$

Using the definitions of M and M_B , see (3.41), using also (3.43), (3.45), the definition of a_M (see (2.12)), and dividing both sides of the inequality (3.63) by $||u||_{L^{\infty}(\Omega)}$, we obtain that

$$1 \le Ca_{M}^{\sigma} \left(\frac{\left(1 + \|u\|_{L^{2^{*}}(\Omega)}^{2^{*}\left(\frac{1}{q} - \frac{1}{r}\right)\sigma}\right)}{h^{\frac{1}{2^{*}/r^{-1}}}(\|u\|_{L^{\infty}(\Omega)})} + \frac{\left(1 + \|u\|_{L^{2^{*}}\left(\frac{1}{q_{B}} - \frac{1}{r_{B}}\right)\sigma}\right)}{h_{B}^{\frac{1}{2^{*},N/r_{B}}^{-1}}(\|u\|_{L^{\infty}(\Omega)})}\right) \|u\|_{L^{2^{*}(\Omega)}}^{(1-\sigma)}$$

Then,

$$h_{m}^{\frac{1}{2_{N/r}^{*}-1}}(\|u\|_{L^{\infty}(\Omega)}) \leq Ca_{M}^{\sigma}\left(2 + \|u\|_{L^{2^{*}}(\Omega)}^{2^{*}\left(\frac{1}{q}-\frac{1}{r}\right)\sigma} + \|u\|_{L^{2^{*}}(\partial\Omega)}^{2^{*}\left(\frac{1}{q_{B}}-\frac{1}{r_{B}}\right)\sigma}\right)\|u\|_{L^{2^{*}}(\Omega)}^{(1-\sigma)}$$

where h_m is defined in (2.13). See the analogy with (3.47). Clearing h_m , we get

$$h_{m}(\|u\|_{L^{\infty}(\Omega)}) \leq Ca_{M}^{\sigma(2^{*}_{N/r}-1)} \left(1 + \|u\|_{L^{2^{*}}(\Omega)}^{2^{*}(\frac{1}{q}-\frac{1}{r})(2^{*}_{N/r}-1)\sigma} + \|u\|_{L^{2^{*}}(\partial\Omega)}^{2^{*}(\frac{1}{q_{B}}-\frac{1}{r_{B}})(2^{*}_{N/r}-1)\sigma}\right) \|u\|_{L^{2^{*}}(\Omega)}^{(1-\sigma)(2^{*}_{N/r}-1)\sigma}$$

Substituting in the exponents the parameter θ (see (3.50)), we get

(3.64)
$$h_m(\|u\|_{L^{\infty}(\Omega)}) \leq Ca_M^{\theta} \left(1 + \|u\|_{L^{2^*}(\Omega)}^{\left[2^*\left(\frac{1}{q} - \frac{1}{r}\right) + \frac{1-\sigma}{\sigma}\right]\theta} + \|u\|_{L^{2^*}(\partial\Omega)}^{2^*\left(\frac{1}{q_B} - \frac{1}{r_B}\right)\theta} \|u\|_{L^{2^*}(\Omega)}^{\frac{1-\sigma}{\sigma}\theta}\right).$$

Let us define the function $\theta_1 = \theta_1(q)$ as the first exponent inside the brackets. Using the definitions of $2^*_{N/q}$, see (2.5), and of σ , see (3.39), we get

$$\theta_1(q) := \left[2^* \left(\frac{1}{q} - \frac{1}{r}\right) + \frac{1 - \sigma}{\sigma}\right] \theta(q)$$
$$= (2^*_{N/r} - 2)\theta(q),$$

note that this value is equal to β in case I of Theorem (2.2), see (3.54)–(3.55).

We define the function $\theta_2 = \theta_2(q)$ as the second exponent. By the definition of $2_{*,N/q_B}$, see (2.5), and the equivalence (3.34)

$$\theta_2(q) := 2_* \left(\frac{1}{q_B} \mp 1 - \frac{1}{r_B} \right) \theta(q) = (2_{*,N/r_B} - 2_{N/q}^*) \theta(q).$$

We define the function $\theta_3 = \theta_3(q)$ as the third exponent. Using the expression (3.39) for σ ,

$$\theta_3(q) := \left(\frac{1-\sigma}{\sigma}\right)\theta(q) = (2^*_{N/q} - 2)\theta(q).$$

As before, let q_B , θ be defined by (3.33), and (3.50) respectively. The function

$$(\theta_1 + \theta_2 + \theta_3)(q) = (2^*_{N/r} + 2_{*,N/rB} - 4)\theta(q),$$

is decreasing, and we look for their infimum for q in the interval (3.35). Thus, as before, we consider the previous two cases.

Case (I) Either $r \ge N$, or N/2 < r < N and $r^* \ge \frac{Nr_B}{N-1}$.

In this case, $q \in \left(\frac{N}{2}, \frac{Nr_B}{N-1+r_B}\right)$. For A is defined in (2.15), the exponents are given by

$$A'_{1} := \theta_{1} \left(\frac{Nr_{B}}{N - 1 + r_{B}} \right) = (2^{*}_{N/r} - 2)A,$$

$$A'_{2} := \theta_{2} \left(\frac{Nr_{B}}{N - 1 + r_{B}} \right) = \frac{2}{N - 2} \left(\frac{N - 1 + r_{B}}{r_{B}} - 1 - \frac{N - 1}{r_{B}} \right) A = 0,$$
and by

and by

bv

$$\begin{aligned} A'_3 &:= \theta_3 \left(\frac{Nr_B}{N - 1 + r_B} \right) = \left(2^* \left(1 - \frac{N - 1 + r_B}{Nr_B} \right) - 2 \right) A \\ &= \left(\frac{2(N - 2)}{N - 2} \left(\frac{r_B - 1}{r_B} \right) - 2 \right) A \\ &= (2_{*,N/r_B} - 2) A \end{aligned}$$

Hence, the inequality (3.64) can be rewritten as

$$h_{m}(\|u\|_{L^{\infty}(\Omega)}) \leq Ca_{M}^{A+\varepsilon} \left(1 + \|u\|_{L^{2^{*}}(\Omega)}^{(2^{*}_{N/r}-2)A+\varepsilon} + \|u\|_{L^{2^{*}}(\partial\Omega)}^{\varepsilon} \|u\|_{L^{2^{*}}(\Omega)}^{(2_{*,N/r_{B}}-2)A+\varepsilon}\right),$$

where A is defined in (2.15).

Case (II) N/2 < r < N and $r^* \leq \frac{Nr_B}{N-1}$.

In that case, $q \in \left(\frac{N}{2}, r\right)$ (see (3.35)). The exponents are given by $A_1'' := \theta_1(r) = 2_{N/r}^* - 2,$ $A_2'' := \theta_2(r) = \frac{2}{N-2} \left(\frac{N}{r} - 1 \mp N - \frac{N-1}{r_B} \right) = 2_{*,N/r_B} - 2_{N/r}^*,$ $A_3'' := \theta_3(r) = 2_{N/r}^* - 2.$

Therefore, the inequality (3.64) is rewritten as

$$h_{m}(\|u\|_{L^{\infty}(\Omega)}) \leq C_{\varepsilon} a_{M}^{1+\varepsilon} \left(1 + \|u\|_{L^{2^{*}}(\Omega)}^{2^{*}_{N/r}-2+\varepsilon} + \|u\|_{L^{2^{*}}(\partial\Omega)}^{2^{*}_{N/r}+\varepsilon} \|u\|_{L^{2^{*}}(\Omega)}^{2^{*}_{N/r}-2+\varepsilon}\right).$$

The next corollary proves that any sequence $\{u_k\} \subset H^1(\Omega)$ of weak solution to (1.1), uniformly bounded in the $L^{2^*}(\Omega)$ norm and in the $L^{2_*}(\partial\Omega)$ norm, is also uniformly bounded in the $C(\overline{\Omega})$ -norm.

Corollary 3.3. Let $f : \Omega \times \mathbb{R} \to \mathbb{R}$ and $f_B : \partial \Omega \times \mathbb{R} \to \mathbb{R}$ be Carathéodory functions, satisfying (f1)-(f2) and $(f_B1)-(f_B2)$, respectively. Let $\{u_k\} \subset H^1(\Omega)$ be a sequence of weak solutions to (1.1) satisfying that, there exist $C_0 > 0$, such that

$$||u_k||_{L^{2^*}(\Omega)} \le C_0$$
 and $||u_k||_{L^{2_*}(\partial\Omega)} \le C_0$.

Then, there exist C > 0 such that,

$$\|u_k\|_{C(\overline{\Omega})} \le C.$$

Proof. We proceed by contradiction, assuming that $||u_k||_{L^{\infty}(\Omega)} \to \infty$. By the Theorem (2.2) and the remark 3.1, we get

$$(3.65) h_m(||u_k||_{C(\overline{\Omega})}) \le C, \quad \text{for} \quad C > 0,$$

where, h_m is defined in (2.13).

Using (2.11), we deduce that $h_m(||u_k||_{C(\overline{\Omega})}) \to \infty$ as $k \to \infty$, which contradicts (3.65).

Corollary 3.4. Let $f : \Omega \times \mathbb{R} \to \mathbb{R}$ and $f_B : \partial\Omega \times \mathbb{R} \to \mathbb{R}$ be Carathéodory functions, satisfying (f1)–(f2) and (f_B1)–(f_B2) respectively. Let $\{u_k\} \subset H^1(\Omega)$ be a sequence of weak solutions to (1.1). Then, the following statements are equivalent

- *i*): $||u_k||_{L^{2^*}(\Omega)} \le C_1$ and $||u_k||_{L^{2^*}(\partial\Omega)} \le C_1$.
- *ii)*: $||u_k||_{C(\overline{\Omega})} \leq C_3$.
- *iii)*: $||u_k||_{H^1(\Omega)} \leq C_2$.

for some constants C_i independent of k, i = 1, 2, 3.

Proof. We prove that $i \Rightarrow ii \Rightarrow iii \Rightarrow iii \Rightarrow i$.

The proof of i) \Rightarrow ii) follows directly from the Corollary 3.3.

Now, using the elliptic regularity result, see the estimate (B.1) in the Theorem B.1, and the Gagliardo-Nirenberg interpolation, the proof of $ii \Rightarrow iii$ is done.

Finally, Sobolev's embedding and the continuity of the trace operator, proves that $iii \Rightarrow i$.

Appendix A. Regularity for the Neumann non homogeneous linear problem

In this appendix, we recall the regularity of weak solution to the linear problem with non homogeneous data both at the interior and on the boundary.

Let us consider the linear nonhomogeneous Neumann problem

(A.1)
$$\begin{cases} -\Delta u + u = g(x), & x \in \Omega, \\ \frac{\partial u}{\partial \eta} = g_B(x), & x \in \partial\Omega, \end{cases}$$

where $\Omega \subset \mathbb{R}^N$, (N > 2), is an open, connected and bounded domain with C^2 boundary.

Theorem A.1. Let us consider the problem (A.1), there exist a positive constant C > 0 independent of u, h and g_B such that the following holds:

(i) If $\partial \Omega \in C^{0,1}$, $g \in L^q(\Omega)$ and $g_B \in L^{q_B}(\partial \Omega)$ with $q \geq 1$ and $q_B \geq 1$, then there exist a unique $u \in W^{1,m}(\Omega)$ and

(A.2)
$$||u||_{W^{1,m}(\Omega)} \leq C \left(||g||_{L^q(\Omega)} + ||g_B||_{L^{q_B}(\partial\Omega)} \right),$$

where $m = \min\{\frac{Nq}{N-q}, \frac{Nq_B}{N-1}\}$ whenever $1 \leq q < N$, or m = $\min\{q, \frac{Nq_B}{N-1}\}$ whenever $q \geq N$. Furthermore, if $q > \frac{N}{2}$ and $q_B > \frac{N}{2}$ N-1, then

$$\|u\|_{C^{\nu}(\overline{\Omega})} \leq C\left(\|g\|_{L^{q}(\Omega)} + \|g_{B}\|_{L^{q_{B}}(\partial\Omega)}\right),$$

- where $\nu = 1 \frac{N}{m}$, (m > N). (ii) If $\partial \Omega \in C^{1,1}$, $g \in C^{\nu}(\Omega) \cap L^q(\Omega)$ and $g_B \in L^{q_B}(\partial \Omega)$ with $q > \frac{N}{2}$ and $q_B > N-1$, then there exist a unique $u \in C^{\nu}(\overline{\Omega}) \cap C^{2,\nu}(\Omega)^2$.
- (iii) If $\partial \Omega \in C^{2,\nu}$, $g \in C^{\nu}(\overline{\Omega})$ and $g_B \in C^{1,\nu}(\partial \Omega)$ with $\nu \in (0,1)$, then there exist a unique $u \in C^{2,\nu}(\overline{\Omega})$ and

$$\|u\|_{C^{2,\nu}(\overline{\Omega})} \leq C\left(\|g\|_{C^{\nu}(\overline{\Omega})} + \|g_B\|_{C^{1,\nu}(\partial\Omega)}\right),$$

where C is a positive constant independent of u, g and g_B .

(iv) If $\partial \Omega \in C^2$, $g \in L^p(\Omega)$ and $g_B \in W^{1-\frac{1}{p},p}(\partial \Omega)$, then $u \in W^{2,p}(\Omega)$ and

$$||u||_{W^{2,p}(\Omega)} \le C\left(||g||_{L^{p}(\Omega)} + ||g_{B}||_{W^{1-\frac{1}{p},p}(\partial\Omega)}\right),$$

where C is a positive constant independent of u, q and $q_{\rm B}$.

- (v) If $\partial \Omega \in C^{1,\nu}$ with $\nu \in (0,1]$, $g \in C^{\nu}(\Omega)$ and $g_B \in C^{\nu}(\partial \Omega) \cap$ $L^{\infty}(\partial\Omega)$ then if u is a bounded weak solution to (A.1), then $u \in C^{1,\beta}(\overline{\Omega}) \cap C^{2,\beta}(\Omega)$, where β depends on ν and N.
- Proof. (i): It follows from [12, Ch.3 Sec. 6] or [16, Lem. 2.2] that there exists a unique $u \in W^{1,p}(\Omega)$ solving (A.1). Now if p > N, using the Sobolev embedding theorem, one has $u \in C^{\alpha}(\overline{\Omega})$. Then by applying [10, Thm. 6.13] for the corresponding nonhomogeneous Dirichlet problem, we have that $u \in C^{1,\alpha}(\overline{\Omega})$, see also [16].
 - (ii): From part (i) we have that $u \in C^{\alpha}(\overline{\Omega})$. Since $\partial \Omega \in C^{1,1}$, Ω satisfies the exterior sphere condition at every point on the boundary and using the fact that $g \in C^{\alpha}(\overline{\Omega})$, reasoning as above it follows from [10, Thm. 6.13] that $u \in C^{\alpha}(\overline{\Omega}) \cap C^{2,\alpha}(\overline{\Omega})$.
 - (iii): See [1, Page 55] or [12, Chap.3 Sec. 3].
 - (iv): See [1, Page 55] or [12, Chap.3 Sec. 9].

(v): By [14, Thm. 2], one has $u \in C^{1,\beta}(\overline{\Omega})$. Then using the bootstrap for the differential equation in Ω , we get the desired regularity in Ω .

APPENDIX B. REGULARITY OF WEAK SOLUTIONS

In this section, we establish auxiliary results on further regularity of weak solutions to (1.1), by assuming that conditions on the growth of the nonlinearities are subcritical or even critical. Using a Moser type procedure, it is known that $u \in L^q(\Omega) \cap L^q(\partial\Omega)$ for all $q < \infty$ (see [15, Theorem 3.1]). Moreover, using elliptic regularity theory, we state the following result that guarantees, in particular, Hölder regularity of any weak solution to (1.1).

Theorem B.1. Let $\Omega \subset \mathbb{R}^N$, $f : \Omega \times \mathbb{R} \to \mathbb{R}$ and $f_B : \partial\Omega \times \mathbb{R} \to \mathbb{R}$ be Carathéodory functions, such that

$$|f(x,s)| \le |a(x)| \left(1 + |s|^{2^*_{N/r}-1}\right) \quad and$$

$$|f_B(x,s)| \le |a_B(x)| \left(1 + |s|^{2_{*,N/r_B}-1}\right),$$

where

$$a(x) \in L^{r}(\Omega),$$
 with $\frac{N}{2} < r \le \infty,$ and $a_{B}(x) \in L^{r_{B}}(\partial\Omega),$ with $N-1 < r_{B} \le \infty.$

Let $u \in H^1(\Omega)$ be a weak solution to (1.1), then $u \in L^q(\Omega) \cap L^q(\partial\Omega)$ for all $1 \leq q < \infty$.

Moreover, $u \in W^{1,m}(\Omega) \cap C^{\nu}(\overline{\Omega})$, and the following estimates holds

(B.1)
$$||u||_{W^{1,m}(\Omega)} \le C \left(||f(\cdot, u)||_{L^r(\Omega)} + ||f_B(\cdot, u)||_{L^{r_B}(\partial\Omega)} \right)$$

and

(B.2)
$$\|u\|_{C^{\nu}(\overline{\Omega})} \leq C \left(\|f(\cdot, u)\|_{L^{r}(\Omega)} + \|f_{B}(\cdot, u)\|_{L^{r_{B}}(\partial\Omega)} \right),$$

where $m = \min\left\{r^*, \frac{Nr_B}{N-1}\right\}$, if $1 \le r < N$, or $m = \min\left\{r, \frac{Nr_B}{N-1}\right\}$, if $r \ge N$ and $\nu = 1 - \frac{N}{m}$.

Besides,

$$||u||_{L^{\infty}(\partial\Omega)} \le ||u||_{C(\overline{\Omega})} = ||u||_{L^{\infty}(\Omega)}.$$

Proof. Let $u \in H^1(\Omega)$ be a weak solution to (1.1). Then $u \in L^q(\Omega) \cap L^q(\partial\Omega)$ for all $q < \infty$ (see [15, Theorem 3.1]).

Next, we use elliptic regularity theory. Since Hölder's inequality,

 $f(\cdot, u) \in L^q(\Omega),$ for every 1 < q < r,

and

 $f_B(x, u) \in L^{q_B}(\partial \Omega),$ for every $1 < q_B < r_B,$

and by elliptic regularity (see Theorem (A.1)), $u \in W^{1,m}(\Omega)$ for $m = \min\left\{q^*, \frac{Nq_B}{N-1}\right\}$ whenever $1 \leq q < N$, or $m = \min\left\{q, \frac{Nq_B}{N-1}\right\}$, whenever $q \geq N$.

Thanks to r > N/2 and $r_B > N - 1$, we can always choose

$$q \in (N/2, r), \quad q_B \in (N-1, r_B).$$

and then m > N, so $u \in C^{\nu}(\overline{\Omega})$ for $\nu = 1 - \frac{N}{m}$.

Moreover, since $u \in C^{\nu}(\overline{\Omega})$, $a \in L^{r}(\Omega)$ and $\tilde{f} \in C(\overline{\Omega})$, using the Hölder inequality, then, the product $|a(\cdot)||\tilde{f}(u(\cdot))| \in L^{r}(\Omega)$. Hence, $f(\cdot, u(\cdot)) \in L^{r}(\Omega)$.

Similarly, if $u \in C^{\nu}(\overline{\Omega})$, $a_B \in L^{r_B}(\partial\Omega)$ and $\tilde{f}_B \in C(\partial\Omega)$, by Hölder inequality, then, the product $|a_B(\cdot)||\tilde{f}_B(u(\cdot))| \in L^{r_B}(\partial\Omega)$. Hence, we can conclude that $f_B(\cdot, u(\cdot)) \in L^{r_B}(\partial\Omega)$.

Then (B.1) and (B.2) hold, ending the proof.

References

- H. Amann. Fixed point equations and nonlinear eigenvalue problems in ordered banach spaces. SIAM Review, 18(4):620–709, 1976.
- H. Brezis. Functional analysis, Sobolev spaces and partial differential equations. Universitext. Springer, New York, 2011.
- [3] A. Castro and R. Pardo. A priori bounds for positive solutions of subcritical elliptic equations. *Rev. Mat. Complut.*, 28(3):715–731, 2015.
- [4] A. Castro and R. Pardo. A priori estimates for positive solutions to subcritical elliptic problems in a class of non-convex regions. *Discrete Contin. Dyn. Syst. Ser. B*, 22(3):783–790, 2017.
- [5] M. Chhetri, N. Mavinga, and R. Pardo. An interpolation approach to L[∞] a priori estimates for elliptic problems with nonlinearity on the boundary. *Proc. Amer. Math. Soc.*, Accepted Manuscript, 2024.
- [6] L. Damascelli and R. Pardo. A priori estimates for some elliptic equations involving the p-Laplacian. Nonlinear Anal. Real World Appl., 41:475–496, 2018.
- [7] D. G. de Figueiredo, P.-L. Lions, and R. D. Nussbaum. A priori estimates and existence of positive solutions of semilinear elliptic equations. J. Math. Pures Appl. (9), 61(1):41–63, 1982.

- [8] P. Drábek, A. Kufner, and F. Nicolosi. Quasilinear elliptic equations with degenerations and singularities, volume 5 of De Gruyter Series in Nonlinear Analysis and Applications. Walter de Gruyter & Co., Berlin, 1997.
- [9] B. Gidas and J. Spruck. A priori bounds for positive solutions of nonlinear elliptic equations. *Comm. Partial Differential Equations*, 6(8):883–901, 1981.
- [10] D. Gilbarg and N. S. Trudinger. Elliptic partial differential equations of second order, volume Vol. 224 of Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin-New York, 1977.
- [11] A. Kufner, O. John, and S. Fučík. *Function spaces*. Monographs and Textbooks on Mechanics of Solids and Fluids, Mechanics: Analysis. Noordhoff International Publishing, Leiden; Academia, Prague, 1977.
- [12] O. A. Ladyzhenskaya and N. N. Ural'tseva. Linear and quasilinear elliptic equations. Academic Press, New York-London, 1968. Translated from the Russian by Scripta Technica, Inc, Translation editor: Leon Ehrenpreis.
- [13] J. Leray and J. Schauder. Topologie et équations fonctionnelles. Ann. Sci. École Norm. Sup. (3), 51:45–78, 1934.
- [14] G. M. Lieberman. Boundary regularity for solutions of degenerate elliptic equations. Nonlinear Analysis: Theory, Methods & Applications, 12(11):1203-1219, 1988.
- [15] G. Marino and P. Winkert. Moser iteration applied to elliptic equations with critical growth on the boundary. *Nonlinear Anal.*, 180:154–169, 2019.
- [16] N. Mavinga and R. Pardo. Bifurcation from infinity for reaction-diffusion equations under nonlinear boundary conditions. Proc. Roy. Soc. Edinburgh Sect. A, 147(3):649–671, 2017.
- [17] N. Mavinga and R. Pardo. A priori bounds and existence of positive solutions for semilinear elliptic systems. J. Math. Anal. Appl., 449(2):1172–1188, 2017.
- [18] J. Moser. A new proof of De Giorgi's theorem concerning the regularity problem for elliptic differential equations. *Comm. Pure Appl. Math.*, 13:457–468, 1960.
- [19] L. Nirenberg. On elliptic partial differential equations. Annali della Scuola Normale Superiore di Pisa - Scienze Fisiche e Matematiche, Ser. 3, 13(2):115– 162, 1959.
- [20] R. Pardo. On the existence of a priori bounds for positive solutions of elliptic problems, I. Rev. Integr. Temas Mat., 37(1):77–111, 2019.
- [21] R. Pardo. On the existence of a priori bounds for positive solutions of elliptic problems, II. Rev. Integr. Temas Mat., 37(1):113–148, 2019.
- [22] R. Pardo. L[∞](Ω) a priori estimates for subcritical semilinear elliptic equations with a Carathéodory non-linearity. J. Fixed Point Theory Appl., 25(2):Paper No. 44, 22, 2023.
- [23] R. Pardo. L[∞] a-priori estimates for subcritical p-laplacian equations with a Carathéodory non-linearity. Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM, 118(2):Paper No. 66, 21, 2024.
- [24] M. Struwe. Variational methods, volume 34 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, fourth edition, 2008. Applications to nonlinear partial differential equations and Hamiltonian systems.

26E. ANTONIO, M. ARCIGA-ALEJANDRE, R. PARDO, AND J. SANCHEZ-ORTIZ

(E. Antonio) UNIVERSIDAD AUTÓNOMA DE GUERRERO, GUERRERO, MÉXICO *Email address*: eaam0207130gmail.com

(M. Arciga-Alejandre) UNIVERSIDAD AUTÓNOMA DE GUERRERO, GUERRERO, MÉXICO

Email address: mpargica@uagro.mx

(R. Pardo) Universidad Complutense de Madrid, Madrid, Spain $\mathit{Email}\ address:\ \mathtt{rpardoQucm.es}$

(J. Sanchez-Ortiz) Universidad Autónoma de Guerrero, Guerrero, México

Email address: jsanchez@uagro.mx