LIPSCHITZ CONTINUITY AND EQUIVALENCE OF POSITIVE VISCOSITY AND WEAK SOLUTIONS TO TRUDINGER'S EQUATION

PETER LINDQVIST, MIKKO PARVIAINEN, AND JARKKO SILTAKOSKI

ABSTRACT. We show that stricty positive viscosity supersolutions to Trudinger's equation are local weak supersolutions when 1 . As an application, we show using theIshii-Lions method that not only viscosity but also weak solutions are Lipschitz continuousin the space variable. To the best of our knowledge, there is no preceding valid Lipschitzproofs in the literature.

Contents

| 1. Introduction | 1 |
|---|----|
| 1.1. Viscosity solutions | 3 |
| 2. Positive, uniformly continuous viscosity supersolutions are weak supersolutions | 4 |
| 2.1. Regularization via inf-convolution | 4 |
| 2.2. Local energy estimates and convergence of Du_{ε} in $L^{r}_{loc}(\Xi)$. | 14 |
| 2.3. Passing to the limit | 20 |
| 3. Local Lipschitz continuity in space | 22 |
| Appendix A. Basic properties | 32 |
| References | 35 |

1. INTRODUCTION

We study Trudinger's equation

$$(p-1)u^{p-2}\partial_t u - \Delta_p u = 0 \tag{1.1}$$

in an open and bounded domain $\Xi \subset \mathbb{R}^{N+1}$, with p > 1 and $\Delta_p u = \operatorname{div}(|Du|^{p-2}Du)$. We restrict ourselves to the (strictly) positive u and show that viscosity supersolutions are local weak supersolutions (Theorem 2.8). The reverse implication also holds so that positive weak and viscosity solutions actually coincide (Theorem 2.9). In the past, such questions

Date: February 21, 2025.

²⁰²⁰ Mathematics Subject Classification. 35K65,35K55, 35B65, 35D30,35D4.

Key words and phrases. Equivalence of solutions, Ishii-Lions method, Lipschitz, parabolic p-Laplace, Trudinger's equation, Viscosity solutions, Weak solutions.

MP is supported by the Research Council of Finland, project 360185. JS is supported by the Emil Aaltonen foundation. The authors would like to thank Juha Kinnunen for useful discussions.

have been studied for example by Ishii [Ish95] for linear elliptic equations and by Juutinen, Lindqvist, and Manfredi for the elliptic and parabolic *p*-Laplace equation in [JLM01]. As an application, we show that not only viscosity solutions but also weak solutions are Lipschitz continuous in the space variable (Theorem 3.1 and Corollary 3.2), which so far has been an open problem.

To prove that viscosity supersolutions are weak supersolutions, we use the technique introduced by Julin and Juutinen [JJ12] based on approximation by inf-convolution. However, many problems appear since there is u^{p-2} in Trudinger's equation. Problems caused by this factor in the theory are serious and, for example, full uniqueness without strict positivity is still an open problem. The more detailed idea of the proof in [JJ12] is that

- Step 1 It holds that inf-convolution is still a viscosity supersolution and by Alexandroff's theorem twice differentiable a.e.
- Step 2 Thus the *p*-Laplacian of the inf-convolution exist and supersolution property holds pointwise a.e. without difficulties at least when $p \ge 2$.
- Step 3 Then the idea is to multiply the *p*-Laplacian of the inf-convolution by a test function and integrate by parts. However, to justify this step further regularization is required.
- Step 4 Finally, one needs to be able to pass to a limit under the integral.

In our case, the counterpart of Step 1 is established in Lemma 2.4, which is the key lemma of the paper. For Trudinger's equation, this step is problematic since the term $u^{p-2}\partial_t u$ has u-dependence, and translation in the inf-convolution causes an extra error which will be difficult to control. Controlling the error requires careful balancing out with the gradient, and this idea is described right before the lemma. Step 2 and Step 3 are taken care of in Lemma 2.5, and again the error term will have to be taken into account. The counterpart of Step 4 is established in Section 2.2. To be able to pass to a limit under the integral sign, we need a uniform bound for the gradients of uniformly bounded weak supersolutions with the error. This is established in Lemma 2.6. Also observe that the weak convergence is not sufficient here. In order to pass to a limit under the integral sign, we need stronger convergence for the gradients obtained in Lemma 2.7. Using this, we finally obtain in Theorem 2.8 that a viscosity supersolution is a weak supersolution under our assumptions. We also establish the results in the case 1 using a special inf-convolution. Finally, the standard argument using Theorem 3 in [LL22] completes the picture and gives that weak solutions are viscosity solutions, and thus the equivalence result in Theorem 2.9.

As an application of the preceding equivalence result, we establish a qualitative Lipschitz result for positive viscosity and weak solutions in Section 3.3, see Theorem 3.1. In order to obtain this, we use the Ishii-Lions technique [IL90] from the theory of viscosity solutions twice. Since we showed above that viscosity solutions are weak solutions, this also implies Lipschitz continuity of positive weak solutions.

The equation (1.1) was originally suggested by Trudinger in [Tru68] as an example of the equation that could have simpler Harnack's inequality than the standard parabolic *p*-Laplace equation. Despite the homogeneity of the equation, the regularity theory turned out to be quite problematic. In [KK07], Kinnunen and Kuusi proved Harnack's inequality for Trudinger's equation using Moser's iteration, whereas Gianazza and Vespri established Harnack's inequality in [GV06] using De Giorgi's method. In [KSU12] Kuusi, Siljander and Urbano obtained Hölder continuity in the case 2 < p for nonnegative solutions, and in [KLSU12] the case 1 was treated by Kuusi, Laleoglu, Siljander and Urbano.In [DU20] Diehl and Urbano establish a sharp Hölder regularity for the inhomogeneousTrudinger's equation. Recently, the asymptotic behaviour and comparison principle forpositive solutions were studied by Hynd and Lindgren [HL21] as well as Lindgren andLindqvist [LL22] respectively.

Also the doubly nonlinear equation,

$$\partial_t (|u|^{q-1}u) - \Delta_p u = 0$$

has received a lot of attention. In [Hen20], Henriques established Hölder regularity for nonnegative weak solutions. Bögelein, Duzaar, Liao and Schätzler studied the Hölder regularity results for the doubly nonlinear equation in a series of papers [BDL21, BDLS22, LS22], where the first paper covers the Trudinger's equation also with sign-changing solutions. Misawa [Mis23] considered the expansion of positivity for doubly nonlinear parabolic equations including Trudinger's equation. In a recent work [BDG⁺23], Bögelein, Duzaar, Gianazza, Liao and Scheven consider continuity of the gradient of solutions to doubly non-linear equations in a certain range of parameters. However, this does not cover Trudinger's equation: To the best of our knowledge, there is no preceding correct Lipschitz proofs in the literature. It follows that the gradient of a solution exists a.e. by Rademacher's theorem in the usual sense in addition to the weak sense, and also that the gradient is locally bounded.

As for the alternative point of view, Bhattacharya and Marazzi studied viscosity solutions to Trudinger's equation and asymptotic behaviour of the viscosity solutions in [BM15] and [BM16]. They proved existence and uniqueness for positive solutions. However, their proof is based on logarithmic change of variables and is quite different from the techniques of this paper. As mentioned, we follow the approach in [JJ12]. Their technique was extended to the parabolic case in [PV20, Appendix A] and [Sil21]. This approach does not rely on the uniqueness theory of viscosity solution, which could also be used to study the equivalence of weak and viscosity solutions as for example in [JLM01]. The aforementioned article [LL22] also briefly discusses viscosity solutions and establishes that positive weak solutions are viscosity solution. Moreover, in [HL21] Hynd and Lindgren study the asymptotic benaviour and show that a related quantity converges to an extremal of Poincaré's inequality. To conclude, we note that Hynd and Lindgren investigated the Lipschitz regularity of viscosity solutions for a different class of doubly nonlinear parabolic equations in [HL19].

1.1. Viscosity solutions. We consider positive, continuous viscosity sub- and supersolutions, which are defined below.

Definition 1.1 (Viscosity solutions). Let $u : \Xi \to (0, \infty)$ be continuous. We say that u is a viscosity supersolution to (1.1) in Ξ if whenever $\phi \in C^2(\Xi)$ touches u from below at $(x,t) \in \Xi$, $D\phi(y,s) \neq 0, y \neq x$, we have

$$\lim_{(y,s)\to(x,t),y\neq x} \sup_{y\neq x} \left(\partial_t \left(\left| \phi(y,s) \right|^{p-2} \phi(y,s) \right) - \Delta_p \phi(y,s) \right) \ge 0.$$

Analogously, we say that u is a viscosity subsolution to (1.1) in Ξ if whenever $\varphi \in C^2(\Xi)$ touches u from above at $(x,t) \in \Xi$, $D\phi(y,s) \neq 0, y \neq x$, we have

$$\liminf_{y,s)\to(x,t),y\neq x} \left(\partial_t \left(\left| \phi(y,s) \right|^{p-2} \phi(y,s) \right) - \Delta_p \phi(y,s) \right) \le 0.$$

Finally, u is a viscosity solution if it is both viscosity sub- and supersolution.

Since we consider positive solutions, the above definition takes the form

$$\lim_{(y,s)\to(x,t),y\neq x} \sup \left((p-1)\partial_t \phi(y,s) - u^{2-p}(y,s)\Delta_p \phi(y,s) \right) \ge 0.$$

in the case of the supersolutions, and analogously in the case of the subsolutions. In the case $p \geq 2$, the definition of the viscosity solutions to Trudinger's equation in terms of parabolic jets is in [BM15]; here we allow the whole range $1 . When using the theorem on sums, we also need the notion of parabolic jets. For example we denote the parabolic subjet as <math>(\theta, \eta, X) \in \mathcal{P}^{2,-}u(x,t)$. For more details, see [Cra97, CIL92, Lin12, DFO14]. Weak solutions to (1.1) are defined below, where dz = dx dt.

Definition 1.2 (Weak solutions). Let $u \in L^p_{loc}(\Xi)$ with $Du \in L^p_{loc}(\Xi)$, where Du is the distributional gradient of u in space variable. We say that u is a *weak supersolution* to (1.1) if for all non-negative $\varphi \in C_0^{\infty}(\Xi)$ we have

$$\int_{\Xi} \left(-u^{p-1} \partial_t \varphi + |Du|^{p-2} Du \cdot D\varphi \right) dz \ge 0$$

For weak subsolutions, the above inequality is reversed and u is a weak solution if it is both weak sub- and supersolution.

2. Positive, uniformly continuous viscosity supersolutions are weak supersolutions

2.1. Regularization via inf-convolution. We use a special form of inf-convolution where the penalty term in time has different weight than the penalty term in space. Given $u: \Xi \to \mathbb{R}$, we denote

$$u_{\varepsilon}(x,t) := \inf_{(y,s)\in\Xi} \left\{ u(y,s) + \frac{|y-x|^q}{q\varepsilon^{q-1}} + \frac{|s-t|^2}{2\delta_{\varepsilon}} \right\},\tag{2.1}$$

where δ_{ε} is a suitable function of ε , q > p/(p-1) if 1 and <math>q = 2 if $p \ge 2$. By choosing small δ_{ε} compared to ε , we may ensure that the time penalty term is small compared to ε . This allows us to control an error term that appears in the inequality satisfied by the inf-convolution of a supersolution.

The following lemma collects some well known properties of inf-convolution. We postpone its proof into the appendix.

Lemma 2.1. Assume that $u : \Xi \to \mathbb{R}$ is lower semicontinuous and bounded. Suppose also that $\delta_{\varepsilon} \to 0$ as $\varepsilon \to 0$. Then u_{ε} has the following properties.

(i) We have $u_{\varepsilon} \leq u$ in Ξ and $u_{\varepsilon} \rightarrow u$ pointwise as $\varepsilon \rightarrow 0$.

(*ii*) Denote $r(\varepsilon) := (q\varepsilon^{q-1} \operatorname{osc}_{\Xi} u)^{\frac{1}{q}}, t(\varepsilon) := (2\delta_{\varepsilon} \operatorname{osc}_{\Xi} u)^{\frac{1}{2}} \text{ and set}$ $\Xi_{\varepsilon} := \left\{ (x,t) \in \Xi : B_{r(\varepsilon)}(x) \times (t-t(\varepsilon),t+t(\varepsilon)) \Subset \Xi \right\}.$

Then, for any $(x,t) \in \Xi_{\varepsilon}$ there exists $(x_{\varepsilon},t_{\varepsilon}) \in \overline{B}_{r(\varepsilon)}(x) \times [t-t(\varepsilon),t+t(\varepsilon)]$ such that

$$u_{\varepsilon}(x,t) = u(x_{\varepsilon},t_{\varepsilon}) + \frac{|x-x_{\varepsilon}|^{q}}{q\varepsilon^{q-1}} + \frac{|t-t_{\varepsilon}|^{2}}{2\delta_{\varepsilon}}.$$

- (iii) The function u_{ε} is semi-concave in Ξ_{ε} . In particular, the function $u_{\varepsilon}(x,t) (C|x|^2 + t^2/\delta_{\varepsilon})$ is concave in Ξ_{ε} , where $C = (q-1)r(\varepsilon)^{q-2}/\varepsilon^{q-1}$.
- (iv) Suppose that u_{ε} is twice differentiable in space and time at $(x,t) \in \Xi_{\varepsilon}$. Then

$$\partial_t u_{\varepsilon}(x,t) = \frac{t - t_{\varepsilon}}{\delta_{\varepsilon}},$$

$$Du_{\varepsilon}(x,t) = (x - x_{\varepsilon}) \frac{|x - x_{\varepsilon}|^{q-2}}{\varepsilon^{q-1}},$$

$$D^2 u_{\varepsilon}(x,t) \le (q-1) \frac{|x - x_{\varepsilon}|^{q-2}}{\varepsilon^{q-1}} I.$$

The following lemma gives an estimate for the penalty term appearing in inf-convolution.

Lemma 2.2. Suppose that $u : \Xi \to \mathbb{R}$ is uniformly continuous. Let ω_t and ω_x be bounded moduli of continuity of u in time and space, respectively. Suppose moreover that ω_t is strictly increasing. Let u_{ε} be defined by (2.1), where $\delta_{\varepsilon} = \frac{(\omega_t^{-1}(\varepsilon^{q-1}))^2}{2 \operatorname{osc}_{\Xi} u}$. Let $(x_{\varepsilon}, t_{\varepsilon})$ be such that

$$u_{\varepsilon}(x,t) = u(x_{\varepsilon},t_{\varepsilon}) + \frac{|x-x_{\varepsilon}|^{q}}{q\varepsilon^{q-1}} + \frac{|t_{\varepsilon}-t|^{2}}{2\delta_{\varepsilon}}.$$

Then

$$\frac{\left|t-t_{\varepsilon}\right|^{2}}{2\delta_{\varepsilon}} \leq \varepsilon^{q-1} \quad and \quad |x-x_{\varepsilon}| \leq q^{1/q} \varepsilon^{\frac{q-1}{q}} \omega_{x}^{1/q}(|x-x_{\varepsilon}|).$$

Proof. By the definition of inf-convolution, we have

$$\frac{|t-t_{\varepsilon}|^2}{2\delta_{\varepsilon}} \le u(x_{\varepsilon}, t_{\varepsilon}) - u(x_{\varepsilon}, t),$$
$$\frac{|x-x_{\varepsilon}|^q}{q\varepsilon^{q-1}} \le u(x_{\varepsilon}, t_{\varepsilon}) - u(x, t_{\varepsilon}).$$

From the first inequality, we estimate $|t - t_{\varepsilon}| \leq \sqrt{2\delta_{\varepsilon} \operatorname{osc}_{\Xi} u}$. Using the first inequality again and the definition of δ_{ε} , we obtain

$$\frac{|t-t_{\varepsilon}|^2}{2\delta_{\varepsilon}} \le \omega_t(|t-t_{\varepsilon}|) \le \omega_t(\sqrt{2\delta_{\varepsilon} \operatorname{osc} u}) \le \omega_t(\omega_t^{-1}(\varepsilon^{q-1})) = \varepsilon^{q-1}.$$

Similarly, from the second inequality we obtain that $|x - x_{\varepsilon}| \leq (q \varepsilon^{q-1} \omega_x (|x - x_{\varepsilon}|))^{1/q}$. \Box

The next elementary inequality immediately follows by integrating the derivative of $h \mapsto h^{\frac{2-p}{2}}$. For the convenience of the reader this is written down in the appendix.

Lemma 2.3 (Elementary inequality). Let p > 1. Suppose that $a \ge m > 0$. Then for all $s \in (0, 1)$ we have

$$\left| (a+s)^{\frac{2-p}{2}} - a^{\frac{2-p}{2}} \right| \le C(p,m)s.$$

Next we prove a key lemma: inf-convolution of a positive, uniformly continuous viscosity supersolution to Trudinger's equation solves the same equation with an additional first-order error term. Let us briefly highlight key arguments of the proof before the full details. Observe that a viscosity solution u to the Trudinger's equation solves

$$(p-1)\partial_t u - u^{2-p}\Delta_p u \ge 0.$$

This form is useful for estimating the error term generated by the inf-convolution u_{ε} , as the error generating *u*-dependence is now decoupled from the time derivative to the elliptic term and the Jensen-Ishii lemma can be used to control the behavior of the elliptic term. Using also the assumption that $0 < m \leq u \leq M$, the argument in [Ish95] and for the *p*-Laplacian in [Sil18] readily gives that u_{ε} is a viscosity supersolution to

$$(p-1)u_{\varepsilon}^{p-2}\partial_t u_{\varepsilon} - \Delta_p u_{\varepsilon} \ge -\mathcal{E} \quad \text{in} \quad \Xi_{\varepsilon},$$
(2.2)

where Ξ_{ε} is an in Lemma 2.1 and the error term has the form (assume for simplicity here that $p \ge 2$ so that q = 2)

$$\mathcal{E} = C \frac{|Du_{\varepsilon}|^{p-2}}{\varepsilon} \left(\frac{|x - x_{\varepsilon}|^4}{\varepsilon^2} + \frac{|t - t_{\varepsilon}|^4}{\delta_{\varepsilon}^2} \right)$$

Here, roughly speaking, the term $|Du_{\varepsilon}|^{p-2}/\varepsilon$ is generated by $\Delta_p u$ and the term in brackets comes from u^{2-p} . The term with $|t - t_{\varepsilon}|$ can be estimated via Lemma 2.2 and the choice of δ_{ε} . The term with $|x - x_{\varepsilon}|^4/\varepsilon^3$ is trickier to deal with, as this ratio may blow up as $\varepsilon \to 0$ unless u is known to be, say, Lipschitz. To deal with the problem, we use the fact that $Du_{\varepsilon} = \frac{x - x_{\varepsilon}}{\varepsilon}$. This together with the uniform continuity of u lets us absorb some of the problematic ratio to the gradient part in (2.9). This way, we arrive at the estimate

$$\mathcal{E} \leq E_{\varepsilon} \left(|Du_{\varepsilon}|^p h(|Du_{\varepsilon}|) + |Du_{\varepsilon}|^{p-2} \right),$$

where $h: [0, \infty) \to [0, \infty)$ is continuous (and thus bounded at 0) and $h(s) \to 0$ as $s \to \infty$ and E_{ε} is a number that converges to zero as $\varepsilon \to 0$. The term $h(|Du_{\varepsilon}|)$ is crucial in (2.18), as its presence lets us prove an energy estimate that ensures the weak convergence of Du_{ε} in L_{loc}^p up to a subsequence.

Lemma 2.4. Let u be a uniformly continuous viscosity supersolution to

$$(p-1)u^{p-2}\partial_t u - \Delta_p u \ge 0$$
 in Ξ .

Suppose that $0 < m \leq u \leq M$ in Ξ and let u_{ε} be defined by (2.1). Then, if $(\theta, \eta, X) \in \mathcal{P}^{2,-}u_{\varepsilon}(x,t)$ with $\eta \neq 0$, and $(x,t) \in \Xi_{\varepsilon}$, we have

$$(p-1)u_{\varepsilon}^{p-2}\theta - |\eta|^{p-2}\operatorname{tr}((I+(p-2)\frac{\eta\otimes\eta}{|\eta|^2})X) \ge -E_{\varepsilon}(|\eta|^p h(|\eta|) + |\eta|^{\max(p-2,0)}),$$

where $E_{\varepsilon} \geq 0$ is a quantity that converges to zero as $\varepsilon \to 0$, $h: [0, \infty) \to \mathbb{R}$ is continuous, strictly decreasing and $h(s) \to 0$ as $s \to \infty$.

Furthermore, in the case that $(\theta, 0, X) \in \mathcal{P}^{2,-}u_{\varepsilon}(x,t)$, we have $\theta \geq 0$.

Proof. Let us first consider the case $\eta \neq 0$. Since $(x, t) \in \Xi_{\varepsilon}$, by Lemma 2.1, there is $(x_{\varepsilon}, t_{\varepsilon})$ such that

$$u_{\varepsilon}(x,t) = u(x_{\varepsilon},t_{\varepsilon}) + \frac{|x-x_{\varepsilon}|^{q}}{q\varepsilon^{q-1}} + \frac{|t-t_{\varepsilon}|^{2}}{2\delta_{\varepsilon}} =: u(x_{\varepsilon},t_{\varepsilon}) + A.$$
(2.3)

It was assumed that $(\theta, \eta, X) \in \mathcal{P}^{2,-}u_{\varepsilon}(x,t)$, so there exists $\varphi \in C^2$ such that it touches u_{ε} from below at (x,t) and $(\partial_t \varphi(x,t), D\varphi(x,t), D^2\varphi(x,t)) = (\theta, \eta, X)$. By direct computation

$$\theta = \frac{t - t_{\varepsilon}}{\delta_{\varepsilon}}$$
 and $\eta = \frac{x - x_{\varepsilon}}{\varepsilon^{q-1}} |x - x_{\varepsilon}|^{q-2}$.

Using the definition of inf-convolution and that φ touches u_{ε} from below at (x, t), we have for all $(y, s), (z, \tau) \in \Xi$ that

$$-u(y,s) + \varphi(z,\tau) - \frac{|y-z|^q}{q\varepsilon^{q-1}} - \frac{|s-\tau|^2}{2\delta_{\varepsilon}} \le -u_{\varepsilon}(z,\tau) + \varphi(z,\tau) \le 0$$

$$= -u_{\varepsilon}(x,t) + \varphi(x,t)$$

$$= -u(x_{\varepsilon},t_{\varepsilon}) + \varphi(x,t) - \frac{|x-x_{\varepsilon}|^q}{q\varepsilon^{q-1}} - \frac{|t_{\varepsilon}-t|^2}{2\delta_{\varepsilon}}.$$

$$(2.4)$$

That is, if we denote

$$\Phi(y,s,z,\tau) := \frac{|y-z|^q}{q\varepsilon^{q-1}} + \frac{|s-\tau|^2}{2\delta_{\varepsilon}},$$

the function $(y, s, z, \tau) \mapsto -u(y, s) + \varphi(z, \tau) - \Phi(y, s, z, \tau)$ has maximum at the point $(y, s, z, \tau) = (x_{\varepsilon}, t_{\varepsilon}, x, t) =: a$.

Invoking the Jensen-Ishii lemma, we conjure symmetric $N \times N$ matrices Z and Y such that

$$(\partial_s \Phi(a), D_y \Phi(a), Y) \in \overline{\mathcal{P}}^{2,+}(-u(x_{\varepsilon}, t_{\varepsilon})), (-\partial_\tau \Phi(a), -D_z \Phi(a), Z) \in \overline{\mathcal{P}}^{2,-}(-\varphi(x, t)).$$

Since $\partial_s \Phi(a) = \frac{t_{\varepsilon} - t}{\delta_{\varepsilon}} = -\theta = \partial_{\tau} \Phi(a)$ and $D_y \Phi(a) = \frac{x_{\varepsilon} - x}{\varepsilon^{q-1}} |x - x_{\varepsilon}|^{q-2} = -\eta = D_z \Phi(x, x_{\varepsilon})$, these can be written as

$$(\theta, \eta, -Y) \in \overline{\mathcal{P}}^{2,-}u(x_{\varepsilon}, t_{\varepsilon}) \text{ and } (\theta, \eta, -Z) \in \overline{\mathcal{P}}^{2,+}\varphi(x, t).$$

Moreover, we have

$$\begin{pmatrix} Y & 0\\ 0 & -Z \end{pmatrix} \leq D^2_{(y,z)} \Phi(x_{\varepsilon}, x) + \varepsilon^{q-1} (D^2_{(y,z)} \Phi(x_{\varepsilon}, x))^2$$
$$= \begin{pmatrix} B & -B\\ -B & B \end{pmatrix} + 2\varepsilon^{q-1} \begin{pmatrix} B^2 & -B^2\\ -B^2 & B^2 \end{pmatrix},$$
(2.5)

where

$$B = \frac{1}{\varepsilon^{q-1}} |x - x_{\varepsilon}|^{q-4} \left((q-2)(x_{\varepsilon} - x) \otimes (x_{\varepsilon} - x) + |x_{\varepsilon} - x|^2 I \right).$$

This implies in particular $Y \leq Z \leq -D^2 \varphi(x,t) = -X$. Denote below

$$F(\eta, X) := -|\eta|^{p-2} \left(\operatorname{tr} X + (p-2) \frac{\eta' X \eta}{|\eta|^2} \right),$$

where η' denotes the transpose of the column vector η . Since u is a viscosity supersolution, we have by the degenerate ellipticity of F that

$$\begin{aligned} &(p-1)\theta + \frac{1}{u_{\varepsilon}^{p-2}(x,t)}F(\eta,X) \\ &= (p-1)\theta + \frac{|\eta|^{p-2}}{u^{p-2}(x_{\varepsilon},t_{\varepsilon})}F(\eta,-Y) - \frac{|\eta|^{p-2}}{u^{p-2}(x_{\varepsilon},t_{\varepsilon})}F(\eta,-Y) + \frac{|\eta|^{p-2}}{u_{\varepsilon}^{p-2}(x,t)}F(\eta,X) \\ &\geq -\frac{|\eta|^{p-2}}{u^{p-2}(x_{\varepsilon},t_{\varepsilon})}F(\eta,-Y) + \frac{|\eta|^{p-2}}{u_{\varepsilon}^{p-2}(x,t)}F(\eta,-Z) \\ &= \frac{|\eta|^{p-2}}{u_{\varepsilon}^{p-2}(x,t)}(\operatorname{tr}(I+(p-2)\frac{\eta\otimes\eta}{|\eta|^{2}})Z) - \frac{|\eta|^{p-2}}{u^{p-2}(x_{\varepsilon},t_{\varepsilon})}(\operatorname{tr}(I+(p-2)\frac{\eta\otimes\eta}{|\eta|^{2}})Y). \end{aligned}$$

Assuming that ε is small enough, we may suppose that $|x - x_{\varepsilon}| \leq 1$. We multiply the inequality (2.5) by the vector $(u^{\frac{2-p}{2}}(x_{\varepsilon}, t_{\varepsilon})\frac{\eta'}{|\eta|}, u_{\varepsilon}^{\frac{2-p}{2}}(x, t)\frac{\eta'}{|\eta|})$ from left and right, and use the elementary inequality in Lemma 2.3 to obtain

$$\begin{split} u_{\varepsilon}^{2-p}(x,t) \frac{\eta' Z\eta}{|\eta|^2} - u^{2-p}(x_{\varepsilon},t_{\varepsilon}) \frac{\eta' Y\eta}{|\eta|^2} &\geq -\left(u_{\varepsilon}^{\frac{2-p}{2}}(x,t) - u^{\frac{2-p}{2}}(x_{\varepsilon},t_{\varepsilon})\right)^2 \frac{\eta'}{|\eta|} \left(B + 2\varepsilon^{q-1}B^2\right) \frac{\eta'}{|\eta|} \\ &\geq -\left(u_{\varepsilon}^{\frac{2-p}{2}}(x,t) - u^{\frac{2-p}{2}}(x_{\varepsilon},t_{\varepsilon})\right)^2 \left(\|B\| + 2\varepsilon^{q-1}\|B^2\|\right) \\ &\geq -\frac{C(q)}{\varepsilon^{q-1}}|x - x_{\varepsilon}|^{q-2} \left(u_{\varepsilon}^{\frac{2-p}{2}}(x,t) - u^{\frac{2-p}{2}}(x_{\varepsilon},t_{\varepsilon})\right)^2 \\ &= -\frac{C(q)}{\varepsilon^{q-1}}|x - x_{\varepsilon}|^{q-2} \left((u(x_{\varepsilon},t_{\varepsilon}) + A)^{\frac{2-p}{2}} - u^{\frac{2-p}{2}}(x_{\varepsilon},t_{\varepsilon})\right)^2 \\ &\geq -C(m,p,q) \frac{|x - x_{\varepsilon}|^{q-2}}{\varepsilon^{q-1}}A^2, \end{split}$$

where the inequality $|x - x_{\varepsilon}| \leq 1$ was used in the estimate of $||B^2||$, and $||B|| = \max\{|\lambda| : \lambda \text{ is an eigenvalue of } B\}$. Similarly, multiplying the matrix inequality (2.5) by the vectors $(u_{\varepsilon}^{2-p}(x,t)e_i, u^{2-p}(x_{\varepsilon},t_{\varepsilon})e_i), i \in \{1,\ldots,N\}$, we see that

$$u_{\varepsilon}^{2-p}(x,t)\operatorname{tr}(Z) - u^{2-p}(x_{\varepsilon},t_{\varepsilon})\operatorname{tr}(Y) \ge -C(N,m,p,q)\frac{|x-x_{\varepsilon}|^{q-2}}{\varepsilon^{q-1}}A^{2}.$$

Thus, combining the last three displays, we arrive at

$$(p-1)\theta + \frac{1}{u_{\varepsilon}^{p-2}(x,t)}F(\eta,X) \ge -C(N,p,q,m) \left|\eta\right|^{p-2} \frac{|x-x_{\varepsilon}|^{q-2}}{\varepsilon^{q-1}} A^{2}.$$
 (2.6)

Now we proceed to estimate the error term. For the sake of exposition, let us consider the cases $1 and <math>p \ge 2$ separately.

Case $p \ge 2$: We use the definition of A, the choice q = 2, the inequality $(c+d)^2 \le C(c^2+d^2)$, and the estimate $\frac{|t-t_{\varepsilon}|^2}{\delta_{\varepsilon}} \le C\varepsilon$ from Lemma 2.2 to obtain

$$\frac{A^2}{\varepsilon} \le \frac{C}{\varepsilon} \left(\frac{|x - x_{\varepsilon}|^4}{\varepsilon^2} + \frac{|t - t_{\varepsilon}|^4}{\delta_{\varepsilon}^2} \right) \le C \frac{|x - x_{\varepsilon}|^4}{\varepsilon^3} + C\varepsilon.$$
(2.7)

Next, recall that by Lemma 2.2 we have

$$|x - x_{\varepsilon}| \le 2^{1/2} \omega_x^{1/2} (|x - x_{\varepsilon}|) \varepsilon^{1/2} \le C_0 \varepsilon^{1/2}$$
(2.8)

We define the function $h: [0, \infty) \to [0, \infty)$ by $h(s) := \omega_x^{1/2} (C_0^2 s^{-1})$ when $s \in (0, \infty)$ and $h(0) := \lim_{s \to 0} h(s)$. Observe that h is well defined, continuous, strictly decreasing and $h(s) \to 0$ as $s \to \infty$. Now, using that $|\eta| = \frac{|x - x_{\varepsilon}|}{\varepsilon}$, we estimate using both inequalities in (2.8)

$$\frac{|x - x_{\varepsilon}|^{4}}{\varepsilon^{3}} = |\eta|^{2} h(|\eta|) \frac{|x - x_{\varepsilon}|^{2}}{\varepsilon} \cdot \frac{1}{h(\frac{|x - x_{\varepsilon}|}{\varepsilon})} \leq 2 |\eta|^{2} h(|\eta|) \frac{\omega_{x}(|x - x_{\varepsilon}|)}{h(\frac{C_{0}}{\varepsilon^{1/2}})}$$
$$\leq 2 |\eta|^{2} h(|\eta|) \frac{\omega_{x}(C_{0}\varepsilon^{1/2})}{h(\frac{C_{0}}{\varepsilon^{1/2}})}$$
$$= 2 |\eta|^{2} h(|\eta|) \frac{\omega_{x}(C_{0}\varepsilon^{1/2})}{\omega_{x}^{1/2}(C_{0}^{2}\frac{\varepsilon^{1/2}}{C_{0}})}$$
$$= 2 |\eta|^{2} h(|\eta|) \omega_{x}^{1/2}(C_{0}\varepsilon^{1/2}). \tag{2.9}$$

Combining (2.6), (2.7) and (2.9), we have

$$(p-1)\theta - \frac{1}{u_{\varepsilon}^{p-2}(x,t)}F(\eta,X) \ge -E_{\varepsilon}(|\eta|^p h(|\eta|) + |\eta|^{p-2}),$$

where we denoted

$$E_{\varepsilon} := C(N, p, m, M) \max(\omega_x^{1/2}(C_0 \varepsilon^{1/2}), \varepsilon).$$

This yields the desired estimate in the case $p \ge 2$ since $C^{-1}(p, m, M) \le u_{\varepsilon}^{p-2} \le C(p, m, M)$.

Case $1 : Now <math>q \neq 2$, so we have

$$\begin{aligned} |\eta|^{p-2} \frac{|x-x_{\varepsilon}|^{q-2}}{\varepsilon^{q-1}} A^2 &\leq C \, |\eta|^{p-2} \, \frac{|x-x_{\varepsilon}|^{q-2}}{\varepsilon^{q-1}} \left(\frac{|x-x_{\varepsilon}|^{2q}}{q^2 \varepsilon^{2(q-1)}} + \frac{|t-t_{\varepsilon}|^4}{4\delta_{\varepsilon}^2} \right) \\ &= C(q) \, |\eta|^{p-2} \, \frac{|x-x_{\varepsilon}|^{3q-2}}{q^2 \varepsilon^{3(q-1)}} + |\eta|^{p-2} \, \frac{|x-x_{\varepsilon}|^{q-2}}{\varepsilon^{q-1}} \frac{|t-t_{\varepsilon}|^4}{4\delta_{\varepsilon}^2}. \\ &=: T_1 + T_2. \end{aligned}$$

Recall that by Lemma 2.2

$$|x - x_{\varepsilon}| \le \omega_x^{1/q} (|x - x_{\varepsilon}|) \varepsilon^{\frac{q-1}{q}} \le C_0(q, M) \varepsilon^{\frac{q-1}{q}}.$$
(2.10)

To estimate T_2 we use that q > p/(p-1) implies (q-1)(p-1) - 1 > 0. We also use the estimate $\frac{|t-t_{\varepsilon}|^2}{\delta_{\varepsilon}} \leq \varepsilon^{q-1}$ from Lemma 2.2, and inequality (2.10). We obtain

$$T_{2} = \frac{|\eta|^{p-2} |x - x_{\varepsilon}|^{q-2}}{4\varepsilon^{q-1}} \frac{|t - t_{\varepsilon}|^{4}}{\delta_{\varepsilon}^{2}} = \frac{|x - x_{\varepsilon}|^{(q-1)(p-1)-1}}{4\varepsilon^{q-1}} \frac{|t - t_{\varepsilon}|^{4}}{\delta_{\varepsilon}^{2}}$$
$$\leq C(q, M) \frac{\varepsilon^{\frac{q-1}{q}((q-1)(p-1)-1)}\varepsilon^{2(q-1)}}{\varepsilon^{q-1}}}{\varepsilon^{q-1}}$$

To estimate T_1 , we define $h(s) := \omega_x^{1/2} (C_0^2 s^{-1})$ like in the case $p \ge 2$, where $C_0 = C_0(q, M)$ is the constant in (2.10). Then we have using also both inequalities in (2.10)

$$\begin{aligned} \frac{|x - x_{\varepsilon}|^{3q-2}}{\varepsilon^{3(q-1)}} &= |\eta|^2 h(|\eta|) \frac{|x - x_{\varepsilon}|^{-2(q-1)}}{\varepsilon^{-2(q-1)}} \frac{|x - x_{\varepsilon}|^{3q-2}}{\varepsilon^{3(q-1)}} \frac{1}{h(\frac{|x - x_{\varepsilon}|^{q-1}}{\varepsilon^{q-1}})} \\ &= |\eta|^2 h(|\eta|) \frac{|x - x_{\varepsilon}|^q}{\varepsilon^{q-1}} \frac{1}{h(\frac{|x - x_{\varepsilon}|^{q-1}}{\varepsilon^{q-1}})} \\ &\leq q |\eta|^2 h(|\eta|) \frac{\omega_x(|x - x_{\varepsilon}|)}{h(C_0 \varepsilon^{\frac{(q-1)^2}{q} - (q-1)})} \\ &\leq q |\eta|^2 h(|\eta|) \frac{\omega_x(C_0 \varepsilon^{\frac{q-1}{q}})}{h(C_0 \varepsilon^{-\frac{q-1}{q}})} \\ &= q |\eta|^2 h(|\eta|) \frac{\omega_x(C_0 \varepsilon^{\frac{q-1}{q}})}{\omega_x^{1/2}(C_0^2 \varepsilon^{\frac{q-1}{q}})} \\ &= q |\eta|^2 h(|\eta|) \omega_x^{1/2}(C_0 \varepsilon^{\frac{q-1}{q}}). \end{aligned}$$

Therefore, we obtain

$$T_1 = C(q) |\eta|^{p-2} \frac{|x - x_{\varepsilon}|^{3q-2}}{\varepsilon^{3(q-1)}} \le C(q) |\eta|^p h(|\eta|) \omega_x^{1/2} (C_0 \varepsilon^{\frac{q-1}{q}}).$$

Combining the estimates of T_1 and T_2 with (2.6), we obtain

$$(p-1)\theta - \frac{1}{u_{\varepsilon}^{p-2}(x,t)}F(\eta,X) \ge -E_{\varepsilon}(|\eta|^p h(|\eta|) + 1),$$

where we have denoted

$$E_{\varepsilon} := C(N, p, q, m, M) \max(\omega_x^{1/2}(C_0 \varepsilon^{\frac{q-1}{q}}), \varepsilon^{q-1}).$$

This yields the desired estimate in the case 1 .

It remains to consider the case $\eta = 0$. Now $x = x_{\varepsilon}$ and so taking $(z, \tau) = (x, t)$ in (2.4), we see that the mapping $\phi(y, s) := u(x, t_{\varepsilon}) - \frac{|y-x|^q}{q\varepsilon^{q-1}} - \frac{|s-t|^2}{2\delta_{\varepsilon}} + \frac{|t_{\varepsilon}-t|^2}{2\delta_{\varepsilon}}$ touches u from below at $(y, s) = (x, t_{\varepsilon})$. Hence, since u is a viscosity supersolution and $D\phi(y, s) \neq 0$ when $y \neq x$, we have

$$\limsup_{(y,s)\to(x,t_{\varepsilon}),y\neq x} \left((p-1)\partial_s \phi(y,s) - u^{2-p}(y,s)\Delta_p \phi(y,s) \right) \ge 0.$$

On the other hand, since q > p/(p-1) or $p \ge 2$, we have $\Delta_p \phi(y,s) \to 0$ as $y \to x$. Therefore, we obtain $0 \le (p-1)\partial_s \phi(x,t_{\varepsilon}) = \frac{t-t_{\varepsilon}}{\delta_{\varepsilon}} = \theta$.

Next, we use the semi-concavity of inf-convolution to show that in the context of Lemma 2.4, the function u_{ε} is also a weak supersolution with some error term.

Lemma 2.5. Let u be a uniformly continuous viscosity supersolution to

$$(p-1)u^{p-2}\partial_t u - \Delta_p u \ge 0 \quad in \quad \Xi.$$

Suppose that $0 < m \leq u \leq M$ in Ξ and let u_{ε} be as in Lemma 2.2. Then u_{ε} is a weak supersolution to

$$\partial_t u_{\varepsilon}^{p-1} - \Delta_p u_{\varepsilon} \ge G(Du_{\varepsilon}) \quad in \quad \Xi_{\varepsilon}, \tag{2.11}$$

where

$$G(Du_{\varepsilon}) = \begin{cases} 0 & \text{if } |Du_{\varepsilon}| = 0, \\ -E_{\varepsilon} \left(|Du_{\varepsilon}|^{p} h(|Du_{\varepsilon}|) + |Du_{\varepsilon}|^{\max(0, p-2)} \right) & \text{if } |Du_{\varepsilon}| \neq 0. \end{cases}$$
(2.12)

Here E_{ε} and h are as in Lemma 2.4.

Proof. Fix non-negative $\varphi \in C_0^{\infty}(\Xi_{\varepsilon})$. By Lemma 2.1, the function

$$\phi(x,t) := u_{\varepsilon}(x,t) - C(|x|^2 + t^2),$$

where $C = C(q, \varepsilon)$, is concave in Ξ_{ε} . Therefore, we can approximate it by smooth concave functions ϕ_i so that

$$(\partial_t \phi_j, D\phi_j, D^2 \phi_j) \to (\partial_t \phi, D\phi, D^2 \phi)$$
 a.e. in Ξ_{ε} .

Above $D^2\phi$ denotes the Alexandrov derivative, see an unpublished version of [Lin12] or [EG92, Section 6.4, pp. 242–245]. We define

$$u_{\varepsilon,j}(x,t) := \phi_j(x,t) + C(|x|^2 + t^2).$$

We regularize the equation in order to remove the singularities, and to this end let $\delta \in (0, 1)$. Then we compute using integration by parts

$$\begin{split} &\int_{\Xi_{\varepsilon}} \varphi(p-1) u_{\varepsilon,j}^{p-2} \partial_t u_{\varepsilon,j} \, dz \\ &+ \int_{\Xi_{\varepsilon}} -\varphi(\delta + |Du_{\varepsilon,j}|^2)^{\frac{p-2}{2}} (\Delta u_{\varepsilon,j} + \frac{p-2}{\delta + |Du_{\varepsilon,j}|^2} Du_{\varepsilon,j}' D^2 u_{\varepsilon,j} Du_{\varepsilon,j}) - \varphi G(Du_{\varepsilon,j}) \, dz \\ &= \int_{\Xi_{\varepsilon}} \varphi \partial_t u_{\varepsilon,j}^{p-1} - \varphi \operatorname{div} \left((\delta + |Du_{\varepsilon,j}|^2)^{\frac{p-2}{2}} Du_{\varepsilon,j} \right) - \varphi G(Du_{\varepsilon,j}) \, dz \\ &= \int_{\Xi_{\varepsilon}} -u_{\varepsilon,j}^{p-1} \partial_t \varphi + (\delta + |Du_{\varepsilon,j}|^2)^{\frac{p-2}{2}} Du_{\varepsilon,j} \cdot D\varphi - \varphi G(Du_{\varepsilon,j}) \, dz. \end{split}$$

We denote

$$\Delta_{p,\delta} u_{\varepsilon,j} := (\delta + |Du_{\varepsilon,j}|^2)^{\frac{p-2}{2}} (\Delta u_{\varepsilon,j} + \frac{p-2}{\delta + |Du_{\varepsilon,j}|^2} Du'_{\varepsilon,j} D^2 u_{\varepsilon,j} Du_{\varepsilon,j}),$$

so that by the previous computation we have

$$\begin{split} &\lim_{j\to\infty} \inf_{\Xi_{\varepsilon}} \varphi((p-1)u_{\varepsilon,j}^{p-2}\partial_t u_{\varepsilon,j} - \Delta_{p,\delta} u_{\varepsilon,j} - G(Du_{\varepsilon,j})) \, dz \\ &= \lim_{j\to\infty} \int_{\Xi_{\varepsilon}} -u_{\varepsilon,j}^{p-1}\partial_t \varphi + (\delta + |Du_{\varepsilon,j}|^2)^{\frac{p-2}{2}} Du_{\varepsilon,j} \cdot D\varphi - \varphi G(Du_{\varepsilon,j}) \, dz. \end{split}$$

We use Fatou's lemma at the left-hand side and dominated convergence at the right-hand side. This way, we arrive at

$$\int_{\Xi_{\varepsilon}} \varphi((p-1)u_{\varepsilon}^{p-2}\partial_{t}u_{\varepsilon} - \Delta_{p,\delta}u_{\varepsilon} - G(Du_{\varepsilon})) dz$$

$$\leq \int_{\Xi_{\varepsilon}} -u_{\varepsilon}^{p-1}\partial_{t}\varphi + (\delta + |Du_{\varepsilon}|^{2})^{\frac{p-2}{2}}Du_{\varepsilon} \cdot D\varphi - \varphi G(Du_{\varepsilon}) dz.$$
(2.13)

To justify the use of Fatou's lemma, observe that by concavity of ϕ_j we have $D^2 u_{\varepsilon,j} \leq CI$. Thus the integral at the left-hand side of (2.13) has a lower bound when $Du_{\varepsilon,j} = 0$. When $Du_{\varepsilon,j} \neq 0$, we have

$$\begin{split} -\Delta_{p,\delta}u_{\varepsilon,j} &= -\left(\delta + |Du_{\varepsilon,j}|^2\right)^{\frac{p-2}{2}} \left(\Delta u_{\varepsilon,j} + \frac{p-2}{\delta + |Du_{\varepsilon,j}|^2} Du'_{\varepsilon,j} D^2 u_{\varepsilon,j} D^2 u_{\varepsilon,j}\right) \\ &= -\frac{\left(\delta + |Du_{\varepsilon,j}|^2\right)^{\frac{p-2}{2}}}{\delta + |Du_{\varepsilon,j}|^2} \left(|Du_{\varepsilon,j}|^2 \left(\Delta u_{\varepsilon,j} + \frac{p-2}{|Du_{\varepsilon,j}|^2} Du'_{\varepsilon,j} D^2 u_{\varepsilon,j} Du_{\varepsilon,j}\right) + \delta \Delta u_{\varepsilon,j}\right) \\ &\geq -\frac{\left(\delta + |Du_{\varepsilon,j}|^2\right)^{\frac{p-2}{2}}}{\delta + |Du_{\varepsilon,j}|^2} C(|Du_{\varepsilon,j}|^2 (N+p-2) + \delta N) \end{split}$$

$$\geq -C \frac{\left(\delta + |Du_{\varepsilon,j}|^2\right)^{\frac{p-2}{2}}}{\delta + |Du_{\varepsilon,j}|^2} (|Du_{\varepsilon,j}|^2 + \delta) \max(N, N+p-2)$$
$$= -C(\delta + |Du_{\varepsilon,j}|^2)^{\frac{p-2}{2}} \max(N, N+p-2),$$

where the right-hand side is bounded independently of j since Du_{ε} is bounded. Therefore, our use of Fatou's lemma is justified. Next, we let $\delta \to 0$. By the dominated convergence theorem we have

$$\lim \inf_{\delta \to 0} \int_{\Xi_{\varepsilon}} -u_{\varepsilon}^{p-1} \partial_t \varphi + (\delta + |Du_{\varepsilon}|^2)^{\frac{p-2}{2}} Du_{\varepsilon} \cdot D\varphi - \varphi G(Du_{\varepsilon}) dz$$
$$= \int_{\Xi_{\varepsilon}} -u_{\varepsilon}^{p-1} \partial_t \varphi + |Du_{\varepsilon}|^{p-2} Du_{\varepsilon} \cdot D\varphi - \varphi G(Du_{\varepsilon}) dz,$$

where the term $|Du_{\varepsilon}|^{p-2} Du_{\varepsilon}$ is read as zero wherever $Du_{\varepsilon} = 0$. On the other hand, by Fatou's lemma we have

$$\begin{split} & \liminf_{\delta \to 0} \int_{\Xi_{\varepsilon}} \left((p-1) u_{\varepsilon}^{p-2} \partial_{t} u_{\varepsilon} - \Delta_{p,\delta} u_{\varepsilon} - G(Du_{\varepsilon}) \right) \varphi \, dz \\ & \geq \int_{\Xi_{\varepsilon}} \liminf_{\delta \to 0} \left((p-1) u_{\varepsilon}^{p-2} \partial_{t} u_{\varepsilon} - \Delta_{p,\delta} u_{\varepsilon} - G(Du_{\varepsilon}) \right) \varphi \, dz \\ & = \int_{\Xi_{\varepsilon} \cap \{Du_{\varepsilon} \neq 0\}} \liminf_{\delta \to 0} \left((p-1) u_{\varepsilon}^{p-2} \partial_{t} u_{\varepsilon} - \Delta_{p,\delta} u_{\varepsilon} - G(Du_{\varepsilon}) \right) \varphi \, dz \\ & + \int_{\Xi_{\varepsilon} \cap \{Du_{\varepsilon} = 0\}} \liminf_{\delta \to 0} \left((p-1) u_{\varepsilon}^{p-2} \partial_{t} u_{\varepsilon} \right) \varphi \, dz \\ & \geq 0, \end{split}$$

where the last estimate follows from Lemma 2.4. We still need to justify our use of Fatou's lemma. When $Du_{\varepsilon} = 0$, this follows directly from the inequality

$$D^2 u_{\varepsilon} \leq \frac{q-1}{\varepsilon} \left| D u_{\varepsilon} \right|^{\frac{q-2}{q-1}} I,$$

which holds by Lemma 2.1. When $Du_{\varepsilon} \neq 0$, we estimate

$$\begin{split} -\Delta_{p,\delta} u_{\varepsilon} &= -\left(\delta + |Du_{\varepsilon}|^{2}\right)^{\frac{p-2}{2}} \left(\Delta u_{\varepsilon} + \frac{p-2}{\delta + |Du_{\varepsilon}|^{2}} Du_{\varepsilon}' D^{2} u_{\varepsilon} Du_{\varepsilon}\right) \\ &= -\frac{\left(\delta + |Du_{\varepsilon}|^{2}\right)^{\frac{p-2}{2}}}{\delta + |Du_{\varepsilon}|^{2}} \left(|Du_{\varepsilon}|^{2} \left(\Delta u_{\varepsilon} + \frac{p-2}{|Du_{\varepsilon}|} Du_{\varepsilon}' D^{2} u_{\varepsilon} Du_{\varepsilon}\right) + \delta \Delta u_{\varepsilon}\right) \\ &\geq -\frac{\left(\delta + |Du_{\varepsilon}|^{2}\right)^{\frac{p-2}{2}}}{\delta + |Du_{\varepsilon}|^{2}} \frac{q-1}{\varepsilon} \left(|Du_{\varepsilon}|^{\frac{q-2}{q-1}+2} \left(N+p-2\right) + |Du_{\varepsilon}|^{\frac{q-2}{q-1}} \delta N\right) \\ &\geq -\frac{\left(\delta + |Du_{\varepsilon}|^{2}\right)^{\frac{p-2}{2}}}{\delta + |Du_{\varepsilon}|^{2}} \frac{q-1}{\varepsilon} \left((|Du_{\varepsilon}|^{2}+\delta) |Du_{\varepsilon}|^{\frac{q-2}{q-1}+2} \max(N,N+p-2)\right) \end{split}$$

$$\geq -(\delta + |Du_{\varepsilon}|^2)^{\frac{p-2}{2}} |Du_{\varepsilon}|^{\frac{q-2}{q-1}} \frac{(q-1)}{\varepsilon} \max(N, N+p-2)$$

$$\geq -(\delta + |Du_{\varepsilon}|^2)^{\frac{1}{2}(p-2+\frac{q-2}{q-1})} \frac{(q-1)}{\varepsilon} \max(N, N+p-2),$$

which is bounded independently of δ since $p - 2 + \frac{q-2}{q-1} > 0$.

2.2. Local energy estimates and convergence of Du_{ε} in $L^r_{loc}(\Xi)$. Next we establish an energy estimate for weak supersolutions to the equation (2.11). Observe that the growth of the first-order term $|\eta|^p h(\eta)$ may be stronger than $|\eta|^q$ for any q < p. As in the proof of an energy estimate one usually tests with the solution itself, the term $u |Du|^p h(Du)$ has to be dealt with. This could be difficult if we only knew that u is in some Sobolev space. However, since we are dealing with locally bounded supersolutions, this is not a problem for us.

If u_{ε} is the inf-convolution of a viscosity supersolution to (2.11), then the energy estimate together with Lemma 2.5 guarantees that Du_{ε} converges weakly in $L_{loc}^{p}(\Xi)$ as $\varepsilon \to 0$ up to a subsequence. However, eventually we need to pass to the limit under the integral sign of

$$\int_{\Xi} -u_{\varepsilon}^{p-1} \partial_t \varphi + \left| Du_{\varepsilon} \right|^{p-2} Du_{\varepsilon} \cdot D\varphi + \varphi G(Du_{\varepsilon}) \, dz,$$

where φ is an arbitrary fixed test function and G is as in Lemma 2.5. For this we need stronger convergence, which we establish in Lemma 2.7. Since we apply this lemma to infconvolutions that are locally Lipschitz, the Lipschitz assumption in the next lemma does not impose any additional restriction for our purposes.

Lemma 2.6 (Caccioppoli's estimate). Assume that u is a locally Lipschitz continuous weak supersolution to

$$\partial_t u^{p-1} - \Delta_p u \ge G(Du) \quad in \ \Xi,$$

where G is defined by the formula (2.12) for some number $E_{\varepsilon} \in (0,1]$ and continuous, decreasing function $h: [0,\infty) \to [0,\infty)$ such that $h(s) \to 0$ as $s \to \infty$. Suppose moreover that $0 \le u \le M$ in Ξ . Then

$$\int_{\Xi} |Du|^p \,\xi^p \,dz \le C(p, M, h) \int_{\Xi} |\partial_t \xi^p| + |D\xi|^p + \xi^p \,dz$$

for any cut-off function $\xi \in C_0^{\infty}(\Xi), \ \xi \geq 0.$

Proof. Define $\varphi := \xi^p(M - u) \ge 0$. By local Lipschitz continuity (both in space and time) φ is an admissible test function whose time derivative exist a.e., and we obtain using the definition of G

$$\int_{\Xi} \xi^{p} |Du|^{p} dz \leq \int_{\Xi} u^{p-1} \partial_{t} \varphi + p \xi^{p-1} (M-u) |Du|^{p-1} |D\xi| dz + \int_{\Xi \cap \{Du \neq 0\}} \varphi E_{\varepsilon} (|Du|^{p} h(|Du|) + |Du|^{\max(p-2,0)}) dz.$$
(2.14)

We proceed to estimate the terms on the right-hand side. We have by integration by parts

$$\int_{\Xi} u^{p-1} \partial_t ((M-u)\xi^p) dz = \int_{\Xi} -\frac{1}{p} \partial_t u^p \xi^p + u^{p-1} (M-u) \partial_t \xi^p dz$$
$$= \int_{\Xi} \frac{1}{p} u^p \partial_t \xi^p + u^{p-1} (M-u) \partial_t \xi^p dz$$
$$\leq \int_{\Xi} C(p) M^p |\partial_t \xi^p| dz.$$
(2.15)

Using Young's inequality we obtain

$$\int_{\Xi} p\xi^{p-1}(M-u) |Du|^{p-1} |D\xi| \, dz \le \int_{\Xi} \frac{1}{4} \xi^p |Du|^p \, dz + C(p) \int_{\Xi} M^p |D\xi|^p \, dz.$$
(2.16)

Further, if p > 2, using Young's inequality with the exponents p/(p-2) and p/2, we get

$$\int_{\Xi \cap \{Du_{\varepsilon} \neq 0\}} \varphi E_{\varepsilon} |Du|^{\max(p-2,0)} dz = \int_{\Xi} E_{\varepsilon} \xi^{p-2} |Du|^{p-2} \xi^{2} (M-u) dz$$
$$\leq \int_{\Xi} \frac{1}{4} \xi^{p} |Du|^{p} dz + \int_{\Xi} \xi^{p} E_{\varepsilon}^{p/2} M^{p/2} dz.$$
(2.17)

Finally, since $h(s) \to 0$ as $s \to \infty$, there exists S > 0 such that $h(s) \leq \frac{1}{4M}$ when s > S. Using this, we estimate

$$\int_{\Xi} E_{\varepsilon} \varphi |Du|^{p} h(|Du|) dz \leq \int_{\Xi \cap \{|Du| \leq S\}} E_{\varepsilon} M |Du|^{p} h(0) \xi^{p} dz + \int_{\Xi \cap \{|Du| > S\}} E_{\varepsilon} M |Du|^{p} \frac{1}{4M} \xi^{p} dz \leq \int_{\Xi \cap \{|Du| \leq S\}} E_{\varepsilon} M S^{p} h(0) \xi^{p} dz + \int_{\Xi \cap \{|Du| > S\}} \frac{1}{4} |Du|^{p} \xi^{p} dz.$$

$$(2.18)$$

Combining the estimates (2.15)-(2.18) with (2.14) and absorbing the term with $\frac{1}{4} |Du|^p$ to the left-hand side, we arrive at

$$\begin{split} \int_{\Xi} \frac{1}{2} \left| Du \right|^p \xi^p \, dz &\leq \int_{\Xi} C(p) M^p \left| \partial_t \xi^p \right| + C(p) M^p \left| D\xi \right|^p + E_{\varepsilon} M S^p h(0) \xi^p \, dz \\ &+ \begin{cases} \int_{\Xi} \xi^p M E_{\varepsilon} \, dz, & 1 2. \end{cases} \end{split}$$

In the case p > 2 we absorb the additional term with $\frac{1}{4} |Du|^p$ to the left-hand side. The claimed estimate follows.

Lemma 2.7. Let (u_l) be a sequence of locally Lipschitz continuous weak supersolutions to

$$\partial_t u^{p-1} - \Delta_p u \ge G(Du) \quad in \quad \Xi$$

where G is defined by the formula (2.12) for some number $E_{\varepsilon} \in (0,1]$ and continuous, decreasing function $h: [0,\infty) \to [0,\infty)$ such that $h(s) \to 0$ as $s \to \infty$. Suppose that $u_l \to u$ pointwise almost everywhere in Ξ . Assume moreover that for all $l \in \mathbb{N}$ we have

$$0 < m \leq u_l \leq M < \infty$$
 in Ξ

Then (Du_l) is a Cauchy sequence in $L^r_{loc}(\Xi)$ for any 1 < r < p.

Proof. Let $U \in \Xi$ and take a cut-off function $\zeta \in C_0^{\infty}(\Xi)$ such that $0 \leq \zeta \leq 1$ and $\zeta \equiv 1$ in U. For $\delta > 0$, set

$$w_{jk} := \begin{cases} \delta, & u_j^{p-1} - u_k^{p-1} > \delta. \\ u_j^{p-1} - u_k^{p-1}, & -\delta < u_j^{p-1} - u_k^{p-1} < \delta, \\ -\delta, & u_j^{p-1} - u_k^{p-1} < -\delta. \end{cases}$$

Then $(\delta - w_{jk})\zeta \ge 0$ is an admissible test function by local Lipschitz continuity and the fact that $w_{jk} \in [-\delta, \delta]$. Moreover, the test function has time derivatives almost everywhere. Using it as a test function for u_j , we obtain

$$0 \leq \int_{\Xi} -u_{j}^{p-1} \partial_{t} ((\delta - w_{jk})\zeta) + |Du_{j}|^{p-2} Du_{j} \cdot D((\delta - w_{jk})\zeta) dz + \int_{\Xi \cap \{Du_{j} \neq 0\}} -E_{\varepsilon} (\delta - w_{jk})\zeta (|Du_{j}|^{p} h(|Du_{j}|) + |Du_{j}|^{\max(0, p-2)}) dz = \int_{\Xi} -\zeta |Du_{j}|^{p-2} Du_{j} \cdot Dw_{jk} + (\delta - w_{jk}) |Du_{j}|^{p-2} Du_{j} \cdot D\zeta dz + \int_{\Xi \cap \{Du_{j} \neq 0\}} -E_{\varepsilon} (\delta - w_{jk})\zeta (|Du_{j}|^{p} h(|Du_{j}|) + |Du_{j}|^{\max(0, p-2)}) dz + \int_{\Xi} \zeta u_{j}^{p-1} \partial_{t} w_{jk} - u_{j}^{p-1} (\delta - w_{jk}) \partial_{t} \zeta dz.$$
(2.19)

Then, since

$$Dw_{jk} = (p-1)\chi_{\{|u_j^{p-1} - u_k^{p-1}| < \delta\}}(u_j^{p-2}Du_j - u_k^{p-2}Du_k),$$

and $|w_{jk}| \leq \delta$, it follows from (2.19) that

$$\begin{split} &\int_{\Xi \cap \{|u_{j}^{p-1}-u_{k}^{p-1}| < \delta\}} \zeta(p-1) |Du_{j}|^{p-2} Du_{j} \cdot (u_{j}^{p-2} Du_{j} - u_{k}^{p-2} Du_{k}) dz \\ &\leq \int_{\Xi \cap \{Du_{j} \neq 0\}} 2\delta |Du_{j}|^{p-1} |D\zeta| + 2\delta E_{\varepsilon} \zeta(|Du_{j}|^{p} h(|Du_{j}|) + |Du_{j}|^{\max(0,p-2)}) dz \\ &+ \int_{\Xi} \zeta u_{j}^{p-1} \partial_{t} w_{jk} + 2\delta u_{j}^{p-1} |\partial_{t} \zeta| dz. \end{split}$$

Since u_k is a weak supersolution, the same arguments except testing now with $(\delta + w_{jk})\zeta \ge 0$ yields

$$\begin{split} &\int_{\Xi \cap \{ |u_{j}^{p-1} - u_{k}^{p-1}| < \delta \}} -\zeta(p-1) |Du_{k}|^{p-2} Du_{k} \cdot (u_{j}^{p-2} Du_{j} - u_{k}^{p-2} Du_{k}) dz \\ &\leq \int_{\Xi \cap \{ Du_{k} \neq 0 \}} 2\delta |Du_{k}|^{p-1} |D\zeta| + 2\delta E_{\varepsilon} \zeta(|Du_{k}|^{p} h(|Du_{k}|) + |Du_{k}|^{\max(0,p-2)}) dz \\ &\quad + \int_{\Xi} -\zeta u_{k}^{p-1} \partial_{t} w_{jk} + 2\delta u_{k}^{p-1} |\partial_{t} \zeta| dz. \end{split}$$

The idea of choosing the above test functions is to get, combining the last two inequalities, the following estimate where the first term contains a useful difference

$$\begin{split} &(p-1)\int_{\Xi\cap\{|u_{j}^{p-1}-u_{k}^{p-1}|<\delta\}}\zeta(|Du_{k}|^{p-2}Du_{k}-|Du_{j}|^{p-2}Du_{j})\cdot(u_{j}^{p-2}Du_{j}-u_{k}^{p-2}Du_{k})\,dz\\ &\leq \int_{\Xi}\zeta(u_{j}^{p-1}-u_{k}^{p-1})\partial_{t}w_{jk}\,dz\\ &+\int_{\Xi}2\delta(u_{j}^{p-1}+u_{k}^{p-1})\,|\partial_{t}\zeta|\,dz\\ &+\sum_{l=j,k}\int_{\Xi\cap\{Du_{l}\neq0\}}2\delta\,|D\zeta|\,|Du_{l}|^{p-1}+2\delta E_{\varepsilon}\zeta(|Du_{l}|^{p}\,h(|Du_{l}|)+|Du_{l}|^{\max(0,p-2)})\,dz. \end{split}$$

At the left-hand side of the above estimate we have

$$\int_{\Xi \cap \{|u_j^{p-1} - u_k^{p-1}| < \delta\}} \zeta(|Du_j|^{p-2} Du_j - |Du_k|^{p-2} Du_k) \cdot (u_j^{p-2} Du_j - u_k^{p-2} Du_k) dz$$

=
$$\int_{\Xi \cap \{|u_j^{p-1} - u_k^{p-1}| < \delta\}} \zeta u_j^{p-2} (|Du_j|^{p-2} Du_j - |Du_k|^{p-2} Du_k) \cdot (Du_j - Du_k) dz$$

$$- \int_{\Xi \cap \{|u_j^{p-1} - u_k^{p-1}| < \delta\}} \zeta (u_k^{p-2} - u_j^{p-2}) (|Du_j|^{p-2} Du_j - |Du_k|^{p-2} Du_k) \cdot Du_k dz.$$

Combining the last two displays we have

$$(p-1) \int_{\Xi \cap \{|u_{j}^{p-1}-u_{k}^{p-1}| < \delta\}} \zeta u_{j}^{p-2} (|Du_{j}|^{p-2} Du_{j} - |Du_{k}|^{p-2} Du_{k}) \cdot (Du_{j} - Du_{k}) dz$$

$$\leq \int_{\Xi} \zeta (u_{j}^{p-1} - u_{k}^{p-1}) \partial_{t} w_{jk} dz$$

$$+ \int_{\Xi} 2\delta (u_{j}^{p-1} + u_{k}^{p-1}) |\partial_{t}\zeta| dz$$

$$+ \sum_{l=j,k} \int_{\Xi \cap \{Du_{l} \neq 0\}} 2\delta |D\zeta| |Du_{l}|^{p-1} + 2\delta E_{\varepsilon} \zeta (|Du_{l}|^{p} h(|Du_{l}|) + |Du_{l}|^{\max(0,p-2)}) dz$$

$$+ (p-1) \int_{\Xi \cap \{ |u_j^{p-1} - u_k^{p-1}| < \delta \}} \zeta (u_k^{p-2} - u_j^{p-2}) (|Du_j|^{p-2} Du_j - |Du_k|^{p-2} Du_k) \cdot Du_k \, dz$$

=: $I_1 + I_2 + I_3 + I_4$, (2.20)

where the definition of I_1, \ldots, I_4 is again apparent. We now proceed to estimate the right-hand side.

Estimate of I_1 : Using the definition of w_{jk} and integrating by parts we get

$$I_{1} = \int_{\Xi \cap \left\{ \left| u_{j}^{p-1} - u_{k}^{p-1} \right| < \delta \right\}} \zeta(u_{j}^{p-1} - u_{k}^{p-1}) \partial_{t}(u_{j}^{p-1} - u_{k}^{p-1}) dz$$
$$= \int_{\Xi \cap \left\{ \left| u_{j}^{p-1} - u_{k}^{p-1} \right| < \delta \right\}} \frac{1}{2} \zeta \partial_{t}(u_{j}^{p-1} - u_{k}^{p-1})^{2} dz \leq \int_{\Xi} \frac{1}{2} w_{jk}^{2} \left| \partial_{t} \zeta \right| dz \leq \delta^{2} C(\zeta)$$

Estimate of I_2 : Since $|u_l| \leq M$ in Ξ for all $l \in \mathbb{N}$, we have

$$I_{2} = \int_{\Xi} 2\delta(|u_{j}|^{p-1} + |u_{k}|^{p-1}) |\partial_{t}\zeta| \ dz \le \delta C(p, M, \zeta).$$

Estimate of I_3 : Observe that by Lemma 2.6 we have

$$\sup_{l \in \mathbb{N}} \int_{\operatorname{spt} \zeta} |Du_l|^p \, dz \le C(p, h, M, \zeta).$$
(2.21)

Since h is a bounded function and $E_{\varepsilon} \in (0, 1]$, the estimate (2.21) together with Hölder's inequality readily implies

$$I_{3} = \sum_{l=j,k} \int_{\Xi \cap \{Du_{l} \neq 0\}} 2\delta |D\zeta| |Du_{l}|^{p-1} + 2\delta E_{\varepsilon} \zeta (|Du_{l}|^{p} h(|Du_{l}|) + |Du_{l}|^{\max(0,p-2)}) dz$$

$$\leq \delta C(p,h,M,\zeta).$$

Estimate of I_4 : Observe that since $0 < m \le u_l \le M$ in Ξ for all $l \in \mathbb{N}$, we have

$$\left|u_{k}^{p-2}-u_{j}^{p-2}\right| \leq C(p,m,M) \left|u_{j}^{p-1}-u_{k}^{p-1}\right|$$
 in Ξ ,

using the mean value theorem. Using this and (2.21), we estimate

$$I_{4} = (p-1) \int_{\Xi \cap \{ |u_{j}^{p-1} - u_{k}^{p-1}| < \delta \}} \zeta(u_{k}^{p-2} - u_{j}^{p-2}) (|Du_{j}|^{p-2} Du_{j} - |Du_{k}|^{p-2} Du_{k}) \cdot Du_{k} dz$$

$$\leq C(p, m, M) \delta \left(\left(\int_{\operatorname{spt} \zeta} |Du_{j}|^{p} dz \right)^{1/p} \left(\int_{\operatorname{spt} \zeta} |Du_{k}|^{p} dz \right)^{1/p} + \int_{\operatorname{spt} \zeta} |Du_{k}|^{p} dz \right)$$

$$\leq C(p, h, m, M, \zeta) \delta.$$

Combining the estimates of I_1, \ldots, I_4 with (2.20), we obtain

$$\int_{\Xi \cap \left\{ \left| u_j^{p-1} - u_k^{p-1} \right| < \delta \right\}} \zeta u_j^{p-2} (\left| Du_j \right|^{p-2} Du_j - \left| Du_k \right|^{p-2} Du_k) \cdot (Du_j - Du_k) \, dz \le \delta C_0, \quad (2.22)$$

18

where $C_0 = C_0(p, h, M, m, \zeta)$. We now estimate the left-hand side. If 1 , we use the vector inequality (see [Lin17, p98])

$$\left(\left|a\right|^{p-2}a - \left|b\right|^{p-2}b\right) \cdot (a-b) \ge (p-1)\left|a-b\right|^2 \left(1 + \left|a\right|^2 + \left|b\right|^2\right)^{\frac{p-2}{2}}$$

estimate (2.22) and Hölder's inequality to get that (recall $\zeta \equiv 1$ in U)

$$\int_{U \cap \{|u_{j}^{p-1}-u_{k}^{p-1}| < \delta\}} |Du_{j} - Du_{k}|^{r} dz
\leq \left(\int_{U \cap \{|u_{j}^{p-1}-u_{k}^{p-1}| < \delta\}} \left(1 + |Du_{j}|^{2} + |Du_{k}|^{2} \right)^{\frac{r(2-p)}{2(2-r)}} dz \right)^{\frac{2-r}{2}}
\cdot \left(\int_{\Xi \cap \{|u_{j}^{p-1}-u_{k}^{p-1}| < \delta\}} \frac{\zeta |Du_{j} - Du_{k}|^{2}}{\left(1 + |Du_{j}|^{2} + |Du_{k}|^{2} \right)^{\frac{2-p}{2}}} dz \right)^{\frac{r}{2}}
\leq C(p,h,r,m,M,\zeta) C_{0} \delta^{\frac{r}{2}},$$
(2.23)

where it was also used that r(2-p)/(2-r) < p. If $p \ge 2$, using the inequality (see [Lin17, p95])

$$(|a|^{p-2}a - |b|^{p-2}b) \cdot (a-b) \ge 2^{2-p}|a-b|^p$$

and (2.22) we get

$$\int_{U \cap \{|u_{j}^{p-1}-u_{k}^{p-1}| < \delta\}} |Du_{j}-Du_{k}|^{r} dz$$

$$\leq C |\operatorname{spt}\zeta|^{\frac{1-r}{p}} \left(\int_{U \cap \{|u_{j}^{p-1}-u_{k}^{p-1}| < \delta\}} \zeta |Du_{j}-Du_{k}|^{p} dz \right)^{\frac{r}{p}}$$

$$\leq C(p,r,m) \left(\int_{U \cap \{|u_{j}^{p-1}-u_{k}^{p-1}| < \delta\}} \zeta u_{j}^{p-2} |Du_{j}-Du_{k}|^{p} dz \right)^{\frac{r}{p}}$$

$$\leq C(p,r,m)C_{0}\delta^{\frac{r}{p}}.$$
(2.24)

Combining the estimates (2.23) and (2.24), we arrive at

$$\int_{U \cap \{ |u_j^{p-1} - u_k^{p-1}| < \delta \}} |Du_j - Du_k|^r \, dz \le C(p, h, r, m, M, \zeta) \delta^{\frac{r}{\max(p, 2)}}.$$

On the other hand, by Hölder's and Chebyshev's inequalities and estimate (2.21),

$$\int_{U \cap \{ |u_j^{p^{-1}} - u_k^{p^{-1}}| \ge \delta \}} |Du_j - Du_k|^r \, dz$$

$$\leq \left| U \cap \{ \left| u_j^{p^{-1}} - u_k^{p^{-1}} \right| \ge \delta \} \right|^{\frac{p-r}{p}} \left(\int_{U \cap \{ |u_j^{p^{-1}} - u_k^{p^{-1}}| \ge \delta \}} |Du_j - Du_k|^p \, dz \right)^{\frac{r}{p}}$$

$$\leq \left(\frac{1}{\delta} \int_U \left| u_j^{p^{-1}} - u_k^{p^{-1}} \right| \, dz \right)^{\frac{p-r}{p}} C(p, r, h, M, \zeta).$$

By the last two displays, we have now established that

$$\int_{U} |Du_j - Du_k|^r \, dz \le \left(\left(\frac{1}{\delta} \int_{U} \left| u_j^{p-1} - u_k^{p-1} \right| \, dz \right)^{\frac{p-r}{p}} + \delta^{\frac{r}{\max(p,2)}} \right) C(p,h,r,m,M,\zeta).$$

Taking first small δ and then large j and k, we see that the right-hand side can be made arbitrarily small. Thus (Du_l) is a Cauchy sequence in $L^r_{loc}(\Xi)$.

2.3. Passing to the limit. It remains to pass to the limit $\varepsilon \to 0$ to show that viscosity supersolutions are weak supersolutions.

Theorem 2.8. Let u be a continuous viscosity supersolution to (1.1) in Ξ . Suppose that $0 < m \le u \le M$. Then u is a local weak supersolution to (1.1) in Ξ .

Proof. First, we may suppose that u is uniformly continuous in Ξ by taking a smaller domain if necessary. Then, fix domains $U \Subset U' \Subset \Xi$. Let u_{ε} be the inf-convolution of u_{ε} and let $\varepsilon > 0$ be so small that $U' \Subset \Xi_{\varepsilon}$. By Lemma 2.5 the inf-convolution is a weak supersolution to

$$\partial_t u_{\varepsilon}^{p-1} - \Delta_p u_{\varepsilon} + G(Du_{\varepsilon}) \ge 0 \quad \text{in } U'.$$
(2.25)

Then by Lemma 2.6 we have Du_{ε} bounded in $L^p(U)$. Thus, passing to a subsequence if necessary, we have $Du_{\varepsilon} \to g$ weakly in $L^p(U)$ for some $g \in L^p(U)$. Then for any $i \in \{1, \ldots, N\}$ and $\varphi \in C_0^{\infty}(U)$ we have

$$\int_{U} u \partial_i \varphi \, dz = \lim_{\varepsilon \to 0} \int_{U} u_\varepsilon \partial_i \varphi \, dz = -\lim_{\varepsilon \to 0} \int_{U} \varphi \partial_i u_\varepsilon \, dz = -\int_{U} \varphi \partial_i u \, dz$$

so that u has a weak distributional space derivative $Du = g \in L^p(U)$.

Let $\varphi \in C_0^\infty(U)$. It now remains to pass to the limit in

$$\int_{U} -u_{\varepsilon}^{p-1} \partial_{t} \varphi + |Du_{\varepsilon}|^{p-2} Du_{\varepsilon} \cdot D\varphi \, dz + \int_{U \cap \{Du_{\varepsilon} \neq 0\}} E_{\varepsilon}(|Du_{\varepsilon}|^{p} h(|Du_{\varepsilon}|) + |Du_{\varepsilon}|^{\max(0, p-2)})\varphi \, dz.$$

Convergence of the first order terms is clear since $u_{\varepsilon} \to u$ pointwise, $E_{\varepsilon} \to 0$ and $|Du_{\varepsilon}|$ is bounded in $L^p(U)$. For the second-order term, one uses that $Du_{\varepsilon} \to Du$ in $L^r(U)$ for any 1 < r < p from Lemma 2.7 together with the vector inequality

$$\left| |a|^{p-2} a - |b|^{p-2} b \right| \le \begin{cases} 2^{2-p} |a-b|^{p-1}, & 1$$

which can be found on [Lin17, pp97-98]. This way, by (2.25), we arrive at

$$\int_{U} \left(-u^{p-1} \partial_t \varphi + |Du|^{p-2} Du \cdot D\varphi \right) \, dz \ge 0.$$

Theorem 2.9. Suppose that $0 < m \le u \le M$. Then a viscosity solution u to (1.1) is a weak solution and vice versa.

Proof. The theorem follows by the previous result and by recalling Theorem 3 in [LL22] where they show that weak solutions are viscosity solutions. In [LL22] they assume $p \ge 2$ but under our positivity assumptions this is not needed.

For the convenience of the reader, we provide the details in the case 1 . Suppose $on the contrary that there is <math>\varphi \in C^2$ that touches u from below at (x,t), $D\varphi(y,s) \neq 0$ for $y \neq x$, and

$$\limsup_{(s,s)\to(x,t),y\neq x} \left(\partial_t \varphi(y,s) - \frac{1}{u^{p-2}(x,t)} \Delta_p \varphi(y,s)\right) < -\varepsilon.$$

It follows that

(y

$$(p-1)\partial_t \varphi - \frac{1}{u^{p-2}(x,t)}\Delta_p \varphi < -\delta$$

in a neighborhood of (x,t). Since φ is C^2 , $\varphi(x,t) = u(x,t)$ and $0 < m \le u \le M$, taking r > 0 to be small enough we may ensure that

$$(p-1)\varphi^{2-p}\partial_t\varphi - \Delta_p\varphi < 0$$

in a cylindrical neighborhood $(B_r(x) \setminus \{x\}) \times (t-r, t+r)$. Multiplying this by non-negative test function $\phi \in C_0^{\infty}((B_r(x) \setminus \{x\}) \times (t-r, t+r))$ and integrating, we obtain for any $0 < \epsilon < r$ that

$$0 > \int_{t-r}^{t+r} \int_{B_r(x)\setminus B_\epsilon(x)} \left(\phi\partial_t \varphi^{p-1} - \phi\Delta_p \varphi\right) \, dy \, ds$$

= $\int_{t-r}^{t+r} \int_{B_r(x)\setminus B_\epsilon(x)} \left(-\varphi^{p-1}\partial_t \phi + |D\varphi|^{p-2} \, D\varphi \cdot D\phi\right) \, dy \, ds$
+ $\int_{\partial B_\epsilon(x)} \left(\varphi \, |D\phi|^{p-2} \, D\phi \cdot \frac{y-x}{\epsilon}\right) \, dS(y) \, ds.$

Letting $\epsilon \to 0$, this yields

$$\int_{B_r(x_0)\times(t-r,t+r)} \left(-\varphi^{p-1}\partial_t\phi + |D\varphi|^{p-2}\,D\varphi\cdot D\phi\right)\,dz < 0$$

so that φ is a weak subsolution in $B_r(x) \times (t-r, t+r)$. But then so is

$$\tilde{\varphi}(x,t) := h\varphi(y,s), \quad \text{where } h := \inf_{(y,s)\in\partial_{\mathcal{P}}B_r(x_0)\times(t-r,t+r)} \frac{u(y,s)}{\varphi(y,s)} > 1.$$

Now by comparison principle of weak solutions [LL22, Corollary 1], we have $\tilde{\varphi} \leq u$ in $B_r(x) \times (t - r, t + r)$, but this contradicts the fact that $\tilde{\varphi}(x, t) = h\varphi(x, t) = hu(x, t) > u(x, t)$.

As an application we also get that the limit of a positive, equicontinuous and bounded sequence of supersolutions (either viscosity or weak) is a supersolution. For the p-parabolic equation this is a special case of Theorem 5.3 in [KKP10], see also [LM07].

Corollary 2.10. Suppose that u_i , i = 1, 2, ... is a sequence of equicontinuous (weak or viscosity) supersolutions such that $0 < m \le u_i \le M$ and u_i converges to u almost everywhere. Then u is a (weak and viscosity) supersolution.

Proof. Let $\Xi' \in \Xi$ and u_i be a sequence of weak supersolutions satisfying the assumptions. By [LL22], each u_i is a viscosity supersolution to (1.1) in Ξ . Therefore, for small enough ε it follows from Lemma 2.5 that the inf-convolution $u_{i,\varepsilon}$ is a weak supersolution to (2.11) in Ξ' , where the function h that appears in the definition of G in (2.12) depends on the modulus of continuity of u_i . Since the functions u_i are equicontinuous, this h can be taken to be independent of i. As $u_{i,\varepsilon}$ are Lipschitz, the Caccioppoli's estimate in Lemma 2.6 implies the local boundedness of $Du_{i,\varepsilon}$ in $L^p_{loc}(\Xi')$. Taking the diagonal, that is, setting $v_i = u_{i,i^{-1}}$, we have by Lemma 2.7 that Dv_i converges for any 1 < r < p in $L^r_{loc}(\Xi')$ up to a subsequence. An argument similar to the proof of Theorem 2.9 then shows that the pointwise limit of v_i is a local weak supersolution in Ξ' and this limit is u, as by equicontinuity $u_{i,\varepsilon}$ converges locally uniformly in Ξ to u_i as $\varepsilon \to 0$ with a speed independent of i.

3. LOCAL LIPSCHITZ CONTINUITY IN SPACE

We prove local Lipschitz continuity of positive viscosity solutions to (1.1) in the space variable using the Ishii-Lions method. While this method can sometimes yield explicit Lipschitz estimates, our result here is only qualitative. This is because in the proof, the solution is essentially frozen, and therefore the Hölder and Lipschitz estimates produced by the method will depend on the modulus of continuity of the solution.

Let $u: \Xi \to \mathbb{R}$, where $\Xi \subset \mathbb{R}^{N+1}$ is a set. We say that

$$\omega(s) := \sup_{(x,t),(y,t)\in\Xi, |x-y|\leq s} |u(x,t) - u(y,t)|$$

is the optimal modulus of continuity of u in Ξ in space variable. We carefully observe that we do not assume Lipschitz continuity at this point, and for brevity we drop the subindex x from the notation of ω . We also denote

$$Q_r := B_r(0) \times (-r, 0] \text{ for } r > 0$$

and its parabolic boundary by $\partial_{\mathcal{P}}Q_r := (\overline{B}_r(0) \times \{-r\}) \cup (\partial B_r(0) \times (-r, 0]).$

Theorem 3.1 (Lipschitz continuity). Let u be a uniformly continuous viscosity solution to (1.1) in $B_2 \times (-2, 0)$. Suppose that $0 < m \le u \le M$. Then there exists L > 0 such that

$$|u(x_0, t_0) - u(y_0, t_0)| \le L |x_0 - y_0|$$
 for all $(x_0, t_0), (y_0, t_0) \in Q_{1/2}$

The constant L depends only on N, p, m and the optimal modulus of continuity of u in $B_2 \times (-2, 0)$.

Using the previous equivalence result Theorem 2.9, we immediately have the Lipschitz result for weak solutions.

Corollary 3.2 (Lipschitz continuity for weak solutions). Suppose that $0 < m \le u \le M$. If u is a weak (or viscosity solutions) to (1.1) in $B_2 \times (-2,0)$, then u is locally Lipschitz continuous.

Since the equation (1.1) is parabolic, it is natural to establish the Lipschitz continuity of solution u in $B_2 \times (-2, 0)$ up to the top of the cylinder $Q_{1/2}$. For this, we need the following lemma that extends the solution property to the set $B_2 \times \{0\}$. For equations that are proper in the language of [CIL92], a proof of such a result can be found in [DFO14, p209]. However, (1.1) is not proper and therefore we rely on the equivalence result and the comparison principle of weak solutions to prove the lemma. The technique resembles the one in [Juu01].

For $(x,t) \in Q_r$, we define the second-order subjet relative to Q_r by

$$\mathcal{P}_{Q_r}^{2,-}u(x,t) := \left\{ (\theta,\eta,X) : u(y,s) \le u(x,t) + (s-t)\theta + (x-y) \cdot \eta + \frac{1}{2}(x-y)'X(x-y) + o(|x-y|^2 + |s-t|) \text{ as } (y,s) \to (x,t), (y,s) \in Q_r \right\}$$

and (θ, η, X) is in $\overline{\mathcal{P}}_{Q_r}^{2,-}u(x,t)$ if there exists $(\theta_i, \eta_i, X_i) \in \mathcal{P}_{Q_r}^{2,-}u(x_i, t_i)$ such that $(x_i, t_i, \theta_i, \eta_i, X_i)$ converges to (x, t, θ, η, X) . The second-order superjet relative to Q_r is denoted by $\mathcal{P}_{Q_r}^{2,+}u(x,t)$ and defined in the analogous way. Finally, note that these subjets coincide with the usual definition within the interior of Q_r .

Lemma 3.3. Let u be a uniformly continuous viscosity supersolution to (1.1) in $B_1 \times (-1,0)$ such that u > m > 0. Suppose that $(\theta, \eta, X) \in \overline{\mathcal{P}}_{Q_1}^{2,-}u(x,0), \eta \neq 0$, where $u(x,0) = \lim_{(y,s)\to(x,0),(y,s)\in Q} u(y,s)$. Then

$$(p-1)\theta - u^{2-p}(x,0)(\operatorname{tr} X + (p-2)\frac{\eta' X\eta}{|\eta|^2}) \ge 0.$$

The analogous result holds for viscosity subsolutions.

Proof. The idea is that if the claim does not hold, then we can construct a C^2 function φ that touches u from below at the top of the cylinder, and φ has to be a strict subsolution near the point of touching. By lifting this function a little, we can get a contradiction since u satisfies the comparison principle.

To be more precise, by uniform continuity of u, it is enough to consider $(\theta, \eta, X) \in \mathcal{P}_{Q_1}^{2,-}u(x,0)$ instead of the closure. Thriving for a contradiction, suppose that there exists $(\theta, \eta, X) \in \mathcal{P}_{Q_1}^{2,-}u(x,0), \eta \neq 0$, such that

$$(p-1)\theta - u^{2-p}(x,0)(\operatorname{tr} X + (p-2)\frac{\eta' X\eta}{|\eta|^2}) < -\varepsilon$$

for some $\varepsilon > 0$. Then there is $\varphi \in C^2(Q_1)$ such that $(\partial_t \varphi(x,0), D\varphi(x,0), D^2 \varphi(x,0)) = (\theta, \eta, X), \varphi(x,0) = u(x,0), \varphi(y,s) < u(y,s)$ when $(y,s) \in Q_1 \setminus \{(x,0)\}$, and

$$(p-1)\partial_t\varphi(x,0) - \varphi^{2-p}(x,0)\Delta_p\varphi(x,0) < -\varepsilon.$$

Since φ is C^2 , it follows from above that there exists $\delta > 0$ such that

$$\partial_t \varphi^{p-1}(y,s) - \Delta_p \varphi(y,s) < -\varepsilon/2 \quad \text{in } B_\delta$$

and also $\varphi > 0$, $D\varphi \neq 0$ in Q_{δ} . From this it follows that φ is, in particular, a weak subsolution to (1.1) in $B_{\delta} \times (-\delta, 0)$. Then so is

$$\tilde{\varphi}(y,s) := h\varphi(y,s), \text{ where } h := \inf_{(y,s)\in\partial_{\mathcal{P}}Q_{\delta}} \frac{u(y,s)}{\varphi(y,s)} > 1.$$

On the other hand, by Theorem 2.8 u is a local weak supersolution to (1.1) in $B_{\delta} \times (-\delta, 0)$. Since by construction of $\tilde{\varphi}$ we have $\tilde{\varphi} \leq u$ on $\partial_{\mathcal{P}}Q_{\delta}$, it follows from the comparison principle [LL22, Corollary 1] and continuity that $\tilde{\varphi} \leq u$ in Q_{δ} , which leads to a contradiction since $\tilde{\varphi}(x,0) = h\varphi(x,0) = hu(x,0) > u(x,0)$.

To prove Theorem 3.1, we first prove the following technical lemma, which we will apply twice: first to obtain Hölder continuity, and then Lipschitz continuity. This lemma encapsulates the use of the theorem on sums, and thus makes the full proof cleaner, since then we avoid repeating that part in both the Hölder and and Lipschitz proofs.

In the following discussion, we drop the localization terms for expository reasons similar to [AMP]. The idea in the Ishii-Lions method when aiming at

$$u(x,t) - u(y,t) \le L\varphi(|x-y|),$$

where φ reflects the desired regularity (Hölder or Lipschitz as detailed below in (3.1)), is thriving for a contradiction to assume that there is a maximum point such that

$$u(x_0, t_0) - u(y_0, t_0) - L\varphi(|x_0 - y_0|) > c > 0.$$

Then using the maximum point property, we have information about the derivatives and second derivatives, which we can use to contradict the opposite information coming from the equation in the same way as in the usual uniqueness machinery for viscosity solutions.

The proof needs to be done in two stages since in the Hölder proof at the maximum point, we can utilize the larger negative curvature of φ , in other words larger negativite impact of the second derivative to a suitable direction, as done in (3.17) and in the computation after the estimate. In the Lipscitz proof the second derivative at the maximum point is not as negative but there (see (3.20)) the positive (bad) term depends on the modulus of continuity of the solution, and this is sufficiently small by the previous Hölder result.

Since the core of the argument is the same in both the stages it is stated separately in the following lemma. It might be instructive to observe that in the proofs of Hölder and Lipschitz continuity, the lemma will be utilized respectively with the functions

$$\varphi(s) := s^{\alpha} \quad \text{and} \quad \varphi(s) := s - \kappa s^{\beta},$$
(3.1)

where α, κ and β are suitable constants.

Lemma 3.4. Suppose that u is a uniformly continuous viscosity solution to (1.1) in $B_1 \times (-1,0)$ and $0 < m \le u \le M$. Let $(x_0,t_0), (y_0,t_0) \in Q_{1/2}$ and define the function

$$\Psi(x,y,t) := u(x,t) - u(y,t) - L\varphi(|x-y|) - \frac{K}{2}|x-x_0|^2 - \frac{K}{2}|y-y_0|^2 - \frac{K}{2}|t-t_0|^2, \quad (3.2)$$

where $K := 8 \operatorname{osc}_{Q_1} u$, and $\varphi : [0, 2] \to [0, \infty)$ is a C^2 -function such that

$$\varphi(0) = 0, \quad |\varphi''(s)| < \frac{\varphi'(s)}{s} \quad and \quad \varphi'' < 0 < c_{\varphi} < \varphi' \quad for \ some \ c_{\varphi} > 0.$$
 (3.3)

Then, if L > L' for some L' that depends only on c_{φ} and $\operatorname{osc}_{Q_1} u$, the following holds: If Ψ has a positive maximum at $(\hat{x}, \hat{y}, \hat{t}) \in \overline{B}_1 \times \overline{B}_1 \times [-1, 0]$, we have

$$-K \le (L\varphi'(|z|))^{p-2} \left(C^{-2}L\varphi''(|z|) + L\frac{\omega(|z|)\varphi'(|z|)}{|z|} + \sqrt{K}\frac{\omega^{1/2}(|z|)}{|z|} + K \right),$$

where $|z| = |\hat{x} - \hat{y}|$ and ω is the optimal modulus of continuity of u in $B_1 \times (-1,0)$ in space variable. Here C depends only on N, p, m and M. Finally, we have the estimate

$$\varphi'(|z|) \le \frac{\omega(|z|)}{L|z|}.$$
(3.4)

Proof. First, we must have $|z| \neq 0$ since otherwise the maximum at $(\hat{x}, \hat{y}, \hat{t})$ would be non-positive. Secondly, we have

$$0 < \left| u(\hat{x}, \hat{t}) - u(\hat{y}, \hat{t}) \right| - L\varphi(|\hat{x} - \hat{y}|) - \frac{K}{2} \left| \hat{x} - x_0 \right|^2 - \frac{K}{2} \left| \hat{y} - y_0 \right|^2 - \frac{K}{2} \left| \hat{t} - t_0 \right|^2$$
(3.5)

so that

$$\left| \hat{t} - t_0 \right|, \left| \hat{x} - x_0 \right|, \left| \hat{y} - y_0 \right| \le \sqrt{\frac{2}{K} \left| u(\hat{x}, \hat{t}) - u(\hat{y}, \hat{t}) \right|} \le \sqrt{\frac{1}{4}} = \frac{1}{2}.$$
(3.6)

Since $x_0, y_0 \in B_{1/2}$ and $t_0 \in (-1/2, 0]$, this implies $\hat{x}, \hat{y} \in B_1$ and $\hat{t} \in (-1, 0]$, which means that the maximum point $(\hat{x}, \hat{y}, \hat{t})$ is in $B_1 \times B_1 \times (-1, 0]$.

By the definition of ω and the uniform continuity of u, we have

$$|u(x,s) - u(y,s)| \le \omega(|x-y|) \quad \text{for all } (x,s), (y,s) \in B_1 \times (0,1],$$
(3.7)

Therefore, (3.6) implies that

$$K\left|\hat{t} - t_{0}\right|, K\left|\hat{x} - x_{0}\right|, K\left|\hat{y} - y_{0}\right| \le \sqrt{K}\omega^{1/2}(\left|\hat{x} - \hat{y}\right|).$$
(3.8)

We also have by concavity of φ and (3.5) that $|z| \varphi'(|z|) \leq \int_0^{|z|} \varphi'(s) ds \leq \frac{\omega(|z|)}{L}$ so that (3.4) holds.

Since $\hat{x} \neq \hat{y}$, the function $\phi(x, y) \mapsto \varphi(|x - y|)$ is C^2 in a neighborhood of (\hat{x}, \hat{y}) . Therefore, we may invoke the parabolic theorem on sums [DFO14, Theorem 9] (see also [CIL92, Theorem 8.3]). For any $\mu > 0$, there exist matrices $X, Y \in S^N$ and $b_1, b_2 \in \mathbb{R}$ such that

$$b_1 + b_2 \ge \partial_t (L\varphi(|x - y|))(\hat{x}, \hat{y}) = 0$$

and

$$(b_1, D_x(L\varphi(|x-y|))(\hat{x}, \hat{y}), X) \in \overline{\mathcal{P}}_{Q_1}^{2,+}(u - \frac{K}{2}|x-x_0|^2 - \frac{K}{2}|t-t_0|^2)(\hat{x}, \hat{t}),$$

$$(-b_2, D_y(L\varphi(|x-y|))(\hat{x}, \hat{y}), Y) \in \overline{\mathcal{P}}_{Q_1}^{2,-}(u + \frac{K}{2}|y-y_0|^2)(\hat{x}, \hat{y}).$$

Denoting $z := \hat{x} - \hat{y}$ and

$$\eta_{1} := L\varphi'(|z|)\frac{z}{|z|} + K(\hat{x} - x_{0}),$$

$$\eta_{2} := L\varphi'(|z|)\frac{z}{|z|} - K(\hat{y} - y_{0}),$$

$$\theta_{1} := b_{1} + K(\hat{t} - t_{0}),$$

$$\theta_{2} := -b_{2},$$

(3.9)

these can be written as

$$(\theta_1, \eta_1, X + KI) \in \overline{\mathcal{P}}_{Q_1}^{2,+} u(\hat{x}, \hat{t}), \quad (\theta_2, \eta_2, Y - KI) \in \overline{\mathcal{P}}_{Q_1}^{2,-} u(\hat{y}, \hat{t}).$$

Assuming L is large enough depending on \sqrt{K} , $\operatorname{osc}_{Q_1} u$ and c_{φ} , and using (3.8), (3.9), we have

$$\begin{aligned} |\eta_1|, |\eta_2| &\leq L\varphi'(|z|) + \sqrt{K}\omega^{1/2}(|\hat{x} - \hat{y}|) \leq L\varphi'(|z|) + Lc_{\varphi} \leq 2L\varphi'(|z|) \\ |\eta_1|, |\eta_2| &\geq \frac{1}{2}L\varphi'(|z|) + \frac{1}{2}Lc_{\varphi} - \sqrt{K}\omega^{1/2}(|\hat{x} - \hat{y}|) \geq \frac{1}{2}L\varphi'(|z|) \geq 1, \end{aligned}$$
(3.10)

Furthermore, by the theorem on sums, we have

$$-(\mu + 4 \|B\|)I \le \begin{pmatrix} X & 0\\ 0 & -Y \end{pmatrix} \le \begin{pmatrix} B & -B\\ -B & B \end{pmatrix} + \frac{2}{\mu} \begin{pmatrix} B^2 & -B^2\\ -B^2 & B^2 \end{pmatrix},$$
(3.11)

where

$$B = L\varphi''(|z|)\frac{z\otimes z}{|z|^2} + L\frac{\varphi'(|z|)}{|z|}\left(I - \frac{z\otimes z}{|z|^2}\right),$$

$$B^2 = BB = L^2 \left|\varphi''(|z|)\right|^2 \frac{z\otimes z}{|z|^2} + L^2 \frac{|\varphi'(|z|)|^2}{|z|^2}\left(I - \frac{z\otimes z}{|z|^2}\right).$$

By assumptions on φ , we have

$$|\varphi''(|z|)| \le \varphi'(|z|)/|z|$$
 and $\varphi''(|z|) < 0 < \varphi'(|z|).$

Thus we deduce that

$$||B|| \le \frac{L\varphi'(|z|)}{|z|}$$
 and $||B^2|| \le L^2 \frac{(\varphi'(|z|))^2}{|z|^2}.$ (3.12)

Moreover, we choose

$$\mu := \frac{2L\varphi'(|z|)}{|z|},$$

so that using the explicit formulae of B and $B^2,$ and the fact that the terms with φ' vanish, we obtain

$$\left\langle B\frac{z}{|z|}, \frac{z}{|z|} \right\rangle + \frac{2}{\mu} \left\langle B^2 \frac{z}{|z|}, \frac{z}{|z|} \right\rangle \le \frac{L}{2} \varphi''(|z|). \tag{3.13}$$

Denote below

$$F((z,s),\zeta,Z) := u^{2-p}(z,s) |\zeta|^{p-2} (\operatorname{tr} Z + (p-2) \frac{\zeta' Z \zeta}{|\zeta|^2}).$$

Since u is a viscosity solution to (1.1), we have (using also Lemma 3.3 if $\hat{t} = 0$)

$$\theta_1 \leq F((\hat{x}, \hat{t}), \eta_1, X + KI)$$
 and $\theta_2 \geq F((\hat{y}, \hat{t}), \eta_2, Y - KI).$

We denote $\tilde{\eta} := L\varphi'(|z|)z/|z|$. Then we subtract the equations and use that $\theta_1 - \theta_2 = b_1 + b_2 + K(\hat{t} - t_0) \ge K(\hat{t} - t_0)$ as well as add and subtract terms to obtain

$$\begin{split} K(\hat{t}-t_0) &\leq F((\hat{x},\hat{t}),\eta_1,X+KI) - F((\hat{y},\hat{t}),\eta_2,Y-KI) \\ &= F((\hat{x},\hat{t}),\eta_1,X) - F((\hat{y},\hat{t}),\eta_2,Y) \\ &+ F((\hat{x},\hat{t}),\eta_1,X+KI) - F((\hat{x},\hat{t}),\eta_1,X) - F((\hat{y},\hat{t}),\eta_2,Y-KI) + F((\hat{y},\hat{t}),\eta_2,Y) \\ &= F((\hat{x},\hat{t}),\tilde{\eta},X) - F((\hat{y},\hat{t}),\tilde{\eta},Y) \\ &+ F((\hat{x},\hat{t}),\eta_1,X) - F((\hat{x},\hat{t}),\tilde{\eta},X) + F((\hat{y},\hat{t}),\tilde{\eta},Y) - F((\hat{y},\hat{t}),\eta_2,Y) \\ &+ F((\hat{x},\hat{t}),\eta_1,X+KI) - F((\hat{x},\hat{t}),\eta_1,X) - F((\hat{y},\hat{t}),\eta_2,Y-KI) + F((\hat{y},\hat{t}),\eta_2,Y) \\ &=: T_1 + T_2 + T_3. \end{split}$$

Estimate of T_1 : By the definition of $\tilde{\eta}$, the matrix inequality (3.11), and estimate (3.13), we have

$$\begin{split} \frac{\tilde{\eta}'}{|\tilde{\eta}|}(X-Y)\frac{\tilde{\eta}}{|\tilde{\eta}|} &= \frac{z'}{|z|}(X-Y)\frac{z}{|z|} = \binom{z/|z|}{-z/|z|}' \begin{pmatrix} X & 0\\ 0 & -Y \end{pmatrix} \begin{pmatrix} z/|z|\\ -z/|z| \end{pmatrix} \\ &\leq \binom{z/|z|}{-z/|z|}' \left(\begin{pmatrix} B & -B\\ -B & B \end{pmatrix} + \frac{2}{\mu} \begin{pmatrix} B^2 & -B^2\\ -B^2 & B^2 \end{pmatrix} \right) \begin{pmatrix} z/|z|\\ -z/|z| \end{pmatrix} \\ &= 2\left\langle B\frac{z}{|z|}, \frac{z}{|z|} \right\rangle + \frac{4}{\mu} \left\langle B^2\frac{z}{|z|}, \frac{z}{|z|} \right\rangle \\ &\leq L\varphi''(|z|). \end{split}$$

Furthermore, (3.11) readily implies $X - Y \leq 0$ by multiplying (3.11) by (ξ, ξ) left and right. In particular, this implies $tr(X - Y) \leq 0$. Moreover, (3.11) also gives that

$$||X||, ||Y|| \le ||B|| + \frac{2}{\mu} ||B||^2 \le C \frac{L\varphi'(|z|)}{|z|}.$$
(3.14)

Using the previous two estimates and that $\varphi'' < 0$, it follows that

$$\begin{split} T_1 &= F((\hat{x}, \hat{t}), \tilde{\eta}, X) - F((\hat{y}, \hat{t}), \tilde{\eta}, Y) \\ &= u(\hat{x}, \hat{t})^{2-p} \left| \tilde{\eta} \right|^{p-2} \left(\operatorname{tr} X + (p-2) \frac{\tilde{\eta}' X \tilde{\eta}}{\left| \tilde{\eta} \right|^2} \right) - u(\hat{y}, \hat{t})^{2-p} \left| \tilde{\eta} \right|^{p-2} \left(\operatorname{tr} Y + (p-2) \frac{\tilde{\eta}' Y \tilde{\eta}}{\left| \tilde{\eta} \right|^2} \right) \\ &= \left(u^{2-p}(\hat{x}, \hat{t}) - u^{2-p}(\hat{y}, \hat{t}) \right) \left| \tilde{\eta} \right|^{p-2} \left(\operatorname{tr} X + (p-2) \frac{\tilde{\eta}' X \tilde{\eta}}{\left| \tilde{\eta} \right|^2} \right) \\ &+ u(\hat{y}, \hat{t})^{2-p} \left| \tilde{\eta} \right|^{p-2} \left(\operatorname{tr} (X - Y) + (p-2) \frac{\tilde{\eta}' (X - Y) \tilde{\eta}}{\left| \tilde{\eta} \right|^2} \right) \\ &\leq C(N, p, m, M) \left| u(\hat{x}, \hat{t}) - u(\hat{y}, \hat{t}) \right| \left| \tilde{\eta} \right|^{p-2} \| X \| + C(p) u(\hat{y}, \hat{t})^{2-p} \left| \tilde{\eta} \right|^{p-2} \frac{\tilde{\eta}' (X - Y) \tilde{\eta}}{\left| \tilde{\eta} \right|^2} \\ &\leq C(N, p, m, M) \omega(|z|) \left| \tilde{\eta} \right|^{p-2} L \frac{\varphi'(|z|)}{|z|} + C^{-1}(N, p, m, M) L \left| \tilde{\eta} \right|^{p-2} \varphi''(|z|), \end{split}$$

for some constant $C(N, p, m, M) \ge 1$.

Estimate of T_2 : First we show the following algebraic inequality

$$\left| |\xi_1|^{p-2} \left(\operatorname{tr} Z + (p-2) |\xi_1|^{-2} \xi_1' Z \xi_1 \right) - |\xi_2|^{p-2} \left(\operatorname{tr} Z + (p-2) |\xi_2|^{-2} \xi_2' Z \xi_2 \right) \right|$$

$$\leq C(N,p) \max(|\xi_1|^{p-3}, |\xi_2|^{p-3}) |\xi_1 - \xi_2| ||Z||$$

$$(3.15)$$

for all $\xi_1, \xi_2 \in \mathbb{R}^N \setminus \{0\}$ and $Z \in S(N)$. To this end, we observe that by [The75]

$$\begin{aligned} \left| \xi_1' Z \xi_1 - \xi_2 Z \xi_2 \right| &= \left| \operatorname{tr} ((\xi_1 \otimes \xi_1 - \xi_2 \otimes \xi_2) Z) \right| \\ &\leq N \left\| \xi_1 \otimes \xi_1 - \xi_2 \otimes \xi_2 \right\| \left\| Z \right\| \\ &= N \left\| (\xi_1 - \xi_2) \otimes \xi_1 - \xi_2 \otimes (\xi_2 - \xi_1) \right\| \left\| Z \right\| \\ &\leq N (|\xi_1| + |\xi_2|) \left\| \xi_1 - \xi_2 \right\| \left\| Z \right\|. \end{aligned}$$

We also use the elementary inequality

$$|a^{q} - b^{q}| \le |q| \max(a^{q-1}, b^{q-1}) |a - b|, \quad a, b > 0 \text{ and } q \in \mathbb{R}$$

Furthermore, may assume that $|\xi_2| \leq |\xi_1|$ by changing notation, if necessary. Using these, we estimate

$$\begin{aligned} \left| |\xi_{1}|^{p-2} - |\xi_{2}|^{p-2} \right| |\operatorname{tr} Z| + (p-2) |\xi_{1}|^{p-4} \left| \xi_{1}' Z \xi_{1} - \xi_{2}' Z \xi_{2} \right| + (p-2) \left| |\xi_{1}|^{p-4} - |\xi_{2}|^{p-4} \right| \left| \xi_{2}' Z \xi_{2} \right| \\ &\leq N |p-2| \max(|\xi_{1}|^{p-3}, |\xi_{2}|^{p-3}) |\xi_{1} - \xi_{2}| ||Z|| + N |p-2| |\xi_{1}|^{p-4} (|\xi_{1}| + |\xi_{2}|) ||\xi_{1} - \xi_{2}| ||Z|| \\ &+ |p-2| |p-4| \max(|\xi_{1}|^{p-5}, |\xi_{2}|^{p-5}) |\xi_{1} - \xi_{2}| ||\xi_{2}|^{2} ||Z|| \\ &\leq C(N, p) |p-2| \max(|\xi_{1}|^{p-3}, |\xi_{2}|^{p-3}) |\xi_{1} - \xi_{2}| ||Z|| , \end{aligned}$$

which proves the algebraic inequality (3.15). Observe now that

$$|\eta_1 - \tilde{\eta}| + |\eta_2 - \tilde{\eta}| = K(|x_0 - \hat{x}| + |y_0 - \hat{y}|) \le 2\sqrt{K}\omega^{1/2}(|z|).$$
(3.16)

Therefore, using (3.15), we obtain

$$\begin{split} T_2 &= F((\hat{x}, \hat{t}), \eta_1, X) - F((\hat{x}, \hat{t}), \tilde{\eta}, X) + F((\hat{y}, \hat{t}), \tilde{\eta}, Y) - F((\hat{y}, \hat{t}), \eta_2, Y) \\ &= u(\hat{x}, \hat{t})^{2-p} |\eta_1|^{p-2} \mathrm{tr}(X - (p-2) \frac{\eta_1' X \eta_1}{|\eta_1|^2}) - u(\hat{x}, \hat{t})^{2-p} |\tilde{\eta}|^{p-2} \mathrm{tr}(X - (p-2) \frac{\tilde{\eta}' X \tilde{\eta}}{|\tilde{\eta}|^2}) \\ &+ u(\hat{y}, \hat{t})^{2-p} |\tilde{\eta}|^{p-2} \mathrm{tr}(Y - (p-2) \frac{\tilde{\eta}' Y \tilde{\eta}}{|\tilde{\eta}|^2}) - u(\hat{y}, \hat{t})^{2-p} |\eta_2|^{p-2} \mathrm{tr}(Y - (p-2) \frac{\eta_2' Y \eta_2}{|\eta_2|^2}) \\ &\leq C(N, p, m, M) \max(|\eta_1|^{p-3}, |\eta_2|^{p-3}, |\tilde{\eta}|^{p-3}) (||X|| + ||Y||) (|\eta_1 - \tilde{\eta}| + |\eta_2 - \tilde{\eta}|) \\ &\leq C(N, p, m, M) |\tilde{\eta}|^{p-3} \sqrt{K} \omega^{1/2} (|z|) L \frac{\varphi'(|z|)}{|z|} \end{split}$$

where in the last estimate we used (3.14), (3.16) and that (3.10) implies $|\eta_1|, |\eta_2| \in [\frac{1}{2} |\tilde{\eta}|, 2 |\tilde{\eta}|]$.

Estimate of T_3 : Since F is quasilinear, we have

$$T_{3} = F((\hat{x}, \hat{t}), \eta_{1}, X + KI) - F((\hat{x}, \hat{t}), \eta_{1}, X) - F((\hat{y}, \hat{t}), \eta_{2}, Y - KI) + F((\hat{y}, \hat{t}), \eta_{2}, Y)$$

= $F((\hat{x}, \hat{t}), \eta_{1}, KI) + F((\hat{y}, \hat{t}), \eta_{2}, KI)$
 $\leq C(N, p)K(|\eta_{1}|^{p-2} + |\eta_{2}|^{p-2})$
 $\leq C(N, p)K |\tilde{\eta}|^{p-2},$

where we also used (3.10) and the definition of $\tilde{\eta}$.

Combining the estimates of T_1, \ldots, T_3 and using (3.10) we have for some $C(N, p, m, M) \ge 1$ that

$$-K$$

$$\leq C^{-1}L \,|\tilde{\eta}|^{p-2} \,\varphi''(|z|) + C\omega(|z|) \,|\tilde{\eta}|^{p-2} \,L \frac{\varphi'(|z|)}{|z|} + |\tilde{\eta}|^{p-3} \,\sqrt{K}\omega^{1/2}(|z|) L \frac{\varphi'(|z|)}{|z|} + CK \,|\tilde{\eta}|^{p-2} \\ = C(L\varphi'(|z|))^{p-2} \left(C^{-2}L\varphi''(|z|) + L \frac{\omega(|z|)\varphi'(|z|)}{|z|} + \sqrt{K} \frac{\omega^{1/2}(|z|)}{|z|} + K \right),$$

where we also used the definition of $\tilde{\eta}$. This yields the desired inequality.

The next lemma gives Hölder continuity with an explicit power.

Lemma 3.5 (Hölder continuity in space). Let u be a uniformly continuous viscosity solution to (1.1) in $B_2 \times (-2, 0)$. Suppose that $0 < m \le u \le M$. Then there exist constants $C_H > 0$ and $\alpha \in (0, 1)$ such that

$$|u(x_0,t_0) - u(y_0,t_0)| \le C_H |x_0 - y_0|^{\alpha}$$
 for all $(x_0,t_0), (y_0,t_0) \in Q_1$.

Moreover, $\alpha = \frac{1}{4}\min(p-2,0) + \frac{1}{2}$ and C_H depends only on N, p, m and the optimal modulus of continuity of u in $B_2 \times (-2,0)$.

Proof. Following the notation in Lemma 3.4, we fix $K := 8 \operatorname{osc}_{Q_1} u$. We also fix $(x_0, t_0), (y_0, t_0) \in Q_{1/2}$ and consider the auxiliary function

$$\Psi(x,y,t) := u(x,t) - u(y,t) - C_H \varphi(|x-y|) - \frac{K}{2} |x-x_0|^2 - \frac{K}{2} |y-y_0|^2 - \frac{K}{2} |t-t_0|^2$$

for some large $C_H > 0$ to be selected later. We define $\varphi : [0,2] \to [0,\infty)$ by $\varphi(s) := s^{\alpha}$, where $\alpha := \frac{1}{4} \min(p-2,0) + \frac{1}{2}$. Since $\alpha \in (0,1)$, we have

$$\varphi''(s) = \alpha(\alpha - 1)s^{\alpha - 2} < 0 < \alpha 2^{\alpha - 1} < \varphi'(s) \text{ and } |\varphi''(s)| \le \frac{\varphi'(s)}{s}$$

so that condition (3.3) in Lemma 3.4 holds. It now suffices to prove that Ψ is non-positive, so suppose on the contrary that Ψ has a positive maximum at $(\hat{x}, \hat{y}, \hat{t}) \in \overline{B}_1 \times \overline{B}_1 \times [-1, 1]$. Provided that $C_H \geq C(\alpha, \operatorname{osc}_{Q_1} u)$, which we now assume, we may apply Lemma 3.4. There exists C_0 , depending only on N, p, m, and M, such that

$$-K$$

$$\leq (C_H \varphi'(|z|))^{p-2} \left(C_0^{-2} C_H \varphi''(|z|) + C_H \omega(|z|) \frac{\varphi'(|z|)}{|z|} + \sqrt{K} \frac{\omega^{1/2}(|z|)}{|z|} + K \right)$$
(3.17)
$$= (\varphi'(|z|) C_H)^{p-2} \left(-C_0^{-2} C_H \alpha |\alpha - 1| |z|^{\alpha - 2} + \alpha C_H \omega(|z|) |z|^{\alpha - 2} + \sqrt{K} \frac{\omega^{1/2}(|z|)}{|z|} + K \right),$$

where ω is the optimal modulus of continuity of u in Q_1 in space variable and $z = \hat{x} - \hat{y}$ for some $(\hat{x}, \hat{t}), (\hat{y}, \hat{t}) \in B_1 \times (-1, 0)$. By the estimate (3.4) we have

$$\omega(|z|) \le \omega \left(\frac{\omega(|z|)}{\varphi'(|z|)C_H}\right) \le \omega \left(\frac{\operatorname{osc}_Q u}{\alpha 2^{\alpha-1}C_H}\right).$$

Then by assuming that C_H is even larger, depending on $\operatorname{osc}_{Q_1} u$, α , C_0 , and ω , we may ensure that

$$\omega(|z|) \le C_0^{-2} \frac{|\alpha - 1|}{2}.$$

Therefore, continuing the estimate, we have

$$-K$$

$$\leq (\varphi'(|z|)C_{H})^{p-2} \left(-C_{0}^{-2}C_{H}\alpha |\alpha - 1| |z|^{\alpha - 2} + C_{0}^{-2}\alpha |\alpha - 1| C_{H} |z|^{\alpha - 2} + \sqrt{K} \frac{\omega^{1/2}(|z|)}{|z|} + K \right)$$

$$= (\varphi'(|z|)C_{H})^{p-2} \left(-\frac{C_{0}^{-2}}{2}C_{H}\alpha |\alpha - 1| |z|^{\alpha - 2} + \sqrt{K} \frac{(\operatorname{osc}_{Q_{1}} u)^{1/2}}{|z|} + K \right)$$

$$\leq (\varphi'(|z|)C_{H})^{p-2} |z|^{\alpha - 2} \left(-\frac{C_{0}^{-2}\alpha |\alpha - 1|}{2}C_{H} + C\sqrt{K} (\operatorname{osc}_{Q_{1}} u)^{1/2} + CK \right),$$

where in the last estimate we used that $|z|^{-1} \leq C |z|^{\alpha-2}$ since |z| < 2 and $\alpha - 2 \in (-2, -1)$. Then we take C_H so large that $C_H \geq C \frac{4C_0^2}{\alpha(\alpha-1)} (\sqrt{K}(\operatorname{osc}_Q u)^{1/2} + K)$ so that by the above display (using also (3.4) to estimate $|z|^{\alpha-2}$), we have

$$-K \leq -(\varphi'(|z|)C_{H})^{p-2} |z|^{\alpha-2} \frac{C_{0}^{-2} \alpha |\alpha-1|}{4} C_{H}$$

$$\leq -\frac{C_{0}^{-2} \alpha |\alpha-1|}{4} (\varphi'(|z|)C_{H})^{p-2} \frac{\omega^{\alpha-2}(|z|)}{C_{H}^{\alpha-2} \varphi'(|z|)^{\alpha-2}} C_{H}$$

$$\leq -\frac{C_{0}^{-2} \alpha |\alpha-1|}{4} (\varphi'(|z|)C_{H})^{p-\alpha} (\operatorname{osc} u)^{\alpha-2} C_{H}, \qquad (3.18)$$

where $p - \alpha > 0$. Therefore, since $\varphi'(|z|) > \alpha 2^{\alpha-1}$, taking large enough C_H in (3.18) yields a contradiction. Thus Ψ is non-positive, which by the definition of φ yields

$$u(x_0, t_0) - u(y_0, t_0) \le C_H \varphi(|x_0 - y_0|) \le C_H |x_0 - y_0|^{\alpha}$$

Since x_0, y_0 and t_0 were arbitrary, u is Hölder continuous in space in $Q_{1/2}$. Repeating the argument, we see that u is Hölder continuous in Q_1 .

We can now finish the proof of Theorem 3.1.

Proof of Theorem 3.1. By Lemma 3.5, the function u is Hölder continuous in particular in Q_1 . Therefore, denoting by ω the optimal modulus of continuity of u in Q_1 in space variable, we have

$$|u(x,t) - u(y,t)| \le \omega(|x-y|) \le C_H |x-y|^{\alpha} \quad \text{for all } (x,t), (y,t) \in Q_1,$$
(3.19)

where $\alpha = \frac{1}{4} \min(p-2,0) + \frac{1}{2}$ and C_H depends only on N, p, m, and the optimal modulus of continuity of u in $B_2 \times (-2,0)$.

We define $\varphi : [0,2] \to [0,\infty)$ by $\varphi(s) := s - \kappa s^{\beta}$, where $\beta := \frac{\alpha}{2} + 1$ and $\kappa := \beta^{-1} 2^{-\beta-1}$. Then

$$\varphi'(s) = 1 - 2^{-\beta - 1} s^{\beta - 1}$$
 and $\varphi''(s) = -2^{-\beta - 1} (\beta - 1) s^{\beta - 2}$

so that, since $\beta \in (1,2)$, we have for $s \in (0,2)$

$$\begin{aligned} \left|\varphi''(s)\right| &- \frac{\left|\varphi'(s)\right|}{s} = 2^{-\beta-1}(\beta-1)s^{\beta-2} - \frac{1-2^{-\beta-1}s^{\beta-1}}{s} \\ &= 2^{-\beta-1}\beta s^{\beta-2} - s^{-1} \\ &= s^{-1}(2^{-\beta-1}\beta s^{\beta-1} - 1) \\ &\leq s^{-1}(2^{-2}\beta - 1) < 0 \end{aligned}$$

and thus the requirement (3.3) of Lemma 3.4 holds. We fix $K := 8 \operatorname{osc}_{Q_1} u$ and $(x_0, t_0), (y_0, t_0) \in B_{1/2} \times (-1/2, 1/2)$ and define

$$\Psi(x,y,t) := u(x,t) - u(y,t) - L\varphi(|x-y|) - \frac{K}{2}|x-x_0|^2 - \frac{K}{2}|y-y_0|^2 - \frac{K}{2}|t-t_0|^2.$$

It now suffices to prove that Ψ is non-positive. Suppose on the contrary that Ψ has a positive maximum at $(\hat{x}, \hat{y}, \hat{t}) \in \overline{B}_1 \times \overline{B}_1 \times [-1, 1]$. Then, assuming that $L \ge C(\alpha, \operatorname{osc}_Q u)$

it follows from Lemma 3.4 that

$$-K \le (L\varphi'(|z|))^{p-2} \left(C_0^{-2} L\varphi''(|z|) + L \frac{\omega(|z|)\varphi'(|z|)}{|z|} + \sqrt{K} \frac{\omega^{1/2}(|z|)}{|z|} + K \right)$$
(3.20)

for some C_0 that depends only on N, p, m and M, and $z = \hat{x} - \hat{y}$, for some $(\hat{x}, \hat{t}), (\hat{y}, \hat{t}) \in Q_1$. We estimate the right-hand side using (3.19), definition of β , formula of φ'' and the fact that $\varphi' \in [3/4, 1]$. We obtain

$$-K \leq CL^{p-2}(-\kappa\beta(\beta-1)C_0^{-2}L|z|^{\beta-2} + C_0C_HL|z|^{\alpha-1} + \sqrt{K}C_H^{1/2}|z|^{\frac{\alpha}{2}-1} + K)$$

= $CL^{p-2}(-\kappa\beta(\beta-1)C_0^{-2}L|z|^{\frac{\alpha}{2}-1} + C_0C_HL|z|^{\alpha-1} + \sqrt{K}C_H^{1/2}|z|^{\frac{\alpha}{2}-1} + K)$
= $CL^{p-2}|z|^{\frac{\alpha}{2}-1}(-C(\alpha)C_0^{-2}L + C_0C_HL|z|^{\frac{\alpha}{2}} + \sqrt{K}C_H^{1/2} + K),$ (3.21)

where we also used that $\kappa\beta(\beta-1) = C(\alpha)$. By (3.4) we have

$$|z|^{\frac{\alpha}{2}} \le \left(\frac{\omega(|z|)}{\varphi'(|z|)L}\right)^{\frac{\alpha}{2}} \le C(\alpha, \underset{Q_1}{\operatorname{osc}} u)L^{-\frac{\alpha}{2}}$$

and thus continuing the estimate (3.21) we get

$$-K \le CL^{p-2} |z|^{\frac{\alpha}{2}-1} \left(-C(\alpha)C_0^{-2}L + C_0C_HC(\alpha, \underset{Q_1}{\operatorname{osc}} u)L^{1-\frac{\alpha}{2}} + \sqrt{K}C_H^{1/2} + K\right).$$

Observe that the negative term inside the parentheses has the highest power of L. Therefore, assuming L to be large enough depending on α , C_0 , C_H , K and $\operatorname{osc}_{Q_1} u$, we can absorb the positive terms to the negative term. This way, we arrive at

$$-K \leq -C(\alpha, C_0) |z|^{\frac{\alpha}{2}-1} L^{p-1}$$

Since $|z|^{\frac{\alpha}{2}-1} > C$, a contradiction follows by taking even larger L if necessary.

Therefore, Ψ is non-positive, which by the definition of φ means that

$$u(x_0, t_0) - u(y_0, t_0) \le L\varphi(|x_0 - y_0|) \le L |x_0 - y_0|$$

Since x_0, y_0 and t_0 were arbitrary, this proves the Lipschitz continuity of u in $Q_{1/2}$.

APPENDIX A. BASIC PROPERTIES

For the benefit of the reader, we prove the basic properties of inf-convolutionin Lemma 2.1 even if they are rather standard. For the convenience of the reader we repeat the lemma here.

Lemma. Assume that $u : \Xi \to \mathbb{R}$ is lower semicontinuous and bounded. Suppose also that $\delta_{\varepsilon} \to 0$ as $\varepsilon \to 0$. Then u_{ε} has the following properties.

- (i) We have $u_{\varepsilon} \leq u$ in Ξ and $u_{\varepsilon} \to u$ pointwise as $\varepsilon \to 0$.
- (ii) Denote $r(\varepsilon) := (q\varepsilon^{q-1} \operatorname{osc}_{\Xi} u)^{\frac{1}{q}}, t(\varepsilon) := (2\delta_{\varepsilon} \operatorname{osc}_{\Xi} u)^{\frac{1}{2}}$ and set

$$\Xi_{\varepsilon} := \left\{ (x,t) \in \Xi : B_{r(\varepsilon)}(x) \times (t - t(\varepsilon), t + t(\varepsilon)) \Subset \Xi \right\}.$$

Then, for any $(x,t) \in \Xi_{\varepsilon}$ there exists $(x_{\varepsilon},t_{\varepsilon}) \in \overline{B}_{r(\varepsilon)}(x) \times [t-t(\varepsilon),t+t(\varepsilon)]$ such that

$$u_{\varepsilon}(x,t) = u(x_{\varepsilon},t_{\varepsilon}) + \frac{|x-x_{\varepsilon}|^{q}}{q\varepsilon^{q-1}} + \frac{|t-t_{\varepsilon}|^{2}}{2\delta_{\varepsilon}}.$$

- (iii) The function u_{ε} is semi-concave in Ξ_{ε} . In particular, the function $u_{\varepsilon}(x,t) (C|x|^2 + t^2/\delta_{\varepsilon})$ is concave in Ξ_{ε} , where $C = (q-1)r(\varepsilon)^{q-2}/\varepsilon^{q-1}$.
- (iv) Suppose that u_{ε} is twice differentiable in space and time at $(x,t) \in \Xi_{\varepsilon}$. Then

$$\partial_t u_{\varepsilon}(x,t) = \frac{t - t_{\varepsilon}}{\delta_{\varepsilon}},$$

$$Du_{\varepsilon}(x,t) = (x - x_{\varepsilon}) \frac{|x - x_{\varepsilon}|^{q-2}}{\varepsilon^{q-1}},$$

$$D^2 u_{\varepsilon}(x,t) \le (q-1) \frac{|x - x_{\varepsilon}|^{q-2}}{\varepsilon^{q-1}} I.$$

Proof of Lemma 2.1. We consider the different properties given in the lemma separately.

Proof of (ii). Let $(x,t) \in \Xi_{\varepsilon}$. Let $(y,s) \in \Xi$ be such that $y \in \mathbb{R}^N \setminus \overline{B}_{r(\varepsilon)}(x)$ or $s \in \mathbb{R}^N \setminus [t - t(\varepsilon), t + t(\varepsilon)]$. Then by the definition of $r(\varepsilon)$ and $t(\varepsilon)$, we have that

$$\frac{|y-x|^q}{q\varepsilon^{q-1}} + \frac{|s-t|^2}{2\delta_{\varepsilon}} > \underset{\Xi}{\operatorname{osc}} u$$

so that

$$u(y,s) + \frac{|y-x|^q}{q\varepsilon^{q-1}} + \frac{|s-t|^2}{2\delta_{\varepsilon}} > u(y,s) + u(x,t) - u(x,t) + \underset{\Xi}{\operatorname{osc}} u \ge u(x,t) \ge u_{\varepsilon}(x,t).$$

Hence by lower semi-continuity

$$\begin{aligned} u_{\varepsilon}(x,t) &= \inf_{(y,s)\in\overline{B}_{r(\varepsilon)}(x)\times[t-t(\varepsilon),t+t(\varepsilon)]} \left\{ u(y,s) + \frac{|y-x|^{q}}{q\varepsilon^{q-1}} + \frac{|s-t|^{2}}{2\delta_{\varepsilon}} \right\} \\ &= u(x_{\varepsilon},t_{\varepsilon}) + \frac{|x-x_{\varepsilon}|^{q}}{q\varepsilon^{q-1}} + \frac{|t-t_{\varepsilon}|^{2}}{2\delta_{\varepsilon}} \end{aligned}$$

for some $(x_{\varepsilon}, t_{\varepsilon}) \in \overline{B}_{r(\varepsilon)} \times [t - t(\varepsilon), t + t(\varepsilon)].$

Proof of (i). Let $(x,t) \in \Xi$ and let $\varepsilon > 0$ be so small that $(x,t) \in \Xi_{\varepsilon}$. From (ii) and boundedness of u, it is clear that $(x_{\varepsilon}, t_{\varepsilon}) \to (x, t)$. Therefore, by lower semi-continuity

$$\limsup_{\varepsilon \to 0} \frac{|x - x_{\varepsilon}|^{q}}{q^{\varepsilon^{q-1}}} + \frac{|t - t_{\varepsilon}|^{2}}{2\delta_{\varepsilon}} = \limsup_{\varepsilon \to 0} (u_{\varepsilon}(x, t) - u(x_{\varepsilon}, t_{\varepsilon})) \le \limsup_{\varepsilon \to 0} (u(x, t) - u(x_{\varepsilon}, t_{\varepsilon})) \le 0$$

and so $u_{\varepsilon}(x,t) \to u(x,t)$.

Proof of (iii). For all $(y,s) \in \Xi$, define the function $\varphi_{(y,s)} : \mathbb{R}^{N+1} \to \mathbb{R}$ by

$$\varphi_{(y,s)}(x,t) := u(y,s) + \frac{|y-x|^q}{q\varepsilon^{q-1}} + \frac{|s-t|^2}{2\delta_{\varepsilon}}.$$

Observe that for any h > 0, whenever $|y - x| \le r(\varepsilon) + h$, we have that

$$D^{2}\varphi_{(y,s)}(x,t) \leq \frac{(q-1)|y-x|^{q-2}}{\varepsilon^{q-1}} \leq \frac{(q-1)(r(\varepsilon)+h)^{q-2}}{\varepsilon^{q-1}},$$
$$\partial_{tt}\varphi_{(y,s)}(x,t) \leq \frac{1}{\delta_{\varepsilon}}.$$

It follows that for $C_h := (q-1)(r(\varepsilon)+h)^{q-2}/\varepsilon^{q-1}$ the function $(x,t) \mapsto \varphi_{(y,s)}(x,t) - (C_h |x|^2 + t^2/\delta_{\varepsilon})$ is concave in the set $\{(x,t) \in \mathbb{R}^{N+1} : |x-y| \le r(\varepsilon) + h\}$. Now, if $(z,\tau) \in \Xi_{\varepsilon}$, then for small h > 0 we have $B_h(z) \times (\tau - h, \tau + h) \subset \Xi_{\varepsilon}$. Therefore, by property *(ii)* and the definition of inf-convolution, we have for all $(x,t) \in B_h(z) \times (\tau - h, \tau + h)$ that

$$u_{\varepsilon}(x,t) - (C_{h}|x|^{2} + t^{2}/\delta_{\varepsilon}) = \inf_{\substack{(y,s)\in\overline{B}_{r(\varepsilon)}(x)\times[t-t(\varepsilon),t+t(\varepsilon)]}} \left\{\varphi_{(y,s)}(x,t)\right\} - C_{h}(|x|^{2} + |t|^{2})$$
$$= \inf_{\substack{(y,s)\in\Xi\cap\overline{B}_{r(\varepsilon)+h}(z)\times[\tau-t(\varepsilon)-h,\tau+t(\varepsilon)+h]}} \left\{\varphi_{(y,s)}(x,t)\right\} - C_{h}(|x|^{2} + |t|^{2})$$
$$= \inf_{\substack{(y,s)\in\Xi\cap\overline{B}_{r(\varepsilon)+h}(z)\times[\tau-t(\varepsilon)-h,\tau+t(\varepsilon)+h]}} \left\{\varphi_{(y,s)}(x,t) - C_{h}(|x|^{2} + |t|^{2})\right\}.$$

It follows that as the infimum of concave functions $u_{\varepsilon}(x,t) - (C_h |x|^2 + t^2/\delta_{\varepsilon})$ is concave in $B_h(z) \times (\tau - h, \tau + h)$, and thus in Ξ_{ε} , since $(z,\tau) \in \Xi_{\varepsilon}$ was arbitrary. Taking $h \to 0$, we see that also $u_{\varepsilon}(x,t) - (C_0 |x|^2 + t^2/\delta_{\varepsilon})$ is concave.

Proof of (iv). If u_{ε} is twice differentiable at $(x,t) \in \Xi_{\varepsilon}$, then there is a C^2 -function φ that touches u_{ε} from below at (x,t) and $\partial_t \varphi(x,t) = \partial_t u_{\varepsilon}(x,t)$, $D\varphi(x,t) = Du_{\varepsilon}(x,t)$ as well as $D^2\varphi(x,t) = D^2u_{\varepsilon}(x,t)$. By (ii) and the definition of inf-convolution, we have

$$\varphi(x,t) = u_{\varepsilon}(x,t) = u(x_{\varepsilon},t_{\varepsilon}) + \frac{|x-x_{\varepsilon}|^{q}}{q\varepsilon^{q-1}} + \frac{|t-t_{\varepsilon}|^{2}}{2\delta_{\varepsilon}},$$

$$\varphi(z,\tau) \le u_{\varepsilon}(z,\tau) \le u(y,s) + \frac{|y-z|^{q}}{q\varepsilon^{q-1}} + \frac{|\tau-s|^{2}}{2\delta_{\varepsilon}} \quad \text{for all } (z,\tau), (y,s) \in \Xi.$$

But this means that the function $\psi(z,\tau) := \varphi(z,\tau) - \frac{|z-x_{\varepsilon}|^{q}}{q^{\varepsilon^{q-1}}} - \frac{|\tau-t_{\varepsilon}|^{2}}{2\delta_{\varepsilon}}$ has a maximum at $(z,\tau) = (x,t)$. Indeed, the first display above implies that $\psi(x,t) = u(x_{\varepsilon},t_{\varepsilon})$, and selecting $(y,s) = (x_{\varepsilon},t_{\varepsilon})$ in the second one we get that $\psi(z,\tau) \leq u(x_{\varepsilon},t_{\varepsilon})$ for all $(z,\tau) \in \Xi$. The claim then follows by direct computation.

Next we restate and prove Lemma 2.3 for the convenience of the reader.

Lemma (Elementary inequality). Let p > 1. Suppose that $a \ge m > 0$. Then for all $s \in (0, 1)$ we have

$$\left| (a+s)^{\frac{2-p}{2}} - a^{\frac{2-p}{2}} \right| \le C(p,m)s.$$

Proof of Lemma 2.3. Clearly

$$\left| (a+s)^{\frac{2-p}{2}} - a^{\frac{2-p}{2}} \right| = \left| \int_{a}^{a+s} \frac{d}{dh} h^{\frac{2-p}{2}} dh \right| = \left| \int_{a}^{a+s} \frac{p-2}{2} h^{-\frac{p-2}{2}-1} dh \right|$$
$$\leq \frac{|p-2|}{2} \max(a^{-\frac{p-2}{2}-1}, (a+s)^{-\frac{p-2}{2}-1})s$$
$$= C(p,m)s.$$

References

- [AMP] R. Anttila, J.J. Manfredi, and M. Parviainen. Connections between coupling and Ishii-Lions methods for tug-of-war with noise stochastic games. *arXiv preprint arXiv:2501.13897*.
- [BDG⁺23] V. Bögelein, F. Duzaar, U. Gianazza, N. Liao, and C. Scheven. Hölder continuity of the gradient of solutions to doubly non-linear parabolic equations. arXiv preprint arXiv:2305.08539, 2023.
- [BDL21] V. Bögelein, F. Duzaar, and N. Liao. On the hölder regularity of signed solutions to a doubly nonlinear equation. *Journal of Functional Analysis*, 281(9):109173, 2021.
- [BDLS22] V. Bögelein, F. Duzaar, N. Liao, and L. Schätzler. On the Hölder regularity of signed solutions to a doubly nonlinear equation. part ii. *Revista Matemática Iberoamericana*, 39(3):1005–1037, 2022.
- [BM15] T. Bhattacharya and L. Marazzi. On the viscosity solutions to Trudinger's equation. Nonlinear Differential Equations and Applications NoDEA, 22:1089–1114, 2015.
- [BM16] T. Bhattacharya and L. Marazzi. Asymptotics of viscosity solutions to some doubly nonlinear parabolic equations. *Journal of Evolution Equations*, 16(4):759–788, 2016.
- [CIL92] M. G. Crandall, H. Ishii, and P.-L. Lions. User's guide to viscosity solutions of second order partial differential equations. Bull. Amer. Math. Soc., 27(1):1–67, 1992.
- [Cra97] M. G. Crandall. Viscosity solutions: a primer. In Viscosity solutions and applications, pages 1–43. Springer, 1997.
- [DF014] J. Diehl, P. K. Friz, and H. Oberhauser. Regularity theory for rough partial differential equations and parabolic comparison revisited. In D. Crisan, B. Hambly, and T. Zariphopoulou, editors, *Stochastic Analysis and Applications 2014*, pages 203–238. Springer International Publishing, 2014.
- [DU20] N. M. L. Diehl and J. M. Urbano. Sharp Hölder regularity for the inhomogeneous Trudinger's equation. Nonlinearity, 33(12):7054, 2020.
- [EG92] L. C. Evans and R. F. Gariepy. Measure theory and fine properties of functions. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1992.
- [GV06] U. Gianazza and V. Vespri. A Harnack inequality for solutions of doubly nonlinear parabolic equations. J. Appl. Funct. Anal., 1:271–284, 2006.
- [Hen20] E. Henriques. Local Hölder regularity for a doubly singular PDE. Commun. Contemp. Math., 22(3):1850054, 30, 2020.
- [HL19] R. Hynd and E. Lindgren. Lipschitz regularity for a homogeneous doubly nonlinear pde. SIAM Journal on Mathematical Analysis, 51(4):3606–3624, 2019.
- [HL21] R. Hynd and E. Lindgren. Large time behavior of solutions of Trudinger's equation. Journal of Differential Equations, 274:188–230, 2021.
- [IL90] H. Ishii and P.-L. Lions. Viscosity solutions of fully nonlinear second-order elliptic partial differential equations. J. Differential Equations, 83(1):26–78, 1990.
- [Ish95] H. Ishii. On the equivalence of two notions of weak solutions, viscosity solutions and distribution solutions. *Funkcial. Ekvac.*, 38(1):101–120, 1995.
- [JJ12] V. Julin and P. Juutinen. A new proof for the equivalence of weak and viscosity solutions for the p-Laplace equation. Comm. Partial Differential Equations, 37(5):934–946, 2012.

| [JLM01] | P. Juutinen, P. Lindqvist, and J. J. Manfredi. On the equivalence of viscosity solutions and weak solutions for a quasi-linear equation. <i>SIAM J. Math. Anal.</i> , 33(3):699–717, 2001. |
|----------|---|
| [Juu01] | P. Juutinen. On the definition of viscosity solutions for parabolic equations. Proc. Amer. Math. Soc., 129(10):2907–2911, 2001. |
| [KK07] | J. Kinnunen and T. Kuusi. Local behaviour of solutions to doubly nonlinear parabolic equations. <i>Math. Ann.</i> , 337(3):705–728, 2007. |
| [KKP10] | R. Korte, T. Kuusi, and M. Parviainen. A connection between a general class of superparabolic functions and supersolutions. J. Evol. Eq., 10(1):1–20, 2010. |
| [KLSU12] | T. Kuusi, R. Laleoglu, J. Siljander, and J. M. Urbano. Hölder continuity for Trudinger's equation in measure spaces. <i>Calculus of Variations and Partial Differential Equations</i> , 45:193–229, 2012. |
| [KSU12] | T. Kuusi, J. Siljander, and J. M. Urbano. Local Hölder continuity for doubly nonlinear parabolic equations. <i>Indiana University Mathematics Journal</i> , pages 399–430, 2012. |
| [Lin12] | P. Lindqvist. Regularity of supersolutions. In <i>Regularity estimates for nonlinear elliptic and parabolic problems</i> , volume 2045 of <i>Lecture Notes in Math.</i> , pages 73–131. Springer, Heidelberg, 2012. |
| [Lin17] | P. Lindqvist. Notes on the <i>p</i> -Laplace equation (second edition). Univ. Jyväskylä, Report 161, 2017. |
| [LL22] | E. Lindgren and P. Lindqvist. On a comparison principle for Trudinger's equation. <i>Adv. Calc. Var.</i> , 15(3):401–415, 2022. |
| [LM07] | P. Lindqvist and J. J. Manfredi. Viscosity supersolutions of the evolutionary <i>p</i> -Laplace equation. <i>Differential Integral Equations</i> , 20(11):1303–1319, 2007. |
| [LS22] | N. Liao and L. Schätzler. On the Hölder regularity of signed solutions to a doubly nonlinear equation. part iii. <i>International Mathematics Research Notices</i> , 2022(3):2376–2400, 2022. |
| [Mis23] | M. Misawa. Expansion of positivity for doubly nonlinear parabolic equations and its application. <i>Calculus of Variations and Partial Differential Equations</i> , 62(9):265, 2023. |
| [PV20] | M. Parviainen and J. L. Vázquez. Equivalence between radial solutions of different parabolic gradient-diffusion equations and applications. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 21:303–359, 2020. |
| [Sil18] | J. Siltakoski. Equivalence of viscosity and weak solutions for the normalized $p(x)$ -Laplacian. Calc. Var. Partial Differential Equations, 57(95), 2018. |
| [Sil21] | J. Siltakoski. Equivalence of viscosity and weak solutions for a <i>p</i> -parabolic equation. J. Evol. Equ., 21(2):2047–2080, 2021. |
| [The75] | C. M. Theobald. An inequality for the trace of the product of two symmetric matrices. <i>Math. Proc. Philos. Soc.</i> , 77(2):265-267, 1975. |
| [Tru68] | N. S. Trudinger. Pointwise estimates and quasilinear parabolic equations. Comm. Pure Appl. Math., 21:205–226, 1968. |
| Depart | ment of Mathematical Sciences, Norwegian University of Science and Technology |
| NO-7491 | TRONDHEIM, NORWAY |

Email address: lqvist@math.ntnu.no

Department of Mathematics and Statistics, University of Jyväskylä, PO Box 35, 40014 Jyväskylä, Finland

Email address: mikko.j.parviainen@jyu.fi

Department of Mathematics and Statistics, University of Helsinki, P.O. Box 68, FI-00014 Helsinki, Finland

Email address: jarkko.siltakoski@helsinki.fi

36

LINDQVIST, PARVIAINEN, AND SILTAKOSKI