## Nonadiabatic quantum geometry and optical conductivity

Raffaele Resta<sup>1, 2, \*</sup>

<sup>1</sup>CNR-IOM Istituto Officina dei Materiali, Strada Costiera 11, 34149 Trieste, Italy <sup>2</sup>Donostia International Physics Center, 20018 San Sebastián, Spain (Dated: run through LATEX on 21-Feb-25 at 1:56)

The ground-state quantum geometry is at the root of several static and adiabatic properties, while genuinely dynamic properties are routinely addressed via Kubo formulæ, whose essential entries are the excited states. It is shown here that the ground-state metric-curvature tensor evolves in time by means of a causal unitary operator, which by construction elucidates the geometrical effect of the excited states in compact form. In the condensed-matter case the generalized tensor encompasses the whole conductivity tensor at arbitrary frequencies in both insulators and metals, with the exception of the Drude term in the metallic case; the latter is shown to be eminently nongeometrical.

Quantum geometry made its appearance in electronicstructure theory the early 1980s, in the form of an entity nowadays called Berry curvature [1, 2]. Since then, the quantum geometry of the ground state constitutes a conceptual foundation and a computational framework for several—apparently disparate—static and adiabatic properties in condensed-matter [3–6] and in molecules [7]. It has been challenging, instead, to relate quantum geometry to genuinely nonadiabatic properties, which in a Kubo formulation involve excited states as well. Some pioneering results in this direction appeared recently, all of them in a band-structure framework addressing the mean-field crystalline case [8–11]. I adopt instead the many-body formulation of quantum geometry [2, 4], which allows for compact and very transparent notations (besides addressing in principle even systems with disorder and correlation). The geometrical nature of a dynamical response property is perspicuous in the time domain, where the metric-curvature tensor evolves by means of a unitary causal operator, Eq. (5) below, which accounts for the effect of the excited states in a meaningful form. The geometrical tensor so defined allows retrieving the full conductivity tensor—real and imaginary parts, longitudinal and transverse terms—in both insulators and metals, with one exception in the metallic case: the Drude term. The latter is shown to be in fact the nongeometrical term in the adiabatic response to a dc field.

Let us consider a system of N interacting d-dimensional electrons in a cubic box of volume  $L^d$ , and the family of many-body Hamiltonians parametrized by  $\kappa$ , called "flux" or "twist":

$$\hat{H}_{\kappa} = \frac{1}{2m} \sum_{i=1}^{N} \left( \mathbf{p}_i + \frac{e}{c} \mathbf{A}(\mathbf{r}_i) + \hbar \kappa \right)^2 + \hat{V}, \qquad (1)$$

where  $\hat{V}$  includes the one-body potential (possibly disordered) and electron-electron interaction; the vector potential summarizes all intrinsic time-reversal breaking terms, as e.g. those due to a coupling to a background of local moments. We assume the system to be macroscopically homogeneous; the eigenstates are normalized to one

in the hypercube of volume  $L^{Nd}$ . The thermodynamic limit  $N \to \infty$ ,  $L \to \infty$ ,  $N/L^d$  constant is understood throughout this work.

We impose Born-von-Kàrmàn periodic boundary conditions: the many-body wavefunctions are periodic with period L over each electron coordinate  $\mathbf{r}_i$  independently; the potential  $\hat{V}$  and the vector potential  $\mathbf{A}(\mathbf{r})$  enjoy the same periodicity. If the eigenvalue problem is instead solved for the same Hamiltonian by adopting standard  $\mathcal{L}^2$  boundary conditions, one may address some geometrical properties other than conductivity in molecules [7]. The flux  $\kappa$ —cast into inverse-length dimensions for convenience—corresponds to perturbing the Hamiltonian with a vector potential  $\hbar c \kappa / e$ , constant in space; if it is constant even in time, then its occurrence is a pure gauge.

The symbols  $\hat{H}$ ,  $|\Psi_n\rangle$  and  $E_n$  will refer to  $\kappa=0$  quantities; the velocity operator at  $\kappa=0$  is

$$\hat{\mathbf{v}} = \frac{1}{m} \sum_{i=1}^{N} \left[ \mathbf{p}_i + \frac{e}{c} \mathbf{A}(\mathbf{r}_i) \right] = \frac{1}{\hbar} \partial_{\kappa} \hat{H}.$$
 (2)

The Berry curvature at  $\kappa = 0$  is

$$\Omega_{\alpha\beta} = -2 \operatorname{Im} \langle \partial_{\kappa_{\alpha}} \Psi_0 | \partial_{\kappa_{\beta}} \Psi_0 \rangle, \tag{3}$$

where Greek subscripts indicate Cartesian coordinates; sum over repeated subscripts is implicitly understood in the following. The intrinsic anomalous Hall conductivity is

$$\sigma_{\alpha\beta}^{(-)}(0) = -\frac{e^2}{\hbar L^d} \Omega_{\alpha\beta}; \tag{4}$$

the expression holds for metals and insulators, in either 2d or 3d. The equivalence of Eqs. (3) and (4) to the corresponding Kubo formula can be straightforwardly proved [12].

For d=2 and in the insulating case Eq. (4) is topologically quantized, although in the  $L\to\infty$  limit only. The single- $\kappa$  expression  $2\pi\Omega_{xy}/L^2$  converges in fact to the Chern number [13], while instead the  $\kappa$ -averaged value is quantized at any L [2, 4].

The Berry curvature is naturally endowed with a symmetric part and with a time dependence by defining

$$F_{\alpha\beta}(t) = \langle \partial_{\kappa_{\alpha}} \Psi_{0} | \tilde{U}(t) | \partial_{\kappa_{\beta}} \Psi_{0} \rangle,$$
  
$$\tilde{U}(t) = \theta(t) \sum_{n} |\Psi_{n}\rangle e^{-i\omega_{0n}t} \langle \Psi_{n} |.$$
 (5)

where  $\omega_{0n} = (E_n - E_0)/\hbar$ . This expression is not gauge-invariant; therefore it is mandatory to evaluate the  $\kappa$ -derivatives in the parallel-transport gauge, thus yielding

$$F_{\alpha\beta}(t) = \theta(t) \sum_{n \neq 0} \frac{\langle \Psi_0 | \hat{v}_\alpha | \Psi_n \rangle \langle \Psi_n | \hat{v}_\beta | \Psi_0 \rangle}{\omega_{0n}^2} e^{-i\omega_{0n}t}. \quad (6)$$

By construction, the ground-state Berry curvature is retrieved as

$$\Omega_{\alpha\beta} = -2 \operatorname{Im} F_{\alpha\beta}(0^+), \tag{7}$$

while Re  $F_{\alpha\beta}(0^+)$  is the quantum metric evaluated in the parallel-transport gauge (i.e. setting  $\langle \Psi_0 | \partial_{\kappa} \Psi_0 \rangle = 0$ ). The metric is divergent in metals [14, 15], and therefore only the insulating case is addressed for the time being; metals will be considered again below. In the present approach, both the metric and the curvature evolve causally by means of the unitary operator  $\tilde{U}(t)$ .

The Fourier transform of Eq. (6) is

$$\mathcal{F}_{\alpha\beta}(\omega) = \sum_{n \neq 0} \frac{\langle \Psi_0 | \hat{v}_\alpha | \Psi_n \rangle \langle \Psi_n | \hat{v}_\beta | \Psi_0 \rangle}{\omega_{0n}^2} \frac{i}{\omega - \omega_{0n} + i\eta},$$
(8)

where  $\eta \to 0^+$ . In order to compare  $\mathcal{F}_{\alpha\beta}(\omega)$  to the Kubo formulæ for optical conductivity it is expedient to adopt the notation

$$\langle \Psi_0 | \hat{v}_{\alpha} | \Psi_n \rangle \langle \Psi_n | \hat{v}_{\beta} | \Psi_0 \rangle = \mathcal{R}_{n,\alpha\beta} + i \mathcal{I}_{n,\alpha\beta}, \tag{9}$$

where the real term  $\mathcal{R}_{n,\alpha\beta}$  is symmetric and the imaginary one  $\mathcal{I}_{n,\alpha\beta}$  is antisymmetric.

The symmetric and antisymmetric parts of the Fourier transform of  $F_{\alpha\beta}(t)$  are:

$$\mathcal{F}_{\alpha\beta}^{(+)}(\omega) = \sum_{n \neq 0} \frac{\mathcal{R}_{n,\alpha\beta}}{\omega_{0n}^2} \left[ \pi \delta(\omega - \omega_{0n}) + \frac{i}{\omega - \omega_{0n}} \right] (10)$$

$$\mathcal{F}_{\alpha\beta}^{(-)}(\omega) = \sum_{n \neq 0} \frac{\mathcal{I}_{n,\alpha\beta}}{\omega_{0n}^2} \left[ i\pi \delta(\omega - \omega_{0n}) - \frac{1}{\omega - \omega_{0n}} \right] (11)$$

Two of the above expressions are immediately related to standard Kubo formulæ for  $\omega > 0$  [12]:

Re 
$$\sigma_{\alpha\beta}^{(+)}(\omega) = \frac{e^2}{\hbar L^d} \omega \operatorname{Re} \mathcal{F}_{\alpha\beta}^{(+)}(\omega);$$
 (12)

Im 
$$\sigma_{\alpha\beta}^{(-)}(\omega) = \frac{e^2}{\hbar L d} \omega \operatorname{Im} \mathcal{F}_{\alpha\beta}^{(-)}(\omega).$$
 (13)

The independent-electron version of Eqs. (10) and (12) was previously arrived at by following a very different logical path [8–11].

The remaining Kubo formulæ are also obtained in terms of  $\mathcal{F}_{\alpha\beta}(\omega)$  as  $\omega$ -integrals via the Kramers-Kronig relationships. The real part of the transverse conductivity is thus

Re 
$$\sigma_{\alpha\beta}^{(-)}(\omega) = \frac{e^2}{\hbar L^d} \frac{2}{\pi} \int_0^\infty d\omega' \, \frac{\omega'^2 \text{Im } \mathcal{F}_{\alpha\beta}^{(-)}(\omega')}{\omega'^2 - \omega^2},$$
 (14)

yielding in the dc case

$$\sigma_{\alpha\beta}^{(-)}(0) = \frac{2e^2}{\pi\hbar L^d} \int_0^\infty d\omega \operatorname{Im} \mathcal{F}_{\alpha\beta}^{(-)}(\omega), \qquad (15)$$

consistently with Eqs. (4) and (7).

The imaginary part of the longitudinal conductivity is

Im 
$$\sigma_{\alpha\beta}^{(+)}(\omega) = -\frac{e^2}{\hbar L^d} \frac{2\omega}{\pi} \int_0^\infty d\omega' \frac{\omega' \operatorname{Re} \mathcal{F}_{\alpha\beta}^{(+)}(\omega')}{\omega'^2 - \omega^2};$$
 (16)

in insulators this quantity is related to the static polarizability [10] as

$$\chi_{\alpha\beta}(0) = \lim_{\omega \to 0} \frac{\operatorname{Im} \, \sigma_{\alpha\beta}^{(+)}(\omega)}{\omega}$$
$$= -\frac{2e^2}{\pi\hbar L^d} \int_0^\infty d\omega \, \frac{\operatorname{Re} \, \mathcal{F}_{\alpha\beta}^{(+)}(\omega)}{\omega}. \tag{17}$$

Even this expression is proved to be equivalent to the corresponding Kubo formula [12].

In the metallic case a comparison with the Kubo formula for longitudinal conductivity shows that Eq. (12) yields the regular term only and lacks the Drude contribution. It has been observed above that Re  $F_{\alpha\beta}(0^+)$ —i.e. the quantum metric—diverges in metals: this divergence owes in fact to the Drude term in the Souza-Wilkens-Martin sum rule [14, 15]. If the same sum rule is evaluated by excluding the Drude term, one gets a regular metric which coincides indeed with Re  $F_{\alpha\beta}(0^+)$ .

The Drude term accounts for the longitudinal adiabatic response of the many-electron system to a dc field; it can be expressed in the same  $\kappa$ -space whose geometry we are investigating here, but it will be shown that in the present framework it is eminently nongeometrical. Quite generally, the adiabatic evolution of an observable is comprised of a nongeometrical and a geometrical term: in the present case of dc conductivity, the longitudinal one is the nongeometrical term and the transverse one is the geometrical term.

The adiabatic observable is the macroscopic current density, whose operator is

$$\hat{\mathbf{j}} = -\frac{e}{L^d}\hat{\mathbf{v}} = -\frac{e}{\hbar L^d}\partial_{\kappa}\hat{H}.$$
 (18)

If the Hamiltonian acquires a time-dependence, the  $\kappa$ -derivative of the energy is

$$\partial_{\kappa_{\alpha}} \langle \Psi_{t} | \hat{H}_{t} | \Psi_{t} \rangle = \langle \Psi_{t} | \partial_{\kappa_{\alpha}} \hat{H}_{t} | \Psi_{t} \rangle$$

$$+ i\hbar (\langle \partial_{\kappa_{\alpha}} \Psi_{t} | \dot{\Psi}_{t} \rangle - \langle \dot{\Psi}_{t} | \partial_{\kappa_{\alpha}} \Psi_{t} \rangle).$$
(19)

The adiabatic theorem states that "when the change of the Hamiltonian in time is made infinitely slow, the system, when started from a stationary state, passes through the corresponding stationary states for all times" [16] . If we set  $|\Psi_t\rangle = |\Psi_0\rangle$  at t=0, the adiabatic limit of Eq. (20) yields

$$\langle \Psi_0 | \partial_{\kappa_\alpha} \hat{H} | \Psi_0 \rangle = \partial_{\kappa_\alpha} E_0 - \hbar \Omega_{\alpha\beta} \, \dot{\kappa}_\beta, \tag{20}$$

where the Hellmann-Feynman theorem has also been used. The two terms in the adiabatic evolution are clearly the nongeometrical and geometrical one, respectively.

It has been observed above that  $\kappa$  is actually a vector potential; therefore  $\dot{\kappa} = -e\mathcal{E}/\hbar$  when a macroscopic field  $\mathcal{E}$  is switched on. By considering its geometrical term only, Eq. (20) yields

$$j_{\alpha} = -\frac{e}{\hbar L^{d}} \frac{1}{\hbar} \langle \Psi_{0} | \partial_{\kappa_{\alpha}} \hat{H} | \Psi_{0} \rangle = -\frac{e^{2}}{\hbar L^{d}} \Omega_{\alpha\beta} \mathcal{E}_{\beta}, \quad (21)$$

which indeed coincides with Eq. (4). If we consider instead the nongeometrical term only in Eq. (20) the time-derivative of the current is

$$\partial_t j_{\alpha}(t) = -\frac{e}{\hbar L^d} \frac{\partial^2 E_0}{\partial t \partial \kappa_{\alpha}} = \frac{e^2}{\hbar^2 L^d} \frac{\partial^2 E_0}{\partial \kappa_{\alpha} \partial \kappa_{\beta}} \mathcal{E}_{\beta}, \tag{22}$$

clearly showing that the many-electron system in a dc field undergoes free acceleration (in absence of extrinsic dissipation mechanisms). The causal Fourier transform of Eq. (22) leads to the famous Kohn's formula for longitudinal dc conductivity [17]:

$$\sigma_{\alpha\beta}^{\text{(Drude)}}(\omega) = \frac{D_{\alpha\beta}}{\pi} \frac{i}{\omega + i\eta}, \quad D_{\alpha\beta} = \frac{\pi e^2}{\hbar^2 L^d} \frac{\partial^2 E_0}{\partial \kappa_\alpha \partial \kappa_\beta}.$$
(23)

Therefore—as stated above—the transverse dc conductivity is geometrical and is in fact proportional to Im  $F_{\alpha\beta}(0^+)$ ; the longitudinal one is nongeometrical and out of reach of our tensor  $F_{\alpha\beta}(t)$  as a matter of principle.

Despite being eminently a nongeometrical quantity, the Drude weight  $D_{\alpha\beta}$  can nonetheless be retrieved from our geometric tensor by means of the f-sum rule [12]:

$$D_{\alpha\beta} = D_{\alpha\beta}^{\text{(free)}} - \frac{2e^2}{\hbar L^d} \int_0^\infty d\omega \, \omega \text{Re } \mathcal{F}_{\alpha\beta}^{(+)}(\omega), \qquad (24)$$

where

$$D_{\alpha\beta}^{\text{(free)}} = \frac{\pi e^2 N}{mL^d} \delta_{\alpha\beta} \tag{25}$$

is the Drude weight of an electron gas, i.e. an interacting many-electron system in a flat potential.

In the special case of a crystalline system of non-interacting electrons in a Kohn-Sham framework the ground-state  $|\Psi_0\rangle$  is a Slater determinant of Bloch orbitals  $|\psi_{j\mathbf{k}}\rangle = \mathrm{e}^{i\mathbf{k}\cdot\mathbf{r}}|u_{j\mathbf{k}}\rangle$  with eigenvalues  $\epsilon_{j\mathbf{k}}$ , normalized

in the crystal cell of volume  $V_{\text{cell}}$ . The  $|u_{j\mathbf{k}}\rangle$  orbitals are eigenstates of the one-body Hamiltonian

$$H_{\mathbf{k}} = \frac{1}{2m} \left( \mathbf{p} + \frac{e}{c} \mathbf{A}(\mathbf{r}) + \hbar \mathbf{k} + \hbar \kappa \right)^2 + V(\mathbf{r}), \quad (26)$$

hence a  $\kappa$ -derivative coincides with a **k**-derivative. The n-th state is a monoexcited determinant where the occupied orbital  $|u_{j\mathbf{k}}\rangle$  is replaced by the unoccupied  $|u_{j'\mathbf{k}}\rangle$  one; the vertical transition frequency is  $\omega_{jj'\mathbf{k}}=(\epsilon_{j'\mathbf{k}}-\epsilon_{j\mathbf{k}})/\hbar$ . The many-body matrix elements of  $\hat{\mathbf{v}}$  are transformed—owing to the Slater-Condon rules—into the matrix element of the corresponding one-body velocity  $\mathbf{v}=\partial_{\mathbf{k}}H_{\mathbf{k}}/\hbar$ , evaluated at  $\mathbf{k}=0$ . If  $\epsilon_{\mathrm{F}}$  is the Fermi level, Eq. (8) becomes

$$\mathcal{F}_{\alpha\beta}(\omega) = \sum_{\substack{\epsilon_{j\mathbf{k}} < \epsilon_{\mathrm{F}} \\ \epsilon_{j'\mathbf{k}} > \epsilon_{\mathrm{F}}}} \frac{\langle u_{j\mathbf{k}} | v_{\alpha} | u_{j'\mathbf{k}} \rangle \langle u_{j'\mathbf{k}} | v_{\beta} | u_{j\mathbf{k}} \rangle}{\omega_{jj'\mathbf{k}}^2} \frac{i}{\omega - \omega_{jj'\mathbf{k}} + i\eta};$$
(27)

the formula is given here per spin channel.

As usual when dealing with conductivities, the singular function  $\mathcal{F}_{\alpha\beta}(\omega)$  becomes a regular one in the thermodynamic limit  $L \to \infty$ . The limit can be explicitly performed in the crystalline case, where the sum over  $\mathbf{k}$  converges to a Brillouin-zone (BZ) integral:

$$\mathcal{F}_{\alpha\beta}(\omega) = \frac{V_{\text{cell}}}{(2\pi)^d} \sum_{jj'} \int_{BZ} d\mathbf{k} \frac{i f_{j\mathbf{k}} (1 - f_{j'\mathbf{k}})}{\omega - \omega_{jj'\mathbf{k}} + i\eta} \times \frac{\langle u_{j\mathbf{k}} | v_{\alpha} | u_{j'\mathbf{k}} \rangle \langle u_{j'\mathbf{k}} | v_{\beta} | u_{j\mathbf{k}} \rangle}{\omega_{jj'\mathbf{k}}^2}, \qquad (28)$$

where  $f_{j\mathbf{k}}$  is the Fermi occupancy factor.

Analogously to the many-body original definition, Eq. (5), the geometrical content of the tensor is clearly apparent in the time domain:

$$F_{\alpha\beta}(t) = \frac{V_{\text{cell}}}{(2\pi)^d} \sum_{j} \int_{\text{BZ}} d\mathbf{k} \, f_{j\mathbf{k}} \langle \partial_{k_{\alpha}} u_{j\mathbf{k}} | \, U_{j\mathbf{k}}(t) \, | \partial_{k_{\beta}} u_{j\mathbf{k}} \rangle,$$

$$U_{j\mathbf{k}}(t) = \theta(t) \sum_{j'} |u_{j'\mathbf{k}}\rangle e^{-i\omega_{jj'\mathbf{k}}t} \langle u_{j'\mathbf{k}} |. \tag{29}$$

In the insulating case this is apparently similar to—and conceptually different from—the time-dependent quantum geometric tensor of Refs. [10, 11]..

By construction, at time  $t=0^+$  Eq. (29) yields the standard band-structure expression for the metric-curvature tensor [3] evaluated in the parallel-transport gauge:

$$F_{\alpha\beta}(0^{+}) = \frac{V_{\text{cell}}}{(2\pi)^{d}} \sum_{j} \int_{\text{BZ}} d\mathbf{k} \ f_{j\mathbf{k}} \langle \partial_{k_{\alpha}} u_{j\mathbf{k}} | \partial_{k_{\beta}} u_{j\mathbf{k}} \rangle; \quad (30)$$

this is subsequently evolved by the unitary matrix  $U_{j\mathbf{k}}(t)$ . Notice that Eq. (30) applies even to the metallic case: its imaginary part is then proportional to the intrinsic anomalous Hall conductivity [5], while its real part is proportional to the Souza-Wilkens-Martin integral [14, 15], where only the regular part of the longitudinal conductivity is included.

A final comment concerns dissipation. Insofar as L stays finite,  $\sigma_{\alpha\beta}(\omega)$  is causal but nondissipative. It becomes dissipative after the  $L \to \infty$  limit is performed, because an infinite system can be regarded as its own thermostat [18]. Owing to Ohm's law the Joule heating Q is proportional to the squared electric field [19]. When the spectrum is continuous—as in Eq. (28)—its geometrical expression in terms of  $\mathcal{F}_{\alpha\beta}(\omega)$  becomes [5]:

$$Q = \frac{e^{2}\omega}{2\hbar L^{d}} \left\{ \operatorname{Re} \mathcal{F}_{\alpha\beta}^{(+)}(\omega) \operatorname{Re} \left[ \mathcal{E}_{\alpha}(\omega) \mathcal{E}_{\beta}^{*}(\omega) \right] + \operatorname{Im} \mathcal{F}_{\alpha\beta}^{(-)}(\omega) \operatorname{Im} \left[ \mathcal{E}_{\alpha}(\omega) \mathcal{E}_{\beta}^{*}(\omega) \right] \right\}.$$
(31)

In conclusion, I have shown that both the metric and the curvature are naturally endowed with a time dependence, which elucidates the geometrical nature of electron transport even beyond the adiabatic regime. The causal evolution of the metric is at the root of the longitudinal optical conductivity and provides its explicit exact expression at any frequency, with the exception of the Drude term in the metallic case.. Analogously, the evolution of the curvature is at the root of the transverse optical conductivity and provides its explicit exact expression. The presentation is made in a many-body framework, which allows dealing in principle with cases including interaction and disorder, but has also the virtue of showing the main concepts by means of simple notations and formulæ. The special case of a crystalline system of noninteracting electrons, in a band-structure formalism, is also presented.

I thank Raquel Queiroz for the fruitful conversations we had about this topic. Work supported by the Office of Naval Research (USA) Grant No. N00014-20-1-2847.

- \* resta@iom.cnr.it.it
- M. V. Berry, Proc. Roy. Soc. Lond. A 392, 45 (1984).
- [2] Q. Niu and D. J. Thouless, J. Phys A 17, 2453 (1984).
- [3] D. Vanderbilt, Berry Phases in Electronic Structure Theory (Cambridge University Press, Cambridge, 2018).
- [4] D. Xiao, M.-C. Chang, and Q. Niu, Rev. Mod. Phys. 82, 1959 (2010).
- [5] N. Nagaosa, J. Sinova, S. Onoda, A. H. MacDonald, and N. P. Ong, Rev. Mod. Phys. 82, 1539 (2010).
- [6] R. Resta, in: Encyclopedia of Condensed Matter Physics, 2nd edition, T. Chakraborty ed. (Elsevier, 2023). p. 670.
- [7] R. Resta, J. Chem. Phys. 158, 024105 (2023).
- [8] J. Ahn, G.-Y. Guo, N. Nagaosa, and A. Vishwanath, Nat. Phys. 18, 290 (2022).
- [9] N. Nagaosa, Annals of Physics 447, 169146 (2022).
- [10] I. Komissarov, T. Holder, and R. Queiroz, Nat. Commun. 15, 4621 (2024).

- [11] N. Verma and R. Queiroz, https://arxiv.org/pdf/2403.07052.
- [12] See the full set of Kubo formulæ in the Supplemental Material.
- [13] D. Ceresoli and R. Resta, Phys. Rev. B 76, 012405 (2007).
- [14] I. Souza, T. Wilkens, and R. M. Martin, Phys. Rev. B 62, 1666 (2000).
- [15] R. Resta, Riv. Nuovo Cimento 41, 463 (2018).
- [16] T. Kato, J. Phys. Soc. Jpn. 5, 435 (1950).
- [17] W. Kohn, Phys. Rev. 133, A171 (1964).
- [18] G. F. Giuliani and G. Vignale, Quantum Theory of the Electron Liquid (Cambridge University Press, Cambridge, 2005).
- [19] L. D. Landau, E. N. Lifshitz, and L. Pitaevskii, *Electrodynamics of Continuous Media*, 2nd Ed. (Elsevier, Oxford, 2008).

## Supplemental Material: Kubo formulæ

I adopt here the notations of the main text in order to explicitly display the relevant Kubo formulæ.

Longitudinal (symmetric) conductivity:

$$\sigma_{\alpha\beta}^{(+)}(\omega) = D_{\alpha\beta} \left[ \delta(\omega) + \frac{i}{\pi\omega} \right] + \sigma_{\alpha\beta}^{(\text{regular})}(\omega), \quad (32)$$

$$D_{\alpha\beta} = \frac{\pi e^2}{L^d} \left( \frac{N}{m} \delta_{\alpha\beta} - \frac{2}{\hbar} \sum_{n \neq 0} \frac{\mathcal{R}_{n,\alpha\beta}}{\omega_{0n}} \right), \quad (33)$$

where for  $\omega > 0$ :

Re 
$$\sigma_{\alpha\beta}^{(\text{regular})}(\omega) = \frac{\pi e^2}{\hbar L^d} \sum_{n \neq 0} \frac{\mathcal{R}_{n,\alpha\beta}}{\omega_{0n}} \delta(\omega - \omega_{0n}), \quad (34)$$

Im 
$$\sigma_{\alpha\beta}^{\text{(regular)}}(\omega) = \frac{2e^2}{\hbar L^d} \sum_{n \neq 0} \frac{\mathcal{R}_{n,\alpha\beta}}{\omega_{0n}} \frac{\omega}{\omega_{0n}^2 - \omega^2};$$
 (35)

the Drude weight  $D_{\alpha\beta}$  vanishes in insulators. At  $\omega=0$  Eq. (33) can be actually summed, to yield the famous Kohn's (1964) formula:

$$D_{\alpha\beta} = \frac{\pi e^2}{\hbar^2 L^d} \frac{\partial^2 E_0}{\partial \kappa_{\alpha} \partial \kappa_{\beta}}.$$
 (36)

In the main text the same quantity occurs as the nongeometrical term of the adiabatic response.

The real part of longitudinal conductivity obeys the

f-sum rule

$$\int_{0}^{\infty} d\omega \operatorname{Re} \, \sigma_{\alpha\beta}(\omega) = \frac{D_{\alpha\beta}}{2} + \int_{0}^{\infty} d\omega \operatorname{Re} \, \sigma_{\alpha\beta}^{(\text{regular})}(\omega)$$
$$= \frac{\pi e^{2} N}{2mL^{d}} \delta_{\alpha\beta}, \tag{37}$$

Transverse (antisymmetric) conductivity:

The intrinsic (geometric) contribution to the anomalous Hall conductivity requires spontaneous breaking of timereversal symmetry. Its expression is:

Re 
$$\sigma_{\alpha\beta}^{(-)}(\omega) = \frac{2e^2}{\hbar L^d} \sum_{n\neq 0} \frac{\mathcal{I}_{n,\alpha\beta}}{\omega_{0n}^2 - \omega^2}$$
 (38)

Im 
$$\sigma_{\alpha\beta}^{(-)}(\omega) = \frac{\pi e^2}{\hbar L^d} \sum_{n \neq 0} \frac{\mathcal{I}_{n,\alpha\beta}}{\omega_{0n}} \delta(\omega - \omega_{0n}), \ \omega > 0.(39)$$

The dc transverse conductivity is easily recast in terms of the many-body Berry curvature at  $\kappa = 0$ :

Re 
$$\sigma_{\alpha\beta}^{(-)}(0) = -\frac{e^2}{\hbar L^d} \Omega_{\alpha\beta}$$
. (40)

The expression holds for metals and insulators, in either 2d or 3d.

Polarizability (symmetric):

The polarizability tensor in insulators is defined as  $\chi_{\alpha\beta}(\omega) = \partial P_{\alpha}/\partial \mathcal{E}_{\beta}$ . Since the current is the time derivative of the polarization, one has  $j_{\alpha}(\omega) = -i\omega P_{\alpha}(\omega) = -i\omega \chi_{\alpha\beta}(\omega)\mathcal{E}_{\beta}(\omega)$  and

Re 
$$\chi_{\alpha\beta}(\omega) = \frac{\text{Im }\sigma_{\alpha\beta}^{(+)}(\omega)}{\omega} = \frac{2e^2}{\hbar L^d} \sum_{n\neq 0} \frac{\mathcal{R}_{n,\alpha\beta}}{\omega_{0n}(\omega_{0n}^2 - \omega^2)}.$$
 (41)

In the  $\omega \to 0$  one retrieves the Kubo formula for the static polarizability:

$$\chi_{\alpha\beta}(0) = \frac{2e^2}{\hbar L^d} \sum_{n \neq 0} \frac{\langle \Psi_0 | \hat{v}_\alpha | \Psi_n \rangle \langle \Psi_n | \hat{v}_\beta | \Psi_0 \rangle}{\omega_{0n}^3}, \tag{42}$$