ASYMPTOTIC BEHAVIOR OF SOLUTIONS OF A TIME-SPACE FRACTIONAL DIFFUSIVE VOLTERRA EQUATION

SOFWAH AHMAD AND MOKHTAR KIRANE

ABSTRACT. In this paper, we study the time-space fractional differential equation of the Volterra type:

$$\mathcal{D}^{\alpha}_{0|t}(u) + (-\Delta_N)^{\sigma} u = u(1 + au - bu^2) - au \int_0^t \mathcal{K}(t-s)u(\cdot) \, ds,$$

where a, b > 0 are given constants, $\alpha, \sigma \in (0, 1)$, equipped with a homogeneous Neumann's boundary condition and a positive initial data. The boundedness and uniform continuity of the solution on the entire \mathbb{R}^+ are established. Moreover, the asymptotic behavior of the positive solution is investigated.

Keywords: time–space fractional differential equation; asymptotic behavior **MSC Classification (2020)**: 26A33, 35B09, 35B40, 34K37, 35R09, 35R11.

1. Introduction

In this paper, we are interested to study the time asymptotic behavior of the bounded solution of the following problem

(1)
$$\mathcal{D}^{\alpha}_{0|t}(u) + (-\Delta_N)^{\sigma} u = u(1 + au - bu^2) - au\mathcal{K} * u,$$

$$(2) u(x,0) = u_0(x)$$

posed on $Q = \Omega \times (0, T)$ where $\Omega \subset \mathbb{R}^d$ (where d = 1, 2 in practice) is a bounded domain with smooth boundary $\partial \Omega$ with outward normal η ; the asterisk (*) denotes the time convolution, i.e

$$\mathcal{K} * u(x,t) = \int_0^t \mathcal{K}(t-s)u(\cdot,s)ds;$$

the constants a and b in the problem are assumed to be positive and \mathcal{K} is the delay kernel function satisfying

$$\mathcal{K} \ge 0, \ \mathcal{K} \in C^1(0,\infty) \cap L^1(0,\infty), \ \int_0^\infty \mathcal{K}(t)dt = 1$$

The derivative operator $\mathcal{D}_{0|t}^{\alpha}$ is the Caputo fractional derivative of order $0 \leq \alpha \leq 1$, the operator $(-\Delta_N)^{\sigma}, 0 < \sigma < 1$, is the fractional power of the classical Laplacian operator with its specified domain and $u_0(x) \geq 0$ is a given bounded function.

Equation (1) describes the evolution of a single species population whose density is u at time t in the bounded domain Ω . As in the existing model with classical

derivatives, the first term in the right-hand side represents the non-delay term with growth rate a due to advantages of aggregating as a group (e.g protected against predator and increased reproduction success) and the decay rate b is due to interspecies competition; the delay represented by convolution term in the Volterra equation (1) represents the regulatory term with the assumption that population growth depends on a smooth average over the past history of the population rather than only information at a certain instantaneous time. More discussion about various population models can be found in [1] and [2].

In this paper, we use the time fractional derivative $\mathcal{D}_{0|t}^{\alpha} u$ rather than the usual derivative to represent the accumulation of the information on the population density variation within the specified time interval (0, T); the fractional Laplacian term $(-\Delta_N)^{\sigma} u$ describes spatial distribution of the species with its possibility to move to remote sites inside the region Ω within a certain period of time (see [3]). With the fractional Laplacian, it is implied that the model under consideration is equipped with the Neumann's boundary condition

(3)
$$\nabla u \cdot \eta = 0, \ (x,t) \in \partial \Omega \times (0,T),$$

where η is the outer normal of $\partial \Omega$, which says that no species migration into or out of Ω .

Before proceeding further, let us dwell on the existing literature on the classical differential equations related to our problem (i.e $\alpha = \sigma = 1$). In [4], the classical differential equation version of (1) is studied with the delay term $(1 + a - b)u\mathcal{K} * u$ instead of $au\mathcal{K} * u$, where the authors showed that for 0 < b < 1 + a and $4a < (3(b-1) + \sqrt{b^2 + 6b + 1})$, u(x, t) goes to 1 uniformly for $x \in \overline{\Omega}$. Schiaffino [5], Redlinger [6], and Yamada [7], studied the same model with intraspecific competion term $u\mathcal{K} * u$ subject to Neumann's boundary condition. The asymptotic properties of solutions of the problem with Dirichlet's boundary condition is also studied by Yamada [8]. In Yoshida and Yamada [9], the study of the spatial inhomogeneity is taken into consideration. Kirane, Guedda, and Tatar [10] also studied problem (1); by means of an entropy functional, they showed that the solution tends to $1/\sqrt{b}$ as $t \to \infty$.

In this work, using the method of Langlais-Phillips, that is outlined in the paper of Kirane [11], by noting that the orbits of the dynamical system are precompact in $C^{\mu}(\Omega)$, for some $\mu > 0$ that will be precised later, we show that the solution of the problem tends to $1/\sqrt{b}$ which is the positive solution of the limiting problem of (1)-(2).

2. Preliminaries

2.1. **Definition and notation.** First, we give the definitions of Riemann-Liouville fractional integral, Riemann-Liouville fractional derivative and Caputo fractional derivative (see e.g. [12])

Definition 1. Let $f \in L^1([0,T])$ and $\alpha > 0$. The left and right Riemann-Liouville fractional integrals of order $\alpha > 0$ are defined by

$$(I_{0|t}^{\alpha}f)(t) := \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(\tau)}{(t-\tau)^{1-\alpha}} d\tau,$$
$$(I_{t|T}^{\alpha}f)(t) := \frac{1}{\Gamma(\alpha)} \int_t^T \frac{f(\tau)}{(\tau-t)^{1-\alpha}} d\tau,$$

respectively.

Definition 2. The Riemann-Liouville fractional derivative of order $\alpha \in (0, 1]$ of an absolutely continuous function f is defined by

$$({}^{RL}D^{\alpha}_{0|t}y)(t) := \frac{d}{dt}(I^{1-\alpha}_{0|t}y)(t) = \frac{1}{\Gamma(1-\alpha)}\frac{d}{dt}\int_0^t \frac{y(\tau)}{(t-\tau)^{\alpha}}\,d\tau, \quad (t>0),$$

$${}^{(RL}D^{\alpha}_{t|T}y)(t) := -\frac{d}{dt}(I^{1-\alpha}_{t|T}y)(t) = \frac{-1}{\Gamma(1-\alpha)}\frac{d}{dt}\int_{t}^{T}\frac{y(\tau)}{(\tau-t)^{\alpha}}\,d\tau, \quad (t < T).$$

The following relation between Riemann-Liouville integral and derivative of order $0 < \alpha \leq 1$ holds:

Lemma 1. [12, Lemma 2.4] If $\alpha > 0$ and $f(t) \in L^p(0,T)$ $(1 \le p \le \infty)$, then the following equalities

 $({}^{RL}D^{\alpha}_{0|t}I^{\alpha}_{0|t}f)(t) = f(t), \quad and \quad ({}^{RL}D^{\alpha}_{t|T}I^{\alpha}_{t|T}f)(t) = f(t),$

hold almost everywhere on [0, T].

Definition 3. The Caputo fractional derivative of order $0 < \alpha \leq 1$ of a differentiable function f is defined by

$$(\mathcal{D}^{\alpha}_{0|t}f)(t) := \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-\tau)^{-\alpha} f'(\tau) d\tau \quad 0 \le t \le T.$$

Next, we consider the definition of the fractional Laplacian given via the spectrum of the classical Laplacian operator $-\Delta$ on the same domain [13]. Let $\{\lambda_n\}_{n=0}^{\infty}$ with $\lambda_0 = 0 < \lambda_1 < \lambda_2 < \ldots$ be the sequence of eigenvalues of the operator $-\Delta$ on Ω that is equipped with Neumann boundary condition on $\partial\Omega$, and $\{e_n\}_{n=0}^{\infty}$ be the corresponding normalized eigenfunctions in Ω , i.e,

$$-\Delta e_n = \lambda_n e_n, \quad x \in \Omega,$$
$$\frac{\partial e_n}{\partial \eta} = 0, \qquad x \in \partial \Omega$$

Consider the space

$$H^{\sigma}(\Omega) := \left\{ u = \sum_{n=0}^{\infty} u_n e_n \in L^2(\Omega); \quad \|u\|_{H^{\sigma}(\Omega)} = \sum_{n=1}^{\infty} \lambda_n^{\sigma} u_n^2 < +\infty \right\}.$$

Defining the inner product

$$\langle u, v \rangle_{H^{\sigma}(\Omega)} := \sum_{n=1}^{\infty} \lambda_n^{\sigma} u_n v_n, \quad \text{for} \quad u = \sum_{n=0}^{\infty} u_n e_n, \ v = \sum_{n=0}^{\infty} v_n e_n,$$

we see that the space $H^{\sigma}(\Omega)$, equipped with the inner product $\langle u, v \rangle_{H^{\sigma}(\Omega)}$, is a Hilbert space.

Definition 4. The fractional Laplacian operator $(-\Delta_N)^{\sigma}$ is defined by

$$(-\Delta_N)^{\sigma}u(x) := \sum_{n=1}^{\infty} \lambda_n^{\sigma} u_n e_n,$$

with domain

$$D((-\Delta_N)^{\sigma}) = \left\{ u \in H^{\sigma}(\Omega); \quad \frac{\partial u}{\partial \eta} = 0 \right\}.$$

Definition 5. The two parameters Mittag-Leffler function $E_{\alpha,\beta}(z)$ is given by

$$E_{\alpha,\beta}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)} \quad (z,\beta \in \mathbb{C}; \ Re(\alpha) > 0).$$

2.2. **Preliminary lemmas.** In this subsection, we give the preliminary lemmas that will be used in proving our result.

Lemma 2. [12, Lemma 2.22] For a continuous function f(t), the following relationship between fractional integral and Caputo fractional derivative holds

(4)
$$I_{0|t}^{\alpha} \mathcal{D}_{0|t}^{\alpha} f(t) = f(t) - f(0).$$

Lemma 3. If $\varphi \in C^1(\mathbb{R})$ is a convex function and $x \in C([0,T]) \cap C^1((0,T])$, for some T > 0, then

$$\mathcal{D}^{\alpha}_{0|t}\varphi(x)(t) \le \varphi'(x(t))\mathcal{D}^{\alpha}_{0|t}x(t), \qquad 0 < \alpha \le 1.$$

Proof. Since φ is convex function, we have

$$\varphi(x) - \varphi(y) - \varphi'(y)(x - y) \ge \frac{1}{2}\varphi''(\lambda x + \mu y)(x - y)^2$$
, where $0 \le \lambda, \mu < 1$.

Define

$$\Psi(s,t) = \varphi(x(s)) - \varphi(x(t)) - \varphi'(x(t)) \left(x(s) - x(t)\right),$$

then

$$\frac{\partial \Psi(s,t)}{\partial s} = \varphi'(x(s))x'(s) - \varphi'(x(t))x'(s).$$

Thus

$$\begin{split} \mathcal{D}_{0|t}^{\alpha}\varphi(x)(t) &- \varphi'(x(t))\mathcal{D}_{0|t}^{\alpha}x(t) \\ &= \frac{1}{\Gamma(1-\alpha)} \left(\int_{0}^{t} \frac{\varphi'(x(s)) \ x'(s)}{(t-s)^{\alpha}} ds - \varphi'(x(t)) \int_{0}^{t} \frac{x'(s)}{(t-s)^{\alpha}} ds \right) \\ &= \frac{1}{\Gamma(1-\alpha)} \left(\int_{0}^{t} \frac{(\varphi'(x(s)) - \varphi'(x(t))) \ x'(s)}{(t-s)^{\alpha}} ds \right) \\ &= \frac{1}{\Gamma(1-\alpha)} \left(\int_{0}^{t} \frac{\partial \Psi(s,t)}{\partial s} (t-s)^{-\alpha} ds \right) \\ &= \frac{1}{\Gamma(1-\alpha)} \left(\lim_{s \to t} \frac{\Psi(s,t)}{(t-s)^{\alpha}} - \frac{\Psi(0,t)}{t^{\alpha}} - \alpha \int_{0}^{t} \frac{\Psi(s,t)}{(t-s)^{\alpha+1}} ds \right) \\ &= \frac{1}{\Gamma(1-\alpha)} \left(\lim_{s \to t} \frac{\partial \Psi(s,t)/\partial s}{\alpha(t-s)^{\alpha-1}} - \frac{\Psi(0,t)}{t^{\alpha}} - \alpha \int_{0}^{t} \frac{\Psi(s,t)}{(t-s)^{\alpha+1}} ds \right) \\ &= \frac{1}{\Gamma(1-\alpha)} \left(-\frac{\Psi(0,t)}{t^{\alpha}} - \alpha \int_{0}^{t} \frac{\Psi(s,t)}{(t-s)^{\alpha+1}} ds \right), \end{split}$$

and the result follows.

Remark. In the case $\varphi(u) = u^p$, we obtain the generalisation of J.I. Diaz, T. Pierantozi and L. Vazquez conjectured inequality, which was obtained by Alsaedi, Ahmad and Kirane [14].

We need the following result on weakly singular Gronwall's inequality.

Lemma 4. [15, Lemma 6.19] Let $\alpha, C, T \in \mathbb{R}^+$. If $\psi : [0, T] \to \mathbb{R}$ is a non-negative continuous function that satisfies

$$\psi(t) \le \psi(0) + \frac{C}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \psi(s) ds,$$

for all $t \in [0,T]$, then

$$\psi(t) \le \psi(0) E_{\alpha,1}(Ct^{\alpha}),$$

where $E_{\alpha,\beta}(z)$ is the two parameters Mittag-Leffler function.

Using the properties of scalar products, the following integration by parts formula for the fractional Laplacian immediately follows.

Lemma 5. For $0 < \sigma \leq 1$ and $u, v \in D((-\Delta_N)^{\sigma})$, it holds

$$\int_{\Omega} u(-\Delta_N)^{\sigma} v \ dx = \int_{\Omega} v(-\Delta_N)^{\sigma} u \ dx.$$

We also need the following inequality of Stroock and Varopoulos.

Lemma 6. [16, Theorem 1] Let $0 < \sigma \leq 2$ and p > 1. For nonnegative $u \in L^p(\Omega)$ with $(-\Delta_N)^{\frac{\sigma}{2}}u \in L^p(\Omega)$, we have

$$\int_{\Omega} u^{p-1} (-\Delta_N)^{\frac{\sigma}{2}} u \, dx \ge \frac{4(p-1)}{p^2} \int_{\Omega} \left| (-\Delta_N)^{\frac{\sigma}{4}} u^{\frac{p}{2}} \right|^2 dx.$$

We also have the following relation for time-fractional integration by parts.

Lemma 7. [17, Theorem 11] Let $0 < \alpha < 1$. The following relations hold:

$$\int_0^T y(t) \mathcal{D}_{0|t}^{\alpha} x(t) dt = \int_0^T x(t) \,^{RL} D_{t|T}^{\alpha} y(t) dt + \left[x(t) \, I_{t|T}^{1-\alpha} y(t) \right]_{t=0}^{t=T}$$

3. Local Existence

In this section, we prove the existence and uniqueness of local (in time) solution u(x,t) on some interval [0,T], for $T < \infty$. Let

$$f(x,t,u) = u(1 + au - bu^2) - au\mathcal{K} * u.$$

Furthermore, let the operators $S_{\alpha}(t), P_{\alpha}(t) : H^{\sigma}(\Omega) \to H^{\sigma}(\Omega)$ be defined as

(5)
$$S_{\alpha}(t)v := \int_{0}^{\infty} \Phi_{\alpha}(\tau) S(\tau t^{\alpha}) v \, d\tau,$$

and

(6)
$$P_{\alpha}(t)v := \alpha t^{\alpha-1} \int_{0}^{\infty} \tau \Phi_{\alpha}(\tau) S\left(\tau t^{\alpha}\right) v \ d\tau,$$

with S(t) is the strongly continuous semigroup generated by the operator $A := (-\Delta_N)^{\sigma}$ and $\Phi_{\alpha}(z)$ is the Wright type function (see Kilbas et al. [12])

$$\Phi_{\alpha}(z) := \sum_{n=0}^{\infty} \frac{(-z)^n}{n! \Gamma(-\alpha n + 1 - \alpha)}, \quad 0 < \alpha < 1, \quad z \in \mathbb{C}.$$

Then, we have the following definition for the representation of the solution of the problem (1)-(3).

Definition 6. Let $u_0 \in L^{\infty}(\Omega)$ and T > 0. We say that $u \in C([0, T], L^{\infty}(\Omega))$ is a mild solution of the problem (1)-(3) if it can be expressed as

(7)
$$u(t) = S_{\alpha}(t)u_0 + \int_0^t P_{\alpha}(t-s)f(s) \, ds,$$

for $t \in [0, T]$.

It is useful to first notice the boundedness of operators S_{α} and P_{α} .

Lemma 8. [13, Proposition 2.1.3] There exists constants $C_s, C_p > 0$ such that for all t > 0 and $v \in H^{\sigma}(\Omega)$, we have

- (1) $||S_{\alpha}(t)v||_{H^{\sigma}(\Omega)} \leq C_s ||v||_{H^{\sigma}(\Omega)},$
- (2) $||t^{1-\alpha}P_{\alpha}(t)v||_{H^{\sigma}(\Omega)} \leq C_p ||v||_{H^{\sigma}(\Omega)}.$

Theorem 1. (Local existence) Let $u_0 \in C(\Omega)$; then there exists a maximal time $T_{max} > 0$ and a unique mild solution $u \in C([0, T_{max}); L^{\infty}(\Omega))$ of the problem (1)-(2) with the alternative

• either
$$T_{max} = +\infty$$
;

• or $T_{max} < +\infty$ and

$$\lim_{t \to T_{max}} \|u(t)\|_{\infty} = \lim_{t \to T_{max}} \sup_{0 < t < T_{max}} |u(t)| = +\infty.$$

Proof. We prove the local existence through Banach's fixed point theorem. Let $u_0 : \Omega \to \mathbb{R}^+$ be such that $||u_0||_{L^{\infty}(\Omega)} \equiv \rho_0$ small enough. Choose ρ such that $\rho - C_s \rho_0 > 0$ and let

$$B_{\rho} := \{ u \in C([0,T]; L^{\infty}(\Omega)) : \|u\|_{B_{\rho}} := \sup_{t \in [0,T]} \|u(\cdot,t)\|_{L^{\infty}(\Omega)} \le \rho \}.$$

We define the operator F on B_{ρ} as

$$(Fu)(t) := S_{\alpha}(t)u_0 + \int_0^t P_{\alpha}(t-s)(f)(x,s,u(s)) \ ds.$$

To use Banach's fixed point theorem, we shall prove that F is a bounded operator and it is a contraction on B_{ρ} .

• We first show that $F: B_{\rho} \to B_{\rho}$. With the aid of Lemma 8, we have

$$\begin{aligned} \|Fu\|_{L^{\infty}(\Omega)} &\leq \|S_{\alpha}(t)u_{0}\|_{L^{\infty}(\Omega)} + \int_{0}^{t} \|P_{\alpha}(t-s)\|\|f(u(s))\|_{L^{\infty}(\Omega)} \, ds \\ &\leq C_{s}\|u_{0}\|_{L^{\infty}(\Omega)} + \int_{0}^{t} (t-s)^{\alpha-1}C_{p}\|f(u)\|_{L^{\infty}(\Omega)} \, ds. \end{aligned}$$

We note that

$$\begin{split} \|f(u)\|_{L^{\infty}(\Omega)} &\leq \|u + au^{2} - bu^{3}\|_{L^{\infty}(\Omega)} + \left\|au \int_{0}^{t} K(t - s)u(x, s)ds\right\|_{L^{\infty}(\Omega)} \\ &\leq \|u\|_{B_{\rho}} + a\|u\|_{B_{\rho}}^{2} + b\|u\|_{B_{\rho}}^{3} + a\|u\|_{B_{\rho}}^{2} \int_{0}^{\infty} \mathcal{K}(s)ds \\ &\leq \rho + 2a\rho^{2} + b\rho^{3}, \end{split}$$

 \mathbf{SO}

$$\|Fu\|_{B_{\rho}} \leq C_{s}\rho_{0} + C_{p}(\rho + 2a\rho^{2} + b\rho^{3}) \sup_{t \in [0,T]} \left\{ \int_{0}^{t} (t-s)^{\alpha-1} ds \right\}$$

$$\leq C_{s}\rho_{0} + C_{p}(\rho + 2a\rho^{2} + b\rho^{3}) \sup_{t \in [0,T]} \left\{ \frac{t^{\alpha}}{\alpha} \right\}$$

$$\leq C_{s}\rho_{0} + C_{p}(\rho + 2a\rho^{2} + b\rho^{3}) \frac{T^{\alpha}}{\alpha}$$

$$\leq \rho,$$

by choosing $T \leq T_1$ where

$$T_1 = \left[\frac{\alpha(\rho - C_s \rho_0)}{C_p(\rho + 2a\rho + b\rho^2)}\right]^{1/\alpha}.$$

Therefore, we conclude that $F: B_{\rho} \to B_{\rho}$. • Next we show that F is a contraction on B_{ρ} . We observe that f is Lipschitz continuous on B_{ρ} since for any $u, v \in B_{\rho}$, we have

$$f(u) - f(v) = (u - v) + a(u^{2} - v^{2}) - b(u^{3} - v^{3})$$

- $a[(u - v)(\mathcal{K} * u) - v(\mathcal{K} * (u - v))]$
= $(u - v)(1 + a(u + v) - b(u^{2} + uv + v^{2}))$
- $a[(u - v)(\mathcal{K} * u) - v(\mathcal{K} * (u - v))],$

 \mathbf{SO}

$$\|f(u) - f(v)\|_{B_{\rho}} \leq \|u - v\|_{B_{\rho}} \|1 + a(u + v) - b(u^{2} + uv + v^{2})\|_{B_{\rho}} + a\|u - v\|_{B_{\rho}} \|\mathcal{K} * u\|_{B_{\rho}} + a\|v\|_{B_{\rho}} \|\mathcal{K} * (u - v)\|_{B_{\rho}}.$$

Whereupon,

$$\|f(u) - f(v)\|_{B_{\rho}} \le \|u - v\|_{B_{\rho}} \left(1 + a(2\rho) + b(3\rho^2) + 2a\rho \int_0^\infty \mathcal{K}(s) \, ds\right)$$
$$:= C_f \|u - v\|_{B_{\rho}}.$$

Therefore,

$$\|Fu - Fv\|_{B_{\rho}} = \left\| \int_{0}^{t} P_{\alpha}(t-s)(f(u(s) - f(v(s)) \ ds \right\|_{B_{\rho}}$$

$$\leq C_{p} \ C_{f} \|u - v\|_{B_{\rho}} \sup_{t \in [0,T]} \left\{ \int_{0}^{t} (t-s)^{\alpha-1} \ ds \right\}$$

$$\leq C_{p} \ C_{f} \|u - v\|_{B_{\rho}} \frac{T^{\alpha}}{\alpha}$$

$$\leq \frac{1}{2} \|u - v\|_{B_{\rho}},$$

for $T \leq T_2$ where $T_2 = \left[\frac{\alpha}{2C_p C_f}\right]^{1/\alpha}$. By letting $T_3 = \min\{T_1, T_2\}$, we see that F is a contraction for $t \in [0, T_3]$ and hence, by Banach's fixed point theorem, the problem has a unique solution on $[0, T_3]$. • Let

 $T_{\max} := \sup\{T > 0 : \text{ there exists a mild solution } u \in B_{\rho}\} \le +\infty.$

We show that either $T_{\max} = +\infty$, or $T_{\max} < +\infty$ and (in the latter case) $\|u\|_{L^{\infty}(\Omega)} \to +\infty$ as $t \to T_{\max}$. If the former holds, we say that the solution is global. To prove the latter part, we proceed by contradiction. Suppose that u is a solution on the interval [0, T] with $T < T_{\max} = +\infty$ but $\|u\|_{L^{\infty}(\Omega)} < \infty$. For some $\tau > 0$, we define $w \in C((0, T_{\max} + \tau), C(\Omega))$ as

$$w(t) = \begin{cases} u(t), & t \in [0, T_{\max}], \\ u(t-\tau), & t \in [T_{\max}, T_{\max} + \tau]. \end{cases}$$

We see that w(t) is also a mild solution of the problem (1)-(2). This contradicts the definition of T_{max} .

• We prove that the solution is unique on the whole interval of existence. Let $u, \bar{u} \in B_{\rho}$ be two mild solutions of problem (1)-(3) on $[0, T_{\max})$, then

$$\begin{aligned} \|u - \bar{u}\|_{L^{\infty}(\Omega)} &\leq \int_{0}^{t} \|P_{\alpha}(t - s)(f(u(s) - f(\bar{u}(s)))\|_{L^{\infty}(\Omega)} \, ds \\ &\leq C_{p} \, C_{f} \int_{0}^{t} (t - s)^{\alpha - 1} \, \|u(s) - \bar{u}(s)\|_{L^{\infty}(\Omega)} \, ds. \end{aligned}$$

Let $\Psi(t) = ||u(\cdot, t) - \bar{u}(\cdot, t)||_{L^{\infty}(\Omega)}$. Note that $\Psi(0) = 0$. By Lemma 4, we conclude that

 $u - \bar{u} = 0 \quad \Longleftrightarrow \quad u = \bar{u}.$

4. Positivity, Boundedness and Uniform Continuity of the Solution

We have established that the solution is bounded on $0 < t < T < T_{max}$. In this section, we show that the solution is positive and bounded for all $t \in \mathbb{R}$.

Theorem 2. Assuming $u_0 \ge 0, u_0 \not\equiv 0$ and $u_0 \in C(\overline{\Omega})$, the solution of problem (1)-(2) satisfies

$$0 \le u(x,t) \le \max\{\|u_0\|_{\infty}, R\} \text{ on } \overline{\Omega} \times (0,\infty),$$

where R is the positive solution of $1 + aw - bw^2 = 0$. Observe that $R \ge \frac{1}{\sqrt{b}}$.

Proof. First we prove the positivity. Let $u^+ = max(u, 0)$ and $u^- = min(u, 0)$. Write $u = u^+ + u^-$. We note that $u^+u^- = 0$, $u^2 = (u^+ + u^-)(u^+ + u^-) = u^+ + u^ (u^+)^2 + (u^-)^2$, and $u^3 = u \cdot u^2 = (u^+)^3 + (u^-)^3$.

Taking the scalar product of (1) by u^- in $L^2(\Omega)$, we obtain

$$\int_{\Omega} u^{-} \mathcal{D}^{\alpha}_{0|t} u^{-} + u^{-} (-\Delta_{N})^{\sigma} (u^{-}) \, dx = \int_{\Omega} (u^{-})^{2} + a(u^{-})^{3} - b(u^{-})^{4} \, dx$$
$$- a \int_{\Omega} (u^{-})^{2} \mathcal{K} * u^{-} \, dx.$$

Using integration by parts, we get

$$\int_{\Omega} u^{-} \mathcal{D}_{0|t}^{\alpha} u^{-} dx = -\int_{\Omega} |(-\Delta_{N})^{(\sigma/2)} (u^{-})|^{2} dx + \int_{\Omega} (u^{-})^{2} + a(u^{-})^{3} - b(u^{-})^{4} - a \int_{\Omega} (u^{-})^{2} \mathcal{K} * u^{-} dx;$$

which implies

$$\int_{\Omega} u^{-} \mathcal{D}^{\alpha}_{0|t} u^{-} dx \leq \int_{\Omega} (u^{-})^{2} - a(u^{-})^{2} \mathcal{K} * u^{-} dx$$

Since the local existence is established, there exists T_{max} such that u satisfies the fractional differential equation (1) on $[0, T_{max})$. This means that the solution u is bounded on [0,T] for all $T < T_{max}$, i.e. there exists a continuous function C(t)such that $||u(t)||_{L^{\infty}(\Omega)} \leq C(t)$. Thus

$$\int_{\Omega} u^{-} \mathcal{D}^{\alpha}_{0|t} u^{-} dx \leq \int_{\Omega} (u^{-})^{2} + aC(u^{-})^{2} \left(\int_{0}^{t} K(s) ds \right) dx$$
$$\leq \int_{\Omega} (u^{-})^{2} + aC(u^{-})^{2} \left(\int_{0}^{\infty} K(s) ds \right) dx$$
$$\leq (1 + aC) \int_{\Omega} (u^{-})^{2} dx.$$

where $C = \sup_{t \in [0,T]} C(t)$. Since by Lemma 3, $\frac{1}{2} \mathcal{D}^{\alpha}_{0|t}(u^-)^2 \leq u^- \mathcal{D}^{\alpha}_{0|t} u^-$, we then have

$$\frac{1}{2} \int_{\Omega} \mathcal{D}^{\alpha}_{0|t}(u^{-})^2 dx \le \left(1 + aC\right) \int_{\Omega} (u^{-})^2 dx.$$

Let $\psi(t) = \int_{\Omega} (u^{-})^{2}$. As $u(x, 0) = u_{0} \ge 0$, then $\psi(0) = \int_{\Omega} u^{-}(x, 0)^{2} dx = 0$. By exchanging the integral and $\mathcal{D}_{0|t}^{\alpha}$ on left-hand side, we get

$$\begin{cases} \mathcal{D}^{\alpha}_{0|t}\psi(t) &\leq 2(1+aC)\psi(t),\\ \psi(0) &= 0. \end{cases}$$

By applying (4), we have

$$\psi(t) \le \psi(0) + 2(1 + aC)I_{0|t}^{\alpha} \psi(t),$$

which by Lemma 4 leads to

$$\psi(t) = 0 \quad \Longleftrightarrow \quad \int_{\Omega} (u^{-})^2 dx = 0 \quad \Longrightarrow \quad u^{-} = 0 \quad \Longrightarrow \quad u = u^{+} \ge 0.$$

To prove the boundedness of u, we consider all possible cases. If $0 \le u \le R$ then the proof is done. Otherwise let $u \ge R$. Multiplying equation (1) by u^{p-1} and integrating over Ω , we get

$$\int_{\Omega} u^{p-1} \mathcal{D}^{\alpha}_{0|t} u \, dx + \int_{\Omega} u^{p-1} (-\Delta_N)^{\sigma} u \, dx = \int_{\Omega} u^{p-1} f(u) \, dx.$$

For u > R, we have $u + au^2 - bu^3 < 0$. As for the convolution term we have

$$\mathcal{K} * u \ge \mathcal{K} * R \implies u\mathcal{K} * u \ge R\mathcal{K} * R \implies -au\mathcal{K} * u \le -aR\mathcal{K} * R < 0.$$

Hence, by using Lemma 3 and (6), we obtain

$$\frac{1}{p} \int_{\Omega} \mathcal{D}_{0|t}^{\alpha}(u^{p}) \, dx + \frac{4(p-1)}{p^{2}} \int_{\Omega} \left| (-\Delta_{N})^{\frac{\sigma}{2}} u^{\frac{p}{2}} \right|^{2} \, dx \le 0;$$

whereupon,

$$\int_{\Omega} \mathcal{D}^{\alpha}_{0|t}(u^p) \, dx \le 0.$$

Integrating with respect to $t \alpha$ -times and using (4), we obtain

$$\int_{\Omega} u^p \, dx \le \int_{\Omega} u_0^p \, dx.$$

Taking $p \to \infty$, we get $||u||_{L^{\infty}(\Omega)} \leq ||u_0||_{L^{\infty}(\Omega)}$.

4.1. Uniform Continuity of u. To prove the uniform continuity of the solution of (1)-(3), we notice that A is a sectorial operator (see Gal and Warma [13]). The following lemma shall be useful (see Guswanto and Suzuki [18]).

Lemma 9. [18, Theorems 2.1, 2.2] Let A be a sectorial operator and let $S_{\alpha}(t)$ and $P_{\alpha}(t)$ be the operators defined by (5) and (6), respectively. Then, the following statements hold:

(i) $S_{\alpha}(t), P_{\alpha}(t) \in B((0, \infty); H^{\sigma}(\Omega))$ for t > 0, and if $x \in H^{\sigma}(\Omega)$ then

(8)
$$AS_{\alpha}(t)x = S_{\alpha}(t)Ax, \quad AP_{\alpha}(t)x = P_{\alpha}(t)Ax;$$

- (ii) the functions $t \mapsto S_{\alpha}(t), t \mapsto P_{\alpha}(t)$ belong to $C^{\infty}((0,\infty); B(H^{\sigma}(\Omega)))$;
- (iii) for $x \in H^{\sigma}(\Omega)$ and t > 0,

(9)
$$\frac{d}{dt}S_{\alpha}(t)x = AP_{\alpha}(t)x.$$

For $0 < \beta < 1$, we consider the fractional power $A^{-\beta}$ (see Henry [19])

$$A^{-\beta} = \frac{1}{\Gamma(\beta)} \int_0^t t^{\beta-1} e^{-At} dt,$$

and A^{β} is considered as the inverse of $A^{-\beta}$ with $D(A^{\beta}) = R(A^{-\beta})$. Moreover, for a fixed p > d, the following embedding holds true:

(10)
$$D(A^{\beta}) \hookrightarrow C^{\nu}(\Omega), \quad \text{if } 0 \le \nu < 2\beta - \frac{d}{p};$$

with $2\beta - \frac{d}{p} < 1$.

In the next lemma, we consider some estimates involving the fractional operator A^{β} and the families of operators $\{S_{\alpha}(t)\}_{t>0}, \{P_{\alpha}(t)\}_{t>0}$ (see Guswanto and Suzuki [18]).

Lemma 10. [18, Theorem 2.6] For each $0 < \beta < 1$, there exist positive constants $C_1 = C_1(\alpha, \beta)$ and $C_2 = C_2(\alpha, \beta)$ such that, for all $x \in H^{\sigma}(\Omega)$,

(11)
$$\begin{aligned} \left\| A^{\beta} S_{\alpha}(t) x \right\|_{H^{\sigma}(\Omega)} &\leq C_{1} t^{-\alpha} \left(t^{\alpha(1-\beta)} + 1 \right) \| x \|_{H^{\sigma}(\Omega)}, \quad t > 0, \\ \left\| A^{\beta} P_{\alpha}(t) x \right\|_{H^{\sigma}(\Omega)} &\leq C_{2} t^{\alpha(1-\beta)-1} \| x \|_{H^{\sigma}(\Omega)}, \quad t > 0. \end{aligned}$$

Theorem 3. For any $\delta > 0$, the function $t \mapsto u(x,t)$ is uniformly continuous with respect to $t \in [\delta, \infty)$ in the topology of $L^2(\Omega)$.

Proof. Applying A^{β} to u(x, t+h) - u(x, t) of (7) for any h > 0 gives

$$\begin{split} A^{\beta}u(x,t+h) - A^{\beta}u(x,t) &= A^{\beta} \left(S_{\alpha}(t+h) - S_{\alpha}(t) \right) u_{0} \\ &+ \int_{0}^{t+h} A^{\beta}P_{\alpha}(t+h-\tau)f(x,\tau,u(\tau)) \ d\tau \\ &- \int_{0}^{t} A^{\beta}P_{\alpha}(t-\tau)f(x,\tau,u(\tau)) \ d\tau \\ &= A^{\beta} \left(S_{\alpha}(t+h) - S_{\alpha}(t) \right) u_{0} \\ &+ \int_{t}^{t+h} A^{\beta}P_{\alpha}(t+h-\tau)f(x,\tau,u(\tau)) \ d\tau \\ &- \int_{0}^{t} A^{\beta} \left(P_{\alpha}(t+h-\tau) - P_{\alpha}(t-\tau) \right) f(x,\tau,u(\tau)) \ d\tau. \end{split}$$

The first term tends to 0 as $h \to 0$, since

$$\begin{split} \|A^{\beta} \left(S_{\alpha}(t+h) - S_{\alpha}(t) \right) u_{0} \|_{L^{2}(\Omega)} &= \|A^{\beta}hS_{\alpha}'(t+\delta h)u_{0}\|_{L^{2}(\Omega)} \\ &= h\|A^{\beta}AP_{\alpha}(t+\delta h)u_{0}\|_{L^{2}(\Omega)} \\ &= h\|A^{\beta}P_{\alpha}(t+\delta h)Au_{0}\|_{L^{2}(\Omega)} \\ &= hC_{2}(\alpha,\beta)(t+\delta h)\|Au_{0}\|_{L^{2}(\Omega)}, \end{split}$$

for $\delta \in (0, 1)$, where we have used (8), (9) and (11), respectively.

For the second term, we observe that

$$\begin{split} \left\| \int_{t}^{t+h} A^{\beta} P_{\alpha}(t+h-\tau) f(u(\tau)) d\tau \right\|_{L^{2}(\Omega)} \\ &= \left\| \int_{0}^{h} A^{\beta} P_{\alpha}(\tau) f(u(t+h-\tau)) d\tau \right\|_{L^{2}(\Omega)} \\ &\leq \int_{0}^{h} \left\| A^{\beta} P_{\alpha}(\tau) f(u(t+h-\tau)) \right\|_{L^{2}(\Omega)} d\tau. \end{split}$$

Note that, by Theorem 2, u is bounded by R. Hence, for t > 0, we have

$$\begin{split} \|f(u(\tau)\|_{L^{2}(\Omega)} &\leq \|u+au^{2}-bu^{3}\|_{L^{2}(\Omega)} + \left\|au\int_{0}^{t}K(t-s)u(x,s)ds\right\|_{L^{2}(\Omega)} \\ &\leq R+2aR^{2}+bR^{3}:=R^{*}. \end{split}$$

Therefore, using (11), we have

$$\int_{0}^{h} \left\| A^{\beta} P_{\alpha}(\tau) f(u(t+h-\tau)) \right\|_{L^{2}(\Omega)} d\tau$$

$$\leq \int_{0}^{h} C_{2} \tau^{\alpha(1-\beta)-1} \left\| f(u(t+h-\tau)) \right\|_{L^{2}(\Omega)} d\tau$$

$$\leq \frac{C_{2} R^{*}}{\alpha(1-\beta)} t^{\alpha(1-\beta)} \Big|_{0}^{h} = \frac{C_{2} R^{*}}{\alpha(1-\beta)} h^{\alpha(1-\beta)},$$

which tends to 0 as $h \to 0$.

Since

$$\begin{split} \left\| A^{\beta} \left(P_{\alpha}(t+h-\tau) - P_{\alpha}(t-\tau) \right) f(u(\tau)) \right\|_{L^{2}(\Omega)} \\ &\leq \left\| A^{\beta} P_{\alpha}(t+h-\tau) f(u(\tau)) \right\| + \left\| A^{\beta} P_{\alpha}(t-\tau) f(u(\tau)) \right\|_{L^{2}(\Omega)} \\ &\leq C_{2} R^{*} \left((t+h-\tau)^{-\alpha(\beta-1)-1} + (t-\tau)^{-\alpha(\beta-1)-1} \right), \end{split}$$

and for any h > 0, we have

$$C_{2}R^{*} \int_{0}^{t} (t+h-\tau)^{-\alpha(\beta-1)-1} + (t-\tau)^{-\alpha(\beta-1)-1} d\tau$$

= $\frac{C_{2}R^{*}}{\alpha(1-\beta)} \left[(t+h-\tau)^{\alpha(1-\beta)} + (t-\tau)^{\alpha(1-\beta)} \right]_{0}^{t}$
 $\leq \frac{C_{2}R^{*}}{\alpha(1-\beta)} \left[h^{\alpha(1-\beta)} + (t+h)^{\alpha(1-\beta)} + t^{\alpha(1-\beta)} \right]$
 $< \infty.$

Thus, by the dominated convergence theorem, we have

(12)
$$\lim_{h \to 0} \left\| \int_0^t A^\beta \left(P_\alpha(t+h-\tau) - P_\alpha(t-\tau) \right) f(u(\tau)) \, d\tau \right\|_{L^2(\Omega)} = 0.$$

In view of (10) and (12), we have

$$\lim_{h \to 0} \|u(x, t+h) - u(x, t)\|_{L^2(\Omega)} = 0.$$

5. Asymptotic stability of the positive equilibrium

In this section, we prove the global attractivity of the positive equilibrium. We denote by $\omega(u_0)$ the ω -limit set for u_0 :

$$\omega(u_0) = \{ \varphi \in H^{\sigma}(\Omega) \cap L^{\infty}(\Omega) \ s.t \ \exists t_n \to \infty, \ u(\cdot, t_n) \to \varphi(\cdot) \ \text{in} \ L^2(\Omega) \}.$$

Due to the compact embedding $C^{\nu}(\Omega) \hookrightarrow C^{\mu}(\Omega)$ for $0 < \mu < \nu < 1$, and the boundedness of the solution, the set $\omega(u_0)$ is nonempty as the orbit $\bigcup_{t\geq 0} \{u(t)\}$ is

precompact in $C^{\mu}(\overline{\Omega})$. We shall see that any non-negative $\varphi \in \omega(u_0)$ solves the limiting problem associated to problem (1)-(2), namely

(13)
$$(-\Delta_N)^{\sigma}\varphi = \varphi(1 - b\varphi^2) \quad \text{in } \Omega.$$

Our result is contained in the following theorem:

Theorem 4. Let $u \in L^{\infty}(\Omega \times (0,1))$ be a weak solution of the problem (1)-(2). Then u tends to the constant stationary solution of the elliptic problem (13) uniformly, that is, for any $\varphi \in \omega(u_0)$

$$(-\Delta_N)^{\sigma}\varphi = \varphi(1 - b\varphi^2)$$
 in Ω

Moreover, $\varphi = 1/\sqrt{b}$.

Proof. Let $\xi \in C^1(\overline{\Omega})$ be such that $\nabla \xi \cdot \eta = 0$ on $\partial \Omega$, and $\rho \in C_c^1(\mathbb{R})$ be such that supp $\rho \subset [0,1]$. Next, for any fixed $n \in \mathbb{N}$, we let $v(x,t) = \rho(t-t_n)\xi(x)$. Furthermore, we assume that ρ satisfies the following conditions

(14)
$$\rho(t-t_n) := (I_{t|t_n+1}^{\alpha} \hat{\rho}')(t-t_n), \text{ for some } \hat{\rho} \text{ with } \hat{\rho}(0) = \hat{\rho}(1);$$

(15) $\tilde{\rho}(t-t_n) := (I_{t|t_n+1}^{1-\alpha} \rho)(t-t_n), \qquad \tilde{\rho}(0) = \tilde{\rho}(1) = 0.$

Multiplying equation (1) by v and integrating over $(t_n - 1, t_n + 1) \times \Omega$ gives

$$\mathcal{I}+\mathcal{J}=\mathcal{K},$$

where,

$$\mathcal{I} := \int_{t_n}^{t_n+1} \int_{\Omega} v \ \mathcal{D}_{0|t}^{\alpha} u(t) \ dx \ dt;$$
$$\mathcal{J} := \int_{t_n}^{t_n+1} \int_{\Omega} v \ (-\Delta_N)^{\sigma} u \ dx \ dt;$$
$$\mathcal{K} := \int_{t_n}^{t_n+1} \int_{\Omega} v \ f(u) \ dx \ dt.$$

First, we prove that the first term in the left-hand side, i.e \mathcal{I} tends to zero as $n \to \infty$. Observe that using fractional integration by parts given in Lemma 7, we have

$$\mathcal{I} = \int_{t_n}^{t_n+1} \rho(t-t_n) \, \mathcal{D}_{0|t}^{\alpha} u(t) \, dt$$

= $\int_{t_n}^{t_n+1} u(t) \, {}^{RL} D_{t|t_n+1}^{\alpha} \rho(t-t_n) \, dt \, + \left[u(t) \, I_{t|t_n+1}^{1-\alpha} \rho(t-t_n) \right]_{t=t_n}^{t=t_n+1}.$

From (15), we can see that ρ is chosen such that $\tilde{\rho}(t - t_n) = (I_{t|t_n+1}^{1-\alpha}\rho)(t - t_n)$ satisfies $\tilde{\rho}(t_n) = \tilde{\rho}(t_n + 1) = 0$, thus we have

$$\mathcal{I} = \int_{t_n}^{t_n+1} u(t) \,^{RL} D_{t|t_n+1}^{\alpha} \rho(t-t_n) \, dt.$$

Notice that, $\binom{RL}{t}D_{t|t_n+1}^{\alpha}\rho(t-t_n) = \binom{RL}{t}D_{t|t_n+1}^{\alpha}I_{t|t_n+1}^{\alpha}\hat{\rho}'(t-t_n) = \hat{\rho}'(t-t_n)$. Thus, we have

$$\mathcal{I} = \int_{t_n}^{t_n+1} u(t) \ ({}^{RL}D^{\alpha}_{t|t_n+1}I^{\alpha}_{t|t_n+1}\hat{\rho}')(t-t_n) \ dt$$
$$= \int_{t_n}^{t_n+1} u(t) \ \hat{\rho}'(t-t_n) \ dt$$
$$(16) \qquad = \int_0^1 u(\tau+t_n) \ \hat{\rho}'(\tau) \ d\tau,$$

where we have used the change of variable $\tau = t - t_n$. Since u(x, t) is uniformly continuous for $t \in [\delta, \infty)$ for $\delta > 0$, by Lebesgue's dominated convergence theorem, we conclude that the integrand of the right-hand side of (16) is bounded and goes to zero for large n. Indeed,

$$\lim_{n \to \infty} \int_0^1 u(\tau + t_n) \ \hat{\rho}'(\tau) \ d\tau = u_\infty(\hat{\rho}(1) - \hat{\rho}(0)) = 0$$

For the remaining part of the equation, using Lemma 5, we notice that \mathcal{J} can be written as

$$\mathcal{J} = \int_{t_n}^{t_n+1} \rho(t-t_n) \int_{\Omega} u \ (-\Delta_N)^{\sigma} \xi(x) \ dx \ dt.$$

Hence, $\mathcal{J} + \mathcal{K} = 0$ is equivalent to

$$\int_{-1}^{1} \rho(\tau) \ d\tau \left(\int_{\Omega} u(x,\tau+t_n) \ (-\Delta_N)^{\sigma} \xi(x) - \xi(x) f(u(x,\tau+t_n)) \ dx \right) = 0.$$

Using Lebesgue's theorem, taking the limit as $n \to \infty$, we have

$$\int_{-1}^{1} \rho(\tau) \ d\tau \left(\int_{\Omega} u_{\infty} \ (-\Delta_N)^{\sigma} \xi(x) - \xi(x) f(u_{\infty}) \ dx \right) = 0,$$

where we have used the uniform continuity result of Theorem 3. Consequently, using (14), we get

$$\int_{\Omega} u_{\infty} (-\Delta_N)^{\sigma} \xi(x) - \xi(x) f(u_{\infty}) dx = 0,$$

or

$$\int_{\Omega} \xi(x) \ (-\Delta_N)^{\sigma} u_{\infty} - \xi(x) f(u_{\infty}) \ dx = 0.$$

which is the variational formulation of the elliptic problem (13).

To show that u tends to $u_{\infty} = 1/\sqrt{b}$, we multiply (13) by $(1 - \sqrt{b}u_{\infty})$ and integrate over Ω to get

$$\int_{\Omega} (-\Delta_N)^{\sigma} u_{\infty} (1 - \sqrt{b} u_{\infty}) dx = \int_{\Omega} u_{\infty} (1 + \sqrt{b} u_{\infty}) (1 - \sqrt{b} u_{\infty})^2 dx.$$

Using integration by parts, we get

$$\int_{\Omega} (-\Delta_N)^{\sigma/2} u_{\infty} (-\Delta_N)^{\sigma/2} (1 - \sqrt{b}u_{\infty}) \, dx = \int_{\Omega} u_{\infty} (1 + \sqrt{b}u_{\infty}) (1 - \sqrt{b}u_{\infty})^2 \, dx,$$

or,

$$-\sqrt{b}\int_{\Omega} |(-\Delta_N)^{\sigma/2} u_{\infty}|^2 dx = \int_{\Omega} u_{\infty} (1+\sqrt{b}u_{\infty})(1-\sqrt{b}u_{\infty})^2 dx$$

As we have established the positivity of our solution, we must have

$$\int_{\Omega} |(-\Delta_N)^{\sigma/2} u_{\infty}|^2 dx = 0 \quad \text{and} \quad \int_{\Omega} u_{\infty} (1 + \sqrt{b} u_{\infty}) (1 - \sqrt{b} u_{\infty})^2 dx = 0.$$

This means

 $(-\Delta_N)^{\sigma/2}u_\infty = 0 \implies u_\infty \text{ is a constant},$

and from $\int_{\Omega} u_{\infty}(1+\sqrt{b}u_{\infty})(1-\sqrt{b}u_{\infty})^2 dx = 0$, we have

$$u_{\infty}(1+\sqrt{b}u_{\infty})(1-\sqrt{b}u_{\infty})^2 = 0 \implies u_{\infty} = 0 \text{ or } u_{\infty} = 1/\sqrt{b}.$$

Lastly, we show that $u_{\infty} = 0$ is unstable. Let $u = u_{\infty} + \bar{u} = \bar{u}$. Substituting in (1) we get the following linearized equation

$$\mathcal{D}_{a|t}^{\alpha}\bar{u} + (-\Delta_N)^{\sigma/2}\bar{u} = \bar{u}.$$

Whose solution is given by

$$\bar{u}(x,t) = E_{\alpha} \left(t^{\alpha} \mathcal{L} \right) u_0(x); \qquad \mathcal{L} = I - (-\Delta_N)^{\sigma/2}.$$

If there exist $\lambda \in spec(\mathcal{L})$ with $\arg(\lambda) \leq \alpha \pi/2$ then $\bar{u}(x,t) \not\rightarrow 0$ or $u_{\infty} = 0$ is unstable. In our case, it is sufficient to show the existence of positive $\lambda \in spec(\mathcal{L})$.

To do that consider the eigenvalue problem

(17)
$$\left(1 - (-\Delta_N)^{\sigma/2}\right)\psi = \lambda\psi.$$

If there exist $\lambda > 0$ such that $(1 - \lambda) > 0$ then the zero solution is unstable. Indeed it is the case; we see that $\lambda = 1$ is the eigenvalue that corresponds to the eigenfunction $\psi = const$ of the problem (17). Hence our assertion follows.

Acknowledgments

The authors would like to express their gratitude to Khalifa University of Science and Technology for their continuous support and resources that made this research possible.

References

- N. Britton, Spatial structures and periodic travelling waves in an integro-differential reaction-diffusion population model, SIAM Journal on Applied Mathematics, vol. 50, Dec. 1990. doi:10.1137/0150099.
- [2] J. M. Cushing, Integrodifferential Equations and Delay Models in Population Dynamics, Lecture Notes in Mathematics, vol. 20, Springer Berlin, Heidelberg, New York, 1977.
- C. Pozrikidis, The Fractional Laplacian, CRC Press, Taylor & Francis Group, 2016. URL: https://books.google.ae/books?id=02lLjgEACAAJ.
- [4] S. Gourley and N. Britton, On a modified Volterra population equation with diffusion, Nonlinear Analysis: Theory, Methods & Applications, vol. 21, Sep. 1993, pp. 389–395. doi:10.1016/0362-546X(93)90082-4.
- [5] A. Schiaffino, On a diffusion Volterra equation, Nonlinear Analysis: Theory, Methods & Applications, vol. 3, Aug. 1979, pp. 595–600. doi:10.1016/0362-546X(79)90088-9.
- [6] R. Redlinger, On Volterra's population equation with diffusion, SIAM Journal on Mathematical Analysis, vol. 16, Jan. 1985. doi:10.1137/0516008.
- [7] Y. Yamada, On a certain class of semilinear Volterra diffusion equations, Journal of Mathematical Analysis and Applications, vol. 88, Aug. 1982, pp. 433–451. doi:10.1016/0022-247X(82)90205-0.
- [8] Y. Yamada, Asymptotic behavior of solutions for semilinear Volterra diffusion equations, Nonlinear Analysis: Theory, Methods & Applications, vol. 21, Aug. 1993, pp. 227–239. doi:10.1016/0362-546X(93)90113-7.
- [9] Y. Yoshida and Y. Yamada, Asymptotic behavior of solutions for semilinear Volterra diffusion equations with spatial inhomogeneity and advection, Tokyo Journal of Mathematics, vol. 39, Mar. 2016. doi:10.3836/tjm/1459367268.
- [10] M. Guedda, M. Kirane, and N. Tatar, Asymptotic behavior of solutions to a diffusive Volterra equation, Communications in Applied Analysis, vol. 1, 1997, pp. 91–102.
- [11] M. Kirane, Global bounds and asymptotics for a system of reaction-diffusion equations, Journal of Mathematical Analysis and Applications, vol. 138, Mar. 1989, pp. 328–342.
- [12] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, Theory and Applications of Fractional Differential Equations, North-Holland Mathematics Studies, vol. 204, 2006. URL: https://www.sciencedirect.com/science/article/pii/S0304020806800101.
- C. Gal and M. Warma, Fractional-in-Time Semilinear Parabolic Equations and Applications, Springer International Publishing, 2020. URL: https://books.google.ae/books?id=odX-DwAAQBAJ.
- [14] A. Alsaedi, B. Ahmad, and M. Kirane, Maximum principle for certain generalized time and space fractional diffusion equations, Quarterly of Applied Mathematics, vol. 73, no. 1, 2015, pp. 163–175. URL: http://www.jstor.org/stable/43639148.
- [15] K. Diethelm, TheAnalysis ofFractional Differential Equations: AnApplication-Oriented Exposition Using Differential **Operators** ofCaputo Type,Berlin, Lecture Notes in Mathematics, Springer Heidelberg, 2010.URL: https://books.google.ae/books?id=K4vKvX98SM8C.
- [16] V. A. Liskevich and Y. A. Semenov, Some inequalities for sub-Markovian generators and their applications to the perturbation theory, Proceedings of the American Mathematical Society, vol. 119, Jun. 1993, pp. 1171–1177. doi:10.1090/s0002-9939-1993-1160303-9.
- [17] R. Almeida, D. Tavares, and D. F. M. Torres, The Variable-Order Fractional Calculus of Variations, Springer International Publishing, 2019. doi:10.1007/978-3-319-94006-9.

- [18] B. H. Guswanto and T. Suzuki, Existence and uniqueness of mild solutions for fractional semilinear differential equations, Electronic Journal of Differential Equations, Jun. 2015, pp. 1–16.
- [19] D. Henry, Geometric Theory of Semilinear Parabolic Equations, Lecture Notes in Mathematics, Springer Berlin, Heidelberg, 1981. doi: https://doi.org/10.1007/BFb0089647.

(S. Ahmad) DEPARTMENT OF MATHEMATICS, COLLEGE OF COMPUTING AND MATHEMAT-ICAL SCIENCES, KHALIFA UNIVERSITY OF SCIENCE AND TECHNOLOGY, P.O. BOX 127788, ABU DHABI, UAE, ORCID.ORG/0000-0001-7641-7759

Email address: 100059797@ku.ac.ae

(M. Kirane) DEPARTMENT OF MATHEMATICS, COLLEGE OF COMPUTING AND MATHEMAT-ICAL SCIENCES, KHALIFA UNIVERSITY OF SCIENCE AND TECHNOLOGY, P.O. BOX 127788, ABU DHABI, UAE, ORCID.ORG/0000-0002-4867-7542

Email address: mokhtar.kirane@ku.ac.ae URL: https://www.ku.ac.ae/college-people/mokhtar-kirane