SHARP THRESHOLDS FOR SPANNING REGULAR SUBGRAPHS

MAKSIM ZHUKOVSKII

ABSTRACT. We prove that $(1 + o(1))\sqrt{e/n}$ is the sharp threshold for the appearance of the square of a Hamilton cycle in G(n, p), confirming the conjecture of Kahn, Narayanan, and Park. We also find the exact asymptotics of the threshold for the emergence of a spanning subgraph isomorphic to a fixed graph F for a wide family of d-regular graphs F. This family includes almost all d-regular graphs.

1. INTRODUCTION

A graph property is called *increasing* if it is closed under the addition of edges. If \mathcal{Q} is a non-trivial increasing property and $\mathbf{G} \sim G(n, p)$ is a binomial random graph, then, for every fixed n, the function $\mathbb{P}(\mathbf{G} \in \mathcal{Q})$ increases in p. In particular, there exists a unique solution $p_c(\mathcal{Q})$ of the equation $\mathbb{P}(\mathbf{G} \in \mathcal{Q}) = 1/2$, which is called the *probability threshold for* \mathcal{Q} . In 1987, Bollobás and Thomason [6] proved that, for any non-trivial increasing property \mathcal{Q} , whp¹ $\mathbf{G} \in \mathcal{Q}$ if $p \gg p_c(\mathcal{Q})^2$ and whp $\mathbf{G} \notin \mathcal{Q}$ if $p \ll p_c(\mathcal{Q})$.

Since the original paper of Erdős and Rényi [8] the task of determining the asymptotic behaviour of $p_c(Q)$ for increasing properties Q has been a central topic in probabilistic combinatorics. While the asymptotic order of the probability threshold has been determined for many natural increasing graph properties, a general solution remains unknown, and determining the *exact asymptotics* of $p_c(Q)$ is even more challenging. In this paper we address the latter question, explicitly posed in [29, Question 2]. We provide an answer for a class of increasing properties generated by *d*-regular graphs and resolve a notable conjecture in this area regarding the asymptotics of p_c for the appearance of the second power of a Hamilton cycle.

Despite this challenge, for any arbitrary increasing \mathcal{Q} , the value of $p_c(\mathcal{Q})$ can be established up to a log *n*-factor if the so called *expectation threshold* is known — thanks to the remarkable result of Park and Pham [28], which resolved the renowned conjecture of Kahn and Kalai. Let us recall the definition of the expectation threshold and then state the theorem of Park and Pham. Roughly speaking, the expectation threshold $p_e(\mathcal{Q})$ is the moment when the expected number of graphs generating \mathcal{Q} hits 1/2. Formally, $p_e(\mathcal{Q})$ is the maximum p such that there exists a set of graphs \mathcal{Q}' satisfying $\sum_{G \in \mathcal{Q}'} p^{|E(G)|} \leq \frac{1}{2}$ and $\mathcal{Q} \subseteq \langle \mathcal{Q}' \rangle$, where $\langle \mathcal{Q}' \rangle$ is the upwards closure of \mathcal{Q}' . Due to Markov's inequality, $p_e(\mathcal{Q}) \leq p_c(\mathcal{Q})$. Let $n_{\mathcal{Q}}$ be the number of edges in the largest minimal element of \mathcal{Q} .

Theorem 1.1 (Park, Pham [28]). There exists a universal constant C > 0 such that the inequality $p_c(\mathcal{Q}) \leq Cp_e(\mathcal{Q}) \cdot \log n_{\mathcal{Q}}$ holds for any increasing property \mathcal{Q} .

The upper bound in this theorem is tight: there exists an increasing \mathcal{Q} such that $p_c(\mathcal{Q}) = \Theta(p_e(\mathcal{Q}) \cdot \log n_{\mathcal{Q}})$. For example, let \mathcal{Q} be the property of being Hamiltonian. It is a routine

¹With high probability, that is, with probability tending to 1 as $n \to \infty$.

²For positive sequences $a = (a_n, n \in \mathbb{N})$ and $b = (b_n, n \in \mathbb{N})$, we write $a \ll b$ when $\lim_{n\to\infty} a_n/b_n = 0$. In this case, we also write $b \gg a$.

to check that $p_e = \Theta(1/n)$, while $p_c = (1 + o(1))\frac{\ln n}{n}$ [21, 30]. It is fair to say that the main remaining challenge is to classify properties based on the value of the ratio $\frac{p_c(Q)}{p_e(Q)} \ge 1$, see, e.g., [3]. In particular, for which properties Q is $p_c(Q) = (1 + o(1))p_e(Q)$? In this paper, we explore this question for the property of containing a given (unlabelled) regular spanning subgraph.

1.1. Regular spanning subgraphs. Let $d \geq 3$ be a fixed constant. Let F = F(n) be a sequence of *d*-regular graphs on the vertex set $[n] := \{1, \ldots, n\}$. For $\tilde{F} \subset F$, we call a vertex $v \in V(\tilde{F})$ boundary, if it has degree strictly less than *d* in \tilde{F} . We call the vertex boundary of \tilde{F} the set $\partial_v(\tilde{F})$ of all boundary vertices and the edge boundary the set $\partial_e(\tilde{F})$ of all edges of *F* between $\partial_v(\tilde{F})$ and $V(F) \setminus V(\tilde{F})$. Let the increasing property \mathcal{Q}_F be generated by the family \mathcal{F}_n of all isomorphic copies of *F* on [n]. For simplicity, we denote $p_c(F) := p_c(\mathcal{Q}_F)$ and $p_e(F) := p_e(\mathcal{Q}_F)$. We also let X_F be the number of graphs from \mathcal{F}_n that are subgraphs of $\mathbf{G} \sim G(n, p)$. Since

$$\mathbb{E}X_F = |\mathcal{F}_n| p^{dn/2} = \frac{n!}{|\operatorname{Aut}(F)|} p^{dn/2} \le n! \cdot p^{dn/2}$$

we get that $p_c(F) \ge p_e(F) \ge (1 - o(1))(e/n)^{2/d}$. On the other hand, if every subgraph of F has edge boundary of size at least d, then $p_e(F) = O(n^{-2/d})$ since it is not hard to see that \mathcal{F}_n is $O(n^{-2/d})$ -spread, i.e. for every graph $H \subset F$,

 $|\mathcal{F}_n \cap \langle \{H\} \rangle| \le (Cn^{-2/d})^{|E(H)|} \cdot |\mathcal{F}_n|$ for some constant C = C(d) > 0.

Therefore, due to Theorem 1.1, $p_c(F) = O(n^{-2/d} \cdot \log n)$.

Riordan proved [31] that the logarithmic factor can be removed, i.e.

$$p_c(F) = \Theta(p_e(F)) = \Theta(n^{-2/d}), \tag{1}$$

if F satisfies a stronger condition on the edge boundary: every $\tilde{F} \subset F$ with $3 \leq |V(\tilde{F})| = o(\log n)$ satisfies $|\partial_e(\tilde{F})| \geq 2d$. In particular, for powers of a Hamilton cycle³, this result implies the following: for every $k \geq 3$, the threshold probability for containing the k-th power of a Hamilton cycle equals $\Theta(n^{-1/k})$. However, the proof of Riordan does not work for k = 2 since the minimum size of the edge boundary of a non-trivial subgraph of the second power of a Hamilton cycle equals d + 2 = 6 < 2d. We also note that, for a d-regular graph to satisfy the condition of Riordan, d should be at least 4. In [23], Kühn and Osthus proved that $n^{-1/2+o(1)}$ is the threshold probability for containing the second power of a Hamilton cycle and conjectured that the threshold is actually $\Theta(n^{-1/2})$. In [27], Nenadov and Škorić proved the upper bound $n^{-1/2}(\log n)^4$, which was improved to $n^{-1/2}(\log n)^3$ by Fischer, Škorić, Steger, and Trujić in [11], and to $n^{-1/2}(\log n)^2$ in an unpublished work of Montgomery (see [16]). From Theorem 1.1 (as well as from its weaker fraction version [12]), it follows that the threshold is $O(n^{-1/2} \log n)$. The conjecture of Kühn and Osthus was eventually resolved by Kahn, Narayanan, and Park in 2020 [18]. They further conjectured that $p_c(F) = (1 + o(1))\sqrt{e/n}$. This paper resolves the conjecture, see Section 1.3.

On the other hand, if F has many small subgraphs with smaller edge boundaries, then it may happen that $p_c(F) \gg p_e(F)$. For instance, assume that there exists $v = v(n) = o(\log n)$ and a graph H = H(n) on [v] with $\frac{dv}{2} - \frac{d}{2}$ edges such that every vertex of F belongs to a subgraph which is isomorphic to H (clearly, its edge boundary equals d). Then, an increasing

³For a graph G, its k-th power G^k is obtained by adding to G edges between vertices that are at distance at most k in the graph metric induced by G.

(at most logarithmic) factor arises — for constant v see the proof in [32] which can be directly generalised to all $v = o(\log n)$. Actually, if we allow even smaller edge boundaries, then the threshold probability may increase by a power of n, as well as the expectation threshold.

In [7], Chen, Han, and Luo showed that the proof of Kahn, Narayanan, and Park [18] can be generalised to show that (1) holds for all *d*-regular graphs F satisfying $|\partial_e(\tilde{F})| \ge d+1$ for every $\tilde{F} \subset F$ with $3 \le |V(\tilde{F})| \le \varepsilon n$. Notice that, for large enough $|V(\tilde{F})|$, the bound on the edge boundary in this result is weaker than the condition of Riordan. We show that the weakest of the two conditions always implies (1).

Theorem 1.2. Let $d \ge 3$, $\varepsilon > 0$ be constants, and let F = F(n) be a sequence of d-regular graphs on [n], $n \in \mathbb{N}$, such that $|\partial_e(\tilde{F})| \ge d + 1$, for every $\tilde{F} \subset F$ with $3 \le |V(\tilde{F})| \le \varepsilon \ln n$. Then (1) holds.

As follows from the above discussion, our bound on edge boundaries is tight, that is, in some sense, the condition on the edge boundary is the only obstacle in getting the equality (1). Indeed, for every $3 \le v = o(\log n)$ and every $0 \le \ell \le d$ (such that $dv - \ell$ is even), there exists a graph F that have a subgraph on v vertices with edge boundary of size ℓ such that (1) does not hold. The proof of Theorem 1.2 uses similar techniques as in [7, 18] and does not require accurate spread approximations, for this reason we postpone it to Appendix A.

Remark 1.3. Our proof allows to get an explicit upper bound on the threshold probability, which we did not try to optimise. However, for graphs with small number of automorphisms, a refined argument gives a better upper bound: Assume, for instance, that, for every proper connected subgraph of F, the number of automorphisms that preserve boundary vertices is bounded and that every subgraph on at least 3 and at most n/2 vertices has edge boundary of size at least d+1. Then $p_c(F) \leq \left(e^{2/d} + \frac{d}{1+1_d \text{ is even}} - 1 + o(1)\right) n^{-2/d}$. The proof of this bound follows the same lines as the proof of Theorem 1.2, but requires more accurate computations. In particular, in the proof of Theorem 1.2, we do not need very accurate estimates for the number of subgraphs in F and for the number of ways to embed such a subgraph into F, provided by Claim 3.1 in Section 3. However, the refined bound requires the full power of Claim 3.1. For instance, for the square of a Hamilton cycle F, the refined bound implies that the threshold probability $p_c(F)$ belongs to a fairly tight interval $\left[\frac{\sqrt{e}-o(1)}{\sqrt{n}}, \frac{\sqrt{e}+1+o(1)}{\sqrt{n}}\right]$. As we will see further, it actually coincides asymptotically with the left boundary of this interval.

1.2. Sharp thresholds. Let $\mathbf{G} \sim G(n, p)$. The threshold probability $p_c(\mathcal{Q})$ is called *sharp*, if, for every constant $\varepsilon > 0$, the following is true: if $p > (1 + \varepsilon)p_c(\mathcal{Q})$, then whp $\mathbf{G} \in \mathcal{Q}$; if $p < (1 - \varepsilon)p_c(\mathcal{Q})$, then whp $\mathbf{G} \notin \mathcal{Q}$. Friedgut proved [14] that all increasing (and isomorphismclosed) properties that do not have sharp thresholds are essentially determined by the presence of subgraphs of bounded sizes. Omitting technical details, it means that, for some $p = \Theta(p_c(\mathcal{Q}))$, $\mathbf{G} \sim G(n, p)$ sprinkled with $\mathbf{G}' \sim G(n, \varepsilon p)$, for some $\varepsilon > 0$, is less likely to have the property \mathcal{Q} , than the union of \mathbf{G} with an independently sampled random clique of some bounded size (see, e.g., [13, Theorem 2.3]). With this remarkable result, it is fairly straightforward to check that the threshold probabilities in Theorem 1.2 are sharp. To the best of our knowledge, there are no general results describing asymptotics of sharp p_c for a wide class of increasing properties (say, for \mathcal{Q}_F , where F are d-regular). In this paper, we find asymptotics of $p_c(F)$ for most d-regular graphs F. **Theorem 1.4.** Let $d \ge 3$ and let F = F(n) be a sequence of d-regular graphs on [n], $n \in \mathbb{N}$, such that one of the following two conditions holds.

- (1) There exists $\delta \in (0, 1/d)$ and $w = w(n) = \omega(1)$ such that (a) $|\operatorname{Aut}(F)| \le e^{o(n^{1-\delta})}$; and (b) every $\tilde{F} \subset F$ with $3 \le |V(\tilde{F})| \le n-3$ has $|\partial_e(\tilde{F})| \ge d+1 + \left| |V(\tilde{F})| \frac{w}{\ln n} \right| \cdot \mathbb{1}_{\ln n/w \le |V(\tilde{F})| \le n^{1-\delta}}.$
- (2) Let $\gamma \in (0,1)$ be a fixed constant, let $C = C(d,\gamma)$ be large enough, and let $D \ge 1$ be a fixed constant.

(a) for every subgraph $\tilde{F} \subset F$ with $|E(\tilde{F})| \leq Cn^{2/d}$ and $|\partial_v(\tilde{F})| \leq \gamma |V(\tilde{F})|$, the number of automorphisms of \tilde{F} that fix all vertices of $\partial_v(\tilde{F})$ is at most D;

(b) for every subgraph $\tilde{F} \subseteq F$ with $|\partial_v(\tilde{F})| \leq \gamma |V(\tilde{F})|$, the number of automorphisms of \tilde{F} that fix all vertices of $\partial_v(\tilde{F})$ is at most $e^{o(|V(\tilde{F})|)}$;

(c) every
$$F \subset F$$
 has $|\partial_e(F)| \ge 2d$, if $3 \le |V(F)| \le n-3$.

Then

$$p_c(F) = (1 + o(1))p_e(F) = (1 + o(1))\left(\frac{e}{n}\right)^{2/d}$$
 is sharp. (2)

We note that the condition 2.(c) implies $d \ge 4$. Though the second condition in Theorem 1.4 is strictly weaker than the condition of Riordan, observe that the first condition is not such. For instance, a random triangle-free 3-regular graph has linearly many subgraphs on 3 vertices with edge boundary of size 5 < 6 = 2d — so, it does not satisfy the condition of Riordan, while it satisfies 1.(a) and 1.(b) whp, see below.

Note that, for every $d \ge 4$, almost all d-regular graphs satisfy the first condition in Theorem 1.4 (see [5, 19, 24]). Indeed, for every $d \geq 3$, letting \mathbf{G}_d be a random d-regular graph on [n], whp \mathbf{G}_d is asymmetric [24], implying 1.(a). Moreover, whp \mathbf{G}_d has Cheeger constant $\Theta(1)$ and, whenever $d \geq 4$, whp \mathbf{G}_d does not have subgraphs with $3 \leq v \leq n-3$ vertices and the edge boundary of size at most d, implying 1.(b). If d = 3, then whp there are no subgraphs with $3 \le v \le n-3$ vertices and the edge boundary of size at most d other than K_3 and their vertex-complements. Therefore, whp \mathbf{G}_d satisfies the first condition, if it does not contain triangles. Also, k-th powers of Hamilton cycles for every $k \geq 3$ and toroidal square grids $T_{m,n/m} \cong C_m \square C_{n/m}$ for $m \ge 4$ satisfy the second condition as well. It is also easy to see that the second condition implies sharp thresholds for square and triangular lattices (by considering their regular completions). It is worth recalling that thresholds for lattices were investigated in the past: the question of which random graphs contain spanning lattices was raised by Venkatesan and Levin [36]. Thresholds for lattices, up to a polylog-factor, were proved in [10] and, up to a constant factor, were proved in [31] (they also follow from Theorem 1.2 and [7, Corollary 1.5]). Summing up, we get the following corollary that demonstrates that the set of regular graphs F satisfying the restrictions in Theorem 1.4 is quite broad.

Corollary 1.5. The following graphs F satisfy (2):

- k-th power of a cycle, $k \geq 3$,
- toroidal grid $T_{m \times n/m}$, $m \ge 4$ (assuming that n is divisible by m),
- square and triangular lattices with d = 4 and d = 6 respectively.

Moreover, (2) holds for every $d \ge 4$ and (asymptotically) almost all d-regular graphs F on [n] (assuming dn is even) and for (asymptotically) almost all triangle-free 3-regular graphs F on [n] (assuming n is odd).

Remark 1.6. Actually, using our methods, we are able to establish the same sharp threshold for almost all 3-regular graphs — the condition of the absence of triangles is redundant, since the number of triangles converges in probability to a Poisson random variable [37], and so it is bounded in probability. In other words, we may allow F to have a bounded number of subgraphs with a smaller edge boundary. However, we do not want to overload the proof with technical details, and so we formulate Theorem 1.4 as well as Corollary 1.5 in their current forms.

Through private communication, we learned that Tamás Makai, Matija Pasch, Kalina Petrova, and Leon Schiller are independently working on determining the asymptotics of $p_c(F)$ for k-th powers of cycles F. They proved (2) for all $k \ge 4$ and plan to upload this result.

Examples of graphs that do not satisfy restrictions from Theorem 1.4 include the second power of a cycle, toroidal grid $T_{3,n/3}$, and, so-called, 2-overlapping 4-cycles. The latter graph, denoted by $C_{4,n}^e$, is obtained from n/2 cyclically order copies of C_4 , where two consecutive C_4 overlap in exactly one edge, whereby each C_4 overlaps with two copies of C_4 in opposite edges. Recall that Kahn, Narayanan, and Park in [18] proved (1) for the second power of a cycle F, resolving the conjecture of Kühn and Osthus [23]. Moreover, using the same method, Espuny Díaz and Person in [9] proved (1) for $F = C_{4,n}^e$ answering the question of Frieze [15] (we note that (1) for these graphs also follow both from Theorem 1.2 and [7, Corollary 1.5]).

1.3. Second powers of Hamilton cycles. In this section, we present results that complement Theorem 1.4 and resolve the conjecture of Kahn, Narayanan, and Park. The novelty of its proof technique constitutes our main contribution.

In 2020, Kahn, Narayanan, and Park [18] conjectured that (2) holds when F is the second power of a cycle. Due to the natural barrier in the application of the fragmentation technique that was used to prove (1) for the second power of a Hamilton cycle [18], to resolve the conjecture of Kahn and Kalai [28], and to prove the so called spread lemma [2] (see details in Section 1.4), this conjecture attracted significant interest and was reiterated in [16, 29]. In this paper, we prove the conjecture.

Theorem 1.7. Let F = F(n) be the second power of an n-cycle on [n]. Then F satisfies (2).

We then show that the method that we use to prove this conjecture can be used to prove a more general result for all d-regular graphs that have "cyclic" structure.

Theorem 1.8. Let $d \ge 3$ be a constant, $r = r(n) = o(\log n)$, and F = F(n) be a sequence of graphs on [n] satisfying the following.

- Every subgraph $\tilde{F} \subset F$ with $3 \leq |V(\tilde{F})| \leq r$ has $|\partial_e(\tilde{F})| \geq d+1$.
- The bijection that maps each $v \in [n]$ to $v + r^4$ is an automorphism of F.
- Every u, v ∈ [n], such that the distance between u and v in the cyclic order on [n] is more than r, are not adjacent in F.
- There are no automorphisms of F that fix vertices from [r].

Then F satisfies (2).

⁴Vertices here are treated as elements of the cyclic group $(\mathbb{Z}_n, +)$.

In particular, (2) holds for the two other 'naive' cases that Theorem 1.4 does not cover — $F = C_{4,n}^e$ and $F = T_{3,n/3}$.

1.4. **Proof strategy.** The crux of our proofs is the *fragmentation trick* that in different forms appeared in many applications. One of them is the famous *spread lemma* [2] which in particular gives good sunflower bounds [35]; in probabilistic terms the application of the trick for the spread lemma is described in [25]. This trick was also the main ingredient in the proof of Theorem 1.1 in [28] and in the proof of Frankston, Kahn, Narayanan, and Park [12] of Talagrand's fractional version of the conjecture of Kahn and Kalai [34]. Our proofs essentially rely on *typical* fragments, similarly to [18], rather than *minimal* fragments, that were used by Park and Pham in [28] to prove Theorem 1.1 — see below. Furthermore, in order to prove Theorem 1.7, we introduce a novel technique of choosing fragments — we show that the "cyclic" structure of second powers of Hamilton cycles allows to choose rare fragments that are sparser than typical ones, from sufficiently many copies of F.

The key idea from [18] that proves (1) for second powers of cycles is as follows. Fix any $F \in \mathcal{F}_n$, before the edges of the uniformly random graph $\mathbf{W} \sim G(n, m)$ are exposed, where $m = \lfloor p \binom{n}{2} \rfloor$. Consider $F \cup \mathbf{W}$ and show that whp almost every $F' \subset F \cup \mathbf{W}$ from \mathcal{F}_n is such that $|F' \setminus \mathbf{W}| = O(\sqrt{n})$. It means that we may replace almost every $F \in \mathcal{F}_n$ with its fragment $F' \setminus \mathbf{W} \subset F$ for typical $F' \subset F \cup \mathbf{W}$ of size $O(\sqrt{n})$ and expose another $\tilde{\mathbf{W}} \sim G(n, \tilde{m})$ independently where \tilde{m} is large enough to cover at least one fragment whp. Then whp $\mathbf{W} \cup \tilde{\mathbf{W}}$ contains a graph from \mathcal{F}_n .

However, if a fragment contains a subgraph with edge boundary of size d + 2 = 6 with $\Omega(\ln n)$ edges — such a subgraph is exactly the second power of a path, we call it *closed* — then $\tilde{m} = \lceil \delta n \sqrt{n} \rceil$ additional uniformly random edges with a certain bounded from zero $\delta > 0$ is not enough to cover it. Unfortunately, we suspect that typical fragments do contain closed subgraphs of size $\Omega(\ln n)$. So, this approach does not allow to get a sharp bound on the threshold. This is the main complication that does not allow to apply directly the fragmentation technique to resolve the conjecture of Kahn, Narayanan, and Park. A careful implementation of the technique gives upper bound on the threshold $(\sqrt{e} + 1 + o(1))n^{-1/2}$, see Remark 1.3.

Nevertheless, we show that there exists a large family of fragments that do not contain closed subgraphs of size $\Omega(\ln n)$ and which is still $n^{-\gamma}$ -spread for some small enough $\gamma > \frac{1}{2}$. A direct approach would be, for a typical $F \in \mathcal{F}_n$ (in the W-measure), to search for a fragment $H \subset F$ that does not have closed subgraphs — indeed, since typically, the set of fragments, for a given F, is large, we suspect that such a fragment exists. There might be some clever decoupling argument that allows to implement this approach, but we could not find it. Instead, we improve a typical fragment manually: for a fragment $H \subset F$ we distribute evenly vertices of maximal closed subgraphs between them. This is possible due to the following crucial observation. Let $P_1 = (u_1^1 u_2^1 v_1 \dots v_p u_3^1 u_4^1)$ and $P_2 = (u_1^2 u_2^2 v_{p+1} \dots v_{p+p'} u_3^2 u_4^2)$ be disjoint closed subgraphs of a fragment $H \subset F \in \mathcal{F}_n$. Then the graph H' obtained from H by applying any permutation to "internal" vertices v_i of $P_1 \sqcup P_2$ remains a fragment of some other $F' \in \mathcal{F}_n$. More formally,

- let π be any permutation on [p + p'],
- let $t \in \{0, 1, \dots, p + p'\}$, and
- let H' be obtained from H by replacing P_1 with $P_1^{\pi} := (u_1^1 u_2^1 v_{\pi(1)} \dots v_{\pi(t)} u_3^1 u_4^1)$ and P_2 with $P_2^{\pi} := (u_1^2 u_2^2 v_{\pi(t+1)} \dots v_{\pi(p+p')} u_3^2 u_4^2)$.

Then, if $H \cup \mathbf{W}$ contains a graph from \mathcal{F}_n , we get that $H' \cup \mathbf{W}$ contains a graph from \mathcal{F}_n as well. Moreover, H and H' have exactly the same number of edges, vertices, and connected components.

Although it allows to get a graph H' that does not have large closed subgraphs, this graph is no longer a subgraph of F. Since the 'improving' function that maps each F to the modified fragment may have large pre-images, this modification of fragments may affect spreadness properties of the multiset of fragments. We implement a probabilistic approach to show that a suitable way to modify fragments exists: we distribute vertices among closed subgraphs according to a rule that is described by a perfect matching in a binomial random bipartite graph with an appropriate edge probability, sampled independently of $\mathbf{W} \sim G(n,m)$. Using this rule, we prove that the maximum cardinality of a pre-image of the 'improving' function differs by a sufficiently small factor from the maximum cardinality of a pre-image of the original function that maps every F to its fragment.

Another complication is that, in order to implement this improvement, we need to have sufficiently many disjoint maximal closed subgraphs with at least 4 vertices, so that after vertices are distributed evenly between them, each closed subgraph has size $o(\log n)$. Although it seems plausible that typical fragments contain sufficiently many closed subgraphs whp, we did not find a way to show this. Instead, we force typical fragments to contain a fixed set of $\omega(\sqrt{n}/\log n)$ closed subgraphs with 4 vertices — diamonds. This is implemented via restricting the family \mathcal{F}_n to the set of all cycles that contain fixed diamonds on specific positions. Due to symmetry and linearity of expectation, we then are able extend the multiset of fragments to the entire \mathcal{F}_n .

The proof of Theorem 1.4 is much more straightforward — typical fragments are good enough to derive (2). Indeed, both conditions do not permit closed subgraph of size $\Omega(\log n)$. We prove the first part of Theorem 1.4 via $1/\delta$ fragmentation steps. The proof of the second part of Theorem 1.4 requires only two fragmentations. Actually the usual second moment method works in this case — for the uniform model instead of the binomial, as in the paper of Riordan [31]. It is possible to show that $\operatorname{Var} X_F = O((\mathbb{E} X_F)^2)$ and then to apply the powerful results of Friedgut [13, 14]. In particular, this strategy was used by Narayanan and Schacht to determine sharp thresholds for nonlinear Hamiltonian cycles in random hypergraphs [27]. However, we give the proof of the second part of Theorem 1.4 using fragmentation for the sake of convenience and coherence.

We believe that our analysis is essentially optimal and significant improvements of conditions in Theorem 1.4 require new ideas. Indeed, we obtain fairly optimal bounds on the number of graphs $|\mathcal{F}_n \cap \langle I \rangle|$ containing a given set of edges I and on the number of subgraphs of F with fixed numbers of vertices, edges, and components (see Claim 3.1). The main novel ingredient in the proof of Claim 3.1 is a very nice property of d-regular graphs that do not have non-trivial subgraphs with edge boundary of size at most d: for every v, there are 1) constantly many closed subgraphs on v vertices that share a vertex, 2) at most linearly many closed subgraphs on v vertices in F, see Claim 2.2 and Claim 2.3.

Let us finally say a few words about our proof of Theorem 1.2. Kahn, Narayanan, and Park in [18] noted that the key observation enabling their proof that the threshold for the appearance of the second power of a Hamilton cycle F equals $p_c(F) = \Theta(n^{-1/2})$ is that \mathcal{F}_n is $(1+o(1))\sqrt{e/n}$ -spread. They refined the concept of spreadness by incorporating the number of components in a subgraph. This refinement was later distilled by Espuny Díaz and Person in [9] under the name of superspreadness and was used to extend the result of Kahn, Narayanan, and Park to a broader class of spanning subgraphs in G(n, p). Somewhat surprisingly, this property allows to prove the weaker result [7, Corollary 1.5] but does not allow to prove Theorem 1.2, as it treats contributions from connected components of both bounded and growing sizes in the same manner. It is also worth noting that Spiro [33] proposed another generalisation of spreadness, which, in particular, recovers the results from [9] and served as a key ingredient in the proofs of [7].

In our approach, we employ the same fragmentation technique but with a more refined analysis of subgraphs of growing sizes, which enables us to prove Theorem 1.2. Since the proof does not introduce any substantially new ideas, we defer it to Appendix A.

1.5. **Organisation.** Properties of closed subgraphs in *d*-regular graphs F are studied in Section 2. We further use them in Section 3 to get tight estimates on the number of subgraphs and their extensions in F. The fragmentation trick and the main Lemma 4.1, that allows to apply it, are described in Section 4. Section 5 is devoted to the proof of Theorem 1.4. Then, in Section 6 we prove Theorem 1.7. It generalises directly to Theorem 1.8. For the sake of clarity of presentation instead of providing a full proof of the more general result, we give a detailed proof of Theorem 1.7 (in Section 6), which is more transparent and avoids unnecessary technical details, and then sketch the proof of Theorem 1.8 in Section 7. In Section 8 we discuss some remaining challenges. The proof of Theorem 1.2 is presented in Appendix A.

1.6. Notation. For every positive integer n, we denote $[n] := \{1, \ldots, n\}$. The number of edges in the clique on [n] is denoted by $N = N(n) := \binom{n}{2}$. We denote by $\partial_e(\tilde{F})$ and $\partial_v(\tilde{F})$ the edge and the vertex boundaries of a subgraph \tilde{F} in a fixed graph F (see definitions in Section 1.1). For a graph G, we denote its minimum degree, its number of vertices, its number of edges, and its number of connected components by $\delta(G)$, x(G), $\ell(G)$, and c(G), respectively. We also denote the automorphism group of G by $\operatorname{Aut}(G)$. For a set $U \subset V(G)$, we let G[U] be the subgraph of G induced by U. For a vertex $v \in V(G)$, its degree in G is denoted by $\deg_G(v)$.

2. Linearly many closed subgraphs

In this section, we consider *d*-regular graphs with good enough expansion properties (edge boundaries have sizes at least d + 1) and prove that they have a limited amount of closed subgraphs. Two main results of this section — Claims 2.2 and 2.3 are used in Section 3 to prove Claim 3.1. The latter claim is essential in the proof of Theorem 1.4.

Let us call a graph F locally sparse if the edge boundary of every subgraph $\tilde{F} \subset F$ with $3 \leq |V(\tilde{F})| \leq n-3$ is of size at least d+1. Clearly d+1 can be replaced with d+2 for even d since in this case $|\partial_e(\tilde{F})|$ cannot be odd. Let $\Delta := d+1$ for odd d and $\Delta := d+2$ for even d. It is easy to see that the condition $|\partial_e(\tilde{F})| \geq \Delta$ holds for all \tilde{F} with $2 \leq |V(\tilde{F})| \leq d-1$ just due to the d-regularity of F. Let us call an *induced* subgraph \tilde{F} of a locally sparse d-regular graph F with the edge boundary of size exactly Δ closed (note that a closed subgraph is always connected — otherwise, it has a connected component with a smaller boundary).

Let F be a locally sparse d-regular graph on [n].

Claim 2.1. Every closed subgraph of F with at least 3 vertices has minimum degree at least d/2.

Proof. Assume that \tilde{F} is a closed subgraph of F with at least 3 vertices and with a vertex w having degree d' < d/2. If we remove the vertex w from \tilde{F} , then we get the graph $\tilde{F} \setminus w$ with

edge boundary of size $|\partial_e(\tilde{F})| + 2d' - d < |\partial_e(\tilde{F})| = \Delta$. This contradicts the local sparsity of F when $|V(\tilde{F})| \ge 4$. Otherwise it contradicts the fact that a subgraph on 2 vertices has the edge boundary of size at least $2d - 2 \ge \Delta$.

Claim 2.2. For any pair of adjacent vertices x, y in F and for every $3 \le v \le n-3$, there are at most two closed subgraphs in F on v vertices containing x and not containing y.

Proof. Fix adjacent vertices x, y and $3 \le v \le n-3$.

A closed subgraph $\tilde{F} \subset F$ sends exactly Δ edges to $F \setminus \tilde{F}$ implying that $F \setminus \tilde{F}$ is also closed. Assume that $v \ge n/2$, and that there are at least 3 closed graphs on v vertices that share x and do not contain y. Then their complements are closed graphs on $n - v \le n/2$ vertices that share y and do not contain x. Therefore, it suffices to prove the claim for $v \le n/2$.

Let H_1, H_2 be different closed subgraphs of F on v vertices that contain x and do not contain y. Note that H_1, H_2 should have at least one other common vertex since otherwise the degree of x is bigger than d due to Claim 2.1. Then $|V(H_1) \cup V(H_2)| \le n-2$.

Let $H_0 = H_1 \cap H_2$. Note that $|E(H_0)| \leq \frac{d}{2}|V(H_0)| - \frac{\Delta}{2}$ implying that $|E(H_j) \setminus E(H_0)| \geq \frac{d}{2}|V(H_j \setminus H_0)|$ for both j = 1 and j = 2 since H_1, H_2 are closed. On the other hand, if, say $|E(H_2) \setminus E(H_0)| > \frac{d}{2}|V(H_2 \setminus H_0)|$, then $|E(H_1 \cup H_2)| > \frac{d}{2}|V(H_1 \cup H_2)| - \frac{\Delta}{2}$ which contradicts the local sparsity of F since $|V(H_1 \cup H_2)| \leq n-2$. Therefore, $|E(H_j) \setminus E(H_0)| = \frac{d}{2}|V(H_j \setminus H_0)|$ for both j = 1 and j = 2, but then $|E(H_0)| = \frac{d}{2}|V(H_0)| - \frac{\Delta}{2}$, i.e. H_0 is closed.

Then, there are exactly Δ edges between H_0 and $F \setminus H_0$, and one of them is the edge between x and y. It means that $H_j \setminus H_0$, $j \in \{1, 2\}$, send at most $\Delta - 1$ edges (in total) to H_0 . This may happen only if $|V(H_j \setminus H_0)| = 1$ for both j = 1 and j = 2. Indeed, $|V(H_1 \setminus H_0)| = |V(H_2 \setminus H_0)|$. Moreover, the number of edges that $H_j \setminus H_0$ sends to H_0 equals

$$|E(H_j) \setminus E(H_0)| - |E(H_j \setminus H_0)| \ge \frac{d}{2}|V(H_j \setminus H_0)| - \left(\frac{d}{2}|V(H_j \setminus H_0)| - \frac{\Delta}{2}\right) = \frac{\Delta}{2}$$

whenever $|V(H_j \setminus H_0)| \ge 2$.

Assume that there exists a closed graph $H_3 \not\subset H_1 \cup H_2$ on v vertices that contains x and does not contain y. From the above it follows that $|V(H_1) \cap V(H_3)| = |V(H_2) \cap V(H_3)| = v - 1$. If $H_0 \not\subset H_3$, then H_3 has to contain both vertices from $(V(H_1) \cup V(H_2)) \setminus V(H_0)$. Therefore, there are at least two vertices in $V(H_0) \setminus V(H_3)$ and then $|V(H_1) \cap V(H_3)| \leq v - 2$ — a contradiction. We get $H_3 \cap H_1 = H_3 \cap H_2 = H_0$. Each vertex of $H_j \setminus H_0$, $j \in \{1, 2, 3\}$, sends at least $\frac{d}{2}$ edges to H_0 due to Claim 2.1. But then the vertices from $H_j \setminus H_0$ send at least $\frac{3d}{2} \geq \Delta$ edges to H_0 — contradiction again, since there is one additional edge $\{x, y\}$ in the edge boundary of H_0 .

Therefore, any other closed graph that contains x and does not contain y should be entirely inside $H_1 \cup H_2$. Assume that such a graph H_3 exists. Let $w_1 \in H_1 \setminus H_0$, $w_2 \in H_2 \setminus H_0$. Clearly, H_3 contains w_1, w_2 and all but one vertex of H_0 . In the same way as above we get that $H_1 \cap H_2 = H_0$, $H_1 \cap H_3$ and $H_2 \cap H_3$ are three closed graphs on v - 1 vertices that contain xand do not contain y. These three closed graphs on v - 1 vertices have the property that none of them is inside the union of the other two — this is only possible when v - 1 = 2, i.e. v = 3. The only possible closed graph on 3 vertices is a triangle. Moreover, a triangle is closed only when d = 4. So, H_1, H_2 are triangles sharing an edge, but then H_3 adds another edge to the union $H_1 \cup H_2$ implying that $H_1 \cup H_2 \cup H_3$ is a 4-clique. We get a contradiction with the local sparsity since the edge boundary of a 4-clique in a 4-regular graph is of size $4 < \Delta = 6$.

From this, it immediately follows that, for every v, there are at most 2dn closed subgraphs on v vertices in F — since F is connected, every subgraph contains a vertex that is incident to an edge than does not belong to this subgraph. The following claim gives a slightly better bound.

Claim 2.3. Let $k \in \mathbb{N}$, and let F' := F[[k]] be the induced subgraph of F on [k]. For every $3 \le v \le n-3$, the number of closed subgraphs of F' with v vertices is at most $\frac{2dk}{3}$.

Proof. Fix a vertex w in F' and let us bound the number — denoted by $\mu(w)$ — of closed subgraphs $\tilde{F} \subset F'$ on v vertices containing w such that $\deg_{\tilde{F}} w < d$. Due to Claim 2.2, $\mu(w) \leq d$ 2d. On the other hand, Claim 2.1 implies that every closed subgraph has vertex boundary of size at least 3. Letting f to be the number of closed subgraphs in F' on v vertices, by double counting, we get that $3f \leq \sum_{w \in V(F')} \mu(w) \leq 2dk$ as needed.

3. SUBGRAPHS AND SPREAD

In this section we estimate two quantities: 1) the number of subgraphs with given numbers of vertices, edges, and connected components in a fixed d-regular locally sparse graph F, 2) the number of ways to extend such a subgraph to an isomorphic copy of F. It is split into two subsections: In Section 3.1, we prove Claim 3.1, that gives fairly tight estimations of both quantities. These estimations are used in the first fragmentation step in the proof of Theorem 1.4. Section 3.2 proves much more straightforward and coarse bounds that hold for any d-regular graph and that we use in later fragmentation steps as well as in the proof of Theorem 1.2 in Appendix A.

Let F be an arbitrary d-regular locally sparse graph on [n]. Let \mathcal{F}_n be the set of all isomorphic copies of F on [n]. Fix $\ell \in [dn/2]$, $c \in [\ell]$, and $x \in \left[\frac{2\ell}{d} + \frac{\Delta}{d}c, \ell + c\right]$. The following quantity will play a crucial role in our proofs:

$$\sigma := \frac{d}{2}x - \left(\ell + \frac{\Delta}{2}c\right).$$

We will call σ the *excess* of a graph with x vertices, ℓ edges, and c connected components. It measures the edit distance to the closest union of c closed subgraphs on x vertices.

3.1. Sharp estimates. This section is devoted to the proof of the following claim.

Claim 3.1. There exist constants $A^1_{\alpha}, A^2_{\alpha} > 0$ such that the number of subgraphs in F without isolated vertices with x vertices, ℓ edges, and c components is

$$\alpha(\ell, x, c) \le \binom{n}{c} \binom{x}{c} e^{A_{\alpha}^{1}c + A_{\alpha}^{2}\sigma} \max_{o \le (\Delta+2)\sigma} \binom{x}{o}.$$
(3)

Moreover, there exist constants $A^1_{\beta}, A^2_{\beta} > 0$ such that, given $H \subset F$ with x vertices, ℓ edges, and c components, the number of ways to extend H to a graph from \mathcal{F}_n is at most

$$\beta(\ell, x, c) = (n - x + c)! e^{A_{\beta}^{1} c + A_{\beta}^{2} \sigma} \min\left\{ \left(\frac{x}{c}\right)^{c} (d - 1)^{x} / |\operatorname{Aut}(F)|, 1 \right\} \max_{o \le (\Delta + 2)\sigma} \binom{x}{o}.$$
(4)

Proof. We first prove (40). There are at most $\binom{x}{c}\binom{\sigma+c}{c}$ ways to choose positive integers ℓ_1, \ldots, ℓ_c and x_1, \ldots, x_c such that

- $\frac{2\ell_i}{d} + \frac{\Delta}{d} \le x_i \le \ell_i + 1$ for all $i \in [c]$, $\sum_{i=1}^c \ell_i = \ell$, and $\sum_{i=1}^c x_i = x$.

For every $i \in [c]$, fix such x_i and ℓ_i , and set

$$\sigma_i = \frac{d}{2}x_i - \left(\ell_i + \frac{\Delta}{2}\right). \tag{5}$$

We first choose closed subgraphs of F that correspond to $\sigma_i = 0$ one by one. The number of ways to choose the first closed subgraph is at most $\frac{2d}{3} \cdot n$, due to Claim 2.3 (if $\ell_1 = 1$, then the number of choices is $\frac{dn}{2} < \frac{2dn}{3}$). In the same way, if some set of \tilde{n} vertices is already included in the subgraph under construction, then the next closed component can be chosen in at most $\frac{2d}{3}(n-\tilde{n})$ ways.

We now switch to not closed components. Note that the *i*-th component z_i has vertex boundary of size $o_i \leq \Delta + 2\sigma_i$. Indeed, let ℓ'_i be the number of edges in $E(F[V(z_i)]) \setminus E(z_i)$, that is the number of missing edges in z_i . Then $dx_i = 2\ell_i + 2\ell'_i + |\partial_e(z_i)|$, implying $|\partial_e(z_i)| + 2\ell'_i = \Delta + 2\sigma_i$ due to (5). Since every missing edge in z_i is incident to two boundary vertices, we get that $|\partial_v(z_i)| \leq |\partial_e(z_i)| + 2\ell'_i$. The desired bound on o_i follows. Assume that \tilde{n} vertices have been already included in the subgraph and we now describe the procedure of choosing the *i*-th component z_i :

- (1) choose the size of the vertex boundary $o_i \leq \Delta + 2\sigma_i \leq (\Delta + 2)\sigma_i \leq (\Delta + 2)^{\sigma_i}$;
- (2) choose a set $\mathcal{O} \in {[x_i] \choose o_i}$ that identifies the labels of boundary vertices in the *i*-th component;
- (3) choose a vertex w out of the set of remaining $n \tilde{n}$ vertices and *activate* it we treat this vertex as the minimum vertex in the component under construction;
- (4) at every step $j \ge 1$, consider the minimum vertex v_j among active vertices:
 - if $j \in \mathcal{O}$ (i.e. v_j should be boundary), then add to the component some set of edges E_j incident to v_j (in at most 2^d ways), deactivate v_j , and *activate* all the vertices incident to edges of the set E_j that have not been considered,
 - if $j \notin \mathcal{O}$, then add to the component all the edges incident to v_j , deactivate v_j , and *activate* all the neighbours of v_j that have not been considered.

The set of edges that have been added during this process forms the desired component z_i . So, the number of ways to choose the *i*-th component is at most $(n - \tilde{n})(\Delta + 2)^{\sigma_i} \max_{\substack{o_i \leq (\Delta + 2)\sigma_i \\ o_i}} {x_i \choose o_i} 2^{do_i}$.

Eventually we get that the number of *ordered* choices of components with parameters $\ell_i, x_i, i \in [c]$, in F is at most

$$c!\binom{n}{c}\left(\frac{2d}{3}\right)^{c}\prod_{i=1}^{c}(\Delta+2)^{\sigma_{i}}\max_{o_{i}\leq(\Delta+2)\sigma_{i}}\binom{x_{i}}{o_{i}}2^{do_{i}}\leq c!\binom{n}{c}d^{c}2^{2(\Delta+1)\sigma}\max_{o\leq(\Delta+2)\sigma}\binom{x}{o}.$$

Note that this bound does not depend on the order of the choice of components, thus

$$\alpha(\ell, x, c) \le \binom{n}{c} \binom{x}{c} \binom{\sigma+c}{c} d^c \cdot 2^{2(\Delta+1)\sigma} \max_{\substack{o \le (\Delta+2)\sigma}} \binom{x}{o}$$
$$\le \binom{n}{c} \binom{x}{c} (2d)^c \cdot 2^{(2\Delta+3)\sigma} \max_{\substack{o \le (\Delta+2)\sigma}} \binom{x}{o}$$

as needed.

Let us now fix $H \subset F$ with x vertices, ℓ edges, and c components. Let us bound the number of ways to extend H to an $F' \in \mathcal{F}_n$. We construct such an extension in the following way.

First of all, we add to H all the isolated vertices from [n] that it misses, and then we forget the labels of all the n vertices of H. Fix some $F' \in \mathcal{F}_n$ such that each vertex (but the first one) has a smaller neighbour (in the linear order on [n]). We will compute the number of ways to embed the unlabelled H into F', i.e. the number $mon(H \to F')$ of monomorphisms $H \to F'$. Clearly, the desired number of extensions is exactly $\frac{\operatorname{mon}(H \to F')}{|\operatorname{Aut}(F')|}$. Let \mathcal{Z} be the set of all n - x + cconnected components of H. We should compute the number of ways to embed the elements of \mathcal{Z} in F' disjointly.

Let z_1, \ldots, z_{n-x+c} be an arbitrary ordering of \mathcal{Z} (there are (n-x+c)! ways to order the elements of \mathcal{Z}). We embed sequentially each z_i in F' in a way such that all vertices of z_i are bigger than all the i-1 minimum vertices of z_1, \ldots, z_{i-1} . At every step $i = 1, \ldots, n-x+c$, consider the minimum vertex κ_i of F' such that none of the embedded elements of \mathcal{Z} in F'contain this vertex. If z_i is a single vertex, then we assign κ_i with z_i and proceed to the next step. Otherwise, we let κ_i be the minimum vertex of the embedding of z_i and, then, distinguish between the following cases.

First, we assume that z_i is closed. If $|V(z_i)| = 2$, then there are at most d ways to choose this edge and at most two ways to place it (two rotations). Thus there are at most 2d ways to embed z_i . If $|V(z_i)| \ge 3$, then there are at most $2d|\operatorname{Aut}(z_i)|$ ways to choose a (labelled) copy of z_i in F' with the minimum vertex κ_i , due to Claim 2.2. Indeed, there are at most 2d ways to choose a subgraph in F' that is isomorphic to z_i : first, choose an edge that is adjacent to κ_i , and then choose a closed subgraph on $|V(z_i)|$ vertices that does not contain the selected edge. Moreover, $|Aut(z_i)| \le |V(z_i)|(d-1)^{|V(z_i)|}$, due to [22, Theorem 2].

Second, let z_i be not closed with $|\partial_v z_i| = o_i$. Choose a set \mathcal{O} from $\binom{[|V(z_i)|]}{o_i}$ that will identify the labels of boundary vertices in the embedding of z_i into F'. Activate $v_1 := \kappa_i$. At every step $j \geq 1$, consider the minimum active vertex v_j in F' and

- if $j \in \mathcal{O}$, then add to the image of z_i under construction some set of edges E_j incident to v_j (in at most 2^d ways), deactivate v_j , and *activate* all the vertices incident to edges of the set E_j that have not been considered,
- if $j \notin \mathcal{O}$, then add to the image of z_i all the edges incident to v_j , deactivate v_j , and activate all the neighbours of v_i that have not been considered.

The image is constructed. However, we have not yet mapped the vertices of z_i to the vertices of the image. The number of such mappings ρ_i is exactly the number of automorphisms of z_i . As above, it is bounded by $|V(z_i)|(d-1)^{|V(z_i)|}$.

We also notice that every automorphism of every z_i respects the property of a vertex to be boundary. Moreover, any automorphism that preserves boundary vertices of z_i extends trivially to an automorphism of the entire F'. Thus, $\prod_{i=1}^{c} |\operatorname{Aut}(z_i)| \leq |\operatorname{Aut}(F')| \prod_{i=1}^{c} d^{o_i}$, where d^{o_i} is the upper bound on the number of automorphisms of z_i that preserve all non-boundary vertices.

We conclude that there are at most

$$(n-x+c)!(2d)^{c} \min\left\{\left(\frac{x}{c}\right)^{c} (d-1)^{x} \prod_{i=1}^{c} \binom{x_{i}}{o_{i}} 2^{do_{i}}, |\operatorname{Aut}(F')| \prod_{i=1}^{c} \binom{x_{i}}{o_{i}} (d2^{d})^{o_{i}}\right\}$$

$$\leq (n-x+c)!(2d)^{c} \min\left\{\left(\frac{x}{c}\right)^{c} (d-1)^{x} 2^{d(2+\Delta)\sigma}, |\operatorname{Aut}(F')| (d2^{d})^{(2+\Delta)\sigma}\right\} \max_{o \leq (2+\Delta)\sigma} \binom{x}{o}$$
aves to embed H into F'. It remains to divide the final bound by $|\operatorname{Aut}(F')|$.

ways to embed H into F'. It remains to divide the final bound by $|\operatorname{Aut}(F')|$.

3.2. Coarse estimates. We will also use the following coarse version of the first part of Claim 3.1. For a graph H with maximum degree at most d, let us denote by $\mathcal{J}_{\ell,x,c}^H$ the set

of all subgraphs $J \subset H$ with ℓ edges, x non-isolated vertices, and c connected components (excluding isolated vertices).

Claim 3.2. For every graph H with maximum degree at most d,

$$|\mathcal{J}_{\ell,x,c}^{H}| \le {\binom{\min\{|E(H)|, |V(H)|\}}{c}} (16d)^{\ell}.$$
 (6)

Proof. We first choose c edges (or vertices, if |V(H)| < |E(H)|) in H that belong to different components of J. Then, the number of ways to assign to each of the c components its number of edges is at most $\binom{\ell+c-1}{c-1} \leq 2^{2\ell}$ since $\ell \geq c$. Finally, as soon as, for every $i \in [c]$, a vertex in the *i*-th component of size ℓ_i is fixed, the number of ways to explore it sequentially, edge by edge, is at most $(4d)^{\ell_i}$: at every step, first decide whether the current edge is incident to the vertex considered at the previous step or to the next explored but unconsidered vertex — two choices (the order of vertices in H is arbitrary and fixed in advance). If the second choice is made, consider the minimum vertex in the set of explored unconsidered vertices. Second, choose the new edge incident to the considered vertex in at most d ways. Finally, if a new vertex has been explored, decide whether it has additional edges in J or not (two choices). If it does not have additional edges, remove it from the pool of unconsidered vertices. It completes the proof of (6).

Moreover, for any *d*-regular graph F on [n] and any $H \subset F$ with ℓ edges, x non-isolated vertices, and c connected components (excluding isolated vertices), by counting mon $(H \to F)$ using the same strategy as in the proof of the second part of Claim 3.1, we get

Claim 3.3. The number of ways to extend H to an isomorphic copy of F on [n] is at most $e^{O(\ell)}(n-x+c)!/|\operatorname{Aut}(F)|.$

Proof. As in the proof of Claim 3.1, we let \mathcal{Z} be the set of all n - x + c connected components of H. We order the elements of \mathcal{Z} and then, at every step $i \in \{1, \ldots, n - x + c\}$, we consider the minimum vertex κ_i of F such that none of the embedded elements of \mathcal{Z} contain this vertex and map some vertex of z_i to κ_i . After that, we embed z_i edge by edge in an arbitrary order that respects connectivity. We then get at most $(n - x + c)! \left(\frac{x}{c}\right)^c d^\ell = e^{O(\ell)}(n - x + c)!$ embeddings since

$$\left(\frac{x}{c}\right)^c = e^{c\ln(x/c)} = e^{x \cdot \frac{\ln(x/c)}{x/c}} \le e^x.$$
(7)

4. FRAGMENTATION: ARBITRARY FAMILIES

This section presents our main tool — fragmentation trick via typical fragments. We, first, use it in the proof of Theorem 1.4 in Section 5, and then we use this tool to show the existence of rare fragments without closed subgraphs of size $\Omega(\log n)$ in our proof of Theorem 1.7 in Section 6. Although the trick appeared in several papers (in particular, in [18], see also Section 1.4), we describe it here in full detail for the sake of completeness. Here we present it in a slightly different form, without explicitly distinguishing between pathological and non-pathological pairs, which is more transparent for us.

Let $\delta_n = o(1)$ be a slowly decreasing function. Let $d \ge 3$ be a constant,

$$p = p(n) = \Theta(n^{-2/d}), \qquad m = m(n) = pN, \qquad f = f(n) = O(n).$$

Let \mathcal{F}_n be an arbitrary multiset of graphs on [n], each graph has f edges. Let \mathcal{B} be an arbitrary graph property (not necessarily isomorphism-closed). For $F \in \mathcal{F}_n$ and $W \subset {[n] \choose 2}$, let $\mathcal{M}(F, W)$ be the multiset of all $F' \in \mathcal{F}_n$ such that $F' \subset F \cup W$. Let $\mathcal{M}_{\mathcal{B}}(F, W)$ be the multiset of all $F' \in \mathcal{M}(F, W)$ such that $F' \cap F \in \mathcal{B}$. Let us say that the pair (F, W) is \mathcal{B} -bad, if $|\mathcal{M}_{\mathcal{B}}(F, W)| > \delta_n |\mathcal{M}(F, W)|$. Let \mathbf{F} be a uniformly random element of \mathcal{F}_n sampled independently of a uniformly random $\mathbf{W} \in {\binom{[n]}{2}}$. We shall prove sufficient conditions for (\mathbf{F}, \mathbf{W}) not being \mathcal{B} -bad whp. That would mean that whp we may replace most of $F \in \mathcal{F}_n$ with fragments $F \cap F'$ that do not have the property \mathcal{B} . Let \mathcal{B}_ℓ contain all graphs with the property \mathcal{B} and ℓ edges. For $F \in \mathcal{F}_n$, let

$$\Pi_{\mathcal{B}}^{F} := \mathbb{P}(F \cap \mathbf{F} \in \mathcal{B}).$$
(8)

Lemma 4.1. If

$$\max_{F \in \mathcal{F}_n} \sum_{\ell=0}^{f} \Pi_{\mathcal{B}_\ell}^F \left(\left(1 + \frac{3f}{m} \right) \frac{N}{m} \right)^\ell e^{-f^2/m + f^3/(3m^2)} = o(1),$$
(9)

then whp (\mathbf{F}, \mathbf{W}) is not \mathcal{B} -bad.

Proof. For $t \in \{0, 1, ..., f\}$, let

$$M(t) := |\mathcal{F}_n| \binom{N-f}{m-t} / \binom{N}{m+f-t}$$

be the expected number of $F \in \mathcal{F}_n$ such that F belongs to a uniformly random subset of $\binom{[n]}{2}$ of size m + f - t.

Let $t \in \{0, 1, \ldots, f\}$. Each pair $\left\{F \in \mathcal{F}_n, W \in \binom{\binom{n}{2}}{m}\right\}$ such that $|\mathcal{M}(F, W)| < \delta_n M(t)$ and $|F \cap W| = t$ can be obtained by choosing a set $A \subset \binom{\binom{n}{2}}{2}$ of size m + f - t, on the role of $F \cup W$ (in at most $\binom{N}{m+f-t}$ ways), choosing an $F \subset A, F \in \mathcal{F}_n$ (in less than $\delta_n M(t)$ ways), and choosing the intersection $F \cap W$ (in $\binom{f}{t}$ ways). We conclude that

$$\mathbb{P}\left(|\mathcal{M}(\mathbf{F},\mathbf{W})| < \delta_n M(t) \mid |\mathbf{F} \cap \mathbf{W}| = t\right) \le \frac{\delta_n M(t) \binom{N}{m+f-t} \binom{f}{t}}{|\mathcal{F}_n| \binom{N-f}{m-t} \binom{f}{t}} = \delta_n$$

Then,

$$\mathbb{P}((\mathbf{F}, \mathbf{W}) \text{ is } \mathcal{B}\text{-bad}, |\mathcal{M}(\mathbf{F}, \mathbf{W})| < \delta_n M(t) \mid |\mathbf{F} \cap \mathbf{W}| = t) \le \delta_n = o(1).$$

Thus, it is sufficient to prove that, uniformly over t,

$$\mathbb{P}((\mathbf{F}, \mathbf{W}) \text{ is } \mathcal{B}\text{-bad}, |\mathcal{M}(\mathbf{F}, \mathbf{W})| \ge \delta_n M(t) | |\mathbf{F} \cap \mathbf{W}| = t) = o(1).$$

The latter probability is at most

$$\mathbb{P}\left(\left|\mathcal{M}_{\mathcal{B}}(\mathbf{F}, \mathbf{W})\right| > \delta_n^2 M(t) \mid |\mathbf{F} \cap \mathbf{W}| = t\right) \le \frac{\mathbb{E}(X \mid |\mathbf{F} \cap \mathbf{W}| = t)}{\delta_n^2 M(t)},\tag{10}$$

where $X = X(\mathbf{F}, \mathbf{W})$ counts the number of $F' \in \mathcal{M}(\mathbf{F}, \mathbf{W})$ such that $F' \cap \mathbf{F} \in \mathcal{B}$. Fix $F \in \mathcal{F}_n$, let $\mathbf{W}'_F = \mathbf{W}'_F(t)$ be a uniformly random (m - t)-subset of $\binom{[n]}{2} \setminus F$, and let $X'_F = X'_F(t)$ be the number of $F' \in \mathcal{M}(F, \mathbf{W}'_F)$ such that $F' \cap F \in \mathcal{B}$. We get

$$\mathbb{E}(X \mid |\mathbf{F} \cap \mathbf{W}| = t) = \sum_{F \in \mathcal{F}_n} \mathbb{E}(X \cdot \mathbb{1}_{\mathbf{F} = F} \mid |\mathbf{F} \cap \mathbf{W}| = t) = \sum_{F \in \mathcal{F}_n} \frac{\mathbb{E}\left(X \cdot \mathbb{1}_{\mathbf{F} = F, |F \cap \mathbf{W}| = t\right)}{\mathbb{P}(|\mathbf{F} \cap \mathbf{W}| = t)}.$$

Since all $F \in \mathcal{F}_n$ have the same number of edges, $\mathbb{P}(|F \cap \mathbf{W}| = t)$ does not depend on $F \in \mathcal{F}_n$. In particular $\mathbb{P}(|\mathbf{F} \cap \mathbf{W}| = t) = \mathbb{P}(|F \cap \mathbf{W}| = t)$ for every $F \in \mathcal{F}_n$. Therefore,

$$\mathbb{E}(X \mid |\mathbf{F} \cap \mathbf{W}| = t) = \sum_{F \in \mathcal{F}_n} \mathbb{E}(X \cdot \mathbb{1}_{\mathbf{F} = F} \mid |F \cap \mathbf{W}| = t).$$

Recall that X(F, W) is the number of $F' \in \mathcal{M}(F, W)$ such that $F' \cap F \in \mathcal{B}$. We get

$$\mathbb{E}(X \mid |\mathbf{F} \cap \mathbf{W}| = t) = \sum_{F \in \mathcal{F}_n} \mathbb{E}(X(F, \mathbf{W}) \cdot \mathbb{1}_{\mathbf{F} = F} \mid |F \cap \mathbf{W}| = t)$$
$$= \sum_{F \in \mathcal{F}_n} \mathbb{E}(X(F, \mathbf{W}) \mid |F \cap \mathbf{W}| = t) \cdot \mathbb{P}(\mathbf{F} = F) = \sum_{F \in \mathcal{F}_n} \mathbb{E}X'_F \cdot \mathbb{P}(\mathbf{F} = F).$$
(11)

Then it remains to prove that $\frac{\mathbb{E}X'_F}{M(t)} = o(1)$ uniformly over $F \in \mathcal{F}_n$ and over t. By the definition of M(t),

$$\frac{\mathbb{E}X'_F}{M(t)} = \sum_{\ell} |\mathcal{F}_n| \frac{\Pi_{\mathcal{B}_{\ell}}^F}{M(t)} \binom{m-t}{f-\ell} / \binom{N-f}{f-\ell}$$
$$= \sum_{\ell} \Pi_{\mathcal{B}_{\ell}}^F \frac{\binom{m-t}{f-\ell} / \binom{N-f}{f-\ell}}{\binom{N-f}{m-t} / \binom{N}{m+f-t}} = \sum_{\ell} \Pi_{\mathcal{B}_{\ell}}^F \frac{\binom{m-t}{f-\ell} \binom{N}{m+f-t}}{\binom{N-f}{m-t} \binom{N-f}{f-\ell}}.$$

Applying Stirling's approximation, we get

$$\begin{split} \frac{\mathbb{E}X'_{F}}{M(t)} &\sim \sum_{\ell} \Pi_{\mathcal{B}_{\ell}}^{F} \frac{(m-t)^{2(m-t)}(N-2f+\ell)^{N-2f+\ell}N^{N}}{(m-t-f+\ell)^{m-t-f+\ell}(m+f-t)^{m+f-t}(N-f)^{2N-2f}} \\ &= \sum_{\ell} \Pi_{\mathcal{B}_{\ell}}^{F} \left(\frac{N-2f+\ell}{m-t-f+\ell} \right)^{\ell} \frac{\left(1+\frac{f-\ell}{m-t-f+\ell}\right)^{m-t-f}\left(1-\frac{f}{m-t+f}\right)^{m-t+f}}{\left(1-\frac{f}{N}\right)^{N}\left(1+\frac{f-\ell}{N-2f+\ell}\right)^{N-2f}} \\ &\leq \sum_{\ell} \Pi_{\mathcal{B}_{\ell}}^{F} \left(\frac{N}{m} \left(1+\frac{t+f-\ell}{m-t-f+\ell}\right) \right)^{\ell} \\ &\qquad \times e^{f-\ell-\frac{(f-\ell)^{2}}{2m}+\frac{(f-\ell)^{2}\ell}{2(m-t-f+\ell)^{2}}+\frac{(f-\ell)^{3}}{3(m-t-f+\ell)^{2}}-f-\frac{f^{2}}{2m}+f-f+\ell+O(1)} \\ &\leq \sum_{\ell} \Pi_{\mathcal{B}_{\ell}}^{F} \left(\left(1+\frac{3f}{m}\right)\frac{N}{m} \right)^{\ell} e^{-f^{2}/m+f^{3}/(3m^{2})+O(1)} = o(1), \end{split}$$

completing the proof.

5. Sharp thresholds

We prove the two parts of Theorem 1.4 separately in the next two sections.

5.1. Better expansion of large subgraphs. Here, we prove the following.

Theorem 5.1. Let $d \ge 3$, $\delta \in (0, 1/d)$, and let F = F(n) be a sequence of d-regular graphs on [n], $n \in \mathbb{N}$, such that

- the number of automorphisms of F is at most $e^{o(n^{1-\delta})}$;
- for some $w = w(n) = \omega(1)$, every $\tilde{F} \subset F$ with $3 \leq |V(\tilde{F})| \leq n^{1-\delta}$ has $|\partial_e(\tilde{F})| \geq d+1+\lfloor |V(\tilde{F})|w/\log n\rfloor$ and every $\tilde{F} \subset F$ with $n^{1-\delta} < |V(\tilde{F})| \leq n-3$ has $|\partial_e(\tilde{F})| \geq d+1$.

Let $\varepsilon > 0$. If $p > (1 + \varepsilon) \left(\frac{e}{n}\right)^{2/d}$, then, assuming that dn is even, whp $\mathbf{G} \sim G(n, p)$ contains an isomorphic copy of F.

We note that Theorem 5.1 indeed implies the first part of Theorem 1.4 since $p_F^e \ge (1 - o(1)) \left(\frac{e}{n}\right)^{2/d}$. Indeed, due to the first requirement, $|\operatorname{Aut}(F)| = e^{o(n)}$, implying

$$\mathbb{E}X_F = \frac{n^n}{e^{n(1+o(1))}} p^{dn/2} = \Theta(1) \quad \Rightarrow \quad p = (1-o(1)) \left(\frac{e}{n}\right)^{2/d}.$$
 (12)

The rest of the section is devoted to the proof of Theorem 5.1. Fix a d-regular graph F satisfying the requirements of the theorem.

5.1.1. Fragmentation: $1/\delta$ days. Consider $\lfloor 1/\delta \rfloor$ independent samples

$$\mathbf{W}_{1} \sim G(n,m), \quad m = \left\lfloor (1+\varepsilon) \left(\frac{e}{n}\right)^{2/d} N \right\rfloor, \quad \text{and}$$
$$\mathbf{W}_{2}, \dots, \mathbf{W}_{\lfloor 1/\delta \rfloor} \sim G(n,m_{0}), \quad m_{0} = \left\lfloor \varepsilon \cdot n^{-2/d} \cdot N \right\rfloor.$$

Recall that we denote by \mathcal{F}_n the family of all isomorphic copies of F on [n].

Claim 5.2. Whp there exists $F' \subset F \cup \mathbf{W}_1, F' \in \mathcal{F}_n$, such that $|F \cap F'| \leq n^{1-\delta}$.

Proof. Let $f = \frac{dn}{2}$, and \mathcal{B} be the property of graphs on n vertices to have more than $n^{1-\delta}$ edges. Due to Lemma 4.1, it is sufficient to prove (9). More precisely, due to symmetry, it is sufficient to prove

$$\sum_{\ell=0}^{f} \Pi_{\mathcal{B}_{\ell}}^{F} \left(\left(1 + \frac{3f}{m} \right) \frac{N}{m} \right)^{\ell} e^{-f^{2}/m + f^{3}/(3m^{2})} = o(1),$$
(13)

for the fixed $F \in \mathcal{F}_n$. Fix non-negative integers c and x. As in Claim 3.2, we denote by $\mathcal{J}_{\ell,x,c}$ the set of all subgraphs $J \subset F$ with ℓ edges, x non-isolated vertices, and c connected components (excluding isolated vertices), and let $p(\ell, x, c) := \mathbb{P}(\mathbf{F} \cap F \in \mathcal{J}_{\ell,x,c})$. Due to Claim 3.1, for some constants $A_1, A_2 > 0$,

$$p(\ell, x, c) \leq \frac{\binom{n}{c}\binom{x}{c}e^{A_1c + A_2\sigma}\max_{o \leq (\Delta+2)\sigma}\binom{x}{o}^2(n - x + c)!}{|\mathcal{F}_n|}$$

Let $\ell > n^{1-\delta}$. Recalling that $x = \frac{2}{d}\ell + \frac{c\Delta}{d} + \frac{2}{d}\sigma$, we get

$$\frac{(n-x+c)!}{|\mathcal{F}_n|} \le e^{o(n^{1-\delta})} \cdot \frac{(n-\frac{2\ell+c(\Delta-d)+2\sigma}{d})!}{n!} \le e^{o(n^{1-\delta})} \cdot \frac{e^{(x-c)^2/n}}{n^{(2\ell+c+2\sigma)/d}}.$$
(14)

We first choose $0 < \varepsilon' \ll \varepsilon''$ small enough. From the bound $\binom{n}{c} \leq \left(\frac{en}{c}\right)^c$, we get

$$p(\ell, x, c) \le e^{o(n^{1-\delta})} \cdot \frac{\left(\frac{e^{A_1+1}n^{1-1/d}}{c}\right)^c {\binom{x}{c}} e^{A_2\sigma} {\binom{x}{o}}^2 e^{(x-c)^2/n}}{n^{(2\ell+2\sigma)/d}},$$
(15)

where $o = o(x) \le 8\sigma$ is chosen in such a way that $\binom{x}{o}$ achieves its maximum. We get that

$$\left(\frac{e^{A_1+1}n^{1-1/d}}{c}\right)^c \le e^{A_1n^{1-1/d}} \le e^{o(\ell)}.$$

We further consider separately several different cases.

1. If $\sigma > \varepsilon' x$, then

$$\binom{x}{c}\binom{x}{o}^{2}e^{A_{2}\sigma}e^{(x-c)^{2}/n} < 8^{x}e^{x}e^{A_{2}\sigma} < n^{2\sigma/d}.$$

Therefore,

$$p(\ell, x, c) \le \left((1+\varepsilon)/n \right)^{2\ell/d}.$$
(16)

2. Let $\sigma \leq \varepsilon' x$. **2.1.** If $c < \varepsilon' x$ and $x < \varepsilon' n$, then

$$\binom{x}{c}\binom{x}{o}^2 e^{A_2\sigma} e^{(x-c)^2/n} < e^{(\varepsilon''/d)\ell}$$

implying (16) as well.

2.2. If $c < \varepsilon' x$ and $x \ge \varepsilon' n$, then $e^{A_2\sigma} {x \choose c} {a \choose c}^2 < e^{(\varepsilon''/d)\ell}$. Thus, (15) implies

$$p(\ell, x, c) \le \frac{e^{(x-c)^2/n}}{n^{2\ell/d}} e^{(\varepsilon''/d + o(1))\ell} \le (e^{1+\varepsilon''}/n)^{2\ell/d},$$

since $2\ell/d = x - c\Delta/d - 2\sigma/d \ge x(1 - 5\varepsilon'/2)$.

2.3. Finally, let $c \ge \varepsilon' x$. Since $x \ge \frac{2}{d}\ell$, we get that $x \gg n^{1-1/d}$. In particular,

$$\left(\frac{e^{A_1+1}n^{1-1/d}}{c}\right)^c \le e^{e^{A_1n^{1-1/d}}} \le e^{-10x}$$

Since $e^{A_2\sigma} < n^{2\sigma/d}$, we get

$$p(\ell, x, c) \le e^{-10x + o(\ell)} \frac{\binom{x}{c} \binom{x}{o}^2 e^{(x-c)^2/n}}{n^{2\ell/d}} \le e^{-10x + x + o(\ell)} 8^x n^{-2\ell/d} \le n^{-2\ell/d}.$$

Summing up,

$$\begin{split} \sum_{\ell} \Pi_{\mathcal{B}_{\ell}}^{F} \left(\left(1 + \frac{3f}{m} \right) \frac{N}{m} \right)^{\ell} e^{-f^{2}/m + f^{3}/(3m^{2})} &\leq \sum_{\ell > n^{1-\delta}} \sum_{x,c} p(\ell, x, c) \left(\frac{1 + o(1)}{1 + \varepsilon} \cdot \left(\frac{n}{e} \right)^{2/d} \right)^{\ell} \\ &\leq n^{2} \sum_{\ell > n^{1-\delta}} \left(\frac{e^{2\varepsilon''/d} + o(1)}{1 + \varepsilon} \right)^{\ell} = o\left(\frac{1}{n^{2}} \right), \end{split}$$
pleting the proof of the claim.

completing the proof of the claim.

For every $F \in \mathcal{F}_n$ (with a slight abuse of notation, for the sake of simplicity of presentation, we now denote by F an arbitrary graph from \mathcal{F}_n) such that there exists F' as in the statement of Claim 5.2, we choose one such F' and put $F \cap F'$ into a multiset $\mathcal{F}_n^{(1)}$. Due to Claim 5.2 and Markov's inequality, we get that whp $|\mathcal{F}_n^{(1)}| = (1 - o(1))|\mathcal{F}_n|$. We then proceed by induction. Assume that, for $i \in [\lceil 1/\delta \rceil - 1]$, we have a multiset $\mathcal{F}_n^{(i)}$ comprising a single $H \subset F$ from almost every $F \in \mathcal{F}_n$ such that the graph $H \cup \mathbf{W}_1 \cup \ldots \cup \mathbf{W}_i$ contains some $F' \in \mathcal{F}_n$ and $|H| = \lfloor n^{1-i\delta} \rfloor$. Let **H** be a uniformly random element of $\mathcal{F}_n^{(i)}$.

Claim 5.3. Whp there exists $H' \subset \mathbf{H} \cup \mathbf{W}_{i+1}$, where $H' \in \mathcal{F}_n^{(i)}$, such that $|\mathbf{H} \cap H'| \leq \mathbf{W}_{i+1}$ $\max\big\{n^{1-(i+1)\delta},\ln n\big\}.$

Proof. Let $f = \lfloor n^{1-i\delta} \rfloor$, $m = m_0$, and \mathcal{B} be the property of graphs on n vertices to have more than $\max\{n^{1-(i+1)\delta}, \ln n\}$ edges. Due to Lemma 4.1, it is sufficient to prove (9).

Fix non-negative integers c and x. Recall that, for $H \in \mathcal{F}_n^{(i)}$, we denote by $\mathcal{J}_{\ell,x,c}^H$ the set of all subgraphs $J \subset H$ with ℓ edges, x non-isolated vertices, and c connected components (excluding isolated vertices), and denote $p^H(\ell, x, c) := \mathbb{P}(\mathbf{H} \cap H \in \mathcal{J}^H_{\ell, x, c})$.

Due to Claim 3.2 and Claim 3.3, for some constant A > 0 that does not depend on ε ,

$$p^{H}(\ell, x, c) \leq \frac{\binom{\lfloor n^{1-i\delta} \rfloor}{c} e^{A\ell} (n-x+c)! / |\operatorname{Aut}(F)|}{|\mathcal{F}_{n}^{(i)}|} = (1+o(1)) \frac{\binom{\lfloor n^{1-i\delta} \rfloor}{c} e^{A\ell} (n-x+c)!}{n!}.$$

Due to (14) and since $x \leq |V(H)| = o(n)$,

$$p^{H}(\ell, x, c) \leq \frac{\left(\frac{en^{1-i\delta}}{c}\right)^{c} e^{(A+o(1))\ell}}{n^{2\ell/d+c/d+2\sigma/d}}.$$

Let C > 0 be a large constant. Since every subgraph $J \subset F$ with $2 \leq |V(H)| \leq n^{1-\delta}$ has $|\partial_e(J)| \geq d+1 + \lfloor |V(J)|w/\log n \rfloor$, for a graph $J \in \mathcal{J}_{\ell,x,c}^H$ consisting of c components that have ℓ_1, \ldots, ℓ_c edges and x_1, \ldots, x_c vertices, we have $\ell_i = \frac{d}{2}x_i - \frac{\Delta}{2} - \sigma_i$, where

$$\sigma_i \ge \frac{dC\ell_i}{2\ln n} \cdot I\left(\ell_i \ge \frac{\ln n(1-d\delta)}{dC}\right).$$

Therefore, for any admissible triple (ℓ, x, c) , we get $\ell = \frac{d}{2}x - \frac{\Delta}{2}c - \sigma$, where

$$\sigma = \min_{J \in \mathcal{J}_{\ell,x,c}^{H}} \sum_{i=1}^{c} \sigma_{i} \ge \min_{J \in \mathcal{J}_{\ell,x,c}^{H}} \sum_{i: \ell_{i} \ge \frac{\ln n(1-d\delta)}{dC}} \frac{dC\ell_{i}}{2\ln n}$$
$$> \left(\ell - c \cdot \frac{\ln n(1-d\delta)}{dC}\right) \frac{dC}{2\ln n} = \frac{dC}{2\ln n} \cdot \ell - \frac{1-d\delta}{2} \cdot c.$$
(17)

Therefore,

$$\begin{split} \sum_{\ell} \Pi_{\mathcal{B}_{\ell}}^{H} \left(\left(1 + \frac{3f}{m} \right) \frac{N}{m} \right)^{\ell} &\leq \sum_{\ell > \max\{n^{1-(i+1)\delta}, \ln n\}} \sum_{x,c} p^{H}(\ell, x, c) \left(\frac{1 + o(1)}{\varepsilon} \cdot n^{2/d} \right)^{\ell} \\ &\leq n^{2} \sum_{\ell > \max\{n^{1-(i+1)\delta}, \ln n\}} \frac{\left(\frac{en^{1-i\delta}}{c} \right)^{c}}{n^{\delta c + \frac{C}{\ln n} \cdot \ell}} \left(\frac{e^{A} + o(1)}{\varepsilon} \right)^{\ell} \\ &\leq n^{2} \sum_{\ell > \max\{n^{1-(i+1)\delta}, \ln n\}} \frac{e^{n^{1-(i+1)\delta}}}{e^{C\ell}} \left(\frac{e^{A} + o(1)}{\varepsilon} \right)^{\ell} = o\left(\frac{1}{n^{2}} \right), \end{split}$$

whenever $C > 2(A + \ln(1/\varepsilon))$, completing the proof.

By induction, whp we get a multiset $\mathcal{F}_n^{(\lceil 1/\delta \rceil)}$ of size $(1 - o(1))|\mathcal{F}_n|$ comprising a single $H \subset F$ from almost every $F \in \mathcal{F}_n$ such that the graph $H \cup \mathbf{W}_1 \cup \ldots \cup \mathbf{W}_{\lceil 1/\delta \rceil}$ contains some $F' \in \mathcal{F}_n$ and $|H| = \lfloor \ln n \rfloor$.

5.1.2. Proof of Theorem 5.1. Let

$$\mathbf{G} \sim G(n, p), \quad p = (1 + \lceil 1/\delta \rceil \varepsilon)(e/n)^{2/d}$$

By [17, Corollary 1.16], whp there exists a multiset $\mathcal{F}_n^{(\lceil 1/\delta \rceil)} = \mathcal{F}_n^{(\lceil 1/\delta \rceil)}(\mathbf{G})$ of graphs of size $\lfloor \ln n \rfloor$ comprising a single subgraph H of almost every $F \in \mathcal{F}_n$ so that $H \cup \mathbf{G}$ contains some $F' \in \mathcal{F}_n$.

Let X be the number of $H \in \mathcal{F}_n^{(\lceil 1/\delta \rceil)}$ that are subgraphs of $\mathbf{G}' \sim G(n, p' = \varepsilon n^{-2/d})$, sampled independently of **G**. We get

$$\mathbb{E}X = \left| \mathcal{F}_n^{(\lceil 1/\delta \rceil)} \right| \cdot p^{\prime \lfloor \ln n \rfloor} = (1 - o(1)) |\mathcal{F}_n| p^{\prime \lfloor \ln n \rfloor} = \omega(1).$$

Let \mathcal{B} be the set of non-empty graphs. Due to the definition (8) of $\Pi^{H}_{\mathcal{B}_{\ell}} = \Pi^{H}_{\mathcal{B}_{\ell}}(\mathcal{F}_{n}^{(\lceil 1/\delta \rceil)}),$

$$\frac{\operatorname{Var} X}{(\mathbb{E} X)^2} \leq \frac{\max_{H \in \mathcal{F}_n^{(\lceil 1/\delta \rceil)}} \sum_{\ell \geq 1} \left(\Pi_{\mathcal{B}_\ell}^H \cdot \left| \mathcal{F}_n^{(\lceil 1/\delta \rceil)} \right| \cdot p'^{\lfloor \ln n \rfloor - \ell} \right)}{\mathbb{E} X} = \max_{H \in \mathcal{F}_n^{(\lceil 1/\delta \rceil)}} \sum_{\ell \geq 1} \Pi_{\mathcal{B}_\ell}^H p'^{-\ell}.$$

Therefore, due to Claim 3.2, Claim 3.3, estimates (17), and the inequality

$$\frac{(n-x+c)!}{n!} \le \frac{e^{(x-c)^2/n}}{n^{(2\ell+c+2\sigma)/d}},$$
(18)

we get that, for some constant A > 0,

$$\frac{\operatorname{Var} X}{(\mathbb{E}X)^2} \le \sum_{\ell \ge 1} \sum_{x,c} (1 - o(1)) \frac{\binom{\lfloor \ln n \rfloor}{c} e^{A\ell}}{n^{2\ell/d + c/d + 2\sigma/d}} \left(\frac{1}{\varepsilon} \cdot n^{2/d}\right)^{\ell}$$
$$\le \sum_{\ell \ge 1} \sum_{c \ge 1} O(\ell) \left(\frac{e \lfloor \ln n \rfloor}{cn^{\delta}}\right)^c \left(\frac{e^A}{\varepsilon \cdot e^C}\right)^{\ell} = O\left(\frac{\ln n}{n^{\delta}}\right),$$

whenever $C > 2(A + \ln(1/\varepsilon))$. This completes the proof of Theorem 5.1, due to Chebyshev's inequality.

5.2. Better expansion of small subgraphs. It remains to prove the second part of Theorem 1.4:

Theorem 5.4. Let $d \ge 4$. Let $\gamma \in (0,1)$ be a fixed constant, let $C = C(d,\gamma)$ be large enough, and let $D \ge 1$ be a fixed constant. Let F = F(n) be a sequence of d-regular graphs on [n], $n \in \mathbb{N}$, such that

- for every subgraph $\tilde{F} \subset F$ with $|E(\tilde{F})| \leq Cn^{2/d}$ and $|\partial_v(\tilde{F})| \leq \gamma |V(\tilde{F})|$, the number of automorphisms of \tilde{F} that fix all vertices of $\partial_v(\tilde{F})$ is at most D;
- for every subgraph $\tilde{F} \subseteq F$ with $|\partial_v(\tilde{F})| \leq \gamma |V(\tilde{F})|$, the number of automorphisms of \tilde{F} that fix all vertices of $\partial_v(\tilde{F})$ is at most $e^{o(|V(\tilde{F})|)}$;
- every $\tilde{F} \subset F$ with $3 \leq |V(\tilde{F})| \leq n-3$ has $|\partial_e(\tilde{F})| \geq 2d$.

Let $\varepsilon > 0$. If $p > (1 + \varepsilon) \left(\frac{e}{n}\right)^{2/d}$, then whp (assuming that dn is even) $\mathbf{G} \sim G(n, p)$ contains an isomorphic copy of F.

We notice that Theorem 5.4 indeed implies the second part of Theorem 1.4 since $p_F^e \ge (1 - o(1)) \left(\frac{e}{n}\right)^{2/d}$. Indeed, due to the second requirement applied to $\tilde{F} = F$ with $\partial_v(\tilde{F}) = \emptyset$, we get $|\operatorname{Aut}(F)| = e^{o(n)}$, implying (12). The rest of the section is devoted to the proof of Theorem 5.4. Fix a *d*-regular graph F satisfying the requirements of the theorem.

5.2.1. Fragmentation: two days. Consider two independent samples

$$\mathbf{W}_1 \sim G(n,m), \ m := \left\lfloor (1+\varepsilon) \left(\frac{e}{n}\right)^{2/d} N \right\rfloor, \quad \text{and} \quad \mathbf{W}_2 \sim G(n,m_0), \ m_0 := \left\lfloor \varepsilon \cdot n^{-2/d} \cdot N \right\rfloor.$$

Let \mathcal{F}_n be the family of all isomorphic copies of F on [n].

Claim 5.5. Whp there exists $F' \subset F \cup \mathbf{W}_1$ satisfying the following: $F' \in \mathcal{F}_n$, $F \cap F'$ has at most $\sqrt{\ln n}$ non-isolated edges, and $|F \cap F'| \leq Cn^{2/d}$.

Proof. Let $f = \frac{dn}{2}$, and let a graph G on n vertices has the property \mathcal{B} if either G has more than $Cn^{2/d}$ edges or G has more than $\sqrt{\ln n}$ non-isolated edges. Due to Lemma 4.1 and due to symmetry, it is sufficient to prove (13).

In contrast to the proof of Claims A.1 and 5.2, here we will distinguish the contribution to $\Pi_{B_{\ell}}^{F}$ between components consisting of a single edge and all the others. Fix non-negative integers c, c', and x. Denote by $\mathcal{J}_{\ell,x,c,c'}$ the set of all subgraphs $J \subset F$ with ℓ edges, xnon-isolated vertices, c isolated edges, and c' other connected components, excluding isolated vertices. Denote by $p(\ell, x, c, c')$ the probability that $\mathbf{F} \cap F \in \mathcal{J}_{\ell, x, c, c'}$. We now refer to the proof of Claim 3.1 and consider separately connected components of some $J \in \mathcal{J}_{\ell, x, c, c'}$ consisting of single edges. The number of ways to embed c disjoint edges into F is at most

$$n(n-2)\dots(n-2(c-1)) \cdot d^{c} = 2^{c} \frac{(n/2)!}{(n/2-c)!} d^{c}$$

$$\leq 2\sqrt{\frac{n}{n-2c+1}} \cdot \frac{(2d)^{c}(n/2)^{c}}{e^{c}} \left(1 + \frac{c}{n/2-c}\right)^{n/2-c}$$

$$\leq 2\sqrt{\frac{n}{n-2c+1}} \cdot (dn)^{c} =: \varphi_{1}.$$

We also note that J does not have closed components. After we identify the image z'_i in F of a non-trivial connected component z_i of J with x_i vertices and o_i boundary vertices, the number of ways to map the vertices of z_i to the vertices of the image is bounded by $(D \cdot \mathbb{1}_{\ell_i \leq Cn^{2/d}} + e^{o(x_i)} \cdot \mathbb{1}_{\ell_i > Cn^{2/d}}) \cdot d^{o_i}$ whenever $o_i \leq \gamma x_i$ (since every automorphism of z'_i acts bijectively on the vertex boundary). If $o_i > \gamma x_i$, then we may bound the number of automorphisms by

$$x_i(d-1)^{x_i} \le (o_i/\gamma)(d-1)^{o_i/\gamma} < (2d)^{o_i/\gamma},$$

in the same way as in the proof of Claim 3.1.

Let $\sigma := \frac{d}{2}(x-2c) - (\ell - c + \frac{\Delta}{2} \cdot c')$ be the excess of the union of components of J with more than 2 vertices. Following the proof of the second part of Claim 3.1, we get that there exist constants $A^1_{\beta}, A^2_{\beta} > 0$ such that the number of ways to embed J into F is at most

$$\beta(\ell, x, c, c') = \varphi_1 \cdot (n - x + c')! e^{A_\beta^1 c' + A_\beta^2 \sigma} \left(1 + e^{o(\ell)} \cdot \mathbb{1}_{\ell > Cn^{2/d}} \right) \times \max_{o \le (\Delta + 2)\sigma} \binom{x - 2c}{o} (2d)^{o/\gamma}.$$

Now, let us bound the number of ways to choose a subgraph $J \subset F$ such that $J \in \mathcal{J}_{\ell,x,c,c'}$. The number of ways to choose c disjoint edges in F is at most

$$\frac{n(n-2)\dots(n-2(c-1))\cdot d^c}{c!\cdot 2^c} \le \frac{\varphi_1}{\sqrt{c}\cdot (2c/e)^c} =:\varphi_2.$$

According to the proof of the first part of Claim 3.1, there exist constants $A^1_{\alpha}, A^2_{\alpha} > 0$ such that the number of ways to choose the other c' components is at most

$$\alpha(\ell, x, c, c') = \binom{n-2c}{c'} \binom{x-2c}{c'} e^{A_{\alpha}^{1}c' + A_{\alpha}^{2}\sigma} \max_{o \le (\Delta+2)\sigma} \binom{x-2c}{o}.$$

We conclude that, for some constants $A_1, A_2 > 0$,

$$p(\ell, x, c, c') \leq \frac{\varphi_2 \cdot \alpha(\ell, x, c, c') \beta(\ell, x, c, c')}{|\mathcal{F}_n| \cdot |\operatorname{Aut}(\mathcal{F})|} \leq \frac{\varphi_1 \varphi_2 \binom{n-2c}{c'} \binom{x-2c}{c'} e^{A_1 c' + A_2 \sigma} (n - x + c')!}{n!} \left(1 + e^{o(\ell)} \cdot \mathbb{1}_{\ell > Cn^{2/d}}\right) \max_{o \leq (\Delta + 2)\sigma} \binom{x - 2c}{o}^2.$$
(19)

Note that every connected $\tilde{F} \subset F$ with $3 \leq |V(\tilde{F})| \leq n-3$, has $2d+2|E(\tilde{F})| \leq d|V(\tilde{F})|$. Therefore, the *i*-th connected component of J with $\ell_i \geq 2$ and $x_i \leq n-3$ has $\sigma_i \geq \frac{2d-\Delta}{2} \geq \frac{d-2}{2}$. In particular, for J consisting of such connected components, we get $\sigma = \sum_i \sigma_i \geq \frac{2d-\Delta}{2} \cdot c'$. Since

$$x - c' = 2\ell/d + (2 - 2/d)c + 2\sigma/d + \frac{\Delta - d}{d}c' \ge 2\ell/d + (2 - 2/d)c + c',$$
(20)

in a similar way as in (14), we get that

$$\frac{(n-x+c')!}{n!} \le \frac{e^{\frac{(x-c')^2}{n}}}{n^{x-c'}} \le \frac{e^{\frac{(x-c')^2}{n}}}{n^{2\ell/d+(2-2/d)c+c'+2(\sigma-(d-\Delta/2)c')/d}}.$$
(21)

We then conclude that

$$\frac{\varphi_1\varphi_2(n-x+c')!}{n!} \le 4c^{-1/2} \frac{n}{n-2c+1} \cdot \frac{\left(\frac{ed^2n^2}{2c}\right)^c}{n^{2\ell/d+(2-2/d)c+c'+2(\sigma-(d-\Delta/2)c')/d}} \cdot e^{\frac{(x-c')^2}{n}}.$$

The latter implies

$$\frac{\varphi_1\varphi_2(n-x+c')!}{n!}e^{-\frac{(nd/2)^2}{m}}\left(\frac{N}{m}\right)^c \le \frac{4n}{n-2c+1} \cdot \frac{c^{-1/2}\left(\frac{ed^2n^2n^{2/d}}{4cm}\right)^c}{n^{2\ell/d+c'+2(\sigma-(d-\Delta/2)c')/d}} \cdot e^{-\frac{(nd/2)^2}{m} + \frac{(x-c')^2}{n}} \le \frac{4n}{n-2c} \cdot \frac{c^{-1/2}}{n^{2(\ell-c)/d+c'+2(\sigma-(d-\Delta/2)c')/d}} \cdot e^{\frac{(x-c')^2}{n}}.$$
(22)

The last expression is $O(1/\sqrt{c})$ when $\ell = c = O(\sqrt{n})$ (and thus $\sigma = c' = 0$ and x = 2c), and, as we will see soon, it gives the main contribution to the sum in (13).

Let us recall that we have to prove that

$$\left(\sum_{\ell > Cn^{2/d}} \sum_{x,c,c'} + \sum_{\ell \le Cn^{2/d}} \sum_{x-2c > \sqrt{\ln n}} \sum_{c'}\right) p(\ell, x, c, c') e^{-\frac{(nd/2)^2}{m}} \left(\frac{N}{m}\right)^\ell \left(1 + \frac{3dn}{2m}\right)^\ell = o(1).$$
(23)

We choose $0 < \varepsilon' \ll \varepsilon'' \ll \varepsilon$ small enough and consider separately three cases: $\ell \leq Cn^{2/d}$, $Cn^{2/d} < \ell \leq \varepsilon' n$, and $\ell > \varepsilon' n$.

1. $\ell \leq Cn^{2/d}$. We get

$$p(\ell, x, c, c')e^{-\frac{(nd/2)^2}{m}} \left(\frac{N}{m}\right)^{\ell} \left(1 + \frac{3dn}{2m}\right)^{\ell} = O\left(\frac{\binom{n-2c}{c'}\binom{x-2c}{c'}e^{A_1c' + A_2\sigma}\binom{x-2c}{o}^2\binom{N}{m}^{\ell-c}}{\sqrt{c} \cdot n^{2(\ell-c)/d+c'+2(\sigma-(d-\Delta/2)c')/d}}\right), \quad (24)$$

where $o \leq (\Delta + 2)\sigma$ maximises the above expression.

1.1. If $\sigma < \varepsilon'(x - 2c)$, then, since $c' \le \frac{2}{2d - \Delta}\sigma$ and $\ell \ge x - c - c'$, we get

$$\frac{\binom{n-2c}{c'}\binom{x-2c}{c'}e^{A_1c'+A_2\sigma}\binom{x-2c}{o}^2\binom{N}{m}^{\ell-c}}{\sqrt{c}\cdot n^{2(\ell-c)/d+c'}} \le \frac{\binom{n-2c}{c'}e^{\varepsilon''\cdot(x-2c)}}{\sqrt{c}\cdot n^{c'}e^{2(\ell-c)/d}} \le \frac{\left(\frac{e^{1+2/d}}{c'}\right)^{c'}e^{\varepsilon''\cdot(x-2c)}}{\sqrt{c}\cdot e^{2(x-2c)/d}} < \left(\frac{e^{1+2/d}}{c'}\right)^{c'}e^{-\frac{x-2c}{d}}\cdot c^{-1/2}.$$

This bound is not sufficient to prove that summation over all x, c with the same value of x - 2c is small. Nevertheless, letting $\tau := \frac{d^2n^2}{4m}$ and $c =: \tau + t$, we get

$$\begin{split} \sum_{c=0}^{(1+\varepsilon')\tau} \left(\frac{ed^2n^2}{4cm}\right)^c &= \sum_t \left(\frac{e\tau}{\tau+t}\right)^{\tau+t} = e^\tau \sum_t e^t \left(\frac{1}{1+t/\tau}\right)^{\tau+t} \\ &\leq e^\tau \sum_t e^{t-(\tau+t)t/\tau+(t/\tau)^2(\tau+t)/2} \leq e^\tau \sum_t e^{-(1-\varepsilon')t^2/(2\tau)} < 3\sqrt{\tau} \cdot e^\tau. \end{split}$$

Moreover, for $c \notin [(1 - \varepsilon')\tau, (1 + \varepsilon')\tau]$, we get

$$\left(\frac{ed^2n^2}{4cm}\right)^c e^{-\frac{(nd/2)^2}{m}} = e^{-\Omega(n^{2/d})}.$$

Summing up, we get that, for every fixed c' and x - 2c,

$$\sum_{c} p(\ell, x, c, c') e^{-\frac{(nd/2)^2}{m}} \left(\frac{N}{m}\right)^{\ell} \left(1 + \frac{3dn}{2m}\right)^{\ell} = O\left(\left(\frac{e^{1+2/d}}{c'}\right)^{c'} \cdot e^{-(x-2c)/d}\right) + e^{-\Omega(n^{2/d})}.$$

Since for a fixed c and fixed x - 2c, the value of ℓ ranges between c + (x - 2c - c') and $c + (x - 2c - c')\frac{d}{2}$, we get

$$\sum_{\ell \le Cn^{2/d}} \sum_{c} \sum_{x,c': \sigma < \varepsilon'(x-2c)} p(\ell, x, c, c') e^{-\frac{(nd/2)^2}{m}} \left(\frac{N}{m}\right)^{\ell} \left(1 + \frac{3dn}{2m}\right)^{\ell} \\ \le \sum_{c'} \sum_{x-2c > \sqrt{\ln n}} \frac{d}{2} (x - 2c) \left(O\left(\left(\frac{e^{1+2/d}}{c'}\right)^{c'} e^{-(x-2c)/d}\right) + e^{-\Omega(n^{2/d})}\right) = e^{-\Omega(\sqrt{\log n})}.$$
(25)

1.2. If $\sigma \geq \varepsilon'(x-2c)$ and $c' \leq \frac{1}{2d-\Delta}\sigma$, then

$$\frac{\binom{n-2c}{c'}\binom{x-2c}{c'}e^{A_{1}c'+A_{2}\sigma}\binom{x-2c}{o}^{2}\binom{N}{m}^{\ell-c}}{n^{2(\ell-c)/d+c'+2(\sigma-(d-\Delta/2)c')/d}} \leq \frac{\binom{n-2c}{c'}e^{O(x-2c)}}{n^{c'+\sigma/d}} \leq \frac{\binom{e}{c'}^{2'}e^{O(x-2c)}}{n^{\Omega(x-2c)}} = n^{-\Omega(x-2c)}.$$
(26)

1.3. Finally, if $\sigma \geq \varepsilon'(x-2c)$ and $c' \geq \frac{1}{2d-\Delta}\sigma$, then

$$\frac{\binom{n-2c}{c'}\binom{x-2c}{c'}e^{A_1c'+A_2\sigma}\binom{x-2c}{o}^2\binom{N}{m}^{\ell-c}}{n^{2(\ell-c)/d+c'+2(\sigma-(d-\Delta/2)c')/d}} \leq \frac{\binom{n-2c}{c'}e^{O(x-2c)}}{n^{c'}} \leq \frac{\binom{e}{c'}}{c'}e^{O(x-2c)} = (c')^{-c'/2}(x-2c)^{-\Omega(x-2c)}.$$

Applying here exactly the same argument as in the case 1.1, we get

$$\sum p(\ell, x, c, c') e^{-\frac{(nd/2)^2}{m}} \left(\frac{N}{m}\right)^{\ell} \left(1 + \frac{3dn}{2m}\right)^{\ell} \le e^{-\Omega\left(\sqrt{\log n} \cdot \log \log n\right)},\tag{27}$$

where the sum is over all $\ell \leq Cn^{2/d}$, x, c, and c', such that $x - 2c > \sqrt{\ln n}, \sigma \geq \varepsilon'(x - 2c)$, and $c' \geq \frac{1}{2d-\Delta}\sigma.$ **2.** $Cn^{2/d} \leq \ell \leq \varepsilon' n$. From (19) and (22), we get

$$p(\ell, x, c, c')e^{-\frac{(nd/2)^2}{m}} \left(\frac{N}{m}\right)^{\ell} \left(1 + \frac{3dn}{2m}\right)^{\ell} \\ \leq e^{\varepsilon''\ell} \cdot \left(\frac{ed^2n^{2/d}}{2ce^{2/d}}\right)^c \cdot e^{-\frac{(nd/2)^2}{m}} \cdot \frac{\binom{n-2c}{c'}\binom{x-2c}{c'}e^{A_1c'+A_2\sigma}\binom{x-2c}{o}^2 \left(\frac{N}{m}\right)^{\ell-c}}{n^{2(\ell-c)/d+c'+2(\sigma-(d-\Delta/2)c')/d}},$$

where $o \leq (\Delta + 2)\sigma$ maximises the above expression. If $c \leq \frac{1}{4}x$, then $x - 2c > \sqrt{\ln n}$, meaning that the argument from the previous case can be applied here as well, implying exactly the same bound. If $c > \frac{1}{4}x$, then the latter expression is at most

$$e^{-x\frac{\ln C}{10}} \cdot \frac{\binom{n-2c}{c'}\binom{x-2c}{c'}e^{A_1c'+A_2\sigma}\binom{x-2c}{o}^2\binom{N}{m}^{\ell-c}}{n^{2(\ell-c)/d+c'+2(\sigma-(d-\Delta/2)c')/d}} < e^{-x\frac{\ln C}{10}} \cdot \frac{8^{x}e^{(A_1+A_2)x}}{n^{2(\ell-c)/d}\binom{M}{m}^{\ell-c}} \le e^{-x\frac{\ln C}{20}}, \quad (28)$$

whenever $\ln C > 20(A_1 + A_2 + 3).$

3. Finally, let $\ell > \varepsilon' n$.

3.1. We separately consider the case that either $x \ge n-2$, c' = 1, and c = 0, or x' = n, c' = 1, and c = 1. In this case, we cannot apply (20). Nevertheless, since c and c' are bounded, we have the following bound, due to (19):

$$p(\ell, x, c, c')e^{-\frac{(nd/2)^2}{m}} \left(\frac{N}{m}\right)^\ell \left(1 + \frac{3dn}{2m}\right)^\ell \le e^{\varepsilon''\ell} \cdot e^{\frac{(x-c')^2}{n}} \cdot \frac{e^{A_2\sigma} {\binom{x-2c}{o}}^2 \left(\frac{N}{m}\right)^\ell}{n^{2\ell/d + 2\sigma/d}}.$$

The latter bound is $n^{-\Omega(n)}$ when $\sigma > \varepsilon' x$. On the other hand, if $\sigma \leq \varepsilon' x$, then $e^{A_2 \sigma} {\binom{x-2c}{\sigma}}^2 < e^{\varepsilon'' \ell}$ and $x = 2\ell/d + 2\sigma/d + O(1)$. Thus,

$$p(\ell, x, c, c')e^{-\frac{(nd/2)^2}{m}} \left(\frac{N}{m}\right)^{\ell} \left(1 + \frac{3dn}{2m}\right)^{\ell} \le e^{2\varepsilon''\ell} \cdot e^x \left(\frac{e^{-2/d}}{1+\varepsilon}\right)^{\ell} \le e^{3\varepsilon''\ell} \cdot e^{2\ell/d} \left(\frac{e^{-2/d}}{1+\varepsilon}\right)^{\ell} = \left(\frac{e^{3\varepsilon''}}{1+\varepsilon}\right)^{\ell} < e^{-\varepsilon\ell/2}.$$
 (29)

3.2. Otherwise, we can apply (20). Due to (19) and (21),

$$p(\ell, x, c, c')e^{-\frac{(nd/2)^2}{m}} \left(\frac{N}{m}\right)^{\ell} \left(1 + \frac{3dn}{2m}\right)^{\ell} \\ \leq e^{\varepsilon''\ell} \cdot e^{\frac{(x-c')^2}{n}} \cdot \left(\frac{ed^2n^{2/d}}{2ce^{2/d}}\right)^c \cdot \frac{\binom{n-2c}{c'}\binom{x-2c}{c'}e^{A_1c'+A_2\sigma}\binom{x-2c}{o}^2 \left(\frac{N}{m}\right)^{\ell-c}}{n^{2(\ell-c)/d+c'+2(\sigma-(d-\Delta/2)c')/d}}, \quad (30)$$

where $o \leq (\Delta + 2)\sigma$ maximises the above expression. If $c' > \varepsilon' x$, then $n^{c'} = n^{\Omega(n)}$ and $n^{2(\ell-c)/d} > \left(\frac{N}{m}\right)^{\ell-c}$, implying that the right-hand side of (30) is $n^{-\Omega(n)}$. If $c' \leq \varepsilon' x$ and $\sigma > (2d - \Delta)\varepsilon' x$, then $n^{2(\sigma - (d - \Delta/2)c')/d} = n^{\Omega(n)}$, implying the bound $n^{-\Omega(n)}$ as well. Finally, if $(d - \Delta/2)c' \le \sigma \le (2d - \Delta)\varepsilon' x$, then consider separately $c < x/\sqrt{\ln n}$ and $c \ge x/\sqrt{\ln n}$. In the latter case, $\left(\frac{ed^2n^{2/d}}{2ce^{2/d}}\right)^c = e^{-\Omega(n\sqrt{\ln n})}$, implying the bound $e^{-\Omega(n\sqrt{\ln n})}$ for the right-hand side of (30). If $c < x/\sqrt{\ln n}$, then $\ell - c = (1 - o(1))\ell$. Thus, recalling (20) and that $\left(\frac{ed^2n^{2/d}}{2ce^{2/d}}\right)^c = 0$ $e^{O(n^{2/d})} = e^{o(\ell)},$

$$p(\ell, x, c, c')e^{-\frac{(nd/2)^2}{m}} \left(\frac{N}{m}\right)^{\ell} \left(1 + \frac{3dn}{2m}\right)^{\ell} \le \frac{e^{2\varepsilon''\ell + 2\ell/d + (2-2/d)c + 2\sigma/d + \frac{\Delta - d}{d}c'}}{((1+\varepsilon)e^{2/d})^{\ell - c}} \le \frac{e^{3\varepsilon''\ell + 2\ell/d}}{((1+\varepsilon)e^{2/d})^{\ell}} = \frac{e^{3\varepsilon''\ell}}{(1+\varepsilon)^{\ell}} < e^{-\varepsilon\ell/2}.$$
(31)

From (24)–(31), we get that the left-hand side in (23) is at most

$$e^{-\Omega(\sqrt{\log n})} + \sum_{x-2c>\sqrt{\ln n}} \sum_{c',\ell} n^{-\Omega(x-2c)} + e^{-\Omega\left(\sqrt{\log n} \cdot \log \log n\right)} + \\ + \sum_{Cn^{2/d} < \ell \le \varepsilon' n} \sum_{x,c,c'} e^{-x \cdot (\ln C)/20} + \sum_{\ell > \varepsilon' n} \sum_{x,c,c'} \left(e^{-\varepsilon \ell/2} + e^{-\Omega(n\sqrt{\ln n})} \right) = o(1),$$
completing the proof.

completing the proof.

For every $F \in \mathcal{F}_n$ such that there exists F' as in the statement of Claim 5.5, we choose one such F' and put $F \cap F'$ into a multiset \mathcal{F}_n^* . Let **H** be a uniformly random element of \mathcal{F}_n^* .

Claim 5.6. Whp there exists $H' \subset \mathbf{H} \cup \mathbf{W}_2, H' \in \mathcal{F}_n^*$, such that $|\mathbf{H} \cap H'| \leq \ln n$.

Proof. Let $f = \lfloor Cn^{2/d} \rfloor$, $m = m_0$, and \mathcal{B} be the property of graphs on n vertices to have more than $\ln n$ edges. Due to Lemma 4.1, it is sufficient to prove (9).

Fix non-negative integers c and x. Recall that, for $H \in \mathcal{F}_n^*$, we denote by $\mathcal{J}_{\ell,x,c}^H$ the set of all subgraphs $J \subset H$ with ℓ edges, x non-isolated vertices, and c connected components (excluding isolated vertices). Denote by $p^{H}(\ell, x, c)$ the probability that $\mathbf{H} \cap H \in \mathcal{J}_{\ell, x, c}^{H}$.

Since H consists of isolated edges and at most $\sqrt{\ln n}$ other edges, we get that $\ell \leq c + \sqrt{\ln n}$. Due to Claim 3.2 and Claim 3.3, for some constant A > 0 that does not depend on ε ,

$$p^{H}(\ell, x, c) \leq \frac{\binom{\lfloor Cn^{2/d} \rfloor}{c} e^{A\ell} (n - x + c)! / |\operatorname{Aut}(F)|}{|\mathcal{F}_{n}^{*}|} = (1 + o(1)) \frac{\binom{\lfloor Cn^{2/d} \rfloor}{c} e^{A\ell} (n - x + c)!}{n!}.$$

Since $d \ge 4$ and since every non-trivial subgraph of F has edge boundary of size at least 2d-2,

$$p^{H}(\ell, x, c) \stackrel{(18)}{\leq} \frac{\left(\frac{eCn^{2/d}}{c}\right)^{c} e^{(A+o(1))\ell}}{n^{2\ell/d+(1-2/d)c}} \leq \frac{\left(\frac{e^{1+A+o(1)}C}{c}\right)^{c} e^{(A+o(1))\sqrt{\ln n}}}{n^{2\ell/d}}.$$

Therefore,

$$\begin{split} \sum_{\ell} \Pi_{\mathcal{B}_{\ell}}^{H} \left(\left(1 + \frac{3f}{m} \right) \frac{N}{m} \right)^{\ell} &\leq \sum_{\ell > \ln n} \sum_{x,c} p^{H}(\ell, x, c) \left(\frac{1 + o(1)}{\varepsilon} \cdot n^{2/d} \right)^{\ell} \\ &\leq n^{2} \sum_{\ell > \ln n} \left(\frac{e^{1 + A}C}{\ell - \sqrt{\ln n}} \right)^{\ell - \sqrt{\ln n}} e^{A\sqrt{\ln n}} \left(\frac{1 + o(1)}{\varepsilon} \right)^{\ell} = o\left(\frac{1}{n^{2}} \right), \end{split}$$
mpleting the proof.

completing the proof.

5.2.2. Proof of Theorem 5.4. Let $\mathbf{G} \sim G(n,p)$, where $p = (1+2\varepsilon)(e/n)^{2/d}$. By [17, Corollary 1.16], whp there exists a multiset $\mathcal{F}_n^* = \mathcal{F}_n^*(\mathbf{G})$ of graphs of size $\lfloor \ln n \rfloor$ comprising a single subgraph H of almost every $F \in \mathcal{F}_n$ so that $H \cup \mathbf{G}$ contains some $F' \in \mathcal{F}_n$ and H has at most $\sqrt{\ln n}$ non-isolated edges.

Let X be the number of $H \in \mathcal{F}_n^*$ that are subgraphs of $\mathbf{G}' \sim G(n, p' = \varepsilon n^{-2/d})$, sampled independently of \mathbf{G} . We get

$$\mathbb{E}X = |\mathcal{F}_n^*| \cdot p^{\prime \lfloor \ln n \rfloor} = (1 - o(1))|\mathcal{F}_n| \cdot p^{\prime \lfloor \ln n \rfloor} = \omega(1).$$

Let \mathcal{B} be the set of non-empty graphs. Due to the definition (8), the fact that every $H \in \mathcal{F}_n^*$ has at most $\sqrt{\ln n}$ non-isolated edges, Claim 3.2, Claim 3.3, and estimate (18), for some constant A > 0,

$$\frac{\operatorname{Var} X}{(\mathbb{E}X)^2} \leq \frac{\max_{H \in \mathcal{F}_n^*} \sum_{\ell \geq 1} \left(\Pi_{\mathcal{B}_{\ell}}^H \cdot |\mathcal{F}_n^*| \cdot p'^{\lfloor \ln n \rfloor - \ell} \right)}{\mathbb{E}X} = \max_{H \in \mathcal{F}_n^*} \sum_{\ell \geq 1} \Pi_{\mathcal{B}_{\ell}}^H p'^{-\ell} \\
\leq \sum_{\ell \geq 1} \sum_{x,c} (1 - o(1)) \frac{\binom{\lfloor \ln n \rfloor}{c} e^{A\ell}}{n^{2\ell/d + (1 - 2/d)c}} \left(\frac{1}{\varepsilon} \cdot n^{2/d} \right)^{\ell} \\
\leq \sum_{\ell \geq 1} \sum_{c \geq 1} O(\ell) \left(\frac{e^{A + 1} \lfloor \ln n \rfloor}{\varepsilon c \sqrt{n}} \right)^c e^{A\sqrt{\ln n}} \left(\frac{1}{\varepsilon} \right)^{\sqrt{\ln n}} = n^{-1/2 + o(1)}.$$

This completes the proof of Theorem 5.4, due to Chebyshev's inequality.

Remark 5.7. As noted in Section 1.4 (and evident from the current proof), we have $VarX_F =$ $O((\mathsf{E}X_F)^2)$. Consequently, Theorem 5.4 follows from Paley–Zygmund inequality and [13, Theorem 2.3]. However, this holds only when there are very few subgraphs whose edge boundary is smaller than 2d. In particular, for d = 3, there are at least linearly many subgraphs that do not satisfy this condition. Specifically, any pair of incident edges forms a subgraph on three vertices with an edge boundary of at most five. This observation prevents the direct application of both Riordan's second-moment argument [31] and the fragmentation method in its current form, necessitating new ideas.

6. Proof of Theorem 1.7

The aim of this section is to prove the following.

Theorem 6.1. For every $\varepsilon > 0$, whp $\mathbf{G} \sim G\left(n, (1+\varepsilon)\sqrt{e/n}\right)$ contains the second power of a Hamilton cycle.

Theorem 6.1 immediately implies Theorem 1.7.

Everywhere in this section F is a second power of a cycle on [n] and \mathcal{F}_n is the set of all isomorphic copies of F on [n]. Let us recall that we call a subgraph $H \subset F \in \mathcal{F}_n$ closed if it is the second power of a path.

The proof of Theorem 6.1 is organised as follows. In Section 6.1, we define a subfamily of \mathcal{F}_n that is convenient for improving fragments: we fix $\omega(\sqrt{n}/\ln n)$ diamonds (4-vertex graphs obtained by removing a single edge from a 4-clique) and include into the family all $F \in \mathcal{F}_n$ that contain these diamonds at equal distances. Then we prove that typical graphs from this subfamily have a fragment of size $O(\sqrt{n})$. In Section 6.2, we improve fragments by moving closed subgraphs of length $\Omega(\ln n)$ 'inside' the diamonds and then prove that there exists a way to assign closed subgraph to the diamonds so that the multiset of improved fragments have "small spreads" — see Claim 6.6. Due to this claim, in the next fragmentation round in Section 6.3 we get fragments of size $O(n^{1/5})$. In Section 6.4, we show that sprinkling $\varepsilon p {n \choose 2}$ extra edges is enough to cover at least one fragment of size $O(n^{1/5})$ whp.

6.1. Day 0: getting fragments of size $O(\sqrt{n})$. Let

$$\mathbf{W} \sim G(n,m), \quad \text{where} \quad m = \left\lfloor (1+\varepsilon)\sqrt{e/n} \cdot N \right\rfloor.$$

In what follows, for the sake of convenience, \mathcal{F}_n is the family of all rooted oriented second powers of a cycle on [n], that is, each cycle $F \in \mathcal{F}_n$ has a specified vertex — its root — and is oriented, starting from the root, in one of the two directions. Clearly, $|\mathcal{F}_n| = n!$ and each $F \in \mathcal{F}_n$ induces a linear order on [n]; we denote the respective permutation acting on [n] by π_F .

Let $w = w(n) = \omega(1)$ be a slowly increasing function. Let

$$\chi := \left\lfloor \frac{w\sqrt{n}}{\ln n} \right\rfloor \tag{32}$$

and let $n = (n_1 + 4) + \ldots + (n_{\chi} + 4)$, where n_1, \ldots, n_{χ} are positive integers that differ by at most 1. Fix a tuple $\overrightarrow{D} = (D_1, \ldots, D_{\chi})$ of disjoint diamonds (K_4 minus an edge) D_1, \ldots, D_{χ} on [n]. We then consider the family $\mathcal{F}_n(\overrightarrow{D}) \subset \mathcal{F}_n$ of all F such that,

- for every $i \in [\chi], D_i \subset F$,
- the order of the diamonds as well as the order of vertices in the diamonds is aligned with π_F and the first vertex of D_1 coincides with the root of F,
- for every $i \in [\chi]$, the number of vertices between D_i and D_{i+1} in F equals n_i (in the cyclic order, i.e. we here set $D_{\chi+1} := D_1$).

Fix $F \in \mathcal{F}_n(\overrightarrow{D})$ and $W \in \binom{\binom{[n]}{2}}{m}$. An (F, W)-fragment is a subgraph H of some $F' \in \mathcal{F}_n(\overrightarrow{D})$, such that $F' \subset F \cup W$ and $H = F \cap F'$. We will show that still whp there exists an (F, \mathbf{W}) fragment H of size $O(\sqrt{n})$. In particular, this fragment has to contain all diamonds D_1, \ldots, D_{χ} . This will allow us to refine the fragment H by spreading evenly the edges of large closed subgraphs in H among the diamonds and, thus, to get rid of all the closed subgraphs of size $\Omega(\ln n)$.

We first expose the edges of \mathbf{W} that belong to $D_1 \cup \ldots \cup D_{\chi}$ and treat them as fixed. Since $|D_1 \cup \ldots \cup D_{\chi}| = 5\chi = o(\sqrt{n})$, whp $\mathbf{W} \cap (D_1 \cup \ldots \cup D_{\chi})$ is empty. So, we may further condition on this event, or, in other words, assume that \mathbf{W} is a uniformly random *m*-element subset of $\binom{[n]}{2} \setminus (D_1 \cup \ldots \cup D_{\chi})$. We also let $\mathcal{F}_n^*(\overrightarrow{D})$ be the set of all $F \setminus (D_1 \cup \ldots \cup D_{\chi})$, $F \in \mathcal{F}_n(\overrightarrow{D})$.

For $F \in \mathcal{F}_n^*(\overrightarrow{D})$ and $W \subset {[n] \choose 2} \setminus (D_1 \cup \ldots \cup D_{\chi})$, let $\mathcal{M}(F, W)$ be the set of all $F' \in \mathcal{F}_n^*(\overrightarrow{D})$ such that $F' \subset F \cup W$. In particular, $F \in \mathcal{M}(F, W)$. An easy counting argument shows that typically $\mathcal{M}(F, \mathbf{W})$ is large. Let

$$M := M(t) = |\mathcal{F}_n^*(\overrightarrow{D})| \binom{N-2n}{m-t} / \binom{N-5\chi}{m+(2n-5\chi)-t}$$

be the expected number of $F' \in \mathcal{F}_n^*(\overrightarrow{D})$ such that F' belongs to a uniformly random subset of $\binom{[n]}{2} \setminus (D_1 \cup \ldots \cup D_{\chi})$ of size $m + (2n - 5\chi) - t$. Notice that M(t) is actually large for every t:

$$M = \frac{n!}{e^{O(\sqrt{n}w)}} \frac{(m-t+(2n-5\chi))^{m-t+(2n-5\chi)} (N-2n)^{N-2n}}{(m-t)^{m-t} (N-5\chi)^{N-5\chi}}$$
$$= \frac{n^n}{e^{n+O(\sqrt{n}w)}} \left(\frac{m-t}{N-5\chi}\right)^{2n-5\chi} \left(1 - \frac{2n-5\chi}{m+(2n-5\chi)-t}\right)^{-m-(2n-5\chi)+t} \left(1 + \frac{2n-5\chi}{N-2n}\right)^{-N+2n}$$
$$= \frac{(1+\varepsilon)^{2n}}{e^{O(\sqrt{n}w)}} \left(\frac{m-t}{m} \cdot \frac{N}{N-5d}\right)^{2n} e^{\Omega(\sqrt{n})} = (1+\varepsilon)^{(2-o(1))n}.$$

We claim that whp $|\mathcal{M}(F, \mathbf{W})| = \Omega(M)$ and that typically most (F, \mathbf{W}) -fragments have size $O(\sqrt{n})$. Although the conclusion of Lemma 4.1 is sufficient for our goals, below we state its stronger version which holds for symmetric families \mathcal{F}_n (i.e. when all graphs in the family are isomorphic). In this version, we replace a random element of $\mathcal{F}_n^*(\vec{D})$ with any fixed element. Since the proofs are almost identical, we postpone them to Appendix B.

Claim 6.2. Let $F \in \mathcal{F}_n^*(\overrightarrow{D})$ and let **W** be a uniformly random *m*-element subset of $\binom{[n]}{2} \setminus (D_1 \cup \dots \cup D_{\chi})$. Let $\delta(n) \to 0$ as $n \to \infty$. Then, for every $t \in \{0, 1, \dots, 2n - 5\chi\}$,

$$\mathbb{P}\left(|\mathcal{M}(F,\mathbf{W})| < \delta(n)M \mid |F \cap \mathbf{W}| = t\right) \le \delta(n).$$

Let C > 0 be large enough. Set $\ell_0 = \lfloor C\sqrt{n} \rfloor$. Let \mathcal{B} be the property of graphs on n vertices to have more than ℓ_0 edges. We claim that (9) holds.

Claim 6.3. Sample a uniformly random $\mathbf{F} \in \mathcal{F}_n^*(\overrightarrow{D})$. For $\ell \in \mathbb{N}$, let

$$\Pi_{\ell} := \mathbb{P}(F \cap \mathbf{F} \text{ has } \ell \text{ edges}).$$
(33)

Then

$$\sum_{\ell \ge \ell_0 + 1} \Pi_{\ell} \cdot e^{-\frac{4n^2}{m}} \left((1 + o(1)) \frac{N}{m} \right)^{\ell} = o\left(\frac{1}{n}\right).$$
(34)

The proof of Claim 6.3 relies on estimations similar to those provided in Claim 3.1 (see Claim 6.8 in Section 6.5). However, we cannot apply Claim 3.1 here directly because of the fixed diamonds, that slightly affect our calculations. The proof occupies Section 6.5. Despite obvious similarities and repetitions, we decided to keep the entire proof of Claim 6.8 to make the proof of Theorem 6.1 self-contained.

For $\ell \in \mathbb{N}$, let $\mathcal{M}_{\ell}(F, W)$ be the set of all $F' \in \mathcal{M}(F, W)$ such that $|F' \cap F| > \ell$. Let us call (F, W) ℓ -bad, if $|\mathcal{M}_{\ell}(F, W)| > \frac{1}{\sqrt{n}} |\mathcal{M}(F, W)|$. Claim 6.3 implies that (F, \mathbf{W}) is not ℓ_0 -bad whp.

Claim 6.4. Let $F \in \mathcal{F}_n^*(\overrightarrow{D})$ and let $\mathbf{W} \subset {\binom{[n]}{2}} \setminus (D_1 \cup \ldots \cup D_{\chi})$ be a uniformly random *m*-subset. Then whp (F, \mathbf{W}) is not ℓ_0 -bad.

That means that whp we may replace most of $F \in \mathcal{F}_n(\vec{D})$ with fragments of sizes at most ℓ_0 (and each fragment contains all $D_i(F)$). The proof of Claim 6.4 is presented in Appendix B.

Let $Y_{\overrightarrow{D}}$ be the number of $F \in \mathcal{F}_n(\overrightarrow{D})$ that have an (F, \mathbf{W}) -fragment of size at most $\ell_0 + 5\chi$. From Claim 6.2 and Claim 6.4, we get that $\mathbb{E}Y_{\overrightarrow{D}} = |\mathcal{F}_n(\overrightarrow{D})|(1 - o(1))$. By Markov's inequality, for every $\delta > 0$,

$$\mathbb{P}\left(Y_{\overrightarrow{D}} < (1-\delta)|\mathcal{F}_n(\overrightarrow{D})|\right) = \mathbb{P}\left(|\mathcal{F}_n(\overrightarrow{D})| - Y_{\overrightarrow{D}} > \delta|\mathcal{F}_n(\overrightarrow{D})|\right) \le \frac{\mathbb{E}\left(|\mathcal{F}_n(\overrightarrow{D})| - Y_{\overrightarrow{D}}\right)}{\delta|\mathcal{F}_n(\overrightarrow{D})|} = o(1).$$
(35)

Therefore, whp $Y_{\overrightarrow{D}} \ge (1 - o(1)) |\mathcal{F}_n(\overrightarrow{D})|.$

6.2. Improving fragments. For $F \in \mathcal{F}_n(\vec{D})$ that have an (F, \mathbf{W}) -fragment of size at most $\ell_0 + 5\chi$, let $H = H(F) \subset F$ be a typical (for reasons that will be explained later) fragment of such a size. All graphs from $\mathcal{F}_n(\vec{D})$ contain every D_i , $i \in [\chi]$. Since $H = F \cap F'$ for some $F' \in \mathcal{F}_n(\vec{D})$, we get that every D_i is a subgraph of H. Since the distance in F between any two D_i is bigger than $2C\sqrt{n}$, we get that the connected components of H that contain different $D_i(F)$ are disjoint. Consider all inclusion-maximal closed subgraphs in H with at least $\ln n/\sqrt{w}$ vertices that do not overlap with any of the diamonds $D_i(F)$. For each such second power of a path $v_1 \dots v_h$, we "cut" it into pieces of length $\mu := \lfloor \ln n/\sqrt{w} \rfloor$ in the following way: v_1v_2 ; $v_3 \dots v_{2+\mu}$; $v_{3+\mu} \dots v_{2+2\mu}$; \dots We let the last piece $v_{2+(j+1)\mu}v_{2+(j+1)\mu+1} \dots$ to be of length at least 2 and at most $\mu + 1$. From every such inclusion-maximal closed subgraph, remove all pieces (i.e. all the listed vertices as well as the edges that contain them) but the first and the last one and glue the first and the last so that they make up the second power of a path $(v_1v_2v_{2+(j+1)\mu+1}\dots)$. Let $\mathcal{P}_H = \{P_1, \dots, P_s\}$ be the set of all pieces of length μ that we have removed $(P_1, \dots, P_s$ respect the order induced by F). Clearly, the number of pieces s is less than $2\ell_0/\mu < \chi/3$, where χ is defined in (32).

We then choose different $j_1, \ldots, j_s \in [\chi]$, according to some rule, that will be explained later. For every $i \in [s]$, we insert the piece $P_i = (v_1^i \ldots v_{\mu}^i)$ between the vertices $u_1^{j_i}, u_2^{j_i}$ and $u_3^{j_i}, u_4^{j_i}$ of $D_{j_i} = (u_1^{j_i}u_2^{j_i}u_3^{j_i}u_4^{j_i})$ making up the square of a path $(u_1^{j_i}u_2^{j_i}v_1^i \ldots v_{\mu}^iu_3^{j_i}u_4^{j_i})$. We get a *smoothed* fragment H' with exactly the same number of edges as H has. The crucial observation is that $H' \cup \mathbf{W}$ still has a subgraph from \mathcal{F}_n . Moreover, this smoothed fragment satisfies the important 'smoothness' requirement: all powers of paths that it contains have length $o(\ln n)$.

So, this modification of fragments allows to avoid problematic large closed subgraphs. However, it may reduce the total number of *different* fragments, that will eventually affect moments calculation. Let $\mathcal{H}_n = \{H(F), F \in \mathcal{F}_n(\overrightarrow{D})\}$ be the (**W**-random) multiset of original (F, \mathbf{W}) fragments (we choose one typical fragment per F), and let \mathcal{H}'_n be the set of smoothed (F, \mathbf{W}) fragments. Although each $H \in \mathcal{H}_n$ maps to its smoothed version $H' \in \mathcal{H}'_n$, this correspondence φ is not necessarily injective. Luckily, every pre-image is not large if we choose the rule of how P_i are matched to D_{j_i} randomly. We now explain how this matching is chosen. Consider a binomial random bipartite graph **B** with parts $V = [\chi]$ and U = [n] and with edge probability $\beta := n^{-1/3}/w$. Then, we insert every $P_i = (v_1^i \dots v_{\mu}^i)$ into some D_{j_i} so that there exists an edge between $j_i \in V$ and $\pi_F(v_1^i) \in U$ in **B**. Clearly, to make it possible for all P_i simultaneously, we need the graph $\mathbf{B}[V \cup \{\pi_F(v_1^i), i \in [s]\}]$ to have a matching that covers all vertices $\pi_F(v_1^i), i \in [s]$ (recall that $s < \chi$). This is typically possible due to the following claim.

Claim 6.5. Let $(\mathbf{F}, \mathbf{W}, \mathbf{F}')$ be a random vector independent of \mathbf{B} , where

- **F** is a uniformly random element of $\mathcal{F}_n(\vec{D})$,
- W is a uniformly random graph from $\binom{\binom{[n]}{2}}{m} \binom{\binom{[n]}{2}}{m}$ chosen independently of F, and
- \mathbf{F}' is a uniformly random subgraph of $\mathbf{F} \cup \mathbf{W}$ that belongs to $\mathcal{F}_n(\overrightarrow{D})$.

Consider the fragment $H := \mathbf{F} \cap \mathbf{F}'$ and the respective pieces $P_i = (v_1^i \dots v_{\mu}^i), i \in [s]$, defined above. Then whp the graph $\mathbf{B}[V \cup \{\pi_{\mathbf{F}}(v_1^i), i \in [s]\}]$ has a matching that covers all vertices $\pi_{\mathbf{F}}(v_1^i), i \in [s]$.

Proof. Let Σ be the set of all (F, W, F') such that $\mathbb{P}(\mathbf{F} = F, \mathbf{W} = W, \mathbf{F}' = F') > 0$ and $|F \cap F'| \leq \ell_0 + 5\chi$. For $(F, W, F') \in \Sigma$ and a bipartite graph B on $V \times U$, let us call the tuple (B, F, W, F') well-mixed, if $B[V \cup \{\pi_F(v_1^i), i \in [s]\}]$ has a matching that covers all vertices $\pi_F(v_1^i), i \in [s]$, that are defined by $H := F \cap F'$. For a fixed set $U' \subset U$ of size exactly χ , whp there exists a perfect matching between V and U' in $\mathbf{B}[V \cup U']$ (see, e.g., [17, Theorem 4.1]). Therefore, recalling that the bound $|F \cap F'| \leq \ell_0 + 5\chi$ implies that the number of pieces that we moved in H is less than χ , we get that, for any fixed $(F, W, F') \in \Sigma$, whp (\mathbf{B}, F, W, F') is well-mixed. Also, due to Claim 6.2 and Claim 6.4, whp $(\mathbf{F}, \mathbf{W}, \mathbf{F}') \in \Sigma$.

Then,

$$\mathbb{P}((\mathbf{B}, \mathbf{F}, \mathbf{W}, \mathbf{F}') \text{ is well-mixed})$$

$$\geq \sum_{(F,W,F')\in\Sigma} \mathbb{P}((\mathbf{B}, F, W, F') \text{ is well-mixed} \mid \mathbf{F} = F, \mathbf{W} = W, \mathbf{F}' = F')$$

$$\times \mathbb{P}(\mathbf{F} = F, \mathbf{W} = W, \mathbf{F}' = F')$$

$$= \sum_{(F,W,F')\in\Sigma} \mathbb{P}((\mathbf{B}, F, W, F') \text{ is well-mixed}) \times \mathbb{P}(\mathbf{F} = F, \mathbf{W} = W, \mathbf{F}' = F')$$

$$= (1 - o(1))\mathbb{P}((\mathbf{F}, \mathbf{W}, \mathbf{F}') \in \Sigma) = 1 - o(1),$$

completing the proof.

From Claim 6.5, we immediately get that there exists a bipartite graph B on $V \times U$ so that every vertex in V has degree $(1 + o(1))n^{2/3}/w$ (due to the Chernoff bound, say) and that whp $B[V \cup \{\pi_{\mathbf{F}}(v_1^i), i \in [s]\}\]$ has a matching that covers all vertices $\pi_{\mathbf{F}}(v_1^i), i \in [s]$, that correspond to $H = \mathbf{F} \cap \mathbf{F}'$, where $(\mathbf{F}, \mathbf{F}')$ is defined in the statement of the claim.

Having this rule of matching the diamonds with the pieces of closed subgraphs, we are now ready to prove that φ is close enough to an injection. In what follows, we consider only those (F, W) that generate a fragment that satisfies the matching rule. Formally, for $F \in \mathcal{F}_n(\overrightarrow{D})$ and $W \in \binom{\binom{[n]}{2} \setminus (D_1 \cup \ldots \cup D_{\chi})}{m}$, we say that $F' \in \mathcal{F}_n(\overrightarrow{D})$ is (F, W)-nice, if

•
$$F' \subset F \cup W$$
,
• $|F' \cap F| \le \ell_0 + 5\chi$,

• the graph $B[V \cup \{\pi_F(v_1^i), i \in [s]\}]$ has a matching that covers all vertices $\pi_F(v_1^i), i \in [s]$, defined by the fragment $H = F \cap F'$.

Let Σ be the set of all pairs $\left(F \in \mathcal{F}_n(\overrightarrow{D}), W \in \binom{\binom{[n]}{2} \setminus (D_1 \cup \ldots \cup D_{\chi})}{m}\right)$ that have an (F, W)-nice F'. Due to the choice of B and due to Claims 6.2, 6.4, and 6.5, whp $(\mathbf{F}, \mathbf{W}) \in \Sigma$. Indeed,

$$\mathbb{P}((\mathbf{F}, \mathbf{W}) \in \Sigma) = \mathbb{P}(\text{there exists an } (\mathbf{F}, \mathbf{W})\text{-nice } F')$$
$$\geq \mathbb{P}(\mathbf{F}' \text{ is } (\mathbf{F}, \mathbf{W})\text{-nice}) = 1 - o(1), \tag{36}$$

where $(\mathbf{F}, \mathbf{W}, \mathbf{F}')$ is defined in the statement of Claim 6.5.

We now notice that the there exists a natural bijection between *disjoint* families $\mathcal{F}_n(\overrightarrow{D})$, $\mathcal{F}_n(\overrightarrow{D'})$ for different tuples of diamonds \overrightarrow{D} and $\overrightarrow{D'}$. In particular, for $Y := \sum_{\overrightarrow{D}} Y_{\overrightarrow{D}}$ we get that

$$\mathbb{E}Y = \sum_{\overrightarrow{D}} \mathbb{E}Y_{\overrightarrow{D}} = (1 - o(1)) \sum_{\overrightarrow{D}} |\mathcal{F}_n(\overrightarrow{D})| = (1 - o(1))|\mathcal{F}_n|$$

In the same way as in (35), we get that whp $Y \ge (1 - o(1))|\mathcal{F}_n|$. Since any $F \in \mathcal{F}_n$ belongs to $\mathcal{F}_n(\overrightarrow{D})$ for some uniquely defined \overrightarrow{D} , we get the definition of Σ naturally extends to pairs (F, W) for an arbitrary $F \in \mathcal{F}_n$. Let a uniformly random element \mathbf{F} of \mathcal{F}_n , uniformly random elements $\mathbf{F}_{\overrightarrow{D}} \in \mathcal{F}_n(\overrightarrow{D})$, a uniformly random *m*-subset $\mathbf{W} \in \binom{\binom{[n]}{2}}{m}$, and uniformly random *m*-subsets $\mathbf{W}_{\overrightarrow{D}} \in \binom{\binom{[n]}{2} \setminus (D_1 \cup \ldots \cup D_{\chi})}{m}$ be chosen independently. Due to (36),

$$\mathbb{P}((\mathbf{F}, \mathbf{W}) \in \Sigma) = \sum_{\overrightarrow{D}} \mathbb{P}((\mathbf{F}, \mathbf{W}) \in \Sigma \mid \mathbf{F} \in \mathcal{F}_n(\overrightarrow{D})) \cdot \mathbb{P}(\mathbf{F} \in \mathcal{F}_n(\overrightarrow{D}))$$
$$= \sum_{\overrightarrow{D}} \mathbb{P}((\mathbf{F}_{\overrightarrow{D}}, \mathbf{W}) \in \Sigma) \cdot \mathbb{P}(\mathbf{F} \in \mathcal{F}_n(\overrightarrow{D}))$$

For a fixed \overrightarrow{D} , we get

$$\mathbb{P}((\mathbf{F}, \mathbf{W}) \in \Sigma) = \mathbb{P}((\mathbf{F}_{\overrightarrow{D}}, \mathbf{W}) \in \Sigma, \mathbf{W} \cap (D_1 \cup \ldots \cup D_{\chi}) = \emptyset)$$
$$= (1 - o(1)) \cdot \mathbb{P}((\mathbf{F}_{\overrightarrow{D}}, \mathbf{W}) \in \Sigma \mid \mathbf{W} \cap (D_1 \cup \ldots \cup D_{\chi}) = \emptyset)$$
$$= (1 - o(1)) \cdot \mathbb{P}((\mathbf{F}_{\overrightarrow{D}}, \mathbf{W}_{\overrightarrow{D}}) \in \Sigma) \stackrel{(36)}{=} 1 - o(1).$$

Let us fix any $W \in \binom{\binom{[n]}{2}}{m}$. Let Σ_W be the set of all $F \in \mathcal{F}_n$ such that $(F, W) \in \Sigma$. For every $F \in \Sigma_W$, consider an (F, W)-nice $F' := F'(F) \in \mathcal{F}_n$. Let $H(F) := F \cap F'$ and H'(F) be the smoothed version of H(F) (with respect to the matching rule B).

Claim 6.6. For any graph S, let $\mathcal{S}(S)$ be the set of all graphs $F \in \Sigma_W$ such that H'(F) contains S as a subgraph. Then

$$|\mathcal{S}(S)| \le 16^{\ell(S)} (n - x(S) + c(S))! n^{c(S)/6}.$$

Proof of Claim 6.6. Let $T_1, \ldots, T_{c(S)}$ be the connected components of S. We fix an arbitrary root in every connected component of S and order edges in every connected component in a way such that edges that are closer to the root appear earlier. Let

- $\alpha: E(S) \to [4]$ be an arbitrary function;
- $A \subset V(S)$ be such that each T_i has at most two vertices in common with A.

The pair (α, A) identifies a non-negative integer $a \leq \max\{c(S), \chi\}$ that will be explicitly defined later. We also let

• $\tau_1: [a] \to [\lfloor 2n^{2/3}/w \rfloor], \ \tau_2: [a] \to [\lfloor \ln n/4 \rfloor]$ be arbitrary functions;

- $\rho: [a] \to [\chi]$ be in injection;
- $\pi: [n x(S) + c(S) a] \rightarrow [n x(S) + c(S) a]$ be a bijection.

Let us show that the tuple $\mathbf{x} := (A, \alpha, \tau_1, \tau_2, \rho, \pi)$ identifies a graph $F \in \Sigma_W$ such that $S \subset H'(F)$. It would immediately imply the desired assertion. Below, we describe the algorithm of reconstruction of F.

- (1) Reconstruct the order (and their position in F[V(S)]) of vertices in every component of S according to α : for every $i \in [c(S)]$, start from the root of T_i and follow the edges according to the fixed order; at every step, one of the ends of the current edge e is already placed, so there are at most four possibilities to place this edge, and then $\alpha(e)$ defines its position.
- (2) Choose the pieces in S that were moved in H(F): For every connected component T_i , if $|A \cap T_i| \leq 1$, then T_i does not contain vertices of a piece that was moved. Otherwise, let $A \cap V(T_i) = \{v_i, u_i\}$, where $v_i < u_i$, according to the linear order of vertices in T_i identified at the previous step. Let the unique "piece" $P_i \subset T_i$ have all edges of T_i induced by the set of vertices that are between v_i and u_i .
- (3) Let \mathcal{I}_A be the set of all $i \in [c(S)]$ such that P_i is defined and $P_i \neq T_i$. We also let $a = |\mathcal{I}_A|$. For every $i \in \mathcal{I}_A$, we choose the index $\eta(i)$ of the respective diamond in F according to the injection ρ . Then, we move pieces P_i , $i \in \mathcal{I}_A$, to their original places in F. For every $i \in \mathcal{I}_A$, the position $\eta(i)$ of the respective diamond in $[\chi]$ is identified. Therefore, the position of the first vertex of the entire piece in F containing P_i is adjacent to $\eta(i)$ in B, where $\eta(i)$ has degree at most $(1 + o(1))n^{2/3}/w$. Then v_i is at most $(0.1 \cdot \ln n)$ -far from this vertex in F and we may identify its position in F according to $(\tau_1(i), \tau_2(i))$.
- (4) Finally, we order all the remaining $n x(S) + c(S) |\mathcal{I}(A)|$ connected components of S (including the remaining singletons) according to the permutation π and complete the construction of F.
- Thus, $|\mathcal{S}(S)|$ is at most the number of all possible tuples **x** which, in turn, does not exceed

$$\max_{a \le \max\{c(S),\chi\}} 4^{\ell(S)} 2^{x(S)} (n^{2/3} \ln n/(2w))^a \frac{\chi!}{(\chi - a)!} (n - x(S) + c(S) - a)!$$

$$\le 16^{\ell(S)} (n - x(S) + c(S))! \max_{a \le c(S)} \left(\frac{\chi n^{2/3} \ln n/(2w)}{n - x(S) + c(S) - a} \right)^a$$

$$\le 16^{\ell(S)} (n - x(S) + c(S))! \left(\frac{\chi n^{2/3} \ln n/(2w)}{n - x(S)} \right)^{c(S)}.$$

Due to (32),

$$\begin{aligned} |\mathcal{S}(S)| &\leq 16^{\ell(S)} (n - x(S) + c(S))! \left(\frac{n^{7/6}}{2(n - 2\ell_0 - 4\chi)}\right)^{c(S)} \\ &\leq 16^{\ell(S)} (n - x(S) + c(S))! n^{c(S)/6}, \end{aligned}$$

completing the proof.

6.3. Day 1: getting fragments of size at most $n^{1/5}$. We are now ready for the second fragmentation round. From the previous round, we have a sufficiently large multiset of smoothed fragments: For all $F \in \mathcal{F}_n$ such that $(F, \mathbf{W}) \in \Sigma$, we choose an (F, W)-nice F', and add the

smoothed version of the fragment $F \cap F'$ to the new multiset $\mathcal{H}^{(1)}$ (the cardinality of $\mathcal{H}^{(1)}$ equals the number of all F such that $(F, \mathbf{W}) \in \Sigma$). Clearly, we may assume that every member of this family has size exactly $\ell_0 + 5\chi$. Next, we expose $\mathbf{W}^{(1)} \sim G(n, m')$ with $m' = \lfloor \varepsilon {n \choose 2} / \sqrt{n} \rfloor$. For $H \in \mathcal{H}^{(1)}$ and $W \subset {[n] \choose 2}$, let $\mathcal{M}^*(H, W)$ be the set of all $H' \in \mathcal{H}^{(1)}$ such that $H' \subset H \cup W$. For $\ell \in \mathbb{N}$, let $\mathcal{M}^*_{\ell}(H, W)$ be the set of all $H' \in \mathcal{M}^*(F, W)$ such that $|H' \cap H| > \ell$. Let $\delta(n) = o(1)$ be a slowly decreasing function. Let us call (H, W) ℓ -bad, if $|\mathcal{M}^*_{\ell}(H, W)| > \delta_n |\mathcal{M}^*(H, W)|$. Let \mathbf{H} be a uniformly random element of $\mathcal{H}^{(1)}$ sampled independently of $\mathbf{W}^{(1)}$. We shall prove that $(\mathbf{H}, \mathbf{W}^{(1)})$ is not $\lceil n^{1/5} \rceil$ -bad whp. That would mean that whp we may replace most of $H \in \mathcal{H}^{(1)}$ with fragments of sizes at most $\lceil n^{1/5} \rceil$.

Claim 6.7. Whp $(\mathbf{H}, \mathbf{W}^{(1)})$ is not $\lceil n^{1/5} \rceil$ -bad.

Proof. For $\ell \in \mathbb{N}$ and $H \in \mathcal{H}^{(1)}$, let $\Pi_{\ell}^{H} := \mathbb{P}(H \cap \mathbf{H} \text{ has } \ell \text{ edges})$. We let $f = \ell_0 + 5\chi$ and \mathcal{B} be the set of all graphs on n vertices with more than $\lceil n^{1/5} \rceil$ edges. We then apply Lemma 4.1 with these f and \mathcal{B} . We get that, in order to prove Claim 6.7, it is sufficient to show that, uniformly in $H \in \mathcal{H}^{(1)}$,

$$\sum_{\ell \ge \lceil n^{1/5} \rceil + 1} \Pi_{\ell}^{H} \left((1 + o(1)) \frac{N}{m'} \right)^{\ell} = o\left(\frac{1}{n}\right).$$

Fix non-negative integers c and x. Let us denote by $\mathcal{J}_{\ell,x,c}^H$ the set of all subgraphs $J \subset H$ with ℓ edges, x non-isolated vertices, and c connected components (excluding isolated vertices). Denote by $p(\ell, x, c)$ the probability that $\mathbf{H} \cap H \in \mathcal{J}_{\ell,x,c}^H$. Due to Claim 6.6,

$$p(\ell, x, c) \le \frac{|\mathcal{J}_{\ell, x, c}^{H}| \cdot 16^{\ell} (n - x + c)! n^{c/6}}{|\mathcal{H}^{(1)}|}.$$

Moreover, due to Claim 3.2, $|\mathcal{J}_{\ell,x,c}^H| \leq {\binom{2\ell_0+5\chi}{c}} 2^{6\ell}$. Therefore,

$$\Pi_{\ell}^{H} = \sum_{c=1}^{\ell} \sum_{\ell/2+3c/2 \le x \le \ell+c} p_{\ell,x,c} \le \sum_{c,x} \frac{\binom{2\ell_0+5\chi}{c} 2^{10\ell} (n-x+c)! n^{c/6}}{(1-o(1))n!}.$$

Now, notice that, for a graph $J \in \mathcal{J}_{\ell,x,c}^H$ consisting of c components that have sizes ℓ_1, \ldots, ℓ_c , we have $\ell_i = 2x_i - 3 - \sigma_i$, where

$$\sigma_i \ge \frac{4C\ell_i}{\ln n} \cdot I\left(\ell_i \ge \frac{\ln n}{100C}\right)$$

and C is a large positive constant. Indeed, since every closed subgraph has size at most 4μ , we get that every set of $x' \ge \frac{\ln n}{200C}$ vertices contributes at most 2x' - 4 edges. Therefore, for any admissible triple (ℓ, x, c) , we get $\ell = 2x - 3c - \sigma$, where

$$\sigma = \min_{J \in \mathcal{J}_{\ell,x,c}^H} \sum_{i=1}^c \sigma_i \ge \min_{J \in \mathcal{J}_{\ell,x,c}^H} \sum_{i: \ell_i \ge \frac{\ln n}{100C}} \frac{4C\ell_i}{\ln n} > \left(\ell - c \cdot \frac{\ln n}{100C}\right) \frac{4C}{\ln n} = \frac{4C}{\ln n} \cdot \ell - \frac{c}{25}.$$
 (37)

In particular, we get

$$\frac{(n-x+c)!}{n!} \le \frac{1}{(n-x+c)^{x-c}} = \frac{1}{n^{x-c}(1-\frac{x-c}{n})^{x-c}} = O\left(n^{-x+c}\right) = O\left(n^{-\ell/2-c/2-\sigma/2}\right).$$
(38)

This and (37) imply

$$\sum_{\ell \ge \lceil n^{1/5} \rceil + 1} \Pi_{\ell}^{H} \left((1 + o(1)) \frac{N}{m'} \right)^{\ell} \le \sum_{c,\ell,x} \frac{\binom{2^{\ell_0 + 5\chi}}{c} 2^{10\ell} n^{c/6}}{n^{\ell/2 + c/2 + \sigma/2}} \left((1 + o(1)) \frac{N}{m'} \right)^{\ell}$$
$$\le \sum_{c,\ell,x} \frac{(e(2\ell_0 + 5\chi)/c)^c \left((1/\varepsilon + o(1)) 2^{10} \right)^{\ell}}{n^{c/3 + \sigma/2}}$$
$$\le \sum_{c,\ell,x} (8Cn^{1/2 - 1/3 + 1/50}/c)^c \left((1/\varepsilon + o(1)) 2^{10} \right)^{\ell} n^{-2C\ell/\ln n}.$$

We can choose $C = C(\varepsilon)$ so large that

$$((1/\varepsilon + o(1))2^{10})^{\ell} n^{-2C\ell/\ln n} \le e^{-C\ell}.$$

Then,

$$\begin{split} \sum_{\ell \ge \lceil n^{1/5} \rceil + 1} \Pi_{\ell}^{H} \left((1 + o(1)) \frac{N}{m'} \right)^{\ell} &\leq \sum_{c,\ell,x} (8Cn^{0.19}/c)^{c} e^{-C\ell} \\ &\leq \sum_{\ell \ge n^{0.2}} \sum_{x} \left(\sum_{c \le n^{0.19}} (8Cn^{0.19})^{n^{0.19}} e^{-C\ell} + \sum_{c > n^{0.19}} (8C)^{c} e^{-C\ell} \right) \\ &\leq n^{3} e^{-\Omega(n^{0.2})} + \sum_{\ell \ge n^{0.2}} \sum_{x} \sum_{c > n^{0.19}} (8Ce^{-C})^{\ell} = e^{-\Omega(n^{1/5})}. \end{split}$$

This completes the proof.

Let $Y^{(1)}$ be the number of $H \in \mathcal{H}^{(1)}$ (with multiplicities) such that $(H, \mathbf{W}^{(1)})$ is not $\lceil n^{1/5} \rceil$ bad. Due to Markov's inequality, $Y^{(1)} = (1 - o(1))n!$ whp. For every such H we choose $H' \subset H \cup \mathbf{W}^{(1)}$ such that $H' \in \mathcal{H}^{(1)}$ and the fragment $H \cap H'$ has size at most $\lceil n^{1/5} \rceil$ and put this fragment into the new multiset $\mathcal{H}^{(2)}$. Without loss of generality, we may assume that each graph in this multiset has exactly $\lceil n^{1/5} \rceil$ edges.

6.4. Day 2: covering a fragment. It is well known that increasing properties that hold whp in the uniform model hold whp in the respective binomial model as well (see, e.g., [17, Corollary 1.16]). In particular, letting $p = (1 + 2\varepsilon)\sqrt{e/n}$, we get that, for $\mathbf{G} \sim G(n, p)$ and every $H \in \mathcal{H}^{(2)} = \mathcal{H}^{(2)}(\mathbf{G})$, whp $H \cup \mathbf{G}$ contains some $F \in \mathcal{F}_n$.

Next, we expose

 $\mathbf{G}' \sim G(n, p), \quad \text{where} \quad p' = \varepsilon \cdot n^{-1/2}.$

It remains to prove that whp there exists $H \in \mathcal{H}^{(2)}$ such that $H \subset \mathbf{G}'$. Let $Y^{(2)}$ be the number of such H. We get

$$\mathbb{E}Y^{(2)} = (p')^{\lceil n^{1/5} \rceil} |\mathcal{H}^{(2)}| = (1 - o(1))(p')^{\lceil n^{1/5} \rceil} n! = \omega(1)$$

On the other hand, recalling that $\mathcal{J}_{\ell,x,c}^H$ is defined in Section 6.3, due to Claim 6.6,

$$\operatorname{Var}Y^{(2)} \le \mathcal{H}^{(2)} \cdot \sum_{\ell, x, c} \max_{H \in \mathcal{H}^{(2)}} \max_{H' \subset H} |\mathcal{J}_{\ell, x, c}^{H'}| \cdot 16^{\ell} (n - x + c)! n^{c/6} (p')^{2\lceil n^{1/5} \rceil - \ell} - (\mathbb{E}Y^{(2)})^2,$$

implying

$$\frac{\operatorname{Var}Y^{(2)}}{(\mathbb{E}Y^{(2)})^2} \le \sum_{\ell,x,c} \max_{H \in \mathcal{H}^{(2)}} \max_{H' \subset H} \frac{|\mathcal{J}_{\ell,x,c}^{H'}| \cdot 16^{\ell} (n-x+c)! n^{c/6}}{(1-o(1))n! p'^{\ell}} - 1$$

Due to Claim 3.2, estimates (37) and (38), and recalling that $x \leq \ell + c$ and that $C = C(\varepsilon)$ is large, we get

$$\begin{split} \frac{\operatorname{Var} Y^{(2)}}{(\mathbb{E} Y^{(2)})^2} &\leq \sum_{c \geq 0, \ell, x} \frac{\binom{2\lceil n^{1/5} \rceil}{c} 2^{10\ell} n^{c/6}}{n^{\ell/2 + c/2 + \sigma/2} (\varepsilon \cdot n^{-1/2})^{\ell}} - 1 \\ &\leq \sum_{c \geq 1, \ell, x} \frac{(2e\lceil n^{-2/15} \rceil/c)^c \left(2^{10}/\varepsilon\right)^{\ell}}{n^{\sigma/2}} \\ &\leq \sum_{c \geq 1, \ell, x} n^{-c/15} \left(2^{10}/e^C\varepsilon\right)^{\ell} \\ &\leq \sum_{c \geq 1, \ell} (\ell + c) n^{-c/15} e^{-\ell} = O(n^{-1/15}), \end{split}$$

completing the proof due to Chebyshev's inequality.

6.5. Proof of Claim 6.3. Let $\ell \ge \ell_0 + 1$ be an integer.

We first note that the square of a path with ℓ edges has $\frac{\ell+3}{2}$ vertices. Therefore, a disjoint union of c squares on ℓ edges has $\frac{\ell+3c}{2}$ vertices. We then fix integers $c \in [\ell]$ and $x \in [\frac{\ell}{2} + \frac{3}{2}c, \ell + c]$. For a subgraph $H \subset F \in \mathcal{F}_n$ with x vertices, ℓ edges, and c components,

$$\sigma := 2x - \ell - 3c$$

is the difference between the number of edges of a disjoint union of squares of a path with x vertices and c components and the number of edges of H.

Let $p(\ell, x, c)$ be the probability that $(\mathbf{F} \cap F) \cup (D_1 \cup \ldots \cup D_{\chi})$ is a graph on $x + 4\chi$ vertices with $\ell + 5\chi$ edges, and $c + \chi$ connected components (as usual, we think about graphs as about sets of their edges, so there are no isolated vertices in $\mathbf{F} \cap F$). The following claim immediately implies that, for sufficiently large $A_1, A_2 > 0$,

$$p(\ell, x, c) \leq \frac{\binom{n}{c}\binom{x}{c+\chi}\left(\frac{2x}{\chi}\right)^{\chi}e^{A_1c+A_2\sigma}\left(\max_{o\leq 8\sigma}\binom{x}{o}\right)^2(n-x-4\chi+c)!}{|\mathcal{F}_n(\overrightarrow{D})|}.$$
(39)

Claim 6.8. There exist $A_{\alpha}^1, A_{\alpha}^2 > 0$ such that the number of subgraphs in F without isolated vertices with $x + 4\chi$ vertices, $\ell + 5\chi$ edges, and $c + \chi$ components is

$$\alpha(\ell, x, c) \le \binom{n}{c} \binom{x}{c+\chi} \left(\frac{2x}{\chi}\right)^{\chi} e^{A_{\alpha}^{1}c+A_{\alpha}^{2}\sigma} \max_{o\le 8\sigma} \binom{x}{o}.$$
(40)

There exist $A_{\beta}^{1}, A_{\beta}^{2} > 0$ such that, given $H \subset F$ with $x + 4\chi$ vertices, $\ell + 5\chi$ edges, and $c + \chi$ components, the number of ways to extend H to a graph from $\mathcal{F}_{n}(\overrightarrow{D})$ is at most

$$\beta(\ell, x, c) = (n - x - 4\chi + c)! e^{A_{\beta}^{1}c + A_{\beta}^{2}\sigma} \max_{o \le 8\sigma} \binom{x}{o}.$$
(41)

Proof. There are at most $\binom{x}{c+\chi}\binom{\sigma+c+\chi}{c+\chi}$ ways to choose positive integers $\ell_1, \ldots, \ell_{c+\chi}$ as well as $x_1, \ldots, x_{c+\chi}$ such that

• $\frac{\ell_i}{2} + \frac{3}{2} \le x_i \le \ell_i + 1$ for all $i \in [c]$ and $\frac{\ell_i}{2} \le x_i \le \ell_i$ for all $i \in \{c+1, \dots, c+\chi\}$,

•
$$\sum_{i=1}^{c+\chi} \ell_i = \ell$$
, and $\sum_{i=1}^{c+\chi} x_i = x$.

Fix such x_i and ℓ_i , $i \in [c + \chi]$. For $i \in [c + \chi]$, set $\sigma_i = 2x_i - \ell_i - 3$.

We first choose closed subgraphs of F that correspond to $\sigma_i = 0$ one by one. The number of ways to choose the first closed subgraph is at most 2n (the heaviest case is $x_i = 2$ that correspond to an edge). In the same way, if some set of \tilde{n} vertices is already included in the subgraph under construction, then the next closed component can be chosen in at most $2(n-\tilde{n})$ ways. On the other hand, if the *i*-th component contains one of the diamonds (i.e. $i \ge c+1$), then the number of ways to choose it is at most x_i .

We now switch to not closed components. Let us call a vertex *free* in a subgraph of F, if its degree is less than 4. Note that the *i*-th component has $o_i \leq 6 + 2\sigma_i$ free vertices. Assume that \tilde{n} vertices have been already included in the subgraph and we now describe the procedure of choosing the *i*-th component:

- (1) choose the number of free vertices $o_i \leq 6 + 2\sigma_i \leq 8\sigma_i \leq 8^{\sigma_i}$;
- (2) choose a set $\mathcal{O} \in {[x_i] \choose o_i}$ that identifies the labels of free vertices in the *i*-th component;
- (3) if $i \leq c$ choose a vertex w out of the set of remaining $n \tilde{n}$ vertices and, if $i \geq c+1$, choose w such that $0 \leq \pi_F(u_{i-c}) \pi_F(w) \leq x_i 1$, where u_{i-c} is the minimum vertex of the diamond D_{i-c} that this component has to contain; *activate* w we treat this vertex as the π_F -minimum vertex in the component under construction;
- (4) at every step $j \ge 1$, consider the π_F -minimum vertex v_j among active vertices:
 - if $j \in \mathcal{O}$ (i.e. v_j should be free), then add to the component some of the neighbours of v_j (in at most 2^4 ways), deactivate v_j , and *activate* all its chosen neighbours,
 - if $j \notin \mathcal{O}$, then add to the component all the neighbours of v_j , deactivate v_j , and *activate* all its neighbours.

We get that, if $i \leq c$, then the number of ways to choose the component is at most $(n - \tilde{n})8^{\sigma_i} \max_{o_i \leq 8\sigma_i} {x_i \choose o_i} 2^{4o_i}$, and at most $x_i 8^{\sigma_i} \max_{o_i \leq 8\sigma_i} {x_i \choose o_i} 2^{4o_i}$, otherwise. Eventually we get that the number of *ordered* choices of components with parameters ℓ_i, x_i ,

Eventually we get that the number of *ordered* choices of components with parameters ℓ_i, x_i , $i \in [c + \chi]$, in F is at most

$$c! \binom{n}{c} 2^{c} (x/\chi)^{\chi} \prod_{i=1}^{c+\chi} 8^{\sigma_{i}} \max_{o_{i} \leq 8\sigma_{i}} \binom{x_{i}}{o_{i}} 2^{4o_{i}} \leq c! \binom{n}{c} (x/\chi)^{\chi} 2^{c} 2^{35\sigma} \max_{o \leq 8\sigma} \binom{x}{o}.$$

Note that this bound does not depend on the order of the choice of components that do not contain diamonds, thus

$$\alpha(\ell, x, c) \leq \binom{n}{c} \binom{x}{c+\chi} \binom{\sigma+c+\chi}{c+\chi} \left(\frac{x}{\chi}\right)^{\chi} 2^{c+35\sigma} \max_{o\leq 8\sigma} \binom{x}{o}$$
$$\leq \binom{n}{c} \binom{x}{c+\chi} \left(\frac{2x}{\chi}\right)^{\chi} 2^{2c+36\sigma} \max_{o\leq 8\sigma} \binom{x}{o}$$

as needed.

Let us now fix $H \subset F$ with $x + 4\chi$ vertices, $\ell + 5\chi$ edges, and $c + \chi$ components. Let us bound the number of ways to extend H to an $F' \in \mathcal{F}_n(\overrightarrow{D})$. We construct such an extension in the following way.

We forget the labels of the vertices from $V(H) \setminus V(D_1 \cup \ldots \cup D_{\chi})$ and assume that the desired $F' \in \mathcal{F}_n$ is the second power of the cycle $(12 \ldots n)$. Let \mathcal{Z} be the union of the set of the c connected components of H that do not contain diamonds with the set of remaining $n - x - 4\chi$ isolated vertices. We should compute the number of ways to embed the elements of Z in F' disjointly.

Let $z_1, \ldots, z_{n-x-4\chi+c}$ be an ordering of Z (there are $(n-x-4\chi+c)!$ ways to order the elements of Z). We embed sequentially each z_i in F' in a way such that all vertices of z_i are bigger than the minimum vertices of z_1, \ldots, z_{i-1} . At every step $i = 1, \ldots, n-x-4\chi+c$,

consider the minimum vertex κ_i of F' such that none of the embedded elements of Z in F'contain this vertex. If z_i is a single vertex, then we assign κ_i with z_i and proceed with the next step. Otherwise, we distinguish between the following cases.

First, we assume that z_i is closed. If $|V(z_i)| = 2$, then there are 2 ways to choose this edge (two edges to smaller vertices are prohibited) and at most two ways to place it (two rotations). Thus there are 4 ways to embed z_i . If $|V(z_i)| = 3$, then there are 6 ways to choose a (labelled) copy of z_i in F' with the minimum vertex κ_i . If $|V(z_i)| = 4$, then the number of ways equals 4. If $|V(z_i)| > 4$, then there are 2 ways to choose a copy of z_i in F' with the minimum vertex κ_i .

Second, let z_i be not closed with o free vertices. Choose a set \mathcal{O} from $\binom{[|V(z_i)|]}{o}$ that will identify the labels of free vertices in the embedding of z_i in F'. Activate κ_i . At every step $j \ge 1$, choose the $\pi_{F'}$ -minimum active vertex v_i in F' and

- if $j \in \mathcal{O}$, then add to the image of z_i under construction some neighbours of v_i (in at most 2^3 ways), deactivate v_i , and *activate* all its neighbours,
- if $j \notin \mathcal{O}$, then add all the neighbours of v_i , deactivate the vertex, and *activate* all its neighbours.

The image is constructed. However, we have not yet mapped the vertices of z_i to the vertices of the image. Clearly, z_i is union of a 'path' P of inclusion-maximum 2-connected graphs joined by (usual) paths with 'fringe' trees, growing from P. There are two ways to orient P, unless Pis a cycle. Every rooted 'fringe' tree T has at most $2^{|V(T)|-1}$ automorphism since T is a subtree of a perfect 3-ary tree. Moreover, the number of free non-root vertices in T is at least |V(T)|/2. In total, there are at most $2 \cdot 2^{2o-1} = 2^{2o}$ automorphisms of z_i and this bound holds true even when P is a cycle. We get that the number of ways to construct the image of z_i is at most $\binom{|V(z_i)|}{2} 2^{5o}$.

It is easy to see that, for a component of H that contains some D_i and has o free vertices, there are at most 6° ways to embed it in F'. We conclude that there are at most

$$(n - x - 4\chi + c)!6^{c} \prod_{i=1}^{c} {x_{i} \choose o_{i}} 2^{5o_{i}} \prod_{i=c+1}^{c+\chi} 6^{o_{i}} \le (n - x - 4\chi + c)!6^{c} 2^{40\sigma} \max_{o \le 8\sigma} {x \choose o}$$
expose F' as needed.

ways to expose F' as needed.

Recalling that $|\mathcal{F}_n(\vec{D})| = (n-4\chi)!$ and $x-c = \ell/2 + c/2 + \sigma/2$, we get

$$\frac{(n-x-4\chi+c)!}{|\mathcal{F}_n(\overrightarrow{D})|} = \frac{(n-\frac{\ell+c+\sigma}{2}-4\chi)!}{(n-4\chi)!} \le \sqrt{2\pi n} \cdot \frac{e^{\frac{\ell+c+\sigma}{2}}}{(n-4\chi)^{\frac{\ell+c+\sigma}{2}}} \left(\frac{n-\frac{\ell+c+\sigma}{2}-4\chi}{n-4\chi}\right)^{n-4\chi-\frac{\ell+c+\sigma}{2}} \le \sqrt{2\pi n} \cdot \frac{e^{(\ell+c+\sigma)^2/(4(n-4\chi))}}{(n-4\chi)^{(\ell+c+\sigma)/2}} = e^{o(\ell)} \cdot \frac{e^{(x-c)^2/n}}{n^{(\ell+c+\sigma)/2}}.$$

We also notice that $\ln(x/\chi) = o(x/\chi)$ since $x/\chi = \Omega(\ln n/w) = \omega(1)$. Therefore,

$$\left(\frac{2x}{\chi}\right)^{\chi} = e^{\chi \ln(2x/\chi)} = e^{\chi \ln(x/\chi) + \chi \ln 2} = e^{o(x)} = e^{o(\ell)}.$$

We first choose $0 < \varepsilon' \ll \delta$ small enough. From (39) and the bound $\binom{n}{c} \leq \left(\frac{en}{c}\right)^c$, we get

$$p(\ell, x, c) \le e^{o(\ell)} \cdot \frac{\left(\frac{e^{A_1 + 1}\sqrt{n}}{c}\right)^c {\binom{x}{c+\chi}} e^{A_2\sigma} {\binom{x}{o}}^2 e^{(x-c)^2/n}}{n^{\ell/2 + \sigma/2}},$$
(42)

where $o = o(x) \leq 8\sigma$ is chosen in such a way that $\binom{x}{o}$ achieves its maximum. Since $\left(\frac{e^{A_1+1}\sqrt{n}}{c}\right)^c \leq e^{A_1\sqrt{n}}$, we get that

$$\left(\frac{e^{A_1+1}\sqrt{n}}{c}\right)^c \le e^{A_1\sqrt{n}} \le e^{(A_1/C)\ell}$$

Let us recall that $\ell > \ell_0$, and thus $x > \ell_0/2$. Here $\ell_0 = \lfloor C\sqrt{n} \rfloor$, and $C \gg \max\{\frac{1}{\epsilon'}, A_1/\delta\}$ can be chosen arbitrarily large.

We further consider separately several different cases.

1. If $\sigma > \varepsilon' x$, then

$$\binom{x}{c+\chi}\binom{x}{o}^2 e^{A_2\sigma} e^{(x-c)^2/n} < 8^x e^x e^{A_2\sigma} < n^{\sigma/2}.$$

Therefore,

$$p(\ell, x, c) \le \left(e^{1+\delta}/n\right)^{\ell/2}.$$
(43)

2. Let $\sigma \leq \varepsilon' x$.

2.1. If $c < \varepsilon' x$ and $x < \varepsilon' n$, then

$$\binom{x}{c+\chi}\binom{x}{o}^2 e^{A_2\sigma} e^{(x-c)^2/n} < e^{(\delta/2)\ell}$$

implying (43) as well.

2.2. If $c < \varepsilon' x$ and $x \ge \varepsilon' n$, then

$$\left(\frac{e^{A_1+1}\sqrt{n}}{c}\right)^c e^{A_2\sigma} \binom{x}{c+\chi} \binom{x}{o}^2 < e^{(\delta/2)\ell}.$$

Thus, (42) implies

$$p(\ell, x, c) \le \frac{e^{(x-c)^2/n}}{n^{\ell/2}} e^{(\delta/2 + o(1))\ell} \le (e^{1+\delta}/n)^{\ell/2}$$

since $\ell/2 = x - 3c/2 - \sigma/2 \ge x(1 - 5\varepsilon'/2).$

2.3. Finally, let $c \ge \varepsilon' x$. Let us choose C so large that $\left(\frac{e^{A_1+1}\sqrt{n}}{c}\right)^c \le e^{-10x}$ (taking any $C > 10e^{A_1+1}e^{1/\varepsilon'}/\varepsilon'$ is enough for that). Since $e^{A_2\sigma} \le n^{\sigma/2}$, we get

$$p(\ell, x, c) \le e^{-10x + o(\ell)} \frac{\binom{x}{c+\chi} \binom{x}{o}^2 e^{(x-c)^2/n}}{n^{\ell/2}} \le e^{-10x + x + o(\ell)} 8^x n^{-\ell/2} \le n^{-\ell/2}.$$

Summing up,

$$\sum_{\ell \ge \ell_0 + 1} \Pi_\ell \left((1 + o(1)) \frac{N}{m} \right)^\ell \le \sum_{\ell \ge \ell_0 + 1} \sum_{x,c} p(\ell, x, c) \left(\frac{1 + o(1)}{1 + \varepsilon} \cdot \sqrt{\frac{n}{e}} \right)^\ell \le n^2 \sum_{\ell \ge \ell_0 + 1} \left(\frac{e^{\delta/2} + o(1)}{1 + \varepsilon} \right)^\ell = o\left(\frac{1}{n}\right)$$

completing the proof of Claim 6.3.

7. Generalisation: proof of Theorem 1.8

Let \mathcal{F}_n be the family of all isomorphic copies of F on [n] rooted in the first r vertices. Let λ be the number of edges in $F_0 := F[[r]]$. We have that $|\mathcal{F}_n| = n!$ and each graph from \mathcal{F}_n induces a linear order on [n].

We then proceed in the same way as in the proof of Theorem 1.7. Here, $w = o(\ln n/r)$ and $\chi = \left\lfloor \frac{wn^{(2d-\Delta)/d}}{\ln n} \right\rfloor$. We replace diamonds with (2r)-subgraphs isomorphic to two consecutive isomorphic copies of F_0 , that is, $F_0^* \cong F[[2r]]$. Note that F_0^* has $\lambda^* \ge 2\lambda - \Delta/2$ edges. In the same way as in Section 6, we define the family $\mathcal{F}_n(\overrightarrow{D}) \subset \mathcal{F}_n$ for a *d*-tuple \overrightarrow{D} of disjoint copies of F_0^* . We then fix C > 0 large enough and sample an *m*-subset $\mathbf{W} \subset {[n] \choose 2} \setminus (D_1 \cup \ldots \cup D_d)$ uniformly at random, where $m = \lfloor (1+\varepsilon)(e/n)^{2/d} \cdot N \rfloor$. We also let $\mathcal{F}_n^*(\overrightarrow{D})$ be the set of all $F \setminus (D_1 \cup \ldots \cup D_d)$, $F \in \mathcal{F}_n(\overrightarrow{D})$. In the same way, for $F \in \mathcal{F}_n^*(\overrightarrow{D})$ and an *m*-set $W \subset {[n] \choose 2} \setminus (D_1 \cup \ldots \cup D_d)$, we say that the pair (F, W) is ℓ -bad, if at least $n^{-1/2}$ -fraction of $F' \subset F \cup W$ that belong to $\mathcal{F}_n^*(\overrightarrow{D})$ have $|F \cap F'| > \ell$. We have that Claim 6.4 holds for these \overrightarrow{D} , $\mathcal{F}_n^*(\overrightarrow{D})$, m, and $\ell_0 = \lfloor Cn^{(2d-\Delta)/d} \rfloor$, as well. Indeed, the proof of Claim 6.2 remains unchanged. Claim 6.4 follows from 1) the fact that, for every $t \in \{0, 1, \ldots, dn/2 - \lambda^*\chi\}$,

$$\frac{\mathbb{E}X'}{M(t)} \le \sum_{\ell \ge \ell_0 + 1} \prod_{\ell} e^{-\frac{(dn/2)^2}{m}(1 - o(1))} \left((1 + o(1)) \frac{N}{m} \right)^{\ell},$$

where

- X' is the number of $F' \subset F \cup \mathbf{W}', F' \in \mathcal{F}_n^*(\overrightarrow{D})$, such that $|F' \cap F| \ge \ell_0 + 1$, and \mathbf{W}' is a uniformly random (m-t)-subset of $\binom{[n]}{2} \setminus (F \cup D_1 \cup \ldots \cup D_{\chi})$;
- M(t) is the expected number of $F' \in \mathcal{F}_n^*(\overrightarrow{D})$ such that F' belongs to a uniformly random subset of $\binom{[n]}{2} \setminus (D_1 \cup \ldots \cup D_{\chi})$ of size $m + (dn/2 \lambda^* \chi) t$.

and 2) the following analogue of Claim 6.3:

Claim 7.1. Sample a uniformly random $\mathbf{F} \in \mathcal{F}_n^*(\overrightarrow{D})$. For $\ell \in \mathbb{N}$, let $\Pi_\ell := \mathbb{P}(F \cap \mathbf{F} \text{ has } \ell \text{ edges})$. Then

$$\sum_{\ell \ge \ell_0 + 1} \prod_{\ell} e^{-\frac{(dn/2)^2}{m}(1 - o(1))} \left((1 + o(1)) \frac{N}{m} \right)^{\ell} = o\left(\frac{1}{n}\right).$$

The proof of the latter claim follows the same reasoning as the proof of Claim 6.3; therefore, we omit it for brevity and to avoid repetitions. We only note that the bound (39) becomes

$$p(\ell, x, c) \leq \frac{\binom{n}{c}\binom{x}{c+\chi} \left(\frac{dx/2}{\chi}\right)^{\chi} e^{A_1 c + A_2 \sigma} \left(\max_{o \leq (d+4)\sigma} \binom{x}{o}\right)^2 (n - x - 2r\chi + c)!}{|\mathcal{F}_n(\overrightarrow{D})|}$$

In particular, the bound on the number of extensions of a subgraph $H \subset F$ carries over to this setting, as every subgraph $H \subset F$ that contains D_1 has no non-trivial automorphisms that fix the vertices of D_1 as well as the boundary vertices of H. Since $|\mathcal{F}_n(\vec{D})| = (n - 2r\chi)!$ and $x - c = \frac{2\ell}{d} + \frac{c(\Delta - d)}{d} + \frac{2\sigma}{d}$, the rest of the proof of Claim 6.3 remains unchanged.

The main novel ingredient in the proof of Theorem 1.7 — the improvement of fragments, described in Section 6.2 — applies here as well. The main difference is that diamonds are replaced here with isomorphic copies of F_0^* . Let $\mu := \left\lfloor \frac{\ln n}{r\sqrt{w}} \right\rfloor$. We also consider all inclusion-maximal closed subgraphs of a typical (F, \mathbf{W}) -fragment H of size at most $\ell_0 + \lambda^* \chi$ with at least $\ln n/\sqrt{w}$ vertices that do not overlap with any D_i , for $F \in \mathcal{F}_n(\overrightarrow{D})$. Here we cannot cut a closed subgraph in an arbitrary place since we have to keep pieces of size r entirely. For each considered closed subgraph $P_i = (v_1^i v_2^i \dots)$, we find the minimum r' such that $H[\{v_{r'+1}^i \dots v_{r'+r}^i\}] \cong F_0$, the bijection preserving the order of vertices is an isomorphism, and $v_{r'+1}^i, \dots, v_{r'+r}^i \notin \partial_v(H)$. Then we remove from H all the consecutive pieces $(v_{r'+1}^i \dots v_{r'+r\mu}^i), (v_{r'+r\mu+1}^i \dots v_{r'+2r\mu}^i), \dots$, of size $r\mu$. We also have to make sure that all vertices of the last removed piece are not boundary.

Finally, we glue the two remaining pieces of every H, and insert each removed piece between the two isomorphic copies of F_0 of the respective $D_i \cong F_0^*$.

As in the proof of Theorem 1.7, we match every piece P_i with D_{j_i} randomly: Consider a binomial random bipartite graph **B** with parts $V = [\chi]$ and U = [n] and edge probability $\beta := n^{(\Delta-d)/(3d)-(2d-\Delta)/d}/w$; insert every P_i into some D_{j_i} so that there exists an edge in **B** between $j_i \in V$ and the position of the first vertex of P_i in F. This is typically possible due to Claim 6.5, that holds in this setting as well. For a bipartite graph B with typical degrees, we consider the set $\Sigma = \Sigma(B)$ of all pairs $(F \in \mathcal{F}_n, W \in \binom{\binom{[n]}{2}}{m})$ such that there exists $F' \in \mathcal{F}_n(\overrightarrow{D}(F))$ satisfying

- (1) $F' \subset F \cup W$,
- (2) $|F' \cap F| \le \ell_0 + \lambda^* \chi$,
- (3) B has a matching that covers all vertices in V that represent positions of first vertices of the removed pieces of the fragment $H = F \cap F'$.

Due to symmetry and linearity of expectation, we get that there exists B such that (asymptotically) almost all pairs (F, W) belong to Σ . In order to complete analysis of smoothed fragments, we state the following analogue of Claim 6.6.

Let $W \in \binom{\binom{[n]}{2}}{m}$. Let Σ_W be the set of all $F \in \mathcal{F}_n$ such that $(F, W) \in \Sigma$. For every $F \in \Sigma_W$, consider an $F' := F'(F) \in \mathcal{F}_n$ satisfying (1)–(3). Let $H(F) := F \cap F'$ and H'(F) be the smoothed version of H(F).

Claim 7.2. For any graph S, let $\mathcal{S}(S)$ be the set of all graphs $F \in \Sigma_W$ such that H'(F) contains S as a subgraph. Then

$$|\mathcal{S}(S)| \le r^{c(S)} (4d)^{\ell(S)} (n - x(S) + c(S))! n^{c(S)(\Delta - d)/(3d)}.$$

We omit the proof since it is almost identical with the proof of Claim 6.6. Actually, the only difference is that when we perform the step (1) and reconstruct the order of vertices in every connected component of S, we choose a vertex of F_0 that corresponds to the root of the connected component (r choices for every component).

Here, since ℓ_0 is much bigger than \sqrt{n} unless d = 4, one additional fragmentation step is not enough. We perform $\frac{2(2d-\Delta)}{\Delta-d} - 1$ steps, where at each step we reduce the size of a fragment by the factor of $\Omega(n^{\frac{\Delta-d}{2d}})$. Assume that, for $i \in \left[\frac{2(2d-\Delta)}{\Delta-d} - 1\right]$, we have a multiset $\mathcal{H}^{(i)}$ of size $(1 - o(1))|\mathcal{F}_n|$ such that almost every $F \in \mathcal{F}_n$ has a fragment H satisfying the following:

- a subgraph $J^{(i)}$ of its smoothed version H' of size $\left\lfloor n^{\frac{(2d-\Delta)}{d}-(i-1)\frac{\Delta-d}{2d}} \right\rfloor$ belongs to $\mathcal{H}^{(i)}$ (if i = 1, then the size should be $\ell_0 + \lambda^* \chi$);
- $J^{(i)} \cup \mathbf{W} \cup \mathbf{W}^{(1)} \cup \ldots \cup \mathbf{W}^{(i-1)}$ contains an isomorphic copy of F, where random graphs $\mathbf{W}^{(1)}, \ldots, \mathbf{W}^{(i-1)}$ were sprinkled at the previous steps.

Let **H** be a uniformly random element of $\mathcal{H}^{(i)}$ and let $\mathbf{W}^{(i)} \sim G(n, m')$, $m' = \lfloor \varepsilon n^{-2/d} \cdot N \rfloor$, be sampled independently of $\mathbf{W}, \mathbf{W}^{(1)}, \ldots, \mathbf{W}^{(i-1)}$.

Claim 7.3. Whp there exists $J \subset \mathbf{H} \cup \mathbf{W}^{(i)}$, where $J \in \mathcal{H}^{(i)}$, such that $|\mathbf{H} \cap J| \le n^{\frac{(2d-\Delta)}{d} - i\frac{\Delta-d}{2d}}$.

In order to prove this claim, we apply Lemma 4.1 with $f = f(i) = \left\lfloor n^{\frac{(2d-\Delta)}{d} - (i-1)\frac{\Delta-d}{2d}} \right\rfloor$ and \mathcal{B} being the set of all graphs on n vertices with more than $n^{\frac{(2d-\Delta)}{d} - i\frac{\Delta-d}{2d}}$ edges. It is then sufficient

to show that, uniformly in $J \in \mathcal{H}^{(i)}$,

$$\sum_{\ell \ge f(i+1)+1} \Pi_{\ell}^{J} \left((1+o(1)) \frac{N}{m'} \right)^{\ell} = o\left(\frac{1}{n}\right),$$

where $\Pi_{\ell}^{J} = \mathbb{P}(J \cap \mathbf{H} \text{ has } \ell \text{ edges})$. The rest of the proof is similar to the proof of Claim 6.7 and follows from Claim 7.2. Therefore, we only outline the differences. We first notice that the bound (38) here becomes

$$\frac{(n-x+c)!}{n!} \le \frac{1}{n^{x-c}(1-\frac{x-c}{n})^{x-c}} = n^{-x+c} \cdot e^{(1-o(1))\frac{(x-c)^2}{n}}.$$

Therefore, the upper bound on Π_{ℓ}^{J} contains an extra factor $r^{c}e^{(1-o(1))\frac{(x-c)^{2}}{n}} \leq r^{c}e^{\ell}$. We choose C large enough and use the bound (37) that holds here as well. Since $r^{c} = n^{o(c)}$, we finally get the required bound

$$\sum_{\ell \ge f(i+1)+1} \Pi_{\ell}^{J} \left((1+o(1)) \frac{N}{m'} \right)^{\ell} \le \sum_{c,\ell,x} \frac{\binom{f(i)}{c} n^{((\Delta-d)/(3d)+o(1))c}}{n^{(\Delta-d)c/d-2c/(25d)}} e^{-C\ell} = e^{-\Omega(f(i+1))}.$$

It remains to cover at least one fragment of size $\lfloor n^{\frac{\Delta-d}{2d}} \rfloor$ from the multiset $\mathcal{H}^{(\frac{2(2d-\Delta)}{\Delta-d})}$ by the last sample $\mathbf{G}' \sim G(n,p), p = \varepsilon \cdot n^{-2/d}$, whp. The proof follows from Claim 7.2, Chebyshev's inequality, and is verbatim as the argument in Section 6.4. Therefore, we omit it.

8. Remaining challenges

Although the expectation threshold conjecture of Kahn and Kalai has been resolved [28], the asymptotics of $p_c(\mathcal{Q})$ remain unknown — even up to a constant factor — for an arbitrary increasing property \mathcal{Q} . In particular, it would be very interesting to determine this asymptotics when \mathcal{Q}_F is generated by all isomorphic copies of any fixed *d*-regular graph F on [n]. Theorem 1.2 gives only a partial answer and the problem becomes significantly more challenging when considering graphs F that contain many subgraphs with smaller edge boundaries.

We suspect that the requirements in Theorem 1.4 are far from optimal. In particular, Theorem 1.8 asserts that (2) is true for a wide class of d-regular graphs F that is not covered by Theorem 1.4. We actually believe that the following is true.

Conjecture 8.1. Let $d \ge 3$ and let F = F(n) be a sequence of d-regular graphs on [n], $n \in \mathbb{N}$, such that $|\partial_e(\tilde{F})| \ge d+1$, for every $\tilde{F} \subset F$ with $3 \le |V(\tilde{F})| \le n-3$. Then $p_c(F) = (1+o(1))p_e(F)$.

A notable special case of graphs F covered by this conjecture is a hexagonal lattice. Although the hexagonal lattice is not a regular graph, it can be extended to a 3-regular graph by adding missing edges between the boundary vertices of degree 2. Since the boundary has $O(\sqrt{n})$ vertices, it does not contribute to the threshold asymptotically. This specific F was mentioned by Riordan in [31], where he considered it as a special case to which his methods do not apply. It is worth noting that Fernandez de la Vega and Manoussakis [10] proved that $p_c(F) =$ $O((\log n/n)^{2/3})$ and that Theorem 1.2, as well as [7, Corollary 1.5], implies (1) for such F.

It is also interesting to establish stronger versions of Theorems 1.4, 1.7, and 1.8. Although it is plausible to refine the asymptotics of $p_c(F)$ to some extent using similar methods, achieving results as precise as those for Hamilton cycles seems very challenging. Let us recall that, for an *n*-cycle F, $p_c(F) = \frac{\ln n + \ln \ln n + O(1)}{n}$ [20] and that the following hitting time result is known [1, 4]: in the random graph process $\mathbf{W} = \mathbf{W}_m$, m = 0, 1, ..., N, the hitting time for \mathcal{Q}_F coincides with the hitting time for $\delta(\mathbf{W}) \geq 2$ whp. It would be very interesting to prove hitting time versions of Theorems 1.4, 1.7, and 1.8. It is clear that for denser regular graphs F hitting times for such local properties, that involve only neighbourhoods of bounded radii, cannot coincide with the hitting times for \mathcal{Q}_F . Nevertheless, one may ask the following question: is there k = o(n) such that the hitting time for containing the second power of a Hamilton cycle coincides with the hitting time for the property that every vertex belongs to the second power of a path of length k whp?

Acknowledgements

This work originated during the author's visit to Tel Aviv University in the summer of 2022. The author is grateful to Wojciech Samotij for his warm hospitality and for helpful discussions throughout the visit. The author would like to thank Asaf Cohen Antonir, Sahar Diskin, Ilay Hoshen, and Michael Krivelevich for valuable feedback on the presentation of the paper. The author would also like to thank József Balogh and Robert A. Krueger for helpful discussions at the Banff International Research Station during the workshop "Bootstrap Percolation and its Applications", as well as the organisers of this event.

An early version of this paper, uploaded to arXiv in January 2023, contained a crucial error that invalidated the proof of Theorem 1.7. The author sincerely thanks Ashwin Sah and Mehtaab Sawhney for identifying the mistake and for subsequent discussions and valuable comments.

References

- M. Ajtai, J. Komlós, E. Szemerédi, The first occurrence of Hamilton cycles in random graphs, Annals of Discrete Mathematics 27 (1985) 173–178.
- [2] R. Alweiss, S. Lovett, K. Wu, J. Zhang, Improved bounds for the sunflower lemma, Ann. of Math. (2), 194:3 (2021) 795–815.
- [3] J. Balogh, A. Bernshteyn, M. Delcourt, A. Ferber, H. T. Pham, Sunflowers in set systems with small VCdimension, Prepring (2024) arXiv:2408.04165.
- [4] B. Bollobás, The evolution of sparse graphs, Graph theory and combinatorics (Cam- bridge, 1983), Academic Press, London, (1984) 35–57.
- [5] B. Bollobás, The isoperimetric number of random regular graphs, European Journal of Combinatorics 9 (1988) 241–244.
- [6] B. Bollobás, A. Thomason, Threshold functions, Combinatorica 7 (1987) 35–38.
- [7] Y. Chen, J. Han, H. Luo, On the thresholds of degenerate hypergraphs, Preprint (2024) arXiv:2411.18596.
- [8] P. Erdős, A. Rényi, On the evolution of random graphs, Publ. Math. Inst. Hung. Acad. Sci 5:1 (1960) 17–60.
- [9] A. Espuny Díaz, Y. Person, Spanning F-cycles in random graphs, Combin. Probab. Comput. 32:5 (2023) 833–850.
- [10] W. Fernandez de la Vega, Y. Manoussakis, Grids in random graphs, Random Structures & Algorithms 5:2 (1994) 329–336.
- [11] M. Fischer, N. Škorić, A. Steger, M. Trujić, Triangle resilience of the square of a Hamilton cycle in random graphs, J. Comb. Theory, Ser. B 152 (2018) 171–220.
- [12] K. Frankston, J. Kahn, B. Narayanan, J. Park, Thresholds versus fractional expectation-thresholds, Ann. of Math. 194:2 (2021) 475–495.
- [13] E. Friedgut, Hunting for sharp thresholds, Random Structures & Algorithms, 26:1–2 (2005) 37–51.
- [14] E. Friedgut, Sharp thresholds of graph properties, and the k-sat problem, Journal of the American Mathematical Society, 12:4 (1999) 1017–1054, With an appendix by Jean Bourgain.
- [15] A. Frieze, A note on spanning K_r -cycles in random graphs, AIMS Mathematics 5:5 (2020) 4849–4852.

- [16] A. Frieze, Hamilton cycles in random graphs: a bibliography, Preprint, arXiv:1901.07139.
- [17] S. Janson, T. Łuczak, A. Ruciński, Random graphs, J. Wiley & Sons Inc., 2000.
- [18] J. Kahn, B. Narayanan, J. Park, The threshold for the square of a Hamilton cycle, Proc. Amer. Math. Soc. 149 (2021) 3201–3208.
- [19] J. H. Kim, B. Sudakov, V. Vu, Small subgraphs of random regular graphs, Discrete Mathematics 307 (2007) 1961–1967.
- [20] J. Komlós, E. Szemerédi, Limit distributions for the existence of Hamilton circuits in a random graph, Discrete Mathematics 43 (1983) 55–63.
- [21] A. D. Korshunov, Solution of a problem of Erdős and Rényi on hamiltonian cycles in nonoriented graphs, Soviet Math. Doklady 17 (1976) 760–764.
- [22] I. Krasikov, A. Lev, B. D. Thatte, Upper bounds on the automorphism group of a graph, Discrete Mathematics 256 (2002) 489–493.
- [23] D. Kühn, D. Osthus, On Pósa's conjecture for random graphs, SIAM J. Discrete math. 26:3 (2012) 1440– 1457.
- [24] B. D. McKay, N. C. Wormald, Automorphisms of random graphs with specified degrees, Combinatorica 4 (1984) 325–338.
- [25] E. Mossel, J. Niles-Weed, N. Sun, I. Zadik, A second moment proof of the spread lemma, Preprint (2022) arXiv:2209.11347.
- [26] B. Narayanan, M. Schacht, Sharp thresholds for nonlinear Hamiltonian cycles in hypergraphs, Random Structures & Algorithms 57:1 (2020) 244–255.
- [27] R. Nenadov, N. Škorić, Powers of Hamilton cycles in random graphs and tight Hamilton cycles in random hypergraphs, Random Structures & Algorithms 54:5 (2019) 187–208.
- [28] J. Park, H. Pham, A Proof of the Kahn-Kalai conjecture, Journal of the American Mathematical Society 37:1 (2024) 235–243.
- [29] W. Perkins, Searching for (sharp) thresholds in random structures: where are we now? Preprint (2024) arXiv:2401.01800.
- [30] L. Pósa, Hamiltonian circuits in random graphs, Discrete Math. 14 (1976) 359–364.
- [31] O. Riordan, Spanning subgraphs of random graphs, Combinatorics, Probability & Computing 9:2 (2000) 125–148.
- [32] J. Spencer, Threshold functions for extension statements, Journal of Combinatorial Theory Ser. A 53 (1990) 286–305.
- [33] S. Spiro, A smoother notion of spread hypergraphs, Combin. Probab. Comput. 32:5 (2023) 809-818.
- [34] M. Talagrand, Are many small sets explicitly small?, Proceedings of the 2010 ACM International Symposium on Theory of Computing (STOC '2010), 13–35.
- [35] T. Tao, The sunflower lemma via Shannon entropy, Online post, 2020.
- [36] R. Venkatesan, L. Levin, Random instances of a graph coloring problem are hard, In Proc. 20th Annual Symposium on the Theory of Computing (1988) 217–222.
- [37] N. C. Wormald, The asymptotic distribution of short cycles in random regular graphs, Journal of Combinatorial Theory Ser. B 31:2 (1981) 168–182.

APPENDIX A. COARSE BOUND: PROOF OF THEOREM 1.2

Let F be as in the statement of Theorem 1.2. Let B > 0 be a large enough constant. Consider d independent samples

$$\mathbf{W}_1, \mathbf{W}_2, \dots, \mathbf{W}_d \sim G(n, m'), \text{ where } m' = |B \cdot n^{-2/d} \cdot N|.$$

Let \mathcal{F}_n be the family of all isomorphic copies of F on [n].

Claim A.1. Whp there exists $F' \subset F \cup \mathbf{W}_1$, $F' \in \mathcal{F}_n$, such that $|F \cap F'| \leq n^{1-1/d}$.

Proof. Let $f = \frac{dn}{2}$, m = m', and \mathcal{B} be the property of graphs on n vertices to have more than $n^{1-1/d}$ edges. Due to Lemma 4.1 and due to symmetry, it is sufficient to prove (13).

Fix non-negative integers c and x. Let us denote by $\mathcal{J}_{\ell,x,c}$ the set of all subgraphs $J \subset F$ with ℓ edges, x non-isolated vertices, and c connected components (excluding isolated vertices). Denote by $p(\ell, x, c)$ the probability that $\mathbf{F} \cap F \in \mathcal{J}_{\ell,x,c}$. Due to Claim 3.2 and Claim 3.3, for some constant A > 0,

$$p(\ell, x, c) \le \frac{\binom{n}{c} e^{A\ell} (n - x + c)! / |\operatorname{Aut}(F)|}{|\mathcal{F}_n|}.$$

Let $\ell > n^{1-1/d}$. Recalling that $x = \frac{2}{d}\ell + \frac{c\Delta}{d} + \frac{2}{d}\sigma$, assuming $n - x + c \ge 1$, we get

$$\frac{(n-x+c)!/|\operatorname{Aut}(F)|}{|\mathcal{F}_n|} = \frac{(n-\frac{2\ell+c(\Delta-d)+2\sigma}{d})!}{n!}$$

$$\leq (1+o(1))\frac{e^{\frac{2\ell+c(\Delta-d)+2\sigma}{d}}}{n^{\frac{2\ell+c(\Delta-d)+2\sigma}{d}}} \left(\frac{n-\frac{2\ell+c(\Delta-d)+2\sigma}{d}}{n}\right)^{n-\frac{2\ell+c(\Delta-d)+2\sigma}{d}}$$

$$\leq (1+o(1))\frac{e^{\frac{2\ell+c(\Delta-d)+2\sigma}{d^2n}}}{n^{\frac{2\ell+c(\Delta-d)+2\sigma}{d}}} \leq (1+o(1)) \cdot \frac{e^{\frac{(x-c)^2}{n}}}{n^{\frac{2\ell+c+2\sigma}{d}}}.$$
(44)

Clearly, the same bound holds when n - x + c = 0 as well. Therefore, since $x - c \leq \min\{\ell, n\}$,

$$p(\ell, x, c) \le \frac{\left(\frac{en^{1-1/d}}{c}\right)^c e^{(A+2)\ell}}{n^{\frac{2\ell+2\sigma}{d}}} \le \frac{e^{n^{1-1/d}}e^{(A+2)\ell}}{n^{\frac{2\ell+2\sigma}{d}}}$$

The requirement on the edge boundary from the statement of Theorem 1.2 implies $\sigma \geq -\frac{2d\ell}{\varepsilon \ln n}$. Indeed, if a component of $J \in \mathcal{J}_{\ell,x,c}$ has size $x_i \leq \varepsilon \ln n$, then the respective $\sigma_i := \frac{d}{2}x_i - \ell_i - \frac{\Delta}{2}$ is non-negative. If $x_i > \varepsilon \ln n$, then its number of edges ℓ_i satisfies $2\ell_i \leq dx_i$. Thus, recalling that $x_i \leq 2\ell_i$, we get

$$\sigma_i = \frac{d}{2}x_i - \ell_i - \frac{\Delta}{2} \ge -\frac{\Delta}{2} \ge -\frac{\Delta x_i}{2\varepsilon \ln n} \ge -\frac{dx_i}{\varepsilon \ln n} \ge -\frac{2d\ell_i}{\varepsilon \ln n}.$$

So,

$$\sum_{\ell} \Pi_{\mathcal{B}_{\ell}}^{F} \left(\left(1 + \frac{3f}{m} \right) \frac{N}{m} \right)^{\ell} e^{-f^{2}/m + f^{3}/(3m^{2})} \leq \sum_{\ell > n^{1-1/d}} \sum_{x,c} p(\ell, x, c) \left(\frac{1 + o(1)}{B} \cdot n^{2/d} \right)^{\ell}$$
$$\leq n^{2} \sum_{\ell > n^{1-1/d}} e^{n^{1-1/d}} \left(\frac{e^{A + 2 + 4/\varepsilon} + o(1)}{B} \right)^{\ell}$$
$$= o\left(\frac{1}{n^{2}} \right),$$

completing the proof of the claim.

For every $F \in \mathcal{F}_n$ such that there exists F' as in the statement of Claim A.1, we choose one such F' and put $F \cap F'$ into a multiset $\mathcal{F}_n^{(1)}$. Due to Claim A.1 and Markov's inequality, whp $|\mathcal{F}_n^{(1)}| = (1 - o(1))|\mathcal{F}_n|$. We then proceed by induction. Assume that, for $i \in [d - 1]$, whp we have a multiset $\mathcal{F}_n^{(i)}$ of graphs with exactly $\lfloor n^{1-i/d} \rfloor$ edges that is obtained by adding a single $H \subset F$ from almost every $F \in \mathcal{F}_n$ such that the graph $H \cup \mathbf{W}_1 \cup \ldots \cup \mathbf{W}_i$ contains some $F' \in \mathcal{F}_n$. We have that $|\mathcal{F}_n^{(i)}| = (1 - o(1))|\mathcal{F}_n|$ whp. Let **H** be a uniformly random element of $\mathcal{F}_n^{(i)}$.

Claim A.2. Whp there exists $H' \subset \mathbf{H} \cup \mathbf{W}_{i+1}$ such that $H' \in \mathcal{F}_n^{(i)}$ and $|\mathbf{H} \cap H'| \leq \max\left\{n^{1-(i+1)/d}, \ln n\right\}.$

Proof. Let $f = \lfloor n^{1-i/d} \rfloor$, m = m', and \mathcal{B} be the property of graphs on n vertices to have more than $n^{1-(i+1)/d}$ edges. Due to Lemma 4.1, it is sufficient to prove (9).

Fix non-negative integers c and x. For $H \in \mathcal{F}_n^{(i)}$, let us recall that $\mathcal{J}_{\ell,x,c}^H$ is the set of all subgraphs $J \subset H$ with ℓ edges, x non-isolated vertices, and c connected components (excluding isolated vertices). Denote by $p^H(\ell, x, c)$ the probability that $\mathbf{H} \cap H \in \mathcal{J}_{\ell,x,c}^H$. Due to Claim 3.2, Claim 3.3, and estimate (44), for some constant A > 0,

$$p^{H}(\ell, x, c) \leq \frac{\left(\frac{\lfloor n^{1-i/d} \rfloor}{c}\right)e^{A\ell}(n-x+c)!/|\operatorname{Aut}(F)|}{|\mathcal{F}_{n}^{(i)}|} \\ \leq (1-o(1))\frac{\left(\frac{en^{1-(i+1)/d}}{c}\right)^{c}e^{(A+2)\ell}}{n^{2(\ell+\sigma)/d}} \leq (1-o(1))\frac{e^{n^{1-(i+1)/d}}e^{(A+2)\ell}}{n^{2(\ell+\sigma)/d}}$$

Recalling that $\sigma \geq -\frac{C\ell}{2\ln n}$, for every $H \in \mathcal{F}_n^{(i)}$,

$$\begin{split} \sum_{\ell} \Pi_{\mathcal{B}_{\ell}}^{H} \left(\left(1 + \frac{3f}{m} \right) \frac{N}{m} \right)^{\ell} e^{-f^{2}/m + f^{3}/(3m^{2})} \\ &\leq \sum_{\ell > \max\{n^{1-(i+1)/d}, \ln n\}} \sum_{x,c} p^{H}(\ell, x, c) \left(\frac{1 + o(1)}{B} \cdot n^{2/d} \right)^{\ell} \\ &\leq n^{2} \sum_{\ell > \max\{n^{1-(i+1)/d}, \ln n\}} e^{n^{1-(i+1)/d}} \left(\frac{e^{A+2+4/\varepsilon} + o(1)}{B} \right)^{\ell} = o\left(\frac{1}{n^{2}}\right), \end{split}$$

as desired.

By induction, whp we get a multiset $\mathcal{F}_n^{(d)} = \mathcal{F}_n^{(d)}(\mathbf{W}_1 \cup \ldots \cup \mathbf{W}_d)$ of graphs of size $\lfloor \ln n \rfloor$ comprising a single $H \subset F$ from almost every $F \in \mathcal{F}_n$ such that the graph $H \cup \mathbf{W}_1 \cup \ldots \cup \mathbf{W}_d$ contains some $F' \in \mathcal{F}_n$. It is well known that increasing properties that hold whp in the uniform model hold whp in the respective binomial model as well (see, e.g., [17, Corollary 1.16]). In particular, let

$$\mathbf{G} \sim G(n, p), \quad \text{where} \quad p = (1 + \varepsilon) dB n^{-2/d}.$$

Whp there exists a multiset $\mathcal{F}_n^{(d)} = \mathcal{F}_n^{(d)}(\mathbf{G})$ of graphs of size $\lfloor \ln n \rfloor$ comprising a single subgraph H of almost every $F \in \mathcal{F}_n$ so that $H \cup \mathbf{G}$ contains some $F' \in \mathcal{F}_n$.

Let X be the number of $H \in \mathcal{F}_n^{(d)}$ that belong to $\mathbf{G}' \sim G(n, p' = Bn^{-2/d})$, sampled independently of **G**. We get

$$\mathbb{E}X = |\mathcal{F}_n^{(d)}| p^{\prime \lfloor \ln n \rfloor} = (1 - o(1)) |\mathcal{F}_n| p^{\prime \lfloor \ln n \rfloor} = \omega(1).$$

Let \mathcal{B} be the set of non-empty graphs. Due to the definition (8) of $\Pi^{H}_{\mathcal{B}_{\ell}} = \Pi^{H}_{\mathcal{B}_{\ell}}(\mathcal{F}_{n}^{(d)}),$

$$\frac{\operatorname{Var} X}{(\mathbb{E} X)^2} \le \frac{\max_{H \in \mathcal{F}_n^{(d)}} \sum_{\ell \ge 1} \Pi_{\mathcal{B}_\ell}^H |\mathcal{F}_n^{(d)}| p'^{\lfloor \ln n \rfloor - \ell}}{\mathbb{E} X} = \max_{H \in \mathcal{F}_n^{(d)}} \sum_{\ell > 1} \Pi_{\mathcal{B}_\ell}^H p'^{-\ell}.$$

Therefore, due to Claim 3.2, Claim 3.3, and estimate (44), for some constant A > 0,

$$\frac{\operatorname{Var} X}{(\mathbb{E}X)^2} \leq \sum_{\ell \geq 1} \sum_{x,c} \frac{\binom{\lfloor \ln n \rfloor}{c} e^{(A+2)\ell}}{n^{2\ell/d+c/d+2\sigma/d}} \left(\frac{1}{B} \cdot n^{2/d}\right)^{\ell} \\
\leq \sum_{\ell \geq 1} \sum_{c \geq 1} O(\ell) \left(\frac{e \ln n}{cn^{1/d}}\right)^c \left(\frac{e^{A+2+4/\varepsilon}}{B}\right)^{\ell} = O\left(\frac{\ln n}{n^{1/d}}\right),$$

completing the proof of Theorem 1.2, due to Chebyshev's inequality.

Appendix B. Proofs of Claims 6.2, 6.4

Claim B.1. Let $F \in \mathcal{F}_n^*(\overrightarrow{D})$ and let **W** be a uniformly random *m*-element subset of $\binom{[n]}{2} \setminus (D_1 \cup \ldots \cup D_d)$. Let $\delta(n) \to 0$ as $n \to \infty$. Then, for every $t \in \{0, 1, \ldots, 2n - 5d\}$,

$$\mathbb{P}\left(|\mathcal{M}(F,\mathbf{W})| < \delta(n)M \mid |F \cap \mathbf{W}| = t\right) \le \delta(n)$$

Proof. Sample a uniformly random $\mathbf{F} \in \mathcal{F}_n^*(\overrightarrow{D})$ independently of \mathbf{W} . Then

$$\begin{split} \mathbb{P}\left(|\mathcal{M}(\mathbf{F},\mathbf{W})| < \delta(n)M \mid |\mathbf{F} \cap \mathbf{W}| = t\right) = \\ & \sum_{F \in \mathcal{F}_n^*(\overrightarrow{D})} \mathbb{P}\left(\mathbf{F} = F, |\mathcal{M}(F,\mathbf{W})| < \delta(n)M \mid |\mathbf{F} \cap \mathbf{W}| = t\right) \\ = & \sum_{F \in \mathcal{F}_n^*(\overrightarrow{D})} \mathbb{P}\left(|\mathcal{M}(F,\mathbf{W})| < \delta(n)M \mid \mathbf{F} = F, |\mathbf{F} \cap \mathbf{W}| = t\right) \\ & \times \mathbb{P}(\mathbf{F} = F \mid |\mathbf{F} \cap \mathbf{W}| = t) \\ = & \sum_{F \in \mathcal{F}_n^*(\overrightarrow{D})} \mathbb{P}\left(|\mathcal{M}(F,\mathbf{W})| < \delta(n)M \mid |F \cap \mathbf{W}| = t\right) \cdot \mathbb{P}(\mathbf{F} = F \mid |\mathbf{F} \cap \mathbf{W}| = t). \end{split}$$

By symmetry, $\mathbb{P}(|F \cap \mathbf{W}| = t)$ and $\mathbb{P}(|\mathcal{M}(F, \mathbf{W})| < \delta(n)M, |F \cap \mathbf{W}| = t)$ do not depend on $F \in \mathcal{F}_n^*(\overrightarrow{D})$. Therefore,

$$\begin{split} \mathbb{P}(\mathbf{F} = F \mid |\mathbf{F} \cap \mathbf{W}| = t) &= \frac{\mathbb{P}(\mathbf{F} = F, |F \cap \mathbf{W}| = t)}{\mathbb{P}(|\mathbf{F} \cap \mathbf{W}| = t)} \\ &= \frac{\mathbb{P}(\mathbf{F} = F)\mathbb{P}(|F \cap \mathbf{W}| = t)}{\sum_{\tilde{F} \in \mathcal{F}_n^*(\vec{D})} \mathbb{P}(\mathbf{F} = \tilde{F})\mathbb{P}(|\tilde{F} \cap \mathbf{W}| = t)} \\ &= \frac{\mathbb{P}(|F \cap \mathbf{W}| = t)}{\sum_{\tilde{F} \in \mathcal{F}_n^*(\vec{D})} \mathbb{P}(|\tilde{F} \cap \mathbf{W}| = t)} = \frac{1}{|\mathcal{F}_n^*(\vec{D})|} \end{split}$$

and

$$\mathbb{P}\left(|\mathcal{M}(F,\mathbf{W})| < \delta(n)M \mid |F \cap \mathbf{W}| = t\right) = \mathbb{P}\left(|\mathcal{M}(F',\mathbf{W})| < \delta(n)M \mid |F' \cap \mathbf{W}| = t\right)$$

for all $F' \in \mathcal{F}_n^*(\overrightarrow{D})$. We also notice that each pair $\{F \in \mathcal{F}_n^*(\overrightarrow{D}), W \in \binom{E(K_n) \setminus (D_1 \cup \ldots \cup D_d)}{m}\}$ such that $|\mathcal{M}(F, W)| < \delta(n)M$ and $|F \cap W| = t$ can be obtained by

- first, choosing a set of edges $A \subset [n]$ of size m + (2n 5d) t, on the role of $F \cup W$, that contains less than $\delta(n)M$ graphs $F \in \mathcal{F}_n^*(\overrightarrow{D})$ — in at most $\binom{N-5d}{m+(2n-5d)-t}$ ways,
- then, choosing an $F \in \mathcal{F}_n^*(\overrightarrow{D})$ such that F is a subgraph of A in less than $\delta(n)M$ ways, and,
- finally, choosing the intersection $F \cap W$ in $\binom{2n-5d}{t}$ ways.

We conclude that

$$\mathbb{P}\left(|\mathcal{M}(F,\mathbf{W})| < \delta(n)M \mid |F \cap \mathbf{W}| = t\right) = \mathbb{P}\left(|\mathcal{M}(\mathbf{F},\mathbf{W})| < \delta(n)M \mid |\mathbf{F} \cap \mathbf{W}| = t\right)$$
$$\leq \frac{\delta(n)M\binom{N-5d}{m+(2n-5d)-t}\binom{2n-5d}{t}}{|\mathcal{F}_n^*(\overrightarrow{D})|\binom{N-2n}{m-t}\binom{2n-5d}{t}} = \delta(n),$$

completing the proof.

Claim B.2. Let $F \in \mathcal{F}_n^*(\overrightarrow{D})$ and let $\mathbf{W} \subset {\binom{[n]}{2}} \setminus (D_1 \cup \ldots \cup D_d)$ be a uniformly random *m*-subset. Then whp (F, \mathbf{W}) is not ℓ_0 -bad.

Proof. Let $\delta(n)$ be a slowly decreasing function and let $t \in \{0, \ldots, 2n - 5d\}$. Due to Claim 6.2, we have

$$\mathbb{P}((F, \mathbf{W}) \text{ is } \ell_0 \text{-bad}, |\mathcal{M}(F, \mathbf{W})| < \delta(n)M(t) | |F \cap \mathbf{W}| = t) = o(1).$$

Thus, it is sufficient to prove that, uniformly over t,

$$\mathbb{P}((F, \mathbf{W}) \text{ is } \ell_0\text{-bad}, |\mathcal{M}(F, \mathbf{W})| \ge \delta(n)M(t) \mid |F \cap \mathbf{W}| = t) = o(1).$$

The latter probability equals

$$\mathbb{P}\left(|\mathcal{M}_{\ell_0}(F, \mathbf{W})| > \frac{1}{\sqrt{n}} |\mathcal{M}(F, \mathbf{W})|, |\mathcal{M}(F, \mathbf{W})| \ge \delta(n)M(t) | |F \cap \mathbf{W}| = t\right) \\
\le \mathbb{P}\left(|\mathcal{M}_{\ell_0}(F, \mathbf{W})| > \frac{\delta(n)}{\sqrt{n}}M(t) | |F \cap \mathbf{W}| = t\right) \le \frac{\sqrt{n} \cdot \mathbb{E}(X | |F \cap \mathbf{W}| = t)}{\delta(n)M(t)},$$

where X counts the number of $F' \in \mathcal{M}(F, \mathbf{W})$ such that $|F' \cap F| \geq \ell_0 + 1$. Let \mathbf{W}' be a uniformly random (m-t)-subset of $\binom{[n]}{2} \setminus (F \cup D_1 \cup \ldots \cup D_d)$, and let X' be the number of $F' \in \mathcal{M}(F, \mathbf{W}')$ such that $|F' \cap F| \geq \ell_0 + 1$. Clearly,

$$\mathbb{E}(X \mid |F \cap \mathbf{W}| = t) = \mathbb{E}X'.$$

Then

$$\mathbb{E}X' = \sum_{\ell \ge \ell_0 + 1} |\mathcal{F}_n^*(\overrightarrow{D})| \Pi_\ell \binom{m - t}{(2n - 5d) - \ell} / \binom{N - 2n}{(2n - 5d) - \ell}.$$
(45)

We have

$$\frac{\mathbb{E}X'}{M} = \sum_{\ell \ge \ell_0 + 1} |\mathcal{F}_n^*(\vec{D})| \cdot \frac{\Pi_\ell}{M} \cdot \frac{\binom{m-t}{(2n-5d)-\ell}}{\binom{N-2n}{(2n-5d)-\ell}} = \sum_{\ell \ge \ell_0 + 1} \Pi_\ell \cdot \frac{\binom{m-t}{(2n-5d)-\ell} / \binom{N-2n}{(2n-5d)-\ell}}{\binom{N-2n}{m-t} / \binom{N-5d}{(m-t+(2n-5d))}}.$$
 (46)

By Stirling's approximation,

$$\frac{\binom{m-t}{(2n-5d)-\ell}/\binom{N-2n}{(2n-5d)-\ell}}{\binom{N-2n}{m-t}\binom{N-5d}{m-t+(2n-5d)}} = \frac{\binom{m-t}{(2n-5d)-\ell}\binom{N-5d}{m-t+(2n-5d)}}{\binom{N-2n}{(2n-5d)-\ell}} \\ \sim \frac{(m-t)^{2(m-t)}(N-4n+5d+\ell)^{N-4n+5d+\ell}(N-5d)^{N-5d}}{(m-t-2n+5d+\ell)^{m-t-2n+5d+\ell}(m-t+2n-5d)^{m-t+2n-5d}(N-2n)^{2N-4n}}.$$

Therefore, for $\ell \geq \ell_0 + 1$

$$\begin{aligned} \frac{\binom{m-t}{(2n-5d)-\ell}/\binom{N-2n}{(2n-5d)-\ell}}{\binom{N-2n}{m-t+(2n-5d)}} &\sim \left(\frac{N-4n+5d+\ell}{m-t-2n+5d+\ell}\right)^{\ell} \\ &\times \frac{\left(1+\frac{2n-5d-\ell}{m-t-2n+5d+\ell}\right)^{m-t-2n+5d} \left(1-\frac{2n-5d}{m-t+2n-5d}\right)^{m-t+2n-5d}}{\left(1-\frac{2n-5d}{N-5d}\right)^{N-5d} \left(1+\frac{2n-5d-\ell}{N-4n+5d+\ell}\right)^{N-4n+5d}} \\ &< \left(\left(1+o(1)\right)\frac{N}{m}\right)^{\ell} e^{2n-5d-\ell-\frac{\ell(2n-5d-\ell)}{m}-\frac{(2n-5d-\ell)^2}{2m}-(2n-5d)-\frac{(2n-5d)^2}{2m}+2n-5d-(2n-5d-\ell)} \\ &= e^{-\frac{4n^2}{m}} \left(\left(1+o(1)\right)\frac{N}{m}\right)^{\ell}. \end{aligned}$$

Claim 6.3 completes the proof.

SCHOOL OF COMPUTER SCIENCE, UNIVERSITY OF SHEFFIELD, UK Email address: m.zhukovskii@sheffield.ac.uk