ON UNIVERSAL-HOMOGENEOUS HYPERBOLIC GRAPHS AND SPACES AND THEIR ISOMETRY GROUPS

KATRIN TENT

ABSTRACT. The Urysohn space is the unique separable metric space that is universal and homogeneous for finite metric spaces, i.e., it embeds any finite metric space any isometry between finite subspaces extends to an isometry of the whole space. We here consider the existence of a universal-homogeneous hyperbolic space. We show that that for $\delta > 0$ there is no δ -hyperbolic space which is universal and homogeneous in the above sense for all finite δ -hyperbolic spaces.

We then show that for any $\delta \geq 0$ and any countable class C of δ -hyperbolic spaces with countably many distinguished δ -closed subspaces there exists a δ hyperbolic metric space $\mathbb{H}_{\mathcal{C}} = \mathbb{H}(\mathcal{C}, \delta)$ such that every $X \in \mathcal{C}$ can be embedded into $\mathbb{H}_{\mathcal{C}}$ as a δ -closed subspace and any isometry between distinguished closed subspaces extends to an isometry of $\mathbb{H}_{\mathcal{C}}$.

If C consists of δ -hyperbolic geodesic spaces, then $\mathbb{H}_{\mathcal{C}}$ contains the *quasi-tree* of spaces as defined in [1].

For C_{δ} the class of all finite δ -hyperbolic spaces with rational distances or the class of finite δ -hyperbolic graphs, the limit \mathbb{H}_{δ} is a δ -hyperbolic space (or graph, respectively) universal for all finite δ -hyperbolic spaces with rational distances (or finite δ -hyperbolic graphs) and such that any isometry between δ -closed subspaces extends to an isometry of \mathbb{H}_{δ} .

We show that the isometry group of \mathbb{H}_{δ} does not contain elements of bounded displacement and has no dense conjugacy class.

1. INTRODUCTION AND NON-EXISTENCE

The Urysohn space is the unique separable metric space that is universal and homogeneous for finite metric spaces, i.e., it embeds any finite metric space and any isometry between finite subspaces extends to an isometry of the whole space. The Urysohn space can easily be constructed by amalgamating all finite metric spaces with rational distances and taking the completion of the resulting metric space, see e.g. [8]. As is often the case with very homogeneous structures, the isometry group of the Urysohn space has a natural maximal normal subgroup, namely the group of isometries of bounded displacement and the quotient group is boundedly simple. In this note we consider the question whether there is an analog for hyperbolic metric spaces. This note is motivated by the non-existence result in Theorem 1.3.

We recall some terminology: Let \mathcal{C} be a countable class of structures. Then we say that \mathcal{C} has the

(1) (AP) Amalgamation Property if for any $A, B_1, B_2 \in \mathcal{C}$ and embeddings $i_j : A \longrightarrow B_j, j = 1, 2$, there is some $D \in \mathcal{C}$ and embeddings $k_j : B_j \longrightarrow D$ such that the diagram commutes, i.e. $i_1 \circ k_1 = i_2 \circ k_2$.

Note that if the empty structure is in \mathcal{C} then any class \mathcal{C} with (AP) also satisfies the Joint Embedding Property (JEP), i.e. any for any $B_1, B_2 \in \mathcal{C}$ there is some $D \in \mathcal{C}$ such that B_2, B_2 embed into D.

If C is a countable class of structures with (AP) and (JEP), there is a unique limit structure \mathbb{M} , called the *Fraissé limit* of C (see e.g. [9], Theorem 4.4.4). This limit \mathbb{M} is characterized by the following two properties:

- (1) (universal) \mathbb{M} is universal for \mathcal{C} , i.e. every $A \in \mathcal{C}$ embeds into \mathbb{M} , and
- (2) (homogeneous) \mathbb{B} is homogeneous for \mathcal{C} , i.e. any isomorphism between substructures $A, B \subset \mathbb{M}$ with $A, B \in \mathcal{C}$ extends to an isomorphism of \mathbb{M} .

We record the following well-known observation:

Lemma 1.1. Let \mathbb{M} be a structure which is universal and homogeneous for C. Then C has (AP).

Proof. Let $A, B_1, B_2 \in \mathcal{C}$ and consider embeddings $i_j : A \longrightarrow B_j, j = 1, 2$. Then by universality, B_1 and B_2 embed into \mathbb{M} and the composition $i_2 \circ i_1^{-1}$ to A induces an isomorphism between the substructures of B_1 and B_2 isomorphic to A. By homogeneity, there is an isomorphism α of \mathbb{M} taking the copy of A inside B_1 to the copy of A inside B_2 . Then any substructure of \mathbb{M} containing $\alpha(B_1) \cup B_2$ is an amalgam as required. \Box

We will use the following definition of δ -hyperbolicity:

Definition 1.2. Let (X, d) be a metric space, $\delta \ge 0$. Then X is δ -hyperbolic if and only if for all $x_1, x_2, x_3, x_4 \in X$ the following holds: Put

$$E = d(x_1, x_2) + d(x_3, x_4),$$

$$F = d(x_1, x_3) + d(x_2, x_4),$$

$$G = d(x_1, x_4) + d(x_2, x_3).$$

If $E \leq F \leq G$, then $G - F < 2\delta$.

It is not too hard to see that for $\delta = 0$, the class of 0-hyperbolic (or ultrametric) spaces has (AP), and hence universal-homogeneous ultrametric spaces exist, see e.g. [3, 4, 6, 7, 11]. However, for $\delta > 0$ there is no universal-homogeneous δ -hyperbolic space:

Theorem 1.3. For $\delta > 0$ there does not exist a δ -hyperbolic space such that every finite δ -hyperbolic space with rational distances can be isometrically embedded and such that any isometry between finite subspaces extends to an isometry of the whole space.

Proof. By Lemma 1.1 it suffices to show that finite δ -hyperbolic spaces with rational distances do not have (AP).

Let $A = \{c, d, e\}$ and $B_1 = A \cup \{a\}, B_2 = A \cup \{b\}, x_0, x_1, z_0, z_1, z \in \mathbb{R}_{>0}$ and suppose

$$\begin{split} d(c,d) &= x_0 + x_1 \\ d(e,c) &= z + x_0 \\ d(e,d) &= z + x_1 \\ d(a,e) &= z_0, \\ d(a,d) &= z_0 + z + x_1 \\ d(a,c) &= z_0 + z + x_0 - 2\delta \\ d(b,e) &= z_1, \\ d(b,d) &= z_1 + z + x_1 - 2\delta \\ d(b,c) &= z_1 + z + x_0 \end{split}$$

Then $\{a, c, d, e\}$ and $\{b, c, d, e\}$ are δ -hyperbolic. Clearly $d(a, b) \leq z_0 + z_1$. If $z > 2\delta$ and $z_0 \neq z_1$, then

$$d(a,d) + d(b,c) = z_0 + z + x_1 + z_1 + z + x_0 > d(a,b) + d(c,d) + 2\delta$$

and similarly

$$d(a,d) + d(b,c) = z_0 + z + x_1 + z_1 + z + x_0$$

> $d(a,c) + d(b,d) + 2\delta$
= $z_0 + z + x_0 - 2\delta + z_1 + z + x_1 - 2\delta + 2\delta$
= $z_0 + z + x_0 + z_1 + z + x_1 - 2\delta$

So there is no $r \in \mathbb{R}$ such that defining d(a, b) = r would make a, b, c, d, e hyperbolic.

2. Hrushovski limits of δ -hyperbolic spaces

The previous result shows that we cannot construct a universal-homogeneous hyperbolic space. In order to obtain a universal δ -hyperbolic metric space with a weaker homogeneity condition, we modify the amalgamation by restricting to strong embeddings to obtain the Hrushovski limit, a weaker analog of the Fraïssé limit, .

Hrushovski limits: Let (\mathcal{C}, \leq) be a countable class of structures with a strong embedding relation \leq , defining a partial order on \mathcal{C} . We say that (AP) holds for (\mathcal{C}, \leq) if for all $A, B_1, B_2 \in \mathcal{C}$ with strong embeddings $i_j : A \longrightarrow B_i, i = 1, 2$, there exists some $D \in \mathcal{C}$ and strong embeddings $k_j : B_j \longrightarrow D$ such that the diagram commutes. If the empty structure is strongly embedded into every $A \in \mathcal{C}$, then also (JEP) holds for strong embeddings. In this case there is a unique limit structure \mathbb{M} , the Hrushovski limit for (\mathcal{C}, \leq) . For $A \in \mathcal{C}$ we define $A \leq \mathbb{M}$ if $A \leq C$ for any $C \in \mathcal{C}$ contained in \mathbb{M} . The Hrushovski limit is universal for \mathcal{C} , i.e. every $A \in \mathcal{C}$ strongly embeds into \mathbb{M} and \mathbb{M} is \leq -homogeneous, i.e. any isomorphism between strongly embedded substructures extends to an isomorphism of \mathbb{M} .

We first define a property of metric spaces inspired by buildings:

Definition 2.1. Let (X, d) be a metric space. We call a subset $A \subseteq X$ gated in X if for every $b \in X \setminus A$ there exists a gate $g = g_A(b) \in A$ such that for all $a \in A$ we have d(b, a) = d(b, g) + d(g, a). We call g the gate for b in A.

Remark 2.2. Note that if A is gated in X, then for $b \in X \setminus A$ the gate $g_A(b)$ is uniquely determined: if g and g' are gates for b in A, then d(b,g) = d(b,g') + d(g',g)and d(b,g') = d(b,g) + d(g,g') forcing d(g,g') = 0. In particular, $g_A(b)$ is the unique element in A with minimal distance to b.

Example 2.3. Clearly, any convex set in a tree is gated. On the other hand, suppose that $X = (x_0, x_1, \ldots, x_n = x_0)$ is a cycle of length n in a graph with the graph metric. Then the subset $X = \{x_1, \ldots, x_{n-1}\}$ is not gated in X as $d(x_0, x_1) = d(x_0, x_{n-1}) = 1$, contradicting our previous remark.

For a metric space (X, d) we call a tuple (x_0, \ldots, x_n) geodetic if $d(x_0, x_n) = d(x_0, x_1) + \ldots + d(x_{n-1}, x_n)$. The following is an easy, but useful consequence of the definition:

Remark 2.4. If A is gated in B, then A is convex in B in the following sense: if (a_1, b, a_2) is geodetic with $a_1, a_2 \in A$, then $b \in A$.

Definition 2.5. Fix $\delta \geq 0$ and let X be a metric space, $A \subset X$. Then we say that A is δ -closed (or: strongly embedded) in X and write $A \leq_{\delta} B$ (or just $A \leq B$ if δ is clear from the context) if

- (1) A is gated in B;
- (2) for $b, b' \in B \setminus A$ with $d(g_A(b), g_A(b')) > \delta$ we have

 $d(b,b') = d(b,g_A(b)) + d(g_A(b),g_A(b')) + d(g_A(b'),b').$

Clearly, any gated set is closed in the sense of the topology induced from the metric. Note however that the empty set is not gated and hence never δ -closed.

Remark 2.6. It is easy to see that being δ -closed is transitive, i.e., if $A \leq_{\delta} B$ and $B \leq_{\delta} C$, then $A \leq_{\delta} C$. We also note that single points are δ -closed in any metric space.

Lemma 2.7. If $A, B, C \in C$ such that $A, B \leq_{\delta} C$ and $A \cap B \neq \emptyset$, then $A \cap B \leq_{\delta} B \leq_{\delta} C$.

Proof. Since being δ -closed is transitive by the previous remark, it suffices to show that $A \cap B$ is δ -closed in A and in B. By symmetry we just show $A \cap B \leq_{\delta} B$. Let $b \in B \setminus A$. We need to show that $g_A(b) \in A \cap B$. Let $c \in A \cap B$. Then $(b, g_A(b), c)$ is geodetic with $b, c \in B$. By Remark 2.4 we have $g_A(b) \in B \cap A$.

Corollary 2.8. Any subset A of a metric space X is contained in a unique smallest δ -closed set $cl(A) = \bigcap \{B \leq_{\delta} X : A \subset B\}$.

By Remark 2.4 any δ -closed set is convex. However, the converse does not hold: Suppose $x, y \in X$ and there is some $z \in X$ such that $\{x, y, z\}$ is a non-degenerate triangle. If there is no z' such that both (z, z', x) and (z, z', y) are geodetic, then $z \in cl(\{x, y\})$.

Note also that the union of δ -closed subsets A, B need not be δ -closed. However, we have the following partial result which will be used in Section 5.

ON UNIVERSAL-HOMOGENEOUS HYPERBOLIC GRAPHS AND SPACES AND THEIR ISOMETRY GROUPS

Lemma 2.9. Suppose A, B are δ -closed subsets of a metric space $X, A \cap B \neq \emptyset$ and $B_{\delta}(A \cap B) \cap (B \cup A) = B \cap A$ where $B_{\delta}(A \cap B)$ denotes the ball of radius δ around $A \cap B$. Then $A \cup B$ is δ -closed in X.

Proof. Let $x \in X \setminus A \cup B$ and let $g_1 = g_A(x), g_2 = g_B(x)$. By symmetry we may assume $d(x, g_1) \leq d(x, g_2)$. We claim that (x, g_1, g_2) is geodesic. By Lemma 2.7 we have $A \cap B \leq \mathbb{H}_d$. Let $c = g_{A \cap B}(x) \in A \cap B$. Then (x, g_1, c) and (x, g_2, c) are geodetic. Furthermore, $c = g_A(g_2) = g_B(g_1)$ and so also (g_1, c, g_2) is geodetic.

Suppose that $d(c, g_2) \ge d(c, g_1) > \delta$. Since $c, g_2 \in B$ and $B \le_{\delta} X$, it follows from the definition of strong embeddings that $d(g_1, x) = d(g_1, c) + d(c, g_2) + d(g_2, x)$, a contradiction and so $d(c, g_1) \le \delta$. By assumption on $A \cap B$ we have $g_1 \in A \cap B \le B$. Thus $g_1 = c$ and since $d(x, g_1) \le d(x, g_2)$ it follows that $g_1 = g_2$ proving the claim.

Definition 2.10. Let A, B_1, B_2 be metric spaces with $A \subseteq B_1, B_2$. Then the canonical amalgam $D = B_1 \otimes_A B_2$ of B_1 and B_2 over A is defined as the disjoint union of A with $B_1 \setminus A$ and $B_2 \setminus A$ with the metric extending the metric on B_1, B_2 defined as follows: for $x \in B_1 \setminus A, y \in B_2 \setminus A$ we put

$$d(x, y) = d(x, g_A(x)) + d(g_A(x), g_A(y)) + d(g_A(y), y)$$

The crucial lemma is the following:

Lemma 2.11. Let A, B_1, B_2 be δ -hyperbolic spaces and assume that $A \leq_{\delta} B_1, B_2$. Then the canonical amalgam $D = B_1 \otimes_A B_2$ of B_1 and B_2 over A is δ -hyperbolic and $B_1, B_2 \leq_{\delta} D$.

Proof. It is clear from the definition of the canonical amalgam that $B_1, B_2 \leq_{\delta} D$. Hence it suffices to show that D is δ -hyperbolic. Let $x_1, \ldots, x_4 \in D$. If x_1, \ldots, x_4 are contained in B_1 or B_2 , there is nothing to show. We now write $g_i = g_A(x_i)$ if $x_i \notin A$ and $g_i = x_i$ if $x_i \in A$

Case 1: Assume $x_1, x_2, x_3 \in B_1, x_4 \in B_2 \setminus A$. Consider

$$E = d(x_1, x_2) + d(x_3, x_4) = d(x_1, x_2) + d(x_3, g_3) + d(g_3, g_4) + d(g_4, x_4),$$

$$F = d(x_1, x_3) + d(x_2, x_4) = d(x_1, x_3) + d(x_2, g_2) + d(g_2, g_4) + d(g_4, x_4)$$

$$G = d(x_1, x_4) + d(x_2, x_3) = d(x_1, g_1) + d(g_1, g_4) + d(g_4, x_4) + d(x_2, x_3)$$

Each of E, F, G contains the summand $d(g_4, x_4)$. Subtracting this summand from each of E, F, G, we get the distances corresponding to the points $x_1, x_2, x_3, g_4 \in B_1$. Since B_1 is hyperbolic, these points satisfy the condition in Definition 1.2, and hence so do x_1, x_2, x_3, x_4 .

Case 2: Assume $x_1, x_2 \in B_1, x_3, x_4 \in B_2 \setminus A$ and let

$$E_1 = d(g_1, g_2) + d(g_3, g_4)$$

$$F_1 = d(g_1, g_3) + d(g_2, g_4)$$

$$G_1 = d(g_1, g_4) + d(g_2, g_3)$$

Then

$$E = d(x_1, x_2) + d(x_3, x_4) \le E_1 + \sum_{i=1}^4 d(x_i, g_i)$$

$$F = d(x_1, x_3) + d(x_2, x_4) = F_1 + \sum_{i=1}^4 d(x_i, g_i)$$

$$G = d(x_1, x_4) + d(x_2, x_3) = F_2 + \sum_{i=1}^4 d(x_i, g_i)$$

If $d(g_3, g_4), d(g_1, g_2) > \delta$, then $E = E_1 + \sum_{i=1}^4 d(x_i, g_i)$. Cancelling $\sum_{i=1}^4 d(x_i, g_i)$ in each of E, F, G we obtain the hyperbolicity condition for the points $g_1, \ldots, g_4 \in A$. Since A is hyperbolic, the result follows.

Now assume (by symmetry) $d(g_3, g_4) \leq \delta$. Then by the triangle inequality we have

$$|d(g_1, g_3) - d(g_1, g_4)| \le \delta$$
 and
 $|d(g_2, g_4) - d(g_2, g_3)| \le \delta.$

Hence

(1) $|F_1 - G_1| - |F - G| \le 2\delta.$

If $E \leq F \leq G$ or $F \leq E \leq G$, the claim follows from (1).

So suppose $F \leq G < E$. Since

$$d(g_1, g_2) \le d(g_1, g_4) + d(g_2, g_3) + \delta = G_1 + \delta$$

we have $E_1 \leq G_1 + 2\delta$. Hence

$$G < E \le E_1 + \sum_{i=1}^4 d(x_i, g_i) \le G_1 + \sum_{i=1}^4 d(x_i, g_i) + 2\delta = G + 2\delta$$

and so $E - G \leq 2\delta$ as required.

Definition 2.12. A marked class of δ -hyperbolic spaces is a countable class $\mathcal{C} = \{(A_k, X_i^k : i \in \mathbb{N}) \mid k \in \mathbb{N}\}$ where each $A_k \in \mathcal{C}$ is δ -hyperbolic, $X_0^k = A_k, X_1^k = \emptyset$, each nonempty X_i^k is δ -closed in A_k , the family of X_i^k is closed under intersections and contains at least one X_i^k consisting of a single point.

Theorem 2.13. Fix δ and let (\mathcal{C}_0, \leq) be a marked class of δ -hyperbolic spaces. Then there is a marked class (\mathcal{C}, \leq) of δ -hyperbolic spaces containing \mathcal{C}_0 and closed under (AP) and (JEP). Hence (\mathcal{C}, \leq) has a Hrushovski limit $\mathbb{H}_{\mathcal{C}} = \mathbb{H}_{\mathcal{C},\delta}$ with at most countably many distinguisehd δ -closed subspaces.

By construction $\mathbb{H}_{\mathcal{C}}$ is universal for δ -hyperbolic spaces in \mathcal{C} and any isometry between marked subspaces of $\mathbb{H}_{\mathcal{C}}$ extends to an isometry of $\mathbb{H}_{\mathcal{C}}$.

Proof. Since any $A \in C$ contains a marked point, we obtain (JEP) by identifying marked points and amalgamating over this point. Hence it suffices to verify (AP).

We build the class C inductively from C_0 as follows: suppose at stage i we have constructed a marked class C_i of δ -hyperbolic spaces with countably many δ -closed subspaces. Then we define

$$C_{i+1} = \mathcal{C}_i \cup \{B_1 \otimes_A B_2 \colon A \leq B_1, B_2 \in C_i, A \text{ marked in } B_1, B_2\}.$$

 $\mathbf{6}$

Furthermore, for any $D = B_1 \otimes_A B_2$ with $A \leq B_1, B_2 \in \mathcal{C}_i$ we define the marked subspaces as the δ -closed subspaces X of D such that $X \cap B_1$ and $X \cap B_2$ are marked in B_1, B_2 , respectively. Then $\mathcal{C} = \bigcup_{i \in \mathbb{N}} \mathcal{C}_i$ has (AP) and (JEP). Hence the Hrushovski limit exists and is as desired and contains countably many marked subspaces and they are closed under intersections.

The previous theorem implies the existence of a universal and \leq -homogeneous δ -hyperbolic space \mathbb{H}_{δ} for the class of finite δ -hyperbolic spaces:

Corollary 2.14. Fix $\delta \geq 0$ and a countable subsemigroup Γ of $(\mathbb{R}_{\geq 0}, +)$. Let $(\mathcal{C}_{\delta,\Gamma}, \leq_{\delta})$ be the class of finite δ -hyperbolic spaces with distances in Γ . Since \mathcal{C}_{δ} is countable and contains only countably δ -closed subsets, we can consider this class as a marked class with all δ -closed subspaces. By Lemma 2.11 the class $(\mathcal{C}_{\delta,\Gamma}, \leq)$ has (AP) and (JEP) and hence the Hrushovski limit $\mathbb{H}_{\delta,\Gamma}$ exists and is universal for finite δ -hyperbolic spaces and \leq_{δ} -homogeneous.

For $\Gamma = \mathbb{Q}_{\geq 0}$ we just write \mathbb{H}_{δ} .

Clearly, this also applies to the class of finite δ -hyperbolic graphs and so we obtain:

Corollary 2.15. For any $\delta \geq 0$ there is a countable δ -hyperbolic graph \mathbb{H}_{δ} universal for finite δ -hyperbolic spaces and \leq_{δ} -homogeneous.

Note that \mathbb{H}_{δ} is a discrete δ -hyperbolic space. However, we can also obtain nondiscrete or geodesic homogeneous δ -hyperbolic spaces as the next examples show:

Examples 2.16. (i) Let T be a regular tree with vertices of valency $k \ge 3$. Note that the set of vertices of T with the graph metric is a countable 0-hyperbolic space. Since every infinite ray in T is a δ -closed subset of T, we see that a countable metric space may have uncountably many δ -closed subspaces.

(ii) Let C_0 consist of the rationals as a metric space. Since there are only countably many δ -closed subsets in the rationals, we can take all δ -closed subsets as distinguished subspaces. This satisfies the assumptions of Theorem 2.13 and $\mathbb{H}_{\mathcal{C}}$ will be a tree infinitely branching at every point with all bi-infinite paths isometric to the rationals and its completion is an \mathbb{R} -tree.

(iii) Let C_0 be the real line as a metric space with a countable collection of δ closed intervals. Then the Hrushovski limit is again an \mathbb{R} -tree infinitely branching at countably many points.

3. Projection subspaces of $\mathbb{H}_{\mathcal{C}}$

In [1] the authors show how to assemble a class \mathbf{Y} of geodesic spaces to a quasitree. The purpose of this section is to show that the quasi-tree of spaces they construct can be retrieved in the corresponding universal space $\mathbb{H}_{\mathcal{C}}$, at least if the class \mathbf{Y} is countable.

We write

 $d(X,Y) = \inf\{d(x,y) \colon x \in X, y \in Y\}.$

For δ -closed subsets, this infinimum is attained:

Lemma 3.1. Let A, B be δ -closed and disjoint subsets of a metric space X and let $G_A(B) = \{g_A(b) : b \in B\}$ and $G_B(A) = \{g_B(a) : a \in A\}$. Then

$$\operatorname{diam}(G_A(B)), \operatorname{diam}(G_A(B)) \le \delta.$$

Furthermore, for all $a \in G_A(B), b \in G_B(A)$ we have

$$d(a, g_B(a)) = d(b, g_A(b)) = d(A, B)$$

and there is an isometry

$$g: G_A(B) \longrightarrow G_B(A), \quad a \mapsto g_B(a).$$

Proof. We first claim that $G_A(B) = \{g_A(b) : b \in G_B(A)\}$ and $G_B(A) = \{g_B(a) : a \in G_A(B)\}$: let $b \in B$ and $a = g_A(b)$. Then $(a, g_B(a), b)$ is geodetic and $a = g_A(g_B(a))$.

Now let $a, a' \in G_A(B)$, say $a = g_A(b), a' = g_A(b'), b, b' \in G_B(A)$. Then $b = g_B(a), b' = g_B(a')$.

If $d(a, a') > \delta$, we have d(b, b') = d(b, a) + d(a, a') + d(a', b') since A and B are δ -closed. Hence $d(b, b') > \delta$ and thus also d(a, a') = d(a, b) + d(b, b') + d(b', a'), a contradiction.

Now suppose d(a, b) < d(a', b'). By symmetry we may assume $d(a, a') \le d(b, b')$. Then since $b' = g_B(a')$ we have d(a', b) = d(a', b') + d(b', b) > d(a', a) + d(a, b), a contradiction. Hence, d(a, b) = d(a', b') is constant and equal to d(A, B).

Finally suppose d(a, a') < d(b, b'). Then as before

$$d(b,a') = d(b,a) + d(a,a') = c + d(a,a') < d + d(b,b') = d(a',b') + d(b,b') = d(a',b),$$

a contradiction. Thus, the gate map is an isometry from $G_A(B)$ to $G_B(A)$.

Remark 3.2. Note that if A, B are closed and disjoint, then the gate sets $G_A(B)$ and $G_B(A)$ need not be δ -closed: let $A = \{x_0, x_1, x_2\}, B = \{y_0, y_2\}$ with $d(x_0, y_0) =$ $d(x_2, y_2) = d(x_0, x_2) = d(y_0, y_2) = \delta$ and $d(x_1, x_0) = 1/3\delta, d(x_1, x_2) = 2/3\delta$. Then $A, B \leq A \cup B$, but $G_A(B) = \{x_0, x_2\}$ is not δ -closed.

We now introduce the projection metric on δ -closed subsets from [1].

Definition 3.3. [see [1], 4.1] Let X, Y, Z be pairwise disjoint gated subsets of a metric space. We put

$$d_Y^{\pi}(X,Z) = \operatorname{diam}(G_Y(X) \cup G_Y(Z)).$$

Lemma 3.4. Let X, Y, Z be pairwise disjoint δ -closed subsets of a metric space and suppose $d_Y^{\pi}(X, Z) > \delta$. Then

$$d(X,Z) > d(X,Y) + d(Y,Z) + \delta.$$

Proof. Since by Lemma 3.1 each set $G_Y(X)$ has diameter $\leq \delta$, there exist $x \in X, z \in Z$ such that $d(g_Y(x), g_Y(z)) > \delta$. Since Y is δ -closed, this implies that the sequence $(x, g_Y(x), g_Y(z), z)$ is geodetic and so $x \in X$ and $g_Y(z) \in Y$ have the same gate in Z. Thus the claim follows from Lemma 3.1.

Proposition 3.5. Let **C** be a class of pairwise disjoint δ -closed subsets of a metric space, $\delta > 0$, and $X, Y, Z, W \in \mathbf{C}$. Then the projection complex axioms of [1] hold, *i.e.* we have the following:

- (PC 1) $d_V^{\pi}(X, Z) = d_V^{\pi}(Z, X);$
- (PC 2) $d_Y^{\pi}(X, Z) + d_Y^{\pi}(Z, W) \ge d_Y^{\pi}(X, W);$
- (PC 3) $\min\{d_V^{\pi}(X, Z), d_Z^{\pi}(X, Y)\} \le \delta;$

(PC 4) for all $X, Z \in \mathbf{C}$ the number of $Y \in \mathbf{C}$ such that $d_Y^{\pi}(X, Z) > \delta$ is finite.

Proof. Clearly (PC 1) holds and (PC 2) follows directly from the triangle inequality in metric spaces.

For (PC 3) suppose X, Y, Z are pairwise disjoint and δ -closed sets with both

 $d_Y^{\pi}(X,Z) > \delta$ and $d_Z^{\pi}(X,Y) > \delta$.

Then by Lemma 3.4 we have

$$d(X,Z) > d(X,Y) + d(Y,Z) + \delta$$
 and $d(X,Y) > d(X,Z) + d(Z,Y) + \delta$,

which is impossible.

For (PC 4) suppose that X, Y_1, Y_2, Z are pairwise disjoint δ -closed sets with both $d_{Y_1}^{\pi}(X, Z), d_{Y_2}^{\pi}(X, Z) > \delta$. Since all gate sets have diameter at most δ , we cannot have both $d_{Y_1}^{\pi}(X, Y_2) \leq \delta$ and $d_{Y_1}^{\pi}(Z, Y_2) \leq \delta$. Now if $d_{Y_1}^{\pi}(X, Y_2) > \delta$, then by Lemma 3.4 we have

$$d(X, Y_2) > d(X, Y_1) + d(Y_1, Y_2) + \delta$$
 and $d(X, Z) > d(X, Y_2) + d(Y_2, Z) + \delta$.

Thus, $d(X, Z) > d(X, Y_1) + d(Y_1, Y_2) + d(Y_2, Z) + 2\delta$. Hence the number of $Y \in \mathbb{C}$ such that $d_Y^{\pi}(X, Z) > \delta$ is bounded by $d(X, Z)/\delta$.

4. Stationary independence relations

In this section we verify that each of the universal \leq -homogeneous spaces $\mathbb{H}_{\mathcal{C}}$ constructed in the previous section supports a stationary independence relation, which will be useful in the study of their isometry groups (see [8].)

Since marked subspaces are closed under intersections, we have the following stronger form of Corollary 2.8:

Corollary 4.1. Any non-empty finite subset A of \mathbb{H}_C is contained in a unique smallest δ -closed marked set $\operatorname{cl}(A) = \bigcap \{B \leq_{\delta} X : A \subset B\}$.

Definition 4.2. For finite subsets $A, B, C \subseteq \mathbb{H}_{\mathcal{C}}, B \neq \emptyset$, we say that A is independent from C over $B, A \perp_B C$, if and only if $cl(A \cup B \cup C)$ is isometric to the canonical amalgam of $cl(A \cup B) \otimes_{cl(B)} cl(B \cup C)$.

Note: the sets $\operatorname{cl}(A \cup B \cup C)$, $\operatorname{cl}(A \cup B)$ and $\operatorname{cl}(A \cup C)$ are marked subsets of $\mathbb{H}_{\mathcal{C}}$. The canonical amalgam $\operatorname{cl}(A \cup B) \otimes_{\operatorname{cl}(B)} \operatorname{cl}(B \cup C)$ is an element in \mathcal{C} . So A and C are independent over B if the distances in $\operatorname{cl}(A \cup B \cup C)$ agree with the distances in the canonical amalgam.

Proposition 4.3. $\ \ defines a local stationary independence relation on <math>\mathbb{H}_{\mathcal{C}}$, *i.e.* the following conditions are satisfied for all finite sets $A, B, C, D \subset \mathbb{H}_{\mathcal{C}}$ where $B \neq \emptyset$:

- (1) $A \bigsqcup_{B} C$ is invariant under automorphisms of M;
- (2)

$$A \underset{B}{\downarrow} CD$$
 if and only if $A \underset{B}{\downarrow} C$ and $A \underset{BC}{\downarrow} D$;

- (3) $A \bigsqcup_{B} C$ if and only if $C \bigsqcup_{B} A$;
- (4) there is some finite set A' such that $cl(A \cup B)$ and $cl(A' \cup B)$ are isometric and $A' \downarrow_B C$;
- (5) if $cl(A \cup B)$ and $cl(D \cup B)$ are isometric and both A and D are independent from C over B, then $cl(A \cup B \cup C)$ and $cl(D \cup B \cup C)$ are isometric.

Proof. All conditions follow easily from the \leq -homogeneity and universality of $\mathbb{H}_{\mathcal{C}}$.

5. The isometry group of \mathbb{H}_{δ}

We now turn our attention to \mathbb{H}_d , the unique δ -hyperbolic space universal and \leq_{δ} -homogeneous for finite δ -hyperbolic spaces with distances in \mathbb{Q} .

Its isometry group $G = \text{Isom}(\mathbb{H}_{\delta})$ is a polish group, where a neighbourhood basis of the identy is given by point stabilizers of finite δ -closed subsets of \mathbb{H}_{δ} . Since every element of \mathbb{H}_{δ} is δ -closed in \mathbb{H}_{δ} , it follows that G acts transitively.

In the case of the Urysohn space \mathbb{U} it is easy to see that the isometry group acts primitively: suppose there is an invariant equivalence relation E with E(x,y) for some $x, y \in \mathbb{U}$ and let $z \in \mathbb{U}$. Then by universality there is a sequence $(x_0 = x, x_1 = y, \ldots, x_n = z)$ for some $n \in \mathbb{N}$ such that $d(x_i, x_{i-1}) = d(x_1, x_0), i = 1, \ldots n$. By homogeneity and invariance of E we see that E(x, z) showing that the equivalence class is all of \mathbb{U} .

It is also easy to see that for any regular tree T the isometry group acts primitively on the vertices.

In contrast to this, we will now show that \mathbb{H}_{δ} has non-trivial invariant equivalence relations, in fact uncountably many:

To show this, we introduce the following auxilliary graph structure on \mathbb{H}_{δ} . Let $R \subset \Gamma_{>\delta}$. We define edges E_R on \mathbb{H}_{δ} in the following way: for $x, y \in \mathbb{H}_{\delta}$ we put $E_R(x, y)$ if and only if $d(x, y) \in R$ and $\{x, y\} \leq \mathbb{H}_{\delta}$.

Lemma 5.1. For any $R \subset \Gamma_{>\delta}$, the graph $(\mathbb{H}_{\delta}, E_R)$ is a forest, i.e. the connected components of this graph are regular trees with infinite valencies and any connected subgraph is δ -closed.

Proof. The fact that every vertex in this graph has infinite valency follows directly from the universality of \mathbb{H}_{δ} : clearly, the set $X = \{x, y_1, \ldots, y_n\}$ with $d(x, y_i) \in R$ and (y_i, x, y_j) geodetic for $1 \leq i \neq j \leq n$ can be strongly embedded into \mathbb{H}_{δ} for any $n \in \mathbb{N}$. Then $\{x, y_i\} \leq X \leq \mathbb{H}_{\delta}$ for any $i \leq n$ and hence $E_R(x, y_i)$ holds. Since the isometry group acts transitively on the points of \mathbb{H}_{δ} , it follows that every point has infinite valency in this graph.

By Lemma 2.9 it follows inductively, that any finite connected subgraph of $(\mathbb{H}_{\delta}, E_R)$ that does not contain a cycle is δ -closed in \mathbb{H}_{δ} : clearly this holds for any edge of the graph. So suppose inductively that A is a finite subgraph without cycles, let $x \in A$ and extend A by an edge $E_R(x, y)$. Then the conditions of Lemma 2.9 are satisfied, and hence $A \cup \{y\} \leq \mathbb{H}_{\delta}$. Furthermore, if A is a finite connected subgraph of $(\mathbb{H}_{\delta}, E_r)$ without cycles, then for $x, y \in A$ the unique path in A from x to y is geodetic, i.e. the distance between points in A in the metric of \mathbb{H}_{δ} is the graph distance in A where an edge $E_R(x, y)$ has length r if $d(x, y) = r \in R$.

Now suppose that $(x_0, \ldots, x_n = x_0)$ is a δ -closed cycle in this graph with n > 2. Then the subpath $X_0 = \{x_1, x_2, x_{n-1}\}$ is geodesic and strong in \mathbb{H}_{δ} by the previous paragraph. Thus, by Remark 2.4, $g_{X_0}(x_0) \in \{x_{n-1}, x_1\}$. However, if $x_1 = g_{X_0}(x_0)$), then (x_0, \ldots, x_{n-1}) is geodesic. By Remark 2.4, the set $\{x_0, x_{n-1}\}$ is not δ -closed in \mathbb{H}_{δ} and so we cannot have $E_R(x_{n-1}, x_0)$. The symmetric argument applies if $x_{n-1} = g_{X_0}(x_0)$). Thus we cannot have both $E_R(x_0, x_1)$ and $E_R(x_{n-1}, x_0)$, so there are no cycles.

To complete the proof, suppose that X is a connected subgraph. Then any finite connected subgraph of X is δ -closed in \mathbb{H}_{δ} . Suppose X is not δ -closed. First assume that X is not gated. Then there exist $a \in \mathbb{H}_{\delta} \setminus X$ for which there is no gate in X. Let $x \in X$ be such that d(a, x) is minimal. Then x is unique: if $y \in X$ with d(a, x) = d(a, y), then the convex closure of $\{x, y\}$ in X is finite and not gated, contradicting the previous paragraph. Furthermore for any $y \in X$ we have d(a, y) = d(a, x) + d(x, y) as this holds in the convex closure of $\{x, y\}$ in X. Now let $a, b \in \mathbb{H}_{\delta} \setminus X$. Then $d(g_X(a), g_X(b)) > \delta$ and since the convex closure of $\{g_X(a), g_X(b)\}$ is δ -closed, it follows that d(a, b) is as required.

Proposition 5.2. Isom (\mathbb{H}_{δ}) acts transitively, but not primitively on \mathbb{H}_{δ} . In fact, there exist uncountably many non-trivial equivalence relations on \mathbb{H}_{δ} invariant under Isom (\mathbb{H}_{δ}) .

Proof. Transitivity of the action follows directly from the \leq -homogeneity and was already noted above.

To see that the action is not primitive, let $R \subset \Gamma_{>\delta}$ and define the equivalence relation T_R on \mathbb{H}_{δ} where $T_R(x, y)$ if and only if x and y lie in the same E_R -connected component. Clearly this is invariant under the action of $\mathrm{Isom}(\mathbb{H}_{\delta})$. Since there are uncountably many subsets R, the claim follows.

Definition 5.3. For $R \subset \Gamma$ we call the E_R connected components of \mathbb{H}_{δ} a T_R -tree.

Remark 5.4. Note that each T_R -tree X is geodetically embedded, i.e. if (x_0, x_1, \ldots, x_n) is a path in a T_R -tree, then (x_0, x_1, \ldots, x_n) is geodetic. Since every finite convex subgraph of X is δ -closed in \mathbb{H}_{δ} by the proof of Lemma 5.1, it follows in particular, that X is δ -closed in \mathbb{H}_{δ} .

Hence we have:

Corollary 5.5. No nontrivial element of $Aut(\mathbb{H}_{\delta})$ has bounded displacement.

Proof. If $g \in \operatorname{Aut}(\mathbb{H}_{\delta})$ has bounded displacement, then so does every commutator [g,h] for $h \in \operatorname{Aut}(\mathbb{H}_{\delta})$. By taking an appropriate commutator we may assume that g fixes a point x and acts nontrivially on the T_R -tree containing x. But no nontrivial isometry of T_R has bounded displacement, proving the corollary. \Box

Corollary 5.6. Suppose $\alpha \in \operatorname{Aut}(\mathbb{H}_{\delta})$ acts as a translation on some T_R -tree X for some R. Then α has no fixed point. In particular, any $\alpha \in \operatorname{Aut}(\mathbb{H}_{\delta})$ for which there exists $x \in \mathbb{H}_{\delta}$ such that $\{x, \alpha(x)\}$ is δ -closed and $d(x, \alpha(x)) > \delta$ is fixed point free.

Proof. Note that X is δ -closed in \mathbb{H}_{δ} . Let $c \in \mathbb{H}_{\delta} \setminus X$ and $g = g_X(c)$. Then $d(c,g) = d(\alpha(c), \alpha(g)) < d(c, \alpha(g))$ by Remark 2.2 and hence $c \neq \alpha(c)$.

If $\{x, \alpha(x)\}$ is δ -closed and $d = d(x, \alpha(x)) > \delta$, then α leaves the T_R -tree X containing x for $R = \{d\}$ invariant and acts on X as a translation. So the result from the first part.

Proposition 5.7. $G = \operatorname{Aut}(\mathbb{H}_{\delta})$ has no dense conjugacy class. For any finite set *B* the pointwise stabilizer $G_{\operatorname{cl}(B)}$ in *G* has a dense conjugacy class.

Proof. We use the criterion from [8], Lemma 2.8: G has a dense conjugacy class if and only if for any finite tuples x, y, a, b such that $cl(x) \cong cl(y)$ and $cl(a) \cong cl(b)$ there are tuples x', y' such that $cl(x'y') \cong cl(xy)$ and $cl(x'a) \cong cl(y'b)$.

To see that this critrion is violated, let (x_0, x_1, x_2) be a path in a $T_{2\delta}$ -tree and put $x = \operatorname{cl}(x) = (x_0, x_1), y = \operatorname{cl}(y) = (x_1, x_2)$ and let $a = b \in \mathbb{H}_{\delta}$ be an arbitrary point. Suppose there are tuples x', y' such that $\operatorname{cl}(x'y') \cong \operatorname{cl}(xy)$ and $\operatorname{cl}(x'a) \cong \operatorname{cl}(y'a)$.

Then $x' = (x'_0, x'_1), y' = (x'_1, x'_2)$ for a path (x'_0, x'_1, x'_2) in some $T_{2\delta}$ -tree and by \leq_{δ} -homogeneity there is some $g \in G$ such that g(x'a) = (y'a). Thus, g acts as a translation on the $T_2\delta$ -connected component containing x'_1 and hence g has no fixed point by Corollary 5.6, a contradiction.

This shows that G does not have a dense conjugacy class. The fact that $G_{cl(B)}$ has a dense conjugacy class for any finite subset B follows from [8], Lemma 2.8 using the fact that \mathbb{H}_{δ} has a local stationary independence relation.

6. FINAL REMARKS AND OPEN QUESTIONS

In the previous sections we tried to point out similarities and dissimilarities between the universal δ -hyperbolic spaces constructed here and the Urysohn space on the one hand and regular trees on the other hand.

In light of the fact that the isometry group of a regular tree and the isometry group of the Urysohn space modulo the normal subgroup of bounded displacement are simple groups, it is natural to ask:

• Is the isometry group of \mathbb{H}_{δ} topologically simple, or even abstractly simple?

Since there \mathbb{H}_{δ} has no isometries of bounded displacement, it seems likely that the answer is 'yes'. However, the methods from [8] do not apply here directly since $\operatorname{Isom}(\mathbb{H}_{\delta})$ has no dense conjugacy class. In the case of trees and more generally right-angled buildings, the automorphism groups also miss a dense conjugacy class. However, the methods from [8] can be applied to stabilizers of maximal flags because these stabilizers have dense conjugacy classes, and the same was shown above for \mathbb{H}_{δ} .

For right-angled buildings, one can then deduce abstract simplicity of the automorphism group using the fact that stabilizers of flags are maximal subgroups. However, as pointed out above, the isometry group of \mathbb{H}_{δ} does not act primitively. Hence point stabilizers are not maximal in this setting and it is harder to approach the simplicity problem.

Another natural question addresses the model theoretic properties of these hyperbolic spaces:

• Is there a natural class of δ -hyperbolic spaces, $\delta > 0$, which is stable?

For 0-hyperbolic spaces (and right-angled buildings) it was shown in [10] that the theory is ω -stable. This can be shown using the notion of independence defined above and verifying that it satisfies the necessary conditions for stability. In general, one cannot expect arbitrary δ -hyperbolic spaces to be stable. It would be interesting to find a natural class of hyperbolic spaces with stable theory.

7. Funding

This research was partially funded through the Cluster of Excellence by the German Research Foundation (DFG) under Germany's Excellence Strategy EXC 2044–390685587, Mathematics Münster: Dynamics–Geometry–Structure and by CRC 1442 Geometry: Deformations and Rigidity.

References

- M. Bestvina, K. Bromberg, K. Fujiwara, Constructing group actions on quasi-trees and applications to mapping class groups. Publ. Math. Inst. Hautes Études Sci. 122 (2015), 1–64.
- [2] S. Gao, Su, Ch. Shao, Polish ultrametric Urysohn spaces and their isometry groups. Topology Appl. 158 (2011), no. 3, 492–508.
- [3] Y. Ishiki, Constructions of Urysohn universal ultrametric spaces. p-Adic Numbers Ultrametric Anal. Appl. 15 (2023), no. 4, 266–283.
- [4] Y. Ishiki, Uniqueness and homogeneity of non-separable Urysohn universal ultrametric spaces. Topology Appl. 342 (2024), Paper No. 108762, 11 pp.
- [5] M. Katetov, On universal metric spaces, in: General Topology and Its Relations to Modern Analysis and Algebra, VI, Prague, 1986, in: Res. Exp. Math., vol. 16, Heldermann, Berlin, 1988, pp. 323–330.
- [6] V. A. Lemin, On a universal ultrametric space. Topology Appl. 103 (2000), no. 3, 339–345.
- [7] V.A. Lemin, On metrically universal ultrametric spaces LV_τ and LW_τ. Ultrametric functional analysis (Nijmegen, 2002), 191–205, Contemp. Math., 319, Amer. Math. Soc., Providence, RI, 2003.
- [8] K. Tent, M. Ziegler, On the isometry group of the Urysohn space, Journal of the London Mathematical Society 2012; doi: 10.1112/jlms/jds027.
- [9] K. Tent, M. Ziegler, A course in model theory. Lecture Notes in Logic, 40. Association for Symbolic Logic, La Jolla, CA; Cambridge University Press, Cambridge, 2012
- [10] K. Tent, The free pseudospace is N-ample, but not (N+1)-ample. J. Symb. Log. 79 (2014), no. 2, 410–428.
- [11] Zh. Wan, A novel construction of Urysohn universal ultrametric space via the Gromov-Hausdorff ultrametric. Topology Appl. 300 (2021), Paper No. 107759, 10 pp.