A Correlation Inequality on Three Functions

Kada K Williams

February 21, 2025

Abstract

Let X and Y be upward closed set systems in the lattice of $\{0, 1\}^n$. The celebrated Harris-Kleitman inequality implies that if $|X| = \alpha 2^n$, $|Y| = \beta 2^n$, the density of the set of points in exactly one of X and Y is maximal when X and Y are independent, meaning $|X \cap Y| = \alpha \beta 2^n$. Is the same true of three upward closed systems, X, Y, and Z? Suppose |X| = |Y| = |Z|. Kahn asked whether the set of points in exactly one of X, Y, Z has density at most $\frac{4}{9}$. We answer this question in the negative.

AMS subject classification: 05A10, 05A20, 05D05

1 Introduction

Let (P, <) be a finite poset. We say that $X \subset P$ is an upward closed system if whenever a < b and $a \in X$, also $b \in X$. Notice that if X and Y are upward closed, then so is $X \cap Y$.

Of particular interest are upward closed systems in the poset Q_n . Our notation is standard, with $[n] = \{1, 2, ..., n\}$ and Q_n is the power set of [n] ordered by \subset .

Theorem 1 (Harris-Kleitman). Let X and Y be upward closed set systems on a ground set of n elements. Then the following inequality holds [1], [2]:

$$\frac{|X \cap Y|}{2^n} \ge \frac{|X|}{2^n} \cdot \frac{|Y|}{2^n}$$

Observe that if X is upward closed, its complement X^c is downward closed, meaning that it is upward closed in the reverse of Q_n . We deduce that if $|X| = \alpha 2^n$, $|Y| = \beta 2^n$, then $|X \cap Y| \ge \alpha \beta \cdot 2^n$ and $|X^c \cap Y^c| \ge (1-\alpha)(1-\beta) \cdot 2^n$. Therefore,

$$|X \cap Y^c| + |X^c \cap Y| = 2^n - |X \cap Y| - |X^c \cap Y^c| \le (\alpha + \beta - 2\alpha\beta) \cdot 2^n$$

Let X, Y, and Z be upward closed systems in Q_n . Applying the Harris-Kleitman inequality to $X \cap Y$ and Z gives a lower bound on the set of points in all three sets,

and similarly for their complements. It is therefore natural to explore the set of points in exactly one of the three systems,

$$S_1 = (X \cap Y^c \cap Z^c) \sqcup (X^c \cap Y \cap Z^c) \sqcup (X^c \cap Y^c \cap Z).$$

Let $|X| = \alpha 2^n$, $|Y| = \beta 2^n$, and $|Z| = \gamma 2^n$. Supposing $Q_n \cong Q_a \times Q_b \times Q_c$, where a + b + c = n, let a set be in X depending only on its Q_a component, and similarly for Y with Q_b and Z with Q_c . Then

$$|X \cap Y^c \cap Z^c| = \alpha(1-\beta)(1-\gamma) \cdot 2^n,$$

and similarly for $X^c \cap Y \cap Z^c$ and $X^c \cap Y^c \cap Z$. Specially, if $\alpha = \beta = \gamma = \rho$, then

$$\frac{|S_1|}{2^n} = 3\rho(1-\rho)^2 \le \frac{4}{9}.$$

Conjecture 2. Let $X, Y, Z \subset Q_n$ be upward closed set systems of density ρ . Then

$$\frac{|S_1|}{2^n} \le 3\rho(1-\rho)^2$$

While this conjecture is false for n = 5, an upper bound is relatively easy to glean for equal-sized systems.

Theorem 3. Let $|X| = |Y| = |Z| = \rho \cdot 2^n$. Then $\frac{|S_1|}{2^n} \le 3\rho \cdot \frac{1-\rho}{1+\rho}$.

The maximum of $\frac{3\rho(1-\rho)}{1+\rho}$ is ≈ 0.515 , achieved when $\rho = \sqrt{2} - 1 \approx 0.414$. This yields a heuristic that if $\rho = \frac{3}{8} \approx 0.375$, the size of S_1 can be nearly maximal.

Conjecture 4 (Kahn). If |X| = |Y| = |Z|, then $\frac{|S_1|}{2^n} \le \frac{4}{9}$.

This conjecture of Jeff Kahn [4] was prompted by the work of Sahi [3]. We provide a counterexample in n = 21 dimensions, where the density of S_1 exceeds 0.447

2 Proof that Conjecture 2 is false

In Q_5 , let X and Y contain the sets that have 1 and 2 as an element, respectively. To begin with, let W contain the sets with at least 3 elements in [5]. Clearly, X, Y, and W are upward closed with density $\frac{1}{2}$. Moreover, $|X \cap Y^c \cap W^c| = |X^c \cap Y \cap W^c| = 4$, counting a singleton and three pairs.

The only set in $W \setminus X \setminus Y$ is $\{3, 4, 5\}$. This can be improved by introducing the sets $\{3, 4\}$ and $\{3, 5\}$ into W, removing the sets $\{1, 4, 5\}$ and $\{2, 4, 5\}$ in return. The resulting set system Z is upward closed, and while $X^c \cap Y^c \cap Z$ increased by 2, there is also a new set contained exactly in X and exactly in Y. Thus, $|S_1| = 5 + 5 + 3 = 13$, which is larger than $\frac{4}{9} \cdot 2^5 = 12$.

3 The upper bound of Theorem 3

For each i = 0, 1, 2, 3, let $s_i \cdot 2^n$ be the number of sets in Q_n that occur in i many of $\{X, Y, Z\}$. We normalised such that

$$s_0 + s_1 + s_2 + s_3 = 1.$$

Theorem 1 implies that $|X \cap Y \cap Z| \ge \rho |X \cap Y|$, and similarly for $Y \cap Z$ and $Z \cap X$. Adding these up yields $3(1 - \rho)s_3 \ge \rho s_2$, where $s_2 \ge 0$. The same argument for the complements informs us that $3\rho s_0 \ge (1 - \rho)s_1$, where $s_1 \ge 0$. Finally, the constraint on the size of X, Y, and Z implies that

$$s_1 + 2s_2 + 3s_3 = 3\rho.$$

Given these constrains on (s_0, s_1, s_2, s_3) , what is the maximum of s_1 ?

The strategy for solving such an optimisation problem is the following. As long as the linear equations for (s_0, s_1, s_2, s_3) are underdetermined, it is possible to move this vector along a line while increasing s_1 , until one of the inequalities becomes an equality. Iterating this observation, we deduce that s_2 is maximal with equalities $3s_0 = \frac{1-p}{p}s_1$ and $s_2 = 0$. Solving the linear system, we obtain $s_1 \leq \frac{3(1-\rho)\rho}{1+\rho}$. The maximum of this bound is $9 - 6\sqrt{2} \approx 0.515$, achieved when $\rho = \sqrt{2} - 1 \approx 0.414$.

Can equality hold in the poset Q_n ? Consider the weighted poset

$$P = \{a, A, p_1, p_2, p_3\},\$$

where $a < p_i$ and $p_i < A$, with weights $\frac{(1-p)^2}{1+p}$, $\frac{(1-p)p}{1+p}$, and $\frac{2p^2}{1+p}$ for a, p_i , and respectively. Theorem 1 holds for upward closed systems in P, the only remarkable instance being of two systems of size three, where $\left(\frac{2p}{1+p}\right)^2 \leq p$. It would seem that P cannot be simulated in Q_n , but we did not find any obvious reason for this.

4 A counterexample for Conjecture 4

We consider the weighted hypercube $Q_n(p)$, which is the measure space of n independent Bernoulli bits B(p). That means that a set of k elements in Q_n has measure $p^k(1-p)^{n-k}$. If X and Y are upward closed, it still holds that the measure of $X \cap Y$ is at least the product of the measures of X and Y, by the FKG inequality [5], or by application of a measurable map $f: Q_{bm} \to Q_m\left(\frac{u}{2^v}\right)$. Thus, we generate a counterexample in Q_{21} of density ≈ 0.447 from an approximate counterexample in $Q_7\left(\frac{3}{8}\right)$.

Consider X and Y to be the sets containing 1 and 2, respectively, in $Q_n(p)$. Let Z contain the sets of size at least l + 1, as well as the sets of size l that are not in X or Y. Then $X \cap Y^c \cap Z^c$ contains $\binom{n-2}{k-1}$ many sets of size k, where $1 \le k \le l$. Also, $X^c \cap Y^c \cap Z$ contains $\binom{n-2}{k}$ many sets of size k, where $l \le k \le n-2$. The total measure of sets in exactly one of X, Y, Z is thus given by

$$q = 2\sum_{k=1}^{l} \binom{n-2}{k-1} p^k (1-p)^{n-k} + \sum_{k=l}^{n-2} \binom{n-2}{k} p^k (1-p)^{n-k}.$$

Compare this to the binomial expansion of $(p + (1 - p))^{n-2}$:

$$\sum_{k=1}^{l} \binom{n-2}{k-1} p^{k-1} (1-p)^{n-k-1} + \sum_{k=l}^{n-2} \binom{n-2}{k} p^k (1-p)^{n-k-2} = 1.$$

The first summand is 2p(1-p) as much, while the second summand is $(1-p)^2$ as much. Specifically, if $p = \frac{1}{3}$, these coefficients both equal $\frac{4}{9}$.

Since $q(p) = \frac{4}{9}$ if $p = \frac{1}{3}$, unless q attains a local maximum at $\frac{1}{3}$, a different value of p could yield a larger value of q. For example, in the case of n = 7 and l = 3, q(p) > 0.447 if $p = \frac{3}{8}$. There, the measure of Z is only ≈ 0.323 , but including all the supersets of $\{1, 2\}$ from $X \cap Y$ boosts the measure of Z up to $\approx 0.377 > \frac{3}{8}$.

A measure-preserving map $f : Q_{21} \to Q_7\left(\frac{3}{8}\right)$ arises from the upward closed $I = \{\{1, 2\}, \{1, 3\}, \{1, 2, 3\}\} \subset Q_3$. Viewing $Q_{21} \cong (Q_3)^7$, we map I to 1 and I^c to 0 at every coordinate. Clearly, the preimages of X, Y, and Z are upward closed.

In Q_{21} , the upward closed system $f^{-1}(Z)$ can be modified by including sets that map to a superset of $\{1, 2\}$. Including only a maximal set in each step, the size of the set system can be incremented until it equals $\frac{3}{8}$ exactly. Meanwhile, the measure of sets contained in exactly one of our three set systems is $> 0.447 > \frac{4}{9}$.

5 Conclusion

Regarding the density of S_1 , there is a wide gap between our upper bound of ≈ 0.515 and our counterexample of ≈ 0.447 , due to the structure of Q_n . Nevertheless, Kahn also asked [4] whether there is a counterexample where the sets $X \cap Y^c \cap Z^c$, $X^c \cap Y \cap Z^c$, and $X^c \cap Y^c \cap Z$ each are of density more than $\frac{4}{27} \approx 0.148$. In particular, can their densities each exceed 0.153? These questions continue to be open.

6 Acknowledgements

The author is grateful to Imre Leader for proofreading and fruitful suggestions. The author is a recipient of the Internal Graduate Studentship in Trinity College, Cambridge, UK.

References

- [1] Harris, T. E. A lower bound for the critical probability in a certain percolation *process*. Proc. Cambridge Philos. Soc. **56**, 13-20, 1960.
- [2] Kleitman, D. J. *Families of non-disjoint subsets*. J. Combinatorial Theory 1, 153-155, 1966.
- [3] Sahi, S. Higher correlation inequalities. Combinatorica 28 (2), 209-227, 2008.
- [4] Narayanan, B. Personal communication, 2022.
- [5] Grimmett, G. R. *Percolation*. Grundlehren der mathematischen Wissenschaften **321**, Springer-Verlag, pp. 34-36, 1999.

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, UNIVERSITY OF CAMBRIDGE, WILBERFORCE ROAD, CAMBRIDGE CB3 0WB.

E-mail address: kkw25@cam.ac.uk