The Fourier coefficients of the holomorphic multiplicative chaos in the limit of large frequency

Joseph Najnudel¹ Elliot Paquette² Nick Simm³ Truong Vu⁴

¹School of Mathematics; University of Bristol, BS8 1UG, United Kingdom; e-mail: *joseph.najnudel@bristol.ac.uk

²Department of Mathematics and Statistics; McGill University, Montreal, Quebec, Canada; e-mail: **elliot.paquette@mcgill.ca

³Department of Mathematics; University of Sussex, BN1 9RH, United Kingdom; e-mail: [†]n.j.simm@sussex.ac.uk

⁴Department of Mathematics, Statistics, and Computer Science, University of Illinois at Chicago, Chicago, Illinois, USA; e-mail: [‡]tvu25@uic.edu

Abstract: The holomorphic multiplicative chaos (HMC) is a holomorphic analogue of the Gaussian multiplicative chaos. It arises naturally as the limit in large matrix size of the characteristic polynomial of Haar unitary, and more generally circular- β -ensemble, random matrices. We consider the Fourier coefficients of the holomorphic multiplicative chaos in the L^1 -phase, and we show that appropriately normalized, this converges in distribution to a complex normal random variable, scaled by the total mass of the Gaussian multiplicative chaos measure on the unit circle. We further generalize this to a process convergence, showing the joint convergence of consecutive Fourier coefficients. As a corollary, we derive convergence in law of the secular coefficients of sublinear index of the circular- β -ensemble for all $\beta > 2$.

MSC2020 subject classifications: Primary 60B20, 60F05; secondary 60F05. Keywords and phrases: random matrix, Circular beta ensemble, Circular unitary ensemble, magic squares, secular coefficient, characteristic polynomial, Gaussian multiplicative chaos, Ewens sampling formula, martingale central limit theorem.

1. Introduction

In this paper we consider the Holomorphic multiplicative chaos. This is a distribution on the unit circle, arising in the theory of random matrices and analytic number theory. See previous work of the first three authors [NPS23] where it was introduced and studied. To define it, let $\{\mathcal{N}_k\}_{k=1}^{\infty}$ be i.i.d. standard complex normal random variables, i.e. having moments

$$\mathbb{E}(\mathcal{N}_k) = 0, \quad \mathbb{E}(\mathcal{N}_k^2) = 0, \quad \text{and} \quad \mathbb{E}(|\mathcal{N}_k|^2) = 1.$$
(1.1)

Let $G^{\mathbb{C}}$ be the Gaussian analytic function on the unit disc \mathbb{D} ,

$$G^{\mathbb{C}}(z) = \sum_{k=1}^{\infty} \frac{z^k}{\sqrt{k}} \mathcal{N}_k.$$
 (1.2)

For trigonometric polynomials ϕ if we let $z \mapsto \phi(z)$ for $z \in \mathbb{D}$ be the continuous harmonic extension to \mathbb{D} , we then define the random distribution

$$(\mathrm{HMC}_{\theta}, \phi) \coloneqq \lim_{r \to 1} \frac{1}{2\pi} \int_{0}^{2\pi} e^{\sqrt{\theta} G^{\mathbb{C}}(re^{i\vartheta})} \overline{\phi(r\vartheta)} \mathrm{d}\vartheta, \qquad (1.3)$$

which is the *holomorphic multiplicative chaos* in terms of the inverse temperature parameter

$$\theta \coloneqq \frac{2}{\beta}.\tag{1.4}$$

We define for $n \in \mathbb{N}_0$ the Fourier coefficient of the HMC

$$c_n \coloneqq (\mathrm{HMC}_{\theta}, \vartheta \mapsto e^{in\vartheta}). \tag{1.5}$$

Equivalently, the coefficients c_n can be extracted from a generating function by the formula

$$c_n = [z^n] e^{\sqrt{\theta} G^{\mathbb{C}}(z)} = [z^n] \exp\left(\sqrt{\theta} \sum_{k=1}^{\infty} \frac{z^k}{\sqrt{k}} \mathcal{N}_k\right), \qquad (1.6)$$

where the notation $[z^n]h(z)$ denotes the coefficient of z^n in the power series expansion of h(z) around the point z = 0. We could as well define this for $n \in \mathbb{Z}$, but for negative integers, this would be 0.

The coefficients (1.6) were studied in detail in [NPS23], where a convergence in distribution result is shown for any $0 < \theta < \frac{1}{2}$. The recent work [GW24a] established analogues of this convergence for a class of random multiplicative functions with the same restriction $0 < \theta < \frac{1}{2}$. Although we do not consider it here, the case $\theta = 1$ has attracted recent attention due to its connections to analytic number theory, in particular the phenomenon of 'better than square root cancellation'. In the critical $\theta = 1$ case, the naïve bound $\mathbb{E}(|c_n|) \leq \sqrt{\mathbb{E}(|c_n|^2)}$ overestimates the true order of magnitude in n. Sharp upper and lower bounds on this quantity are derived in [SZ22]; see also [NPS23] for another proof of the upper bound.

Moreover, for various randomized analytic number theory problems, such as random multiplicative functions, better-than-square-root cancellation has been proven [Har20, Ger22, SZ22, Xu24, GW24b]. See also [Har24] for a recent review and see [GZ24] for related work on non-Gaussian holomorphic multiplicative chaoses.

For $0 < \theta < 1$, the Gaussian multiplicative chaos can be defined via (1.2) as

$$\mathrm{GMC}_{\theta}(\mathrm{d}\vartheta) \coloneqq \lim_{r \to 1} (1 - r^2)^{\theta} |e^{\sqrt{\theta} G^{\mathbb{C}}(re^{i\vartheta})}|^2 \mathrm{d}\vartheta = \lim_{r \to 1} (1 - r^2)^{\theta} e^{\sqrt{\theta} G(re^{i\vartheta})} \mathrm{d}\vartheta, \quad (1.7)$$

where $G(z) = 2\text{Re}(G^{\mathbb{C}}(z))$, with the sense of the limit being an in-probability, weak-* convergence. The existence of this limit as a random measure is shown¹ in [JS17a] where the convergence is shown to hold in L^q for any 0 < q < 1 (c.f. [CN19, Proposition 3.1]). Moreover it is shown in [Rem20] that the total mass of this particular random measure has law characterized by the natural analytic continuation of the Morris integral, i.e. for any p > 0 with $p\theta < 1$,

$$\mathbb{E}(\mathcal{M}^{p}_{\theta}) = \frac{\Gamma(1-p\theta)}{\Gamma(1-\theta)^{p}}, \quad \text{where} \quad \mathcal{M}_{\theta} \coloneqq \frac{1}{2\pi} \int_{0}^{2\pi} \text{GMC}_{\theta}(\mathrm{d}\vartheta). \tag{1.8}$$

The main goal of this paper is to prove:

Theorem 1.1 (L^1 -phase). For any $0 < \theta < 1$, we have the convergence in distribution

$$\frac{c_n}{\sqrt{\mathbb{E}(|c_n|^2)}} \xrightarrow[n \to \infty]{d} \sqrt{\mathcal{M}_{\theta}} \mathcal{Z}$$
(1.9)

where Z and \mathcal{M}_{θ} are independent, Z is standard complex normal, and \mathcal{M}_{θ} has law (1.8).

This improves over [NPS23] which establishes the same theorem in the range θ from $0 < \theta < \frac{1}{2}$, and it now covers the full range of subcritical HMC. In the case $\theta = 1$, we expect that after rescaling the prelimit by $(\log n)^{1/4}$ there is again convergence in distribution to a random variable connected to the critical Gaussian multiplicative chaos; see also [ASV+22] for simulations related to this convergence.

We note there have been recent developments on the Fourier coefficients of the real GMC, which display a formula similar to (1.8); in [GV23], they show the *n*-th Fourier coefficient of $\text{GMC}_{\theta}(d\vartheta)$ converges in law (after renormalization) to a complex normal multiplied by $\sqrt{M_{2\theta}}$ for $2\theta < 1$.

1.1. Process convergence

The proof we give extends to finite-dimensional marginal convergence of the $\{c_{n+k} : k \in \mathbb{Z}\}$ in the limit of $n \to \infty$. To formulate this convergence, we can equivalently consider a fixed complex polynomial p and define the statistic

$$X_n[p] \coloneqq (\mathrm{HMC}_{\theta}, \vartheta \mapsto e^{in\vartheta} p(e^{i\vartheta}))$$

A small modification of the proof of Theorem 1.1 gives rise to the following generalization.

Theorem 1.2 (L^1 -phase-multipoint). For any $0 < \theta < 1$, we have the convergence in distribution

$$\frac{(X_n[p])}{\sqrt{\mathbb{E}(|c_n|^2)}} \xrightarrow[n \to \infty]{d} \sqrt{\mathcal{M}_{\theta}(|p^2|)} \mathcal{Z}$$
(1.10)

¹The Gaussian field used in [JS17a, Eqn 6.1] contains a constant term in its Fourier series definition in comparison to our field G. This turns out not to affect the convergence results of [JS17a], see [NPS23, Appendix B].

where Z is independent of GMC_{θ} , Z is standard complex normal, and where

$$\mathcal{M}_{\theta}(|p^2|) \coloneqq \frac{1}{2\pi} \int_0^{2\pi} |p(e^{i\vartheta})|^2 \mathrm{GMC}_{\theta}(\mathrm{d}\vartheta).$$

We shall prove Theorem 1.2, from which Theorem 1.1 follows as a direct corollary. Another corollary is the joint convergence of the shifted coefficients $\{c_{n+k}\}_{k=1}^{\ell}$.

Corollary 1.3. Fix an $\ell \in \mathbb{N}$. Define the positive-definite Toeplitz matrix

$$\mathcal{H}_{k_1,k_2} = \frac{1}{2\pi} \int_0^{2\pi} e^{i\vartheta(k_1 - k_2)} \mathrm{GMC}_{\theta}(\mathrm{d}\vartheta), \quad 0 \le k_1, k_2 \le \ell.$$

For any $0 < \theta < 1$, we have the convergence in distribution

$$\frac{(c_{n+k}: 0 \le k \le \ell)}{\sqrt{\mathbb{E}|c_n|^2}} \xrightarrow[n \to \infty]{d} \sqrt{\mathcal{H}}(\mathcal{Z}_k: 0 \le k \le \ell),$$

where $(\mathcal{Z}_k : 0 \leq k \leq \ell)$ are *i.i.d.* standard complex normal random variables, independent of GMC_{θ} .

1.2. The circular β -ensemble

One way in which the HMC arises is through the limit of the characteristic polynomial of a random matrix. The *circular* β -ensemble is the joint density N points $\vartheta_j \in [0, 2\pi)$ for $1 \leq j \leq N$ given by

$$C\beta E_N(\vartheta_1,\ldots,\vartheta_N) \propto \prod_{1 \le k < j \le N} |e^{i\vartheta_k} - e^{i\vartheta_j}|^{\beta}$$
 (1.11)

When $\beta = 2$, i.e. $\theta = \frac{2}{\beta} = 1$, this distribution arises as the law of the eigenvalues of a Haar distributed unitary matrix and is better known as the CUE (Circular Unitary Ensemble). When $\beta \neq 2$ it also arises from the eigenvalue distribution of certain sparse random matrix models, see [KN04].

The characteristic polynomial of these points is given by

$$\chi_N(z) = \prod_{j=1}^N (1 - ze^{-i\vartheta_j}),$$
(1.12)

which would have the representation $det(1 - zU_N^*)$ if $\{e^{i\vartheta_j} : 1 \le j \le N\}$ were the eigenvalues of the matrix U_N .

The secular coefficients, defined most simply as the coefficients $c_n^{(N)}$ in the expansion of (1.12) in its Fourier basis,

$$\chi_N(z) = \sum_{n=0}^N c_n^{(N)} z^n \tag{1.13}$$

can be connected to the Fourier coefficients $\{c_n\}$. Moreover, from [CN19], there is a probability space realizing all $\{c_n^{(N)}\}$ so that

$$c_n^{(N)} \xrightarrow[N \to \infty]{a.s.} c_n \quad \text{for all } n \ge 0.$$

From [NPS23, Lemma 7.2], for any $\theta < 1$ there is a constant C_{θ}

$$\mathbb{E}|c_n - c_n^{(N)}|^2 \le C_\theta \left(\mathbb{E}|c_n|^2\right) \frac{n}{N}.$$

Thus the following corollary follows immediately.

Corollary 1.4. For any $\theta < 1$ and any $N_n \to \infty$ with $n/N_n \to 0$ as $n \to \infty$,

$$\frac{(c_{n+k}^{(N_n)}: 0 \le k \le \ell)}{\sqrt{\mathbb{E}|c_n|^2}} \xrightarrow[n \to \infty]{d} \sqrt{\mathcal{H}}(\mathcal{Z}_k: 0 \le k \le \ell).$$

Acknowledgements

The first three authors were supported by the Swedish Research Council under grant no. 2021-06594 while in residence at Institut Mittag-Leffler in Djursholm, Sweden during the fall semester of 2024. EP would like to acknowledge the support from a Canada NSERC Discovery grant. NS would like to acknowledge support from the Royal Society, grant URF\R\231028.

2. Preliminaries from [NPS23] and the Ewens sampling formula

By formula (1.6), the secular coefficients c_n can be expressed as

$$c_n = \sum_{\vec{m} \in S_n} \prod_{k=1}^n \frac{\mathcal{N}_k^{m_k}}{m_k!} \left(\frac{\theta}{k}\right)^{m_k/2}.$$
 (2.1)

The summation is over the set S_n of all compositions of n, that is $\vec{m} = (m_1, \ldots, m_n)$ is such that each m_k is a non-negative integer satisfying

$$\sum_{k=1}^{n} km_k = n. \tag{2.2}$$

From (2.1), we have that c_n is measurable with respect to $\mathscr{G}_n \coloneqq \sigma \{\mathcal{N}_1, \ldots, \mathcal{N}_n\}$. So, going forward, we will make use of the filtration $\mathscr{G} = (\mathscr{G}_n : n \in \mathbb{N})$. We also make use of the following truncated coefficients $c_{n,q}$ defined as in (2.1) but setting $\mathcal{N}_k = 0$ for k > q:

$$c_{n,q} \coloneqq \sum_{\substack{(m_k): 1 \le k \le q \\ \sum_{k=1}^{q} km_k = n}} \prod_{k=1}^{q} \frac{\theta^{m_k/2}}{m_k! k^{m_k/2}} \mathcal{N}_k^{m_k}.$$
 (2.3)

Taking the L^2 -norm of (2.1) gives

$$\mathbb{E}(|c_n|^2) = \sum_{\vec{l} \in S_n, \vec{m} \in S_n} \prod_{k=1}^n \frac{\mathbb{E}(\mathcal{N}_k^{m_k} \overline{\mathcal{N}_k}^{l_k})}{m_k! l_k!} \left(\frac{\theta}{k}\right)^{(m_k+l_k)/2}$$

$$= \sum_{\vec{m} \in S_n} \prod_{k=1}^n \frac{1}{m_k!} \left(\frac{\theta}{k}\right)^{m_k}$$
(2.4)

where we used independence of the family $\{\mathcal{N}_k\}_{k=1}^{\infty}$ and the Gaussian formula

$$\mathbb{E}(\mathcal{N}_k^{m_k}\overline{\mathcal{N}_k}^{l_k}) = \mathbb{1}_{m_k = l_k}(m_k)!.$$
(2.5)

Given a permutation σ on n symbols, we can characterize it using its cycle structure (m_1, \ldots, m_n) where m_j denotes the number of cycles in σ having length j, and $\sum_{j=1}^n jm_j = n$.

Definition 2.1. The Ewens sampling formula is a probability distribution on cycle counts $\vec{M}^{(n)} = (M_1, \ldots, M_n)$ given by

$$\mathbb{P}(\vec{M}^{(n)} = \vec{m}) = \frac{\mathbb{1}_{\sum_{k=1}^{n} km_k = n}}{\binom{n+\theta-1}{\theta-1}} \prod_{k=1}^{n} \left(\frac{\theta}{k}\right)^{m_k} \frac{1}{m_k!}.$$
(2.6)

By the normalization of (2.6) and (2.4), this gives the L^2 norm explicitly

$$\mathbb{E}(|c_n|^2) = \binom{n+\theta-1}{\theta-1}.$$
(2.7)

A straightforward generalization of (2.4) gives, for $0 \le q \le n$,

$$\mathbb{E}(c_{n-q,q_1}\overline{c_{n-q,q_2}}) = \binom{n-q+\theta-1}{\theta-1} \mathbb{P}(L^{(n-q)} \le q_1 \land q_2), \tag{2.8}$$

where $L^{(n)}$ denotes the longest cycle in a Ewens distributed permutation of length *n*. The properties of the Ewens measure are surveyed in detail in the text [ABT03]. A key result used therein is that (2.6) is the joint law of *n* independent random variables (Z_1, \ldots, Z_n) where each Z_k is Poisson distributed with parameter θ/k subject to the condition that $T_{0n} = n$ where $T_{0n} = \sum_{j=1}^{n} jZ_j$. In the following we quote the limiting distributions of the random variables T_{0n} and $L^{(n)}$.

Lemma 2.2. ([ABT03, Section 4]) Suppose that $r = r_n \in \mathbb{N}$ satisfies $r/n \to y \in (0, \infty)$ as $n \to \infty$. Then

$$\lim_{n \to \infty} n \mathbb{P}(T_{0n} = r) = p_{\theta}(y) \tag{2.9}$$

where $p_{\theta}(y)$ is an explicit probability density function with rapid tail decay $\sup_{y>n} p_{\theta}(y) \leq \theta^n/n!$.

The function $p_{\theta}(y)$ also appears in the following result [ABT03, Lemma 4.23], attributed to Kingman (1977).

Lemma 2.3. As $n \to \infty$, we have the convergence in distribution $n^{-1}L^{(n)} \xrightarrow{d} L^{(\infty)}$ where $L^{(\infty)}$ is a random variable with distribution function F_{θ} given by

$$F_{\theta}(x) = e^{\gamma_{\rm E}\theta} x^{\theta-1} \Gamma(\theta) p_{\theta}(1/x) \quad \text{for all } x \in (0,1].$$
(2.10)

We will need the following constant defined in terms of $L^{(\infty)}$.

Lemma 2.4. ([NPS23, Lemma 4.5]) Consider the quantity

$$C_{\delta} \coloneqq \theta \int_{\delta}^{1} (1-x)^{\theta-1} \mathbb{P}\left(L^{(\infty)} \le \frac{x}{1-x}\right) \frac{\mathrm{d}x}{x}$$
(2.11)

where $L^{(\infty)}$ is the limiting random variable from Lemma 2.3. Then C_{δ} can be explicitly computed as

$$C_{\delta} = 1 - \Gamma(\theta) e^{\gamma_{\rm E}\theta} \,\delta^{\theta - 1} p_{\theta}(1/\delta), \qquad (2.12)$$

where $\gamma_{\rm E}$ is the Euler-Mascheroni constant. We have

$$C_{\delta} = 1 + O(\delta), \qquad \delta \to 0.$$
 (2.13)

These facts are used in [NPS23, Lemma 4.1] to prove the following.

Lemma 2.5. Let $0 < \delta < 1$ and assume $\theta \in [0, 1]$. Define

$$\tilde{c}_{n}^{(\delta)} \coloneqq \sum_{q=\lfloor \delta n \rfloor}^{n} \mathcal{N}_{q} \sqrt{\frac{\theta}{q}} c_{n-q,q-1}$$
(2.14)

Then

$$\lim_{\delta \to 0} \lim_{n \to \infty} \frac{\mathbb{E}(|c_n - \tilde{c}_n^{(\delta)}|^2)}{\mathbb{E}(|c_n|^2)} = 0.$$
(2.15)

3. Martingale approximation and convergence

In this section we begin by finding an alternative martingale structure in the sum (2.1). This opens up the possibility of applying a central limit theorem for martingales, and we explain why this is relevant for our proof of Theorem 1.1. In contrast to [NPS23], we show that the bracket process converges in the whole L^1 phase $0 < \theta < 1$.

3.1. Martingale approximation

Note that $\tilde{c}_n^{(\delta)}$ in (2.14) is a martingale, for any fixed n, δ , when viewed as sum over q of martingale increments and with respect to the filtration $\mathscr{G} =$

 $(\sigma\{\mathcal{N}_1,\ldots,\mathcal{N}_q\}: q \in \mathbb{N})$. This follows from the fact that $c_{n-q,q-1}$ only depends on the first q-1 Gaussians. In this section, we make a further simplification of $\tilde{c}_n^{(\delta)}$. Let $\epsilon > 0$ be another fixed parameter and let

$$n_k \coloneqq (\lfloor \delta n \rfloor + k \lfloor \epsilon n \rfloor) \wedge n$$

for $k \in \mathbb{N}_0$. We let K be the smallest k such that $n_k = n$; note that for fixed δ, ϵ, K is almost fixed in that it is nearly the smallest k such that $\delta + k\epsilon \ge 1$, up to ± 1 . Then we define

$$\tilde{c}_n^{(\delta,\epsilon)} \coloneqq \sum_{k=0}^{K-1} \sum_{q=n_k+1}^{n_{k+1}} \mathcal{N}_q \sqrt{\frac{\theta}{n_k}} c_{n-q,n_k}.$$
(3.1)

Note that $\tilde{c}_n^{(\delta,\epsilon)}$ continues to satisfy the martingale property in q. Next we show further:

Lemma 3.1 (Martingale approximation). Let $0 < \delta, \epsilon < 1$ and assume $\theta \in [0, 1]$. We have

$$\lim_{\delta \to 0} \lim_{\epsilon \to 0} \lim_{n \to \infty} \frac{\mathbb{E}(|c_n - \tilde{c}_n^{(\sigma, \epsilon)}|^2)}{\mathbb{E}(|c_n|^2)} = 0.$$
(3.2)

Proof. By Lemma 2.5, it is sufficient to show the same for $\mathbb{E}(|\tilde{c}_n^{(\delta)} - \tilde{c}_n^{(\delta,\epsilon)}|^2)$. We use the martingale property to write

$$\mathbb{E}(|\tilde{c}_{n}^{(\delta)} - \tilde{c}_{n}^{(\delta,\epsilon)}|^{2}) = \theta \sum_{k=0}^{K-1} \sum_{q=n_{k}+1}^{n_{k+1}} \mathbb{E}\left(\left|c_{n-q,q-1}/\sqrt{q} - c_{n-q,n_{k}}/\sqrt{n_{k}}\right|^{2}\right).$$
 (3.3)

We expand the square and compute the three terms

$$S_1 := \theta \sum_{k=0}^{K-1} \sum_{q=n_k+1}^{n_{k+1}} \frac{1}{q} \mathbb{E}(|c_{n-q,q-1}|^2),$$
(3.4)

$$S_2 := \theta \sum_{k=0}^{K-1} \sum_{q=n_k+1}^{n_{k+1}} \frac{1}{\sqrt{q}\sqrt{n_k}} \mathbb{E}(c_{n-q,q-1}\overline{c_{n-q,n_k}}),$$
(3.5)

and

$$S_3 := \theta \sum_{k=0}^{K-1} \sum_{q=n_k+1}^{n_{k+1}} \frac{1}{n_k} \mathbb{E}(|c_{n-q,n_k}|^2).$$
(3.6)

We will prove the Lemma by showing that all three terms, when rescaled, converge to the same constant,

$$\frac{\mathcal{S}_j}{\mathbb{E}(|c_n|^2)} \to C(\delta), \qquad j = 1, 2, 3, \tag{3.7}$$

in the limit $n \to \infty$ followed by $\epsilon \to 0$, where $C(\delta)$ is the explicit constant in (2.11) and where we recall $\mathbb{E}(|c_n|^2) \sim n^{\theta-1}/\Gamma(\theta)$ is the asymptotics of the normalizing factor. Note that we already verified (3.7) for j = 1 in [NPS23, Lemma 4.5]. For j = 2, formula (2.8) gives

$$\mathbb{E}(c_{n-q,q-1}\overline{c_{n-q,n_k}}) = \binom{n-q+\theta-1}{\theta-1} \mathbb{P}(L^{(n-q)} \le n_k),$$
(3.8)

where we used that $n_k \leq q - 1$. So

$$S_2 = \theta \sum_{k=0}^{K-1} \sum_{q=n_k+1}^{n_{k+1}} \frac{1}{\sqrt{q}\sqrt{n_k}} \binom{n-q+\theta-1}{\theta-1} \mathbb{P}(L^{(n-q)} \le n_k).$$
(3.9)

We use the asymptotics of the binomial coefficient for large n-q together with Lemma 2.3 and approximate the sum over q as a Riemann integral. We justify this with dominated convergence arguments as follows. The sum over q in (3.9) is equal to

$$\frac{\sqrt{n}}{\sqrt{n_k}} n^{\theta-1} \int_{n_k/n+1/n}^{n_{k+1}/n} f_n(\lfloor nx \rfloor) \mathrm{d}x \tag{3.10}$$

where

$$f_n(\lfloor nx \rfloor) = \frac{\sqrt{n}}{\sqrt{\lfloor nx \rfloor}} n^{1-\theta} \binom{n-\lfloor nx \rfloor+\theta-1}{\theta-1} \mathbb{P}(L^{(n-\lfloor nx \rfloor)} \le n_k).$$
(3.11)

We make use of a classical bound on the ratio of two Gamma functions, see [DLMF, 5.6.4]

$$\frac{\Gamma(x+\theta)}{\Gamma(x+1)} < x^{\theta-1}, \qquad x > 0, \quad \theta \in (0,1)$$

which gives the uniform bound

$$n^{1-\theta} \binom{n-\lfloor nx \rfloor+\theta-1}{\theta-1} \le (1-x)^{\theta-1}, \quad n \in \mathbb{N}.$$
(3.12)

Then $f_n(\lfloor nx \rfloor)$ is uniformly bounded by $2(1-x)^{\theta-1}$, which is integrable for all $\theta > 0$. Furthermore, for each fixed 0 < x < 1, we have $f_n(x) \to f(x)$ as $n \to \infty$, where

$$f(x) = \frac{1}{\Gamma(\theta)} \frac{1}{\sqrt{x}} (1-x)^{\theta-1} \mathbb{P}\left(L^{(\infty)} \le \frac{\delta+k\epsilon}{1-x}\right).$$
(3.13)

Then using the definition of n_k , the dominated convergence theorem gives

$$S_2 \sim \frac{n^{\theta-1}}{\Gamma(\theta)} \theta \sum_{k=0}^{K-1} \int_{\delta+k\epsilon}^{(\delta+(k+1)\epsilon)\wedge 1} \frac{1}{\sqrt{x}\sqrt{\delta+k\epsilon}} (1-x)^{\theta-1} \mathbb{P}\left(L^{(\infty)} \leq \frac{\delta+k\epsilon}{1-x}\right) \mathrm{d}x.$$
(3.14)

Now similarly, we view the sum over k as approximating a Riemann integral. We write it as

$$\int_{0}^{\epsilon(K-1)} \mathrm{d}y \frac{1}{\epsilon} \int_{\delta+\epsilon \lfloor y/\epsilon \rfloor}^{\delta+\epsilon \lfloor y/\epsilon \rfloor + \epsilon} \mathrm{d}x \, \frac{1}{\sqrt{x}} \frac{1}{\sqrt{\delta+\epsilon \lfloor y/\epsilon \rfloor}} (1-x)_{+}^{\theta-1} \mathbb{P}\left(L^{(\infty)} \le \frac{\delta+\epsilon \lfloor y/\epsilon \rfloor}{1-x}\right)$$

where we have dropped the Riemann integral approximation terms, and where we have incorporated the $(\cdot \wedge 1)$ boundary condition into the integrand. In the x integral make the change of variables $x' = (x - \delta - \epsilon \lfloor y/\epsilon \rfloor)/\epsilon$. Then the sum over k in (3.14) is equal to

$$\int_{0}^{\epsilon(K-1)} \mathrm{d}y \int_{0}^{1} \mathrm{d}x' \frac{1}{\sqrt{\delta + \epsilon \lfloor y/\epsilon \rfloor + x'\epsilon}} \frac{1}{\sqrt{\delta + \epsilon \lfloor y/\epsilon \rfloor}} \times (1 - \delta - \epsilon \lfloor y/\epsilon \rfloor - x'\epsilon)_{+}^{\theta - 1} \mathbb{P}\left(L^{(\infty)} \le \frac{\delta + \epsilon \lfloor y/\epsilon \rfloor}{1 - \delta - \epsilon \lfloor y/\epsilon \rfloor - x'\epsilon}\right).$$
(3.15)

Note that for every fixed $\delta > 0$, the integrand above is bounded by a constant times $(1 - \delta - \epsilon \lfloor y/\epsilon \rfloor - x'\epsilon)^{\theta-1}$, which is integrable for all $\theta > 0$. Note that this is needed because at $\lfloor y/\epsilon \rfloor = (K-1)$ and x' = 1 this term is singular. Furthermore, the integrand converges pointwise as $\epsilon \to 0$ to

$$\frac{1}{\delta+y}(1-\delta-y)^{\theta-1}\mathbb{P}\left(L^{(\infty)} \le \frac{\delta+y}{1-\delta-y}\right).$$
(3.16)

Then application of the dominated convergence theorem shows that

$$S_2 \sim \frac{n^{\theta-1}}{\Gamma(\theta)} \theta \int_0^{1-\delta} \frac{1}{\delta+y} (1-\delta-y)^{\theta-1} \mathbb{P}\left(L^{(\infty)} \leq \frac{\delta+y}{1-\delta-y}\right) \mathrm{d}y.$$
(3.17)

The substitution $u = \delta + y$ shows that this integral is exactly the constant C_{δ} in (2.11). Hence (3.7) is verified for j = 2. The calculation of S_3 is similar and gives, as $n \to \infty$,

$$S_3 \sim \frac{n^{\theta-1}}{\Gamma(\theta)} \theta \sum_{k=0}^{K-1} \int_{\delta+k\epsilon}^{\delta+(k+1)\epsilon} \frac{1}{\delta+k\epsilon} (1-x)_+^{\theta-1} \mathbb{P}\left(L^{(\infty)} \leq \frac{\delta+k\epsilon}{1-x}\right) \,\mathrm{d}x, \quad (3.18)$$

for fixed $\delta, \epsilon > 0$. Then taking $\epsilon \to 0$ by a similar Riemann sum approximation, we obtain (3.7) for j = 3. This completes the proof of the Lemma.

3.2. Martingale convergence

We fix a polynomial

$$p(z) = \sum_{r=0}^{\ell} d_r z^r$$

with complex coefficients throughout this section. Define

$$\mathcal{M}_{\theta}(|p^{2}|) \coloneqq \frac{1}{2\pi} \int_{0}^{2\pi} |p(e^{i\vartheta})|^{2} \mathrm{GMC}_{\theta}(\mathrm{d}\vartheta).$$

As a consequence of Lemma 3.1, it suffices to show

$$\tilde{X}_n^{(\delta,\epsilon)}[p] \coloneqq \sum_{k=0}^{K-1} \sum_{q=n_k+1}^{n_{k+1}} \mathcal{N}_q \sqrt{\frac{\theta}{n_k}} \sum_{s=0}^{\ell} d_s c_{n-q+s,n_k}, \qquad (3.19)$$

converges in law since

$$\lim_{\delta \to 0} \lim_{\epsilon \to 0} \lim_{n \to \infty} \frac{\mathbb{E}(|\tilde{X}_n^{(\delta,\epsilon)}[p] - X_n[p]|^2)}{\mathbb{E}(|c_n|^2)} = 0,$$

Remark 3.2. Note that to properly apply Lemma 3.1 in this case, we would rather need a version in which n_k is replaced by

$$n_k(s) \coloneqq (\lfloor \delta(n+s) \rfloor + k \lfloor \epsilon(n+s) \rfloor) \land (n+s),$$

for $s \in \mathbb{N}_0$ is fixed. The details of the proof are unchanged.

As X_n is a complex martingale (note, not in the index *n* but rather viewing the sum over *q* as a sum of martingale increments), we can define its bracket process

$$\mathcal{M}_{\theta,\delta,\epsilon,n} \coloneqq \frac{1}{\mathbb{E}(|c_n|^2)} \sum_{k=0}^{K-1} \sum_{q=n_k+1}^{n_{k+1}} \frac{\theta}{n_k} \left| \sum_{s=0}^{\ell} d_s c_{n-q+s,n_k} \right|^2.$$
(3.20)

We recall from Parseval's identity

$$\sum_{n=0}^{\infty} \left| \sum_{s=0}^{\ell} d_s c_{n+s,n_k} \right|^2 r^{2n} = \sum_{s_1,s_2} d_{s_1} \overline{d_{s_2}} \sum_{n=0}^{\infty} c_{n+s_1,n_k} \overline{c_{n+s_2,n_k}} r^{2n}$$

$$= \sum_{s_1,s_2} d_{s_1} \overline{d_{s_2}} r^{-2s_2} \sum_{n=0}^{\infty} c_{n+s_1-s_2,n_k} \overline{c_{n,n_k}} r^{2n}$$

$$= \frac{1}{2\pi} \int_0^{2\pi} p(e^{i\vartheta}) \overline{p(r^{-2}e^{i\vartheta})} e^{\sqrt{\theta} G_{n_k}(re^{i\vartheta})} d\vartheta,$$
(3.21)

where

$$G_k(z) \coloneqq 2\Re\left(\sum_{\ell=1}^k \frac{z^\ell}{\sqrt{\ell}} \mathcal{N}_\ell\right). \tag{3.22}$$

We also introduce the variance of this process

$$V_k(r) \coloneqq 2\sum_{\ell=1}^k \frac{r^{2\ell}}{\ell} = \mathbb{E}(G_k(r)^2).$$
 (3.23)

Now we have the following convergence, which holds along any sequence of r_k which tend to infinity:

Lemma 3.3. Let $r_k \in [0,1)$ be any sequence with $r_k \to 1$ as $k \to \infty$. Then

$$\left(\sum_{n=0}^{\infty} \left|\sum_{s=0}^{\ell} d_s c_{n+s,k}\right|^2 r_k^{2n}\right) e^{-\theta V_k(r_k)/2} \xrightarrow{p}{k \to \infty} \mathcal{M}_{\theta}(|p^2|).$$

Proof. We introduce a two-parameter family of random variables

$$X_k(r) \coloneqq \left(\sum_{n=0}^{\infty} \left|\sum_{s=0}^{\ell} d_s c_{n+s,k}\right|^2 r^{2n}\right) e^{-\theta V_k(r)/2}, \quad k \in \mathbb{N}, r \in [0,1].$$

We also let $X_{\infty}(r)$ be the limit of $X_k(r)$ as $k \to \infty$ for fixed $r \in [0, 1)$, i.e.

$$X_{\infty}(r) \coloneqq \left(\sum_{n=0}^{\infty} \left|\sum_{s=0}^{\ell} d_s c_{n+s}\right|^2 r^{2n}\right) e^{-\theta V_{\infty}(r)/2},$$

Our goal is to show that $X_k(r_k) \xrightarrow{p} \mathcal{M}_{\theta}(|p^2|)$. Using Parseval's identity, (c.f. (3.21) and (3.22)), $X_k(r)$ can be connected to

Using Parseval's identity, (c.f. (3.21) and (3.22)), $X_k(r)$ can be connected to GMC with a somewhat non-standard two-parameter regularization (note that the error term $\mathscr{P}(r)$ is killed by the diverging variance). However, by standard theory [Ber17, Sha16, JS17b],

$$X_k(1) \xrightarrow[k \to \infty]{p} \mathcal{M}_{\theta}(|p^2|),$$
$$X_{\infty}(r) \xrightarrow[r \to 1]{p} \mathcal{M}_{\theta}(|p^2|),$$

as the first is the classical Kahane-type Dirichlet-kernel regularization and the second is a Poisson-kernel regularization. As $\theta < 1$, these convergences additionally hold in L¹.

Now, we additionally have a martingale structure. For all r < 1,

$$\mathbb{E}\left(X_{\infty}(r)|\mathscr{F}_{k}\right) = X_{k}(r).$$

Moreover, by L¹–convergence, on taking $r \to 1$ on both sides, we can exchange limit and conditional expectation to conclude that $\{X_k(1):k\}$ is a uniformlyintegrable martingale and

$$X_k(1) = \mathbb{E}\left(\mathcal{M}_{\theta}(|p^2|) | \mathscr{F}_k\right).$$

We have that

$$X_k(r_k) = \mathbb{E}\left(X_{\infty}(r_k)|\mathscr{F}_k\right),\,$$

and hence

$$\mathbb{E}|X_k(r_k) - X_k(1)| = \mathbb{E}\left|\mathbb{E}\left(X_{\infty}(r_k) - \mathcal{M}_{\theta}(|p^2|)\big|\mathscr{F}_k\right)\right| \\ \leq \mathbb{E}\left|X_{\infty}(r_k) - \mathcal{M}_{\theta}(|p^2|)\right| \xrightarrow[k \to \infty]{} 0.$$

On the other hand, from the L¹-convergence of $X_k(1) \to \mathcal{M}_{\theta}(|p^2|)$, we conclude

$$X_k(r_k) \xrightarrow[k \to \infty]{L^1} \mathcal{M}_{\theta}(|p^2|)$$

As a corollary, we derive:

Corollary 3.4. For any integer $m \ge 0$, along any sequence $n_k \to \infty$,

$$\frac{1}{n_k^{\theta}} \sum_{q} e^{-qm/n_k} \left| \sum_{s=0}^{\ell} d_s c_{q+s,n_k} \right|^2 \xrightarrow{p} \mathcal{M}_{\theta}(|p^2|) \times \left(e^{\theta \gamma_E} \int_0^{\infty} e^{-my} p_{\theta}(y) \mathrm{d}y \right),$$

with γ_E the Euler-Mascheroni constant.

Proof. Set $r_k = e^{-m/(2n_k)}$. Then we have

$$\begin{split} \frac{1}{n_k^{\theta}} &\sum_q e^{-mq/n_k} \left| \sum_{s=0}^{\ell} d_s c_{q+s,n_k} \right|^2 \\ &= \frac{1}{n_k^{\theta}} e^{\theta V_{n_k}(r_k)/2} \times \left(\sum_q r_k^{2q} \left| \sum_{s=0}^{\ell} d_s c_{q+s,n_k} \right|^2 e^{-\theta V_{n_k}(r_k)/2} \right). \end{split}$$

Thus applying Lemma 3.3, it suffices to find the limit of the constant,

$$\frac{1}{n_k^{\theta}} e^{\theta V_{n_k}(r_k)/2} = \exp\left(\theta(\gamma_E + o_{n_k}(1)) + \theta \sum_{\ell=1}^{n_k} \frac{e^{-\ell m/n_k} - 1}{\ell}\right),$$

where $\gamma_E = \lim_{n \to \infty} \left(-\log n + \sum_{\ell=1}^n \frac{1}{\ell} \right)$ is the Euler-Mascheroni constant. Recalling T_{0n} from (2.9), we have that the probability generating function of T_{0n} is given by

$$\mathbb{E}\left(z^{T_{0n}}\right) = \prod_{j=1}^{n} \mathbb{E}\left(z^{jZ_j}\right) = \prod_{j=1}^{n} e^{\frac{\theta}{j}(z^j-1)}.$$

Hence taking $z = e^{-m/n}$ and using the weak-convergence of T_{0n}/n ,

$$\exp\left(\theta \sum_{\ell=1}^{n} \frac{e^{-\ell m/n} - 1}{\ell}\right) = \mathbb{E}\left(e^{-m(T_{0n}/n)}\right) \xrightarrow[n \to \infty]{} \int_{0}^{\infty} e^{-my} p_{\theta}(y) \mathrm{d}y,$$

which completes the proof.

Now using Tauberian theory, we prove the following main result:

Lemma 3.5. For any $0 < \delta, \epsilon < 1$, there is a (deterministic) constant $A(\theta, \delta, \epsilon)$ so that

$$\mathcal{M}_{\theta,\delta,\epsilon,n} \xrightarrow{p} A(\theta,\delta,\epsilon)\mathcal{M}_{\theta}(|p^2|).$$

Moreover, we have that

$$\lim_{\delta\to 0}\lim_{\epsilon\to 0}A(\theta,\delta,\epsilon)=1.$$

Proof. This follows the proof of the in-probability version of Karamata's Tauberian theorem ([NPS23, Theorem A.2]), with minor changes, and so we give a sketch of the proof. We recall (3.20), in terms of which

$$\mathcal{M}_{\theta,\delta,\epsilon,n} = \frac{1}{\mathbb{E}(|c_n|^2)} \sum_{k=0}^{K-1} \sum_{q=n_k+1}^{n_{k+1}} \frac{\theta}{n_k} \left| \sum_{s=0}^{\ell} d_s c_{n-q+s,n_k} \right|^2$$
$$=: \frac{1}{\mathbb{E}(|c_n|^2)} \sum_{k=0}^{K-1} \theta n_k^{\theta-1} X_{\epsilon,k,n}.$$

Now we let $\tilde{h}^{(k,n)}$ be the function

$$\tilde{h}^{(k,n)}(x) \coloneqq \mathbf{1}\{e^{-(n-n_k-1)/n_k} \le x < e^{-(n-n_{k+1})/n_k}\}.$$

Then

$$X_{\epsilon,k,n} = \frac{1}{n_k^{\theta}} \sum_{q=0}^{\infty} \tilde{h}^{(k,n)}(e^{-q/n_k}) \left| \sum_{s=0}^{\ell} d_s c_{q+s,n_k} \right|^2.$$

The function $\tilde{h}^{(k,n)}(x)$ converges almost everywhere as $n \to \infty$

$$\tilde{h}^{k,n}(x) \xrightarrow[n \to \infty]{a.e.} \mathbf{1}\{\exp\left(-\frac{1-\delta-k\epsilon}{\delta+k\epsilon}\right) \le x < \exp\left(-\frac{1-\delta-k\epsilon-\epsilon}{\delta+k\epsilon}\right)\}.$$

So we define

$$\tilde{h}^{k,\pm}(x) \coloneqq \tilde{h}^{k,\pm,\eta}(x) \coloneqq \mathbf{1}\{\exp\left(-\frac{1-\delta-k\epsilon}{\delta+k\epsilon}\right) \mp \eta \le x < \exp\left(-\frac{1-\delta-k\epsilon-\epsilon}{\delta+k\epsilon}\right) \pm \eta\},$$

so that for all $\eta > 0$ and all n sufficiently large (depending on η) and all $0 \le x \le 1$

$$\tilde{h}^{k,-}(x) \le \tilde{h}^{k,n}(x) \le \tilde{h}^{k,+}(x).$$

The strategy now follows standard Tauberian theory methods. For any polynomial P on [0, 1] we have from Corollary 3.4

$$\frac{1}{n_k^{\theta}} \sum_q P(e^{-q/n_k}) \left| \sum_{s=0}^{\ell} d_s c_{q+s,n_k} \right|^2 \xrightarrow{p}{k \to \infty} \mathcal{M}_{\theta}(|p^2|) \times \left(e^{\theta \gamma_E} \int_0^{\infty} P(e^{-y}) p_{\theta}(y) \mathrm{d}y \right) \right|$$

Using a combination of piecewise linear approximation and Weierstrass approximation, we can now sandwich \tilde{h} by polynomials

$$P^{-,\eta}(x) \le \tilde{h}^{k,-}(x) \le \tilde{h}^{k,n}(x) \le \tilde{h}^{k,+}(x) \le P^{+,\eta}(x),$$

with the property that

$$\int_0^\infty \left(P^{+,\eta}(e^{-y}) - P^{-,\eta}(e^{-y}) \right) p_\theta(y) \mathrm{d}y \xrightarrow[\eta \to 0]{} 0,$$

where we have used the density of polynomials in the L¹-space on [0, 1] with probability measure $p_{\theta}(\log(1/x))\frac{dx}{x}$, the density of which vanishes at 0 by the estimate in Lemma 2.2.

We conclude taking $n \to \infty$ followed by $\eta \to 0$ that

$$X_{\epsilon,k,n} \xrightarrow[n \to \infty]{p} \mathcal{M}_{\theta}(|p^2|) \times \left(e^{\theta \gamma_E} \int_0^\infty \mathbf{1} \{ \frac{1 - \delta - k\epsilon - \epsilon}{\delta + k\epsilon} \le y < \frac{1 - \delta - k\epsilon}{\delta + k\epsilon} \} p_{\theta}(y) \mathrm{d}y \right).$$

Hence

$$\mathcal{M}_{\theta,\delta,\epsilon,n} = \sum_{k=0}^{K-1} \frac{\theta n_k^{\theta-1}}{\mathbb{E}(|c_n|^2)} X_{\epsilon,k,n}$$

$$\xrightarrow{p}{n \to \infty} \Gamma(\theta+1) \mathcal{M}_{\theta}(|p^2|) e^{\theta \gamma_E} \sum_{k=0}^{K-1} (\delta+k\epsilon)^{\theta-1} \int_{\frac{1-\delta-k\epsilon}{\delta+k\epsilon}}^{\frac{1-\delta-k\epsilon}{\delta+k\epsilon}} p_{\theta}(y) \mathrm{d}y.$$
(3.24)
(3.25)

Now make the substitution $y = \frac{1-x}{\delta+k\epsilon}$ and apply (2.10). This shows that the right-hand side of (3.25) is $\mathcal{M}_{\theta}(|p|^2) A(\theta, \delta, \epsilon)$ where

$$A(\theta, \delta, \epsilon) := \theta \sum_{k=0}^{K-1} \frac{1}{\delta + k\epsilon} \int_{\delta + k\epsilon}^{\delta + k\epsilon + \epsilon} (1-x)^{\theta - 1} \mathbb{P}\left(L^{(\infty)} \le \frac{\delta + k\epsilon}{1-x}\right) \mathrm{d}x \quad (3.26)$$

Note that this is the same constant given in (3.18). Comparing with (3.7) with j = 3, we get $A(\theta, \delta, \epsilon) \to C(\delta)$ as $\epsilon \to 0$ and from Lemma 2.4 we have $C(\delta) \to 1$ as $\delta \to 0$. We conclude that

$$\lim_{\delta \to 0} \lim_{\epsilon \to 0} A(\theta, \delta, \epsilon) = 1.$$
(3.27)

3.3. Proof of main theorem using the martingale CLT

Proof of Theorem 1.2. The details of the proof now follow in the same way as [NPS23]. We apply martingale central limit theorem of [HH80]. The Lindeberg condition of the CLT follows from computation of the sum of conditional fourth moments of the martingale increments (see [NPS23, Lemma 3.7]). Hence it follows that for any δ, ϵ

$$\frac{\tilde{X}_n^{(\delta,\epsilon)}[p]}{\sqrt{\mathbb{E}(|c_n|^2)}} \xrightarrow[n \to \infty]{d} \sqrt{A(\theta, \delta, \epsilon)} \mathcal{M}_{\theta}(|p^2|) N_{\mathbb{C}},$$

for a complex normal random variable $N_{\mathbb{C}}$ independent of the mass \mathcal{M}_{θ} . Hence, letting ϱ be a metric which metrizes weak convergence, we have

$$\lim_{\delta \to 0} \limsup_{\epsilon \to 0} \limsup_{n \to \infty} \lim_{n \to \infty} \varrho\left(\frac{\tilde{X}_n^{(\delta,\epsilon)}[p]}{\sqrt{\mathbb{E}(|c_n|^2)}}, \sqrt{\mathcal{M}_{\theta}(|p^2|)}N_{\mathbb{C}}\right) = 0.$$

On the other hand, from Lemma 3.1, we also have

$$\lim_{\delta \to 0} \limsup_{\epsilon \to 0} \limsup_{n \to \infty} \rho\left(\frac{\tilde{X}_n^{(\delta,\epsilon)}[p]}{\sqrt{\mathbb{E}(|c_n|^2)}}, \frac{X_n[p])}{\sqrt{\mathbb{E}(|c_n|^2)}}\right) = 0,$$

from which it follows

$$\frac{X_n[p]}{\sqrt{\mathbb{E}(|c_n|^2)}} \xrightarrow[n \to \infty]{d} \sqrt{\mathcal{M}_{\theta}(|p^2|)} N_{\mathbb{C}}.$$

3.4. Joint convergence of the shifted coefficients

Proof of Corollary 1.3. By the Cramer-Wold device, it suffices to consider the convergence of the characteristic function, for real constants $\{a_k\}$ and $\{b_k\}$.

$$\varphi_n \coloneqq \mathbb{E}\left(\exp\left(\frac{i}{\sqrt{\mathbb{E}(|c_n|^2)}}\sum_{k=0}^{\ell}(a_k \Re c_{n+k} + b_k \Im c_{n+k})\right)\right).$$

Now suppose $p(z) = \sum_{k=0}^{\ell} d_k z^k$ for d_k to be determined, and we look to make

$$(\Re X_n[p] + \Im X_n[p]) = \sum_{k=0}^{\ell} (a_k \Re c_{n+k} + b_k \Im c_{n+k})$$
$$= \sum_{k=0}^{\ell} \frac{a_k - ib_k}{2} c_{n+k} + \frac{a_k + ib_k}{2} \overline{c_{n+k}}$$

This is possible if we take

 $(\Re d_k + \Im d_k) = a_k$ and $(-\Im d_k + \Re d_k) = b_k \iff 2d_k = (a_k + b_k) + i(a_k - b_k).$ Thus, with this p, applying Theorem 1.2, we conclude

$$\varphi_n = \mathbb{E}\left(\exp\left(i\frac{(\Re X_n[p] + \Im X_n[p])}{\sqrt{\mathbb{E}(|c_n|^2)}}\right)\right) \to \mathbb{E}\left(\exp\left(i(\sqrt{\mathcal{M}_{\theta}(|p^2|)})(\Re \mathcal{Z} + \Im \mathcal{Z})\right)\right)$$
$$= \mathbb{E}\left(\exp\left(-\mathcal{M}_{\theta}(|p^2|)/2\right)\right) =: \varphi_{\infty}.$$

Now we have that

$$\mathcal{M}_{\theta}(|p^{2}|) = \frac{1}{2\pi} \int_{0}^{2\pi} \sum_{k_{1},k_{2}} d_{k_{1}} \overline{d_{k_{2}}} e^{i\vartheta(k_{1}-k_{2})} \mathrm{GMC}_{\theta}(\mathrm{d}\vartheta).$$

Note that

$$d_{k_1}\overline{d_{k_2}} = \frac{1}{4} \left((a_{k_1} + b_{k_1}) + i(a_{k_1} - b_{k_1}) \right) \left((a_{k_2} + b_{k_2}) - i(a_{k_2} - b_{k_2}) \right)$$
$$= \frac{1}{2} \left(a_{k_1} - ib_{k_1} \right) \left(a_{k_2} + ib_{k_2} \right).$$

Now let, as in the statement, \mathcal{H} be the Toeplitz matrix

$$\mathcal{H}_{k_1,k_2} = \frac{1}{2\pi} \int_0^{2\pi} e^{i\vartheta(k_1 - k_2)} \mathrm{GMC}_{\theta}(\mathrm{d}\vartheta), \quad 0 \le k_1, k_2 \le \ell,$$

which is positive semidefinite. Then define the random vector

$$(\widehat{c}_0, \widehat{c}_1, \dots, \widehat{c}_\ell)^\top \coloneqq \sqrt{\mathcal{H}}(\mathcal{Z}_0, \mathcal{Z}_1, \dots, \mathcal{Z}_\ell)^\top.$$

Now if we use the complex normal structure, and the computations derived above

$$\mathbb{E} \exp\left(i\sum_{k=0}^{\ell} (a_k \Re \widehat{c}_k + b_k \Im \widehat{c}_k)\right)$$

$$= \mathbb{E} \exp\left(i\sum_{k=0}^{\ell} \frac{a_k - ib_k}{2} \widehat{c}_k + \frac{a_k + ib_k}{2} \overline{\widehat{c}_k}\right)$$

$$= \mathbb{E} \exp\left(\mathbb{E}\left[\left(i\sum_{k=0}^{\ell} \frac{a_k - ib_k}{2} \widehat{c}_k + \frac{a_k + ib_k}{2} \overline{\widehat{c}_k}\right)^2 \middle| \mathrm{GMC}_{\theta}\right]\right)$$

$$= \mathbb{E} \exp\left(-\sum_{k_1, k_2}^{\ell} \frac{(a_{k_1} - ib_{k_1})(a_{k_2} + ib_{k_2})}{4} \mathcal{H}_{k_1, k_2}\right)$$

$$= \mathbb{E} \exp\left(-\mathcal{M}_{\theta}(|p^2|)/2\right).$$

This completes the proof.

References

- [ABT03] Richard Arratia, Andrew D. Barbour, and Simon Tavaré. Logarithmic Combinatorial Structures: a Probabilistic Approach. European Mathematical Society, 2003.
- [ASV⁺22] Daksh Aggarwal, Unique Subedi, William Verreault, Asif Zaman, and Chenghui Zheng. A conjectural asymptotic formula for multiplicative chaos in number theory. *Research in Number Theory*, 8(2):35, 2022.
- [Ber17] Nathanaël Berestycki. An elementary approach to Gaussian multiplicative chaos. *Electronic Communications in Probability*, 22(none):1 – 12, 2017.
- [CN19] Reda Chhaibi and Joseph Najnudel. On the circle, $GMC^{\gamma} = \lim_{\alpha \to \infty} C\beta E_n$ for $\gamma = \sqrt{\frac{2}{\beta}}$, $(\gamma \leq 1)$. arXiv:1904.00578, 2019.
- [DLMF] NIST Digital Library of Mathematical Functions. http://dlmf.nist.gov/, Release 1.0.20 of 2018-09-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller and B. V. Saunders, eds.

- [Ger22] Maxim Gerspach. Almost sure lower bounds for a model problem for multiplicative chaos in number theory. *Mathematika*, 68(4):1331– 1363, 2022.
- [GV23] Christophe Garban and Vincent Vargas. Harmonic analysis of Gaussian multiplicative chaos on the circle. *arXiv preprint arXiv:2311.04027*, 2023.
- [GW24a] Ofir Gorodetsky and Mo Dick Wong. Martingale central limit theorem for random multiplicative functions. *arXiv preprint arXiv:2405.20311*, 2024.
- [GW24b] Ofir Gorodetsky and Mo Dick Wong. A short proof of helson's conjecture. arXiv preprint arXiv:2405.19151, 2024.
- [GZ24] Haotian Gu and Zhenyuan Zhang. Universality and phase transitions in low moments of secular coefficients of critical holomorphic multiplicative chaos. arXiv preprint arXiv:2401.05681, 2024.
- [Har20] Adam J. Harper. Moments of random multiplicative functions, I: Low moments, better than squareroot cancellation, and critical multiplicative chaos. *Forum Math. Pi*, 8:e1, 95, 2020.
- [Har24] Adam J Harper. Moments of random multiplicative functions, iii: A short review. arXiv preprint arXiv:2410.11523, 2024.
- [HH80] Peter Hall and Christopher C. Heyde. Martingale Limit Theory and Its Application. Academic Press, 1980.
- [JS17a] Janne Junnila and Eero Saksman. Uniqueness of critical Gaussian chaos. *Electron. J. Probab.*, 22:Paper No. 11, 31, 2017.
- [JS17b] Janne Junnila and Eero Saksman. Uniqueness of critical Gaussian chaos. *Electronic Journal of Probability*, 22(none):1 31, 2017.
- [KN04] Rowan Killip and Irina Nenciu. Matrix models for circular ensembles. Int. Math. Res. Not., (50):2665–2701, 2004.
- [NPS23] Joseph Najnudel, Elliot Paquette, and Nick Simm. Secular coefficients and the holomorphic multiplicative chaos. The Annals of Probability, 51(4):1193–1248, 2023.
- [Rem20] Guillaume Remy. The Fyodorov-Bouchaud formula and Liouville conformal field theory. *Duke Math. J.*, 169(1):177–211, 2020.
- [Sha16] Alexander Shamov. On Gaussian multiplicative chaos. Journal of Functional Analysis, 270(9):3224–3261, 2016.
- [SZ22] Kannan Soundararajan and Asif Zaman. A model problem for multiplicative chaos in number theory. *Enseign. Math.*, 68(3-4):307–340, 2022.
- [Xu24] Max Wenqiang Xu. Better than square-root cancellation for random multiplicative functions. Transactions of the American Mathematical Society, Series B, 11(14):482–507, 2024.