DO WE REALLY NEED THE RADEMACHER COMPLEXITIES?

DANIEL BARTL AND SHAHAR MENDELSON

ABSTRACT. We study the fundamental problem of learning with respect to the squared loss in a convex class. The state-of-the-art sample complexity estimates in this setting rely on Rademacher complexities, which are generally difficult to control. We prove that, contrary to prevailing belief and under minimal assumptions, the sample complexity is not governed by the Rademacher complexities but rather by the behaviour of the limiting gaussian process. In particular, all such learning problems that have the same L_2 structure—even those with heavy-tailed distributions—share the same sample complexity. This constitutes the first universality result for general convex learning problems.

The proof is based on a novel learning procedure, and its performance is studied by combining optimal mean estimation techniques for real-valued random variables with Talagrand's generic chaining method.

1. INTRODUCTION

On the face of it, the question of whether the Rademacher complexities are truly needed seems preposterous: the Rademacher complexities have had a central role in machine learning and statistical learning theory for more than 25 years; they appear in almost every state-of-the-art sample complexity estimate, and their appearance at each instance is simply natural.

Intuitively, the Rademacher complexities capture the 'empirical oscillations' of the class of functions participating in a statistical learning problem. The idea is that if those oscillations—corresponding to the 'size' of the class—are too large, no learning procedure is able to select the right function from the class. And unfortunately, because the Rademacher complexities capture what is a rather subtle local behaviour within the class, these parameters are often difficult to handle.

At the same time, the uniform central limit theorem shows that the limits of the empirical processes used to define the Rademacher complexities as the sample size N tends to infinity, are determined by the behaviour of the limiting gaussian process, which is simply the gaussian process indexed by the underlying class of functions. But there were no indications that the limiting gaussian process had anything to do with the sample complexity of the problem (beyond special situations in which the empirical oscillations were trivially

Date: February 24, 2025.

dominated by that gaussian process). That is completely natural: unlike the limiting gaussian process, the sample complexity is a 'finite N phenomenon'. In fact, the sharpest known sample complexity estimates—given in terms of the Rademacher complexities—, tend to be much larger than their limiting gaussian versions, especially in learning problems that involve heavy tailed random variables.

The main result of this article is therefore totally unexpected: that the limiting gaussian process determines the error of an optimal learning procedure.

We design a learning procedure that exhibits finite sample errors that depend purely on the behaviour of the underlying gaussian process, even in heavy tailed problems. The (more complicated and often much larger) Rademacher complexities are not needed.

To explain what this means and how it improves the current state-of-the-art, we must first describe the setting of the learning problems we focus on.

1.1. The classical learning problem. In what follows we consider the classical learning scenario: learning with respect to the squared loss, and for an underlying class of functions that is convex and compact. The study of such learning problems has been of central importance in statistical learning theory since the very early days of the area—almost 60 years ago. We refer e.g. to the books [3, 7, 9, 13, 18, 28, 29, 30] and references therein.

A learning problem consists of a triplet: a probability space (Ω, μ) , a compact class of functions $F \subset L_2(\mu)$ and a target random variable Y; if X is distributed according to μ , the triplet is denoted by (F, X, Y). The crucial point is that the learner does not have access to all this information. Usually, all that the learner knows are the identities of Ω and of the class of functions F, but not the underlying distribution X or the target random variable Y. Still, the goal is to find a good approximation of Y in F, and the information that can be used to achieve that goal (other than the identity of F), is a sample $(X_i, Y_i)_{i=1}^N$, selected independently according to the joint distribution (X, Y).

Naturally, 'good approximation' can be understood in a variety of ways, and is dictated by the choice of a loss functional. Here, we focus on what is arguably the most natural choice of a loss — the squared loss: the loss caused by 'guessing' f(x) instead to y is $\ell(f(x) - y) = (f(x) - y)^2$.

For every $f \in F$, the risk of the function f is $\mathbb{E}(f(X) - Y)^2$, and thanks to the convexity and compactness of F, a risk minimizer in F exists and is unique. Set $f^* = \operatorname{argmin}_{f \in F} \mathbb{E}(f(X) - Y)^2$ and let

$$\mathcal{L}_f(X,Y) = (f(X) - Y)^2 - (f^*(X) - Y)^2$$

 $\mathbf{2}$

be the *excess loss* associated with f. It is standard to verify that thanks to the convexity of F,

(1.1)
$$\|f - f^*\|_{L_2}^2 \le \mathbb{E}\mathcal{L}_f;$$

this so-called *convexity condition* plays an important role in what follows.

Remark 1.1. Keep in mind that the learner has no information on \mathcal{L}_f —even on the given sample, the value $\mathcal{L}_f(X_i, Y_i)$ depends on $f^*(X_i)$, but obviously, f^* is not known to the learner.

The two main problems in this setup are estimation and prediction: how to use the sample to produce a function $\hat{f} \in F$ for which $\|\hat{f} - f^*\|_{L_2}^2$ is 'small' (estimation), or that the conditional expectation $\mathbb{E}\mathcal{L}_{\hat{f}}$ is small (prediction). In both cases, the key question is the best accuracy/confidence tradeoff that one can have: given a sample size N, to find the best accuracy ε and confidence parameter δ for which, with probability at least $1 - \delta$,

(1.2)
$$\|\widehat{f} - f^*\|_{L_2}^2 \le \varepsilon \text{ and/or } \mathbb{E}\mathcal{L}_{\widehat{f}} \le \varepsilon.$$

Naturally, identifying the *learning procedure* \hat{f} that achieves that tradeoff is essential.

An alternative and equivalent formulation of the same problem is to obtain the optimal sample complexity estimate: that is, given the desired accuracy and confidence levels, find the smallest sample size N for which (1.2) holds.

Identifying the optimal tradeoff/sample complexity in this setup has been the topic of hundreds of articles and dozens of books (see for example the references mentioned previously and [6, 11, 16, 20, 22]). And what was believed to be an (almost) complete solution in the setup studied here was established in [16] (and then extended to the non-convex case in [23]): it was shown that there is a learning procedure for which the tradeoff is governed by the behaviour of the empirical oscillations, leading to the *Rademacher complexities* of the triplet.

1.2. The Rademacher complexities. Over the years, the (localized) Rademacher averages have been the key ingredient in the analysis of learning problems. Among their roles, they are used to define the Rademacher complexities of a triplet (F, X, Y).

Let $(\varepsilon_i)_{i=1}^N$ be independent, symmetric, $\{-1, 1\}$ -valued random variables that are independent of $(X_i, Y_i)_{i=1}^N$. For r > 0, set

$$\Phi_N(r) = \mathbb{E} \sup_{u \in (F-F) \cap rD} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i u(X_i) \right|,$$

where D is the L_2 unit ball and $F - F = \{f - h : f, h \in F\}$.

In a similar fashion, let $\xi = f^*(X) - Y$, set $\xi_i = f^*(X_i) - Y_i$, and define

$$\Phi_{N,\xi}(r) = \mathbb{E} \sup_{u \in (F-F) \cap rD} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^{N} \varepsilon_i \xi_i u(X_i) \right|.$$

Definition 1.2. For $\kappa > 0$ let

(1.3)
$$r_{\mathbb{Q}}^{\mathrm{rad}}(\kappa) = \inf\left\{r > 0 : \Phi_N(r) \le \kappa \sqrt{N}r\right\}.$$

Clearly, the set F - F is star-shaped around 0: if $u \in F - F$ then for any $\lambda \in [0, 1]$, $\lambda u \in F - F$. As a result, it is straightforward to verify that $\frac{1}{r}\Phi_N(r)$ is decreasing for r > 0—because the sets $(F - F) \cap rD$ become 'richer' as r decreases. Moreover, $r_{\mathbb{Q}}^{\mathrm{rad}}(\kappa)$ is a solution to the equation $\Phi_N(r) = \kappa \sqrt{Nr}$; when $r \geq r_{\mathbb{Q}}^{\mathrm{rad}}(\kappa)$ we have that $\Phi_N(r) \leq \kappa \sqrt{Nr}$; and when $r \leq r_{\mathbb{Q}}^{\mathrm{rad}}(\kappa)$ the reverse inequality holds.

It was shown in [22] that for a suitable choice of the constant κ , $r_{\mathbb{Q}}^{\mathrm{rad}}(\kappa)$ captures the 'intrinsic complexity' of the class, and in particular, is the dominating factor in the sample complexity of 'low noise' problems. The noise level σ is defined by

$$\sigma^2 = \mathbb{E}(f^*(X) - Y)^2 = \|f^* - Y\|_{L_2}^2,$$

and the problem is considered to have *low noise* if $\sigma \leq r_{\mathbb{Q}}^{\mathrm{rad}}(\kappa)$ (for a more detailed explanation, see [6] and [22]).

As it happens, when the noise level exceeds $r_{\mathbb{Q}}^{\mathrm{rad}}(\kappa)$, the dominating factor is a different fixed point.

Definition 1.3. Set $\xi = f^*(X) - Y$, and for $\kappa > 0$ define

(1.4)
$$r_{\mathbf{M}}^{\mathrm{rad}}(\kappa) = \inf\left\{r > 0 : \Phi_{N,\xi}(r) \le \kappa \sqrt{N}r^{2}\right\}.$$

Once again, using the star-shape property of F - F, it is evident that $r_{\mathbb{M}}^{\mathrm{rad}}(\kappa)$ is given by a fixed point condition; that for $r \geq r_{\mathbb{M}}^{\mathrm{rad}}(\kappa)$ we have that $\Phi_{N,\xi}(r) \leq \kappa \sqrt{N}r^2$; and when $r \leq r_{\mathbb{M}}^{\mathrm{rad}}(\kappa)$ the reverse inequality holds.

The fixed points $r_{\mathbb{Q}}^{\mathrm{rad}}(\kappa)$ and $r_{\mathbb{M}}^{\mathrm{rad}}(\kappa)$ are the *Rademacher complexities* associated with the triplet (F, X, Y).

The sharpest bound on the accuracy/confidence tradeoff (and on the sample complexity) in our setting was established in [16], and is based on the Rademacher complexities.

The other two complexity parameters that were used in [16] are fixed points given in terms of packing numbers of the class F with respect to the L_2 norm. Denote by $\mathcal{M}(A, rD)$ the cardinality of a maximal r-separated subset of A with respect to the L_2 norm.

Definition 1.4. For $\kappa, \gamma > 0$ let

(1.5)
$$\lambda_{\mathbb{Q}}(\kappa,\gamma) = \inf\{r > 0 : \log \mathcal{M}((F-F) \cap rD,\gamma rD) \le \kappa^2 N\},\$$

and

(1.6)
$$\lambda_{\mathbb{M}}(\kappa,\gamma) = \inf\{r > 0 : \log \mathcal{M}((F-F) \cap rD,\gamma rD) \le \kappa^2 N r^2\}.$$

To formulate the current estimate from [16] we require the following:

Assumption 1.5.

- For every $f, h \in F \cup \{0\}, \|f h\|_{L_4} \le L\|f h\|_{L_2};$
- for every $f \in F$, $||f Y||_{L_4} \le L||f Y||_{L_2}$;
- $||f^* Y||_{L_2} \leq \overline{\sigma}$ for some known constant $\overline{\sigma} > 0$.

The final condition in Assumption 1.5 means that the learner has some apriori information on the noise level of the problem and the resulting accuracy/confidence tradeoff depends on $\overline{\sigma}$.

The first two components of Assumption 1.5 are that the class $F \cup \{0\}$ satisfies $L_4 - L_2$ norm equivalence with constant L, as does the class $\{f - Y : f \in F\}$; neither assumption is really restrictive.

Example 1.6. There are many scenarios in which $X \in \mathbb{R}^d$ is a centred random vector and satisfies that for every $z \in \mathbb{R}^d$, $\|\langle X, z \rangle \|_{L_4} \leq L \|\langle X, z \rangle \|_{L_2}$; therefore, every class of linear functionals of the random vector satisfies the first part of Assumption 1.5. Here we present two such examples. Let x be a mean-zero, variance 1 random variable, for which $\mathbb{E}x^4 \leq cL^4$ for a well-chosen absolute constant c. The random vector $X = (x_1, ..., x_d)$ that has independent copies of x as coordinates satisfies $L_4 - L_2$ norm equivalence with constant L.

At the other, light-tailed end of the spectrum, one can show that for any log-concave random vector, every $p \ge 2$ and every $z \in \mathbb{R}^d$,

(1.7)
$$\|\langle X, z \rangle\|_{L_p} \le Lp \|\langle X, z \rangle\|_{L_2}$$

for L that is an absolute constant. Indeed, (1.7) follows from Borell's inequality (see, e.g. [4]). Recalling that for a random variable x and $\alpha \in [1, 2]$,

$$\|x\|_{\psi_{\alpha}} = \sup_{p \ge 2} \frac{\|x\|_{L_p}}{p^{1/\alpha}},$$

(1.7) means that a log-concave random vector satisfies a $\psi_1 - L_2$ norm equivalence for an absolute constant L.

The following theorem is the main result in [16] and is the current 'gold standard' estimate on learning in convex classes with respect to the squared loss.

Theorem 1.7. There are constants c, c_0, c_1 and c_2 that depend only on L for which the following holds. Let

(1.8)
$$r_{\mathrm{rad}}^* = \max\left\{\lambda_{\mathbb{Q}}(c_1, c_2), \lambda_{\mathbb{M}}(c_1/\overline{\sigma}, c_2), r_{\mathbb{Q}}^{\mathrm{rad}}(c_1), r_{\mathbb{M}}^{\mathrm{rad}}(c_1)\right\} ,$$

and fix $r \geq 2r_{rad}^*$. There exists a procedure that, based on the data $\mathcal{D}_N = (X_i, Y_i)_{i=1}^N$ and the values of $L, \overline{\sigma}$ and r, selects a function $\widehat{f} \in F$ which satisfies that with probability at least

$$1 - \exp\left(-c_0 N \min\left\{1, \frac{r^2}{\overline{\sigma}^2}\right\}\right),$$
$$|\widehat{f} - f^*||_{L_2} \le cr \quad \text{and} \quad \mathbb{E}\left(\mathcal{L}_{\widehat{f}} \mid \mathcal{D}_N\right) \le cr^2$$

An alternative formulation of Theorem 1.7 (that can be found in [23]) is the corresponding sample complexity estimate; it uses the fact that $r_{\rm rad}^*$ decreases in N.

Corollary 1.8. Given ε and δ , let $N_0(\varepsilon)$ be the smallest integer for which $r_{\rm rad}^* \leq c_3 \sqrt{\varepsilon}$. If

$$N \ge N_0(\varepsilon) + c_4\left(\frac{\overline{\sigma}^2}{\varepsilon} + 1\right)\log\left(\frac{2}{\delta}\right),$$

then with probability at least $1 - \delta$,

$$\|\widehat{f} - f^*\|_{L_2} \le \sqrt{\varepsilon} \text{ and } \mathbb{E}\left(\mathcal{L}_{\widehat{f}} \mid \mathcal{D}_N\right) \le \varepsilon.$$

Remark 1.9. It is important to stress that the most natural learning procedure, *empirical* risk minimization (ERM)—that selects a function in F that minimizes the empirical risk $\frac{1}{N}\sum_{i=1}^{N}(f(X_i)-Y_i)^2$ —, is far from optimal. In heavy tailed situations its performance does not come close to that of the tournament procedure introduced in the proof of Theorem 1.7.

Example 1.10 (Linear regression in \mathbb{R}^d). Let $T \subset \mathbb{R}^d$ be convex (and compact) and set $F_T = \{\langle \cdot, t \rangle : t \in T\}$ to be the class of linear functionals associated with T. Let X be a random vector that satisfies $L_4 - L_2$ norm equivalence, and set w to be a centred random variable that is independent of X and for which $||w||_{L_4} \leq L||w||_{L_2}$. Let $z_0 \in \mathbb{R}^d$ and put $Y = \langle X, z_0 \rangle + w$.

It is standard to verify that the triplet (F, X, Y) satisfies the first two components in Assumption 1.5 with a constant that depends only on L. As a result, Theorem 1.7 holds, and the Rademacher complexities are the fixed points associated with the functions $\Phi_N(r)$ and $\Phi_{N,\xi}(r)$.

Thanks to the linear structure this problem has, the Rademacher complexities have a (seemingly) simpler description. Note that T - T is a convex, centrally symmetric set, and therefore so is $(T - T) \cap rD$. As such, $(T - T) \cap rD$ is the unit ball of a norm on \mathbb{R}^d (or on

a subspace of \mathbb{R}^d), and denote that norm by $\|\cdot\|_{K_r}$. Thus, the Rademacher complexities are determined by the behaviour of the functions

(1.9)
$$\Phi_N(r) = \mathbb{E} \left\| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i X_i \right\|_{K_r} \text{ and } \Phi_{N,\xi}(r) = \mathbb{E} \left\| \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i \xi_i X_i \right\|_{K_r}$$

Unfortunately, controlling the expected norms of the two random vectors in terms of the geometry of the set K_r is highly nontrivial and far from understood—other than in rather special situations. These are far more complex objects than, say, $\mathbb{E}||G||_{K_r}$, where G is the gaussian random vector in \mathbb{R}^d that has the same covariance as X. Indeed, while there are sophisticated methods that lead to a complete characterization of $\mathbb{E}||G||_{K_r}$ purely in terms of the geometry of the set K_r (see, for example [27]), there are no known characterizations for (1.9). At the same time, it follows from the central limit theorem that $\Phi_N(r) \to \mathbb{E}||G||_{K_r}$ as $N \to \infty$.

1.3. If not the Rademacher complexities, then what? The one obvious, but at the same time highly unlikely candidate that one might consider as a potential replacement for the Rademacher complexities are the fixed points associated with the limiting gaussian process. In other words, instead of using parameters that capture the oscillations of empirical/multiplier processes, one may consider the behaviour of the oscillations of the gaussian process indexed by F. Of course, at this point there is no hint that these oscillations have something to do with the learning problem, and the idea of considering them as an option is speculative at best.

Definition 1.11. Let $\sigma^2 = \inf_{f \in F} \mathbb{E}(f(X) - Y)^2$, set $\{G_h : h \in F - F\}$ to be the gaussian process indexed by F - F whose covariance is endowed by $L_2(\mu)$, and put

(1.10)
$$r_{\mathbb{Q}}(\kappa) = \inf\left\{r > 0 : \mathbb{E}\sup_{h \in (F-F)\cap rD} G_h \le \kappa \sqrt{N}r\right\},$$
$$r_{\mathbb{M}}(\kappa) = \inf\left\{r > 0 : \mathbb{E}\sup_{h \in (F-F)\cap rD} G_h \le \kappa \sqrt{N}\frac{r^2}{\sigma}\right\}.$$

Using that F - F is star-shaped around 0, it is evident that if $r \ge r_{\mathbb{Q}}(\kappa)$ then

$$\mathbb{E} \sup_{h \in (F-F) \cap rD} G_h \le \kappa \sqrt{N}r$$

and if $r \leq r_{\mathbb{Q}}(\kappa)$ then the reverse inequality holds. A similar phenomenon is true for $r_{\mathbb{M}}(\kappa)$.

With the two new parameters, the notions of 'low noise' and 'high noise' have to be adjusted. The meaning of low noise is that $\sigma \leq r_{\mathbb{Q}}(\kappa)$, while high noise means that $\sigma \geq r_{\mathbb{Q}}(\kappa)$ —which implies that $\sigma \geq r_{\mathbb{M}}(\kappa)$ as well. Indeed, if $\sigma \geq r_{\mathbb{Q}}(\kappa)$ then

$$\mathbb{E} \sup_{h \in (F-F) \cap \sigma D} G_h \le \kappa \sqrt{N} \sigma = \kappa \sqrt{N} \frac{\sigma^2}{\sigma},$$

and therefore $r = \sigma$ is an 'eligible candidate' in the definition of $r_{\mathbb{M}}(\kappa)$.

A conjecture that for a well chosen κ ,

$$r^*(\kappa) = \max\{r_{\mathbb{O}}(\kappa), r_{\mathbb{M}}(\kappa)\}$$

might bound the optimal tradeoff / sample complexity seems widely optimistic. Firstly, the empirical oscillation functions $\Phi_N(r)$ and $\Phi_{N,\xi}(r)$ tend to be much bigger than their gaussian counterpart—especially in heavy tailed situation (see Appendix B for an example). Secondly, a bound that is based solely on the behaviour of the limiting gaussian process does not 'see' all the fluctuations caused by the random sample. Such a bound would be the same for all distributions that endow the same L_2 structure, and would coincide with the light tailed estimate even if the problem is heavy tailed.

Taking all that into account, such a conjecture seems be too good to be true.

Despite that, our main result, presented in the next section, is that essentially, $\max\{r_{\mathbb{Q}}(\kappa), r_{\mathbb{M}}(\kappa)\}\$ is an upper bound on the optimal tradeoff even in heavy tailed problems, once we assume that the learner is given a little more information on the problem.

1.4. The main result. The 'little more information' that one requires is an *isomorphic* distance oracle on $F \cup \{0\}$.

Assumption 1.12. The learner has access to a symmetric¹ function d that satisfies, for every $f, h \in F \cup \{0\}$,

$$\frac{1}{\eta} \|f - h\|_{L_2} \le d(f, h) \le \eta \|f - h\|_{L_2},$$

for some $\eta \geq 1$.

Assumption 1.12 implies that the learner has access to *crude* information on L_2 distances between functions in F, as well as their L_2 norms. Crucially, the leaner does not have more information on Y or on functions that depend on f - Y, nor is there a need for accurate information on X. A situation in which Assumption 1.12 is natural is presented in Example 1.18.

Remark 1.13. As we explain in what follows, thanks to the distance oracle and to the majorizing measures theorem, the learner can identify $\mathbb{E}\sup_{h\in(F-F)\cap rD} G_h$ for every r > 0 (up to multiplicative constants that depend on η). As a result, the learner knows the value of $r_{\mathbb{Q}}(\kappa)$ —again, up to multiplicative constants in η . The situation regarding $r_{\mathbb{M}}(\kappa)$ is more subtle, as its definition is based on the unknown noise level σ . Thus, a preliminary step in the learning procedure that will be used here, is obtaining a rough estimate on the noise level.

¹That is, d(f,h) = d(h,f).

We also assume that the first two parts of Assumption 1.5 hold: that $||f - h||_{L_4} \leq L||f - h||_{L_2}$ for every $f, h \in F \cup \{0\}$, and that $||f - Y||_{L_4} \leq L||f - Y||_{L_2}$ for every $f \in F$.

The first ingredient we need is an 'isomorphic' estimate on the noise level $\sigma^2 = \mathbb{E}(f^*(X) - Y)^2$.

Lemma 1.14. There are constants c_1, c_2, c_3 that depend only on η and L such that the following holds. If r satisfies that $\log \mathcal{M}(F, c_1 r D) \leq c_2 N$, then there is a procedure that, upon receiving the data $\mathcal{D}_N = (X_i, Y_i)_{i=1}^N$, the distance oracle d, and the values of L, η, r , selects $\hat{\sigma}$ which satisfies that with probability at least $1 - 2 \exp(-c_3 N)$,

$$\widehat{\sigma}^2 \leq 2 \max\{\sigma^2, r^2\}, \text{ and if } \sigma > r, \text{ then } \widehat{\sigma}^2 \geq \frac{1}{2}\sigma^2.$$

Remark 1.15. We will show in what follows that the role of the entropic condition $\log \mathcal{M}(F, c_1 r D) \leq c_2 N$ is minimal—see Remark 1.19.

Lemma 1.14 gives a *data dependent* way of controlling r^* : it is straightforward to verify that

$$\min\left\{1,\frac{r}{\sigma}\right\} \sim \min\left\{1,\frac{r}{\widehat{\sigma}}\right\},\,$$

and thus,

$$\widehat{r}^*(\kappa) = \inf\left\{r > 0 : \mathbb{E}\sup_{h \in (F-F)\cap rD} G_h \le \kappa \sqrt{N}r \min\left\{1, \frac{r}{\widehat{\sigma}}\right\}\right\}$$

is equivalent to $r^* = \max\{r_{\mathbb{Q}}, r_{\mathbb{M}}\}.$

The last ingredient we need is an entropic fixed point. Let

(1.11)
$$\lambda^*(\kappa) = \inf\left\{r > 0 : \log \mathcal{M}(F, rD) \le \kappa N \min\left\{1, \frac{r^2}{\sigma^2}\right\}\right\}.$$

Again, replacing σ by $\hat{\sigma}$ in the definition of $\lambda^*(\kappa)$ results in an equivalent condition. Also note that the condition that $\log \mathcal{M}(F, c_1 r D) \leq c_2 N$ appearing in Lemma 1.14 is satisfied if $r > \frac{1}{c_1} \lambda^*(c_2)$.

With all the ingredients set in place, let us formulate our main result.

Theorem 1.16. There are constants c_0, c_1, c_2 that depend only on L and η for which the following holds. If $r > c_0 \max\{r^*(c_1), \lambda^*(c_1)\}$, then there exists a procedure that, based on the data $\mathcal{D}_N = (X_i, Y_i)_{i=1}^N$, the distance oracle d, and the values of L, η and r, selects a function $\hat{f} \in F$ which satisfies that with probability at least

$$1 - 2 \exp\left(-c_2 N \min\left\{1, \frac{r^2}{\sigma^2}\right\}\right),$$
$$\|\widehat{f} - f^*\|_{L_2} \le r \quad and \quad \mathbb{E}\left(\mathcal{L}_{\widehat{f}} \mid \mathcal{D}_N\right) \le r^2.$$

Thanks to Theorem 1.16, it is straightforward to derive the corresponding sample complexity bounds:

Corollary 1.17. There are constants c_1, c_2, c_3 that depend only on L and η for which the following holds. Given ε and δ , let $N_0(\varepsilon)$ be the smallest integer for which

$$\max\{r^*(c_1), \lambda^*(c_1)\} \le c_2\sqrt{\varepsilon}.$$

If

(1.12)
$$N \ge N_0(\varepsilon) + c_3 \left(\frac{\sigma^2}{\varepsilon} + 1\right) \log\left(\frac{2}{\delta}\right),$$

then with probability at least $1 - \delta$,

$$\|\widehat{f} - f^*\|_{L_2} \le \sqrt{\varepsilon} \text{ and } \mathbb{E}(\mathcal{L}_{\widehat{f}} | \mathcal{D}_N) \le \varepsilon.$$

And as in the fixed point conditions before, σ in (1.12) may be replaced with $\hat{\sigma}$.

Example 1.18 (Linear regression in \mathbb{R}^d , revised). Just as in Example 1.10 and using its notation, set $X \in \mathbb{R}^d$ to be the random vector, let $T \subset \mathbb{R}^d$ to be a convex and compact set, put $F = \{\langle \cdot, t \rangle : t \in T\}$ and consider $Y = \langle X, z_0 \rangle + w$. As noted in Example 1.10, the triplet (F, X, Y) satisfies the required norm equivalence assumptions.

Assumption 1.12 is satisfied if there is some additional crude information on the covariance matrix of X (denoted by Σ_X in what follows). For example, it suffices that the learner has access to a positive semi-definite matrix A for which

$$\eta^{-2}\Sigma_X \preceq A \preceq \eta^2 \Sigma_X,$$

and in that case one may set

$$d^{2}(u,v) = \langle A(u-v), u-v \rangle.$$

There are many natural linear regression problems where the covariance matrix Σ_X is known, and then Assumption 1.12 is trivially satisfied with $\eta = 1$.

Remark 1.19. The entropy condition that $r \geq \lambda^*(\kappa)$ should not be alarming. Indeed, set $d_F = \operatorname{diam}(F, L_2)$, and consider the high noise regime—when $\sigma \geq r_{\mathbb{Q}}(\kappa)$. As noted previously, in that case, $\sigma \geq r_{\mathbb{M}}(\kappa)$, and therefore, $\sigma \geq r^*(\kappa)$. Clearly, if $r^*(\kappa) \geq 2d_F$, then $\lambda^*(\kappa) \leq r^*(\kappa)$ because $\mathcal{M}(F, 2d_F) = 1$; and otherwise one can show that if σ is a little larger than r^* , for example, if $r^*(c\kappa)\sqrt{\log(\frac{4d_F}{r^*(c\kappa)})} \leq \sigma$, then

$$\lambda^*(\kappa) \le r^*(c\kappa) \sqrt{\log\left(\frac{4d_F}{r^*(c\kappa)}\right)}.$$

Moreover, for any σ , we have that $\lambda^*(\kappa) \leq \tilde{r}^*(c\kappa)$, where

$$\widetilde{r}^*(\kappa) = \inf\left\{r > 0: \mathbb{E}\sup_{h \in (F-F) \cap rD} G_h \le \kappa \sqrt{N}\theta(r)r\min\left\{1, \frac{r}{\sigma}\right\}\right\}$$

and $\theta(r) = 1/\max\{1, \sqrt{\log(2d_F/r)}\}$. The proof of both facts can be found in Appendix A.

It should be stressed that the procedure we use in the proof of Theorem 1.16 is not intuitive, nor is it feasible from a computational perspective. Still, the fact that such a procedure even exists—and that the accuracy/confidence tradeoff can be controlled using the limiting gaussian process even in heavy tailed situations—, is a complete surprise.

The proof of Theorem 1.16 is based on an unorthodox chaining argument, combined with an arbitrary optimal mean estimation procedure for real-valued random variables. We believe that this type of argument will have far reaching implications (see for example, the results in [5]). The unorthodox chaining argument is described in Section 3.

Remarks and notation. The procedures we use are functions Ψ that are given the values $(X_i, Y_i)_{i=1}^N$ as data. In addition, they have access to functions in F (or in $F - F = \{f - h : f, h \in F\}$), and therefore can 'see' the values $(f(X_i))_{i=1}^N$ as well. At times, the procedures also have access to the wanted confidence parameter δ and to the distance oracle d. However, the procedures never know the identities of the target function Y or of the risk minimizer f^* .

In what follows and with a slight abuse of notation, if we write, say, for $f \in F$ and $\xi = h(X) - Y$, " $\Psi(f,\xi)$ " it means that Ψ "sees" the values $(f(X_i))_{i=1}^N$, $(h(X_i))_{i=1}^N$, and $(h(X_i) - Y_i)_{i=1}^N$. It also knows the identities of f and h, but not that of Y or of f^* . All that is possible given the data the learner has access to.

Throughout this article, c, c_0, c_1 denote strictly positive absolute constants that may change their value in each appearance. If a constant c depends on a parameter α , that is denoted by $c = c(\alpha)$. We write $\alpha \leq \beta$ if there is an absolute constant c for which $\alpha \leq c\beta$ and $\alpha \sim \beta$ if both $\alpha \leq \beta$ and $\beta \leq \alpha$.

With another minor abuse of notation we write f instead of f(X) and f - Y instead of f(X) - Y. Also, we may assume without loss of generality that $0 \in F$. If not, fix an arbitrary $f_0 \in F$ and consider the shifted class $F' = \{f - f_0 : f \in F\}$ and the shifted target $Y - f_0$; the given data will now be $(X_i, Y_i - f_0(X_i))_{i=1}^N$. Clearly, F - F = F' - F', and $\sup_{f' \in F'} ||f'||_{L_2} \sim \operatorname{diam}(F, L_2)$.

DANIEL BARTL AND SHAHAR MENDELSON

2. Proof of Theorem 1.16

The learning procedure used in the proof of Theorem 1.16 has several components that are described here. Each one of the components relies on the theorem's assumptions.

The first part of the procedure is a pre-processing step that is not random. It will be presented in Section 3.1.

The two main components of the procedure are a *crude risk oracle*, denoted by $\Psi_{\mathbb{C}}$, and a *fine risk oracle*, denoted by $\Psi_{\mathcal{L}}$. Let $(X_i, Y_i)_{i=1}^{2N}$ be the given sample and set $r \geq c_0 \max\{r^*(c_1), \lambda^*(c_1)\}$ for constants c_0 and c_1 that depend only on L and η and are specified in what follows.

The crude risk oracle receives as input the first half of the sample, $(X_i, Y_i)_{i=1}^N$ and the parameter r, and for every $f \in F$ returns $\Psi_{\mathbb{C}}(f)$. Set

$$\widehat{\sigma}^2 = \inf_{f \in F} \Psi_{\mathbb{C}}(f), \quad \sigma_* = \max{\{\widehat{\sigma}, r\}},$$

and define

$$\widehat{V} = \left\{ f \in F : \Psi_{\mathbb{C}}(f) \le 4\sigma_*^2 \right\}.$$

Creating the set \hat{V} is the crude oracle's main task. The idea behind \hat{V} is to exclude functions in F that have an unreasonably large risk, while at the same time ensuring that $f^* \in \hat{V}$.

The second component—the fine risk oracle, receives as input the second half of the sample $(X_i, Y_i)_{i=N+1}^{2N}$, the set \hat{V} , and the values r and σ_* . For every ordered pair $(f, h) \in \hat{V} \times \hat{V}$ it returns $\Psi_{\mathcal{L}}(f, h)$. Based on those values, a tournament is played between functions in \hat{V} , in a way that will be clarified.

The procedure selects any winner of this tournament.

Let us describe the performance of the two components.

2.1. The crude risk oracle. The first component of the learning procedure is a crude risk oracle. While its performance is far from the wanted one, it allows the learner to exclude functions that have large risks.

Theorem 2.1. There are constants c_1, c_2, c_3 that depend only on L and η such that the following holds. Set r that satisfies $\log \mathcal{M}(F, c_1 r D) \leq c_2 N$. Then with probability at least

$$1-2\exp\left(-c_3N\right),$$

for every $f \in F$,

(2.1)
$$\left| \Psi_{\mathbb{C}}(f) - \mathbb{E}(f(X) - Y)^2 \right| \leq \frac{1}{2} \max\left\{ r^2, \mathbb{E}(f(X) - Y)^2 \right\}.$$

Note that the entropy condition $\log \mathcal{M}(F, c_1 r D) \leq c_2 N$ in the theorem is satisfied when $r \geq \frac{1}{c_1} \lambda^*(c_2)$, which we may assume in the context of Theorem 1.16.

The function $\Psi_{\mathbb{C}}$ is a mean estimation procedure for functions in the class $\{(f - Y)^2 : f \in F\}$, but as a mean estimator it is rather inaccurate: if $\mathbb{E}(f - Y)^2 \ge r^2$ then $\Psi_{\mathbb{C}}(f)$ is an 'isomorphic' guess of the mean, satisfying that

$$\frac{1}{2}\mathbb{E}(f-Y)^2 \le \Phi_{\mathbb{C}}(f) \le \frac{3}{2}\mathbb{E}(f-Y)^2.$$

And if $\mathbb{E}(f-Y)^2 < r^2$ then $\Psi_{\mathbb{C}}(f)$ only exhibits that fact without providing a better guess of what $\mathbb{E}(f-Y)^2$ really is.

The proof of Theorem 2.1 is presented in Section 4.

With Theorem 2.1 at hand, we can immediately obtain the isomorphic estimator for the noise level stated in Lemma 1.14.

Proof of Lemma 1.14. : Recall that $\sigma^2 = \inf_{f \in F} \mathbb{E}(f - Y)^2$ and that

$$\widehat{\sigma}^2 = \inf_{f \in F} \Psi_{\mathbb{C}}(f).$$

Note that in the high probability event in which (2.1) holds, $\hat{\sigma}^2 \leq 2 \max\{r^2, \sigma^2\}$, and if $\sigma > r$ then $\hat{\sigma}^2 \geq \frac{1}{2}\sigma^2$. Hence, if $\sigma > r$ then $\frac{1}{2}\sigma^2 \leq \hat{\sigma}^2 \leq 2\sigma^2$, and if $\sigma < r$ then $\hat{\sigma}^2 \leq 2r^2$. \Box

In particular, for $\sigma_* = \max\{r, \hat{\sigma}\}$ it is evident that

$$\min\left\{1, \frac{r^2}{\sigma_*^2}\right\} \sim \min\left\{1, \frac{r^2}{\sigma^2}\right\}.$$

Recall that

$$\widehat{V} = \left\{ f \in F : \Psi_{\mathbb{C}}(f) \le 4\sigma_*^2 \right\}$$

is an output of the crude risk oracle. We will show the following:

Corollary 2.2. In the event in which (2.1) holds, $f^* \in \widehat{V}$ and for every $f \in \widehat{V}$,

$$\mathbb{E}(f(X) - Y)^2 \le 16 \max\left\{r^2, \sigma^2\right\}.$$

The proof of Corollary 2.2 follows from a straightforward application of Theorem 2.1, and is outlined in Section 4.

2.2. The fine risk oracle. The second main component is an accurate mean estimation procedure for functions of the form $(f(X) - Y)^2 - (h(X) - Y)^2$.

Theorem 2.3. There are constants c_1 and c_2 that depend only on L and η for which the following holds. For $r > \max\{r^*(c_1), \lambda^*(c_1)\}$, there is a mean estimation procedure $\Psi_{\mathcal{L}}$ for which the following hold. With probability at least

$$1 - 2\exp\left(-c_2N\min\left\{1,\frac{r^2}{\sigma^2}\right\}\right),\,$$

for every $f, h \in F$ that satisfy

(2.2)
$$\mathbb{E}(h(X) - Y)^2 \le 16 \max\{r^2, \sigma^2\}$$

we have that

(2.3)
$$\left|\Psi_{\mathcal{L}}(f,h) - \left(\mathbb{E}(f-Y)^2 - \mathbb{E}(h-Y)^2\right)\right| \leq \frac{1}{2} \max\left\{r^2, \|f-h\|_{L_2}^2\right\}.$$

The proof of Theorem 2.3 is presented in Section 5.

With the main components set in place, the learning procedure is a tournament played between functions in F—following a similar path to the one used in [16] and [23].

2.3. The tournament. Recall that the symmetric distance oracle d satisfies that for every $f, h \in F \cup \{0\}$,

(2.4)
$$\eta^{-1} \|f - h\|_{L_2} \le d(f, h) \le \eta \|f - h\|_{L_2}.$$

The tournament consists of 'matches' between functions in F. Every match has a 'home function' and a 'visitor function' and the result of a match is that one of the sides wins.

- Only functions that belong to \hat{V} participate in the tournament.
- h wins its home match against f if

$$\Psi_{\mathcal{L}}(f,h) \ge \begin{cases} 0 & \text{if } d(f,h) \ge \eta r \\ -\frac{\eta^4}{2}r^2 & \text{otherwise.} \end{cases}$$

- Let V^* be the set of functions in \hat{V} that won all their home matches.
- The procedure selects any function in V^* .

To prove Theorem 1.16, it suffices to show that with the wanted probability, $V^* \neq \emptyset$ and that any $h \in V^*$ satisfies

$$||h - f^*||_{L_2} \le c(\eta)r$$
 and $\mathbb{E}(h - Y)^2 \le \mathbb{E}(f^* - Y)^2 + (c(\eta)r)^2$.

Proof of Theorem 1.16. Fix a realization of $(X_i, Y_i)_{i=1}^{2N}$ for which the assertions of both Theorem 2.1 and Theorem 2.3 hold—using the sample as described previously. By Corollary 2.2, every function $h \in \hat{V}$ satisfies (2.2), and it follows from (2.3) that for every

$$f, h \in \widehat{V},$$

(2.5)
$$\left|\Psi_{\mathcal{L}}(f,h) - \left(\mathbb{E}(f-Y)^2 - \mathbb{E}(h-Y)^2\right)\right| \leq \frac{1}{2} \max\left\{r^2, \|f-h\|_{L_2}^2\right\}.$$

For every $f, h \in F$, set

$$\mathcal{L}(f,h) = \mathbb{E}(f-Y)^2 - \mathbb{E}(h-Y)^2,$$

and note that $\mathcal{L}(f, f^*) = \mathbb{E}\mathcal{L}_f$. Thus, by the convexity of F,

(2.6)
$$\mathcal{L}(f, f^*) \ge \|f - f^*\|_{L_2}^2 \quad \text{for every } f \in F.$$

To prove that $f^* \in V^*$, recall that by Corollary 2.2, $f^* \in \widehat{V}$; therefore f^* participates in the tournament. Set $f \in \widehat{V}$ and consider the home match of f^* against f. If $d(f, f^*) \ge \eta r$, then (2.4) implies that $\|f - f^*\|_{L_2} \ge r$, and by (2.5) and the convexity condition (2.6),

$$\Psi_{\mathcal{L}}(f, f^*) \ge \mathcal{L}(f, f^*) - \frac{1}{2} \|f - f^*\|_{L_2}^2 \ge 0$$

hence, f^* wins its home match against f.

Otherwise, if $d(f, f^*) < \eta r$, then $||f - f^*||_{L_2} < \eta^2 r$ by (2.4). It follows from (2.5) that

$$\Psi_{\mathcal{L}}(f, f^*) \ge \mathcal{L}(f, f^*) - \frac{1}{2} \|f - f^*\|_{L_2}^2 \ge -\frac{\eta^4}{2} r^2,$$

and again f^* wins its home match against f.

Now set $h \in V^*$ to be any other winner of the tournament. In particular h won its home match against f^* . The first observation is that necessarily $d(f^*, h) < \eta r$. Indeed, if not and $d(f^*, h) \ge \eta r$, then $\Psi_{\mathcal{L}}(f^*, h) \ge 0$. At the same time, by (2.4), $||f^* - h||_{L_2} \ge r$, and thanks to (2.5) and (2.6),

$$\begin{split} \Psi_{\mathcal{L}}(f^*,h) &\leq \mathcal{L}(f^*,h) + \frac{1}{2} \|f^* - h\|_{L_2}^2 \\ &= -\mathcal{L}(h,f^*) + \frac{1}{2} \|f^* - h\|_{L_2}^2 \\ &\leq -\|f^* - h\|_{L_2} + \frac{1}{2} \|f^* - h\|_{L_2} < 0, \end{split}$$

which is impossible. Thus $d(f^*, h) < \eta r$, and in particular $||f^* - h||_{L_2} < \eta^2 r$. Now, since h won its home match against f^* , $\Psi_{\mathcal{L}}(f^*, h) \ge -\frac{1}{2}\eta^4 r^2$. It follows from (2.5) that

$$\mathcal{L}(f^*,h) \ge \Psi_{\mathcal{L}}(f^*,h) - \frac{1}{2} \max\left\{r^2, \|f^* - h\|_{L_2}^2\right\}$$
$$\ge -\eta^4 r^2,$$

implying that $\mathbb{E}(h-Y)^2 \leq \mathbb{E}(f^*-Y)^2 + \eta^4 r^2$.

It is clear that at the heart of the proof of Theorem 1.16 are the two functions $\Psi_{\mathbb{C}}$ and $\Psi_{\mathcal{L}}$. The rest of this article is devoted to their construction, which is based on a surprising combination: optimal mean estimation procedures for real valued random variables and Talagrand's generic chaining mechanism. We expect that the impact of this combination will go well beyond the proof of Theorem 1.16.

3. UNORTHODOX CHAINING

Generic Chaining was introduced by M. Talagrand for the study of certain stochastic processes. Let $\{Z_v : v \in V\}$ be a (centred) random process, and consider the problem of controlling $\mathbb{E} \sup_{v \in V} Z_v$. By creating increasingly fine approximating sets $V_s \subset V$ and setting $\pi_s v$ to be the best approximation (in some appropriate sense) of v in V_s , it follows that

(3.1)
$$Z_{v} - Z_{\pi_{0}v} = \sum_{s \ge 0} (Z_{\pi_{s+1}v} - Z_{\pi_{s}v}).$$

If $|V_0| = 1$ then $\mathbb{E} \sup_{v \in V} Z_v = \mathbb{E} \sup_{v \in V} (Z_v - Z_{\pi_0 v})$; therefore it suffices to control the behaviour of each 'link' $Z_{\pi_{s+1}v} - Z_{\pi_s v}$ in the chain corresponding to v to obtain a bound on $\sup_{v \in V} (Z_v - Z_{\pi_0 v})$. Intuitively, the number of links in the s stage is at most $|V_s| \cdot |V_{s+1}|$, and that cardinality has to be 'balanced' with the probability that the link $Z_{\pi_{s+1}v} - Z_{\pi_s v}$ is 'large'. As a result, if there is some compatibility between the notion of approximation used in V and that probability, the chaining argument leads to an upper bound on $\mathbb{E} \sup_{v \in V} Z_v$ that is based on metric features of the indexing set V.

For a detailed exposition on chaining methods we refer the reader to Talagrand's treasured book [27].

Even with this somewhat vague description of generic chaining, it is clear that the key to an upper bound on $\mathbb{E} \sup_{v \in V} Z_v$ is the *increment condition*: an estimate on the probability that $|Z_v - Z_w|$ is much larger than the 'distance' between v and w.

Definition 3.1. An admissible sequence of a metric space (V, ρ) is a collection $(V_s)_{s\geq 0}$ of subsets of V that satisfy $|V_s| \leq 2^{2^s}$ and $|V_0| = 1$. For $v \in V$, denote by $\pi_s v$ a nearest element to v in V_s , and set

$$\gamma_2(V,\rho) = \inf_{(V_s)_{s \ge 0}} \sup_{v \in V} \sum_{s \ge 0} 2^{s/2} \rho(\pi_s v, \pi_{s+1} v),$$

with the infimum taken over all admissible sequences.

What is a remarkable justification of the generic chaining mechanism is that it leads to a complete metric characterization of the expected supremum of gaussian processes:

Theorem 3.2. There are absolute constants c_1 and c_2 for which the following hold. Let $\{G_v : v \in V\}$ be a centred gaussian process and set $\rho(v, w) = ||G_v - G_w||_{L_2}$. Then

$$c_1\gamma_2(V,\rho) \le \mathbb{E}\sup_{v\in V} G_v \le c_2\gamma_2(V,\rho).$$

The upper bound follows by filling the gaps in our vague description of generic chaining, while the lower bound is Talagrand's celebrated majorizing measures theorem [26].

3.1. A preprocessing step. The key ingredient in the proof of the upper estimate in Theorem 3.2 is the subgaussian tail estimate. Crucially, such a tail estimate is simply false when considering other processes—for example—an empirical process in a heavy tailed situation. What is true, however, is that there are mean estimation procedures that exhibit subgaussian tails even for general random variables.

For the remainder of this article, denote by $\psi_{\delta} : \mathbb{R}^N \to \mathbb{R}$ an optimal mean-estimation procedure, i.e., a data dependent procedure that satisfies the following.

Theorem 3.3. There are absolute constants c_0, c_1 and for any $\delta \geq 2 \exp(-c_0 N)$ and any random variable Z that has a finite mean and variance,

(3.2)
$$|\psi_{\delta}(Z_1,...,Z_N) - \mathbb{E}Z| \le c_1 \sigma_Z \sqrt{\frac{\log(2/\delta)}{N}}$$
 with probability at least $1 - \delta$,

where σ_Z^2 is the variance of Z.

The fact that (3.2) is best possible behaviour of a mean estimation procedure is standard; and among the examples of such optimal mean estimators are the *median-of-means* (see, e.g. [2, 12, 25]) and the *trimmed mean* (see, e.g. [17]). For more results on optimal meanestimation procedures see e.g., [1, 8, 10, 24], and for a survey see [15].

Remark 3.4. The identity of the mean estimation procedure that will be used in what follows is of no importance. As far as the proof of Theorem 1.16 goes, all that matters is that ψ_{δ} satisfies (3.2).

The next ingredient is an almost optimal admissible sequence $(H_s)_{s\geq 0}$ whose existence is guaranteed by Theorem 3.2. Such a sequence can be constructed by solving an optimization problem–something that can, in-principle, be done if one knows the underlying metric. Although the learner does not have access to the L_2 metric, the functional d works equally well, as it satisfies that

$$\eta^{-1} \| f - h \|_{L_2} \le d(f, h) \le \eta \| f - h \|_{L_2}$$
 for every $f, h \in H \cup \{0\}$.

Therefore, solving the optimization problem for d would lead to an admissible sequence $(H_s)_{s>0}$ for which

(3.3)
$$\sum_{s\geq 0} 2^{s/2} \|\Delta_s h\|_{L_2} \leq c(\eta) \mathbb{E} \sup_{h\in H} G_h.$$

In a similar fashion, an almost optimal admissible sequence of $(H - H) \cap rD$ can be constructed for every $r \ge 0$.

Having access to the L_2 distance oracle d leads to suitable admissible sequences for the classes we care about, namely F and its localizations $(F - F) \cap rD$. Moreover, there is these admissible sequences and the majorizing measures the

Moreover, thanks to those admissible sequences and the majorizing measures theorem, it is possible to identify $\mathbb{E}\sup_{h\in (F-F)\cap rD} G_h$ (up to multiplicative constants that depend on η). Therefore, one may derive estimates on the fixed point condition $\mathbb{E}\sup_{h\in (F-F)\cap rD} G_h \leq \kappa r \sqrt{N}$.

3.2. Subgaussian mean estimators for multiplier classes. Let $\xi \in L_4$ be a random variable (that need not be independent of X).

Our goal here is to design a uniform mean estimation procedure for $\mathbb{E}\xi h$: the procedure receives as data the values $(\xi_i, h(X_i))_{i=1}^N$ and returns its 'guess' of $\mathbb{E}\xi h$.

The procedure should exhibit a subgaussian accuracy/confidence tradeoff: if H were L-subgaussian, that is, if for every $f, h \in H \cup \{0\}$, $||f - h||_{\psi_2} \leq L||f - h||_{L_2}$, and if $||\xi||_{\psi_2} < \infty$, a nontrivial chaining argument (see [21] for the proof) shows that the following holds. Let H be a centrally symmetric² set, put

$$R_H = \sup_{h \in H} \|h\|_{L_2}$$

and set

$$d_*(H) = \left(\frac{\mathbb{E}\sup_{h \in H} G_h}{R_H}\right)^2,$$

to be the *critical dimension* of the set H. Then for $N \ge c_0 d_*(H)$, with probability at least $1 - 2 \exp(-c_1 d_*(H))$,

(3.4)
$$\sup_{h \in H} \left| \frac{1}{N} \sum_{i=1}^{N} \xi_i h(X_i) - \mathbb{E} \xi h \right| \le c_2(L) \frac{\|\xi\|_{\psi_2}}{\sqrt{N}} \cdot \mathbb{E} \sup_{h \in H} G_h.$$

We would like to design a mean estimator that exhibits a similar error as the empirical mean in (3.4), but holds for a heavy-tailed class of functions H and a heavy-tailed multiplier ξ . Obviously, that estimator cannot be the empirical mean, whose performance deteriorates dramatically when the indexing class is not subgaussian.

As it happens, the subgaussian assumption can be relaxed considerably.

²That is, if $h \in H$ then $-h \in H$.

Assumption 3.5. *H* is a centrally symmetric class of mean zero functions that satisfies for every $f, h \in H \cup \{0\}$,

$$||f - h||_{L_4} \le L||f - h||_{L_2}.$$

Also, assume as always that the learner has access to a symmetric functional d that satisfies

$$\eta^{-1} \|f - h\|_{L_2} \le d(f, h) \le \eta \|f - h\|_{L_2}$$

for every $f, h \in H \cup \{0\}$.

Remark 3.6. Clearly the $L_4 - L_2$ norm equivalence condition in Assumption 3.5 is significantly weaker than the $\psi_2 - L_2$ norm equivalence that holds for subgaussian classes. In fact, functions in H need not have $L_{4+\varepsilon}$ moments, let alone arbitrary high moments, and thus can heavy tailed.

Theorem 3.7. If H satisfies Assumption 3.5, there is an absolute constant c_1 and a constant $c_2 = c_2(L,\eta)$ for which the following holds. Let $\xi \in L_4$ and set $2^{s_0} \leq c_1 N$. There exists a procedure $\Phi_{\mathbb{M}}$ for which, with probability at least $1 - 2 \exp(-2^{s_0})$,

$$\sup_{h \in H} |\Phi_{\mathbb{M}}(h) - \mathbb{E}\xi h| \le c_2 \frac{\|\xi\|_{L_4}}{\sqrt{N}} \left(\mathbb{E} \sup_{h \in H} G_h + 2^{s_0/2} R_H \right).$$

In particular, if $N \ge c_1 d_*(H)$, then setting $2^{s_0} \sim d_*(H)$ we have that with probability at least $1 - 2 \exp(-c_3 d_*(H))$,

$$\sup_{h \in H} |\Phi_{\mathbb{M}}(h) - \mathbb{E}\xi h| \le c_4(L,\eta) \frac{\|\xi\|_{L_4}}{\sqrt{N}} \mathbb{E} \sup_{h \in H} G_h,$$

thus exhibiting the wanted subgaussian behaviour as in (3.4).

The proof of Theorem 3.7 follows a standard chaining argument—with one significant twist: the subgaussian increment condition holds thanks to the mean estimation procedure ψ_{δ} that satisfies (3.2) for the right choices of δ .

Proof of Theorem 3.7. Let $\alpha \geq 5$ be a well-chosen absolute constant and set s_1 to be the largest integer satisfying $\alpha 2^{s_1} \leq c_0 N$, where c_0 is the constant appearing in Theorem 3.3. In particular $s_0 < s_1$. Let $(H_s)_{s\geq 0}$ be an admissible sequence of H that satisfies (3.3).

The proof consists of two steps. First, we design a mean estimator for $\xi \pi_{s_1} h$, and later show that the means of ξh and $\xi \pi_{s_1} h$ are sufficiently close.

Step 1: Write

$$\xi \pi_{s_1} h = \xi \pi_{s_0} h + \sum_{s=s_0}^{s_1 - 1} \xi \Delta_s h,$$

and note that for every $s \leq s_1$, $\delta_s = 2 \exp(-\alpha 2^s)$ is a legal choice of δ in Theorem 3.3. Moreover, the cardinality of the set $\{\xi \Delta_s h : h \in H\}$ is at most $2^{2^{s+2}}$. Define

$$\Phi_{\mathbb{M}}(h) = \psi_{\delta_{s_0}} \left(\left(\xi_i \pi_{\delta_{s_0}} h(X_i) \right)_{i=1}^N \right) + \sum_{s=s_0}^{s_1-1} \psi_{\delta_s} \left(\left(\xi_i \Delta_s h(X_i) \right)_{i=1}^N \right),$$

and the key is to show that with probability at least $1 - 2\exp(-2^{s_0})$, for every $h \in H$,

(3.5)
$$|\Phi_{\mathbb{M}}(h) - \mathbb{E}\xi \pi_{s_1}h| \leq c_1(L) \frac{\|\xi\|_{L_4}}{\sqrt{N}} \left(\sum_{s=s_0}^{s_1-1} 2^{s/2} \|\Delta_s h\|_{L_2} + 2^{s_0/2} \|\pi_{s_0}h\|_{L_2} \right)$$

To establish (3.5), fix $s_0 \leq s < s_1$ and examine each term $\psi_{\delta_s}((\xi_i \Delta_s h(X_i))_{i=1}^N)$ —which is an optimal mean estimator of $\xi \Delta_s h$. Hence, each random variable of the form $\xi \Delta_s h$ satisfies that with probability at least $1 - 2 \exp(-\alpha 2^s)$,

(3.6)
$$\left|\psi_{\delta_s}\left(\left(\xi_i\Delta_s h(X_i)\right)_{i=1}^N\right) - \mathbb{E}\xi\Delta_s h\right| \le c_2 \|\xi\Delta_s h\|_{L_2} \sqrt{\frac{2^s}{N}},$$

and clearly,

$$\|\xi\Delta_s h\|_{L_2} \le \|\xi\|_{L_4} \|\Delta_s h\|_{L_4} \le L \|\xi\|_{L_4} \|\Delta_s h\|_{L_2}.$$

Moreover, thanks to the high probability estimate with which (3.6) holds and the fact that there are at most $2^{2^{s+2}} \leq \exp(-4 \cdot 2^s)$ functions of the form $\xi \Delta_s h$, we have that with probability $1 - 2\exp(-(\alpha - 4)2^s)$, every such function satisfies (3.6).

Repeating this argument for every $s_0 \leq s \leq s_1 - 1$, and also for the functions $\{\xi \pi_{s_0} h : h \in H\}$, it is evident by the union bound that with probability at least $1 - 2 \exp(-2^{s_0})$, for every $h \in H$ and every $s_0 \leq s \leq s_1 - 1$,

(3.7)
$$\left| \psi_{\delta_s} \left((\xi_i \Delta_s h(X_i))_{i=1}^N \right) - \mathbb{E} \xi \Delta_s h \right| \le c_2 L \|\xi\|_{L_4} \|\Delta_s h\|_{L_2} \sqrt{\frac{2^s}{N}} \quad \text{and} \\ \left| \psi_{\delta_{s_0}} \left((\xi_i \pi_{s_0} h(X_i))_{i=1}^N \right) - \mathbb{E} \xi \pi_{s_0} h \right| \le c_2 L \|\xi\|_{L_4} \|\pi_{s_0} h\|_{L_2} \sqrt{\frac{2^{s_0}}{N}}.$$

Finally,

$$\mathbb{E}\xi\pi_{s_1}h = \mathbb{E}\xi\pi_{s_0}h + \sum_{s=s_0}^{s_1-1}\mathbb{E}\xi\Delta_sh,$$

and the wanted estimate in (3.5) follows from (3.7).

Step 2: Observe that for every $h \in H$,

(3.8)
$$|\mathbb{E}\xi h - \mathbb{E}\xi \pi_{s_1}h| \le \frac{c_3 \|\xi\|_{L_2}}{\sqrt{N}} \sum_{s \ge s_1} 2^{s/2} \|\Delta_s h\|_{L_2}.$$

Indeed,

$$\|\mathbb{E}\xi h - \mathbb{E}\xi \pi_{s_1}h\| \le \|\xi(h - \pi_{s_1}h)\|_{L_1} \le \|\xi\|_{L_2}\|h - \pi_{s_1}h\|_{L_2}$$

Since $h - \pi_{s_1} h = \sum_{s \ge s_1} \Delta_s h$, it follows from the triangle inequality and the choice of s_1 that

$$\|h - \pi_{s_1}h\|_{L_2} \le \frac{1}{2^{s_1/2}} \sum_{s \ge s_1} 2^{s_1/2} \|\Delta_s h\|_{L_2} \le \frac{c_3}{\sqrt{N}} \sum_{s \ge s_1} 2^{s/2} \|\Delta_s h\|_{L_2},$$

and (3.8) follows.

The proof is completed by combining the estimates in (3.7) and (3.8) and recalling that $(H_s)_{s\geq 0}$ satisfies

$$\sum_{s \ge s_0} 2^{s/2} \|\Delta_s h\|_{L_2} \le c_4(\eta) \mathbb{E} \sup_{h \in H} G_h.$$

Corollary 3.8. Let H be a class of mean zero functions that satisfies $L_4 - L_2$ norm equivalence with constant L. Let $2^{s_0} \leq c_1 N$ and set $\mathcal{U} = \{\xi_j : 1 \leq j \leq 2\exp(-2^{s_0-1})\} \subset L_4$ to be a collection of functions. Then there is a procedure $\Phi_{\mathbb{M}}$ for which, with probability $1 - 2\exp(-2^{s_0-1})$, for every $\xi \in \mathcal{U}$,

$$\sup_{h \in H} |\Phi_{\mathbb{M}}(h,\xi) - \mathbb{E}\xi h| \le c(L) \frac{\|\xi\|_{L_4}}{\sqrt{N}} \left(\mathbb{E}\sup_{h \in H} G_h + 2^{s_0/2} R_H \right)$$

Remark 3.9. The procedure $\Phi_{\mathbb{M}}$ has access to the values $(h(X_i))_{i=1}^N$ and $(\xi_i)_{i=1}^N$ for every $\xi \in \mathcal{U}$, and it is constructed using an admissible sequence of H. However, it has no access to the identity of each ξ (or even to the identity of the set \mathcal{U}) beyond the given sample.

3.3. A subgaussian mean estimator for a product class. A somewhat more involved problem is dealing with products of classes. Let $H^1, H^2 \subset L_2(\mu)$ be centrally symmetric classes of mean zero functions, and one would like to obtain a uniform estimate on $\mathbb{E}fh$ for $f \in H^1$ and $h \in H^2$. The benchmark is the performance of the empirical mean when H^1 and H^2 are subgaussian with constant L. In such a scenario the right estimate requires a rather involved chaining argument.

Theorem 3.10. Assume that H^1 and H^2 are centrally symmetric L-subgaussian classes of mean zero functions and set $2^{s_0} \leq c_1 N$. Then with probability at least $1 - 2 \exp(-2^{s_0})$, for every $f \in H^1$ and $h \in H^2$,

$$\left|\frac{1}{N}\sum_{i=1}^{N} f(X_i)h(X_i) - \mathbb{E}fh\right| \le \frac{c_2(L)}{\sqrt{N}} \left(R_{H^1}\mathbb{E}\sup_{h\in H^2} G_h + R_{H^2}\mathbb{E}\sup_{f\in H^1} G_f + 2^{s_0/2}R_{H^1}R_{H^2} \right).$$

Theorem 3.10 is an immediate outcome of Theorem 1.13 in [19].

As it happens, one may replace the empirical mean by a better mean estimator and obtain the same subgaussian bound — even if the classes merely satisfy $L_4 - L_2$ norm equivalence rather than the $\psi_2 - L_2$ norm equivalence needed in Theorem 3.10.

Theorem 3.11. If H^1 and H^2 satisfy Assumption 3.5, then there is an absolute constant c_1 and a constant $c_2 = c_2(L,\eta)$ for which the following holds. For every $2^{s_0} \leq c_1 N$, there

is a procedure $\Psi_{\mathbb{Q}}$ for which, with probability at least $1 - 2\exp(-2^{s_0})$, for every $f \in H^1$ and $h \in H^2$,

$$|\Psi_{\mathbb{Q}}(f,h) - \mathbb{E}fh| \le \frac{c_2}{\sqrt{N}} \left(R_{H^1} \mathbb{E} \sup_{h \in H^2} G_h + R_{H^2} \mathbb{E} \sup_{f \in H^1} G_f + 2^{s_0/2} R_{H^1} R_{H^2} \right).$$

Proof. To simplify notation, for any function a = a(X), we write $\psi_{\delta}(a)$ instead of $\psi_{\delta}((a(X_i)_{i=1}^N))$.

Set $\alpha \geq 10$ to be a well-chosen absolute constant and let s_1 be the largest integer satisfying $\alpha 2^{s_1} \leq c_0 N$; hence $s_0 < s_1$. Set $\delta_s = 2 \exp(-\alpha 2^s)$ and let $(H_s^1)_{s\geq 0}$ and $(H_s^2)_{s\geq 0}$ be admissible sequences of H^1 and H^2 that satisfy (3.3).

Step 1: We begin by designing a uniform mean estimator for the products $\pi_{s_1} f \pi_{s_1} h$. To that end, set

$$\Psi(f,h) = \psi_{\delta_{s_0}} \left(\pi_{s_0} f \pi_{s_0} h \right) + \sum_{s=s_0}^{s_1-1} \psi_{\delta_s} \left(\pi_{s+1} f \pi_{s+1} h - \pi_s f \pi_s h \right)$$

Let us show that with probability at least $1 - 2\exp(-2^{s_0})$, for every $f \in H^1$ and $h \in H^2$,

$$(3.9) \quad |\Psi(f,h) - \mathbb{E}\pi_{s_1} f \pi_{s_1} h| \le \frac{c}{\sqrt{N}} \left(R_{H^1} \mathbb{E} \sup_{h \in H^2} G_h + R_{H^2} \mathbb{E} \sup_{f \in H^1} G_f + \sqrt{2^{s_0}} R_{H^1} R_{H^2} \right),$$

where $c = c(L, \eta)$.

Let $s \leq s_1$ and note that δ_s is a legal choice of δ in the context of Theorem 3.3. Therefore, with probability at least $1 - 2 \exp(-\alpha 2^s)$,

(3.10)
$$\begin{aligned} |\psi_{\delta_s} \left(\pi_{s+1} f \pi_{s+1} h - \pi_s f \pi_s h \right) - \mathbb{E} \left(\pi_{s+1} f \pi_{s+1} h - \pi_s f \pi_s h \right)| \\ &\leq c_1 \|\pi_{s+1} f \pi_{s+1} h - \pi_s f \pi_s h\|_{L_2} \sqrt{\frac{2^s}{N}}. \end{aligned}$$

There are at most $2^{2^{s+4}}$ functions of the form $\pi_{s+1}f\pi_{s+1}h - \pi_sf\pi_sh$, and it follows from the union bound that with probability at least $1 - 2\exp(-2 \cdot 2^{s_0})$, (3.10) holds uniformly for every $f \in H^1$, $h \in H^2$ and $s_0 \leq s < s_1$. Moreover,

$$\pi_{s+1}f\pi_{s+1}h - \pi_s f\pi_s h = (\Delta_s f)\pi_{s+1}h - (\Delta_s h)\pi_s f,$$

and by the Cauchy-Schwartz inequality and the $L_4 - L_2$ norm-equivalence,

$$\begin{aligned} \|\pi_{s+1}f\pi_{s+1}h - \pi_s f\pi_s h\|_{L_2} &\leq \|\Delta_s f\|_{L_4} \|\pi_{s+1}h\|_{L_4} + \|\Delta_s h\|_{L_4} \|\pi_s f\|_{L_4} \\ &\leq L^2 \left(\|\Delta_s f\|_{L_2} R_{H^2} + \|\Delta_s h\|_{L_2} R_{H^1}\right). \end{aligned}$$

Therefore, in the high probability event in which (3.10) holds,

$$\sum_{s=s_0}^{s_1-1} |\psi_{\delta_s} \left(\pi_{s+1} f \pi_{s+1} h - \pi_s f \pi_s h \right) - \mathbb{E} \left(\pi_{s+1} f \pi_{s+1} h - \pi_s f \pi_s h \right)|$$

$$\leq \frac{c_1 L^2}{\sqrt{N}} \sum_{s=s_0}^{s_1-1} 2^{s/2} \left(R_{H^2} \|\Delta_s f\|_{L_2} + R_{H^1} \|\Delta_s h\|_{L_2} \right)$$

$$\leq \frac{c_2(L,\eta)}{\sqrt{N}} \left(R_{H^2} \mathbb{E} \sup_{f \in H^1} G_f + R_{H^1} \mathbb{E} \sup_{h \in H^2} G_h \right),$$

where the last inequality follows thanks to the choices of the admissible sequences $(H_s^1)_{s\geq 0}$ and $(H_s^2)_{s\geq 0}$.

The same arguments can be used to show that with probability at least $1-2\exp(-2\cdot 2^{s_0})$, for every $f \in H^1$ and $h \in H^2$,

$$\left|\psi_{\delta_{s_0}}\left(\pi_{s_0}f\pi_{s_0}h\right) - \mathbb{E}\pi_{s_0}f\pi_{s_0}h\right| \le c_3(L)R_{H^1}R_{H^2}\sqrt{\frac{2^{s_0}}{N}},$$

and (3.9) follows.

Step 2: To complete the proof, it suffices to show that for every $f \in H^1$ and $h \in H^2$,

(3.11)
$$|\mathbb{E}fh - \mathbb{E}\pi_{s_1}f\pi_{s_1}h| \leq \frac{c_4(\eta)}{\sqrt{N}} \left(R_{H^1}\mathbb{E}\sup_{h \in H^2} G_h + R_{H^2}\mathbb{E}\sup_{f \in H^1} G_f \right).$$

To that end, note that

$$|\mathbb{E}fh - \mathbb{E}\pi_{s_1}f\pi_{s_1}h| \le ||f - \pi_{s_1}f||_{L_2}||h||_{L_2} + ||h - \pi_{s_1}h||_{L_2}||\pi_{s_1}f||_{L_2},$$

and $\|h\|_{L_2} \leq R_{H^2}$, $\|\pi_{s_1} f\|_{L_2} \leq R_{H^1}$. Finally, using that $2^{s_1} \sim N$ and (3.3),

$$\|f - \pi_{s_1} f\|_{L_2} \le \frac{1}{2^{s_1/2}} \sum_{s \ge s_1} 2^{s/2} \|\Delta_s f\|_{L_2} \le \frac{c_5(\eta)}{\sqrt{N}} \mathbb{E} \sup_{f \in H^1} G_f.$$

The analogue estimate on $||h - \pi_{s_1}h||_{L_2}$ clearly follows from the same argument, showing that (3.11) holds.

4. A CRUDE RISK ORACLE

From now on, we shall always assume the norm equivalence condition and the existence of a distance oracle, namely:

- For every $f, h \in F \cup \{0\}, \|f h\|_{L_4} \le L \|f h\|_{L_2}$.
- For every $f \in F$, $||f Y||_{L_4} \le L||f Y||_{L_2}$.
- d is symmetric and $\eta^{-1} ||f h||_{L_2} \le d(f, h) \le \eta ||f h||_{L_2}$ for every $f, h \in F \cup \{0\}$.

Proof of Theorem 2.1. Let θ and γ be (small) constants that depend on η and L and that are chosen in what follows. Moreover, set $2^{s_0} = \theta^2 N$, and assume that $\log \mathcal{M}(F, \gamma r D) \leq 2^{s_0-1}$.

First we use the distance oracle d to construct a partition of F to at most $\exp(2^{s_0-1})$ sets in the following way: let (u_j) be a maximal $\eta\gamma r$ separated subset of F with respect to d. Therefore, it is an γr -separated set with respect to L_2 and hence its cardinality is at most $\exp(2^{s_0-1})$. Set U_j to be the corresponding partition of F to disjoint sets whose 'centres' are u_j . It follows that every $f \in F$ belongs to a unique set U_j ; thus $d(u_j, f) \leq \eta\gamma r$, implying that

$$\|f - u_j\|_{L_2} \le \eta^2 \gamma r.$$

Let $\psi_{\delta_{s_0}}$ be the optimal mean estimator as in Theorem 3.3 with $\delta_{s_0} = 2 \exp(-2^{s_0})$. It follows from the union bound that with probability at least $1 - 2 \exp(-2^{s_0-1})$, for every $1 \le j \le \exp(2^{s_0-1})$,

(4.1)
$$\left|\psi_{\delta_{s_0}}\left((u_j - Y)^2\right) - \mathbb{E}(u_j - Y)^2\right| \le c_1 \sqrt{\frac{2^{s_0}}{N}} \|(u_j - Y)^2\|_{L_2} \le \frac{1}{8} \|u_j - Y\|_{L_2}^2,$$

where we have also used norm equivalence in the second inequality and that $2^{s_0} = \theta^2 N$ for $\theta = 1/(8c_1L)$. For $f \in F$, let u_j be the center of the unique U_j where f belongs to and set

$$\Psi_{\mathbb{C}}(f) = \psi_{\delta_{s_0}} \left((u_j - Y)^2 \right)$$

Thus, in the event in which (4.1) holds, we have that for every $f \in F$,

$$\begin{aligned} \left| \Psi_{\mathbb{C}}(f) - \mathbb{E}(f-Y)^2 \right| &\leq \frac{1}{8} \mathbb{E}(u_j - Y)^2 + \left| \mathbb{E}(u_j - Y)^2 - \mathbb{E}(f-Y)^2 \right| \\ &\leq \frac{1}{8} \mathbb{E}(f-Y)^2 + \frac{9}{8} \left| \mathbb{E}(u_j - Y)^2 - \mathbb{E}(f-Y)^2 \right| = (1). \end{aligned}$$

Therefore, if $\mathbb{E}(f-Y)^2$ and $\mathbb{E}(u_j-Y)^2$ are sufficiently close, namely

(4.2)
$$\left| \mathbb{E}(u_j - Y)^2 - \mathbb{E}(f - Y)^2 \right| \leq \frac{1}{9} \mathbb{E}(f - Y)^2 + \frac{1}{4}r^2,$$

then $(1) \leq \frac{1}{2} \max\{r^2, \mathbb{E}(f-Y)^2\}.$

To show that (4.2) holds, observe that

$$(u_j - Y)^2 - (f - Y)^2 = (u_j - f)^2 + 2(u_j - f)(f - Y),$$

and thus

$$\begin{aligned} \left| \mathbb{E}(u_j - Y)^2 - \mathbb{E}(f - Y)^2 \right| &\leq \|u_j - f\|_{L_2}^2 + 2\|u_j - f\|_{L_2} \|f - Y\|_{L_2} \\ &\leq (\eta^2 \gamma r)^2 + 2\eta^2 \gamma r \|f - Y\|_{L_2} \\ &\leq (\eta^2 \gamma r)^2 + 9(\eta^2 \gamma r)^2 + \frac{\|f - Y\|_{L_2}^2}{9}, \end{aligned}$$

where we have used the elementary fact $ab \leq \frac{t}{4}a^2 + \frac{1}{t}b^2$ for a, b, t > 0 in the last inequality. Thus, if $\gamma = \gamma(\eta)$ is a sufficiently small constant, (4.2) follows

Proof of Corollary 2.2. Recall that $\sigma^2 = \inf_{f \in F} \mathbb{E}(f - Y)^2$. Fix a realization of $(X_i, Y_i)_{i=1}^N$ for which the assertion of Theorem 2.1 holds, i.e., for every $f \in F$,

$$|\Psi_{\mathbb{C}}(f) - \mathbb{E}(f(X) - Y)^2| \le \frac{1}{2} \max\{r^2, \mathbb{E}(f(X) - Y)^2\}.$$

It follows that if $\mathbb{E}(f(X) - Y)^2 > r^2$, then

$$\frac{1}{2}\mathbb{E}(f(X) - Y)^2 \le \Psi_{\mathbb{C}}(f) \le 2\mathbb{E}(f(X) - Y)^2,$$

and if $\mathbb{E}(f(X) - Y)^2 \leq r^2$, then

$$\Psi_{\mathbb{C}}(f) \le 2r^2.$$

Therefore, setting

$$\widehat{\sigma}^2 = \inf_{f \in F} \Psi_{\mathbb{C}}(f),$$

it is evident that if $\sigma \geq r$ we have that $\frac{1}{2}\sigma^2 \leq \hat{\sigma}^2 \leq 2\sigma^2$, and if $\sigma < r$, then $\hat{\sigma}^2 \leq 2r^2$. Let $\sigma_* = \max\{r, \hat{\sigma}\}$ and observe that $f^* \in \hat{V}$, i.e. $\Phi_{\mathbb{C}}(f^*) \leq 4\sigma_*^2$. Indeed, if $\sigma > r$, then

$$\Psi_{\mathbb{C}}(f^*) \le 2\sigma^2 \le 4\widehat{\sigma}^2$$

and if $\sigma \leq r$, then $\Psi_{\mathbb{C}}(f^*) \leq 2r^2$.

Finally, to show that if $f \in \hat{V}$, then $\mathbb{E}(f - Y)^2 \leq 16 \max\{r^2, \sigma^2\}$, we may assume that $\mathbb{E}(f - Y)^2 > r^2$ and therefore

$$\mathbb{E}(f-Y)^2 \le 2\Psi_{\mathbb{C}}(f) \le 2 \cdot 4 \max\{r^2, \widehat{\sigma}^2\} \le 16 \max\{r^2, \sigma^2\},$$

as claimed.

5. A MIXTURE MEAN ESTIMATION PROCEDURE

Finally we have all the ingredients needed for the proof of Theorem 2.3. But before we proceed we need to make a short detour: Set U = W = F - F, and V = F. For every $f, h \in F$, let u = w = f - h and note that $u \in U$ and w = W. Also, set v = h. Hence,

$$(f - Y)^{2} - (h - Y)^{2} = u(w + 2(v - Y)).$$

We will design a mean estimation procedure that holds uniformly for all functions in the class

$$\{u(w+2(v-Y)): u \in U, w \in W, v \in V\}.$$

The procedure is given the identities of u, w, v and the second half of the sample $(X_i, Y_i)_{i=N+1}^{2N}$, and returns its best guess of the mean $\mathbb{E}u(w+2(v-Y))$.

As will become clear immediately, that leads to a uniform mean estimation procedure for all the functions of the form $(f - Y)^2 - (h - Y)^2$.

Theorem 5.1. There are absolute constants c_1, c_2 and a constant $c_3 = c_3(L, \eta)$ such that the following holds. Let $\theta \in (0, 1)$, set $2^{s_0} \leq c_1 \theta^2 N \min\{1, \frac{r^2}{\sigma^2}\}$ and consider r for which

(5.1)
$$\mathbb{E}\sup_{h\in(F-F)\cap rD}G_h\leq\theta\sqrt{N}r\min\left\{1,\frac{r}{\sigma}\right\}\quad and\quad \log\mathcal{M}(F,rD)\leq2^{s_0-1}.$$

Then there exists a mean estimation procedure Ψ_* that satisfies, with probability at least $1 - 2\exp(-c_22^{s_0})$, for every $u, w \in F - F$ and $v \in F$,

$$\Psi_*(u, w, v) - \mathbb{E}u(w + 2(v - Y))|$$

$$\leq c_3 \theta \left(\max\{r, \|u\|_{L_2}\} \max\{r, \|w\|_{L_2}\} + r \max\{r, \|u\|_{L_2}\} \frac{r + \|v - Y\|_{L_2}^2}{\max\{r, \sigma\}} \right)$$

Remark 5.2. Note that the first condition in (5.1) just means that $r \ge \max\{r_{\mathbb{Q}}(\theta), r_{\mathbb{M}}(\theta)\}$.

Let us show how the assertion of Theorem 5.1 leads to the proof of Theorem 2.3.

Proof of Theorem 2.3. Let $(X_i, Y_i)_{i=1}^{2N}$ be the given sample. Its first half is used compute the crude risk estimator (considered in Theorem 2.1 and Lemma 1.14), and it outputs the value of σ_* . The second half of the sample is used to compute $\Psi_{\mathcal{L}}$, defined in what follows, with a choice of 2^{s_0} that is determined by σ_* .

Fix a realization of the sample $(X_i, Y_i)_{i=1}^N$ for which the assertion of Theorem 2.1 holds. This event has probability $1 - \exp(-c_0 N)$, and we have that

$$\min\left\{1, \frac{r^2}{\sigma_*^2}\right\} \sim \min\left\{1, \frac{r^2}{\sigma^2}\right\}.$$

Next, in the setting of Theorem 5.1 and using its notation, let $\theta = 1/(12c_3)$ and set

$$2^{s_0} \sim \theta^2 N \min\left\{1, \frac{r^2}{\sigma_*^2}\right\} \le c_1 \theta^2 N \min\left\{1, \frac{r^2}{\sigma^2}\right\};$$

Thus r satisfies (5.1).

For $f, h \in F$ let

$$\Psi_{\mathcal{L}}(f,h) = \Psi_*(f-h,f-h,h),$$

where Ψ_* is computed using $(X_i, Y_i)_{i=N+1}^{2N}$. If we set u = w = f - h and v = h, then

$$\mathbb{E}u(w+2(v-Y)) = \mathbb{E}(f-Y)^2 - \mathbb{E}(h-Y)^2,$$

and in the high probability event in which the assertion of Theorem 5.1 holds,

$$\begin{aligned} \left| \Psi_{\mathcal{L}}(f,h) - \left(\mathbb{E}(f-Y)^2 - \mathbb{E}(h-Y)^2 \right) \right| \\ &\leq \frac{1}{12} \left(\max\{r^2, \|f-h\|_{L_2}^2\} + r \max\{r, \|f-h\|_{L_2}\} \frac{r+\|h-Y\|_{L_2}}{\max\{r,\sigma\}} \right). \end{aligned}$$

Finally, if $||h-Y||_{L_2}^2 \leq 16 \max\{r^2, \sigma^2\}$, then using the inequality $a \max\{a, b\} \leq \max\{a^2, b^2\}$, it follows that

$$\begin{split} \left| \Psi_{\mathcal{L}}(f,h) - \left(\mathbb{E}(f-Y)^2 - \mathbb{E}(h-Y)^2 \right) \right| \\ &\leq \frac{1}{12} \max\{r^2, \|f-h\|_{L_2}^2\} \left(1 + \frac{r+4\max\{r,\sigma\}}{\max\{r,\sigma\}} \right) \\ &\leq \frac{1}{2} \max\{r^2, \|f-h\|_{L_2}^2\}, \end{split}$$

as claimed.

Let us turn to the definition of the procedure Ψ_* . The starting point is to use the distance oracle and the entropy estimate in (5.1) to construct a partition of F to at most $\exp(2^{s_0-1})$ sets V_j with centres v_j . Every $v \in F$ belongs to a unique V_j , and

$$\|v - v_j\|_{L_2} \le \eta^2 r = r_0.$$

Moreover, for every u, w and v,

$$u(w + 2(v - Y)) = uw + 2u(v - v_j) + 2u(v_j - Y).$$

Set $\Psi_{\mathbb{Q}}$ to be the product mean estimation procedure for the classes $H^1 = H^2 = (F - F) \cap r_0 D$, and let $\Psi_{\mathbb{M}}$ be the mean estimation procedure for the class $H = (F - F) \cap r_0 D$ and the $\exp(2^{s_0-1})$ multipliers $v_i - Y$.

Next, recall that $r \leq r_0$, for $\tilde{u} \in F - F$ set $\alpha(\tilde{u}) = \max\{r, d(\tilde{u}, 0)\}$, and note that

$$\frac{r\widetilde{u}}{\alpha(\widetilde{u})} \in (F - F) \cap r_0 D.$$

With that in mind, for $\widetilde{w}, \widetilde{u} \in F - F$ set

$$u = \frac{r\widetilde{u}}{\alpha(\widetilde{u})}, \text{ and } w = \frac{r\widetilde{w}}{\alpha(\widetilde{w})},$$

and define the following mean estimation procedures for $\widetilde{w}, \widetilde{u} \in F - F$ and $v \in F$. Let

$$\Psi_1(\widetilde{u},\widetilde{w}) = \frac{\alpha(\widetilde{u})}{r} \cdot \frac{\alpha(\widetilde{w})}{r} \Psi_{\mathbb{Q}}(u,w).$$

In other words, scale-down \tilde{u} and \tilde{w} into $(F - F) \cap r_0 D$; then use the product mean estimation procedure in $(F - F) \cap r_0 D$; and finally, re-scale using the inverse factor. Since $\mathbb{E}uw$ is homogeneous, the result will be an estimator of $\mathbb{E}\tilde{u}\tilde{w}$.

In a similar fashion, set

$$\Psi_2(\widetilde{u},v) = rac{lpha(\widetilde{u})}{r} \Psi_{\mathbb{Q}}(u,v-v_j),$$

with the scaling and re-scaling only taking place for \tilde{u} . Finally, let

$$\Psi_3(\widetilde{u}, v) = \frac{\alpha(\widetilde{u})}{r} \Psi_{\mathbb{M}}(u, v_j - Y)$$

be a uniform estimator for functions of the form $\tilde{u}\xi_j$ for the $\exp(2^{s_0-1})$ multipliers $\xi_j = v_j - Y$.

Now set for every $\widetilde{u}, \widetilde{w} \in F - F$ and $v \in F$,

$$\Psi_*(\widetilde{u},\widetilde{w},v) = \Psi_1(\widetilde{u},\widetilde{w}) + 2\Psi_2(\widetilde{u},v) + 2\Psi_3(\widetilde{u},v)$$

The key to the performance of Ψ_* is its behaviour on the scaled down versions, which we explore next.

Proposition 5.3. In the setting of Theorem 5.1 and using its notation, we have that with probability at least $1 - 2\exp(-c_22^{s_0})$, for every $u, w \in (F - F) \cap r_0D$ and $v \in F$,

$$\begin{aligned} |\Psi_{\mathbb{Q}}(u,w) - \mathbb{E}uw| &\leq c_3 \theta r^2, \\ |\Psi_{\mathbb{Q}}(u,v-v_j) - \mathbb{E}u(v-v_j)| &\leq c_3 \theta r^2, \quad and \\ |\Psi_{\mathbb{M}}(u,v_j-Y) - u(v_j-Y)| &\leq c_3 \theta r^2 \cdot \frac{r + \|v-Y\|_{L_2}}{\max\{r,\sigma\}}. \end{aligned}$$

Proof. Observe that the choice $2^{s_0} \leq c_1 N$ is valid in the context of Theorem 3.11. Therefore, with probability at least $1 - 2 \exp(-2^{s_0})$, for every $u, w \in (F - F) \cap r_0 D$,

$$|\Psi_{\mathbb{Q}}(u,w) - \mathbb{E}uw| \le \frac{c_2}{\sqrt{N}} \left(r_0 \mathbb{E} \sup_{h \in (F-F) \cap r_0 D} G_h + 2^{s_0/2} r_0^2 \right) = (1),$$

for $c_2 = c_2(L,\eta)$. Moreover, for every $v \in F$, $v - v_j \in (F - F) \cap r_0 D$ —with v_j being the centre in the unique set in the partition (V_j) to which v belongs. Hence, on the same event, for every $u \in (F - F) \cap r_0 D$ and $v \in F$,

$$|\Psi_{\mathbb{Q}}(u, v - v_j) - \mathbb{E}u(v - v_j)| \le (1).$$

Next, by Corollary 3.8 and since

$$||v_j - Y||_{L_4} \le c_3(L) \left(r_0 + ||v - Y||_{L_2} \right),$$

we have that with probability at least $1 - 2\exp(-2^{s_0-1})$, for every $u \in (F-F) \cap r_0 D$ and $v_j, 1 \le j \le \exp(2^{s_0-1})$,

$$\begin{aligned} |\Psi_{\mathbb{M}}(u, v_j - Y) - \mathbb{E}u(v_j - Y)| &\leq c_4 \frac{\|v_j - Y\|_{L_4}}{\sqrt{N}} \left(\mathbb{E} \sup_{h \in (F - F) \cap r_0 D} G_h + 2^{s_0/2} r_0 \right) \\ &\leq c_5 \frac{r_0 + \|v - Y\|_{L_2}}{\sqrt{N}} \left(\mathbb{E} \sup_{h \in (F - F) \cap r_0 D} G_h + 2^{s_0/2} r_0 \right) = (2), \end{aligned}$$

for constants c_4 and c_5 that depend on L, η .

To control the error terms (1) and (2), by the condition on r in (5.1) and since $r \leq r_0 \leq \eta^2 r$, it is evident that

(5.2)
$$\mathbb{E}\sup_{h\in(F-F)\cap r_0D}G_h\leq c_6(\eta)\theta\sqrt{N}r\min\left\{1,\frac{r}{\sigma}\right\},$$

and by the choice of s_0 ,

(5.3)
$$2^{s_0} \lesssim \theta^2 N \min\left\{1, \frac{r^2}{\sigma^2}\right\}.$$

Hence,

$$(1) \le c_7(L,\eta)\theta r^2.$$

Finally, using (5.2), (5.3), the fact that $r_0 \leq \eta^2 r$, and the elementary fact $\min\{1, \frac{r}{\sigma}\} = \frac{r}{\max\{r, \sigma\}}$,

$$(2) \le c_8(r_0 + \|v - Y\|_{L_2}) \cdot \theta r \min\left\{1, \frac{r}{\sigma}\right\} \\ \le c_9 \theta r^2 \cdot \frac{r + \|v - Y\|_{L_2}}{\max\{r, \sigma\}}$$

for constants c_8 and c_9 that depend only on L, η .

Proof of Theorem 5.1. Fix a realization of the high probability event in which the assertion of Proposition 5.3 holds.

To control the error of Ψ_* (when $\alpha(\tilde{u}) > r$ and/or $\alpha(\tilde{w}) > r$), recall the scaling-down

$$u = \frac{r\widetilde{u}}{\alpha(\widetilde{u})}$$
 and $w = \frac{r\widetilde{w}}{\alpha(\widetilde{w})}$

implying that $u, w \in (F - F) \cap r_0 D$. Hence

$$\left|\frac{\alpha(\widetilde{u})\alpha(\widetilde{w})}{r^2} \cdot (\Psi_{\mathbb{Q}}(u,w) - \mathbb{E}uw)\right| \leq \frac{\alpha(\widetilde{u})\alpha(\widetilde{w})}{r^2} \cdot c(L)\theta r^2$$
$$= c(L)\theta\alpha(\widetilde{u})\alpha(\widetilde{w}).$$

But

$$\frac{\alpha(\widetilde{u})\alpha(\widetilde{w})}{r^2}\mathbb{E}uw = \mathbb{E}\widetilde{u}\widetilde{w}, \text{ and } \frac{\alpha(\widetilde{u})\alpha(\widetilde{w})}{r^2}\Psi_{\mathbb{Q}}(u,w) = \Psi_1(\widetilde{u},\widetilde{w}).$$

Thus, for every $\widetilde{u}, \widetilde{w} \in F - F$,

$$\begin{aligned} |\Psi_1(\widetilde{u},\widetilde{w}) - \mathbb{E}\widetilde{u}\widetilde{w}| &\leq c_1(L,\eta)\theta\alpha(\widetilde{u})\alpha(\widetilde{w}) \\ &= c_1(L,\eta)\theta\max\{r,\|\widetilde{u}\|_{L_2}\}\cdot\max\{r,\|\widetilde{w}\|_{L_2}\} \end{aligned}$$

A similar analysis for the other two terms, (keeping in mind the $v-v_j$ is not scaled-down), shows that

$$|\Psi_2(\widetilde{u}, v) - \mathbb{E}\widetilde{u}(v - v_j)| \le c_2(L, \eta)\theta r \cdot \max\{r, \|\widetilde{u}\|_{L_2}\} \quad \text{and}$$
$$|\Psi_3(\widetilde{u}, v) - \mathbb{E}\widetilde{u}(v_j - Y)| \le c_2(L, \eta)\theta r \cdot \max\{r, \|\widetilde{u}\|_{L_2}\}\frac{r_0 + \|v - Y\|_{L_2}}{\max\{r, \sigma\}}.$$

The proof follows because

$$\mathbb{E}\widetilde{u}(\widetilde{w}+2(v-Y)) = \mathbb{E}\widetilde{u}\widetilde{w}+2\mathbb{E}\widetilde{u}(v-v_j)+2\mathbb{E}\widetilde{u}(v_j-Y).$$

Acknowledgement: This research was funded in whole or in part by the Austrian Science Fund (FWF) [doi: 10.55776/P34743 and 10.55776/ESP31] and the Austrian National Bank [Jubiläumsfond, project 18983].

References

- P. Abdalla and N. Zhivotovskiy. Covariance estimation: Optimal dimension-free guarantees for adversarial corruption and heavy tails. *Journal of the European Mathematical Society*, arXiv:2205.08494v3, to appear.
- [2] N. Alon, Y. Matias, and M. Szegedy. The space complexity of approximating the frequency moments. Journal of Computer and System Sciences, 58(1):137–147, 1999.
- [3] M. Anthony and P. L. Bartlett. Neural network learning: theoretical foundations. Cambridge University Press, Cambridge, 1999.
- [4] S. Artstein-Avidan, A. Giannopoulos, and V. Milman. Asymptotic Geometric Analysis, Part I, volume 202. American Mathematical Society, 2015.
- [5] D. Bartl and S. Mendelson. Uniform mean estimation via generic chaining. preprint.
- [6] P. Bartlett, O. Bousquet, and S. Mendelson. Localized Rademacher complexities. Ann. Statist., 33:1497–1537, 2005.
- [7] P. Bühlmann and S. van de Geer. Statistics for high-dimensional data. Springer Series in Statistics. Springer, Heidelberg, 2011. Methods, theory and applications.
- [8] O. Catoni. Challenging the empirical mean and empirical variance: a deviation study. 48(4):1148–1185, 2012.
- [9] L. Devroye, L. Györfi, and G. Lugosi. A probabilistic theory of pattern recognition, volume 31 of Applications of Mathematics (New York). Springer-Verlag, New York, 1996.
- [10] L. Devroye, M. Lerasle, G. Lugosi, and R. Oliveira. Sub-gaussian mean estimators. Annals of Statistics, 44:2695–2725, 2016.

- [11] L. Györfi, M. Kohler, A. Krzyżak, and H. Walk. A Distribution-Free Theory of Nonparametric Regression. Springer, New York, 2002.
- [12] M. Jerrum, L. Valiant, and V. Vazirani. Random generation of combinatorial structures from a uniform distribution. *Journal of Computer and System Sciences*, 43(1-2):169–188, 1986.
- [13] V. Koltchinskii. Oracle inequalities in empirical risk minimization and sparse recovery problems, volume 2033 of Lecture Notes in Mathematics.
- [14] M. Ledoux and M. Talagrand. Probability in Banach Spaces: isoperimetry and processes, volume 23. Springer Science & Business Media, 1991.
- [15] G. Lugosi and S. Mendelson. Mean estimation and regression under heavy-tailed distributions: A survey. Foundations of Computational Mathematics, 19(5):1145–1190, 2019.
- [16] G. Lugosi and S. Mendelson. Risk minimization by median-of-means tournaments. Journal of the European Mathematical Society, 22(3):925–965, 2019.
- [17] G. Lugosi and S. Mendelson. Robust multivariate mean estimation: The optimality of trimmed mean. The Annals of Statistics, 49(1):393 – 410, 2021.
- [18] P. Massart. Concentration inequalities and model selection. Ecole d'été de Probabilités de Saint-Flour 2003. Lecture Notes in Mathematics. Springer, 2006.
- [19] S. Mendelson. Empirical processes with a bounded ψ_1 diameter. Geometric and Functional Analysis, 20(4):988–1027, 2010.
- [20] S. Mendelson. Learning without concentration. Journal of the ACM, 62:21, 2015.
- [21] S. Mendelson. Upper bounds on product and multiplier empirical processes. Stochastic Processes and their Applications, 126(12):3652–3680, 2016.
- [22] S. Mendelson. "Local" vs. "global" parameters—breaking the Gaussian complexity barrier. The Annals of Statistics, 45(5):1835 – 1862, 2017.
- [23] S. Mendelson. An unrestricted learning procedure. Journal of the ACM (JACM), 66(6):1–42, 2019.
- [24] S. Minsker. Geometric median and robust estimation in Banach spaces. Bernoulli, 21:2308–2335, 2015.
- [25] A. S. Nemirovskij and D. B. Yudin. Problem complexity and method efficiency in optimization. 1983.
- [26] M. Talagrand. Regularity of gaussian processes. Acta Mathematica, 159:99–149, 1987.
- [27] M. Talagrand. Upper and lower bounds for stochastic processes: decomposition theorems, volume 60. Springer Nature, 2022.
- [28] A. Tsybakov. Introduction to nonparametric estimation. Springer Series in Statistics.
- [29] S. van de Geer. Applications of empirical process theory, volume 6 of Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge, 2000.
- [30] V. Vapnik and A. Chervonenkis. Theory of Pattern Recognition. Nauka, Moscow, 1974. (in Russian); German translation: Theorie der Zeichenerkennung, Akademie Verlag, Berlin, 1979.

APPENDIX A. THE ENTROPY ESTIMATE

Recall that $d_F = \operatorname{diam}(F, L_2)$ and that for $\kappa \in (0, 1)$,

$$r^{*}(\kappa) = \inf\left\{r > 0: \mathbb{E}\sup_{h \in (F-F)\cap rD} G_{h} \le \kappa\sqrt{N}r\min\left\{1, \frac{r}{\sigma}\right\}\right\},\$$
$$\lambda^{*}(\kappa) = \inf\left\{r > 0: \log\mathcal{M}(F, rD) \le c_{1}N\min\left\{1, \frac{r^{2}}{\sigma^{2}}\right\}\right\}, \text{ and }$$
$$\tilde{r}^{*}(\kappa) = \inf\left\{r > 0: \mathbb{E}\sup_{h \in (F-F)\cap rD} G_{h} \le \kappa\sqrt{N}\theta(r)r\min\left\{1, \frac{r}{\sigma}\right\}\right\},$$

where $\theta(r) = 1/\max\{1, \sqrt{\log(2d_F/r)}\}.$

Let us prove the claim made previously—that these quantities are not too far from each other. To that end recall that trivially $\lambda^*(\kappa) \leq 2d_F$ for any κ because $\mathcal{M}(F, 2d_F D) = 1$. Hence, the focus is only on cases when $r^* \leq 2d_F$.

Lemma A.1. There is an absolute constant c such that for every $\kappa \in (0,1)$, $\lambda^*(\kappa) \leq \tilde{r}^*(c\kappa)$. Moreover, if $r^*(c\kappa) \leq 2d_F$ satisfies

$$r^*(c\kappa)\sqrt{\log\left(\frac{4d_F}{r^*(c\kappa)}\right)} \le \sigma$$

then

$$\lambda^*(\kappa) \le r^*(c\kappa) \sqrt{\log\left(\frac{4d_F}{r^*(c\kappa)}\right)}.$$

Proof. For every s, t > 0, set

$$H(s,t) = \sup_{f \in F} \mathcal{N}(F \cap (f+sD), tD),$$

i.e. H(s,t) is the t covering number with respect to the L_2 norm of the 'largest' localization of F of radius s. Using that F is convex, one may show that

$$\log H(s,t) \le c_0 \log \left(\frac{2s}{t}\right) \cdot \log H(4t,t),$$

see [22, Lemma 3.1]. Therefore, setting $s = d_F$,

$$\log \mathcal{N}(F, tD) \le c_0 \log \left(\frac{2d_F}{t}\right) \cdot \log H(4t, t).$$

Next, note that for every $f \in F$, if $h \in F \cap (f + 4tD)$, then h = f + (h - f), and $h - f \in (F - F) \cap 4tD$. Therefore,

$$F \cap (f + 4tD) \subset f + \left[(F - F) \cap 4tD \right],$$

implying that

$$\mathcal{N}(F \cap (f + 4tD), tD) \le \mathcal{N}((F - F) \cap 4tD, tD)$$

By Sudakov's inequality (see, e.g., [14])

$$\log \mathcal{N}((F-F) \cap 4tD, tD) \le c_1 \left(\frac{\mathbb{E}\sup_{h \in (F-F) \cap 4tD} G_h}{t}\right)^2,$$

and thus,

(A.1)
$$\log \mathcal{N}(F, tD) \le c_2 \log \left(\frac{2d_F}{t}\right) \cdot \left(\frac{\mathbb{E}\sup_{h \in (F-F) \cap 4tD} G_h}{t}\right)^2.$$

The claim that $\lambda^*(\kappa) \leq \tilde{r}^*(c\kappa)$ follows from (A.1), the definitions of λ^* and \tilde{r}^* , and the standard relations between packing numbers and covering numbers.

As for the other claim, let $r^*(c\kappa) \leq 2d_F$ satisfy that

$$r^*(c\kappa)\sqrt{\log\left(\frac{4d_F}{r^*(c\kappa)}\right)} \le \sigma$$

Set $r = 4r^*(c\kappa)\sqrt{\log(\frac{4d_F}{r^*(c\kappa)})}$ so that $r \leq 4\sigma$. Moreover, $\frac{1}{2}r \geq r^*(c\kappa)$, and hence $\mathcal{M}(F, rD) \leq \mathcal{N}(F, r^*(c\kappa)D)$. Thus, by (A.1) and the definition of $r^*(c\kappa)$,

$$\log \mathcal{N}(F, r^*(c\kappa)D) \le c_2 \log \left(\frac{2d_F}{r^*(c\kappa)}\right) \cdot (c\kappa)^2 N \frac{(r^*(c\kappa))^2}{\sigma^2}$$
$$\le c_3(c\kappa)^2 N \frac{r^2}{\sigma^2} \le 16c_3(c\kappa)^2 N \min\left\{1, \frac{r^2}{\sigma^2}\right\},$$

implying that $\lambda^*(\kappa) \leq r$, as claimed.

Appendix B. Some of the weaknesses of the Rademacher complexities

The one obvious advantage in a complexity parameter that depends on the expected supremum of the gaussian process is that it is determined solely by the L_2 -structure of the indexing class. In contrast, the Rademacher complexities depend on the Rademacher averages, that is, the expected supremum of a Bernoulli process indexed by localizations of the random set $\{f(X_i)_{i=1}^N : f \in F\}$, making them inherently harder to handle. On top of that, the expected supremum of the gaussian process is likely to be much smaller than the corresponding Rademacher averages. While (under minimal assumptions of H), we have that

$$\mathbb{E}\sup_{h\in H}\frac{1}{\sqrt{N}}\sum_{i=1}^{N}\varepsilon_{i}h(X_{i})\to\mathbb{E}\sup_{h\in H}G_{h},$$

the gap between the two expectations can be substantial for a fixed value of N—especially in heavy-tailed situations, as the next example shows.

Example B.1. Fix k to be specified in what follows. Let $\tilde{x} = \varepsilon |\tilde{x}|$ be a symmetric random variable for ε that is a random sign, independent of $|\tilde{x}|$ and $|\tilde{x}|$ that takes the values $k^{1/4}$ and 1. If

$$\mathbb{P}(|\tilde{x}| = k^{1/4}) = \frac{1}{k}$$
 and $\mathbb{P}(|\tilde{x}| = 1) = 1 - \frac{1}{k}$,

then $1 \leq \|\tilde{x}\|_{L_2} \leq 2$ and $\|\tilde{x}\|_{L_4} \leq 2$. Thus, setting $x = \varepsilon \frac{|\tilde{x}|}{c_0}$ for a suitable absolute constant $1 \leq c_0 \leq 2$, it follows that x is symmetric; has variance 1; satisfies that $\|x\|_{L_4} \leq 2$; and takes the values $\pm 1/c_0$ and $\pm k^{1/4}/c_0$.

Let $x_1, ..., x_d$ be independent copies of x and let $X = (x_1, ..., x_d)$. Using the independence and symmetry of the coordinates of X, it is standard to verify that X is isotropic and satisfies $L_4 - L_2$ norm equivalence with an absolute constant L.

Let $T = B_1^d$ and recall that if G is the standard gaussian random vector in \mathbb{R}^d , then

$$\mathbb{E}\sup_{t\in B_1^d} \langle G, t \rangle = \mathbb{E}\max_{1 \le j \le d} |g_j| \sim \sqrt{\log d}$$

On the other hand, if $X_1, ..., X_N$ are independent copies of X, then

$$\mathbb{E}\sup_{t\in B_1^d} \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i \langle X_i, t \rangle = \mathbb{E}\max_{1\leq j\leq d} |U_j|,$$

where

$$U_j = \frac{1}{\sqrt{N}} \sum_{i=1}^N \langle X_i, e_j \rangle = \frac{1}{\sqrt{N}} \sum_{i=1}^N x_{ij}.$$

Let $U = \frac{1}{\sqrt{N}} \sum_{i=1}^{N} x_i$, and thus the random variables $(U_j)_{j=1}^d$ are independent and distributed according to U. It is straightforward to verify that if $\mathbb{P}(|U| \ge M) \ge \frac{10}{d}$, then $\mathbb{E} \max_{1 \le j \le d} |U_j| \ge c_1 M$ for an absolute constant c_1 .

To identify a suitable M, fix a constant c_2 and consider the event

$$\mathcal{A} = \left\{ \text{There is a single index } 1 \le s \le N \text{ s.t. } |x_s| = \frac{k^{1/4}}{c_0} \text{ and } \left| \sum_{i=1}^N \varepsilon_i \right| \le c_2 \sqrt{N} \right\}.$$

On that event,

$$\left|\sum_{i=1}^{N} x_i\right| \ge |x_s| - \left|\sum_{i \ne s} \varepsilon_i \cdot \frac{1}{c_0}\right| \ge |x_s| - \frac{1}{c_0} \left(\left|\sum_{i=1}^{N} \varepsilon_i\right| + 1\right) \ge \frac{1}{c_0} (k^{1/4} - c_3 \sqrt{N}).$$

Therefore, if $k^{1/4} \ge 2c_3\sqrt{N}$ and $\mathbb{P}(\mathcal{A}) \ge \frac{10}{d}$, then $\mathbb{E}\max_{1\le j\le d} |U_j| \ge c_4\frac{k^{1/4}}{\sqrt{N}}$.

To control $\mathbb{P}(\mathcal{A})$, observe that by independence and for c_2 sufficiently large,

$$\mathbb{P}(\mathcal{A}) \ge \frac{N}{k} \cdot \left(1 - \frac{1}{k}\right)^{N-1} \cdot \frac{1}{2} \ge \frac{10}{d},$$

provided that $N \leq k \leq c_5 N d$.

With that in mind, let $\alpha > 1$ and set $N = d^{1/\alpha}$ and $k \sim N^{1+\alpha}$. As all the conditions are verified with these choices, we have that

$$\mathbb{E} \sup_{t \in B_1^d} \frac{1}{\sqrt{N}} \sum_{i=1}^N \varepsilon_i \langle X_i, t \rangle \ge c_4 \frac{k^{1/4}}{\sqrt{N}} \gtrsim N^{\frac{1}{4}(\alpha-1)},$$

which is much bigger than $\mathbb{E}\sup_{t\in B_1^d} \langle G,t\rangle \sim \sqrt{\log d} = \sqrt{\alpha \log N}.$

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIENNA, AUSTRIA *Email address*: daniel.bartl@univie.ac.at

DEPARTMENT OF MATHEMATICS, ETH ZURICH, SWITZERLAND *Email address*: shahar.mendelson@gmail.com