Dimension-free bounds in high-dimensional linear regression via error-in-operator approach

Fedor Noskov*

Nikita Puchkin[†]

Vladimir Spokoiny [‡]

Abstract

We consider a problem of high-dimensional linear regression with random design. We suggest a novel approach referred to as error-in-operator which does not estimate the design covariance Σ directly but incorporates it into empirical risk minimization. We provide an expansion of the excess prediction risk and derive non-asymptotic dimension-free bounds on the leading term and the remainder. This helps us to show that auxiliary variables do not increase the effective dimension of the problem, provided that parameters of the procedure are tuned properly. We also discuss computational aspects of our method and illustrate its performance with numerical experiments.

Contents

I	Introduction	3
2	Statistical properties of the error-in-operator estimate	6
3	Computational aspects	11
4	Numerical experiments	13
5	Proof of Theorem 2.3	13
6	Proof of Theorem 2.4	16
A	Auxiliary results: properties of the Hessian of $\mathcal{L}(v)$ A.1 Proof of Lemma A.1 A.2 Proof of Proposition A.2 A.3 Proof of Lemma A.3 A.4 Proof of Lemma A.4	26 28 29 32 35
B	Properties of the target functional minimizerB.1Proof of Lemma B.1B.2Proof of Lemma B.2	36 38 38

^{*}HSE University, Russian Federation, fnoskov@hse.ru

[†]HSE University, Russian Federation, npuchkin@hse.ru

[‡]HSE University, Russian Federation; WIAS Berlin and Humboldt University, Germany, spokoiny@wias-berlin.de

	B.3 Proof of Lemma B.3	41
	B.4 Proof of Lemma B.4	44
~		
С	Results from the proof of Theorem 2.3	46
	C.1 Proof of Lemma 5.1	46
	C.2 Proof of Proposition C.1	53
	C.3 Proof of Proposition C.2	54
р	Posults from the proof of Theorem 2.4	51
ν	D 1 Preliminaries	54
	D.2 Proof of Lemma 6.1	56
	D.2 Proof of Lemma 6.2	50 60
	D.5 Froof of Lemma 6.2 \dots	62
	D.4 F1001 01 Lemma 6.4	65
	D.5 Proof of Lemma 0.4	60
	D.0 Proof of Lemma 0.5	09 70
		12
	D.8 Auxiliary results	15
	D.8.1 Proof of Lemma D.2	75
	D.8.2 Proof of Lemma D.3	76
	D.8.3 Proof of Lemma D.4	78
	D.8.4 Proof of Lemma D.5	79
F	Concentration inequalities	80
112	E 1 Proof of Theorem 2.5	80
	E.1 Proof of Theorem 2.6	83
	E.2 FIOOD OF THEOREM 2.0 \ldots	0 <i>3</i> 0 <i>1</i>
		04
F	Proof of Corollary 2.7	86
	F.1 Proof of Lemma F.1	88
G	Properties of the bias \mathbf{b}_{λ}	89
н	Proof of Lemma 3.1 and auxiliary results	90
	H.1 Proof of Lemma 3.1.	90
	H.2 Proof of Lemma H.1	91
	H.3 Proof of Lemma H.3	93
	H4 Proof of Proposition H 2	96
		20
Ι	Proof of Theorem 3.3	97

1 Introduction

Recent advances in supervised machine learning devoted to understanding of deep neural networks revealed surprising effects going beyond the classical statistics theory. In contrast to the standard intuition that a learner should search for a trade-off between approximation and estimation errors, researchers empirically observed that large interpolating rules may still have small test error. Moreover, when the number of parameters exceeds sample size the prediction risk of neural networks passes a U-shaped curve and decreases again [Zhang et al., 2017, Nakkiran et al., 2020]. A bit later it became clear that benign overfitting and double descent are not distinctive features of deep learning. Similar phenomena are ubiquitous for overparametrized models such as random forests and random feature models [Belkin et al., 2019, Mei and Montanari, 2022], kernel methods [Belkin et al., 2018, Liang and Rakhlin, 2020], and linear regression [Bartlett et al., 2020, Hastie et al., 2022] to name a few. In [Belkin et al., 2018], the authors reasonably suggested that we must study more tractable "shallow" methods better before diving into deep learning theory.

In the present paper, we consider a classical linear regression problem, where a learner aims to estimate an unknown vector $\theta^{\circ} \in \mathbb{R}^d$ from i.i.d. pairs $\{(\mathbf{X}_i, Y_i) : 1 \leq i \leq n\} \subset \mathbb{R}^d \times \mathbb{R}$ generated from the model

$$Y_i = \mathbf{X}_i^{\top} \boldsymbol{\theta}^{\circ} + \varepsilon_i, \quad 1 \leqslant i \leqslant n.$$
⁽¹⁾

We do not impose any assumptions on the ambient dimension d (in particular, it may be much larger than n) or on the structure of θ° , but we require $\mathbb{E} ||\mathbf{X}_1||^2$ to be finite. Given an independent copy (\mathbf{X}, Y) of (\mathbf{X}_1, Y_1) , the quality of an estimate $\tilde{\theta}$ is measured with its quadratic risk

$$R(\widetilde{\boldsymbol{\theta}}) = \mathbb{E}_{(\mathbf{X},Y)}(Y - \mathbf{X}^{\top}\widetilde{\boldsymbol{\theta}})^{2}$$
$$= \mathbb{E}_{(\mathbf{X},Y)}(Y - \mathbf{X}^{\top}\boldsymbol{\theta}^{\circ})^{2} + \left\|\Sigma^{1/2}(\widetilde{\boldsymbol{\theta}} - \boldsymbol{\theta}^{\circ})\right\|^{2}, \text{ where } \Sigma = \mathbb{E}\mathbf{X}\mathbf{X}^{\top} \in \mathbb{R}^{d \times d}.$$

We can rewrite (1) in a more compact vector form

$$\mathbf{Y} = \mathbb{X}^{\top} \boldsymbol{\theta}^{\circ} + \boldsymbol{\varepsilon}, \tag{2}$$

where $\mathbf{Y} = (Y_1, \dots, Y_n)^\top$, $\boldsymbol{\varepsilon} = (\varepsilon_1, \dots, \varepsilon_n)^\top$, and $\mathbb{X} \in \mathbb{R}^{d \times n}$ is a matrix with columns $\mathbf{X}_1, \dots, \mathbf{X}_n$. The problem (2) provides a convenient framework for studying complex high-dimensional phenomena. For this reason, it has recently got considerable attention. Researchers put a lot of efforts into careful and subtle analysis of popular minimum norm least squares and ridge regression estimates defined as

$$\widehat{oldsymbol{ heta}}^{(LS)} = (\mathbb{X}\mathbb{X}^{ op})^{\dagger} \mathbb{X} \mathbf{Y} \quad ext{and} \quad \widehat{oldsymbol{ heta}}^{(R)} = (\mathbb{X}\mathbb{X}^{ op} + au I_d)^{-1} \mathbb{X} \mathbf{Y},$$

respectively. Here and further in the paper, B^{\dagger} stands for the Moore-Penrose pseudoinverse of a matrix B. In [Bartlett et al., 2020, Chinot and Lerasle, 2025+, Chinot et al., 2022], the authors relied on concentration of measure and statistical learning theory and established dimension-free rates of convergence for $\hat{\theta}^{(LS)}$ depending on the eigenvalues of $\Sigma = \mathbb{E} \mathbf{X}_1 \mathbf{X}_1^{\top}$ rather than on the ambient dimension d. In the same spirit, Tsigler and Bartlett [2023] derived non-asymptotic dimension-free bounds on the squared prediction risk of $\hat{\theta}^{(R)}$ and supported observations of Kobak et al. [2020] that the optimal value of the penalization parameter τ may be negative due to the implicit regularization. Another line of research (see, for instance, [Dobriban and Wager, 2018, El Karoui, 2018, Taheri et al., 2021, Richards et al., 2021, Mahdaviyeh and Naulet, 2019, Derezinski et al., 2020, Hastie et al., 2022, Wu and Xu, 2020, Bach, 2024]) examined the least squares and ridge regression estimates through the lens of random matrix theory and studied $\hat{\theta}^{(LS)}$ and $\hat{\theta}^{(R)}$ in the socalled proportional regime, that is, when the ratio d/n tends to some constant $\gamma \in (0, \infty)$ as n approaches infinity. We would like to emphasize that the aforementioned approaches complement each other. While Bartlett et al. [2020] argue that $\mathbb{E} || \mathbf{X}_1 ||^2 = \operatorname{Tr}(\Sigma) = o(n), n \to \infty$, is a necessary condition for benign overfitting, in the proportional regime one usually assumes the opposite: $\operatorname{Tr}(\Sigma) = \Omega(n)$. However, in a recent paper [Cheng and Montanari, 2024], the authors combined resolvent analysis with concentration inequalities and obtained non-asymptotic risk expansion for the ridge regression estimate.

In the context of overparametrized models, explicit regularization has a very nice property previously unseen in the classical underparametrized setup. In [Nakkiran et al., 2021], the author noticed that optimally regularized least squares estimate can mitigate double descent. This fact is of great importance, because in the presence of double (or, more generally, multiple descent, see, for instance, [Chen et al., 2021, Meng et al., 2024]) adding new batches to the training data may lead to worse generalization error (see [Nakkiran et al., 2020, Section 7]). In the present paper, we study properties of an estimate with double regularization. Let us note that both $\hat{\theta}^{(LS)}$ and $\hat{\theta}^{(R)}$ have a form $(\check{\Sigma})^{\dagger}\mathbf{Z}$, where $\mathbf{Z} = \mathbb{X}\mathbf{Y}/n$ and $\check{\Sigma}$ is an estimate of $\Sigma = \mathbb{E}\mathbf{X}_1\mathbf{X}_1^{\top}$. Instead of estimating Σ directly, we incorporate learning of Σ into an optimization problem. To be more specific, we introduce an objective function

$$L(\boldsymbol{\upsilon}) = L(\boldsymbol{\theta}, \boldsymbol{\eta}, A) = \frac{1}{2} \|\mathbf{Z} - \boldsymbol{\eta}\|^2 + \frac{1}{2} \|\boldsymbol{\eta} - A\boldsymbol{\theta}\|^2 + \frac{\mu^2}{2} \|\widehat{\boldsymbol{\Sigma}} - A\|_{\mathrm{F}}^2 + \frac{\lambda}{2} \|\boldsymbol{\theta}\|^2,$$
(3)

where

$$\mathbf{Z} = \frac{1}{n} \mathbb{X} \mathbf{Y}, \quad \widehat{\Sigma} = \frac{1}{n} \mathbb{X} \mathbb{X}^{\top},$$

and consider a regularized empirical risk minimizer

$$\widehat{\boldsymbol{\upsilon}} = (\widehat{\boldsymbol{\theta}}, \widehat{\boldsymbol{\eta}}, \widehat{A}) \in \operatorname*{argmin}_{(\boldsymbol{\theta}, \boldsymbol{\eta}, A)} L(\boldsymbol{\theta}, \boldsymbol{\eta}, A).$$
(4)

Similar ideas were used in linear inverse problems [Hoffmann and Reiss, 2008, Trabs, 2018] and Bayesian inference [Spokoiny, 2023]. In the context of kernel ridge regression, Chen et al. [2023] replaced the standard Euclidean distances ||x-x'|| by $||\Sigma^{1/2}(x-x')||$ and considered a family of Gram matrices parametrized by Σ . However, this is very different from our setup, because, in contrast to [Hoffmann and Reiss, 2008, Trabs, 2018, Chen et al., 2023], we do not assume any low-dimensional parametrization of the operator $A \in \mathbb{R}^{d \times d}$. This means that \hat{v} estimates a parameter $v^{\circ} = (\theta^{\circ}, \Sigma \theta^{\circ}, \Sigma)$ of much larger dimension than d. At the first glance, we dramatically increased the problem complexity. However, let us recall that we are interested in the target parameter $\theta^{\circ} \in \mathbb{R}^d$ rather than in the whole vector $v^{\circ} \in \mathbb{R}^{2d+d^2}$. In the simple case $\mu = +\infty$, it is straightforward to check that the estimate $\hat{\theta}$ corresponds to the plug-in one

$$\widehat{\boldsymbol{\theta}}_{\infty} = (\widehat{\Sigma}^2 + 2\lambda I_d)^{-1}\widehat{\Sigma}\mathbf{Z}, \quad \text{where} \quad \mathbf{Z} = \frac{1}{n}\mathbb{X}\mathbf{Y} \quad \text{and} \quad \widehat{\Sigma} = \frac{1}{n}\mathbb{X}\mathbb{X}^{\top} = \frac{1}{n}\sum_{i=1}^{n}\mathbf{X}_i\mathbf{X}_i^{\top},$$

which is nothing but the ridge regression estimate applied to the model

$$\mathbf{Z} = \widehat{\Sigma} \boldsymbol{\theta}^{\circ} + \mathbf{U}, \quad \text{where} \quad \mathbf{U} = \frac{1}{n} \mathbb{X} \boldsymbol{\varepsilon}.$$
 (5)

Hence, if we take the parameter μ sufficiently large, the rate of convergence of $\hat{\theta}$ will not suffer (though a rigorous proof of this fact is quite challenging). Moreover, additional regularization with respect to operator helps $\hat{\theta}$ to be less susceptible to double descent. Following [Spokoiny, 2023], we call $\hat{\theta}$ error-in-operator estimate.

Contribution. The main contributions can be summarized as follows.

- We suggest a novel approach for the problem of random design linear regression. We specify leading terms in the bias and variance expansions of the error-in-operator estimate $\hat{\theta}$ and derive an explicit high-probability upper bound on the remainder. In contrast to the standard ridge regression estimate, $\hat{\theta}$ does not have a simple closed-form expression. For this reason, we have to use more sophisticated tools for its analysis.
- We establish non-asymptotic dimension-free bounds on the excess quadratic risk $R(\hat{\theta}) R(\theta^{\circ}) = \|\Sigma^{1/2}(\hat{\theta} \theta^{\circ})\|^2$ and show that the influence of the introduced operator A is negligible once the parameter μ is tuned properly.
- We discuss minimization the non-convex objective (3) and prove that a simple alternating optimization procedure converges to the error-in-operator estimate $\hat{\theta}$.
- We illustrate the validity of statistical properties of $\hat{\theta}$ with numerical experiments. We also show that the generalization error of the error-in-operator estimate is smaller than the ones of $\hat{\theta}^{(R)}$ and $\hat{\theta}_{\infty}$ and that the risk of $\hat{\theta}$ is less susceptible to double descent.

Notation. Throughout the paper, $||B||_F$ and ||B|| stand for the Frobenius and the operator norm of B, respectively. If the matrix B is symmetric positive semidefinite and $B \neq O$, where O is the matrix with zero entries, we denote its effective rank by

$$\mathbf{r}(B) = \mathrm{Tr}(B) / \|B\|.$$

For any matrices $A \in \mathbb{R}^{p \times q}$ and $B \in \mathbb{R}^{r \times s}$, their Kronecker product $A \otimes B$ is a matrix of size $pr \times qs$, defined as

$$\begin{pmatrix} a_{11}B & \dots & a_{1q}B \\ \vdots & \ddots & \vdots \\ a_{p1}B & \dots & a_{pq}B. \end{pmatrix}$$

For any matrix $U \in \mathbb{R}^{p \times q}$ with columns $\mathbf{u}_1, \ldots, \mathbf{u}_q$, the vectorization operator is given by $\mathbf{vec}(U) = (\mathbf{u}_1^\top, \ldots, \mathbf{u}_q^\top)^\top \in \mathbb{R}^{pq}$. Here and further in the paper, the bold font is reserved for vectors, while matrices and scalars are displayed in regular font. The operator ∇_A stands for the gradient with respect to $\mathbf{vec}(A^\top)$. Thus, if $\mathbf{a}_1, \ldots, \mathbf{a}_d$ are the rows of A and f(A) is a smooth function, then $\nabla_A f = (\nabla_{\mathbf{a}_1} f^\top, \ldots, \nabla_{\mathbf{a}_d} f^\top)^\top \in \mathbb{R}^{d^2}$. For a random variable ξ and a random vector $\boldsymbol{\nu}$, their Orlicz ψ_s -norms, $s \ge 1$, are defined as

$$\|\xi\|_{\psi_s} = \inf\left\{t > 0: \mathbb{E}e^{|\xi|^s/t^s} \leqslant 2\right\} \quad \text{and} \quad \|\boldsymbol{\nu}\|_{\psi_s} = \sup_{\|\mathbf{u}\|=1} \|\mathbf{u}^\top \boldsymbol{\nu}\|_{\psi_s}.$$

The expressions $(a \lor b)$ and $(a \land b)$ denote $\max\{a, b\}$ and $\min\{a, b\}$, respectively. Sometimes, instead of the standard \mathcal{O} notation, we use $f \leq g$ or $g \geq f$, which mean that there is a universal constant c > 0, such that $f \leq cg$.

Paper organization. The rest of the paper is organized as follows. In Section 2, we present our main results concerning statistical properties of the error-in-operator estimate $\hat{\theta}$. In Section 3, we discuss computational aspects of optimizing the non-convex objective (3). We illustrate performance of the suggested procedure with numerical experiments in Section 4. Sections 5 and 6 are devoted to the proofs of bias and variance expansions of the estimate $\hat{\theta}$. Derivations of numerous auxiliary results are moved to appendix.

2 Statistical properties of the error-in-operator estimate

In the present section, we discuss main statistical properties of the error-in-operator estimate (4). In particular, we specify leading terms in expansion of $\hat{\theta}$ and provide an upper bound on its quadratic risk. We assume that the covariates $\mathbf{X}, \mathbf{X}_1, \dots, \mathbf{X}_n$ meet the following requirements.

Assumption 2.1. There exists a positive constant C_X such that, for any matrix M and any λ satisfying the inequality $C_X|\lambda| \leq 1/||M||_{\rm F}$, it holds that

$$\log \mathbb{E} \exp \left\{ \lambda (\Sigma^{-1/2} \mathbf{X})^\top M (\Sigma^{-1/2} \mathbf{X}) \right\} \leqslant \lambda \operatorname{Tr}(M) + C_X^2 \lambda^2 \|M\|_{\mathrm{F}}^2, \quad \text{where} \quad \Sigma = \mathbb{E} \mathbf{X} \mathbf{X}^\top.$$

In [Puchkin and Rakhuba, 2024], the authors imposed the same assumption on the distribution of X_1, \ldots, X_n X_n in the problem of covariance estimation, and an equivalent condition appeared in [Puchkin et al., 2025]. We refer a reader to these two papers for a comprehensive list of distributions satisfying Assumption 2.1. In particular, it is fulfilled when X has a Gaussian distribution or when $\Sigma^{-1/2}X$ has independent sub-Gaussian components. Such conditions often appear in the context of high-dimensional linear regression (see, for instance, [Bartlett et al., 2020, Nakkiran et al., 2021, Chinot and Lerasle, 2025+, Bach, 2024]). More interesting examples include situations when X has a convex concentration property, like in [Cheng and Montanari, 2024], or when the underlying distribution meets the logarithmic Sobolev inequality (such as the uniform distribution on a sphere [Meng et al., 2024]). Finally, Assumption 2.1 holds in nonlinear feature models of the form $\mathbf{X} = \varphi(W\boldsymbol{\xi})$, where φ is a Lipschitz map and $\boldsymbol{\xi}$ has either a standard Gaussian distribution [Hastie et al., 2022] or the uniform distribution on a unit sphere [Mei and Montanari, 2022] (the distribution of X should be considered conditionally on W). This follows from the fact that $\boldsymbol{\xi}$ with Gaussian or uniform distribution on a sphere satisfies the logarithmic Sobolev inequality. Then, due to the Herbst argument, it possesses a Lipschitz concentration property. This yields that X has the same property as well, and, hence, it satisfies the Hanson-Wright inequality (see [Adamczak, 2015]). Let us note that it is enough to check Assumption 2.1 for symmetric matrices only. Indeed, due to the convexity of the squared Frobenius norm, we have

$$\left\| 0.5M + 0.5M^{\top} \right\|_{\mathbf{F}}^{2} \leq 0.5 \|M\|_{\mathbf{F}}^{2} + 0.5\|M^{\top}\|_{\mathbf{F}}^{2} = \|M\|_{\mathbf{F}}^{2}$$

and then it holds that

$$\begin{split} \log \mathbb{E} \exp \left\{ \lambda (\Sigma^{-1/2} \mathbf{X})^\top M (\Sigma^{-1/2} \mathbf{X}) \right\} &= \log \mathbb{E} \exp \left\{ \lambda (\Sigma^{-1/2} \mathbf{X})^\top (0.5M + 0.5M^\top) (\Sigma^{-1/2} \mathbf{X}) \right\} \\ &\leqslant \lambda \mathrm{Tr} (0.5M + 0.5M^\top) + C_X^2 \lambda^2 \| 0.5M + 0.5M^\top \|_{\mathrm{F}}^2 \\ &\leqslant \lambda \mathrm{Tr} (M) + C_X^2 \lambda^2 \| M \|_{\mathrm{F}}^2. \end{split}$$

Assumption 2.1 means that the vector $\Sigma^{-1/2}\mathbf{X}$ is sub-Gaussian and fulfils the ψ_2 - L_2 -equivalence condition (see [Puchkin and Rakhuba, 2024, Lemma B.2]) often arising in high-dimensional statistics. However, we would like to mention that in [Puchkin et al., 2025, Proposition 2.3] the authors gave an example of a random vector satisfying the ψ_2 - L_2 -equivalence requirement, which does not meet Assumption 2.1. We also impose the following condition on the noise distribution.

Assumption 2.2. The *i.i.d.* pairs $\{(\mathbf{X}_i, \varepsilon_i) : 1 \leq i \leq n\}$ are such that

$$\mathbb{E}(\varepsilon_1 \mid \mathbf{X}_1) = 0, \quad \mathbb{E}\mathbf{X}_1\mathbf{X}_1^\top = \Sigma, \quad and \quad \|\Sigma^{-1/2}\mathbf{X}_1\varepsilon_1\|_{\psi_1} = \sigma < +\infty.$$

For instance, Assumption 2.2 holds if ε_1 is Gaussian (or, more generally, sub-Gaussian) and independent of \mathbf{X}_1 [Hsu et al., 2012, Nakkiran et al., 2021]. However, we do not need independence of ε_i from \mathbf{X}_i for our purposes.

Two main results of this section are devoted to bias and variance of the estimate $\hat{\theta}$ defined in (4). For this reason, let us define a population counterpart of the objective (3) given by

$$\mathcal{L}(\boldsymbol{\upsilon}) = \frac{1}{2} \|\boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ} - \boldsymbol{\eta}\|^2 + \frac{1}{2} \|\boldsymbol{\eta} - \boldsymbol{A}\boldsymbol{\theta}\|^2 + \frac{\mu^2}{2} \|\boldsymbol{\Sigma} - \boldsymbol{A}\|_{\mathrm{F}}^2 + \frac{\lambda}{2} \|\boldsymbol{\theta}\|^2$$

and introduce the best parametric fit

$$\boldsymbol{v}^* = (\boldsymbol{\theta}^*, \boldsymbol{\eta}^*, A^*) \in \operatorname*{argmin}_{\boldsymbol{v}} \mathcal{L}(\boldsymbol{v}).$$
(6)

One of the key properties of the functional $\mathcal{L}(v)$ we will often exploit in our analysis is that its gradient differs from $\nabla L(v)$ by a parameter-free vector \mathfrak{z} :

$$oldsymbol{
abla}
abla {L}(oldsymbol{v}) = oldsymbol{
abla} \mathcal{L}(oldsymbol{v}) - oldsymbol{\mathfrak{z}}, ext{ where } oldsymbol{\mathfrak{z}} = egin{pmatrix} oldsymbol{0} & \ oldsymbol{Z} - \mathbb{E} oldsymbol{Z} \ oldsymbol{vec}(\widehat{\Sigma} - \Sigma) \end{pmatrix} \in \mathbb{R}^{2d+d^2}.$$

In other words, for any \boldsymbol{v} the difference $L(\boldsymbol{v}) - \mathcal{L}(\boldsymbol{v})$ is an affine function of $(\mathbf{Z} - \mathbb{E}\mathbf{Z})$ and $(\widehat{\Sigma} - \Sigma)$. We are ready to present bias and variance expansions of the error-in-operator estimate $\widehat{\boldsymbol{\theta}}$.

Theorem 2.3. Let the parameters μ and λ be non-negative. Assume that the following inequalities hold:

$$\|\boldsymbol{\theta}^{\circ}\| \leq \frac{\mu}{420}, \quad and \quad \|\boldsymbol{\theta}^{\circ}\|\|\Sigma\| \leq \frac{\mu\sqrt{\lambda}}{168}.$$

Then the vector θ^* *defined in* (6) *satisfies*

$$\|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ - \mathbf{b}_\lambda\| \leqslant 210 \left(\frac{\|\boldsymbol{\theta}^\circ\|}{\mu} + \frac{\|\boldsymbol{\Sigma}\|\|\mathbf{b}_\lambda\|}{\mu\sqrt{\lambda}}\right) \|\mathbf{b}_\lambda\|, \quad where \quad \mathbf{b}_\lambda = -\lambda(\boldsymbol{\Sigma}^2/2 + \lambda I_d)^{-1}\boldsymbol{\theta}^\circ.$$

Theorem 2.4. Grant Assumptions 2.1 and 2.2. Let us fix an arbitrary $\delta \in (0, 1)$. Suppose that $4\mathbf{r}(\Sigma)^2 + \log(4/\delta)$ and $2^{12}(1+C_X)^2(\mathbf{r}(\Sigma) + \log(2/\delta))$ do not exceed the sample size n. Assume that the parameters $\mu \ge 112 \|\boldsymbol{\theta}^\circ\|$ and $\lambda > 0$ satisfy the inequalities

$$\|\Sigma\|\|\boldsymbol{\theta}^{\circ}\| \leqslant \frac{\mu\sqrt{\lambda}}{8\cdot 18\cdot 56}, \quad 5\sigma\|\Sigma\|^{1/2}\sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \leqslant \|\boldsymbol{\theta}^{\circ}\|(\sqrt{\lambda} \wedge \|\Sigma\|), \tag{7}$$

$$17(1+C_X)\|\Sigma\|\sqrt{\frac{4\mathbf{r}(\Sigma)+\log(2/\delta)}{n}} \leqslant \sqrt{\lambda},\tag{8}$$

and

$$\mu\sqrt{\lambda} \ge 395 \cdot 8\sigma \|\Sigma\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}, \quad \sqrt{\lambda} \ge 395 \cdot 8C_X \|\Sigma\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}. \tag{9}$$

Then, with probability at least $(1 - 13\delta/2)$ *, the vector*

$$\boldsymbol{\zeta} = \Sigma \mathbf{U} - \Sigma (\widehat{\Sigma} - \Sigma) \mathbf{b}_{\lambda} - (\widehat{\Sigma} - \Sigma) \Sigma \mathbf{b}_{\lambda} \quad \text{with} \quad \mathbf{U} = \frac{1}{n} \mathbb{X} \boldsymbol{\varepsilon} = \frac{1}{n} \sum_{i=1}^{n} \mathbf{X}_{i} \boldsymbol{\varepsilon}_{i}$$

satisfies the inequality

$$\begin{split} \left\| \Sigma^{1/2} \left(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^* - \left(\Sigma^2 + 2\lambda I_d \right)^{-1} \boldsymbol{\zeta} \right) \right\| &\leq \frac{19 \|\boldsymbol{\theta}^\circ\|}{\sqrt{3}\lambda^{3/4}} \left(1 + \sqrt{\frac{\Psi(n,\delta)}{27\lambda}} \right) \Psi(n,\delta) \\ &+ \frac{\|\boldsymbol{\theta}^\circ\|}{\lambda^{1/4}} \left(\left(\frac{17 \|\boldsymbol{\theta}^\circ\|}{\mu} \right)^2 + \frac{\|\mathbf{b}_\lambda\|}{160\mu} \right) \sqrt{\Psi(n,\delta)} \end{split}$$

where

$$\Psi(n,\delta) = 196 \left(\frac{\sigma \|\Sigma\|^{1/2}}{\|\boldsymbol{\theta}^{\circ}\|} + 2C_X \|\Sigma\|\right)^2 \cdot \frac{\mathbf{r}(\Sigma)^2 + \log(4/\delta)}{n}.$$

The proofs of Theorem 2.3 and Theorem 2.4 are postponed to Sections 5 and 6, respectively. These results ensure that

$$\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^{\circ} \approx \mathbf{b}_{\lambda} + (\Sigma^2 + 2\lambda I_d)^{-1} \left(\Sigma \mathbf{U} - \Sigma (\widehat{\Sigma} - \Sigma) \mathbf{b}_{\lambda} - (\widehat{\Sigma} - \Sigma) \Sigma \mathbf{b}_{\lambda} \right)$$
(10)

with high probability provided that the sample size n and the penalization parameter μ are large enough. It is not surprising that the leading terms in the expansion (10) correspond to the ones in decomposition of the estimate $\hat{\theta}_{\infty}$ defined as

$$(\widehat{oldsymbol{ heta}}_{\infty}, \widehat{oldsymbol{\eta}}_{\infty}) \in \operatorname*{argmin}_{(oldsymbol{ heta}, oldsymbol{\eta})} \left\{ rac{1}{2} \| \mathbf{Z} - oldsymbol{\eta} \|^2 + rac{1}{2} \| oldsymbol{\eta} - \widehat{\Sigma} oldsymbol{ heta} \|^2 + rac{\lambda}{2} \| oldsymbol{ heta} \|^2
ight\}.$$

Hence, the insertion of the operator A has no negative influence on the quadratic risk $R(\hat{\theta})$ provided that the parameter μ is sufficiently large. Theorems 2.3 and 2.4 essentially reduce the analysis of the excess risk $R(\hat{\theta}) - R(\theta^{\circ}) = \|\Sigma^{1/2}(\hat{\theta} - \theta^{\circ})\|^2$ to large deviation bounds on the right-hand side of (10), which easily follow from the next theorems.

Theorem 2.5. Grant Assumption 2.1 and let $A \in \mathbb{R}^{p \times d}$ and $B \in \mathbb{R}^{q \times d}$ be arbitrary matrices. Let us fix any $\delta \in (0, 1)$ satisfying the inequality

$$\mathbf{r}(\Sigma^{1/2}A^{\top}A\Sigma^{1/2})\,\mathbf{r}(\Sigma^{1/2}B^{\top}B\Sigma^{1/2}) + \log(2/\delta) \leqslant 4n.$$

Then, with probability at least $(1 - \delta)$ *, it holds that*

$$\|B(\widehat{\Sigma} - \Sigma)A^{\top}\|_{\mathbf{F}} \leqslant 4C_X \|A\Sigma^{1/2}\| \|B\Sigma^{1/2}\| \sqrt{\frac{\mathbf{r}(\Sigma^{1/2}A^{\top}A\Sigma^{1/2})\,\mathbf{r}(\Sigma^{1/2}B^{\top}B\Sigma^{1/2}) + \log(2/\delta)}{n}}$$

Theorem 2.6. Suppose that a sample $\{(\mathbf{X}_i, \varepsilon_i) : 1 \leq i \leq n\} \subset \mathbb{R}^d \times \mathbb{R}$ consists of *i.i.d.* pairs of random elements satisfying Assumption 2.2. Let us fix an arbitrary matrix $B \in \mathbb{R}^{q \times d}$ and any $\delta \in (0, 1)$ such that

$$\mathbf{r}(\Sigma^{1/2}B^{\top}B\Sigma^{1/2}) + \log(2/\delta) \leqslant n.$$

Then, with probability at least $(1 - \delta)$ *, it holds that*

$$\left\|\frac{1}{n}\sum_{i=1}^{n} B\mathbf{X}_{i}\varepsilon_{i}\right\| \leq 8\sigma \left\|B\Sigma^{1/2}\right\| \sqrt{\frac{\mathbf{r}(\Sigma^{1/2}B^{\top}B\Sigma^{1/2}) + \log(2/\delta)}{n}}$$

The proofs of Theorems 2.5 and 2.6 are moved to Appendix E. They rely on the PAC-Bayesian variational inequality (see, e.g., [Catoni and Giulini, 2017, Proposition 2.1]). In [Zhivotovskiy, 2024, Puchkin and Rakhuba, 2024, Abdalla and Zhivotovskiy, 2025+], the authors use similar arguments in the context of covariance estimation. Let us note that the upper bound in Theorem 2.5 agrees with the results of Bunea and Xiao [2015] and Puchkin et al. [2025] in the case $A = B = I_d$ and with Theorem 16 of Hsu et al. [2012] when $A = B = (\Sigma + \lambda I_d)^{-1/2}$.

Theorems 2.5 and 2.6 applied to the vector ζ from Theorem 2.4 help us to quantify the excess risk of the error-in-operator estimate in terms of

$$r_q(k) = \sum_{j>k} \left(\frac{\sigma_j}{\sigma_{k+1}}\right)^q, \quad k \in \{1, \dots, d\},\tag{11}$$

where $\sigma_1 \ge \sigma_2 \ge \ldots \ge \sigma_d$ denote the eigenvalues of Σ . In Appendix F, we prove the following upper bound on the norm of $\Sigma^{1/2}(\hat{\theta} - \theta^{\circ})$ with explicit constants.

Corollary 2.7. Assume the conditions of Theorem 2.3 and Theorem 2.4. Let $\sigma_1, \ldots, \sigma_d$ stand for the eigenvalues of Σ and define

$$k^* = k^*(\lambda) = \max\left\{k \in \mathbb{N} : \sigma_k^2 \ge 2\lambda\right\}.$$
(12)

Fix an arbitrary $\delta \in (0, 1)$ *such that*

$$\left(1 + \frac{2\lambda}{\|\Sigma\|^2}\right)^2 \left(k^* + r_4(k^*)\right) + \log(4/\delta) \leqslant n \quad and \quad k^* + r_2(k^*) + 0.25\log(4/\delta) \leqslant n \tag{13}$$

with $r_2(k)$ and $r_4(k)$ given by (11). Then, with probability at least $(1 - 8\delta)$, it holds that

$$\left\| \Sigma^{1/2}(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^{\circ}) \right\| \leq \left\| \Sigma^{1/2} \mathbf{b}_{\lambda} \right\| + 4 \left(2\sigma + C_X \| \Sigma^{1/2} \mathbf{b}_{\lambda} \| \right) \sqrt{\frac{k^* + r_4(k^*) + \log(4/\delta)}{n}} + \frac{2C_X \| \Sigma^{3/2} \mathbf{b}_{\lambda} \|}{\sqrt{2\lambda}} \sqrt{\frac{4k^* + 4r_2(k^*) + \log(4/\delta)}{n}} + \diamondsuit,$$
(14)

where $\mathbf{b}_{\lambda} = -\lambda (\Sigma^2/2 + \lambda I_d)^{-1} \boldsymbol{\theta}^{\circ}$ and

$$\diamond = 210 \left(\frac{\|\boldsymbol{\theta}^{\circ}\|}{\mu} + \frac{\|\boldsymbol{\Sigma}\|\|\mathbf{b}_{\lambda}\|}{\mu\sqrt{\lambda}} \right) \|\boldsymbol{\Sigma}\|^{1/2} \|\mathbf{b}_{\lambda}\| + \frac{\|\boldsymbol{\theta}^{\circ}\|}{\lambda^{1/4}} \left(\left(\frac{17\|\boldsymbol{\theta}^{\circ}\|}{\mu} \right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{160\mu} \right) \sqrt{\Psi(n,\delta)} + \frac{19\|\boldsymbol{\theta}^{\circ}\|}{\sqrt{3}\lambda^{3/4}} \left(1 + \sqrt{\frac{\Psi(n,\delta)}{\lambda}} \right) \Psi(n,\delta).$$

$$(15)$$

Corollary 2.7 gives an intuition for the choice of μ . It should be chosen in such a way that $\diamondsuit = O(1/\mu)$, $\mu \to \infty$, does not exceed the leading terms in the risk bound (14). We would like to note that the expression in the right-hand side of (14) can be simplified in view Lemma G.1 from Appendix G below. Let

$$\Sigma = \sum_{j=1}^d \sigma_j \mathbf{v}_j \mathbf{v}_j^{\mathsf{T}}$$

be the eigenvalue decomposition of Σ . For any $k \in \{1, ..., d\}$, we define $\beta_k = \mathbf{v}_k^\top \boldsymbol{\theta}^\circ$ and the projections of $\boldsymbol{\theta}$ onto the linear span of $\mathbf{v}_1, ..., \mathbf{v}_k$ and its orthogonal complement, respectively:

$$\boldsymbol{\theta}_{\leqslant k}^{\circ} = \sum_{j=1}^{k} \beta_j \mathbf{v}_j \text{ and } \boldsymbol{\theta}_{>k}^{\circ} = \boldsymbol{\theta}^{\circ} - \boldsymbol{\theta}_{\leqslant k}^{\circ}.$$

Then, under the conditions of Corollary 2.7, with probability at least $(1 - 8\delta)$ it holds that

$$\left\| \Sigma^{1/2}(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^{\circ}) \right\| \leq \left(\sigma_{k^{*}}^{2} \left\| \Sigma^{-1/2} \boldsymbol{\theta}_{\leq k^{*}}^{\circ} \right\|^{2} + \left\| \Sigma^{1/2} \boldsymbol{\theta}_{>k^{*}}^{\circ} \right\|^{2} \right)^{1/2} \left(1 + 8C_{X} \sqrt{\frac{k^{*} + r_{2}(k^{*}) + \log(4/\delta)}{n}} \right) \\ + 8\sigma \sqrt{\frac{k^{*} + r_{4}(k^{*}) + \log(4/\delta)}{n}} + \diamondsuit,$$
(16)

where \Diamond is defined in (15). The first term in the right-hand side of (16) corresponds to the bias and the random design effect while the second one reflects the noise influence. It is worth mentioning that the variance of the standard ridge regression estimate

$$\widehat{\boldsymbol{\theta}}^{(R)} = \underset{\boldsymbol{\theta}}{\operatorname{argmin}} \left\{ \left\| \mathbf{Y} - \mathbf{X}^{\top} \boldsymbol{\theta} \right\|^{2} + \tau \| \boldsymbol{\theta} \|^{2} \right\}$$
(17)

is of order (see, for instance, [Hsu et al., 2012, Cheng and Montanari, 2024, Bach, 2024])

$$\operatorname{Var}\left(\widehat{\boldsymbol{\theta}}^{(R)} \,\big| \, \mathbb{X}\right) = \mathcal{O}\left(\frac{\operatorname{Tr}\left(\Sigma^2 (\Sigma + \tau I_d)^2\right)}{n}\right) = \mathcal{O}\left(\frac{\widetilde{k} + r_2(\widetilde{k})}{n}\right), \quad n \to \infty, \tag{18}$$

where

$$\widetilde{k} = \widetilde{k}(\tau) = \max\left\{k \in \mathbb{N} : \sigma_k \ge \tau\right\}.$$

Remark 2.8. To be more precise, in [Cheng and Montanari, 2024, Bach, 2024], the variance is expressed through the effective regularization parameter $\varkappa(\tau)$. In the present paper, we focus on the case $\mathbf{r}(\Sigma) = \mathcal{O}(\sqrt{n}) = o(n)$ (see Theorem 2.4). In this situation, $\varkappa(\tau)$ and τ do not differ too much. In contrast, Cheng and Montanari [2024] are interested in the opposite scenario.

Let us note that $\tilde{k}(\tau)$ coincides with $k^*(\lambda)$ (see (12)) when $\tau^2 = 2\lambda$. This suggests us to compare the performance of the standard ridge estimate (17) with the prediction risk of the error-in-operator estimate (4) with $\lambda = \tau^2/2$. First, observe that the bound (16) yields that

$$\operatorname{Var}\left(\widehat{\boldsymbol{\theta}} \mid \mathbb{X}\right) = \mathcal{O}\left(\frac{k^* + r_4(k^*)}{n} + \diamondsuit^2\right), \quad n \to \infty.$$

Since $r_q(k)$ monotonously decreases in q > 0, $\operatorname{Var}\left(\widehat{\boldsymbol{\theta}} \mid \mathbb{X}\right)$ exhibits a better behaviour than $\operatorname{Var}\left(\widehat{\boldsymbol{\theta}}^{(R)} \mid \mathbb{X}\right)$. At the same time, the first term in the right-hand side of (16) is of the same order as the upper bound on the bias of $\widehat{\boldsymbol{\theta}}^{(R)}$ obtained in [Cheng and Montanari, 2024, Proposition 2.2]. Moreover, one can easily show that the leading bias term $\|\Sigma^{1/2}\mathbf{b}_{\lambda}\|$, $\lambda = \tau^2/2$ of the error-in-operator estimate $\widehat{\boldsymbol{\theta}}$ does not exceed $\sqrt{2}\tau \|\Sigma^{1/2}(\Sigma + \tau I_d)^{-1}\boldsymbol{\theta}^{\circ}\|$ (see Lemma G.2).

3 Computational aspects

Since the objective (3) of the error-in-operator estimate (4) is non-convex, we must elaborate on computational aspects of the suggested procedure. We defined the estimator $\hat{\theta}$ as a minimizer of a multivariate polynomial of degree 4. Optimizing a multivariate fourth-degree polynomial is intractable in general. Therefore, to find $\hat{\theta}$, we should harness the inner structure of our problem. First, we reduce the inference to a convex optimization problem. Let us fix an arbitrary ρ_0 from (0, 1/3] and define

$$\Upsilon(\rho_0) = \left\{ \boldsymbol{\upsilon} = (\boldsymbol{\theta}, \boldsymbol{\eta}, A) : (1 - \rho_0^2) \|\boldsymbol{\theta}\|^2 \leqslant \rho_0^2 \mu^2, \|\boldsymbol{\eta} - A\boldsymbol{\theta}\|^2 \leqslant \rho_0^2 \mu^2 \lambda \right\}.$$
(19)

The introduced set plays a crucial role in our analysis, because, the function $L(\theta, \eta, A)$ has a positive definite Hessian on $\Upsilon(\rho_0)$ (see Lemma A.1 below). However, note that $\Upsilon(\rho_0)$ is not convex. For this reason, we restrict our attention on its convex subset of the form (see Proposition H.4)

$$\Theta \times \mathsf{H} \times \mathsf{A} \subset \Upsilon(\rho_0),$$

where

$$\begin{split} \Theta &= \left\{ \boldsymbol{\theta} \in \mathbb{R}^d : \|\boldsymbol{\theta}\| \leqslant \rho_0 \mu \text{ and } \|\boldsymbol{\Sigma}\| \|\boldsymbol{\theta} - \boldsymbol{\theta}^*\| \leqslant 5\rho_0 \mu \sqrt{\lambda}/96 \right\}, \\ \mathsf{H} &= \left\{ \boldsymbol{\eta} \in \mathbb{R}^d : \|\boldsymbol{\eta} - \mathbf{Z}\| \leqslant \rho_0 \mu \sqrt{\lambda}/3 \right\}, \\ \mathsf{A} &= \left\{ A \in \mathbb{R}^{d \times d} : \|A\| \leqslant 3 \|\boldsymbol{\Sigma}\| + \sqrt{\lambda}/3 \right\}. \end{split}$$

Next, we show that under mild assumptions the estimate \hat{v} belongs to $\Theta \times H \times A$.

Lemma 3.1. Grant Assumptions 2.1 and 2.2 and fix an arbitrary $\delta \in (0,1)$ such that $n \ge 2^{12}(1 + C_X)^2(\mathbf{r}(\Sigma) + \log(2/\delta))$. Let $\rho_0 \le 1/8$ and let the sets Θ , H, A be as defined above. Assume that the non-negative parameters λ and μ satisfy the inequalities

$$\|\boldsymbol{\theta}^{\circ}\| \leqslant \frac{\rho_{0}\mu}{7}, \quad \|\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^{\circ}\| \leqslant \frac{\rho_{0}\mu\sqrt{\lambda}}{56\cdot 18}, \quad \frac{2^{11}\sigma\|\boldsymbol{\Sigma}\|^{1/2}}{\rho_{0}\mu}\sqrt{\frac{\mathbf{r}(\boldsymbol{\Sigma}) + \log(4/\delta)}{n}} \leqslant \frac{\lambda}{\|\boldsymbol{\Sigma}\|} \wedge \sqrt{\lambda},$$

and

$$\frac{2^{13}(1+C_X)\|\Sigma\|^2\|\boldsymbol{\theta}^{\circ}\|}{\rho_0\mu}\sqrt{\frac{4\mathbf{r}(\Sigma)+\log(2/\delta)}{n}} \leqslant \lambda$$

Then, $(\theta^*, \eta^*, A^*) \in \Theta \times H \times A$ and, with probability at least $(1 - \delta)$, the triplets $(\mathbf{0}, \mathbf{Z}, \widehat{\Sigma})$ and $(\widehat{\theta}, \widehat{\eta}, \widehat{A})$ belong to the set $\Theta \times H \times A$ as well.

Remark 3.2. Careful examination of the proof of Lemma 3.1 reveals that one can replace Assumption 2.1 with a bit milder ψ_2 - L_2 -equivalence condition.

Lemma 3.1 implies that $L(\cdot)$ possesses local convexity property, which is widely used in non-convex optimization (see the survey [Jain and Kar, 2017]). Thus, instead of (4), we can study the problem

$$L(\theta, \eta, A) \to \min_{\theta, \eta, A}$$
 subject to $(\theta, \eta, A) \in \Theta \times H \times A.$ (20)

The problem (20) could be solved using constrained convex optimization methods, if the sets Θ , H and A were observable. Unfortunately, it is not the case, but we prove below that the following procedure converges to $(\hat{\theta}, \hat{A})$:

$$\begin{aligned} \boldsymbol{\theta}_t &\in \operatorname*{argmin}_{\boldsymbol{\theta} \in \mathbb{R}^d} \mathsf{L}(\boldsymbol{\theta}, (A_{t-1}\boldsymbol{\theta} + \mathbf{Z})/2, A_{t-1}), \\ A_t &\in \operatorname*{argmin}_{A \in \mathbb{R}^{d \times d}} \mathsf{L}(\boldsymbol{\theta}_t, (A\boldsymbol{\theta}_t + \mathbf{Z})/2, A) \end{aligned}$$

The estimates θ_t and A_t admit explicit expressions, and $\hat{\eta}$ can be computed as

$$\widehat{\boldsymbol{\eta}} = \frac{1}{2} (\mathbf{Z} + \widehat{A}\widehat{\boldsymbol{\theta}}).$$

We summarize the optimization procedure in Algorithm 1.

Algorithm 1 Error-in-operator regression

Require: observations **Y**, feature matrix \mathbb{X} , regularization parameters $\mu, \lambda > 0$, number of iterations T. 1: Initialization: set $\mathbf{Z} = \mathbb{X}^{\top} \mathbf{Y}/n$, $\widehat{\Sigma} = \mathbb{X}^{\top} \mathbb{X}/n$, and $A_0 = \widehat{\Sigma}$.

- 2: for t = 1, ..., T do
- 3: Compute

$$\boldsymbol{\theta}_t = (A_{t-1}^\top A_{t-1} + 2\lambda I_d)^{-1} A_{t-1}^\top \mathbf{Z}$$

4: and

$$A_t = \mathbf{Z} \boldsymbol{\theta}_t^{ op} \left(2\mu^2 I_d + \boldsymbol{\theta}_t \boldsymbol{\theta}_t^{ op}
ight)^{-1} + \widehat{\Sigma} \left(I_d + rac{\boldsymbol{\theta}_t \boldsymbol{\theta}_t^{ op}}{2\mu^2}
ight)^{-1}.$$

5: end for

6: return θ_T .

The following theorem justifies convergence of Algorithm 1 to the error-in-operator estimate $\hat{\theta}$ defined in (4).

Theorem 3.3. Grant Assumptions 2.1 and 2.2. Fix arbitrary positive numbers $\delta < 1$ and $\rho_0 \leq 1/8$. Suppose that $n \geq 2^{12}(1 + C_X)^2 (\mathfrak{r}(\Sigma) + \log(2/\delta))$ and assume that the parameters μ and λ satisfy the conditions

$$\|\boldsymbol{\theta}^{\circ}\| \leq \rho_0 \mu/7, \quad \|\boldsymbol{\Sigma}\| \|\boldsymbol{\theta}^{\circ}\| \leq \rho_0 \mu \sqrt{\lambda}/(56 \cdot 18), \tag{21}$$

$$\lambda \ge \frac{2^{13}(1+C_X) \|\Sigma\|^2 \|\boldsymbol{\theta}^\circ\|}{\rho_0 \mu} \sqrt{\frac{4\mathbf{r}(\Sigma) + \log(2/\delta)}{n}},\tag{22}$$

and

$$\frac{\lambda}{\|\Sigma\|} \wedge \sqrt{\lambda} \ge \frac{2^{11} \sigma \|\Sigma\|^{1/2}}{\rho_0 \mu} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}.$$
(23)

Then, with probability at least $1 - \delta$, Algorithm 1 outputs θ_T such that

$$\|\boldsymbol{\theta}_T - \widehat{\boldsymbol{\theta}}\| \leq \rho_0^{2(T-1)} \left(\|\boldsymbol{\theta}^\circ\| + 8\sigma \|\boldsymbol{\Sigma}\|^{1/2} \sqrt{\frac{\mathbf{r}(\boldsymbol{\Sigma}) + \log(4/\delta)}{\lambda n}} \right).$$

Let us note that larger values of μ yield weaker conditions (21)–(23) on admissible ρ_0 . This means that the algorithm converges faster as μ increases. This is the behavior one should expect, because in the case $\mu = \infty$ the objective L(·) can be minimized explicitly in one step.

4 Numerical experiments

In this section, we illustrate our studies with numerical experiments. First, we show that Theorems 2.3,2.4 present correct leading terms in the expansion of $R(\hat{\theta}) - R(\theta^{\circ})$. We fix the following parameters of the random design linear regression problem:

$$d = 200,$$

$$\mathbf{X}_{i} = \left(k^{-1/8} \cdot \sin(\pi k \cdot \xi_{k})\right)_{k=1}^{d},$$

$$\boldsymbol{\theta}_{k}^{\circ} = k^{-3}, \quad k = 1, \dots, d,$$
(24)

where ξ_k are i.i.d. samples from Uniform[-1; 1]. It is clear, that Σ is diagonal with eigenvalues $k^{-1/4}/2$, $k = 1, \ldots, d$. We set $\varepsilon_1, \ldots, \varepsilon_n$ to be i.i.d. Gaussian random variables with the standard deviation 0.09.

For this set of parameters, we plot the dependence of ratio $\|\Sigma^{1/2}\mathbf{b}_{\lambda}\|/\|\Sigma^{1/2}(\boldsymbol{\theta}^*-\boldsymbol{\theta}^\circ)\|$ on μ, λ in Figure 1 (left). As implied by Theorem 2.3, this ratio is close to 1 for large values of μ .

To illustrate Theorem 2.4, we fix $\mu = 10^8$. We denote

$$\tilde{\boldsymbol{\zeta}} = (\Sigma^2 + 2\lambda I_d)^{-1} \boldsymbol{\zeta}$$

If Theorem 2.4 yields the correct leading term of $\Sigma^{1/2}(\hat{\theta} - \theta^*)$, then the ratio $\|\Sigma^{1/2}\tilde{\zeta}\|/\|\Sigma^{1/2}(\hat{\theta} - \theta^*)\|$ should be close to 1 for large enough *n*. We plot the dependence of $\|\Sigma^{1/2}\tilde{\zeta}\|/\|\Sigma^{1/2}(\hat{\theta} - \theta^*)\|$ on λ and *n* in Figure 1 (middle and right). To obtain smoother curves, we averaged the studied ratio over 40 generations. On the right picture, $\lambda_{opt}(n)$ is estimated by the grid search for each *n*. As one can see, our experiments agree with suggested theory.

Then, we compare the performance of our estimator $\hat{\theta}$ and the ridge estimator $\hat{\theta}^{(R)}$ defined in (17). We estimate the optimal hyperparameters μ , λ and τ by the grid search. Due to (16), the variance leading term $(\Sigma^2 + 2\lambda I_d)\zeta$ of $\hat{\theta}$ depends on the fourth powers of covariance eigenvalues, while the performance of $\hat{\theta}^{(R)}$ depends on the second powers. Thus, we expect that our estimator $\hat{\theta}$ should outperform $\hat{\theta}^{(R)}$ for Σ with heavy-tailed eigenvalues. We support these observations by Figure 2 (left plot). A reader can observe that our estimator $\hat{\theta}$ outperforms the ridge estimator $\hat{\theta}^{(R)}$ for the suggested setup (24).

Next, we study the impact of finite μ on the performance of $\hat{\theta}$ in the considered setup (24). Empirically, we show that finite μ can smooth the double descent curve. Figure 2 (right) displays the dependence of the risk $R(\hat{\theta}) - R(\theta^{\circ})$ on the sample size $n \in \{50, 100, \ldots, 500\}$. For each n and $\mu = \infty$, we find $\lambda_{opt}(n) \in \{1.3^{-40}, 1.3^{-39}, \ldots, 1.3^{39}\}$ by the grid search. Then, for each n and $\lambda \in \lambda_{opt}(n) \cdot \{10^{-6}, 10^{-4}, 10^{-2}, 1, 2\}$, we estimate $\mu_{opt}(\lambda, n)$ by the grid search on $\{2^0, 2^1, \ldots, 2^{29}\}$. We plot the dependence of the excess risk $R(\hat{\theta}) - R(\theta^{\circ})$ on n and λ for $\mu = \infty$ by the dashed line. The solid line corresponds to $\mu = \mu_{opt}(\lambda, n)$. We did not obtain the impact of finite μ on the risk curve when $\lambda \ge \lambda_{opt}(n)$, but finite μ can significantly mitigate the effect of double descent if $\lambda < \lambda_{opt}$.

5 Proof of Theorem 2.3

We start with studying the difference $v^* - v^\circ$. For any $t \in [0, 1]$, let us define $\mathbf{u}^{(t)} = (1 - t)v^\circ + tv^*$. Due to the Newton-Leibniz formula, we have

$$\boldsymbol{\nabla}\mathcal{L}(\boldsymbol{\upsilon}^*) - \boldsymbol{\nabla}\mathcal{L}(\boldsymbol{\upsilon}^\circ) = \boldsymbol{\nabla}\mathcal{L}(\mathbf{u}^{(1)}) - \boldsymbol{\nabla}\mathcal{L}(\mathbf{u}^{(0)}) = \int_0^1 \nabla^2 \mathcal{L}(\mathbf{u}^{(t)})(\boldsymbol{\upsilon}^* - \boldsymbol{\upsilon}^\circ) \, \mathrm{d}t.$$



Figure 1: Left: The ratio between the bias component $\|\Sigma^{1/2}(\theta^* - \theta^\circ)\|$ of the risk and its leading term established in Theorem 2.3. Middle and Right: The ratio between the variance component of the risk $\|\Sigma^{1/2}(\hat{\theta} - \theta^*)\|$ and its leading term established in Theorem 2.4.



Figure 2: Numerical studies of Error-in-Operator estimator $\hat{\theta}$. *Left:* The comparison of the Error-in-Operator estimator and the standard ridge estimator. *Right:* The effect of finite μ on the double descent curve.

Let us denote $H(\mathbf{u}) = \nabla^2 \mathcal{L}(\mathbf{u})$ and $\langle H \rangle = \int_0^1 H(\mathbf{u}^{(t)}) dt$ for brevity. Generally, for a matrix-valued function $f(\mathbf{u}) = f(\boldsymbol{\theta}, \boldsymbol{\eta}, A)$, we let

$$\langle f \rangle = \int_0^1 f(\mathbf{u}^{(t)}) \mathrm{d}t.$$

We are going to apply Lemma B.1 with $\rho_0 = 1/7$. To satisfy its conditions, we bound $\|\Sigma\| \|\theta^\circ\|$ as follows:

$$\|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| \leqslant \mu \sqrt{\lambda} / (7 \cdot 24) = \frac{\rho_0 \mu \sqrt{\lambda}}{24}.$$
(25)

Then, Lemma B.1 and Lemma A.1 imply that the matrix $\langle H \rangle$ is invertible. With the introduced notations we have

$$\boldsymbol{v}^* - \boldsymbol{v}^\circ = \langle H \rangle^{-1} \left(\boldsymbol{\nabla} \mathcal{L}(\boldsymbol{v}^*) - \boldsymbol{\nabla} \mathcal{L}(\boldsymbol{v}^\circ) \right).$$

Since v^* minimizes $\mathcal{L}(v)$, we obtain that

$$abla \mathcal{L}(oldsymbol{v}^*) - oldsymbol{
abla} \mathcal{L}(oldsymbol{v}^\circ) = -oldsymbol{
abla} \mathcal{L}(oldsymbol{v}^\circ)$$

It is straightforward to check that the only non-zero components of $\nabla \mathcal{L}(\boldsymbol{v}^{\circ})$ correspond to the first *d* coordinates. For this reason, we can restrict our attention on $\boldsymbol{\theta}^* - \boldsymbol{\theta}^{\circ}$. Let

$$\left\langle H\right\rangle_{\chi\chi} = \begin{pmatrix} \langle H\rangle_{\eta\eta} & \langle H\rangle_{\etaA} \\ \langle H\rangle_{A\eta} & \langle H\rangle_{AA} \end{pmatrix}$$

stand for the block of $\langle H \rangle$ corresponding to the nuisance parameter $\chi = (\eta, A)$. Due to the block-inversion formula (53), we have

$$\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ = (\langle H \rangle / \langle H \rangle_{\chi\chi})^{-1} (\boldsymbol{\nabla}_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\upsilon}^*) - \boldsymbol{\nabla}_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\upsilon}^\circ)) = -\lambda (\langle H \rangle / \langle H \rangle_{\chi\chi})^{-1} \boldsymbol{\theta}^\circ,$$
(26)

where $(\langle H \rangle / \langle H \rangle_{\chi\chi})$ is the Schur complement of the submatrix $\langle H \rangle_{\chi\chi}$. Rearranging the terms, we obtain that

$$\|\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ} - \mathbf{b}_{\lambda}\| = \|\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ} + \lambda(\Sigma^{2}/2 + \lambda I_{d})^{-1}\boldsymbol{\theta}^{\circ}\|$$

$$= \lambda \left\| \left[\left(\langle H \rangle / \langle H \rangle_{\chi\chi} \right)^{-1} (\Sigma^{2}/2 + \lambda I_{d}) - I_{d} \right] (\Sigma^{2}/2 + \lambda I_{d})^{-1}\boldsymbol{\theta}^{\circ} \right\|$$

$$\leq \lambda \left\| \left(\langle H \rangle / \langle H \rangle_{\chi\chi} \right)^{-1} (\Sigma^{2}/2 + \lambda I_{d}) - I_{d} \right\| \| (\Sigma^{2}/2 + \lambda I_{d})^{-1}\boldsymbol{\theta}^{\circ} \|$$

$$= \left\| (\langle H \rangle / \langle H \rangle_{\chi\chi})^{-1} (\Sigma^{2}/2 + \lambda I_{d}) - I_{d} \right\| \| \mathbf{b}_{\lambda} \|$$
(27)

The next lemma ensures that the rightmost expression is small compared to the norm of $\lambda (\Sigma^2/2 + \lambda I_d)^{-1} \theta^{\circ}$. For the sake of brevity, let us define $\rho = 7 \|\theta^{\circ}\|/\mu$. By the assumptions of Theorem 2.3, we have

$$\rho \leqslant 1/60. \tag{28}$$

Lemma 5.1. Suppose that the assumptions of Theorem 2.3 hold. Let $\mathfrak{m}(\cdot)$ be an arbitrary probability measure supported on [0, 1]. Define

$$\langle H \rangle^{\mathfrak{m}} = \int_0^1 H((1-t)\boldsymbol{v}^\circ + t\boldsymbol{v}^*) \mathrm{d}\mathfrak{m}(t).$$

Then it holds that

$$\left\| (\Sigma^2/2 + \lambda I_d)^{-1} (\langle H \rangle^{\mathfrak{m}} / \langle H \rangle_{\chi\chi}^{\mathfrak{m}}) - I_d \right\| \leq 15\rho + \frac{5\|\Sigma\| \|\mathbf{b}_\lambda\|}{\mu\sqrt{\lambda}}$$

Let us denote $(\Sigma^2/2 + \lambda I_d)^{-1} (\langle H \rangle / \langle H \rangle_{\chi\chi}) - I_d$ by *E*. Applying Lemma 5.1 with m equal to the uniform measure on [0, 1], we obtain that

$$||E|| \leq 15\rho + \frac{5||\Sigma|| ||\mathbf{b}_{\lambda}||}{\mu\sqrt{\lambda}} \leq 15\rho + \frac{5||\Sigma|| ||\boldsymbol{\theta}^{\circ}||}{\mu\sqrt{\lambda}} \leq \frac{1}{2},$$

where we used $\rho \leq 1/60$ and (25). It yields

$$\left\| (\langle H \rangle / \langle H \rangle_{\chi\chi})^{-1} (\Sigma^2 / 2 + \lambda I_d) - I_d \right\|$$

= $\| (I_d + E)^{-1} - I_d \| = \| E (I_d + E)^{-1} \| \leq \frac{\|E\|}{1 - \|E\|}$
 $\leq 30\rho + \frac{10 \|\Sigma\| \|\mathbf{b}_\lambda\|}{\mu\sqrt{\lambda}}.$

Substituting this bound into (27), we get that

$$\|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ - \mathbf{b}_\lambda\| \leq 30\rho \|\mathbf{b}_\lambda\| + \frac{10\|\Sigma\|\|\mathbf{b}_\lambda\|^2}{\mu\sqrt{\lambda}}.$$

6 Proof of Theorem 2.4

Before we dive into details, let us discuss the main idea of the proof. We split it into four major steps. First, we check that the conditions of Lemma 3.1 are satisfied with high probability. This yields that both \hat{v} and v^* belong to a convex set around v° where, according to Lemma A.1, the Hessian $\nabla^2 \mathcal{L}(v)$ is positive definite. Hence, if the penalization parameters μ and λ are chosen in agreement with (7), the difference $(\hat{v} - v^*)$ can be approximated by the standardized score $H_*^{-1}\mathfrak{z}$ where H_* stands for $\nabla^2 \mathcal{L}(v^*)$ (see Lemma 6.1). On the second step, we show that $\hat{\theta} - \theta^*$ admits an expansion with a leading term s_{θ}^* equal to the first d components of $H_*^{-1}\mathfrak{z}$. After that, we quantify the remainder $(\hat{\theta} - \theta^* - s_{\theta}^*)$ handling the subvectors of the standardized score corresponding to the target and the nuisance parameters separately (see Lemma 6.2 and Lemma 6.3, respectively). Summing up the bounds obtained on the second and the third steps, we provide an upper bound of the form

$$\left\|\Sigma^{1/2}(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^* - \mathbf{s}_{\boldsymbol{\theta}}^*)\right\| \lesssim \lambda^{-3/4} \left(1 + \sqrt{\frac{\Psi(n,\delta)}{27\lambda}}\right) \Psi(n,\delta)$$
⁽²⁹⁾

holding with high probability. Finally, on the fourth step we elaborate on s_{θ}^* and argue that it is close to the vector ζ defined in the statement of Theorem 2.4.

Step 1: localization. To start with, let us define

$$\rho_0 = \frac{7 \|\boldsymbol{\theta}^\circ\|}{\mu} \left(1 \vee \frac{72 \|\boldsymbol{\Sigma}\|}{\sqrt{\lambda}} \right).$$

Note that $\rho_0 \leq 1/16$, because of the condition (7) and the inequality $\mu \geq 112 \|\boldsymbol{\theta}^{\circ}\|$. Moreover, (7) implies that

$$\frac{2^{11}\sigma \|\Sigma\|^{1/2}}{\rho_0 \mu} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \leqslant \frac{2^{11}\sigma \|\Sigma\|^{1/2}\sqrt{\lambda}}{7 \cdot 72 \|\Sigma\|\|\boldsymbol{\theta}^\circ\|} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ \leqslant \frac{5\sigma \|\Sigma\|^{1/2}\sqrt{\lambda}}{\|\Sigma\|\|\boldsymbol{\theta}^\circ\|} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \leqslant \frac{\sqrt{\lambda}}{\|\Sigma\|\|\boldsymbol{\theta}^\circ\|} \cdot \|\boldsymbol{\theta}^\circ\|(\sqrt{\lambda} \wedge \|\Sigma\|) = \sqrt{\lambda} \wedge \frac{\lambda}{\|\Sigma\|}.$$

while (8) yields

$$\frac{2^{13}(1+C_X)\|\Sigma\|^2\|\boldsymbol{\theta}^\circ\|}{\rho_0\mu}\sqrt{\frac{4\mathbf{r}(\Sigma)+\log(2/\delta)}{n}}$$
$$\leqslant \frac{2^{13}(1+C_X)\|\Sigma\|^2\|\boldsymbol{\theta}^\circ\|\sqrt{\lambda}}{7\cdot72\|\Sigma\|\|\boldsymbol{\theta}^\circ\|}\sqrt{\frac{4\mathbf{r}(\Sigma)+\log(2/\delta)}{n}}$$
$$\leqslant 17(1+C_X)\|\Sigma\|\sqrt{\lambda}\cdot\sqrt{\frac{4\mathbf{r}(\Sigma)+\log(2/\delta)}{n}}\leqslant \lambda.$$

This means that the assumptions of Lemma 3.1 are satisfied. Hence, there is an event \mathcal{E}_0 of probability at least $1 - \delta$, such that \hat{v} and v^* belong to $\Theta \times H \times A$ on \mathcal{E}_0 , where

$$\begin{split} \Theta &= \left\{ \boldsymbol{\theta} \in \mathbb{R}^d : \|\boldsymbol{\theta}\| \leqslant \rho_0 \mu \text{ and } \|\boldsymbol{\Sigma}\| \|\boldsymbol{\theta} - \boldsymbol{\theta}^*\| \leqslant 5\rho_0 \mu \sqrt{\lambda}/96 \right\}, \\ \mathsf{H} &= \left\{ \boldsymbol{\eta} \in \mathbb{R}^d : \|\boldsymbol{\eta} - \mathbf{Z}\| \leqslant \rho_0 \mu \sqrt{\lambda}/3 \right\}, \\ \mathsf{A} &= \left\{ A \in \mathbb{R}^{d \times d} : \|A\| \leqslant 3 \|\boldsymbol{\Sigma}\| + \sqrt{\lambda}/3 \right\}. \end{split}$$

In the rest of the proof, we restrict our attention on the event \mathcal{E}_0 .

Step 2: expansion of $\hat{v} - v^*$. The result obtained on the previous step allows us to exploit geometric properties of $\mathcal{L}(v)$. To be more precise, Proposition H.4 implies that

$$\widehat{\boldsymbol{v}}, \boldsymbol{v}^* \in \Theta imes \mathsf{H} imes \mathsf{A} \subseteq \Upsilon(\rho_0) \quad \text{on } \mathcal{E}_0,$$

where the set $\Upsilon(\rho_0)$ is defined in (19). Then, due to Lemma A.1 and Lemma A.3, $\nabla^2 \mathcal{L}(\boldsymbol{v})$ is positive definite and smooth on the segment $\{t\hat{\boldsymbol{v}} + (1-t)\boldsymbol{v}^* : 0 \leq t \leq 1\}$. These properties play a crucial role in the proof of the following technical result.

Lemma 6.1. Let S, D, and D_0 be $(2d + d^2) \times (2d + d^2)$ block-diagonal matrices of the form

$$S = \operatorname{diag}\left(\Sigma^{1/2}, I_{d+1} \otimes O_d\right),$$

$$\mathsf{D}_*^2 = \operatorname{diag}\left((A^*)^\top A^* + \lambda I_d, 2I_d, I_d \otimes \left(\mu^2 I_d + \boldsymbol{\theta}^*(\boldsymbol{\theta}^*)^\top\right)\right),$$

$$\mathsf{D}_0^2 = \operatorname{diag}\left((A^*)^\top A^* + \lambda I_d, 2I_d, I_d \otimes \left(\mu_0^2 I_d + \boldsymbol{\theta}^*(\boldsymbol{\theta}^*)^\top\right)\right),$$
(30)

and

where $\mu_0 \in (0, \mu]$ is an arbitrary constant. Fix an arbitrary $\rho_0 \in (0, 1/16]$ and consider an event \mathcal{E} such that v^* and \hat{v} belong to a convex set $\mathcal{U} \subset \Upsilon(\rho_0)$ and $395 \|\mathbb{D}_*^{-1}\mathfrak{z}\| \leq \mu \sqrt{\lambda}$ on \mathcal{E} . Then, on this event, it holds that

$$\left\|S\big(\widehat{\boldsymbol{\upsilon}}-\boldsymbol{\upsilon}^*-H_*^{-1}\mathfrak{z}\big)\right\| \leqslant \frac{16}{\mu_0\sqrt{\lambda}} \left\|\mathsf{D}_0H_*^{-1}\mathfrak{z}\right\|^2 \left(\lambda^{-1/4} + \sqrt{\frac{\|\Sigma-A^*\|}{\lambda}}\right) \left(1 + \frac{\|\mathsf{D}_0H_*^{-1}\mathfrak{z}\|}{3\mu_0\sqrt{\lambda}}\right).$$

The proof of Lemma 6.1 is postponed to Section D.2. Let us note that the first *d* components of the vector $S(\hat{\boldsymbol{v}} - \boldsymbol{v}^* - H_*^{-1}\boldsymbol{z})$ coincide with $\Sigma^{1/2}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^* - \mathbf{s}_{\boldsymbol{\theta}}^*)$ while the other entries of $S(\hat{\boldsymbol{v}} - \boldsymbol{v}^* - H_*^{-1}\boldsymbol{z})$ are equal to zero. This yields that

$$\left\|\Sigma^{1/2}(\widehat{\boldsymbol{\theta}}-\boldsymbol{\theta}^*-\mathbf{s}_{\boldsymbol{\theta}}^*)\right\|=\left\|S(\widehat{\boldsymbol{\upsilon}}-\boldsymbol{\upsilon}^*-H_*^{-1}\mathfrak{z})\right\|.$$

Moreover, according to Theorems 2.5 and 2.6, with probability at least $(1 - 3\delta/2)$, we simultaneously have

$$\|\widehat{\Sigma} - \Sigma\|_{\mathrm{F}} \leqslant 4C_X \|\Sigma\| \sqrt{\frac{\mathbf{r}(\Sigma)^2 + \log(4/\delta)}{n}}, \quad \|(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^\circ\| \leqslant 4C_X \|\Sigma\| \|\boldsymbol{\theta}^\circ\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}, \quad (31)$$

and

$$\|\mathbf{U}\| \leqslant 8\sigma \|\Sigma\|^{1/2} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}.$$
(32)

We denote the corresponding event by \mathcal{E}_1 . Then the inequalities (9) ensure that

$$395\|\widehat{\Sigma} - \Sigma\|_{\mathrm{F}} \leqslant \frac{\sqrt{\lambda}}{2}, \quad 395\|\mathbf{U}\| \leqslant \mu\sqrt{\lambda}, \quad \text{and} \quad 395\|(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^{\circ}\| \leqslant \frac{\|\boldsymbol{\theta}^{\circ}\|\sqrt{\lambda}}{2}.$$

Taking into account the triangle inequality $\|\mathbf{Z} - \mathbb{E}\mathbf{Z}\| \leq \|\mathbf{U}\| + \|(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^{\circ}\|$, we obtain that

$$395^2 \|\mathbf{D}_*^{-1}\mathfrak{z}\|^2 \leqslant \frac{395^2}{2} \|\mathbf{Z} - \mathbb{E}\mathbf{Z}\|^2 + 395^2 \mu^2 \|\widehat{\boldsymbol{\Sigma}} - \boldsymbol{\Sigma}\|_{\mathrm{F}}^2 \leqslant \frac{\mu^2 \lambda}{2} \left(1 + \frac{\|\boldsymbol{\theta}^\circ\|}{2\mu}\right)^2 + \frac{\mu^2 \lambda}{4} \leqslant \mu^2 \lambda.$$

Thus, the conditions of Lemma 6.1 are satisfied on the intersection of \mathcal{E}_0 and \mathcal{E}_1 with $\mu_0 = \|\boldsymbol{\theta}^\circ\|\sqrt{3}$. Due to the union bound $\mathbb{P}(\mathcal{E}_0 \cap \mathcal{E}_1) \ge 1 - 5\delta/2$. Moreover, applying Lemma B.3 with $14\|\boldsymbol{\theta}^\circ\|/\mu \le 1/8$, we obtain that

$$\|A^* - \Sigma\| \leqslant \frac{\sqrt{\lambda}}{64} + \frac{5\|\Sigma\| \|\mathbf{b}_{\lambda}\|}{16\mu}.$$

Since

$$\mu\sqrt{\lambda} \geqslant 8 \cdot 18 \cdot 56 \|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| \quad \text{and} \quad \|\mathbf{b}_{\lambda}\| = \lambda \left\| \left(\frac{1}{2}\Sigma^{2} + \lambda I_{d}\right)^{-1} \boldsymbol{\theta}^{\circ} \right\| \leqslant \|\boldsymbol{\theta}^{\circ}\|,$$

it holds that

$$\|A^* - \Sigma\| \leqslant \frac{\sqrt{\lambda}}{64} + \frac{5\sqrt{\lambda}}{16 \cdot 8 \cdot 18 \cdot 56} = \frac{4\sqrt{\lambda}}{16^2} + \frac{5\sqrt{\lambda}}{16^2 \cdot 9 \cdot 56} \leqslant \frac{9\sqrt{\lambda}}{16^2}.$$

Hence, we have just proved that the remainder $(\hat{\theta} - \theta^* - \mathbf{s}^*_{\theta})$ satisfies

$$\begin{split} \left\| \Sigma^{1/2} (\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^* - \mathbf{s}_{\boldsymbol{\theta}}^*) \right\| &= \left\| S (\widehat{\boldsymbol{v}} - \boldsymbol{v}^* - H_*^{-1} \mathfrak{z}) \right\| \\ &\leqslant \frac{16}{\mu_0 \sqrt{\lambda}} \left\| \mathsf{D}_0 H_*^{-1} \mathfrak{z} \right\|^2 \left(\lambda^{-1/4} + \frac{3\lambda^{-1/4}}{16} \right) \left(1 + \frac{\left\| \mathsf{D}_0 H_*^{-1} \mathfrak{z} \right\|}{3\mu_0 \sqrt{\lambda}} \right) \\ &\leqslant \frac{19}{\mu_0 \lambda^{3/4}} \left\| \mathsf{D}_0 H_*^{-1} \mathfrak{z} \right\|^2 \left(1 + \frac{\left\| \mathsf{D}_0 H_*^{-1} \mathfrak{z} \right\|}{3\mu_0 \sqrt{\lambda}} \right) \end{split}$$
(33)

on the event $\mathcal{E}_0 \cap \mathcal{E}_1$ of probability at least $(1 - 5\delta/2)$. We proceed with the analysis of $D_0 H_*^{-1}\mathfrak{z}$.

Step 3: bound on the norm of $D_0 H_*^{-1}\mathfrak{z}$. The proof of the upper bound on $||D_0 H_*^{-1}\mathfrak{z}||$ is based on a block representation of $\nabla^2 \mathcal{L}(\boldsymbol{v})$. Let $\chi = (\boldsymbol{\eta}, A) \in \mathbb{R}^{d+d^2}$ stand for the nuisance parameter. For any $\boldsymbol{v} = (\boldsymbol{\theta}, \chi) = (\boldsymbol{\theta}, \boldsymbol{\eta}, A) \in \mathbb{R}^{2d+d^2}$, we denote

$$H_{\theta\theta} = \nabla^2_{\theta\theta} \mathcal{L}(\boldsymbol{\upsilon}), \quad H_{\theta\chi} = H_{\chi\theta}^\top = \nabla^2_{\theta\chi} \mathcal{L}(\boldsymbol{\upsilon}), \quad \text{and} \quad H_{\chi\chi} = \nabla^2_{\chi\chi} \mathcal{L}(\boldsymbol{\upsilon}).$$

With the introduced notations, the Hessian of $\mathcal{L}(v)$ is equal to

$$H = H(\boldsymbol{v}) = \begin{pmatrix} H_{\boldsymbol{\theta}\boldsymbol{\theta}} & H_{\boldsymbol{\theta}\chi} \\ H_{\chi\boldsymbol{\theta}} & H_{\chi\chi} \end{pmatrix}$$

Let $H_{\theta\theta} = H/H_{\chi\chi}$ and $H_{\chi\chi} = H/H_{\theta\theta}$ stand for the Schur complements of $H_{\chi\chi}$ and $H_{\theta\theta}$, respectively. Using the block-matrix inversion formula (see (53))

$$\begin{pmatrix} H_{\theta\theta} & H_{\theta\chi} \\ H_{\chi\theta} & H_{\chi\chi} \end{pmatrix}^{-1} = \begin{pmatrix} \breve{H}_{\theta\theta}^{-1} & -\breve{H}_{\theta\theta}^{-1}H_{\theta\chi}H_{\chi\chi}^{-1} \\ -\breve{H}_{\chi\chi}^{-1}H_{\chi\theta}H_{\theta\theta}^{-1} & \breve{H}_{\chi\chi}^{-1} \end{pmatrix}$$

and taking into account that \mathfrak{z} has a form $(\mathbf{0},\mathfrak{z}_{\chi})$, where $\mathfrak{z}_{\chi} = (\mathbf{Z} - \mathbb{E}\mathbf{Z}, \mu^2 \operatorname{vec}(\widehat{\Sigma} - \Sigma))$, we obtain that

$$H^{-1}\mathfrak{z} = \begin{pmatrix} \breve{H}_{\theta\theta}^{-1} & -\breve{H}_{\theta\theta}^{-1}H_{\theta\chi}H_{\chi\chi}^{-1} \\ -\breve{H}_{\chi\chi}^{-1}H_{\chi\theta}H_{\theta\theta}^{-1} & \breve{H}_{\chi\chi}^{-1} \end{pmatrix} \begin{pmatrix} \mathbf{0} \\ \mathfrak{z}_{\chi} \end{pmatrix} = \begin{pmatrix} -\breve{H}_{\theta\theta}^{-1}H_{\theta\chi}H_{\chi\chi}^{-1}\mathfrak{z}_{\chi} \\ \breve{H}_{\chi\chi}^{-1}\mathfrak{z}_{\chi} \end{pmatrix}$$
(34)

Then, recalling the definition of D_0 (see (30)), we observe that

$$\|\mathbb{D}_{0}H_{*}^{-1}\mathfrak{z}\|^{2} = \left\| \left(\frac{1}{2} (A^{*})^{\top}A^{*} + \lambda I_{d} \right)^{1/2} \breve{H}_{\theta\theta}^{-1}(\boldsymbol{\upsilon}^{*})H_{\theta\chi}(\boldsymbol{\upsilon}^{*})H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*})\mathfrak{z}_{\chi} \right\|^{2} + \left\|\mathbb{D}_{0\chi}\breve{H}_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*})\mathfrak{z}_{\chi}\right\|^{2}, \quad (35)$$

where we introduced

$$\mathsf{D}_{0\chi}^{2} = \operatorname{diag}\left(2I_{d}, I_{d}\otimes(\mu_{0}^{2}I_{d}+\boldsymbol{\theta}^{*}(\boldsymbol{\theta}^{*})^{\top})\right).$$

The two terms in the right-hand side of (35) should be studied separately. We start with the analysis of the subvector of $D_0 H_*^{-1}\mathfrak{z}$ corresponding to the target parameter.

Lemma 6.2. Grant Assumptions 2.1 and 2.2 and let $v \in \Upsilon(\rho)$ for some $\rho \leq 1/5$ (see (19) for the definition of $\Upsilon(\rho)$). Let $\delta \in (0, 1)$ and suppose that

$$\mathbf{r}(\Sigma) + \log(4/\delta) \leqslant n.$$

Let \mathcal{E}_1 be the event where (31) and (32) hold. Then there exists an event \mathcal{E}_2 of probability at least $1 - 5\delta/2$ such that

(i) the following inequalities hold simultaneously on \mathcal{E}_2 :

$$\|(\widehat{\Sigma} - \Sigma)(\boldsymbol{\theta} - \boldsymbol{\theta}^{\circ})\| \leq 4C_X \|\Sigma\|^{1/2} \|\Sigma^{1/2}(\boldsymbol{\theta} - \boldsymbol{\theta}^{\circ})\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}},$$
(36)

$$\|(\widehat{\Sigma} - \Sigma)(A\boldsymbol{\theta} - \boldsymbol{\eta})\| \leq 4C_X \|\Sigma\|^{1/2} \|\Sigma^{1/2}(A\boldsymbol{\theta} - \boldsymbol{\eta})\| \sqrt{\frac{\mathtt{r}(\Sigma) + \log(4/\delta)}{n}}, \tag{37}$$

$$|(A\boldsymbol{\theta}-\boldsymbol{\eta})^{\top}(\widehat{\boldsymbol{\Sigma}}-\boldsymbol{\Sigma})\boldsymbol{\theta}| \leqslant 4C_X \|\boldsymbol{\Sigma}^{1/2}\boldsymbol{\theta}\| \|\boldsymbol{\Sigma}^{1/2}(A\boldsymbol{\theta}-\boldsymbol{\eta})\| \sqrt{\frac{1+\log(4/\delta)}{n}},$$
(38)

$$|(A\boldsymbol{\theta}-\boldsymbol{\eta})^{\top}(\widehat{\boldsymbol{\Sigma}}-\boldsymbol{\Sigma})\boldsymbol{\theta}^{\circ}| \leq 4C_X \|\boldsymbol{\Sigma}^{1/2}\boldsymbol{\theta}^{\circ}\| \|\boldsymbol{\Sigma}^{1/2}(A\boldsymbol{\theta}-\boldsymbol{\eta})\| \sqrt{\frac{1+\log(4/\delta)}{n}},$$
(39)

$$\left| (A\boldsymbol{\theta} - \boldsymbol{\eta})^{\mathsf{T}} \mathbf{U} \right| \leq 8\sigma \| \Sigma^{1/2} (A\boldsymbol{\theta} - \boldsymbol{\eta}) \| \sqrt{\frac{1 + \log(4/\delta)}{n}};$$
(40)

(ii) on the intersection of \mathcal{E}_1 and \mathcal{E}_2 , one has

$$\begin{aligned} \left\| \left(\frac{1}{2} A^{\top} A + \lambda I_{d} \right)^{1/2} \breve{H}_{\theta\theta}^{-1} H_{\theta\chi} H_{\chi\chi}^{-1} \mathfrak{z}_{\chi} \right\| \\ &\leqslant \frac{1}{1 - 10\rho^{2}} \left\| \left(\frac{1}{2} A^{\top} A + \lambda I_{d} \right)^{-1/2} H_{\theta\chi} H_{\chi\chi}^{-1} \mathfrak{z}_{\chi} \right\| \\ &\leqslant \frac{2\sqrt{2} \|\Sigma\|^{1/2}}{1 - 10\rho^{2}} \left(2\sigma + C_{X} \|\Sigma^{1/2} (\theta - \theta^{\circ})\| + \frac{\sqrt{2}C_{X} \|\Sigma^{1/2} (A\theta - \eta)\|}{\sqrt{\lambda}} \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ \frac{4\sqrt{2}\rho^{2} \|\Sigma\|^{1/2}}{1 - 10\rho^{2}} \left(2\sigma + 3C_{X} \|\Sigma^{1/2} \theta^{\circ}\| \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ \frac{8\rho \|\Sigma^{1/2} (A\theta - \eta)\|}{(1 - 10\rho^{2})\mu\sqrt{\lambda}} \left(C_{X} \|\Sigma^{1/2} \theta\| + \sigma \right) \sqrt{\frac{1 + \log(4/\delta)}{n}}. \end{aligned}$$

We provide the proof of Lemma 6.2 in Section D.3. Let us elaborate on the result of this lemma when $v = v^*$. According to Lemma B.2 (see (ii) and (iv)), we have

$$||A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*|| \leq \sqrt{\lambda} ||\boldsymbol{\theta}^\circ||, \quad ||\boldsymbol{\theta}^*|| \leq ||\boldsymbol{\theta}^\circ||,$$

and

$$\|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ\| \leq \|\mathbf{b}_{\lambda}\| \left(1 + \frac{3\|\boldsymbol{\theta}^\circ\|}{\mu}\right) \leq \|\boldsymbol{\theta}^\circ\| \left(1 + \frac{3}{111}\right) = \frac{38\|\boldsymbol{\theta}^\circ\|}{37}.$$

This means that θ^* belongs to $\Upsilon(\rho^*)$ with $\rho^* = \|\theta^\circ\|/\mu \le 1/112$. Hence, there exists an event \mathcal{E}_2^* such that $\mathbb{P}(\mathcal{E}_2^*) \ge (1 - 5\delta/2)$ and the inequalities

$$\|(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^*\| \leq 4C_X \|\Sigma\|^{1/2} \|\Sigma^{1/2}\boldsymbol{\theta}^*\| \sqrt{\frac{\mathtt{r}(\Sigma) + \log(4/\delta)}{n}},\tag{41}$$

$$\|(\widehat{\Sigma} - \Sigma)(A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)\| \leq 4C_X \|\Sigma\|^{1/2} \|\Sigma^{1/2}(A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}},$$
(42)

$$|(A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)^{\top}(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^*| \leq 4C_X \|\Sigma^{1/2}\boldsymbol{\theta}^*\| \|\Sigma^{1/2}(A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)\| \sqrt{\frac{1 + \log(4/\delta)}{n}}, \tag{43}$$

$$|(A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)^{\top}(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^\circ| \leq 4C_X \|\Sigma^{1/2}\boldsymbol{\theta}^\circ\| \|\Sigma^{1/2}(A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)\| \sqrt{\frac{1 + \log(4/\delta)}{n}}, \tag{44}$$

$$\left| (A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^*)^\top \mathbf{U} \right| \leq 8\sigma \| \Sigma^{1/2} (A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^*) \| \sqrt{\frac{1 + \log(4/\delta)}{n}}$$
(45)

hold simultaneously on $\mathcal{E}_2^*.$ Moreover, on $\mathcal{E}_1\cap \mathcal{E}_2^*$ it holds that

$$\left\| \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{1/2} \breve{H}_{\theta\theta}^{-1}(\boldsymbol{v}^*) H_{\theta\chi}(\boldsymbol{v}^*) H_{\chi\chi}^{-1}(\boldsymbol{v}^*) \mathfrak{z}_{\chi} \right\|$$

$$\leq 1.001 \left\| \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1/2} H_{\theta\chi}(\boldsymbol{v}^*) H_{\chi\chi}^{-1}(\boldsymbol{v}^*) \mathfrak{z}_{\chi} \right\|$$
(46)

and

$$\begin{aligned} \left\| \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\| \\ &\leqslant 2\sqrt{2} \|\Sigma\|^{1/2} \left(2\sigma + \frac{38C_{X} \|\Sigma\|^{1/2} \|\boldsymbol{\theta}^{\circ}\|}{37} + \sqrt{2}C_{X} \|\Sigma\|^{1/2} \|\boldsymbol{\theta}^{\circ}\| \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ \frac{4\sqrt{2} \|\Sigma\|^{1/2}}{112^{2}} \left(2\sigma + 3C_{X} \|\Sigma\|^{1/2} \|\boldsymbol{\theta}^{\circ}\| \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ \frac{8 \|\Sigma\|^{1/2}}{112^{2}} \left(C_{X} \|\Sigma\|^{1/2} \|\boldsymbol{\theta}^{\circ}\| + \sigma \right) \sqrt{\frac{1 + \log(4/\delta)}{n}} \\ &\leqslant \left(4\sqrt{2} + \frac{8\sqrt{2}}{112^{2}} + \frac{8}{112^{2}} \right) \|\Sigma\|^{1/2} \left(\sigma + 2C_{X} \|\Sigma\|^{1/2} \|\boldsymbol{\theta}^{\circ}\| \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}. \end{aligned}$$

Finally, the next lemma concerns the norm of $(D_0 H_*^{-1}\mathfrak{z})_{\chi}$ and finishes the third step of the proof. Lemma 6.3. Let $\rho \leq 1/16$ and fix an arbitrary $\upsilon \in \Upsilon(\rho)$, where $\Upsilon(\rho)$ is defined in (19). Introduce

$$\widetilde{\mathsf{D}}^2 = \operatorname{diag}\left(2I_d, I_d \otimes (\widetilde{\mu}^2 I_d + \boldsymbol{\theta} \boldsymbol{\theta}^\top)\right) \in \mathbb{R}^{(d+d^2) \times (d+d^2)},$$

where $\tilde{\mu}$ is an arbitrary number from $[0, \mu]$. Finally, let \mathcal{E}_1 be the event where (31) and (32) hold. Then, on this event, we have

$$\begin{split} \left\| \widetilde{\mathsf{D}} \breve{H}_{\chi\chi}^{-1} \mathfrak{z}_{\chi} \right\| &\leqslant 8 \left(1 + \frac{9\rho}{2} \right) \sigma \|\Sigma\|^{1/2} \sqrt{\frac{\mathsf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ 4 \left(1 + \frac{9\rho}{2} \right) C_X \|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| \sqrt{\frac{\mathsf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ 4 \left(\sqrt{\widetilde{\mu}^2 + \|\boldsymbol{\theta}\|^2} + \frac{9\rho\mu}{2} \right) C_X \|\Sigma\| \sqrt{\frac{\mathsf{r}(\Sigma)^2 + \log(4/\delta)}{n}}. \end{split}$$

The proof of Lemma 6.3 is moved to Section D.4. Similarly to Lemma 6.2, we are going to apply Lemma 6.3 to $\boldsymbol{v} = \boldsymbol{v}^*$. As we discussed earlier, \boldsymbol{v}^* lies in $\Upsilon(\rho)$, where $\rho^* = \|\boldsymbol{\theta}^\circ\|/\mu \leq 1/112$. Taking $\tilde{\mu} = \mu_0 = \|\boldsymbol{\theta}^\circ\|/\sqrt{3}, \rho = \rho^* \leq 1/112$, and $\boldsymbol{v} = \boldsymbol{v}^*$, we obtain that

$$\begin{aligned} \left\| \mathsf{D}_{0\chi} \breve{H}_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \mathfrak{z}_{\chi} \right\| &\leq 8 \left(1 + \frac{9}{224} \right) \sigma \|\Sigma\|^{1/2} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ 4 \left(1 + \frac{9}{224} \right) C_X \|\Sigma\| \|\boldsymbol{\theta}^\circ\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ 4 \left(2\|\boldsymbol{\theta}^\circ\| + \frac{9\|\boldsymbol{\theta}^\circ\|}{224} \right) C_X \|\Sigma\| \sqrt{\frac{\mathbf{r}(\Sigma)^2 + \log(4/\delta)}{n}} \\ &\leq \frac{17}{2} \left(\sigma \|\Sigma\|^{1/2} + 2C_X \|\Sigma\| \|\boldsymbol{\theta}^\circ\| \right) \sqrt{\frac{\mathbf{r}(\Sigma)^2 + \log(4/\delta)}{n}}. \end{aligned}$$
(48)

on the event \mathcal{E}_1 . Here we used the bound $\|\boldsymbol{\theta}^*\| \leq \|\boldsymbol{\theta}^\circ\|$ following from the definition of $\boldsymbol{\theta}^*$ (see Lemma B.2(iv)). The inequalities (35), (46), (47), and (48) immediately imply that

$$\begin{split} \| \mathbb{D}_{0} H_{*}^{-1} \mathfrak{z} \| &\leq 1.001 \left(4\sqrt{2} + \frac{8\sqrt{2}}{112^{2}} + \frac{8}{112^{2}} \right) \| \Sigma \|^{1/2} \left(\sigma + 2C_{X} \| \Sigma \|^{1/2} \| \boldsymbol{\theta}^{\circ} \| \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ 8 \left(1 + \frac{9}{224} \right) \left(\sigma \| \Sigma \|^{1/2} + 2C_{X} \| \Sigma \| \| \boldsymbol{\theta}^{\circ} \| \right) \sqrt{\frac{\mathbf{r}(\Sigma)^{2} + \log(4/\delta)}{n}} \\ &\leq 14 \left(\sigma \| \Sigma \|^{1/2} + 2C_{X} \| \Sigma \| \| \boldsymbol{\theta}^{\circ} \| \right) \sqrt{\frac{\mathbf{r}(\Sigma)^{2} + \log(4/\delta)}{n}} \equiv \| \boldsymbol{\theta}^{\circ} \| \sqrt{\Psi(n, \delta)} \end{split}$$

on the intersection of the events \mathcal{E}_0 , \mathcal{E}_1 , and \mathcal{E}_2^* (that is, with probability at least $1 - 5\delta$). In view of (33), on the same event, we have

$$\begin{split} \left\| \Sigma^{1/2} (\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^* - \mathbf{s}_{\boldsymbol{\theta}}^*) \right\| &= \left\| S(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^* - H_*^{-1} \mathfrak{z}) \right\| \\ &\leqslant \frac{19}{\mu_0 \lambda^{3/4}} \left\| \mathsf{D}_0 H_*^{-1} \mathfrak{z} \right\|^2 \left(1 + \frac{\|\mathsf{D}_0 H_*^{-1} \mathfrak{z}\|}{3\mu_0 \sqrt{\lambda}} \right) \\ &\leqslant \frac{19 \|\boldsymbol{\theta}^\circ\|}{\sqrt{3} \lambda^{3/4}} \left(1 + \sqrt{\frac{\Psi(n, \delta)}{27\lambda}} \right) \Psi(n, \delta). \end{split}$$
(49)

and (29) follows.

Step 4: elaborating on s_{θ}^* . It remains to quantify the difference between s_{θ}^* and

$$\frac{1}{2}\left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1} \left(\Sigma(\mathbf{Z} - \mathbb{E}\mathbf{Z}) - \Sigma(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^* - (\widehat{\Sigma} - \Sigma)\Sigma\mathbf{b}_\lambda\right).$$

We perform this task in three tiny steps. First, we note that \mathbf{s}^*_{θ} can be approximated by

$$-\left(\Sigma^2+\lambda I_d\right)^{-1}H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^*)H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*)\boldsymbol{\mathfrak{z}}_{\chi}.$$

We support this claim by the following result.

Lemma 6.4. Assume that the parameters μ and λ fulfil (7) and that $\mu \ge 112 \|\boldsymbol{\theta}^{\circ}\|$. Then, on the event where (47) holds, we also have

$$\begin{split} \left\| \Sigma^{1/2} \mathbf{s}_{\boldsymbol{\theta}}^* + \Sigma^{1/2} \left(\Sigma^2 + \lambda I_d \right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \mathfrak{z}_{\chi} \right\| \\ &\leqslant \frac{\|\boldsymbol{\theta}^\circ\|}{2\lambda^{1/4}} \left(\left(\frac{17\|\boldsymbol{\theta}^\circ\|}{\mu} \right)^2 + \frac{\|\mathbf{b}_{\lambda}\|}{160\mu} \right) \sqrt{\Psi(n,\delta)}, \end{split}$$

where

$$\Psi(n,\delta) = 196 \left(\frac{\sigma \|\Sigma\|^{1/2}}{\|\boldsymbol{\theta}^{\circ}\|} + 2C_X \|\Sigma\|\right)^2 \cdot \frac{\mathbf{r}(\Sigma)^2 + \log(4/\delta)}{n}.$$

We postpone the proof of Lemma 6.4 to Section D.5 and finish the proof of Theorem 2.4 first. The next auxiliary lemmata ensure that $H_{\theta\chi}(\boldsymbol{v}^*)H_{\chi\chi}^{-1}(\boldsymbol{v}^*)\mathfrak{z}_{\chi}$ is close to

$$-\frac{1}{2}\left(\Sigma(\mathbf{Z}-\mathbb{E}\mathbf{Z})-\Sigma(\widehat{\Sigma}-\Sigma)\boldsymbol{\theta}^*-(\widehat{\Sigma}-\Sigma)\Sigma\mathbf{b}_{\lambda}\right).$$

Lemma 6.5. Assume that the parameters μ and λ fulfil (7) and that $\mu \ge 112 \|\boldsymbol{\theta}^{\circ}\|$. Then, on the event $\mathcal{E}_1 \cap \mathcal{E}_2^*$, the vector

$$\boldsymbol{\tau} = (A^*)^{\top} \mathbf{U} - (A^*)^{\top} (\widehat{\Sigma} - \Sigma) (\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) - 2(\widehat{\Sigma} - \Sigma) (A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^*)$$

satisfies the inequality

$$\left\| \Sigma^{1/2} \left(\Sigma^2/2 + \lambda I_d \right)^{-1} \left(H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \boldsymbol{\mathfrak{z}}_{\chi} + 0.5\boldsymbol{\tau} \right) \right\| \leqslant \frac{\|\boldsymbol{\theta}^\circ\|^3}{\mu^2 \lambda^{1/4}} \sqrt{\Psi(n,\delta)}.$$

Lemma 6.6. Assume that the parameters $\mu \ge 112 \|\boldsymbol{\theta}^{\circ}\|$ and $\lambda > 0$ fulfil (7) and that $4n \ge \mathbf{r}(\Sigma) + \log(4/\delta)$. Let $\boldsymbol{\tau}$ be as defined in Lemma 6.6 and let

$$\boldsymbol{\zeta} = \Sigma \mathbf{U} - \Sigma (\widehat{\Sigma} - \Sigma) \mathbf{b}_{\lambda} - (\widehat{\Sigma} - \Sigma) \Sigma \mathbf{b}_{\lambda}.$$

Then there exists an event \mathcal{E}_3 of probability at least $(1 - 3\delta/2)$ such that on $\mathcal{E}_1 \cap \mathcal{E}_3$, it holds that

$$\left\|\Sigma^{1/2}\left(\Sigma^{2}+2\lambda I_{d}\right)^{-1}\left(\boldsymbol{\tau}-\boldsymbol{\zeta}\right)\right\| \leqslant \frac{12\|\boldsymbol{\theta}^{\circ}\|}{25\lambda^{1/4}} \left(\left(\frac{17\|\boldsymbol{\theta}^{\circ}\|}{\mu}\right)^{2}+\frac{\|\mathbf{b}_{\lambda}\|}{160\mu}\right)\sqrt{\Psi(n,\delta)}.$$

The proofs of Lemma 6.5 and Lemma 6.6 are moved to Sections D.6 and D.7, respectively. Summing up Lemmata 6.4, 6.5, and 6.6, we obtain that

$$\left\| \Sigma^{1/2} \mathbf{s}_{\boldsymbol{\theta}}^{*} - \Sigma^{1/2} \left(\Sigma^{2} + 2\lambda I_{d} \right)^{-1} \boldsymbol{\zeta} \right\| \leq \left(\frac{1}{2} + \frac{12}{25} + \frac{1}{17^{2}} \right) \frac{\|\boldsymbol{\theta}^{\circ}\|}{\lambda^{1/4}} \left(\left(\frac{17\|\boldsymbol{\theta}^{\circ}\|}{\mu} \right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{160\mu} \right) \sqrt{\Psi(n,\delta)}$$

$$\leq \frac{\|\boldsymbol{\theta}^{\circ}\|}{\lambda^{1/4}} \left(\left(\frac{17\|\boldsymbol{\theta}^{\circ}\|}{\mu} \right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{160\mu} \right) \sqrt{\Psi(n,\delta)}$$
(50)

on the event $\mathcal{E}_0 \cap \mathcal{E}_1 \cap \mathcal{E}_2^* \cap \mathcal{E}_3$. Finally, the triangle inequality and the bounds (49), (50), imply that, with probability at least $(1 - 13\delta/2)$,

$$\begin{split} \left\| \Sigma^{1/2} \left(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^* - \left(\Sigma^2 + 2\lambda I_d \right)^{-1} \boldsymbol{\zeta} \right) \right\| &\leq \frac{19 \|\boldsymbol{\theta}^\circ\|}{\sqrt{3}\lambda^{3/4}} \left(1 + \sqrt{\frac{\Psi(n,\delta)}{27\lambda}} \right) \Psi(n,\delta) \\ &+ \frac{\|\boldsymbol{\theta}^\circ\|}{\lambda^{1/4}} \left(\left(\frac{17 \|\boldsymbol{\theta}^\circ\|}{\mu} \right)^2 + \frac{\|\mathbf{b}_\lambda\|}{160\mu} \right) \sqrt{\Psi(n,\delta)}. \end{split}$$

References

- P. Abdalla and N. Zhivotovskiy. Covariance estimation: Optimal dimension-free guarantees for adversarial corruption and heavy tails. Journal of the European Mathematical Society (to appear), arXiv:2205.08494, 2025+.
- R. Adamczak. A note on the Hanson-Wright inequality for random vectors with dependencies. *Electronic Communications in Probability*, 20:1–13, 2015.

- F. Bach. High-dimensional analysis of double descent for linear regression with random projections. *SIAM Journal on Mathematics of Data Science*, 6(1):26–50, 2024.
- P. L. Bartlett, P. M. Long, G. Lugosi, and A. Tsigler. Benign overfitting in linear regression. Proceedings of the National Academy of Sciences, 117(48):30063–30070, 2020.
- M. Belkin, S. Ma, and S. Mandal. To understand deep learning we need to understand kernel learning. In Proceedings of the 35th International Conference on Machine Learning, volume 80 of Proceedings of Machine Learning Research, pages 541–549. PMLR, 2018.
- M. Belkin, D. Hsu, S. Ma, and S. Mandal. Reconciling modern machine-learning practice and the classical bias-variance trade-off. *Proceedings of the National Academy of Sciences of the United States of America*, 116(32):15849–15854, 2019.
- F. Bunea and L. Xiao. On the sample covariance matrix estimator of reduced effective rank population matrices, with applications to fPCA. *Bernoulli*, 21(2):1200–1230, 2015.
- O. Catoni and I. Giulini. Dimension-free PAC-Bayesian bounds for matrices, vectors, and linear least squares regression. Preprint, arXiv:1712.02747, 2017.
- L. Chen, Y. Min, M. Belkin, and A. Karbasi. Multiple descent: Design your own generalization curve. In *Advances in Neural Information Processing Systems*, volume 34, pages 8898–8912. Curran Associates, Inc., 2021.
- Y. Chen, Y. Li, K. Liu, and F. Ruan. Kernel learning in ridge regression "automatically" yields exact low rank solution. Preprint. ArXiv:2310.11736, 2023.
- C. Cheng and A. Montanari. Dimension free ridge regression. *The Annals of Statistics*, 52(6):2879–2912, 2024.
- G. Chinot and M. Lerasle. On the robustness of the minimum ℓ_2 interpolator. Bernoulli (to appear), arXiv:2003.05838, 2025+.
- G. Chinot, M. Löffler, and S. van de Geer. On the robustness of minimum norm interpolators and regularized empirical risk minimizers. *The Annals of Statistics*, 50(4):2306–2333, 2022.
- M. Derezinski, F. T. Liang, and M. W. Mahoney. Exact expressions for double descent and implicit regularization via surrogate random design. *Advances in neural information processing systems*, 33:5152–5164, 2020.
- E. Dobriban and S. Wager. High-dimensional asymptotics of prediction: Ridge regression and classification. *The Annals of Statistics*, 46(1):247–279, 2018.
- N. El Karoui. On the impact of predictor geometry on the performance on high-dimensional ridgeregularized generalized robust regression estimators. *Probability Theory and Related Fields*, 170:95–175, 2018.
- T. Hastie, A. Montanari, S. Rosset, and R. J. Tibshirani. Surprises in high-dimensional ridgeless least squares interpolation. *Annals of statistics*, 50(2):949, 2022.
- M. Hoffmann and M. Reiss. Nonlinear estimation for linear inverse problems with error in the operator. *The Annals of Statistics*, 36(1):310–336, 2008.
- D. Hsu, S. M. Kakade, and T. Zhang. Random design analysis of ridge regression. In *Proceedings of the* 25th Annual Conference on Learning Theory, volume 23 of Proceedings of Machine Learning Research,

pages 9.1-9.24. PMLR, 2012.

- P. Jain and P. Kar. Non-convex Optimization for Machine Learning. *Foundations and Trends*® *in Machine Learning*, 10(3-4):142–336, 2017.
- D. Kobak, J. Lomond, and B. Sanchez. The optimal ridge penalty for real-world high-dimensional data can be zero or negative due to the implicit ridge regularization. *Journal of Machine Learning Research*, 21 (169):1–16, 2020.
- C. Leng and G. Pan. Covariance estimation via sparse Kronecker structures. *Bernoulli*, 24(4B):3833 3863, 2018.
- T. Liang and A. Rakhlin. Just interpolate: kernel "ridgeless" regression can generalize. *The Annals of Statistics*, 48(3):1329–1347, 2020.
- Y. Mahdaviyeh and Z. Naulet. Risk of the least squares minimum norm estimator under the spike covariance model. *arXiv preprint arXiv:1912.13421*, 2019.
- S. Mei and A. Montanari. The generalization error of random features regression: Precise asymptotics and the double descent curve. *Communications on Pure and Applied Mathematics*, 75(4):667–766, 2022.
- X. Meng, J. Yao, and Y. Cao. Multiple descent in the multiple random feature model. *Journal of Machine Learning Research*, 25(44):1–49, 2024.
- P. Nakkiran, G. Kaplun, Y. Bansal, T. Yang, B. Barak, and I. Sutskever. Deep double descent: Where bigger models and more data hurt. In *International Conference on Learning Representations*, 2020.
- P. Nakkiran, P. Venkat, S. Kakade, and T. Ma. Optimal regularization can mitigate double descent. In *The Ninth International Conference on Learning Representations*, 2021.
- N. Puchkin and M. Rakhuba. Dimension-free structured covariance estimation. In *Proceedings of Thirty Seventh Conference on Learning Theory*, volume 247 of *Proceedings of Machine Learning Research*, pages 4276–4306. PMLR, 2024.
- N. Puchkin, F. Noskov, and V. Spokoiny. Sharper dimension-free bounds on the frobenius distance between sample covariance and its expectation. *Bernoulli*, 31(2):1664–1691, 2025.
- D. Richards, J. Mourtada, and L. Rosasco. Asymptotics of ridge (less) regression under general source condition. In *International Conference on Artificial Intelligence and Statistics*, pages 3889–3897. PMLR, 2021.
- V. Spokoiny. Mixed laplace approximation for marginal posterior and bayesian inference in error-in-operator model. Preprint. ArXiv:2305.09336, 2023.
- H. Taheri, R. Pedarsani, and C. Thrampoulidis. Fundamental limits of ridge-regularized empirical risk minimization in high dimensions. In *International Conference on Artificial Intelligence and Statistics*, pages 2773–2781. PMLR, 2021.
- M. Trabs. Bayesian inverse problems with unknown operators. *Inverse Problems*, 34(8):085001, 2018.
- A. Tsigler and P. L. Bartlett. Benign overfitting in ridge regression. *Journal of Machine Learning Research*, 24(123):1–76, 2023.
- D. Wu and J. Xu. On the optimal weighted ℓ_2 regularization in overparameterized linear regression. In *Advances in Neural Information Processing Systems*, volume 33, pages 10112–10123. Curran Associates, Inc., 2020.

- C. Zhang, S. Bengio, M. Hardt, B. Recht, and O. Vinyals. Understanding deep learning requires rethinking generalization. In *International Conference on Learning Representations*, 2017.
- N. Zhivotovskiy. Dimension-free bounds for sums of independent matrices and simple tensors via the variational principle. *Electronic Journal of Probability*, 29:1–28, 2024.

A Auxiliary results: properties of the Hessian of $\mathcal{L}(v)$

Appendix A contains some useful properties of the Hessian of $\mathcal{L}(\boldsymbol{v})$. Throughout this section, denote $\nabla^2 \mathcal{L}(\boldsymbol{v})$ by $H = H(\boldsymbol{v})$ for brevity. The value of \boldsymbol{v} will always be clear from context. We will extensively use the block form of H:

$$H = \begin{pmatrix} H_{\theta\theta} & H_{\theta\eta} & H_{\thetaA} \\ H_{\eta\theta} & H_{\eta\eta} & H_{\etaA} \\ H_{A\theta} & H_{A\eta} & H_{AA} \end{pmatrix}.$$
 (51)

Here the diagonal blocks correspond to

$$H_{\theta\theta} = A^{\top}A + \lambda I_d, \quad H_{\eta\eta} = 2I_d, \text{ and } H_{AA} = \operatorname{diag}(H_{\mathbf{a}_1\mathbf{a}_1}, \dots, H_{\mathbf{a}_d\mathbf{a}_d}),$$

while the non-diagonal ones are given by

$$H_{\theta\eta} = H_{\eta\theta}^{\top} = -A^{\top}, \quad H_{\theta A} = H_{A\theta}^{\top} = \begin{pmatrix} H_{\theta \mathbf{a}_1} & \dots & H_{\theta \mathbf{a}_d} \end{pmatrix}, \quad H_{\eta A} = H_{A\eta}^{\top} = \begin{pmatrix} H_{\eta \mathbf{a}_1} & \dots & H_{\eta \mathbf{a}_d} \end{pmatrix},$$

where, for any $j \in \{1, \ldots, d\}$,

$$H_{\mathbf{a}_j \mathbf{a}_j} = \mu^2 I_d + \boldsymbol{\theta} \boldsymbol{\theta}^\top, \quad H_{\boldsymbol{\theta} \mathbf{a}_j} = (\mathbf{a}_j^\top \boldsymbol{\theta} - \boldsymbol{\eta}_j) I_d + \mathbf{a}_j \boldsymbol{\theta}^\top \text{ and } H_{\boldsymbol{\eta} \mathbf{a}_j} = -\mathbf{e}_j \boldsymbol{\theta}^\top.$$

Throughout the proofs, we will often use a representation of H in terms of the Kronecker product. Let us recall that, for any matrices $V \in \mathbb{R}^{p \times q}$ and $W \in \mathbb{R}^{r \times s}$, their Kronecker product $U \otimes W$ is a matrix of size $pq \times rs$, defined as

$$\begin{pmatrix} V_{11}W & \dots & V_{1q}W \\ \vdots & \ddots & \vdots \\ V_{p1}W & \dots & V_{pq}W \end{pmatrix}$$

Then we can rewrite *H* in the following form:

$$H = \begin{pmatrix} A^{\top}A + \lambda I_d & -A^{\top} & (A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} \otimes I_d + A^{\top} \otimes \boldsymbol{\theta}^{\top} \\ -A & 2I_d & -I_d \otimes \boldsymbol{\theta}^{\top} \\ (A\boldsymbol{\theta} - \boldsymbol{\eta}) \otimes I_d + A \otimes \boldsymbol{\theta} & -I_d \otimes \boldsymbol{\theta} & I_d \otimes (\mu^2 I_d + \boldsymbol{\theta} \boldsymbol{\theta}^{\top}) \end{pmatrix}.$$
 (52)

The first result indicates that $\nabla^2 \mathcal{L}(\boldsymbol{v})$ is positive definite on a set which is sufficiently large for our purposes. In particular, according to Lemma 3.1, the proper choice of the parameters μ , λ , and ρ yields that both $\hat{\boldsymbol{v}}$ and \boldsymbol{v}^* belong to this set.

Lemma A.1. Let us fix any $\rho \in [0, 1/7]$ and let $\boldsymbol{v} = (\boldsymbol{\theta}, \boldsymbol{\eta}, \mathbf{vec}(A))$ be any triplet from

$$\Upsilon(\rho) = \left\{ \boldsymbol{\upsilon} = (\boldsymbol{\theta}, \boldsymbol{\eta}, \operatorname{vec}(A)) : (1 - \rho^2) \|\boldsymbol{\theta}\|^2 \leqslant \rho^2 \mu^2, \|\boldsymbol{\eta} - A\boldsymbol{\theta}\|^2 \leqslant \rho^2 \mu^2 \lambda \right\}.$$

Then it holds that

$$\nabla^2 \mathcal{L}(\boldsymbol{\upsilon}) = H \succeq \frac{1 - 2\rho}{4} \, \mathrm{D}^2,$$

where

$$\mathsf{D}^{2} = \mathsf{D}^{2}(\boldsymbol{v}) = \operatorname{diag}\left(H_{\boldsymbol{\theta}\boldsymbol{\theta}}, H_{\boldsymbol{\eta}\boldsymbol{\eta}}, H_{AA}\right) = \operatorname{diag}\left(A^{\top}A + \lambda I_{d}, 2I_{d}, I_{d} \otimes (\mu^{2}I_{d} + \boldsymbol{\theta}\boldsymbol{\theta}^{\top})\right).$$

We postpone the proof of Lemma A.1 to Section A.1 and move to the following auxiliary result. The error-in-operator estimate $\hat{\theta}$ obviously depends on the nuisance parameter $\chi = (\eta, A)$. Let

$$H_{\theta\chi} = H_{\chi\theta}^{\top} = \begin{pmatrix} H_{\theta\eta} & H_{\theta A} \end{pmatrix} \text{ and } H_{\chi\chi} = \begin{pmatrix} H_{\eta\eta} & H_{\eta A} \\ H_{A\eta} & H_{AA} \end{pmatrix}$$

stand for the corresponding blocks H. Our analysis of bias and variance of the error-in-operator estimate $\hat{\theta}$ requires examination of H^{-1} . According to the block-matrix inversion formula, it holds that

$$H^{-1} = \begin{pmatrix} \breve{H}_{\theta\theta}^{-1} & -\breve{H}_{\theta\theta}^{-1}H_{\theta\chi}H_{\chi\chi}^{-1} \\ -\breve{H}_{\chi\chi}^{-1}H_{\chi\theta}H_{\theta\theta}^{-1} & \breve{H}_{\chi\chi}^{-1} \end{pmatrix} = \begin{pmatrix} \breve{H}_{\theta\theta}^{-1} & -H_{\theta\theta}^{-1}H_{\theta\chi}\breve{H}_{\chi\chi}^{-1} \\ -H_{\chi\chi}^{-1}H_{\chi\theta}\breve{H}_{\theta\theta}^{-1} & \breve{H}_{\chi\chi}^{-1} \end{pmatrix}.$$
 (53)

We will be particularly interested in

$$\check{H}_{\theta\theta} = H_{\theta\theta} - H_{\theta\chi} H_{\chi\chi}^{-1} H_{\chi\theta} = H/H_{\chi\chi};$$

which is nothing but the Schur complement of $H_{\chi\chi}$. Before we proceed, we have to introduce additional notation. Let $\mathfrak{m}(\cdot)$ be an arbitrary probability measure on the space of triplets $\boldsymbol{v} = (\boldsymbol{\theta}, \boldsymbol{\eta}, A)$. For a function $f(\boldsymbol{v})$ (scalar, vector- or matrix-valued), we denote

$$\langle f \rangle^{\mathfrak{m}} = \int f(\boldsymbol{v}) \, \mathrm{d}\mathfrak{m}(\boldsymbol{v})$$

In particular, $\langle H \rangle_{\chi\chi}^{\mathfrak{m}}$ stands for

$$\langle H \rangle_{\chi\chi}^{\mathfrak{m}} = \int H_{\chi\chi}(\boldsymbol{v}) \,\mathrm{d}\mathfrak{m}(\boldsymbol{v}).$$
 (54)

The next proposition shows that the inverse of $\langle H \rangle_{\chi\chi}^{\mathfrak{m}}$ (and, hence, $H_{\chi\chi}^{-1}$ as well) has a nice structure. **Proposition A.2.** Let $\rho \leq 1/2$ and let $\mathfrak{m}(\cdot)$ be an arbitrary probability measure supported on

$$\{(\boldsymbol{\theta}, \boldsymbol{\eta}, A) \in \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^{d \times d} : \|\boldsymbol{\theta}\| \leqslant \rho \mu\}.$$
(55)

Denote the inverse of $\langle H \rangle_{\chi\chi}^{\mathfrak{m}}$ defined in (54) by J. Then the matrix J can be represented in the form

$$J = \begin{pmatrix} J_{\eta\eta} & J_{\eta A} \\ J_{A\eta} & J_{AA} \end{pmatrix} = \begin{pmatrix} (0.5 + r_1)I_d & r_2I_d \otimes \widetilde{\boldsymbol{\theta}}^\top \\ r_2I_d \otimes \widetilde{\boldsymbol{\theta}} & I_d \otimes R_3 \end{pmatrix}, \quad \text{where } \widetilde{\boldsymbol{\theta}} = \mu^2 \left(\mu^2 I_d + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^\top \right\rangle^{\mathfrak{m}} \right)^{-1} \left\langle \boldsymbol{\theta} \right\rangle^{\mathfrak{m}},$$

and the non-negative numbers r_1, r_2 and the matrix $R_3 \in \mathbb{R}^{d \times d}$ satisfy the inequalities

$$0 \leqslant r_1 \leqslant \rho^2$$
, $0 \leqslant r_2 \leqslant \frac{1}{\mu^2}$, and $||R_3|| \leqslant \frac{2}{\mu^2}$.

In addition, there exists $r'_2 \in [0, \rho^2/\mu^2]$, $r'_3 \in [0, \mu^{-4}]$, and $R'_3 \in \mathbb{R}^{d \times d}$, $||R'_3|| \leq \mu^{-4}$, such that

$$r_2 = \frac{1}{2\mu^2} + r'_2, \quad and \quad R_3 = \mu^{-2}I_d + R'_3 \left\langle \boldsymbol{\theta}\boldsymbol{\theta}^\top \right\rangle^{\mathfrak{m}} + r'_3 \widetilde{\boldsymbol{\theta}}\widetilde{\boldsymbol{\theta}}^\top.$$

We move the proof of Proposition A.2 to Section A.2 and proceed with a lemma allowing us to control the smoothness of the Hessian of $\mathcal{L}(v)$.

Lemma A.3. Let us fix arbitrary $\lambda_0 > 0$ and $\mu_0 > 0$ and denote

$$\mathsf{D}_0^2 = \operatorname{diag}\left((A^*)^\top A^* + \lambda_0 I_d, 2I_d, I_d \otimes \left(\mu_0^2 I_d + \boldsymbol{\theta}^* (\boldsymbol{\theta}^*)^\top \right) \right).$$

Let v^* be as defined in (6). Then, for any vectors **u** and v of dimension $2d + d^2$, it holds that

$$\left|\mathbf{u}^{\top}\left(\nabla^{2}\mathcal{L}(\boldsymbol{\upsilon}^{*}+\boldsymbol{\upsilon})-\nabla^{2}\mathcal{L}(\boldsymbol{\upsilon}^{*})\right)\mathbf{u}\right| \leqslant \frac{4}{\mu_{0}\sqrt{\lambda_{0}}} \|\mathbf{D}_{0}\boldsymbol{\upsilon}\|\|\mathbf{D}_{0}\mathbf{u}\|^{2}+\frac{2}{\mu_{0}^{2}\lambda_{0}}\|\mathbf{D}_{0}\boldsymbol{\upsilon}\|^{2}\|\mathbf{D}_{0}\mathbf{u}\|^{2}.$$

The proof of Lemma A.3 is moved to Section A.3.

A.1 Proof of Lemma A.1

Note that, for any $\mathbf{u}, \mathbf{v} \in \mathbb{R}^d$ and $\mathbf{w} \in \mathbb{R}^{d^2}$, the quadratic form

$$\begin{pmatrix} \mathbf{u}^\top & \mathbf{v}^\top & \mathbf{w}^\top \end{pmatrix} \begin{pmatrix} H_{\boldsymbol{\theta}\boldsymbol{\theta}} & H_{\boldsymbol{\theta}\boldsymbol{\eta}} & H_{\boldsymbol{\theta}\boldsymbol{A}} \\ H_{\boldsymbol{\eta}\boldsymbol{\theta}} & H_{\boldsymbol{\eta}\boldsymbol{\eta}} & H_{\boldsymbol{\eta}\boldsymbol{A}} \\ H_{\boldsymbol{A}\boldsymbol{\theta}} & H_{\boldsymbol{A}\boldsymbol{\eta}} & H_{\boldsymbol{A}\boldsymbol{A}} \end{pmatrix} \begin{pmatrix} \mathbf{u} \\ \mathbf{v} \\ \mathbf{w} \end{pmatrix}$$

is equal to

$$\mathbf{u}^{\top} H_{\boldsymbol{\theta}\boldsymbol{\theta}} \mathbf{u} + \mathbf{v}^{\top} H_{\boldsymbol{\eta}\boldsymbol{\eta}} \mathbf{v} + \mathbf{w}^{\top} H_{AA} \mathbf{w} + 2\mathbf{u}^{\top} H_{\boldsymbol{\theta}\boldsymbol{\eta}} \mathbf{v} + 2\mathbf{u}^{\top} H_{\boldsymbol{\theta}A} \mathbf{w} + 2\mathbf{v}^{\top} H_{\boldsymbol{\eta}A} \mathbf{w}$$
$$= \|A\mathbf{u}\|^{2} + \lambda \|\mathbf{u}\|^{2} + 2\|\mathbf{v}\|^{2} + \|H_{AA}^{1/2} \mathbf{w}\|^{2} + 2\mathbf{u}^{\top} A^{\top} \mathbf{v} + 2\mathbf{u}^{\top} H_{\boldsymbol{\theta}A} \mathbf{w} + 2\mathbf{v}^{\top} H_{\boldsymbol{\eta}A} \mathbf{w}.$$
(56)

Our goal is to show that the right-hand side of (56) is not smaller than

$$\frac{1-2\rho}{4} \left(\|A\mathbf{u}\|^2 + \lambda \|\mathbf{u}\|^2 + 2\|\mathbf{v}\|^2 + \|H_{AA}^{1/2}\mathbf{w}\|^2 \right).$$

For this purpose, we prove the following lemma in Appendix A.4.

Lemma A.4. Under the conditions of Lemma A.1, it holds that

$$\|H_{\boldsymbol{\eta}A}H_{AA}^{-1/2}\| \leqslant \rho \quad and \quad \|H_{\boldsymbol{\theta}\boldsymbol{\theta}}^{-1/2}H_{\boldsymbol{\theta}A}H_{AA}^{-1/2}\| \leqslant \rho\sqrt{2}.$$

The desired lower bound easily follows from Lemma A.4. Indeed, using to this lemma and the Cauchy-Schwarz inequality $2\sqrt{2}ab \leq 3a^2/4 + 8b^2/3$, we obtain that

$$2\mathbf{u}^{\top}H_{\theta A}\mathbf{w} \ge -2\|H_{\theta \theta}^{1/2}\mathbf{u}\|\|H_{\theta \theta}^{-1/2}H_{\theta A}H_{AA}^{1/2}\|\|H_{AA}^{1/2}\mathbf{w}\| \ge -\rho\left(\frac{3}{4}\|A\mathbf{u}\|^{2} + \frac{3\lambda}{4}\|\mathbf{u}\|^{2} + \frac{8}{3}\|H_{AA}^{1/2}\mathbf{w}\|^{2}\right)$$

and

$$2\mathbf{v}^{\top}H_{\boldsymbol{\eta}A}\mathbf{w} \geq -2\|\mathbf{v}\|\|H_{\boldsymbol{\eta}A}H_{AA}\|\|H_{AA}^{1/2}\mathbf{w}\| \geq -\rho\left(\|\mathbf{v}\|^2 + \|H_{AA}^{1/2}\mathbf{w}\|^2\right).$$

This yields that

$$\begin{aligned} \|A\mathbf{u}\|^{2} + \lambda \|\mathbf{u}\|^{2} + 2\|\mathbf{v}\|^{2} + \mathbf{w}^{\top} H_{AA}\mathbf{w} + 2\mathbf{u}^{\top} A^{\top}\mathbf{v} + 2\mathbf{u}^{\top} H_{\boldsymbol{\theta}A}\mathbf{w} + 2\mathbf{v}^{\top} H_{\boldsymbol{\eta}A}\mathbf{w} \\ \geqslant \left(1 - \frac{3\rho}{4}\right) \left(\|A\mathbf{u}\|^{2} + \lambda \|\mathbf{u}\|^{2}\right) + 2\left(1 - \frac{\rho}{2}\right) \|\mathbf{v}\|^{2} + \left(1 - \frac{8\rho}{3} - \rho\right) \|H_{AA}^{1/2}\mathbf{w}\|^{2} + 2\mathbf{u}^{\top} A^{\top}\mathbf{v}. \end{aligned}$$

Taking into account the lower bound

$$\|A\mathbf{u}\|^{2} + 2\|\mathbf{v}\|^{2} + 2\mathbf{u}^{\top}A^{\top}\mathbf{v} \ge \|A\mathbf{u}\|^{2} + 2\|\mathbf{v}\|^{2} - \left(\frac{2}{3}\|A\mathbf{u}\|^{2} + \frac{3}{2}\|\mathbf{v}\|^{2}\right) \ge \frac{1}{3}\|A\mathbf{u}\|^{2} + \frac{1}{2}\|\mathbf{v}\|^{2},$$

we conclude that

$$\|A\mathbf{u}\|^{2} + \lambda \|\mathbf{u}\|^{2} + 2\|\mathbf{v}\|^{2} + \mathbf{w}^{\top} H_{AA}\mathbf{w} + 2\mathbf{u}^{\top} A^{\top}\mathbf{v} + 2\mathbf{u}^{\top} H_{\theta A}\mathbf{w} + 2\mathbf{v}^{\top} H_{\eta A}\mathbf{w}$$

$$\geq \left(\frac{1}{3} - \frac{3\rho}{4}\right) \left(\|A\mathbf{u}\|^{2} + \lambda \|\mathbf{u}\|^{2}\right) + 2\left(\frac{1}{4} - \frac{\rho}{2}\right) \|\mathbf{v}\|^{2} + \left(1 - \frac{8\rho}{3} - \rho\right) \|H_{AA}^{1/2}\mathbf{w}\|^{2}.$$

Finally, since $\rho \in [0, 1/7]$, it holds that

$$\frac{1}{3} - \frac{3\rho}{4} \ge \frac{1}{4} - \left(\frac{3}{4} - \frac{1}{4}\right)\rho = \frac{1}{4} - \frac{\rho}{2}, \quad \text{and} \quad 1 - \frac{11\rho}{3} \ge \frac{10}{21} \ge \frac{1}{4} - \frac{\rho}{2},$$

and we obtain the desired bound:

$$\begin{aligned} \|A\mathbf{u}\|^{2} + \lambda \|\mathbf{u}\|^{2} + 2\|\mathbf{v}\|^{2} + \mathbf{w}^{\top} H_{AA}\mathbf{w} + 2\mathbf{u}^{\top} A^{\top}\mathbf{v} + 2\mathbf{u}^{\top} H_{\theta A}\mathbf{w} + 2\mathbf{v}^{\top} H_{\eta A}\mathbf{w} \\ \geqslant \frac{1-2\rho}{4} \left(\|A\mathbf{u}\|^{2} + \lambda \|\mathbf{u}\|^{2} + 2\|\mathbf{v}\|^{2} + \|H_{AA}^{1/2}\mathbf{w}\|^{2} \right), \end{aligned}$$

which is equivalent to

$$\begin{pmatrix} \mathbf{u}^{\top} & \mathbf{v}^{\top} & \mathbf{w}^{\top} \end{pmatrix} \begin{pmatrix} H_{\theta\theta} & H_{\theta\eta} & H_{\thetaA} \\ H_{\eta\theta} & H_{\eta\eta} & H_{\etaA} \\ H_{A\theta} & H_{A\eta} & H_{AA} \end{pmatrix} \begin{pmatrix} \mathbf{u} \\ \mathbf{v} \\ \mathbf{w} \end{pmatrix}$$

$$\geq \frac{1-2\rho}{4} \begin{pmatrix} \mathbf{u}^{\top} & \mathbf{v}^{\top} & \mathbf{w}^{\top} \end{pmatrix} \begin{pmatrix} H_{\theta\theta} & O & O \\ O & H_{\eta\eta} & O \\ O & O & H_{AA} \end{pmatrix} \begin{pmatrix} \mathbf{u} \\ \mathbf{v} \\ \mathbf{w} \end{pmatrix}.$$

The proof is finished.

A.2 Proof of Proposition A.2

Since the measure \mathfrak{m} is clear from the context, we omit it throughout the proof and write $\langle H \rangle$ and $\langle \theta \rangle$, instead of $\langle H \rangle^{\mathfrak{m}}$ and $\langle \theta \rangle^{\mathfrak{m}}$ for brevity. Let us note that, according to (52), the matrix $H_{\chi\chi}(\boldsymbol{v})$ admits the following form:

$$H_{\chi\chi} = \begin{pmatrix} 2I_d & I_d \otimes \boldsymbol{\theta}^\top \\ I_d \otimes \boldsymbol{\theta} & I_d \otimes (\mu^2 I_d + \boldsymbol{\theta} \boldsymbol{\theta}^\top) \end{pmatrix}.$$

Then, due to the definition of $\langle H \rangle$ it holds that

$$\langle H \rangle_{\chi\chi} = \begin{pmatrix} 2I_d & I_d \otimes \langle \boldsymbol{\theta} \rangle^\top \\ I_d \otimes \langle \boldsymbol{\theta} \rangle & I_d \otimes (\mu^2 I_d + \langle \boldsymbol{\theta} \boldsymbol{\theta}^\top \rangle) \end{pmatrix}.$$

Due to the blockwise inversion formula, we have

$$J_{\eta\eta} = \left(\langle H \rangle_{\eta\eta} - \langle H \rangle_{\eta A} \langle H \rangle_{AA}^{-1} \langle H \rangle_{A\eta} \right)^{-1} \\ = \left(2I_d - \left(I_d \otimes \langle \boldsymbol{\theta} \rangle^\top \right) \left[I_d \otimes \left(\mu^2 I_d + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^\top \right\rangle \right) \right]^{-1} \left(I_d \otimes \langle \boldsymbol{\theta} \rangle \right) \right)^{-1}.$$

The latter expression can significantly simplified using the Kronecker product properties:

$$J_{\eta\eta} = \left(2 - \langle \boldsymbol{\theta} \rangle^{\top} \left(\mu^2 I_d + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right\rangle\right)^{-1} \langle \boldsymbol{\theta} \rangle\right)^{-1} I_d.$$

We are going to show that the remainder

$$r_1 = \left(2 - \langle \boldsymbol{\theta} \rangle^\top \left(\mu^2 I_d + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^\top \right\rangle\right)^{-1} \langle \boldsymbol{\theta} \rangle\right)^{-1} - \frac{1}{2}$$

is small and, hence, $J_{\eta\eta}$ is close to $0.5I_d$. Indeed, taking into account that m is supported on the set defined in (55), we observe that

$$\|\langle \boldsymbol{\theta} \rangle\|^2 \leqslant \rho^2 \mu^2.$$

Then, according to the Sherman-Morrison formula,

$$\begin{split} \langle \boldsymbol{\theta} \rangle^{\top} \left(\mu^{2} I_{d} + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right\rangle \right)^{-1} \langle \boldsymbol{\theta} \rangle &\leq \langle \boldsymbol{\theta} \rangle^{\top} \left(\mu^{2} I_{d} + \left\langle \boldsymbol{\theta} \right\rangle \left\langle \boldsymbol{\theta} \right\rangle^{\top} \right)^{-1} \langle \boldsymbol{\theta} \rangle \\ &= \langle \boldsymbol{\theta} \rangle^{\top} \left(\frac{1}{\mu^{2}} I_{d} - \frac{\left\langle \boldsymbol{\theta} \right\rangle \left\langle \boldsymbol{\theta} \right\rangle^{\top}}{\mu^{2} (\mu^{2} + \| \left\langle \boldsymbol{\theta} \right\rangle \|^{2})} \right) \left\langle \boldsymbol{\theta} \rangle \\ &= \frac{\| \left\langle \boldsymbol{\theta} \right\rangle \|^{2}}{\mu^{2} + \| \left\langle \boldsymbol{\theta} \right\rangle \|^{2}} \leq \rho^{2}. \end{split}$$

This yields that

$$r_{1} = \frac{\langle \boldsymbol{\theta} \rangle^{\top} (\mu^{2} I_{d} + \langle \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \rangle)^{-1} \langle \boldsymbol{\theta} \rangle}{2(2 - \langle \boldsymbol{\theta} \rangle^{\top} (\mu^{2} I_{d} + \langle \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \rangle)^{-1} \langle \boldsymbol{\theta} \rangle)} \leqslant \frac{\rho^{2}}{4 - 2\rho^{2}}.$$
(57)

Since $\rho \leq 1/2$, the expression in the right-hand side is at most $2\rho^2/7 < \rho^2$.

We proceed with the block-inversion formula and compute:

$$J_{\eta A} = -J_{\eta \eta} \langle H \rangle_{\eta A} \langle H \rangle_{AA}^{-1} = \left(\frac{1}{2} + r_1\right) \left(I_d \otimes \langle \boldsymbol{\theta} \rangle^{\top}\right) \left[I_d \otimes \left(\mu^2 I_d + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right\rangle\right)^{-1}\right] = r_2 I_d \otimes \widetilde{\boldsymbol{\theta}}^{\top},$$
(58)

where

$$\widetilde{oldsymbol{ heta}} = \mu^2 \left(\mu^2 I_d + \left\langle oldsymbol{ heta} oldsymbol{ heta}^{ op}
ight
angle^{-1} \left\langle oldsymbol{ heta}
ight
angle \quad ext{and} \quad r_2 = \mu^{-2} \left(rac{1}{2} + r_1
ight) \leqslant \mu^{-2}.$$

Note that

$$\|\widetilde{\boldsymbol{\theta}}\| \leqslant \| \langle \boldsymbol{\theta} \rangle \| \leqslant \rho \mu, \tag{59}$$

because of $(1 - \rho^2) \| \langle \boldsymbol{\theta} \rangle \|^2 \leq \rho^2 \mu^2$.

It remains to consider J_{AA} . Since the product $\langle H \rangle J$ equals to the identity matrix, it holds that

$$\left\langle H\right\rangle_{A\boldsymbol{\eta}} J_{\boldsymbol{\eta}A} + \left\langle H\right\rangle_{AA} J_{AA} = I_{d^2}.$$

This yields that $J_{AA} = \langle H \rangle_{AA}^{-1} - \langle H \rangle_{AA}^{-1} \langle H \rangle_{A\eta} J_{\eta A}$ and, applying (58), we obtain that

$$J_{AA} = I_d \otimes \left(\mu^2 I_d + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^\top \right\rangle \right)^{-1} + r_2 \left[I_d \otimes \left(\mu^2 I_d + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^\top \right\rangle \right) \right]^{-1} (I_d \otimes \left\langle \boldsymbol{\theta} \right\rangle) (I_d \otimes \widetilde{\boldsymbol{\theta}}^\top)$$

= $I_d \otimes R_3$,

where

$$R_{3} = \left(\mu^{2}I_{d} + \left\langle \boldsymbol{\theta}\boldsymbol{\theta}^{\top} \right\rangle \right)^{-1} + r_{2} \left(\mu^{2}I_{d} + \left\langle \boldsymbol{\theta}\boldsymbol{\theta}^{\top} \right\rangle \right)^{-1} \left\langle \boldsymbol{\theta} \right\rangle \widetilde{\boldsymbol{\theta}}^{\top}$$
$$= \left(\mu^{2}I_{d} + \left\langle \boldsymbol{\theta}\boldsymbol{\theta}^{\top} \right\rangle \right)^{-1} + \mu^{-4} \left(\frac{1}{2} + r_{1}\right) \widetilde{\boldsymbol{\theta}}\widetilde{\boldsymbol{\theta}}^{\top}.$$

Since the norm of $\tilde{\theta}$ does not exceed $\rho\mu$ (see (59)), we have

$$||R_3|| \leq \mu^{-2} + \left(\frac{1}{2} + r_1\right)\mu^{-4}||\widetilde{\boldsymbol{\theta}}||^2 \leq \mu^{-2} + \rho^2 \left(\frac{1}{2} + r_1\right)\mu^{-2} \leq 2\mu^{-2}.$$

Finally, we study the leading terms of $J_{\eta A}$ and J_{AA} . Set $r'_1 = r_1/\mu^2$. Clearly, r'_1 is not greater than $\rho^2 \mu^{-2}$, and, due to (58), it holds that

$$J_{\boldsymbol{\eta}A} = r_2 I_d \otimes \widetilde{\boldsymbol{\theta}}^{\top} = \mu^{-2} \left(\frac{1}{2} + r_1 \right) I_d \otimes \widetilde{\boldsymbol{\theta}}^{\top} = \frac{1}{2} I_d \otimes \widetilde{\boldsymbol{\theta}}^{\top} + r_1' I_d \otimes \widetilde{\boldsymbol{\theta}}^{\top}.$$

Next, we determine the leading term in the expression for R_3 :

$$\begin{split} R_{3} &= \left(\mu^{2} I_{d} + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right\rangle \right)^{-1} + \mu^{-4} \left(\frac{1}{2} + r_{1}\right) \widetilde{\boldsymbol{\theta}} \widetilde{\boldsymbol{\theta}}^{\top} \\ &= \frac{1}{\mu^{2}} \left(\mu^{2} I_{d} + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right\rangle \right)^{-1} \left(\mu^{2} I_{d} + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right\rangle - \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right\rangle \right) + \mu^{-4} \left(\frac{1}{2} + r_{1}\right) \widetilde{\boldsymbol{\theta}} \widetilde{\boldsymbol{\theta}}^{\top} \\ &= \frac{1}{\mu^{2}} I_{d} + R_{3}' \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right\rangle + r_{3}' \widetilde{\boldsymbol{\theta}} \widetilde{\boldsymbol{\theta}}^{\top}, \end{split}$$

where the introduced matrix

$$R_3' = \frac{1}{\mu^2} \left(\mu^2 I_d + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^\top \right\rangle \right)^{-1}$$

and the coefficient $r_3'=\mu^{-4}(1/2+r_1)$ satisfy the bounds

$$\|R'_3\| = \left\|\mu^{-2}\left(\mu^2 I_d + \left\langle \boldsymbol{\theta} \boldsymbol{\theta}^\top \right\rangle\right)\right\| \leqslant \mu^{-4} \quad \text{and} \quad r'_3 = \mu^{-4}\left(\frac{1}{2} + r_1\right) \leqslant \mu^{-4}.$$

A.3 Proof of Lemma A.3

Throughout the proof, we will represent v^* , v, and u as triplets:

$$\boldsymbol{v} = (\boldsymbol{\theta}, \boldsymbol{\eta}, \operatorname{vec}(A)), \quad \boldsymbol{v}^* = (\boldsymbol{\theta}^*, \boldsymbol{\eta}^*, \operatorname{vec}(A^*)), \quad \mathbf{u} = (\mathbf{t}, \boldsymbol{\nu}, \operatorname{vec}(B)).$$

Then, calculating the Hessian of $\mathcal{L}(\boldsymbol{v})$ (see, for instance, the proof of Lemma A.1), we can rewrite the quadratic form $\mathbf{u}^{\top} \nabla^2 \mathcal{L}(\boldsymbol{v}^*) \mathbf{u}$ in the following way:

$$\mathbf{u}^{\top} \nabla^{2} \mathcal{L}(\boldsymbol{\upsilon}^{*}) \mathbf{u} = \|A^{*} \mathbf{t}\|^{2} + \lambda \|\mathbf{t}\|^{2} + 2\|\boldsymbol{\nu}\|^{2} + \|B\boldsymbol{\theta}^{*}\|^{2} + \mu \|B\|_{\mathrm{F}}^{2}$$
$$- 2\boldsymbol{\nu}^{\top} A^{*} \mathbf{t} + 2\mathbf{t}^{\top} (A^{*})^{\top} B\boldsymbol{\theta}^{*} + 2(A^{*}\boldsymbol{\theta}^{*} - \boldsymbol{\eta}^{*})^{\top} B\mathbf{t} - 2\boldsymbol{\nu}^{\top} B\boldsymbol{\theta}^{*}.$$

Note that the parameters λ and μ , appearing in the definition of $\mathcal{L}(\boldsymbol{v})$, may differ from λ_0 and μ_0 . It does not affect the proof, because λ and μ vanish, once we subtract $\mathbf{u}^\top \nabla^2 \mathcal{L}(\boldsymbol{v}^*)\mathbf{u}$ from $\mathbf{u}^\top \nabla^2 \mathcal{L}(\boldsymbol{v}^* + \boldsymbol{v})\mathbf{u}$. Indeed, it holds that

$$\mathbf{u}^{\top} \left(\nabla^{2} \mathcal{L}(\boldsymbol{v}^{*} + \boldsymbol{v}) - \nabla^{2} \mathcal{L}(\boldsymbol{v}^{*}) \right) \mathbf{u} = 2\mathbf{t}^{\top} A^{\top} A^{*} \mathbf{t} + \|A\mathbf{t}\|^{2} + 2\boldsymbol{\theta}^{\top} B^{\top} B \boldsymbol{\theta}^{*} + \|B\boldsymbol{\theta}\|^{2} - 2\boldsymbol{\nu}^{\top} A \mathbf{t} \\ + 2\boldsymbol{\theta}^{\top} B^{\top} A^{*} \mathbf{t} + 2\mathbf{t}^{\top} A^{\top} B \boldsymbol{\theta}^{*} + 2\mathbf{t}^{\top} A^{\top} B \boldsymbol{\theta} \\ + 2(A^{*} \boldsymbol{\theta} + A \boldsymbol{\theta}^{*} + A \boldsymbol{\theta} - \boldsymbol{\eta})^{\top} B \mathbf{t} - 2\boldsymbol{\nu}^{\top} B \boldsymbol{\theta}.$$

It is straightforward to observe that the difference of $\mathbf{u}^{\top}\nabla^{2}\mathcal{L}(\boldsymbol{v}^{*}+\boldsymbol{v})\mathbf{u}$ and $\mathbf{u}^{\top}\nabla^{2}\mathcal{L}(\boldsymbol{v}^{*})\mathbf{u}$ consists of the third order and the fourth order terms (in \mathbf{u} and \boldsymbol{v}):

$$\mathbf{u}^{\top} \left(\nabla^2 \mathcal{L} (\boldsymbol{v}^* + \boldsymbol{v}) - \nabla^2 \mathcal{L} (\boldsymbol{v}^*) \right) \mathbf{u} = \mathcal{T} + \mathcal{Q}, \tag{60}$$

where

$$\mathcal{T} = 2\mathbf{t}^{\top}A^{\top}A^{*}\mathbf{t} + 2\boldsymbol{\theta}^{\top}B^{\top}B\boldsymbol{\theta}^{*} - 2\boldsymbol{\nu}^{\top}A\mathbf{t} + 2\boldsymbol{\theta}^{\top}B^{\top}A^{*}\mathbf{t} + 2\mathbf{t}^{\top}A^{\top}B\boldsymbol{\theta}^{*} + 2(A^{*}\boldsymbol{\theta} + A\boldsymbol{\theta}^{*} - \boldsymbol{\eta})^{\top}B\mathbf{t} - 2\boldsymbol{\nu}^{\top}B\boldsymbol{\theta}$$

and

$$\mathcal{Q} = \|A\mathbf{t}\|^2 + \|B\boldsymbol{\theta}\|^2 + 2\mathbf{t}^\top A^\top B\boldsymbol{\theta} + 2\boldsymbol{\theta}^\top A^\top B\mathbf{t}$$

It is more convenient to study \mathcal{T} and \mathcal{Q} separately. For this reason, we split the rest of the proof into two steps and show that

$$|\mathcal{T}|\leqslant rac{4}{\mu_0\sqrt{\lambda_0}}\|\mathtt{D}_0oldsymbol{v}\|\|\mathtt{D}_0oldsymbol{u}\|^2 \quad ext{and} \quad |\mathcal{Q}|\leqslant rac{2}{\mu_0^2\lambda_0}\|\mathtt{D}_0oldsymbol{v}\|^2\|\mathtt{D}_0oldsymbol{u}\|^2.$$

Step 1: upper bound on |Q|. We start with a simpler inequality

$$|\mathcal{Q}| \leqslant rac{2}{\mu_0^2 \lambda_0} \| \mathtt{D}_0 \mathbf{u} \|^2 \| \mathtt{D}_0 oldsymbol{v} \|^2$$

and postpone the proof of the upper bound on the absolute value \mathcal{T} to the next step. It is enough to show that

$$\frac{2}{\mu_0^2\lambda_0} \|\mathsf{D}_0\mathbf{u}\|^2 \|\mathsf{D}_0\boldsymbol{\upsilon}\|^2 \leqslant \mathcal{Q} \leqslant \frac{2}{\mu_0^2\lambda_0} \|\mathsf{D}_0\mathbf{u}\|^2 \|\mathsf{D}_0\boldsymbol{\upsilon}\|^2.$$

The lower bound trivially follows from the inequalities

$$\mathcal{Q} = \|A\mathbf{t} + B\boldsymbol{\theta}\|^2 + 2\boldsymbol{\theta}^\top A^\top B\mathbf{t} \ge -2\|\boldsymbol{\theta}\| \|A\|_{\mathrm{F}} \|B\|_{\mathrm{F}} \|\mathbf{t}\| \ge -\frac{2}{\mu^2 \lambda} \|\mathsf{D}_0 \mathbf{u}\|^2 \|\mathsf{D}_0 \boldsymbol{v}\|^2.$$

From now on, we focus on the upper bound on Q. It holds that

$$\mathcal{Q} = \|A\mathbf{t}\|^2 + \|B\boldsymbol{\theta}\|^2 + 2\mathbf{t}^\top A^\top B\boldsymbol{\theta} + 2\boldsymbol{\theta}^\top A^\top B\mathbf{t}$$

$$\leq 2\|A\|_{\mathrm{F}}^2 \|\mathbf{t}\|^2 + 2\|B\|_{\mathrm{F}}^2 \|\boldsymbol{\theta}\|^2 + 2\|A\|_{\mathrm{F}} \|\boldsymbol{\theta}\| \|B\|_{\mathrm{F}} \|\mathbf{t}\|.$$
(61)

The right-hand side can be viewed as a quadratic form of $\mu_0 ||A||_F$ and $\sqrt{\lambda_0} ||\theta||$:

$$2\|A\|_{\rm F}^{2}\|\mathbf{t}\|^{2} + 2\|B\|_{\rm F}^{2}\|\boldsymbol{\theta}\|^{2} + 2\|A\|_{\rm F}\|\boldsymbol{\theta}\|\|B\|_{\rm F}\|\mathbf{t}\| = \frac{1}{\mu_{0}^{2}\lambda_{0}}\left(\mu_{0}\|A\|_{\rm F} - \sqrt{\lambda_{0}}\|\boldsymbol{\theta}\|\right) \begin{pmatrix} 2\lambda_{0}\|\mathbf{t}\|^{2} & \mu_{0}\sqrt{\lambda_{0}}\|B\|_{\rm F}\|\mathbf{t}\| \\ \mu_{0}\sqrt{\lambda_{0}}\|B\|_{\rm F}\|\mathbf{t}\| & 2\mu_{0}^{2}\|B\|_{\rm F} \end{pmatrix} \begin{pmatrix} \mu\|A\|_{\rm F} \\ \sqrt{\lambda}\|\boldsymbol{\theta}\| \end{pmatrix}.$$
(62)

Let us introduce $\varphi \in [0, \pi/2]$, such that

$$\sin\varphi = \frac{\mu_0 \|B\|_{\rm F}}{\sqrt{\lambda_0 \|t\|^2 + \mu_0^2 \|B\|_{\rm F}^2}}, \quad \cos\varphi = \frac{\sqrt{\lambda_0} \|t\|}{\sqrt{\lambda_0 \|t\|^2 + \mu_0^2 \|B\|_{\rm F}^2}}$$

and consider the matrix

$$\frac{1}{\mu_0^2 \|B\|_{\mathrm{F}}^2 + \lambda_0 \|t\|^2} \begin{pmatrix} 2\lambda_0 \|\mathbf{t}\|^2 & \mu_0 \sqrt{\lambda_0} \|B\|_{\mathrm{F}} \|\mathbf{t}\| \\ \mu_0 \sqrt{\lambda_0} \|B\|_{\mathrm{F}} \|\mathbf{t}\| & 2\mu_0^2 \|B\|_{\mathrm{F}} \end{pmatrix} = \begin{pmatrix} 2\cos^2\varphi & \sin\varphi\cos\varphi \\ \sin\varphi\cos\varphi & 2\sin^2\varphi \end{pmatrix}.$$

Direct calculations show that both its eigenvalues belong to [0, 2]. Hence, the operator norm of this matrix does not exceed 2 and we obtain that

$$\begin{aligned} \left(\mu_{0}\|A\|_{\mathrm{F}} \quad \sqrt{\lambda_{0}}\|\boldsymbol{\theta}\|\right) \begin{pmatrix} 2\lambda_{0}\|\mathbf{t}\|^{2} & \mu_{0}\sqrt{\lambda_{0}}\|B\|_{\mathrm{F}}\|\mathbf{t}\| \\ \mu_{0}\sqrt{\lambda_{0}}\|B\|_{\mathrm{F}}\|\mathbf{t}\| & 2\mu_{0}^{2}\|B\|_{\mathrm{F}} \end{pmatrix} \begin{pmatrix} \mu_{0}\|A\|_{\mathrm{F}} \\ \sqrt{\lambda_{0}}\|\boldsymbol{\theta}\| \end{pmatrix} \\ &\leqslant \left(\mu_{0}^{2}\|A\|_{\mathrm{F}}^{2} + \lambda_{0}\|\boldsymbol{\theta}\|^{2}\right) \left\| \begin{pmatrix} 2\lambda_{0}\|\mathbf{t}\|^{2} & \mu_{0}\sqrt{\lambda_{0}}\|B\|_{\mathrm{F}}\|\mathbf{t}\| \\ \mu_{0}\sqrt{\lambda_{0}}\|B\|_{\mathrm{F}}\|\mathbf{t}\| & 2\mu_{0}^{2}\|B\|_{\mathrm{F}} \end{pmatrix} \right\| \\ &= \left(\mu_{0}^{2}\|A\|_{\mathrm{F}}^{2} + \lambda_{0}\|\boldsymbol{\theta}\|^{2}\right) \left(\mu_{0}^{2}\|B\|_{\mathrm{F}}^{2} + \lambda_{0}\|t\|^{2}\right) \left\| \begin{pmatrix} 2\cos^{2}\varphi & \sin\varphi\cos\varphi \\ \sin\varphi\cos\varphi & 2\sin^{2}\varphi \end{pmatrix} \right\| \\ &\leqslant 2\left(\mu_{0}^{2}\|A\|_{\mathrm{F}}^{2} + \lambda_{0}\|\boldsymbol{\theta}\|^{2}\right) \left(\mu_{0}^{2}\|B\|_{\mathrm{F}}^{2} + \lambda_{0}\|t\|^{2}\right). \end{aligned}$$
(63)

Taking (61), (62), and (63) into account, we obtain that

$$\mathcal{Q} \leq \frac{2}{\mu_0^2 \lambda_0} \left(\mu_0^2 \|A\|_{\rm F}^2 + \lambda_0 \|\boldsymbol{\theta}\|^2 \right) \left(\mu_0^2 \|B\|_{\rm F}^2 + \lambda_0 \|t\|^2 \right) \leq \frac{2}{\mu_0^2 \lambda_0} \|\mathsf{D}_0 \boldsymbol{\upsilon}\|^2 \|\mathsf{D}_0 \boldsymbol{\upsilon}\|^2 \|\mathsf{D}_0 \boldsymbol{\upsilon}\|^2.$$
(64)

Step 2: upper bound on $|\mathcal{T}|$. It remains to bound the absolute value of \mathcal{T} . First, note that

$$\frac{\mu_{0}\sqrt{\lambda_{0}}|\mathcal{T}|}{2} \leqslant \mu_{0}\sqrt{\lambda_{0}}\|\mathbf{t}\|\|A\|_{\mathrm{F}}\|A^{*}\mathbf{t}\| + \mu_{0}\sqrt{\lambda_{0}}\|\boldsymbol{\theta}\|\|B\|_{\mathrm{F}}\|B\boldsymbol{\theta}^{*}\| + \mu_{0}\sqrt{\lambda_{0}}\|\boldsymbol{\nu}\|\|A\|_{\mathrm{F}}\|\mathbf{t}\| \\
+ \mu_{0}\sqrt{\lambda_{0}}\|\boldsymbol{\theta}\|\|B\|_{\mathrm{F}}\|A^{*}\mathbf{t}\| + \mu_{0}\sqrt{\lambda_{0}}\|\mathbf{t}\|\|A\|_{\mathrm{F}}\|B\boldsymbol{\theta}^{*}\| + \mu_{0}\sqrt{\lambda_{0}}\|\mathbf{t}\|\|B\|_{\mathrm{F}}\|A^{*}\boldsymbol{\theta}\| \quad (65) \\
+ \mu_{0}\sqrt{\lambda_{0}}\|\mathbf{t}\|\|B\|_{\mathrm{F}}\|A\boldsymbol{\theta}^{*}\| + \mu_{0}\sqrt{\lambda_{0}}\|\mathbf{t}\|\|B\|_{\mathrm{F}}\|\boldsymbol{\eta}\| + \mu_{0}\sqrt{\lambda_{0}}\|\boldsymbol{\nu}\|\|B\|_{\mathrm{F}}\|\boldsymbol{\theta}\|.$$

The right-hand side of (65) is quite massive. Let us first examine the terms $\sqrt{\lambda_0} \|\boldsymbol{\theta}\| \|A^* \mathbf{t}\| + \sqrt{\lambda_0} \|\mathbf{t}\| \|A^* \boldsymbol{\theta}\|$ and $\mu_0 \|A\|_{\mathrm{F}} \|B\boldsymbol{\theta}^*\| + \mu_0 \|B\|_{\mathrm{F}} \|A\boldsymbol{\theta}^*\|$. Due to the Cauchy-Schwarz inequality, it holds that

$$\sqrt{\lambda_0} \|\boldsymbol{\theta}\| \|A^* \mathbf{t}\| + \sqrt{\lambda_0} \|\mathbf{t}\| \|A^* \boldsymbol{\theta}\| \leq \sqrt{\|A^* \mathbf{t}\|^2 + \lambda_0 \|\mathbf{t}\|^2} \sqrt{\|A^* \boldsymbol{\theta}\|^2 + \lambda_0 \|\boldsymbol{\theta}\|^2}$$

and

$$\mu_0 \|A\|_{\mathbf{F}} \|B\boldsymbol{\theta}^*\| + \mu_0 \|B\|_{\mathbf{F}} \|A\boldsymbol{\theta}^*\| \leq \sqrt{\mu_0^2 \|B\|_{\mathbf{F}}^2 + \|B\boldsymbol{\theta}^*\|^2} \sqrt{\mu_0^2 \|A\|_{\mathbf{F}}^2 + \|A\boldsymbol{\theta}^*\|^2}.$$

Thus,

$$\mu_{0}\sqrt{\lambda_{0}}\|\boldsymbol{\theta}\|\|B\|_{F}\|A^{*}\mathbf{t}\| + \mu_{0}\sqrt{\lambda_{0}}\|\mathbf{t}\|\|A\|_{F}\|B\boldsymbol{\theta}^{*}\| + \mu_{0}\sqrt{\lambda_{0}}\|\mathbf{t}\|\|B\|_{F}\|A^{*}\boldsymbol{\theta}\|
+ \mu_{0}\sqrt{\lambda_{0}}\|\mathbf{t}\|\|B\|_{F}\|A\boldsymbol{\theta}^{*}\| + \mu_{0}\sqrt{\lambda_{0}}\|\mathbf{t}\|\|B\|_{F}\|\boldsymbol{\eta}\|
\leqslant \mu_{0}\|B\|_{F}\sqrt{\|A^{*}\mathbf{t}\|^{2} + \lambda_{0}}\|\mathbf{t}\|^{2}\sqrt{\|A^{*}\boldsymbol{\theta}\|^{2} + \lambda_{0}}\|\boldsymbol{\theta}\|^{2}}
+ \sqrt{\lambda_{0}}\|\mathbf{t}\|\sqrt{\mu_{0}^{2}}\|B\|_{F}^{2} + \|B\boldsymbol{\theta}^{*}\|^{2}}\sqrt{\mu_{0}^{2}}\|A\|_{F}^{2} + \|A\boldsymbol{\theta}^{*}\|^{2}} + \mu_{0}\sqrt{\lambda_{0}}\|\mathbf{t}\|\|B\|_{F}\|\boldsymbol{\eta}\|.$$
(66)

Applying the Cauchy-Schwarz inequality again, we obtain that

$$\mu_{0} \|B\|_{\mathrm{F}} \sqrt{\|A^{*}\mathbf{t}\|^{2} + \lambda_{0}\|\mathbf{t}\|^{2}} \sqrt{\|A^{*}\boldsymbol{\theta}\|^{2} + \lambda_{0}\|\boldsymbol{\theta}\|^{2}} + \sqrt{\lambda_{0}} \|\mathbf{t}\| \sqrt{\mu_{0}^{2}} \|B\|_{\mathrm{F}}^{2} + \|B\boldsymbol{\theta}^{*}\|^{2}} \sqrt{\mu_{0}^{2}} \|A\|_{\mathrm{F}}^{2} + \|A\boldsymbol{\theta}^{*}\|^{2}} + \mu_{0} \sqrt{\lambda_{0}} \|\mathbf{t}\| \|B\|_{\mathrm{F}} \|\boldsymbol{\eta}\| \\ \leq \sqrt{\|A^{*}\mathbf{t}\|^{2} + \lambda_{0}} \|\mathbf{t}\|^{2}} \sqrt{\mu_{0}^{2}} \|B\|_{\mathrm{F}}^{2} + \|B\boldsymbol{\theta}^{*}\|^{2}} \\ \cdot \left(\sqrt{\|A^{*}\boldsymbol{\theta}\|^{2} + \lambda_{0}} \|\boldsymbol{\theta}\|^{2}} + \sqrt{\mu_{0}^{2}} \|A\|_{\mathrm{F}}^{2} + \|A\boldsymbol{\theta}^{*}\|^{2}} + \frac{1}{\sqrt{2}} \cdot \sqrt{2} \|\boldsymbol{\eta}\|\right) \\ \leq \sqrt{\|A^{*}\mathbf{t}\|^{2} + \lambda_{0}} \|\mathbf{t}\|^{2}} \sqrt{\mu_{0}^{2}} \|B\|_{\mathrm{F}}^{2} + \|B\boldsymbol{\theta}^{*}\|^{2}} \cdot \sqrt{\frac{5}{2}} \|D_{0}\boldsymbol{v}\|.$$
(67)

The inequalities (66) and (67) help us to simplify the upper bound (65) drastically. In particular, they yield that

$$\begin{aligned} \frac{\mu_0 \sqrt{\lambda_0} |\mathcal{T}|}{2} &\leqslant \mu_0 \sqrt{\lambda_0} \|\mathbf{t}\| \|A\|_{\mathrm{F}} \|A^* \mathbf{t}\| + \mu_0 \sqrt{\lambda_0} \|\boldsymbol{\theta}\| \|B\|_{\mathrm{F}} \|B\boldsymbol{\theta}^*\| + \mu_0 \sqrt{\lambda_0} \|\boldsymbol{\nu}\| \|A\|_{\mathrm{F}} \|\mathbf{t}\| \\ &+ \mu_0 \sqrt{\lambda_0} \|\boldsymbol{\nu}\| \|B\|_{\mathrm{F}} \|\boldsymbol{\theta}\| + \sqrt{\|A^* \mathbf{t}\|^2 + \lambda_0} \|\mathbf{t}\|^2} \sqrt{\mu_0^2 \|B\|_{\mathrm{F}}^2 + \|B\boldsymbol{\theta}^*\|^2} \cdot \sqrt{\frac{5}{2}} \|\mathsf{D}_0 \boldsymbol{\upsilon}\|. \end{aligned}$$

The expression in the right-hand side can be simplified even further. Using the inequalities

$$\mu_0 \|A\|_{\mathrm{F}} \leqslant \|\mathtt{D}_0 \boldsymbol{v}\|, \quad \text{and} \quad \sqrt{\lambda_0} \|\boldsymbol{\theta}\| \leqslant \|\mathtt{D}_0 \boldsymbol{v}\|,$$

we obtain that

$$\begin{split} \frac{\mu_0 \sqrt{\lambda_0} |\mathcal{T}|}{2} &\leqslant \mu_0 \sqrt{\lambda_0} \|\mathbf{t}\| \|A\|_{\mathrm{F}} \|A^* \mathbf{t}\| + \mu_0 \sqrt{\lambda_0} \|\boldsymbol{\theta}\| \|B\|_{\mathrm{F}} \|B\boldsymbol{\theta}^*\| + \mu_0 \sqrt{\lambda_0} \|\boldsymbol{\nu}\| \|A\|_{\mathrm{F}} \|\mathbf{t}\| \\ &+ \mu_0 \sqrt{\lambda_0} \|\boldsymbol{\nu}\| \|B\|_{\mathrm{F}} \|\boldsymbol{\theta}\| + \sqrt{\|A^* \mathbf{t}\|^2 + \lambda_0 \|\mathbf{t}\|^2} \sqrt{\mu_0^2 \|B\|_{\mathrm{F}}^2 + \|B\boldsymbol{\theta}^*\|^2} \cdot \sqrt{\frac{5}{2}} \|\mathbf{D}_0 \boldsymbol{v}\| \\ &\leqslant \sqrt{\lambda_0} \|\mathbf{t}\| \|A^* \mathbf{t}\| \|\mathbf{D}_0 \boldsymbol{v}\| + \mu_0 \|B\|_{\mathrm{F}} \|B\boldsymbol{\theta}^*\| \|\mathbf{D}_0 \boldsymbol{v}\| + \sqrt{\lambda_0} \|\boldsymbol{\nu}\| \|\mathbf{t}\| \|\mathbf{D}_0 \boldsymbol{v}\| \\ &+ \mu_0 \|\boldsymbol{\nu}\| \|B\|_{\mathrm{F}} \|\mathbf{D}_0 \boldsymbol{v}\| + \sqrt{\|A^* \mathbf{t}\|^2 + \lambda \|\mathbf{t}\|^2} \sqrt{\mu^2 \|B\|_{\mathrm{F}}^2 + \|B\boldsymbol{\theta}^*\|^2} \cdot \sqrt{\frac{5}{2}} \|\mathbf{D}_0 \boldsymbol{v}\|. \end{split}$$

Moreover, introducing

$$\mathbf{w} = \left(\sqrt{\|A^*\mathbf{t}\|^2 + \lambda\|\mathbf{t}\|^2}, \sqrt{2}\|\boldsymbol{\nu}\|, \sqrt{\mu^2\|B\|_{\mathrm{F}}^2 + \|B\boldsymbol{\theta}^*\|^2}\right)^{\top} \quad \text{and} \quad \Phi = \begin{pmatrix} 1 & \sqrt{2}/4 & \sqrt{10}/4\\ \sqrt{2}/4 & 0 & \sqrt{2}/4\\ \sqrt{10}/4 & \sqrt{2}/4 & 1 \end{pmatrix},$$

we observe that

$$\begin{split} \frac{\mu_0 \sqrt{\lambda_0} |\mathcal{T}|}{2} &\leqslant \sqrt{\lambda_0} \|\mathbf{t}\| \|A^* \mathbf{t}\| \|\mathbf{D}_0 \boldsymbol{v}\| + \mu_0 \|B\|_{\mathrm{F}} \|B\boldsymbol{\theta}^*\| \|\mathbf{D}_0 \boldsymbol{v}\| + \sqrt{\lambda_0} \|\boldsymbol{\nu}\| \|\mathbf{t}\| \|\mathbf{D}_0 \boldsymbol{v}\| \\ &+ \mu_0 \|\boldsymbol{\nu}\| \|B\|_{\mathrm{F}} \|\mathbf{D}_0 \boldsymbol{v}\| + \sqrt{\|A^* \mathbf{t}\|^2 + \lambda \|\mathbf{t}\|^2} \sqrt{\mu^2 \|B\|_{\mathrm{F}}^2 + \|B\boldsymbol{\theta}^*\|^2} \cdot \sqrt{\frac{5}{2}} \|\mathbf{D}_0 \boldsymbol{v}\| \\ &\leqslant \|\mathbf{D}_0 \boldsymbol{v}\| \|\mathbf{w}^\top \Phi \mathbf{w}. \end{split}$$

Finally, direct calculations show that $\|\Phi\| < 2$, and then

$$\frac{\mu_0 \sqrt{\lambda_0} |\mathcal{T}|}{2} \leqslant 2 \| \mathbf{D}_0 \boldsymbol{v} \| \left(\|A^* \mathbf{t}\|^2 + \lambda_0 \|\mathbf{t}\|^2 + 2 \|\boldsymbol{\nu}\|^2 + \mu_0^2 \|B\|_{\mathrm{F}}^2 + \|B\boldsymbol{\theta}^*\|^2 \right) = 2 \| \mathbf{D}_0 \boldsymbol{v} \| \|\mathbf{D}_0 \mathbf{u}\|^2.$$
(68)

Hence, summing up (60), (64), and (68), we obtain that

$$\left|\mathbf{u}^{\top}\left(\nabla^{2}\mathcal{L}(\boldsymbol{\upsilon})-\nabla^{2}\mathcal{L}(\boldsymbol{\upsilon}^{*})\right)\mathbf{u}\right| \leq |\mathcal{T}|+|\mathcal{Q}| \leq \frac{4}{\mu_{0}\sqrt{\lambda_{0}}} \|\mathbf{D}_{0}\boldsymbol{\upsilon}\|\|\mathbf{D}_{0}\mathbf{u}\|^{2}+\frac{2}{\mu_{0}^{2}\lambda_{0}}\|\mathbf{D}_{0}\boldsymbol{\upsilon}\|^{2}\|\mathbf{D}_{0}\mathbf{u}\|^{2}.$$

A.4 Proof of Lemma A.4

The proofs of the upper bounds $||H_{\eta A}H_{AA}^{-1/2}|| \leq \rho$ and $||H_{\theta\theta}^{-1/2}H_{\theta A}H_{AA}^{-1/2}|| \leq \rho\sqrt{2}$ is based on a simple observation that the operator norm of a matrix B does not exceed ρ if and only if $B^{\top}B \leq \rho^2 I_d$. In our case, it is enough to show that

$$H_{AA}^{-1/2}H_{\eta A}H_{A\eta}H_{AA}^{-1/2} \preceq \rho^2 I_d \quad \text{and} \quad H_{\theta\theta}^{-1/2}H_{\theta A}H_{AA}^{-1}H_{A\theta}H_{\theta\theta}^{-1/2} \preceq 2\rho^2 I_d.$$

For convenience, we split the derivations into two steps.

Step 1: upper bound on $||H_{\eta A}H_{AA}||$. Let us recall that (see (51) and (52))

$$H_{\boldsymbol{\eta}A} = H_{A\boldsymbol{\eta}}^{\top} = -I_d \otimes \boldsymbol{\theta}^{\top} \text{ and } H_{AA} = I_d \otimes \left(\mu^2 I_d + \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right).$$

According to the conditions of the lemma, the triplet $\boldsymbol{v} = (\boldsymbol{\theta}, \boldsymbol{\eta}, \mathbf{vec}(A))$ belongs to the set $\Upsilon(\rho)$. Then it is straightforward to observe that

$$H_{A\boldsymbol{\eta}}H_{\boldsymbol{\eta}A} = I_d \otimes \left(\boldsymbol{\theta}\boldsymbol{\theta}^{\top}\right) \preceq I_d \otimes \left(\rho^2 \boldsymbol{\theta}\boldsymbol{\theta}^{\top} + (1-\rho^2) \|\boldsymbol{\theta}\|^2 I_d\right) \preceq I_d \otimes \left(\rho^2 \boldsymbol{\theta}\boldsymbol{\theta}^{\top} + \rho^2 \mu^2 I_d\right) = \rho^2 H_{AA}.$$

This yields that

$$\left\|H_{\eta A}H_{AA}^{-1/2}\right\|^{2} = \left\|H_{AA}^{-1/2}H_{A\eta}H_{\eta A}H_{AA}^{-1/2}\right\| \leq \rho^{2}.$$

Step 2: upper bound on $||H_{\theta\theta}^{-1/2}H_{\theta A}H_{AA}^{-1/2}||$. The proof of the upper bound on the operator norm of $H_{\theta\theta}^{-1/2}H_{\theta A}H_{AA}^{-1/2}$ relies on the explicit representation of H_{AA}^{-1} . According to the Woodbury matrix identity, it holds that

$$\left(\mu^2 I_d + \boldsymbol{\theta} \boldsymbol{\theta}^{\top}\right)^{-1} = \frac{1}{\mu^2} I_d - \frac{1}{\mu^2 (\mu^2 + \|\boldsymbol{\theta}\|^2)} \boldsymbol{\theta} \boldsymbol{\theta}^{\top}$$

$$H_{AA}^{-1} = rac{1}{\mu^2} \left(I_d \otimes \left(I_d - rac{1}{\mu^2 + \| oldsymbol{ heta} \|^2} oldsymbol{ heta} oldsymbol{ heta}^{ op}
ight)
ight),$$

and we obtain that

Then

$$\begin{aligned} H_{\boldsymbol{\theta}A}H_{AA}^{-1}H_{A\boldsymbol{\theta}} &= \frac{1}{\mu^{2}}\left((A\boldsymbol{\theta}-\boldsymbol{\eta})^{\top}\otimes I_{d}+A^{\top}\otimes\boldsymbol{\theta}^{\top}\right)\left(I_{d}\otimes\left(I_{d}-\frac{1}{\mu^{2}+\|\boldsymbol{\theta}\|^{2}}\boldsymbol{\theta}\boldsymbol{\theta}^{\top}\right)\right)\left((A\boldsymbol{\theta}-\boldsymbol{\eta})\otimes I_{d}+A\otimes\boldsymbol{\theta}\right)\\ &= \frac{\|A\boldsymbol{\theta}-\boldsymbol{\eta}\|^{2}}{\mu^{2}}\left(I_{d}-\frac{1}{\mu^{2}+\|\boldsymbol{\theta}\|^{2}}\boldsymbol{\theta}\boldsymbol{\theta}^{\top}\right)+\frac{A^{\top}(A\boldsymbol{\theta}-\boldsymbol{\eta})\boldsymbol{\theta}^{\top}}{\mu^{2}+\|\boldsymbol{\theta}\|^{2}}+\frac{\boldsymbol{\theta}(A\boldsymbol{\theta}-\boldsymbol{\eta})^{\top}A}{\mu^{2}+\|\boldsymbol{\theta}\|^{2}}+\frac{A^{\top}A\|\boldsymbol{\theta}\|^{2}}{\mu^{2}+\|\boldsymbol{\theta}\|^{2}}\\ &\leq \frac{\|A\boldsymbol{\theta}-\boldsymbol{\eta}\|^{2}}{\mu^{2}}+\frac{A^{\top}(A\boldsymbol{\theta}-\boldsymbol{\eta})\boldsymbol{\theta}^{\top}}{\mu^{2}+\|\boldsymbol{\theta}\|^{2}}+\frac{\boldsymbol{\theta}(A\boldsymbol{\theta}-\boldsymbol{\eta})^{\top}A}{\mu^{2}+\|\boldsymbol{\theta}\|^{2}}+\frac{A^{\top}A\|\boldsymbol{\theta}\|^{2}}{\mu^{2}+\|\boldsymbol{\theta}\|^{2}}.\end{aligned}$$

Thus, for any unit vector $\mathbf{u} \in \mathbb{R}^d$, it holds that

$$\mathbf{u}^{\top} H_{\boldsymbol{\theta}A} H_{AA}^{-1} H_{A\boldsymbol{\theta}} \mathbf{u} \leqslant \frac{\|A\boldsymbol{\theta} - \boldsymbol{\eta}\|^2}{\mu^2} + \frac{2\mathbf{u}^{\top} \boldsymbol{\theta} (A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} A \mathbf{u}}{\mu^2 + \|\boldsymbol{\theta}\|^2} + \frac{\|A\mathbf{u}\|^2 \|\boldsymbol{\theta}\|^2}{\mu^2 + \|\boldsymbol{\theta}\|^2}$$
$$\leqslant \frac{\|A\boldsymbol{\theta} - \boldsymbol{\eta}\|^2}{\mu^2} + \frac{2\mathbf{u}^{\top} \boldsymbol{\theta} (A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} A \mathbf{u}}{\mu^2 + \|\boldsymbol{\theta}\|^2} + \frac{\|A\mathbf{u}\|^2 \|\boldsymbol{\theta}\|^2}{\mu^2 + \|\boldsymbol{\theta}\|^2}$$
$$\leqslant \frac{2\|A\boldsymbol{\theta} - \boldsymbol{\eta}\|^2}{\mu^2} + \frac{\|A\mathbf{u}\|^2 (\mathbf{u}^{\top} \boldsymbol{\theta})^2 + \|A\mathbf{u}\|^2 \|\boldsymbol{\theta}\|^2}{\mu^2 + \|\boldsymbol{\theta}\|^2}.$$

By the definition of the set $\Upsilon(\rho)$, θ , η , and A satisfy the inequalities $||A\theta - \eta||^2 \leq \rho^2 \mu^2 \lambda$ and $||\theta||^2 \leq \rho^2 (\mu^2 + ||\theta||^2)$. This yields that

$$\mathbf{u}^{\top} H_{\boldsymbol{\theta}A} H_{AA}^{-1} H_{A\boldsymbol{\theta}} \mathbf{u} \leqslant 2\rho^2 \lambda + 2\rho^2 \|A\mathbf{u}\|^2 = 2\rho^2 \mathbf{u}^{\top} H_{\boldsymbol{\theta}\boldsymbol{\theta}} \mathbf{u}.$$

Hence, it holds that

$$H_{\theta\theta}^{-1/2}H_{\theta A}H_{AA}^{-1}H_{A\theta}H_{\theta\theta}^{-1/2} \leq 2\rho^2 I_d$$

and then

$$\|H_{\boldsymbol{\theta}\boldsymbol{\theta}}^{-1/2}H_{\boldsymbol{\theta}A}H_{AA}^{-1/2}\| \leqslant \rho\sqrt{2}$$

B Properties of the target functional minimizer

In this section, we list auxiliary properties of the triplet $v^* = (\theta^*, \eta^*, A^*)$ minimizing the objective function $\mathcal{L}(v)$. They play a fundamental role in studying the bias of the estimate \hat{v} defined in (4). Our first observation is that v^* belongs to a convex subset of $\Upsilon(\rho_0)$, $\rho_0 \in [0, 1/7]$ (see (19)), under mild assumptions. In view of Lemma A.1, this means that the functional $\mathcal{L}(v)$ is strongly convex in a vicinity of v^* .
Lemma B.1. Let us fix an arbitrary $\rho_0 \in [0, 1/7]$ and let $\Upsilon(\rho_0)$ be as defined in (19). Let

$$\begin{split} \Theta^* &= \left\{ \boldsymbol{\theta} \in \mathbb{R}^d : \|\boldsymbol{\theta}\| \leqslant \frac{\rho_0 \mu}{2} \text{ and } \|\boldsymbol{\Sigma}\boldsymbol{\theta} - \boldsymbol{\Sigma}\boldsymbol{\theta}^\circ\| \leqslant \frac{\rho_0 \mu \sqrt{\lambda}}{3} \right\}, \\ \mathsf{A}^* &= \left\{ A \in \mathbb{R}^{d \times d} : \|A - \boldsymbol{\Sigma}\| \leqslant \frac{\sqrt{\lambda}}{3} \right\}, \\ \mathsf{H}^* &= \left\{ \boldsymbol{\eta} \in \mathbb{R}^d : \|\boldsymbol{\eta} - \boldsymbol{\Sigma}\boldsymbol{\theta}^\circ\| \leqslant \frac{\rho_0 \mu \sqrt{\lambda}}{3} \right\} \end{split}$$

and assume that Σ and θ° satisfy the inequalities

$$\|\boldsymbol{\theta}^{\circ}\| \leq \frac{\rho_{0}\mu}{7} \quad and \quad \|\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^{\circ}\| \leq \frac{\rho_{0}\mu\sqrt{\lambda}}{24}.$$
 (69)

Then it holds that

$$\boldsymbol{v}^* = (\boldsymbol{\theta}^*, \boldsymbol{\eta}^*, A^*) \in \Theta^* \times \mathsf{H}^* \times \mathsf{A}^* \subseteq \Upsilon(\rho_0).$$

We provide the proof of Lemma B.1 in Appendix B.1 below. Its main ingredient is the next result following from the definition of v^* .

Lemma B.2. Assume that $\|\Sigma\| \|\theta^{\circ}\| \leq 4\mu\sqrt{\lambda}$ and $\|\theta^{\circ}\| \leq \mu/7$. Then we have

- (i) $||A^* \Sigma||_{\mathrm{F}} \leq \mu^{-1} \sqrt{\lambda} ||\boldsymbol{\theta}^{\circ}||$ and $||A^* \Sigma|| \leq \mu^{-2} ||\Sigma|| ||\boldsymbol{\theta}^{\circ}||^2$;
- (*ii*) $\|\boldsymbol{\eta}^* \Sigma \boldsymbol{\theta}^\circ\| \leqslant \sqrt{\lambda} \|\boldsymbol{\theta}^\circ\|;$
- (iii) $||A^*\theta^* \eta^*|| \leq \sqrt{\lambda} ||\theta^\circ||;$
- (iv) $\|\boldsymbol{\theta}^*\| \leq \|\boldsymbol{\theta}^\circ\|$ and $\|\boldsymbol{\theta}^* \boldsymbol{\theta}^\circ\| \leq \|\mathbf{b}_{\lambda}\| + \|\boldsymbol{\theta}^\circ\|^2/\mu$.

The proof of Lemma B.2 is moved to Section B.2. The inequalities (i) will often be enough for our purposes. However, sometimes we will need sharper bounds on the operator norm of $A^* - \Sigma$ and on the Euclidean norm of $A^*\theta^* - \eta^*$.

Lemma B.3. Assume that $\|\theta^{\circ}\| \leq \mu/49$ and $\|\Sigma\|\|\theta^{\circ}\| \leq \mu\sqrt{\lambda}/24$. Then it holds that

$$\|A^* - \Sigma\| \leqslant \left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 \sqrt{\lambda} + \frac{35\|\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^\circ\|\|\mathbf{b}_\lambda\|}{\mu^2}$$

Lemma B.4. Assume that $\|\theta^{\circ}\| \leq \mu/49$ and $\|\Sigma\|\|\theta^{\circ}\| \leq \mu\sqrt{\lambda}/24$. Then it holds that

$$\left\|A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^* - \frac{\Sigma \mathbf{b}_{\lambda}}{2}\right\| \leqslant 3\|\boldsymbol{\theta}^\circ\|\sqrt{\lambda}\left(\left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{35\|\Sigma\|\|\boldsymbol{\theta}^\circ\|\|\mathbf{b}_{\lambda}\|}{\mu^2\sqrt{\lambda}}\right),$$

where $\mathbf{b}_{\lambda} = -\lambda (\Sigma^2/2 + \lambda I_d)^{-1} \boldsymbol{\theta}^{\circ}$.

We provide the proofs of Lemma B.3 and Lemma B.4 in Sections B.3 and B.4, respectively.

B.1 Proof of Lemma B.1

First, let us show that $\theta^* \in \Theta^*$. Applying Lemma B.2(iv) with $\rho_0 \leq 1/7$ in place of ρ , we obtain that

$$\|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ\| \leqslant 2\|\boldsymbol{\theta}^\circ\|.$$
(70)

Next, taking the condition (ii) of the lemma into account and using Lemma B.2(iv), we observe that

$$\|\Sigma\boldsymbol{\theta}^* - \Sigma\boldsymbol{\theta}^\circ\| \leqslant \|\Sigma\|(\|\boldsymbol{\theta}^*\| + \|\boldsymbol{\theta}^\circ\|) \leqslant 2\|\Sigma\|\|\boldsymbol{\theta}^\circ\| \leqslant \rho_0 \mu \sqrt{\lambda}/3.$$
(71)

The inequalities (70) and (71) yield that the vector θ^* belongs to Θ^* . Our next goal is to prove that $A^* \in A^*$. Due to Lemma B.2(i), we have

$$\|A^* - \Sigma\| \leqslant \frac{\sqrt{\lambda} \|\boldsymbol{\theta}^\circ\|}{\mu} \leqslant \frac{\rho_0 \sqrt{\lambda}}{7} \leqslant \frac{\sqrt{\lambda}}{3},\tag{72}$$

and, hence, $A^* \in A^*$. An upper bound on $\|\eta^* - \Sigma \theta^\circ\|$ easily follows from (71) and (72):

$$\begin{split} \|\boldsymbol{\eta}^* - \boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ}\| &= \frac{1}{2} \|\boldsymbol{A}^*\boldsymbol{\theta}^* - \boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ}\| \leqslant \frac{1}{2} (\|\boldsymbol{A}^*\boldsymbol{\theta}^* - \boldsymbol{\Sigma}\boldsymbol{\theta}^*\| + \|\boldsymbol{\Sigma}\boldsymbol{\theta}^* - \boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ}\|) \\ &\leqslant \frac{1}{2} (\|\boldsymbol{A}^* - \boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^*\| + \|\boldsymbol{\Sigma}\boldsymbol{\theta}^* - \boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ}\|) \leqslant \frac{\rho_0 \mu \sqrt{\lambda}}{3}, \end{split}$$

where the last inequality is due to the fact that $\theta^* \in \Theta^*$ and $A \in A^*$. Thus, $\eta^* \in H^*$. It only remains to prove that the product $\Theta^* \times A^* \times H^*$ is a subset of $\Upsilon(\rho_0)$. For this purpose, let us fix an arbitrary $(\theta, A, \eta) \in \Theta^* \times A^* \times H^*$. Then, due to the definition of Θ^*, A^* and H^* , it holds that

$$\|A\boldsymbol{\theta}-\boldsymbol{\eta}\| \leq \|A-\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}\| + \|\boldsymbol{\Sigma}\boldsymbol{\theta}-\boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ}\| + \|\boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ}-\boldsymbol{\eta}\| \leq \rho_{0}\mu\sqrt{\lambda}.$$

This implies that the triplet (θ, A, η) belongs to $\Upsilon(\rho_0)$ as well. Hence, $\Theta^* \times A^* \times H^* \subseteq \Upsilon(\rho_0)$, and the proof is finished.

B.2 Proof of Lemma **B.2**

Let us remind the reader that

$$\mathcal{L}(\boldsymbol{\upsilon}) = \mathcal{L}(\boldsymbol{\theta}, \boldsymbol{\eta}, A) = \frac{1}{2} \|\boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ} - \boldsymbol{\eta}\|^{2} + \frac{1}{2} \|\boldsymbol{\eta} - A\boldsymbol{\theta}\|^{2} + \frac{\mu^{2}}{2} \|\boldsymbol{\Sigma} - A\|_{\mathrm{F}}^{2} + \frac{\lambda}{2} \|\boldsymbol{\theta}\|^{2}.$$

Since (θ^*, η^*, A^*) minimizes $\mathcal{L}(\theta, \eta, A)$, we have

$$\mathcal{L}(\boldsymbol{\theta}^*, A^*, \boldsymbol{\eta}^*) \leqslant \mathcal{L}(\boldsymbol{\theta}^\circ, \Sigma, \Sigma \boldsymbol{\theta}^\circ) = rac{\lambda \|\boldsymbol{\theta}^\circ\|^2}{2}$$

and then

$$\|A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*\| \leqslant \sqrt{\lambda} \|\boldsymbol{\theta}^\circ\|, \quad \|A^* - \Sigma\|_{\mathrm{F}} \leqslant \mu^{-1}\sqrt{\lambda} \|\boldsymbol{\theta}^\circ\|, \\ \|\boldsymbol{\eta}^* - \Sigma\boldsymbol{\theta}^\circ\| \leqslant \sqrt{\lambda} \|\boldsymbol{\theta}^\circ\|, \quad \|\boldsymbol{\theta}^*\| \leqslant \|\boldsymbol{\theta}^\circ\|.$$
(73)

This yields the statements (ii) and (iii) of the lemma and the first part of (i) and (iv). In contrary, the derivation of the inequalities

 $\|A^* - \Sigma\| \leqslant \|\Sigma\| \|\boldsymbol{\theta}^\circ\|^2 / \mu^2 \quad ext{and} \quad \|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ\| \leqslant \|\mathbf{b}_\lambda\| + 3\|\boldsymbol{\theta}^\circ\| \|\mathbf{b}_\lambda\| / \mu$

is not straightforward. For the ease of exposure, we split the rest of the proof into several steps.

Step 1. An intermediate bound on $\|\theta^* - \theta^\circ\|$. To bound the norm of $\theta^* - \theta^\circ$, we consider the gradient of \mathcal{L} . Since $\nabla_{\theta} \mathcal{L}(\theta^*, \eta^*, A^*) = 0$ and

$$\nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}, \boldsymbol{\eta}, A) = A^{\top} (A\boldsymbol{\theta} - \boldsymbol{\eta}) + \lambda \boldsymbol{\theta},$$

we have

$$-\lambda\boldsymbol{\theta}^{\circ} = \boldsymbol{\nabla}_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}^{*}, \boldsymbol{\eta}^{*}, A^{*}) - \boldsymbol{\nabla}_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}^{\circ}, \boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ}, \boldsymbol{\Sigma}) = (A^{*})^{\top} (A^{*}\boldsymbol{\theta}^{*} - \boldsymbol{\eta}^{*}) + \lambda(\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ})$$

Let us note that $\eta^* = (A^* \theta^* + \Sigma \theta^\circ)/2$. This yields that

$$\frac{1}{2}(A^*)^{\top}(A^*\boldsymbol{\theta}^* - \boldsymbol{\Sigma}\boldsymbol{\theta}^\circ) + \lambda(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) = -\lambda\boldsymbol{\theta}^\circ$$

or, equivalently,

$$\left(\frac{1}{2}(A^*)^{\top}A^* + \lambda I_d\right)(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) = -\lambda\boldsymbol{\theta}^\circ - \frac{1}{2}(A^*)^{\top}(A^* - \Sigma)\boldsymbol{\theta}^\circ = -\lambda\boldsymbol{\theta}^\circ - \frac{1}{2}\Sigma(A^* - \Sigma)\boldsymbol{\theta}^\circ - \frac{1}{2}(A^* - \Sigma)^{\top}(A^* - \Sigma)\boldsymbol{\theta}^\circ.$$
(74)

Thus, due to the triangle inequality, we obtain that

$$\begin{aligned} \|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ\| &\leq \left\| \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} \left(\lambda \boldsymbol{\theta}^\circ + \frac{1}{2} \Sigma (A^* - \Sigma) \boldsymbol{\theta}^\circ \right) \right\| \\ &+ \frac{1}{2} \left\| \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} (A^* - \Sigma)^\top (A^* - \Sigma) \boldsymbol{\theta}^\circ \right\| \\ &\leq \left\| \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} \left(\lambda \boldsymbol{\theta}^\circ + \frac{1}{2} \Sigma (A^* - \Sigma) \boldsymbol{\theta}^\circ \right) \right\| + \frac{\|A^* - \Sigma\|^2 \|\boldsymbol{\theta}^\circ\|}{2\lambda}. \end{aligned}$$

Let us introduce $\tilde{\rho} = \|\boldsymbol{\theta}^{\circ}\|/\mu \leq 1/7$. This, together with the bound (73), implies that

$$\frac{\|A^* - \Sigma\|^2 \|\boldsymbol{\theta}^\circ\|}{2\lambda} \leqslant \frac{\|\boldsymbol{\theta}^\circ\|}{2\lambda} \cdot \frac{\lambda \|\boldsymbol{\theta}^\circ\|^2}{\mu^2} = \frac{\|\boldsymbol{\theta}^\circ\|^3}{2\mu^2} \leqslant \frac{\tilde{\rho}^2 \|\boldsymbol{\theta}^\circ\|}{2}$$

Hence, it holds that

$$\|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ\| \leq \left\| \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} \left(\lambda \boldsymbol{\theta}^\circ + \frac{1}{2} \Sigma (A^* - \Sigma) \boldsymbol{\theta}^\circ \right) \right\| + \frac{\tilde{\rho}^2 \|\boldsymbol{\theta}^\circ\|}{2}.$$
(75)

.

Step 2. Final bound on $\|\theta^* - \theta^\circ\|$. In what follows, we are going to show that

$$\left\| \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} \left(\lambda \boldsymbol{\theta}^\circ + \frac{1}{2} \Sigma (A^* - \Sigma) \boldsymbol{\theta}^\circ \right) \right\| \leq \|\mathbf{b}_\lambda\| + \frac{35\tilde{\rho} \|\boldsymbol{\theta}^\circ\|}{12}.$$
(76)

Then the inequalities (75) and (76) will yield the desired bound on $\|\theta^* - \theta^\circ\|$. Let us introduce

$$S = \frac{1}{2} (A^*)^{\top} A^* - \frac{1}{2} \Sigma^2 \quad \text{and} \quad B = \left(\frac{1}{2} \Sigma^2 + \lambda I_d\right)^{-1/2} S\left(\frac{1}{2} \Sigma^2 + \lambda I_d\right)^{-1/2}.$$

Note that, according to Lemma D.4 and (73), the operator norm of B does not exceed

$$||B|| \leq ||A^* - \Sigma|| \sqrt{\frac{2}{\lambda}} + \frac{||A^* - \Sigma||^2}{\lambda} \leq \frac{||\theta^\circ||\sqrt{2}}{\mu} + \frac{||\theta^\circ||^2}{\mu^2} = \tilde{\rho}\sqrt{2} + \tilde{\rho}^2 \leq \frac{1}{4}.$$
 (77)

Here we used the inequalities $\|\boldsymbol{\theta}^{\circ}\| = \tilde{\rho}\mu$ and $0 \leq \tilde{\rho} \leq 1/7$ holding by the definition of $\tilde{\rho}$ and conditions of the lemma. In particular, (77) ensures that the matrix $(I_d + B)$ is invertible and

$$\left\| (I_d + B)^{-1} \right\| \leqslant \frac{1}{1 - \|B\|} \leqslant \frac{4}{3}.$$

Using the relation

$$(I_d + B)^{-1} = (I_d + B)^{-1}(I_d + B - B) = I_d - (I_d + B)^{-1}B,$$

we obtain that

$$\left(\frac{1}{2}(A^*)^{\top}A^* + \lambda I_d\right)^{-1} \left(\lambda \theta^{\circ} + \frac{1}{2}\Sigma(A^* - \Sigma)\theta^{\circ}\right) \\
= \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \left(I_d + B\right)^{-1} \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \left(\lambda \theta^{\circ} + \frac{1}{2}\Sigma(A^* - \Sigma)\theta^{\circ}\right) \\
= -\mathbf{b}_{\lambda} - \lambda \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \left(I_d + B\right)^{-1} B \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \theta^{\circ} \\
+ \frac{1}{2} \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \left(I_d + B\right)^{-1} \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \Sigma(A^* - \Sigma)\theta^{\circ}.$$
(78)

The norm of the second term in the right-hand side of (78) in not greater than

$$\left\|\lambda\left(\frac{1}{2}\Sigma^{2}+\lambda I_{d}\right)^{-1/2}(I_{d}+B)^{-1}B\left(\frac{1}{2}\Sigma^{2}+\lambda I_{d}\right)^{-1/2}\boldsymbol{\theta}^{\circ}\right\|$$
$$\leqslant\lambda\cdot\frac{1}{\sqrt{\lambda}}\cdot\frac{\|B\|}{1-\|B\|}\cdot\frac{\|\boldsymbol{\theta}^{\circ}\|}{\sqrt{\lambda}}=\frac{4}{3}(\tilde{\rho}\sqrt{2}+\tilde{\rho}^{2})\|\boldsymbol{\theta}^{\circ}\|\leqslant\frac{9\tilde{\rho}\|\boldsymbol{\theta}^{\circ}\|}{4},$$
(79)

while the third one does not exceed

$$\left\| \frac{1}{2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} (I_d + B)^{-1} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \Sigma (A^* - \Sigma) \boldsymbol{\theta}^{\circ} \right\|$$

$$\leq \frac{1}{2\sqrt{\lambda}} \cdot \frac{1}{1 - \|B\|} \cdot \left\| \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \Sigma \right\| \|A^* - \Sigma\| \|\boldsymbol{\theta}^{\circ}\|$$

$$\leq \frac{1}{2\sqrt{\lambda}} \cdot \frac{1}{1 - \|B\|} \cdot \frac{\|\boldsymbol{\theta}^{\circ}\|^2 \sqrt{\lambda}}{\mu} \leq \frac{4}{3} \cdot \frac{\tilde{\rho} \|\boldsymbol{\theta}^{\circ}\|}{2} = \frac{2\tilde{\rho} \|\boldsymbol{\theta}^{\circ}\|}{3}.$$
(80)

The inequalities (78), (79), and (80) imply that

$$\left\| \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} \left(\lambda \boldsymbol{\theta}^\circ + \frac{1}{2} \Sigma (A^* - \Sigma) \boldsymbol{\theta}^\circ \right) \right\|$$

$$\leqslant \| \mathbf{b}_\lambda \| + \frac{9\tilde{\rho} \| \boldsymbol{\theta}^\circ \|}{4} + \frac{2\tilde{\rho} \| \boldsymbol{\theta}^\circ \|}{3} = \| \mathbf{b}_\lambda \| + \frac{35\tilde{\rho} \| \boldsymbol{\theta}^\circ \|}{12}.$$

Hence, (76) holds and then (75) yields

$$\|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ\| \leq \|\mathbf{b}_{\lambda}\| + \frac{35\tilde{\rho}\|\boldsymbol{\theta}^\circ\|}{12} + \frac{\tilde{\rho}^2\|\boldsymbol{\theta}^\circ\|}{2} \leq \|\mathbf{b}_{\lambda}\| + 3\tilde{\rho}\|\boldsymbol{\theta}^\circ\| = \|\mathbf{b}_{\lambda}\| + \|\boldsymbol{\theta}^\circ\|^2/\mu,$$

where the last inequality is due to the fact that $\tilde{\rho} \leq 1/7$.

Step 3. The bound on $||A^* - \Sigma||$. Finally, we are going to prove that $||A^* - \Sigma|| \leq ||\Sigma|| ||\theta^\circ||^2 / \mu^2$ using the derived bound on $||\theta^* - \theta^\circ||$. Note that $||\mathbf{b}_{\lambda}|| \leq ||\theta^\circ||$ and then

$$\|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ\| \leq \|\boldsymbol{\theta}^\circ\| + 3\tilde{\rho}\|\boldsymbol{\theta}^\circ\| \leq \|\boldsymbol{\theta}^\circ\| + \frac{3\|\boldsymbol{\theta}^\circ\|}{7} \leq \frac{3\|\boldsymbol{\theta}^\circ\|}{2}.$$
(81)

Similarly to the analysis of $\theta^* - \theta^\circ$, our study of $A^* - \Sigma$ relies on the first-order optimality condition:

$$O_d = \nabla_A \mathcal{L}(\boldsymbol{\theta}^*, \boldsymbol{\eta}^*, A^*) = (A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^*)(\boldsymbol{\theta}^*)^\top + \mu^2 (A^* - \Sigma)$$

Since $\eta^* = (A^* \theta^* + \Sigma \theta^\circ)/2$, it holds that

$$\mu^{2}(A^{*}-\Sigma) = \frac{1}{2}(\Sigma\boldsymbol{\theta}^{\circ} - A^{*}\boldsymbol{\theta}^{*})(\boldsymbol{\theta}^{*})^{\top}.$$

Thus, we have

$$\left(\mu^2 I_d + \boldsymbol{\theta}^*(\boldsymbol{\theta}^*)^{\top}\right) (A^* - \Sigma) = \frac{1}{2} \Sigma (\boldsymbol{\theta}^\circ - \boldsymbol{\theta}^*) (\boldsymbol{\theta}^*)^{\top}.$$

Since the smallest eigenvalue of $\mu^2 I_d + \boldsymbol{\theta}^* (\boldsymbol{\theta}^*)^\top$ is equal to μ^2 , we obtain that

$$\|A^* - \Sigma\| \leqslant \frac{\|\Sigma\| \|\boldsymbol{\theta}^*\| \|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ\|}{2\mu^2} \leqslant \frac{3\|\Sigma\| \|\boldsymbol{\theta}^\circ\|^2}{4\mu^2} \leqslant \|\Sigma\| \|\boldsymbol{\theta}^\circ\|^2/\mu^2.$$

B.3 Proof of Lemma **B.3**

Similarly to the proof of Theorem 2.3, the analysis of $A^* - \Sigma$ starts with the Newton-Leibniz formula:

$$-\boldsymbol{\nabla}\mathcal{L}(\boldsymbol{v}^{\circ}) = \boldsymbol{\nabla}\mathcal{L}(\boldsymbol{v}^{\ast}) - \boldsymbol{\nabla}\mathcal{L}(\boldsymbol{v}^{\circ}) = \int_{0}^{1} \nabla^{2}\mathcal{L}(t\boldsymbol{v}^{\ast} + (1-t)\boldsymbol{v}^{\circ})(\boldsymbol{v}^{\ast} - \boldsymbol{v}^{\circ}) \,\mathrm{d}t = \langle H \rangle \, (\boldsymbol{v}^{\ast} - \boldsymbol{v}^{\circ}).$$

Here we used the fact that $abla \mathcal{L}(v^*) = 0$ and introduced the notation

$$\langle H \rangle = \int_0^1 \nabla^2 \mathcal{L} (t \boldsymbol{v}^* + (1-t) \boldsymbol{v}^\circ) \, \mathrm{d}t.$$

More generally, throughout the proof, for a matrix-valued function $f(\boldsymbol{v}) = f(\boldsymbol{\theta}, \boldsymbol{\eta}, A), \langle f \rangle$ stands for

$$\langle f \rangle = \int_0^1 f(t \boldsymbol{v}^* + (1-t) \boldsymbol{v}^\circ) \,\mathrm{d}t.$$

Note that, according to Lemma B.1 and Lemma A.1, the matrix $\langle H \rangle$ is invertible. Then it holds that $v^* - v^\circ = -\langle H \rangle^{-1} \nabla \mathcal{L}(v^\circ)$, and we have to study the last d^2 components of $\langle H \rangle^{-1} \nabla \mathcal{L}(v^\circ)$. For this purpose, we use the block form representation of the matrix $\langle H \rangle$ and its inverse:

$$\langle H \rangle = \begin{pmatrix} \langle H \rangle_{\theta\theta} & \langle H \rangle_{\theta\eta} & \langle H \rangle_{\thetaA} \\ \langle H \rangle_{\eta\theta} & \langle H \rangle_{\eta\eta} & \langle H \rangle_{\etaA} \\ \langle H \rangle_{A\theta} & \langle H \rangle_{A\eta} & \langle H \rangle_{AA} \end{pmatrix} = \begin{pmatrix} \langle H \rangle_{\theta\theta} & \langle H \rangle_{\theta\eta} & \langle H \rangle_{\theta\mathbf{a}_{1}} & \dots & \langle H \rangle_{\theta\mathbf{a}_{d}} \\ \langle H \rangle_{\eta\theta} & \langle H \rangle_{\eta\eta} & \langle H \rangle_{\eta\mathbf{a}_{1}} & \dots & \langle H \rangle_{\eta\mathbf{a}_{d}} \\ \langle H \rangle_{\mathbf{a}_{1}\theta} & \langle H \rangle_{\mathbf{a}_{1}\eta} & \langle H \rangle_{\mathbf{a}_{1}\mathbf{a}_{1}} & \dots & \langle H \rangle_{\mathbf{a}_{1}\mathbf{a}_{d}} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \langle H \rangle_{\mathbf{a}_{d}\theta} & \langle H \rangle_{\mathbf{a}_{d}\eta} & \langle H \rangle_{\mathbf{a}_{d}\mathbf{a}_{1}} & \dots & \langle H \rangle_{\mathbf{a}_{d}\mathbf{a}_{d}} \end{pmatrix}.$$

In what follows, we denote $\langle H \rangle^{-1}$ by F and, for any $\nu, \gamma \in \{\theta, \eta, \mathbf{a}_1, \dots, \mathbf{a}_d\}$, $F_{\nu\gamma}$ stands for the corresponding block of F. Let us note that all the components of $\nabla \mathcal{L}(\boldsymbol{v}^\circ)$, except for the first d ones, are equal to zero. Therefore, for any $j \in \{1, \dots, d\}$ we may write

$$\mathbf{a}_{j}^{*} - \Sigma \mathbf{e}_{j} = -\left(\langle H \rangle^{-1} \nabla \mathcal{L}(\boldsymbol{v}^{\circ})\right)_{\mathbf{a}_{j}} = -\left(F \nabla \mathcal{L}(\boldsymbol{v}^{\circ})\right)_{\mathbf{a}_{j}}$$
$$= -F_{\mathbf{a}_{j}\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{v}^{\circ}) - F_{\mathbf{a}_{j}\boldsymbol{\eta}} \nabla_{\boldsymbol{\eta}} \mathcal{L}(\boldsymbol{v}^{\circ}) - \sum_{k=1}^{d} F_{\mathbf{a}_{j}\mathbf{a}_{k}} \nabla_{\mathbf{a}_{k}} \mathcal{L}(\boldsymbol{v}^{\circ})$$
$$= -F_{\mathbf{a}_{j}\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{v}^{\circ}) = -\lambda F_{\mathbf{a}_{j}\boldsymbol{\theta}} \boldsymbol{\theta}^{\circ}.$$

Let us elaborate on $F_{\mathbf{a}_{i}\boldsymbol{\theta}}$. Denote the nuisance parameter by $\chi = (\boldsymbol{\eta}, A)$ and let

$$\langle H \rangle_{\chi\chi} = \begin{pmatrix} \langle H \rangle_{\eta\eta} & \langle H \rangle_{\etaA} \\ \langle H \rangle_{A\eta} & \langle H \rangle_{AA} \end{pmatrix} = \begin{pmatrix} \langle H \rangle_{\eta\eta} & \langle H \rangle_{\eta\mathbf{a}_{1}} & \dots & \langle H \rangle_{\eta\mathbf{a}_{d}} \\ \langle H \rangle_{\mathbf{a}_{1}\eta} & \langle H \rangle_{\mathbf{a}_{1}\mathbf{a}_{1}} & \dots & \langle H \rangle_{\mathbf{a}_{1}\mathbf{a}_{d}} \\ \vdots & \vdots & \ddots & \vdots \\ \langle H \rangle_{\mathbf{a}_{d}\eta} & \langle H \rangle_{\mathbf{a}_{d}\mathbf{a}_{1}} & \dots & \langle H \rangle_{\mathbf{a}_{d}\mathbf{a}_{d}} \end{pmatrix}.$$

stand for the corresponding block of $\langle H \rangle$. Introducing $J = \langle H \rangle_{\chi\chi}^{-1}$ and using the blockwise inversion formula (53), we obtain that

$$F_{\chi \theta} = -\langle H \rangle_{\chi \chi}^{-1} H_{\chi \theta} \left(\langle H \rangle / \langle H \rangle_{\chi \chi} \right)^{-1} = J H_{\chi \theta} \left(\langle H \rangle / \langle H \rangle_{\chi \chi} \right)^{-1}$$

Thus, for any $j \in \{1, \ldots, d\}$, we have

$$F_{\mathbf{a}_{j}\boldsymbol{\theta}} = -J_{\mathbf{a}_{j}\boldsymbol{\eta}}H_{\boldsymbol{\eta}\boldsymbol{\theta}}\left(\left\langle H\right\rangle/\left\langle H\right\rangle_{\chi\chi}\right)^{-1} - \sum_{k=1}^{d} J_{\mathbf{a}_{j}\mathbf{a}_{k}}H_{\mathbf{a}_{k}\boldsymbol{\theta}}\left(\left\langle H\right\rangle/\left\langle H\right\rangle_{\chi\chi}\right)^{-1}.$$

According to Proposition A.2, it holds that $J_{\mathbf{a}_j \mathbf{a}_k} = O_d$ for all $j \neq k$. Then the expression for $F_{\mathbf{a}_j \theta}$ simplifies to

$$F_{\mathbf{a}_{j}\boldsymbol{\theta}} = -J_{\mathbf{a}_{j}\boldsymbol{\eta}}H_{\boldsymbol{\eta}\boldsymbol{\theta}}\left(\left\langle H\right\rangle / \left\langle H\right\rangle_{\chi\chi}\right)^{-1} - J_{\mathbf{a}_{j}\mathbf{a}_{j}}H_{\mathbf{a}_{j}\boldsymbol{\theta}}\left(\left\langle H\right\rangle / \left\langle H\right\rangle_{\chi\chi}\right)^{-1}$$

Let us recall that in the proof of Theorem 2.3, we showed (see (26))

$$\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ} = -\lambda \left(\left\langle H \right\rangle / \left\langle H \right\rangle_{\chi\chi} \right)^{-1} \boldsymbol{\theta}^{\circ}.$$

This yields that

$$\mathbf{a}_{j}^{*} - \Sigma \mathbf{e}_{j} = -J_{\mathbf{a}_{j}\boldsymbol{\eta}}H_{\boldsymbol{\eta}\boldsymbol{\theta}}(\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ}) - J_{\mathbf{a}_{j}\mathbf{a}_{j}}H_{\mathbf{a}_{j}\boldsymbol{\theta}}(\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ}) \quad \text{for all } j \in \{1, \dots, d\}$$

and, hence,

$$A^{*} - \Sigma = \sum_{j=1}^{d} \left(\mathbf{e}_{j} (\mathbf{a}_{j}^{*})^{\top} - \Sigma \mathbf{e}_{j} \mathbf{e}_{j}^{\top} \right)$$

$$= -\sum_{j=1}^{d} \mathbf{e}_{j} (\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ})^{\top} \left(\left\{ \left\langle (\mathbf{a}_{j}^{\top} \boldsymbol{\theta} - \boldsymbol{\eta}_{j}) \right\rangle I_{d} + \left\langle \mathbf{a}_{j} \boldsymbol{\theta}^{\top} \right\rangle \right\} R_{3} + r_{2} \left\langle A \right\rangle^{\top} \mathbf{e}_{j} \widetilde{\boldsymbol{\theta}}^{\top} \right)$$

$$= - \left\langle (A\boldsymbol{\theta} - \boldsymbol{\eta}) \right\rangle (\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ})^{\top} R_{3} - \left\langle A (\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ}) \boldsymbol{\theta}^{\top} \right\rangle R_{3} - r_{2} \left\langle A \right\rangle (\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ}) \widetilde{\boldsymbol{\theta}}^{\top}.$$
(82)

Let us consider the first term in the right-hand side of (82). For the sake of brevity, we introduce

$$\rho := \frac{7 \|\boldsymbol{\theta}^{\circ}\|}{\mu}.$$

Due to Proposition C.2, Lemma B.2(iv) and the assumption $\|\theta^{\circ}\| \leq \rho \mu/7$, we have

$$\|\langle A\boldsymbol{\theta} - \boldsymbol{\eta} \rangle \| \leq \rho \mu \sqrt{\lambda}/2 + 3 \|\Sigma\| \|\mathbf{b}_{\lambda}\|$$
 and $\|\boldsymbol{\theta}^*\| \leq \|\boldsymbol{\theta}^\circ\| \leq \rho \mu/7.$

This implies that

$$\| \langle (A\boldsymbol{\theta} - \boldsymbol{\eta}) \rangle (\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ)^\top R_3 \| \leq \left(\rho \mu \sqrt{\lambda} / 2 + 3 \|\Sigma\| \|\mathbf{b}_\lambda\| \right) \cdot (2\rho\mu) \cdot \|R_3\| / 7$$

$$\leq \rho^2 \sqrt{\lambda} / 3 + 2\rho \|\Sigma\| \|\mathbf{b}_\lambda\| / \mu,$$
(83)

where we use $||R_3|| \leq 2\mu^{-2}$ due to Proposition A.2. Then we bound last two terms of (82). We have

$$\left\| \left\langle A(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \boldsymbol{\theta}^\top \right\rangle \right\| \leq \left\| \Sigma(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \left\langle \boldsymbol{\theta} \right\rangle^\top \right\| + \left\| \left\langle (A - \Sigma)(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \boldsymbol{\theta}^\top \right\rangle \right\|,$$
$$\| \left\langle A \right\rangle (\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \widetilde{\boldsymbol{\theta}}^\top \| \leq \left\| \Sigma(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \widetilde{\boldsymbol{\theta}}^\top \right\| + \left\| \left\langle A - \Sigma \right\rangle (\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \widetilde{\boldsymbol{\theta}}^\top \right\|.$$

Using Lemma B.2(iv) and the triangle inequality, we deduce that

$$\left\| \left\langle A(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \boldsymbol{\theta}^\top \right\rangle \right\| \leq \|\Sigma\| \left\{ \|\mathbf{b}_{\lambda}\| + 3\rho \|\boldsymbol{\theta}^\circ\| \right\} \| \left\langle \boldsymbol{\theta} \right\rangle^\top \| + \int_0^1 \| (A^{(t)} - \Sigma)(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ)(\boldsymbol{\theta}^{(t)})^\top \| \mathrm{d}\mathfrak{m}(t), \\ \left\| \left\langle A \right\rangle (\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \widetilde{\boldsymbol{\theta}}^\top \right\| \leq \|\Sigma\| \left\{ \|\mathbf{b}_{\lambda}\| + 3\rho \|\boldsymbol{\theta}^\circ\| \right\} \| \left\langle \boldsymbol{\theta} \right\rangle^\top \| + \int_0^1 \| (A^{(t)} - \Sigma)(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \widetilde{\boldsymbol{\theta}}^\top \| \mathrm{d}\mathfrak{m}(t).$$

Since $A^{(t)} \in A^*$, we have $||A^{(t)} - \Sigma|| \leq \sqrt{\lambda}/3$. On the other hand, the vectors θ^* , θ° , and $\theta^{(t)}$ belong to Θ^* . Thus, we can bound their norms by $\rho\mu$, as well as the norm of $\tilde{\theta}$ due to (59), and obtain that

$$\left\| \left\langle A(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \boldsymbol{\theta}^\top \right\rangle \right\| \leq \rho \mu \|\Sigma\| \left\{ \|\mathbf{b}_{\lambda}\| + 3\rho \|\boldsymbol{\theta}^\circ\| \right\} + 2\rho^2 \mu^2 \sqrt{\lambda}/3, \\ \| \left\langle A \right\rangle (\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \widetilde{\boldsymbol{\theta}} \| \leq \rho \mu \|\Sigma\| \left\{ \|\mathbf{b}_{\lambda}\| + 3\rho \|\boldsymbol{\theta}^\circ\| \right\} + 2\rho^2 \mu^2 \sqrt{\lambda}/3.$$

It remains to bound the first terms of expressions above. By the assumptions of the lemma, we have

$$\|\Sigma\|\|\boldsymbol{\theta}^{\circ}\| \leqslant \mu\sqrt{\lambda}/24$$

This yields that

$$\left\| \left\langle A(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) \boldsymbol{\theta}^\top \right\rangle \right\| \leq \rho^2 \mu^2 \sqrt{\lambda} + \rho \mu \| \boldsymbol{\Sigma} \| \| \mathbf{b}_{\lambda} \|, \tag{84}$$

$$\left\| \langle A \rangle \left(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ \right) \widetilde{\boldsymbol{\theta}}^\top \right\| \leqslant \rho^2 \mu^2 \sqrt{\lambda} + \rho \mu \| \boldsymbol{\Sigma} \| \| \mathbf{b}_\lambda \|.$$
(85)

Combining the inequalities (83), (84), (85) with (82), we finally get that

$$\|A^* - \Sigma\| \leq \rho^2 \sqrt{\lambda}/3 + \frac{2\rho \|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu} + \|R_3\| \cdot (\rho^2 \mu^2 \sqrt{\lambda} + \|\Sigma\| \|\mathbf{b}_{\lambda}\|) + r_2(\rho^2 \mu^2 \sqrt{\lambda} + \|\Sigma\| \|\mathbf{b}_{\lambda}\|),$$

nich is at most $4\rho^2 \sqrt{\lambda} + 5\rho \|\Sigma\| \|\mathbf{b}_{\lambda}\|/\mu$ due to Proposition A.2.

which is at most $4\rho^2\sqrt{\lambda} + 5\rho \|\Sigma\| \|\mathbf{b}_{\lambda}\|/\mu$ due to Proposition A.2.

B.4 Proof of Lemma **B.4**

Since $\boldsymbol{\eta}^* = (A^*\boldsymbol{\theta}^* - \Sigma\boldsymbol{\theta}^\circ)/2$, it holds that

$$\left\| A^{*}\boldsymbol{\theta}^{*} - \boldsymbol{\eta}^{*} - \frac{1}{2}\Sigma \mathbf{b}_{\lambda} \right\| = \frac{1}{2} \left\| A^{*}\boldsymbol{\theta}^{*} - \Sigma \boldsymbol{\theta}^{\circ} - \Sigma \mathbf{b}_{\lambda} \right\|$$
$$\leq \frac{1}{2} \left\| A^{*} - \Sigma \right\| \left\| \boldsymbol{\theta}^{*} \right\| + \frac{1}{2} \left\| \Sigma (\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ} - \mathbf{b}_{\lambda}) \right\|.$$
(86)

Using Lemma B.3 and Lemma B.2(iv) we can easily bound the first term in the right-hand side:

$$\|A^* - \Sigma\| \|\boldsymbol{\theta}^*\| \leqslant \left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 \|\boldsymbol{\theta}^\circ\|\sqrt{\lambda} + \frac{35\|\Sigma\|\|\boldsymbol{\theta}^\circ\|^2\|\mathbf{b}_\lambda\|}{\mu^2}.$$
(87)

To bound the second term in the right-hand side of (86), we use the first-order optimality condition $\nabla_{\theta} \mathcal{L}(\theta^*, \eta^*, A^*) =$ **0**. Similarly to the proof of Lemma B.2 (see (74)), we obtain that

$$\left(\frac{1}{2}(A^*)^{\top}A^* + \lambda I_d\right)(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) = -\lambda\boldsymbol{\theta}^\circ - \frac{1}{2}(A^*)^{\top}(A^* - \Sigma)\boldsymbol{\theta}^\circ.$$
(88)

Let us introduce

$$B = \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \left((A^*)^\top A^* - \Sigma^2\right) \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2}$$

According to Lemma D.4, the operator norm of B does not exceed

$$\|B\| \leqslant \|A^* - \Sigma\| \sqrt{\frac{2}{\lambda}} + \frac{\|A^* - \Sigma\|^2}{2\lambda}.$$

Due to Lemma **B.3**, we have

$$\frac{\|A^* - \Sigma\|}{\sqrt{\lambda}} \leqslant \left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{35\|\Sigma\|\|\boldsymbol{\theta}^\circ\|\|\mathbf{b}_\lambda\|}{\mu^2\sqrt{\lambda}}.$$

The conditions of the lemma yield that $||A^* - \Sigma|| / \sqrt{\lambda}$ is not greater than 1. Indeed, since $||\theta^\circ|| / \mu = \rho / 7 \leq 1/49$ and $\mu \sqrt{\lambda} \geq 24 ||\Sigma|| ||\theta^\circ||$, it holds that

$$\frac{\|A^* - \Sigma\|}{\sqrt{\lambda}} \le \left(\frac{2}{7}\right)^2 + \frac{35\|\mathbf{b}_{\lambda}\|}{24\mu} \le \left(\frac{2}{7}\right)^2 + \frac{35\|\boldsymbol{\theta}^\circ\|}{24\mu} \le \left(\frac{2}{7}\right)^2 + \frac{5}{24\cdot7} < \frac{1}{8}.$$

This implies that

$$\|A^* - \Sigma\|\sqrt{\frac{2}{\lambda}} + \frac{\|A^* - \Sigma\|^2}{2\lambda} \leq \frac{\|A^* - \Sigma\|}{\sqrt{\lambda}} \left(\sqrt{2} + \frac{1}{2}\right) \leq \frac{2\|A^* - \Sigma\|}{\sqrt{\lambda}}.$$

and then

$$\|B\| \leqslant \frac{2\|A^* - \Sigma\|}{\sqrt{\lambda}} \leqslant 2\left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{70\|\Sigma\|\|\boldsymbol{\theta}^\circ\|\|\mathbf{b}_\lambda\|}{\mu^2\sqrt{\lambda}} \leqslant \frac{1}{4}.$$
(89)

Using the identity (88), we obtain that

$$\begin{split} \|\Sigma\left(\boldsymbol{\theta}^{*}-\boldsymbol{\theta}^{\circ}-\mathbf{b}_{\lambda}\right)\| &= \left\|\lambda\Sigma\left(\frac{1}{2}\Sigma^{2}+\lambda I_{d}\right)^{-1}\boldsymbol{\theta}^{\circ}-\lambda\Sigma\left(\frac{1}{2}(A^{*})^{\top}A^{*}+\lambda I_{d}\right)^{-1}\boldsymbol{\theta}^{\circ}\right.\\ &\left.-\frac{1}{2}\Sigma\left(\frac{1}{2}(A^{*})^{\top}A^{*}+\lambda I_{d}\right)^{-1}(A^{*})^{\top}(A^{*}-\Sigma)\boldsymbol{\theta}^{\circ}\right\|\\ &= \left\|\lambda\Sigma\left(\frac{1}{2}\Sigma^{2}+\lambda I_{d}\right)^{-1/2}\left(I_{d}-(I_{d}+B)^{-1}\right)\left(\frac{1}{2}\Sigma^{2}+\lambda I_{d}\right)^{-1/2}\boldsymbol{\theta}^{\circ}\right.\\ &\left.-\frac{1}{2}\Sigma\left(\frac{1}{2}(A^{*})^{\top}A^{*}+\lambda I_{d}\right)^{-1}(A^{*})^{\top}(A^{*}-\Sigma)\boldsymbol{\theta}^{\circ}\right\|. \end{split}$$

Due to the triangle inequality and

$$||I_d - (I_d + B)^{-1}|| = ||B(I_d + B)^{-1}|| \le \frac{||B||}{1 - ||B||} \le \frac{4||B||}{3},$$

it holds that

$$\begin{split} \|\Sigma \left(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ - \mathbf{b}_{\lambda}\right)\| &\leq \frac{4\lambda \|B\|}{3} \left\|\Sigma \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \left\| \left\| \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \boldsymbol{\theta}^\circ \right\| \right. \\ &+ \frac{1}{2} \left\|\Sigma \left(\frac{1}{2}(A^*)^\top A^* + \lambda I_d\right)^{-1} (A^*)^\top (A^* - \Sigma) \boldsymbol{\theta}^\circ \right\|. \end{split}$$

An upper bound on the first term in the right-hand side immediately follows from the inequalities

$$\left\| \Sigma \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \right\| \leqslant \sqrt{2} \quad \text{and} \quad \left\| \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \boldsymbol{\theta}^{\circ} \right\| \leqslant \frac{\|\boldsymbol{\theta}^{\circ}\|}{\sqrt{\lambda}},$$

while the second one does not exceed

$$\begin{aligned} \frac{1}{2} \left\| \Sigma \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} (A^*)^\top (A^* - \Sigma) \theta^{\circ} \right\| \\ &\leq \frac{1}{2} \left\| A^* \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} (A^*)^\top (A^* - \Sigma) \theta^{\circ} \right\| \\ &+ \frac{1}{2} \left\| (A^* - \Sigma) \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} (A^*)^\top (A^* - \Sigma) \theta^{\circ} \right\| \\ &\leq \| (A^* - \Sigma) \theta^{\circ} \| + \frac{\| A^* - \Sigma \|^2 \| \theta^{\circ} \|}{2\lambda} \left\| \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} A^* \right\| \\ &\leq \| (A^* - \Sigma) \theta^{\circ} \| + \frac{\| A^* - \Sigma \|^2 \| \theta^{\circ} \|}{\sqrt{2\lambda}} \end{aligned}$$

Thus, we obtain that

$$\begin{split} &\|\Sigma\left(\boldsymbol{\theta}^{*}-\boldsymbol{\theta}^{\circ}-\mathbf{b}_{\lambda}\right)\| \leqslant \frac{4\|B\|\|\boldsymbol{\theta}^{\circ}\|\sqrt{2\lambda}}{3} + \|(A^{*}-\Sigma)\boldsymbol{\theta}^{\circ}\| + \frac{\|A^{*}-\Sigma\|^{2}\|\boldsymbol{\theta}^{\circ}\|}{2\lambda} \\ &\leqslant \frac{8\|\boldsymbol{\theta}^{\circ}\|\sqrt{2\lambda}}{3} \left(\left(\frac{14\|\boldsymbol{\theta}^{\circ}\|}{\mu}\right)^{2} + \frac{35\|\Sigma\|\|\boldsymbol{\theta}^{\circ}\|\|\mathbf{b}_{\lambda}\|}{\mu^{2}\sqrt{\lambda}}\right) + \|A^{*}-\Sigma\|\|\boldsymbol{\theta}^{\circ}\|\left(1+\frac{\|A^{*}-\Sigma\|}{\sqrt{2\lambda}}\right) \\ &\leqslant \frac{8\|\boldsymbol{\theta}^{\circ}\|\sqrt{2\lambda}}{3} \left(\left(\frac{14\|\boldsymbol{\theta}^{\circ}\|}{\mu}\right)^{2} + \frac{35\|\Sigma\|\|\boldsymbol{\theta}^{\circ}\|\|\mathbf{b}_{\lambda}\|}{\mu^{2}\sqrt{\lambda}}\right) + \|A^{*}-\Sigma\|\|\boldsymbol{\theta}^{\circ}\|\left(1+\frac{1}{8\sqrt{2}}\right) \\ &\leqslant \left(\frac{8\sqrt{2}}{3} + 1 + \frac{1}{8\sqrt{2}}\right) \left(\left(\frac{14\|\boldsymbol{\theta}^{\circ}\|}{\mu}\right)^{2} + \frac{35\|\Sigma\|\|\boldsymbol{\theta}^{\circ}\|\|\mathbf{b}_{\lambda}\|}{\mu^{2}\sqrt{\lambda}}\right) \|\boldsymbol{\theta}^{\circ}\|\sqrt{\lambda} \\ &\leqslant 5\|\boldsymbol{\theta}^{\circ}\|\sqrt{\lambda} \left(\left(\frac{14\|\boldsymbol{\theta}^{\circ}\|}{\mu}\right)^{2} + \frac{35\|\Sigma\|\|\boldsymbol{\theta}^{\circ}\|\|\mathbf{b}_{\lambda}\|}{\mu^{2}\sqrt{\lambda}}\right). \end{split}$$

This and the inequalities (86) and (87) yield that

$$\left\|A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^* - \frac{1}{2}\Sigma \mathbf{b}_{\lambda}\right\| \leqslant 3\|\boldsymbol{\theta}^\circ\|\sqrt{\lambda} \left(\left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{35\|\Sigma\|\|\boldsymbol{\theta}^\circ\|\|\mathbf{b}_{\lambda}\|}{\mu^2\sqrt{\lambda}}\right).$$

C Results from the proof of Theorem 2.3

C.1 Proof of Lemma 5.1

Since the measure \mathfrak{m} is fixed, we omit the superscript \mathfrak{m} and simply write $\langle H \rangle$ instead of $\langle H \rangle^{\mathfrak{m}}$. Similarly, for a matrix-valued function f, we define

$$\langle f \rangle = \int_0^1 f(\mathbf{u}^{(t)}) \,\mathrm{d}\mathfrak{m}(t), \quad \text{where} \quad \mathbf{u}^{(t)} = (\boldsymbol{\theta}^{(t)}, \boldsymbol{\eta}^{(t)}, A^{(t)}) = (1-t)\boldsymbol{v}^\circ + t\boldsymbol{v}^*. \tag{90}$$

Our analysis heavily relies on the block-form representation of H(v):

$$H = H(\boldsymbol{\upsilon}) = \begin{pmatrix} H_{\boldsymbol{\theta}\boldsymbol{\theta}} & H_{\boldsymbol{\theta}\boldsymbol{\eta}} & H_{\boldsymbol{\theta}A} \\ H_{\boldsymbol{\eta}\boldsymbol{\theta}} & H_{\boldsymbol{\eta}\boldsymbol{\eta}} & H_{\boldsymbol{\eta}A} \\ H_{A\boldsymbol{\theta}} & H_{A\boldsymbol{\eta}} & H_{AA} \end{pmatrix}.$$

Here, for any ν and γ from $\{\theta, \eta, A\}$, the block $H_{\nu\gamma} = H_{\nu\gamma}(\upsilon)$ corresponds to the second partial derivative of L with respect to ν and γ . In particular, the diagonal blocks are equal to

$$H_{\theta\theta} = A^{\top}A + \lambda I_d, \quad H_{\eta\eta} = 2I_d, \quad \text{and} \quad H_{AA} = \text{diag}(H_{\mathbf{a}_1\mathbf{a}_1}, \dots, H_{\mathbf{a}_d\mathbf{a}_d}),$$

while the off-diagonal ones are given by

$$H_{\boldsymbol{\theta}\boldsymbol{\eta}} = H_{\boldsymbol{\eta}\boldsymbol{\theta}}^{\top} = -A^{\top}, \quad H_{\boldsymbol{\theta}A} = H_{A\boldsymbol{\theta}}^{\top} = \begin{pmatrix} H_{\boldsymbol{\theta}\mathbf{a}_1} & \dots & H_{\boldsymbol{\theta}\mathbf{a}_d} \end{pmatrix}, \quad H_{\boldsymbol{\eta}A} = H_{A\boldsymbol{\eta}}^{\top} = \begin{pmatrix} H_{\boldsymbol{\eta}\mathbf{a}_1} & \dots & H_{\boldsymbol{\eta}\mathbf{a}_d} \end{pmatrix},$$

where, for any $j \in \{1, \ldots, d\}$,

$$H_{\mathbf{a}_j \mathbf{a}_j} = \mu^2 I_d + \boldsymbol{\theta} \boldsymbol{\theta}^\top, \quad H_{\boldsymbol{\theta} \mathbf{a}_j} = (\mathbf{a}_j^\top \boldsymbol{\theta} - \boldsymbol{\eta}_j) I_d + \mathbf{a}_j \boldsymbol{\theta}^\top \text{ and } H_{\boldsymbol{\eta} \mathbf{a}_j} = -\mathbf{e}_j \boldsymbol{\theta}^\top.$$

Since the proof of Lemma 5.1 is quite cumbersome, we split it into several steps.

Step 1. Decomposing the Schur complement. Let us recall that we are interested in properties of the Schur complement of $\langle H \rangle_{\chi\chi}$ corresponding to the nuisance parameter $\chi = (\eta, A)$:

$$\langle H \rangle_{\chi\chi} = \begin{pmatrix} \langle H \rangle_{\eta\eta} & \langle H \rangle_{\etaA} \\ \langle H \rangle_{A\eta} & \langle H \rangle_{AA} \end{pmatrix} = \begin{pmatrix} \langle H \rangle_{\eta\eta} & \langle H \rangle_{\eta\mathbf{a}_1} & \dots & \langle H \rangle_{\eta\mathbf{a}_d} \\ \langle H \rangle_{\mathbf{a}_1\eta} & \langle H \rangle_{\mathbf{a}_1\mathbf{a}_1} & \dots & \langle H \rangle_{\mathbf{a}_1\mathbf{a}_d} \\ \vdots & \vdots & \ddots & \vdots \\ \langle H \rangle_{\mathbf{a}_d\eta} & \langle H \rangle_{\mathbf{a}_d\mathbf{a}_1} & \dots & \langle H \rangle_{\mathbf{a}_d\mathbf{a}_d} \end{pmatrix}$$

In what follows, we denote the inverse of $\langle H \rangle_{\chi\chi}$ by J:

$$\langle H \rangle_{\chi\chi}^{-1} = J = \begin{pmatrix} J_{\eta\eta} & J_{\eta A} \\ J_{A\eta} & J_{AA} \end{pmatrix} = \begin{pmatrix} J_{\eta\eta} & J_{\eta \mathbf{a}_1} & \dots & J_{\eta \mathbf{a}_d} \\ J_{\mathbf{a}_1\eta} & J_{\mathbf{a}_1\mathbf{a}_1} & \dots & J_{\mathbf{a}_1\mathbf{a}_d} \\ \vdots & \vdots & \ddots & \vdots \\ J_{\mathbf{a}_d\eta} & J_{\mathbf{a}_d\mathbf{a}_1} & \dots & J_{\mathbf{a}_d\mathbf{a}_d} \end{pmatrix}.$$

Then it holds that

$$\left(\left\langle H \right\rangle / \left\langle H \right\rangle_{\chi\chi} \right) = \left\langle H \right\rangle_{\theta\theta} - \left\langle H \right\rangle_{\theta\chi} J \left\langle H \right\rangle_{\chi\theta}$$

$$= \left\langle A^{\top} A \right\rangle + \lambda I_d - \left\langle H \right\rangle_{\theta\eta} J_{\eta\eta} \left\langle H \right\rangle_{\eta\theta} - \sum_{j=1}^d \left\langle H \right\rangle_{\theta\eta} J_{\eta\mathbf{a}_j} \left\langle H \right\rangle_{\mathbf{a}_j\theta}$$

$$- \sum_{j=1}^d \left\langle H \right\rangle_{\theta\mathbf{a}_j} J_{\mathbf{a}_j\eta} \left\langle H \right\rangle_{\eta\theta} - \sum_{j=1}^d \left\langle H \right\rangle_{\theta\mathbf{a}_j} J_{\mathbf{a}_j\mathbf{a}_j} \left\langle H \right\rangle_{\mathbf{a}_j\theta} .$$

$$(91)$$

We simplify the terms in the right-hand side one by one using Proposition A.2, which ensures that J admits a nice decomposition. First, according to Proposition A.2, we have

$$\langle H \rangle_{\boldsymbol{\theta}\boldsymbol{\eta}} J_{\boldsymbol{\eta}\boldsymbol{\eta}} \langle H \rangle_{\boldsymbol{\eta}\boldsymbol{\theta}} = \left(\frac{1}{2} + r_1\right) \langle A \rangle^\top \langle A \rangle, \quad \text{where} \quad 0 \leqslant r_1 \leqslant \rho^2.$$

Next, using the identity

$$A = \sum_{j=1}^{d} \mathbf{e}_j \mathbf{a}_j^\top,$$

we observe that the same proposition yields that the third term in the right-hand side of (91) is equal to

$$\sum_{j=1}^{d} \langle H \rangle_{\boldsymbol{\theta}\boldsymbol{\eta}} J_{\boldsymbol{\eta}\mathbf{a}_{j}} \langle H \rangle_{\mathbf{a}_{j}\boldsymbol{\theta}} = -r_{2} \langle A \rangle^{\top} \sum_{j=1}^{d} \mathbf{e}_{j} \widetilde{\boldsymbol{\theta}}^{\top} \left\langle (\mathbf{a}_{j}^{\top} \boldsymbol{\theta} - \boldsymbol{\eta}_{j}) \right\rangle - r_{2} \langle A \rangle^{\top} \sum_{j=1}^{d} \mathbf{e}_{j} \widetilde{\boldsymbol{\theta}}^{\top} \left\langle \boldsymbol{\theta}\mathbf{a}_{j}^{\top} \right\rangle$$
$$= -r_{2} \langle A \rangle^{\top} \left\langle (A\boldsymbol{\theta} - \boldsymbol{\eta}) \right\rangle \widetilde{\boldsymbol{\theta}}^{\top} - r_{2} \langle A \rangle^{\top} \left\langle A \widetilde{\boldsymbol{\theta}}^{\top} \boldsymbol{\theta} \right\rangle, \tag{92}$$

where

$$\widetilde{\boldsymbol{\theta}} = \mu^2 \left(\mu^2 I_d + \left(\left\langle \boldsymbol{\theta} \boldsymbol{\theta}^\top \right\rangle \right) \right)^{-1} \left\langle \boldsymbol{\theta} \right\rangle.$$

Let us denote the rightmost side of (92) by $R_{(92)}$. The sum

$$\sum_{j=1}^{d} \langle H \rangle_{\boldsymbol{\theta} \mathbf{a}_{j}} J_{\mathbf{a}_{j} \boldsymbol{\eta}} \langle H \rangle_{\boldsymbol{\eta} \boldsymbol{\theta}}$$

is nothing but the transpose of the left-hand side of (92). Hence, it remains simplify the last term in (91):

$$\sum_{j=1}^{d} \langle H \rangle_{\boldsymbol{\theta} \mathbf{a}_{j}} J_{\mathbf{a}_{j} \mathbf{a}_{j}} \langle H \rangle_{\mathbf{a}_{j} \boldsymbol{\theta}} = \sum_{j=1}^{d} \left\langle (\mathbf{a}_{j}^{\top} \boldsymbol{\theta} - \boldsymbol{\eta}_{j}) \right\rangle^{2} R_{3} + \sum_{j=1}^{d} \left\langle (\mathbf{a}_{j}^{\top} \boldsymbol{\theta} - \boldsymbol{\eta}_{j}) \right\rangle R_{3} \left\langle \boldsymbol{\theta} \mathbf{a}_{j}^{\top} \right\rangle$$
$$+ \sum_{j=1}^{d} \left\langle \mathbf{a}_{j} \boldsymbol{\theta}^{\top} \right\rangle R_{3} \left\langle (\mathbf{a}_{j}^{\top} \boldsymbol{\theta} - \boldsymbol{\eta}_{j}) \right\rangle + \sum_{j=1}^{d} \left\langle \mathbf{a}_{j} \boldsymbol{\theta}^{\top} \right\rangle R_{3} \left\langle \boldsymbol{\theta} \mathbf{a}_{j}^{\top} \right\rangle \qquad(93)$$
$$= \left\| \left\langle A \boldsymbol{\theta} - \boldsymbol{\eta} \right\rangle \right\|^{2} R_{3} + R_{3} \left\langle \boldsymbol{\theta} \left\langle (A \boldsymbol{\theta} - \boldsymbol{\eta}) \right\rangle^{\top} A \right\rangle$$
$$+ \left\langle A^{\top} \left\langle (A \boldsymbol{\theta} - \boldsymbol{\eta}) \right\rangle \boldsymbol{\theta}^{\top} \right\rangle R_{3} + \left\langle A^{\top} \boldsymbol{\theta}^{\top} \right\rangle R_{3} \left\langle \boldsymbol{\theta} A \right\rangle. \qquad(94)$$

We denote the latter sum by $R_{(93)}$. Summing up (91), (92), and (93), we obtain the following identity:

$$(\Sigma^{2}/2 + \lambda I_{d})^{-1} \left(\langle H \rangle / \langle H \rangle_{\chi\chi} \right)$$

= $(\Sigma^{2}/2 + \lambda I_{d})^{-1} \left[\left(\langle A^{\top}A \rangle - \frac{1}{2} \langle A \rangle^{\top} \langle A \rangle + \lambda I_{d} \right) - r_{1} \langle A \rangle^{\top} \langle A \rangle - \mathbb{R}_{(92)} - \mathbb{R}_{(92)}^{\top} - \mathbb{R}_{(93)}^{\top} \right].$ (95)

Step 2. The leading term expansion. On this step, we focus on the leading term in the right-hand side of (95)

$$(\Sigma^2/2 + \lambda I_d)^{-1} \left(\left\langle A^\top A \right\rangle - \frac{1}{2} \left\langle A \right\rangle^\top \left\langle A \right\rangle + \lambda I_d \right)$$
(96)

leaving analysis of the remainders for future. We are going to show that (96) approximately equals I_d . To do so, we use the following proposition.

Proposition C.1. Assume that

$$\rho := \frac{7 \|\boldsymbol{\theta}^{\circ}\|}{\mu} \leqslant 1/7 \quad and \quad 7\rho \|\boldsymbol{\Sigma}\| \leqslant \sqrt{\lambda}.$$

Then the following holds. For any $s, t \in [0, 1]$ *, we have*

$$\|A^{(s)} - \Sigma\| \leq 4\rho^2 \sqrt{\lambda} + 5\rho \|\Sigma\| \|\mathbf{b}_{\lambda}\|/\mu \quad and \quad \|(A^{(s)})^{\top} A^{(t)} - \Sigma^2\| \leq 3\rho\lambda + 3\|\Sigma\| \|\mathbf{b}_{\lambda}\|\sqrt{\lambda}/\mu.$$

Applying Proposition C.1 and using the bound $\|(\Sigma^2/2 + \lambda I_d)^{-1}\| \leq \lambda^{-1}$, we infer

$$\begin{split} \left\| (\Sigma^2/2 + \lambda I_d)^{-1} \left(\left\langle A^\top A \right\rangle - \Sigma^2 \right) \right\| &\leq \int_0^1 \left\| (\Sigma^2/2 + \lambda I_d)^{-1} ((A^{(t)})^\top A^{(t)} - \Sigma^2) \right\| \mathrm{d}\mathfrak{m}(t) \\ &\leq \left(3\rho\lambda + \frac{3\|\Sigma\| \|\mathbf{b}_\lambda\| \sqrt{\lambda}}{\mu} \right) \| (\Sigma^2/2 + \lambda I_d)^{-1} \| \leq 3\rho + \frac{3\|\Sigma\| \|\mathbf{b}_\lambda\|}{\mu\sqrt{\lambda}}. \end{split}$$

Similarly, it holds that

$$\begin{split} \left\| (\Sigma^2/2 + \lambda I_d)^{-1} (\langle A \rangle^\top \langle A \rangle - \Sigma^2) \right\| &\leq \int_0^1 \int_0^1 \| (\Sigma^2/2 + \lambda I_d)^{-1} \| \| (A^{(s)})^\top A^{(t)} - \Sigma^2 \| \mathrm{d}\mathfrak{m}^{\otimes 2}(s, t) \\ &\leq 3\rho + \frac{3 \| \Sigma \| \| \mathbf{b}_\lambda \|}{\mu \sqrt{\lambda}}. \end{split}$$

Substituting these bounds into (95), we obtain that

$$\begin{split} \left\| (\Sigma^2/2 + \lambda I_d)^{-1} (\langle H \rangle / \langle H \rangle_{\chi\chi}) - I_d \right\| &= \left\| (\Sigma^2/2 + \lambda I_d)^{-1} \left[(\langle H \rangle / \langle H \rangle_{\chi\chi}) - (\Sigma^2/2 + \lambda I_d) \right] \right\| \\ &\leq \frac{9\rho}{2} + \frac{3\|\Sigma\| \|\mathbf{b}_\lambda\|}{2\mu\sqrt{\lambda}} + r_1 \| (\Sigma^2/2 + \lambda I_d)^{-1}\Sigma^2 \| \\ &+ \left(3\rho + \frac{3\|\Sigma\| \|\mathbf{b}_\lambda\|}{\mu\sqrt{\lambda}} \right) \cdot r_1 + \left\| (\Sigma^2/2 + \lambda I_d)^{-1} \mathbf{R}_{(92)} \right\| \\ &+ \left\| (\Sigma^2/2 + \lambda I_d)^{-1} \mathbf{R}_{(92)}^\top \| + \| (\Sigma^2/2 + \lambda I_d)^{-1} \mathbf{R}_{(93)} \right\|. \end{split}$$

Let us recall that, according to Proposition A.2, $r_1 \leq \rho^2$. This inequality, together with the bound

$$\left\| (\Sigma^2/2 + \lambda I_d)^{-1} \Sigma^2 \right\| \leqslant 2,$$

yields that

$$\left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} (\langle H \rangle / \langle H \rangle_{\chi\chi}) - I_{d} \right\| \leq 6\rho + \frac{2 \|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu \sqrt{\lambda}} + \left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \,\mathbf{R}_{(92)} \right\| + \left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \,\mathbf{R}_{(93)} \right\|.$$
(97)

Step 3. Bounding the remainder I. We start bounding the term $\|(\Sigma^2/2 + \lambda I_d)^{-1} \mathbb{R}_{(92)}\|$, and then proceed with $\|(\Sigma^2/2 + \lambda I_d)^{-1} \mathbb{R}_{(92)}^{\top}\|$. We have

$$\left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \mathbf{R}_{(92)} \right\| \leq r_{2} \left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \right\| \left\| \langle A \rangle^{\top} \left\langle (A\boldsymbol{\theta} - \boldsymbol{\eta}) \right\rangle \widetilde{\boldsymbol{\theta}}^{\top} \right\| + r_{2} \left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \left\langle A \right\rangle^{\top} \left\langle A \boldsymbol{\theta}^{\top} \widetilde{\boldsymbol{\theta}} \right\rangle \right\|.$$
(98)

We bound r_2 by μ^{-2} . Next, we bound the first term of the expression above. We have

$$r_2 \left\| (\Sigma^2/2 + \lambda I_d)^{-1} \right\| \left\| \langle A \rangle^\top \left\langle (A\boldsymbol{\theta} - \boldsymbol{\eta}) \right\rangle \widetilde{\boldsymbol{\theta}}^\top \right\| \leq \frac{\| \langle A \rangle \| \| \langle A\boldsymbol{\theta} - \boldsymbol{\eta} \rangle \| \| \boldsymbol{\theta} \|}{\mu^2 \lambda} \leq \frac{\| \langle A \rangle \| \| \langle A\boldsymbol{\theta} - \boldsymbol{\eta} \rangle \| \| \boldsymbol{\theta} \|}{\mu^2 \lambda}.$$

Since $(\theta^{(t)}, \eta^{(t)}, A^{(t)}) \in \Theta^* \times H^* \times A^*$, we have $||\langle A \rangle || \leq ||\Sigma|| + \sqrt{\lambda}/3$ by the definition of A^{*}, and $||\langle \theta \rangle || \leq ||\theta^\circ|| \leq \rho \mu$ since $||\theta^*|| \leq ||\theta^\circ||$ by Lemma B.2(iv). To bound $||\langle A\theta - \eta \rangle||$, we use the following proposition.

Proposition C.2. Assume that

$$\rho := \frac{7 \|\boldsymbol{\theta}^{\circ}\|}{\mu} \leqslant 1 \quad and \quad \|\boldsymbol{\Sigma}\| \|\boldsymbol{\theta}^{\circ}\| \leqslant \mu \sqrt{\lambda}/24.$$

For any $t \in [0, 1]$, we have

$$\|A^{(t)}\boldsymbol{\theta}^{(t)} - \boldsymbol{\eta}^{(t)}\| \leq \rho \mu \sqrt{\lambda}/2 + 3\|\boldsymbol{\Sigma}\| \|\mathbf{b}_{\lambda}\|.$$

It implies

$$r_{2} \left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \right\| \cdot \left\| \langle A \rangle^{\top} \left\langle (A\boldsymbol{\theta} - \boldsymbol{\eta}) \right\rangle \widetilde{\boldsymbol{\theta}}^{\top} \right\| \leq \frac{\left(\|\Sigma\| + \sqrt{\lambda}/3 \right) \cdot (\rho\mu\sqrt{\lambda}/2 + \|\Sigma\| \|\mathbf{b}_{\lambda}\|) \cdot \rho\mu}{\mu^{2}\lambda}$$

$$\leq \frac{\rho^{2} \|\Sigma\|}{2\sqrt{\lambda}} + \rho^{2}/6 + \frac{\rho\|\Sigma\|^{2} \|\mathbf{b}_{\lambda}\|}{\mu\lambda} + \frac{\rho\|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu\sqrt{\lambda}}$$

$$\leq \rho/14 + \rho^{2}/6 + \frac{\|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu\sqrt{\lambda}}$$

$$\leq \rho/6 + \frac{\|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu\sqrt{\lambda}}, \qquad (99)$$

where we used the condition $7\rho \|\Sigma\| \leq \sqrt{\lambda}$.

Next, we bound
$$r_2 \left\| (\Sigma^2/2 + \lambda I_d)^{-1} \langle A \rangle^\top \left\langle A \boldsymbol{\theta}^\top \widetilde{\boldsymbol{\theta}} \right\rangle \right\|$$
. We have

$$\begin{aligned} r_2 \left\| (\Sigma^2/2 + \lambda I_d)^{-1} \langle A \rangle^\top \left\langle A \boldsymbol{\theta}^\top \widetilde{\boldsymbol{\theta}} \right\rangle \right\| \\ &\leqslant \mu^{-2} \int_0^1 \int_0^1 \left\| (\Sigma^2/2 + \lambda I_d)^{-1} (A^{(s)})^\top A^{(t)} \cdot \widetilde{\boldsymbol{\theta}}^\top \boldsymbol{\theta}^{(t)} \right\| \, \mathrm{d}\mathfrak{m}^{\otimes 2}(s, t) \\ &\leqslant \left\| (\Sigma^2/2 + \lambda I_d)^{-1} \Sigma^2 \right\| \cdot \mu^{-2} \int_0^1 |\widetilde{\boldsymbol{\theta}}^\top \boldsymbol{\theta}^{(t)}| \, \mathrm{d}\mathfrak{m}(t) \\ &+ \mu^{-2} \left\| (\Sigma^2/2 + \lambda I_d)^{-1} \right\| \int_0^1 \int_0^1 \| (A^s)^\top A^{(t)} - \Sigma^2 \| \cdot |\widetilde{\boldsymbol{\theta}}^\top \boldsymbol{\theta}^{(t)}| \, \mathrm{d}\mathfrak{m}^{\otimes 2}(s, t). \end{aligned}$$

By the Cauchy-Schwarz inequality and bound $\|\tilde{\boldsymbol{\theta}}\| \leq \rho \mu$ from Proposition A.2, we have $\mu^{-2}|\tilde{\boldsymbol{\theta}}^{\top}\boldsymbol{\theta}^{(t)}| \leq \rho \mu^{-1} \|\boldsymbol{\theta}^{(t)}\|$. The latter is at most $\rho^2/7$, since $\|\boldsymbol{\theta}^{(t)}\| \leq \max\{\|\boldsymbol{\theta}^*\|, \|\boldsymbol{\theta}^\circ\|\} \leq \|\boldsymbol{\theta}^\circ\| \leq \rho \mu/7$ from Lemma B.2(iv). Bounding $\|(\Sigma^2/2 + \lambda I_d)^{-1}\Sigma^2\|$ by 1 and applying Proposition C.1, we obtain

$$r_{2} \left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \langle A \rangle^{\top} \left\langle A \boldsymbol{\theta}^{\top} \widetilde{\boldsymbol{\theta}} \right\rangle \right\| \leq \rho^{2} + \rho^{2} \cdot \left(3\rho\lambda + \frac{3\|\Sigma\| \|\mathbf{b}_{\lambda}\|\sqrt{\lambda}}{\mu} \right) \left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \right\| \leq \rho^{2} + 3\rho^{3} + \rho^{2}/4 \leq 3\rho,$$
(100)

where we used $\|\Sigma\| \|\mathbf{b}_{\lambda}\| \leq \|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| \leq \mu \sqrt{\lambda}/24$ from (25). Substituting bounds (99), (100) in (98), we get

$$\|(\Sigma^2/2 + \lambda I_d)^{-1} \mathbf{R}_{(92)}\| \leqslant 7\rho/2 + \frac{\|\Sigma\| \|\mathbf{b}_\lambda\|}{\mu\sqrt{\lambda}}.$$
(101)

Then, we proceed with the term $\|(\Sigma^2/2 + \lambda I_d)^{-1} \mathbf{R}_{(92)}^{\top}\|$. We have

$$\| (\Sigma^2/2 + \lambda I_d)^{-1} \mathbf{R}_{(92)}^\top \| \leq r_2 \| (\Sigma^2/2 + \lambda I_d)^{-1} \| \| \langle A \rangle^\top \langle A \boldsymbol{\theta} - \boldsymbol{\eta} \rangle \widetilde{\boldsymbol{\theta}}^\top \| + r_2 \| (\Sigma^2/2 + \lambda I_d)^{-1} \left\langle \boldsymbol{\theta}^\top \widetilde{\boldsymbol{\theta}} A^\top \right\rangle \langle A \rangle^\top \|.$$

The first term can be bounded by (99). We bound the second term using the fact that for each $s \in [0, 1]$, the quantity $(\boldsymbol{\theta}^{(s)})^{\top} \widetilde{\boldsymbol{\theta}}$ is scalar. Hence, we have

$$\begin{split} \|(\Sigma^2/2 + \lambda I_d)^{-1} \left\langle \boldsymbol{\theta}^\top \widetilde{\boldsymbol{\theta}} A^\top \right\rangle \left\langle A \right\rangle \| &= \left\| \int_0^1 \int_0^1 \widetilde{\boldsymbol{\theta}}^\top \boldsymbol{\theta}^{(s)} \left(\Sigma^2/2 + \lambda I_d \right)^{-1} (A^{(s)})^\top A^{(t)} \mathrm{d} \mathfrak{m}^{\otimes 2}(s, t) \right\| \\ &\leq \rho^2 \mu^2/7 \int_0^1 \int_0^1 \int_0^1 \left\| (\Sigma^2/2 + \lambda I_d)^{-1} (A^{(s)})^\top A^{(t)} \right\| \mathrm{d} \mathfrak{m}^{\otimes 2}(s, t), \end{split}$$

where we used $\|\boldsymbol{\theta}^{(s)}\| \leq \max\{\|\boldsymbol{\theta}^*\|, \|\boldsymbol{\theta}^\circ\|\} \leq \rho\mu/7$ from Lemma B.2(iv) and $\|\widetilde{\boldsymbol{\theta}}\| \leq \rho\mu$ from Proposition A.2. Then, we bound $\|(\Sigma^2/2 + \lambda I_d)^{-1}(A^{(s)})^\top A^{(t)}\|$ as before:

$$\begin{split} \left\| (\Sigma^2/2 + \lambda I_d)^{-1} (A^{(s)})^\top A^{(t)} \right\| &\leq \left\| (\Sigma^2/2 + \lambda I_d)^{-1} \Sigma^2 \right\| + \left\| (\Sigma^2/2 + \lambda I_d)^{-1} \left((A^{(s)})^\top A^{(t)} - \Sigma^2 \right) \right\| \\ &\leq 1 + \frac{1}{\lambda} \left(3\rho\lambda + \frac{3\|\Sigma\| \|\mathbf{b}_\lambda\| \sqrt{\lambda}}{\mu} \right) = 1 + 3\rho + \frac{3\|\Sigma\| \|\mathbf{b}_\lambda\|}{\mu\sqrt{\lambda}}, \end{split}$$

where we used $\|(\Sigma^2/2 + \lambda I_d)^{-1}\| \leq \lambda^{-1}$ and Proposition C.1 for the second inequality. Hence, we have

$$\|(\Sigma^2/2 + \lambda I_d)^{-1} \mathbf{R}_{(92)}^{\top}\| \leq \rho/6 + \frac{\|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu\sqrt{\lambda}} + \rho^2/7 + 3\rho^3/7 + \frac{3\rho^2 \|\Sigma\| \|\mathbf{b}_{\lambda}\|}{7\mu\sqrt{\lambda}}$$

Bounding $\|\Sigma\| \|\mathbf{b}_{\lambda}\| \leq \|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| \leq \mu \sqrt{\lambda}/24$ due to (25), we derive

$$\|(\Sigma^2/2 + \lambda I_d)^{-1} \mathbf{R}_{(92)}^{\top}\| \leqslant 7\rho/2 + \frac{\|\Sigma\| \|\mathbf{b}_\lambda\|}{\mu\sqrt{\lambda}}.$$
(102)

Step 4. Bounding the remainder II. Next, we analyze $\|(\Sigma^2/2 + \lambda I_d)^{-1} \mathbb{R}_{(93)}\|$. The first three terms of (93), can be bounded in the same way:

$$\left\| (\Sigma^2/2 + \lambda I_d)^{-1} \left\{ \left\| \left\langle A\boldsymbol{\theta} - \boldsymbol{\eta} \right\rangle \right\|^2 R_3 + R_3 \left\langle \boldsymbol{\theta} \left\langle (A\boldsymbol{\theta} - \boldsymbol{\eta}) \right\rangle^\top A \right\rangle + \left\langle A^\top \left\langle (A\boldsymbol{\theta} - \boldsymbol{\eta}) \right\rangle \boldsymbol{\theta}^\top \right\rangle R_3 \right\} \right\| \\ \leqslant \frac{1}{\lambda} \left\{ \left\| \left\langle A\boldsymbol{\theta} - \boldsymbol{\eta} \right\rangle \right\|^2 \cdot \left\| R_3 \right\| + 2 \| R_3 \| \int_0^1 \| \boldsymbol{\theta}^{(t)} \| \left\| \left\langle A\boldsymbol{\theta} - \boldsymbol{\eta} \right\rangle \| \| A^{(t)} \| \operatorname{d} \mathfrak{m}(t) \right\}.$$

Due to Proposition C.2, the quantity $\|\langle A\theta - \eta \rangle\|$ is at most $\rho \mu \sqrt{\lambda}/2 + \|\Sigma\| \|\mathbf{b}_{\lambda}\|$, and

$$\|\langle A\boldsymbol{\theta} - \boldsymbol{\eta} \rangle \|^2 \leqslant \rho^2 \mu^2 \lambda / 4 + \rho \mu \sqrt{\lambda} \|\Sigma\| \|\mathbf{b}_{\lambda}\| + \|\Sigma\|^2 \|\mathbf{b}_{\lambda}\|^2.$$

Next, the norm $||R_3||$ is at most $2\mu^{-2}$ due to Proposition A.2 and $||\Sigma|| ||\mathbf{b}_{\lambda}|| \leq ||\Sigma|| ||\boldsymbol{\theta}^{\circ}|| \leq \mu \sqrt{\lambda}/24$ due to (25), therefore, the above is at most

$$\frac{1}{\lambda} \left\{ \rho^2 \lambda/2 + \frac{\|\Sigma\| \|\mathbf{b}_{\lambda}\| \sqrt{\lambda}}{\mu} + \left(\rho \sqrt{\lambda} \mu^{-1} + \frac{2\|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu^2} \right) \int_0^1 \|\boldsymbol{\theta}^{(t)}\| \|A^{(t)}\| \mathrm{d}\mathfrak{m}(t) \right\}$$

Again, we use $A^{(t)} \in A^*$, and bound $||A^{(t)}|| \leq ||\Sigma|| + \sqrt{\lambda}/3$. Since $||\theta^*|| \leq ||\theta^\circ||$ due to Lemma B.2(iv), $||\theta^{(t)}|| \leq \rho \mu/7$. Hence, the above is at most

$$\begin{split} \rho^2/2 + \left(\frac{\rho^2}{\sqrt{\lambda}} + \frac{2\rho \|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu\lambda}\right) \left\{ \|\Sigma\| + \sqrt{\lambda}/3 \right\} &\leq \rho^2/2 + \frac{\rho}{7} + \rho^2/3 + \frac{\|\Sigma\| \|\mathbf{b}_{\lambda}\|}{7\mu\sqrt{\lambda}} + \frac{2\rho \|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu\sqrt{\lambda}} \\ &\leq \rho + \frac{\|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu\sqrt{\lambda}}, \end{split}$$

where we used the condition $7\rho \|\Sigma\| \leq \sqrt{\lambda}$. Thus, the bound

$$\left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \left\{ \| \langle A\boldsymbol{\theta} - \boldsymbol{\eta} \rangle \|^{2} R_{3} + R_{3} \left\langle \boldsymbol{\theta} \langle (A\boldsymbol{\theta} - \boldsymbol{\eta}) \rangle^{\top} A \right\rangle + \left\langle A^{\top} \langle (A\boldsymbol{\theta} - \boldsymbol{\eta}) \rangle \boldsymbol{\theta}^{\top} \right\rangle R_{3} \right\} \right\|$$

$$\leq \rho + \frac{\|\Sigma\| \| \mathbf{b}_{\lambda} \|}{\mu \sqrt{\lambda}}$$

$$(103)$$

holds. Finally, we bound the norm of $(\Sigma^2/2 + \lambda I_d)^{-1} \langle A^\top \boldsymbol{\theta}^\top \rangle R_3 \langle \boldsymbol{\theta} A \rangle$. Applying the triangle inequality, we derive

As before, we use the definition of A^* , to get bound on the norm of $A^{(t)} \in A^*$ and $A^{(t)} - \Sigma$, and, as a consequence, obtain

$$\|\langle \boldsymbol{\theta} A \rangle \| \leq \int_0^1 \|\boldsymbol{\theta}^{(t)}\| \|A^{(t)}\| \mathrm{d}\mathfrak{m}(t) \leq \rho \mu \left\{ \|\Sigma\| + \sqrt{\lambda}/3 \right\}.$$

Next, we have $\|R_3\| \leqslant 2\mu^{-2}$ and $\|A^{(t)} - \Sigma\| \leqslant \sqrt{\lambda}/3$, hence, it yields

$$\left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \left\langle A^{\top} \boldsymbol{\theta}^{\top} \right\rangle R_{3} \left\langle \boldsymbol{\theta} A \right\rangle \right\| \leq 2\rho^{2} \left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \Sigma \right\| \left\{ \|\Sigma\| + \sqrt{\lambda}/3 \right\} + \frac{2\rho^{2}\sqrt{\lambda}}{3} \left\| (\Sigma^{2}/2 + \lambda I_{d})^{-1} \right\| \left\{ \|\Sigma\| + \sqrt{\lambda}/3 \right\}.$$
(104)

Let $\lambda_1(\Sigma) \ge \ldots \ge \lambda_1(\Sigma)$ stand for the eigenvalues of Σ . Then, due to the Cauchy-Schwarz inequality, it holds that

$$\left| (\Sigma^2/2 + \lambda I_d)^{-1} \Sigma \right\| = \max_{1 \leq j \leq d} \left\{ \frac{\lambda_j(\Sigma)}{\lambda_j(\Sigma)^2/2 + \lambda} \right\} \leq \sqrt{\frac{2}{\lambda}}$$

Applying this bound and the inequality $\|(\Sigma^2/2 + \lambda I_d)^{-1}\| \leq 1/\lambda$ to (104), we deduce

$$\left\| (\Sigma^2/2 + \lambda I_d)^{-1} \left\langle A^\top \boldsymbol{\theta}^\top \right\rangle R_3 \left\langle \boldsymbol{\theta} A \right\rangle \right\| \leq \frac{3\rho^2}{\sqrt{\lambda}} \left\{ \|\Sigma\| + \sqrt{\lambda}/3 \right\} \leq \frac{3\rho}{7} + \rho^2 \leq \rho.$$
(105)

As before, we took into account that $7\rho \|\Sigma\| \leq \sqrt{\lambda}$ and $\rho \leq 1/2$ due to the proposition conditions. Thus, (103) and (105) imply that

$$\|(\Sigma^2/2 + \lambda I_d)^{-1} \mathbf{R}_{(93)}\| \leq 2\rho + \frac{\|\Sigma\| \|\mathbf{b}_\lambda\|}{\mu\sqrt{\lambda}}.$$
(106)

Substituting the bounds (101), (102), and (106) into (97), we obtain the desired bound:

$$\left\| (\Sigma^2/2 + \lambda I_d)^{-1} (\langle H \rangle^{\mathfrak{m}} / \langle H \rangle_{\chi\chi}^{\mathfrak{m}}) - I_d \right\| \leq 15\rho + \frac{5 \|\Sigma\| \|\mathbf{b}_\lambda\|}{\mu \sqrt{\lambda}}.$$

The proof is finished.

C.2 Proof of Proposition C.1

Due to the definition of $A^{(t)}$ (see (90)), we clearly have $||A^{(t)} - \Sigma|| = t ||A^* - \Sigma||$ for all $t \in [0, 1]$. Therefore, according to Lemma B.3, it holds that

$$\|A^{(t)} - \Sigma\| \leq \|A^* - \Sigma\| \leq 4\rho^2 \sqrt{\lambda} + \frac{5\rho \|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu}.$$

It only remains to consider $||(A^{(s)})^{\top}A^{(t)} - \Sigma^2||$. The triangle yields that

$$\|(A^{(s)})^{\top}A^{(t)} - \Sigma^{2}\| = \|(A^{(s)} - \Sigma)A^{(t)} + \Sigma(A^{(t)} - \Sigma)\|$$

$$\leq \|A^{(s)} - \Sigma\|\|A^{(t)}\| + \|\Sigma\|\|A^{(t)} - \Sigma\|$$

$$\leq \|A^{*} - \Sigma\|(\|A^{*}\| + \|\Sigma\|).$$

Since, according to Lemma B.1 applied with $\rho_0 = 1/7$, A^* and Σ belong to the set A^* , we have $||A^* - \Sigma|| \leq \sqrt{\lambda}/3$ and then

$$\|\Sigma\| + \|A^*\| \leq 2\|\Sigma\| + \|A^* - \Sigma\| \leq 2\|\Sigma\| + \frac{\sqrt{\lambda}}{3}$$

This yields that

$$\begin{split} \|(A^{(s)})^{\top}A^{(t)} - \Sigma^2\| &= \|A^* - \Sigma\| \left(\|A^*\| + \|\Sigma\| \right) \\ &\leq \left(4\rho^2 \sqrt{\lambda} + \frac{5\rho \|\Sigma\| \|\mathbf{b}_{\lambda}\|}{\mu} \right) \left(2\|\Sigma\| + \sqrt{\lambda}/3 \right) \\ &\leq 8\rho^2 \|\Sigma\| \sqrt{\lambda} + 4\rho^2 \lambda/3 + \frac{10\rho \|\Sigma\|^2 \|\mathbf{b}_{\lambda}\|}{\mu} + \frac{5\rho \|\Sigma\| \|\mathbf{b}_{\lambda}\| \sqrt{\lambda}}{3\mu}. \end{split}$$

Since $7\rho \|\Sigma\| \leq \sqrt{\lambda}$ and $\rho \leq 1/7$, the latter is at most $3\rho\lambda + 3\|\Sigma\| \|\mathbf{b}_{\lambda}\| \sqrt{\lambda}/\mu$.

C.3 Proof of Proposition C.2

Due to the triangle inequality, $||A^{(t)}\theta^{(t)} - \eta^{(t)}||$ satisfies

$$\begin{aligned} \|A^{(t)}\boldsymbol{\theta}^{(t)} - \boldsymbol{\eta}^{(t)}\| &\leq \|A^{(t)} - \boldsymbol{\Sigma}\| \|\boldsymbol{\theta}^{(t)}\| + \|\boldsymbol{\Sigma}\boldsymbol{\theta}^{(t)} - \boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ}\| + \|\boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ} - \boldsymbol{\eta}^{(t)}\| \\ &= t\|A^{*} - \boldsymbol{\Sigma}\| \|\boldsymbol{\theta}^{(t)}\| + t\|\boldsymbol{\Sigma}\boldsymbol{\theta}^{*} - \boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ}\| + t\|\boldsymbol{\Sigma}\boldsymbol{\theta}^{\circ} - \boldsymbol{\eta}^{*}\|. \end{aligned}$$

Since $\eta^* = (A^* \theta^* + \Sigma \theta^\circ)/2$, it holds that

$$\begin{aligned} \|A^{(t)}\boldsymbol{\theta}^{(t)} - \boldsymbol{\eta}^{(t)}\| &\leq \|A^* - \boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^{(t)}\| + \|\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ\| + \frac{1}{2}\|\boldsymbol{\Sigma}\boldsymbol{\theta}^\circ - A^*\boldsymbol{\theta}^*\| \\ &\leq \|A^* - \boldsymbol{\Sigma}\|\left(\|\boldsymbol{\theta}^{(t)}\| + \frac{\|\boldsymbol{\theta}^*\|}{2}\right) + \frac{3}{4}\|\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^\circ - \boldsymbol{\theta}^*\| \end{aligned}$$

Let us recall that $\theta^{(t)}$ is a convex combination of θ^* and θ° , and $\|\theta^*\| \leq \|\theta^\circ\|$ in view of Lemma B.2(iv). Moreover, we have $\|\theta^{(t)}\| \leq \|\theta^\circ\| = \rho\mu/7$ due to the conditions of the proposition. Applying Lemma B.2(iv) to $\|\theta^* - \theta^\circ\|$, we obtain that

$$\|A^{(t)}\boldsymbol{\theta}^{(t)} - \boldsymbol{\eta}^{(t)}\| \leq \frac{3\rho\mu\|A^* - \Sigma\|}{14} + 3\|\Sigma\|\|\mathbf{b}_{\lambda}\| + 3\rho\|\Sigma\|\|\boldsymbol{\theta}^\circ\|.$$

Using the fact that $\|\Sigma\| \| \boldsymbol{\theta}^{\circ} \| \leq \mu \sqrt{\lambda}/24$ by the assumptions of Lemma B.1, we obtain that

 $3\rho \|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| \leqslant \rho \mu \sqrt{\lambda}/8.$

Finally, the chain of the inequalities

$$\|A^* - \Sigma\| \leqslant \mu^{-1} \sqrt{\lambda} \|\boldsymbol{\theta}^{\circ}\| \leqslant \rho \sqrt{\lambda} / 7$$

following from Lemma B.2, implies that

$$\|A^{(t)}\boldsymbol{\theta}^{(t)} - \boldsymbol{\eta}^{(t)}\| \leq \rho \mu \sqrt{\lambda}/2 + 3 \|\Sigma\| \|\mathbf{b}_{\lambda}\|.$$

D Results from the proof of Theorem 2.4

This section contained proof of auxiliary results related to Theorem 2.4.

D.1 Preliminaries

We start with the following result playing an important role in the proof of Lemma 6.1.

Lemma D.1. Let us fix any $\rho \in [0, 1/16]$ and introduce a block diagonal matrix

$$\mathsf{D}^2_* = \operatorname{diag}\left((A^*)^\top A^* + \lambda I_d, 2I_d, I_d \otimes \left(\mu^2 I_d + \boldsymbol{\theta}^*(\boldsymbol{\theta}^*)^\top\right)\right).$$

Consider an event \mathcal{E} such that v^* and \hat{v} belong to a convex set $\mathcal{U} \subset \Upsilon(\rho)$ and $171 \| \mathsf{D}_*^{-1} \mathfrak{z} \| \leq \mu \sqrt{\lambda}$ on \mathcal{E} . Then, on this event, it holds that

$$\|\mathsf{D}_*(\widehat{\boldsymbol{\upsilon}}-\boldsymbol{\upsilon}^*)\| \leqslant 9\|\mathsf{D}_*^{-1}\mathfrak{z}\|.$$

Proof of Lemma D.1. Let us denote $r = 9 \| \mathbb{D}_*^{-1} \mathfrak{z} \|$ for brevity and show that $\| \mathbb{D}(\widehat{\boldsymbol{v}} - \boldsymbol{v}^*) \| \leq r$ on the event \mathcal{E} . For this purpose, we fix an arbitrary vector $\mathbf{u} \in \mathbb{R}^{2d+d^2}$ of the norm $\| \mathbf{u} \| = r$ and consider a function

$$f(t) = \mathbf{L}(\boldsymbol{\upsilon}^* + t \, \mathbf{D}_*^{-1} \mathbf{u}), \quad t > 0.$$

We are going to prove that $f'(t) \neq 0$ for any t > 1 such that $v^* + t D^{-1} u \in U$. This yields the desired bound, because \hat{v} is a stationary point of L. Our approach is based on the analysis of the first and the second order derivatives:

$$f'(t) = \boldsymbol{\nabla} \mathbf{L}(\boldsymbol{\upsilon}^* + t \, \mathbf{D}_*^{-1} \mathbf{u})^\top \mathbf{D}_*^{-1} \mathbf{u} \quad \text{and} \quad f''(t) = \mathbf{u}^\top \mathbf{D}_*^{-1} \nabla^2 \mathbf{L}(\boldsymbol{\upsilon}^* + t \, \mathbf{D}_*^{-1} \mathbf{u}) \mathbf{D}_*^{-1} \mathbf{u}.$$

According to Taylor's formula with the integral remainder term, it holds that

$$f'(1) - f'(0) - f''(0) = \int_{0}^{1} \left(f''(s) - f''(0) \right) \mathrm{d}s.$$
(107)

Let us fix any $s \in (0, 1)$ and consider (f''(s) - f''(0)). Applying Lemma A.3 with $\lambda_0 = \lambda$ and $\mu_0 = \mu$, we obtain that

$$\begin{split} \left| f''(s) - f''(0) \right| &= \left| \mathbf{u}^\top \mathsf{D}_*^{-1} \left(\nabla^2 \mathsf{L}(\boldsymbol{\upsilon}^* + ts \, \mathsf{D}_*^{-1} \mathbf{u}) - \nabla^2 \mathsf{L}(\boldsymbol{\upsilon}^*) \right) \mathsf{D}_*^{-1} \mathbf{u} \right| \\ &\leqslant \frac{4s \, \|\mathbf{u}\|^3}{\mu \sqrt{\lambda}} + \frac{2s^2 \, \|\mathbf{u}\|^4}{\mu^2 \lambda} \\ &= \frac{4s \, r^3}{\mu \sqrt{\lambda}} + \frac{2s^2 \, r^4}{\mu^2 \lambda}. \end{split}$$

Substituting this bound into (107), we observe that

$$\left| f'(1) - f'(0) - f''(0) \right| \leqslant \frac{4r^3}{\mu\sqrt{\lambda}} \int_0^1 s \, \mathrm{d}s + \frac{2r^4}{\mu^2\lambda} \int_0^1 s^2 \, \mathrm{d}s = \frac{2r^3}{\mu\sqrt{\lambda}} \left(1 + \frac{r}{3\mu\sqrt{\lambda}} \right)$$

Since $r = 9 \|\mathbf{D}_*^{-1}\mathfrak{z}\|$ and $171 \|\mathbf{D}_*^{-1}\mathfrak{z}\| \leqslant \mu \sqrt{\lambda}$ on the event \mathcal{E} , it holds that

$$\left|f'(1) - f'(0) - f''(0)\right| \leq \frac{2r^2}{19}\left(1 + \frac{1}{57}\right) = \frac{r^2}{19} \cdot \frac{58}{57}.$$

Note that Lemma A.1 yields

$$f''(0) = \mathbf{u}^{\top} \mathbf{D}_*^{-1} \nabla^2 \mathbf{L}(\boldsymbol{\upsilon}^*) \mathbf{D}_*^{-1} \mathbf{u} \ge \frac{(1-2\rho) \|\mathbf{u}\|^2}{4} = \frac{(1-2\rho)r^2}{4}$$

Taking into account this inequality and the relation

$$|f'(0)| = \left| \boldsymbol{\nabla} \mathbf{L}(\boldsymbol{\upsilon}^*)^\top \mathbf{D}_*^{-1} \mathbf{u} \right| = \left| \boldsymbol{\mathfrak{z}}^\top \mathbf{D}_*^{-1} \mathbf{u} \right| \leqslant \| \mathbf{D}_*^{-1} \boldsymbol{\mathfrak{z}} \| \| \mathbf{u} \| \leqslant \frac{r^2}{9},$$

we obtain that

$$f'(1) \ge \frac{(1-2\rho)r^2}{4} - \frac{r^2}{10} - \frac{2r^2}{19} \cdot \frac{58}{57}$$

It is straightforward to check that

$$\frac{2}{19} \cdot \frac{58}{57} = \frac{6 \cdot 58}{57^2} \leqslant \frac{6 \cdot 58}{57^2 - 1} = \frac{6}{56} = \frac{3}{28}.$$

Thus, it holds that

$$f'(1) = \left(\frac{1}{4} - \frac{1}{9} - \frac{3}{28} - \frac{\rho}{2}\right)r^2 = \left(\frac{5}{36} - \frac{3}{28} - \frac{\rho}{2}\right)r^2 = \left(\frac{2}{63} - \frac{\rho}{2}\right)r^2 \ge \frac{(1 - 16\rho)r^2}{32} \ge 0.$$

Since f''(t) > 0 for any t > 0 such that $\boldsymbol{v}^* + t \, \mathsf{D}_*^{-1} \mathbf{u} \in \mathcal{U}$, the Lagrange mean value theorem yields that $f'(t) \ge f'(1) > 0$ for any $t \ge 1$ satisfying $\boldsymbol{v}^* + t \, \mathsf{D}_*^{-1} \mathbf{u} \in \mathcal{U}$. Hence, we deduce that $\|\mathsf{D}_*(\widehat{\boldsymbol{v}} - \boldsymbol{v}^*)\| \le r$.

D.2 Proof of Lemma 6.1

For convenience, we split the proof into several steps.

Step 1: Taylor's expansion. Let us remind a reader that $H_* = \nabla^2 \mathcal{L}(\boldsymbol{v}^*) = \nabla^2 \mathcal{L}(\boldsymbol{v}^*)$. Due to the definition of $\hat{\boldsymbol{v}}$, it holds that $\nabla \mathcal{L}(\hat{\boldsymbol{v}}) = \mathbf{0}$. Using the Newton-Leibniz rule, we obtain that

$$\begin{aligned} \nabla \mathsf{L}(\boldsymbol{\upsilon}^* + H_*^{-1}\boldsymbol{\mathfrak{z}}) &= \nabla \mathsf{L}(\boldsymbol{\upsilon}^* + H_*^{-1}\boldsymbol{\mathfrak{z}}) - \nabla \mathsf{L}(\widehat{\boldsymbol{\upsilon}}) \\ &= \int_0^1 \nabla^2 \mathsf{L}\big(t\widehat{\boldsymbol{\upsilon}} + (1-t)(\boldsymbol{\upsilon}^* + H_*^{-1}\boldsymbol{\mathfrak{z}}\big)(\boldsymbol{\upsilon}^* - \widehat{\boldsymbol{\upsilon}} + H_*^{-1}\boldsymbol{\mathfrak{z}}) \mathrm{d}t \\ &= \int_0^1 \nabla^2 \mathsf{L}\big(\boldsymbol{\upsilon}^* + t(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^*) + (1-t)H_*^{-1}\boldsymbol{\mathfrak{z}}\big)(\boldsymbol{\upsilon}^* - \widehat{\boldsymbol{\upsilon}} + H_*^{-1}\boldsymbol{\mathfrak{z}}) \mathrm{d}t. \end{aligned}$$

Thus, it holds that

$$\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^* - H_*^{-1} \boldsymbol{\mathfrak{z}} = - (H_* + R)^{-1} \, \boldsymbol{\nabla} \mathsf{L}(\boldsymbol{\upsilon}^* + H_*^{-1} \boldsymbol{\mathfrak{z}}),$$

where we introduced

$$R = \int_{0}^{1} \left[\nabla^{2} \mathsf{L} \left(\boldsymbol{\upsilon}^{*} + t(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^{*}) + (1 - t)H_{*}^{-1} \boldsymbol{\mathfrak{z}} \right) - H_{*} \right] \mathrm{d}t$$

The representation

$$-(H_*+R)^{-1} = -H_*^{-1}H_*(H_*+R)^{-1} = -H_*^{-1} + H_*^{-1}R(H_*+R)^{-1}$$

yields the equality

$$\widehat{\boldsymbol{v}} - \boldsymbol{v}^* - H_*^{-1}\boldsymbol{z} = -H_*^{-1}\boldsymbol{\nabla} L(\boldsymbol{v}^* + H_*^{-1}\boldsymbol{z}) + H_*^{-1}R(H_* + R)^{-1}\boldsymbol{\nabla} L(\boldsymbol{v}^* + H_*^{-1}\boldsymbol{z}).$$

Then it holds that

$$\begin{split} \left\| S \big(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^* - H_*^{-1} \mathfrak{z} \big) \right\| &\leq \left\| S H_*^{-1} \nabla \mathsf{L} (\boldsymbol{\upsilon}^* + H_*^{-1} \mathfrak{z}) \right\| + \left\| S H_*^{-1} R (H_* + R)^{-1} \nabla \mathsf{L} (\boldsymbol{\upsilon}^* + H_*^{-1} \mathfrak{z}) \right\| \\ &\leq \left\| S H_*^{-1} \mathsf{D}_0 \right\| \left\| \mathsf{D}_0^{-1} \nabla \mathsf{L} (\boldsymbol{\upsilon}^* + H_*^{-1} \mathfrak{z}) \right\| \\ &+ \left\| S H_*^{-1} \mathsf{D}_* \right\| \left\| \mathsf{D}_*^{-1} R (H_* + R)^{-1} \mathsf{D}_* \right\| \left\| \mathsf{D}_*^{-1} \nabla \mathsf{L} (\boldsymbol{\upsilon}^* + H_*^{-1} \mathfrak{z}) \right\|. \end{split}$$

Let us recall that

$$\mathsf{D}^2_* = \operatorname{diag}\left((A^*)^\top A^* + \lambda I_d, 2I_d, I_d \otimes \left(\mu^2 I_d + \boldsymbol{\theta}^* (\boldsymbol{\theta}^*)^\top \right) \right)$$

and that μ_0 does not exceed μ by the definition. Then it is straightforward to observe that $D_*^2 \succeq D_0^2$ and $D_*^{-2} \preceq D_0^{-2}$. Taking these inequalities into account, we obtain that

$$\|S(\hat{\boldsymbol{v}} - \boldsymbol{v}^* - H_*^{-1}\boldsymbol{\mathfrak{z}})\| \leq \|SH_*^{-1}\mathsf{D}_*\| \|\mathsf{D}_0^{-1}\boldsymbol{\nabla}\mathsf{L}(\boldsymbol{v}^* + H_*^{-1}\boldsymbol{\mathfrak{z}})\| (1 + \|\mathsf{D}_*^{-1}R(H_* + R)^{-1}\mathsf{D}_*\|)$$
(108)

In the rest of the proof, we bound three terms in the right-hand side of (108) one by one.

Step 2: upper bound on $\|D_0^{-1}\nabla L(v^* + H_*^{-1}\mathfrak{z})\|$. We start with an upper bound on the norm of $D_0^{-1}\nabla L(v^* + H_*^{-1}\mathfrak{z})$. Note that

$$\begin{split} \mathsf{D}_0^{-1} \nabla \mathsf{L}(\boldsymbol{\upsilon}^* + H_*^{-1} \boldsymbol{\mathfrak{z}}) &= \mathsf{D}_0^{-1} \nabla \mathcal{L}(\boldsymbol{\upsilon}^* + H_*^{-1} \boldsymbol{\mathfrak{z}}) - \mathsf{D}_0^{-1} \boldsymbol{\mathfrak{z}} \\ &= \mathsf{D}_0^{-1} \left(\nabla \mathcal{L}(\boldsymbol{\upsilon}^* + H_*^{-1} \boldsymbol{\mathfrak{z}}) - \nabla \mathcal{L}(\boldsymbol{\upsilon}^*) \right) - \mathsf{D}_0^{-1} \boldsymbol{\mathfrak{z}}. \end{split}$$

Using Taylor's expansion with the integral remainder term, we obtain that

$$\begin{split} \left\| \mathbf{D}_{0}^{-1} \nabla \mathbf{L} (\boldsymbol{\upsilon}^{*} + H_{*}^{-1} \boldsymbol{\mathfrak{z}}) \right\| &= \left\| -\mathbf{D}_{0}^{-1} \boldsymbol{\mathfrak{z}} + \mathbf{D}_{0}^{-1} \int_{0}^{1} \nabla^{2} \mathcal{L} (\boldsymbol{\upsilon}^{*} + t \, H_{*}^{-1} \boldsymbol{\mathfrak{z}}) \, H_{*}^{-1} \boldsymbol{\mathfrak{z}} \, \mathrm{d}t \right\| \\ &= \left\| \int_{0}^{1} \mathbf{D}_{0}^{-1} \left(\nabla^{2} \mathcal{L} (\boldsymbol{\upsilon}^{*} + t \, H_{*}^{-1} \boldsymbol{\mathfrak{z}}) - H_{*} \right) \mathbf{D}_{0}^{-1} \mathbf{D}_{0} H_{*}^{-1} \boldsymbol{\mathfrak{z}} \, \mathrm{d}t \right\| \\ &\leq \left\| \mathbf{D}_{0} H_{*}^{-1} \boldsymbol{\mathfrak{z}} \right\| \int_{0}^{1} \left\| \mathbf{D}_{0}^{-1} \left(\nabla^{2} \mathcal{L} (\boldsymbol{\upsilon}^{*} + t \, H_{*}^{-1} \boldsymbol{\mathfrak{z}}) - H_{*} \right) \mathbf{D}_{0}^{-1} \right\| \, \mathrm{d}t \end{split}$$

Lemma A.3 implies that

$$\left\| \mathsf{D}_{0}^{-1} \left(\boldsymbol{\nabla}^{2} \mathcal{L} (\boldsymbol{\upsilon}^{*} + t \, H_{*}^{-1} \boldsymbol{\mathfrak{z}}) - H_{*} \right) \mathsf{D}_{0}^{-1} \right\| \leq \frac{4t}{\mu_{0} \sqrt{\lambda}} \left\| \mathsf{D}_{0} H_{*}^{-1} \boldsymbol{\mathfrak{z}} \right\| + \frac{2t^{2}}{\mu_{0}^{2} \lambda} \left\| \mathsf{D}_{0} H_{*}^{-1} \boldsymbol{\mathfrak{z}} \right\|^{2},$$

Hence, it holds that

$$\begin{aligned} \left\| \mathsf{D}_{0}^{-1} \nabla \mathsf{L}(\boldsymbol{v}^{*} + H_{*}^{-1} \boldsymbol{\mathfrak{z}}) \right\| &\leq \frac{4}{\mu_{0} \sqrt{\lambda}} \left\| \mathsf{D}_{0} H_{*}^{-1} \boldsymbol{\mathfrak{z}} \right\|^{2} \int_{0}^{1} t \, \mathrm{d}t + \frac{2}{\mu_{0}^{2} \lambda} \left\| \mathsf{D}_{0} H_{*}^{-1} \boldsymbol{\mathfrak{z}} \right\|^{3} \int_{0}^{1} t^{2} \, \mathrm{d}t \\ &= \frac{2}{\mu_{0} \sqrt{\lambda}} \left\| \mathsf{D}_{0} H_{*}^{-1} \boldsymbol{\mathfrak{z}} \right\|^{2} \left(1 + \frac{\left\| \mathsf{D}_{0} H_{*}^{-1} \boldsymbol{\mathfrak{z}} \right\|}{3\mu_{0} \sqrt{\lambda}} \right). \end{aligned}$$
(109)

.

Step 3: upper bound on $\|D_*^{-1}R(H_* + R)^{-1}D_*\|$. We proceed with an upper bound on the norm of $D_*^{-1}R(H_* + R)^{-1}D_*$. Representing the matrix of interest as the product of $D_*^{-1}RD_*^{-1}$ and $D_*(H_* + R)^{-1}D_*$, we obtain that

$$\left\| \mathsf{D}_{*}^{-1} R(H_{*} + R)^{-1} \mathsf{D}_{*} \right\| \leqslant \left\| \mathsf{D}_{*}^{-1} R \, \mathsf{D}_{*}^{-1} \right\| \left\| \mathsf{D}_{*}(H_{*} + R)^{-1} \mathsf{D}_{*} \right\|$$

Due to the Jensen inequality, the first term in the right-hand side does not exceed

$$\begin{split} \left\| \mathbf{D}_{*}^{-1} R \, \mathbf{D}_{*}^{-1} \right\| &= \left\| \int_{0}^{1} \mathbf{D}_{*}^{-1} \left[\nabla^{2} \mathbf{L} \left(\boldsymbol{\upsilon}^{*} + t(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^{*}) + (1 - t) H_{*}^{-1} \boldsymbol{\mathfrak{z}} \right) - H_{*} \right] \mathbf{D}_{*}^{-1} \, \mathrm{d}t \right\| \\ &\leq \int_{0}^{1} \left\| \mathbf{D}_{*}^{-1} \left[\nabla^{2} \mathbf{L} \left(\boldsymbol{\upsilon}^{*} + t(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^{*}) + (1 - t) H_{*}^{-1} \boldsymbol{\mathfrak{z}} \right) - H_{*} \right] \mathbf{D}_{*}^{-1} \right\| \, \mathrm{d}t. \end{split}$$

Next, we can bound the expression under the integral using Lemmata D.1, A.1, and A.3. Indeed, in view of Lemma A.3, for any $t \in [0, 1]$, we have

$$\begin{split} \left\| \mathsf{D}_*^{-1} \left(\nabla^2 \mathsf{L} \left(\boldsymbol{\upsilon}^* + t(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^*) + (1 - t) H_*^{-1} \mathfrak{z} \right) - H_* \right) \mathsf{D}_*^{-1} \right\| \\ \leqslant \frac{4}{\mu \sqrt{\lambda}} \left\| t \, \mathsf{D}_* (\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^*) + (1 - t) \mathsf{D}_* H_*^{-1} \mathfrak{z} \right\| + \frac{2}{\mu^2 \lambda} \left\| t \, \mathsf{D}_* (\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^*) + (1 - t) \, \mathsf{D}_* H_*^{-1} \mathfrak{z} \right\|^2. \end{split}$$

Lemma A.1 implies that

$$\left\|\mathsf{D}_{*}H_{*}^{-1}\mathsf{D}_{*}\right\| \leqslant \frac{4}{1-2\rho_{0}} \quad \text{and} \quad \left\|\mathsf{D}_{*}H_{*}^{-1}\mathfrak{z}\right\| \leqslant \frac{4}{1-2\rho_{0}}\left\|\mathsf{D}_{*}^{-1}\mathfrak{z}\right\|.$$

Combining this bound with the result of Lemma D.1, we get that

$$\begin{split} \left\| t \, \mathsf{D}_{*}(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^{*}) + (1 - t) \mathsf{D}_{*} H_{*}^{-1} \mathfrak{z} \right\| &\leq t \, \|\mathsf{D}_{*}(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^{*})\| + (1 - t) \, \big\| \mathsf{D}_{*} H_{*}^{-1} \mathfrak{z} \big\| \\ &\leq 9t \| \mathsf{D}_{*}^{-1} \mathfrak{z} \| + \frac{4(1 - t)}{1 - 2\rho_{0}} \| \mathsf{D}_{*}^{-1} \mathfrak{z} \| \\ &\leq 9 \| \mathsf{D}_{*}^{-1} \mathfrak{z} \|. \end{split}$$

The last inequality follows from the fact that $\rho \in [0, 1/16]$. Similarly, it holds that

$$\left\| t \, \mathsf{D}_{*}(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^{*}) + (1 - t) \mathsf{D}_{*} H_{*}^{-1} \mathfrak{z} \right\|^{2} \leqslant t \, \|\mathsf{D}_{*}(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^{*})\|^{2} + (1 - t) \, \|\mathsf{D}_{*} H_{*}^{-1} \mathfrak{z}\|^{2} \leqslant 81 \|\mathsf{D}_{*}^{-1} \mathfrak{z}\|^{2}.$$

Hence, we obtain that

$$\left\| \mathsf{D}_{*}^{-1} R \, \mathsf{D}_{*}^{-1} \right\| \leqslant \frac{36}{\mu\sqrt{\lambda}} \left\| \mathsf{D}_{*}^{-1} \mathfrak{z} \right\| \left(1 + \frac{9\|\mathsf{D}_{*}^{-1}\mathfrak{z}\|}{2\mu\sqrt{\lambda}} \right) \leqslant \frac{36}{\mu\sqrt{\lambda}} \left\| \mathsf{D}_{*}^{-1}\mathfrak{z} \right\| \left(1 + \frac{1}{36} \right) = \frac{37}{\mu\sqrt{\lambda}} \left\| \mathsf{D}_{*}^{-1}\mathfrak{z} \right\|.$$
(110)

The inequality (110) and Lemma A.1 immediately yield an upper bound on $\|D_*(H_* + R)^{-1}D_*\|$. Indeed, the smallest eigenvalue of $D_*^{-1}(H_* + R)D_*^{-1}$ is at least

$$\frac{1-2\rho_0}{4} - \left\| \mathsf{D}_*^{-1} R \, \mathsf{D}^{-1} \right\| \ge \frac{1}{4} - \frac{\rho_0}{2} - \frac{37}{\mu\sqrt{\lambda}} \left\| \mathsf{D}_*^{-1} \mathfrak{z} \right\| \ge \frac{1}{4} - \frac{1}{32} - \frac{3}{32} = \frac{1}{8}.$$

Thus,

$$\left\|\mathbf{D}_{*}(H_{*}+R)^{-1}\mathbf{D}_{*}\right\|\leqslant8$$

and, as a consequence, we have

$$\left\| \mathsf{D}_{*}^{-1} R(H_{*} + R)^{-1} \mathsf{D}_{*} \right\| \leq 8 \cdot \frac{37}{\mu \sqrt{\lambda}} \left\| \mathsf{D}_{*}^{-1} \mathfrak{z} \right\| = \frac{296}{\mu \sqrt{\lambda}} \left\| \mathsf{D}_{*}^{-1} \mathfrak{z} \right\|.$$
(111)

Step 4: upper bound on $||SH_*^{-1}D_*||$. It only remains to bound $||SH_*^{-1}D_*||$ to finish the proof. According to Lemma A.1, $4H_* \succeq (1-2\rho_0)D_*^2$. This yields that $(1-2\rho_0)H_*^{-1} \preceq 4D_*^{-2}$ and then

$$\|SH_*^{-1}\mathsf{D}_*\| = \|S\,\mathsf{D}_*^{-1}\,\mathsf{D}_*H_*^{-1}\mathsf{D}_*\| \leqslant \|S\,\mathsf{D}_*^{-1}\|\|\mathsf{D}_*H_*^{-1}\mathsf{D}_*\| \leqslant \frac{4}{1-2\rho_0}\|S\,\mathsf{D}_*^{-1}\|.$$

Let us recall that

$$S = \operatorname{diag}(\Sigma^{1/2}, I_{d+1} \otimes O_d) \quad \text{and} \quad \mathsf{D}^2_* = \operatorname{diag}\left((A^*)^\top A^* + \lambda I_d, 2I_d, I_d \otimes \left(\mu^2 I_d + \boldsymbol{\theta}^*(\boldsymbol{\theta}^*)^\top\right)\right).$$

Hence, it holds that

$$\begin{split} \|S \mathsf{D}_{*}^{-1}\| &= \left\| \Sigma^{1/2} \left((A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} \right\| = \left\| \left((A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} \Sigma \left((A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} \right\|^{1/2} \\ &\leq \left\| \left((A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} A^{*} \left((A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} \right\|^{1/2} \\ &+ \left\| \left((A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} (\Sigma - A^{*}) \left((A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} \right\|^{1/2} \\ &\leq \left\| \left((A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} \right\|^{1/2} + \sqrt{\frac{\|\Sigma - A^{*}\|}{\lambda}} \leqslant \lambda^{-1/4} + \sqrt{\frac{\|\Sigma - A^{*}\|}{\lambda}} \end{split}$$

and then

$$\|SH_*^{-1}\mathsf{D}_*\| \leq \frac{4}{1-2\rho_0} \left(\lambda^{-1/4} + \sqrt{\frac{\|\Sigma - A^*\|}{\lambda}}\right).$$
 (112)

Step 5: final bound. Summing up the inequalities (108), (109), (111), and (112), we obtain that

$$\begin{split} \left\| S \left(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^* - H^{-1} \mathfrak{z} \right) \right\| &\leqslant \frac{4}{1 - 2\rho_0} \left(\lambda^{-1/4} + \sqrt{\frac{\|\boldsymbol{\Sigma} - A^*\|}{\lambda}} \right) \cdot \frac{2}{\mu_0 \sqrt{\lambda}} \, \| \mathbf{D}_0 H_*^{-1} \mathfrak{z} \|^2 \\ & \cdot \left(1 + \frac{\|\mathbf{D}_0 H_*^{-1} \mathfrak{z}\|}{3\mu_0 \sqrt{\lambda}} \right) \left(1 + \frac{296}{\mu \sqrt{\lambda}} \, \| \mathbf{D}_*^{-1} \mathfrak{z} \| \right). \end{split}$$

Since, due to the conditions of Lemma 6.1, $395 \|D_*^{-1}\mathfrak{z}\| \le \mu \sqrt{\lambda}$ and $\rho \le 1/16$, the expression in the right-hand side simplifies to

$$\begin{split} \left\| S \big(\widehat{\boldsymbol{\upsilon}} - \boldsymbol{\upsilon}^* - H_*^{-1} \mathfrak{z} \big) \right\| &\leq \frac{8 \| \mathbf{D}_0 H_*^{-1} \mathfrak{z} \|^2}{(1 - 1/8)\mu_0 \sqrt{\lambda}} \left(\lambda^{-1/4} + \sqrt{\frac{\| \boldsymbol{\Sigma} - A^* \|}{\lambda}} \right) \left(1 + \frac{\| \mathbf{D}_0 H_*^{-1} \mathfrak{z} \|}{3\mu_0 \sqrt{\lambda}} \right) \left(1 + \frac{296}{395} \right) \\ &\leq \frac{16}{\mu_0 \sqrt{\lambda}} \| \mathbf{D}_0 H_*^{-1} \mathfrak{z} \|^2 \left(\lambda^{-1/4} + \sqrt{\frac{\| \boldsymbol{\Sigma} - A^* \|}{\lambda}} \right) \left(1 + \frac{\| \mathbf{D}_0 H_*^{-1} \mathfrak{z} \|}{3\mu_0 \sqrt{\lambda}} \right). \end{split}$$

In the last line, we used the fact that $64/7 \cdot (1 + 296/395) < 16$. The proof is finished.

D.3 Proof of Lemma 6.2

The main idea of the proof of Lemma 6.2 is to represent $\check{H}_{\theta\theta}^{-1}H_{\theta\chi}H_{\chi\chi}^{-1}\mathfrak{z}_{\chi}$ as a linear transform of $(\mathbf{Z} - \mathbb{E}\mathbf{Z})$ and $(\hat{\Sigma} - \Sigma)$ and then apply the large deviation inequalities from Theorems 2.5 and 2.6. Despite the simple idea, the proof is quite technical, so we split it into several steps for convenience.

Step 1. We start with an observation that the matrix $\breve{H}_{\theta\theta}$ is close to $(A^{\top}A/2 + \lambda I_d)$.

Lemma D.2. Assume that $\rho \leq 1/5$. Let us fix an arbitrary $v \in \Upsilon(\rho)$ and let $\check{H}_{\theta\theta} = H_{\theta\theta} - H_{\theta\chi}H_{\chi\chi}^{-1}H_{\chi\theta}$ stand for the Schur complement of $H_{\chi\chi}$. Then it holds that

$$\left\| \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{-1/2} \breve{H}_{\theta\theta} \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{-1/2} - I_d \right\| \leq 10\rho^2.$$

The proof of Lemma D.2 is postponed to Appendix D.8.1. Lemma D.2 yields that the smallest eigenvalue of $(A^{\top}A/2 + \lambda I_d)^{-1/2} \breve{H}_{\theta\theta} (A^{\top}A/2 + \lambda I_d)^{1/2}$ is at least $1 - 11\rho^2$. Then we deduce that the operator norm of

$$\left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{1/2} \breve{H}_{\theta\theta}^{-1} \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{1/2}$$

does not exceed $(1-10\rho^2)^{-1}$ and then

$$\left\| \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{1/2} \check{H}_{\theta\theta}^{-1} H_{\theta\chi} H_{\chi\chi}^{-1} \mathfrak{z}_{\chi} \right\| \leq \frac{1}{1 - 10\rho^2} \left\| \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{-1/2} H_{\theta\chi} H_{\chi\chi}^{-1} \mathfrak{z}_{\chi} \right\|$$

Thus, we can restrict our attention on a bit simpler term

$$\left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{-1/2} H_{\boldsymbol{\theta}\chi} H_{\boldsymbol{\chi}\chi}^{-1} \boldsymbol{\mathfrak{z}}_{\boldsymbol{\chi}}.$$

Step 2: explicit expression for $H_{\theta\chi}H_{\chi\chi}^{-1}\mathfrak{z}_{\chi}$. Let us elaborate on $H_{\theta\chi}H_{\chi\chi}^{-1}\mathfrak{z}_{\chi}$. The next lemma shows that $H_{\theta\chi}H_{\chi\chi}^{-1}\mathfrak{z}_{\chi}$ can be represented as a sum of linear transforms of $(\mathbf{Z} - \mathbb{E}\mathbf{Z})$ and $(\widehat{\Sigma} - \Sigma)$. Lemma D.3. Let $\rho \leq 1/2$ and let us fix any $\boldsymbol{v} \in \Upsilon(\rho)$. With the introduced notations, it holds that

$$\begin{split} H_{\theta\chi}H_{\chi\chi}^{-1}\mathfrak{z}_{\chi} &= -\frac{1}{2}A^{\top}\mathbf{U} + \frac{1}{2}A^{\top}(\widehat{\Sigma} - \Sigma)(\theta - \theta^{\circ}) + (\widehat{\Sigma} - \Sigma)(A\theta - \eta) \\ &- \left(r_{1} - \frac{r_{2}\mu^{2}\|\theta\|^{2}}{\mu^{2} + \|\theta\|^{2}}\right)A^{\top}(\mathbf{Z} - \mathbb{E}\mathbf{Z}) + \frac{r_{2}\mu^{2}}{\mu^{2} + \|\theta\|^{2}}\left((A\theta - \eta)^{\top}(\mathbf{Z} - \mathbb{E}\mathbf{Z})\right)\theta \\ &- \left(\frac{\|\theta\|^{2}}{2\mu^{2} + \|\theta\|^{2}} - \frac{\|\theta\|^{2}}{2(\mu^{2} + \|\theta\|^{2})} + \frac{r_{2}'\mu^{4}}{\mu^{2} + \|\theta\|^{2}}\right)A^{\top}(\widehat{\Sigma} - \Sigma)\theta \\ &- \frac{1}{2\mu^{2} + \|\theta\|^{2}}\left((A\theta - \eta)^{\top}(\widehat{\Sigma} - \Sigma)\theta\right)\theta, \end{split}$$

where

$$\mathbf{U} = \mathbf{Z} - \mathbb{E}\mathbf{Z} - (\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^{\circ} = \frac{1}{n} \mathbb{X}\boldsymbol{\varepsilon}$$

and the constants $r_1 \in [0, \rho^2]$, $r_2 = 0.5\mu^{-2} + r'_2$, $r'_2 \in [0, \rho^2/\mu^2]$ are defined in Proposition A.2.

The proof of Lemma D.3 is moved to Appendix D.8.2. The inequalities $0 \le r_1 \le \rho^2$ and $0 \le r'_2 \le \rho^2/\mu^2$ yield that

$$\left|r_{1} - \frac{r_{2}\mu^{2} \|\boldsymbol{\theta}\|^{2}}{\mu^{2} + \|\boldsymbol{\theta}\|^{2}}\right| \leqslant \rho^{2} \quad \text{and} \quad \left|\frac{\|\boldsymbol{\theta}\|^{2}}{2\mu^{2} + \|\boldsymbol{\theta}\|^{2}} - \frac{\|\boldsymbol{\theta}\|^{2}}{2(\mu^{2} + \|\boldsymbol{\theta}\|^{2})} + \frac{r_{2}'\mu^{4}}{\mu^{2} + \|\boldsymbol{\theta}\|^{2}}\right| \leqslant 2\rho^{2}.$$

Then, according to Lemma D.3 and the triangle inequality, we have

$$\begin{aligned} \left\| \left(\frac{1}{2} A^{\mathsf{T}} A + \lambda I_d \right)^{-1/2} H_{\boldsymbol{\theta}\chi} H_{\boldsymbol{\chi}\chi}^{-1} \mathfrak{z}_{\boldsymbol{\chi}} \right\| \\ &\leq \frac{1}{2} \left\| \left(\frac{1}{2} A^{\mathsf{T}} A + \lambda I_d \right)^{-1/2} A^{\mathsf{T}} \mathbf{U} \right\| + \frac{1}{2} \left\| \left(\frac{1}{2} A^{\mathsf{T}} A + \lambda I_d \right)^{-1/2} A^{\mathsf{T}} (\widehat{\Sigma} - \Sigma) (\boldsymbol{\theta} - \boldsymbol{\theta}^{\circ}) \right\| \\ &+ \rho^2 \left\| \left(\frac{1}{2} A^{\mathsf{T}} A + \lambda I_d \right)^{-1/2} A^{\mathsf{T}} (\mathbf{Z} - \mathbb{E} \mathbf{Z}) \right\| + 2\rho^2 \left\| \left(\frac{1}{2} A^{\mathsf{T}} A + \lambda I_d \right)^{-1/2} A^{\mathsf{T}} (\widehat{\Sigma} - \Sigma) \boldsymbol{\theta} \right\|$$
(113)
$$&+ \frac{\lambda^{-1/2} \|\boldsymbol{\theta}\|}{\mu^2 + \|\boldsymbol{\theta}\|^2} \left| (A\boldsymbol{\theta} - \boldsymbol{\eta})^{\mathsf{T}} (\mathbf{Z} - \mathbb{E} \mathbf{Z}) \right| + \frac{\lambda^{-1/2} \|\boldsymbol{\theta}\|}{2\mu^2 + \|\boldsymbol{\theta}\|^2} \left| (A\boldsymbol{\theta} - \boldsymbol{\eta})^{\mathsf{T}} (\widehat{\Sigma} - \Sigma) \boldsymbol{\theta} \right| \\ &+ \left\| \left(\frac{1}{2} A^{\mathsf{T}} A + \lambda I_d \right)^{-1/2} (\widehat{\Sigma} - \Sigma) (A\boldsymbol{\theta} - \boldsymbol{\eta}) \right\|. \end{aligned}$$

Here we used the fact that

$$\left\| (A^{\top}A + \lambda I_d)^{-1/2} \boldsymbol{\theta} \right\| \leq \frac{\|\boldsymbol{\theta}\|}{\sqrt{\lambda}}.$$

Since

$$\left\| \left(\frac{1}{2} A^{\top} A + \lambda I_d \right)^{-1/2} A^{\top} \right\| \leqslant \sqrt{2} \quad \text{and} \quad \left\| \left(\frac{1}{2} A^{\top} A + \lambda I_d \right)^{-1/2} \right\| \leqslant \frac{1}{\sqrt{\lambda}},$$

the inequality (113) can be simplified even further:

$$\left\| \left(\frac{1}{2} A^{\top} A + \lambda I_d \right)^{-1/2} H_{\boldsymbol{\theta}\chi} H_{\boldsymbol{\chi}\chi}^{-1} \mathfrak{z}_{\chi} \right\| \leq \frac{\|\mathbf{U}\|}{\sqrt{2}} + \frac{\|(\widehat{\Sigma} - \Sigma)(\boldsymbol{\theta} - \boldsymbol{\theta}^{\circ})\|}{\sqrt{2}} + \frac{\|(\widehat{\Sigma} - \Sigma)(A\boldsymbol{\theta} - \boldsymbol{\eta})\|}{\sqrt{\lambda}} + \sqrt{2}\rho^2 \|\mathbf{Z} - \mathbb{E}\mathbf{Z}\| + 2\sqrt{2}\rho^2 \left\| (\widehat{\Sigma} - \Sigma)\boldsymbol{\theta} \right\| + \frac{\lambda^{-1/2} \|\boldsymbol{\theta}\|}{\mu^2 + \|\boldsymbol{\theta}\|^2} \left| (A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} (\mathbf{Z} - \mathbb{E}\mathbf{Z}) \right| + \frac{\lambda^{-1/2} \|\boldsymbol{\theta}\|}{2\mu^2 + \|\boldsymbol{\theta}\|^2} \left| (A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} (\widehat{\Sigma} - \Sigma)\boldsymbol{\theta} \right|.$$

$$(114)$$

In other words, Lemma D.3 allows us to reduce the study of a complicated expression for $\check{H}_{\theta\theta}^{-1}H_{\theta\chi}H_{\chi\chi}^{-1}\mathfrak{z}_{\chi}$ to analysis of much more tractable terms. In the rest of the proof, we bound the summands in the right-hand side of (113) one by one applying Theorems 2.5 and 2.6.

Step 3: large deviation bounds. Let us recall that on the event \mathcal{E}_1 we have

$$\|(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^{\circ}\| \leqslant 4C_X \|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \quad \text{and} \quad \|\mathbf{U}\| \leqslant 8\sigma \|\Sigma\|^{1/2} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}.$$

Moreover, according to Theorem 2.5, each of the inequalities (36)–(39) holds with probability at least $1 - \delta/2$. Note that the conditions of Theorem 2.5 are fulfilled, because $\mathbf{r}(\Sigma) + \log(4/\delta) \leq n \leq 4n$ by our assumptions. Similarly, due to Theorem 2.6, with probability at least $1 - \delta/2$, it holds that

$$\left| (A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} \mathbf{U} \right| \leq 8\sigma \| \Sigma^{1/2} (A\boldsymbol{\theta} - \boldsymbol{\eta}) \| \sqrt{\frac{1 + \log(4/\delta)}{n}}$$

The union bound yields that there exists an event \mathcal{E}_2 of probability at least $1 - 5\delta/2$ where the inequalities (36)–(40) hold simultaneously. On the intersection of the events \mathcal{E}_1 and \mathcal{E}_2 , the right-hand side of (114) does not exceed

$$\begin{aligned} \left\| \left(\frac{1}{2} A^{\top} A + \lambda I_d \right)^{-1/2} H_{\boldsymbol{\theta}\chi} H_{\chi\chi}^{-1} \mathfrak{z}_{\chi} \right\| \\ &\leqslant 2\sqrt{2} \|\Sigma\|^{1/2} \left(2\sigma + C_X \|\Sigma^{1/2} (\boldsymbol{\theta} - \boldsymbol{\theta}^\circ)\| + \frac{\sqrt{2}C_X \|\Sigma^{1/2} (A\boldsymbol{\theta} - \boldsymbol{\eta})\|}{\sqrt{\lambda}} \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ 4\sqrt{2}\rho^2 \|\Sigma\|^{1/2} \left(2\sigma + 3C_X \|\Sigma^{1/2} \boldsymbol{\theta}^\circ\| \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ \frac{8\|\boldsymbol{\theta}\|}{\mu^2 + \|\boldsymbol{\theta}\|^2} \cdot \frac{\|\Sigma^{1/2} (A\boldsymbol{\theta} - \boldsymbol{\eta})\|}{\sqrt{\lambda}} \left(C_X \|\Sigma^{1/2} \boldsymbol{\theta}\| + \sigma \right) \sqrt{\frac{1 + \log(4/\delta)}{n}}. \end{aligned}$$

Finally, since $\boldsymbol{v} \in \Upsilon(\rho)$, we have

$$\|oldsymbol{ heta}\|^2\leqslant
ho^2(\mu^2+\|oldsymbol{ heta}\|^2)\quad ext{and}\quad rac{\|oldsymbol{ heta}\|}{\mu^2+\|oldsymbol{ heta}\|^2}\leqslantrac{\|oldsymbol{ heta}\|}{\mu\sqrt{\mu^2+\|oldsymbol{ heta}\|^2}}\leqslantrac{
ho}{\mu}.$$

This yields the desired bound:

$$\begin{split} & \left\| \left(\frac{1}{2} A^{\top} A + \lambda I_{d} \right)^{1/2} \breve{H}_{\theta\theta}^{-1} H_{\theta\chi} H_{\chi\chi}^{-1} \mathfrak{z}_{\chi} \right\| \\ & \leq \frac{1}{1 - 10\rho^{2}} \left\| \left(\frac{1}{2} A^{\top} A + \lambda I_{d} \right)^{-1/2} H_{\theta\chi} H_{\chi\chi}^{-1} \mathfrak{z}_{\chi} \right\| \\ & \leq \frac{2\sqrt{2} \|\Sigma\|^{1/2}}{1 - 10\rho^{2}} \left(2\sigma + C_{X} \|\Sigma^{1/2}(\theta - \theta^{\circ})\| + \frac{\sqrt{2}C_{X} \|\Sigma^{1/2}(A\theta - \eta)\|}{\sqrt{\lambda}} \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ & + \frac{4\sqrt{2}\rho^{2} \|\Sigma\|^{1/2}}{1 - 10\rho^{2}} \left(2\sigma + 3C_{X} \|\Sigma^{1/2}\theta^{\circ}\| \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ & + \frac{8\rho}{(1 - 10\rho^{2})\mu} \cdot \frac{\|\Sigma^{1/2}(A\theta - \eta)\|}{\sqrt{\lambda}} \left(C_{X} \|\Sigma^{1/2}\theta\| + \sigma \right) \sqrt{\frac{1 + \log(4/\delta)}{n}}. \end{split}$$

D.4 Proof of Lemma 6.3

As before, we will use the blockwise representation of the Hessian of $\mathcal{L}(\boldsymbol{v})$. We would like to recall that $\check{H}_{\chi\chi} = H_{\chi\chi} - H_{\chi\theta}H_{\theta\theta}^{-1}H_{\chi\theta}$, where

$$H_{\theta\theta} = A^{\top}A + \lambda I_d, \quad H_{\chi\chi} = \begin{pmatrix} 2I_d & H_{\eta A} \\ H_{A\eta} & H_{AA} \end{pmatrix} \text{ and } H_{\theta\chi} = \begin{pmatrix} -A^{\top} & H_{\theta A} \end{pmatrix}.$$

The key observation in the proof of Lemma 6.3 is that the off-diagonal blocks of $\breve{H}_{\chi\chi}$ are small. To be more precise, let us introduce a block-diagonal matrix

$$\mathsf{D}_{\chi}^{2} = \operatorname{diag}\left(2I_{d} - A(A^{\top}A + \lambda I_{d})^{-1}A^{\top}, H_{AA}\right),$$

and consider the difference

$$Q = \breve{H}_{\chi\chi} - \mathsf{D}_{\chi}^2 = \begin{pmatrix} O_d & -AH_{\theta\theta}^{-1}H_{\theta A} + H_{\eta A} \\ -H_{A\theta}H_{\theta\theta}^{-1}A^\top + H_{A\eta} & -H_{A\theta}H_{\theta\theta}^{-1}H_{\theta A} \end{pmatrix}$$

We are going to show that $\|\mathbf{D}_{\chi}^{-1}Q\mathbf{D}_{\chi}^{-1}\| \leq (1+\sqrt{2})\rho + 2\rho^2$. As usual, we split the rest of the proof into several steps for convenience.

Step 1: upper bound on $\|D_{\chi}^{-1}Q D_{\chi}^{-1}\|$. Since $A(A^{\top}A + \lambda I_d)^{-1}A^{\top} \leq I_d$ and $D_{\chi}^2 \geq \text{diag}(I_d, H_{AA})$, it is enough to check that

$$\left\| \operatorname{diag}(I_d, H_{AA}^{-1/2}) Q \operatorname{diag}(I_d, H_{AA}^{-1/2}) \right\| \leq (1 + \sqrt{2})\rho + 2\rho^2.$$

For this purpose, we fix arbitrary $\mathbf{v} \in \mathbb{R}^d$ and $\mathbf{w} \in \mathbb{R}^{d^2}$ and study

$$\begin{pmatrix} \mathbf{v}^\top & \mathbf{w}^\top \end{pmatrix} Q \begin{pmatrix} \mathbf{v} \\ \mathbf{w} \end{pmatrix}.$$

Due to the Cauchy-Schwarz inequality, it holds that

$$\begin{pmatrix} \mathbf{v}^{\top} & \mathbf{w}^{\top} \end{pmatrix} Q \begin{pmatrix} \mathbf{v} \\ \mathbf{w} \end{pmatrix} = 2 \mathbf{v}^{\top} A H_{\theta\theta}^{-1} H_{\theta A} \mathbf{w} + 2 \mathbf{v}^{\top} H_{\eta A} \mathbf{w} + \mathbf{w}^{\top} H_{A\theta} H_{\theta\theta}^{-1} H_{\theta A} \mathbf{w}$$

$$\leq 2 \| \mathbf{v} \| \| A H_{\theta\theta}^{-1/2} \| \| H_{\theta\theta}^{-1/2} H_{\theta A} H_{AA}^{-1/2} \| \| H_{AA}^{1/2} \mathbf{w} \|$$

$$+ 2 \| \mathbf{v} \| \| H_{\eta A} H_{AA}^{-1/2} \| \| H_{AA}^{1/2} \mathbf{w} \| + \| H_{\theta\theta}^{-1/2} H_{\theta A} H_{AA}^{-1/2} \|^2 \| H_{AA}^{1/2} \mathbf{w} \|^2.$$

According to Lemma A.4, we have

$$\|H_{\eta A}H_{AA}^{-1/2}\| \leq \rho \quad \text{and} \quad \|H_{\theta\theta}^{-1/2}H_{\theta A}H_{AA}^{-1/2}\| \leq \rho\sqrt{2}.$$

This, together with the inequality

$$\|AH_{\boldsymbol{\theta}\boldsymbol{\theta}}^{-1/2}\| = \left\|A(A^{\top}A + \lambda I_d)^{-1/2}\right\| \leq 1,$$

yields that

$$\begin{aligned} \left(\mathbf{v}^{\top} \quad \mathbf{w}^{\top} \right) Q \begin{pmatrix} \mathbf{v} \\ \mathbf{w} \end{pmatrix} &\leq 2\rho (1 + \sqrt{2}) \|\mathbf{v}\| \, \|H_{AA}^{1/2} \mathbf{w}\| + 2\rho^2 \|H_{AA}^{1/2} \mathbf{w}\|^2 \\ &\leq \rho (1 + \sqrt{2}) \left(\|\mathbf{v}\|^2 + \|H_{AA}^{1/2} \mathbf{w}\|^2 \right) + 2\rho^2 \|H_{AA}^{1/2} \mathbf{w}\|^2 \\ &\leq \left((1 + \sqrt{2})\rho + 2\rho^2 \right) \left(\|\mathbf{v}\|^2 + \|H_{AA}^{1/2} \mathbf{w}\|^2 \right). \end{aligned}$$

Hence, we proved that

$$\|\mathsf{D}_{\chi}^{-1}Q\,\mathsf{D}_{\chi}^{-1}\| \leqslant \left\|\operatorname{diag}(I_d, H_{AA}^{-1/2})\,Q\,\operatorname{diag}(I_d, H_{AA}^{-1/2})\right\| \leqslant (1+\sqrt{2})\rho + 2\rho^2.$$
(115)

Step 2: upper bound on $\|\widetilde{D}\breve{H}_{\chi\chi}^{-1}\mathfrak{z}_{\chi}\|$. With the inequality (115) the upper bound on the norm of $\widetilde{D}\breve{H}_{\chi\chi}^{-1}\mathfrak{z}_{\chi}$ is straightforward. Indeed, it holds that

$$\begin{split} \breve{H}_{\chi\chi}^{-1}\mathfrak{z}_{\chi} &= \left(\mathsf{D}_{\chi}^{2} + Q\right)^{-1}\mathfrak{z}_{\chi} = \mathsf{D}_{\chi}^{-1}\left(I_{d} + \mathsf{D}_{\chi}^{-1}Q\,\mathsf{D}_{\chi}^{-1}\right)^{-1}\mathsf{D}_{\chi}^{-1}\mathfrak{z}_{\chi} \\ &= \mathsf{D}_{\chi}^{-1}\left(I_{d} - \left(I_{d} + \mathsf{D}_{\chi}^{-1}Q\,\mathsf{D}_{\chi}^{-1}\right)^{-1}\right)\mathsf{D}_{\chi}^{-1}\mathfrak{z}_{\chi} \\ &= \mathsf{D}_{\chi}^{-2}\mathfrak{z}_{\chi} - \mathsf{D}_{\chi}^{-1}\left(I_{d} + \mathsf{D}_{\chi}^{-1}Q\,\mathsf{D}_{\chi}^{-1}\right)^{-1}\mathsf{D}_{\chi}^{-1}\mathcal{Q}\,\mathsf{D}_{\chi}^{-2}\mathfrak{z}_{\chi}. \end{split}$$

Then (115) and the triangle inequality imply that

$$\begin{split} \left\| \widetilde{\mathsf{D}} \breve{H}_{\chi\chi}^{-1} \mathfrak{z}_{\chi} \right\| &\leq \left\| \widetilde{\mathsf{D}} \, \mathsf{D}_{\chi}^{-2} \mathfrak{z}_{\chi} \right\| + \left\| \widetilde{\mathsf{D}} \, \mathsf{D}_{\chi}^{-1} \left(I_{d} + \mathsf{D}_{\chi}^{-1} Q \, \mathsf{D}_{\chi}^{-1} \right)^{-1} \mathsf{D}_{\chi}^{-1} Q \, \mathsf{D}_{\chi}^{-2} \mathfrak{z}_{\chi} \right\| \\ &\leq \left\| \widetilde{\mathsf{D}} \, \mathsf{D}_{\chi}^{-2} \mathfrak{z}_{\chi} \right\| + \left\| \widetilde{\mathsf{D}} \, \mathsf{D}_{\chi}^{-1} \right\| \cdot \frac{\left\| \mathsf{D}_{\chi}^{-1} Q \, \mathsf{D}_{\chi}^{-1} \right\|}{1 - \left\| \mathsf{D}_{\chi}^{-1} Q \, \mathsf{D}_{\chi}^{-1} \right\|} \cdot \left\| \mathsf{D}_{\chi}^{-1} \mathfrak{z}_{\chi} \right\| \\ &\leq \left\| \widetilde{\mathsf{D}} \, \mathsf{D}_{\chi}^{-2} \mathfrak{z}_{\chi} \right\| + \frac{\left((1 + \sqrt{2}) \rho + 2\rho^{2} \right) \sqrt{2}}{1 - (1 + \sqrt{2}) \rho - 2\rho^{2}} \cdot \left\| \mathsf{D}_{\chi}^{-1} \mathfrak{z}_{\chi} \right\|. \end{split}$$

Here we used the fact that

$$\mathsf{D}_{\chi}^{2} \succeq \operatorname{diag}(I_{d}, H_{AA}) = \operatorname{diag}(I_{d}, I_{d} \otimes (\mu^{2}I_{d} + \boldsymbol{\theta}\boldsymbol{\theta}^{\top})) \succeq \frac{1}{2} \operatorname{diag}(2I_{d}, I_{d} \otimes (\widetilde{\mu}^{2}I_{d} + \boldsymbol{\theta}\boldsymbol{\theta}^{\top})) \succeq \frac{1}{2} \widetilde{\mathsf{D}}^{2}$$

which yields $\|\widetilde{\mathtt{D}}\,\mathtt{D}_{\chi}^{-1}\|\leqslant\sqrt{2}.$ Since $\rho\leqslant1/16$, we obtain that

$$\left\|\widetilde{\mathsf{D}}\breve{H}_{\chi\chi}^{-1}\mathfrak{z}_{\chi}\right\| \leqslant \left\|\widetilde{\mathsf{D}}\,\mathsf{D}_{\chi}^{-2}\mathfrak{z}_{\chi}\right\| + \frac{((1+\sqrt{2})\rho + \rho/8)\sqrt{2}}{1-(1+\sqrt{2})/16 - 1/128} \,\left\|\mathsf{D}_{\chi}^{-1}\mathfrak{z}_{\chi}\right\| \leqslant \left\|\mathsf{D}_{\chi}^{-2}\mathfrak{z}_{\chi}\right\| + \frac{9\rho}{2} \,\left\|\mathsf{D}_{\chi}^{-1}\mathfrak{z}_{\chi}\right\|.$$

It only remains to note that

$$\begin{split} \left\| \widetilde{\mathsf{D}} \, \mathsf{D}_{\chi}^{-2} \mathfrak{z}_{\chi} \right\|^{2} &\leq \left\| \operatorname{diag} \left(I_{d}, \frac{\sqrt{\widetilde{\mu}^{2} + \|\boldsymbol{\theta}\|^{2}}}{\mu^{2}} I_{d^{2}} \right) \mathfrak{z}_{\chi} \right\|^{2} \\ &= \left\| \operatorname{diag} \left(I_{d}, \frac{\sqrt{\widetilde{\mu}^{2} + \|\boldsymbol{\theta}\|^{2}}}{\mu^{2}} I_{d^{2}} \right) \left(\frac{\mathbf{Z} - \mathbb{E}\mathbf{Z}}{\mu^{2} \operatorname{vec}(\widehat{\Sigma} - \Sigma)} \right) \right\|^{2} \\ &= \left\| \mathbf{Z} - \mathbb{E}\mathbf{Z} \right\|^{2} + \left(\widetilde{\mu}^{2} + \|\boldsymbol{\theta}\|^{2} \right) \left\| \widehat{\Sigma} - \Sigma \right\|_{\mathrm{F}}^{2} \end{split}$$

and, similarly,

$$\left\| \mathsf{D}_{\chi}^{-1} \mathfrak{z}_{\chi} \right\|^{2} \leqslant \left\| \operatorname{diag} \left(I_{d}, \frac{1}{\mu} I_{d^{2}} \right) \left(\frac{\mathbf{Z} - \mathbb{E} \mathbf{Z}}{\mu^{2} \operatorname{vec}(\widehat{\Sigma} - \Sigma)} \right) \right\|^{2} = \left\| \mathbf{Z} - \mathbb{E} \mathbf{Z} \right\|^{2} + \mu^{2} \left\| \widehat{\Sigma} - \Sigma \right\|_{\mathrm{F}}^{2}.$$

Then

$$\left\|\widetilde{\mathsf{D}}\breve{H}_{\chi\chi}^{-1}\mathfrak{z}_{\chi}\right\| \leqslant \left(1 + \frac{9\rho}{2}\right) \left(\|\mathbf{U}\| + \|(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^{\circ}\|\right) + \left(\sqrt{\widetilde{\mu}^{2} + \|\boldsymbol{\theta}\|^{2}} + \frac{9\rho\mu}{2}\right) \left\|\widehat{\Sigma} - \Sigma\right\|_{\mathrm{F}}$$

and, taking into account that

$$\|\widehat{\Sigma} - \Sigma\|_{\mathrm{F}} \leqslant 4C_X \|\Sigma\| \sqrt{\frac{\mathbf{r}(\Sigma)^2 + \log(4/\delta)}{n}}, \quad \|(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^\circ\| \leqslant 4C_X \|\Sigma\| \|\boldsymbol{\theta}^\circ\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}},$$

and

$$\|\mathbf{U}\| \leq 8\sigma \|\Sigma\|^{1/2} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}$$

on the event \mathcal{E}_1 (see (31) and (32)), we deduce the desired bound:

$$\begin{split} \left\| \widetilde{\mathsf{D}} \breve{H}_{\chi\chi}^{-1} \mathfrak{z}_{\chi} \right\| &\leq 8 \left(1 + \frac{9\rho}{2} \right) \sigma \|\Sigma\|^{1/2} \sqrt{\frac{\mathsf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ 4 \left(1 + \frac{9\rho}{2} \right) C_X \|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| \sqrt{\frac{\mathsf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ 4 \left(\sqrt{\widetilde{\mu}^2 + \|\boldsymbol{\theta}\|^2} + \frac{9\rho\mu}{2} \right) C_X \|\Sigma\| \sqrt{\frac{\mathsf{r}(\Sigma)^2 + \log(4/\delta)}{n}}. \end{split}$$

D.5 Proof of Lemma 6.4

We would like to remind a reader that (see (34))

$$\mathbf{s}_{\boldsymbol{\theta}}^* = -\breve{H}_{\boldsymbol{\theta}\boldsymbol{\theta}}^{-1}(\boldsymbol{\upsilon}^*)H_{\boldsymbol{\theta}\boldsymbol{\chi}}(\boldsymbol{\upsilon}^*)H_{\boldsymbol{\chi}\boldsymbol{\chi}}^{-1}(\boldsymbol{\upsilon}^*)\boldsymbol{\mathfrak{z}}_{\boldsymbol{\chi}},$$

where, as before, $\breve{H}_{\theta\theta}(\upsilon^*) = H_*/H_{\chi\chi}(\upsilon^*)$ stands for the Schur complement of $H_{\chi\chi}(\upsilon^*)$. The proof of Lemma 6.4 consists of two steps. First, using Lemma D.2, we show that \mathbf{s}^*_{θ} is close to

$$\left(\frac{1}{2}(A^*)^{\top}A^* + \lambda I_d\right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \boldsymbol{\mathfrak{z}}_{\chi}$$

After that, we quantify the difference between

$$\left(\frac{1}{2}(A^*)^{\top}A^* + \lambda I_d\right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \mathfrak{z}_{\chi} \quad \text{and} \quad \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \mathfrak{z}_{\chi}$$

Step 1. Let us introduce

$$R = \left(\frac{1}{2}(A^*)^\top A^* + \lambda I_d\right)^{-1/2} \breve{H}_{\boldsymbol{\theta}\boldsymbol{\theta}}(\boldsymbol{\upsilon}^*) \left(\frac{1}{2}(A^*)^\top A^* + \lambda I_d\right)^{-1/2} - I_d$$

and note that

$$\begin{aligned} \left\| \Sigma^{1/2} \mathbf{s}_{\boldsymbol{\theta}}^{*} + \Sigma^{1/2} \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\| \\ &= \left\| \Sigma^{1/2} \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} R(I_{d} + R)^{-1} \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\| \\ &\leq \left\| R(I_{d} + R)^{-1} \right\| \left\| \Sigma^{1/2} \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} \right\| \left\| \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\|. \end{aligned}$$

According to Lemma B.2 (see (ii) and (iv)), it holds that

$$||A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^*|| \leq \sqrt{\lambda} ||\boldsymbol{\theta}^\circ||$$
 and $||\boldsymbol{\theta}^*|| \leq ||\boldsymbol{\theta}^\circ||.$

This means that θ^* belongs to $\Upsilon(\rho^*)$ with $\rho^* = \|\theta^\circ\|/\mu \leq 1/112$. Then, due to Lemma D.2, the operator norm of R does not exceed $10(\rho^*)^2$. This yields that

$$\left\| R(I_d + R)^{-1} \right\| \leq \|R\| \left\| (I_d + R)^{-1} \right\| \leq \frac{\|R\|}{1 - \|R\|} \leq \frac{10(\rho^*)^2}{1 - 10(\rho^*)^2} \leq 1.001(\rho^*)^2 = \frac{1.001 \|\boldsymbol{\theta}^\circ\|^2}{\mu^2},$$

and, as a consequence, we obtain that

$$\left\| \Sigma^{1/2} \mathbf{s}_{\boldsymbol{\theta}}^{*} + \Sigma^{1/2} \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\|$$

$$\leq \frac{1.001 \|\boldsymbol{\theta}^{\circ}\|^{2}}{\mu^{2}} \left\| \Sigma^{1/2} \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} \left\| \left\| \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\|.$$
(116)

We are going to show that, under the conditions of the lemma, the operator norm of

$$\Sigma^{1/2} \left((A^*)^\top A^*/2 + \lambda I_d \right)^{-1/2}$$

is not greater than $\lambda^{-1/4}$. The proof of this fact is based on the following lemma.

Lemma D.4. Let us fix arbitrary $A \in \mathbb{R}^{d \times d}$ and $\lambda > 0$. Then it holds that

$$\frac{1}{2} \left\| \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \left(A^\top A - \Sigma^2 \right) \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \right\| \leq \|A - \Sigma\| \sqrt{\frac{2}{\lambda}} + \frac{\|A - \Sigma\|^2}{2\lambda}.$$

We postpone the proof of Lemma D.4 to Appendix D.8.3 and proceed with the proof of Lemma 6.4. Taking into account (7) and applying Lemma B.3, we note that

$$\frac{\|A^* - \Sigma\|}{\sqrt{\lambda}} \leqslant \left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{5\|\mathbf{b}_\lambda\|}{64 \cdot 18\mu} \leqslant \left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{\|\mathbf{b}_\lambda\|}{230\mu}$$

Due to the definition of \mathbf{b}_{λ} and the condition $112 \|\boldsymbol{\theta}^{\circ}\| \leq \mu$, the right-hand side does not exceed

$$\left(\frac{14\|\boldsymbol{\theta}^{\circ}\|}{\mu}\right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{230\mu} \leqslant \frac{1}{64} + \frac{1}{112 \cdot 230} < \frac{1}{63}.$$

Then, due to Lemma D.4, it holds that

$$\left\| \left(\frac{1}{2}\Sigma^{2} + \lambda I_{d}\right)^{-1/2} \left(\frac{1}{2} (A^{*})^{\top} A^{*} - \frac{1}{2}\Sigma^{2}\right) \left(\frac{1}{2}\Sigma^{2} + \lambda I_{d}\right)^{-1/2} \right\|$$

$$\leq \left[\left(\frac{14\|\boldsymbol{\theta}^{\circ}\|}{\mu}\right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{230\mu} \right] \left(\sqrt{2} + \frac{1}{2\cdot 63}\right) \leq \left(\frac{17\|\boldsymbol{\theta}^{\circ}\|}{\mu}\right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{160\mu} \leq \frac{1}{43}.$$
(117)

In the last line, we used the inequalities

$$\sqrt{2} + 1/126 < (17/14)^2$$
, $(\sqrt{2} + 1/126)/230 < 1/160$, and $\frac{\|\mathbf{b}_{\lambda}\|}{\mu} \leq \frac{\|\boldsymbol{\theta}^{\circ}\|}{\mu} \leq \frac{1}{112}$.

Let us denote the matrix in the left-hand side by B:

$$B = \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \left(\frac{1}{2}(A^*)^\top A^* - \frac{1}{2}\Sigma^2\right) \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2}.$$
 (118)

Then it is straightforward to observe that

$$\left\| \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{1/2} \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1/2} \right\|^2$$

$$= \left\| \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d + \frac{1}{2} (A^*)^\top A^* - \frac{1}{2} \Sigma^2 \right) \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{1/2} \right\|$$

$$= \left\| (I_d + B)^{-1} \right\| \leqslant \frac{1}{1 - \|B\|} \leqslant \frac{43}{42}.$$

$$(119)$$

This inequality and (116) immediately imply that

$$\left\| \Sigma^{1/2} \mathbf{s}_{\boldsymbol{\theta}}^{*} + \Sigma^{1/2} \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\|$$

$$\leq \frac{1.001 \|\boldsymbol{\theta}^{\circ}\|^{2}}{\mu^{2}} \sqrt{\frac{43}{42}} \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^{2} + \lambda I_{d} \right)^{-1/2} \left\| \left\| \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\|.$$

Due to the Cauchy-Schwarz inequality,

$$\left\|\Sigma^{1/2} \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2}\right\|^2 = \left\|\left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \Sigma \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2}\right\| \leqslant \frac{1}{\sqrt{2\lambda}}.$$
 (120)

This yields that

$$\left\| \Sigma^{1/2} \mathbf{s}_{\boldsymbol{\theta}}^{*} + \Sigma^{1/2} \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\|$$

$$\leq \frac{1.001 \|\boldsymbol{\theta}^{\circ}\|^{2}}{\mu^{2}} \sqrt{\frac{43}{42}} \cdot (2\lambda)^{-1/4} \left\| \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\|.$$
(121)

Hence, we proved that $\Sigma^{1/2} \mathbf{s}_{\boldsymbol{\theta}}^*$ is close to $\Sigma^{1/2} \left((A^*)^\top A^*/2 + \lambda I_d \right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{v}^*) H_{\chi\chi}^{-1}(\boldsymbol{v}^*) \mathfrak{z}_{\chi}$. Our next goal is to show that

$$\Sigma^{1/2} \left((A^*)^\top A^*/2 + \lambda I_d \right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \mathfrak{z}_{\chi} \quad \text{and} \quad \Sigma^{1/2} \left(\Sigma^2/2 + \lambda I_d \right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \mathfrak{z}_{\chi}$$

are close as well.

Step 2. Using (118) we rewrite the difference between

$$-\Sigma^{1/2} \left((A^*)^\top A^*/2 + \lambda I_d \right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{v}^*) H_{\chi\chi}^{-1}(\boldsymbol{v}^*) \mathfrak{z}_{\chi} \quad \text{and} \quad -\Sigma^{1/2} \left(\Sigma^2/2 + \lambda I_d \right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{v}^*) H_{\chi\chi}^{-1}(\boldsymbol{v}^*) \mathfrak{z}_{\chi}$$

in the following form:

$$\begin{aligned} & \left\| \Sigma^{1/2} \left[\left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} - \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \right] H_{\theta\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \mathfrak{z}_{\chi} \right\| \\ & = \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \left((I_d + B)^{-1} - I_d \right) \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} H_{\theta\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \mathfrak{z}_{\chi} \right\| \\ & = \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} B(I_d + B)^{-1} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} H_{\theta\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \mathfrak{z}_{\chi} \right\|. \end{aligned}$$

Since, due to (117), the operator norm of $B(I_d + B)^{-1}$ does not exceed

$$||B(I_d + B)^{-1}|| \leq \frac{||B||}{1 - ||B||} \leq \frac{43}{42} \left(\left(\frac{17||\boldsymbol{\theta}^\circ||}{\mu} \right)^2 + \frac{||\mathbf{b}_\lambda||}{160\mu} \right),$$

we obtain that

$$\begin{aligned} \left\| \Sigma^{1/2} \left[\left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} - \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \right] H_{\boldsymbol{\theta}\chi}(\boldsymbol{v}^*) H_{\chi\chi}^{-1}(\boldsymbol{v}^*) \mathfrak{z}_{\chi} \right\| \\ &\leqslant \frac{43}{42} \left(\left(\frac{17 \|\boldsymbol{\theta}^\circ\|}{\mu} \right)^2 + \frac{\|\mathbf{b}_{\lambda}\|}{160\mu} \right) \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \right\| \\ &\cdot \left\| \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} H_{\boldsymbol{\theta}\chi}(\boldsymbol{v}^*) H_{\chi\chi}^{-1}(\boldsymbol{v}^*) \mathfrak{z}_{\chi} \right\|. \end{aligned}$$

Combining this inequality with (119) and (120), we obtain that

$$\left\| \Sigma^{1/2} \left[\left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1} - \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \right] H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \mathfrak{z}_{\chi} \right\|$$
(122)
$$\leq \frac{43}{42} \left(\left(\frac{17 \|\boldsymbol{\theta}^\circ\|}{\mu} \right)^2 + \frac{\|\mathbf{b}_{\lambda}\|}{160\mu} \right) \cdot (2\lambda)^{-1/4} \cdot \sqrt{\frac{43}{42}} \left\| \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d \right)^{-1/2} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^*) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*) \mathfrak{z}_{\chi} \right\|.$$

Finally, summing up the inequalities (121) and (122), we obtain the desired bound:

$$\begin{split} \left\| \Sigma^{1/2} \mathbf{s}_{\boldsymbol{\theta}}^{*} + \Sigma^{1/2} \left(\Sigma^{2} + \lambda I_{d} \right)^{-1} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\| \\ & \leq \frac{1.001 \|\boldsymbol{\theta}^{\circ}\|^{2}}{\mu^{2}} \sqrt{\frac{43}{42}} \cdot (2\lambda)^{-1/4} \left\| \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\| \\ & + \frac{43}{42} \left(\left(\frac{17 \|\boldsymbol{\theta}^{\circ}\|}{\mu} \right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{160\mu} \right) \cdot (2\lambda)^{-1/4} \cdot \sqrt{\frac{43}{42}} \left\| \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\| \\ & \leq \lambda^{-1/4} \left(\left(\frac{17 \|\boldsymbol{\theta}^{\circ}\|}{\mu} \right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{160\mu} \right) \left\| \left(\frac{1}{2} (A^{*})^{\top} A^{*} + \lambda I_{d} \right)^{-1/2} H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^{*}) H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^{*}) \mathfrak{z}_{\chi} \right\|. \end{split}$$

It only remains to note that, due to (47), the expression in the right-hand side does not exceed

$$\left(4\sqrt{2} + \frac{8\sqrt{2}}{112^2} + \frac{8}{112^2} \right) \frac{\|\boldsymbol{\theta}^\circ\|}{14\lambda^{1/4}} \left(\left(\frac{17\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{\|\mathbf{b}_\lambda\|}{160\mu} \right) \sqrt{\Psi(n,\delta)}$$

$$\leq \frac{\|\boldsymbol{\theta}^\circ\|}{2\lambda^{1/4}} \left(\left(\frac{17\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{\|\mathbf{b}_\lambda\|}{160\mu} \right) \sqrt{\Psi(n,\delta)},$$

and the claim of the lemma follows.

D.6 Proof of Lemma 6.5

The proof of the lemma is based on the expansion of $H_{\theta\chi}(\boldsymbol{v}^*)H_{\chi\chi}^{-1}(\boldsymbol{v}^*)\mathfrak{z}_{\chi}$ from Lemma D.3. According to Lemma B.2 (see (ii) and (iv)), it holds that

$$\|A^* oldsymbol{ heta}^* - oldsymbol{\eta}^*\| \leqslant \sqrt{\lambda} \|oldsymbol{ heta}^\circ\| \quad ext{and} \quad \|oldsymbol{ heta}^*\| \leqslant \|oldsymbol{ heta}^\circ\|.$$

This means that θ^* belongs to $\Upsilon(\rho^*)$ with $\rho^* = \|\theta^\circ\|/\mu \leq 1/112$. Applying Lemma D.3 with $\upsilon = \upsilon^*$ and $\rho = \rho^*$, we obtain that

$$\begin{split} H_{\boldsymbol{\theta}\chi}(\boldsymbol{v}^*)H_{\chi\chi}^{-1}(\boldsymbol{v}^*)\boldsymbol{\mathfrak{z}}_{\chi} &+ \frac{1}{2}(A^*)^{\top}\mathbf{U} - \frac{1}{2}(A^*)^{\top}(\widehat{\Sigma} - \Sigma)(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) - (\widehat{\Sigma} - \Sigma)(A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*) \\ &= -\left(r_1 - \frac{r_2\mu^2 \|\boldsymbol{\theta}^*\|^2}{\mu^2 + \|\boldsymbol{\theta}^*\|^2}\right)(A^*)^{\top}(\mathbf{Z} - \mathbb{E}\mathbf{Z}) + \frac{r_2\mu^2}{\mu^2 + \|\boldsymbol{\theta}^*\|^2}\left((A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)^{\top}(\mathbf{Z} - \mathbb{E}\mathbf{Z})\right)\boldsymbol{\theta}^* \\ &- \left(\frac{\|\boldsymbol{\theta}\|^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} - \frac{\|\boldsymbol{\theta}\|^2}{2(\mu^2 + \|\boldsymbol{\theta}\|^2)} + \frac{r'_2\mu^4}{\mu^2 + \|\boldsymbol{\theta}\|^2}\right)(A^*)^{\top}(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^* \\ &- \frac{1}{2\mu^2 + \|\boldsymbol{\theta}^*\|^2}\left((A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)^{\top}(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^*\right)\boldsymbol{\theta}^*, \end{split}$$

where $r_1 \in [0, (\rho^*)^2]$, $r_2 = 0.5\mu^{-2} + r'_2$, and $r'_2 \in [0, (\rho^*)^2/\mu^2]$ are defined in Proposition A.2. Let us denote

$$\mathbf{R} = -\left(r_1 - \frac{r_2 \mu^2 \|\boldsymbol{\theta}^*\|^2}{\mu^2 + \|\boldsymbol{\theta}^*\|^2}\right) (A^*)^\top (\mathbf{Z} - \mathbb{E}\mathbf{Z}) - \left(\frac{\|\boldsymbol{\theta}\|^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} - \frac{\|\boldsymbol{\theta}\|^2}{2(\mu^2 + \|\boldsymbol{\theta}\|^2)} + \frac{r'_2 \mu^4}{\mu^2 + \|\boldsymbol{\theta}\|^2}\right) (A^*)^\top (\widehat{\Sigma} - \Sigma) \boldsymbol{\theta}^*$$

and

$$\mathbf{Q} = \frac{r_2 \mu^2}{\mu^2 + \|\boldsymbol{\theta}^*\|^2} \left((A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^*)^\top (\mathbf{Z} - \mathbb{E}\mathbf{Z}) \right) \boldsymbol{\theta}^* - \frac{1}{2\mu^2 + \|\boldsymbol{\theta}^*\|^2} \left((A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^*)^\top (\widehat{\Sigma} - \Sigma) \boldsymbol{\theta}^* \right) \boldsymbol{\theta}^*.$$

With the introduced notations, it holds that

$$H_{\boldsymbol{\theta}\chi}(\boldsymbol{\upsilon}^*)H_{\chi\chi}^{-1}(\boldsymbol{\upsilon}^*)\mathfrak{z}_{\chi}+0.5\boldsymbol{\tau}=\mathtt{R}+\mathtt{Q}.$$
(123)

Let us recall that on the event \mathcal{E}_2^* we have (see (43)–(45))

$$|(A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)^{\top}(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^*| \leqslant 4C_X \|\Sigma^{1/2}\boldsymbol{\theta}^*\| \|\Sigma^{1/2}(A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)\| \sqrt{\frac{1 + \log(4/\delta)}{n}},$$
$$|(A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)^{\top}(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^\circ| \leqslant 4C_X \|\Sigma^{1/2}\boldsymbol{\theta}^\circ\| \|\Sigma^{1/2}(A^*\boldsymbol{\theta}^* - \boldsymbol{\eta}^*)\| \sqrt{\frac{1 + \log(4/\delta)}{n}},$$

and

$$\left| (A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^*)^\top \mathbf{U} \right| \leq 8\sigma \| \Sigma^{1/2} (A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^*) \| \sqrt{\frac{1 + \log(4/\delta)}{n}}.$$

Since $||A^*\theta^* - \eta^*|| \leq \sqrt{\lambda} ||\theta^\circ||$ and $||\theta^*|| \leq ||\theta^\circ||$ due to Lemma B.2, it holds that

$$\begin{aligned} \left| \frac{r_{2}\mu^{2}}{\mu^{2} + \|\boldsymbol{\theta}^{*}\|^{2}} \left((A^{*}\boldsymbol{\theta}^{*} - \boldsymbol{\eta}^{*})^{\top} (\mathbf{Z} - \mathbb{E}\mathbf{Z}) \right) - \frac{1}{2\mu^{2} + \|\boldsymbol{\theta}^{*}\|^{2}} \left((A^{*}\boldsymbol{\theta}^{*} - \boldsymbol{\eta}^{*})^{\top} (\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^{*} \right) \right| \\ &\leq \frac{4\|\Sigma\|^{1/2} \|A^{*}\boldsymbol{\theta}^{*} - \boldsymbol{\eta}^{*}\|}{\mu^{2}} \left((1 + (\rho^{*})^{2}) \left(\sigma + C_{X} \|\Sigma\|^{1/2} \|\boldsymbol{\theta}^{\circ}\| \right) \sqrt{\frac{1 + \log(4/\delta)}{n}} \right. \\ &\leq 1.001 \cdot \frac{4\|\boldsymbol{\theta}^{\circ}\|^{2} \sqrt{\lambda}}{\mu^{2}} \left(\frac{\sigma\|\Sigma\|^{1/2}}{\|\boldsymbol{\theta}^{\circ}\|} + C_{X} \|\Sigma\| \right) \sqrt{\frac{1 + \log(4/\delta)}{n}} \\ &\leq 1.001 \cdot \frac{2\|\boldsymbol{\theta}^{\circ}\|^{2}}{7\mu^{2}} \sqrt{\lambda\Psi(n, \delta)}. \end{aligned}$$

Let $\lambda_1(\Sigma), \ldots, \lambda_d(\Sigma)$ be the eigenvalues of Σ . Let us fix an arbitrary $j \in \{1, \ldots, d\}$. Applying the Young inequality

$$ab \leqslant \frac{a^p}{p} + \frac{b^q}{q}$$
 with $a = \sqrt{\lambda_j(\Sigma)}$, $b = (2\lambda/3)^{3/4}$, $p = 4$, and $q = 4/3$,

we obtain that

$$\frac{\sqrt{\lambda_j(\Sigma)}}{\lambda_j(\Sigma)^2/2 + \lambda} \leqslant \frac{(2\lambda/3)^{-3/4}}{2} \quad \text{for any } j \in \{1, \dots, d\},$$

and then

$$\left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \right\| = \max_{1 \le j \le d} \left\{ \frac{\sqrt{\lambda_j(\Sigma)}}{\lambda_j(\Sigma)^2/2 + \lambda} \right\} \le \frac{(2\lambda/3)^{-3/4}}{2}.$$
 (124)

Then, using the inequality

$$\left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \boldsymbol{\theta}^* \right\| \leqslant \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \right\| \| \boldsymbol{\theta}^* \| \leqslant \frac{1}{2} \left(\frac{2\lambda}{3} \right)^{-3/4} \| \boldsymbol{\theta}^\circ \|,$$

we conclude that

$$\left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \mathbb{Q} \right\| \leq \frac{1}{2} \left(\frac{2\lambda}{3} \right)^{-3/4} \|\boldsymbol{\theta}^\circ\| \cdot 1.001 \cdot \frac{2\|\boldsymbol{\theta}^\circ\|^2}{7\mu^2} \sqrt{\lambda \Psi(n, \delta)}$$

$$\leq \frac{\|\boldsymbol{\theta}^\circ\|^3}{5\mu^2 \lambda^{1/4}} \sqrt{\Psi(n, \delta)}.$$
(125)

On the other hand, since $0 \le r_1 \le (\rho^*)^2$ and $0 \le r'_2 \le (\rho^*)^2/\mu^2$, it holds that

$$\left| r_1 - \frac{r_2 \mu^2 \|\boldsymbol{\theta}^*\|^2}{\mu^2 + \|\boldsymbol{\theta}^*\|^2} \right| \leq (\rho^*)^2 = \frac{\|\boldsymbol{\theta}^\circ\|^2}{\mu^2}$$

and

$$\left|\frac{\|\boldsymbol{\theta}\|^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} - \frac{\|\boldsymbol{\theta}\|^2}{2(\mu^2 + \|\boldsymbol{\theta}\|^2)} + \frac{r'_2\mu^4}{\mu^2 + \|\boldsymbol{\theta}\|^2}\right| \leqslant 2(\rho^*)^2 = \frac{2\|\boldsymbol{\theta}^\circ\|^2}{\mu^2}.$$

Then

$$\begin{aligned} \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \mathbf{R} \right\| &\leq \frac{\|\boldsymbol{\theta}^\circ\|^2}{\mu^2} \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} (A^*)^\top (\mathbf{Z} - \mathbb{E}\mathbf{Z}) \right\| \\ &+ \frac{2\|\boldsymbol{\theta}^\circ\|^2}{\mu^2} \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} (A^*)^\top (\widehat{\Sigma} - \Sigma) \boldsymbol{\theta}^* \right\|. \end{aligned}$$

Let us consider the operator norm of

$$\Sigma^{1/2} \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1} (A^*)^\top.$$

Due to the triangle inequality, we have

$$\left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} (A^*)^\top \right\| \leq \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \Sigma \right\| + \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \right\| \| \Sigma - A^* \|.$$

Let us note that we have already bounded the operator norm of $\Sigma^{1/2}(\Sigma^2/2 + \lambda I_d)^{-1}$ (see (124)). Similarly, using Young's inequality

$$ab \leqslant \frac{a^p}{p} + \frac{b^q}{q}$$
 with $a = \lambda_j(\Sigma)^{3/2}$, $b = (6\lambda)^{1/4}$, $p = 4/3$, and $q = 4$,

we observe that

$$\left\|\Sigma^{1/2} \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1} \Sigma\right\| = \max_{1 \le j \le d} \left\{\frac{\lambda_j(\Sigma)^{3/2}}{\lambda_j(\Sigma)^2/2 + \lambda}\right\} \leqslant \frac{3}{2(6\lambda)^{1/4}}.$$

Thus, it holds that

$$\left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} (A^*)^\top \right\| \leq \frac{3}{2(6\lambda)^{1/4}} + \frac{(2\lambda/3)^{-3/4}}{2} \|A^* - \Sigma\|.$$

Moreover, Lemma B.3 and the condition (7) imply that

$$\frac{\|A^* - \Sigma\|}{\sqrt{\lambda}} \leqslant \left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{\|\mathbf{b}_\lambda\|}{230\mu} \leqslant \frac{1}{64} + \frac{1}{112 \cdot 230} < \frac{1}{63},$$

and we obtain that

$$\left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} (A^*)^\top \right\| \leqslant \frac{3}{2(6\lambda)^{1/4}} + \frac{(2\lambda/3)^{-3/4}}{126} \leqslant \lambda^{1/4}.$$
(126)

This yields that

$$\begin{split} \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \mathbf{R} \right\| &\leq \frac{\|\boldsymbol{\theta}^\circ\|^2}{\mu^2} \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} (A^*)^\top (\mathbf{Z} - \mathbb{E} \mathbf{Z}) \right\| \\ &+ \frac{2\|\boldsymbol{\theta}^\circ\|^2}{\mu^2} \left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} (A^*)^\top (\widehat{\Sigma} - \Sigma) \boldsymbol{\theta}^* \right\| \\ &\leq \frac{\|\boldsymbol{\theta}^\circ\|^2}{\mu^2 \lambda^{1/4}} \left(\|\mathbf{U}\| + \|(\widehat{\Sigma} - \Sigma) \boldsymbol{\theta}^\circ\| + 2\|(\widehat{\Sigma} - \Sigma) \boldsymbol{\theta}^*\| \right). \end{split}$$

Recall that, on the intersection of \mathcal{E}_1 and \mathcal{E}_2^* , we have (see (31), (32), and (36))

$$\|(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^{\circ}\| \leq 4C_X \|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}, \quad \|(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^{*}\| \leq 4C_X \|\Sigma\| \|\boldsymbol{\theta}^{*}\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}},$$

and

and

$$\|\mathbf{U}\| \leqslant 8\sigma \|\Sigma\|^{1/2} \sqrt{\frac{\mathtt{r}(\Sigma) + \log(4/\delta)}{n}}.$$

Taking into account that, due to Lemma B.2, $\|\theta^*\| \leq \|\theta^\circ\|$, we finally get that

$$\left\| \Sigma^{1/2} \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1} \mathbb{R} \right\| \leq \frac{8 \|\boldsymbol{\theta}^\circ\|^3}{\mu^2 \lambda^{1/4}} \left(\frac{\sigma \|\Sigma\|^{1/2}}{\|\boldsymbol{\theta}^\circ\|} + \frac{3C_X \|\Sigma\|}{2} \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}$$

$$\leq \frac{4 \|\boldsymbol{\theta}^\circ\|^3}{7\mu^2 \lambda^{1/4}} \sqrt{\Psi(n,\delta)}.$$
(127)

The identity (123) and the inequalities (125) and (127) yield that

$$\left\|H_{\boldsymbol{\theta}\chi}(\boldsymbol{v}^*)H_{\chi\chi}^{-1}(\boldsymbol{v}^*)\boldsymbol{\mathfrak{z}}_{\chi}+0.5\boldsymbol{\tau}\right\| \leqslant \frac{\|\boldsymbol{\theta}^\circ\|^3}{5\mu^2\lambda^{1/4}}\sqrt{\Psi(n,\delta)} + \frac{4\|\boldsymbol{\theta}^\circ\|^3}{7\mu^2\lambda^{1/4}}\sqrt{\Psi(n,\delta)} \leqslant \frac{\|\boldsymbol{\theta}^\circ\|^3}{\mu^2\lambda^{1/4}}\sqrt{\Psi(n,\delta)}.$$

D.7 Proof of Lemma 6.6

Note that, due to the definitions of τ and ζ , we have

$$\boldsymbol{\tau} - \boldsymbol{\zeta} = (A^* - \Sigma)\mathbf{U} + (A^*)^\top (\widehat{\Sigma} - \Sigma)(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ - \mathbf{b}_\lambda) - (A^* - \Sigma)^\top (\widehat{\Sigma} - \Sigma)\mathbf{b}_\lambda - (\widehat{\Sigma} - \Sigma)(2A^*\boldsymbol{\theta}^* - 2\boldsymbol{\eta}^* - \mathbf{b}_\lambda).$$
Then the triangle inequality yields that

$$\begin{split} \left\| \Sigma^{1/2} \left(\Sigma^{2} + 2\lambda I_{d} \right)^{-1} \left(\boldsymbol{\tau} - \boldsymbol{\zeta} \right) \right\| &\leq \left\| \Sigma^{1/2} \left(\Sigma^{2} + 2\lambda I_{d} \right)^{-1} \right\| \left\| A^{*} - \Sigma \right\| \left(\left\| \mathbf{U} \right\| + \left\| (\widehat{\Sigma} - \Sigma) \mathbf{b}_{\lambda} \right\| \right) \\ &+ \left\| \Sigma^{1/2} \left(\Sigma^{2} + 2\lambda I_{d} \right)^{-1} \left(A^{*} \right)^{\top} \right\| \left\| (\widehat{\Sigma} - \Sigma) (\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ} - \mathbf{b}_{\lambda}) \right\| \\ &+ \left\| \Sigma^{1/2} \left(\Sigma^{2} + 2\lambda I_{d} \right)^{-1} \right\| \left\| (\widehat{\Sigma} - \Sigma) (2A^{*}\boldsymbol{\theta}^{*} - 2\boldsymbol{\eta}^{*} - \Sigma \mathbf{b}_{\lambda}) \right\| . \end{split}$$

In view of (124) and (126), we obtain that

$$\begin{split} \left\| \Sigma^{1/2} \left(\Sigma^2 + 2\lambda I_d \right)^{-1} \left(\boldsymbol{\tau} - \boldsymbol{\zeta} \right) \right\| &\leq \frac{(2\lambda/3)^{-3/4}}{4} \left\| A^* - \Sigma \right\| \left(\|\mathbf{U}\| + \left\| (\widehat{\Sigma} - \Sigma) \mathbf{b}_\lambda \right\| \right) \\ &+ \frac{1}{2\lambda^{1/4}} \left\| (\widehat{\Sigma} - \Sigma) (\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ - \mathbf{b}_\lambda) \right\| \\ &+ \frac{(2\lambda/3)^{-3/4}}{2} \left\| (\widehat{\Sigma} - \Sigma) \left(A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^* - \frac{1}{2} \Sigma \mathbf{b}_\lambda \right) \right\|. \end{split}$$

According to Theorem 2.5, there exists an event \mathcal{E}_3 , $\mathbb{P}(\mathcal{E}_3) \ge 1 - 3\delta/2$ such that the following inequalities holds simultaneously on \mathcal{E}_3 :

$$\left\| (\widehat{\Sigma} - \Sigma) \mathbf{b}_{\lambda} \right\| \leqslant 4C_X \|\Sigma\|^{1/2} \|\Sigma^{1/2} \mathbf{b}_{\lambda}\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}, \\ \left\| (\widehat{\Sigma} - \Sigma)(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ - \mathbf{b}_{\lambda}) \right\| \leqslant 4C_X \|\Sigma\|^{1/2} \|\Sigma^{1/2}(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ - \mathbf{b}_{\lambda})\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}, \\ \left\| (\widehat{\Sigma} - \Sigma) \left(A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^* - \frac{1}{2} \Sigma \mathbf{b}_{\lambda} \right) \right\| \leqslant 4C_X \|\Sigma\| \left\| A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^* - \frac{1}{2} \Sigma \mathbf{b}_{\lambda} \right\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}.$$

From now on, we restrict our attention on the intersection of \mathcal{E}_1 and \mathcal{E}_3 . Since on \mathcal{E}_1 (see (32))

$$\|\mathbf{U}\| \leqslant 8\sigma \|\Sigma\|^{1/2} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}},$$

it holds that

$$\begin{aligned} \left\| \Sigma^{1/2} \left(\Sigma^{2} + 2\lambda I_{d} \right)^{-1} \left(\boldsymbol{\tau} - \boldsymbol{\zeta} \right) \right\| \\ &\leqslant (2\lambda/3)^{-3/4} \|A^{*} - \Sigma\| \left(2\sigma \|\Sigma\|^{1/2} + C_{X} \|\Sigma\| \|\mathbf{b}_{\lambda}\| \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ \frac{2C_{X}}{\lambda^{1/4}} \|\Sigma\| \|\boldsymbol{\theta}^{*} - \boldsymbol{\theta}^{\circ} - \mathbf{b}_{\lambda}\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &+ 2(2\lambda/3)^{-3/4} C_{X} \|\Sigma\| \left\| A^{*}\boldsymbol{\theta}^{*} - \boldsymbol{\eta}^{*} - \frac{1}{2} \Sigma \mathbf{b}_{\lambda} \right\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}. \end{aligned}$$
(128)

It remains to bound the term in the right-hand side one by one. Applying Lemma B.3 and taking into account the inequalities (7) and $\|\mathbf{b}_{\lambda}\| \leq \|\boldsymbol{\theta}^{\circ}\|$, we obtain that

$$(2\lambda/3)^{-3/4} \|A^* - \Sigma\| \left(8\sigma \|\Sigma\|^{1/2} + C_X \|\Sigma\| \|\mathbf{b}_\lambda\| \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ \leqslant (2\lambda/3)^{-3/4} \cdot \sqrt{\lambda} \left(\left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu} \right)^2 + \frac{\|\mathbf{b}_\lambda\|}{230\mu} \right) \left(2\sigma \|\Sigma\|^{1/2} + C_X \|\Sigma\| \|\boldsymbol{\theta}^\circ\| \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}.$$

Due to Lemma B.4, it holds that

$$2(2\lambda/3)^{-3/4}C_X \|\Sigma\| \left\| A^* \boldsymbol{\theta}^* - \boldsymbol{\eta}^* - \frac{1}{2} \Sigma \mathbf{b}_\lambda \right\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}$$

$$\leq \frac{6 \cdot (3/2)^{3/4}}{\lambda^{1/4}} C_X \|\Sigma\| \|\boldsymbol{\theta}^\circ\| \left(\left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{\|\mathbf{b}_\lambda\|}{230\mu} \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}.$$

Finally, we use the following result to bound the second term in the right-hand side of (128). Lemma D.5. Assume that $\|\theta^{\circ}\| \leq \mu/49$ and $\|\Sigma\| \|\theta^{\circ}\| \leq \mu\sqrt{\lambda}/24$. Then it holds that

$$\|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ - \mathbf{b}_\lambda\| \leqslant \frac{7\|\boldsymbol{\theta}^\circ\|}{2} \left(\left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{70\|\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^\circ\|\|\mathbf{b}_\lambda\|}{\mu^2\sqrt{\lambda}} \right).$$

We provide the proof of Lemma D.5 in Appendix D.8.4. It immediately implies that

$$\begin{split} &\frac{2C_X}{\lambda^{1/4}} \|\Sigma\| \|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ - \mathbf{b}_\lambda\| \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &\leqslant \frac{7C_X \|\Sigma\| \|\boldsymbol{\theta}^\circ\|}{\lambda^{1/4}} \left(\left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{70\|\Sigma\| \|\boldsymbol{\theta}^\circ\| \|\mathbf{b}_\lambda\|}{\mu^2 \sqrt{\lambda}} \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &\leqslant \frac{7C_X \|\Sigma\| \|\boldsymbol{\theta}^\circ\|}{\lambda^{1/4}} \left(\left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{\|\mathbf{b}_\lambda\|}{230\mu} \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}. \end{split}$$

Hence, we obtain that

$$\begin{split} \left\| \Sigma^{1/2} \left(\Sigma^{2} + 2\lambda I_{d} \right)^{-1} \left(\boldsymbol{\tau} - \boldsymbol{\zeta} \right) \right\| \\ &\leqslant \frac{2 \|\boldsymbol{\theta}^{\circ}\|}{(2/3)^{3/4} \lambda^{1/4}} \left(\frac{\sigma \|\Sigma\|^{1/2}}{\|\boldsymbol{\theta}^{\circ}\|} + \left(1 + 3 + \frac{7}{2 \cdot (3/2)^{3/4}} \right) C_{X} \|\Sigma\| \right) \\ &\cdot \left(\left(\frac{14 \|\boldsymbol{\theta}^{\circ}\|}{\mu} \right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{230\mu} \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &\leqslant \frac{2 \|\boldsymbol{\theta}^{\circ}\|}{(2/3)^{3/4} \lambda^{1/4}} \left(\frac{\sigma \|\Sigma\|^{1/2}}{\|\boldsymbol{\theta}^{\circ}\|} + 7C_{X} \|\Sigma\| \right) \left(\left(\frac{14 \|\boldsymbol{\theta}^{\circ}\|}{\mu} \right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{230\mu} \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \\ &\leqslant \frac{7 \|\boldsymbol{\theta}^{\circ}\|}{(2/3)^{3/4} \lambda^{1/4}} \left(\frac{\sigma \|\Sigma\|^{1/2}}{\|\boldsymbol{\theta}^{\circ}\|} + 2C_{X} \|\Sigma\| \right) \left(\left(\frac{14 \|\boldsymbol{\theta}^{\circ}\|}{\mu} \right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{230\mu} \right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}. \end{split}$$

It remains to note that

$$\left(\frac{14\|\boldsymbol{\theta}^{\circ}\|}{\mu}\right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{230\mu} \leqslant \frac{1}{\sqrt{2}} \left(\left(\frac{17\|\boldsymbol{\theta}^{\circ}\|}{\mu}\right)^{2} + \frac{\|\mathbf{b}_{\lambda}\|}{160\mu} \right)$$

and

$$\left(\frac{\sigma \|\Sigma\|^{1/2}}{\|\boldsymbol{\theta}^{\circ}\|} + 2C_X \|\Sigma\|\right) \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}} \leqslant \frac{\sqrt{\Psi(n,\delta)}}{14}.$$

Then it holds that

$$\begin{split} \left\| \Sigma^{1/2} \left(\Sigma^2 + 2\lambda I_d \right)^{-1} \left(\boldsymbol{\tau} - \boldsymbol{\zeta} \right) \right\| &\leq \frac{(3/2)^{3/4} \|\boldsymbol{\theta}^\circ\|}{2\sqrt{2}\lambda^{1/4}} \left(\left(\frac{17\|\boldsymbol{\theta}^\circ\|}{\mu} \right)^2 + \frac{\|\mathbf{b}_\lambda\|}{160\mu} \right) \sqrt{\Psi(n,\delta)} \\ &\leq \frac{12\|\boldsymbol{\theta}^\circ\|}{25\lambda^{1/4}} \left(\left(\frac{17\|\boldsymbol{\theta}^\circ\|}{\mu} \right)^2 + \frac{\|\mathbf{b}_\lambda\|}{160\mu} \right) \sqrt{\Psi(n,\delta)}. \end{split}$$

D.8 Auxiliary results

D.8.1 Proof of Lemma D.2

Applying Proposition A.2, we obtain that

$$\breve{H}_{\theta\theta} - \left(\frac{1}{2}A^{\top}A + \lambda I_d\right) = \left[-r_1 + \frac{2r_2\mu^2 \|\theta\|^2}{\mu^2 + \|\theta\|^2} - \frac{\|\theta\|^2}{\mu^2} \left(1 + r'_3\mu^2 \|\theta\|^2\right)\right] A^{\top}A$$

$$- \frac{1}{\mu^2} \|A\theta - \eta\|^2 I_d + \frac{r_2\mu^2}{\mu^2 + \|\theta\|^2} \left(\theta(A\theta - \eta)^{\top}A + A^{\top}(A\theta - \eta)\theta^{\top}\right),$$
(129)

where

$$0 \leqslant r_1 \leqslant \rho^2$$
, $0 \leqslant r_2 \leqslant \frac{1}{\mu^2}$, and $0 \leqslant r'_3 \leqslant \frac{2}{\mu^4}$.

In the rest of the proof, we study the terms in the right-hand side of (129) one by one. First, let us note that the inequalities $\|\theta\|^2 \leq \rho^2(\mu^2 + \|\theta\|^2)$ and $\rho^2 \leq 1/2$ yield that

$$\left| -r_1 + \frac{2r_2\mu^2 \|\boldsymbol{\theta}\|^2}{\mu^2 + \|\boldsymbol{\theta}\|^2} - \frac{\|\boldsymbol{\theta}\|^2}{\mu^2} \left(1 + r_3'\mu^2 \|\boldsymbol{\theta}\|^2 \right) \right| \leq \rho^2 + \frac{2\|\boldsymbol{\theta}\|^2}{\mu^2 + \|\boldsymbol{\theta}\|^2} + \frac{\|\boldsymbol{\theta}\|^2(\mu^2 + 2\|\boldsymbol{\theta}\|^2)}{\mu^4} \\ \leq \rho^2 + 2\rho^2 + 2\rho^4 \leq 3\rho^2 + \frac{2\rho^2}{25}.$$

Then the first term in the right-hand side of (129) does not exceed

$$\left(3\rho^{2} + \frac{2\rho^{2}}{25}\right) \left\| \left(\frac{1}{2}A^{\top}A + \lambda I_{d}\right)^{-1/2} A^{\top}A \left(\frac{1}{2}A^{\top}A + \lambda I_{d}\right)^{-1/2} \right\| \leq 6\rho^{2} + 2\rho^{4} \leq 6\rho^{2} + \frac{4\rho^{2}}{25}.$$

Second, it holds that

$$\frac{\|A\boldsymbol{\theta} - \boldsymbol{\eta}\|^2}{\mu^2} \left\| \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{-1} \right\| \leq \rho^2 \lambda \cdot \frac{1}{\lambda} = \rho^2.$$

Finally, let us consider the latter term in the right-hand side of (129). The triangle inequality and the submultiplicativity of the operator norm imply that

$$\begin{split} & \frac{r_2\mu^2}{\mu^2 + \|\boldsymbol{\theta}\|^2} \left\| \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{-1/2} \left(\boldsymbol{\theta}(A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top}A + A^{\top}(A\boldsymbol{\theta} - \boldsymbol{\eta})\boldsymbol{\theta}^{\top}\right) \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{-1/2} \right\| \\ & \leqslant \frac{2}{\mu^2 + \|\boldsymbol{\theta}\|^2} \left\| \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{-1/2} \boldsymbol{\theta} \right\| \left\| A^{\top} \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{-1/2} \right\| \left\| A\boldsymbol{\theta} - \boldsymbol{\eta} \right\| \\ & \leqslant \frac{2}{\mu^2 + \|\boldsymbol{\theta}\|^2} \cdot \frac{\|\boldsymbol{\theta}\|}{\sqrt{\lambda}} \cdot \sqrt{2} \cdot \rho \mu \sqrt{\lambda} \leqslant \frac{2\sqrt{2}\rho\mu}{\sqrt{\mu^2 + \|\boldsymbol{\theta}\|^2}} \cdot \frac{\|\boldsymbol{\theta}\|}{\sqrt{\mu^2 + \|\boldsymbol{\theta}\|^2}} \leqslant \frac{2\sqrt{2}\rho\mu}{\mu} \cdot \rho = 2\sqrt{2}\rho^2. \end{split}$$

Hence, it holds that

$$\left\| \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{-1/2} \breve{H}_{\theta\theta} \left(\frac{1}{2}A^{\top}A + \lambda I_d\right)^{-1/2} - I_d \right\| \leq 6\rho^2 + \frac{4\rho^2}{25} + \rho^2 + 2\sqrt{2}\rho^2 \leq 10\rho^2.$$

D.8.2 Proof of Lemma D.3

The proof of Lemma D.3 relies on the expression (52) for the Hessian of $\nabla^2 \mathcal{L}(\boldsymbol{v})$ and Proposition A.2. Let us denote the inverse of $H_{\chi\chi}$ by J. It holds that

$$\begin{aligned} H_{\theta\chi} J_{\mathfrak{Z}\chi} &= \begin{pmatrix} H_{\theta\eta} & H_{\theta A} \end{pmatrix} \begin{pmatrix} J_{\eta\eta} & J_{\eta A} \\ J_{A\eta} & J_{AA} \end{pmatrix} \begin{pmatrix} \mathfrak{z}\eta \\ \mathfrak{z}A \end{pmatrix} \\ &= H_{\theta\eta} J_{\eta\eta} \mathfrak{z}\eta + H_{\theta\eta} J_{\etaA} \mathfrak{z}A + H_{\theta A} J_{A\eta} \mathfrak{z}\eta + H_{\theta A} J_{AA} \mathfrak{z}A. \end{aligned}$$

In the rest of the proof, we study the terms in the right-hand-side one by one. Using Proposition A.2, we obtain that

$$\begin{aligned} H_{\boldsymbol{\theta}A}J_{A\boldsymbol{\eta}} &= r_2 \left((A\boldsymbol{\theta} - \boldsymbol{\eta})^\top \otimes I_d + A^\top \otimes \boldsymbol{\theta}^\top \right) \left(I_d \otimes \widetilde{\boldsymbol{\theta}} \right) \\ &= r_2 (A\boldsymbol{\theta} - \boldsymbol{\eta})^\top \otimes \widetilde{\boldsymbol{\theta}} + r_2 A^\top \otimes \left(\boldsymbol{\theta}^\top \widetilde{\boldsymbol{\theta}} \right) \\ &= r_2 \widetilde{\boldsymbol{\theta}} (A\boldsymbol{\theta} - \boldsymbol{\eta})^\top + r_2 \left(\boldsymbol{\theta}^\top \widetilde{\boldsymbol{\theta}} \right) A^\top, \end{aligned}$$

where $\tilde{\boldsymbol{\theta}} = \mu^2 (\mu^2 I_d + \boldsymbol{\theta} \boldsymbol{\theta}^\top)^{-1} \boldsymbol{\theta}$. The Woodbury matrix identity yields that

$$\widetilde{\boldsymbol{ heta}} = rac{\mu^2}{\mu^2 + \|\boldsymbol{ heta}\|^2} \boldsymbol{ heta} \quad ext{and} \quad \boldsymbol{ heta}^{ op} \widetilde{\boldsymbol{ heta}} = rac{\mu^2 \|\boldsymbol{ heta}\|^2}{\mu^2 + \|\boldsymbol{ heta}\|^2}.$$

Then we have

$$H_{\boldsymbol{\theta}A}J_{A\boldsymbol{\eta}\boldsymbol{\vartheta}\boldsymbol{\eta}} = \frac{r_2\mu^2}{\mu^2 + \|\boldsymbol{\theta}\|^2} \left((A\boldsymbol{\theta} - \boldsymbol{\eta})^\top (\mathbf{Z} - \mathbb{E}\mathbf{Z}) \right) \boldsymbol{\theta} + \frac{r_2\mu^2 \|\boldsymbol{\theta}\|^2}{\mu^2 + \|\boldsymbol{\theta}\|^2} A^\top (\mathbf{Z} - \mathbb{E}\mathbf{Z}).$$
(130)

Similarly, we note that

$$H_{\theta\eta}J_{\eta\eta\mathfrak{z}\eta} = -A^{\top}J_{\eta\eta\mathfrak{z}\eta} = -\left(\frac{1}{2} + r_1\right)A^{\top}(\mathbf{Z} - \mathbb{E}\mathbf{Z}).$$
(131)

Due to the properties of the Kronecker product and the vectorization operator (see, for instance, [Puchkin and Rakhuba, 2024, Appendix A, eq. (25)]), it holds that

$$(I_d \otimes \widetilde{\boldsymbol{\theta}}^{\top}) \mathbf{vec}(\widehat{\Sigma} - \Sigma) = \mathbf{vec} \left(I_d(\widehat{\Sigma} - \Sigma) \widetilde{\boldsymbol{\theta}} \right) = (\widehat{\Sigma} - \Sigma) \widetilde{\boldsymbol{\theta}}.$$

Thus, we obtain that

$$H_{\theta\eta}J_{\eta A\mathfrak{Z}A} = -r_2\mu^2 A^{\top}(\widehat{\Sigma}-\Sigma)\widetilde{\theta} = -\frac{r_2\mu^4}{\mu^2 + \|\theta\|^2}A^{\top}(\widehat{\Sigma}-\Sigma)\theta.$$
(132)

It remains to consider $H_{\theta A}J_{AA\mathfrak{z}A}$. Let us focus on J_{AA} . Note that, due to (52), the matrix $H_{\chi\chi}$ can be represented in the following compact form using the Kronecker product:

$$H_{\chi\chi} = \begin{pmatrix} 2I_d & -I_d \otimes \boldsymbol{\theta}^\top \\ -I_d \otimes \boldsymbol{\theta} & I_d \otimes (\mu^2 I_d + \boldsymbol{\theta} \boldsymbol{\theta}^\top) \end{pmatrix}.$$

Then, according to the block-inversion formula (see (53) in Appendix A), it holds that

$$J_{AA} = \left(I_d \otimes (\mu^2 I_d + \boldsymbol{\theta} \boldsymbol{\theta}^\top) - \frac{1}{2} \left(I_d \otimes \boldsymbol{\theta}^\top\right) (I_d \otimes \boldsymbol{\theta})\right)^{-1}$$
$$= \left(I_d \otimes \left(\mu^2 I_d + \frac{1}{2} \boldsymbol{\theta} \boldsymbol{\theta}^\top\right)\right)^{-1} = I_d \otimes \left(\mu^2 I_d + \frac{1}{2} \boldsymbol{\theta} \boldsymbol{\theta}^\top\right)^{-1}$$

Using the Woodbury matrix identity, we obtain that

$$J_{AA} = I_d \otimes \left(\frac{1}{\mu^2} I_d - \frac{1}{\mu^2 (2\mu^2 + \|\boldsymbol{\theta}\|^2)} \boldsymbol{\theta} \boldsymbol{\theta}^\top\right).$$

Since $\mathfrak{z}_A = \mu^2 \mathbf{vec}(\widehat{\Sigma} - \Sigma)$, we have

$$H_{\boldsymbol{\theta}A}J_{AA\boldsymbol{\vartheta}A} = \mu^{2} \left((A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} \otimes I_{d} \right) \left(I_{d} \otimes \left(\frac{1}{\mu^{2}} I_{d} - \frac{1}{\mu^{2}(2\mu^{2} + \|\boldsymbol{\theta}\|^{2})} \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right) \right) \operatorname{vec} \left(\widehat{\Sigma} - \Sigma \right) + \mu^{2} \left(A^{\top} \otimes \boldsymbol{\theta}^{\top} \right) \left(I_{d} \otimes \left(\frac{1}{\mu^{2}} I_{d} - \frac{1}{\mu^{2}(2\mu^{2} + \|\boldsymbol{\theta}\|^{2})} \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right) \right) \operatorname{vec} \left(\widehat{\Sigma} - \Sigma \right) = \left((A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} \otimes \left(I_{d} - \frac{1}{2\mu^{2} + \|\boldsymbol{\theta}\|^{2}} \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right) \right) \operatorname{vec} \left(\widehat{\Sigma} - \Sigma \right) + \left(A^{\top} \otimes \boldsymbol{\theta}^{\top} \left(I_{d} - \frac{1}{2\mu^{2} + \|\boldsymbol{\theta}\|^{2}} \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right) \right) \operatorname{vec} \left(\widehat{\Sigma} - \Sigma \right).$$
(133)

Using the properties of the Kronecker product and the vectorization operator **vec** one more time, we obtain that

$$\begin{pmatrix} (A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} \otimes \left(I_d - \frac{1}{2\mu^2 + \|\boldsymbol{\theta}\|^2} \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right) \end{pmatrix} \operatorname{vec} \left(\widehat{\Sigma} - \Sigma \right) = \operatorname{vec} \left((A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} (\widehat{\Sigma} - \Sigma) \left(I_d - \frac{1}{2\mu^2 + \|\boldsymbol{\theta}\|^2} \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right) \right)$$

$$= \left(I_d - \frac{1}{2\mu^2 + \|\boldsymbol{\theta}\|^2} \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right) (\widehat{\Sigma} - \Sigma) (A\boldsymbol{\theta} - \boldsymbol{\eta})$$

$$(134)$$

and

$$\left(A^{\top} \otimes \boldsymbol{\theta}^{\top} \left(I_d - \frac{1}{2\mu^2 + \|\boldsymbol{\theta}\|^2} \boldsymbol{\theta} \boldsymbol{\theta}^{\top} \right) \right) \operatorname{\mathbf{vec}} \left(\widehat{\Sigma} - \Sigma \right) = \frac{2\mu^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} \left(A^{\top} \otimes \boldsymbol{\theta}^{\top} \right) \operatorname{\mathbf{vec}} \left(\widehat{\Sigma} - \Sigma \right)$$
$$= \frac{2\mu^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} \operatorname{\mathbf{vec}} \left(A^{\top} (\widehat{\Sigma} - \Sigma) \boldsymbol{\theta} \right)$$
$$= \frac{2\mu^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} A^{\top} (\widehat{\Sigma} - \Sigma) \boldsymbol{\theta}.$$
(135)

Substituting the equalities (134) and (135) into (133), we obtain that

$$H_{\boldsymbol{\theta}A}J_{AA\boldsymbol{\mathfrak{J}}A} = (\widehat{\Sigma} - \Sigma)(A\boldsymbol{\theta} - \boldsymbol{\eta}) - \frac{1}{2\mu^2 + \|\boldsymbol{\theta}\|^2} ((A\boldsymbol{\theta} - \boldsymbol{\eta})^\top (\widehat{\Sigma} - \Sigma)\boldsymbol{\theta})\boldsymbol{\theta} + \frac{2\mu^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} A^\top (\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}.$$
(136)

Summing up (130), (131), (132), and (136), we conclude that

$$\begin{aligned} H_{\boldsymbol{\theta}\chi} J_{\boldsymbol{\mathfrak{z}}\chi} &= -\left(\frac{1}{2} + r_1 - \frac{r_2 \mu^2 \|\boldsymbol{\theta}\|^2}{\mu^2 + \|\boldsymbol{\theta}\|^2}\right) A^\top (\mathbf{Z} - \mathbb{E}\mathbf{Z}) + \frac{r_2 \mu^2}{\mu^2 + \|\boldsymbol{\theta}\|^2} \left((A\boldsymbol{\theta} - \boldsymbol{\eta})^\top (\mathbf{Z} - \mathbb{E}\mathbf{Z}) \right) \boldsymbol{\theta} \\ &+ \left(\frac{2\mu^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} - \frac{r_2 \mu^4}{\mu^2 + \|\boldsymbol{\theta}\|^2} \right) A^\top (\widehat{\boldsymbol{\Sigma}} - \boldsymbol{\Sigma}) \boldsymbol{\theta} - \frac{1}{2\mu^2 + \|\boldsymbol{\theta}\|^2} \left((A\boldsymbol{\theta} - \boldsymbol{\eta})^\top (\widehat{\boldsymbol{\Sigma}} - \boldsymbol{\Sigma}) \boldsymbol{\theta} \right) \boldsymbol{\theta} \\ &+ (\widehat{\boldsymbol{\Sigma}} - \boldsymbol{\Sigma}) (A\boldsymbol{\theta} - \boldsymbol{\eta}). \end{aligned}$$

Due to the definition of r_2 (see Proposition A.2), it holds that $r_2 = 0.5\mu^{-2} + r'_2$, where $0 \le r'_2 \le \rho^2/\mu^2$. Using this identity, we can simplify the coefficient before $A^{\top}(\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}$:

$$\frac{2\mu^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} - \frac{r_2\mu^4}{\mu^2 + \|\boldsymbol{\theta}\|^2} = 1 - \frac{\|\boldsymbol{\theta}\|^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} - \frac{\mu^2}{2(\mu^2 + \|\boldsymbol{\theta}\|^2)} - \frac{r'_2\mu^4}{\mu^2 + \|\boldsymbol{\theta}\|^2}$$
$$= 1 - \frac{\|\boldsymbol{\theta}\|^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} - \frac{1}{2} + \frac{\|\boldsymbol{\theta}\|^2}{2(\mu^2 + \|\boldsymbol{\theta}\|^2)} - \frac{r'_2\mu^4}{\mu^2 + \|\boldsymbol{\theta}\|^2}$$
$$= \frac{1}{2} - \frac{\|\boldsymbol{\theta}\|^2}{2\mu^2 + \|\boldsymbol{\theta}\|^2} + \frac{\|\boldsymbol{\theta}\|^2}{2(\mu^2 + \|\boldsymbol{\theta}\|^2)} - \frac{r'_2\mu^4}{\mu^2 + \|\boldsymbol{\theta}\|^2}.$$

Finally, taking into account the representation $\mathbf{Z} - \mathbb{E}\mathbf{Z} = \mathbf{U} + (\widehat{\Sigma} - \Sigma)\boldsymbol{\theta}^{\circ}$, we obtain that

$$\begin{aligned} H_{\boldsymbol{\theta}\chi} J_{\boldsymbol{\mathfrak{Z}}\chi} &= -\frac{1}{2} A^{\top} \mathbf{U} + \frac{1}{2} A^{\top} (\widehat{\Sigma} - \Sigma) (\boldsymbol{\theta} - \boldsymbol{\theta}^{\circ}) + (\widehat{\Sigma} - \Sigma) (A\boldsymbol{\theta} - \boldsymbol{\eta}) \\ &- \left(r_{1} - \frac{r_{2} \mu^{2} \|\boldsymbol{\theta}\|^{2}}{\mu^{2} + \|\boldsymbol{\theta}\|^{2}} \right) A^{\top} (\mathbf{Z} - \mathbb{E}\mathbf{Z}) + \frac{r_{2} \mu^{2}}{\mu^{2} + \|\boldsymbol{\theta}\|^{2}} \left((A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} (\mathbf{Z} - \mathbb{E}\mathbf{Z}) \right) \boldsymbol{\theta} \\ &- \left(\frac{\|\boldsymbol{\theta}\|^{2}}{2\mu^{2} + \|\boldsymbol{\theta}\|^{2}} - \frac{\|\boldsymbol{\theta}\|^{2}}{2(\mu^{2} + \|\boldsymbol{\theta}\|^{2})} + \frac{r_{2}' \mu^{4}}{\mu^{2} + \|\boldsymbol{\theta}\|^{2}} \right) A^{\top} (\widehat{\Sigma} - \Sigma) \boldsymbol{\theta} \\ &- \frac{1}{2\mu^{2} + \|\boldsymbol{\theta}\|^{2}} \left((A\boldsymbol{\theta} - \boldsymbol{\eta})^{\top} (\widehat{\Sigma} - \Sigma) \boldsymbol{\theta} \right) \boldsymbol{\theta}. \end{aligned}$$

D.8.3 Proof of Lemma D.4

Step 1: a bound on the operator norm. Let us start with an upper bound on the operator norm of

$$\frac{1}{2} \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \left(A^{\top}A - \Sigma^2\right) \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2}.$$

Due to the definition of S and the triangle inequality, it holds that

$$\frac{1}{2} \left\| \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \left(A^\top A - \Sigma^2 \right) \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \right\|$$

$$\leq \left\| \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \Sigma (A - \Sigma) \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \right\|$$

$$+ \frac{1}{2} \left\| \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} (A^\top - \Sigma) (A - \Sigma) \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \right\|.$$

Let us note that

$$\left\| \left(\frac{1}{2}\Sigma^2 + \lambda I_d \right)^{-1/2} \Sigma \right\| \leqslant \sqrt{2} \quad \text{and} \quad \left\| \left(\frac{1}{2}\Sigma^2 + \lambda I_d \right)^{-1/2} \right\| \leqslant \frac{1}{\sqrt{\lambda}}.$$
(137)

Taking into account submultiplicativity of the operator norm, we obtain that

$$\left\| \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \Sigma(A - \Sigma) \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \right\| \leq \|A - \Sigma\|\sqrt{\frac{2}{\lambda}}$$

and

$$\frac{1}{2} \left\| \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} (A^\top - \Sigma) (A - \Sigma) \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \right\| \leq \frac{\|A - \Sigma\|^2}{2\lambda}.$$

Hence, it holds that

$$\frac{1}{2} \left\| \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \left(A^\top A - \Sigma^2 \right) \left(\frac{1}{2} \Sigma^2 + \lambda I_d \right)^{-1/2} \right\| \leqslant \|A - \Sigma\| \sqrt{\frac{2}{\lambda}} + \frac{\|A - \Sigma\|^2}{2\lambda}.$$

D.8.4 Proof of Lemma D.5

The proof relies on the identity

$$\left(\frac{1}{2}(A^*)^{\top}A^* + \lambda I_d\right)(\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ) = -\lambda\boldsymbol{\theta}^\circ - \frac{1}{2}(A^*)^{\top}(A^* - \Sigma)\boldsymbol{\theta}^\circ$$

following from the first-order optimality condition $\nabla_{\theta} \mathcal{L}(\theta^*, \eta^*, A^*) = \mathbf{0}$ (see (74) in the proof of Lemma **B**.2 for the details). Let us introduce

$$B = \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \left((A^*)^\top A^* - \Sigma^2 \right) \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2}.$$

Then we can rewrite the difference $(\theta^* - \theta^\circ - \mathbf{b}_\lambda)$ in the following form:

$$\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ - \mathbf{b}_{\lambda} = \lambda \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right) \boldsymbol{\theta}^\circ - \lambda \left(\frac{1}{2}(A^*)^\top A^* + \lambda I_d\right)^{-1} \boldsymbol{\theta}^\circ$$
$$- \frac{1}{2} \left(\frac{1}{2}(A^*)^\top A^* + \lambda I_d\right)^{-1} (A^*)^\top (A^* - \Sigma) \boldsymbol{\theta}^\circ$$
$$= \lambda \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \left(I_d - (I_d + B)^{-1}\right) \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \boldsymbol{\theta}^\circ$$
$$- \frac{1}{2} \left(\frac{1}{2}(A^*)^\top A^* + \lambda I_d\right)^{-1} (A^*)^\top (A^* - \Sigma) \boldsymbol{\theta}^\circ.$$

The inequalities

$$\left\| \left(\frac{1}{2}\Sigma^2 + \lambda I_d\right)^{-1/2} \right\| \leqslant \frac{1}{\sqrt{\lambda}}, \quad \left\| \left(\frac{1}{2} (A^*)^\top A^* + \lambda I_d\right)^{-1} (A^*)^\top \right\| \leqslant \frac{1}{\sqrt{2\lambda}},$$

and

$$||I_d - (I_d + B)^{-1}|| = ||B(I_d + B)^{-1}|| \le \frac{||B||}{1 - ||B||}$$

yield that

$$\|\boldsymbol{\theta}^* - \boldsymbol{\theta}^\circ - \mathbf{b}_{\lambda}\| \leqslant \left(\frac{\|B\|}{1 - \|B\|} + \frac{\|A^* - \Sigma\|}{2\sqrt{2\lambda}}\right) \|\boldsymbol{\theta}^\circ\|$$

Similarly to the proof of Lemma B.4, we deduce that (see (89))

$$\|B\| \leqslant \frac{2\|A^* - \Sigma\|}{\sqrt{\lambda}} \leqslant 2\left(\frac{14\|\boldsymbol{\theta}^\circ\|}{\mu}\right)^2 + \frac{70\|\Sigma\|\|\boldsymbol{\theta}^\circ\|\|\mathbf{b}_\lambda\|}{\mu^2\sqrt{\lambda}} \leqslant \frac{1}{4}.$$

Then it holds that

$$\begin{split} \|\boldsymbol{\theta}^{*}-\boldsymbol{\theta}^{\circ}-\mathbf{b}_{\lambda}\| &\leqslant \left(\frac{4\|B\|}{3} + \frac{\|A^{*}-\boldsymbol{\Sigma}\|}{2\sqrt{2\lambda}}\right)\|\boldsymbol{\theta}^{\circ}\|\\ &\leqslant \left(\frac{8\|A^{*}-\boldsymbol{\Sigma}\|}{3\sqrt{\lambda}} + \frac{\|A^{*}-\boldsymbol{\Sigma}\|}{2\sqrt{2\lambda}}\right)\|\boldsymbol{\theta}^{\circ}\|\\ &\leqslant \frac{7\|\boldsymbol{\theta}^{\circ}\|}{2} \cdot \frac{\|A^{*}-\boldsymbol{\Sigma}\|}{\sqrt{\lambda}}\\ &\leqslant \frac{7\|\boldsymbol{\theta}^{\circ}\|}{2} \left(\left(\frac{14\|\boldsymbol{\theta}^{\circ}\|}{\mu}\right)^{2} + \frac{70\|\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^{\circ}\|\|\mathbf{b}_{\lambda}\|}{\mu^{2}\sqrt{\lambda}}\right) \end{split}$$

E Concentration inequalities

E.1 Proof of Theorem 2.5

Following the ideas of Zhivotovskiy [2024], we use the tools of PAC-Bayesian analysis to prove a large deviation inequality. The core of this approach is the following variational inequality (see, e. g., [Catoni and Giulini, 2017, Proposition 2.1]).

Lemma E.1. Let $\mathbf{X}, \mathbf{X}_1, \ldots, \mathbf{X}_n$ be i.i.d. random elements on a measurable space \mathcal{X} . Let Θ be a parameter space equipped with a measure μ (which is also referred to as prior). Let $f : \mathcal{X} \times \Theta \to \mathbb{R}$. Then, with probability at least $1 - \delta$, it holds that

$$\mathbb{E}_{\boldsymbol{\theta} \sim \rho} \frac{1}{n} \sum_{i=1}^{n} f(\mathbf{X}_{i}, \boldsymbol{\theta}) \leqslant \mathbb{E}_{\boldsymbol{\theta} \sim \rho} \log \mathbb{E}_{\mathbf{X}} e^{f(X, \boldsymbol{\theta})} + \frac{\mathcal{KL}(\rho, \mu) + \log(1/\delta)}{n}$$

simultaneously for all $\rho \ll \mu$.

In what follows, we show that a high probability upper bound on $||B(\widehat{\Sigma} - \Sigma)||_{\rm F}$ can be deduced from the PAC-Bayes variational inequality. We split the proof into several steps for convenience.

Step 0: auxiliary constants. Before we move to the proof, let us introduce auxiliary constants. We define

$$\beta = 2 \operatorname{r}(\Sigma^{1/2} A^{\top} A \Sigma^{1/2}) \operatorname{r}(\Sigma^{1/2} B^{\top} B \Sigma^{1/2}), \qquad (138)$$

$$R = \sqrt{\frac{2}{\beta} \operatorname{Tr}(\Sigma^{1/2} A^{\top} A \Sigma^{1/2}) \operatorname{Tr}(\Sigma^{1/2} B^{\top} B \Sigma^{1/2})},$$
(139)

and

$$\lambda = \frac{1}{C_X \sqrt{\|\Sigma^{1/2} A^{\top} A \Sigma^{1/2} \| \|\Sigma^{1/2} B^{\top} B \Sigma^{1/2} \|}} \cdot \sqrt{\frac{\mathbf{r}(\Sigma^{1/2} A^{\top} A \Sigma^{1/2}) \mathbf{r}(\Sigma^{1/2} B^{\top} B \Sigma^{1/2}) + \log(2/\delta)}{4n}},$$
(140)

where C_X is defined in Assumption 2.1. We move to the proof of Theorem 2.5.

Step 1: a variational inequality. On this step, we make some preparations. We specify a prior μ and a family of measures ρ from Lemma E.1. A reader will immediately see the result of these efforts on the second step where we relate the variational approach with a high probability upper bound on the Frobenius norm of $B(\hat{\Sigma} - \Sigma)$. Let us introduce $\boldsymbol{\xi} = \Sigma^{-1/2} \mathbf{X}$, $\boldsymbol{\xi}_i = \Sigma^{-1/2} \mathbf{X}_i$, $1 \leq i \leq n$, and a random matrix

$$\Psi = \frac{1}{\sqrt{\beta}} \Sigma^{1/2} B^{\top} Z A \Sigma^{1/2}.$$

where Z is a matrix of size $(q \times d)$ with i.i.d. $\mathcal{N}(0, 1)$ entries and β is defined in (138). Note that

$$\operatorname{vec}(\Psi) \sim \mathcal{N}\left(\mathbf{0}, \beta^{-1} \left(\Sigma^{1/2} A^{\top} A \Sigma^{1/2}\right) \otimes \left(\Sigma^{1/2} B^{\top} B \Sigma^{1/2}\right)\right)$$

in this case (see, for instance, [Leng and Pan, 2018] for the details). Denote its density (with respect to the volume measure on the image of $(\Sigma^{1/2}A^{\top}A\Sigma^{1/2}) \otimes (\Sigma^{1/2}B^{\top}B\Sigma^{1/2})$) by p and, for any $U \in \mathbb{R}^{d \times d}$, define

$$\mathsf{p}_U(\mathbf{x}) = \frac{1}{\pi_R} \, \mathsf{p}\big(\mathbf{x} - \mathbf{vec}(U)\big) \,\mathbb{1}(\|\mathbf{x} - \mathbf{vec}(U)\| \leqslant R),$$

where $\mathbf{vec}(U) \in \mathbb{R}^{d^2}$ is a vector obtained by reshaping of U and $\pi_R \in (0, 1)$ is a normalizing constant. Then $(\boldsymbol{\xi}_1^\top U \boldsymbol{\xi}_1 + \dots \boldsymbol{\xi}_n^\top U \boldsymbol{\xi}_n)/n - \mathbb{E} \boldsymbol{\xi}^\top U \boldsymbol{\xi}$ can be represented in the following form:

$$\frac{1}{n}\sum_{i=1}^{n}\boldsymbol{\xi}_{i}^{\top}U\boldsymbol{\xi}_{i} - \mathbb{E}\boldsymbol{\xi}^{\top}U\boldsymbol{\xi} = \mathbb{E}_{\mathbf{vec}(\Phi)\sim\mathbf{p}_{U}}\left(\frac{1}{n}\sum_{i=1}^{n}\boldsymbol{\xi}_{i}^{\top}\Phi\boldsymbol{\xi}_{i} - \mathbb{E}_{X}\boldsymbol{\xi}^{\top}\Phi\boldsymbol{\xi}\right).$$

According to Lemma E.1, for any $\delta \in (0,1)$ there exists an event \mathcal{E} , such that $\mathbb{P}(\mathcal{E}) \ge 1 - \delta$ and

$$\begin{split} \lambda \left(\frac{1}{n} \sum_{i=1}^{n} \boldsymbol{\xi}_{i}^{\top} U \boldsymbol{\xi}_{i} - \mathbb{E} \boldsymbol{\xi}^{\top} U \boldsymbol{\xi} \right) &= \lambda \mathbb{E}_{\mathbf{vec}(\Phi) \sim \mathbf{p}_{U}} \left(\frac{1}{n} \sum_{i=1}^{n} \boldsymbol{\xi}_{i}^{\top} \Phi \boldsymbol{\xi}_{i} - \mathbb{E}_{X} \boldsymbol{\xi}^{\top} \Phi \boldsymbol{\xi} \right) \\ &\leqslant -\lambda \mathbb{E}_{\mathbf{vec}(\Phi) \sim \mathbf{p}_{U}} \mathbb{E}_{\mathbf{X}} \boldsymbol{\xi}^{\top} \Phi \boldsymbol{\xi} + \mathbb{E}_{\mathbf{vec}(\Phi) \sim \mathbf{p}_{U}} \log \mathbb{E}_{\mathbf{X}} e^{\lambda \boldsymbol{\xi}^{\top} \Phi \boldsymbol{\xi}} \\ &+ \frac{\mathcal{K} \mathcal{L}(\mathbf{p}_{U}, \mathbf{p}) + \log(1/\delta)}{n} \end{split}$$

simultaneously for all $U \in \mathbb{R}^{d \times d}$ on \mathcal{E} . From now on, we restrict our attention on this event.

Step 2: relating the variational inequality with the Frobenius norm. Rewrite the Frobenius norm of $B(\hat{\Sigma} - \Sigma)A^{\top}$ as follows:

$$\begin{split} \|B(\widehat{\Sigma} - \Sigma)A^{\top}\|_{\mathrm{F}} &= \sup_{\substack{V \in \mathbb{R}^{d \times q}, \\ \|V\|_{\mathrm{F}} = 1}} \operatorname{Tr} \left(VB(\widehat{\Sigma} - \Sigma)A^{\top} \right) = \sup_{\substack{V \in \mathbb{R}^{d \times q}, \\ \|V\|_{\mathrm{F}} = 1}} \left(\frac{1}{n} \sum_{i=1}^{n} \mathbf{X}_{i}^{\top} A^{\top} VB \mathbf{X}_{i} - \mathbb{E} \mathbf{X}^{\top} A^{\top} VB \mathbf{X} \right) \\ &= \sup_{\substack{U = \Sigma^{1/2} A^{\top} VB \Sigma^{1/2}, \\ \|V\|_{\mathrm{F}} = 1}} \left(\frac{1}{n} \sum_{i=1}^{n} \boldsymbol{\xi}_{i}^{\top} U \boldsymbol{\xi}_{i} - \mathbb{E} \boldsymbol{\xi}^{\top} U \boldsymbol{\xi} \right). \end{split}$$

On the event \mathcal{E} , defined on Step 1, it holds that

$$\lambda \| B(\widehat{\Sigma} - \Sigma) A^{\top} \|_{\mathrm{F}} \leq \sup_{\substack{U = \Sigma^{1/2} V B \Sigma^{1/2}, \\ \|V\|_{\mathrm{F}} = 1}} \left(-\lambda \mathbb{E}_{\mathbf{vec}(\Phi) \sim \mathsf{p}_{U}} \mathbb{E}_{\mathbf{X}} \boldsymbol{\xi}^{\top} \Phi \boldsymbol{\xi} + \mathbb{E}_{\mathbf{vec}(\Phi) \sim \mathsf{p}_{U}} \log \mathbb{E}_{\mathbf{X}} e^{\lambda \boldsymbol{\xi}^{\top} \Phi \boldsymbol{\xi}} + \frac{\mathcal{KL}(\mathsf{p}_{U}, \mathsf{p}) + \log(1/\delta)}{n} \right).$$
(141)

It only remains to bound the terms in the right hand side.

Step 3: a bound on $\mathcal{KL}(p_U, p)$. Applying Lemma E.2 with $P = \Sigma^{1/2} A^{\top}$, $Q = \Sigma^{1/2} B^{\top}$, and $U = \Sigma^{1/2} A^{\top} V B \Sigma^{1/2}$, $\|V\|_{\rm F} = 1$, we obtain that

$$\mathcal{KL}(\mathbf{p}_{U},\mathbf{p}) \leq \log 2 + \frac{\beta}{2} \left\| (\Sigma^{1/2}A^{\top})^{\dagger} U(B\Sigma^{1/2})^{\dagger} \right\|_{\mathrm{F}}^{2} \leq \log 2 + \frac{\beta}{2} \|V\|_{\mathrm{F}}^{2} = \log 2 + \frac{\beta}{2}.$$
(142)

Step 4: a bound on $\mathbb{E}_{\mathbf{vec}(\Phi)\sim \mathsf{p}_U} \log \mathbb{E}_{\mathbf{X}} e^{\lambda \boldsymbol{\xi}^\top \Phi \boldsymbol{\xi}}$. Note that, since $U = \Sigma^{1/2} A^\top V B \Sigma^{1/2}$, $\|V\|_{\mathrm{F}} = 1$, the following inequality holds almost surely:

$$\begin{split} \|\Phi\|_{\mathrm{F}}^{2} &\leqslant 2\|U\|_{\mathrm{F}}^{2} + 2R^{2} \leqslant 2\|\Sigma^{1/2}A^{\top}\|^{2}\|V\|_{\mathrm{F}}^{2}\|B\Sigma^{1/2}\|^{2} + 2R^{2} \\ &= 2\|\Sigma^{1/2}A^{\top}A\Sigma^{1/2}\|\|\Sigma^{1/2}B^{\top}B\Sigma^{1/2}\| + 2R^{2}. \end{split}$$

By the definition of R (see (139) and (138)), we have

$$\|\Phi\|_{\mathrm{F}}^{2} \leqslant 4 \|\Sigma^{1/2} A^{\top} A \Sigma^{1/2} \| \|\Sigma^{1/2} B^{\top} B \Sigma^{1/2} \|$$
 almost surely.

Note that this inequality yields that λ , defined in (140), does not exceed $1/(C_X \|\Phi\|_F)$. Hence, due to Assumption 2.1, it holds that

$$-\lambda \mathbb{E}_{\boldsymbol{\xi}}(\boldsymbol{\xi}^{\top} \Phi \boldsymbol{\xi}) + \log \mathbb{E}_{\boldsymbol{\xi}} e^{\lambda \boldsymbol{\xi}^{\top} \Phi \boldsymbol{\xi}} \leqslant C_X^2 \lambda^2 \|\Phi\|_{\mathrm{F}}^2 \leqslant 4C_X^2 \lambda^2 \|\Sigma^{1/2} A^{\top} A \Sigma^{1/2}\| \|\Sigma^{1/2} B^{\top} B \Sigma^{1/2}\|.$$
(143)

Summing up the inequalitites (141), (142), and (143), we obtain that

$$\lambda \| B(\widehat{\Sigma} - \Sigma) A^\top \|_{\mathcal{F}} \leqslant 4C_X^2 \lambda^2 \| \Sigma^{1/2} A^\top A \Sigma^{1/2} \| \| \Sigma^{1/2} B^\top B \Sigma^{1/2} \| + \frac{\beta/2 + \log(2/\delta)}{n}$$

on the event \mathcal{E} . Dividing both parts by λ and taking into account (138) and (140), we conclude that

$$\|B(\widehat{\Sigma} - \Sigma)A^{\top}\|_{\mathrm{F}} \leq 4C_X \|\Sigma^{1/2}A^{\top}A\Sigma^{1/2}\| \|\Sigma^{1/2}B^{\top}B\Sigma^{1/2}\|$$
$$\cdot \sqrt{\frac{\mathbf{r}(\Sigma^{1/2}A^{\top}A\Sigma^{1/2})\mathbf{r}(\Sigma^{1/2}B^{\top}B\Sigma^{1/2}) + \log(2/\delta)}{n}}$$

with probability at least $1 - \delta$.

E.2 Proof of Theorem 2.6

The proof of this statement is also based on the PAC-Bayesian approach. Note that

$$\left\|\frac{1}{n}\sum_{i=1}^{n}B\mathbf{X}_{i}\varepsilon_{i}\right\| = \sup_{\|\mathbf{v}\|=1}\frac{1}{n}\sum_{i=1}^{n}\mathbf{v}^{\top}B\mathbf{X}_{i}\varepsilon_{i} = \sup_{\substack{\mathbf{u}=\sum^{1/2}B^{\top}\mathbf{v},\\\|\mathbf{v}\|=1}}\frac{1}{n}\sum_{i=1}^{n}\mathbf{u}^{\top}\Sigma^{-1/2}\mathbf{X}_{i}\varepsilon_{i}.$$

Let μ be the probability measure on \mathbb{R}^d corresponding to the Gaussian distribution

$$\mathcal{N}\left(\mathbf{0}, \frac{1}{2\mathbf{r}(\Sigma^{1/2}B^{\top}B\Sigma^{1/2})}\Sigma^{1/2}B^{\top}B\Sigma^{1/2}\right).$$

For any u, such that $\mathbf{u} = \Sigma^{1/2} B^{\top} \mathbf{v}$ for some unit vector v, define a probability measure $\rho_{\mathbf{u}}$ by the formula

$$\mathrm{d}\rho_{\mathbf{u}}(\mathbf{x}) = \frac{1}{\pi} \mathbb{1}\left(\|\mathbf{x} - \mathbf{u}\| \leqslant \|B\Sigma^{1/2}\| \right) \mathrm{d}\mu(\mathbf{x} - \mathbf{u}),$$

where

$$\pi = \mu\left(\left\{\mathbf{x} : \|\mathbf{x} - \mathbf{u}\| \leqslant \|B\Sigma^{1/2}\|\right\}\right)$$

is a normalizing constant. Then it holds that

$$\sup_{\substack{\mathbf{u}=\Sigma^{1/2}B^{\top}\mathbf{v},\\\|\mathbf{v}\|=1}}\frac{1}{n}\sum_{i=1}^{n}\mathbf{u}^{\top}\Sigma^{-1/2}\mathbf{X}_{i}\varepsilon_{i} = \sup_{\substack{\mathbf{u}=\Sigma^{1/2}B^{\top}\mathbf{v},\\\|\mathbf{v}\|=1}}\frac{1}{n}\mathbb{E}_{\boldsymbol{\xi}\sim\rho_{\mathbf{u}}}\sum_{i=1}^{n}\boldsymbol{\xi}^{\top}\Sigma^{-1/2}\mathbf{X}_{i}\varepsilon_{i}.$$

Let us fix

$$\lambda = \frac{1}{4\sigma} \sqrt{\frac{\mathbf{r}(\Sigma^{1/2} B^{\top} B \Sigma^{1/2}) + \log(2/\delta)}{\|\Sigma\| n}}.$$

.

According to Lemma E.1, for any $\delta \in (0, 1)$, with probability at least $(1 - \delta)$, we have

$$\lambda \sup_{\substack{\mathbf{u}=\sum^{1/2}B^{\top}\mathbf{v},\\ \|\mathbf{v}\|=1}} \frac{1}{n} \mathbb{E}_{\boldsymbol{\xi}\sim\rho_{\mathbf{u}}} \sum_{i=1}^{n} \boldsymbol{\xi}^{\top} \Sigma^{-1/2} \mathbf{X}_{i} \varepsilon_{i} \leqslant \mathbb{E}_{\boldsymbol{\xi}\sim\rho_{\mathbf{u}}} \log \mathbb{E}_{\mathbf{X},\varepsilon} e^{\lambda \boldsymbol{\xi}^{\top} \Sigma^{-1/2} \mathbf{X}\varepsilon} + \frac{\mathcal{K}\mathcal{L}(\rho_{\mathbf{u}},\mu) + \log(1/\delta)}{n}$$

simultaneously for all $\rho_{\mathbf{u}}$, such that $\mathbf{u} = \Sigma^{1/2} B^{\top} \mathbf{v}$ for some unit vector \mathbf{v} . Applying Lemma E.2 with $P = \Sigma^{1/2} B^{\top}$, Q = 1, and $\beta = 2\mathbf{r}(\Sigma^{1/2} B^{\top} B \Sigma^{1/2})$, we obtain that

$$\mathcal{KL}(\rho_{\mathbf{u}},\mu) \leq \log 2 + \mathbf{r}(\Sigma^{1/2}B^{\top}B\Sigma^{1/2}) \left\| (\Sigma^{1/2}B^{\top})^{\dagger}\mathbf{u} \right\|^{2} \leq \log 2 + \mathbf{r}(\Sigma^{1/2}B^{\top}B\Sigma^{1/2}).$$

Besides, since $\|\Sigma^{-1/2} \mathbf{X}_1 \varepsilon_1\|_{\psi_1} = \sigma$ and

$$\lambda = \frac{1}{4\sigma} \sqrt{\frac{\mathbf{r}(\Sigma^{1/2} B^\top B \Sigma^{1/2}) + \log(2/\delta)}{\|\Sigma\| n}} \leqslant \frac{1}{4\sigma \sqrt{\|\Sigma^{1/2} B^\top B \Sigma^{1/2}\|}} \leqslant \frac{1}{2\sigma \|\xi\|},$$

Lemma 2 from Zhivotovskiy [2024] yields that

$$\log \mathbb{E}_{\mathbf{X},\varepsilon} e^{\lambda \boldsymbol{\xi}^{\top} \boldsymbol{\Sigma}^{-1/2} \mathbf{X}\varepsilon} \leq 4\lambda^2 \sigma^2 \|\boldsymbol{\xi}\|^2 \leq 8\lambda^2 \sigma^2 \left(\|\mathbf{u}\|^2 + \|B\boldsymbol{\Sigma}^{1/2}\|^2 \right) \leq 16\lambda^2 \sigma^2 \|B\boldsymbol{\Sigma}^{1/2}\|^2.$$

Here we used the fact that $\|\boldsymbol{\xi}\|^2 \leq 2\|\mathbf{u}\|^2 + 2\|B\Sigma^{1/2}\|^2 \leq 4\|B\Sigma^{1/2}\|^2$ almost surely. Hence, with probability at least $(1 - \delta)$, it holds that

$$\begin{split} \|B(\mathbf{Z} - \mathbb{E}\mathbf{Z})\| &= \frac{1}{n} \left\| \sum_{i=1}^{n} B\mathbf{X}_{i} \varepsilon_{i} \right\| \leq 16\lambda \sigma^{2} \|\Sigma^{1/2} B^{\top} B\Sigma^{1/2}\| + \frac{\mathbf{r}(\Sigma^{1/2} B^{\top} B\Sigma^{1/2}) + \log(2/\delta)}{\lambda n} \\ &= 8\sigma \left\| B\Sigma^{1/2} \right\| \sqrt{\frac{\mathbf{r}(\Sigma^{1/2} B^{\top} B\Sigma^{1/2}) + \log(2/\delta)}{n}}. \end{split}$$

E.3 Auxiliary results

Lemma E.2. Let β be a positive number and let $P \in \mathbb{R}^{p \times r}$ and $Q \in \mathbb{R}^{q \times s}$ be arbitrary matrices. Let p be the probability density on $\operatorname{Im}((QQ^{\top}) \otimes (PP^{\top}))$ (with respect to the volume measure \mathcal{V}) of the Gaussian distribution $\mathcal{N}(0, \beta^{-1}(QQ^{\top}) \otimes (PP^{\top}))$, where \otimes stands for the Kronecker product. That is,

$$\mathbf{p}(\mathbf{x}) \propto \exp\left\{-\frac{\beta}{2}\mathbf{x}^{\top} \left((QQ^{\top}) \otimes (PP^{\top})\right)^{\dagger} \mathbf{x}\right\}, \quad \mathbf{x} \in \operatorname{Im}((QQ^{\top}) \otimes (PP^{\top})).$$

Fix a matrix $U \in \mathbb{R}^{p \times q}$, such that $\mathbf{vec}(U) \in \mathrm{Im}((QQ^{\top}) \otimes (PP^{\top}))$, and set

$$R = \sqrt{\frac{2}{\beta}} \operatorname{Tr}(PP^{\top}) \operatorname{Tr}(QQ^{\top}).$$

Define the density p_U *as follows:*

$$\mathsf{p}_U(\mathbf{x}) = \frac{1}{\pi_R} \mathsf{p}(\mathbf{x} - \mathbf{vec}(U)) \mathbb{1}(\|\mathbf{x} - \mathbf{vec}(U)\| \leq R), \quad \mathbf{x} \in \mathrm{Im}((QQ^{\top}) \otimes (PP^{\top})),$$

where

$$\pi_R = \int_{\|\mathbf{x}\| \leq R} \mathbf{p}(\mathbf{x}) \mathrm{d}\mathcal{V}(\mathbf{x}) = \mathbb{P}_{\boldsymbol{\xi} \sim \mathbf{p}} \big(\|\boldsymbol{\xi}\| \leq R \big)$$

is a normalizing constant. Then it holds that

$$\mathcal{KL}(\mathbf{p}_U, \mathbf{p}) = \log \frac{1}{\pi_R} + \frac{\beta}{2} \|P^{\dagger} U(Q^{\dagger})^{\top}\|_{\mathbf{F}}^2 \leqslant \log 2 + \frac{\beta}{2} \|P^{\dagger} U(Q^{\dagger})^{\top}\|_{\mathbf{F}}^2$$

Proof. Denote $S = (QQ^{\top}) \otimes (PP^{\top})$ and $\mathbf{u} = \mathbf{vec}(U)$ for brevity. By the definition of $\mathcal{KL}(\mathbf{p}_U, \mathbf{p})$, it holds that

$$\begin{split} \mathcal{KL}(\mathsf{p}_U,\mathsf{p}) &= \int \log \frac{\mathsf{p}_U(\mathbf{x})}{\mathsf{p}(\mathbf{x})} \, \mathsf{p}_U(\mathbf{x}) \mathrm{d}\mathcal{V}(\mathbf{x}) \\ &= \int \log \frac{1}{\pi_R} \, \mathsf{p}_U(\mathbf{x}) \mathrm{d}\mathcal{V}(\mathbf{x}) + \frac{\beta}{2} \int\limits_{\|\mathbf{x}-\mathbf{u}\| \leqslant R} \left[\mathbf{x}^\top S^\dagger \mathbf{x} - (\mathbf{x}-\mathbf{u})^\top S^\dagger (\mathbf{x}-\mathbf{u}) \right] \mathrm{d}\mathcal{V}(\mathbf{x}) \\ &= \log \frac{1}{\pi_R} + \frac{\beta}{2} \int\limits_{\|\mathbf{x}-\mathbf{u}\| \leqslant R} \left[2\mathbf{x}^\top S^\dagger \mathbf{u} - \mathbf{u}^\top S^\dagger \mathbf{u} \right] \mathrm{d}\mathcal{V}(\mathbf{x}) \\ &= \log \frac{1}{\pi_R} + \frac{\beta}{2} \left(2\mathbf{u}^\top S^\dagger \mathbf{u} - \mathbf{u}^\top S^\dagger \mathbf{u} \right) = \log \frac{1}{\pi_R} + \frac{\beta}{2} \mathbf{u}^\top S^\dagger \mathbf{u}. \end{split}$$

Let us recall the following properties of the Kronecker product:

$$((QQ^{\top}) \otimes (PP^{\top}))^{\dagger} = (QQ^{\top})^{\dagger} \otimes (PP^{\top})^{\dagger}, ((QQ^{\top})^{\dagger} \otimes (PP^{\top})^{\dagger}) \mathbf{vec}(U) = \mathbf{vec} ((PP^{\top})^{\dagger} U (QQ^{\top})^{\dagger}).$$

Using these two inequalities, we obtain that

$$\mathbf{u}^{\top} S^{\dagger} \mathbf{u} = \mathbf{vec}(U)^{\top} \mathbf{vec} \left((PP^{\top})^{\dagger} U(QQ^{\top})^{\dagger} \right) = \mathrm{Tr} \left(U^{\top} (PP^{\top})^{\dagger} U(QQ^{\top})^{\dagger} \right)$$
$$= \mathrm{Tr} \left(U^{\top} (P^{\dagger})^{\top} P^{\dagger} U(Q^{\dagger})^{\top} Q^{\dagger} \right) = \mathrm{Tr} \left(Q^{\dagger} U^{\top} (P^{\dagger})^{\top} P^{\dagger} U(Q^{\dagger})^{\top} \right)$$
$$= \| P^{\dagger} U(Q^{\dagger})^{\top} \|_{\mathrm{F}}^{2}.$$

Hence, it holds that

$$\mathcal{KL}(\rho_U,\mu) = \log \frac{1}{\pi_R} + \frac{\beta}{2} \|P^{\dagger} U(Q^{\dagger})^{\top}\|_{\mathrm{F}}^2.$$

To prove the second part of the lemma, note that

$$\pi_R = \mathbb{P}(\|\boldsymbol{\xi}\| \leq R), \quad \text{where } \boldsymbol{\xi} \sim \mathcal{N}(0, \beta^{-1} (QQ^{\top}) \otimes (PP^{\top})).$$

Using Chebyshev's inequality, we obtain that

$$\begin{aligned} \pi_R &\ge 1 - \frac{\mathbb{E} \|\boldsymbol{\xi}\|^2}{R^2} = 1 - \frac{\mathbb{E} \mathrm{Tr}(\boldsymbol{\xi}\boldsymbol{\xi}^\top)}{R^2} = 1 - \frac{\mathrm{Tr}((QQ^\top) \otimes (PP^\top))}{\beta R^2} \\ &= 1 - \frac{\mathrm{Tr}((QQ^\top))\mathrm{Tr}((PP^\top))}{\beta R^2} \ge \frac{1}{2}, \end{aligned}$$

and thus,

$$\mathcal{KL}(\mathsf{p}_U,\mathsf{p}) = \log \frac{1}{\pi_R} + \frac{\beta}{2} \|P^{\dagger} U(Q^{\dagger})^{\top}\|_{\mathrm{F}}^2 \leqslant \log 2 + \frac{\beta}{2} \|P^{\dagger} U(Q^{\dagger})^{\top}\|_{\mathrm{F}}^2.$$

F Proof of Corollary 2.7

According to Theorem 2.3 and 2.4, with probability at least $(1 - 13\delta/2)$, it holds that

$$\left\|\Sigma^{1/2}(\widehat{\boldsymbol{\theta}}-\boldsymbol{\theta}^{\circ})\right\| \leqslant \left\|\Sigma^{1/2}\mathbf{b}_{\lambda}\right\| + \left\|\Sigma^{1/2}(\Sigma^{2}+2\lambda I_{d})^{-1}\boldsymbol{\zeta}\right\| + \diamondsuit,$$
(144)

where $\boldsymbol{\zeta} = \Sigma \mathbf{U} - \Sigma (\widehat{\Sigma} - \Sigma) \mathbf{b}_{\lambda} - (\widehat{\Sigma} - \Sigma) \Sigma \mathbf{b}_{\lambda}$ and \diamondsuit defined in (15). In the rest of the proof, we show that the norm of $\Sigma^{1/2} (\Sigma^2 + 2\lambda I_d)^{-1} \boldsymbol{\zeta}$ does not exceed

$$4\left(2\sigma + C_X \|\Sigma^{1/2}\mathbf{b}_{\lambda}\|\right) \sqrt{\frac{k^* + r_4(k^*) + \log(4/\delta)}{n}} + \frac{2C_X \|\Sigma^{3/2}\mathbf{b}_{\lambda}\|}{\sqrt{2\lambda}} \sqrt{\frac{4k^* + 4r_2(k^*) + \log(4/\delta)}{n}}.$$

In order to do so, we use the triangle inequality

$$\left\| \Sigma^{1/2} (\Sigma^{2} + 2\lambda I_{d})^{-1} \boldsymbol{\zeta} \right\| \leq \left\| \Sigma^{1/2} (\Sigma^{2} + 2\lambda I_{d})^{-1} \Sigma \mathbf{U} \right\| + \left\| \Sigma^{1/2} (\Sigma^{2} + 2\lambda I_{d})^{-1} \Sigma (\widehat{\Sigma} - \Sigma) \mathbf{b}_{\lambda} \right\| + \left\| \Sigma^{1/2} (\Sigma^{2} + 2\lambda I_{d})^{-1} (\widehat{\Sigma} - \Sigma) \Sigma \mathbf{b}_{\lambda} \right\|$$
(145)

and bound the three terms in the right-hand side one by one using Theorems 2.5 and 2.6. Before we proceed, we formulate an auxiliary result.

Lemma F.1. Let $r_q(k)$ and $k^* = k^*(\lambda)$ be as defined in (11) and (12), respectively. Then it holds that

$$\mathbf{r}\left(\Sigma^4(\Sigma^2+2\lambda I_d)^{-2}\right) \leqslant \left(1+\frac{2\lambda}{\|\Sigma\|^2}\right)^2 \left(k^*+r_4(k^*)\right)$$

and

$$r\left(\Sigma^2(\Sigma^2+2\lambda I_d)^{-2}\right) \leqslant 4k^*+4r_2(k^*).$$

The proof of Lemma F.1 is postponed to Section F.1. It helps us to verify the conditions of Theorems 2.5 and 2.6. Indeed, taking into account that $r_4(k) \leq r_2(k)$ for all $k \in \mathbb{N}$, we observe that the condition (13) and Lemma F.1 yield

$$\mathbf{r}\left(\Sigma^4(\Sigma^2 + 2\lambda I_d)^{-2}\right) + \log(4/\delta) \leqslant \left(1 + \frac{2\lambda}{\|\Sigma\|^2}\right)^2 \left(k^* + r_4(k^*)\right) + \log(4/\delta) \leqslant n$$

Then, according to Theorems 2.5 and 2.6, with probability at least $(1 - \delta)$, we simultaneously have

$$\begin{split} & \left\| \Sigma^{1/2} (\Sigma^2 + 2\lambda I_d)^{-1} \Sigma (\widehat{\Sigma} - \Sigma) \mathbf{b}_\lambda \right\| \\ & \leqslant 4C_X \left\| \Sigma^2 (\Sigma^2 + 2\lambda I_d)^{-1} \right\| \left\| \Sigma^{1/2} \mathbf{b}_\lambda \right\| \sqrt{\frac{\mathbf{r} (\Sigma^4 (\Sigma^2 + 2\lambda I_d)^{-2}) + \log(4/\delta)}{n}} \\ & = \frac{4C_X \|\Sigma\|^2}{\|\Sigma\|^2 + 2\lambda} \left\| \Sigma^{1/2} \mathbf{b}_\lambda \right\| \sqrt{\frac{\mathbf{r} (\Sigma^4 (\Sigma^2 + 2\lambda I_d)^{-2}) + \log(4/\delta)}{n}} \end{split}$$

and

$$\begin{split} \left\| \Sigma^{1/2} (\Sigma^2 + 2\lambda I_d)^{-1} \Sigma \mathbf{U} \right\| &\leq 8\sigma \left\| \Sigma^2 (\Sigma^2 + 2\lambda I_d)^{-1} \right\| \sqrt{\frac{\mathbf{r} (\Sigma^4 (\Sigma^2 + 2\lambda I_d)^{-2}) + \log(4/\delta)}{n}} \\ &= \frac{8\sigma \|\Sigma\|^2}{\|\Sigma\|^2 + 2\lambda} \sqrt{\frac{\mathbf{r} (\Sigma^4 (\Sigma^2 + 2\lambda I_d)^{-2}) + \log(4/\delta)}{n}}. \end{split}$$

In view of Lemma F.1, we obtain that, on the same event, it holds that

$$\left\|\Sigma^{1/2}(\Sigma^2 + 2\lambda I_d)^{-1}\Sigma(\widehat{\Sigma} - \Sigma)\mathbf{b}_\lambda\right\| \leqslant 4C_X \|\Sigma^{1/2}\mathbf{b}_\lambda\| \sqrt{\frac{k^* + r_4(k^*) + \log(4/\delta)}{n}}$$
(146)

and

$$\left\|\Sigma^{1/2}(\Sigma^2 + 2\lambda I_d)^{-1}\Sigma \mathbf{U}\right\| \leqslant 8\sigma \sqrt{\frac{k^* + r_4(k^*) + \log(4/\delta)}{n}}.$$
(147)

It remains to bound the norm of $\Sigma^{1/2}(\Sigma^2 + 2\lambda I_d)^{-1}(\widehat{\Sigma} - \Sigma)\Sigma \mathbf{b}_{\lambda}$ to finish the proof. Note that Lemma F.1 and (13) imply that

$$\mathbf{r}\left(\Sigma^{2}(\Sigma^{2}+2\lambda I_{d})^{-2}\right) + \log(4/\delta) \leqslant 4k^{*} + 4r_{2}(k^{*}) + \log(4/\delta) \leqslant 4n.$$

Then, due to Theorem 2.5, with probability at least $(1 - \delta/2)$, we have

$$\begin{aligned} \left\| \Sigma^{1/2} (\Sigma^{2} + 2\lambda I_{d})^{-1} (\widehat{\Sigma} - \Sigma) \Sigma \mathbf{b}_{\lambda} \right\| \\ &\leq 4C_{X} \left\| \Sigma (\Sigma^{2} + 2\lambda I_{d})^{-1} \right\| \left\| \Sigma^{3/2} \mathbf{b}_{\lambda} \right\| \sqrt{\frac{\mathbf{r} (\Sigma^{2} (\Sigma^{2} + 2\lambda I_{d})^{-2}) + \log(4/\delta)}{n}} \\ &\leq \frac{2C_{X} \left\| \Sigma^{3/2} \mathbf{b}_{\lambda} \right\|}{\sqrt{2\lambda}} \sqrt{\frac{4k^{*} + 4r_{2}(k^{*}) + \log(4/\delta)}{n}}. \end{aligned}$$
(148)

Here we used the fact that

$$\left\|\Sigma(\Sigma^2 + 2\lambda I_d)^{-1}\right\| = \max_{1 \le j \le d} \frac{\sigma_j}{\sigma_j^2 + 2\lambda} \le \frac{1}{2\sqrt{2\lambda}}.$$

Summing up the inequalities (145)–(148) and using the union bound, we obtain that, with probability at least $(1 - 3\delta/2)$,

$$\left\| \Sigma^{1/2} (\Sigma^{2} + 2\lambda I_{d})^{-1} \boldsymbol{\zeta} \right\| \leq 4 \left(2\sigma + C_{X} \| \Sigma^{1/2} \mathbf{b}_{\lambda} \| \right) \sqrt{\frac{k^{*} + r_{4}(k^{*}) + \log(4/\delta)}{n}} + \frac{2C_{X} \| \Sigma^{3/2} \mathbf{b}_{\lambda} \|}{\sqrt{2\lambda}} \sqrt{\frac{4k^{*} + 4r_{2}(k^{*}) + \log(4/\delta)}{n}}$$
(149)

Finally, (144) and (149) yield that

$$\begin{split} \left\| \Sigma^{1/2}(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^{\circ}) \right\| &\leq \left\| \Sigma^{1/2} \mathbf{b}_{\lambda} \right\| + 4 \left(2\sigma + C_X \| \Sigma^{1/2} \mathbf{b}_{\lambda} \| \right) \sqrt{\frac{k^* + r_4(k^*) + \log(4/\delta)}{n}} \\ &+ \frac{2C_X \| \Sigma^{3/2} \mathbf{b}_{\lambda} \|}{\sqrt{2\lambda}} \sqrt{\frac{4k^* + 4r_2(k^*) + \log(4/\delta)}{n}} + \diamondsuit \end{split}$$

with probability at least $(1 - 8\delta)$.

- 6		
-1		

F.1 Proof of Lemma F.1

Due to the definition of the effective rank, we have

$$\mathbf{r}\left(\Sigma^4(\Sigma^2 + 2\lambda I_d)^{-2}\right) = \frac{\mathrm{Tr}(\Sigma^4(\Sigma^2 + 2\lambda I_d)^{-2})}{\|\Sigma^4(\Sigma^2 + 2\lambda I_d)^{-2}\|} = \left(\frac{\|\Sigma\|^2 + 2\lambda}{\|\Sigma\|^2}\right)^2 \sum_{j=1}^d \frac{\sigma_j^4}{(\sigma_j^2 + 2\lambda)^2}.$$

Taking into account that the operator norm of $\Sigma^4 (\Sigma^2 + 2\lambda I_d)^{-2}$ is equal to $\|\Sigma\|^4 / (\|\Sigma\|^2 + 2\lambda)^2$, we obtain that

$$\mathbf{r} \left(\Sigma^4 (\Sigma^2 + 2\lambda I_d)^{-2} \right) = \left(\frac{\|\Sigma\|^2 + 2\lambda}{\|\Sigma\|^2} \right)^2 \left(\sum_{j=1}^{k^*} \frac{\sigma_j^4}{(\sigma_j^2 + 2\lambda)^2} + \sum_{j>k^*} \frac{\sigma_j^4}{(\sigma_j^2 + 2\lambda)^2} \right)$$

$$\leq \left(\frac{\|\Sigma\|^2 + 2\lambda}{\|\Sigma\|^2} \right)^2 \left(\sum_{j=1}^{k^*} 1 + \sum_{j>k^*} \frac{\sigma_j^4}{\sigma_{k^*+1}^4} \right)$$

$$= \left(1 + \frac{2\lambda}{\|\Sigma\|^2} \right)^2 (k^* + r_4(k^*)) .$$

Similarly, it holds that

$$\begin{split} \mathbf{r} \left(\Sigma^2 (\Sigma^2 + 2\lambda I_d)^{-2} \right) &= \left(\max_{1 \leqslant k \leqslant d} \frac{\sigma_k^2}{(\sigma_k^2 + 2\lambda)^2} \right)^{-1} \sum_{j=1}^d \frac{\sigma_j^2}{(\sigma_j^2 + 2\lambda)^2} \\ &= \min_{1 \leqslant k \leqslant d} \frac{(\sigma_k^2 + 2\lambda)^2}{\sigma_k^2} \left(\sum_{j=1}^{k^*} \frac{\sigma_j^2}{(\sigma_j^2 + 2\lambda)^2} + \sum_{j > k^*} \frac{\sigma_j^2}{(\sigma_j^2 + 2\lambda)^2} \right) \\ &\leqslant \frac{(\sigma_{k^*}^2 + 2\lambda)^2}{\sigma_{k^*}^2} \sum_{j=1}^{k^*} 1 + \frac{(\sigma_{k^*+1}^2 + 2\lambda)^2}{\sigma_{k^*+1}^2} \sum_{j > k^*} \frac{\sigma_j^2}{(\sigma_j^2 + 2\lambda)^2}. \end{split}$$

Since the definition of k^* (see (12)) yields that

$$\frac{(\sigma_{k^*}^2 + 2\lambda)^2}{\sigma_{k^*}^2} \leqslant \frac{(\sigma_{k^*}^2 + \sigma_{k^*}^2)^2}{\sigma_{k^*}^2} = 4\sigma_{k^*}^2$$

and

$$\frac{(\sigma_{k^*+1}^2 + 2\lambda)^2}{\sigma_{k^*+1}^2} \sum_{j > k^*} \frac{\sigma_j^2}{(\sigma_j^2 + 2\lambda)^2} \leqslant \frac{(\sigma_{k^*+1}^2 + 2\lambda)^2}{4\lambda^2} \sum_{j > k^*} \frac{\sigma_j^2}{\sigma_{k^*+1}^2} \leqslant \frac{(2\lambda + 2\lambda)^2}{4\lambda^2} \sum_{j > k^*} \frac{\sigma_j^2}{\sigma_{k^*+1}^2} = 4r_2(k^*),$$

we get the desired bound:

$$r\left(\Sigma^2(\Sigma^2+2\lambda I_d)^{-2}\right) \leqslant 4k^*+4r_2(k^*).$$

G Properties of the bias \mathbf{b}_{λ}

In this section, we provide rigorous proofs of some properties of $\mathbf{b}_{\lambda} = -2\lambda(\Sigma^2 + 2\lambda I_d)^{-1}\boldsymbol{\theta}^{\circ}$ mentioned in Section 2. Let us introduce auxiliary notation to simplify further derivations. Throughout this section, $\sigma_1 \ge \sigma_2 \ge \ldots \ge \sigma_d$ stand for the eigenvalues of Σ and

$$\Sigma = \sum_{j=1}^d \sigma_j \mathbf{v}_j \mathbf{v}_j^{\mathsf{T}}$$

is its eigenvalue decomposition. For any $k \in \{1, ..., d\}$, we define $\beta_k = \mathbf{v}_k^\top \boldsymbol{\theta}^\circ$ and the projections of $\boldsymbol{\theta}$ onto the linear span of $\mathbf{v}_1, ..., \mathbf{v}_k$ and its orthogonal complement, respectively:

$$oldsymbol{ heta}_{\leqslant k}^{\circ} = \sum_{j=1}^k eta_j \mathbf{v}_j \quad ext{and} \quad oldsymbol{ heta}_{>k}^{\circ} = oldsymbol{ heta}^{\circ} - oldsymbol{ heta}_{\leqslant k}^{\circ}.$$

Similarly to the statement of Corollary 2.7, we denote

$$k^* = k^*(\lambda) = \max\left\{k \in \mathbb{N} : \sigma_k^2 \ge 2\lambda\right\}$$

The first result relates the squared norms of $\Sigma^{1/2} \mathbf{b}_{\lambda}$ and $\Sigma^{3/2} \mathbf{b}_{\lambda}$ with the ones of $\boldsymbol{\theta}_{\leq k^*}^{\circ}$ and $\boldsymbol{\theta}_{>k^*}^{\circ}$. Lemma G.1. With the notation introduced above, it holds that

$$\left\|\Sigma^{1/2}\mathbf{b}_{\lambda}\right\|^{2} \leqslant \frac{\sigma_{k^{*}}^{2}}{4} \left\|\Sigma^{-1/2}\boldsymbol{\theta}_{\leqslant k^{*}}^{\circ}\right\|^{2} + \left\|\Sigma^{1/2}\boldsymbol{\theta}_{>k^{*}}^{\circ}\right\|^{2}$$

and

$$\frac{\left\|\Sigma^{3/2}\mathbf{b}_{\lambda}\right\|^{2}}{2\lambda} \leqslant \sigma_{k^{*}}^{2} \left\|\Sigma^{-1/2}\boldsymbol{\theta}_{\leqslant k^{*}}^{\circ}\right\|^{2} + \frac{1}{4} \left\|\Sigma^{1/2}\boldsymbol{\theta}_{>k^{*}}^{\circ}\right\|^{2}.$$

Proof. Let us first elaborate on the upper bound on $\|\Sigma^{1/2}\mathbf{b}_{\lambda}\|^2$. It holds that

$$\begin{split} \left\| \Sigma^{1/2} \mathbf{b}_{\lambda} \right\|^{2} &= (2\lambda)^{2} \left\| \Sigma^{1/2} (\Sigma^{2} + 2\lambda I_{d})^{-1} \boldsymbol{\theta}^{\circ} \right\|^{2} = 4\lambda^{2} \sum_{j=1}^{d} \frac{\sigma_{j} \beta_{j}^{2}}{(\sigma_{j}^{2} + 2\lambda)^{2}} \\ &= 4\lambda^{2} \sum_{j=1}^{k^{*}} \frac{\sigma_{j} \beta_{j}^{2}}{(\sigma_{j}^{2} + 2\lambda)^{2}} + 4\lambda^{2} \sum_{j>k^{*}} \frac{\sigma_{j} \beta_{j}^{2}}{(\sigma_{j}^{2} + 2\lambda)^{2}}. \end{split}$$

Due to the Cauchy inequality, we have $8\lambda\sigma_j^2 \leq (\sigma_j^2 + 2\lambda)^2$ for any $j \in \{1, \ldots, d\}$. This yields that

$$\begin{split} \left\| \Sigma^{1/2} \mathbf{b}_{\lambda} \right\|^{2} &= 4\lambda^{2} \sum_{j=1}^{k^{*}} \frac{\sigma_{j} \beta_{j}^{2}}{(\sigma_{j}^{2} + 2\lambda)^{2}} + 4\lambda^{2} \sum_{j > k^{*}} \frac{\sigma_{j} \beta_{j}^{2}}{(\sigma_{j}^{2} + 2\lambda)^{2}} \\ &\leq \frac{\lambda}{2} \sum_{j=1}^{k^{*}} \sigma_{j}^{-1} \beta_{j}^{2} + \sum_{j > k^{*}} \sigma_{j} \beta_{j}^{2} = \frac{\lambda}{2} \left\| \Sigma^{-1/2} \boldsymbol{\theta}_{\leq k^{*}}^{\circ} \right\|^{2} + \left\| \Sigma^{1/2} \boldsymbol{\theta}_{>k^{*}}^{\circ} \right\|^{2}. \end{split}$$

Due to the definition of k^* , the right-hand side does not exceed $\sigma_{k^*}^2 \|\Sigma^{-1/2} \boldsymbol{\theta}_{\leqslant k^*}^\circ\|^2 / 4 + \|\Sigma^{1/2} \boldsymbol{\theta}_{>k^*}^\circ\|^2$. Similarly, we obtain that

$$\frac{\left\| \Sigma^{3/2} \mathbf{b}_{\lambda} \right\|^{2}}{2\lambda} = 2\lambda \sum_{j=1}^{k^{*}} \frac{\sigma_{j}^{3} \beta_{j}^{2}}{(\sigma_{j}^{2} + 2\lambda)^{2}} + 2\lambda \sum_{j>k^{*}} \frac{\sigma_{j}^{3} \beta_{j}^{2}}{(\sigma_{j}^{2} + 2\lambda)^{2}} \leqslant 2\lambda \sum_{j=1}^{k^{*}} \sigma_{j}^{-1} \beta_{j}^{2} + \frac{1}{4} \sum_{j>k^{*}} \sigma_{j} \beta_{j}^{2}$$
$$= 2\lambda \left\| \Sigma^{-1/2} \boldsymbol{\theta}_{\leqslant k^{*}}^{\circ} \right\|^{2} + \frac{1}{4} \left\| \Sigma^{1/2} \boldsymbol{\theta}_{>k^{*}}^{\circ} \right\|^{2} \leqslant \sigma_{k^{*}}^{2} \left\| \Sigma^{-1/2} \boldsymbol{\theta}_{\leqslant k^{*}}^{\circ} \right\|^{2} + \frac{1}{4} \left\| \Sigma^{1/2} \boldsymbol{\theta}_{>k^{*}}^{\circ} \right\|^{2}.$$

Lemma G.2. Let $\tau^2 = 2\lambda$. Then it holds that

$$\max\left\{\left\|\Sigma^{1/2}\mathbf{b}_{\lambda}\right\|^{2}, \frac{\left\|\Sigma^{3/2}\mathbf{b}_{\lambda}\right\|^{2}}{2\lambda}\right\} \leqslant 2\tau^{2}\left\|\Sigma^{1/2}(\Sigma+\tau I_{d})^{-1}\boldsymbol{\theta}^{\circ}\right\|^{2}.$$

Proof. The proof is trivial. First, note that

$$\begin{split} \left\| \Sigma^{1/2} \mathbf{b}_{\lambda} \right\|^{2} &= (2\lambda)^{2} \left\| \Sigma^{1/2} (\Sigma^{2} + 2\lambda I_{d})^{-1} \boldsymbol{\theta}^{\circ} \right\|^{2} = 4\lambda^{2} \sum_{j=1}^{d} \frac{\sigma_{j} \beta_{j}^{2}}{(\sigma_{j}^{2} + 2\lambda)^{2}} \\ &= \tau^{4} \sum_{j=1}^{d} \frac{\sigma_{j} \beta_{j}^{2}}{(\sigma_{j}^{2} + \tau^{2})^{2}} \leqslant \tau^{2} \sum_{j=1}^{d} \frac{\sigma_{j} \beta_{j}^{2}}{\sigma_{j}^{2} + \tau^{2}} \leqslant 2\tau^{2} \sum_{j=1}^{d} \frac{\sigma_{j} \beta_{j}^{2}}{(\sigma_{j} + \tau)^{2}}. \end{split}$$

The expression in the right-hand side is nothing but $2\tau^2 \|\Sigma^{1/2} (\Sigma + \tau I_d)^{-1} \theta^{\circ}\|^2$. This concludes the first part of the proof. The second one follows from the inequalities

$$\frac{\left\|\boldsymbol{\Sigma}^{3/2}\mathbf{b}_{\lambda}\right\|^{2}}{2\lambda} = 2\lambda \sum_{j=1}^{d} \frac{\sigma_{j}^{3}\beta_{j}^{2}}{(\sigma_{j}^{2}+2\lambda)^{2}} = \tau^{2} \sum_{j=1}^{d} \frac{\sigma_{j}^{3}\beta_{j}^{2}}{(\sigma_{j}^{2}+\tau^{2})^{2}}$$
$$\leqslant \tau^{2} \sum_{j=1}^{d} \frac{\sigma_{j}\beta_{j}^{2}}{\sigma_{j}^{2}+\tau^{2}} \leqslant 2\tau^{2} \sum_{j=1}^{d} \frac{\sigma_{j}\beta_{j}^{2}}{(\sigma_{j}+\tau)^{2}} = 2\tau^{2} \left\|\boldsymbol{\Sigma}^{1/2}(\boldsymbol{\Sigma}+\tau\boldsymbol{I}_{d})^{-1}\boldsymbol{\theta}^{\circ}\right\|^{2}.$$

H Proof of Lemma 3.1 and auxiliary results

H.1 Proof of Lemma 3.1

We start the proof with the following non-probabilistic lemma.

Lemma H.1. For some $\rho_0 \leq 1/8$, define sets

$$\begin{split} \Theta &= \left\{ \boldsymbol{\theta} \in \mathbb{R}^d \mid \|\boldsymbol{\theta}\| \leqslant \rho_0 \mu \text{ and } \|\boldsymbol{\Sigma}\| \|\boldsymbol{\theta} - \boldsymbol{\theta}^*\| \leqslant 5\rho_0 \mu \sqrt{\lambda}/96 \right\}, \\ \mathsf{H} &= \left\{ \boldsymbol{\eta} \in \mathbb{R}^d \mid \|\boldsymbol{\eta} - \mathbf{Z}\| \leqslant \rho_0 \mu \sqrt{\lambda}/3 \right\}, \\ \mathsf{A} &= \left\{ A \in \mathbb{R}^{d \times d} \mid \|A\| \leqslant 3 \|\boldsymbol{\Sigma}\| + \sqrt{\lambda}/3 \right\}. \end{split}$$

Suppose that

- (i) $\|\boldsymbol{\theta}^{\circ}\| \leq \rho_0 \mu/7$ and $\|\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^{\circ}\| \leq \rho_0 \mu \sqrt{\lambda}/(18 \cdot 16);$
- (ii) $\|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| \|\widehat{\Sigma} \Sigma\| \leq \rho_0 \mu \lambda / (2 \cdot 96)$ and $\|\widehat{\Sigma} \Sigma\| \leq \|\Sigma\|$;
- (iii) $\|\Sigma\| \|\mathbf{U}\| \leq \rho_0 \mu \lambda / (2 \cdot 96)$ and $\|\mathbf{U}\| \leq \rho_0 \mu \sqrt{\lambda} / (2 \cdot 96)$.

Then, we have the following:

$$(\mathbf{0}, \mathbf{Z}, \widehat{\Sigma}), (\boldsymbol{\theta}^*, A^*, \boldsymbol{\eta}^*), (\widehat{\boldsymbol{\theta}}, \widehat{A}, \widehat{\boldsymbol{\eta}}) \in \Theta \times \mathsf{A} \times \mathsf{H}.$$

and $\Theta \times \mathsf{H} \times \mathsf{A} \subset \Upsilon(\rho_0)$. Moreover, $\|\widehat{A} - \widehat{\Sigma}\| \leq \sqrt{\lambda}/3$.

Next, we will choose λ , to satisfy conditions (ii)-(iii) using the following proposition proved in Section H.4.

Proposition H.2. Fix $\delta \in (0; 1)$ and $\rho_0 \leq 1/8$. Grant Assumptions 2.1 and 2.2. Assume that

- (i) $\|\boldsymbol{\theta}^{\circ}\| \leq \rho_{0}\mu/7;$ (ii) $\|\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^{\circ}\| \leq \rho_{0}\mu\sqrt{\lambda};$
- (iii)

$$\begin{aligned} \frac{\lambda}{\|\Sigma\|} \wedge \sqrt{\lambda} &\geq \frac{2^{11} \sigma \|\Sigma\|^{1/2}}{\rho_0 \mu} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}, \\ \lambda &\geq \frac{2^{13} \|\Sigma\|^2 \|\boldsymbol{\theta}^\circ\|}{\rho_0 \mu} (1 + C_X) \sqrt{\frac{4\mathbf{r}(\Sigma) + \log(2/\delta)}{n}}, \\ n &\geq 2^{12} (1 + C_X)^2 \left(\mathbf{r}(\Sigma) + \log(2/\delta)\right). \end{aligned}$$

Then, conditions (i)-(iii) of Lemma H.1 are satisfied and the upper bound

$$\|\mathbf{U}\| \leq 8\sigma \|\Sigma\|^{1/2} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}$$

holds with probability at least $1 - \delta$.

Clearly, Lemma H.1 combined with Proposition H.2 implies the desired result.

H.2 Proof of Lemma H.1

Step 1. Bounding the norm of $\hat{\theta}$. We start from the following chain of inequalities:

$$\lambda/2\|\widehat{\boldsymbol{\theta}}\|^2 \leqslant \mathcal{L}(\widehat{\boldsymbol{\theta}},\widehat{\boldsymbol{\eta}},\widehat{\boldsymbol{\Sigma}}) \leqslant \mathcal{L}(\mathbf{0},\mathbf{Z},\widehat{\boldsymbol{\Sigma}}) = \frac{1}{2}\|\mathbf{Z}\|^2.$$

It implies $\|\widehat{\boldsymbol{\theta}}\|^2 \leqslant \|\mathbf{Z}\|^2/\lambda$. Next, we have

$$\|\mathbf{Z}\| \leq \|\widehat{\boldsymbol{\Sigma}}\boldsymbol{\theta}^{\circ}\| + \|\mathbf{U}\| \leq \|\widehat{\boldsymbol{\Sigma}} - \boldsymbol{\Sigma}\|\|\boldsymbol{\theta}\| + \|\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^{\circ}\| + \|\mathbf{U}\|$$

Using $\|\widehat{\Sigma} - \Sigma\| \leq \|\Sigma\|$ from condition (ii), $\|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| \leq \rho_0 \mu \sqrt{\lambda}/(18 \cdot 8)$ from condition (i) and $\|\mathbf{U}\| \leq \rho_0 \mu \sqrt{\lambda}/18$ from condition (iii), we infer

$$\|\mathbf{Z}\| \leq 2\|\Sigma\| \|\boldsymbol{\theta}^{\circ}\| + \|\mathbf{U}\| \leq \rho_0 \mu \sqrt{\lambda}/6.$$
(150)

It yields

$$\|\widehat{\boldsymbol{\theta}}\| \leqslant \|\mathbf{Z}\|/\sqrt{\lambda} \leqslant \rho_0 \mu/6.$$
(151)

Note that bound (150) also implies that $(\mathbf{0}, \mathbf{Z}, \widehat{\Sigma}) \in \Theta \times \mathsf{H} \times \mathsf{A}$.

Step 2. Weak perturbation bounds. A bound on $\|\widehat{\theta}\|$ allows us applying the following lemma:

Lemma H.3. Assume that $\|\Sigma\| \|\theta^{\circ}\| \leq 4\mu\sqrt{\lambda}$. Then, if $\max\{\|\widehat{\theta}\|, \|\theta^{*}\|, \|\theta^{\circ}\|\} \leq \rho_{0}\mu$ for $\rho_{0} \leq 1/2$, then

$$\|\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\| \leq \frac{8\|\Sigma - \widehat{\Sigma}\|\|\boldsymbol{\theta}^\circ\|}{\sqrt{\lambda}} + \frac{2\|\mathbf{U}\|}{\sqrt{\lambda}}.$$

Applying Lemma B.2 with ρ_0 in place of ρ , we get $\|\theta^*\| \leq \|\theta^\circ\|$, and the latter is at most $\rho_0 \mu/7$ by the conditions of the lemma to be proved. Therefore, assumptions of Lemma H.3 are satisfied, and its conclusion holds.

Step 3. Proving that $\hat{\theta}, \theta^* \in \Theta$. Since $\|\theta^*\| \leq \|\theta^\circ\| \leq \rho_0 \mu$ due to Lemma B.2(iv), the vector θ^* belongs to Θ obviously. Next, we check that $\hat{\theta} \in \Theta$. It requires to bound

$$\|\Sigma\|\|\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\| \leqslant \frac{8\|\Sigma\|\|\widehat{\Sigma} - \Sigma\|\|\boldsymbol{\theta}^\circ\|}{\sqrt{\lambda}} + \frac{2\|\Sigma\|\|\mathbf{U}\|}{\sqrt{\lambda}}.$$
(152)

Using conditions (ii),(iii), we bound the latter by $4\rho_0\mu\sqrt{\lambda}/96 + \rho_0\mu\sqrt{96} = 5\rho_0\mu\sqrt{\lambda}/96$, so $\hat{\theta} \in \Theta$. Step 4. Proving that $\hat{A}, A^* \in A$. Using

$$\frac{\mu^2}{2} \|\widehat{A} - \widehat{\Sigma}\|^2 \leqslant \mathcal{L}(\widehat{\theta}, \widehat{\eta}, \widehat{A}) \leqslant \mathcal{L}(\mathbf{0}, \mathbf{Z}, \widehat{\Sigma}) = \|\mathbf{Z}\|^2 / 2,$$

we get $\|\widehat{A} - \widehat{\Sigma}\| \leq \|\mathbf{Z}\|/\mu$. Using (150), we establish $\|\widehat{A} - \widehat{\Sigma}\| \leq \rho_0 \sqrt{\lambda}/6 \leq \sqrt{\lambda}/3$. This implies

$$\|\widehat{A}\| \leqslant 3\|\Sigma\| + \sqrt{\lambda}/3,$$

since $\|\widehat{\Sigma}\| \leq 2\|\Sigma\|$ due to condition (ii). Thus, we have $\widehat{A} \in A$.

Next, applying Lemma B.2(i), we get with ρ_0 in place of ρ , we obtain

$$\|A^* - \widehat{\Sigma}\| \leq \mu^{-1} \sqrt{\lambda} \|\boldsymbol{\theta}^{\circ}\| + \|\Sigma - \widehat{\Sigma}\| \leq \rho_0 \sqrt{\lambda}/7 + \|\Sigma\| \leq \|\Sigma\| + \sqrt{\lambda}/3,$$

where we used conditions (i),(ii) in the second inequality. Hence, $||A^*|| \leq 3||\Sigma|| + \sqrt{\lambda}/3$ and $A^* \in A$. Step 5. Proving that $\eta^*, \widehat{\eta} \in H$. To prove that $\widehat{\eta} \in H$, note that $\widehat{\eta} = \frac{1}{2}(\widehat{A}\widehat{\theta} - \mathbf{Z})$. It yields

$$\begin{aligned} \|\widehat{\boldsymbol{\eta}} - \mathbf{Z}\| &= \frac{1}{2} \|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\| \leqslant \|\widehat{A}\| \|\widehat{\boldsymbol{\theta}}\|/2 + \|\mathbf{Z}\|/2 \\ &\leqslant (3\|\Sigma\| + \sqrt{\lambda}/3) \|\widehat{\boldsymbol{\theta}}\|/2 + \rho_0 \mu \sqrt{\lambda}/12 \\ &\leqslant 3\|\Sigma\| \|\boldsymbol{\theta}^*\|/2 + 3\|\Sigma\| \|\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|/2 + \sqrt{\lambda} \|\widehat{\boldsymbol{\theta}}\|/6 + \rho_0 \mu \sqrt{\lambda}/12 \\ &\leqslant 3\|\Sigma\| \|\boldsymbol{\theta}^\circ\|/2 + 3\|\Sigma\| \|\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|/2 + \sqrt{\lambda} \|\widehat{\boldsymbol{\theta}}\|/6 + \rho_0 \mu \sqrt{\lambda}/12 \\ &\leqslant \frac{\rho_0 \mu \sqrt{\lambda}}{6 \cdot 16} + \frac{15\rho_0 \mu \sqrt{\lambda}}{96} + \frac{\rho_0 \mu \sqrt{\lambda}}{6} + \frac{\rho_0 \mu \sqrt{\lambda}}{12} \leqslant \rho_0 \mu \sqrt{\lambda}/3. \end{aligned}$$

where we used $\hat{A} \in A$ and $\|\mathbf{Z}\| \leq \rho_0 \mu \sqrt{\lambda}/6$ from (150) for the second inequality, $\|\boldsymbol{\theta}^*\| \leq \|\boldsymbol{\theta}^\circ\|$ from Lemma B.2(iv) for the fourth inequality, $\hat{\boldsymbol{\theta}} \in \Theta$ and $\|\boldsymbol{\Sigma}\| \|\boldsymbol{\theta}^\circ\| \leq \mu \sqrt{\lambda}/(18 \cdot 16)$ from condition (i) for the fifth inequality.

To prove $\eta^* \in \mathsf{H}$, we bound it as follows:

$$\begin{aligned} \|\boldsymbol{\eta}^* - \mathbf{Z}\| &= \left\| \frac{1}{2} (A^* \boldsymbol{\theta}^* + \Sigma \boldsymbol{\theta}^\circ) - \mathbf{Z} \right\| \\ &\leq \left(\frac{\|A^*\| \|\boldsymbol{\theta}^*\| + \|\Sigma\| \|\boldsymbol{\theta}^\circ\|}{2} \right) + \|\mathbf{Z}\| \\ &\leq \frac{\|A^*\| + \|\Sigma\|}{2} \cdot \|\boldsymbol{\theta}^\circ\| + \|\mathbf{Z}\| \\ &\leq \|\Sigma\| \|\boldsymbol{\theta}^\circ\| + \|\boldsymbol{\theta}^\circ\| \sqrt{\lambda}/3 + \|\mathbf{Z}\| \\ &\leq \rho_0 \mu \sqrt{\lambda}/(18 \cdot 16) + \rho_0 \mu \sqrt{\lambda}/21 + \rho_0 \mu \sqrt{\lambda}/6 \leq \rho_0 \mu \sqrt{\lambda}/3 \end{aligned}$$

where we used $A^* \in A$ in the second inequality. To establish the fourth inequality, we applied bound (150) and condition (i).

Step 6. Endgame. We are left to prove that $\Theta \times H \times A \subset \Upsilon(\rho_0)$. We state it as the following proposition. **Proposition H.4.** *We have*

$$\Theta \times \mathsf{H} \times \mathsf{A} \subset \Upsilon(\rho_0).$$

Proof. Consider arbitrary $(\theta, \eta, A) \in \Theta \times H \times A$. It is enough to prove that $\|\theta\| \leq \rho_0 \mu$ and $\|A\theta - \eta\| \leq \rho_0 \mu \sqrt{\lambda}$. The former inequality is satisfied by the definition of Θ . To satisfy the second inequality, we bound $\|A\theta - \eta\|$ as follows:

$$\begin{split} \|A\boldsymbol{\theta} - \boldsymbol{\eta}\| &\leq \|A\| \|\boldsymbol{\theta}\| + \|\mathbf{Z}\| + \|\boldsymbol{\eta} - \mathbf{Z}\| \\ &\leq (\|\Sigma\| + \sqrt{\lambda}/3) \|\boldsymbol{\theta}\| + \|\mathbf{Z}\| + \rho_0 \mu \sqrt{\lambda} \\ &\leq \|\Sigma\| \|\boldsymbol{\theta}^*\| + \|\Sigma\| \|\boldsymbol{\theta} - \boldsymbol{\theta}^*\| + \|\boldsymbol{\theta}\| \sqrt{\lambda}/3 + \|\mathbf{Z}\| + \rho_0 \mu \sqrt{\lambda}/3 \\ &\leq \|\Sigma\| \|\boldsymbol{\theta}^*\| + 5\rho_0 \mu \sqrt{\lambda}/96 + \rho_0 \mu \sqrt{\lambda}/3 + \|\mathbf{Z}\| + \rho_0 \mu \sqrt{\lambda}. \end{split}$$

where the second inequality holds by the definitions of A, H, and the fourth inequality holds by the definition of Θ . Due to (150), we have $\|\mathbf{Z}\| \leq \rho_0 \mu \sqrt{\lambda}/6$, hence, we have

$$\|A\boldsymbol{\theta} - \boldsymbol{\eta}\| \leq (1 - 11/96) \rho_0 \mu \sqrt{\lambda} + \|\boldsymbol{\Sigma}\| \|\boldsymbol{\theta}^*\|.$$

Due to Lemma B.2(iv), we have $\|\boldsymbol{\theta}^*\| \leq \|\boldsymbol{\theta}^\circ\|$. Finally, the quantity $\|\boldsymbol{\Sigma}\|\|\boldsymbol{\theta}^\circ\|$ is at most $11\rho_0\mu\sqrt{\lambda}/96$ due to condition (i), so $\|A\boldsymbol{\theta} - \boldsymbol{\eta}\| \leq \rho_0\mu\sqrt{\lambda}$.

H.3 Proof of Lemma H.3

Step 1. Constructing a finite difference equation. The idea is to employ the equation

$$abla \mathtt{L}(oldsymbol{ heta},\widehat{oldsymbol{\eta}},\overline{A}) - oldsymbol{
abla} \mathtt{L}(oldsymbol{ heta}^*,oldsymbol{\eta}^*,A^*) = oldsymbol{0},$$

which holds by definitions of $\widehat{\boldsymbol{v}} = (\widehat{\boldsymbol{\theta}}, \widehat{\boldsymbol{\eta}}, \widehat{A})$ and $\boldsymbol{v}^* = (\boldsymbol{\theta}^*, \boldsymbol{\eta}^*, A^*)$. We have

$$\boldsymbol{\nabla} \mathbf{L}(\boldsymbol{\theta}, \boldsymbol{\eta}, A) = \begin{pmatrix} -A^{\top} \boldsymbol{\eta} + A^{\top} A \boldsymbol{\theta} + \lambda \boldsymbol{\theta} \\ -\mathbf{Z} + 2 \boldsymbol{\eta} - A \boldsymbol{\theta} \\ A \boldsymbol{\theta} \boldsymbol{\theta}^{\top} - \boldsymbol{\eta} \boldsymbol{\theta}^{\top} + \mu^{2} (A - \widehat{\Sigma}) \end{pmatrix} \text{ and } \nabla \mathcal{L}(\boldsymbol{\theta}, \boldsymbol{\eta}, A) = \begin{pmatrix} -A^{\top} \boldsymbol{\eta} + A^{\top} A \boldsymbol{\theta} + \lambda \boldsymbol{\theta} \\ -\mathbb{E} \mathbf{Z} + 2 \boldsymbol{\eta} - A \boldsymbol{\theta} \\ A \boldsymbol{\theta} \boldsymbol{\theta}^{\top} - \boldsymbol{\eta} \boldsymbol{\theta}^{\top} + \mu^{2} (A - \widehat{\Sigma}) \end{pmatrix}.$$

Define an operator Δ , acting on a matrix-valued functions $f(\theta, \eta, A)$ as $\Delta(f) = f(\widehat{A}, \widehat{\eta}, \widehat{\theta}) - f(A^*, \eta^*, \theta^*)$. Note that it possesses the following identities for any two matrix-valued functions f, g such that their product is defined:

$$\Delta(fg) = \Delta(f) g(\widehat{\theta}, \widehat{\eta}, \widehat{A}) + f(\theta^*, \eta^*, A^*) \Delta(g);$$

$$\Delta(fg) = f(\widehat{\theta}, \widehat{\eta}, \widehat{A}) \Delta(g) + \Delta(f) g(\theta^*, \eta^*, A^*).$$
 (153)

Using $\nabla \mathtt{L}(\widehat{\pmb{ heta}},\widehat{\pmb{\eta}},\widehat{A}) - \nabla \mathcal{L}(\pmb{ heta}^*,\pmb{\eta}^*,A^*) = 0$, we obtain

$$\begin{cases} -\Delta(A^{\top}\boldsymbol{\eta}) + \Delta(A^{\top}A\boldsymbol{\theta}) + \lambda\Delta(\boldsymbol{\theta}) &= \mathbf{0} \\ -(\mathbf{Z} - \mathbb{E}\mathbf{Z}) + 2\Delta(\boldsymbol{\eta}) - \Delta(A\boldsymbol{\theta}) &= \mathbf{0} \\ \Delta(A\boldsymbol{\theta}\boldsymbol{\theta}^{\top}) - \Delta(\boldsymbol{\eta}\boldsymbol{\theta}^{\top}) + \mu^{2}\Delta(A) &= \mu^{2}(\widehat{\Sigma} - \Sigma) \end{cases}$$

The second equation implies $\Delta(\eta) = \frac{1}{2}(\Delta(A\theta) + (\mathbf{Z} - \mathbb{E}\mathbf{Z}))$. Substituting it in the above system, we get

$$\begin{split} \Delta(A^{\top}\boldsymbol{\eta}) &= \widehat{A}^{\top}\Delta(\boldsymbol{\eta}) + \Delta^{\top}(A)\boldsymbol{\eta}^{*} \\ &= \frac{1}{2}\widehat{A}^{\top}\Delta(A\boldsymbol{\theta}) + \frac{1}{2}\widehat{A}^{\top}\Delta_{\mathbf{Z}} + \Delta^{\top}(A)\boldsymbol{\eta}^{*}, \\ \Delta(\boldsymbol{\eta}\boldsymbol{\theta}^{\top}) &= \Delta(\boldsymbol{\eta})\widehat{\boldsymbol{\theta}}^{\top} + \boldsymbol{\eta}^{*}\Delta^{\top}(\boldsymbol{\theta}) \\ &= \frac{1}{2}\Delta(A\boldsymbol{\theta})\widehat{\boldsymbol{\theta}}^{\top} + \frac{1}{2}\Delta_{\mathbf{Z}}\widehat{\boldsymbol{\theta}}^{\top} + \boldsymbol{\eta}^{*}\Delta^{\top}(\boldsymbol{\theta}), \end{split}$$

where we denote $\mathbf{Z} - \mathbb{E}\mathbf{Z}$ by $\Delta_{\mathbf{Z}}$ for brevity. Since $\Delta(A\boldsymbol{\theta}\boldsymbol{\theta}^{\top}) = \Delta(A\boldsymbol{\theta})\widehat{\boldsymbol{\theta}}^{\top} + A^*\boldsymbol{\theta}^*\Delta^{\top}(\boldsymbol{\theta})$ and $\Delta(A^{\top}A\boldsymbol{\theta}) = \widehat{A}^{\top}\Delta(A\boldsymbol{\theta}) + \Delta^{\top}(A)A^*\boldsymbol{\theta}^*$ due to (153), we have

$$\begin{cases} \frac{1}{2}\widehat{A}^{\top}\Delta(A\boldsymbol{\theta}) - \frac{1}{2}\widehat{A}^{\top}\Delta_{\mathbf{Z}} + \Delta^{\top}(A)(A^{*}\boldsymbol{\theta}^{*} - \boldsymbol{\eta}^{*}) + \lambda\Delta(\boldsymbol{\theta}) &= \mathbf{0}, \\ \frac{1}{2}\Delta(A\boldsymbol{\theta})\widehat{\boldsymbol{\theta}}^{\top} - \frac{1}{2}\Delta_{\mathbf{Z}}\widehat{\boldsymbol{\theta}}^{\top} + (A^{*}\boldsymbol{\theta}^{*} - \boldsymbol{\eta}^{*})\Delta^{\top}(\boldsymbol{\theta}) + \mu^{2}\Delta(A) &= \mu^{2}(\widehat{\Sigma} - \Sigma). \end{cases}$$
(154)

Step 2. Bounding the norm of $||\Delta(A)||$. Deriving $\Delta(A\theta) = \Delta(A)\widehat{\theta} + A^*\Delta(\theta)$ from (153) and substituting it into the last equation of (154), we obtain

$$\Delta(A)\left(\mu^2 I_d + \frac{1}{2}\widehat{\theta}\widehat{\theta}^{\top}\right) = \mu^2(\widehat{\Sigma} - \Sigma) - (A^*\theta^* - \eta^*)\Delta^{\top}(\theta) + \frac{1}{2}\Delta_{\mathbf{Z}}\widehat{\theta}^{\top}.$$

For brevity, denote $A^*\theta^* - \eta^*$ by \mathbf{b}_{η} . Rearranging terms, we get

$$\begin{split} \Delta(A) &= \mu^2 (\widehat{\Sigma} - \Sigma) \left(\mu^2 + \frac{1}{2} \widehat{\boldsymbol{\theta}} \widehat{\boldsymbol{\theta}}^\top \right)^{-1} - \mathbf{b}_{\boldsymbol{\eta}} \Delta^\top (\boldsymbol{\theta}) \left(\mu^2 + \frac{1}{2} \widehat{\boldsymbol{\theta}} \boldsymbol{\theta}^\top \right)^{-1} \\ &+ \frac{1}{2} \Delta_{\mathbf{Z}} \widehat{\boldsymbol{\theta}}^\top \left(\mu^2 + \frac{1}{2} \widehat{\boldsymbol{\theta}} \boldsymbol{\theta}^\top \right)^{-1}, \end{split}$$

and, therefore, we bound

$$\|\Delta(A)\| \leq \|\Sigma - \widehat{\Sigma}\| + \mu^{-2} \|\mathbf{b}_{\eta}\| \|\Delta(\boldsymbol{\theta})\| + \frac{1}{2}\mu^{-2} \|\Delta_{\mathbf{Z}}\| \|\widehat{\boldsymbol{\theta}}\|.$$
(155)

Step 3. Establishing a bound on $\|\Delta(\theta)\|$. We bound $\|\Delta(\theta)\|$ using $\|\Delta(A)\theta^*\|$ and $\|\Delta^\top(A)\mathbf{b}_{\eta}\|$. From the first equation of (154), we obtain

$$\frac{1}{2}\widehat{A}^{\top}\Delta(A\boldsymbol{\theta}) - \frac{1}{2}\widehat{A}^{\top}\Delta_{\mathbf{Z}} + \Delta^{\top}(A)\mathbf{b}_{\boldsymbol{\eta}} + \lambda\Delta(\boldsymbol{\theta}) = \mathbf{0}$$

Since $\Delta(A\theta) = \widehat{A}\Delta(\theta) + \Delta(A)\theta^*$, we have

$$\left(\frac{1}{2}\widehat{A}^{\top}\widehat{A} + \lambda I_d\right)\Delta(\boldsymbol{\theta}) = \frac{1}{2}\widehat{A}^{\top}\Delta_{\mathbf{Z}} - \Delta^{\top}(A)\mathbf{b}_{\boldsymbol{\eta}} - \frac{1}{2}\widehat{A}^{\top}\Delta(A)\boldsymbol{\theta}^*.$$
(156)

Next, we bound the norms $\left\| \left(\frac{1}{2}\widehat{A}^{\top}\widehat{A} + \lambda I_d\right)^{-1} \right\|$ and $\left\| \left(\frac{1}{2}\widehat{A}^{\top}\widehat{A} + \lambda I_d\right)^{-1}\widehat{A} \right\|$. While the first norm is clearly at most $1/\lambda$, for the second norm we need to maximize

$$\left\| \left(\frac{1}{2}\widehat{A}^{\top}\widehat{A} + \lambda I_d\right)^{-1}\widehat{A} \right\| = \max_i \frac{\sigma_i}{\sigma_i^2/2 + \lambda}$$

over all possible singular values $\sigma_1, \ldots, \sigma_n$ of \widehat{A} . However, we have

$$\max_{i} \frac{\sigma_{i}}{\sigma_{i}^{2}/2 + \lambda} \leqslant \max_{x \in \mathbb{R}_{+}} \frac{x}{x^{2}/2 + \lambda} = \frac{\sqrt{2\lambda}}{2\lambda} \leqslant \frac{1}{\sqrt{2\lambda}}$$

by elementary calculus. Thus, multiplying both sides of (156) by $\left(\frac{1}{2}\widehat{A}^{\top}\widehat{A} + \lambda I_{d}\right)^{-1}$ and applying bounds on $\left\| \left(\frac{1}{2}\widehat{A}^{\top}\widehat{A} + \lambda I_{d}\right)^{-1} \right\|$ and $\left\| \left(\frac{1}{2}\widehat{A}^{\top}\widehat{A} + \lambda I_{d}\right)^{-1}\widehat{A} \right\|$, we derive $\|\Delta(\boldsymbol{\theta})\| \leqslant \frac{\|\Delta \mathbf{z}\|}{\sqrt{8\lambda}} + \frac{1}{\lambda} \|\Delta^{\top}(A)\mathbf{b}_{\boldsymbol{\eta}}\| + \frac{1}{2\sqrt{\lambda}} \|\Delta(A)\boldsymbol{\theta}^{*}\|$ $\leqslant \frac{\|\Delta \mathbf{z}\|}{\sqrt{8\lambda}} + \frac{\|\Delta(A)\|}{\sqrt{\lambda}} \left(\frac{1}{2}\|\boldsymbol{\theta}^{*}\| + \frac{\|\mathbf{b}_{\boldsymbol{\eta}}\|}{\sqrt{\lambda}}\right)$ $\leqslant \frac{\|\Delta \mathbf{z}\|}{\sqrt{8\lambda}} + \frac{1}{\sqrt{\lambda}} \left(\|\Sigma - \widehat{\Sigma}\| + \frac{\mu^{-2}}{2}\|\Delta \mathbf{z}\|\|\widehat{\boldsymbol{\theta}}\|\right) \left(\frac{1}{2}\|\boldsymbol{\theta}^{*}\| + \frac{\|\mathbf{b}_{\boldsymbol{\eta}}\|}{\sqrt{\lambda}}\right)$ $+ \frac{\|\mathbf{b}_{\boldsymbol{\eta}}\|\|\Delta(\boldsymbol{\theta})\|}{\mu^{2}\sqrt{\lambda}} \left(\frac{1}{2}\|\boldsymbol{\theta}^{*}\| + \frac{\|\mathbf{b}_{\boldsymbol{\eta}}\|}{\sqrt{\lambda}}\right),$

where the third inequality holds due to (155). According to Lemma B.2(iii), we have $\|\mathbf{b}_{\eta}\| \leq \sqrt{\lambda} \|\boldsymbol{\theta}^{\circ}\|$. It implies

$$\begin{split} \|\Delta(\boldsymbol{\theta})\| &\leqslant \frac{\Delta_{\mathbf{Z}}}{\sqrt{8\lambda}} + \frac{1}{\sqrt{\lambda}} \left(\|\Sigma - \widehat{\Sigma}\| + \frac{\mu^{-2}}{2} \|\Delta_{\mathbf{Z}}\| \|\widehat{\boldsymbol{\theta}}\| \right) \left(\frac{1}{2} \|\boldsymbol{\theta}^*\| + \|\boldsymbol{\theta}^\circ\| \right) \\ &+ \mu^{-2} \|\Delta(\boldsymbol{\theta})\| \|\boldsymbol{\theta}^\circ\| \left(\frac{1}{2} \|\boldsymbol{\theta}^*\| + \|\boldsymbol{\theta}^\circ\| \right). \end{split}$$

By assumptions of the lemma, we have $\|\boldsymbol{\theta}^{\circ}\|, \|\boldsymbol{\theta}^{*}\|$ and $\|\widehat{\boldsymbol{\theta}}\|$ are at most $\rho_{0}\mu$, which implies

$$\begin{aligned} \|\Delta(\boldsymbol{\theta})\| &\leqslant \frac{\|\Delta_{\mathbf{Z}}\|}{\sqrt{8\lambda}} + \frac{1}{\sqrt{\lambda}} \left(\|\Sigma - \widehat{\Sigma}\| + \rho_0 \mu^{-1}/2 \|\Delta_{\mathbf{Z}}\| \right) \left(\frac{1}{2} \|\boldsymbol{\theta}^*\| + \|\boldsymbol{\theta}^\circ\| \right) \\ &+ 2\rho_0^2 \|\Delta(\boldsymbol{\theta})\|. \end{aligned}$$

Since $\Delta_{\mathbf{Z}} = (\Sigma - \widehat{\Sigma}) \boldsymbol{\theta}^{\circ} + \mathbf{U}$, we may simplify the inequality above and obtain

$$(1 - 2\rho_0^2) \|\Delta(\boldsymbol{\theta})\| \leq \frac{\|\Delta_{\mathbf{Z}}\|}{2\sqrt{\lambda}} \cdot (1 + 2\rho_0^2) + \frac{\|\Sigma - \Sigma\|}{\sqrt{\lambda}} (\|\boldsymbol{\theta}^*\| + \|\boldsymbol{\theta}^\circ\|)$$
$$\leq \frac{2\|\Sigma - \widehat{\Sigma}\|}{\sqrt{\lambda}} (\|\boldsymbol{\theta}^*\| + \|\boldsymbol{\theta}^\circ\|) + \frac{\|\mathbf{U}\|}{\sqrt{\lambda}},$$

where we used $\rho_0 \leq 1/2$. It implies the desired bound on $\Delta(\theta) = \hat{\theta} - \theta^*$:

$$\|\Delta(oldsymbol{ heta})\| \leqslant rac{4\|\Sigma - \widehat{\Sigma}\|}{\sqrt{\lambda}} (\|oldsymbol{ heta}^*\| + \|oldsymbol{ heta}^\circ\|) + rac{2\|\mathbf{U}\|}{\sqrt{\lambda}}.$$

Applying bound $\|\theta^*\| \leq \|\theta^\circ\|$ from Lemma B.2(iv), we obtain the lemma.

H.4 Proof of Proposition H.2

Step 1. Bound on $||\mathbf{U}||$. To ensure conditions (ii),(iii), we shall obtain bounds on $||\widehat{\Sigma} - \Sigma||$ and $||\mathbf{U}||$ with high probability. For the latter quantity, we apply Theorem 2.6 with $B = I_d$, and obtain

$$\|\mathbf{U}\| \leqslant 8\sigma \|\Sigma\|^{1/2} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}},\tag{157}$$

with probability at least $1 - \delta/2$, provided $n \ge \mathbf{r}(\Sigma) + \log(4/\delta)$.

Step 2. Bound on $\|\widehat{\Sigma} - \Sigma\|$. First, we will show that under Assumption 2.1, a random vector \mathbf{X}_1 satisfies $\psi_2 - L_2$ -equivalence. In what follows, $\|\cdot\|_{\psi_1}, \|\cdot\|_{\psi_2}$ stand for ψ_1 and ψ_2 Orlicz norms.

Lemma H.5 (Lemma B.3 of Puchkin and Rakhuba [2024]). Suppose that a random vector \mathbf{X} satisfies Assumption 2.1. Then, for any vector $\mathbf{u} \in \mathbb{R}^d$, it holds that

$$\|\mathbf{u}^{\top}\mathbf{X}\|_{\psi_2}^2 \leqslant \frac{1+C_X}{\log 2} \cdot \mathbf{u}^{\top} \Sigma \mathbf{u}$$

Then, we deduce the concentration bound on $\widehat{\Sigma}$ from the following theorem.

Lemma H.6 (Theorem 1 of Zhivotovskiy [2024]). Assume that M_1, \ldots, M_n are independent copies of a d by d positive semi-definite symmetric matrix M with mean $\mathbb{E}M = \Sigma$. Let M satisfy for some $\kappa \ge 1$

$$\|\mathbf{u}^{\top} M \mathbf{u}\|_{\psi_1} \leqslant \kappa^2 \mathbf{u}^{\top} \Sigma \mathbf{u}$$

for all $\mathbf{u} \in \mathbb{R}^d$. Then, for any $\delta > 0$, with probability $1 - \delta$, it holds that

$$\left|\frac{1}{n}\sum_{i=1}^{n}M_{i}-\Sigma\right\| \leq 20\kappa^{2}\|\Sigma\|\sqrt{\frac{4\mathbf{r}(\Sigma)+\log(1/\delta)}{n}}$$

whenever $n \ge 4\mathbf{r}(\Sigma) + \log(1/\delta)$

Combining the above two lemmata and using an inequality $\|\xi^2\|_{\psi_1} \leq \|\xi\|_{\psi_2}^2$ suitable for any random variable ξ , we derive

$$\|\widehat{\Sigma} - \Sigma\| \leqslant \frac{20(1+C_X)}{\log 2} \cdot \|\Sigma\| \sqrt{\frac{4\mathbf{r}(\Sigma) + \log(2/\delta)}{n}}$$
(158)

with probability at least $1 - \delta/2$.

Step 3. Satisfiability of conditions (ii),(iii) of Lemma H.1. Note that if

$$\lambda \geqslant \frac{2^{13} \|\Sigma\|^2 \|\boldsymbol{\theta}^\circ\|}{\rho_0 \mu} (1 + C_X) \sqrt{\frac{4\mathbf{r}(\Sigma) + \log(2/\delta)}{n}},\tag{159}$$

$$n \ge 2^{12} (1 + C_X)^2 \left(\mathbf{r}(\Sigma) + \log(2/\delta) \right),$$
 (160)

then condition (ii) of Lemma H.1 is satisfied with probability at least $1 - \delta/2$ due to (158). Similarly, due to (157), condition (iii) of Lemma H.1 is satisfied with probability at least $1 - \delta/2$ if

$$\frac{\lambda}{\|\Sigma\|} \wedge \sqrt{\lambda} \ge \frac{2^{11}\sigma \|\Sigma\|^{1/2}}{\rho_0 \mu} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}$$
(161)

holds. The union bound implies that conditions (ii)-(iii) simultaniously hold with probability at least $1 - \delta$, provided inequalities (159)-(161) are satisfied.

I Proof of Theorem 3.3

Step 1. Establishing a finite difference equation. We consider the following three stationary point conditions:

$$\begin{aligned} \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}_t, (A_{t-1}\boldsymbol{\theta}_t + \mathbf{Z})/2, A_{t-1}) &= \mathbf{0}, \\ \nabla_A \mathcal{L}(\boldsymbol{\theta}_t, (A_t\boldsymbol{\theta}_t + \mathbf{Z})/2, A_t) &= \mathbf{0}, \\ \nabla_{\boldsymbol{\eta}} \mathcal{L}(\widehat{\boldsymbol{\theta}}, (\widehat{A}\widehat{\boldsymbol{\theta}} + \mathbf{Z})/2, \widehat{A}) &= \mathbf{0}, \end{aligned}$$

The first two equations can be expanded as follows:

$$\begin{cases} \left(\frac{1}{2}A_{t-1}^{\top}A_{t-1} + \lambda I_{d}\right)\boldsymbol{\theta}_{t} - \frac{1}{2}A_{t-1}^{\top}\mathbf{Z} = \mathbf{0} \\ A_{t}\boldsymbol{\theta}_{t}\boldsymbol{\theta}_{t}^{\top}/2 - \frac{1}{2}\mathbf{Z}\boldsymbol{\theta}^{\top} + \mu^{2}(A_{t} - \widehat{\Sigma}) = \mathbf{0}. \end{cases}$$
(162)

For a matrix-valued function $f(\theta, \eta, A)$, we define an operator

$$\Delta_t(f) = f(\boldsymbol{\theta}_t, \boldsymbol{\eta}_t, A_t) - f(\widehat{\boldsymbol{\theta}}, \widehat{\boldsymbol{\eta}}, \widehat{A}).$$

Note that for product of any matrix-valued functions f, g of appropriate shapes, we can write

$$\Delta_t(fg) = f(\boldsymbol{\theta}_t, \boldsymbol{\eta}_t, A_t) \Delta_t(g) + \Delta_t(f) g(\widehat{\boldsymbol{\theta}}, \widehat{\boldsymbol{\eta}}, \widehat{A})$$

$$\Delta_t(fg) = \Delta_t(f) g(\boldsymbol{\theta}_t, \boldsymbol{\eta}_t, A_t) + f(\widehat{\boldsymbol{\theta}}, \widehat{\boldsymbol{\eta}}, \widehat{A}) \Delta_t(g).$$
(163)

Using this notation, we subtract $\nabla_{\boldsymbol{\theta}} \mathcal{L}(\widehat{\boldsymbol{\theta}}, (\widehat{A}\widehat{\boldsymbol{\theta}} + \mathbf{Z})/2, \widehat{A}) = \mathbf{0}$ from the first equation of (162) and $\nabla_A \mathcal{L}(\widehat{\boldsymbol{\theta}}, (\widehat{A}\widehat{\boldsymbol{\theta}} + \mathbf{Z})/2, \widehat{A}) = \mathbf{0}$ from the second equation of (162), and obtain

$$\begin{cases} \left(\frac{1}{2}A_{t-1}^{\top}A_{t-1}+\lambda I_{d}\right)\Delta_{t}(\boldsymbol{\theta})+\frac{1}{2}\Delta_{t-1}(A^{\top}A)\widehat{\boldsymbol{\theta}}-\frac{1}{2}\Delta_{t-1}^{\top}(A)\mathbf{Z} &=\mathbf{0},\\ \Delta_{t}(A)(\boldsymbol{\theta}_{t}\boldsymbol{\theta}_{t}^{\top}/2+\mu^{2}I_{d})+\frac{1}{2}\widehat{A}\Delta_{t}(\boldsymbol{\theta}\boldsymbol{\theta}^{\top})-\frac{1}{2}\mathbf{Z}\Delta_{t}^{\top}(\boldsymbol{\theta}) &=\mathbf{0}. \end{cases}$$

Expanding $\Delta_{t-1}(A^{\top}A) = A_{t-1}^{\top}\Delta_{t-1}(A) + \Delta_{t-1}^{\top}(A)\widehat{A}, \ \Delta_t(\boldsymbol{\theta}\boldsymbol{\theta}^{\top}) = \Delta_t(\boldsymbol{\theta})\boldsymbol{\theta}_t^{\top} + \widehat{\boldsymbol{\theta}}\Delta_t^{\top}(\boldsymbol{\theta})$ due to (163), and rearranging terms, we obtain

$$\begin{cases} \Delta_t(\boldsymbol{\theta}) &= -\frac{1}{2} (A_{t-1}^\top A_{t-1}/2 + \lambda I_d)^{-1} \left[A_{t-1}^\top \Delta_{t-1}(A) \widehat{\boldsymbol{\theta}} + \frac{1}{2} \Delta_{t-1}(A) (\widehat{A} \widehat{\boldsymbol{\theta}} - \mathbf{Z}) \right], \\ \Delta_t(A) &= -\frac{1}{2} \left[\widehat{A} \Delta_t(\boldsymbol{\theta}) \boldsymbol{\theta}_t^\top + (\widehat{A} \widehat{\boldsymbol{\theta}} - \mathbf{Z}) \Delta_t(\boldsymbol{\theta}) \right] (\mu^2 I_d + \boldsymbol{\theta}_t \boldsymbol{\theta}_t^\top/2)^{-1}. \end{cases}$$
(164)

Step 2. Recursively bounding the norm of $\Delta_t(\theta)$ and $\Delta_t(A)$. Let us start with the first equation of the system above. We have

$$\|\Delta_{t}(\boldsymbol{\theta})\| \leq \|\Delta_{t-1}(A)\| \left[\|(A_{t-1}^{\top}A_{t-1} + \lambda I_{d})^{-1}A_{t-1}^{\top}\| \|\widehat{\boldsymbol{\theta}}\| + \|(A_{t-1}^{\top}A_{t-1} + 2\lambda I_{d})^{-1}\| \|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\| \right].$$

Let $\sigma_1, \ldots, \sigma_d$ be the singular values of A_{t-1} . Using elementary calculus, we obtain

$$\|(A_{t-1}^{\top}A_{t-1} + \lambda I_d)^{-1}A_{t-1}^{\top}\| = \max_i \frac{\sigma_i}{\sigma_i^2 + \lambda} \leqslant \max_{x \in \mathbb{R}_+} \frac{x}{x^2 + \lambda} \leqslant \frac{1}{\sqrt{\lambda}}$$

Then, using $\|(A_{t-1}^{\top}A_{t-1}+2\lambda I_d)^{-1}\|\leqslant 1/(2\lambda)$, we elaborate

$$\|\Delta_t(\boldsymbol{\theta})\| \leq \|\Delta_{t-1}(A)\| \left[\frac{\|\widehat{\boldsymbol{\theta}}\|}{\sqrt{\lambda}} + \frac{\|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\|}{2\lambda} \right].$$
(165)

Next, we analyze the norm of $\Delta_t(A)$. From the second equation (164), we infer

$$\|\Delta_t(A)\| \leq \mu^{-2} \left[\|\widehat{A}\| \|\boldsymbol{\theta}_t\| + \|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\| \right] \cdot \|\Delta_t(\boldsymbol{\theta})\|/2$$

Bounding $\|\boldsymbol{\theta}_t\| \leq \|\widehat{\boldsymbol{\theta}}\| + \|\Delta_t(\boldsymbol{\theta})\|$ and applying bound (165), we obtain

$$\begin{split} \|\Delta_t(A)\| &\leqslant \frac{\mu^{-2}}{2} \left[\|\widehat{A}\| \|\widehat{\boldsymbol{\theta}}\| + \|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\| \right] \left[\frac{\|\widehat{\boldsymbol{\theta}}\|}{\sqrt{\lambda}} + \frac{\|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\|}{2\lambda} \right] \|\Delta_{t-1}(A)\| \\ &+ \frac{\mu^{-2}}{2} \|\widehat{A}\| \left[\frac{\|\widehat{\boldsymbol{\theta}}\|}{\sqrt{\lambda}} + \frac{\|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\|}{2\lambda} \right]^2 \|\Delta_{t-1}(A)\|^2. \end{split}$$

Step 3. Exploiting the recursion. We will prove by induction, that

$$\|\Delta_t(A)\| \leqslant \rho_0^{2t} \|\widehat{A} - \widehat{\Sigma}\|.$$
(166)

Clearly, the base t = 0 holds, by the definition of $A_0 = \hat{\Sigma}$. Assume that (166) holds for t, and let us prove it for t + 1. We have

$$\begin{split} \|\Delta_{t+1}(A)\| &\leq \frac{\mu^{-2}}{2} \left[\|\widehat{A}\| \|\widehat{\theta}\| + \|\widehat{A}\widehat{\theta} - \mathbf{Z}\| + \|\widehat{A}\| \left[\frac{\|\widehat{\theta}\|}{\sqrt{\lambda}} + \frac{\|\widehat{A}\widehat{\theta} - \mathbf{Z}\|}{2\lambda} \right] \cdot \|\Delta_t(A)\| \right] \\ &\times \left[\frac{\|\widehat{\theta}\|}{\sqrt{\lambda}} + \frac{\|\widehat{A}\widehat{\theta} - \mathbf{Z}\|}{2\lambda} \right] \|\Delta_t(A)\| \\ &\leq \frac{\mu^{-2}}{2} \left[\|\widehat{A}\| \|\widehat{\theta}\| + \|\widehat{A}\widehat{\theta} - \mathbf{Z}\| + \|\widehat{A}\| \left[\frac{\|\widehat{\theta}\|}{\sqrt{\lambda}} + \frac{\|\widehat{A}\widehat{\theta} - \mathbf{Z}\|}{2\lambda} \right] \cdot \|\widehat{A} - \widehat{\Sigma}\| \right] \\ &\times \left[\frac{\|\widehat{\theta}\|}{\sqrt{\lambda}} + \frac{\|\widehat{A}\widehat{\theta} - \mathbf{Z}\|}{2\lambda} \right] \|\Delta_t(A)\|. \end{split}$$

Next, we use the fact that $\widehat{A} \in A$ and $\widehat{\theta} \in \Theta$. The definitions of sets Θ, A imply $\|\widehat{A} - \widehat{\Sigma}\| \leq \sqrt{\lambda}/3$, $\|\widehat{\theta}\| \leq \rho_0 \mu$ and

$$\|\widehat{A}\widehat{\theta} - \mathbf{Z}\| \leq \|\widehat{A} - \widehat{\Sigma}\|\|\widehat{\theta}\| + \|\widehat{\Sigma}\widehat{\theta} - \mathbf{Z}\| \leq \sqrt{\lambda}\|\widehat{\theta}\|/3 + \rho_0\mu\sqrt{\lambda}/3 \leq 2\rho_0\mu\sqrt{\lambda}/3.$$
(167)

It yields

$$\begin{split} \|\Delta_{t+1}(A)\| &\leq \frac{\mu^{-2}}{2} \left[\|\widehat{A}\| \|\widehat{\boldsymbol{\theta}}\| + \rho_0 \mu \sqrt{\lambda}/3 + \|\widehat{A}\| \|\widehat{\boldsymbol{\theta}}\|/3 + \frac{\|\widehat{A}\| \|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\|}{3\sqrt{\lambda}} \right] \cdot \frac{2\rho_0 \mu}{3\sqrt{\lambda}} \cdot \|\Delta_t(A)\| \\ &= \rho_0 \cdot \left[\rho_0/3 + \frac{4\|\widehat{A}\| \|\widehat{\boldsymbol{\theta}}\|}{9\mu\sqrt{\lambda}} + \frac{\|\widehat{A}\| \|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\|}{9\mu\lambda} \right] \|\Delta_t(A)\|. \end{split}$$

Bound (167) does not allow establishing $\|\Delta_t(A)\| \leq \rho_0^2 \|\Delta_t(A)\|$ from the above inequality, which would imply (166). Instead, we use the definition of $(\widehat{\theta}, \widehat{\eta}, \widehat{A})$:

$$\mathcal{L}(\widehat{\boldsymbol{\theta}},\widehat{\boldsymbol{\eta}},\widehat{A}) \leqslant \mathcal{L}(\boldsymbol{\theta}^{\circ},(\widehat{\boldsymbol{\Sigma}}\boldsymbol{\theta}^{\circ}+\mathbf{Z})/2,\widehat{\boldsymbol{\Sigma}}) = \frac{1}{4}\|\widehat{\boldsymbol{\Sigma}}\boldsymbol{\theta}^{\circ}-\mathbf{Z}\|^{2} + \lambda/2\|\boldsymbol{\theta}^{\circ}\|^{2}.$$

Using $\widehat{\boldsymbol{\eta}} = (\widehat{A}\widehat{\boldsymbol{\theta}} + \mathbf{Z})/2$, we get

$$\mathcal{L}(\widehat{\boldsymbol{\theta}},\widehat{\boldsymbol{\eta}},\widehat{A}) = \frac{1}{4} \|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\|^2 + \frac{\mu^2}{2} \|\widehat{A} - \widehat{\Sigma}\| + \frac{\lambda}{2} \|\widehat{\boldsymbol{\theta}}\|,$$

so we have

$$\|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\| \leqslant \|\widehat{\Sigma}\boldsymbol{\theta}^{\circ} - \mathbf{Z}\| + \sqrt{2\lambda}\|\boldsymbol{\theta}^{\circ}\| = \|\mathbf{U}\| + \sqrt{2\lambda}\|\boldsymbol{\theta}^{\circ}\|,$$
(168)

where we used decomposition $\mathbf{Z} = \widehat{\Sigma} \boldsymbol{\theta}^{\circ} + \mathbf{U}$. Therefore, it holds that

$$\|\Delta_{t+1}(A)\| \leq \rho_0 \cdot \left[\rho_0/3 + \frac{4\|\widehat{A}\|\|\widehat{\boldsymbol{\theta}}\|}{9\mu\sqrt{\lambda}} + \frac{\|\widehat{A}\|\|\mathbf{U}\|}{9\mu\lambda} + \frac{2\|\widehat{A}\|\|\boldsymbol{\theta}^\circ\|}{9\mu\sqrt{\lambda}}\right] \cdot \|\Delta_t(A)\|.$$

Under assumptions of the theorem, Proposition H.2 implies the satisfaction of conditions (i)-(iii) of Lemma H.1 and the upper bound

$$\|\mathbf{U}\| \leq 8\sigma \|\Sigma\|^{1/2} \sqrt{\frac{\mathbf{r}(\Sigma) + \log(4/\delta)}{n}}$$
(169)

with probability at least $1 - \delta$. So, we assume that these conditions hold. Then, we have

$$\|\widehat{A}\| \leq \|\widehat{\Sigma}\| + \|\widehat{A} - \widehat{\Sigma}\| \leq 2\|\Sigma\| + \sqrt{\lambda}/3,$$

where we used condition (ii) of Lemma H.1 and the fact that $\widehat{A} \in A$. It yields

$$\|\Delta_{t+1}(A)\| \leq \rho_0 \cdot \left[\rho_0/3 + \frac{8\|\Sigma\|\|\widehat{\boldsymbol{\theta}}\|}{9\mu\sqrt{\lambda}} + \frac{4\|\widehat{\boldsymbol{\theta}}\|}{27\mu} + \frac{2\|\Sigma\|\|\mathbf{U}\|}{9\mu\lambda} + \frac{\|\mathbf{U}\|}{27\mu\sqrt{\lambda}} + \frac{2\|\Sigma\|\|\boldsymbol{\theta}^\circ\|}{9\mu\sqrt{\lambda}}\right] \cdot \|\Delta_t(A)\|.$$

Using the fact that $\hat{\theta} \in \Theta$ and conditions (ii),(iii) of Lemma H.1, we obtain

$$\|\Delta_{t+1}(A)\| \leq \rho_0 \cdot \left[\rho_0/2 + \frac{8\|\Sigma\|\|\widehat{\boldsymbol{\theta}}\|}{9\mu\sqrt{\lambda}}\right] \cdot \|\Delta_t(A)\|$$

From Lemma H.3, we have

$$\|\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\| \leqslant \frac{4\|\widehat{\boldsymbol{\Sigma}} - \boldsymbol{\Sigma}\| (\|\boldsymbol{\theta}^*\| + \|\boldsymbol{\theta}^\circ\|)}{\sqrt{\lambda}} + \frac{2\|\mathbf{U}\|}{\sqrt{\lambda}} \leqslant \frac{8\|\boldsymbol{\theta}^\circ\|}{7} + \frac{2\|\mathbf{U}\|}{\sqrt{\lambda}}$$

where we used $\|\widehat{\Sigma} - \Sigma\| \leq \sqrt{\lambda}/7$ from condition (ii) and $\|\theta^*\| \leq \|\theta^\circ\|$ from Lemma B.2(iv). Thus, we have

$$\|\widehat{\boldsymbol{\theta}}\| \leq \|\boldsymbol{\theta}^*\| + \frac{8\|\boldsymbol{\theta}^\circ\|}{7} + \frac{2\|\mathbf{U}\|}{\sqrt{\lambda}} \leq \frac{15\|\boldsymbol{\theta}^\circ\|}{7} + \frac{2\|\mathbf{U}\|}{\sqrt{\lambda}},\tag{170}$$

and

$$\begin{aligned} \|\Delta_{t+1}(A)\| &\leq \rho_0 \cdot \left[\rho_0/2 + \frac{120\|\Sigma\| \|\boldsymbol{\theta}^\circ\|}{63\mu\sqrt{\lambda}} + \frac{16\|\Sigma\| \|\mathbf{U}\|}{9\mu\lambda}\right] \|\Delta_t(A)\| \\ &\leq \rho_0^2 \|\Delta_t(A)\|, \end{aligned}$$

where we used conditions (i),(iii) of Lemma H.1. Hence, the inequality (166) indeed holds.

Step 4. Final bound. Substituting (166) into (165), we obtain

$$\|\Delta_t(\boldsymbol{\theta})\| \leqslant \rho_0^{2(t-1)} \|\widehat{A} - \widehat{\Sigma}\| \cdot \left[\frac{\|\widehat{\boldsymbol{\theta}}\|}{\sqrt{\lambda}} + \frac{\|\widehat{A}\widehat{\boldsymbol{\theta}} - \mathbf{Z}\|}{2\lambda}\right]$$

Using bounds (168),(170) and $\|\widehat{A} - \widehat{\Sigma}\| \leqslant \sqrt{\lambda}/3$ from the fact that $\widehat{A} \in A$, we obtain

$$\|\Delta_t(\boldsymbol{\theta})\| \leqslant \rho_0^{2(t-1)} \left[2\|\boldsymbol{\theta}^\circ\| + \frac{\|\mathbf{U}\|}{3\sqrt{\lambda}}\right].$$

Replacing t in the inequality above with T and bounding $\|\mathbf{U}\|$ by (169) complete the proof.