

Renormalization group of the gravity coupled with the scalar theory and its effective normalization

Daiki Yamaguchi *

¹*Department of Physics, Kindai University, Higashi-Osaka, Osaka 577-8502, Japan*

²*Research Institute for Science and Technology,
Kindai University, Higashi-Osaka, Osaka 577-8502, Japan*

³*Related with Research Institution of Science and its Departments,
Kyosan University, Kamikaga-Kyoto, Kyoto, Japan*

Abstract

The normalization of the quantum corrected action is resolving the equation divergent dependence of the cutoff towards the system apparent result in quantum gravity. Here we consider the normalization to Einstein R twice scalar action with the cutoff runs from apparent infrared momentu to the ultraviolet momentum. These gravitational actions, Einstein R twice and Einstein R twice with the scalar theory, are come to the ensured apparent system recommendations in classical quantum gravity.

1 Introduction

The quantum correction renormalization is the problem in modern physics. The 1 loop quantum correction and anomalous dimension dependent is researched in many works such as [3] these have a lot of methods for thinking about the renormalization with the functional renormalization group equation. From [3], the gravitational renormalization is used with the local heat kernel expansion and its coefficients are list. The quantum gravity meaned as the field theory with gravitational perturbative expansion is renormalized with the heat kernel expansion of the renormalization group equation, and so, the equation divergence cutoff scale k^2 is absorbed into the running coupling as $G(k \rightarrow \Lambda)$. In moder research [17] is the representative work for renomalized the quantum gravity with matters numbers towards Newton coupling goes to asymptotic safe from the classic G_0 . Fixed point behaves $0 < g_* < 1$.

In this study, we consider the nonlocal heat kernel expansion of the functional renormalization group equation and its expanded result of the quantum gravity which is the gravity coupled with the scalar theory [18], and then, we give the normalize partition function lead the effective action towards the gravitational effective action with background metric and scalar field. Since the gravity and scalar actions are perturbatively expanded to fields quadratic forms, These are summerized in to the field $\delta\Phi$. The Hessians these are gravity coupled scalar and ghost are derived with the decided background metric and scalar. The functional renormalization group equation with non local heat kernel expansion lead to

*e-mail: daichanqg@gmail.com

the effective action with verifications of the heat kernel coefficients and structure functions. Therefore, the gravitational effective actions these are scalar coupled Einstein R^2 action and Einstein R^2 action with no scalar matter.

The normalized actions (3.73) and (3.74) are the gravitational effective action the constants are positively decided as to be sure the appropriate actions with the proofs of the gravity. From these reasons, we are ensure that the normalized gravitational actions are having possibilities to find fits of the gravity datas. Einstein R^2 scalar action or Einstein R^2 action are matched to the needed sophisticated systems. The cosmologies and gravitation proplems are anticipated to be resolved with these Einstein R^2 types actions as the classical quantum gravity.

This paper is list contents as followings. In section 2, we consider the non local heat kernel expansion of the functional renormalization group equation with the scalar theory on the Einstein manifold. The generally also formally master equation and structure functions are verified in this section. In section 3, we consider the scalar gravity expanded to the fields theory as dependent of the laplace operator and that's ghost operator. The non local heat kernel expansion with heat kernel coefficients and structure functions are to the renormalization group equation with gravity coupled with the scalar. The effective action is got by the cutoff scale integration these are $\mu(> \sqrt{z}/2) \rightarrow \Lambda$ [18]. Over the reference, we could give the partition function as Z . The given effective action is normalized with the partition function goes towards Einstein R^2 scalar system with Lugendre transformation on $k[\mu \rightarrow 0^+(> \sqrt{z}/2), \Lambda]$.

2 Functional renormalization group and non local heat kernel expansion

Functional renormalization group equation is the method for verify the couplings beta functions with the local heat kernel expansions [3]. The functional rfenormalization group equation is the equation with the cutoff function. The equation is found by the Wetterich [11]. Wetterich equation or called as the functional renormalization group equation is writte as,

$$\frac{d\Gamma_k}{dt} = \frac{1}{2} \text{Tr} \left(\frac{\delta^2 \Gamma_k}{\delta \phi \delta \phi} + R_k \right)^{-1} \frac{dR_k}{dt}, \quad t = \log k. \quad (2.1)$$

Where we denote the values and functions. k is the cutoff momentum, R_k is the cutoff function, the function $\phi(x)$ is the scalar function on the theoretical space which means the field expression of the theory at stand. This functional renormalization group equation is carry up the field with the integration of the cutoff momentum. We here consider the laplace operator with the factor \mathbf{U} ,

$$\Delta = -D^2 \mathbf{1} + \mathbf{U}, \quad (2.2)$$

where D_μ is the covariant derivative in the space. To consider the expansion of the functional renormalization group equation, we begin with the action as,

$$S(\phi) = \frac{1}{2} \int d^d x \sqrt{g} \phi(x) (\Delta + \omega) \phi(x). \quad (2.3)$$

Be care for we thinking about the field theory as Euclidian space. The value ω is the constant. The cutoff function R_k is given by the eigen value of the laplace operator,

$$R_k = R_k(z \leftrightarrow \Delta). \quad (2.4)$$

$R_k(z)$ runs with $0 < k < \Lambda \rightarrow \infty$. This function is cutoff function set by $k^2 > z$ region. The cutoff action is written as,

$$\Delta S_k = \frac{1}{2} \int d^d x \sqrt{g} \phi(x) R_k(z) \phi(x). \quad (2.5)$$

Therefore total action is given as follows,

$$S(\phi) + \Delta S(\phi) = \frac{1}{2} \int d^d x \sqrt{g} \phi(x) (\Delta + \omega + R_k(\Delta)) \phi(x). \quad (2.6)$$

We use the Eigen value z at the needed situations. The weight \sqrt{g} is apparent written here. The average effective action is given as,

$$\Gamma_k = \frac{1}{2} \log \left[\det(\Delta + \omega + R_k(\Delta)) \right]. \quad (2.7)$$

This derivation is done by the definition of the effective action from the field $\phi(x)$. The functional renormalization group equation is give as,

$$\frac{d\Gamma_k}{dt} = \frac{1}{2} \text{Tr} \frac{\partial_k R_k(\Delta)}{\Delta + \omega + R_k(\Delta)}. \quad (2.8)$$

The inverse propagator with the cutoff function is,

$$h_k(\Delta, \omega) = \frac{\partial_k R_k(\Delta)}{\Delta + \omega + R_k(\Delta)}. \quad (2.9)$$

Here we introduce the laplace parameter z . The (2.8) with (2.9) is the laplace transformed by,

$$\frac{d\Gamma_k}{dt} = \frac{1}{2} \int_0^\infty ds \tilde{h}_k(s) \text{Tr} e^{-s\Delta}. \quad (2.10)$$

z is the real value from the sight of the eigenvalue. From the expression in [18], the heat kernel is expressed by,

$$\begin{aligned} \text{Tr} e^{-s\Delta} = & \frac{1}{(4\pi s)^{d/2}} \int d^d x \sqrt{g} \text{Tr} \left[1 - s\mathbf{U} + s\mathbf{1} \frac{R}{6} \right. \\ & + s^2 [\mathbf{1} R_{\mu\nu} f_{Ric}(s\Delta) R^{\mu\nu} + \mathbf{1} R f_R(s\Delta) R \\ & \left. + R f_{RU}(s\Delta) \mathbf{U} + \mathbf{U} f_U(s\Delta) \mathbf{U} + \Omega_{\mu\nu} f_\Omega(s\Delta) \Omega^{\mu\nu} \right] + \dots \end{aligned} \quad (2.11)$$

Where $\Omega_{\mu\nu}$ is the connection of the gravity coupled field on theoretical space. The space is defined the gravutational manifold S and matter bundle V as $S \times V$ connected with $\Omega_{\mu\nu}$. Structures written as $f(x)$ is given as,

$$f(x) = \int_0^1 d\xi e^{-x\xi(1-\xi)}, \quad (2.12)$$

and,

$$\begin{aligned}
f_{Ric}(x) &= \frac{1}{6x} + \frac{f(x) - 1}{x^2}, \\
f_R(x) &= \frac{f(x)}{32} + \frac{f(x)}{8x} - \frac{7}{48x} - \frac{f(x) - 1}{8x^2}, \\
f_{RU}(x) &= -\frac{f(x)}{4} - \frac{f(x) - 1}{2x}, \\
f_U(x) &= \frac{f(x)}{2}, \\
f_\Omega(x) &= -\frac{f(x) - 1}{2x}.
\end{aligned} \tag{2.13}$$

When we consider Taylor series with $x \ll 1$, we could expand $f(x)$ as,

$$f(x) = 1 - \frac{x}{6} + \frac{x^2}{60} + \mathcal{O}(x^4). \tag{2.14}$$

$f(x)$ is computed easily. Here we use the eigen value expression z and so write down the functional renormalization griup equation as follows,

$$\begin{aligned}
\frac{d\Gamma_k}{dt} &= \frac{1}{2(4\pi)^{d/2}} \int d^d x \sqrt{g} \text{Tr} \left(\mathbf{1} \left[\int_0^\infty ds \tilde{h}_{k,\omega}(s) s^{-d/2} \right] - \mathbf{U} \left[\int_0^\infty ds \tilde{h}_{k,\omega}(s) s^{-d/2+1} \right] \right. \\
&\quad + \mathbf{1} \frac{R}{6} \left[\int_0^\infty ds \tilde{h}_{k,\omega}(s) s^{-d/2+1} \right] + \mathbf{1} R_{\mu\nu} \left[\int_0^\infty ds \tilde{h}_{k,\omega}(s) s^{-d/2+2} f_{Ric}(sz) \right] R^{\mu\nu} \\
&\quad + \mathbf{1} R \left[\int_0^\infty ds \tilde{h}_{k,\omega}(s) s^{-d/2+2} f_R(sz) \right] R + R \left[\int_0^\infty ds \tilde{h}_{k,\omega}(s) s^{-d/2+2} f_{RU}(sz) \right] \mathbf{U} \\
&\quad \left. + \mathbf{U} \left[\int_0^\infty ds \tilde{h}_{k,\omega}(s) s^{-d/2+2} f_U(sz) \right] \mathbf{U} + \Omega_{\mu\nu} \left[\int_0^\infty ds \tilde{h}_{k,\omega}(s) s^{-d/2+2} f_\Omega(sz) \right] \Omega^{\mu\nu} + \dots \right) \tag{2.15}
\end{aligned}$$

Here we introduce $Q_n[f]$ functionals with positive integars $n > 0$. $Q_n[f]$ is given by,

$$Q_n[f] = \int_0^\infty ds s^{-n} \tilde{f}(s) = \frac{1}{\Gamma(n)} \int_0^\infty dw w^{n-1} f(w). \tag{2.16}$$

The last equal is Mellin transformation. We denote that $n = 0$ is,

$$Q_0[f] = f(0). \tag{2.17}$$

In later we understand the $Q_n[f]$ includes resolutions $n = 0, 1, 2, \dots$. Here we back to the definition of Q_n function with $n = 0, 1, 2, 3, \dots$ as parallel transformation a . The functional is given by,

$$Q_n[h_{k,\omega}^a] = \int_0^\infty ds \tilde{h}_{k,\omega}(s) s^{-n} e^{-sa}, \quad n = 0, 1, 2, 3, \dots \tag{2.18}$$

We denote the function $h_{k,\omega}^a$,

$$h_{k,\omega}^a(z) = h_{k,\omega}(z + a). \tag{2.19}$$

The convenient function is introduced as,

$$g_A = \int_0^\infty ds \tilde{h}_{k,\omega}(s) s^{-d/2+2} f_A(sz), \quad n = 0, 1, 2, 3, \dots \quad (2.20)$$

A index is meaning that *Ric*, *R*, *RU*, *U*, Ω . Therefore, the functional renormalization group equation is non local heat kernel expanded as follows,

$$\begin{aligned} \frac{d\Gamma_k}{dt} = & \frac{1}{2} \frac{1}{(4\pi)^{d/2}} \int d^d x \sqrt{g} \left[Q_{d/2}[h_k] \text{Tr} \mathbf{1} + Q_{d/2-1}[h_k] \text{Tr} \left(\frac{R}{6} \mathbf{1} - \mathbf{U} \right) + R_{\mu\nu} g_{Ric} R^{\mu\nu} \text{Tr} \mathbf{1} \right. \\ & \left. + R g_R R \text{Tr} \mathbf{1} + R g_{RU} \text{Tr} \mathbf{U} + \text{Tr}(\mathbf{U} g_U \mathbf{U}) + \text{Tr}(\Omega_{\mu\nu} g_\Omega \Omega^{\mu\nu}) + \dots \right] \end{aligned} \quad (2.21)$$

We wrote $h_k(z) = h_{k,\omega}(z)$. The g_A functions are,

$$\begin{aligned} g_{Ric} &= \frac{1}{6z} Q_{d/2-1}[h_k] - \frac{1}{z^2} Q_{d/2}[h_k] + \frac{1}{z^2} \int_0^1 d\xi Q_{d/2}[h_k^{z\xi(1-\xi)}], \\ g_R &= -\frac{7}{48z} Q_{d/2-1}[h_k] + \frac{1}{8z^2} Q_{d/2}[h_k] + \frac{1}{32} \int_0^1 d\xi Q_{d/2-2}[h_k^{z\xi(1-\xi)}] \\ &+ \frac{1}{8z} \int_0^1 d\xi Q_{d/2-1}[h_k^{z\xi(1-\xi)}] - \frac{1}{8z^2} \int_0^1 d\xi Q_{d/2}[h_k^{z\xi(1-\xi)}], \\ g_{RU} &= \frac{1}{2z} Q_{d/2-1}[h_k] - \frac{1}{4} \int_0^1 d\xi Q_{d/2-2}[h_k^{z\xi(1-\xi)}] - \frac{1}{2z} \int_0^1 d\xi Q_{d/2-1}[h_k^{z\xi(1-\xi)}], \\ g_U &= \frac{1}{2} \int_0^1 d\xi Q_{d/2-2}[h_k^{z\xi(1-\xi)}], \\ g_\Omega &= \frac{1}{2z} Q_{d/2-1}[h_k] - \frac{1}{2z} \int_0^1 d\xi Q_{d/2-1}[h_k^{z\xi(1-\xi)}]. \end{aligned} \quad (2.22)$$

Remaining our work is computing g_A functions with a set of the cutoff functions. To ensure this aim, we use the optimized cutoff function,

$$R_k(z) = (k^2 - z)\theta(k^2 - z) = k^2(1 - \tilde{z})\theta(1 - \tilde{z}). \quad (2.23)$$

Where $\tilde{z} = z/k^2$ is the dimensionless property. When we consider $n > 0$, we could use Mellin transformation. Therefore, $Q_n[h_k]$ is computed as,

$$Q_n[h_k] = \frac{1}{\Gamma(n)} \int_0^\infty dz z^{n-1} \frac{2k^2\theta(k^2 - z)}{z + \omega + (k^2 - z)} = \frac{2k^{2n}}{\Gamma(n+1)(1 + \tilde{\omega})}, \quad n > 0, \quad (2.24)$$

$n = 0$ solution is included the last naturally,

$$Q_n[h_k] = \frac{2k^{2n}}{\Gamma(n+1)(1 + \tilde{\omega})}, \quad n = 0, 1, 2, \dots \quad (2.25)$$

Similarly, we compute $Q_n[h_k^{z\xi(1-\xi)}]$. Thinking about $n > 0$, the functional is,

$$\begin{aligned} Q_n[h_k^{z\xi(1-\xi)}] &= \frac{1}{\Gamma(n)} \int_0^\infty dx x^{n-1} \frac{2k^2}{k^2 + \omega} \theta(k^2 - x - z\xi(1-\xi)) \\ &= \frac{1}{\Gamma(n)} \int_0^{k^2 - z\xi(1-\xi)} dx x^{n-1} \frac{2k^2}{k^2 + \omega} \theta(k^2 - z\xi(1-\xi)) \\ &= \frac{1}{\Gamma(n+1)} \frac{2k^2}{k^2 + \omega} (k^2 - z\xi(1-\xi))^n \theta(k^2 - z\xi(1-\xi)). \end{aligned} \quad (2.26)$$

We know that the $n = 0$ is included in above. The $Q_n[h_k^{z\xi(1-\xi)}]$ with $n = 0, 1, 2, 3, \dots$ is given by,

$$Q_n[h_k^{z\xi(1-\xi)}] = \frac{1}{\Gamma(n+1)} \frac{2k^2}{k^2 + \omega} (k^2 - z\xi(1-\xi))^n \theta(k^2 - z\xi(1-\xi)), \quad n = 0, 1, 2, 3, \dots \quad (2.27)$$

(2.27) is integrated from $\xi = 0$ to $\xi = 1$. To do the integration of (2.27), we consider the integration contour with $\xi - f(\xi)$ coordinate. Cross points are given by,

$$f(\xi) = k^2 - z\xi(1-\xi) = 0, \quad (2.28)$$

we find points α, β as,

$$\alpha = \frac{1}{2} \left[1 - \sqrt{1 - \frac{4k^2}{z} \theta(z - 4k^2)} \right], \quad \beta = \frac{1}{2} \left[1 + \sqrt{1 - \frac{4k^2}{z} \theta(z - 4k^2)} \right]. \quad (2.29)$$

The integration is carried with $k^2 - z\xi(1-\xi) > 0$. This means that the integration region is $0 \leq \xi \leq \alpha$ and $\beta \leq \xi \leq 1$. From these reasons, we find $Q_n[h_k^{z\xi(1-\xi)}]$ integration as follows,

$$\begin{aligned} \int_0^1 d\xi Q_0[h_k^{z\xi(1-\xi)}] &= \frac{2k^2}{k^2 + \omega} \left[1 - \sqrt{1 - \frac{4k^2}{z} \theta(z - 4k^2)} \right], \\ \int_0^1 d\xi Q_1[h_k^{z\xi(1-\xi)}] &= \frac{2k^4}{k^2 + \omega} \left[1 - \frac{z}{6k^2} + \frac{z}{6k^2} \left(1 - \frac{4k^2}{z} \right)^{3/2} \theta(z - 4k^2) \right], \\ \int_0^1 d\xi Q_2[h_k^{z\xi(1-\xi)}] &= \frac{k^6}{k^2 + \omega} \left[1 - \frac{z}{3k^2} + \frac{z^2}{30k^4} - \frac{z^2}{30k^4} \left(1 - \frac{4k^2}{z} \right)^{5/2} \theta(z - 4k^2) \right]. \end{aligned} \quad (2.30)$$

g_A functions are calculated straightforwardly,

$$\begin{aligned} g_{Ric} &= \frac{1}{30} \frac{1}{1 + \tilde{\omega}} \left[1 - \left(1 - \frac{4}{\tilde{z}} \right)^{5/2} \theta(\tilde{z} - 4) \right], \\ g_R &= \frac{1}{1 + \tilde{\omega}} \left[\frac{1}{60} - \frac{1}{16} \left(1 - \frac{4}{\tilde{z}} \right)^{1/2} \theta(\tilde{z} - 4) \right. \\ &\quad \left. + \frac{1}{24} \left(1 - \frac{4}{\tilde{z}} \right)^{3/2} \theta(\tilde{z} - 4) + \frac{1}{240} \left(1 - \frac{4}{\tilde{z}} \right)^{5/2} \theta(\tilde{z} - 4) \right], \\ g_{RU} &= \frac{1}{1 + \tilde{\omega}} \left[-\frac{1}{3} + \frac{1}{2} \left(1 - \frac{4}{\tilde{z}} \right)^{1/2} \theta(\tilde{z} - 4) - \frac{1}{6} \left(1 - \frac{4}{\tilde{z}} \right)^{3/2} \theta(\tilde{z} - 4) \right], \\ g_U &= \frac{1}{1 + \tilde{\omega}} \left[1 - \left(1 - \frac{4}{\tilde{z}} \right)^{1/2} \theta(\tilde{z} - 4) \right], \\ g_\Omega &= \frac{1}{6} \frac{1}{1 + \tilde{\omega}} \left[1 - \left(1 - \frac{4}{\tilde{z}} \right)^{3/2} \theta(\tilde{z} - 4) \right]. \end{aligned} \quad (2.31)$$

We derived master equation of the functional renormalization group equation with the non local heat kernel expansion. The master equation is convenient to verify the equation with the fields setups. The results are anticipated to have the coupled matters fields in the manifold quantum gravity. The sophisticated adjoints of the heat kernel coefficients and structures in stands after discussions.

3 Gravity coupled with the scalar and appatrent normalized effective action

In this section we compute the effective action of the gravity coupled with scalar field theory in [18]. After that, we normalized this effective action towards normalized effective action that would be matched with the appropriate tests confirmations in fine signs. The gravity coupled with the scalar theory is expanded as fluctuation perturbative expansions used in [3, 18]. And so, we consider the fields theory contributions with background metric and scalar field are fixed. The functional renormalization laplace operators are found by the fields setups include the ghost action. Heat kernel coefficients and structures are calculated to results, by the way, the ghost coefficients amplitudes are adjusted as adjoints. The effective action in $k^2 > z/4 = -\nabla^2/4$ to the ultraviolet area with $d = 4$ in [18]. Finally, we could have the normalization partition function makes the effective action runs from $\mu^2 \rightarrow 0^{+2} > z/4$ to Λ^2 normalized into the Einstein R^2 action includes the coupled interaction from Lugendre transformation. We recognized that Einstein R^2 interacted action ensures the gravitational physics as the interacted gravity, parts are given in the interacted terms.

3.1 Running effective action of the gravity and scalar field

We begin with the preparations of the actions. Einstein action is written by,

$$S_H[g + h] = -\frac{1}{\kappa} \int d^d x \sqrt{g + h} R(g + h). \quad (3.1)$$

κ is defined as,

$$\kappa = 16\pi G_0. \quad (3.2)$$

The scalar action is given by,

$$S_m[g + h, \phi] = \int d^d x \sqrt{g + h} \left[\frac{1}{2} (g + h)^{\mu\nu} \nabla_\mu \phi \nabla_\nu \phi + V(\phi) \right]. \quad (3.3)$$

When we consider the fields theory, we should be gauge fixed with the appropriate gauge function. We use the gauge function called as De-Donder gauge function,

$$\chi_\mu = \nabla^\nu h_{\mu\nu} - \frac{1}{2} \nabla_\mu h. \quad (3.4)$$

The gauge fixing is expressed as the action,

$$S_{gf}[h, g] = \frac{1}{2} \int d^d x \sqrt{g} \chi_\mu \chi^\mu. \quad (3.5)$$

The gauge fixing is done with the gauge fixed action above. The gauge invariance is breaking with the selection of the gauge function. To recover the invariance, we introduce the possibility of the ghost action written by,

$$S_{gh}[\bar{C}, C, g] = \int d^d x \sqrt{g} \bar{C}_\mu (-\nabla^2 \delta^\mu_\nu - R^\mu_\nu) C^\nu. \quad (3.6)$$

The ghost laplace operator is,

$$(\Delta_{gh})^\mu{}_\nu = -\nabla^2 \delta^\mu{}_\nu - R^\mu{}_\nu. \quad (3.7)$$

Here we consider perturbative expansion of the scalar coupled gravity. Now we consider the perturbative expansion of the gravity as,

$$S_H(g+h) + S_{gf}(g+h) = S_H(g) + \frac{1}{2\kappa} \int d^d x \sqrt{g} h_{\alpha\beta} \mathcal{H}^{\alpha\beta\mu\nu} h_{\mu\nu}. \quad (3.8)$$

The gravitational hessian is given by,

$$\mathcal{H}^{\mu\nu}{}_{\rho\sigma} = K^{\mu\nu}{}_{\rho\sigma}(-\nabla^2) + u^{\mu\nu}{}_{\rho\sigma}. \quad (3.9)$$

De-Witt metric $K^{\mu\nu}{}_{\rho\sigma}$ and $u^{\mu\nu}{}_{\rho\sigma}$ are given by,

$$K^{\alpha\beta\mu\nu} = \frac{1}{2} \left(\delta^{\alpha\beta, \mu\nu} - \frac{1}{2} g^{\alpha\beta} g^{\mu\nu} \right) = \frac{1}{4} (g^{\mu\alpha} g^{\nu\beta} + g^{\nu\alpha} g^{\mu\beta} - g^{\alpha\beta} g^{\mu\nu}), \quad (3.10)$$

and so on,

$$u^{\mu\nu}{}_{\rho\sigma} = K^{\mu\nu}{}_{\rho\sigma} R + \frac{1}{2} (R^{\mu\nu} g_{\rho\sigma} + R_{\rho\sigma} g^{\mu\nu}) - \delta^{\mu(\rho} R^{\nu)\sigma} - R^{\mu(\rho}{}^{\nu)\sigma}. \quad (3.11)$$

The matter action is expressed as,

$$S_m(g+h, \phi + \delta\phi) = \int d^d x \sqrt{g+h} \left[\frac{1}{2} (g^{\mu\nu} - h^{\mu\nu} + h^\mu{}_\rho h^{\rho\nu}) \nabla_\mu (\phi + \delta\phi) \nabla_\nu (\phi + \delta\phi) + V(\phi) + V'(\phi) \delta\phi + \frac{1}{2} V''(\phi) \delta\phi^2 \right]. \quad (3.12)$$

The quadratic fields terme are remained because of the functional renormalization group equation detected by the twice any fields derivatives. Therefore, we should consider the contributions of the perturbative expansions as the quadratic fields. As we consider the perturbative expansions of the scalar coupled gravity, the recomanded form is the ,

$$\frac{1}{2\kappa} \int d^d x \sqrt{g} \delta\Phi^T \mathbf{H} \delta\Phi, \quad (3.13)$$

with field definition,

$$\delta\Phi = \begin{pmatrix} h_{\mu\nu} \\ \delta\phi \end{pmatrix}. \quad (3.14)$$

Fields are used as the vector notation and The hessian \mathbf{H} is treated as the matrix as,

$$\mathbf{H} = \begin{pmatrix} H^{\alpha\beta\mu\nu} & H^{\alpha\beta\cdot} \\ H^{\cdot\mu\nu} & H^{\cdot\cdot} \end{pmatrix}. \quad (3.15)$$

The hessian is written as,

$$\mathbf{H} = \mathbf{K}(-\nabla^2) + 2\mathbf{V}^\delta \nabla_\delta + \mathbf{U}. \quad (3.16)$$

Matrices \mathbf{K} , \mathbf{V}^δ , \mathbf{U} are computed towards,

$$\mathbf{K} = \begin{pmatrix} K^{\alpha\beta\mu\nu} & 0 \\ 0 & \kappa \end{pmatrix}, \quad (3.17)$$

continually,

$$\mathbf{V}^\delta = \begin{pmatrix} 0 & -\kappa K^{\alpha\beta\gamma\delta} \nabla_\gamma \phi \\ \kappa K^{\mu\nu\gamma\delta} \nabla_\gamma \phi & 0 \end{pmatrix}, \quad (3.18)$$

$$\mathbf{U} = \begin{pmatrix} U^{\alpha\beta\mu\nu} & \frac{1}{2}\kappa g^{\alpha\beta} V'(\phi) \\ 2\kappa K^{\mu\nu\gamma\delta} \nabla_\gamma \nabla_\delta \phi + \frac{1}{2}\kappa g^{\mu\nu} V'(\phi) & \kappa V''(\phi) \end{pmatrix}. \quad (3.19)$$

Where, $K^{\alpha\beta\mu\nu}$ is the De-Witt metric. The $U^{\alpha\beta\mu\nu}$ is expanded as follows,

$$\begin{aligned} U^{\alpha\beta\mu\nu} = & K^{\alpha\beta\mu\nu} R + \frac{1}{2}(g^{\mu\nu} R^{\alpha\beta} + R^{\mu\nu} g^{\alpha\beta}) - \frac{1}{4}(g^{\alpha\mu} R^{\beta\nu} + g^{\alpha\nu} R^{\beta\mu} + g^{\beta\mu} R^{\alpha\nu} + g^{\beta\nu} R^{\alpha\mu}) \\ & - \frac{1}{2}(R^{\alpha\mu\beta\nu} + R^{\alpha\nu\beta\mu}) + \kappa \left[-\frac{1}{2} K^{\alpha\beta\mu\nu} (\nabla\phi)^2 - \frac{1}{4}(g^{\alpha\beta} \nabla^\mu \phi \nabla^\nu \phi + g^{\mu\nu} \nabla^\alpha \phi \nabla^\beta \phi) \right. \\ & \left. + \frac{1}{4}(g^{\alpha\mu} \nabla^\beta \phi \nabla^\nu \phi + g^{\alpha\nu} \nabla^\beta \phi \nabla^\mu \phi + g^{\beta\mu} \nabla^\alpha \phi \nabla^\nu \phi + g^{\beta\nu} \nabla^\alpha \phi \nabla^\mu \phi) - K^{\alpha\beta\mu\nu} V \right]. \end{aligned} \quad (3.20)$$

From this hessian, we find the laplace operator as new relation. \mathbf{K}^{-1} is prepared as,

$$\mathbf{K}^{-1} = \begin{pmatrix} K_{\alpha\beta\mu\nu}^{-1} & 0 \\ 0 & \frac{1}{\kappa} \end{pmatrix}, \quad (3.21)$$

where we write the inverse De-Witt metric,

$$K_{\alpha\beta\mu\nu}^{-1} = 2\delta_{\alpha\beta,\mu\nu} - \frac{2}{d-2} g_{\alpha\beta} g_{\mu\nu}. \quad (3.22)$$

The laplace operator is defined as $\mathbf{K}^{-1}\mathbf{H}$ as,

$$\Delta = \mathbf{I}(-\nabla^2) + 2\mathbf{Y}^\delta \nabla_\delta + \mathbf{W}. \quad (3.23)$$

Also \mathbf{I} is the unit matrix, matrixs \mathbf{Y}^δ and \mathbf{W} are calculated in following results. \mathbf{Y}^δ is written as,

$$\mathbf{Y}^\delta = \begin{pmatrix} 0 & -\kappa \delta_{\alpha\beta} \gamma^\delta \nabla_\gamma \phi \\ K^{\mu\nu\gamma\delta} \nabla_\gamma \phi & 0 \end{pmatrix}. \quad (3.24)$$

\mathbf{W} is computed towards,

$$\mathbf{W} = \begin{pmatrix} W_{\alpha\beta}{}^{\mu\nu} & -\frac{2}{d-2} \kappa g_{\alpha\beta} V'(\phi) \\ 2K^{\mu\nu\gamma\delta} \nabla_\gamma \nabla_\delta \phi + \frac{1}{2} g^{\mu\nu} V'(\phi) & V''(\phi) \end{pmatrix}, \quad (3.25)$$

where we write $W_{\alpha\beta}{}^{\mu\nu}$,

$$W_{\alpha\beta}{}^{\mu\nu} = 2U_{\alpha\beta}{}^{\mu\nu} - \frac{d-4}{d-2} g_{\alpha\beta} \left[R^{\mu\nu} - \frac{1}{2} R g^{\mu\nu} - \frac{\kappa}{2} (\nabla^\mu \phi \nabla^\nu \phi - \frac{1}{2} g^{\mu\nu} (\nabla\phi)^2) \right] - \frac{\kappa}{2} g_{\alpha\beta} g^{\mu\nu} V. \quad (3.26)$$

This laplace operator is the different form from the definition. Then, we redefine the covariant derivative as,

$$\tilde{\nabla}_\mu = \nabla_\mu \mathbf{I} - \mathbf{Y}_\mu. \quad (3.27)$$

The laplace operator is expressed as,

$$\Delta = -\tilde{\nabla}_\mu \tilde{\nabla}^\mu + \tilde{\mathbf{W}}, \quad \tilde{\mathbf{W}} = \mathbf{W} - \nabla_\mu \mathbf{Y}^\mu + \mathbf{Y}_\mu \mathbf{Y}^\mu. \quad (3.28)$$

To compute the $\tilde{\mathbf{W}}$, we prepare some parts of the laplace operator. $\nabla_\mu \mathbf{Y}^\mu$ is,

$$\nabla_\mu \mathbf{Y}^\mu = \begin{pmatrix} 0 & -\kappa \nabla_\alpha \nabla_\beta \phi \\ K^{\mu\nu\gamma\delta} \nabla_\delta \nabla_\gamma \phi & 0 \end{pmatrix}. \quad (3.29)$$

$\mathbf{Y}_\mu \mathbf{Y}^\mu$ is the following result,

$$\mathbf{Y}_\mu \mathbf{Y}^\mu = \begin{pmatrix} -\kappa \delta_{\alpha\beta} \gamma^\delta K^{\mu\nu\epsilon} \nabla_\gamma \phi \nabla_\epsilon \phi & 0 \\ 0 & -\kappa (\nabla \phi)^2 \end{pmatrix}. \quad (3.30)$$

We reach the $\tilde{\mathbf{W}}$ into,

$$\tilde{\mathbf{W}} = \begin{pmatrix} W_{\alpha\beta}{}^{\mu\nu} - \kappa \delta_{\alpha\beta} \gamma^\delta K^{\mu\nu\epsilon} \nabla_\gamma \phi \nabla_\epsilon \phi & -\frac{2}{d-2} \kappa g_{\alpha\beta} V' + \kappa \nabla_\alpha \nabla_\beta \phi \\ K^{\mu\nu\gamma\delta} \nabla_\gamma \nabla_\delta \phi + \frac{1}{2} g^{\mu\nu} V' & V'' - \kappa (\nabla \phi)^2 \end{pmatrix}. \quad (3.31)$$

Components of $\tilde{\mathbf{W}}$ are given by,

$$\begin{aligned} A_{\alpha\beta}{}^{\mu\nu} &= W_{\alpha\beta}{}^{\mu\nu} - \frac{1}{2} \kappa \delta^{(\mu} (\nabla_{\alpha} \nabla_{\beta)} \nabla^{\nu)} \phi + \frac{1}{4} \kappa g^{\mu\nu} \nabla_\alpha \phi \nabla_\beta \phi, \\ B_{\alpha\beta} &= -\frac{2}{d-2} \kappa g_{\alpha\beta} V' + \kappa \nabla_\alpha \nabla_\beta \phi, \\ C^{\mu\nu} &= K^{\mu\nu\alpha\beta} \nabla_\alpha \nabla_\beta \phi + \frac{1}{2} g^{\mu\nu} V', \\ D &= -\kappa (\nabla \phi)^2 + V''. \end{aligned} \quad (3.32)$$

From here, we consider the curvature expression. The curvature is calculated to,

$$\begin{aligned} \tilde{\Omega}_{\mu\nu} &= [\nabla_\mu - \mathbf{Y}_\mu, \nabla_\nu - \mathbf{Y}_\nu] \\ &= \Omega_{\mu\nu} - \nabla_\mu \mathbf{Y}_\nu + \nabla_\nu \mathbf{Y}_\mu + [\mathbf{Y}_\mu, \mathbf{Y}_\nu]. \end{aligned} \quad (3.33)$$

Some parts of terms above are verified towards,

$$\nabla_{[\rho} \mathbf{Y}_{\sigma]} = \begin{pmatrix} 0 & -\kappa \delta_{\alpha\beta} \gamma^\delta g_{[\sigma|\delta} \nabla_{\rho]} \nabla_\gamma \phi \\ K^{\mu\nu\gamma\delta} g_{[\sigma|\delta} \nabla_{\rho]} \nabla_\gamma \phi & 0 \end{pmatrix}, \quad (3.34)$$

$$\mathbf{Y}_{[\rho} \mathbf{Y}_{\sigma]} = \begin{pmatrix} -\kappa \delta_{\alpha\beta, \gamma[\rho} K^{\mu\nu}{}_{|\epsilon|\sigma]} \nabla^\gamma \phi \nabla^\epsilon \phi & 0 \\ 0 & 0 \end{pmatrix}. \quad (3.35)$$

Components of the curvature matrix are to come results as follows,

$$\begin{aligned} (\tilde{\Omega}_{\rho\sigma})_{\alpha\beta}{}^{\mu\nu} &= (\Omega_{\rho\sigma})_{\alpha\beta}{}^{\mu\nu} - 2\kappa \delta_{\alpha\beta, \gamma[\rho} K^{\mu\nu}{}_{|\epsilon|\sigma]} \nabla^\gamma \phi \nabla^\epsilon \phi, \\ (\tilde{\Omega}_{\rho\sigma})_{\alpha\beta} &= 2\kappa \delta_{\alpha\beta} \gamma^\delta g_{[\sigma|\delta} \nabla_{\rho]} \nabla_\gamma \phi, \\ (\tilde{\Omega}_{\rho\sigma})^{\mu\nu} &= -2K^{\mu\nu\gamma\delta} g_{[\sigma|\delta} \nabla_{\rho]} \nabla_\gamma \phi. \end{aligned} \quad (3.36)$$

We prepared the laplace operator properties and curvature properties already. Begin to expand the functional renormalization group equation with the non local heat kernel expressions.

We consider heat kernel coefficients and non local structure terms with calculation results of the needed properties. $d = 4$ is applied in latter here. Remember the laplace operator of scalar coupling gravity means that $\widetilde{\mathbf{W}}$ is the unique tensor. Then, we compute some needs. The trace is,

$$\text{Tr}\widetilde{\mathbf{W}} = A_{\mu\nu}{}^{\mu\nu} + D, \quad (3.37)$$

where,

$$A_{\mu\nu}{}^{\mu\nu} = 6R - \kappa(\nabla\phi)^2 - 10\kappa V. \quad (3.38)$$

$\text{Tr}\widetilde{\mathbf{W}}^2$ is calculated as,

$$\text{Tr}\widetilde{\mathbf{W}}^2 = A_{\mu\nu}{}^{\alpha\beta} A_{\alpha\beta}{}^{\mu\nu} + 2B_{\alpha\beta} C^{\alpha\beta} + D^2, \quad (3.39)$$

where,

$$\begin{aligned} A_{\mu\nu}{}^{\alpha\beta} A_{\alpha\beta}{}^{\mu\nu} &= 3R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 6R_{\mu\nu} R^{\mu\nu} + 5R^2 - \frac{3}{2}\kappa R(\nabla\phi)^2 - 12\kappa V R \\ &+ \frac{7}{4}\kappa^2((\nabla\phi)^2)^2 + 2\kappa^2 V(\nabla\phi)^2 + 10\kappa^2 V^2, \end{aligned} \quad (3.40)$$

$$B_{\alpha\beta} C^{\alpha\beta} = \kappa V' \nabla^2 \phi - 2\kappa V'^2 + \frac{1}{2}\kappa \nabla_\mu \nabla_\nu \phi \nabla^\mu \nabla^\nu \phi - \frac{1}{4}\kappa(\nabla^2 \phi)^2. \quad (3.41)$$

Continually, curvature notations are computed as follows,

$$\text{Tr}\widetilde{\Omega}_{\mu\nu} \widetilde{\Omega}^{\mu\nu} = (\widetilde{\Omega}_{\mu\nu})_{\alpha\beta}{}^{\gamma\delta} (\widetilde{\Omega}^{\mu\nu})_{\gamma\delta}{}^{\alpha\beta} + 2(\widetilde{\Omega}_{\mu\nu})_{\alpha\beta} (\widetilde{\Omega}^{\mu\nu})^{\alpha\beta}, \quad (3.42)$$

where,

$$(\widetilde{\Omega}_{\mu\nu})_{\alpha\beta}{}^{\gamma\delta} (\widetilde{\Omega}^{\mu\nu})_{\gamma\delta}{}^{\alpha\beta} = -6R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} + \kappa R(\nabla\phi)^2 + 2\kappa R_{\mu\nu} \nabla^\mu \phi \nabla^\nu \phi - \frac{3}{2}\kappa^2((\nabla\phi)^2)^2, \quad (3.43)$$

$$2(\widetilde{\Omega}_{\mu\nu})_{\alpha\beta} (\widetilde{\Omega}^{\mu\nu})^{\alpha\beta} = \kappa(\nabla^2 \phi)^2 - 4\kappa \nabla_\mu \nabla_\nu \phi \nabla^\mu \nabla^\nu \phi. \quad (3.44)$$

To simplify the heatkernel coefficients and structures, we use below relation,

$$R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} = 4R_{\mu\nu} R^{\mu\nu} - R^2. \quad (3.45)$$

Heat kernel coefficients of the scalar coupling gravity are list as,

$$\text{Tr}\mathbf{1} = 11, \quad \text{Tr}\mathbf{1} \frac{R}{6} - \text{Tr}\widetilde{\mathbf{W}} = -\frac{25}{6}R + 2\kappa(\nabla\phi)^2 - V'' + 10\kappa V. \quad (3.46)$$

The ghost heat kernel coefficients are computed as,

$$\text{Tr}\mathbf{1}_{gh} = 4, \quad \text{Tr}\mathbf{1}_{gh} \frac{R}{6} - \text{Tr}\mathbf{U}_{gh} = \frac{5}{3}RC_{gh,2} = \frac{1}{3}R. \quad (3.47)$$

The trace is adjusted with the amplitude constant $C_{gh,2} = \frac{1}{5}$. Remaining our work is calculating non local structures. The non local structures of the gravity and scalar are,

$$\begin{aligned}
& R_{\mu\nu}(11f_{Ric} + 6f_U - 24f_\Omega)R^{\mu\nu} + R(11f_R + 6f_{RU} + 2f_U + 6f_\Omega)R \\
& + \kappa R(\nabla\phi)^2 \left(-2f_{RU} - \frac{3}{2}f_U + f_\Omega \right) + 2\kappa R^{\mu\nu} \nabla_\mu \phi \nabla_\nu \phi f_\Omega + RV'' f_{RU} + \kappa RV(-10f_{RU} - 12f_U) \\
& + \kappa \nabla_\mu \nabla_\nu \phi \nabla^\mu \nabla^\nu \phi (f_U - 4f_\Omega) + \kappa (\nabla^2 \phi)^2 \left(-\frac{1}{2}f_U + f_\Omega \right) + 2\kappa \nabla^2 \phi V' f_U - 2\kappa V'' (\nabla\phi)^2 f_U \\
& + 2\kappa^2 V (\nabla\phi)^2 f_U - 4\kappa V' V' f_U + V'' V'' f_U + \kappa^2 ((\nabla\phi)^2)^2 \left(\frac{11}{4}f_U - \frac{3}{2}f_\Omega \right) + 10\kappa^2 V V f_U. \quad (3.48)
\end{aligned}$$

The non local structures of the ghost are,

$$R_{\mu\nu}(4f_{Ric} + f_U - 4f_\Omega)R^{\mu\nu} + R(4f_R - f_{RU} + f_\Omega)R. \quad (3.49)$$

From these results, we know used heat kernel coefficients and non local structure as expansions of the functional renormalization group. Therefore, we expand the functional renormalization group equation below with related properties,

$$\frac{d\Gamma_k}{dt} = \frac{1}{2} \text{Tr} \frac{\partial_t R_k(\Delta)}{\Delta + R_k(\Delta)} - \text{Tr} \frac{\partial_t R_k(\Delta_{gh})}{\Delta_{gh} + R_k(\Delta_{gh})}. \quad (3.50)$$

Using (3.46), (3.47), (3.48) and (3.49), we found the functional renormalization group equation as follows,

$$\begin{aligned}
\frac{d\Gamma_k}{dt} = \frac{1}{32\pi^2} \int d^4x \sqrt{g} & \left[3k^4 + k^2 \left(-\frac{29}{3}R + 4\kappa(\nabla\phi)^2 - 2V'' + 20\kappa V \right) + R_{\mu\nu} g_1 R^{\mu\nu} + R g_2 R \right. \\
& + \kappa R(\nabla\phi)^2 g_3 + \kappa R_{\mu\nu} \nabla^\mu \phi \nabla^\nu \phi g_4 + RV'' g_5 + \kappa RV g_6 + \kappa \nabla_\mu \nabla_\nu \phi \nabla^\mu \nabla^\nu \phi g_7 \\
& + \kappa ((\nabla^2 \phi))^2 g_8 + \kappa^2 ((\nabla\phi)^2)^2 g_9 + \kappa \nabla^2 \phi V' g_{10} - \kappa (\nabla\phi)^2 V'' g_{10} + \kappa^2 (\nabla\phi)^2 V g_{10} \\
& \left. - 2\kappa V' V' g_{10} + \frac{1}{2} V'' V'' g_{10} + 5\kappa^2 V V g_{10} \right]. \quad (3.51)
\end{aligned}$$

Where we defined $g_a, a = 1, 2, 3, 4, 5, 6, 7, 8, 9, 10$ functions. These structures are given by,

$$g_1 = 3f_{Ric} + 4g_U - 16g_\Omega, \quad (3.52)$$

$$g_2 = 3g_R + 2g_U + 8g_{RU} + 4g_\Omega, \quad (3.53)$$

$$g_3 = -2g_{RU} - \frac{3}{2}g_U + g_\Omega, \quad (3.54)$$

$$g_4 = 2g_\Omega, \quad (3.55)$$

$$g_5 = g_{RU}, \quad (3.56)$$

$$g_6 = -10g_{RU} - 12g_U, \quad (3.57)$$

$$g_7 = g_U - 4g_\Omega, \quad (3.58)$$

$$g_8 = -\frac{1}{2}g_U + g_\Omega, \quad (3.59)$$

$$g_9 = \frac{11}{4}g_U - \frac{3}{2}g_\Omega, \quad (3.60)$$

$$g_{10} = 2g_U. \quad (3.61)$$

Here we consider to derive the effective action with results of structures g_a , $a = 1, 2, 3, 4, 5, 6, 7, 8, 9, 10$. To ensure our purposes, g -structures are calculated with $k^2 > z/4$,

$$g_1 = \frac{43}{30}, \quad g_2 = \frac{1}{20}, \quad g_3 = -\frac{2}{3}, \quad g_4 = \frac{1}{3}, \quad g_5 = -\frac{1}{3}, \quad (3.62)$$

$$g_6 = -\frac{26}{3}, \quad g_7 = \frac{1}{3}, \quad g_8 = -\frac{1}{3}, \quad g_9 = \frac{5}{2}, \quad g_{10} = 2. \quad (3.63)$$

We are assuming that the $z > 0$ and $k^2 > z/4$ runnings. We set the integration region $k[\mu \rightarrow 0^+ (> \sqrt{z}/2), \Lambda]$ as the ultraviolet cutoff momentum Λ , and for, we could write down the effective action with needed sources,

$$\begin{aligned} \Gamma_\Lambda[g] = \frac{1}{32\pi^2} \int d^4x \sqrt{g} & \left[3\Lambda^4 + \left(-\frac{29}{3}R + 4\kappa(\nabla\phi)^2 - 2V'' + 20\kappa V \right) \Lambda^2 \right. \\ & + \left(\frac{43}{30}R_{\mu\nu}R^{\mu\nu} + \frac{1}{20}R^2 - \frac{2}{3}\kappa R(\nabla\phi)^2 - \frac{26}{3}\kappa RV - \frac{1}{3}RV'' \right. \\ & \quad \left. + \frac{5}{2}\kappa^2((\nabla\phi)^2)^2 + 2\kappa^2V(\nabla\phi)^2 - 4\kappa V''(\nabla\phi)^2 \right. \\ & \quad \left. + 10\kappa^2V^2 - 4\kappa V'^2 + V''^2 \right) \log\left(\frac{\Lambda^2}{\mu^2}\right) + \text{finites} \left. \right]. \quad (3.64) \end{aligned}$$

Be care for the renormalization scale μ^2 runs to $0^{+2} > z/4$. The full effective action is written by some apparent descriptions from the finite terms.

3.2 Normalization of the effective action

The effective action including renormalization scale and cutoff. We are normalize this effective action (3.64) with the definition of quantum corrected action being equal to the effective action normalized by the cutoff path integration normalization as,

$$Z = \int_{\mu \rightarrow 0^+}^{\Lambda} (dk) N[g, \phi] \exp(-S_k[g, \phi]), \quad \mu^2 \rightarrow 0^{+2} > z/4. \quad (3.65)$$

The path integral is the summed ups of the cutoff scale running from $\mu \rightarrow \Lambda$ as the definition of the total partition function of Z . The quantum corrected action is the effective action (3.64) as,

$$\begin{aligned} S_k[g, \phi] = \frac{1}{32\pi^2} \int d^4x \sqrt{g} & \left[3k^4 + \left(-\frac{29}{3}R + 4\kappa(\nabla\phi)^2 - 2V'' + 20\kappa V \right) k^2 \right. \\ & + \left(\frac{43}{30}R_{\mu\nu}R^{\mu\nu} + \frac{1}{20}R^2 - \frac{2}{3}\kappa R(\nabla\phi)^2 - \frac{26}{3}\kappa RV - \frac{1}{3}RV'' \right. \\ & \quad \left. + \frac{5}{2}\kappa^2((\nabla\phi)^2)^2 + 2\kappa^2V(\nabla\phi)^2 - 4\kappa V''(\nabla\phi)^2 \right. \\ & \quad \left. + 10\kappa^2V^2 - 4\kappa V'^2 + V''^2 \right) \log\left(\frac{k^2}{\mu^2 \rightarrow 0^{+2} (> z/4)}\right) + \text{finites} \left. \right]. \quad (3.66) \end{aligned}$$

The normalize factor $N[g, \phi]$ is the important property to make Z towards the basic fixed exponential part. To do so, we consider the factor as,

$$N[g, \phi] = \frac{N_0[g, \phi]}{N_v[g, \phi]}. \quad (3.67)$$

$N_0[g, \phi]$ is the basic exponential as to include the bare action having Einstein scalar coupling action. $N_v[g, \phi]$ is take $\int_{\mu \rightarrow 0^+ (> \sqrt{z}/2)}^{\Lambda} (dk) \exp(-S_k[g, \phi])$ towards 1. It isn't permitted to make $N_0[g, \phi]$ arbitrarily. $N_0[g, \phi]$ should includes every relating action terms with thinking about the well defined action in gravity. The quantum corrected action is true in (3.66), and on, we could derive the $N_0[g, \phi]$ as follows,

$$N_0[g, \phi] = \exp \left[-\frac{1}{32\pi^2} \int d^4x \sqrt{g} \left[(const.) + \left(-\frac{29}{3}RC_1 + 4\kappa(\nabla\phi)^2C_2 - 2V''C_3 + 20\kappa VC_4 \right) \right. \right. \\ \left. \left. + \left(\frac{43}{30}R_{\mu\nu}R^{\mu\nu}C_a + \frac{1}{20}R^2C_b - \frac{2}{3}\kappa R(\nabla\phi)^2C_c - \frac{26}{3}\kappa RV C_d - \frac{1}{3}RV''C_e \right. \right. \right. \\ \left. \left. \left. + \frac{5}{2}\kappa^2((\nabla\phi)^2)^2C_f + 2\kappa^2V(\nabla\phi)^2C_g - 4\kappa V''(\nabla\phi)^2C_h \right. \right. \right. \\ \left. \left. \left. + 10\kappa^2V^2C_i - 4\kappa V'^2C_j + V''^2C_k \right) \right] \right]. \quad (3.68)$$

Introduced coefficients are C_1, C_2, C_3, C_4 dimensionfull parameters and C_a, \dots, C_k dimensionless parameters. For true Einstein scalar theory, we know that the parameters as,

$$C_n > 0 \quad \text{and} \quad n = 1, 2, 3, 4, \quad (3.69)$$

$$C_m = (-1)^{T,F} C_M (> 0) \quad \text{and} \quad m = M = a, b, c, \dots, k. \quad (3.70)$$

Therefore, the normal factor $N_0[g, \phi]$ is reexpressed by,

$$N_0[g, \phi] = \exp \left[-\frac{1}{32\pi^2} \int d^4x \sqrt{g} \left[(const.) + \left(-\frac{29}{3}RC_1 + 4\kappa(\nabla\phi)^2C_2 - 2V''C_3 + 20\kappa VC_4 \right) \right. \right. \\ \left. \left. + \left(\frac{43}{30}R_{\mu\nu}R^{\mu\nu}(-1)^{T,F}C_A + \frac{1}{20}R^2(-1)^{T,F}C_B - \frac{2}{3}\kappa R(\nabla\phi)^2(-1)^{T,F}C_C - \frac{26}{3}\kappa RV(-1)^{T,F}C_D \right. \right. \right. \\ \left. \left. \left. - \frac{1}{3}RV''(-1)^{T,F}C_E + \frac{5}{2}\kappa^2((\nabla\phi)^2)^2(-1)^{T,F}C_F + 2\kappa^2V(\nabla\phi)^2(-1)^{T,F}C_G \right. \right. \right. \\ \left. \left. \left. - 4\kappa V''(\nabla\phi)^2(-1)^{T,F}C_H + 10\kappa^2V^2(-1)^{T,F}C_I - 4\kappa V'^2(-1)^{T,F}C_J + V''^2(-1)^{T,F}C_K \right) \right] \right]. \quad (3.71)$$

The partition function called by connected is,

$$W = \log Z = -\frac{1}{32\pi^2} \int d^4x \sqrt{g} \left[(const.) + \left(-\frac{29}{3}RC_1 + 4\kappa(\nabla\phi)^2C_2 - 2V''C_3 + 20\kappa VC_4 \right) \right. \\ \left. + \left(\frac{43}{30}R_{\mu\nu}R^{\mu\nu}(-1)^{T,F}C_A + \frac{1}{20}R^2(-1)^{T,F}C_B - \frac{2}{3}\kappa R(\nabla\phi)^2(-1)^{T,F}C_C - \frac{26}{3}\kappa RV(-1)^{T,F}C_D \right. \right. \\ \left. \left. - \frac{1}{3}RV''(-1)^{T,F}C_E + \frac{5}{2}\kappa^2((\nabla\phi)^2)^2(-1)^{T,F}C_F + 2\kappa^2V(\nabla\phi)^2(-1)^{T,F}C_G \right. \right. \\ \left. \left. - 4\kappa V''(\nabla\phi)^2(-1)^{T,F}C_H + 10\kappa^2V^2(-1)^{T,F}C_I - 4\kappa V'^2(-1)^{T,F}C_J + V''^2(-1)^{T,F}C_K \right) \right]. \quad (3.72)$$

From these results, we could find the effective gravitational action as follows,

$$\begin{aligned} \Gamma[g, \phi] = & \frac{1}{32\pi^2} \int d^4x \sqrt{g} \left[(const.) + \left(-\frac{29}{3}RC_1 + 4\kappa(\nabla\phi)^2C_2 - 2V''C_3 + 20\kappa VC_4 \right) \right. \\ & + \left(\frac{43}{30}R_{\mu\nu}R^{\mu\nu}(-1)^{T,F}C_A + \frac{1}{20}R^2(-1)^{T,F}C_B - \frac{2}{3}\kappa R(\nabla\phi)^2(-1)^{T,F}C_C - \frac{26}{3}\kappa RV(-1)^{T,F}C_D \right. \\ & \quad \left. - \frac{1}{3}RV''(-1)^{T,F}C_E + \frac{5}{2}\kappa^2((\nabla\phi)^2)^2(-1)^{T,F}C_F + 2\kappa^2V(\nabla\phi)^2(-1)^{T,F}C_G \right. \\ & \left. \left. - 4\kappa V''(\nabla\phi)^2(-1)^{T,F}C_H + 10\kappa^2V^2(-1)^{T,F}C_I - 4\kappa V'^2(-1)^{T,F}C_J + V''^2(-1)^{T,F}C_K \right) \right]. \quad (3.73) \end{aligned}$$

The effective gravitational action has many important terms of the gravity, scalar field theory and coupled fields. The renormalization is to induced in the $N_v[g, \phi]$ confirmly. From the considerations of the gravity, the gravitational effective action is used as the R^2 gravitation with non scalar field as,

$$\Gamma[g, 0] = \frac{1}{32\pi^2} \int d^4x \sqrt{g} \left((const.) - \frac{29}{3}RC_1 + \frac{43}{30}R_{\mu\nu}R^{\mu\nu}(-1)^{T,F}C_A + \frac{1}{20}R^2(-1)^{T,F}C_B \right). \quad (3.74)$$

This is the Einstein R^2 pure action for consider the gravitation developments in the universal story. The scalar coupling Einstein R^2 action is anticipated in considering the early gravitational ages in physics.

4 Results and future

We derived the gravitational effective action from the quantum corrected action with running the cuoff $\mu(\rightarrow 0^+ > \sqrt{z}/2) \rightarrow \Lambda$. The actions (3.73) and (3.74) were basic actions for gravitation since the Einstein R^2 couplling with scalar matter and potential. Therefore, we could lead the useful results of the gravity from the quantum gravity what is the non local heat kernel expansions in the functional renormalization group equation.

In section 2, we prepared to verify the functional renormalization group equation with the quantum gravity. We considered the scalar action on the curved space. The functional renormalization group equation was expressed by the propagator also expanded to the non local heat kernel expansions these are written by heat kernel coefficients and structure functions. As the result, the nonlocal heat kernel expansion of the functional renormalization group equation was lead towards the expanded results given by Q_n functionals. Q_n functionals were calculated to local result and non local structure dependent result. Then, the master equation was structured formally in this section.

In latter section, we considered that the renormalization group are expanded with the gravity coupled with the scalar matter on Einstein manifold. The perturbative expansions were giving to the scalar coupled gravity hessian, on the other side, the ghost operator were also given. The functional renormalization group was expanded by these laplace operators towards the quantum corrected Wetterich equation. The heat kernel coefficients and structure functions were calculated with this quantum corrected equation, therefore, we reached the effective action included the cutoff momentum Λ with the scale $\mu^2 > z/4$. Since we got the quantum corrected action with the cuoff k , we could normalize it towards non cutoff scale dependent effective aqction as (3.73) and (3.74). Therefore, we knew that the gravitational

effective action lead from normalization of the quantum corrected action in the quantum gravity to ensure the Einstein R^2 scalar coupled or non coupled systems.

References

- [1] S. Hawking and G. Ellis, *The Large-Scale Structure of Space-Time*, Cambridge (1973)
- [2] S. A. Hayward, “Formation and Evaporation of Nonsingular Black Holes,” *Phys. Rev. Lett* **96**, 031103 (2006).
- [3] R. Percacci, “An Introduction to Covariant Quantum Gravity and Asymptotic Safety,” World Scientific, Singapore, 2017, doi:10.1142/10369.
- [4] Ron-Gen. Cai, Li-Ming. Cao, and N. Ohta “Mass and thermodynamics of Kaluza-Klein black holes with aquashed horizons,” *Phys. Lett. B* 639 (2006), 354-361.
- [5] K. Wilson, and J. Kogut, *Phys. Rep.* **12 C** 2 (1974).
- [6] F. J. Wegner and A. Houghton, *Phys. Rev.* **A 8** 410 (1973).
- [7] A. Hasenfratz and P. Hasenfratz, *Nucl. Phys.* **B270** 687-701 (1986).
- [8] J. Polchinski, *Nucl. Phys.* B231, 269-295 (1984).
- [9] C. Bagunuls and C. Berviller. *Phys. Rept.* **348** 91 (2001), [arXiv:0002034[hep-th]].
- [10] J. Berges, N. Tetradis and C. Wetterich, *Phys. Rept.* **363**, 223 (2002), [arXiv:0005122[hep-th]].
- [11] C. Wetterich, *Phys. Lett.* **B 301**, 90 (1993).
- [12] A. O. Barvinski and G. A. Vilkoviski, *Nucl. Phys. B* **282** 163 (1987).
- [13] A. O. Barvinski and G. A. Vilkoviski, *Nucl. Phys. B* **333** 471 (1990).
- [14] I. G. Avramidi, *Phys. Lett. B* **236** 443 (1990).
- [15] I. G. Avramidi, *Phys. Lett. B* **238** 92 (1990).
- [16] A. Codello and O. Zanusso, *J. Math. Phys.* 54 013513 (2013) [arXiv:1203.2034[math-ph]].
- [17] P. Donà, A. Eichhorn and R. Percacci, “Matter matters in asymptotically safe quantum gravity,” *Phys. Rev. D* **89**, 084035 (2014) [arXiv:1311.2898 [hep-th]].
- [18] A. Codello, R. Percacci, L. Rashwal and A. Tonero, “Computing the Effective Action with the Functional Renormalization Group,” *Eur. Phys. J. C* 76, no. 4, 226 (2016) [arXiv:1505.03119[hep-th]].
- [19] J. Meibohm, J. M. Pawłowski and M. Reichert, “Asymptotic safety of gravity-matter systems,” *Phys. Rev. D* **93**, 084035 (2016) [arXiv:1510.07018 [hep-th]].
- [20] M. Reuter and F. Saueressig, “Quantum Gravity and the Functional Renormalization Group: The Road towards Asymptotic Safety,” Cambridge University Press, Cambridge, UK, 2019, doi:10.1017/9781316227596.

- [21] M. Reuter, “Nonperturbative evolution equation for quantum gravity,” *Phys. Rev. D* **57**, 971 (1998) [arXiv:hep-th/9605030 [hep-th]].
- [22] A. Bonanno and M. Reuter, “Renormalization group improved black hole space-times,” *Phys. Rev. D* **62**, 043008 (2000) [arXiv:hep-th/0002196 [hep-th]].
- [23] A. Eichhorn, arXiv[hep-th]:1810.07615 (2019)
- [24] A. Bonanno and M. Reuter, *Phys. Rev. D* **60**, 084011 (1999), arXiv[gr-qc]:9811026
- [25] A. Bonanno and M. Reuter, “Spacetime structure of an evaporating black hole in quantum gravity,” *Phys. Rev. D* **73**, 083005 (2006) [arXiv:hep-th/0602159 [hep-th]].
- [26] M. Reuter and E. Tuiran, “Quantum Gravity Effects in the Kerr Spacetime,” *Phys. Rev. D* **83**, 044041 (2011) [arXiv:1009.3528 [hep-th]].
- [27] K. Falls, D. F. Litim and A. Raghuraman, “Black Holes and Asymptotically Safe Gravity,” *Int. J. Mod. Phys. A* **27**, 1250019 (2012) [arXiv:1002.0260 [hep-th]].
- [28] B. Koch and F. Saueressig, “Structural aspects of asymptotically safe black holes,” *Class. Quant. Grav.* **31**, 015006 (2014) [arXiv:1306.1546 [hep-th]].
- [29] B. Koch and F. Saueressig, “Black holes within Asymptotic Safety,” *Int. J. Mod. Phys. A* **29**, 1430011 (2014) [arXiv:1401.4452 [hep-th]].
- [30] A. Bonanno, B. Koch and A. Platania, “Gravitational collapse in Quantum Einstein Gravity,” *Found. Phys.* **48**, 1393 (2018) [arXiv:1710.10845 [gr-qc]].
- [31] J. M. Pawłowski and D. Stock, “Quantum-improved Schwarzschild-(A)dS and Kerr-(A)dS spacetimes,” *Phys. Rev. D* **98**, 106008 (2018) [arXiv:1807.10512 [hep-th]].
- [32] M. Reuter and H. Weyer, “Running Newton constant, improved gravitational actions, and galaxy rotation curves,” *Phys. Rev. D* **70**, 124028 (2004) [arXiv:hep-th/0410117 [hep-th]].
- [33] A. Platania, “Dynamical renormalization of black-hole spacetimes,” *Eur. Phys. J. C* **79**, 470 (2019) [arXiv:1903.10411 [gr-qc]].
- [34] A. Bonanno, R. Casadio and A. Platania, “Gravitational antiscreening in stellar interiors,” *JCAP* **01**, 022 (2020) [arXiv:1910.11393 [gr-qc]].
- [35] A. Ishibashi, N. Ohta, D. Yamaguchi, “Quantum improved charged black holes,” *Phys. Rev. D* **104**, 066016 (2021) [arXiv:2106.05015 [hep-th]].
- [36] A. Platania, “From renormalization group flows to cosmology,” *Front. in Phys.* **8**, 188 (2020) [arXiv:2003.13656 [gr-qc]].
- [37] A. Eichhorn and S. Lippoldt, “Quantum gravity and Standard-Model-like fermions,” *Phys. Lett. B* **767**, 142 (2017) [arXiv:1611.05878 [gr-qc]].

- [38] J. Biemans, A. Platania and F. Saueressig, “Renormalization group fixed points of foliated gravity-matter systems,” JHEP **05**, 093 (2017) [arXiv:1702.06539 [hep-th]].
- [39] U. Harst and M. Reuter, “QED coupled to QEG,” JHEP **05**, 119 (2011) [arXiv:1101.6007 [hep-th]].
- [40] N. Christiansen and A. Eichhorn, “An asymptotically safe solution to the U(1) triviality problem,” Phys. Lett. B **770**, 154 (2017) [arXiv:1702.07724 [hep-th]].
- [41] A. Eichhorn and F. Versteegen, “Upper bound on the Abelian gauge coupling from asymptotic safety,” JHEP **01**, 030 (2018) [arXiv:1709.07252 [hep-th]].
- [42] R. Alkofer, A. Eichhorn, A. Held, C. M. Nieto, R. Percacci and M. Schröfl, “Quark masses and mixings in minimally parameterized UV completions of the Standard Model,” Annals Phys. **421**, 168282 (2020) [arXiv:2003.08401 [hep-ph]].
- [43] J. L. Lehners and K. S. Stelle, “A Safe Beginning for the Universe?,” Phys. Rev. D **100**, 083540 (2019) [arXiv:1909.01169 [hep-th]].
- [44] J. N. Borissova and A. Eichhorn, “Towards black-hole singularity-resolution in the Lorentzian gravitational path integral,” Universe **7**, 48 (2021) [arXiv:2012.08570 [gr-qc]].
- [45] J. Chojnacki and J. Kwapisz, “Finite Action Principle and Horava-Lifshitz Gravity: early universe, black holes and wormholes,” [arXiv:2102.13556 [gr-qc]]
- [46] C. Jonas, J. L. Lehners and J. Quintin, “Cosmological consequences of a principle of finite amplitudes,” Phys. Rev. D **103**, 103525 (2021) [arXiv:2102.05550 [hep-th]]
- [47] A. Bonanno, A. Eichhorn, H. Gies, J. M. Pawłowski, R. Percacci, M. Reuter, F. Saueressig and G. P. Vacca, “Critical reflections on asymptotically safe gravity,” Front. in Phys. **8**, 269 (2020) [arXiv:2004.06810 [gr-qc]].
- [48] A. Held, “Invariant Renormalization-Group improvement,” [2105.11458 [gr-qc]]
- [49] J.M. Bardeen, “Non-singular general relativistic gravitational collapse,” in Proceedings of International Conference GR5 (Tbilisi, U.S.S.R., 1968).
- [50] *Handbook of Mathematical Functions*, (Dover, New York, 1965) edited by M. Abramowitz, and I.A. Stegun.
- [51] E. Poisson and W. Israel, “Inner-horizon instability and mass inflation in black holes,” Phys. Rev. Lett. **63**, 1663 (1989)
- [52] E. Poisson and W. Israel, “Internal structure of black holes,” Phys. Rev. D **41**, 1796 (1990)
- [53] H. Maeda, T. Torii, and T. Harada, “Novel Cauchy horizon instability,” Phys. Rev. D **71**, 064015 (2005) [arXiv:0501042[gr-qc]].

- [54] G. T. Horowitz and R. Myers, “The Value of Singularities,” *Gen. Rel. Grav.*, **27**, 915 (1995) [arXiv:gr-qc/9503062 [gr-qc]]
- [55] S. de Alwis, A. Eichhorn, A. Held, J. M. Pawłowski, M. Schiffer and F. Versteegen, “Asymptotic safety, string theory and the weak gravity conjecture,” *Phys. Lett. B* **798**, 134991 (2019) [arXiv:1907.07894 [hep-th]].
- [56] C. Gonzalez and B. Koch, “Improved Reissner–Nordström–(A)dS black hole in asymptotic safety,” *Int. J. Mod. Phys. A* **31** 1650141 (2016) [arXiv:1508.01502 [hep-th]]
- [57] M. Sharif and F. Javed, “ On the stability of bardeen thin-shell wormholes”, *Gen. Relativ. Gravit* 48:158 (2016).
- [58] J. Lemos and V. Zanchin, “Regular black holes: Electrically charged solutions, Reissner-Nordstrom outside a de Sitter core”, *Phys. Rev. D* **83**, 124005 (2011).
- [59] E. Poisson, “A RELATIVIST’S TOOLKIT”, Cambridge (2010).