

Claw-free cubic graphs are $(1, 1, 1, 3)$ -packing edge-colorable

Jingxi Hou*

Tao Wang[†]Xiaoqing Yang[‡]

February 24, 2025

Abstract

For a non-decreasing positive integer sequence $S = (s_1, \dots, s_k)$, an S -packing edge-coloring of a graph G is a partition of the edge set of G into subsets E_1, \dots, E_k such that for each $1 \leq i \leq k$, the distance between any two distinct edges $e_1, e_2 \in E_i$ is at least $s_i + 1$. Hocquard et al. conjectured that cubic graphs, except for the Petersen and Tietze graphs, admit $(1, 1, 1, 3)$ -packing edge-colorings. In this paper, we prove that every claw-free cubic graph admits such a coloring.

Keywords: Packing edge-coloring; Cubic graphs; Claw-free graphs

1 Introduction

A *proper edge-coloring* of a graph G is an assignment of colors to its edges such that no two adjacent edges receive the same color. This concept can be extended to *d -distance edge-coloring*, where the *edge-distance* in G , defined as the vertex-distance in its line graph, must be at least $d + 1$ for edges sharing the same color.

A *d -packing edge-coloring* of G partitions $E(G)$ into color classes such that the distance between any two edges within the same class is at least $d + 1$. Generalizing this notion, given a non-decreasing sequence $S = (s_1, \dots, s_k)$, an *S -packing edge-coloring* assigns colors to edges such that any two edges sharing color i have a distance greater than s_i . A graph admitting such a coloring is called *S -packing edge-colorable*. Introduced by Gastineau and Togni [6] as an extension of S -packing coloring (proposed by Goddard and Xu [7, 8]), this concept has inspired significant research (see [1–3, 5, 10]).

In an S -packing edge-coloring, each subset E_i is referred to as an *s_i -packing*. Notably, a 1-packing corresponds to a matching, while a 2-packing forms an induced matching. For brevity, exponents denote repeated elements in S ; e.g., $(1, 2^4)$ represents $(1, 2, 2, 2, 2)$. Hocquard and Lajou [9] proved that subcubic graphs admit $(1, 1, 2^5)$ - and $(1, 2^8)$ -packing edge-colorings. Gastineau and Togni [6] showed that cubic graphs with a 2-factor are $(1, 1, 1, 3, 3)$ -, $(1, 1, 1, 4^5)$ -, and $(1, 1, 2^5)$ -packing edge-colorable. Li et al. [11] established that subcubic outerplanar graphs admit $(1, 2^5)$ - and $(1^2, 2^3)$ -packing edge-colorings, but exclude $(1, 2^4)$ - or $(1^2, 2^2)$ -packing edge-colorings. Liu and Yu [12] further showed that all subcubic graphs are $(1^2, 2^4)$ -packing edge-colorable.

*School of Mathematics and Statistics, Henan University, Kaifeng, 475004, P. R. China.

[†]Center for Applied Mathematics, Henan University, Kaifeng, 475004, P. R. China. **Corresponding author:** wangtao@henu.edu.cn; <https://orcid.org/0000-0001-9732-1617>. Tao Wang was supported by the Natural Science Foundation of Henan Province (No. 242300420238).

[‡]School of Mathematics and Statistics, Henan University, Kaifeng, 475004, P. R. China. Xiaoqing Yang was supported by National Natural Science Foundation of China (No. 12101187).

Gastineau and Togni [6] also studied S -packing edge-colorings of subcubic graphs with specific 1's in S . By Vizing's theorem, every subcubic graph is $(1, 1, 1, 1)$ -packing edge-colorable. Additionally, Fouquet and Vanherpe's result implies:

Theorem 1.1 (Fouquet and Vanherpe [4]). Every subcubic graph admits a $(1, 1, 1, 2)$ -packing edge-coloring.

Gastineau and Togni [6] conjectured that every cubic graph is $(1, 1, 1, 3)$ -packing edge-colorable. However, Hocquard et al. [9] demonstrated exceptions: the Petersen and Tietze graphs are not $(1, 1, 1, 3)$ -packing edge-colorable.

Conjecture 1 (Hocquard et al. [9]). Every cubic graph, except for the Petersen and Tietze graphs, is $(1, 1, 1, 3)$ -packing edge-colorable.

This conjecture has been verified for certain restricted classes of snarks. In this paper, we extend this line of research by focusing on claw-free cubic graphs and proving the following result.

Theorem 1.2. Every claw-free cubic graph is $(1, 1, 1, 3)$ -packing edge-colorable.

2 Proof of Theorem 1.2

A *diamond* is a graph D formed by removing one edge from the complete graph K_4 . In a diamond D , the two degree-3 vertices are called *internal vertices*, while the remaining two are *external vertices*.

For a positive integer k , a *string of k diamonds* is constructed as follows: take k disjoint diamonds D_1, \dots, D_k , where $V(D_i) = \{x_i, y_i, z_i, w_i\}$, and z_i, w_i are the internal vertices of D_i , add two additional isolated vertices x_0 and y_{k+1} , and connect x_{i-1} to y_i for all $i \in [k+1]$. A *ring of diamonds* is a connected claw-free cubic graph in which every vertex belongs to a diamond. Note that a ring of diamonds can be obtained from a string of diamonds by removing x_0 and y_{k+1} , and adding an edge between y_1 and x_k .

For a graph G , *replacing an edge with a string of k diamonds* is defined as follows: for an edge $xy \in E(G)$, remove xy from G , and identify x_0 of a string of k diamonds with x and y_{k+1} of the same string with y .

We rely on the following two results in our proof.

Theorem 2.1 (Oum [13]). A graph G is 2-edge-connected, claw-free, and cubic if and only if it satisfies one of the following:

- (i) $G \cong K_4$;
- (ii) G is a ring of diamonds; or
- (iii) G is constructed from a 2-edge-connected cubic multigraph H by
 - substituting each vertex of H with a triangle, and/or
 - replacing some edges of H with strings of diamonds.

Theorem 2.2 (Plesník [14]). Let G be a 2-edge-connected cubic multigraph. For any two edges $e, f \in E(G)$, the graph G contains a 2-factor that includes e and f .

In this paper, we assign colors to the edges of G from the set $C = \{1_a, 1_b, 1_c, 3_a\}$.




Fig. 1: A $(1, 1, 1)$ -packing edge-coloring of a ring of diamonds

Proposition 1. If G is a ring of diamonds, then G is $(1, 1, 1)$ -packing edge-colorable.

Proof. In each diamond, there are two sets of non-adjacent external edges and one internal edge. For a ring of diamonds, we assign colors as follows:

- color one set of non-adjacent external edges with 1_a ,
- color the remaining set of non-adjacent external edges with 1_b ,
- color the internal edges of each diamond, as well as the edges connecting adjacent diamonds, with 1_c .

This ensures that edges sharing the same color are pairwise non-adjacent (for 1_a and 1_b) or separated by at least one edge (for 1_c), satisfying the distance requirement for a $(1, 1, 1)$ -packing edge-coloring. See Fig. 1 for an illustration of this coloring scheme applied to a ring of diamonds. \square

Theorem 2.3. If G is a 2-edge-connected claw-free cubic graph, then G is $(1, 1, 1, 3)$ -packing edge-colorable.

Proof. By Theorem 2.1, G satisfies one of the following:

- (i) $G \cong K_4$;
- (ii) G is a ring of diamonds; or
- (iii) G is constructed from a 2-edge-connected cubic multigraph H by
 - substituting each vertex of H with a triangle, and/or
 - replacing some edges of H with strings of diamonds.

If $G \cong K_4$, then G is trivially $(1, 1, 1)$ -packing edge-colorable. If G is a ring of diamonds, by Proposition 1, G admits a $(1, 1, 1)$ -packing edge-coloring. Now, consider the case where G is constructed by substituting each vertex of a 2-edge-connected, cubic multigraph H with a triangle, and/or replacing some edges of H with strings of diamonds. (Note that edges may or may not be replaced.) By Theorem 2.2, H contains a 2-factor \mathcal{F}' that is a disjoint union of cycles, say $\mathcal{F}' = C'_1 \cup \dots \cup C'_k$.

Case 1: No edges are replaced by strings of diamonds.

Suppose G is constructed by substituting each vertex of H with a triangle but without replacing any edge of H with strings of diamonds. For each $i \in [k]$, let $C'_i : h_1^i h_2^i \dots h_{m_i}^i h_1^i$ be a cycle




Fig. 2: Creating G by substituting each vertex of H with a triangle.

in \mathcal{F}' . Denote by C_i the subgraph of G with a Hamiltonian cycle $x_1^i h_1^i y_1^i x_2^i h_2^i y_2^i \dots x_{m_i}^i h_{m_i}^i y_{m_i}^i x_1^i$, where $x_j^i h_j^i y_j^i$ represents the substitution of the vertex h_j^i in H with the triangle $x_j^i h_j^i y_j^i$.

Observe that the edge $y_j^i x_{j+1}^i$ on C_i of G corresponds to the edge $h_j^i h_{j+1}^i$ on C'_i of H . Let $\mathcal{F} = C_1 \cup \dots \cup C_k$ (see Fig. 2, which illustrates the construction of G by substituting each vertex of H with a triangle). Since \mathcal{F}' is a 2-factor of H , it follows that \mathcal{F} becomes a 2-factor of G , covering all vertices of G with disjoint cycles.

Since $H - E(\mathcal{F}')$ is a perfect matching in H , denote this matching by M' . (For example, in Fig. 2, $M' = \{h_1^1 h_1^2, h_2^1 h_2^2, h_3^1 h_3^2, h_3^3 h_3^4, h_4^1 h_4^2\}$.) Define M as the set of edges obtained by adding all edges $x_j^i y_j^i$ (from the substituted triangles) to M' . Consequently, G decomposes into M and \mathcal{F} , i.e., $G = M + \mathcal{F}$.

We consider two cases based on the parity of m_i :

- (1) When m_i is even, color the edges of C_i alternately with 1_a and 1_b in sequence.
- (2) When m_i is odd, select an edge $y_j^i x_{j+1}^i$ in C_i and color it with 3_a . Then, color the remaining edges along the path $x_{j+1}^i h_{j+1}^i \dots x_j^i h_j^i y_j^i$ alternately with 1_a and 1_b .

Finally, color all edges in M with 1_c . It is straightforward to verify that the resulting coloring is indeed a $(1, 1, 1, 3)$ -packing edge-coloring of G . (See Fig. 3 for an illustration of a $(1, 1, 1, 3)$ -packing edge-coloring of G after substituting each vertex of H with a triangle.)

Case 2: Some edges are replaced by strings of diamonds.

Suppose G is constructed by replacing some edges of H with strings of diamonds in addition to substituting each vertex of H with a triangle. For each cycle C'_i in \mathcal{F}' , construct C_i in G as in Case 1. Color C_i and M as previously described in Case 1. For edges replaced by strings of diamonds, use the following method to color each string of diamonds.

Type 1 An edge in M' of H is replaced by a string of diamonds. In this case, let the two sets of non-adjacent external edges of the string of diamonds be colored with 1_a and 1_b , while all other edges in the string are colored with 1_c .

Type 2 An edge in the 2-factor of H is replaced by a string of diamonds. Suppose




Fig. 3: A $(1, 1, 1, 3)$ -packing edge-coloring of G by substituting each vertex of H with a triangle.

the edge $y_j^i x_{j+1}^i$ in \mathcal{F} is replaced by a string of diamonds. Let c be its assigned color in Case 1.

Type 2.1 If $c = 1_a$, color the two sets of non-adjacent external edges in the diamond string with 1_b and 1_c , while assigning 1_a to all other edges.

Type 2.2 If $c = 1_b$, color the two sets of non-adjacent external edges in the diamond string with 1_a and 1_c , while assigning 1_b to all other edges.

Type 2.3 If $c = 3_a$, color the two sets of non-adjacent external edges in the diamond string with 1_c and 1_b (or 1_a), while assigning 1_a (or 1_b) to all other edges in the diamond string, except for the edge at one endpoint which is assigned 3_a .

One can easily verify that this is indeed a $(1, 1, 1, 3)$ -packing edge-coloring of G . (See Fig. 4 which depicts a $(1, 1, 1, 3)$ -packing edge-coloring of G obtained by replacing each vertex of H with a triangle and some edges of H with strings of diamonds.) \square

To extend the $(1, 1, 1, 3)$ -packing edge-coloring to claw-free cubic graphs containing bridges, we proceed as follows: Let G be a connected claw-free cubic graph with bridge set $B(G) = \{e_1, \dots, e_b\}$. Removing $B(G)$ decomposes G into components $\{G_i\}_{0 \leq i \leq b}$, where each G_i is a connected induced subgraph of G . Define a new graph T_G , referred to as the bridge tree of G , as follows: The vertex set of T_G is $V(T_G) = \{g_0, g_1, \dots, g_b\}$, where g_i corresponds to component G_i of $G - B(G)$. The edge set of T_G is $E(T_G) = \{g_i g_j \mid \text{there exists a bridge } e \in B(G) \text{ that connects } G_i \text{ and } G_j\}$. Since each edge in $B(G)$ is a bridge, T_G is a tree.

The following lemma is essential for the subsequent proof process.

Lemma 2.4 ([3]). Let G be a connected claw-free cubic graph with $|B(G)| \geq 1$, and let G_i be the components of $G - B(G)$. Then G_i is 2-edge-connected and satisfies one of the following.

- (1) G_i is isomorphic to K_3 (i.e., $\Delta(G_i) = 2$);
- (2) G_i is isomorphic to a diamond D (i.e., $\Delta(G_i) = 3$ and $|V(G_i)| = 4$);
- (3) G_i is isomorphic to a 2-edge-connected graph with $\Delta(G_i) = 3$ and $|V(G_i)| \geq 5$. Moreover, G_i is a leaf of T_G if and only if it contains exactly one degree-2 vertex.




Fig. 4: A $(1, 1, 1, 3)$ -packing edge-coloring of G by replacing each vertex of H with a triangle and some edges of H with strings of diamonds.

Fix a longest path P in T_G of length $d = \text{diam}(T_G)$, and root T_G at a leaf g_0 of P . Partition $V(T_G)$ into levels $A_i = \{g_j \in V(T_G) \mid d_{T_G}(g_0, g_j) = i\}$. By construction:

- $A_0 = \{g_0\}$;
- Each $g_j \in A_i$ ($i > 0$) has a unique parent $g_k \in A_{i-1}$;
- Every vertex of A_d is a leaf in T_G .

Consequently, if $g_j \in A_i$ ($i > 0$), there exists a unique degree-2 vertex $p \in V(G_j)$ with exactly one neighbor $q \in V(G_k)$, where g_k is the parent of g_j in T_G . In this context, q is referred to as the *up-neighbor* of p .

When $G_i \cong K_3$ or $G_i \cong D$, a desired coloring is straightforward. We therefore focus on constructing a 2-edge-connected claw-free cubic graph \widetilde{G}_i when G_i satisfies conclusion (3) of Lemma 2.4. Assume G_i is a 2-edge-connected graph with maximum degree 3, where $|V(G_i)| \geq 5$. Let $V_i = \{v_1^i, v_2^i, \dots, v_{r_i}^i\}$ be the set of all degree-2 vertices in G_i , where v_1^i is uniquely required to have an up-neighbor. For each $v_j^i \in V_i$, where $1 \leq j \leq r_i$, let u_j^i and w_j^i denote the two neighbors of v_j^i in G_i .

Claim 1. Every vertex in V_i lies on a triangle of G_i , i.e., $u_j^i w_j^i \in E(G_i)$. Consequently, V_i is an independent set.

Proof. Since G is claw-free and v_j^i is incident with a bridge, it follows that v_j^i must lie on a triangle of G_i .

Assume without loss of generality that v_1^i and v_2^i are adjacent. As every vertex in V_i lies on a triangle of G_i , we have that v_1^i and v_2^i share a common neighbor z_1^i . If $d_{G_i}(z_1^i) = 2$, this

contradicts the condition $|V(G_i)| \geq 5$. If $d_{G_i}(z_1^i) = 3$, G_i contains a bridge, which is also a contradiction. \square

Let s_j^i be the neighbor of u_j^i distinct from v_j^i and w_j^i , and let b_j^i be the neighbor of w_j^i distinct from u_j^i and v_j^i .

Claim 2. $s_j^i \neq b_j^i$, and $d_{G_i}(s_j^i) = d_{G_i}(b_j^i) = 3$.

Proof. Assume $s_j^i = b_j^i$. In this case, G_i must either contain a bridge or be a diamond, leading to a contradiction. Since $d_{G_i}(u_j^i) = 3$ and $u_j^i b_j^i \notin E(G)$, if $d_{G_i}(b_j^i) = 2$, Claim 1 implies that b_j^i must lie on a triangle of G_i , which is a contradiction. By symmetry, s_j^i must also have degree 3 in G_i . \square

Now, we construct a 2-edge-connected claw-free cubic graph \widetilde{G}_i from G_i . Consider two cases based on the parity of r_i .

Case 1 (r_i even). Construct \widetilde{G}_i from G_i by adding the edges $v_j^i v_{j+1}^i$ for all odd j with $1 \leq j \leq r_i - 1$.

Case 2 (r_i odd). By Claims 1 and 2, we have $u_1^i w_1^i \in E(G_i)$, $s_1^i \neq b_1^i$, and $d_{G_i}(s_1^i) = d_{G_i}(b_1^i) = 3$. Suppose $s_1^i b_1^i \in E(G_i)$. Since G is claw-free, it follows that s_1^i and b_1^i share a common neighbor z_1^i . If $d_{G_i}(z_1^i) = 2$, this contradicts r_i being odd (as it would cause $r_i = 2$). If $d_{G_i}(z_1^i) = 3$, then G_i would contain a bridge, which is also a contradiction. Thus, $s_1^i b_1^i \notin E(G_i)$. Let \widetilde{G}_i be the graph obtained from $G_i - \{v_1^i, u_1^i, w_1^i\}$ by adding edge $s_1^i b_1^i$ and edges $v_j^i v_{j+1}^i$ for all even j with $2 \leq j \leq r_i - 1$.

In [3], it was proved that \widetilde{G}_i is a 2-edge-connected claw-free cubic graph, and if r_i is odd, then $s_1^i b_1^i$ does not lie on a triangle of \widetilde{G}_i unless $\widetilde{G}_i \cong K_4$.

Theorem 2.5. Every connected claw-free cubic graph is $(1, 1, 1, 3)$ -packing edge-colorable.

Proof. Let G be a connected claw-free cubic graph with $|B(G)| = b \geq 0$, where $B(G)$ denotes its bridge set. Let G_i denote the components of $G - B(G)$. When $b = 0$, Theorem 2.3 provides a $(1, 1, 1, 3)$ -packing edge-coloring of G . For $b > 0$, consider the tree T_G defined by $V(T_G) = \{g_0, g_1, \dots, g_b\}$ and $E(T_G) = \{g_i g_j \mid \text{there exists an edge } e \in B(G) \text{ that connects } G_i \text{ and } G_j\}$. Choose a path P in T_G of length $d = \text{diam}(T_G)$ and let g_0 be a leaf of P .

First, we consider the component G_0 . Clearly, g_0 is also a leaf of T_G . By Lemma 2.4, G_0 contains exactly one degree-2 vertex v_1^0 , and in this case, $r_0 = 1$. By Claims 1 and 2, $u_1^0 w_1^0 \in E(G_0)$, $s_1^0 \neq b_1^0$ and $s_1^0 b_1^0 \notin E(G_0)$. We construct \widetilde{G}_0 as follows: apply the method as Case 2 to obtain the graph \widetilde{G}_0 from $G_0 - \{v_1^0, u_1^0, w_1^0\}$ by adding the edge $s_1^0 b_1^0$. The resulting graph \widetilde{G}_0 is a 2-edge-connected claw-free cubic graph.

Claim 3. \widetilde{G}_0 admits a $(1, 1, 1, 3)$ -packing edge-coloring φ' such that $\varphi'(s_1^0 b_1^0) = 1_a$, and no edge incident with s_1^0 or b_1^0 is colored with 3_a .

Proof. Note that the edge $s_1^0 b_1^0$ is not on a triangle of \widetilde{G}_0 unless $\widetilde{G}_0 \cong K_4$. If $\widetilde{G}_0 \cong K_4$ or \widetilde{G}_0 is a ring of diamonds, it is straightforward to verify that there exists a proper edge-coloring using colors $1_a, 1_b, 1_c$ such that $s_1^0 b_1^0$ is colored with 1_a . So we may assume that $s_1^0 b_1^0$ is not on a triangle of \widetilde{G}_0 , and \widetilde{G}_0 is not a ring of diamonds. By Theorem 2.1, let \widetilde{G}_0 be constructed from a 2-edge-connected cubic multigraph H by substituting each vertex of H with a triangle, and possibly replacing some edges of H with strings of diamonds. Let $s_1^0 b_1^0$ correspond to an edge e

in H . By Theorem 2.2, H has a 2-factor \mathcal{F}' containing e , where $e \in C'_i$. Then $s_1^0 b_1^0$ corresponds to $y_j^i x_{j+1}^i$, where $y_j^i x_{j+1}^i$ may or may not be replaced by a string of diamonds, and $1 \leq j \leq m_i$.

We consider two cases:

- (1) m_i is even. By the coloring process described in the proof of Theorem 2.3, \widetilde{G}_0 admits a $(1, 1, 1, 3)$ -packing edge-coloring φ' with $\varphi'(s_1^0 b_1^0) = 1_a$.
- (2) m_i is odd. If necessary, we can relabel the vertices on the cycle $x_1^i h_1^i y_1^i x_2^i h_2^i y_2^i \dots x_{m_i}^i h_{m_i}^i y_{m_i}^i x_1^i$ such that, from the coloring process described in the proof of Theorem 2.3, e is colored with 1_a , and correspondingly $s_1^0 b_1^0$ is also colored with 1_a .

This completes the proof of Claim 3. \square

Let φ' be an edge-coloring from Claim 3. By symmetry, let $b_1^0 w_1^0$ be the edge in $\{b_1^0 w_1^0, s_1^0 u_1^0\}$ farthest from the edges colored with 3_a in \widetilde{G}_0 . Now, define a coloring φ of G_0 as follows: let $\varphi(e) = \varphi'(e)$ for all $e \in E(G_0 - \{v_1^0, u_1^0, w_1^0\})$, color the edges of the path $s_1^0 u_1^0 v_1^0 w_1^0$ alternately with 1_a and 1_b in sequence, and assign $\varphi(b_1^0 w_1^0) = 3_a$ and $\varphi(u_1^0 w_1^0) = 1_c$.

Next, we consider each G_i , where $i \geq 1$. If $G_i \cong K_3$, then the edges of G_i are assigned the colors $1_a, 1_b$ and 1_c , respectively. If $G_i \cong D$, then the two sets of non-adjacent external edges and one internal edge are assigned the colors $1_a, 1_b$, and 1_c , respectively.

Claim 4. Every 2-edge-connected G_i with $\Delta(G_i) = 3$ and $|V(G_i)| \geq 5$ admits a $(1, 1, 1, 3)$ -packing edge-coloring.

Proof. We consider the cases based on the parity of r_i .

- (1) r_i is even. We construct \widetilde{G}_i via Case 1. By Theorem 2.3, \widetilde{G}_i has a $(1, 1, 1, 3)$ -packing edge-coloring φ' . Let φ be a coloring of G_i , and $\varphi(e) = \varphi'(e)$ for all $e \in E(G_i)$. Clearly, φ is a $(1, 1, 1, 3)$ -packing edge-coloring.
- (2) r_i is odd. We construct \widetilde{G}_i via Case 2. By Theorem 2.3, \widetilde{G}_i has a $(1, 1, 1, 3)$ -packing edge-coloring φ' . Let φ be a coloring of G_i , and $\varphi(e) = \varphi'(e)$ for all $e \in E(G_i)$. Color the edges of the path $s_1^0 u_1^0 v_1^0 w_1^0$ alternately with colors 1_a and 1_b in sequence, and assign $\varphi(b_1^0 w_1^0) = 3_a$ and $\varphi(u_1^0 w_1^0) = 1_c$. From the proof of Claim 3, this yields a valid $(1, 1, 1, 3)$ -packing edge-coloring. \square

Claim 5. For the coloring described above on G_i , all edges incident with degree-2 vertices in G_i use colors from $\{1_a, 1_b, 1_c\}$.

Proof. For any degree-2 vertex v_j^i in G_i , let u_j^i and w_j^i denote the two neighbors of v_j^i in G_i . Since G is claw-free, we have $u_j^i w_j^i \in E(G_i)$. According to the coloring process for G_i , edges colored with 3_a cannot lie on a triangle. The coloring process explicitly assigns only $\{1_a, 1_b, 1_c\}$ to edges adjacent to v_j^i , satisfying the claim. \square

The colors in $\{1_a, 1_b, 1_c\}$ can be freely permuted without affecting the validity of the packing condition on G_i . Specifically, if we perform a permutation on the edges that have already been assigned colors, for example, by changing all edges assigned 1_a to 1_b and all edges assigned 1_b to 1_a , this operation does not cause any contradictions or conflicts. This is because the permutation only reassigns the colors of the edges without altering the structure of the edges or the coloring requirements of the graph.

We perform a coloring of T_G using Breadth-First Search (BFS). First, assign $\{1_a, 1_b, 1_c, 3_a\}$ to the edges of G_0 (root component). For each edge pq connecting A_0 and A_1 , where q is the

up-neighbor of p , the edge pq 's color is determined by G_0 's coloring. Note that each component $G_i \neq G_0$ contains exactly one vertex having an up-neighbor. If necessary, we can permute the colors $1_a, 1_b, 1_c$ in G_i to allow pq to use the color missing at q . Next, by appropriately permuting $\{1_a, 1_b, 1_c\}$, assign colors to the edges of G_i in A_1 . The color of the edges between A_1 and A_2 is then determined by the coloring of G_i corresponding to a vertex $g_i \in A_1$. Continuing this process iteratively, we can obtain a $(1, 1, 1, 3)$ -packing edge-coloring of G . \square

References

- [1] B. Brešar, N. Gastineau and O. Togni, Packing colorings of subcubic outerplanar graphs, *Aequationes Math.* 94 (5) (2020) 945–967.
- [2] B. Brešar, S. Klavžar, D. F. Rall and K. Wash, Packing chromatic number, $(1, 1, 2, 2)$ -colorings, and characterizing the Petersen graph, *Aequationes Math.* 91 (1) (2017) 169–184.
- [3] B. Brešar, K. Kuenzel and D. F. Rall, Claw-free cubic graphs are $(1, 1, 2, 2)$ -colorable, arXiv: 2409.15455 (2024) <https://doi.org/10.48550/arXiv.2409.15455>.
- [4] J.-L. Fouquet and J.-M. Vanherpe, On parsimonious edge-colouring of graphs with maximum degree three, *Graphs Combin.* 29 (3) (2013) 475–487.
- [5] N. Gastineau and O. Togni, S -packing colorings of cubic graphs, *Discrete Math.* 339 (10) (2016) 2461–2470.
- [6] N. Gastineau and O. Togni, On S -packing edge-colorings of cubic graphs, *Discrete Appl. Math.* 259 (2019) 63–75.
- [7] W. Goddard and H. Xu, The S -packing chromatic number of a graph, *Discuss. Math. Graph Theory* 32 (4) (2012) 795–806.
- [8] W. Goddard and H. Xu, A note on S -packing colorings of lattices, *Discrete Appl. Math.* 166 (2014) 255–262.
- [9] H. Hocquard, D. Lajou and B. Lužar, Between proper and strong edge-colorings of subcubic graphs, *J. Graph Theory* 101 (4) (2022) 686–716.
- [10] A. Kostochka and X. Liu, Packing $(1, 1, 2, 4)$ -coloring of subcubic outerplanar graphs, *Discrete Appl. Math.* 302 (2021) 8–15.
- [11] S. Li, Y. Li and X. Liu, Packing edge-colorings of subcubic outerplanar graphs, arXiv: 2411.05720 (2024) <https://doi.org/10.48550/arXiv.2411.05720>.
- [12] X. Liu and G. Yu, On the $(1^2, 2^4)$ -packing edge-coloring of subcubic graphs, arXiv: 2402.18353 (2024) <https://doi.org/10.48550/arXiv.2402.18353>.
- [13] S.-i. Oum, Perfect matchings in claw-free cubic graphs, *Electron. J. Combin.* 18 (1) (2011) Paper 62.
- [14] J. Plesník, Connectivity of regular graphs and the existence of 1-factors, *Mat. Časopis Sloven. Akad. Vied* 22 (1972) 310–318.