

On the length over which k -Göbel sequences remain integers

Yuh Kobayashi and Shin-ichiro Seki

ABSTRACT. We prove that the sequence $(N_k)_k$, where each N_k is defined as the smallest positive integer n for which the n th term $g_{k,n}$ of the k -Göbel sequence is not an integer, is unbounded.

1. Introduction

For $k \geq 2$, the k -Göbel sequence $(g_{k,n})_n$ is defined by the initial value $g_{k,1} = 2$ and the recursion $ng_{k,n} = (n-1)g_{k,n-1} + g_{k,n-1}^k$. Let $N_k := \inf\{n \geq 1 \mid g_{k,n} \notin \mathbb{Z}\}$. The original Göbel's sequence (= 2-Göbel sequence, [7, A003504]) with $N_2 = 43$ has attracted interest as an example of *the strong law of small numbers* ([2]). (The growth of $g_{k,n}$ is very fast. In fact, the value $g_{2,43} \approx 5.4 \times 10^{178485291567}$ is very large.)

The behavior of the sequence $(N_k)_k$ ([7, A108394]) is not yet understood well and remains mysterious. In [6], Matsuhira, Matsusaka, and Tsuchida proved that $\min_{k \geq 2} N_k = 19$. As mentioned in [6, Section 3] and [4, Episode 3], the following three questions are fundamental problems about the sequence $(N_k)_k$:

- (1) Why is N_k a prime number for most values of k ? Or rather, in what cases does N_k become a composite number?
- (2) Does N_k always take a finite value for any $k \geq 2$?
- (3) Is the sequence $(N_k)_k$ unbounded?

For Problems (1) and (2), only numerical data has been obtained. In [5], it is shown that N_k is prime for 86.5% of values up to $k \leq 10^7$, and that N_k is finite for $k \leq 10^{14}$.

In this short note, however, we report that Problem (3) can be solved in a very elementary way within the framework of [6]. Let $m\#$ denote the primorial of m or, in other words, the radical of $m!$.

THEOREM. *Let m be a positive integer. If $k \geq 2$ satisfies $k \equiv 1 \pmod{m!/m\#}$, then $N_k > m$. In particular, $\sup_{k \geq 2} N_k = \infty$.*

2. Proof

Let $k \geq 2$, $r \geq 1$ be integers and p a prime. Let $\mathbb{Z}_{(p)}$ be the localization of \mathbb{Z} at (p) and $\nu_p(n)$ be the p -adic valuation of n . For any positive integer n with $\nu_p(n!) \leq r$, we define $g_{k,p,r}(n) \in \mathbb{Z}/p^{r-\nu_p(n!)}\mathbb{Z} \cup \{\mathbf{F}\}$ as in [6, Definition 9]: For $n = 1$, $g_{k,p,r}(1) = 2 \pmod{p^r}$. For $n \geq 2$, when $g_{k,p,r}(n-1) = \mathbf{F}$, $g_{k,p,r}(n) = \mathbf{F}$. When $g_{k,p,r}(n-1) = a \pmod{p^{r-\nu_p((n-1)!)}}$, if $(n-1)a + a^k \not\equiv 0 \pmod{p^{\nu_p(n)}}$, then $g_{k,p,r}(n) = \mathbf{F}$, and otherwise, $g_{k,p,r}(n) = \frac{(n-1)a + a^k}{p^{\nu_p(n)}} \cdot c \pmod{p^{r-\nu_p(n!)}}$, where c is an integer such that $c \cdot (n/p^{\nu_p(n)}) \equiv 1 \pmod{p^{r-\nu_p(n!)}}$. Let $\varphi(n)$ denote Euler's totient function.

2020 *Mathematics Subject Classification.* Primary 11B37; Secondary 11B50.

Key words and phrases. Göbel sequences.

This research was supported by JSPS KAKENHI Grant Numbers JP22K13960 (Kobayashi) and JP21K13762 (Seki).

PROOF OF THEOREM. Let $m \geq 4$ and $k \geq 2$ be integers. Assume that $k \equiv 1 \pmod{m!/m\#}$. For each prime $p \leq m$, set $r_p := \nu_p(m!)$ and $k_p := \varphi(p^{r_p}) + 1$. It is clear that $k_p > r_p \geq 1$. Let us temporarily suppose that for some prime $p \leq m$, we have $g_{k_p, p, r_p}(m) \neq \mathbf{F}$. Since $\varphi(p^{r_p})$ divides $m!/m\#$, we see that $k \equiv k_p \pmod{\varphi(p^{r_p})}$, and thus, by [6, Proposition 12], it follows that $g_{k, p, r_p}(m) = g_{k_p, p, r_p}(m) \neq \mathbf{F}$. Therefore, by [6, Lemma 10], we conclude that $g_{k, n} \in \mathbb{Z}_{(p)}$ for all $1 \leq n \leq m$. By this argument, together with [6, Lemma 4] and the fact that $\bigcap_p \mathbb{Z}_{(p)} = \mathbb{Z}$, in order to prove $N_k > m$, it suffices to show that $g_{k_p, p, r_p}(m) \neq \mathbf{F}$ for each prime $p \leq m$.

Since $k_2 > r_2$, we see that $2 + 2^{k_2} \equiv 2 \pmod{2^{r_2}}$. Hence, we have $g_{k_2, 2, r_2}(2) = 1 \pmod{2^{r_2-1}}$. It is clear that, subsequently, $g_{k_2, 2, r_2}(n) = 1 \pmod{2^{r_2-\nu_2(n)}}$ for $2 \leq n \leq m$. In particular, $g_{k_2, 2, r_2}(m) \neq \mathbf{F}$.

Let $p \leq m$ be an odd prime. For any $1 \leq n \leq m$, we have $g_{k_p, p, r_p}(n) = 2 \pmod{p^{r_p-\nu_p(n)}}$. In fact, if $g_{k_p, p, r_p}(n-1) = 2 \pmod{p^{r_p-\nu_p((n-1)!)}}$, then since $(n-1)2 + 2^{k_p} \equiv 2n \pmod{p^{r_p-\nu_p((n-1)!)}}$ by Euler's theorem, we have $g_{k_p, p, r_p}(n) = 2 \pmod{p^{r_p-\nu_p(n)}}$. In particular, $g_{k_p, p, r_p}(m) \neq \mathbf{F}$. \square

3. Remark

By replacing the initial value in the definition of the k -Göbel sequence with $g_{k,1} = l$, we define the (k, l) -Göbel sequence, which has been investigated in [1, 3, 5]. Generalizing our previous arguments as follows, the theorem holds in the same form for (k, l) -Göbel sequences as well.

Fix l and a prime $p \leq m$, and in the definition of $g_{k, p, r}$, replace the initial condition with $g_{k, p, r}(1) = l \pmod{p^r}$. We use the notation r_p and k_p as in the previous section. We can easily check that by induction on n , for $n \leq p^{\nu_p(l)}$, $g_{k_p, p, r_p}(n) = (l/p^{\nu_p(n)}) \cdot c_{n, p} \pmod{p^{r_p-\nu_p(n)}}$ holds, while for $n \geq p^{\nu_p(l)}$, we have $g_{k_p, p, r_p}(n) = l/p^{\nu_p(l)} \pmod{p^{r_p-\nu_p(n)}}$. Here, $c_{n, p} \in \mathbb{Z}$ satisfies $c_{n, p} \cdot (n/p^{\nu_p(n)}) \equiv 1 \pmod{p^{r_p-\nu_p(n)}}$. Therefore, the same proof works for a general l .

References

- [1] H. Gima, T. Matsusaka, T. Miyazaki, S. Yara, *On integrality and asymptotic behavior of the (k, l) -Göbel sequences*, J. Integer Seq. **27** (2024) Article 24.8.1, 16pp.
- [2] R. K. Guy, *The strong law of small numbers*, Amer. Math. Monthly **95** (1988), 697–712.
- [3] H. Ibstedt, *Some sequences of large integers*, Fibonacci Quart. **28** (1990), 200–203.
- [4] D. Kobayashi, S. Seki, *Seisu-tan 1: A strange tale of integers' world*, Nippon Hyoron Sha (in Japanese), 2023.
- [5] Y. Kobayashi, S. Seki, *A note on non-integrality of the (k, l) -Göbel sequences*, arXiv:2410.23240.
- [6] R. Matsuhira, T. Matsusaka, K. Tsuchida, *How long can k -Göbel sequences remain integers?*, Amer. Math. Monthly **131** (2024), 784–793.
- [7] N. J. A. Sloane, *The on-line encyclopedia of integer sequences*, available at <https://oeis.org>.

DEPARTMENT OF MATHEMATICAL SCIENCES, AOYAMA GAKUIN UNIVERSITY, 5-10-1 FUCHINOBE, CHUO-KU, SAGAMIHARA, KANAGAWA, 252-5258, JAPAN

Email address: kobayashi@math.aoyama.ac.jp

DEPARTMENT OF MATHEMATICAL SCIENCES, AOYAMA GAKUIN UNIVERSITY, 5-10-1 FUCHINOBE, CHUO-KU, SAGAMIHARA, KANAGAWA, 252-5258, JAPAN

Email address: seki@math.aoyama.ac.jp