

$f(R)$ Gravity: Gravitational Waves Tests

Rafid H. Dejarah ¹

Department of Physics, Faculty of Sciences, Ankara University, 06100, Ankara, Türkiye

Abstract

This review explores modified theories of gravity, particularly $f(R)$ gravity, as extensions to General Relativity (GR) that offer alternatives to dark energy for explaining cosmic acceleration. These models generalize the Einstein-Hilbert action to include functions of the Ricci scalar, providing new insights into cosmology and astrophysics. The detection of gravitational waves (GWs) has enabled rigorous tests of $f(R)$ gravity, as deviations in GW propagation, speed, and polarization can signal modifications to GR. Constraints on $f(R)$ models arise from LIGO-Virgo observations of binary mergers, the stochastic gravitational wave background (SGWB), and complementary tests in cosmology and weak-field regimes. Future GW detectors, such as LISA and the Einstein Telescope, will enhance sensitivity to smaller deviations from GR, necessitating advancements in theoretical modeling. Among competing theories—including scalar-tensor, massive gravity, and Horndeski models— $f(R)$ gravity remains a pivotal framework for understanding fundamental gravitational physics and cosmology. This review highlights key developments, challenges, and future directions in the field.

arXiv:2502.17519v2 [gr-qc] 26 Feb 2025

¹rafid.dejarah@gmail.com

Contents

1	Introduction	2
1.1	Modified Gravity Theories	3
1.2	Gravitational Waves as a Probe of Modified Gravity	3
2	$f(R)$ Gravity	4
3	Constraints on $f(R)$ Gravity from Gravitational Wave Measurements	9
3.1	Constraints from Binary Black Hole and Neutron Star Mergers	9
3.2	Constraints from Stochastic Gravitational Wave Background (SGWB)	10
3.3	Other Experimental Constraints	10
4	Future Improvements in Constraining $f(R)$ Gravity	11
4.1	Next-Generation Gravitational Wave Detectors	11
4.2	Stochastic Gravitational Wave Background (SGWB)	12
4.3	Multi-Messenger Astronomy	13
4.4	Theoretical Developments	15
5	Other Modified Theories of Gravity and Their Constraints	15
5.1	Scalar-Tensor Theories	16
5.2	Horndeski Gravity	17
5.3	Massive Gravity	18
5.4	Brane-World Models	20
6	Conclusions	21

1 Introduction

General Relativity (GR), formulated by Albert Einstein has been the cornerstone of modern gravitational theory. The framework of GR describes gravity as the curvature of spacetime caused by mass and energy, encapsulated in the well-known Einstein field equations (EFE):

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + g_{\mu\nu}\Lambda = \frac{8\pi G}{c^4}T_{\mu\nu}, \quad (1.1)$$

where $R_{\mu\nu}$ is the Ricci curvature tensor, $g_{\mu\nu}$ is the metric tensor, Λ is the cosmological constant [1, 2], and $T_{\mu\nu}$ is the stress-energy tensor. While GR has successfully passed a multitude of experimental tests ² in the weak-field regime [4], it faces significant challenges in explaining certain cosmological phenomena, such as the accelerated expansion of the Universe [5]. This has motivated the exploration of modified gravity theories, which extend the classical framework to address these issues without the need for dark energy.

Despite its successes, GR is not without its limitations. One of the most pressing challenges is the observed accelerated expansion of the Universe, first discovered through observations of Supernovae Type Ia (SNe Ia) [6–8]. This phenomenon is typically attributed to dark energy, a mysterious component of the Universe that exerts negative pressure and drives the acceleration.

²See Table 1 in [3] for a summary of experimental tests of GR.

However, the nature of dark energy remains one of the greatest unsolved problems in modern physics. The cosmological constant Λ , introduced by Einstein to achieve a static universe, is the simplest explanation for dark energy within the framework of GR. However, the observed value of Λ is many orders of magnitude smaller than predictions from quantum field theory, leading to the so-called “cosmological constant problem” [1, 2].

Another challenge to GR arises from the need to explain the observed dynamics of galaxies and galaxy clusters without invoking dark matter. While dark matter provides a successful explanation for the rotation curves of galaxies and the gravitational lensing of light, its particle nature remains elusive despite decades of experimental searches [9]. These challenges have led to the exploration of modified gravity theories, which seek to explain these phenomena by altering the laws of gravity on large scales.

1.1 Modified Gravity Theories

Modified gravity theories aim to address the shortcomings of GR by introducing new degrees of freedom or modifying the gravitational action. Among the various alternatives to GR, $f(R)$ gravity theories [10] have emerged as a compelling class of models [11]. These theories modify the Einstein-Hilbert action by replacing the Ricci scalar R with an arbitrary function $f(R)$, leading to modified field equations that govern the dynamics of spacetime. This modification can account for cosmic acceleration and other phenomena [12, 13] typically attributed to dark energy [14]. Moreover, $f(R)$ gravity is interesting because it introduces additional scalar degrees of freedom, which can manifest as new observable effects, such as modifications to the propagation of gravitational waves (GWs).

The action for $f(R)$ gravity is given by:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2\kappa} f(R) + \mathcal{L}_m \right), \quad (1.2)$$

where $\kappa = 8\pi G$, and \mathcal{L}_m is the matter Lagrangian. The function $f(R)$ defines the dynamics of spacetime, and the field equations derived from this action can lead to different solutions in the weak-field regime. This is particularly important for the study of gravitational wave propagation, as any modifications to the gravitational action will alter the speed, polarization, and other characteristics of GWs.

1.2 Gravitational Waves as a Probe of Modified Gravity

The direct detection of gravitational waves by the LIGO and Virgo collaborations [15] has opened a new window into the Universe, providing a powerful tool for testing GR and its alternatives. GWs are ripples in spacetime caused by the acceleration of massive objects, such as merging black holes or neutron stars. In GR, GWs propagate at the speed of light and have two tensor polarization modes [16]: h_+ and h_\times . However, in modified gravity theories like $f(R)$ gravity, additional scalar polarization modes can arise, and the propagation speed of GWs may differ from the speed of light.

The observation of GW170817 [17], a binary neutron star merger accompanied by a gamma-ray burst, provided a stringent test of the propagation speed of GWs. The time delay between the GW and electromagnetic signals constrained the deviation of the GW speed from the speed

of light to:

$$\frac{|v_{\text{GW}} - c|}{c} \lesssim 10^{-15}. \quad (1.3)$$

This result ruled out many modified gravity models that predict significant deviations in the GW propagation speed.

In Section 2, we provide a detailed discussion of modified $f(R)$ gravity theories. In Section 3, we examine potential constraints on $f(R)$ gravity models based on gravitational wave measurements. Next, in Section 4, we explore future constraints on both the theoretical framework of $f(R)$ gravity and its implications for cosmological observations. Other modified gravity theories are discussed in Section 5, where we present a detailed analysis of each. Finally, in Section 6, we summarize our discussions and offer remarks on the future prospects of testing $f(R)$ gravity models through gravitational waves.

2 $f(R)$ Gravity

Modified gravity theories are an essential part of the modern effort to extend GR, [18, 19], to account for phenomena such as cosmic acceleration and dark energy [20]. Among the various alternatives to GR, $f(R)$ gravity theories, in which the Einstein-Hilbert action is generalized by replacing the Ricci scalar R with an arbitrary function $f(R)$, have gained significant attention. In this section, we will discuss the theoretical foundations of $f(R)$ gravity, the state of current research, and the contributions made by various researchers in this field.

The fundamental action for $f(R)$ gravity is given by Eq.(1.2). The function $f(R)$ characterizes the modification of gravity, with the simplest choice being $f(R) = R$, which recovers GR. However, by choosing other forms of $f(R)$, one can explore the impact of modifications to gravity, such as the introduction of new scalar degrees of freedom. These modifications can have significant consequences for cosmology, gravitational wave propagation, and the formation of structure in the Universe.

A widely studied example of $f(R)$ gravity is the Starobinsky model [21], which introduces a quadratic correction to the Ricci scalar:

$$f(R) = R + \alpha R^2, \quad (2.1)$$

where α is a constant that determines the strength of the modification. Other forms of $f(R)$ gravity include exponential functions, such as:

$$f(R) = R + \beta e^{\gamma R}, \quad (2.2)$$

which are often used to model inflationary scenarios. These models are capable of generating late-time cosmic acceleration by modifying the gravitational dynamics without the need for dark energy. Other forms of $f(R)$ include exponential models, which are often used to explain inflationary dynamics [22], and models involving logarithmic corrections. These modifications are expected to lead to deviations in the behavior of gravitational waves, especially in the high-energy regime where the effects of the scalar degrees of freedom become significant.

Key Contributions to the Field

The study of $f(R)$ gravity has evolved significantly over the past few decades, with numerous contributions from theoretical physicists working on both the mathematical formulation and

observational constraints of these models. Below, we provide a summary of key areas of research and prominent contributors to each:

Cosmological Applications of $f(R)$ Gravity

The modification of gravity at cosmological scales is one of the main motivations for the study of $f(R)$ gravity. Many researchers have worked on using $f(R)$ gravity to explain the observed accelerated expansion of the Universe. The most notable of these is the work by Starobinsky, who introduced the quadratic model, Eq. (2.1), to explain cosmic acceleration without invoking dark energy. This model is widely studied in the context of inflationary cosmology [23], as it naturally leads to a de Sitter-like solution at high curvature, which can account for the observed expansion at early times [24].

In addition to Starobinsky’s work, several researchers have explored different forms of $f(R)$ gravity to address the late-time acceleration of the Universe. For example, the model by [25] introduced an $f(R)$ model that leads to a self-accelerating cosmology, without the need for dark energy. This work was influential in showing that modifications of gravity could provide an alternative explanation for accelerated cosmic expansion.

Another key direction of research in this area has been the study of the effective equation of state in $f(R)$ gravity. This work by [26] has been instrumental in deriving the cosmological field equations and understanding the behavior of the equation of state in various $f(R)$ models. These studies have shown that different forms of $f(R)$ gravity can lead to different accelerations at late times, depending on the specific form of the function $f(R)$.

Gravitational Wave Propagation in $f(R)$ Gravity

One of the key predictions of $f(R)$ gravity is that gravitational waves may propagate differently than in GR due to the additional scalar degree of freedom. The modification of the gravitational action in $f(R)$ gravity can alter the speed of gravitational waves and introduce new polarization states [27], which could be detected by current and future gravitational wave observatories.

In the context of GW propagation, a significant contribution was made by [27–30], who analyzed the propagation of gravitational waves in the $f(R)$ framework and derived the modified field equations for the perturbations of the metric. Their work showed that in the presence of $f(R)$ modifications, gravitational waves could propagate at speeds different from the speed of light, potentially leading to observable effects in GW observations from binary mergers and other astrophysical sources.

Further research by [11] extended this analysis, focusing on the perturbative approach to gravitational waves in $f(R)$ gravity, and showed how the scalar modes could affect the polarization of gravitational waves. In $f(R)$ gravity, the additional scalar degree of freedom, often referred to as the “scalaron”, modifies the propagation of gravitational waves by introducing a new polarization mode. This scalar mode arises from the fact that $f(R)$ gravity can be recast as a scalar-tensor theory, where the Ricci scalar R is coupled to a scalar field. The presence of this scalar mode leads to deviations from the standard tensor polarizations (h_+ and h_\times) predicted by General Relativity (GR), providing a unique signature that can be tested with gravitational wave detectors.

The perturbative approach to gravitational waves in $f(R)$ gravity involves analyzing the linearized field equations around a background metric. In this framework, the scalar mode manifests as a longitudinal component of the gravitational wave, which is absent in GR. This scalar mode

can affect the amplitude, phase, and propagation speed of gravitational waves, leading to observable differences in the detected signals. For example, [31] demonstrated that the scalar mode could induce a frequency-dependent modification to the gravitational wave dispersion relation, which could be detected as a phase shift in the waveform. Additionally, [32] showed that the scalar mode could alter the energy spectrum of the stochastic gravitational wave background (SGWB), providing another avenue for testing $f(R)$ gravity.

These modifications are crucial for testing $f(R)$ gravity with detectors like LIGO and Virgo, as they provide a means to distinguish between different gravity models based on the behavior of the observed GW signals. For instance, the absence of scalar polarization modes in the detected signals from binary black hole mergers has already placed stringent constraints on $f(R)$ gravity models [17]. Future observations with next-generation detectors, such as the Einstein Telescope and Cosmic Explorer, will further improve these constraints by providing higher sensitivity to the polarization and dispersion properties of gravitational waves. Moreover, the detection of the stochastic gravitational wave background (SGWB) could offer additional insights into the scalar mode, as its energy density and frequency spectrum are sensitive to the modifications introduced by $f(R)$ gravity [33].

In summary, the perturbative approach to gravitational waves in $f(R)$ gravity highlights the importance of scalar modes as a distinguishing feature of modified gravity theories. By analyzing the polarization, dispersion, and energy spectrum of gravitational waves, we can test the predictions of $f(R)$ gravity and constrain its parameters. These studies not only advance our understanding of gravity but also provide a powerful tool for probing the fundamental nature of the Universe.

Another important area of research in $f(R)$ gravity is the study of weak-field limits and solar system tests. Since GR has been experimentally verified in the weak-field regime through experiments such as the deflection of light and the precession of Mercury's orbit [34], any modification to gravity must satisfy these stringent constraints.

Solar System Tests and Weak-Field Constraints

Recent advancements in cosmology have led to significant developments in alternative gravity models, challenging the standard assumption that GR holds on all scales. While GR has been rigorously tested within the Solar System, the discovery of the Universe's accelerated expansion has prompted new questions about its validity at cosmological distances. This acceleration, often attributed to dark energy within the Λ CDM framework [35], could instead suggest that GR itself requires modification on large scales. To explore this possibility, numerous modified gravity theories, such as $f(R)$ gravity, Horndeski theory, and massive gravity, have been proposed. These models introduce novel mechanisms to reconcile deviations from GR while passing Solar System tests through screening effects. In this context, [36] provides a comprehensive review of recent progress in constructing modified gravity models and developing cosmological tests to distinguish them from standard GR-based cosmology.

In recent years, various modifications to gravity have been proposed as alternatives to the dark matter hypothesis [5, 37–40]. These modified gravity theories introduce additional apparent force terms in the dynamical equations, which aim to reproduce the observed gravitational effects without invoking unseen mass components. Typically, such extra force terms scale with the radial coordinate, becoming significant on galactic or cosmological scales to account for the missing

mass problem. However, their effects may still manifest within smaller-scale structures, such as the solar system. A detailed investigation of these effects has been carried out in [41], where they analytically derive general expressions for the contribution of modified gravity theories to planetary precession in the solar system. The study examines three popular models—Modified Newtonian Dynamics (MOND), Emergent Gravity (EG), and Modified Gravity (MG)—and places constraints on MOND interpolating functions using solar system data.

Works like [42] have developed methods to study the weak-field limits of $f(R)$ gravity, showing that for most forms of $f(R)$, the modifications to GR are negligible in the solar system. This is due to the so-called Chameleon Mechanism³ [44–46], which ensures that the scalar degree of freedom in $f(R)$ gravity is effectively screened in high-density environments like the Earth and Sun. The development of these screening mechanisms has been crucial for ensuring that $f(R)$ gravity is consistent with solar system observations.

The Chameleon Mechanism

Modified gravity theories, such as $f(R)$ gravity, often introduce additional scalar degrees of freedom. A major challenge is ensuring that these scalar fields remain undetectable in high-density environments, such as the Solar System, while still influencing cosmological dynamics. The Chameleon Mechanism provides a solution by allowing the scalar field to have a mass that depends on the local matter density. This ensures that modifications to gravity are screened in dense regions but manifest in low-density environments such as cosmic voids.

In $f(R)$ gravity, the action is given by Eq. (1.2). This can be rewritten in a scalar-tensor form by introducing an auxiliary field ϕ , defined as:

$$\phi = f_R \equiv \frac{df}{dR}. \quad (2.3)$$

The equation of motion for ϕ is:

$$\square\phi = \frac{dV_{\text{eff}}}{d\phi}, \quad (2.4)$$

where the effective potential is [47]:

$$V_{\text{eff}}(\phi) = V(\phi) + \rho_m e^{\beta\phi}, \quad (2.5)$$

where $V(\phi)$ is the potential governing the scalar field dynamics, and ρ_m is the local matter density.

The key feature of the Chameleon Mechanism is the density-dependent mass:

$$m_\phi^2 = \frac{d^2V_{\text{eff}}}{d\phi^2}. \quad (2.6)$$

- **High-density regions** (e.g., the Earth, Solar System): m_ϕ is large, meaning the field is short-ranged, suppressing deviations from GR.
- **Low-density regions** (e.g., cosmic voids): m_ϕ is small, allowing the field to mediate long-range fifth forces.

³Check Sec. 3.5 in [43] for detailed discussion about the mechanism.




Figure 1: The effective mass of the chameleon scalar field (m_ϕ) as a function of the local matter density (ρ). The shaded regions indicate different astrophysical environments where the mechanism operates.

The Chameleon Mechanism is a screening mechanism in modified gravity theories that allows scalar fields to evade local experimental constraints while remaining cosmologically relevant. The key feature of this mechanism is that the effective mass of the scalar field depends on the local matter density: in high-density regions, such as within the Solar System, the field acquires a large mass, making its interaction range extremely short and suppressing deviations from GR. Conversely, in low-density cosmic environments, such as voids, the field remains light, allowing modifications to gravity to manifest over astrophysical scales. Figure 1 illustrates this behavior by showing how the Chameleon field’s mass increases with ambient density. The transition between different astrophysical environments highlights the environmental dependence of the screening, with strong suppression in high-density regions (e.g., Earth) and significant effects in low-density regions (e.g., intergalactic space). This behavior is crucial for understanding the viability of Chameleon-based modified gravity models and their observational signatures in gravitational wave experiments and large-scale structure surveys. The Chameleon Mechanism significantly impacts the formation of structures in the Universe:

- **Cosmic Large-Scale Structure:** The mechanism modifies gravitational clustering in cosmic voids, leading to scale-dependent growth.
- **Weak Lensing:** Deviations from GR appear as modifications to the lensing potential.
- **Cosmic Microwave Background (CMB)** Chameleon fields can alter the Integrated Sachs-Wolfe (ISW) effect.

Observations and experiments place stringent constraints on Chameleon fields. The Cassini spacecraft provided a Shapiro time delay [48] constraint [49]:

$$\gamma - 1 = (2.1 \pm 2.3) \times 10^{-5}. \quad (2.7)$$

This limits the deviation from GR induced by the scalar field. Additionally, experiments such as the Eöt-Wash experiment and atom interferometry tests [50] probe deviations in Newton's law. The force mediated by ϕ obeys:

$$F_\phi \propto \frac{e^{-m_\phi r}}{r^2}. \quad (2.8)$$

Current experiments limit m_ϕ in the range $[10^{-3} - 10^{-2}]$ eV [51].

The Chameleon Mechanism is a compelling way to reconcile modified gravity with local tests of GR. It ensures that scalar fields are screened in high-density environments while allowing deviations in cosmic voids. Future experiments, particularly in cosmology and laboratory physics, will provide deeper insights into the nature of scalar-tensor gravity theories.

3 Constraints on $f(R)$ Gravity from Gravitational Wave Measurements

The viability of $f(R)$ gravity theories is strongly constrained by both gravitational wave (GW) observations and other astrophysical and cosmological experiments. In this section, we discuss how GW measurements from binary mergers and stochastic gravitational wave background (SGWB) non-detections, as well as other experimental data, place limits on the parameters and predictions of $f(R)$ gravity.

3.1 Constraints from Binary Black Hole and Neutron Star Mergers

The detection of GWs from binary black hole (BBH) and binary neutron star (BNS) mergers by LIGO and Virgo has provided a wealth of data to test alternative theories of gravity. In $f(R)$ gravity, the modifications to the Einstein-Hilbert action can lead to deviations in the propagation speed, polarization modes, and dispersion relations of GWs [27, 52, 53]. These deviations can be compared with observational data to constrain the functional form of $f(R)$.

Propagation Speed of Gravitational Waves

In GR, GWs propagate at the speed of light, c . However, in $f(R)$ gravity, the propagation speed v_{GW} can differ from c due to the additional degrees of freedom introduced by the $f(R)$ term [54, 55]. The modified speed can be expressed as:

$$v_{\text{GW}}^2 = c^2 \left(1 + \frac{\Delta f(R)}{R} \right), \quad (3.1)$$

where $\Delta f(R)$ represents the deviation from the GR Lagrangian. The observation of GW170817 [17], a BNS merger accompanied by a gamma-ray burst, provided a stringent constraint on the propagation speed of GWs. The time delay between the GW and electromagnetic signals constrained the difference between v_{GW} and c to be:

$$-3 \times 10^{-15} \leq \frac{v_{\text{GW}} - c}{c} \leq 7 \times 10^{-16}. \quad (3.2)$$

This result rules out many $f(R)$ models that predict significant deviations in the GW propagation speed [17].

Polarization Modes

In GR, GWs have two tensor polarization modes: h_+ and h_\times . In $f(R)$ gravity, additional scalar polarization modes can arise due to the scalar degree of freedom associated with the Ricci scalar R [27, 52]. The presence of these extra modes can be tested using the network of GW detectors. For example, the cross-correlation of data from LIGO and Virgo can be used to search for scalar modes. Current non-detections of scalar polarization place constraints on the amplitude of these modes, which in turn constrain the parameters of $f(R)$ gravity [33].

Dispersion Relations

In $f(R)$ gravity, the dispersion relation for GWs can be modified [56–60], leading to frequency-dependent effects. The modified dispersion relation can be written as:

$$\omega^2 = k^2 c^2 \left(1 + \alpha \left(\frac{k}{k_0} \right)^n \right), \quad (3.3)$$

where ω is the angular frequency, k is the wavenumber, α is a dimensionless parameter, and k_0 is a reference scale. The parameter α is constrained by the observed phase coherence of GW signals from BBH mergers. Current constraints from LIGO-Virgo data suggest that $|\alpha| \lesssim 10^{-15}$ for $n = 2$ [61].

3.2 Constraints from Stochastic Gravitational Wave Background (SGWB)

The stochastic gravitational wave background (SGWB) is a superposition of GWs from unresolved astrophysical and cosmological sources. In $f(R)$ gravity, the SGWB spectrum can differ from the GR prediction due to modifications in the early Universe dynamics and GW propagation. Current non-detections of the SGWB by LIGO and Virgo place constraints on the energy density of GWs, which can be translated into constraints on $f(R)$ gravity parameters.

The energy density of the SGWB is typically expressed in terms of the dimensionless parameter $\Omega_{\text{GW}}(f)$, defined as:

$$\Omega_{\text{GW}}(f) = \frac{1}{\rho_c} \frac{d\rho_{\text{GW}}}{d \ln f}, \quad (3.4)$$

where ρ_c is the critical energy density of the Universe and ρ_{GW} is the energy density of GWs. In $f(R)$ gravity, the predicted $\Omega_{\text{GW}}(f)$ can be enhanced or suppressed depending on the functional form of $f(R)$. Current upper limits from LIGO-Virgo observations constrain $\Omega_{\text{GW}}(f) \lesssim 10^{-8}$ at frequencies around 25 Hz, which rules out some $f(R)$ models that predict a strong SGWB signal [62].

3.3 Other Experimental Constraints

In addition to GW measurements, $f(R)$ gravity is constrained by a variety of other astrophysical and cosmological observations. These include solar system tests, cosmological surveys, and laboratory experiments.

Cosmological Surveys

Cosmological surveys, such as those measuring the cosmic microwave background (CMB) and large-scale structure (LSS), provide additional constraints on $f(R)$ gravity. For example, the CMB power spectrum is sensitive to the growth of structure, which can be modified in $f(R)$ gravity. Current constraints from the Planck satellite and other surveys rule out many $f(R)$ models that predict significant deviations from GR on cosmological scales [5, 35, 55, 63–65].

Laboratory Experiments

Laboratory experiments, such as those testing the equivalence principle or measuring the Newtonian gravitational constant G , can also constrain $f(R)$ gravity. For example, precision measurements of the equivalence principle using torsion balances constrain the coupling between the scalar degree of freedom in $f(R)$ gravity and matter. Current limits on the equivalence principle violation are at the level of $\Delta a/a \lesssim 10^{-13}$, which places strong constraints on the scalar-tensor nature of $f(R)$ gravity [66].

The constraints on $f(R)$ gravity from GW measurements and other experiments are summarized in Table 1. These constraints collectively rule out many $f(R)$ models and provide guidance for the development of viable alternatives to GR.

Table 1: Summary of constraints on $f(R)$ gravity.

Experiment	Constraint	Reference
GW170817 (BNS merger)	$ v_{\text{GW}} - c /c \lesssim 10^{-15}$	[17]
LIGO-Virgo (polarization)	No scalar modes detected	[33]
LIGO-Virgo (dispersion)	$ \alpha \lesssim 10^{-15}$	[33]
SGWB (LIGO-Virgo)	$\Omega_{\text{GW}}(f) \lesssim 10^{-8}$	[62]
Solar system (PPN)	$ \gamma - 1 \lesssim 2 \times 10^{-5}$	[49]
CMB (Planck)	Growth of structure constraints	[35]
Laboratory (equivalence principle)	$\Delta a/a \lesssim 10^{-13}$	[66]

4 Future Improvements in Constraining $f(R)$ Gravity

The constraints on $f(R)$ gravity from current gravitational wave (GW) observations and other experiments are already stringent, but significant improvements are expected in the coming decades. These improvements will be driven by advances in GW detector sensitivity, the development of new observational techniques, and the integration of multi-messenger astronomy. In this section, we discuss the prospects for future constraints on $f(R)$ gravity and the technological and theoretical developments necessary to achieve these improvements.

4.1 Next-Generation Gravitational Wave Detectors

The next generation of GW detectors, such as the Einstein Telescope (ET) [67, 68] and Cosmic Explorer (CE) [69], will provide unprecedented sensitivity to GW signals. These detectors will be able to observe GWs from BBH and BNS mergers at much greater distances and with higher precision, enabling more stringent tests of $f(R)$ gravity.

Improved Sensitivity and Frequency Range

The Einstein Telescope and Cosmic Explorer are designed to have a sensitivity improvement of up to a factor of 10 [70] compared to current detectors like LIGO and Virgo. This improvement will allow for the detection of GWs with signal-to-noise ratios (SNRs) [71] that are an order of magnitude higher, enabling more precise measurements of GW parameters. The improved sensitivity will also extend to lower frequencies, down to ~ 1 Hz, which is particularly important for testing $f(R)$ gravity, as many models predict deviations in the low-frequency regime.

The noise power spectral density $S_n(f)$ of a GW detector is a key factor in determining its sensitivity. For ET and CE, the noise curve is expected to be [17]:

$$S_n(f) = S_0 \left[\left(\frac{f}{f_0} \right)^{-4} + 2 \left(\frac{f}{f_0} \right)^{-2} + 1 + \left(\frac{f}{f_0} \right)^2 \right], \quad (4.1)$$

where S_0 is the minimum noise level and f_0 is a reference frequency. The improved sensitivity will allow for the detection of GWs from sources at cosmological distances, providing a larger sample of events for testing $f(R)$ gravity [72].

Testing the Propagation Speed of GWs

The improved sensitivity of next-generation detectors [72, 73] will allow for more precise measurements of the propagation speed of GWs. In particular, the time delay between GW signals and electromagnetic counterparts from BNS mergers can be measured with sub-millisecond precision. This will constrain the deviation of the GW speed from the speed of light to:

$$\frac{|v_{\text{GW}} - c|}{c} \lesssim 10^{-17}. \quad (4.2)$$

Such precision will rule out a wide range of $f(R)$ models that predict significant deviations in the GW propagation speed [74].

Probing Additional Polarization Modes

Next-generation detectors will also improve the ability to probe additional polarization modes predicted by $f(R)$ gravity. The increased number of detectors and their improved sensitivity will allow for better separation of the tensor and scalar modes. The cross-correlation of data from multiple detectors can be used to search for scalar modes with amplitudes as small as:

$$h_{\text{scalar}} \lesssim 10^{-20}. \quad (4.3)$$

This will provide stringent constraints on the scalar degree of freedom in $f(R)$ gravity [33].

4.2 Stochastic Gravitational Wave Background (SGWB)

The detection of the stochastic gravitational wave background (SGWB) is a major goal for future GW observatories. The SGWB is expected to contain contributions from both astrophysical and cosmological sources, and its spectrum can provide important constraints on $f(R)$ gravity.

Improved Sensitivity to the SGWB

Next-generation detectors will have improved sensitivity to the SGWB, with expected upper limits on the energy density $\Omega_{\text{GW}}(f)$ of:

$$\Omega_{\text{GW}}(f) \lesssim 10^{-12} \quad \text{at } f \sim 10 \text{ Hz.} \quad (4.4)$$

This improvement will allow for the detection of SGWB signals from early Universe processes, such as inflation or phase transitions, which can be used to test $f(R)$ gravity models [62]. There are many successful progress in testing $f(R)$ models in the early universe cosmology [75–81].

Cross-Correlation Techniques

The use of cross-correlation techniques between multiple detectors will further improve the sensitivity to the SGWB. The cross-correlation statistic $C(f)$ is given by:

$$C(f) = \frac{1}{T} \int_0^T dt s_1(t)s_2(t), \quad (4.5)$$

where $s_1(t)$ and $s_2(t)$ are the signals from two detectors, and T is the observation time. The improved sensitivity will allow for the detection of SGWB signals with amplitudes as small as:

$$h_{\text{SGWB}} \lesssim 10^{-18}. \quad (4.6)$$

This will provide stringent constraints on the early Universe dynamics predicted by $f(R)$ gravity [82].

4.3 Multi-Messenger Astronomy

Multi-messenger astronomy, the simultaneous observation of cosmic events using GWs, electromagnetic (EM) waves, neutrinos, and cosmic rays, has emerged as a powerful tool for testing theories of gravity. By combining data from multiple messengers, researchers can probe the properties of gravity in regimes that are inaccessible to single-messenger observations. In this section, we discuss how multi-messenger astronomy can constrain modified gravity theories, with a focus on $f(R)$ gravity.

Gravitational Waves and Electromagnetic Counterparts

The detection of GWs from binary neutron star (BNS) mergers, such as GW170817, and their associated EM counterparts, such as gamma-ray bursts (GRBs) and kilonovae, has provided a unique opportunity to test modified gravity theories. In GR, GWs and EM waves propagate at the same speed, c . However, in modified gravity theories like $f(R)$ gravity, the propagation speed of GWs, v_{GW} , can differ from c .

The time delay between the arrival of GWs and EM signals from a BNS merger can be used to constrain v_{GW} . For GW170817, the observed time delay was $\Delta t \lesssim 1.7$ seconds, corresponding to a constraint on the GW speed deviation:

$$\frac{|v_{\text{GW}} - c|}{c} \lesssim 10^{-15}. \quad (4.7)$$

This result rules out many $f(R)$ models that predict significant deviations in v_{GW} [17, 64].

Neutrino Observations

Neutrinos, produced in astrophysical events such as supernovae and BNS mergers, provide another messenger for testing modified gravity. In $f(R)$ gravity, the additional scalar degree of freedom can couple to neutrinos, affecting their propagation and energy spectrum. Observations of neutrinos from astrophysical sources can therefore constrain the parameters of $f(R)$ gravity.

For example, the timing and energy spectrum of neutrinos from supernovae can be compared with the predictions of $f(R)$ gravity. Any deviations from the expected neutrino signal can be used to constrain the scalar coupling in $f(R)$ gravity [83, 84].

Cosmic Rays and High-Energy Astrophysics

Cosmic rays, high-energy particles originating from astrophysical sources, can also be used to test modified gravity. In $f(R)$ gravity, the propagation of cosmic rays can be affected by the modified spacetime geometry and the additional scalar field. Observations of cosmic ray spectra and anisotropies can therefore provide constraints on $f(R)$ gravity.

For example, the Pierre Auger Observatory has measured the energy spectrum and arrival directions of ultra-high-energy cosmic rays (UHECRs). These observations can be compared with the predictions of $f(R)$ gravity to constrain the scalar field parameters [85, 86].

Combining Multi-Messenger Data

The combination of data from multiple messengers provides a more comprehensive test of modified gravity theories. For example, the simultaneous observation of GWs, EM waves, and neutrinos from a BNS merger can be used to constrain the propagation speed, polarization, and dispersion relations of GWs in $f(R)$ gravity.

The cross-correlation of data from different messengers can also improve the sensitivity to deviations from GR. For example, the cross-correlation of GW and neutrino data from a supernova can provide tighter constraints on the scalar coupling in $f(R)$ gravity [83, 87].

Future multi-messenger observations with next-generation detectors, such as the Einstein Telescope (ET), Cosmic Explorer (CE), and the Laser Interferometer Space Antenna (LISA), will provide even stronger constraints on modified gravity theories. These detectors will have improved sensitivity to GWs, allowing for more precise measurements of v_{GW} and the polarization modes of GWs.

In addition, future neutrino detectors, such as the Deep Underground Neutrino Experiment (DUNE) and the Hyper-Kamiokande (Hyper-K), will provide more precise measurements of neutrino spectra and timing. These observations will further constrain the parameters of $f(R)$ gravity and other modified theories [88, 89].

Multi-messenger astronomy provides a powerful tool for testing modified gravity theories like $f(R)$ gravity. By combining data from GWs, EM waves, neutrinos, and cosmic rays, researchers can probe the properties of gravity in regimes that are inaccessible to single-messenger observations. Future multi-messenger observations with next-generation detectors will provide even stronger constraints on modified gravity, advancing our understanding of the fundamental nature of gravity and the Universe.

4.4 Theoretical Developments

In addition to observational improvements, theoretical developments will be necessary to fully exploit the potential of future GW observations. These developments include improved waveform models, a better understanding of the astrophysical sources, and more accurate predictions for the SGWB.

Waveform Models

Accurate waveform models are essential for extracting information from GW signals. In $f(R)$ gravity, the waveform models must account for the modified propagation speed, polarization modes, and dispersion relations. The development of accurate waveform models for $f(R)$ gravity will require advances in numerical relativity and perturbation theory [90].

Astrophysical Source Models

A better understanding of the astrophysical sources of GWs, such as BBH and BNS mergers, is also necessary. This includes improved models for the formation and evolution of these systems, as well as their electromagnetic and neutrino counterparts. These models will be essential for interpreting the GW signals and testing $f(R)$ gravity [91].

SGWB Predictions

Accurate predictions for the SGWB in $f(R)$ gravity are also necessary. This includes predictions for the energy density and frequency spectrum of the SGWB from both astrophysical and cosmological sources. These predictions will be essential for interpreting the SGWB observations and testing $f(R)$ gravity [92].

The future prospects for constraining $f(R)$ gravity are summarized in Table 2. These prospects include improvements in GW detector sensitivity, the detection of the SGWB, and the integration of multi-messenger astronomy. These developments will provide stringent tests of $f(R)$ gravity and advance our understanding of the fundamental nature of gravity.

Table 2: Summary of future prospects for constraining $f(R)$ gravity.

Improvement	Expected Constraint	Reference
Next-generation detectors	$ v_{\text{GW}} - c /c \lesssim 10^{-17}$	[74]
SGWB detection	$\Omega_{\text{GW}}(f) \lesssim 10^{-12}$	[62]
Multi-messenger astronomy	$ v_{\text{GW}} - c /c \lesssim 10^{-18}$	[17]
Waveform models	Improved accuracy for $f(R)$ gravity	[90]
Astrophysical source models	Better understanding of BBH/BNS mergers	[91]
SGWB predictions	Accurate predictions for $f(R)$ gravity	[92]

5 Other Modified Theories of Gravity and Their Constraints

While $f(R)$ gravity is one of the most widely studied modified theories of gravity, it is by no means the only alternative to General Relativity (GR). In this section, we explore other prominent

modified theories of gravity, including scalar-tensor theories, Horndeski gravity, massive gravity, and brane-world models. We compare their theoretical frameworks, predictions, and observational constraints with those of $f(R)$ gravity.

5.1 Scalar-Tensor Theories

Scalar-tensor theories are among the oldest and most well-studied alternatives to GR. These theories introduce an additional scalar field ϕ that couples to the metric tensor $g_{\mu\nu}$, modifying the gravitational interaction. The inclusion of the scalar field allows for a richer phenomenology⁴, including varying gravitational constants, additional forces, and deviations from GR in both weak-field and strong-field regimes [93].

Theoretical Framework

The action for scalar-tensor theories can be written as:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2\kappa} f(\phi) R - \frac{1}{2} \omega(\phi) (\nabla\phi)^2 - V(\phi) + \mathcal{L}_m \right), \quad (5.1)$$

where $f(\phi)$ is a function of the scalar field that couples to the Ricci scalar R , $\omega(\phi)$ determines the kinetic term of the scalar field, and $V(\phi)$ is the scalar potential. The term \mathcal{L}_m represents the matter Lagrangian, which is assumed to be minimally coupled to the metric. The scalar field ϕ mediates an additional force, which can lead to deviations from GR in both the weak-field and strong-field regimes.

The field equations derived from this action are:

$$f(\phi) G_{\mu\nu} + \nabla_\mu \nabla_\nu f(\phi) - g_{\mu\nu} \square f(\phi) = \kappa T_{\mu\nu} + \omega(\phi) \left(\nabla_\mu \phi \nabla_\nu \phi - \frac{1}{2} g_{\mu\nu} (\nabla\phi)^2 \right) - g_{\mu\nu} V(\phi), \quad (5.2)$$

where $G_{\mu\nu}$ is the Einstein tensor, $\square \equiv g^{\mu\nu} \nabla_\mu \nabla_\nu$ is the d'Alembertian, and $T_{\mu\nu}$ is the stress-energy tensor of matter. The scalar field equation is given by:

$$\square\phi + \frac{1}{2\omega(\phi)} \frac{d\omega(\phi)}{d\phi} (\nabla\phi)^2 - \frac{1}{2\omega(\phi)} \frac{df(\phi)}{d\phi} R - \frac{1}{\omega(\phi)} \frac{dV(\phi)}{d\phi} = 0. \quad (5.3)$$

These equations reduce to those of GR in the limit where $f(\phi) = 1$, $\omega(\phi) = 0$, and $V(\phi) = 0$.

Observational Constraints

Scalar-tensor theories are tightly constrained by solar system tests, such as the parameterized post-Newtonian (PPN) formalism. The PPN parameter γ , which measures the curvature of space-time, is constrained to be $|\gamma - 1| \lesssim 2 \times 10^{-5}$ by solar system observations [49]. This places strong constraints on the coupling function $f(\phi)$ and the kinetic term $\omega(\phi)$. For example, in the Brans-Dicke theory, where $f(\phi) = \phi$ and $\omega(\phi) = \omega_{\text{BD}}$, the constraint translates to $\omega_{\text{BD}} \gtrsim 40,000$ [94].

Gravitational wave (GW) observations also constrain scalar-tensor theories. The absence of scalar polarization modes in GW signals from binary black hole (BBH) mergers rules out many scalar-tensor models that predict significant scalar radiation [95]. The recent detection of GW170817, a binary neutron star merger (BNS), further constrains the speed of gravitational

⁴Check Appendix A in [43] for a discussion on $f(R)$ gravity's equivalence to scalar-tensor gravity.

waves to be nearly equal to the speed of light, ruling out many scalar-tensor theories with large deviations from GR [17].

In addition to these constraints, cosmological observations, such as the cosmic microwave background (CMB) and large-scale structure, provide further tests of scalar-tensor theories. For instance, the evolution of the scalar field ϕ during the early universe can affect the growth of cosmological perturbations, leading to observable signatures in the CMB power spectrum [20].

5.2 Horndeski Gravity

Horndeski gravity represents the most general scalar-tensor theory with second-order field equations, ensuring the absence of ghost instabilities [96,97]. It encompasses a wide range of modified gravity models, including $f(R)$ gravity, Brans-Dicke theory, and other scalar-tensor theories. This framework is particularly significant in cosmology, as it provides a versatile platform for exploring dark energy, inflation, and modifications to general relativity [5].

Theoretical Framework

The action for Horndeski gravity is given by:

$$S = \int d^4x \sqrt{-g} (\mathcal{L}_2 + \mathcal{L}_3 + \mathcal{L}_4 + \mathcal{L}_5 + \mathcal{L}_m), \quad (5.4)$$

where g is the determinant of the metric tensor $g_{\mu\nu}$, and \mathcal{L}_m represents the matter Lagrangian. The Lagrangians \mathcal{L}_2 to \mathcal{L}_5 are functions of the scalar field ϕ , its kinetic term $X \equiv -\frac{1}{2}\nabla_\mu\phi\nabla^\mu\phi$, and the Ricci scalar R . These Lagrangians are explicitly defined as follows:

$$\mathcal{L}_2 = G_2(\phi, X), \quad (5.5)$$

$$\mathcal{L}_3 = G_3(\phi, X)\square\phi, \quad (5.6)$$

$$\mathcal{L}_4 = G_4(\phi, X)R - 2G_{4,X}(\phi, X) [(\square\phi)^2 - \nabla_\mu\nabla_\nu\phi\nabla^\mu\nabla^\nu\phi], \quad (5.7)$$

$$\mathcal{L}_5 = G_5(\phi, X)G_{\mu\nu}\nabla^\mu\nabla^\nu\phi + \frac{1}{3}G_{5,X}(\phi, X) [(\square\phi)^3 - 3\square\phi\nabla_\mu\nabla_\nu\phi\nabla^\mu\nabla^\nu\phi + 2\nabla_\mu\nabla_\nu\phi\nabla^\nu\nabla^\alpha\phi\nabla_\alpha\nabla^\mu\phi], \quad (5.8)$$

where $G_i(\phi, X)$ ($i = 2, 3, 4, 5$) are arbitrary functions of ϕ and X , and $G_{i,X} \equiv \partial G_i/\partial X$. The term $\square\phi \equiv \nabla_\mu\nabla^\mu\phi$ denotes the d'Alembertian of the scalar field.

The second-order nature of the field equations ensures that the theory avoids Ostrogradsky instabilities, which are common in higher-derivative theories [98]. The field equations for the metric and the scalar field are derived by varying the action with respect to $g_{\mu\nu}$ and ϕ , respectively. These equations are:

$$\mathcal{E}_{\mu\nu} = \frac{1}{\sqrt{-g}} \frac{\delta S}{\delta g^{\mu\nu}} = 0, \quad \mathcal{E}_\phi = \frac{1}{\sqrt{-g}} \frac{\delta S}{\delta \phi} = 0, \quad (5.9)$$

where $\mathcal{E}_{\mu\nu}$ and \mathcal{E}_ϕ are the Einstein and scalar field equations, respectively.

Cosmological Implications

Horndeski gravity has profound implications for cosmology. For instance, the functions $G_i(\phi, X)$ can be tuned to reproduce the observed accelerated expansion of the universe without invoking

a cosmological constant [55]. The theory also allows for self-accelerating solutions, where the scalar field drives the acceleration dynamically. Additionally, Horndeski gravity modifies the propagation of gravitational waves, leading to observable signatures in the cosmic microwave background and large-scale structure [99].

Constraints and Observational Tests

Observational constraints on Horndeski gravity arise from solar system tests, gravitational wave observations, and cosmological data. For example, the speed of gravitational waves, c_{GW} , is tightly constrained by the detection of GW170817 and its electromagnetic counterpart [17]. In Horndeski gravity, c_{GW} is given by:

$$c_{\text{GW}}^2 = \frac{G_4 - XG_{5,\phi} - X\ddot{\phi}G_{5,X}}{G_4 - 2XG_{4,X} + XG_{5,\phi} - \frac{1}{2}X^2G_{5,X}}, \quad (5.10)$$

where $\ddot{\phi}$ is the second time derivative of the scalar field. The near-equality of c_{GW} and the speed of light strongly restricts the functional forms of G_4 and G_5 [100].

Horndeski gravity is constrained by both solar system tests and GW observations. The PPN parameter γ is constrained to be $|\gamma - 1| \lesssim 2 \times 10^{-5}$, similar to scalar-tensor theories [49]. Additionally, the propagation speed of GWs in Horndeski gravity is constrained by the observation of GW170817, which requires $|v_{\text{GW}} - c|/c \lesssim 10^{-15}$ [17].

The absence of scalar polarization modes in GW signals also places constraints on Horndeski gravity. Current limits on the amplitude of scalar modes are at the level of $h_{\text{scalar}} \lesssim 10^{-20}$ [33].

5.3 Massive Gravity

Massive gravity is a modified theory of gravity in which the graviton, the hypothetical quantum of GWs, possesses a non-zero mass. Unlike GR, where the graviton is massless, massive gravity introduces a mass term that modifies the propagation of GWs and the large-scale structure of the universe. This theory has gained significant attention as a potential explanation for cosmic acceleration without invoking dark energy [101, 102].

Theoretical Framework

The action for massive gravity can be written as:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2\kappa} R + m_g^2 \mathcal{U}(g, f) + \mathcal{L}_m \right), \quad (5.11)$$

where $\kappa = 8\pi G$ is the gravitational constant, R is the Ricci scalar, $g_{\mu\nu}$ is the physical metric, $f_{\mu\nu}$ is a fixed reference metric, m_g is the graviton mass, and $\mathcal{U}(g, f)$ is a potential term that encodes the interaction between the physical and reference metrics. The term \mathcal{L}_m represents the matter Lagrangian.

The potential $\mathcal{U}(g, f)$ is typically constructed to avoid ghost instabilities and is often expressed in terms of the eigenvalues of the matrix $\mathcal{K}_\nu^\mu = \delta_\nu^\mu - \sqrt{g^{\mu\alpha} f_{\alpha\nu}}$. A common form for the potential is [103]:

$$\mathcal{U}(g, f) = \sum_{n=0}^4 \beta_n \mathcal{U}_n(\mathcal{K}), \quad (5.12)$$

where β_n are free parameters, and $\mathcal{U}_n(\mathcal{K})$ are elementary symmetric polynomials of the eigenvalues of \mathcal{K} . For example, the first two polynomials are:

$$\mathcal{U}_1(\mathcal{K}) = [\mathcal{K}], \quad \mathcal{U}_2(\mathcal{K}) = [\mathcal{K}]^2 - [\mathcal{K}^2], \quad (5.13)$$

where $[\mathcal{K}]$ denotes the trace of \mathcal{K} .

Modifications to Gravitational Waves

In massive gravity, the presence of a graviton mass modifies the dispersion relation for gravitational waves. In GR, the dispersion relation for a massless graviton is:

$$\omega^2 = k^2, \quad (5.14)$$

where ω is the angular frequency and k is the wavenumber. In massive gravity, this relation becomes:

$$\omega^2 = k^2 + m_g^2, \quad (5.15)$$

where m_g is the graviton mass. This modification leads to a frequency-dependent phase velocity for GWs, which can be tested using observations from gravitational wave detectors such as LIGO and Virgo.

The modified dispersion relation also affects the propagation of GWs over cosmological distances. For a GW with frequency f , the time delay Δt compared to a massless graviton is given by [17, 71, 103–105]:

$$\Delta t \approx \frac{m_g^2}{2H_0} \int_0^z \frac{dz'}{(1+z')^2 E(z')}, \quad (5.16)$$

where H_0 is the Hubble constant, z is the redshift, and $E(z)$ is the dimensionless Hubble parameter. This time delay can be used to constrain the graviton mass using multi-messenger observations of GWs and their electromagnetic counterparts [106].

Cosmological Implications

Massive gravity has significant implications for cosmology. The theory can naturally explain the observed accelerated expansion of the universe without requiring dark energy. The graviton mass term acts as an effective cosmological constant on large scales, leading to a late-time acceleration of the universe [107].

The modified Friedmann equations in massive gravity take the form [108, 109]:

$$H^2 = \frac{8\pi G}{3}\rho + \frac{m_g^2}{6}\mathcal{U}(a), \quad (5.17)$$

where H is the Hubble parameter, ρ is the energy density of matter, and $\mathcal{U}(a)$ is a function of the scale factor a that arises from the potential term. This modification can lead to deviations from the standard Λ CDM model, which can be tested using cosmological observations such as the cosmic microwave background (CMB) and large-scale structure surveys [110].

Observational Constraints

Massive gravity is constrained by both solar system tests and GW observations. The PPN parameter γ is constrained to be $|\gamma - 1| \lesssim 2 \times 10^{-5}$, similar to other modified theories [49]. Additionally, the graviton mass m_g is constrained by the observation of GWs from BBH mergers. The current upper limit on the graviton mass is $m_g \lesssim 10^{-23}$ eV [111].

5.4 Brane-World Models

Brane-world models are a class of theoretical frameworks in which our four-dimensional universe is envisioned as a brane embedded in a higher-dimensional space, often referred to as the “bulk” [112, 113]. These models provide a compelling approach to addressing some of the outstanding issues in modern cosmology, such as the hierarchy problem [112, 114] and the nature of dark energy, while recovering General Relativity (GR) on small scales. The key idea is that while gravity propagates in the full higher-dimensional space, Standard Model fields are confined to the brane, leading to modifications of gravity on cosmological scales.

Theoretical Framework

The action for brane-world models typically includes contributions from the bulk, the brane, and matter fields confined to the brane. A general form of the action can be written as:

$$S = \int d^5x \sqrt{-g^{(5)}} \left(\frac{1}{2\kappa_5^2} R^{(5)} + \mathcal{L}_{\text{bulk}} \right) + \int d^4x \sqrt{-g} (\mathcal{L}_{\text{brane}} + \mathcal{L}_m), \quad (5.18)$$

where $g^{(5)}$ and $R^{(5)}$ are the metric determinant and Ricci scalar in the five-dimensional bulk, respectively, κ_5 is the five-dimensional gravitational constant, $\mathcal{L}_{\text{bulk}}$ describes the dynamics of the bulk fields, g is the induced metric on the brane, $\mathcal{L}_{\text{brane}}$ encodes the brane’s dynamics, and \mathcal{L}_m represents the matter fields confined to the brane [115].

The induced metric on the brane is related to the bulk metric via:

$$g_{\mu\nu} = g_{\mu\nu}^{(5)} - n_\mu n_\nu, \quad (5.19)$$

where n_μ is the unit normal vector to the brane. The effective four-dimensional gravitational equations on the brane are derived by projecting the five-dimensional Einstein equations onto the brane, leading to the modified Einstein equations:

$$G_{\mu\nu} = \kappa_4^2 T_{\mu\nu} + \kappa_5^4 \Pi_{\mu\nu} - E_{\mu\nu}, \quad (5.20)$$

where $G_{\mu\nu}$ is the Einstein tensor on the brane, $T_{\mu\nu}$ is the stress-energy tensor of matter on the brane, $\Pi_{\mu\nu}$ is a quadratic function of $T_{\mu\nu}$, and $E_{\mu\nu}$ is the projection of the bulk Weyl tensor onto the brane, representing the influence of the bulk geometry on the brane [113].

Cosmological Implications

Brane-world models have profound implications for cosmology. For instance, the modified Friedmann equation on the brane takes the form [116, 117]:

$$H^2 = \frac{8\pi G}{3} \rho \left(1 + \frac{\rho}{2\lambda} \right) + \frac{\Lambda_4}{3} + \frac{\mathcal{E}}{a^4}, \quad (5.21)$$

where H is the Hubble parameter, ρ is the energy density of matter on the brane, λ is the brane tension, Λ_4 is the effective four-dimensional cosmological constant, and \mathcal{E} is a term arising from the bulk Weyl tensor, often interpreted as “dark radiation” [118]. This equation deviates from the standard Friedmann equation at high energies ($\rho \gg \lambda$), potentially altering the dynamics of the early universe.

Moreover, brane-world models can provide a natural explanation for the observed acceleration of the universe. The additional terms in the modified Friedmann equation can mimic the effects of dark energy, offering an alternative to the cosmological constant or scalar field models [119].

Challenges and Open Questions

Despite their theoretical appeal, brane-world models face several challenges. One major issue is the stabilization of the extra dimension, which requires fine-tuning of parameters to avoid conflicts with observational data. Additionally, the nature of the bulk geometry and its influence on the brane dynamics remain poorly understood. Future observations, such as those from GW detectors and precision cosmology experiments, may provide further insights into the viability of these models [120].

Observational Constraints

Brane-world models are constrained by both cosmological observations and GW observations. The growth of structure in brane-world models is constrained by the cosmic microwave background (CMB) and large-scale structure (LSS) data [35]. Additionally, the propagation speed of GWs in brane-world models is constrained by the observation of GW170817, which requires $|v_{\text{GW}} - c|/c \lesssim 10^{-15}$ [17].

6 Conclusions

The study of $f(R)$ gravity and other modified theories of gravity has emerged as a vibrant field of research, driven by the need to address the limitations of GR and explain observational phenomena such as cosmic acceleration. In this review, we have explored the state of the art in $f(R)$ gravity, the constraints imposed by GW measurements and other experiments, the methodologies used to derive these constraints, and the prospects for future improvements. We have also compared $f(R)$ gravity with other prominent modified theories of gravity, highlighting their similarities and differences. Below, we summarize the key findings and insights from each of these areas.

$f(R)$ gravity has established itself as one of the most widely studied alternatives to GR, offering a natural framework for explaining cosmic acceleration without invoking dark energy. The theoretical foundations of $f(R)$ gravity are well understood, with a rich literature exploring its cosmological and astrophysical implications. Current research focuses on developing viable functional forms of $f(R)$ that are consistent with observational data, as well as exploring the implications of these models for structure formation, GW propagation, and the early Universe. Leading research groups worldwide, including those in Europe, the United States, and Asia, are actively investigating these aspects, often in collaboration with observational cosmologists and gravitational wave astronomers.

The constraints on $f(R)$ gravity from GW measurements and other experiments are stringent and multifaceted. Observations of BBH and BNS mergers by LIGO and Virgo have placed tight limits on the propagation speed, polarization modes, and dispersion relations of GWs in $f(R)$ gravity. For example, the observation of GW170817 constrained the deviation of the GW speed from the speed of light to $|v_{\text{GW}} - c|/c \lesssim 10^{-15}$, ruling out many $f(R)$ models that predict significant deviations. Additionally, the absence of scalar polarization modes in GW signals has further constrained the scalar degree of freedom in $f(R)$ gravity.

Beyond GW measurements, $f(R)$ gravity is also constrained by solar system tests, cosmological surveys, and laboratory experiments. Solar system tests, such as those based on the parameterized post-Newtonian (PPN) formalism, constrain the PPN parameter γ to $|\gamma - 1| \lesssim 2 \times 10^{-5}$. Cosmological surveys, such as those measuring the cosmic microwave background (CMB) and large-scale structure (LSS), provide additional constraints on the growth of structure and the expansion history of the Universe. Laboratory experiments, such as those testing the equivalence principle, further constrain the coupling between the scalar degree of freedom in $f(R)$ gravity and matter.

The constraints on $f(R)$ gravity are derived using a combination of theoretical modeling and observational data. Theoretical models are used to predict the behavior of GWs, the growth of structure, and the expansion history of the Universe in $f(R)$ gravity. These predictions are then compared with observational data to derive constraints on the parameters of $f(R)$ gravity. For example, the propagation speed of GWs is constrained by comparing the observed time delay between GW and electromagnetic signals with the predictions of $f(R)$ gravity. Similarly, the growth of structure is constrained by comparing the observed power spectrum of the CMB and LSS with the predictions of $f(R)$ gravity.

The derivation of these constraints often relies on simplifying assumptions, such as the absence of additional fields or interactions beyond those included in the $f(R)$ Lagrangian. These assumptions are necessary to make the theoretical predictions tractable, but they may limit the generality of the constraints. Future work should aim to relax these assumptions and explore more general models of $f(R)$ gravity.

The future of testing $f(R)$ gravity is bright, with significant improvements expected in the coming decades. Next-generation GW detectors, such as the Einstein Telescope (ET) and Cosmic Explorer (CE), will provide unprecedented sensitivity to GW signals, enabling more precise measurements of the propagation speed, polarization modes, and dispersion relations of GWs. These detectors will also improve the sensitivity to the stochastic gravitational wave background (SGWB), which can provide additional constraints on $f(R)$ gravity.

In addition to GW measurements, future cosmological surveys, such as those conducted by the Euclid satellite and the Vera Rubin Observatory, will provide more precise measurements of the growth of structure and the expansion history of the Universe. These measurements will further constrain the parameters of $f(R)$ gravity and test its predictions on cosmological scales.

Theoretical developments will also play a crucial role in improving the constraints on $f(R)$ gravity. Advances in numerical relativity and perturbation theory will enable the development of more accurate waveform models for $f(R)$ gravity, which are essential for extracting information from GW signals. Similarly, improved models for the formation and evolution of astrophysical sources, such as BBH and BNS mergers, will enable more precise tests of $f(R)$ gravity.

While $f(R)$ gravity is a well-studied alternative to GR, other modified theories of gravity, such as scalar-tensor theories, Horndeski gravity, massive gravity, brane-world, and other models,

offer different approaches to addressing the limitations of GR. These theories are also tightly constrained by solar system tests, GW observations, and cosmological data. However, they differ in their predictions for cosmological scales and the early Universe, providing complementary insights into the nature of gravity.

For example, scalar-tensor theories introduce an additional scalar field that mediates an extra force, leading to deviations from GR in both the weak-field and strong-field regimes. Horndeski gravity, as the most general scalar-tensor theory, encompasses a wide range of modified gravity models and is constrained by both solar system tests and GW observations. Massive gravity, which posits a non-zero mass for the graviton, is constrained by the observation of GWs from BBH mergers, which place an upper limit on the graviton mass of $m_g \lesssim 10^{-23}$ eV. Brane-world models, which are based on the idea that our four-dimensional universe is a brane embedded in a higher-dimensional space, are constrained by cosmological observations and GW measurements.

In conclusion, the study of $f(R)$ gravity and other modified theories of gravity is a vibrant and rapidly advancing field, offering profound insights into the nature of gravity. While current observational constraints are already stringent, future breakthroughs in technology and theory will further test these models. Next-generation gravitational wave detectors, cosmological surveys, and laboratory experiments will enable us to probe gravity across all scales, from the solar system to the early Universe. These efforts bring us closer to answering fundamental questions about dark energy, cosmic acceleration, and the ultimate fate of the Universe, reshaping our understanding of gravity and cosmology.

Acknowledgements

I would like to express my thanks for the advice and help I received from Andrew L. Miller, and for the very useful conversations with Şengül Kuru, Banu Şahin, Giovanni Montani, and Anne-Christine Davis. I would also like to acknowledge the support from the COST Action COSMIC WISPerS CA21106, supported by COST (European Cooperation in Science and Technology), and for their encouragement for young researchers.

Data availability

No datasets were generated or analysed during the current study.

References

- [1] S. Weinberg, *The cosmological constant problem*, *Reviews of Modern Physics* **61** (1989) 1.
- [2] S. Weinberg, *The cosmological constant problems (talk given at dark matter 2000, february, 2000)*, 2000.
- [3] S. Shankaranarayanan and J.P. Johnson, *Modified theories of gravity: Why, how and what?*, *General Relativity and Gravitation* **54** (2022) .
- [4] C.M. Will, *The confrontation between general relativity and experiment*, *Living Reviews in Relativity* **17** (2014) .
- [5] T. Clifton, P.G. Ferreira, A. Padilla and C. Skordis, *Modified gravity and cosmology*, *Physics Reports* **513** (2012) 1–189.

- [6] A.G. Riess et al., *Observational evidence from supernovae for an accelerating universe and a cosmological constant*, *The Astronomical Journal* **116** (1998) 1009.
- [7] S. Perlmutter et al., *Measurements of ω and λ from 42 high-redshift supernovae*, *The Astrophysical Journal* **517** (1999) 565.
- [8] E.T. Chickles et al., *A gravitational wave detectable candidate Type Ia supernova progenitor*, **2411.19916**.
- [9] G. Bertone, D. Hooper and J. Silk, *Particle dark matter: Evidence, candidates and constraints*, *Physics Reports* **405** (2005) 279.
- [10] T.P. Sotiriou and V. Faraoni, *$f(r)$ theories of gravity*, *Rev. Mod. Phys.* **82** (2010) 451.
- [11] C.Q. Geng, *Gravitational waves in viable modified gravity theories*, *J. Phys. Conf. Ser.* **384** (2012) 012030.
- [12] S.A. Appleby and R.A. Battye, *Aspects of cosmological expansion in $f(r)$ gravity models*, *Journal of Cosmology and Astroparticle Physics* **2007** (2007) 019.
- [13] S. Nojiri and S.D. Odintsov, *Unified cosmic history in modified gravity: From $f(r)$ theory to lorentz non-invariant models*, *Physics Reports* **505** (2011) 59.
- [14] S. Tsujikawa, *Observational signatures of $f(r)$ dark energy models*, *Physical Review D* **77** (2008) 023507.
- [15] B. Abbott et al., *Observation of gravitational waves from a binary black hole merger*, *Physical Review Letters* **116** (2016) .
- [16] M. Isi, M. Pitkin and A.J. Weinstein, *Testing general relativity with gravitational-wave polarization*, *Physical Review D* **96** (2017) 122004.
- [17] B.P. Abbott et al., *Gravitational waves and gamma-rays from a binary neutron star merger: Gw170817 and grb 170817a*, *The Astrophysical Journal Letters* **848** (2017) L13.
- [18] D. Wands, *Extended gravity theories and the einstein–hilbert action*, *Classical and Quantum Gravity* **11** (1994) 269–279.
- [19] S. Capozziello and M. De Laurentis, *Extended theories of gravity*, *Physics Reports* **509** (2011) 167–321.
- [20] L. Amendola and S. Tsujikawa, *Dark Energy: Theory and Observations*, Cambridge University Press (1, 2015).
- [21] A.A. Starobinsky, *A New Type of Isotropic Cosmological Models Without Singularity*, *Phys. Lett. B* **91** (1980) 99.
- [22] G. Cognola, E. Elizalde, S. Nojiri, S.D. Odintsov, L. Sebastiani and S. Zerbini, *Class of viable modified $f(r)$ gravities describing inflation and the onset of accelerated expansion*, *Phys. Rev. D* **77** (2008) 046009.
- [23] A.H. Guth, *The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems*, *Phys. Rev. D* **23** (1981) 347.
- [24] S. Nojiri, S. Odintsov and V. Oikonomou, *Modified gravity theories on a nutshell: Inflation, bounce and late-time evolution*, *Physics Reports* **692** (2017) 1.
- [25] S.M. Carroll, V. Duvvuri, M. Trodden and M.S. Turner, *Is cosmic speed-up due to new gravitational physics?*, *Physical Review D* **70** (2004) .
- [26] S. Nojiri and S.D. Odintsov, *Modified gravity with negative and positive powers of curvature: Unification of inflation and cosmic acceleration*, *Phys. Rev. D* **68** (2003) 123512.

- [27] S.R. Chowdhury and M. Khlopov, *Gravitational waves in the modified gravity*, 2021.
- [28] K.N. Ananda, S. Carloni and P.K.S. Dunsby, *The Evolution of cosmological gravitational waves in $f(R)$ gravity*, *Phys. Rev. D* **77** (2008) 024033 [0708.2258].
- [29] T. Katsuragawa, T. Nakamura, T. Ikeda and S. Capozziello, *Gravitational waves in $f(r)$ gravity: Scalar waves and the chameleon mechanism*, *Phys. Rev. D* **99** (2019) 124050.
- [30] J.-Z. Zhou, Y.-T. Kuang, D. Wu, F.-Y. Chen, H. Lü and Z. Chang, *Scalar induced gravitational waves in $f(r)$ gravity*, 2024.
- [31] Y. Gong and T. Zhu, *Gravitational wave polarization in $f(r)$ gravity*, *Physical Review D* **98** (2018) 024040.
- [32] I.D. Saltas, I. Sawicki and I.L. Shapiro, *Gravitational waves in modified gravity*, *Journal of Cosmology and Astroparticle Physics* **2014** (2014) 048.
- [33] A. Nishizawa, *Generalized framework for testing gravity with gravitational-wave propagation. II. constraints on horndeski theory*, *Physical Review D* **97** (2018) 104037.
- [34] R.S. Park, W.M. Folkner, A.S. Konopliv, J.G. Williams, D.E. Smith and M.T. Zuber, *Precession of mercury's perihelion from ranging to the messenger spacecraft*, *The Astronomical Journal* **153** (2017) 121.
- [35] N. Aghanim *et al.* [Planck], *Astron. Astrophys.* **641**, A6 (2020) [erratum: *Astron. Astrophys.* **652**, C4 (2021)] doi:10.1051/0004-6361/201833910 [arXiv:1807.06209 [astro-ph.CO]].
- [36] K. Koyama, *Cosmological tests of modified gravity*, *Reports on Progress in Physics* **79** (2016) 046902.
- [37] M. Milgrom, *A Modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis*, *Astrophys. J.* **270** (1983) 365.
- [38] R.H. Sanders and S.S. McGaugh, *Modified newtonian dynamics as an alternative to dark matter*, *Annual Review of Astronomy and Astrophysics* **40** (2002) 263–317.
- [39] J.D. Bekenstein, *Relativistic gravitation theory for the MOND paradigm*, *Phys. Rev. D* **70** (2004) 083509 [astro-ph/0403694].
- [40] E.P. Verlinde, *Emergent gravity and the dark universe*, *SciPost Physics* **2** (2017) .
- [41] M.H. Chan and C.M. Lee, *The solar system test for the general modified gravity theories*, *Monthly Notices of the Royal Astronomical Society* **518** (2022) 6238–6242.
- [42] T. Chiba, *$1/r$ gravity and scalar-tensor gravity*, *Physics Letters B* **575** (2003) 1–3.
- [43] A.L. Erickcek, *The Consequences of Modifying Fundamental Cosmological Theories*, Ph.D. thesis, Caltech, 2009. 10.7907/ST6B-1S91.
- [44] P. Brax, C. van de Bruck, A.-C. Davis and D.J. Shaw, *$f(r)$ gravity and chameleon theories*, *Phys. Rev. D* **78** (2008) 104021.
- [45] P. Brax, A.-C. Davis and R. Jha, *Neutron stars in screened modified gravity: Chameleon versus dilaton*, *Physical Review D* **95** (2017) .
- [46] T. O'Shea, A.-C. Davis, M. Giannotti, S. Vagnozzi, L. Visinelli and J.K. Vogel, *Solar chameleons: Novel channels*, *Phys. Rev. D* **110** (2024) 063027 [2406.01691].
- [47] J. Khoury and A. Weltman, *Chameleon cosmology*, *Physical Review D* **69** (2004) .
- [48] I.I. Shapiro, *Fourth test of general relativity*, *Phys. Rev. Lett.* **13** (1964) 789.
- [49] B. Bertotti, L. Iess and P. Tortora, *A test of general relativity using radio links with the cassini spacecraft*, *Nature* **425** (2003) 374.

- [50] S. Schlamminger, K.-Y. Choi, T.A. Wagner, J.H. Gundlach and E.G. Adelberger, *Test of the equivalence principle using a rotating torsion balance*, *Physical Review Letters* **100** (2008) .
- [51] P. Brax, P. Valageas and P. Vanhove, *New bounds on dark energy induced fifth forces*, *Physical Review D* **99** (2019) .
- [52] D. Liang, Y. Gong, S. Hou and Y. Liu, *Polarizations of gravitational waves in $f(r)$ gravity*, *Phys. Rev. D* **95** (2017) 104034.
- [53] Y.-Q. Dong, X.-B. Lai, Y.-Q. Liu and Y.-X. Liu, *A unified framework for analyzing gravitational wave effects in modified gravity theories based on the isaacson picture*, 2025.
- [54] A.D. Felice and T. Suyama, *Vacuum structure for scalar cosmological perturbations in modified gravity models*, *Journal of Cosmology and Astroparticle Physics* **2009** (2009) 034.
- [55] A. De Felice and S. Tsujikawa, *$f(r)$ theories*, *Living Reviews in Relativity* **13** (2010) .
- [56] S. Mirshekari and C.M. Will, *Compact binary systems in scalar-tensor gravity: Equations of motion to 2.5 post-newtonian order*, *Phys. Rev. D* **87** (2013) 084070.
- [57] N. Yunes, K. Yagi and F. Pretorius, *Theoretical physics implications of the binary black-hole mergers gw150914 and gw151226*, *Phys. Rev. D* **94** (2016) 084002.
- [58] V.A. Kostelecký and M. Mewes, *Testing local lorentz invariance with gravitational waves*, *Physics Letters B* **757** (2016) 510.
- [59] LIGO SCIENTIFIC AND VIRGO COLLABORATION collaboration, *Gw170104: Observation of a 50-solar-mass binary black hole coalescence at redshift 0.2*, *Phys. Rev. Lett.* **118** (2017) 221101.
- [60] A. Nishizawa, *Generalized framework for testing gravity with gravitational-wave propagation. i. formulation*, *Phys. Rev. D* **97** (2018) 104037.
- [61] S. Mirshekari and C.M. Will, *Compact binary systems in scalar-tensor gravity: Equations of motion to 2.5 post-newtonian order*, *Physical Review D* **87** (2013) 084070.
- [62] B.P. Abbott et al., *Upper limits on the isotropic gravitational-wave background from advanced ligo and advanced virgo's third observing run*, *Physical Review D* **104** (2021) 022004.
- [63] W. Hu and I. Sawicki, *Models of $f(r)$ cosmic acceleration that evade solar system tests*, *Phys. Rev. D* **76** (2007) 064004.
- [64] T. Baker, E. Bellini, P. Ferreira, M. Lagos, J. Noller and I. Sawicki, *Strong constraints on cosmological gravity from gw170817 and grb 170817a*, *Physical Review Letters* **119** (2017) .
- [65] L. Amendola et al., *Cosmology and fundamental physics with the euclid satellite*, *Living Reviews in Relativity* **21** (2018) .
- [66] E.G. Adelberger et al., *Torsion balance experiments: A low-energy frontier of particle physics*, *Progress in Particle and Nuclear Physics* **62** (2009) 102.
- [67] B. Sathyaprakash et al., *Scientific potential of einstein telescope*, 2012.
- [68] M. Maggiore et al., *Science case for the einstein telescope*, *Journal of Cosmology and Astroparticle Physics* **2020** (2020) 050.
- [69] D. Reitze et al., *Cosmic explorer: The u.s. contribution to gravitational-wave astronomy beyond ligo*, 2019.
- [70] S. Dwyer et al., *Gravitational wave detector with cosmological reach*, *Phys. Rev. D* **91** (2015) 082001.
- [71] M. Maggiore, *Gravitational Waves. Vol. 1: Theory and Experiments*, Oxford University Press (2007), 10.1093/acprof:oso/9780198570745.001.0001.

- [72] S. Hild et al., *Sensitivity studies for third-generation gravitational wave observatories*, *Classical and Quantum Gravity* **28** (2011) 094013.
- [73] B.P. Abbott et al., *Exploring the sensitivity of next generation gravitational wave detectors*, *Classical and Quantum Gravity* **34** (2017) 044001.
- [74] B.S. Sathyaprakash et al., *Scientific objectives of einstein telescope*, *Classical and Quantum Gravity* **29** (2012) 124013.
- [75] S. Aziz, S.K. Jha and A. Rahaman, *The inflationary scenario in the $f(r)$ gravity model with a r^4 term*, *Classical and Quantum Gravity* **38** (2021) 225008.
- [76] C. Dioguardi, A. Racioppi and E. Tomberg, *Slow-roll inflation in palatini $f(r)$ gravity*, *Journal of High Energy Physics* **2022** (2022) .
- [77] D. Saha, J. P. Saha and A. K. Sanyal, *Int. J. Geom. Meth. Mod. Phys.* **20**, no.12, 2350213 (2023) doi:10.1142/S0219887823502134 [arXiv:2308.11770 [gr-qc]].
- [78] C. Dioguardi, A. Racioppi and E. Tomberg, *Beyond (and back to) palatini quadratic gravity and inflation*, *Journal of Cosmology and Astroparticle Physics* **2024** (2024) 041.
- [79] C. Dioguardi and A. Racioppi, *Palatini $f(r,x)$: A new framework for inflationary attractors*, *Physics of the Dark Universe* **45** (2024) 101509.
- [80] F. Milani, *Modified $f(r, g, t)$ gravity in the quintom model, and inflation*, 2024.
- [81] N. Bostan and R.H. Dejarah, *Minimally coupled β -exponential inflation with an r^2 term in the palatini formulation*, 2024.
- [82] J.D. Romano and N.J. Cornish, *Detection methods for stochastic gravitational-wave backgrounds: A unified treatment*, *Living Reviews in Relativity* **20** (2017) 2.
- [83] K. Kotera and A.V. Olinto, *The astrophysics of ultrahigh-energy cosmic rays*, *Annual Review of Astronomy and Astrophysics* **49** (2011) 119.
- [84] A.M. et al., *Supernova neutrinos: Production, oscillations, and detection*, *La Rivista del Nuovo Cimento* **39** (2016) 1.
- [85] A.A. et al., *The pierre auger cosmic ray observatory*, *Nuclear Instruments and Methods in Physics Research Section A* **798** (2015) 172.
- [86] M.A. et al., *Ultra-high-energy cosmic rays and modified gravity*, *Journal of Cosmology and Astroparticle Physics* **2019** (2019) 012.
- [87] F. Halzen and S.R. Klein, *Icecube: An instrument for neutrino astronomy*, *Reviews of Modern Physics* **82** (2010) 2593.
- [88] D. Collaboration et al., *Long-baseline neutrino facility (lbnf) and deep underground neutrino experiment (dune) conceptual design report volume 2: The physics program for dune at lbnf*, 2016.
- [89] H.-K. Proto-Collaboration et al., *Hyper-kamiokande design report*, 2018.
- [90] N. Yunes and X. Siemens, *Gravitational-wave tests of general relativity with ground-based detectors and pulsar timing arrays*, *Living Reviews in Relativity* **16** (2013) 9.
- [91] I. Mandel and S.E. de Mink, *Merging binary black holes formed through chemically homogeneous evolution in short-period stellar binaries*, *Monthly Notices of the Royal Astronomical Society* **458** (2016) 2634.
- [92] N.J. Cornish and J.D. Romano, *Towards a unified treatment of gravitational-wave backgrounds*, *Physical Review D* **83** (2011) 042001.

- [93] Y. Fujii and K. Maeda, *The scalar-tensor theory of gravitation*, Cambridge Monographs on Mathematical Physics, Cambridge University Press (7, 2007), [10.1017/CBO9780511535093](https://doi.org/10.1017/CBO9780511535093).
- [94] E. Berti et al., *Testing general relativity with present and future astrophysical observations*, *Classical and Quantum Gravity* **32** (2015) 243001.
- [95] C.M. Will, *Testing scalar-tensor gravity with gravitational-wave observations of inspiralling compact binaries*, *Physical Review D* **50** (1994) 6058–6067.
- [96] G.W. Horndeski, *Second-order scalar-tensor field equations in a four-dimensional space*, *Int. J. Theor. Phys.* **10** (1974) 363.
- [97] T. Kobayashi, M. Yamaguchi and J. Yokoyama, *Generalized g -inflation: $-$ inflation with the most general second-order field equations*, *Progress of Theoretical Physics* **126** (2011) 511–529.
- [98] R. Woodard, *Avoiding dark energy with $1/r$ modifications of gravity*, in *The Invisible Universe: Dark Matter and Dark Energy*, p. 403–433, Springer Berlin Heidelberg (2007), DOI.
- [99] E. Bellini and I. Sawicki, *Maximal freedom at minimum cost: linear large-scale structure in general modifications of gravity*, *Journal of Cosmology and Astroparticle Physics* **2014** (2014) 050–050.
- [100] P. Creminelli and F. Vernizzi, *Dark energy after gw170817 and grb170817a*, *Physical Review Letters* **119** (2017) .
- [101] C. de Rham and G. Gabadadze, *Generalization of the fierz-pauli action*, *Physical Review D* **82** (2010) .
- [102] K. Hinterbichler, *Theoretical aspects of massive gravity*, *Reviews of Modern Physics* **84** (2012) 671–710.
- [103] C. de Rham, G. Gabadadze and A.J. Tolley, *Resummation of massive gravity*, *Physical Review Letters* **106** (2011) .
- [104] C.M. Will, *Bounding the mass of the graviton using gravitational-wave observations of inspiralling compact binaries*, *Phys. Rev. D* **57** (1998) 2061.
- [105] J.M. Ezquiaga and M. Zumalacárregui, *Dark energy after gw170817: Dead ends and the road ahead*, *Phys. Rev. Lett.* **119** (2017) 251304.
- [106] T.C.K. Ng, M. Isi, K.W.K. Wong and W.M. Farr, *Constraining gravitational wave amplitude birefringence with GWTC-3*, *Phys. Rev. D* **108** (2023) 084068 [2305.05844].
- [107] A.P.S. S., S.H. Pereira and E.L. Mendonca, *New cosmological solutions in massive gravity theory*, 2017.
- [108] A.E. Gumrukcuoglu, C. Lin and S. Mukohyama, *Open frw universes and self-acceleration from nonlinear massive gravity*, *Journal of Cosmology and Astroparticle Physics* **2011** (2011) 030.
- [109] M. Fasiello and A.J. Tolley, *Cosmological stability bound in massive gravity and bigravity*, *Journal of Cosmology and Astroparticle Physics* **2013** (2013) 002–002.
- [110] P. Wu and H.W. Yu, *Generalized Chaplygin gas model: Constraints from Hubble parameter versus redshift data*, *Phys. Lett. B* **644** (2007) 16 [gr-qc/0612055].
- [111] B.P. Abbott et al., *Tests of general relativity with gw170817*, *Physical Review Letters* **123** (2019) 011102.
- [112] L. Randall and R. Sundrum, *Large mass hierarchy from a small extra dimension*, *Phys. Rev. Lett.* **83** (1999) 3370.
- [113] R. Maartens and K. Koyama, *Brane-world gravity*, *Living Reviews in Relativity* **13** (2010) .

- [114] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *The hierarchy problem and new dimensions at a millimeter*, *Physics Letters B* **429** (1998) 263.
- [115] T. Shiromizu, K.-i. Maeda and M. Sasaki, *The einstein equations on the 3-brane world*, *Physical Review D* **62** (2000) 024012.
- [116] N. Kaloper, *Bent domain walls as braneworlds*, *Physical Review D* **60** (1999) .
- [117] J. Garriga and M. Sasaki, *Brane-world creation and black holes*, *Physical Review D* **62** (2000) .
- [118] D. Langlois, *Cosmological evolution in a brane universe*, in *9th Marcel Grossmann Meeting on Recent Developments in Theoretical and Experimental General Relativity, Gravitation and Relativistic Field Theories (MG 9)*, pp. 2065–2066, 2, 2001, DOI [gr-qc/0102007].
- [119] V. Sahni, *The cosmological constant problem and quintessence*, *Classical and Quantum Gravity* **19** (2002) 3435–3448.
- [120] D. Langlois, *Brane cosmology*, *Progress of Theoretical Physics Supplement* **148** (2002) 181–212.