

LOCAL MINIMIZERS IN 3D OF VECTOR ALLEN-CAHN WITH A QUADRUPLE JUNCTION.

ABHISHEK ADIMURTHI AND PETER STERNBERG

ABSTRACT. For Ω a perturbation of the unit ball in \mathbb{R}^3 , we establish the existence of a sequence of local minimizers for the vector Allen-Cahn energy. The sequence converges in L^1 to a partition of Ω whose skeleton is given by a tetrahedral cone and thus contains a quadruple point. This is accomplished by proving that the partition is an isolated local minimizer of the associated Γ -limit of the sequence of Allen-Cahn functionals.

1. INTRODUCTION.

For a domain $\Omega \subset \mathbb{R}^3$ described through a slight deformation of a ball, we establish the existence of a sequence of local minimizers $u_\varepsilon : \Omega \rightarrow \mathbb{R}^3$ to a vector Allen-Cahn (or Modica-Mortola) type energy that converges as $\varepsilon \rightarrow 0$ to a partition of Ω exhibiting a quadruple junction. Invoking the well-known machinery of Γ -convergence for Modica-Mortola to be found in [Bal90, KS89], our approach consists of proving that the partition of Ω into four subdomains via a tetrahedral cone represents an isolated local minimizer of perimeter in the L^1 topology.

To be more specific, we define the sequence of energies

$$E_\varepsilon(u) := \int_\Omega \frac{\varepsilon}{2} |\nabla u|^2 + \frac{1}{\varepsilon} W(u) dx, \quad (1.1)$$

where $u : \Omega \rightarrow \mathbb{R}^3$ and $W : \mathbb{R}^3 \mapsto \mathbb{R}$ is a non-negative function that vanishes on a set of four points $P := \{p_1, p_2, p_3, p_4\} \subset \mathbb{R}^3$. For say $C^{1,\alpha}$ potentials W , a local minimizer represents a solution to the problem

$$\varepsilon^2 \Delta u = W_u(u) \text{ in } \Omega, \quad \nabla u \cdot \nu_{\partial\Omega} = 0 \text{ on } \partial\Omega, \quad (1.2)$$

where $\nu_{\partial\Omega}$ denotes the outer unit normal to $\partial\Omega$. In the analogous situation in one lower dimension where $u : \Omega \rightarrow \mathbb{R}^2$ with $\Omega \subset \mathbb{R}^2$ taken as a certain deformation of a disc, and with W given by a triple-well potential, this program is carried out in [SZ94] to produce solutions to (1.2) possessing triple junction structure.

We note that entire solutions $u : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ to this PDE exhibiting quadruple junction structure have been constructed under symmetry assumptions on the potential W in [GS08, AF11, Ali12]. See

[AFS18] for a definitive exposition on entire solutions for symmetric potentials in all dimensions. One motivation for the present article, in work presently being pursued, is to construct an entire solution whose ‘blow-downs’ converge to a quadruple junction partition of \mathbb{R}^3 by first blowing up the local minimizers constructed here on a bounded domain. Such a strategy for non-symmetric triple-well potentials in two dimensions has been carried out in [AG24, AG23, Gen25, SS24] to produce entire solutions with triple junction structure. If successful in three dimensions, the advantage would be two-fold: it would require no symmetry assumption on W and it would lead to a locally minimizing entire solution in the sense of De Giorgi, that is, an entire solution that minimizes the energy on compact sets when compared with competitors sharing its boundary values. The previously constructed entire solutions mentioned above do not come with any guarantee of stability.

This entire program represents an attempt at ‘desingularization’ of all possible minimal structures. From this perspective, the most basic case is a plane whose diffuse counterpart for $\Delta u = W_u(u)$ is a heteroclinic connection, that is, an ODE joining two wells. Next comes the aforementioned triple junction and here we contemplate the other possible limiting cone, the quadruple junction in \mathbb{R}^3 , as per Taylor’s seminal classification in [Tay76].

Before describing the partitioning problem in more detail we recall the relevant work on Γ -convergence. Generalizing the approach in [FT89, Ste88, Ste91] on Γ -convergence for vector Modica-Mortola with double well-type potentials, under some mild assumptions on a multi-well potential W , Baldo establishes in [Bal90] the Γ -convergence of E_ε in the L^1 topology to a functional E_0 given in our setting by

$$E_0(v) := \sum_{1 \leq i < j \leq 4} c_{ij} \mathcal{H}^2(\partial^* R_i \cap \partial^* R_j \cap \Omega) \quad (1.3)$$

for $v \in BV(\mathbb{R}^3; P)$. Here the region R_i is defined as

$$R_i := \{x \in \Omega \mid v(x) = p_i\},$$

so that $\mathcal{R} := (R_1, R_2, R_3, R_4)$ represents a partition of Ω by four sets of finite perimeter. In what follows, we will frequently write $E_0(\mathcal{R})$ rather than $E_0(v)$. Here \mathcal{H}^2 is two-dimensional Hausdorff measure and $\partial^* A$ refers to the reduced boundary of the set A . The coefficients c_{ij} are given by

$$c_{ij} := d(p_i, p_j), \quad (1.4)$$

with $d(p, q)$ denoting the metric defined through

$$d(p, q) := \inf \left\{ \sqrt{2} \int_0^1 \sqrt{W(\gamma(t))} |\gamma'(t)| dt \left| \begin{array}{l} \gamma \in C^1([0, 1]; \mathbb{R}^3), \\ \gamma(0) = p, \\ \gamma(1) = q, \end{array} \right. \right\}$$

for any $p, q \in \mathbb{R}^3$.

To describe the main result of this paper, Theorem 3.1, we consider a tetrahedral cone centered at the origin. The *regular* tetrahedral cone, which corresponds to the case where all c_{ij} 's in (1.4) are equal, is defined as the cone over a regular tetrahedron centered at the origin. Such a cone, which we denote by \mathcal{C} , partitions \mathbb{R}^3 into four regions, say $\tilde{S}_1, \dots, \tilde{S}_4$, each having boundary consisting of three planes. We will denote the (constant) unit normal to the plane separating region \tilde{S}_i from \tilde{S}_j , pointing from \tilde{S}_i to \tilde{S}_j , by n_{ij} .

The only point in common to the boundary of all four regions is the origin, but given any triple of distinct indices $\{i, j, k\} \subset \{1, 2, 3, 4\}$, one has that $\partial\tilde{S}_i \cap \partial\tilde{S}_j \cap \partial\tilde{S}_k$ consists of a ray emanating from the origin. Stationarity of the tetrahedral cone with respect to surface area implies a ‘balance law’ for the normals to the planes meeting at such a ray:

$$n_{ij} + n_{jk} + n_{ki} = 0 \quad \text{for every distinct triple } i, j, k \in \{1, 2, 3, 4\}. \quad (1.5)$$

More generally, one can consider partitions arising as critical points of E_0 with *unequal* weights c_{ij} and the partitions of interest here are induced by non-symmetric tetrahedral cones centered at the origin. Their description is similar to that of the regular tetrahedral cone, but now stationarity for E_0 leads to the requirement that the angles at which three planes meet along the triple-junction like rays are determined through the conditions

$$c_{ij}n_{ij} + c_{jk}n_{jk} + c_{ki}n_{ki} = 0 \quad \text{for every distinct triple } i, j, k \in \{1, 2, 3, 4\}. \quad (1.6)$$

Following [Sch35, Theorem 1] and [LM94], such a non-symmetric cone can be constructed provided the matrix

$$M := \begin{bmatrix} 2c_{12}^2 & c_{12}^2 + c_{13}^2 - c_{23}^2 & c_{12}^2 + c_{14}^2 - c_{24}^2 \\ c_{12}^2 + c_{13}^2 - c_{23}^2 & 2c_{13}^2 & c_{13}^2 + c_{14}^2 - c_{34}^2 \\ c_{12}^2 + c_{13}^2 - c_{23}^2 & c_{13}^2 + c_{14}^2 - c_{34}^2 & 2c_{14}^2 \end{bmatrix} \quad (1.7)$$

is positive definite. Under this assumption, which we make for the remainder of the paper, there exists a tetrahedron with vertices A_1, A_2, A_3 and A_4 lying in \mathbb{R}^3 with the property that the length of the edge joining

A_i and A_j is c_{ij} . Then for all distinct pairs $i, j \in \{1, 2, 3, 4\}$, we define a unit normal vector to the plane separating phase i from phase j via

$$n_{ij} := \frac{1}{c_{ij}}(A_j - A_i).$$

The corresponding six planes with these normals passing through the origin lead to the non-symmetric version of the cone \mathcal{C} satisfying (1.6). We note that positivity of M , in particular, implies strict triangle inequalities between any three of the c_{ij} 's.

In Section 2.3 we will describe a domain $\Omega \subset \mathbb{R}^3$ arising as a perturbation of a ball such that the partition $\mathcal{S} := \{S_1, S_2, S_3, S_4\}$ defined through $S_i := \tilde{S}_i \cap \Omega$ results in an isolated L^1 local minimizer of E_0 in the sense that for some $\delta = \delta(\Omega) > 0$, the function

$$u_0(x) := \sum_{i=1}^4 p_i \chi_{S_i}(x) \tag{1.8}$$

satisfies the condition

$$E_0(u_0) < E_0(v) \quad \text{provided } v \in BV(\Omega; P) \text{ and } 0 < \|v - u_0\|_{L^1(\Omega)} < \delta.$$

This is the content of Theorem 3.1. As we shall see, the crucial property of this deformation of the ball is that in a neighborhood of $\mathcal{C} \cap \partial\Omega$, the vectors n_{ij} satisfy conditions (2.4).

In light of Theorem 4.1 of [KS89], Theorem 3.1 immediately implies:

Theorem 1.1. *There exists a domain $\Omega \subset \mathbb{R}^3$ such that for all ε sufficiently small, the energy E_ε possesses an L^1 local minimizer u_ε . As $\varepsilon \rightarrow 0$, one has that $u_\varepsilon \rightarrow u_0$ in $L^1(\Omega)$ with u_0 given by (1.8).*

The proof of Theorem 3.1 follows in two steps. In the first, phrased separately as Theorem 3.2, we argue that one need only consider partitions that meet $\partial\Omega$ uniformly close to where the partition \mathcal{S} meets $\partial\Omega$. This is accomplished by proving a boundary version of Maggi's infiltration lemma [Mag12, Lemma 30.2]. The second step then consists of a calibration argument in the spirit of [LM94]. We begin in Section 2.1 with some general comments and notation regarding sets of finite perimeter. Section 2.2 contains a few tools we will invoke, and in Section 2.3 we describe the deformation of the ball required for the proof of Theorem 3.1.

2. PRELIMINARIES.

2.1. Partitions with sets of finite perimeter. We recall that a set $E \subset \mathbb{R}^n$ has finite perimeter in an open set $\Omega \subset \mathbb{R}^n$ if the characteristic function χ_E has bounded variation, and we denote the perimeter within

Ω by $\text{Per}(E; \Omega)$. We also recall that sets of finite perimeter possess a measure-theoretic exterior normal which is suitably general to ensure the validity of the Gauss-Green theorem. A unit vector ν^* is defined as the measure theoretic exterior normal to E at x provided

$$\lim_{r \rightarrow 0} \frac{\mathcal{L}^3(B(x, r) \cap \{y : (y - x) \cdot \nu^* < 0, y \notin E\})}{r^n} = 0$$

and

$$\lim_{r \rightarrow 0} \frac{\mathcal{L}^3(B(x, r) \cap \{y : (y - x) \cdot \nu^* > 0, y \in E\})}{r^n} = 0,$$

where $B(x, r)$ denotes the open ball of radius r centered at x and \mathcal{L}^3 denotes three-dimensional Lebesgue measure. We denote the set of such points x by ∂^*E , the reduced boundary of E .

By definition, sets of finite perimeter are determined only up to sets of measure zero. In order to avoid this ambiguity, whenever a set of finite perimeter, E , is considered, we shall always employ the measure-theoretic closure as the set to represent E . Thus, with this convention, we have

$$x \in E \text{ if and only if } \limsup_{r \rightarrow 0} \frac{\mathcal{L}^n(E \cap B(x, r))}{\mathcal{L}^n(B(x, r))} > 0.$$

We also note that

$$\text{Per}(E; \Omega) = \mathcal{H}^{n-1}(\Omega \cap \partial^*E).$$

Throughout, we will employ the concept of a partition of Ω by k sets of finite perimeter.

Definition 2.1. *We say $\mathcal{T} := (T_1, T_2, \dots, T_k)$ is a partition of a bounded domain $\Omega \subset \mathbb{R}^n$ if each $T_i \subset \Omega$ is a set of finite perimeter, $\mathcal{L}^n(\Omega \setminus \cup_{i=1}^k T_i) = 0$, and for each distinct pair $i, j \in \{1, 2, \dots, k\}$, $\mathcal{L}^n(T_i \cap T_j) = 0$.*

If $\mathcal{E} := (E_1, E_2, \dots, E_k)$ is a partition of Ω , then we define the perimeter of \mathcal{E} in Ω to be

$$\text{Per}(\mathcal{E}; \Omega) := \sum_{1 \leq i < j \leq k} \mathcal{H}^{n-1}(\partial^*E_i \cap \partial^*E_j \cap \Omega).$$

Further, if $\mathcal{R} := (R_1, \dots, R_k)$ is another partition of Ω , then we set

$$\mathcal{L}^n(\mathcal{R} \Delta \mathcal{E}) := \sum_{i=1}^k \mathcal{L}^n \left[(R_i \setminus E_i) \cup (E_i \setminus R_i) \right].$$

For more background on sets of finite perimeter we refer the reader to e.g. [EG15, Giu84, Mag12].

2.2. Two helpful tools. In the analysis to follow, we require the following two results.

Lemma 2.2. (cf. [SZ94, Lemma 2.2]) *Let $E \subset \mathbb{R}^n, n \geq 2$ be a set of finite perimeter. Suppose there is a point x_0 in $\partial^* E$ with the property that for some cube Q_0 centered at x_0 , the unit outward normal to E given by $\nu_E(x)$ is a constant ν_0 for all $x \in Q_0 \cap \partial^* E$. Then, $Q_0 \cap \partial^* E = Q_0 \cap \Pi$, where Π is the unique hyperplane in \mathbb{R}^n containing x_0 with normal ν_0 .*

Proposition 2.3 (Relative isoperimetric inequality). *Let $\Omega \subset \mathbb{R}^n, n \geq 2$ be a bounded and connected domain with C^1 boundary. Then, there exist constants $r_0(\Omega) > 0$ and $C_0 = C_0(n, \Omega) > 0$ such that for any $x_0 \in \partial\Omega$ and any $E \subset \Omega \cap B(x_0, r_0)$ of finite perimeter, one has:*

$$\text{Per}(E; \Omega) \geq C_0 \left[\mathcal{L}^n(E) \right]^{\frac{n-1}{n}}. \quad (2.1)$$

A version of Proposition 2.3 for the case when Ω is a ball in \mathbb{R}^n can be found e.g. in [EG15, Section 5.6]. For completeness, we provide a proof of Proposition 2.3 below, though the proof is essentially unchanged.

Proof. We fix any $r_0 > 0$ depending on Ω such that for all $x_0 \in \partial\Omega$ one has

$$\mathcal{L}^n(B(x_0, r_0) \cap \Omega) \leq \mathcal{L}^n(\Omega \setminus B(x_0, r_0)). \quad (2.2)$$

For a function $f : \mathbb{R}^n \mapsto \mathbb{R}$, we set

$$(f)_\Omega := \frac{1}{\mathcal{L}^n(\Omega)} \int_\Omega f(y) dy.$$

For any $x_0 \in \partial\Omega$ and any subset $E \subset B(x_0, r_0) \cap \Omega$, one has

$$\begin{aligned} & \int_\Omega |\chi_E(y) - (\chi_E)_\Omega|^{\frac{n}{n-1}} dy \\ &= \int_\Omega \left| \chi_E(y) - \frac{\mathcal{L}^n(E)}{\mathcal{L}^n(\Omega)} \right|^{\frac{n}{n-1}} dy \\ &= \left(\frac{\mathcal{L}^n(\Omega - E)}{\mathcal{L}^n(\Omega)} \right)^{\frac{n}{n-1}} \mathcal{L}^n(E) + \left(\frac{\mathcal{L}^n(E)}{\mathcal{L}^n(\Omega)} \right)^{\frac{n}{n-1}} \mathcal{L}^n(\Omega - E). \\ &\geq \left(\frac{1}{2} \right)^{\frac{n}{n-1}} \mathcal{L}^n(E), \end{aligned}$$

where the final inequality follows from (2.2). Hence,

$$\|\chi_E - (\chi_E)_\Omega\|_{L^{\frac{n}{n-1}}(\Omega)} > \frac{1}{2} \left[\mathcal{L}^n(E) \right]^{\frac{n-1}{n}}. \quad (2.3)$$

Now by the Poincaré inequality applied to any $f \in W^{1, \frac{n}{n-1}}(\Omega)$, we have

$$\|f - (f)_\Omega\|_{L^{\frac{n}{n-1}}(\Omega)} \leq C_1 \|Df\|_{L^{\frac{n}{n-1}}(\Omega)},$$

for some $C_1 = C_1(\Omega)$. Furthermore, we recall that for any $f \in BV(\Omega)$, there exist functions $\{f_k\}_{k=1}^\infty \subset BV(\Omega) \cap C^\infty(\Omega)$ such that

$$f_k \rightarrow f \text{ in } L^1(\Omega) \quad \text{and} \quad \|Df_k\|(\Omega) \rightarrow \|Df\|(\Omega) \text{ as } k \rightarrow \infty,$$

(see e.g. [Giu84, Chap. 1]). Approximating $f = \chi_E$ by smooth functions we may thus extend the Poincaré inequality to such an f with the right-hand side now denoting total variation (perimeter). Combining this inequality with (2.3), we complete the proof with $C_0 = \frac{1}{2C_1}$. \square

2.3. Description of the Domain. To describe the domain $\Omega \subset \mathbb{R}^3$ in which we will construct a sequence of local minimizers of the vector Allen-Cahn system, the key object is the infinite tetrahedral cone, namely the cone \mathcal{C} described in the introduction. We recall that this cone partitions \mathbb{R}^3 into four regions, $\tilde{S}_1, \dots, \tilde{S}_4$, each having boundary consisting of three planes with the normals satisfying the conditions (1.5) in the symmetric case or (1.6) in the non-symmetric case.

Let us now focus on the restriction of \mathcal{C} to the unit ball in \mathbb{R}^3 . For each distinct triple of indices, we denote by \tilde{q}_{ijk} the point where the ray on \mathcal{C} emanating from the origin lying on $\partial\tilde{S}_i \cap \partial\tilde{S}_j \cap \partial\tilde{S}_k$ meets \mathbb{S}^2 . The intersection of \mathcal{C} with \mathbb{S}^2 consists of four geodesic triangles, having three of the four \tilde{q}_{ijk} 's as vertices. Furthermore, there are a total of six circular arcs, namely the sides of these triangles, comprising $\mathcal{C} \cap \mathbb{S}^2$, and we denote these arcs by $\tilde{\gamma}_{ij}$, so that $\tilde{\gamma}_{ij}$ has endpoints \tilde{q}_{ijk} and \tilde{q}_{ijl} for distinct indices i, j, k and l with

$$\tilde{\gamma}_{ij} = \partial\tilde{S}_i \cap \partial\tilde{S}_j \cap \mathbb{S}^2.$$

The restriction of $\{\tilde{S}_1, \dots, \tilde{S}_4\}$ to the unit ball constitutes a partition of the unit ball, see Figures 1 and 2. We now describe a C^1 deformation of \mathbb{S}^2 in a neighborhood of

$$\mathcal{C} \cap \mathbb{S}^2 = \bigcup_{\substack{i, j \in \{1, 2, 3, 4\} \\ i \neq j}} \tilde{\gamma}_{ij}.$$

The resulting surface forms the boundary of a domain Ω which is partitioned by \mathcal{C} to form an isolated local minimizer of E_0 in the domain. This deformation is carried out so as to create six troughs along neighborhoods of each $\tilde{\gamma}_{ij}$, except near its endpoints \tilde{q}_{ijk} and \tilde{q}_{ijl} . By ‘troughs’ we mean that within this neighborhood of each $\tilde{\gamma}_{ij}$, the closest points to the origin are those on the circular arc itself, with the




FIGURE 1. Partition of the unit ball by the tetrahedral cone.




FIGURE 2. One component of the partition of the unit ball.

deformed surface sloping downwards towards $\tilde{\gamma}_{ij}$ on either side. Then, in a neighborhood of each of the four triple junctions \tilde{q}_{ijk} where three such troughs meet, we depress \mathbb{S}^2 to create small valleys, for example hemispheres. The deepest point of each valley we denote by q_{ijk} where, say, $q_{ijk} = (1 - \lambda)\tilde{q}_{ijk}$ for some small $\lambda > 0$. Finally, away

from these valleys and troughs we glue the surface in a C^1 manner to a slightly enlarged sphere, say of radius $1 + \lambda$, though as we will see, the particular shape away from neighborhoods of the troughs and valleys is unimportant. Of course, such a construction is not unique, but we describe in detail one such construction in Section 4.

This deformation of \mathbb{S}^2 now bounds an open set that we denote by Ω . The tetrahedral cone \mathcal{C} partitions Ω into four subsets that we henceforth denote by S_1, \dots, S_4 , where for each distinct i and j , $\partial S_i \cap \partial S_j$ and $\partial \tilde{S}_i \cap \partial \tilde{S}_j$ lie on the same plane. Finally, we let

$$\gamma_{ij} := \partial S_i \cap \partial S_j \cap \partial \Omega.$$

We note that in the case of highly non-symmetric cones \mathcal{C} , care must be taken to ensure that these valleys are all disjoint, as are the various troughs.

Ultimately, what we require of our domain Ω is that \mathcal{C} determines a partition Ω given by $\mathcal{S} := \{S_1, S_2, S_3, S_4\}$ satisfying the following property:

There exists a real number $\eta = \eta(\Omega) > 0$ such that for each distinct pair of indices $i, j \in \{1, 2, 3, 4\}$ and every point in the set

$$\{x \in \partial S_i \cap \partial \Omega : \text{dist}(x, \gamma_{ij}) < \eta\},$$

one has the property:

$$\begin{aligned} n_{ij} \cdot \nu_{\partial \Omega}(x) &> 0 && \text{if } x \notin \gamma_{ij}, \\ n_{ij} \cdot \nu_{\partial \Omega}(x) &= 0 && \text{if } x \in \gamma_{ij}. \end{aligned} \tag{2.4}$$

Remark 2.4. *We note that the arguments to follow only require that the property (2.4) holds \mathcal{H}^2 a.e. on the specified subset of $\partial \Omega$. In light of this, one could relax the regularity assumption on $\partial \Omega$ to be merely Lipschitz.*

3. THE MAIN THEOREM.

We now present our main result, which in light of previous results on Γ -convergence immediately implies Theorem 1.1. Throughout, we consider a fixed domain Ω satisfying the conditions (2.4) with respect to the normals coming from \mathcal{C} .

Theorem 3.1. *The partition \mathcal{S} is an isolated L^1 local minimizer of E_0 . That is, there exists $\delta > 0$ such that one has*

$$E_0(\mathcal{T}) > E_0(\mathcal{S}) \quad \text{provided } 0 < \mathcal{L}^3(\mathcal{T} \Delta \mathcal{S}) \leq \delta. \tag{3.1}$$

For ease of presentation, we will present the proof for the case of a symmetric tetrahedral cone, that is, for the case of equal weights c_{ij} .

However, in Remark 3.3 we point out the minor adaptation needed to handle the case of non-symmetric cones satisfying (1.6) provided the weights c_{ij} satisfy the condition (1.7).

Before beginning the proof, we consider for any $\delta > 0$ the following constrained partition problem:

$$\inf_{\mathcal{T} \in \mathcal{A}^\delta} E_0(\mathcal{T}), \quad (3.2)$$

where

$$\mathcal{A}^\delta := \{\mathcal{T} := (T_1, T_2, T_3, T_4) \text{ is a partition of } \Omega \text{ s.t. } \mathcal{L}^3(\mathcal{T} \Delta \mathcal{S}) \leq \delta\}. \quad (3.3)$$

The existence of a minimizer of this constrained problem is standard via the Direct Method, in light of the lower-semicontinuity of perimeter under L^1 -convergence. We denote a minimizer of this problem by

$$\mathcal{T}^\delta = (T_1^\delta, T_2^\delta, T_3^\delta, T_4^\delta).$$

The proof of Theorem 3.1 will rely crucially on the following ‘corralling’ result, which shows that for each $i \in \{1, 2, 3, 4\}$, $\partial T_i^\delta \cap \partial \Omega$ lies near $\partial S_i \cap \partial \Omega$. The argument represents a type of boundary ‘infiltration lemma’, and is an adaptation of the interior result [Mag12, Lemma 30.2].

Theorem 3.2. *There exists a number $\delta > 0$ depending on \mathcal{C} and Ω such that the following three conditions hold for the partition \mathcal{T}^δ :*

(I) *For each $i \in \{1, 2, 3, 4\}$, one has :*

$$\mathcal{H}^2 \left(\left\{ x \in \left(\bigcup_{\substack{j \in \{1, 2, 3, 4\} \\ i \neq j}} T_j^\delta \right) \cap \partial \Omega \cap \partial S_i : \text{dist}(x, \mathcal{C}) \geq \eta \right\} \right) = 0.$$

(II) *For distinct indices $i, j, k, l \in \{1, 2, 3, 4\}$, one has:*

$$\mathcal{H}^2(\{x : \text{dist}(x, q_{ijk}) < \eta\} \cap T_l^\delta) = 0.$$

(III) *For the distinct indices $i, j, k, l \in \{1, 2, 3, 4\}$, one has:*

$$\mathcal{H}^2(\{x : \text{dist}(x, \gamma_{ij}) < \eta\} \cap (T_k^\delta \cup T_l^\delta)) = 0.$$

Proof. We begin with the proof of (I), taking for instance, the case $i = 1$; the other cases follow similarly. Let $0 < r_1 < r_0$ be such that for all $x \in \partial S_1 \cap \partial \Omega$ with $\text{dist}(x, \mathcal{C}) \geq \eta$, one has

$$B(x, r_1) \cap \Omega \subset\subset S_1, \quad (3.4)$$

where the value r_0 is defined in Proposition 2.3. Consider any $x \in \partial S_1 \cap \partial \Omega$ such that $\text{dist}(x, \mathcal{C}) \geq \eta$ and suppose first that for some

$0 < r \leq r_1$ one has the condition

$$\mathcal{L}^3 \left([T_2^\delta \cup T_3^\delta \cup T_4^\delta] \cap B(x, r) \right) \leq \varepsilon_0 r^3, \text{ where } \varepsilon_0 := \left(\frac{C_0}{18} \right)^3, \quad (3.5)$$

and where C_0 is defined in Proposition 2.3. Then we will show that

$$\mathcal{L}^3 \left([T_2^\delta \cup T_3^\delta \cup T_4^\delta] \cap B \left(x, \frac{r}{2} \right) \right) = 0. \quad (3.6)$$

To this end, suppressing dependence on δ , we let

$$\begin{aligned} E_s &:= B(x, s) \cap [T_2^\delta \cup T_3^\delta \cup T_4^\delta] \quad \text{and} \\ m(s) &:= \mathcal{L}^3(E_s) = \int_0^s \sum_{j=2}^4 \mathcal{H}^2(T_j^\delta \cap \partial B(x, t)) dt, \end{aligned} \quad (3.7)$$

where the last equality in (3.7) follows from the co-area formula. Further, we have:

$$m'(s) = \sum_{j=2}^4 \mathcal{H}^2(T_j^\delta \cap \partial B(x, s)), \text{ for a.e. } 0 < s < r_1, \quad (3.8)$$

and since the perimeter of $[T_2^\delta \cup T_3^\delta \cup T_4^\delta]$ in Ω is finite, we have

$$\sum_{j=2}^4 \mathcal{H}^2(\partial^* T_j^\delta \cap \partial B(x, s)) = 0 \text{ for a.e. } 0 < s < r_1. \quad (3.9)$$

We will now modify the partition \mathcal{T}^δ by giving E_s to T_1^δ to form a new partition $\mathcal{F}(s) := (F(s, 1), F(s, 2), F(s, 3), F(s, 4))$ given by

$$\begin{aligned} F(s, 1) &:= T_1^\delta \cup E_s, \\ F(s, 2) &:= T_2^\delta - E_s, \\ F(s, 3) &:= T_3^\delta - E_s, \\ F(s, 4) &:= T_4^\delta - E_s, \end{aligned} \quad (3.10)$$

where we again have suppressed the δ dependence of \mathcal{F} . We now make the claim that for a.e. $0 < s < r_1$, one has the inequality,

$$\text{Per}(\mathcal{T}^\delta; \Omega \cap B(x, s)) - \text{Per}(\mathcal{F}(s); \Omega \cap B(x, s)) \leq m'(s). \quad (3.11)$$

To establish (3.11), we first observe that giving E_s to T_1^δ at worst leaves the L^1 distance to \mathcal{S} unchanged; hence the new partition satisfies:

$$\mathcal{L}^3(\mathcal{F}(s) \Delta \mathcal{S}) \leq \delta, \quad \forall 0 < s < r_1.$$

Thus, for all $0 < s < r_1$, $\mathcal{F}(s) \in \mathcal{A}^\delta$. Since, $\mathcal{T}^\delta = (T_1^\delta, T_2^\delta, T_3^\delta, T_4^\delta)$ is the minimizer, for all $0 < s < r_1$, we have

$$\begin{aligned} & \text{Per}(\mathcal{T}^\delta; \Omega \cap B(x, s)) + \sum_{j=1}^4 \mathcal{H}^2(\partial^* T_j^\delta \cap \partial B(x, s)) + \text{Per}(\mathcal{T}^\delta; \Omega - \bar{B}(x, s)) \\ &= \text{Per}(\mathcal{T}^\delta; \Omega) \leq \text{Per}(\mathcal{F}(s); \Omega) = \text{Per}(\mathcal{F}(s); \Omega \cap B(x, s)) \\ &+ \sum_{j=1}^4 \mathcal{H}^2(\partial^* F(s, j) \cap \partial B(x, s)) + \text{Per}(\mathcal{F}(s); \Omega - \bar{B}(x, s)). \end{aligned} \tag{3.12}$$

We observe now that

$$\text{Per}(\mathcal{T}^\delta; \Omega - \bar{B}(x, s)) = \text{Per}(\mathcal{F}(s); \Omega - \bar{B}(x, s))$$

for all $0 < s < r_1$, and also by (3.7) and (3.8), for a.e $0 < s < r_1$, one has

$$\sum_{j=1}^4 \mathcal{H}^2(\partial^* F(s, j) \cap \partial B(x, s)) = \sum_{j=2}^4 \mathcal{H}^2(T_j^\delta \cap \partial B(x, s)) = m'(s).$$

Applying these observations to (3.12) yields the claim (3.11).

We next claim that for a.e. $0 < s < r_1$, one has

$$\text{Per}(\mathcal{T}^\delta; \Omega \cap B(x, s)) - \text{Per}(\mathcal{F}(s); \Omega \cap B(x, s)) \geq \frac{C_0}{2} m(s)^{\frac{2}{3}} - m'(s). \tag{3.13}$$

To establish claim (3.13), we first note that

$$\begin{aligned} \sum_{j=2}^4 \text{Per}(T_j^\delta; \Omega \cap B(x, s)) &\geq \text{Per}(E_s; \Omega \cap B(x, s)) \\ &= \text{Per}(E_s; \Omega) - \mathcal{H}^2(\partial E_s \cap \partial B(x, s)) \\ &= \text{Per}(E_s; \Omega) - \sum_{j=2}^4 \mathcal{H}^2(T_j^\delta \cap \partial B(x, s)) \\ &\geq C_0 m(s)^{\frac{2}{3}} - m'(s), \end{aligned} \tag{3.14}$$

for a.e $0 < s < r_1$, where the last inequality comes from the isoperimetric inequality, Proposition 2.3. Next we observe that

$$\begin{aligned} & 2 [\text{Per}(\mathcal{T}^\delta; \Omega \cap B(x, s)) - \text{Per}(\mathcal{F}(s); \Omega \cap B(x, s))] \\ &= \sum_{j=1}^4 \text{Per}(T_j^\delta; \Omega \cap B(x, s)) - \sum_{j=1}^4 \text{Per}(F(s, j); \Omega \cap B(x, s)), \end{aligned} \tag{3.15}$$

for any $0 < s < r_1$. By the construction of $\mathcal{F}(s)$, the entire set $\Omega \cap B(x, s)$ consists of $F(s, 1)$. Hence, for any $0 < s < r_1$, we have

$$\sum_{j=1}^4 \text{Per}(F(s, j); \Omega \cap B(x, s)) = 0.$$

Consequently, by (3.14) and (3.15), we have

$$\begin{aligned} & 2 \left[\text{Per}(\mathcal{T}^\delta; \Omega \cap B(x, s)) - \text{Per}(\mathcal{F}(s); \Omega \cap B(x, s)) \right] \\ & \geq \text{Per}(T_1^\delta; \Omega \cap B(x, s)) + C_0 m(s)^{\frac{2}{3}} - m'(s) \\ & \geq C_0 m(s)^{\frac{2}{3}} - m'(s), \end{aligned}$$

thus establishing claim (3.13).

The inequalities (3.11) and (3.13) together now yield

$$C_0 m(s)^{\frac{2}{3}} - m'(s) \leq 2m'(s),$$

for a.e. $0 < s < r_1$, or equivalently,

$$\frac{9}{C_0} \frac{d}{ds} \left[\sqrt[3]{m(s)} \right] \geq 1. \quad (3.16)$$

Since the map $s \mapsto m(s)$ is monotonically increasing, we have that the support of m is $[r^*, \infty)$, for some $r^* \geq 0$. If $m(r) = 0$ for r in (3.5), then trivially (3.6) holds. Otherwise $m(r) > 0$ and so necessarily $r > r^*$. Integrating (3.16) on (r^*, r) gives

$$\begin{aligned} r - r^* & \leq \frac{9}{C_0} \left[\sqrt[3]{m(r)} - \sqrt[3]{m(r^*)} \right] \leq \frac{9}{C_0} \sqrt[3]{m(r)} \\ & \leq \frac{9}{C_0} \sqrt[3]{\varepsilon_0 r} = \frac{r}{2}, \end{aligned} \quad (3.17)$$

where we used (3.5) for the last inequality in (3.17). Hence, $r^* \geq r/2 > 0$, and so the assumption (3.5) implies the conclusion (3.6).

On the other hand, consider any $x \in \partial S_1 \cap \partial \Omega$ such that $\text{dist}(x, \mathcal{C}) \geq \eta$ for which assumption (3.5) does not hold. That is, suppose

$$\forall r \leq r_1, \quad \mathcal{L}^3(T^\delta \cap B(x, r)) > \varepsilon_0 r^3. \quad (3.18)$$

In particular, take $r = \sqrt[4]{\delta}$, where we stipulate that

$$\delta < \min \left\{ \left(\frac{r_1}{2} \right)^4, \frac{\varepsilon_0^4}{2} \right\}.$$

Plugging this r in (3.18) and noting that $\mathcal{L}^3(\mathcal{T}^\delta \Delta \mathcal{S}) \leq \delta$, we reach a contradiction to the assumption (3.18) in that

$$\delta \geq \mathcal{L}^3(T^\delta \cap B(x, r_1)) > \varepsilon_0 r^3 = \varepsilon_0 \delta^{\frac{3}{4}}. \quad (3.19)$$

The proof of (I) is complete.

Now, we turn to the proof of (II). We will consider the case $i = 1, j = 2, k = 3, l = 4$, so our goal is to show

$$\mathcal{H}^2(T_4^\delta \cap \{x \in \partial\Omega : \text{dist}(x, q_{123}) < \eta\}) = 0. \quad (3.20)$$

The other cases follow in the similar manner. To this end, we fix any $0 < r_2 < r_0$ such that

$$r_2 < \frac{1}{2} \min \left\{ \text{dist}(B_{i_1 i_2 i_3}, B_{j_1 j_2 j_3}) \left| \begin{array}{l} i_1, j_1, i_2, j_2, i_3, j_3 \in \{1, 2, 3, 4\}, \\ i_k \neq j_k \text{ for all } k = 1, 2, 3. \end{array} \right. \right\} \quad (3.21)$$

where $B_{ijk} := \{x \in \partial\Omega; \text{dist}(x, q_{ijk}) < \eta\}$ and again, the value r_0 comes from Proposition 2.3.

We consider a point $x_0 \in B_{123}$ for $0 < s < r_2$. In analogy with (3.5) from the proof of (I), we assume that for some $0 < r < r_2$, one has

$$\mathcal{L}^3(B(x_0, r) \cap T_4^\delta) \leq \varepsilon_0 r^3.$$

The primary difference between the proof of (II) and (I) is that here we create a competing partition by giving $B(x_0, s) \cap T_4^\delta$ to one of T_1^δ, T_2^δ or T_3^δ , say to T_1^δ , rather than giving $B(x_0, s) \cap (\cup_{j=2}^4 T_j^\delta)$ to T_1^δ . That is, for all $0 < s < r$, we set

$$\begin{aligned} G(s, 1) &:= T_1^\delta \cup [B(x_0, s) \cap T_4^\delta], \\ G(s, 2) &:= T_2^\delta, \\ G(s, 3) &:= T_3^\delta, \\ G(s, 4) &:= T_4^\delta - B(x_0, s), \\ \mathcal{G}(s) &:= (G(s, 1), G(s, 2), G(s, 3), G(s, 4)), \end{aligned} \quad (3.22)$$

where we again have suppressed the δ dependence on \mathcal{G} . Giving $T_4^\delta \cap B(x_0, s)$ to T_1^δ either decreases the L^1 distance to \mathcal{S} or at worst, the L^1 distance remains unchanged. Therefore, we have

$$\forall 0 < s < r, \quad \mathcal{L}^3(\mathcal{S} \Delta \mathcal{G}(s)) \leq \delta \text{ and so } \mathcal{G} \in \mathcal{A}^\delta.$$

Here, the role of $m(s)$ from (3.7) is taken up by say $n(s)$, where for all $0 < s < r$, we set

$$\begin{aligned} n(s) &:= \mathcal{L}^3(T_4^\delta \cap B(x_0, s)) \\ &= \int_0^s \mathcal{H}^2[T_4^\delta \cap \partial B(x_0, t)] dt. \end{aligned} \quad (3.23)$$

Further, for a.e. $0 < s < r$, we have:

$$n'(s) = \mathcal{H}^2[T_4^\delta \cap \partial B(x_0, s)] \text{ and } \mathcal{H}^2[\partial^* T_4^\delta \cap \partial B(x_0, s)] = 0, \quad (3.24)$$

where again we have suppressed the δ dependence on n . Since \mathcal{T}^δ is a minimizer of (3.2), in the same manner in which we established (3.11), for a.e $0 < s < r$, one has

$$\text{Per}(\mathcal{T}^\delta; \Omega \cap B(x_0, s)) - \text{Per}(\mathcal{G}(s); \Omega \cap B(x_0, s)) \leq n'(s). \quad (3.25)$$

To establish the analogue of (3.13), we begin by noting, for any $0 < s < r$ that:

$$\begin{aligned} & 2 [\text{Per}(\mathcal{T}^\delta; \Omega \cap B(x_0, s)) - \text{Per}(\mathcal{G}(s); \Omega \cap B(x_0, s))] \\ &= \sum_{i=1}^4 \text{Per}(T_i^\delta; \Omega \cap B(x_0, s)) - \sum_{i=1}^4 \text{Per}(G(s, i); \Omega \cap B(x_0, s)) \\ &= \sum_{i=1}^4 \mathcal{H}^2(\partial^* T_i^\delta \cap \Omega \cap B(x_0, s)) - \sum_{i=1}^4 \mathcal{H}^2(\partial^* G(s, i) \cap \Omega \cap B(x_0, s)) \\ &= \sum_{i=1,4} \left[\mathcal{H}^2(\partial^* T_i^\delta \cap \Omega \cap B(x_0, s)) - \mathcal{H}^2(\partial^* G(s, i) \cap \Omega \cap B(x_0, s)) \right], \end{aligned} \quad (3.26)$$

where in the final line of (3.26), we have used

$$\mathcal{H}^2(\partial^* T_i^\delta \cap \Omega \cap B(x_0, s)) = \mathcal{H}^2(\partial^* G(s, i) \cap \Omega \cap B(x_0, s)), \text{ for } i = 2, 3.$$

Next, focusing on $i = 1$ in the last line of (3.26), one has

$$\begin{aligned} & \mathcal{H}^2(\partial^* T_1^\delta \cap \Omega \cap B(x_0, s)) - \mathcal{H}^2(\partial^* G(s, 1) \cap \Omega \cap B(x_0, s)) \\ &= \sum_{j=2,3,4} \mathcal{H}^2(\partial^* T_j^\delta \cap \partial^* T_1^\delta \cap \Omega \cap B(x_0, s)) - \sum_{j=2,3} \mathcal{H}^2(\partial^* T_j^\delta \cap \partial^* T_1^\delta \cap \Omega \cap B(x_0, s)) \\ &= \mathcal{H}^2(\partial^* T_4^\delta \cap \partial^* T_1^\delta \cap \Omega \cap B(x_0, s)) \geq 0. \end{aligned} \quad (3.27)$$

Now focusing on $i = 4$ in the last line of (3.26), one has

$$\text{Per}(G(s, 4); \Omega \cap B(x_0, s)) = 0,$$

and so, invoking the isoperimetric inequality, we obtain the following analogue of (3.13):

$$\text{Per}(T_4^\delta; \Omega \cap B(x_0, s)) \geq C_0 n(s)^{\frac{2}{3}} - n'(s). \quad (3.28)$$

Implementing (3.27)-(3.28) in (3.26), we have

$$\text{Per}(\mathcal{T}^\delta; \Omega \cap B(x_0, s)) - \text{Per}(\mathcal{G}(s, 1); \Omega \cap B(x_0, s)) \geq C_0 n(s)^{\frac{2}{3}} - n'(s). \quad (3.29)$$

Combining (3.25) and (3.29), we reach the differential inequality:

$$\frac{9}{C_0} \frac{\partial}{\partial s} \left[\sqrt[3]{n(s)} \right] \geq 1.$$

As seen in the proof of (I), the map $s \mapsto n(s)$ is monotonically increasing and so, we can denote the support of this map $s \mapsto n(s, 1)$ by $[r_2^*, \infty)$, for some $r_2^* > 0$. With the further stipulation on δ that

$$\delta < \min \left\{ \left(\frac{r_2}{2} \right)^4, \frac{\varepsilon_0^4}{2} \right\},$$

an analysis similar as in (3.17) implies that

$$n \left(\frac{r}{2} \right) = 0.$$

Furthermore, by the same reasoning seen in (3.18)-(3.19), one cannot have that for all $0 < r < r_2$, $n(r) > \varepsilon_0 r^3$. This concludes the proof of (II).

We now turn to the proof of (III) and consider, for example, the case $i = 1, j = 2, k = 3, l = 4$; the other cases follow similarly. That is, we need to show that

$$\mathcal{H}^2 \left(\{x \in \partial\Omega : \text{dist}(x, \gamma_{12}) < \eta\} \cap (T_3^\delta \cup T_4^\delta) \right) = 0. \quad (3.30)$$

Proceeding as in the proofs of (I) and (II), we let $0 < r_3 < r_0$ be a number such that

$$\{x \in \partial\Omega : \text{dist}(x, \gamma_{12}) < \eta\} \cap B(x, r_3) \subset\subset (\partial S_1 \cup \partial S_2) \cap \partial\Omega. \quad (3.31)$$

Now, let $x_0 \in \{x \in \partial\Omega : \text{dist}(x, \gamma_{12}) < \eta\}$ and assume that for some $0 < r < r_3$, one has

$$\mathcal{L}^3((T_3^\delta \cup T_4^\delta) \cap B(x_0, r)) < \varepsilon_0 r^3.$$

Then, we give the region $T_3^\delta \cup T_4^\delta$ in $B(x_0, r)$ to one of the regions T_1^δ or T_2^δ - say we give it to T_1^δ . That is, for $0 < s < r_3$, we create a new partition:

$$\begin{aligned} H(s, 1) &:= T_1^\delta \cup \left[B(x_0, s) \cap (T_3^\delta \cup T_4^\delta) \right], \\ H(s, 2) &:= T_2^\delta, \\ H(s, 3) &:= T_3^\delta - B(x_0, s), \\ H(s, 4) &:= T_4^\delta - B(x_0, s), \\ \mathcal{H}(s) &:= (H(s, 1), H(s, 2), H(s, 3), H(s, 4)). \end{aligned} \quad (3.32)$$

Giving $T_3^\delta \cup T_4^\delta$ in $B(x_0, r)$ to T_1^δ either decreases the L^1 distance to \mathcal{S} or at worst, the L^1 distance remains unchanged. Hence $\mathcal{H}(s) \in \mathcal{A}^\delta$. Finally, with the further stipulation on δ that

$$\delta < \min \left\{ \left(\frac{r_3}{2} \right)^4, \frac{\varepsilon_0^4}{2} \right\},$$

an analysis similar to the proofs of (I) and (II) yields

$$\mathcal{L}^3 \left((T_3^\delta \cap T_4^\delta) \cap B \left(x_0, \frac{r}{2} \right) \right) = 0,$$

implying (3.30), and by the same reasoning seen before in the proofs of (I) and (II), one cannot have that for all $0 < r < r_3$,

$$\mathcal{L}^3 \left((T_3^\delta \cap T_4^\delta) \cap B(x_0, r) \right) > \varepsilon_0 r^3.$$

Combining the various conditions on δ , the required δ for Theorem 3.2 to hold true is

$$0 < \delta \leq \min \left\{ \left(\frac{r_1}{2} \right)^4, \left(\frac{r_2}{2} \right)^4, \left(\frac{r_3}{2} \right)^4, \frac{1}{2} \left(\frac{C_0}{18} \right)^{12} \right\}.$$

□

Having argued that the constrained minimizer \mathcal{T}^δ to problem (3.2) has the property that for each $i \in \{1, 2, 3, 4\}$, ∂T_i^δ lies in an η neighborhood of ∂S_i on $\partial\Omega$, we can now apply a calibration argument inspired by that of [LM94].

Proof of Theorem 3.1. With Theorem 3.2 in hand, we now argue that in fact, \mathcal{T}^δ coincides with \mathcal{S} . In what follows, for ease of notation, we will not indicate the δ dependence of \mathcal{T}^δ and write simply $\mathcal{T} = (T_1, T_2, T_3, T_4)$. We recall that n_{ij} denotes the unit normal to $\partial S_i \cap \partial S_j$ pointing from S_i to S_j , $\nu_{\partial\Omega}$ denotes the unit outward normal to $\partial\Omega$, and we let ν_{ij}^* denote the unit normal to $\partial^* T_i \cap \partial^* T_j$ pointing from T_i to T_j ,

We begin with four applications of the Gauss-Green theorem to obtain

$$\begin{aligned} 0 = \int_{T_1} \operatorname{div}(n_{14}) dx &= \int_{\partial^* T_1 \cap \partial^* T_2} n_{14} \cdot \nu_{12}^* d\mathcal{H}^2 + \int_{\partial^* T_1 \cap \partial^* T_3} n_{14} \cdot \nu_{13}^* d\mathcal{H}^2 \\ &\quad + \int_{\partial^* T_1 \cap \partial^* T_4} n_{14} \cdot \nu_{14}^* d\mathcal{H}^2 + \int_{\partial\Omega \cap \partial^* T_1} n_{14} \cdot \nu_{\partial\Omega} d\mathcal{H}^2, \end{aligned} \tag{3.33}$$

$$\begin{aligned} 0 = \int_{T_2} \operatorname{div}(n_{14}) dx &= \int_{\partial^* T_2 \cap \partial^* T_1} n_{14} \cdot \nu_{21}^* d\mathcal{H}^2 + \int_{\partial^* T_2 \cap \partial^* T_3} n_{14} \cdot \nu_{23}^* d\mathcal{H}^2 \\ &\quad + \int_{\partial^* T_2 \cap \partial^* T_4} n_{14} \cdot \nu_{24}^* d\mathcal{H}^2 + \int_{\partial\Omega \cap \partial^* T_2} n_{14} \cdot \nu_{\partial\Omega} d\mathcal{H}^2, \end{aligned} \tag{3.34}$$

$$\begin{aligned}
0 = \int_{T_2} \operatorname{div}(n_{21}) dx &= \int_{\partial^* T_2 \cap \partial^* T_1} n_{21} \cdot \nu_{21}^* d\mathcal{H}^2 + \int_{\partial^* T_2 \cap \partial^* T_3} n_{21} \cdot \nu_{23}^* d\mathcal{H}^2 \\
&\quad + \int_{\partial^* T_2 \cap \partial^* T_4} n_{21} \cdot \nu_{24}^* d\mathcal{H}^2 + \int_{\partial\Omega \cap \partial^* T_2} n_{21} \cdot \nu_{\partial\Omega} d\mathcal{H}^2,
\end{aligned} \tag{3.35}$$

and

$$\begin{aligned}
0 = \int_{T_3} \operatorname{div}(n_{34}) dx &= \int_{\partial^* T_3 \cap \partial^* T_1} n_{34} \cdot \nu_{31}^* d\mathcal{H}^2 + \int_{\partial^* T_3 \cap \partial^* T_2} n_{34} \cdot \nu_{32}^* d\mathcal{H}^2 \\
&\quad + \int_{\partial^* T_3 \cap \partial^* T_4} n_{34} \cdot \nu_{34}^* d\mathcal{H}^2 + \int_{\partial\Omega \cap \partial^* T_3} n_{34} \cdot \nu_{\partial\Omega} d\mathcal{H}^2.
\end{aligned} \tag{3.36}$$

We remark that the choice of the normals and the domains of integration is not unique. For example, we could also have begun with

$$0 = \int_{T_4} \operatorname{div}(n_{41}) dx + \int_{T_4} \operatorname{div}(n_{13}) dx + \int_{T_1} \operatorname{div}(n_{13}) dx + \int_{T_2} \operatorname{div}(n_{23}) dx.$$

Using the cyclic property of the normals (1.5), the fact that $\nu_{ij}^* = -\nu_{ji}^*$ and $n_{ij} = -n_{ji}$, adding the equations (3.33), (3.34), (3.35) and (3.36), we get

$$\begin{aligned}
0 &= \int_{T_1} \operatorname{div}(n_{14}) dx + \int_{T_2} \operatorname{div}(n_{14}) dx + \int_{T_2} \operatorname{div}(n_{21}) dx + \int_{T_3} \operatorname{div}(n_{34}) dx \\
&= \int_{\partial^* T_2 \cap \partial^* T_1} n_{21} \cdot \nu_{21}^* d\mathcal{H}^2 + \int_{\partial^* T_1 \cap \partial^* T_3} n_{31} \cdot \nu_{31}^* d\mathcal{H}^2 \\
&\quad + \int_{\partial^* T_1 \cap \partial^* T_4} n_{14} \cdot \nu_{14}^* d\mathcal{H}^2 + \int_{\partial^* T_3 \cap \partial^* T_2} n_{23} \cdot \nu_{23}^* d\mathcal{H}^2 \\
&\quad + \int_{\partial^* T_2 \cap \partial^* T_4} n_{24} \cdot \nu_{24}^* d\mathcal{H}^2 + \int_{\partial^* T_3 \cap \partial^* T_4} n_{34} \cdot \nu_{34}^* d\mathcal{H}^2 \\
&\quad - \int_{\partial\Omega \cap \partial^* T_1} n_{41} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 - \int_{\partial\Omega \cap \partial^* T_2} n_{42} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 - \int_{\partial\Omega \cap \partial^* T_3} n_{43} \cdot \nu_{\partial\Omega} d\mathcal{H}^2.
\end{aligned} \tag{3.37}$$

Rearranging this identity, we have:

$$\begin{aligned}
& \int_{\partial^* T_2 \cap \partial^* T_1} n_{21} \cdot \nu_{21}^* d\mathcal{H}^2 + \int_{\partial^* T_1 \cap \partial^* T_3} n_{31} \cdot \nu_{31}^* d\mathcal{H}^2 \\
& \quad + \int_{\partial^* T_1 \cap \partial^* T_4} n_{14} \cdot \nu_{14}^* d\mathcal{H}^2 + \int_{\partial^* T_3 \cap \partial^* T_2} n_{23} \cdot \nu_{23}^* d\mathcal{H}^2 \\
& \quad + \int_{\partial^* T_2 \cap \partial^* T_4} n_{24} \cdot \nu_{24}^* d\mathcal{H}^2 + \int_{\partial^* T_3 \cap \partial^* T_4} n_{34} \cdot \nu_{34}^* d\mathcal{H}^2 \\
& = \int_{\partial\Omega \cap \partial^* T_1} n_{41} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial\Omega \cap \partial^* T_2} n_{42} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial\Omega \cap \partial^* T_3} n_{43} \cdot \nu_{\partial\Omega} d\mathcal{H}^2.
\end{aligned} \tag{3.38}$$

Since $n_{ij} \cdot \nu_{ij}^* \leq 1$ for all i, j we conclude that

$$\begin{aligned}
E_0(\mathcal{T}) & := \sum_{1 \leq i < j \leq 4} \mathcal{H}^2(\partial^* T_i \cap \partial^* T_j \cap \Omega) \\
& \geq \int_{\partial^* T_2 \cap \partial^* T_1} n_{21} \cdot \nu_{21}^* d\mathcal{H}^2 + \int_{\partial^* T_1 \cap \partial^* T_3} n_{31} \cdot \nu_{31}^* d\mathcal{H}^2 \\
& \quad + \int_{\partial^* T_1 \cap \partial^* T_4} n_{14} \cdot \nu_{14}^* d\mathcal{H}^2 + \int_{\partial^* T_3 \cap \partial^* T_2} n_{23} \cdot \nu_{23}^* d\mathcal{H}^2 \\
& \quad + \int_{\partial^* T_2 \cap \partial^* T_4} n_{24} \cdot \nu_{24}^* d\mathcal{H}^2 + \int_{\partial^* T_3 \cap \partial^* T_4} n_{34} \cdot \nu_{34}^* d\mathcal{H}^2 \\
& = \int_{\partial\Omega \cap \partial^* T_1} n_{41} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial\Omega \cap \partial^* T_2} n_{42} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial\Omega \cap \partial^* T_3} n_{43} \cdot \nu_{\partial\Omega} d\mathcal{H}^2.
\end{aligned} \tag{3.39}$$

Here we also observe that if we replace \mathcal{T} by \mathcal{S} in the calculation above, then one instead obtains the identity

$$E_0(\mathcal{S}) = \int_{(\partial\Omega \cap \partial S_1)} n_{41} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{(\partial\Omega \cap \partial S_2)} n_{42} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{(\partial\Omega \cap \partial S_3)} n_{43} \cdot \nu_{\partial\Omega} d\mathcal{H}^2. \tag{3.40}$$

We will first argue that \mathcal{T} and \mathcal{S} agree on $\partial\Omega$. To this end, we decompose the domains of integration in the last line of (3.39), by writing

each $\partial\Omega \cap \partial^*T_i$ as $\partial\Omega \cap \partial S_i$ plus corrections, for $i = 1, 2, 3$:

$$\begin{aligned}
E_0(\mathcal{T}) &\geq \int_{\partial\Omega \cap \partial^*T_1} n_{41} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial\Omega \cap \partial^*T_2} n_{42} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial\Omega \cap \partial^*T_3} n_{43} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 \\
&= \int_{(\partial\Omega \cap \partial S_1)} n_{41} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{(\partial\Omega \cap \partial S_2)} n_{42} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{(\partial\Omega \cap \partial S_3)} n_{43} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 \\
&\quad + \int_{(\partial^*T_2 \cup \partial^*T_3 \cup \partial^*T_4) \cap \partial S_1} n_{14} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial^*T_1 \cap (\partial S_2 \cup \partial S_3 \cup \partial S_4)} n_{41} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 \\
&\quad + \int_{(\partial^*T_1 \cup \partial^*T_3 \cup \partial^*T_4) \cap \partial S_2} n_{24} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial^*T_2 \cap (\partial S_1 \cup \partial S_3 \cup \partial S_4)} n_{42} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 \\
&\quad + \int_{(\partial^*T_1 \cup \partial^*T_2 \cup \partial^*T_4) \cap \partial S_3} n_{34} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial^*T_3 \cap (\partial S_1 \cup \partial S_2 \cup \partial S_4)} n_{43} \cdot \nu_{\partial\Omega} d\mathcal{H}^2.
\end{aligned} \tag{3.41}$$

By (3.40), we have

$$\begin{aligned}
E_0(\mathcal{T}) &\geq E_0(\mathcal{S}) \\
&\quad + \int_{(\partial^*T_2 \cup \partial^*T_3 \cup \partial^*T_4) \cap \partial S_1} n_{14} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial^*T_1 \cap (\partial S_2 \cup \partial S_3 \cup \partial S_4)} n_{41} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 \\
&\quad + \int_{(\partial^*T_1 \cup \partial^*T_3 \cup \partial^*T_4) \cap \partial S_2} n_{24} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial^*T_2 \cap (\partial S_1 \cup \partial S_3 \cup \partial S_4)} n_{42} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 \\
&\quad + \int_{(\partial^*T_1 \cup \partial^*T_2 \cup \partial^*T_4) \cap \partial S_3} n_{34} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial^*T_3 \cap (\partial S_1 \cup \partial S_2 \cup \partial S_4)} n_{43} \cdot \nu_{\partial\Omega} d\mathcal{H}^2.
\end{aligned} \tag{3.42}$$

Then, invoking property (1.5), we obtain

$$\begin{aligned}
E_0(\mathcal{T}) &\geq E_0(\mathcal{S}) \\
&\quad + \int_{\partial S_1 \cap \partial^*T_2} n_{12} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial S_1 \cap \partial^*T_3} n_{13} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial S_1 \cap \partial^*T_4} n_{14} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 \\
&\quad + \int_{\partial S_2 \cap \partial^*T_1} n_{21} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial S_2 \cap \partial^*T_3} n_{23} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial S_2 \cap \partial^*T_4} n_{24} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 \\
&\quad + \int_{\partial S_3 \cap \partial^*T_1} n_{31} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial S_3 \cap \partial^*T_2} n_{32} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial S_3 \cap \partial^*T_4} n_{34} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 \\
&\quad + \int_{\partial S_4 \cap \partial^*T_1} n_{41} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial S_4 \cap \partial^*T_2} n_{42} \cdot \nu_{\partial\Omega} d\mathcal{H}^2 + \int_{\partial S_4 \cap \partial^*T_3} n_{43} \cdot \nu_{\partial\Omega} d\mathcal{H}^2.
\end{aligned} \tag{3.43}$$

Next we observe that each of the twelve integrals in (3.43) is non-negative by property (2.4) and Theorem 3.2. To conclude, from (3.41)-(3.43), we see that unless

$$\sum_{i=1}^4 \mathcal{H}^2 \left[(\partial^* T_i \Delta \partial S_i) \cap \partial \Omega \right] = 0, \quad (3.44)$$

we would find that $E_0(\mathcal{T}) > E_0(\mathcal{S})$, contradicting the minimality of \mathcal{T} . Consequently, the last line of (3.39) equals $E_0(\mathcal{S})$, and so we find that

$$\begin{aligned} E_0(\mathcal{S}) \geq E_0(\mathcal{T}) &\geq \int_{\partial^* T_2 \cap \partial^* T_1} n_{21} \cdot \nu_{21}^* d\mathcal{H}^2 + \int_{\partial^* T_1 \cap \partial^* T_3} n_{31} \cdot \nu_{31}^* d\mathcal{H}^2 \\ &+ \int_{\partial^* T_1 \cap \partial^* T_4} n_{14} \cdot \nu_{14}^* d\mathcal{H}^2 + \int_{\partial^* T_3 \cap \partial^* T_2} n_{23} \cdot \nu_{23}^* d\mathcal{H}^2 \\ &+ \int_{\partial^* T_2 \cap \partial^* T_4} n_{24} \cdot \nu_{24}^* d\mathcal{H}^2 + \int_{\partial^* T_3 \cap \partial^* T_4} n_{34} \cdot \nu_{34}^* d\mathcal{H}^2 \\ &= E_0(\mathcal{S}), \end{aligned} \quad (3.45)$$

again contradicting the minimality of \mathcal{T} unless the inequalities in (3.45) are in fact equalities. We conclude that for all distinct $i, j \in \{1, 2, 3, 4\}$, $n_{ij} = \nu_{ij}^*$, \mathcal{H}^2 a.e. on $\partial^* T_i \cap \partial^* T_j \cap \Omega$. Hence, with an appeal to Lemma 2.2 we conclude that $\partial^* T_i \cap \partial^* T_j$ are all planar and by (3.44) they agree with $\partial S_i \cap \partial S_j$ on $\partial \Omega$. Necessarily, $\mathcal{T} = \mathcal{S}$ and the proof is complete. \square

Remark 3.3. *When c_{ij} 's are all not equal, the only change in the proof of Theorem 3.1 is to replace n_{ij} with $c_{ij}n_{ij}$ for each distinct pair $i, j \in \{1, 2, 3, 4\}$, and then to apply the more general stationarity condition (1.6) in place of (1.5). The proof of Theorem 3.2 remains unchanged in this more general setting.*

4. CONSTRUCTION OF A SUITABLE DOMAIN

In this section we describe one procedure by which a perturbation of \mathbb{S}^2 can serve as the boundary of a domain Ω satisfying (2.4). It will suffice to describe the construction of one ‘trough’ and one ‘valley’ since the other five troughs and three valleys can be built similarly.

To this end, let us suppose that, say, $\partial \tilde{S}_1 \cap \partial \tilde{S}_2$ lies within the xz -plane, where as in Section 2.3, the tetradedral cone partitions the unit ball in \mathbb{R}^3 into $\{\tilde{S}_1, \dots, \tilde{S}_4\}$. Suppose further that \tilde{S}_1 lies in the region $y > 0$ so that the normal n_{12} is given by $(0, -1, 0)$. Fixing a small number r_* , we will place a hemispherical ‘valley’ of radius r_* centered

at a point on the z -axis below $(0, 0, 1)$. Then away from this valley, the ‘trough’ will lie in a neighborhood of the circular arc $\tilde{\gamma}_{12}$ given by

$$\tilde{\gamma}_{12} = \{(x, 0, \sqrt{1-x^2}) : 2r_* < x < \bar{x}\}, \quad (4.1)$$

for some $\bar{x} > 0$. We note that within this trough the set $\partial S_1 \cap \partial S_2 \cap \partial \Omega$ will agree with $\tilde{\gamma}_{12}$, where $\partial \Omega$ denotes the region perturbed from \mathbb{S}^2 . The number \bar{x} is unimportant since the focus here is just on constructing one trough, one valley and then connecting them up in a C^1 manner while preserving conditions (2.4).

Construction of the trough. For each $x \in (2r_*, \bar{x})$, consider the plane having normal given by the tangent vector to $\tilde{\gamma}_{12}$. Restricted to each such plane, say Π_x , the trough will be given by a circular arc of radius r_* . To execute this part of the construction, we start with the circular arc in the yz -plane given by:

$$\{(0, y, z) : z = r_* - \sqrt{r_*^2 - y^2}, |y| < r_*/2\}$$

Then we apply the rotation matrix

$$R(x) := \begin{pmatrix} \cos \alpha(x) & 0 & -\sin \alpha(x) \\ 0 & 1 & 0 \\ \sin \alpha(x) & 0 & \cos \alpha(x) \end{pmatrix}$$

to this arc, where

$$\tan \alpha(x) = \left(\sqrt{1-x^2}\right)' = -\frac{x}{\sqrt{1-x^2}}, \quad (4.2)$$

so that the arc lies in a plane parallel to Π_x . Finally, we translate this rotated arc so that it sits on $\tilde{\gamma}_{12}$. This results in a trough, given parametrically by a map F :

$$F(x, y) := \left(x - (r_* - \sqrt{r_*^2 - y^2}) \sin \alpha(x), y, \sqrt{1-x^2} + (r_* - \sqrt{r_*^2 - y^2}) \cos \alpha(x) \right), \quad (4.3)$$

for $2r_* < x < \bar{x}$, $|y| < r_*/2$.

Now for this trough we check the condition (2.4) with $i = 1$, $j = 2$, $n_{12} = (0, -1, 0)$ and

$$\nu_{\partial \Omega}(x, y) = \frac{F_x \times F_y}{|F_x \times F_y|}. \quad (4.4)$$

Hence, property (2.4) will hold along the trough provided the second component of $\nu_{\partial \Omega}$ satisfies the conditions

$$\nu_{\partial \Omega}^{(2)}(x, y) < 0 \text{ for } 2r_* < x < \bar{x}, 0 < y < r_*/2,$$

and

$$\nu_{\partial\Omega}^{(2)}(x, 0) = 0 \text{ for } 2r_* < x < \bar{x}, \quad 0 < y < r_*/2.$$

A routine calculation using (4.2), (4.3) and (4.4) yields that

$$\nu_{\partial\Omega}^{(2)}(x, y) = -y \frac{(r_*^2 - \sqrt{r_*^2 - y^2})(1 + (1 - x^2)^2)}{\sqrt{r_*^2 - y^2} (1 - x^2)^{5/2}}, \quad (4.5)$$

and so these two conditions are met.

Connecting the trough to a valley. The trough just constructed will ultimately connect in a C^1 manner to a lower hemisphere of radius r_* , centered at the point $(0, 0, 1 - 2r_*)$, namely

$$\{(x, y, z) : x^2 + y^2 + (z - (1 - 2r_*))^2 = r_*^2, \quad z \leq 1 - 2r_*\}. \quad (4.6)$$

Within this hemisphere, the outer normal is given by

$$\nu_{\partial\Omega} = \frac{(-x, -y, 1 - 2r_* - z)}{\sqrt{x^2 + y^2 + (z - (1 - 2r_*))^2}},$$

so clearly $\nu_{\partial\Omega}^{(2)} < 0$ for $y > 0$ and $\nu_{\partial\Omega}^{(2)} = 0$ for $y = 0$ and (2.4) is again satisfied.

What remains is to connect the trough to this hemisphere in a C^1 manner. Here it is more convenient to use y and z as parameters, rather than x and y . To this end we introduce a smooth interpolating function $g(z)$ that will serve to define $\partial S_1 \cap \partial S_2 \cap \partial\Omega$ via the curve $(g(z), 0, z)$ as a replacement for (4.1). To ensure C^1 contact between this interpolating trough and the previous trough and between the interpolating trough and the hemisphere, we take $g : [1 - 2r_*, \sqrt{1 - 4r_*^2}] \rightarrow \mathbb{R}$ to be any smooth function satisfying the conditions:

$$\begin{aligned} g(1 - 2r_*) &= r_*, & g(\sqrt{1 - 4r_*^2}) &= 2r_*, \\ g'(1 - 2r_*) &= 0, & g'(\sqrt{1 - 4r_*^2}) &= -\frac{\sqrt{1 - 4r_*^2}}{2r_*}, \end{aligned}$$

with $g'(z) > 0$ and $g''(z) > 0$ on $(1 - 2r_*, \sqrt{1 - 4r_*^2})$.

Then, analogous to the procedure used to obtain the previous portion of the trough, we obtain this interpolation portion by first rotating the circular arc in the xy -plane given by

$$x = -r_* + \sqrt{r_*^2 - y^2}$$

through an angle $\beta(z)$ such that $\tan \beta(z) = g'(z)$ and then translating it by $(g(z), 0, z)$. Again, the resulting surface has cross-sections on

planes with normal in the direction $(g'(z), 0, 1)$ given by circular arcs. The parametrization for this portion of the trough is

$$G(y, z) := \left(g(z) - (r_* - \sqrt{r_*^2 - y^2}) \cos \beta(z), y, z + (r_* - \sqrt{r_*^2 - y^2}) \sin \beta(z) \right), \quad (4.7)$$

for $1 - 2r_* \leq z \leq \sqrt{1 - 4r_*^2}$ and $|y| < r_*/2$. Through a calculation similar to the one yielding the formula (4.5), we find that on this portion of $\partial\Omega$ one has

$$\nu_{\partial\Omega}^{(2)} = -y \frac{[-b(y)\beta'(z) + \cos \beta(z) + g'(z) \sin \beta(z)]}{\sqrt{r_*^2 - y^2}},$$

where $b(y) := -r_* + \sqrt{r_*^2 - y^2}$. Since $b(y) < 0$, $\beta'(z) > 0$, $g'(z) > 0$ and $\beta \in [0, \pi/2)$, we see that the expression in brackets is positive and so the conditions in (2.4) again hold.

We observe that, in particular, for the value $z = 1 - 2r_*$, one has $\beta(1 - 2r_*) = 0$ and consequently,

$$G(y, 1 - 2r_*) = (\sqrt{r_*^2 - y^2}, y, 1 - 2r_*)$$

Comparing this with (4.6) we see that indeed the trough glues on to the hemisphere precisely along the arc $x^2 + y^2 = r_*^2$, $z = 1 - 2r_*$ for $|y| < r_*/2$.

This completes the description of how one can construct one trough connecting to a hemispherical valley. This trough is then rotated by $2\pi/3$ in the symmetric case or by the requisite angles in the general case to obtain three troughs terminating at a valley and the whole procedure is repeated to construct all four valleys and six troughs. The rest of the surface constituting $\partial\Omega$ is then just completed in a C^1 manner on the complement of the troughs and valleys.

Acknowledgments: We wish to thank Michael Novack for suggesting we pursue a boundary version of the infiltration lemma, which was critical to the proof of Theorem 3.2. We thank Matthias Weber for his generous time in producing the figures. The research of P.S. was supported by a Simons Collaboration grant 585520 and an NSF grant DMS 2106516.

REFERENCES

- [AF11] Nicholas D. Alikakos and Giorgio Fusco. Entire solutions to equivariant elliptic systems with variational structure. *Arch. Ration. Mech. Anal.*, 202(2):567–597, 2011.

- [AFS18] Nicholas D. Alikakos, Giorgio Fusco, and Panayotis Smyrnelis. *Elliptic systems of phase transition type*, volume 91 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser/Springer, Cham, 2018.
- [AG23] Nicholas D. Alikakos and Zhiyuan Geng. On the triple junction problem with general surface tension coefficients. *arXiv:2304.13106*, pages Paper No. 24, 58, 2023.
- [AG24] Nicholas D. Alikakos and Zhiyuan Geng. On the triple junction problem without symmetry hypotheses. *Arch. Ration. Mech. Anal.*, 248(2):Paper No. 24, 58, 2024.
- [Ali12] Nicholas D. Alikakos. A new proof for the existence of an equivariant entire solution connecting the minima of the potential for the system $\Delta u - W_u(u) = 0$. *Comm. Partial Differential Equations*, 37(12):2093–2115, 2012.
- [Bal90] Sisto Baldo. Minimal interface criterion for phase transitions in mixtures of Cahn-Hilliard fluids. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 7(2):67–90, 1990.
- [EG15] Lawrence C. Evans and Ronald F. Gariepy. *Measure theory and fine properties of functions*. Textbooks in Mathematics. CRC Press, Boca Raton, FL, revised edition, 2015.
- [FT89] Irene Fonseca and Luc Tartar. The gradient theory of phase transitions for systems with two potential wells. *Proc. Roy. Soc. Edinburgh Sect. A*, 111(1-2):89–102, 1989.
- [Gen25] Zhiyuan Geng. Uniqueness of the blow-down limit for a triple junction problem. *Comm. Pure Appl. Math.*, 78(2):500–534, 2025.
- [Giu84] Enrico Giusti. *Minimal surfaces and functions of bounded variation*, volume 80 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1984.
- [GS08] Changfeng Gui and Michelle Schatzman. Symmetric quadruple phase transitions. *Indiana Univ. Math. J.*, 57(2):781–836, 2008.
- [KS89] Robert V. Kohn and Peter Sternberg. Local minimisers and singular perturbations. *Proc. Roy. Soc. Edinburgh Sect. A*, 111(1-2):69–84, 1989.
- [LM94] Gary Lawlor and Frank Morgan. Paired calibrations applied to soap films, immiscible fluids, and surfaces or networks minimizing other norms. *Pacific J. Math.*, 166(1):55–83, 1994.
- [Mag12] Francesco Maggi. *Sets of finite perimeter and geometric variational problems*, volume 135 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2012. An introduction to geometric measure theory.
- [Sch35] I. J. Schoenberg. Remarks to Maurice Fréchet’s article “Sur la définition axiomatique d’une classe d’espace distanciés vectoriellement applicable sur l’espace de Hilbert” [MR1503246]. *Ann. of Math. (2)*, 36(3):724–732, 1935.
- [SS24] Étienne Sandier and Peter Sternberg. Allen-Cahn solutions with triple junction structure at infinity. *Comm. Pure Appl. Math.*, 77(11):4163–4211, 2024.
- [Ste88] Peter Sternberg. The effect of a singular perturbation on nonconvex variational problems. *Arch. Rational Mech. Anal.*, 101(3):209–260, 1988.

- [Ste91] Peter Sternberg. Vector-valued local minimizers of nonconvex variational problems. volume 21, pages 799–807. 1991. Current directions in nonlinear partial differential equations (Provo, UT, 1987).
- [SZ94] Peter Sternberg and William P. Zeimer. Local minimisers of a three-phase partition problem with triple junctions. *Proc. Roy. Soc. Edinburgh Sect. A*, 124(6):1059–1073, 1994.
- [Tay76] Jean E. Taylor. The structure of singularities in soap-bubble-like and soap-film-like minimal surfaces. *Ann. of Math. (2)*, 103(3):489–539, 1976.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY BLOOMINGTON, IN 47405, USA.

Email address: `abadim@iu.edu, abhishek.adimu@gmail.com`

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY BLOOMINGTON, IN 47405, USA.

Email address: `sternber@iu.edu`