

Cosmological constraints on two vacuum decay models

Yupeng Yang^a, Yicheng Wang and Xinyi Dai

¹School of Physics and Physical Engineering, Qufu Normal University, Qufu, Shandong, 273165, China

Received: date / Accepted: date

Abstract We constrain two vacuum decay models ($\Lambda(t)$ CDM) proposed by the authors of [1]) utilizing the baryon acoustic oscillations (BAO) data released by the Dark Energy Spectroscopic Instrument (DESI), distance prior from the cosmic microwave background (CMB) observed by the Planck satellite, Hubble rate data obtained via the cosmic chronometers (CC) method and type Ia supernova (SNIa) data. The interaction terms between dark matter and dark energy are defined as $Q = 3\varepsilon H\rho_\Lambda$ for model I and $3\varepsilon aH\rho_\Lambda$ for model II. We find that the decay parameter is constrained to be $\varepsilon = 0.0094^{+0.0037}_{-0.0033}$ for model I and $\varepsilon = 0.0119 \pm 0.0045$ for model II, respectively, indicating a potential interaction between dark matter and dark energy at the 2σ confidence level. The current Hubble parameter values are estimated to be $H_0 = 70.30 \pm 0.67$ for model I and $H_0 = 70.28 \pm 0.64$ for model II. These values of H_0 fall between those derived from the Planck and SH0ES data, suggesting that these two vacuum decay models could provide a potential solution to alleviate the Hubble tension problem.

1 Introduction

In the classical standard model of cosmology, based on different astronomical observations, it is well known that the Universe is primarily composed of baryonic matter, dark matter, and dark energy. Unlike baryonic and dark matter, dark energy drives the accelerating expansion of the Universe and its nature remains unknown. Among the various models proposed to explain dark energy, the most extensively studied is the cosmological constant term Λ , which does not vary with time and has an equation of state parameter $w = p_\Lambda/\rho_\Lambda = -1$. A universe with the cosmologi-

cal constant as dark energy is commonly referred to as the Λ CDM cosmology model, where CDM denotes cold dark matter.

While the cosmological constant term represents the simplest approach to characterizing dark energy, it is not without its limitations and challenges [2]. Based on current observational data, the energy density associated with the cosmological constant term is estimated to be $\rho_\Lambda \sim 10^{-47} \text{ GeV}^4$ [3]. If the cosmological constant originates from vacuum energy, quantum field theory predicts an energy density of $\rho_{\text{vac}} \sim 10^{74} \text{ GeV}^4$, derived from the cutoff of vacuum energy at the Planck scale m_{pl} . This results in a discrepancy of approximately 121 orders of magnitude between the two values. Even when considering the energy scale of quantum chromodynamics, the vacuum energy density is estimated to be $\rho_{\text{vac}} \sim 10^{-3} \text{ GeV}^4$, which is about 44 orders of magnitude higher than the observed value. Another critical issue for the Λ CDM model is the Hubble tension [4,5]. The Hubble constant H_0 can be determined through various astronomical observations. For instance, analyses of type Ia supernova, gravitational lensing, and Cepheid variable stars yield a Hubble constant of $H_0 = 73.17 \pm 0.86 \text{ km s}^{-1} \text{ Mpc}^{-1}$, derived from low-redshift data and largely independent of specific cosmological models [6,7]. Conversely, the Hubble constant from the analysis of cosmic microwave background (CMB) radiation, particularly when combined with baryon acoustic oscillations (BAO) and Planck data, is $67.4 \pm 0.5 \text{ km s}^{-1} \text{ Mpc}^{-1}$, originating from high-redshift measurements and based on the Λ CDM cosmological model [8]. The discrepancy between these two values reaches 5.8σ , showing a significant challenge to the validity of the Λ CDM model.

Many approaches have been proposed to address the Hubble tension [4,5]. One possibility is that there may be systematic issues with low-redshift measurements.

^ae-mail: ypyang@aliyun.com

Another approach involves modifying the Λ CDM universe model. For instance, kinetic dark energy models, where the equation of state parameter evolves over time ($w(z)$), have been extensively explored (see, e.g., Refs. [3] for a review). Recent BAO data released by the DESI collaboration support the idea that dark energy evolves over time [9]. In addition to dynamic dark energy models, interactions between dark matter and dark energy have also been proposed as potential solutions to the Hubble tension problem [10, 11, 12, 13, 14, 15]. Early dark energy models have similarly been suggested as a resolution to this issue [16, 17]. Furthermore, the vacuum decay model, in which the cosmological constant term evolves over time, akin to an interaction between vacuum energy and dark matter, has been introduced to address related challenges [18, 19, 20, 21, 22, 23, 24, 25, 26]. Recently, the authors of Ref. [1] proposed two specific forms of vacuum decay models aimed at resolving the Hubble tension. Using Cosmic Microwave Background (CMB) and H0 data, they constrained these models and found that vacuum decays into dark matter, ruling out the non-interacting model at 6σ confidence level. In this paper, we further investigate these two vacuum decay models using additional data sets, finding a potential interaction between dark matter and dark energy at the 2σ confidence level.

This paper is organized as follows. In Sec. 2, we first briefly review the main contents of two vacuum decay models, then describe the datasets utilized and presents the final limiting results in Sec. 3. The conclusions are given in Sec. 4.

2 The basic properties of vacuum decay model

In this section, we briefly review the main contents of the vacuum decay model, and one can refer to, e.g., Refs. [18, 19, 20, 21, 22, 23, 24, 25] for more details. The vacuum decay model has been proposed by previous works. In the classical cosmology model, i.e. Λ CDM model, Λ does not evolve with time. The key point of the vacuum decay model is that vacuum energy density changes over time $\Lambda(t)$. Within the framework of the standard cosmological model under the Friedmann-Robertson-Walker (FRW) metric, the continuity equation can be formulated as follows [1],

$$\dot{\rho}_{\text{DM}} + 3H\rho_{\text{DM}} = Q \quad (1)$$

$$\dot{\rho}_{\Lambda} = -Q \quad (2)$$

where ρ_{DM} and ρ_{Λ} are the energy density of dark matter and vacuum. Q denotes the interaction between dark matter and vacuum dark energy, and $Q = 0$ for Λ CDM

model. Here, following the Ref. [1], we adopt the interaction forms as,

$$Q = \begin{cases} 3\varepsilon H\rho_{\Lambda} & \text{Model I} \\ 3\varepsilon aH\rho_{\Lambda} & \text{Model II} \end{cases}$$

where ε is a constant and $a = 1/(1+z)$ is the scale factor. Based on these two forms of interaction, the expansion rate of the Universe, $H(z)$, can be expressed as follows, for Model I:

$$\frac{H^2}{H_0^2} = \left(\Omega_m - \frac{\varepsilon\Omega_{\Lambda}}{1-\varepsilon} \right) (1+z)^3 + \frac{\Omega_{\Lambda}(1+z)^{3\varepsilon}}{1-\varepsilon} + \Omega_r(1+z)^4 \quad (3)$$

and for Model II:

$$\frac{H^2}{H_0^2} = \frac{\Omega_m}{a^3} + \frac{\Omega_{\Lambda}}{9\varepsilon^3 a^3} \left[P(\varepsilon) - Q(\varepsilon a)e^{3\varepsilon(1-a)} \right] + \Omega_r(1+z)^4 \quad (4)$$

with $P(x) = 9x^3 + 9x^2 + 6x + 2$ and $Q(x) = 9x^2 + 6x + 2$. $\Omega_i \equiv 8\pi G\rho_i/(3H_0^2)$ is the energy density parameter of different component at present. Note that here we consider flat Λ CDM cosmology which has a relation of $\Omega_m + \Omega_{\Lambda} + \Omega_r = 1$.

3 The datasets and constrains

3.1 Baryon acoustic oscillation

The Dark Energy Spectroscopic Instrument (DESI) have released the first-year data (DR1), which includes six different classed of traces [9]: the Bright Galaxy Sample (BGS, $z_{\text{eff}} = 0.30$), the Luminous Red Galaxy Sample (LRG, $z_{\text{eff}} = 0.51$ and 0.71), the Emission Line Galaxy Sample (ELG, $z_{\text{eff}} = 1.32$), the combined LRG and ELG Sample (LRG+ELG, $z_{\text{eff}} = 0.93$), the Quasar Sample (QSO, $z_{\text{eff}} = 1.49$) and the Lyman- α Forest Sample (Ly α , $z_{\text{eff}} = 2.33$). The released data by the DESI collaboration are in the form of $D_{\text{M,H,V}}/r_d$ and here we fix the value $r_d = 147.09$ Mpc obtained using all CMB and CMB lensing information [8]. In the context of a homogeneous and isotropic cosmology, the transverse comoving distance, $D_{\text{M}}(z)$, can be written as [8],

$$D_{\text{M}}(z) = \frac{c}{H_0} \int_0^z \frac{dz'}{H(z')/H_0}, \quad (5)$$

where c is the light speed and $H(z)$ is the Hubble parameter given by Eqs. (3) and (4). The distance variable is defined as $D_{\text{H}}(z) = c/H(z)$, and then the angle-averaged distance D_V can be written as

$$D_V(z) = [z D_M(z)^2 D_H(z)]^{\frac{1}{3}} \quad (6)$$

The χ^2 for DESI BAO data is

$$\chi_{\text{BAO}}^2 = \sum_i \Delta D_i^T \text{Cov}_{\text{BAO}}^{-1} \Delta D_i, \quad (7)$$

where Cov_{BAO} is non unit covariance matrix for the tracers of LGR,LGR+ELG, ELG and Ly α QSO, and a unit matrix for the tracers of BGS and QSO [27].

3.2 Type Ia supernova

For type Ia supernova (SNIa) observations, one of the main relevant parameters is the luminosity distance $d_L(z)$. For a flat Λ CDM universe, the luminosity distance can be written as

$$d_L(z) = (1+z) \int_0^z \frac{dz'}{H(z')/H_0} \quad (8)$$

The distance modulus $\mu(z)$ is given by

$$\mu(z) = 5\log_{10}d_L(z) + 25 \quad (9)$$

For SNIa data, the Hubble constant is a nuisance parameter and here we use the methods given in, e.g., Refs. [28,29], for calculating χ_{SN}^2 ,

$$\chi_{\text{SN}}^2 = A - \frac{B^2}{C} \quad (10)$$

where

$$\begin{aligned} A &= \sum_i^N \frac{(\mu_{\text{obs}}(z_i) - 5\log_{10}(D_L(z_i)))^2}{\sigma_i^2} \\ B &= \sum_i^N \frac{\mu_{\text{obs}}(z_i) - 5\log_{10}(D_L(z_i))}{\sigma_i^2} \\ C &= \sum_i^N \frac{1}{\sigma_i^2} \end{aligned} \quad (11)$$

where $D_L(z) = H_0/c \times d_L(z)$. We use the Pantheon sample consisting of a total of 1048 SNIa ranging of $0.01 < z < 2.3$ [30] for our constraints.

3.3 Hubble rate from cosmic chronometers

The Hubble rate can be determined utilizing the cosmic chronometer (CC) method [31,32,33]. This approach primarily focuses on measuring the differential age evolution of the Universe, dt , within a redshift interval dz [31]. In practice, this is accomplished by studying carefully selected massive and passive galaxies. The Hubble rate using CC method can be written as [31]

$$H(z) = -\frac{1}{1+z} \frac{\Delta z}{\Delta t} \quad (12)$$

The χ^2 for CC data is

$$\chi_{\text{CC}}^2 = \sum_i \frac{(H(z_i) - H_{\text{obs}}(z_i))^2}{\sigma_i^2}, \quad (13)$$

and we use 31 CC data points given in different Refs. [34, 35, 36, 37, 38, 39, 40, 41] for our constraints.

3.4 Cosmic microwave background

In general, the angular power spectrum data of CMB have been used to constraint the cosmological parameters [8]. For our purpose, instead of using the angular power spectrum data, here we utilize the distance prior, including the shift parameter R , the acoustic scale l_A and the baryon density $\Omega_b h^2$, from the Planck 2018 data to constrain our parameters [8,42]. The shift parameter and acoustic scale can be written as [8,43,44,45]

$$R = \frac{1+z_*}{c} D_A(z_*) \sqrt{\Omega_M H_0^2} \quad (14)$$

$$l_A = (1+z_*) \frac{\pi D_A(z_*)}{r_s(z_*)} \quad (15)$$

here z_* is the redshift at the epoch of photon decoupling and we adopt the approximate form of [46]

$$z_* = 1048[1 + 0.00124(\Omega_b h^2)^{-0.738}][1 + g_1(\Omega_m h^2)^{g_2}] \quad (16)$$

with

$$g_1 = \frac{0.0738(\Omega_b h^2)^{-0.238}}{1 + 39.5(\Omega_b h^2)^{0.763}}, \quad g_2 = \frac{0.560}{1 + 21.1(\Omega_b h^2)^{1.81}} \quad (17)$$

r_s is the comoving sound horizon and can be written as [9]

$$r_s(z) = \frac{c}{H_0} \int_0^{1/(1+z)} \frac{da}{a^2 H(a) \sqrt{3(1 + \frac{3\Omega_b h^2}{4\Omega_\gamma h^2} a)}} \quad (18)$$

where $(\Omega_\gamma h^2)^{-1} = 42000(T_{\text{CMB}}/2.7\text{K})^{-4}$ with $T_{\text{CMB}} = 2.7255\text{K}$. The χ^2 for CMB data is

$$\chi_{\text{CMB}}^2 = \Delta X^T \text{Cov}_{\text{CMB}}^{-1} \Delta X \quad (19)$$

where $\Delta X = X - X^{\text{obs}}$ is a vector with $X^{\text{obs}} = (R, l_A, \Omega_b h^2)$, and we use the values of X^{obs} and covariance matrix $\text{Cov}_{\text{CMB}}^{-1}$ from Planck 2018 [47].

3.5 Constraints on the models

For our purpose, we use the MCMC method to find the best fit values and the posterior distributions of parameters: $\{\varepsilon, \Omega_m, H_0, \Omega_b h^2\}$. Since we will use the DESI BAO data, SNIa data, CC data and CMB data, the total χ^2 can be written as

$$\chi_{\text{total}}^2 = \chi_{\text{BAO}}^2 + \chi_{\text{SN}}^2 + \chi_{\text{CC}}^2 + \chi_{\text{CMB}}^2 \quad (20)$$

The total likelihood function is $L \propto e^{-\chi_{\text{total}}^2/2}$, and we employ the public code `emcee` [48] to perform the MCMC samples. We set uniform prior for the parameters as $\varepsilon \in (-1, 1)$, $\Omega_m \in (0, 1)$, $H_0 \in (50, 90)$ and $\Omega_b h^2 \in (0.0001, 0.1)$. We use the public code `GetDist` [49] to analyze the MCMC chains. The best-fit parameters, along with their corresponding 1σ uncertainties, are summarized in Tab. 1. Fig. 1 shows the one-dimensional marginalized probability distributions and two-dimensional contour plots for the parameters derived from DESI+CMB, DESI+CMB+CC, and DESI+CMB+CC+SNIa datasets, for both Model I and Model II.

We have found that the best-fit values of the decay parameter are $\varepsilon = 0.0094^{+0.0037}_{-0.0033}$ (1σ) for model I and $\varepsilon = 0.0119 \pm 0.0045$ (1σ) for model II, respectively. Compared to the results presented in previous work [1], our best-fit values for the decay parameter are smaller. Furthermore, we find that the noninteracting model is excluded at 2.8σ (for model I) and 2.6σ (for model II) confidence levels, which contracts with the 7.3σ (for model I) and 6σ (for model II) confidence levels reported in Refs. [1]. Additionally, we have estimated the current Hubble constant to be $H_0 = 70.30 \text{ kms}^{-1}\text{Mpc}^{-1}$ (for model I) and $H_0 = 70.28 \text{ kms}^{-1}\text{Mpc}^{-1}$ (for model II), which are both lower than the values of $H_0 = 73.1 \text{ kms}^{-1}\text{Mpc}^{-1}$ reported in Refs. [1].

4 Conclusions

We have constrained two vacuum decay models, $\Lambda(t)\text{CDM}$, utilizing datasets from DESI BAO, CMB, CC and SNIa. The interaction forms of dark energy and dark matter

are defined as $Q = 3\varepsilon H\rho_A$ for model I and $3\varepsilon aH\rho_A$ for model II. We have found that the best-fit values of the decay parameter are $\varepsilon = 0.0094^{+0.0037}_{-0.0033}$ (1σ) for model I and $\varepsilon = 0.0119 \pm 0.0045$ (1σ) for model II, respectively. This indicates that the noninteracting model is excluded at 2.8σ (for model I) and 2.6σ (for model II) confidence levels. We have estimated the current Hubble constant to be $H_0 = 70.30 \text{ kms}^{-1}\text{Mpc}^{-1}$ (for model I) and $H_0 = 70.28 \text{ kms}^{-1}\text{Mpc}^{-1}$ (for model II), suggesting that these two vacuum decay models could provide a potential solution to alleviate the Hubble tension problem.

5 Acknowledgements

We thank Fengquan Wu(NAOC), Xin Zhang(NEU), Yan Gong(NAOC), Lei Feng(PMO), Shuangxi Yi(QFNU), Yankun Qu(QFNU) and Fayin Wang(NJU) for their helpful suggestions and comments. This work is supported by the Shandong Provincial Natural Science Foundation (Grant Nos. ZR2021MA021).

References

1. L.S. Brito, J.F. Jesus, A.A. Escobal, S.H. Pereira, Can decaying vacuum solve the H_0 Tension? (2024). [arXiv:2412.06756](https://arxiv.org/abs/2412.06756) [astro-ph.CO]
2. L. Perivolaropoulos, F. Skara, Challenges for Λ CDM: An update. *New Astron. Rev.* **95**, 101659 (2022). <https://doi.org/10.1016/j.newar.2022.101659>. [arXiv:2105.05208](https://arxiv.org/abs/2105.05208) [astro-ph.CO]
3. E.J. Copeland, M. Sami, S. Tsujikawa, Dynamics of dark energy. *Int. J. Mod. Phys. D* **15**, 1753–1936 (2006). <https://doi.org/10.1142/S021827180600942X>. [arXiv:hep-th/0603057](https://arxiv.org/abs/hep-th/0603057)
4. E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri, D.F. Mota, A.G. Riess, J. Silk, In the realm of the Hubble tension—a review of solutions. *Class. Quant. Grav.* **38**(15), 153001 (2021). <https://doi.org/10.1088/1361-6382/ac086d>. [arXiv:2103.01183](https://arxiv.org/abs/2103.01183) [astro-ph.CO]
5. J.P. Hu, F.Y. Wang, Hubble Tension: The Evidence of New Physics. *Universe* **9**(2), 94 (2023). <https://doi.org/10.3390/universe9020094>. [arXiv:2302.05709](https://arxiv.org/abs/2302.05709) [astro-ph.CO]
6. A.G. Riess, L. Macri, S. Casertano, M. Sosey, H. Lampeitl, H.C. Ferguson, A.V. Filippenko, S.W. Jha, W. Li, R. Chornock, D. Sarkar, A Redetermination of the Hubble Constant with the Hubble Space Telescope from a Differential Distance Ladder. *apj* **699**(1), 539–563 (2009). <https://doi.org/10.1088/0004-637X/699/1/539>. [arXiv:0905.0695](https://arxiv.org/abs/0905.0695) [astro-ph.CO]
7. L. Breuval, A.G. Riess, S. Casertano, W. Yuan, L.M. Macri, M. Romaniello, Y.S. Murakami, D. Scolnic, G.S. Anand, I. Soszyński, Small Magellanic Cloud Cepheids Observed with the Hubble Space Telescope Provide a New Anchor for the SH0ES Distance Ladder. *Astrophys. J.* **973**(1), 30 (2024). <https://doi.org/10.3847/1538-4357/ad630e>. [arXiv:2404.08038](https://arxiv.org/abs/2404.08038) [astro-ph.CO]
8. N. Aghanim, et al., Planck 2018 results. VI. Cosmological parameters. *A&A* **641**, A6 (2020). <https://doi.org/10.1051/0004-6361/201833910>. [arXiv:1807.06209](https://arxiv.org/abs/1807.06209) [astro-ph.CO]




Fig. 1 The one-dimensional marginalized probability distribution and two-dimensional contour plots of parameters for model I (left) and model II (right) at the 68% and 95% confidence levels, derived using DESI+CMB, DESI+CMB+CC, and DESI+CMB+CC+SNIa data.

Table 1 Constraints on the parameters of two vacuum decay models with 1σ uncertainties using various observational data combinations: DESI+CMB, DESI+CMB+CC and DESI+CMB+CC+SNIa.

Model/Dataset	ε	Ω_m	H_0	$\Omega_b h^2$
Model I				
DESI+CMB	$0.0142^{+0.0039}_{-0.0032}$	$0.2667^{+0.0087}_{-0.0099}$	71.69 ± 0.92	0.02255 ± 0.00015
DESI+CMB+CC	$0.0131^{+0.0041}_{-0.0034}$	0.2704 ± 0.0093	71.33 ± 0.89	0.02256 ± 0.00015
DESI+CMB+CC+SNIa	$0.0094^{+0.0037}_{-0.0033}$	0.2813 ± 0.0074	70.30 ± 0.67	0.02259 ± 0.00014
Model II				
DESI+CMB	$0.0178^{+0.0052}_{-0.0041}$	$0.2680^{+0.0092}_{-0.0100}$	71.60 ± 0.95	0.02256 ± 0.00015
DESI+CMB+CC	$0.0163^{+0.0051}_{-0.0042}$	0.2719 ± 0.0091	71.22 ± 0.87	0.02256 ± 0.00015
DESI+CMB+CC+SNIa	0.0119 ± 0.0045	0.2819 ± 0.0071	70.28 ± 0.64	0.02259 ± 0.00015

9. A.G. Adame, et al., DESI 2024 VI: Cosmological Constraints from the Measurements of Baryon Acoustic Oscillations (2024). [arXiv:2404.03002](https://arxiv.org/abs/2404.03002) [astro-ph.CO]
10. B. Wang, E. Abdalla, F. Atrio-Barandela, D. Pavon, Dark Matter and Dark Energy Interactions: Theoretical Challenges, Cosmological Implications and Observational Signatures. *Rept. Prog. Phys.* **79**(9), 096901 (2016). <https://doi.org/10.1088/0034-4885/79/9/096901>. [arXiv:1603.08299](https://arxiv.org/abs/1603.08299) [astro-ph.CO]
11. K. Naidoo, M. Jaber, W.A. Hellwing, M. Bilicki, Dark matter solution to the H_0 and S8 tensions, and the integrated Sachs-Wolfe void anomaly. *Phys. Rev. D* **109**(8), 083511 (2024). <https://doi.org/10.1103/PhysRevD.109.083511>. [arXiv:2209.08102](https://arxiv.org/abs/2209.08102) [astro-ph.CO]
12. A. Poutsidou, T. Tram, Reconciling cmb and structure growth measurements with dark energy interactions. *Phys. Rev. D* **94**, 043518 (2016). <https://doi.org/10.1103/PhysRevD.94.043518>. URL <https://link.aps.org/doi/10.1103/PhysRevD.94.043518>
13. E. Di Valentino, A. Melchiorri, O. Mena, Can interacting dark energy solve the H_0 tension? *Phys. Rev. D* **96**, 043503 (2017). <https://doi.org/10.1103/PhysRevD.96.043503>. URL <https://link.aps.org/doi/10.1103/PhysRevD.96.043503>
14. S. Kumar, R.C. Nunes, Echo of interactions in the dark sector. *Phys. Rev. D* **96**, 103511 (2017). <https://doi.org/10.1103/PhysRevD.96.103511>. URL <https://link.aps.org/doi/10.1103/PhysRevD.96.103511>
15. W. Yang, A. Mukherjee, E. Di Valentino, S. Pan, Interacting dark energy with time varying equation of state and the H_0 tension. *Phys. Rev. D* **98**, 123527 (2018). <https://doi.org/10.1103/PhysRevD.98.123527>. URL <https://link.aps.org/doi/10.1103/PhysRevD.98.123527>

- link.aps.org/doi/10.1103/PhysRevD.98.123527
16. M. Kamionkowski, A.G. Riess, The Hubble Tension and Early Dark Energy. *Ann. Rev. Nucl. Part. Sci.* **73**, 153–180 (2023). <https://doi.org/10.1146/annurev-nucl-111422-024107>. arXiv:2211.04492 [astro-ph.CO]
 17. V. Poulin, T.L. Smith, T. Karwal, M. Kamionkowski, Early Dark Energy Can Resolve The Hubble Tension. *Phys. Rev. Lett.* **122**(22), 221301 (2019). <https://doi.org/10.1103/PhysRevLett.122.221301>. arXiv:1811.04083 [astro-ph.CO]
 18. M. Koussour, N. Myrzakulov, J. Rayimbaev, Cosmological constraints on time-varying cosmological terms: A study of FLRW universe models with $\Lambda(t)$ CDM cosmology. *Adv. Space Res.* **74**, 1343–1351 (2024). <https://doi.org/10.1016/j.asr.2024.04.045>. arXiv:2404.15982 [astro-ph.CO]
 19. A.M. Öztaş, The effects of a varying cosmological constant on the particle horizon. *Mon. Not. Roy. Astron. Soc.* **481**(2), 2228–2234 (2018). <https://doi.org/10.1093/mnras/sty2375>
 20. R.G. Vishwakarma, Consequences on variable Lambda models from distant type Ia supernovae and compact radio sources. *Class. Quant. Grav.* **18**, 1159–1172 (2001). <https://doi.org/10.1088/0264-9381/18/7/301>. arXiv:astro-ph/0012492
 21. J.M. Overduin, F.I. Cooperstock, Evolution of the scale factor with a variable cosmological term. *Phys. Rev. D* **58**, 043506 (1998). <https://doi.org/10.1103/PhysRevD.58.043506>. arXiv:astro-ph/9805260
 22. H. Azri, A. Bounames, Cosmological Consequences of a Variable Cosmological Constant Model. *Int. J. Mod. Phys. D* **26**(7), 1750060 (2017). <https://doi.org/10.1142/S0218271817500602>. arXiv:1412.7567 [gr-qc]
 23. H. Azri, A. Bounames, Geometrical Origin of the Cosmological Constant. *Gen. Rel. Grav.* **44**, 2547–2561 (2012). <https://doi.org/10.1007/s10714-012-1413-9>. arXiv:1007.1948 [gr-qc]
 24. M. Szydlowski, Cosmological model with decaying vacuum energy from quantum mechanics. *Phys. Rev. D* **91**(12), 123538 (2015). <https://doi.org/10.1103/PhysRevD.91.123538>. arXiv:1502.04737 [astro-ph.CO]
 25. M. Bruni, R. Maier, D. Wands, Nonsingular cosmology from an interacting vacuum. *Phys. Rev. D* **105**(6), 063532 (2022). <https://doi.org/10.1103/PhysRevD.105.063532>. arXiv:2111.01765 [gr-qc]
 26. H.A.P. Macedo, L.S. Brito, J.F. Jesus, M.E.S. Alves, Cosmological constraints on $\Lambda(t)$ CDM models. *Eur. Phys. J. C* **83**(12), 1144 (2023). <https://doi.org/10.1140/epjc/s10052-023-12321-0>. arXiv:2305.18591 [astro-ph.CO]
 27. J.X. Li, S. Wang, A comprehensive numerical study on four categories of holographic dark energy models (2024). arXiv:2412.09064 [astro-ph.CO]
 28. Y. Gong, X. Chen, Two Component Model of Dark Energy. *Phys. Rev. D* **76**, 123007 (2007). <https://doi.org/10.1103/PhysRevD.76.123007>. arXiv:0708.2977 [astro-ph]
 29. S. Nesseris, L. Perivolaropoulos, Comparison of the legacy and gold type ia supernovae dataset constraints on dark energy models. *Phys. Rev. D* **72**, 123519 (2005). <https://doi.org/10.1103/PhysRevD.72.123519>. URL <https://link.aps.org/doi/10.1103/PhysRevD.72.123519>
 30. D.M. Scolnic, et al., The Complete Light-curve Sample of Spectroscopically Confirmed SNe Ia from Pan-STARRS1 and Cosmological Constraints from the Combined Pantheon Sample. *Astrophys. J.* **859**(2), 101 (2018). <https://doi.org/10.3847/1538-4357/aab9bb>. arXiv:1710.00845 [astro-ph.CO]
 31. R. Jimenez, A. Loeb, Constraining cosmological parameters based on relative galaxy ages. *Astrophys. J.* **573**, 37–42 (2002). <https://doi.org/10.1086/340549>. arXiv:astro-ph/0106145
 32. M. Moresco, et al., Unveiling the Universe with emerging cosmological probes. *Living Rev. Rel.* **25**(1), 6 (2022). <https://doi.org/10.1007/s41114-022-00040-z>. arXiv:2201.07241 [astro-ph.CO]
 33. R. Jimenez, M. Moresco, L. Verde, B.D. Wandelt, Cosmic chronometers with photometry: a new path to $H(z)$. *JCAP* **11**, 047 (2023). <https://doi.org/10.1088/1475-7516/2023/11/047>. arXiv:2306.11425 [astro-ph.CO]
 34. E.K. Li, M. Du, Z.H. Zhou, H. Zhang, L. Xu, Testing the effect of H_0 on $f\sigma_8$ tension using a Gaussian process method. *Mon. Not. Roy. Astron. Soc.* **501**(3), 4452–4463 (2021). <https://doi.org/10.1093/mnras/staa3894>. arXiv:1911.12076 [astro-ph.CO]
 35. D. Stern, R. Jimenez, L. Verde, M. Kamionkowski, S.A. Stanford, Cosmic chronometers: constraining the equation of state of dark energy. I: $H(z)$ measurements. *jcap* **2010**(2), 008 (2010). <https://doi.org/10.1088/1475-7516/2010/02/008>. arXiv:0907.3149 [astro-ph.CO]
 36. M. Moresco, A. Cimatti, R. Jimenez, L. Pozzetti, G. Zamorani, M. Bolzonella, J. Dunlop, F. Lamareille, M. Mignoli, H. Pearce, P. Rosati, D. Stern, L. Verde, E. Zucca, C.M. Carollo, T. Contini, J.P. Kneib, O. Le Fèvre, S.J. Lilly, V. Mainieri, A. Renzini, M. Scoggio, I. Balestra, R. Gobat, R. McLure, S. Bardelli, A. Bondi, K. Caputi, O. Cucciati, S. de la Torre, L. de Ravel, P. Franzetti, B. Garilli, A. Iovino, P. Kampczyk, C. Knobel, K. Kovač, J.F. Le Borgne, V. Le Brun, C. Maier, R. Pelló, Y. Peng, E. Perez-Montero, V. Presotto, J.D. Silverman, M. Tanaka, L.A.M. Tasca, L. Tresse, D. Vergani, O. Almaini, L. Barnes, R. Bordoloi, E. Bradshaw, A. Cappi, R. Chuter, M. Cirasuolo, G. Coppa, C. Diener, S. Foucaud, W. Hartley, M. Kamionkowski, A.M. Koekemoer, C. López-Sanjuan, H.J. McCracken, P. Nair, P. Oesch, A. Stanford, N. Weilakala, Improved constraints on the expansion rate of the Universe up to $z \sim 1$ from the spectroscopic evolution of cosmic chronometers. *jcap* **2012**(8), 006 (2012). <https://doi.org/10.1088/1475-7516/2012/08/006>. arXiv:1201.3609 [astro-ph.CO]
 37. M. Moresco, L. Pozzetti, A. Cimatti, R. Jimenez, C. Maraston, L. Verde, D. Thomas, A. Citro, R. Tojeiro, D. Wilkinson, A 6% measurement of the Hubble parameter at $z \sim 0.45$: direct evidence of the epoch of cosmic re-acceleration. *JCAP* **05**, 014 (2016). <https://doi.org/10.1088/1475-7516/2016/05/014>. arXiv:1601.01701 [astro-ph.CO]
 38. M. Moresco, Raising the bar: new constraints on the Hubble parameter with cosmic chronometers at $z \sim 2$. *Mon. Not. Roy. Astron. Soc.* **450**(1), L16–L20 (2015). <https://doi.org/10.1093/mnrasl/slv037>. arXiv:1503.01116 [astro-ph.CO]
 39. A.L. Ratsimbazafy, S.I. Loubser, S.M. Crawford, C.M. Cress, B.A. Bassett, R.C. Nichol, P. Väistönen, Age-dating Luminous Red Galaxies observed with the Southern African Large Telescope. *Mon. Not. Roy. Astron. Soc.* **467**(3), 3239–3254 (2017). <https://doi.org/10.1093/mnras/stx301>. arXiv:1702.00418 [astro-ph.CO]
 40. C. Zhang, H. Zhang, S. Yuan, S. Liu, T.J. Zhang, Y.C. Sun, Four new observational $H(z)$ data from luminous red galaxies in the Sloan Digital Sky Survey data release seven. *Research in Astronomy and Astrophysics* **14**(10), 1221–1233 (2014). <https://doi.org/10.1088/1674-4527/14/10/002>. arXiv:1207.4541 [astro-ph.CO]
 41. H. Singirikonda, S. Desai, Model comparison of Λ CDM vs $R_h = ct$ using cosmic chronometers. *Eur. Phys.*

-
- J. C **80**(8), 694 (2020). <https://doi.org/10.1140/epjc/s10052-020-8289-8>. arXiv:2003.00494 [astro-ph.CO]
42. L. Chen, Q.G. Huang, K. Wang, Distance Priors from Planck Final Release. JCAP **02**, 028 (2019). <https://doi.org/10.1088/1475-7516/2019/02/028>. arXiv:1808.05724 [astro-ph.CO]
43. Y. Liu, R.Y. Guo, J.F. Zhang, X. Zhang, Revisit of constraints on dark energy with Hubble parameter measurements including future redshift drift observations. JCAP **05**, 016 (2019). <https://doi.org/10.1088/1475-7516/2019/05/016>. arXiv:1811.12131 [astro-ph.CO]
44. Y.Y. Xu, X. Zhang, Comparison of dark energy models after Planck 2015. Eur. Phys. J. C **76**(11), 588 (2016). <https://doi.org/10.1140/epjc/s10052-016-4446-5>. arXiv:1607.06262 [astro-ph.CO]
45. L. Feng, T. Lu, A new equation of state for dark energy model. Journal of Cosmology and Astroparticle Physics **2011**(11), 034–034 (2011). <https://doi.org/10.1088/1475-7516/2011/11/034>. URL <http://dx.doi.org/10.1088/1475-7516/2011/11/034>
46. W. Hu, N. Sugiyama, Small scale cosmological perturbations: An Analytic approach. Astrophys. J. **471**, 542–570 (1996). <https://doi.org/10.1086/177989>. arXiv:astro-ph/9510117
47. Z. Zhai, Y. Wang, Robust and model-independent cosmological constraints from distance measurements. JCAP **07**, 005 (2019). <https://doi.org/10.1088/1475-7516/2019/07/005>. arXiv:1811.07425 [astro-ph.CO]
48. D. Foreman-Mackey, D.W. Hogg, D. Lang, J. Goodman, <tt>emcee</tt>: The mcmc hammer. Publications of the Astronomical Society of the Pacific **125**(925), 306–312 (2013). <https://doi.org/10.1086/670067>. URL <http://dx.doi.org/10.1086/670067>
49. A. Lewis, GetDist: a Python package for analysing Monte Carlo samples (2019). URL <https://getdist.readthedocs.io>. arXiv:1910.13970 [astro-ph.IM]