

On the existence and the stability of solutions in nonsmooth vector optimization[†]

Tran Van Nghi · Le Ngoc Kien · Nguyen Van Tuyen

Received: date / Accepted: date

Abstract The paper is devoted to the existence of weak Pareto solutions and the weak sharp minima at infinity property for a general class of constrained nonsmooth vector optimization problems with unbounded constraint set via asymptotic cones and generalized asymptotic functions. Then we show that these conditions are useful for studying the solution stability of nonconvex vector optimization problems with linear perturbation. We also provide some applications for a subclass of nonsmooth robustly quasiconvex vector optimization problems.

Keywords Vector optimization · Existence · Stability · Weak sharp minima · Asymptotic cone · Asymptotic function · Linear perturbation

Mathematics Subject Classification (2000) 90C29 · 49J30 · 90C31 · 90C26 · 49J52

Tran Van Nghi
Department of Mathematics, Hanoi Pedagogical University 2,
Xuan Hoa, Phuc Yen, Vinh Phuc, Vietnam
tranvanngghi@hpu2.edu.vn

Le Ngoc Kien
Faculty of Fundamental Science, Vietnam-Hungary Industrial University,
Xuan Khanh, Son Tay, Hanoi, Vietnam;
Department of Mathematics, Hanoi Pedagogical University 2,
Xuan Hoa, Phuc Yen, Vinh Phuc, Vietnam
lengockien@viu.edu.vn

Nguyen Van Tuyen, Corresponding author
Department of Mathematics, Hanoi Pedagogical University 2,
Xuan Hoa, Phuc Yen, Vinh Phuc, Vietnam
nguyenvantuyen83@hpu2.edu.vn; tuyensp2@yahoo.com

1 Introduction

Vector optimization is an important part of mathematical programming. It is used to denote a type of optimization problems, where two or more objectives are to be minimized over a given set. Vector optimization problems arise naturally in a variety of theoretical and practical problems and have a wide range of applications. These problems are important tools for problems in economics, management, multicriteria design optimization, water resource planning, medicine, mathematical biology, etc; see, for example, [4, 5, 13, 14, 31, 38, 41].

An important and fundamental issue in theory of vector optimization problems is the investigation of existence and stability of solutions; see, for example, [3, 11, 14, 23–25, 30, 31, 38, 39]. In this paper, we consider the following vector optimization problem

$$\text{Min}_{\mathbb{R}_+^m} \{f(x) := (f_1(x), \dots, f_m(x)) : x \in X\}, \quad (\text{VP})$$

where X is a closed and unbounded subset in \mathbb{R}^n and $f_i: \mathbb{R}^n \rightarrow \mathbb{R}$, $i = 1, \dots, m$, are lower semicontinuous functions.

Some authors have studied theorems of Frank–Wolfe type for polynomial vector optimization problems. Kim et al. [32] showed the existence of Pareto solutions of an unconstrained polynomial vector optimization problem under the assumptions that the Palais–Smale-type condition holds and the image of the objective function has a bounded section. For the case where X is a convex semi-algebraic set and f is convex, Lee et al. [37] proved that (VP) has a Pareto solution if and only if $f(X)$ has a nonempty bounded section. Recently, Flores-Bazán et al. [18] introduced two notions of vector asymptotic functions for vector functions with their image space ordered by the nonnegative orthant and established coercivity properties, coercive and noncoercive existence results for solutions of vector optimization problems.

The main difficulty in investigation the existence of (weak) Pareto solutions is due to directions in unbounded constraint set along which the objective function may decrease. Asymptotic tools have proved to be very useful in order to overcome the above difficulty; see, for example, [1, 9, 18, 36, 44, 47]. The idea of the asymptotic approach is based on the asymptotic behavior of sets and functions. When the constraint set is unbounded, asymptotic tools allow us to obtain coercivity properties and existence results for vector optimization problems.

In this paper, we investigate the existence and stability of weak Pareto solutions of constrained nonsmooth vector optimization problems by using the tool of asymptotic analysis. We make the following contributions to vector optimization problems:

- As a first result, we present new results on existence of weak Pareto solutions and the weak sharp minima at infinity property for a general class

of constrained nonsmooth vector optimization problems with unbounded constraint set via asymptotic cones and generalized asymptotic functions.

- We obtain stability results, including: the nonemptiness and compactness, the upper/lower semicontinuity, and the closedness of the weak Pareto solution map of the considered problem under linear perturbations.

- We introduce the concept of robustly quasiconvex for vector-valued functions and apply our previous results to the particular case when the objective function is quasiconvex or α -robustly quasiconvex. By using the q -asymptotic function, we develop sufficient conditions for the mentioned problems under weaker assumptions.

This paper is organized as follows. Section 2 provides the necessary background required for the subsequent development. In Section 3, we investigate the solution existence and the weak sharp minima at infinity. Section 4 presents stability results under a linear perturbation. Section 5 proposes some applications to the particular case when the objective function is quasiconvex or α -robustly quasiconvex. A summary of conclusions is explained in the last section.

2 Preliminaries

Throughout the paper, the considered spaces are finite-dimensional Euclidean with the inner product and the norm being denoted by $\langle \cdot, \cdot \rangle$ and by $\| \cdot \|$, respectively. The open ball centered at the origin with a radius of $\alpha > 0$ and the set of all positive integer numbers are denoted, respectively, by \mathbb{B}_α and \mathbb{N} . The nonnegative orthant of \mathbb{R}^m is defined by

$$\mathbb{R}_+^m := \{x = (x_1, \dots, x_m) \in \mathbb{R}^m : x_i \geq 0, i = 1, \dots, m\}.$$

Let $\varphi: \mathbb{R}^n \rightarrow \overline{\mathbb{R}} := \mathbb{R} \cup \{\pm\infty\}$ be an extended-real-valued function. The *domain* and *epigraph* of φ are defined, respectively, by

$$\begin{aligned} \text{dom } \varphi &:= \{x \in \mathbb{R}^n : \varphi(x) < +\infty\} \quad \text{and} \\ \text{epi } \varphi &:= \{(x, t) \in \text{dom } \varphi \times \mathbb{R} : \varphi(x) \leq t\}. \end{aligned}$$

We say that φ is proper if $\text{dom } \varphi \neq \emptyset$ and $\varphi(x) > -\infty$ for every $x \in \mathbb{R}^n$. We adopt the usual convention $\inf_\emptyset \varphi := +\infty$ and $\sup_\emptyset \varphi := -\infty$.

For a given $\alpha \in \mathbb{R}$, we denote by

$$\text{lev}(\varphi, \alpha) := \{x \in \mathbb{R}^n : \varphi(x) \leq \alpha\}$$

the sublevel set of φ at value λ . For a nonempty subset X in \mathbb{R}^n , the set of all minimum points of φ on X is defined by

$$\text{argmin}_X \varphi := \{x \in X : \varphi(x) \leq \varphi(y) \quad \forall y \in X\}.$$

Definition 2.1 We say that the function φ is:

- (i) *convex* if $\text{epi } \varphi$ is a convex subset of $\mathbb{R}^n \times \mathbb{R}$.
- (ii) *quasiconvex* if $\text{lev}(\varphi, \alpha)$ is a convex subset of \mathbb{R}^n for all $\alpha \in \mathbb{R}$.

It is well-known that the function φ is convex if and only if its domain is convex and for every $x, y \in \text{dom } \varphi$

$$\varphi(\lambda x + (1 - \lambda)y) \leq \lambda\varphi(x) + (1 - \lambda)\varphi(y) \quad \forall \lambda \in [0, 1]$$

and it is quasiconvex if and only if for every $x, y \in \text{dom } \varphi$,

$$\varphi(\lambda x + (1 - \lambda)y) \leq \max\{\varphi(x), \varphi(y)\} \quad \forall \lambda \in [0, 1].$$

Clearly, every convex function is also quasiconvex, but not vice versa. For example, the function $\varphi(x) := \sqrt{|x|}$ for all $x \in \mathbb{R}$ is quasiconvex but it is nonconvex.

In [42, Lemma 1.1], the authors showed that the quasiconvexity is not stable under linear perturbations and hence, they introduced the so-called α -robustly quasiconvex functions.

Definition 2.2 (see [10, 42]) For $\alpha \geq 0$, a proper function $\varphi : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ is said to be α -robustly quasiconvex if the function $x \mapsto \varphi(x) + \langle u, x \rangle$ is quasiconvex for all $u \in \mathbb{B}_\alpha$.

As shown in [10, p. 1091], every convex function is also α -robustly quasiconvex for all $\alpha \geq 0$, but not vice versa. Furthermore, every local minimum of an α -robustly quasiconvex function is a global one. For a further study we refer the readers to [10, 42, 43].

We now introduce the concept of robustly quasiconvex for vector-valued functions.

Definition 2.3 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $x \mapsto f(x) := (f_1(x), \dots, f_m(x))$ be a mapping and $\alpha \geq 0$. We say that f is:

- (i) \mathbb{R}_+^m -*quasiconvex* if for every $y = (y_1, \dots, y_m) \in \mathbb{R}^m$, the sublevel set

$$\text{lev}(f, y) := \{x \in \mathbb{R}^n : f_i(x) \leq y_i, \quad i = 1, \dots, m\}$$

of f at y is a convex subset of \mathbb{R}^n .

- (ii) \mathbb{R}_+^m - α -*robustly quasiconvex* if the mapping $x \mapsto f(x) + \langle u, x \rangle$ is \mathbb{R}_+^m -quasiconvex for all $u = (u_1, \dots, u_m) \in \mathbb{R}^{m \times n}$ with $\|u_i\| \leq \alpha$, $i = 1, \dots, m$, where

$$\langle u, x \rangle := (\langle u_1, x \rangle, \dots, \langle u_m, x \rangle).$$

It is easy to see that f is \mathbb{R}_+^m -quasiconvex (resp., \mathbb{R}_+^m - α -robustly quasiconvex) if and only if each component f_i , $i = 1, \dots, m$, of f is quasiconvex (resp., \mathbb{R}_+^m - α -robustly quasiconvex).

Next, we recall some basic definitions and properties of asymptotic cones and functions, which can be found in [9].

Definition 2.4 Let X be a subset of \mathbb{R}^n .

(i) The *asymptotic cone* of X , denoted by X^∞ , is the set defined by

$$X^\infty := \left\{ d \in \mathbb{R}^n : \exists t_k \rightarrow +\infty, \exists x_k \in X, \frac{x_k}{t_k} \rightarrow d \right\} \text{ and } \emptyset^\infty = \emptyset.$$

(ii) The (generalized) *recession cone* of X , denoted by $\text{rec } X$, is the set defined by

$$\text{rec } X := \{d \in \mathbb{R}^n : x + td \in X \ \forall x \in X, \forall t \geq 0\} \text{ and } \text{rec } \emptyset = \emptyset.$$

It follows from [9, Propositions 2.11 and 2.12] that X^∞ is a closed cone and X is bounded if and only if $X^\infty = \{0\}$. While $\text{rec } X$ is a convex cone, $\text{rec } X \subset X^\infty$, and if X is bounded, then $\text{rec } X = \{0\}$. When $X^\infty = \text{rec } X$, then X is called regular. In particular, if X is convex, then it is regular and

$$X^\infty = \left\{ d \in \mathbb{R}^n : x_0 + td \in X \ \forall t \geq 0 \right\} \text{ for any } x_0 \in X,$$

see, for example, [9, Proposition 2.1.5].

Definition 2.5 (see [9]) Let $\varphi: \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ be a proper function. The *asymptotic function* $\varphi^\infty: \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ of φ is the function defined by

$$\varphi^\infty(d) = \inf \left\{ \liminf_{k \rightarrow +\infty} \frac{\varphi(t_k d_k)}{t_k} : t_k \rightarrow +\infty, d_k \rightarrow d \right\} \quad \forall d \in \mathbb{R}^n.$$

By definition, it is easy to see that φ^∞ is lower semicontinuous and positively homogeneous, and the following relation holds $\text{epi } \varphi^\infty = (\text{epi } \varphi)^\infty$. Moreover, when φ is lower semicontinuous and convex, then we have

$$\varphi^\infty(d) = \sup_{t > 0} \frac{\varphi(x_0 + td) - \varphi(x_0)}{t} = \lim_{t \rightarrow +\infty} \frac{\varphi(x_0 + td) - \varphi(x_0)}{t}$$

for all $x_0 \in \text{dom } \varphi$; see [9, Proposition 2.5.2]. Recall that φ is called *lower semicontinuous* (lsc) at some $\bar{x} \in \mathbb{R}^n$ if for every real number $r < \varphi(\bar{x})$ there exists a neighborhood V of \bar{x} such that $r < \varphi(x)$ for all $x \in V$. We say that φ is lsc around \bar{x} if it is lsc at any point of some neighborhood of \bar{x} . If φ is lsc at every $x \in X \subset \mathbb{R}^n$, then f is said to be lsc on X . By definition, it is clear that φ is automatically lsc at \bar{x} when $\varphi(\bar{x}) = -\infty$; when $\varphi(\bar{x}) = +\infty$ the lower semicontinuity of φ means that the values of φ remain as large as required, provided one stays in some small neighborhood of \bar{x} .

It is well-known that, if $\mathcal{K}(\varphi) = \{0\}$, where

$$\mathcal{K}(\varphi) := \{d \in \mathbb{R}^n : \varphi^\infty(d) \leq 0\},$$

then φ is coercive, i.e.,

$$\lim_{\|x\| \rightarrow +\infty} \varphi(x) = +\infty.$$

Furthermore, if φ is convex and lsc, then

$$\mathcal{K}(\varphi) = \{0\} \iff \varphi \text{ is coercive} \iff \text{argmin}_{\mathbb{R}^n} \varphi \text{ is nonempty and compact}$$

see [9, Proposition 3.1.3]. However, when φ is nonconvex, the asymptotic function φ^∞ is not good enough for providing adequate information on the behavior of φ at infinity. For example, let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be the function defined by $\varphi(x) = \sqrt{|x|}$. Then φ is coercive and $\operatorname{argmin}_{\mathbb{R}} \varphi = \{0\}$, but $\mathcal{K}(\varphi) = \mathbb{R}$. Therefore, several authors have proposed different notions for dealing with, specially, quasiconvex functions, see [17, 20, 21, 26, 34].

Definition 2.6 (see [17, 21]) Let $\varphi : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ be a proper function. The q -asymptotic function of φ is the function $\varphi_q^\infty : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ given by

$$\varphi_q^\infty(d) := \sup_{x \in \operatorname{dom} \varphi} \sup_{t > 0} \frac{\varphi(x + td) - \varphi(x)}{t} \quad \forall d \in \mathbb{R}^n.$$

As shown in [17] that $\operatorname{epi} \varphi_q^\infty = \operatorname{rec}(\operatorname{epi} \varphi)$ and if φ is lsc and quasiconvex, then

$$\operatorname{argmin}_{\mathbb{R}^n} f \text{ is nonempty and compact iff } \mathcal{K}_q(\varphi) = \{0\},$$

where

$$\mathcal{K}_q(\varphi) := \{d \in \mathbb{R}^n : \varphi_q^\infty(d) \leq 0\}.$$

It can be shown that φ_q^∞ is always positively homogeneous, convex, and the following relation holds

$$\varphi^\infty(d) \leq \varphi_q^\infty(d) \quad \forall d \in \mathbb{R}^n,$$

see [21]. Furthermore, the above inequality could be strict, see, for example, [34].

For a further study on generalized convexity and asymptotic analysis we refer the reader to [2, 6–10, 12, 15–17, 19–22, 26–29, 33–35, 40, 42, 44–46] and references therein.

We now recall definitions of the upper and lower semicontinuity to set-valued mappings. Let $F : \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ be a set-valued mapping and $\bar{x} \in \mathbb{R}^n$. The *Painlevé-Kuratowski outer/upper limit* of F at \bar{x} is defined by

$$\operatorname{Lim sup}_{x \rightarrow \bar{x}} F(x) := \left\{ y \in \mathbb{R}^m : \exists x_k \rightarrow \bar{x}, y_k \rightarrow y \text{ with } y_k \in F(x_k), \forall k = 1, 2, \dots \right\}.$$

Definition 2.7 (see [9]) We say that:

- (i) F is *upper semicontinuous* (usc henceforth) at \bar{x} if, for any open set $V \subset \mathbb{R}^m$ such that $F(\bar{x}) \subset V$ there exists a neighborhood U of \bar{x} in X such that $F(x) \subset V$ for all $x \in U$.
- (ii) F is *lower semicontinuous* (lsc) at \bar{x} if $F(\bar{x}) \neq \emptyset$ and if, for any open set $V \subset \mathbb{R}^m$ such that $F(\bar{x}) \cap V \neq \emptyset$ there exists a neighborhood U of \bar{x} such that $F(x) \cap V \neq \emptyset$ for all $x \in U$.
- (iii) F is *continuous* at \bar{x} if it is both usc and lsc at this point.
- (iv) F is *closed* at \bar{x} if for any sequences $x_k \in \mathbb{R}^n$ and $y_k \in \mathbb{R}^m$ one has

$$x_k \rightarrow \bar{x}, y_k \rightarrow \bar{y}, y_k \in F(x_k) \Rightarrow \bar{y} \in F(\bar{x}).$$

- (v) F is *locally bounded* at \bar{x} if there exists a neighborhood U of \bar{x} such that $\cup_{x \in U} F(x)$ is bounded.

Remark 2.1 (i) By definition, it is easy to see that F is closed at \bar{x} if and only if the following condition holds

$$\text{Lim sup}_{x \rightarrow \bar{x}} F(x) \subset F(\bar{x}).$$

- (ii) By [9, Theorem 1.4.1], if F is usc at \bar{x} and $F(\bar{x})$ is closed, then F is closed at \bar{x} . Conversely, if F is closed and locally bounded at \bar{x} , then F is usc at that point. We note here that the condition of locally boundedness is essential, i.e., if F is closed at \bar{x} but not locally bounded at this point, then F may not be usc at \bar{x} . For example, let $F: \mathbb{R} \rightrightarrows \mathbb{R}$ be defined by

$$F(x) = \begin{cases} \{\frac{1}{x}\} & \text{if } x \neq 0, \\ \{0\} & \text{if } x = 0. \end{cases}$$

Then, it is easy to see that F is closed at $\bar{x} = 0$ but F is not usc at this point.

3 The solution existence and the weak sharp minima at infinity

In this section, we consider the following vector optimization problem:

$$\text{Min}_{\mathbb{R}_+^m} \{f(x) := (f_1(x), \dots, f_m(x)) : x \in X\}, \quad (\text{VP})$$

where X is a closed and unbounded subset in \mathbb{R}^n , and $f_i: \mathbb{R}^n \rightarrow \mathbb{R}$, $i \in I$ with $I := \{1, \dots, m\}$, are lsc functions. The set

$$\Delta := \left\{ \lambda \in \mathbb{R}_+^m : \sum_{i=1}^m \lambda_i = 1 \right\}$$

is called the unit simplex of \mathbb{R}^m . For each $\lambda \in \Delta$, we consider the following scalarization problem

$$\min \left\{ \varphi_\lambda(x) := \langle \lambda, f(x) \rangle : x \in X \right\}. \quad (\text{VP}_\lambda)$$

The solution set of (VP_λ) is denoted by $S(\lambda)$.

Definition 3.1 Let $\bar{x} \in X$. We say that:

- (i) \bar{x} is a *weak Pareto solution* of (VP) if

$$f(X) \cap (f(\bar{x}) - \text{int } \mathbb{R}_+^m) = \emptyset.$$

The set of all weak Pareto solutions of (VP) is denoted by $\text{Sol}^w(\text{VP})$.

(ii) \bar{x} is a *Pareto solution* of (VP) if

$$f(X) \cap (f(\bar{x}) - \mathbb{R}_+^m \setminus \{0\}) = \emptyset.$$

The set of all Pareto solutions of (VP) is denoted by $\text{Sol}(\text{VP})$.

Remark 3.1 By definition, it is clear that

$$\text{Sol}(\text{VP}) \subset \text{Sol}^w(\text{VP}).$$

The following result is well-known but we give a proof here for the sake of the reader.

Proposition 3.1 *The following inclusion holds true*

$$\bigcup_{\lambda \in \Lambda} S(\lambda) \subset \text{Sol}^w(\text{VP}).$$

Proof Fix any $\lambda \in \Lambda$ and let $\bar{x} \in S(\lambda)$. We need to show that $\bar{x} \in \text{Sol}^w(\text{VP})$. If otherwise, then there exists $x_0 \in X$ such that

$$f(x_0) \in f(\bar{x}) - \text{int } \mathbb{R}_+^m,$$

or, equivalently, $f_i(x_0) < f_i(\bar{x})$ for all $i \in I$. This and the fact that $\lambda \in \Lambda$ imply that

$$\sum_{i \in I} \lambda_i f_i(x_0) < \sum_{i \in I} \lambda_i f_i(\bar{x}),$$

contrary to $\bar{x} \in S(\lambda)$. The proof is complete. \square

Theorem 3.1 (The coercivity and the weak sharp minima at infinity)

Assume that

$$X^\infty \cap \left(\bigcup_{i \in I} \mathcal{K}(f_i) \right) = \{0\}, \quad (1)$$

where $\mathcal{K}(f_i) := \{d \in \mathbb{R}^n : f_i^\infty(d) \leq 0\}$, $i \in I$. Then the following statements hold:

- (a) For each $i \in I$, $\text{argmin}_X f_i(x)$ is nonempty and compact. Consequently, f is bounded from below on X .
- (b) For each $\lambda \in \Lambda$, $S(\lambda)$ is nonempty and compact. Consequently, $\text{Sol}^w(\text{VP})$ is nonempty.
- (c) f is \mathbb{R}_+^m -zero-coercive on X , i.e., there exists $\lambda \in \mathbb{R}_+^m \setminus \{0\}$ such that

$$\lim_{x \xrightarrow{X} \infty} \varphi_\lambda(x) := \langle \lambda, f(x) \rangle = +\infty,$$

where $x \xrightarrow{X} \infty$ means that $\|x\| \rightarrow \infty$ and $x \in X$.

- (d) $\text{Sol}^w(\text{VP})$ is compact.
- (e) For each $\lambda \in \Lambda$, the following weak sharp minima at infinity holds true: there exist constants $c > 0$ and $R > 0$ such that

$$\text{dist}(f(x), f(S(\lambda))) \geq c \text{dist}(x, \text{Sol}^w(\text{VP})) \quad \forall x \in X \setminus \mathbb{B}_R.$$

(f) If in addition, f is continuous, then the following weak sharp minima at infinity holds true, i.e., there exist constants $c > 0$ and $R > 0$ such that

$$\text{dist}(f(x), f(\text{Sol}^w(\text{VP}))) \geq c \text{dist}(x, \text{Sol}^w(\text{VP})) \quad \forall x \in X \setminus \mathbb{B}_R.$$

Proof (a): By (1), we have $X^\infty \cap \mathcal{K}(f_i) = \{0\}$ for all $i \in I$. Hence, it follows from [44, Theorem 4.2.1] that $\text{argmin}_X f_i(x)$ is nonempty and compact for every $i \in I$. This implies that f is bounded from below on X .

(b): Fix any $\lambda \in \Lambda$ and let φ_λ be the function defined by

$$\varphi_\lambda(x) := \sum_{i \in I} \lambda_i f_i(x) \quad \forall x \in \mathbb{R}^n.$$

By (1), $f_i^\infty(d) > 0$ for all $d \in X^\infty \setminus \{0\}$. This and [9, Proposition 2.6.1(b)] imply that

$$(\varphi_\lambda)^\infty(d) \geq \sum_{i \in I} \lambda_i (f_i)^\infty(d) > 0 \quad \forall d \in X^\infty \setminus \{0\}.$$

Consequently, $X^\infty \cap \mathcal{K}(\varphi_\lambda) = \{0\}$. Thus, for each $\lambda \in \Lambda$, $S(\lambda)$ is nonempty and compact due to [44, Theorem 4.2.1].

(c): Fix any $\lambda \in \Lambda$. Then we have $X^\infty \cap \mathcal{K}(\varphi_\lambda) = \{0\}$. By [36, Theorem 3.1(c)], φ_λ is coercive on X . This means that f is \mathbb{R}_+^m -zero-coercive on X with respect to λ .

(d): To prove the compactness of $\text{Sol}^w(\text{VP})$, we need to show that $\text{Sol}^w(\text{VP})$ is closed and bounded. Firstly, we claim that $\text{Sol}^w(\text{VP})$ is bounded. Indeed, if otherwise, then there exists a sequence $x_k \in \text{Sol}^w(\text{VP})$ such that $x_k \rightarrow \infty$ as $k \rightarrow \infty$. By passing to subsequences if necessary we may assume that $d_k := \frac{x_k}{\|x_k\|}$ converges to some $d \in \mathbb{R}^n$. Clearly, $d \in X^\infty$ and $\|d\| = 1$. Fix any $x_0 \in X$. Then for each $k \in \mathbb{N}$, due to the fact that $x_k \in \text{Sol}^w(\text{VP})$, there exists $i_k \in I$ such that $f_{i_k}(x_k) \leq f_{i_k}(x_0)$. Since $i_k \in I$ for all $k \in \mathbb{N}$, there exist $i_0 \in I$ such that $i_k = i_0$ for infinite many k . Hence, by passing to a subsequence if necessary, we may assume that $f_{i_0}(x_k) \leq f_{i_0}(x_0)$ for every $k \in \mathbb{N}$. Put $t_k := \|x_k\|$. We have

$$\frac{f_{i_0}(t_k d_k)}{t_k} \leq \frac{f_{i_0}(x_0)}{t_k}.$$

This implies

$$\liminf_{k \rightarrow +\infty} \frac{f_{i_0}(t_k d_k)}{t_k} \leq \liminf_{k \rightarrow +\infty} \frac{f_{i_0}(x_0)}{t_k} = 0.$$

It follows that $f_{i_0}^\infty(d) \leq 0$. This means that $d \in \mathcal{K}(f_{i_0})$. Hence

$$d \in X^\infty \cap \left(\bigcup_{i=1}^m \mathcal{K}(f_i) \right) \quad \text{and} \quad \|d\| = 1,$$

contrary to (1). Thus $\text{Sol}^w(\text{VP})$ is bounded. We now show that $\text{Sol}^w(\text{VP})$ is closed. Assume that x_k is a sequence in $\text{Sol}^w(\text{VP})$ such that x_k converges to

some \bar{x} . We claim that $\bar{x} \in \text{Sol}^w(\text{VP})$ and so $\text{Sol}^w(\text{VP})$ is closed. If otherwise, $\bar{x} \notin \text{Sol}^w(\text{VP})$. Hence, there exists $x_0 \in X$ such that

$$f_i(x_0) < f_i(\bar{x}) \quad \forall i \in I. \quad (2)$$

It follows from the lower semicontinuity of f_i at \bar{x} that

$$f_i(\bar{x}) \leq \liminf_{k \rightarrow \infty} f_i(x_k) \quad \forall i \in I.$$

This and (2) give us that there exists $k \in \mathbb{N}$ large enough such that

$$f_i(x_0) < f_i(x_k) \quad \forall i \in I,$$

contrary to the fact that $x_k \in \text{Sol}^w(\text{VP})$. Hence $\text{Sol}^w(\text{VP})$ is closed, as required.

(e) Let any $\lambda \in \Lambda$, then we have $X^\infty \cap \mathcal{K}(\varphi_\lambda) = \{0\}$. By [36, Theorem 3.1(b)], φ_λ satisfies the weak sharp minima at infinity. This means that there exist $c > 0$ and $R > 0$ such that

$$c \text{dist}(x, S(\lambda)) \leq \varphi_\lambda(x) - \varphi_\lambda^* \quad \forall x \in X \setminus \mathbb{B}_R,$$

where $\varphi_\lambda^* := \min_{x \in X} \varphi_\lambda(x)$. For each $y \in S(\lambda)$, one has

$$\begin{aligned} \varphi_\lambda(x) - \varphi_\lambda^* &= \sum_{i \in I} \lambda_i [f_i(x) - f_i(y)] \\ &\leq \sum_{i \in I} \lambda_i [\max_{j \in I} |f_j(x) - f_j(y)|] \\ &= \max_{j \in I} |f_j(x) - f_j(y)| \\ &\leq \|f(x) - f(y)\|. \end{aligned}$$

Hence

$$c \text{dist}(x, S(\lambda)) \leq \text{dist}(f(x), S(\lambda)) \quad \forall x \in X \setminus \mathbb{B}_R.$$

Clearly, $\text{dist}(x, \text{Sol}^w(\text{VP})) \leq \text{dist}(x, S(\lambda))$. Thus

$$c \text{dist}(x, \text{Sol}^w(\text{VP})) \leq \text{dist}(f(x), S(\lambda)) \quad \forall x \in X \setminus \mathbb{B}_R,$$

as required.

(f): By contradiction, assume that there exists a sequence $x_k \in X$ such that $x_k \rightarrow \infty$ and

$$0 \leq \text{dist}(f(x_k), f(\text{Sol}^w(\text{VP}))) < \frac{1}{k} \text{dist}(x_k, \text{Sol}^w(\text{VP})) \quad \forall k \in \mathbb{N}. \quad (3)$$

Since $x_k \rightarrow \infty$, by passing to subsequences if necessary we may assume that $\frac{x_k}{\|x_k\|}$ converges to some $d \in \mathbb{R}^n$. Clearly, $d \in X^\infty$ and $\|d\| = 1$. By (1), $d \notin \mathcal{K}(f_i)$ for all $i \in I$. This means that

$$f_i^\infty(d) = \inf \left\{ \liminf_{k \rightarrow \infty} \frac{f_i(t_k d_k)}{t_k} : t_k \rightarrow \infty, d_k \rightarrow d \right\} > 0 \quad \forall i \in I. \quad (4)$$

For each $k \in \mathbb{N}$, due to the compactness of $\text{Sol}^w(\text{VP})$ and the continuity of f , there exists a $y^k \in \text{Sol}^w(\text{VP})$ such that

$$\text{dist}(f(x_k), f(\text{Sol}^w(\text{VP}))) = \|f(x_k) - f(y^k)\|.$$

This and (3) imply that

$$\max_{i \in I} |f_i(x_k) - f_i(y_k)| \leq \|f(x_k) - f(y_k)\| < \frac{1}{k} \text{dist}(x_k, \text{Sol}^w(\text{VP})) \quad \forall k \in \mathbb{N}. \quad (5)$$

For any $k \in \mathbb{N}$, we have $\max_{i \in I} \{f_i(x_k) - f_i(y_k)\} \geq 0$ due to the fact that $y_k \in \text{Sol}^w(\text{VP})$. Hence, there exists $i_k \in I$ such that

$$\max_{i \in I} \{f_i(x_k) - f_i(y_k)\} = f_{i_k}(x_k) - f_{i_k}(y_k) \geq 0. \quad (6)$$

By the finiteness of I , there exists $i_0 \in I$ such that the inequality (6) holds true for infinite many $k \in \mathbb{N}$. By passing to subsequences if necessary we may assume that

$$f_{i_0}(x_k) - f_{i_0}(y_k) \geq 0 \quad \forall k \in \mathbb{N}.$$

Combining this with (5) we obtain

$$\begin{aligned} f_{i_0}(x_k) - f_{i_0}(y_k) &= |f_{i_0}(x_k) - f_{i_0}(y_k)| \leq \max_{i \in I} |f_i(x_k) - f_i(y_k)| \\ &< \frac{1}{k} \text{dist}(x_k, \text{Sol}^w(\text{VP})) \end{aligned}$$

for all $k \in \mathbb{N}$. Let $z_k \in \text{Sol}^w(\text{VP})$ be satisfied $\|x_k - z_k\| = \text{dist}(x_k, \text{Sol}^w(\text{VP}))$. Then, by the compactness of $\text{Sol}^w(\text{VP})$, we have

$$\|x_k - z_k\| \leq \|x_k\| + \|z_k\| \leq \|x_k\| + M,$$

where $M := \max\{\|z\| : z \in \text{Sol}^w(\text{VP})\}$. Hence,

$$0 \leq f_{i_0}(x_k) - f_{i_0}(y_k) < \frac{1}{k} (\|x_k\| + M) \quad \forall k \in \mathbb{N},$$

or, equivalently,

$$f_{i_0}(y_k) \leq f_{i_0}(x_k) < f_{i_0}(y_k) + \frac{1}{k} (\|x_k\| + M) \quad \forall k \in \mathbb{N}.$$

Hence

$$\frac{f_{i_0}(y_k)}{\|x_k\|} \leq \frac{f_{i_0}(x_k)}{\|x_k\|} < \frac{f_{i_0}(y_k)}{\|x_k\|} + \frac{1}{k} \left(1 + \frac{M}{\|x_k\|}\right) \quad \forall k \in \mathbb{N}. \quad (7)$$

Since $y_k \in \text{Sol}^w(\text{VP})$ and $\text{Sol}^w(\text{VP})$ is compact, without any loss of generality we may assume that the sequence y_k converges to some $\bar{y} \in \text{Sol}^w(\text{VP})$.

Hence $\lim_{k \rightarrow \infty} \frac{f_{i_0}(y_k)}{\|x_k\|} = 0$. Combining this with (7) we obtain

$$\lim_{k \rightarrow \infty} \frac{f_{i_0}(x_k)}{\|x_k\|} = 0.$$

Since $\|x_k\| \rightarrow \infty$ and $\frac{x_k}{\|x_k\|} \rightarrow d$, the above equality implies that $d \in \mathcal{K}(f_{i_0})$, contrary to (4). The proof is complete. \square

4 The solution stability results under linear perturbation

For every $u = (u_1, \dots, u_m) \in \mathbb{R}^{m \times n}$, we define the function $f^u : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by

$$f^u(x) := f(x) - \langle u, x \rangle := (f_1^{u_1}(x), \dots, f_m^{u_m}(x)) \text{ for all } x \in \mathbb{R}^n,$$

where $f_i^{u_i}(x) := f_i(x) - \langle u_i, x \rangle$, $i \in I$. Consider the following perturbed optimization problem

$$\text{Min}_{\mathbb{R}_+^m} \{f^u(x) : x \in X\}, \quad (\text{VP}_u)$$

where u is the parameter of perturbation. The weak Pareto solution set of (VP_u) is denoted by $\text{Sol}^w(u)$. When $u = 0$, one has $\text{Sol}^w(0) = \text{Sol}^w(\text{VP})$.

Solution stability is an interesting and very useful research field in optimization (see [8,9,22,46] among others) in virtue of its applications on concrete applications since, in practice, we are usually finding the solution of the optimization problem via numerical methods.

By a direct computation, one has

$$(f_i^{u_i})^\infty(y) = (f_i)^\infty(y) - \langle u_i, y \rangle, \quad (8)$$

$$(f_i^{u_i})_q^\infty(y) = (f_i)_q^\infty(y) - \langle u_i, y \rangle. \quad (9)$$

Lemma 4.1 *If the condition (1) is satisfied, then there exists $\varepsilon > 0$ such that for all $u = (u_1, \dots, u_m) \in \mathbb{R}^{m \times n}$ satisfying $\|u\| := \max_{i \in I} \|u_i\| < \varepsilon$, the following condition holds*

$$X^\infty \cap \left(\bigcup_{i \in I} \mathcal{K}(f_i^{u_i}) \right) = \{0\}. \quad (10)$$

Proof It follows from (1) that $X^\infty \cap \mathcal{K}(f_i) = \{0\}$ for all $i \in I$. For each $i \in I$, we show that there exists $\varepsilon_i > 0$ such that

$$X^\infty \cap \mathcal{K}(f_i^{u_i}) = \{0\} \quad \forall u_i \in \mathbb{B}_{\varepsilon_i}. \quad (11)$$

Indeed, if otherwise, then for any $k \in \mathbb{N}$, there is $u_i^k \in \mathbb{B}_{\frac{1}{k}}$ such that

$$X^\infty \cap \mathcal{K}(f_i^{u_i^k}) \neq \{0\}.$$

Hence, there exists $d_k \in X^\infty \setminus \{0\}$ such that $(f_i^{u_i^k})^\infty(d_k) \leq 0$. Since X^∞ is a closed cone, by passing a subsequence if necessary, we may assume that $h_k := \frac{d_k}{\|d_k\|}$ converges to some $h \in X^\infty$ with $\|h\| = 1$. For each $k > 0$, by the positive homogeneity of $(f_i^{u_i^k})^\infty$ and the fact that $(f_i^{u_i^k})^\infty(d_k) \leq 0$, one has $(f_i^{u_i^k})^\infty(h_k) \leq 0$. By (8), we have

$$(f_i)^\infty(h_k) - \langle u_i^k, h_k \rangle = (f_i^{u_i^k})^\infty(h_k) \leq 0.$$

Consequently,

$$(f_i)^\infty(h_k) \leq \langle u_i^k, h_k \rangle \quad \forall k \in \mathbb{N}.$$

This and the lower semicontinuity of $(f_i)^\infty$ imply that

$$(f_i)^\infty(h) \leq \liminf_{k \rightarrow \infty} (f_i)^\infty(h_k) \leq \lim_{k \rightarrow \infty} \langle u_i^k, h_k \rangle = 0.$$

Hence $h \in X^\infty \cap \mathcal{K}(f_i)$, which contradicts the fact that $X^\infty \cap \mathcal{K}(f_i) = \{0\}$. Thus (11) holds. Put $\varepsilon := \min\{\varepsilon_i : i \in I\} > 0$. Then we have $X^\infty \cap \mathcal{K}(f_i^{u_i}) = \{0\}$ for all $i \in I$ and $u_i \in \mathbb{R}^n$ satisfying $\|u_i\| < \varepsilon$. Hence, (10) is satisfied for all $u = (u_1, \dots, u_m) \in \mathbb{R}^{m \times n}$ such that $\|u\| < \varepsilon$. \square

Theorem 4.1 *Assume that the condition (1) is satisfied. Then there exists $\varepsilon > 0$ such that for all $u \in \mathbb{R}^{m \times n}$ satisfying $\|u\| < \varepsilon$, the following statements hold:*

- (a) f^u is bounded from below on X .
- (b) f^u is \mathbb{R}_+^m -zero-coercive on X .
- (c) $\text{Sol}^w(u)$ is nonempty and compact.
- (d) $\text{Lim sup}_{u \rightarrow 0} \text{Sol}^w(u) \subset \text{Sol}^w(0)$, i.e., the solution mapping $\text{Sol}^w(\cdot)$ is closed at 0.
- (e) $\text{Sol}^w(\cdot)$ is usc at 0.

Proof By Lemma 4.1, there exists $\varepsilon > 0$ such that (10) holds for all $u \in \mathbb{R}^{m \times n}$ satisfying $\|u\| < \varepsilon$. Hence, conclusions (a), (b), and (c) follow directly from Theorem 3.1.

(d): Let \bar{x} be an arbitrary element belonging to $\text{Lim sup}_{u \rightarrow 0} \text{Sol}^w(u)$. By definition, there exist sequences $u^k := (u_1^k, \dots, u_m^k) \rightarrow 0$ and $x_k \in \text{Sol}^w(u^k)$ such that $x_k \rightarrow \bar{x}$ as $k \rightarrow \infty$. Fix any $x \in X$. For each $k \in \mathbb{N}$, since $x_k \in \text{Sol}^w(u^k)$, there exists $i_k \in I$ such that

$$f_{i_k}^{u^k}(x) - f_{i_k}^{u^k}(x_k) \geq 0.$$

Since $i_k \in I$ for all $k \in \mathbb{N}$, there exists $i_0 \in I$ such that $i_k = i_0$ for infinite many k . Hence, by passing to subsequences if necessary, we may assume that

$$f_{i_0}^{u^k}(x) - f_{i_0}^{u^k}(x_k) \geq 0 \quad \forall k \in \mathbb{N},$$

or, equivalently,

$$f_{i_0}(x_k) - \langle u_{i_0}^k, x_k \rangle \leq f_{i_0}(x) - \langle u_{i_0}^k, x \rangle \quad \forall k \in \mathbb{N}.$$

This and the lower semicontinuity of f_{i_0} imply that

$$\begin{aligned} f_{i_0}(\bar{x}) &\leq \liminf_{k \rightarrow \infty} f_{i_0}(x_k) = \liminf_{k \rightarrow \infty} (f_{i_0}(x_k) - \langle u_{i_0}^k, x_k \rangle) \\ &\leq \liminf_{k \rightarrow \infty} (f_{i_0}(x) - \langle u_{i_0}^k, x \rangle) = f_{i_0}(x). \end{aligned}$$

Hence, for each $x \in X$, there exists $i_0 \in I$ such that $f_{i_0}(x) - f_{i_0}(\bar{x}) \geq 0$. This means that $\bar{x} \in \text{Sol}^w(0)$, as required.

(e): Suppose on the contrary that $\text{Sol}^w(\cdot)$ is not usc at 0. By definition, there exists an open set $U \subset \mathbb{R}^n$, with $\text{Sol}^w(0) \subset U$, such that for every

neighborhood W of zero, there exists $u \in W$ satisfying $\text{Sol}^w(u) \not\subseteq U$. Thus there exist sequences $u^k := (u_1^k, \dots, u_m^k) \rightarrow 0$ and $x_k \in \text{Sol}^w(u_k) \setminus U$ for all $k \in \mathbb{N}$. We consider two case of the sequence x_k as follows.

Case 1. $\{x_k\}$ is bounded. Then, without loss of generality we may assume that $x_k \rightarrow \bar{x}$. By (d), one has $\bar{x} \in \text{Sol}^w(0) \subset U$. However, since $x_k \notin U$ for all k and U is open, we see that $\bar{x} \notin U$, a contradiction.

Case 2. $\{x_k\}$ is unbounded. By passing to a subsequence if necessary, we may assume that $\|x_k\| \rightarrow \infty$ as $k \rightarrow \infty$. Hence, without loss of generality one can assume that $\frac{x_k}{\|x_k\|} \rightarrow d \in X^\infty$ with $\|d\| = 1$. Fix any $x \in X$. By a similar argument as in the proof of (d), there exists $i_0 \in I$ such that

$$f_{i_0}(x_k) - \langle u_{i_0}^k, x_k \rangle \leq f_{i_0}(x) - \langle u_{i_0}^k, x \rangle \quad \forall k \in \mathbb{N}.$$

We obtain that

$$\begin{aligned} (f_{i_0})^\infty(d) &\leq \liminf_{k \rightarrow \infty} \frac{f_{i_0}(x_k)}{\|x_k\|} = \liminf_{k \rightarrow \infty} \frac{f_{i_0}(x_k) - \langle u_{i_0}^k, x_k \rangle}{\|x_k\|} \\ &\leq \liminf_{k \rightarrow \infty} \frac{f_{i_0}(x) - \langle u_{i_0}^k, x \rangle}{\|x_k\|} = 0. \end{aligned}$$

Hence, $d \in X^\infty \cap \mathcal{K}(f_{i_0})$, contrary to (1). The proof is complete. \square

The following example shows that there exists $u_k \rightarrow 0$ as $k \rightarrow \infty$ such that $\text{Sol}^w(u_k) = \emptyset$ if (1) is not satisfied.

Example 4.1 Consider the following vector optimization problem

$$\text{Min}_{\mathbb{R}_+^2} \{f(x) := (f_1(x), f_2(x)) : x \in X\},$$

where $X = \mathbb{R}$, $f_1(x) = 0$, and $f_2(x) = x$ for all $x \in \mathbb{R}$. Then, $\text{Sol}^w(0) = \mathbb{R}$ and

$$X^\infty \cap (\mathcal{K}(f_1) \cup \mathcal{K}(f_2)) = \mathbb{R}.$$

Thus (1) is not satisfied. For each k , let $u_k = (-\frac{1}{k}, \frac{1}{k}) \in \mathbb{R}^2$. Then $u_k \rightarrow 0$ as $k \rightarrow \infty$. Consider the following perturbed problem

$$\text{Min}_{\mathbb{R}_+^2} \left\{ f^{u_k}(x) := \left(f_1^{-\frac{1}{k}}(x) = \frac{1}{k}x, f_2^{\frac{1}{k}}(x) = x - \frac{1}{k}x \right) : x \in X \right\}.$$

By definition, we obtain that $\text{Sol}^w(u_k) = \emptyset$.

The following theorem presents sufficient conditions for the lower semicontinuity of the weak Pareto solution map $\text{Sol}^w(\cdot)$.

Theorem 4.2 *If the following conditions are satisfied:*

- (a) $\text{Sol}^w(0)$ is a singleton,
- (b) $X^\infty \cap (\bigcup_{i \in I} \mathcal{K}(f_i)) = \{0\}$,

then $\text{Sol}^w(\cdot)$ is lsc at 0.

Proof Suppose that (a) and (b) hold. By (a), $\text{Sol}^w(0) = \{\bar{x}\}$ for some $\bar{x} \in X$. Let U be an open neighborhood containing \bar{x} . Then, by (b) and Lemma 4.1, there exists $\varepsilon_1 > 0$ such that for all $u = (u_1, \dots, u_m) \in \mathbb{R}^{m \times n}$ satisfying $\|u\| = \max_{i \in I} \|u_i\| < \varepsilon_1$, the condition (10) holds. It follows from Theorem 4.1(c) that $\text{Sol}^w(u) \neq \emptyset$ for every u with $\|u\| < \varepsilon_1$. Since $\text{Sol}^w(\cdot)$ is usc at 0, by Theorem 4.1(e), there exists $\varepsilon_2 > 0$ such that $\text{Sol}^w(u) \subset U$ for every u satisfying $\|u\| < \varepsilon_2$. By taking $\varepsilon := \min\{\varepsilon_1, \varepsilon_2\} > 0$, we obtain that $\text{Sol}^w(u) \cap U \neq \emptyset$ for all u satisfying $\|u\| < \varepsilon$. Hence $\text{Sol}^w(\cdot)$ is lsc at 0. \square

Theorem 4.3 *Assume that X and f are \mathbb{R}_+^m -convex. If $\text{Sol}^w(\cdot)$ is lsc at 0, then the following statements are equivalent:*

- (a) $\text{Sol}^w(0)$ is bounded.
- (b) $X^\infty \cap \left(\bigcup_{i \in I} \mathcal{K}(f_i)\right) = \{0\}$.

Proof By assumptions, we first claim that

$$X^\infty \cap \left(\bigcap_{i \in I} \mathcal{K}(f_i)\right) = \{0\}.$$

Indeed, if otherwise, then there exists a nonzero vector $d \in X^\infty$ such that $(f_i)^\infty(d) \leq 0$ for all $i \in I$. For each $k \in \mathbb{N}$, let $u_i^k := \frac{1}{k}d$ for all $i \in I$. Then, $u_i^k \rightarrow 0$ as $k \rightarrow \infty$ and $\langle u_i^k, d \rangle > 0$ for all $k \in \mathbb{N}$ for all $i \in I$. Hence,

$$(f_i^{u_i^k})^\infty(d) = (f_i)^\infty(d) - \langle u_i^k, d \rangle < 0 \quad \forall k \in \mathbb{N}.$$

We show that $\text{Sol}^w(u^k)$ is empty for every $k \in \mathbb{N}$ and so which contradicts the lower semicontinuity of $\text{Sol}^w(\cdot)$ at 0 and the fact that $u^k \rightarrow 0$ as $k \rightarrow \infty$. Indeed, if otherwise, then there exists $k_0 \in \mathbb{N}$ such that $\text{Sol}^w(u^{k_0}) \neq \emptyset$. Fix any $x_0 \in \text{Sol}^w(u^{k_0})$. It follows from the convexity of X that $x_0 + td \in X$ for all $t > 0$. By [9, Proposition 2.5.2] and the convexity of f_i , one has

$$(f_i^{u_i^{k_0}})^\infty(d) = \lim_{t \rightarrow \infty} \frac{f_i^{u_i^{k_0}}(x_0 + td) - f_i^{u_i^{k_0}}(x_0)}{t} = \lim_{t \rightarrow \infty} \frac{f_i^{u_i^{k_0}}(x_0 + td)}{t}.$$

Hence,

$$\lim_{t \rightarrow \infty} \frac{f_i^{u_i^{k_0}}(x_0 + td)}{t} = (f_i^{u_i^{k_0}})^\infty(d) < 0.$$

This implies that $\lim_{t \rightarrow \infty} f_i^{u_i^{k_0}}(x_0 + td) = -\infty$ for all $i \in I$. Hence, there exists t large enough such that

$$f^{u^{k_0}}(x_0 + td) \in f^{u^{k_0}}(x_0) - \text{int } \mathbb{R}_+^m,$$

contrary to the fact that $x_0 \in \text{Sol}^w(u^{k_0})$, as required.

Now, by the lower semicontinuity of $\text{Sol}^w(\cdot)$ at 0, we see that $\text{Sol}^w(0)$ is nonempty. Hence the statement (a) is equivalent to the nonemptiness and compactness of $\text{Sol}^w(0)$. The equivalence of (a) and (b) follows immediately from [48, Theorem 3.1]. The proof is complete. \square

The following example not only illustrates Theorem 4.3 but also shows that the lower semicontinuity of $\text{Sol}^w(\cdot)$ does not imply the singletonness of the set $\text{Sol}^w(0)$.

Example 4.2 Consider the following vector optimization problem

$$\text{Min}_{\mathbb{R}_+^2} \{f(x) := (f_1(x), f_2(x)) : x \in X\},$$

where $X = [-1, +\infty)$, $f_1(x) = x$, and $f_2(x) = x^2$ for all $x \in \mathbb{R}$. Then, $\text{Sol}^w(0) = [-1, 0]$ is not a singleton and

$$X^\infty \cap (\mathcal{K}(f_1) \cup \mathcal{K}(f_2)) = \{0\}.$$

For each $u = (u_1, u_2) \in \mathbb{R}^2$, we have the following perturbed problem

$$\text{Min}_{\mathbb{R}_+^2} \{f^u(x) := (f_1^{u_1}(x) = x - u_1x, f_2^{u_2}(x) = x^2 - u_2x) \mid x \in X = [-1, +\infty)\}.$$

We can check that

$$\text{Sol}^w(u) = \left[-1, \frac{u_2}{2(1 - u_1)} \right]$$

and $\text{Sol}^w(\cdot)$ is lsc at 0.

5 The Quasiconvex Case

In this section, we apply our previous results to the particular case when the objective function in problem (VP) is quasiconvex or α -robustly quasiconvex (see Definition 2.2).

As mentioned in the introduction, when the function f is nonconvex, then the usual asymptotic function f^∞ does not provide adequate information on the behavior of f at infinity. For instance, and in relation to Theorem 3.1, we mention that when the function f is quasiconvex, the assumption

$$X^\infty \cap \mathcal{K}(f) = \{0\},$$

is too restrictive. Indeed, let us consider the one-dimensional real-valued function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = \sqrt{|x|}$ and $X = \mathbb{R}$. Here f is coercive and $\text{argmin}_X f = \{0\}$ is a singleton, but $X^\infty \cap \mathcal{K}(f) = \mathbb{R}$, and Theorem 3.1 cannot be applied to this basic situation.

On the other hand, if we use any of the generalized asymptotic functions f_q^∞ and f_λ^∞ , we obtain that

$$f_q^\infty(u) = +\infty \quad \forall u \neq 0, \quad \text{and} \quad 0 < f_\lambda^\infty(u) \leq 2 \quad \forall u \neq 0.$$

Therefore,

$$X^\infty \cap K_q(f) = \{0\}, \quad \text{and} \quad X^\infty \cap K_\lambda(f) = \{0\},$$

where $K_q(f) := \{d \in \mathbb{R}^n : f_q^\infty(d) \leq 0\}$ and $K_\lambda(f) := \{d \in \mathbb{R}^n : f_\lambda^\infty(d) \leq 0\}$, respectively.

Before stating main results of this section, we give a result on the nonemptiness and compactness of the solution set to constrained optimization problems by using the q -asymptotic function.

Lemma 5.1 *Assume that X is a convex set and f is a \mathbb{R}_+^m -quasiconvex mapping. Then, the following assertions are equivalent:*

- (a) $\text{Sol}^w(\text{VP})$ is nonempty and compact.
- (b) $X^\infty \cap \left(\bigcup_{i \in I} \mathcal{K}_q(f_i)\right) = \{0\}$.

Proof In order to prove the equivalence, we first show that

$$(f_i + \delta_X)_q^\infty(d) = (f_i)_q^\infty(d) + \delta_{X^\infty}(d) \quad \forall i \in I.$$

Indeed, by definition of the q -asymptotic function and the convexity of X , we have

$$\begin{aligned} (f_i + \delta_X)_q^\infty(d) &= \sup_{x \in \text{dom}(f_i + \delta_X)} \sup_{t > 0} \frac{(f_i + \delta_X)(x + td) - (f_i + \delta_X)(x)}{t} \\ &= \sup_{x \in X} \sup_{t > 0} \frac{f_i(x + td) + \delta_X(x + td) - f_i(x)}{t} \\ &= \begin{cases} (f_i)_q^\infty(d), & \text{if } d \in X^\infty, \\ +\infty, & \text{otherwise,} \end{cases} \\ &= (f_i)_q^\infty(d) + \delta_{X^\infty}(d). \end{aligned}$$

Hence,

$$\mathcal{K}_q(f_i + \delta_X) = \{d \in \mathbb{R}^n : (f_i)_q^\infty(d) + \delta_{X^\infty}(d) \leq 0\} = X^\infty \cap \mathcal{K}_q(f_i) \quad \forall i \in I.$$

This implies that

$$\bigcup_{i \in I} \mathcal{K}_q(f_i + \delta_X) = X^\infty \cap \left(\bigcup_{i \in I} \mathcal{K}_q(f_i)\right).$$

Combining this with [18, Proposition 4.5] we obtain the conclusion of the lemma. \square

In the next proposition we improve Theorem 3.1 for proper, lsc, \mathbb{R}_+^m - α -robustly quasiconvex functions.

Proposition 5.1 *Let X be a convex set and f be a proper, lsc, \mathbb{R}_+^m - α -robustly quasiconvex mapping ($\alpha > 0$). If*

$$X^\infty \cap \left(\bigcup_{i \in I} \mathcal{K}_q(f_i)\right) = \{0\}, \quad (12)$$

then there exists $\varepsilon > 0$ such that for all $u \in \mathbb{B}_\varepsilon$, the following statements hold:

- (a) f_u is bounded from below on X .
- (b) $\text{Sol}^w(u)$ is nonempty and compact.
- (c) $\text{Lim sup}_{u \rightarrow 0} \text{Sol}^w(u) \subset \text{Sol}^w(0)$, i.e., the solution mapping $\text{Sol}^w(\cdot)$ is closed at 0.
- (d) $\text{Sol}^w(\cdot)$ is usc at 0.

Proof We first show that there exists $\varepsilon > 0$ such that the following condition holds

$$X^\infty \cap \left(\bigcup_{i \in I} \mathcal{K}_q(f_i^{u_i}) \right) = \{0\} \quad \forall u \in \mathbb{B}_\varepsilon. \quad (13)$$

Indeed, suppose on the contrary that for every $k \in \mathbb{N}$, there exists $u^k \in \mathbb{B}_{\frac{1}{k}}$ such that (13) does not hold, i.e., there exist $i_k \in I$ and $d_k \in X^\infty \setminus \{0\}$ such that

$$\left(f_{i_k}^{u^k} \right)_q^\infty (d_k) \leq 0.$$

By the finiteness of I , there exists $i_0 \in I$ such that the inequality

$$\left(f_{i_0}^{u^k} \right)_q^\infty (d_k) \leq 0 \quad (14)$$

holds true for infinite many $k \in \mathbb{N}$. By passing to subsequences if necessary we may assume that (14) is satisfied for all $k \in \mathbb{N}$.

Since X^∞ is a closed cone, by passing a subsequence if necessary, we may assume that $\left\{ h_k := \frac{d_k}{\|d_k\|} \right\}_k \subset X^\infty$ converges to $h \in X^\infty$ with $\|h\| = 1$.

For every $k \in \mathbb{N}$, since $\left(f_{i_0}^{u^k} \right)_q^\infty$ is positively homogeneous of degree one and $\left(f_{i_0}^{u^k} \right)_q^\infty (h_k) \leq 0$, and by using relation (9), we have

$$\left(f_{i_0}^{u^k} \right)_q^\infty (h_k) = (f_{i_0})_q^\infty (h_k) - \langle u_k, h_k \rangle \leq 0.$$

Since f is lsc, $(f_{i_0})_q^\infty$ is lsc too by [17, p. 118], thus

$$(f_{i_0})_q^\infty (h) \leq \liminf_{k \rightarrow +\infty} (f_{i_0})_q^\infty (h_k) \leq \liminf_{k \rightarrow +\infty} \langle u_k, h_k \rangle = 0.$$

Hence, $h \in X^\infty \cap \mathcal{K}_q(f_{i_0})$ with $h \neq 0$, a contradiction. Therefore, relation (13) holds.

Since X is convex and f is \mathbb{R}_+^m - α -robustly quasiconvex, we obtain that the function $f_u(x) = f(x) - \langle u, x \rangle$ is \mathbb{R}_+^m -quasiconvex for all $u \in \mathbb{B}_\alpha$. Let ε be satisfied (13) and $\varepsilon < \alpha$. Hence, (b) follows from relation (13) and Lemma 5.1.

The relation (13) gives

$$X^\infty \cap \mathcal{K}_q(f_i^{u_i}) = \{0\} \quad \forall u \in \mathbb{B}_\varepsilon.$$

Combining this with [36, Proposition 5.4] yields that $f_i^{u_i}$ is bounded from below on X and (a) follows.

The proof of part (c) is quiet similar to the proof of Theorem 4.1(d), so omitted.

(d): Suppose on the contrary that $\text{Sol}^w(\cdot)$ is not usc at 0. Then, there exist an open set $U \subset \mathbb{R}^n$, with $\text{Sol}^w(0) \subset U$, sequences u_k and x_k such that

$u_k \rightarrow 0, x_k \in \text{Sol}^w(u_k) \setminus U$ for all $k \in \mathbb{N}$. We consider two case of the sequence x_k as follows.

Case 1. $\{x_k\}$ is bounded. Then, without loss of generality we may assume that $x_k \rightarrow \bar{x}$. By (c), one has $\bar{x} \in \text{Sol}^w(0) \subset U$. However, since $x_k \notin U$ for all k and U is open, we see that $\bar{x} \notin U$, a contradiction.

Case 2. $\{x_k\}$ is unbounded. By passing to a subsequence if necessary, we may assume that $\|x_k\| \rightarrow \infty$ as $k \rightarrow \infty$. Hence, without loss of generality one can assume that $\frac{x_k}{\|x_k\|} \rightarrow d \in X^\infty$ with $\|d\| = 1$. Fix any $y \in X$. By a similar argument as in the proof of Theorem 4.1(d), there exists $i_0 \in I$ such that

$$f_{i_0}(x_k) - \langle u_{i_0}^k, x_k \rangle \leq f_{i_0}(y) - \langle u_{i_0}^k, y \rangle \quad \forall k \in \mathbb{N}. \quad (15)$$

Since $u_k \rightarrow 0$, there exists $k_1 \in \mathbb{N}$ such that $u_k \in \mathbb{B}_\alpha$ for all $k \geq k_1$. Furthermore, since $\|x_k\| \rightarrow +\infty$ for every $t > 0$, there exists $k_2 \in \mathbb{N}$ such that $0 < \frac{t}{\|x_k\|} < 1$ for all $k \geq k_2$.

Since X is convex and f is \mathbb{R}_+^m - α -robustly quasiconvex, by (15), we obtain for every $k \geq k_0 := \max\{k_1, k_2\}$ that

$$\begin{aligned} & f_{i_0} \left(\left(1 - \frac{t}{\|x_k\|}\right) y + \frac{t}{\|x_k\|} x_k \right) - \left\langle u_k, \left(1 - \frac{t}{\|x_k\|}\right) y + \frac{t}{\|x_k\|} x_k \right\rangle \\ &= f_{i_0}^{u_k} \left(\left(1 - \frac{t}{\|x_k\|}\right) y + \frac{t}{\|x_k\|} x_k \right) \\ &\leq \max\{f_{i_0}^{u_k}(x_k), f_{i_0}^{u_k}(y)\} \\ &= f_{i_0}(y) - \langle u_k, y \rangle. \end{aligned}$$

Letting $k \rightarrow \infty$, we have

$$f_{i_0}(y + td) \leq f_{i_0}(y) \quad \forall t > 0, \quad \forall y \in X,$$

due to the lower semicontinuity of f_{i_0} . This implies that $(f_{i_0})_q^\infty(d) \leq 0$ and we therefore get $d \in X^\infty \cap \mathcal{K}_q(f_{i_0})$, contrary to (12). The proof is complete. \square

Furthermore, we also adapt the results for solution stability for the quasi-convex case below.

Proposition 5.2 *Let X be a convex set and f be an \mathbb{R}_+^m - α -robustly quasiconvex mapping ($\alpha > 0$). If the following conditions hold:*

- (a) $\text{Sol}^w(0)$ is a singleton;
- (b) $X^\infty \cap \left(\bigcup_{i \in I} \mathcal{K}_q(f_i)\right) = \{0\}$;

then $\text{Sol}^w(\cdot)$ is lsc at 0. Conversely, if $\text{Sol}^w(\cdot)$ is lsc at 0, then the following condition is satisfied

$$X^\infty \cap \left(\bigcap_{i \in I} \mathcal{K}_q(f_i)\right) = \{0\}. \quad (16)$$

Proof The proof of the first conclusion is quiet similar to that of the proof of Theorem 4.2, so omitted. We now suppose on the contrary that $\text{Sol}^w(\cdot)$ is lsc at 0 but (16) does not hold. Then, $\text{Sol}^w(0)$ is nonempty and there exists $d \in X^\infty \setminus \{0\}$ such that $(f_i)_q^\infty(d) \leq 0$ for every $i \in I$. Let $\bar{x} \in \text{Sol}^w(0)$. Clearly, $\bar{x} + td \in X$ for all $t > 0$. Let $u_k := \frac{1}{k}d$. Then $u_k \rightarrow 0$ as $k \rightarrow \infty$ and for each $i \in I$,

$$(f_i^{u_k})_q^\infty(d) = (f_i)_q^\infty(d) - \langle u_k, d \rangle = (f_i)_q^\infty(d) - \frac{1}{k}\|d\|^2 < 0.$$

By definition, we have

$$\sup_{t>0} \frac{f_i^{u_k}(\bar{x} + td) - f_i^{u_k}(\bar{x})}{t} < 0.$$

This implies that $f_i^{u_k}(\bar{x} + td) \rightarrow -\infty$ as $t \rightarrow \infty$ for every $i \in I$. This means that for each $z \in X$ there exists $t > 0$ large enough such that

$$\max_{i \in I} \{f_i^{u_k}(\bar{x} + td) - f_i^{u_k}(z)\} < 0.$$

It follows that $\text{Sol}^w(u_k) = \emptyset$. Hence, $\text{Sol}^w(\cdot)$ is not lsc at 0, a contradiction. The proof is complete. \square

6 Conclusions

In this paper, we have used the tool of asymptotic analysis to investigate existence and stability of weak Pareto solutions of constrained nonsmooth vector optimization problems. Under the proposed assumption that

$$X^\infty \cap \left(\bigcup_{i \in I} \mathcal{K}(f_i) \right) = \{0\},$$

which is a relation between the asymptotic of objective functions and the asymptotic cone of the constraint set, we present new results on existence of weak Pareto solutions, the weak sharp minima at infinity property, the upper/lower semicontinuity, the nonemptiness and compactness, and the closedness of the weak Pareto solution map of the considered problem with linear perturbation. We have used q -asymptotic functions to obtain similar results to the particular case when the objective function is quasiconvex or α -robustly quasiconvex.

Disclosure statement

The authors declare that they have no conflict of interest.

Data availability

There is no data included in this paper.

Funding

This research is funded by Hanoi Pedagogical University 2.

References

1. Ait Mansour, M., Chbani, Z., Riahi, H.: Recession bifunction and solvability of noncoercive equilibrium problems. *Commun. Appl. Anal.* **7**, 369–377 (2003)
2. Amara, C.: Directions de majoration d'une fonction quasiconvexe et applications. *Serdica Math. J.* **24**, 289–306 (1998)
3. Ansari, Q.H., Yao, J.C.: On nondifferentiable and nonconvex vector optimization problems. *J. Optim. Theory Appl.* **106**, 487–500 (2000)
4. Ansari, Q.H., Yao, J.-C.: *Recent Advances in Vector Optimization*. Springer, Berlin (2012)
5. Ansari, Q.H., Köbis, E., Yao, J.C.: *Vector Variational Inequalities and Vector Optimization: Theory and Applications*. Springer (2018)
6. Attouch, H., Buttazzo, G., Michaille, G.: *Variational Analysis in Sobolev and BV Spaces: Applications to PDEs and Optimization*. MPS-SIAM, Philadelphia (2006)
7. Auslender, A.: Noncoercive optimization problems. *Math. Oper. Res.* **21**, 769–782 (1996)
8. Auslender, A.: How to deal with the unbounded in optimization: theory and algorithms. *Math. Program.* **79**, 3–18 (1997)
9. Auslender, A., Teboulle, M.: *Asymptotic Cones and Functions in Optimization and Variational Inequalities*. Springer, Berlin (2003)
10. Barron, E.N., Goebel, R., Jensen, R.R.: Functions which are quasiconvex under linear perturbations. *SIAM J. Optim.* **22**, 1089–1108 (2012)
11. Borwein, J.M.: On the existence of Pareto efficient points. *Math. Oper. Res.* **8**, 64–73 (1983)
12. Cambini, A., Martein, L.: *Generalized Convexity and Optimization*. Springer-Verlag, Berlin Heidelberg (2009)
13. Debreu, G.: *Theory of Value*. John Wiley, New York (1959)
14. Ehrgott, M.: *Multicriteria Optimization*. Springer (2005)
15. Flores-Bazán, F.: Existence theorems for generalized noncoercive equilibrium problems: the quasiconvex case. *SIAM J. Optim.* **11**, 675–690 (2001)
16. Flores-Bazán, F.: Ideal, weakly efficient solutions for vector optimization problems. *Math. Program.* **93**, 453–475 (2002)
17. Flores-Bazán, F., Flores-Bazán, F., Vera, C.: Maximizing and minimizing quasiconvex functions: related properties, existence and optimality conditions via radial epiderivates. *J. Global Optim.* **63**, 99–123 (2015)
18. Flores-Bazán, F., López, R., Vera, C.: Vector asymptotic functions and their application to multiobjective optimization problems. *SIAM J. Optim.* **34**, 1826–1851 (2024)
19. Hadjisavvas, N., Komlosi, S., Schaible, S.: *Handbook of Generalized Convexity and Generalized Monotonicity*. Springer, Boston (2005)
20. Hadjisavvas, N., Lara, F., Luc, D.T.: A general asymptotic function with applications in nonconvex optimization. *J. Global Optim.* **78**, 49–68 (2020)
21. Hadjisavvas, N., Lara, F., Martínez-Legaz, J.E.: A quasi-convex asymptotic function with applications in optimization. *J. Optim. Theory Appl.* **180**, 170–186 (2019)
22. Hernández, E., López, R.: Stability in set-valued optimization problems using asymptotic analysis and epi-convergence. *Appl. Math. Optim.* **86**, 7 (2022)
23. Hung, N.C., Chuong, T.D., Anh, N.L.H.: Solution existence for a class of nonsmooth robust optimization problems. *J. Global Optim.* (2024). <https://doi.org/10.1007/s10898-024-01450-9>
24. Hung, N.C., Chuong, T.D., Anh, N.L.H.: Solution existence and compactness analysis for nonsmooth optimization problems. To appear in *J. Optim. Theory Appl.* (2025)
25. Huy, N.Q., Kim, D.S., Tuyen, N.V.: Existence theorems in vector optimization with generalized order. *J. Optim. Theory Appl.* **174**, 728–745 (2017)

26. Iusem, A., Lara, F.: Second order asymptotic functions and applications to quadratic programming. *J. Convex Anal.* **25**, 271–291 (2018)
27. Iusem, A., Lara, F.: Optimality conditions for vector equilibrium problems with applications. *J. Optim. Theory Appl.* **180**, 187–206 (2019)
28. Iusem, A., Lara, F.: Existence results for noncoercive mixed variational inequalities in finite dimensional spaces. *J. Optim. Theory Appl.* **183**, 122–138 (2019)
29. Iusem, A., Lara, F.: Quasiconvex optimization and asymptotic analysis in Banach spaces. *Optimization* **69**, 2453–2470 (2020)
30. Jahn, J.: Existence theorems in vector optimization. *J. Optim. Theory Appl.* **50**, 397–406 (1986)
31. Khan, A.A., Tammer, C., Zalinescu, C.: *Set-Valued Optimization*. Springer (2016)
32. Kim, D.S., Pham, T.S., Tuyen, N.V.: On the existence of Pareto solutions for polynomial vector optimization problems. *Math. Program.* **177**, 321–341 (2019)
33. Lara, F.: Generalized asymptotic functions in nonconvex multiobjective optimization problems. *Optimization* **66**, 1259–1272 (2017)
34. Lara, F.: Quadratic fractional programming under asymptotic analysis. *J. Convex Anal.* **26**, 15–32 (2019)
35. Lara, F., López, R.: Formulas for asymptotic functions via conjugates, directional derivatives and subdifferentials. *J. Optim. Theory Appl.* **173**, 793–811 (2017)
36. Lara, F., Tuyen, N.V., Nghi, T.V.: Weak sharp minima at infinity and solution stability in mathematical programming via asymptotic analysis. *arXiv preprint* (2024). <https://arxiv.org/abs/2410.04695>
37. Lee, J.H., Sissarat, N., Jiao, L.G.: Multi-objective convex polynomial optimization and semidefinite programming relaxations. *J. Global Optim.* **80**, 117–138 (2021)
38. Luc, D.T.: *Theory of Vector Optimization*. Springer, Berlin (1989)
39. Luc, D.T.: An existence theorem in vector optimization. *Math. Oper. Res.* **14**, 693–699 (1989)
40. Luc, D.T., Penot, J.P.: Convergence of asymptotic directions, *Trans. Amer. Math. Soc.* **353**, 4095–4121 (2001)
41. Mordukhovich, B.S.: *Variational Analysis and Generalized Differentiation, Vol. II: Applications*. Springer (2006)
42. Phu, H.X., An, P.T.: Stable generalization of convex functions. *Optimization* **38**, 309–318 (1996)
43. Phu, H.X., An, P.T.: Stability of generalized convex functions with respect to linear disturbances. *Optimization* **46**, 381–389 (1999)
44. Rele, R., Nedić, A.: On existence of solutions to non-convex minimization problems. *arXiv preprint* (2024). <https://arxiv.org/abs/2405.04688>
45. Rockafellar, R.T.: *Convex Analysis*. Princeton University Press, Princeton, New Jersey (1970)
46. Rockafellar, R.T., Wets, R.J.: *Variational Analysis*. Springer, New York (2009)
47. Steinitz, R.E.: Bedingt konvergente Reihen und konvexe Systeme I, II, III, *J. Reine Angew. Math.* **143**, 128–175 (1913); **144**, 1–40 (1914); **146**, 1–52 (1916)
48. Wang, G., Fu, Y.: Characterizing the nonemptiness and compactness of weakly efficient solution sets for non-convex multiobjective optimization problems. *Oper. Res. Lett.* **53**, 107092 (2024)