Planar graphs with distance of 3-cycles at least 2 and no cycles of lengths 5, 6, 7

Tao Wang*

Ya-Nan Wang[†]

Xiaojing Yang[‡]

June 27, 2024

Abstract

Weak degeneracy of a graph is a variation of degeneracy that has a close relationship to many graph coloring parameters. In this article, we prove that planar graphs with distance of 3-cycles at least 2 and no cycles of lengths 5, 6, 7 are weakly 2-degenerate. Furthermore, such graphs can be vertex-partitioned into two subgraphs, one of which has no edges, and the other is a forest.

Keywords: Weakly degenerate; $(\mathcal{I}, \mathcal{F})$ -partitionable; Planar graphs

1 Introduction

A graph G is k-degenerate if the minimum degree of every subgraph is at most k. Equivalently, the vertices of G can be linearly ordered < such that each vertex v has at most k neighbors u with v < u. Using Euler's formula, we know that every triangle-free planar graph is 3-degenerate. Regarding to degeneracy of some planar graphs, see [4, 9, 12].

Let \mathbb{Z} be the set of integers, and \mathbb{Z}^G be the set of all mappings from V(G) to \mathbb{Z} . For a subset U of V(G), we let $f_{-U}: V(G) - U \longrightarrow \mathbb{Z}$ be a mapping defined as $f_{-U}(x) = f(x) - |N_G(x) \cap U|$ for all $x \in V(G) - U$. In particular, we write f_{-v} for $f_{-\{v\}}$. A graph G is strictly f-degenerate if every subgraph R of G contains a vertex v with $d_R(v) < f(v)$. According to definition, G is strictly 1-degenerate if and only if it has no edges, which is an independent set; G is strictly 2-degenerate if and only if it is a forest.

Motivated by the study of greedy algorithms for graph coloring, Bernshteyn and Lee [1] introduced a weaker version of ordinary degenerate, named weakly degenerate.

Definition 1. Given a graph G and $f \in \mathbb{Z}^G$. For a vertex $u \in V(G)$, the operation $\mathsf{Delete}_u(G, f)$ outputs the graph $G_{-u} = G - u$ and the function f_{-u} . If $f(u) \ge 0$ and $f_{-u} \ge 0$, we say the operation Delete_u is *legal*.

Definition 2. Given a graph G and $f \in \mathbb{Z}^G$. For two adjacent vertices a and b in G, the operation $\mathsf{DeleteSave}_{(a,b)}$ on (G, f) outputs the graph $G_{-a} = G - a$ and the function f' defined as

$$f'(v) \coloneqq \begin{cases} f(v) - 1, & \text{if } v \text{ is a neighbor of } a \text{ and } v \neq b; \\ f(v), & \text{otherwise.} \end{cases}$$

^{*}Center for Applied Mathematics, Henan University, Kaifeng, 475004, P. R. China. Corresponding author: wangtao@henu.edu.cn; https://orcid.org/0000-0001-9732-1617.

[†]School of Mathematics and Statistics, Henan University, Kaifeng, 475004, P. R. China.

[‡]School of Mathematics and Statistics, Henan University, Kaifeng, 475004, P. R. China.



Fig. 1: A set of unavoidable configurations, where a semicircle represents a path of length two or three with interior vertices having degree three in G, a solid point represents a vertex of degree three, and a green square point represents a vertex of degree four in G.

If f(a) > f(b) and $f' \ge 0$, we say the operation $\mathsf{DeleteSave}_{(a,b)}$ is legal on (G, f).

A graph G is weakly f-degenerate if it can be reduced to a null graph through a sequence of legally operations Delete and DeleteSave. For a positive integer k, we define G as weakly k-degenerate if it is weakly f-degenerate, where f is a constant function with a value of k. The degeneracy d(G) of G is the minimum integer k such that G is k-degenerate. The weak degeneracy wd(G) of G is the minimum integer k for which G is weakly k-degenerate. Bernshteyn and Lee [1] established a chain of inequalities:

$$\chi(G) \le \chi_{\ell}(G) \le \chi_{\mathsf{DP}}(G) \le \mathsf{wd}(G) + 1 \le \mathsf{d}(G) + 1, \tag{1}$$

where $\chi_{\mathsf{DP}}(G)$ is the DP-chromatic number of G.

According to Eq. (1), various graph coloring parameters are upper bounded by wd(G) + 1. Consequently, there is significant interest in finding various sufficient conditions for planar graphs to be weakly k-degenerate. Wang [11] provided two such conditions for a plane graph to be weakly 2-degenerate. A best possible condition for a toroidal graph to be weakly 3-degenerate was given in [3].

Han et al. [5] proved that planar graphs of girth 5 are weakly 2-degenerate.

Theorem 1.1 (Han et al. [5]). If G is a triangle-free planar graph in which no 4-cycle is normally adjacent to a cycle of length at most five, then $wd(G) \leq 2$.

Liu et al. [7] proved that planar graphs with triangles at distance greater than two and no 5-, 6-, 7-cycles are DP-3-colorable. This paper strengthens this result. Let $dist^{\Delta}$ be the smallest distance between triangles, i.e., $dist^{\Delta} = \min\{d_G(v_1, v_2) : v_i \in V(T_i) \text{ and } T_i$'s are triangles of $G\}$.

Theorem 1.2. Every planar graph with dist^{Δ} ≥ 2 and without 5-, 6-, 7-cycles is weakly 2-degenerate.

We first prove the following structural result, which can imply Theorem 1.2.

Theorem 1.3. Let G be a plane graph with dist^{Δ} \geq 2 and without cycles of length 5, 6, 7. Then one of the following two conclusions holds:

1. there is a vertex of degree at most two.

2. there exists a degree-restricted subgraph isomorphic to a configuration as depicted in Fig. 1.

Actually, Theorem 1.3 can also be used to prove a result on a special strictly f-degenerate transversal, which can imply an $(\mathcal{I}, \mathcal{F})$ -coloring of G.

A graph G is $(\mathcal{I}, \mathcal{F})$ -partitionable if its vertices set can be divided into two parts, one part is a forest and the other has no edges. Borodin and Glebov [2] confirmed that every planar graph with girth at least 5 is $(\mathcal{I}, \mathcal{F})$ -partitionable. Liu et al. [7] proved that planar graphs without 4-, 6- and 8-cycles are $(\mathcal{I}, \mathcal{F})$ -partitionable, and Kang et al. [6] proved that planar graphs without 4-, 6- and 9-cycles are $(\mathcal{I}, \mathcal{F})$ -partitionable. In this paper, we are interested in the $(\mathcal{I}, \mathcal{F})$ -partition of planar graphs without 5-, 6- and 7-cycles.

Theorem 1.4. Every planar graph with dist^{Δ} \geq 2 and without cycles of length 5, 6, 7 is (\mathcal{I}, \mathcal{F})-partitionable.

To end this section, we give some notations. An *l*-vertex, or an l^+ -vertex, or an l^- -vertex means a vertex with degree l, or at least l, or at most l. The notions of an *l*-face, an l^+ -face and an l^- -face are defined similarly. For a face $f \in F$, if the vertices on f are cyclically listed as $u_1, u_2, u_3, \ldots, u_k$, then we say that f is a $(d(u_1), d(u_2), \ldots, d(u_k))$ -face. A face is light if the vertices on this face are all 3-vertices. If ab is a common edge of a 4⁻-face and a 7⁺-face, then we say ab controls the smaller 4⁻-face. For a given face g, we say a facial (d_1, d_2, \ldots, d_t) -walk on g is a d- or d^- -controlling walk if each edge on the walk controls a face other than g with size d or at most d, respectively. If all vertices on the walk are distinct, we say the walk is a dor d^- -controlling path. Assume abc is a facial path on a 7⁺-face f, we say b is rich with respect to f if neither ab nor bc controls a 4⁻-face, semi-rich with respect to f if exactly one of ab and bc controls a 4⁻-face, and poor with respect to f if each of ab and bc controls a 4⁻-face. We say a facial path is maximal if every edge on f controls a 4⁻-face, but the edges on the path immediately after and before it control 5⁺-faces. Let h be a face with $d(h) \geq 7$, and let $s_0(h)$ denote the number of vertices on h that are not on any 4⁻-controlling path.

2 Proof of Theorem 1.3

Assume that G is a counterexample to Theorem 1.3. Then the minimum degree of G is at least three, and there exists no degree-restricted subgraph isomorphic to a configuration as depicted in Fig. 1.

Lemma 2.1. Assuming G is a plane graph with distance of triangles at least 2 and no cycles of lengths 5, 6, 7. Then the following structural results hold:

- (1) A 3-cycle and a 4^- -cycle have no edges in common.
- (2) There are no 6-faces.
- (3) A 7-face adjacent to a 4^- -face should be as depicted in Fig. 2.
- (4) A 3-face is not adjacent to any 7^{-} -face.
- (5) A 4-face is not adjacent to any 6^- -face.
- (6) If an 8-face has a cut vertex incident to two 4⁻-faces, then it should be as depicted in Fig. 4.



Fig. 2: A 7-face adjacent to a 4-face.



(a) Type I, where at least one of u and v is a 4⁺-vertex. (b) Type II, where w is a 4⁺-vertex.

Fig. 3: Two types of special 8-faces.

Proof. Suppose that f is a 6-face. Since there are no 6-cycles in G, the boundary of f is not a 6-cycle. Then the boundary of f consists of two triangles, this contradicts the hypothesis that triangles have distance at least two from each other. Hence, there are no 6-faces in G.

Let g be a 7-face. Similarly, the boundary of g is not a 7-cycle, so it can only be a triangle and a 4-cycle. Let [xyzxabc] be a 7-face. It follows from Lemma 2.1(1) that none of xy, yz and xz can be incident with a 4⁻-face. Similarly, none of xa, ab, bc and cx can be incident with a 3-face. If the 7-face g is adjacent to a 4-face h, then h have two consecutive common edges with xabc. Hence, it must be as drawn in Fig. 2.

Assume f is an 8-face that is bounded by an 8-cycle. We say that f is *special* if it is incident with seven 3-vertices and a semi-rich 4-vertex, adjacent to two 3-faces and two 4-faces, precisely three of them are light 4⁻-faces, and the other 4⁻-face containing at least two 4⁺-vertices. See an illustration in Fig. 3.

Lemma 2.2. A 4-vertex cannot be incident to two special 8-faces at opposite.

Proof. Let f_1, f_2, f_3 and f_4 be four incident faces of a 4-vertex v. Assume f_2 and f_4 are two special 8-faces. By the definition of special 8-faces, v is a semi-rich 4-vertex of the incident special 8-faces f_2 and f_4 . It follows that exactly one of f_1 and f_3 , say f_1 , is a 4⁻-face. Furthermore, f_1 is a 4-face incident with precisely two 4⁺-vertices. Hence, G contains a copy of Fig. 1(h).

We use the famous discharging method to finish the proof. Let $\mu(v) = 2d(v) - 6$ be the initial charge for each vertex $v \in V(G)$ and $\mu(f) = d(f) - 6$ be the initial charge of each face $f \in F(G)$. By the Handshaking theorem and Euler's formula, we have that

$$\sum_{v \in V(G) \cup F(G)} \mu(x) = \sum_{v \in V(G)} (2d(v) - 6) + \sum_{f \in F(G)} (d(f) - 6) = -12$$

Applying the following discharging rules, we obtain a new charge function μ^* . In this procedure, the total sum of charges is preserved. However, we show that $\mu^* \ge 0$ for each vertex and each face, a contradiction. For convenience, we use $\tau(a \to b)$ to denote the charges transferred from the element a to the element b.

The discharge rules are as follows:

x

R-1. Each 4^- -face gets 1 from each 4^+ -vertex on it.

R-2. Let f and g be two adjacent faces. Then

$$\tau(g \to f) = \begin{cases} 1, & \text{if } f \text{ is a } (3,3,3)\text{-face.} \\ \frac{2}{3}, & \text{if } f \text{ is a } (3,3,4^+)\text{-face.} \\ \frac{1}{3}, & \text{if } f \text{ is a } (3,4^+,4^+)\text{-face.} \\ \frac{1}{2}, & \text{if } f \text{ is a } (3,3,3,3)\text{-face.} \\ \frac{1}{4}, & \text{if } f \text{ is a } (3,3,3,4^+)\text{-face.} \\ \frac{1}{2}, & \text{if } g \text{ is a } (3^+,4^+,4^+,4^+)\text{-face and it has a common } (4^+,4^+)\text{-edge with } f. \end{cases}$$

- **R-3.** Assume v is a 4-vertex incident with four faces f_1, f_2, f_3 and f_4 in a cyclic order, and v is incident to at most one 4⁻-face.
 - (a) If f_1 is a 4⁻-face, f_2 is a special 8-face, and f_4 is not a special 8-face, then v sends $\frac{1}{2}$ to f_2 , and $\frac{1}{4}$ to each of f_3 and f_4 .
 - (b) If f_1 is a 4⁻-face, neither f_2 nor f_4 is a special 8-face, then v sends $\frac{1}{2}$ to f_3 , and $\frac{1}{4}$ to each of f_2 and f_4 .
 - (c) If v is not incident to any 4⁻-face, then v sends $\frac{1}{2}$ to each incident face.

R-4. Let v be a 5-vertex on an 8-face f. Then

$$\tau(v \to f) = \begin{cases} \frac{3}{4}, & \text{if } v \text{ is semi-rich with respect to } f.\\\\ \frac{1}{2}, & \text{otherwise.} \end{cases}$$

R-5. Each 8-face receives 1 from each incident 6^+ -vertex.

Lemma 2.3. For every $z \in V(G) \cup F(G)$, we have $\mu^*(z) \ge 0$.

Proof. Let v be an arbitrary vertex in G. If d(v) = 3, then $\mu^*(v) = \mu(v) = 0$. Let v be a 4-vertex incident with four faces f_1, f_2, f_3 and f_4 in a cyclic order. We have the following scenarios:

• Suppose that v is incident with at least two 4⁻-faces. Since there are no adjacent 4⁻-faces, v is incident with precisely two 4⁻-faces at opposite. By R-1, v gives 1 to each incident 4⁻-face, then $\mu^*(v) = 2 \times 4 - 6 - 1 \times 2 = 0$.

• If f_1 is a 4⁻-face, then by Lemma 2.2, it is impossible that both f_2 and f_4 are special 8-faces.

• Suppose that f_1 is a 4⁻-face, f_2 is a special 8-face, but f_4 is not. By R-1 and R-3a, v sends 1 to f_1 , $\frac{1}{2}$ to f_2 , and $\frac{1}{4}$ to each of f_3 and f_4 . Then $\mu^*(v) = 2 \times 4 - 6 - 1 - \frac{1}{2} - \frac{1}{4} \times 2 = 0$.

• Suppose that f_1 is a 4⁻-face, and neither f_2 nor f_4 is a special 8-face. By R-1 and R-3b, v sends 1 to f_1 , $\frac{1}{2}$ to f_3 , and $\frac{1}{4}$ to each of f_2 and f_4 . Then $\mu^*(v) = \mu(v) - 1 - \frac{1}{2} - \frac{1}{4} \times 2 = 0$.

• If every face incident to v is a 5⁺-face, then it should send $\frac{1}{2}$ to each incident face by R-3c, thus $\mu^*(v) = 2 \times 4 - 6 - \frac{1}{2} \times 4 = 0$.

Let d(v) = 5. Since 4⁻-faces are not adjacent, v is incident to at most two 4⁻-faces. According to R-4, when v is semi-rich with respect to an 8-face, it gives $\frac{3}{4}$ to each such 8-face; if v is a rich or poor 5-vertex on an 8-face, then it gives $\frac{1}{2}$ to each such face. If v is incident with two

4⁻-faces, then $\mu^*(v) \ge 2 \times 5 - 6 - 1 \times 2 - \frac{3}{4} \times 2 - \frac{1}{2} = 0$. If v is incident with precisely one 4⁻-face, then $\mu^*(v) \ge 2 \times 5 - 6 - 1 - \frac{3}{4} \times 2 - \frac{1}{2} \times 2 > 0$. If every face incident with v is a 5⁺-face, then $\mu^*(v) \ge 2 \times 5 - 6 - \frac{1}{2} \times 5 > 0$.

Let $d(v) \ge 6$. According to the discharging rules, v sends at most 1 to each incident face, implying $\mu^*(v) \ge 2d(v) - 6 - 1 \times d(v) \ge 0$.

Next, we check μ^* for every face. Let f be an arbitrary 4⁻-face. By Lemma 2.1(4) and Lemma 2.1(5), all faces adjacent to f are 7⁺-faces. If there is no 4⁺-vertices on f, then it receives 1 from each adjacent face, implying $\mu^*(f) = 3-6+1\times 3 = 0$. If f is a $(3,3,4^+)$ -face, then it receives $\frac{2}{3}$ from each adjacent face, and 1 from the incident 4⁺-vertex, thus $\mu^*(f) = 3-6+1+\frac{2}{3}\times 3 = 0$. If f there are exactly two 4⁺-vertices, then $\mu^*(f) = 3-6+1\times 2+\frac{1}{3}\times 3 = 0$ by R-2. If f is a $(4^+, 4^+, 4^+)$ -face, then $\mu^*(f) = 3-6+1\times 3 = 0$ by R-2. If f is a (3,3,3,3)-face, then it receives $\frac{1}{2}$ from each adjacent face, thus $\mu^*(f) = 4-6+\frac{1}{2}\times 4 = 0$. If f is a (3,3,3,3)-face, then it receives 1 from the incident 4⁺-vertex, and $\frac{1}{4}$ from each adjacent face, thus $\mu^*(f) = 4-6+\frac{1}{4}\times 4+1 = 0$. If f is incident with precisely two 3-vertices, then it receives 1 from each of the two incident 4⁺-vertices, thus $\mu^*(f) = 4-6+1\times 2 = 0$. If f is a $(3,4^+,4^+,4^+)$ -face, then it sends $\frac{1}{2}$ to each adjacent face that has a common $(4^+,4^+)$ -edge with f, and thus $\mu^*(f) = 4-6+1\times 3-\frac{1}{2}\times 2 = 0$. If f is a $(4^+,4^+,4^+,4^+)$ -face, then it sends $\frac{1}{2}$ to each adjacent face that has $\mu^*(f) = 4-6+1\times 3-\frac{1}{2}\times 4 = 0$.

Since there are no 5-cycles, there are no 5-faces. By Lemma 2.1(2), there are no 6-faces in G. For a 7-face f, if it is adjacent to a 4⁻-face, then it must be as depicted in Fig. 2, thus it sends at most $\frac{1}{4} \times 2$ via the incident edges by R-2, thus $\mu^*(f) \ge 7 - 6 - \frac{1}{4} \times 2 = \frac{1}{2} > 0$. Let f be an 8⁺-face.

Claim 1. If P is a 4-controlling path on f, then f sends at most $\frac{1}{2}$ in total via the edges on P.

Proof. Let $P = u_1 u_2 \ldots u_t$. If P has length one, then f sends at most $\frac{1}{2}$ via the edge of $u_1 u_2$. Assume the length of P is at least two. Since there are no adjacent 4⁻-faces, the interior vertices u_2, \ldots, u_{t-1} are all 4⁺ vertices, thus f sends at most $\frac{1}{4}$ via each of $u_1 u_2$ and $u_{t-1} u_t$ and no charge via the edge $u_2 u_3, u_3 u_4, \ldots, u_{t-2} u_{t-1}$, thus it sends at most $\frac{1}{4} \times 2 = \frac{1}{2}$ via the path P.

Claim 2. Let P = xyzu be a 4⁻-controlling path on f. Then f sends at most $\frac{11}{12}$ in total via the edges on P.

Proof. Since there are no adjacent 4^- -faces, we have $d(y), d(z) \ge 4$. If the three controlled faces are 4-faces, then f sends at most $\frac{1}{4}$ via each of xy and zu, and no charge via yz, thus it sends at most $\frac{1}{4} \times 2 = \frac{1}{2} < \frac{11}{12}$ via the path P. Since the distance of triangles is least two, P controls at most one 3-face. If xy or zu, say xy, controls a 3-face, then f sends at most $\frac{2}{3}$ via $xy, \frac{1}{4}$ via zu, and no charge via yz, thus it sends at most $\frac{2}{3} + \frac{1}{4} = \frac{11}{12}$ via the path P. If yz controls a 3-face, then f sends at most $\frac{1}{3}$ via yz, at most $\frac{1}{4}$ via each of xy and zu, thus it sends at most $\frac{1}{3} + \frac{1}{4} \times 2 < \frac{11}{12}$ via the path P.

Similarly, we have the following claims for the 4⁻-controlling path of length at most two.

Claim 3. Let P be a 4⁻-controlling path of length two on f. Then f sends at most $\frac{11}{12}$ in total via the edges on P.

Claim 4. If uv is a 4⁻-controlling path on f, then f sends at most 1 via uv.

Remark 1. In the proof of the above four claims, when we give an upper bound on the total charge that v sends via a 4⁻-controlling path P, we treat all vertices on the controlled 4⁻-faces as 3-vertices except the interior vertices of the path P. Note that all interior vertices on P are 4⁺-vertices. Hence, when we know another 4⁺-vertex is on the controlled face, it will "save" at



Fig. 4: Two 4-faces share edges with a common 8-face.

least $\frac{1}{4}$. For example, if P is a maximal 4-controlling path of length one, then f sends at most $\frac{1}{2}$ via the path P; moreover, if the controlled face contains precisely one 4⁺-vertex, then it "saves" $\frac{1}{4}$, that is, f sends $\frac{1}{2} - \frac{1}{4} = \frac{1}{4}$ via the path P, this is consistent with R-2. If the controlled face is incident with precisely two 4⁺-vertices, then it "saves" $\frac{1}{4} \times 2 = \frac{1}{2}$, so f sends no charge via the path P, this is also consistent with R-2.

Let t'_3 = the number of 3-faces that are adjacent to f. Thus there are at most $\left\lfloor \frac{d(f)-s_0(f)-2t'_3}{2} \right\rfloor$ 4-controlling paths that are disjoint from controlled 3-faces. By Claims 1–4,

$$\mu^*(f) \ge \mu(f) - 1 \times t_3' - \frac{1}{2} \left\lfloor \frac{d(f) - s_0(f) - 2t_3'}{2} \right\rfloor = \left(d(f) - \frac{1}{2} \left\lfloor \frac{d(f) - s_0(f)}{2} \right\rfloor \right) - 6 - \frac{1}{2} t_3'. \quad (\star)$$

Since the distance of triangles is at least two, we have $t'_3 \leq \lfloor \frac{d(f)}{3} \rfloor$. It is easy to verify that $\mu^*(f) \geq 0$ when f is a 10⁺-face or a 9-face with $t'_3 \leq 2$. If d(f) = 9 and $t'_3 = 3$, then there is no adjacent 4-faces disjoint from the adjacent 3-faces, thus $\mu^*(f) \geq 3 - 1 \times t'_3 = 0$.

From now on, we assume that f is an 8-face. First, assume that the boundary of f is not a cycle. Since dist^{Δ} ≥ 2 and there is no 5-cycles, the boundary of f consists of two 4-cycles. In this case, f is not adjacent to any 3-face and is adjacent to at most two 4-faces, see Fig. 4. By R-2, f sends at most $\frac{1}{4} \times 4$ to the adjacent 4-faces, thus $\mu^*(f) \geq 8 - 6 - \frac{1}{4} \times 4 > 0$. Next, suppose f is bounded by an 8-cycle. Now, we have $t'_3 \leq \lfloor \frac{8}{3} \rfloor = 2$.

Suppose f contains at most two maximal 4^- -controlling paths. If f sends at most 1 via each maximal path, then $\mu^*(f) \ge 8 - 6 - 1 \times 2 = 0$. Assume f sends more than 1 via a maximal path P. Then P has length at least four and controls two 3-faces. It is easy to check that f sends at most $\frac{2}{3} \times 2 = \frac{4}{3}$ via the path P, and at most $\frac{1}{2}$ via the other maximal path (if it exists). Hence, $\mu^*(f) \ge 8 - 6 - \frac{4}{3} - \frac{1}{2} > 0$.

Assume f contains at least three maximal 4⁻-controlling paths. By parity, we have $s_0(f) \le 2$ and each maximal 4⁻-controlling path of f has length at most three. By Claims 1–4, it sends at most $\frac{1}{2}$ via each maximal 4-controlling path, and at most 1 to each of the other maximal 4⁻-controlling path. It follows that f sends at most $1 \times 2 + \frac{1}{2} \times 2 = 3$ in total.

If f contains a 6⁺-vertex, then $\mu^*(f) \ge 8 - 6 - 3 + 1 = 0$ by R-5.

Suppose that f is incident to a semi-rich 5-vertex u. Then it receives $\frac{3}{4}$ from each such vertex by R-4. Meanwhile, the 5-vertex u "saves" $\frac{1}{4}$ for the expenditure by Remark 1, thus $\mu^*(f) \ge 8 - 6 - 3 + \frac{1}{4} + \frac{3}{4} = 0$.

Suppose that f is incident to a poor or a rich 5-vertex, namely w. By parity, f must have precisely three maximal 4⁻-controlling paths. Thus, f receives at least $\frac{1}{2}$ from w by R-4, resulting in $\mu^*(f) \ge 8 - 6 - 1 \times 2 - \frac{1}{2} + \frac{1}{2} = 0$.

Assume every vertex on f is a 4⁻-vertex.

Case 1. There exists a rich 4-vertex on f.

By parity, f contains precisely three maximal 4⁻-controlling paths. Similar to the previous arguments, f sends at most $1 \times 2 + \frac{1}{2} = \frac{5}{2}$. If f can receive at least $\frac{1}{2}$ in total from its incident

vertices, then $\mu^*(f) \ge 8 - 6 - \frac{5}{2} + \frac{1}{2} = 0$. So we may assume that f can only receive less than $\frac{1}{2}$ in total from its incident vertices. Let $f = [vu_1u_2xyzw_2w_1]$ be an 8-face, and let v be a rich 4-vertex with four incident faces f_1, f_2, f_3 and f_4 in a cyclic order and $f = f_3$. By Lemma 2.2, at most one of f_2 and f_4 is a special 8-face. According to the discharging rules, we have $\tau(v \to f) = \frac{1}{4}$.

According to the discussions above, we can assume that each rich 4-vertex on f sends $\frac{1}{4}$ to f. It follows that f contains precisely one rich 4-vertex, namely v, for otherwise $\mu^*(f) \geq 1$ $8-6-\frac{5}{2}+\frac{1}{4}\times 2=0$. If f sends at most $\frac{9}{4}$ in total, then $\mu^*(f)\geq 8-6-\frac{9}{4}+\frac{1}{4}=0$. So we may assume that f sends more than $\frac{9}{4}$ in total. Since f contains precisely three maximal 4⁻-controlling paths and f is adjacent to at most two 3-faces, we conclude that there is a maximal 4-controlling path P. Then f sends more than $\frac{1}{4}$ via P, so P controls a (3,3,3,3)-face or two (3,3,3,4)-faces. According to the discharging rules, f sends $\frac{1}{2}$ in total via this maximal 4-controlling path P. Let the other two maximal 4⁻-controlling paths be P_1 and P_2 . Since f sends more than $\frac{9}{4}$ in total, we have that f sends at least $\frac{3}{4}$ via each of P_1 and P_2 . It follows that neither P_1 nor P_2 is a maximal 4-controlling path by Claims 1–4. Note that there are at most seven vertices on maximal 4⁻-controlling paths, and one from $\{P_1, P_2\}$, say P_1 , has length one. Thus, P_1 controls a (3,3,3)-face. Moreover, P_2 controls a (3,3,3)-face, or P_2 is a (3,4,3)-path controlling a (3,3,4)-face and a (3,3,3,4)-face. Since $\tau(v \to f) = \frac{1}{4}$, we have f_2 or f_4 , say f_4 , is a special 8-face. Since f is adjacent to two 3-faces and dist^{Δ} ≥ 2 , the vertex v is not incident with a 3-face. Thus, f_4 is a special 8-face of type I. Now, G contains Fig. 1(i) or Fig. 1(j) as a subgraph.

Case 2. There exists no rich 4-vertices on f.

Recall that f is adjacent to at most two 3-faces, at most two of the maximal 4⁻-controlling paths can control some 3-faces. Similar to the previous cases, f sends at most $1 \times 2 + \frac{1}{2} \times 2 = 3$. Assume that f contains at least two semi-rich 4-vertices. By R-3, f receives at least $\frac{1}{4}$ from each incident semi-rich 4-vertex. Moreover, each semi-rich 4-vertex "saves" at least $\frac{1}{4}$ by Remark 1. Thus, $\mu^*(f) \ge 8 - 6 - 3 + \frac{1}{4} \times 2 + \frac{1}{4} \times 2 = 0$. Assume that there is at most one semi-rich 4-vertex with respect to f.

Subcase 2.1. f contains precisely three maximal 4⁻-controlling paths.

Similar to the above cases, f sends at most $1 \times 2 + \frac{1}{2} = \frac{5}{2}$. If f contains a semi-rich 4-vertex, then it sends $\frac{1}{4}$ to f, and it also "saves" at least $\frac{1}{4}$ for f, yielding $\mu^*(f) \ge 8 - 6 - \frac{5}{2} + \frac{1}{4} \times 2 = 0$. So every 4-vertex on f is poor. Since f contains precisely three maximal 4⁻-controlling paths, it contains at most two poor 4-vertices. If f controls a $(4, 4, 3^+, 4^+)$ -face via a (4, 4)-edge, then it receives $\frac{1}{2}$ from this 4-face by R-2, implying $\mu^*(f) \ge 8 - 6 - \frac{5}{2} + \frac{1}{2} = 0$. If f controls a $(4, 4, 4^+)$ -face, then f contains a maximal controlling path of length three and it sends at most $\frac{1}{2}$ via this path, thus $\mu^*(f) \ge 8 - 6 - 1 - \frac{1}{2} \times 2 = 0$. Hence, every (4, 4)-edge on f controls a (4, 4, 3)-face or (4, 4, 3, 3)-face.

A controlled 4⁻-face g is good if it is controlled by a (3,3)-edge with a 4⁺-vertex on g, or controlled by a (3,4)-edge with at least two 4⁺-vertices on g. By Remark 1, every good face "saves" at least $\frac{1}{4}$ for f. If f controls at least two good faces, then it "saves" at least $\frac{1}{4} \times 2 = \frac{1}{2}$ in total, and then $\mu^*(f) \ge 8 - 6 - \frac{5}{2} + \frac{1}{2} = 0$. So we can assume that f controls at most one good face.

Assume that f contains no poor 4-vertex. Then f is adjacent to at least two light 4⁻-faces. If all the adjacent light faces are 4-faces, then $\mu^*(f) \ge 8-6-1-\frac{1}{2} \times 2 = 0$. Hence, f is adjacent to a light 3-face, and Fig. 1(a) is appeared in G, a contradiction. Assume that f contains precisely one poor 4-vertex. Then G contains a subgraph as depicted in Fig. 1(b) or Fig. 1(c), a contradiction.

Assume that f contains two non-adjacent poor 4-vertices. Then either Fig. 1(d) or Fig. 1(e) is a subgraph of G, a contradiction.

Assume that f contains two adjacent poor 4-vertices. Recall that every (4, 4)-edge on f controls a (4, 4, 3)-face or (4, 4, 3, 3)-face. Therefore, G contains a subgraph as depicted in Fig. 1(f), a contradiction.

Subcase 2.2. f contains precisely four maximal 4⁻-controlling paths.

Then each maximal 4⁻-controlling path has length one. If each controlled face contains a 4⁺-vertex, then each such 4⁺-vertex "saves" at least $\frac{1}{4}$ for f, implying $\mu^*(f) \ge 8 - 6 - 3 + \frac{1}{4} \times 4 = 0$. Hence, f controls a light 4⁻-face. If f contains no semi-rich 4-vertex, then each vertex on f is a 3-vertex, and there is a subgraph as depicted in Fig. 1(a), which leads to a contradiction.

Next, we consider the case that f contains precisely one semi-rich 4-vertex, say x. If f is adjacent to precisely one light 4⁻-face, then x sends $\frac{1}{4}$ to f by R-3, and each of the three non-light 4⁻-face "saves" at least $\frac{1}{4}$ for f, yielding $\mu^*(f) \ge 8 - 6 - 3 + \frac{1}{4} + \frac{1}{4} \times 3 = 0$. Hence, f is adjacent to at least two light 4⁻-faces. Suppose that xy controls a 4⁻-face h = [xyz] or h = [xyzu]. If h contains precisely one 4⁺-vertex, then there is a subgraph as depicted in Fig. 1(c) or Fig. 1(g), a contradiction. So h contains two 4⁺-vertices.

Assume h is a 4-face. If f sends at most $\frac{9}{4}$ in total to the other three controlled faces, then $\mu^*(f) \ge 8-6-\frac{9}{4}+\frac{1}{4}=0$, where the $\frac{1}{4}$ is given from x by R-3. The other case is when f sends more than $\frac{9}{4}$ in total to the other three controlled faces. Then f controls two light 3-faces and one light 4-face. Now, f is a special 8-face, and x sends $\frac{1}{2}$ to f by R-3, thus $\mu^*(f) \ge 8-6-1\times 2-\frac{1}{2}+\frac{1}{2}=0$.

Assume h is a 3-face. Indeed, h is a $(3, 4, 4^+)$ -face, and f sends $\frac{1}{3}$ to h. If f sends at most $\frac{23}{12}$ in total to the other three controlled faces, then $\mu^*(f) \ge 8 - 6 - \frac{23}{12} - \frac{1}{3} + \frac{1}{4} = 0$. The other case is that f sends more than $\frac{23}{12}$ in total to the other three controlled faces. Hence, f controls one light 3-face and two light 4-faces. Now, f must be a special 8-face. Then x sends $\frac{1}{2}$ to f by R-3, and $\mu^*(f) \ge 8 - 6 - 1 - \frac{1}{2} \times 2 - \frac{1}{3} + \frac{1}{2} > 0$.

3 Proof of Theorem 1.2

In this section, we establish the validity of Theorem 1.2. A graph is a *GDP-tree* if it is connected and each block is either a complete graph or a cycle. Bernshteyn and Lee [1] demonstrated the following conclusion.

Theorem 3.1. Assume G is a graph satisfying $wd(G) = d \ge 3$ and $wd(G^*) < d$ for each proper subgraph G^* of G. Then the following statements are true:

- (i) Every vertex has degree at least d.
- (ii) For any subset $X \subseteq \{x \in V(G) : d_G(x) = d\}$, each connected component of G[X] forms a GDP-tree.

Now, we can easily prove Theorem 1.2.

Proof of Theorem 1.2. Suppose that G is a counterexample to Theorem 1.2 with minimum number of |V(G)| + |E(G)|. Thus, G is connected. Moreover, wd(G) = 3 and $wd(G^*) < 3$ for every proper subgraph G^* of G. By Theorem 1.3, G has a vertex of degree at most two, or it



Fig. 5: A set of unavoidable configurations, where a semicircle represents a path of length two or three with interior vertices having degree three in G, a solid point represents a vertex of degree three, and a green square point represents a vertex of degree four in G.

contains a subgraph isomorphic to Fig. 1. According to Theorem 3.1, a 2⁻-vertex is reducible, hence there exists a subgraph isomorphic to Fig. 1. Let f(v) = 2 for each vertex $v \in V(G)$.

We use G_1 to denote the graph induced by all vertices in Fig. 1(a). As G_1 contains the depicted graph as a spanning subgraph, G_1 has exactly one block that is neither a cycle nor a complete graph. From Theorem 3.1, Fig. 1(a) cannot be a subgraph of G, which leads to a contradiction.

Let $G_t, t \in \{2, ..., 7\}$, represent the subgraph induced by vertices depicted in Figs. 1(b)–1(g). The vertices of G_t are labeled as illustrated in Figs. 5(b)–5(g). For each $i \in \{1, ..., 7\}$, if $x_i x_{i+1}$ controls a 4⁻-face, then $x_i x_{i+1}$ controls $x_i u_i v_i x_{i+1}$ or $x_i v_i x_{i+1}$. By the minimality, we remove all vertices from $G - V(G_t)$ by a series of legal operations. Given an ordering:

$$x_2, x_3, \ldots, x_8, x_1, (u_1), v_1,$$

if u_i, v_i $(2 \le i \le 7)$ exist in G_t , we insert the vertices u_i, v_i into the list order following x_i . Let g_t be the resulting function on G_t . Since $g_t(x_2) > g_t(v_1)$, we legally remove the vertices from G_t by a series of Delete operations except the operation DeleteSave (x_2,v_1) . Hence, G is weakly 2-degenerate, which contradicts our assumption.

Let G_8 denote the graph induced by vertices depicted in Fig. 1(h). The vertices in G_8 are labeled as illustrated in Fig. 5(h). We order the vertices of G_8 as:

$$v_6, v_7, u_6, u_7, v, u_1, u_2, \ldots, u_5, w_2, v_1, v_2, \ldots, v_5, w_1.$$

By minimality, starting from (G, f), we remove all vertices from $G - V(G_8)$ by a series of legal operations. Let f' be the resulting function on G_8 . Note that $f'(v_6) > f'(w_1)$ and $f'(u_6) >$ $f'(w_2)$. Beginning from (G_8, f') , following the given order, we can delete all vertices from G_8 , except for the initial operation $\mathsf{DeleteSave}_{(v_6,w_1)}$ and the subsequent operation $\mathsf{DeleteSave}_{(u_6,w_2)}$. Consequently, G is weakly 2-degenerate, which contradicts our assumption.

Let G_9 (resp. G_{10}) denote the graph induced by the vertices depicted in Fig. 1(i) (resp. Fig. 1(j)). The vertices of G_9 and G_{10} are as depicted in Figs. 5(i) and 5(j). The edge u_2u_3 in Fig. 5(j) controls a face $u_2w_4w_5u_3$ or $u_2w_5u_3$. We order all vertices S of G_9 and G_{10} as:

$$v_2, v_1, v, u_6, u_5, u_4, u_3, (w_5), (w_4), u_2, u_1, u, v_6, v_5, w_3, w_2, v_4, v_3, w_1.$$

Similarly, we remove all vertices from G - S by a sequence of legal operations. Let f' be the resulting function on S. Notably, $f'(v_2) > f'(w_1)$, thus we can further legally delete all vertices from S except for the operation $\mathsf{DeleteSave}_{(v_2,w_1)}$. Consequently, G is weakly 2-degenerate, which contradicts our assumption.

4 Proof of Theorem 1.4

In this section, we establish Theorem 1.4 utilizing a concept termed strictly f-degenerate transversals.

Let G be a simple graph, and for each $v \in V(G)$, let $L_v = \{(v, 1), (v, 2), \dots, (v, s)\}$. For each edge $ab \in E(G)$, let \mathscr{M}_{ab} be a matching between L_a and L_b . A cover of G is represented by a graph H with the following conditions:

- (i) the vertex set of H is the union of L_v for all $v \in V(G)$; and
- (ii) the edge set \mathscr{M} of H is the union of all \mathscr{M}_{ab} whenever ab is an edge of G.

Let H be a cover of G, and let $f \in \mathbb{Z}^H$, forming a valued cover (H, f) of G. For any subset S of V(G), the induced subgraph $H[\bigcup_{v \in S} L_v]$ is simply written as H_S . A subset $T \subseteq V(H)$ is a transversal of H if each L_v contains exactly one vertex from T. Furthermore, if H[T] is strictly f-degenerate, then T is called a strictly f-degenerate transversal (SfDT for short).

We establish a more robust result as follows:

Theorem 4.1. Let G be a plane graph with dist^{Δ} \geq 2 and without 5-, 6-, 7-cycles. Assume H is a cover and $f \in \{0, 1, 2\}^{H}$. If each vertex $v \in V(G)$ satisfies $f(v, 1) + \cdots + f(v, s) \geq 3$, then H contains a SfDT.

To establish Theorem 4.1, we require some preliminary results. For a graph G and a valued cover (H, f), it is *minimally non-strictly f-degenerate* if H has no SfDT, yet $H - L_v$ admits a SfDT for any $v \in V(G)$.

Define \mathscr{D} as the set of vertices $v \in V(G)$ satisfying $f(v, 1) + \cdots + f(v, s) \ge d_G(v)$. Lu et al. [8] presented the next result on critical graphs.

Theorem 4.2 (Lu et al. [8]). Let G be a graph and (H, f) be a valued cover of G. Let D' be a nonempty subset of \mathscr{D} with G[D'] having no cut vertices. If (H, f) is a minimally non-strictly f-degenerate pair, then

- (i) G is connected and $f(v, 1) + f(v, 2) + \cdots + f(v, s) \le d_G(v)$ for each $v \in V(G)$, and
- (ii) G[D'] is a cycle, a complete graph, or $d_{G[D']}(v) \leq \max_q \{f(v,q)\}$ for each $v \in D'$.

Theorem 4.3 (Wang et al. [10]). Let k be an integer with $k \ge 3$, and let K be an induced subgraph of G such that the vertices of K can be ordered as u_1, u_2, \ldots, u_m satisfying:

- (i) $k (d_G(u_1) d_K(u_1)) > k (d_G(u_m) d_K(u_m));$
- (ii) $d_G(u_m) \leq k$ and $u_1 u_m$ is an edge in G; and
- (iii) for $2 \le i \le m-1$, u_i has at most k-1 neighbors in $G \{u_{i+1}, \ldots, u_m\}$.

Let *H* be a cover of *G* and $f \in \{0, 1, 2\}^H$. If $f(v, 1) + \cdots + f(v, s) \ge k$ for each vertex $v \in V(G)$, then any SfDT of $H - H_K$ can be extended to that of *H*.

Now, we prove Theorem 4.1.

Proof of Theorem 4.1. Assume that G is a counterexample to Theorem 4.1 such that (H, f) contains no SfDT, yet $(H - L_v, f)$ has a SfDT for any $v \in V(G)$. According to Theorem 1.3, G must contain a 2⁻-vertex, or a subgraph isomorphic to Fig. 1. By Theorem 4.2, every vertex has degree at least three, thereby indicating the presence of a subgraph isomorphic to Fig. 1 in G.

In the following, we still define G_1, G_2, \ldots, G_{10} to be the graph induced by the vertices in Figs. 1(a)-1(j) respectively, as that of in the proof of Theorem 1.2, and the vertices of G_t are labeled as in Figs. 5(a)-5(j).

Note that G_1 is a 2-connected graph that is neither a cycle nor a complete graph. Furthermore, within this subgraph, there exists a vertex v with degree greater than two. This contradicts Theorem 4.2.

Next we consider G_t for $t \in \{2, ..., 7\}$. For each $i \in \{1, ..., 7\}$, if $x_i x_{i+1}$ controls a 4⁻-face, then $x_i x_{i+1}$ controls $x_i u_i v_i x_{i+1}$ or $x_i v_i x_{i+1}$. Given an ordering:

$$x_2, x_3, \ldots, x_8, x_1, (u_1), v_1,$$

if u_i, v_i $(2 \le i \le 7)$ exist in G_t , we insert the vertices of u_i, v_i to the list order following x_i . Denoting the resulting list as S. It is observed that S satisfies the conditions in Theorem 4.3 with k = 3 and pair (x_2, v_1) . By minimality, $(H - H_S, f)$ has a SfDT T. According to Theorem 4.3, we can obtain a SfDT of H by extending T, leading to a contradiction.

When we consider G_8 , we order the vertices of G_8 as follows:

$$v_6, v_7, u_6, u_7, v, u_1, u_2, \ldots, u_5, w_2, v_1, v_2, \ldots, v_5, w_1.$$

It is evident that this ordering satisfies the conditions in Theorem 4.3 with k = 3 and the pairs (v_6, w_1) and (u_6, w_2) . A similar contradiction is derived as above cases.

Now we consider G_9 and G_{10} . Note that the edge u_2u_3 in Fig. 5(j) controls a face $u_2w_4w_5u_3$ or $u_2w_5u_3$. We order all vertices S of G_9 and G_{10} as follows:

 $v_2, v_1, v, u_6, u_5, u_4, u_3, (w_5), (w_4), u_2, u_1, u, v_6, v_5, w_3, w_2, v_4, v_3, w_1.$

It is notable that this list complies with the conditions specified in Theorem 4.3 with k = 3 and the pair (v_2, w_1) . A similar contradiction is derived as above cases.

For a cover H of G, we define H as a good cover of G if it has a property that (u, i) and (v, j) are adjacent iff i = j and $uv \in E(G)$. Observe that a good cover of G is isomorphic to s copies of G.

Proof of Theorem 1.4. Let s = 2, and consider a good cover H of G. Let f be a mapping with f(v, 1) = 1 and f(v, 2) = 2 for every vertex $v \in V(G)$. According to Theorem 4.1, H has a SfDT T. Let $\mathcal{I} = \{v : (v, 1) \in T\}$ and $\mathcal{F} = \{v : (v, 2) \in T\}$. It is observed that \mathcal{I} is an independent set in G, and \mathcal{F} induces a forest in G. Consequently, G is $(\mathcal{I}, \mathcal{F})$ -partitionable. This establishes Theorem 1.4.

Acknowledgments. We thank the two anonymous referees for their valuable comments and constructive suggestions on the manuscript. The first author was supported by the Natural Science Foundation of Henan Province (No. 242300420238). The third author was supported by National Natural Science Foundation of China (No. 12101187).

References

- A. Bernshteyn and E. Lee, Weak degeneracy of graphs, J. Graph Theory 103 (4) (2023) 607–634.
- [2] O. V. Borodin and A. N. Glebov, On the partition of a planar graph of girth 5 into an empty and an acyclic subgraph, Diskretn. Anal. Issled. Oper. Ser. 1 8 (4) (2001) 34–53.
- [3] P. Chen and T. Wang, Toroidal graphs without K_5^- and 6-cycles, Discrete Math. 347 (9) (2024) 114076.
- [4] G. Fijavž, M. Juvan, B. Mohar and R. Škrekovski, Planar graphs without cycles of specific lengths, European J. Combin. 23 (4) (2002) 377–388.
- [5] M. Han, T. Wang, J. Wu, H. Zhou and X. Zhu, Weak degeneracy of planar graphs and locally planar graphs, Electron. J. Combin. 30 (4) (2023) Paper No. 4.18.

- [6] Y. Kang, H. Lu and L. Jin, (I, F)-partition of planar graphs without cycles of length 4, 6, and 9, Discuss. Math. Graph Theory (2024) https://doi.org/10.7151/dmgt.2523.
- [7] R. Liu, S. Loeb, Y. Yin and G. Yu, DP-3-coloring of some planar graphs, Discrete Math. 342 (1) (2019) 178–189.
- [8] F. Lu, Q. Wang and T. Wang, Cover and variable degeneracy, Discrete Math. 345 (4) (2022) 112765.
- [9] P. Sittitrai and K. Nakprasit, Planar graphs without mutually adjacent 3-, 5-, and 6-cycles are 3-degenerate, Discrete Math. 345 (9) (2022) 112942.
- [10] Q. Wang, T. Wang and X. Yang, Variable degeneracy of graphs with restricted structures, arXiv:2112.09334, https://doi.org/10.48550/arXiv.2112.09334.
- [11] T. Wang, Weak degeneracy of planar graphs without 4- and 6-cycles, Discrete Appl. Math. 334 (2023) 110–118.
- [12] W. Wang and K.-W. Lih, Choosability and edge choosability of planar graphs without five cycles, Appl. Math. Lett. 15 (5) (2002) 561–565.