Emergence of Lorentz symmetry from an almost-commutative twisted spectral triple.

G. Nieuviarts¹

Abstract

This article demonstrates how the transition from a twisted spectral triple to a pseudo-Riemannian spectral triple arises within an almost-commutative spectral triple. This suggests that the noncommutative part itself is responsible for the signature change. This opens a new perspective on the signature problem, suggesting that the almost-commutative structure at the heart of the noncommutative standard model of particle physics could be the origin of the emergence of a Lorentzian spectral triple, starting from self-adjoint Dirac operators and conventional inner product structures, within the framework of twisted spectral triples.

•

Contents

4...

I	Intr	oduction	2		
2	Twisted spectral triple				
	2.1	Fundamentals of twisted spectral triple	4		
	2.2	From twisted to indefinite inner products	6		
	2.3	Twisted real structure	7		
3	Pseudo-Riemannian spectral triples				
	3.1	Clifford algebras for even-dimensional vector spaces and associated twist	8		
	3.2	From automorphisms of Clifford algebras to indefinite inner products	13		
	3.3	Even dimensional pseudo-Riemannian manifolds	17		
	3.4	Indefinite inner products and pseudo-Riemannian spectral triples	21		
4	Isomorphism between pseudo-Riemannian and twisted spectral triples				
	4.1	<i>K</i> -morphism of spectral triples	23		
	4.2	Signature change for even-dimensional manifolds	25		
5	K-morphism from an almost-commutative twisted spectral triple				
	5.1	Product of twisted spectral triples and spectral triple	28		
	5.2	<i>K</i> -morphism from the spectral triple product structure	30		
	5.3	Structure of the induced pseudo-Riemannian spectral triple	32		
6	Cor	clusion	37		

¹gaston.nieuviarts.wk@gmail.com

1 Introduction

Noncommutative geometry is a branch of mathematics that generalizes classical geometry to spaces where the algebra of functions is no longer commutative. Inspired by quantum mechanics, where observables (elements in an algebra) do not commute, this theory provides tools to describe "spaces" that are too irregular or singular to be treated with traditional differential geometry. One key realization of noncommutative geometry is its ability to recover Riemannian manifolds through spectral data. A. Connes, a pioneer of the field, developed the concept of spectral triples to encode the geometric information. A spectral triple ($\mathcal{A}, \mathcal{H}, D$) is the data of an involutive unital algebra \mathcal{A} represented by bounded operators on a Hilbert space \mathcal{H} together with a self-adjoint operator D acting on \mathcal{H} . In this framework, Riemannian geometry can be reconstructed from the spectral properties of the Dirac operator by starting from the spectral triple ($\mathcal{C}^{\infty}(\mathcal{M}), L^2(\mathcal{M}, S), D, J, \Gamma$), where $L^2(\mathcal{M}, S)$ is the Hilbert space of square integrable sections of the spinor bundle over the smooth manifold \mathcal{M} . The Dirac operator D is constructed using the Riemannian gamma matrices and the Levi-Civita spin connection. J is a given anti-unitary operator, and Γ is the grading operator.

In theoretical physics, fundamental interactions are described as Yang-Mills and Gravity theories, both based on the concept of connection on vector bundles. The extension of these concepts to noncommutative geometry relies on a crucial duality given by the Serre-Swan theorem. The gauge group emerges from inner automorphisms of \mathcal{A} . According to the noncommutative differential structure we choose, two main frameworks have been used to construct Noncommutative Gauge Field Theories. The first one (historically), proposed by M. Dubois-Violette, R. Kerner, and J. Madore in [15] concerns the use of derivation-based differential structure. The second one, mainly developed by A. Connes, J. Lott, and A. Chamseddine in [7, 5] concerns the use of spectral triples based differential structure. Let focus on this latter approach by considering a spectral triple ($\mathcal{A}, \mathcal{H}, D, J, \Gamma$) (see [11, 9] for the complete definition). The derivative is given by $\delta(.) = [D, .]$ so that oneforms are given by elements $A = \sum_{k} a_k [D, b_k]$ with $a_k, b_k \in A$. The connection with gauge theory goes as follows, if we consider a unitary $u \in \mathcal{U}(\mathcal{A})$ (the set of unitaries in \mathcal{A}), and defines the adjoint unitary $U = \pi(u)J\pi(u)J^{-1}$, then, considering a one-form ω as a gauge field and the associated "fluctuated" Dirac operator $D_{\omega} := D + \omega + \epsilon' J \omega J^{-1}$, inner fluctuations $(D_{\omega})^{u} := UD_{\omega}U^{\dagger}$ are then equivalent to gauge transformations $\omega^{u} := u\omega u^{\dagger} + u\delta(u^{\dagger})$ via the relation $(D_{\omega})^u = D_{\omega^u}$. The spectral action, a central spectral invariant, is given by $S_b[D] := \operatorname{Tr} f(DD^{\dagger}/\Lambda^2)$ (for more details, see [6]).

Based on this construction, we can consider a noncommutative extension of $(C^{\infty}(\mathcal{M}), L^2(\mathcal{M}, S), D, J, \Gamma)$ by taking the Almost-Commutative (AC) algebra $C^{\infty}(\mathcal{M}) \otimes \mathcal{A}_F$ with \mathcal{A}_F a finite dimensional noncommutative algebra. The Dirac operator is $D_{\mathcal{M}} \otimes \mathbb{1} + \Gamma \otimes D_F$, with Hilbert space $L^2(\mathcal{M}, S) \otimes \mathcal{H}_F$, real structure $J \otimes J_F$ and grading $\Gamma \otimes \Gamma_F$. A fundamental aspect of this AC manifold is that its spectral action corresponds to the Lagrangian of an Einstein-Yang-Mills-Higgs model, entirely determined by the product spectral triple. Driven by the constraints coming with the axioms of even real spectral triples, a reformulation of the Standard Model of Particle Physics coupled to gravitation was elaborated, with crucial steps in [10, 5] until its full development in [6] (we refer to [11] for detailed explanations). At the heart of this construction lies the finite geometry encoded in $\mathcal{A}_F := \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ and its associated spectral triple. The symmetry group of the spectral action is given by $\mathcal{G} = Map(\mathcal{M}, G) \rtimes Diff(\mathcal{M})$, where Diff(\mathcal{M}) denotes the diffeomorphism group, and $Map(\mathcal{M}, G)$ represents the gauge group of the second kind,

with $G = U(1) \times SU(2) \times SU(3)$. In this way, noncommutative geometry provides a unified framework to describe both Einstein-Hilbert gravity (in Euclidean signature) and classical gauge theories. This framework is known as the Non-Commutative Standard Model. Notably, it offers an elegant description of the Standard Model, including the Higgs mechanism and neutrino mixing, as "gravity" on an AC-manifold. The fermionic masses are encoded into D_F , so that the masses of the Higgs boson and the ones of fermions became related, thus providing a prediction for the Higgs mass.

The noncommutative approach links geometry to operator algebras, revealing new mathematical and physical insights. But it falls short when describing spacetimes with pseudo-Riemannian signatures that are essential in physics. Indeed, realistic models require a formulation that incorporates time and causality, motivating the search for Lorentzian approaches within noncommutative geometry. This issue has spurred various proposals, from modified axioms for spectral triples to alternative formulations that attempt to reconcile causality with noncommutative structures see [1, 17, 23, 28, 3, 14, 1, 29, 2, 16]. An essential feature of most of these approaches is the replacement of the Hilbert space by a Krein space, turning the inner product into an indefinite inner product $\langle ., . \rangle_K := \langle ., K. \rangle$ with the unitary operator $K = K^{\dagger}$. Such structures are essential in quantum field theory, as a physical description of spinor fields on pseudo-Riemannian manifolds inherently requires Krein spaces. However, developing a fully satisfactory and widely accepted framework for Lorentzian noncommutative geometry remains an open and actively pursued area of research.

In [22], I present a connection between twisted spectral triples and pseudo-Riemannian spectral triples based on a fundamental link between the previous Krein product and the twist defined by $\rho(.) := K(.)K$. A concept of spectral triple morphism was introduced, enabling the transformation of a twisted spectral triple into a pseudo-Riemannian one. This morphism was shown to induce a signature change in the context of the manifold's spectral triples. The signature change transformation was determined solely by the unitary operator K, being the central element of the approach.

The following article proposes to extend these results. The first three sections introduce what are twisted spectral triples, pseudo-Riemannian spectral triples, and the *K*-morphism between them. One of the main contributions is the definition of a specific real structure for twisted spectral triples in subsection 2.3, adapted to the bimodule structure associated with the derivations and related to the special *K*-unitary operators which generates the twisted fluctuations. Interestingly, this real structure also emerges in the setting of pseudo-Riemannian spectral triples and remains invariant under the *K*morphism, thus retaining its significance in both frameworks.

Another important point was the introduction of the twist in the context of Clifford algebras (for even-dimensional vector spaces) in subsection 3.1, showing how this twist played a crucial role in the structure, in particular to characterise the orthochronous spin group and to introduce the Krein product $\langle . , . \rangle_K$, of particular importance in physics.

The main results are presented in section 5. An almost-commutative spectral triple is constructed on a twisted spectral triple, naturally inducing the emergence of the operator K in the product Dirac operator. This leads to the appearence of a Krein structure together with the pseudo-Riemannian Dirac operators, from the very first spectral triple structure, then implementing the K-morphism. We then show how the specific signature of space-time is recovered in the 4-dimensional case. This section follows a top-down approach, revealing the signature change as an intrinsic outcome of the algebraic structure governing the almost-commutative spectral triple.

2 Twisted spectral triple

Spectral triples have been particularly successful in describing noncommutative spaces arising from type I and type II von Neumann algebras, where a well-defined trace exists. However, a major challenge in noncommutative geometry has been the extension of these methods to type III von Neumann algebras, which lack such a finite normal trace. Type III algebras arise naturally in quantum field theory, especially in the algebraic approach to quantum statistical mechanics and the study of vacuum states in curved spacetime.

To address this challenge, twisted spectral triples were introduced by A. Connes and H. Moscovici in their 2006 work, see [12]. These modified spectral triples generalize the standard construction by incorporating an automorphism, which compensates for the absence of a trace and allows for the formulation of geometric invariants in this setting.

Interestingly, the introduction of this twist is related to the modular theory of Tomita-Takesaki, which governs the structure of von Neumann algebras. Tomita-Takesaki theory links the modular structure of von Neumann algebras to time evolution, defining a natural flow on the algebra. Twisted spectral triples were then implemented in the context of the noncommutative Standard Model in [13, 18, 19]. Applications to conformal and noncommutative geometry can also be found in [24, 25].

2.1 Fundamentals of twisted spectral triple

Let \mathcal{A} be a *-algebra with representation π on a Hilbert space \mathcal{H} . The faithful representation π will be omitted by identifying the elements of \mathcal{A} with their representation. The adjoint \dagger in $\mathcal{B}(\mathcal{H})$ is then given by the relation $\pi(a^*) = \pi(a)^{\dagger}$.

Definition 2.1 (Twisted spectral triple) A twisted spectral triple $(\mathcal{A}, \mathcal{H}, D, \rho)$ is obtained by considering an involutive algebra \mathcal{A} acting faithfully as a bounded operator algebra on a Hilbert space \mathcal{H} with a self-adjoint operator D with compact resolvent together with an automorphism $\rho \in \text{Aut}(\mathcal{A})$ called twist which satisfy the regularity condition

$$\rho(a^{\dagger}) = (\rho^{-1}(a))^{\dagger} \qquad \forall a \in \mathcal{A}.$$
(2.1)

The twisted commutator

$$[D,a]_{\rho} := Da - \rho(a)D \tag{2.2}$$

is used instead of [D, a] and we require only the twisted one to be bounded.

In the same way as for spectral triples, we can add a grading Γ and a real structure *J*. The axioms of such even real twisted spectral triples are the same as for spectral triples

$$J^{2} = \epsilon_{0} \qquad JD = \epsilon_{1}DJ \qquad J\Gamma = \epsilon_{2}\Gamma J \qquad \epsilon_{0}, \epsilon_{1}, \epsilon_{2} = \pm 1$$
(2.3)

together with the introduction of another characteristic number $\epsilon = \pm 1$ such that

$$\rho(J) = \epsilon J \tag{2.4}$$

and the replacement of the first order condition by the twisted first order condition

$$[[D, a]_{\rho}, b^{\circ}]_{\rho^{\circ}} = 0$$
(2.5)

 $\forall a \in \mathcal{A} \text{ and } b^{\circ} \in \mathcal{A}^{\circ} \text{ with } \mathcal{A}^{\circ} \text{ the opposite algebra. The opposite twist } \rho^{\circ} \text{ acting on } \mathcal{A}^{\circ} \text{ is defined by}$

$$\rho^{\circ}(a^{\circ}) := (\rho^{-1}(a))^{\circ}.$$
(2.6)

See [18] for further details.

The set of twisted 1-forms $\Omega_D^1(\mathcal{A}, \rho)$ is defined as

$$\Omega_D^1(\mathcal{A},\rho) := \{ \sum_k a_k [D, b_k]_\rho, \ a_k, b_k \in \mathcal{A} \}$$

$$(2.7)$$

so that twisted-fluctuations of *D* are given by

$$D_{A_{\rho}} := D + A_{\rho} + \epsilon_1 J A_{\rho} J^{-1} \qquad \text{with} \qquad A_{\rho} \in \Omega_D^1(\mathcal{A}, \rho).$$
(2.8)

The set $\Omega^1_D(\mathcal{A}, \rho)$ is a \mathcal{A} -bimodule for the following product

$$a \cdot A_{\rho} \cdot b = \rho(a)A_{\rho}b \tag{2.9}$$

for any $a, b \in A$ and $A_{\rho} \in \Omega^{1}_{D}(A, \rho)$. This make the map $a \to \delta_{\rho}(a) := [D, a]_{\rho}$ be a derivation since we will have that

$$\delta_{\rho}(ab) = \delta_{\rho}(a)b + \rho(a)\delta_{\rho}(b) = \delta_{\rho}(a) \cdot b + a \cdot \delta_{\rho}(b).$$
(2.10)

As shown in [18] a way to "minimaly twist" the spectral triple $(\mathcal{A}, \mathcal{H}, D, \Gamma)$ i.e. without changing the Hilbert space and the Dirac operator, is by the use of the decomposition coming with Γ . Using the projections $p_{\pm} := (\mathbb{1} \pm \Gamma)/2$ so that $\mathcal{H} = p_{+}\mathcal{H} \oplus p_{-}\mathcal{H} :=$ $\mathcal{H}_{+} \oplus \mathcal{H}_{-}$, all elements $a = (a_{1}, a_{2}) \in \mathcal{A} \otimes \mathbb{C}^{2}$ are represented accordingly:

$$\pi(a_1, a_2) := p_+ \pi_0(a_1) + p_- \pi_0(a_2) = \begin{pmatrix} \pi_+(a_1) & 0\\ 0 & \pi_-(a_2) \end{pmatrix}$$
(2.11)

with π_0 the representation used for \mathcal{A} and $\pi_{\pm}(.) := p_{\pm}\pi_0(.)_{|\mathcal{H}_{\pm}}$ the restrictions to \mathcal{H}_{\pm} . This will be called the chiral representation.

Definition 2.2 (twist by grading) Taking the graded spectral triple $(\mathcal{A}, \mathcal{H}, D, \Gamma)$, its corresponding twist by gradding is given by the twisted spectral triple $(\mathcal{A} \otimes \mathbb{C}^2, \mathcal{H}, D, \Gamma, \rho)$, using the chiral representation. The twist acts on elements $a = (a_1, a_2) \in \mathcal{A} \otimes \mathbb{C}^2$ as

$$\rho(a_1, a_2) = (a_2, a_1). \tag{2.12}$$

As shown in [18], the twist by the grading is the only way to minimally twist the spectral triple of even-dimensional manifolds based on $C^{\infty}(\mathcal{M}) \otimes \mathbb{C}^2$. The fact that no such procedure for twisting odd-dimensional manifold's spectral triple have been found will be one of the justifications to focus on the study of even-dimensional manifolds. More information on twisted spectral triples can be found in [18, 19, 21].

2.2 From twisted to indefinite inner products

Following [22], any automorphism ρ will be required to be $\mathcal{B}(\mathcal{H})$ -regular, i.e., a regular automorphism on all of $\mathcal{B}(\mathcal{H})$. Since [14], a novelty consists in the possibility of considering a new inner product $\langle . , . \rangle_{\rho}$, called the ρ -inner product, which by definition satisfies

$$\langle \psi, O\psi' \rangle_{\rho} = \langle \rho(O)^{\dagger}\psi, \psi' \rangle_{\rho} \tag{2.13}$$

for all $\psi, \psi' \in \mathcal{H}$ and $O \in \mathcal{B}(\mathcal{H})$.

This new inner product naturally comes with its own notion of adjoint, denoted by "+", defined as $(.)^+ := \rho(.)^\dagger$, along with an associated notion of ρ -unitarity, characterized by elements U_ρ satisfying $U_\rho U_\rho^+ = U_\rho^+ U_\rho = \mathbb{1}$, thereby preserving the ρ -product. As mentioned in [22], the uniqueness of this adjoint (i.e., the equality of the left and right adjoint) is ensured by the $\mathcal{B}(\mathcal{H})$ -regularity condition. We will restrict ourselves to twists that are *-automorphisms, i.e., $\rho(O^\dagger) = \rho(O)^\dagger$. This, together with the regularity condition, implies that $\rho^2 = \mathbb{1}$, which in turn ensures that $(.)^+$ is an involution.

As shown in [19] and [21], there are two ways to generate the twisted fluctuations of equation (2.8), using either usual unitary operators or ρ -unitary operators:

1. Type 1: $D_{A_{\rho}} = UDU^+$ with $U = uJuJ^{-1}$ and u a unitary operator in \mathcal{A} .

2. Type 2: $D_{A_{\rho}} = U_{\rho}DU_{\rho}^{\dagger}$ with $U_{\rho} = u_{\rho}Ju_{\rho}J^{-1}$ and u_{ρ} a ρ -unitary operator in \mathcal{A} .

This justifies the introduction of the notions of ρ -product, along with the adjoint "+" and the notion of ρ -unitarity.

In the case where $\rho \in \text{Inn}(\mathcal{B}(\mathcal{H}))$, there exists a unitary operator *K* acting on \mathcal{H} such that $\rho(O) = KOK^{\dagger}$. In this case, the ρ -product is determined by *K*, so that $\langle ., . \rangle_{\rho} := \langle ., K. \rangle$. Furthermore, the twist is a *-automorphism of \mathcal{A} .

In the following, any twist ρ will be assumed to be an element of $\text{Inn}(\mathcal{B}(\mathcal{H}))$. Since different operators *K* can implement the same twist ρ while inducing different inner products, we will adopt a more precise notation. The twisted spectral triple will be denoted by $(\mathcal{A}, \mathcal{H}, D, K)$, and the ρ -inner product notation $\langle ., . \rangle_{\rho}$ will be replaced by the *K*-inner product notation:

$$\langle \, . \, , \, . \, \rangle_K := \langle \, . \, , K . \, \rangle \tag{2.14}$$

so that the associated adjoint is called the *K*-adjoint and is defined by

$$(\,.\,)^{\dagger_{\mathcal{K}}} := \rho(\,.\,)^{\dagger}. \tag{2.15}$$

We define *K*-unitary operators by the relation $U_K U_K^{\dagger_K} = U_K^{\dagger_K} U_K = 1$, with the corresponding set denoted as $\mathcal{U}_K(\mathcal{B}(\mathcal{H}))$.

Taking the automorphism $\rho(.) = K(.)K^{\dagger}$ to be $\mathcal{B}(\mathcal{H})$ -regular, or regular on \mathcal{A} if the algebra is a simple finite-dimensional algebra, we obtain

$$K = \exp(i\theta)K^{\dagger} \tag{2.16}$$

for some real number θ . As a consequence, $\langle ., . \rangle_K$ is Hermitian and indefinite if and only if $K = K^{\dagger}$ (see [22] for more details). The condition $K = K^{\dagger}$ will be derived from algebraic constraints in the top-down approach presented in section 5.

Definition 2.3 (Fundamental twist) A fundamental twist is a twist $\rho \in \text{Inn}(\mathcal{B}(\mathcal{H}))$, implemented by the unitary K such that $K = K^{\dagger}$.

2.3 Twisted real structure

In non-commutative geometry, the real structure associated with the operator J is used to relate left and right actions on A-bimodules. We have seen in equation (2.9) that $\Omega_D^1(A, \rho)$ is a A-bimodule whose left and right actions are connected by the twist ρ . Here, I propose to introduce a notion of real structure that implements this connection. This particular real structure will play a crucial role throughout the article, appearing notably in the context of Clifford algebras with indefinite (pseudo-Riemannian) metrics in subsection 3.1, and then in the framework of pseudo-Riemannian spectral triples in subsection 3.4.

Taking $a \in A$, the left action of the algebra on bimodules is denoted by l(a). This operator satisfies $l(a^{\dagger}) = l(a)^{\dagger}$. The corresponding right action r(a) is given through J by

$$r(a) = Jl(a)^{\dagger}J^{-1},$$
 (2.17)

with the zero-order condition [l(a), r(a)] = 0, ensuring that the right and left actions on the A-bimodules are connected by the relation

$$a_{.}(.)_{.}b = l(a)r(b)(.).$$
 (2.18)

For the usual action on a bimodule, we have $a \cdot (\cdot) \cdot b = a(\cdot)b$, meaning that the left and right actions correspond to the usual multiplication of operators acting on elements of the bimodule. We denote by \hat{J} the corresponding real structure, i.e., $\hat{l}(a) = a$ and $\hat{r}(a) = \hat{J}\hat{l}(a)^{\dagger}\hat{J}^{-1}$. We define $\hat{r}(a) = a^{\circ} = \hat{J}a^{\dagger}\hat{J}^{-1}$ as the element of the corresponding opposite algebra \mathcal{A}° . This is the real structure used for conventional spectral triples.

Now, to recover the twisted product action (2.9), i.e.,

$$a.(.).b = \rho(a)(.)b,$$
 (2.19)

we have to find *J* such that $l(a) = Jr(a)^{\dagger}J^{-1} = \rho(a)$ with $r(a) = \hat{r}(a) = a$. Noting that

$$l(a) = \rho(\hat{l}(a)) = \rho(\hat{f}(a)^{\dagger}\hat{f}^{-1}) = \rho(\hat{f}r(a)^{\dagger}\hat{f}^{-1}), \qquad (2.20)$$

we can define¹

$$J \mathrel{\mathop:}= KJ \tag{2.21}$$

so that $l(a) = Jr(a)^{\dagger}J^{-1} = \rho(a)$. We define $r(a) = a^{\circ} = \rho(a^{\circ}) = Ja^{\dagger}J^{-1}$ as the element of the corresponding opposite algebra \mathcal{A}° .

In subsection 2.2, we highlight the important role the *K*-unitary operators associated with $\langle ., . \rangle_K$ play in the context of twisted spectral triples. Given any $(\psi, U_K) \in (\mathcal{H}, \mathcal{A})$ with U_K a *K*-unitary operator, then the adjoint action on the bimodule is given by

$$\psi \to U_K \psi U_K^{-1} = U_K \psi U_K^{\dagger_K} = U_K J U_K J^{-1} \psi$$
(2.22)

showing that the adjoint action of K-unitary operators is intrinsically connected with the operator J defining the twisted real structure.

Spectral triples correspond to the special case $K = \mathbb{1}$. The twisted commutator then reduces to the usual one, $U_K = U$ is unitary for $\langle ., . \rangle$, and $J = \hat{J}$, so that we recover the usual adjoint action on the bimodule:

$$\psi \to U\psi U^{-1} = U\psi U^{\dagger} = U\hat{J}U\hat{J}^{-1}\psi.$$
(2.23)

¹In full generality, we may have $J := \alpha K \hat{J}$ with $\alpha \in \mathbb{C}$ such that $\alpha \alpha^* = 1$, but this phase shift is set to 1 to avoid unnecessary terms in the expressions.

3 Pseudo-Riemannian spectral triples

The following subsections aims at introducing Clifford algebras with the so-called three structural automorphisms, their properties and interrelations. These automorphisms are of great importance to characterize Clifford algebras and their automorphisms. The main novelty consists in the introduction of the twist (satisfying the $\mathcal{B}(\mathcal{H})$ -regularity condition; see Proposition 3.6), being one of these automorphisms, deeply connected to the metric and providing the parity operator in the case the chosen basis is constituted of unitary operators. This twist will also plays a significant role in subsection 3.2, to characterize an important class of automorphisms, which holds fundamental importance in physics, i.e., the orthochronous spin group and the associated indefinite inner product. The author has followed similar conventions [20].

3.1 Clifford algebras for even-dimensional vector spaces and associated twist

Let $V \simeq \mathbb{R}^{2m}$ be an even-dimensional (real) vector space over \mathbb{R} of dimension 2m, with $m \in \mathbb{N}$. Taking any vector $v \in V$, a complex linear representation of v is provided by the Clifford multiplication operator c(v) as a $2^m \times 2^m$ complex square matrix. A non-degenerate symmetric bilinear form g on V determines a metric with signature (n, 2m - n), where $n \in \mathbb{N}$, subject to the relation

$$c(u)c(v) + c(v)c(u) = 2g(u,v)\mathbb{1}_{2^m} \qquad \forall u, v \in V.$$
 (3.1)

Unless explicitly mentioned, we will identify elements $v \in V$ with their representation c(v) in what follows, thereby identifying the vector space *V* with its representation.

The (complex) Clifford algebra Cl(V, g) is the algebra generated by the vector space $V \subset Cl(V, g)$ and the identity $\mathbb{1}_{2^m} \subset Cl(V, g)$.

Automorphisms Aut(Cl(V, g)) of the Clifford algebra are bijective maps ϕ : $Cl(V, g) \rightarrow Cl(V, g)$ that preserve the defining relation (3.1). We denote by Aut(V) the automorphism group of the vector space V. Since Cl(V, g) has even dimension, it is isomorphic to $M_{2^m}(\mathbb{C})$, implying that all its automorphisms are inner, i.e., Aut(Cl(V, g)) = Inn(Cl(V, g)). Defining $Cl^*(V, g)$ as the subset of invertible elements in Cl(V, g), then every automorphism of Cl(V, g) can be characterized by an element $x \in Cl^*(V, g)$ through the adjoint action

$$\phi_x(y) := \operatorname{ad}_x(y) = xyx^{-1} \qquad \forall y \in Cl(V,g).$$
(3.2)

Let $\{e_a\}_{1 \le a \le 2m}$ be a *g*-orthonormal basis of *V*, i.e., subject to the relation

$$e_{a}e_{b} + e_{b}e_{a} = 2g(e_{a}, e_{b})\mathbb{1}_{2^{m}} = \begin{cases} 2\delta_{ab}\mathbb{1}_{2^{m}} & \text{for } 1 \le a \le n, \\ -2\delta_{ab}\mathbb{1}_{2^{m}} & \text{for } n+1 \le a \le 2m. \end{cases}$$
(3.3)

Any vector $v \in V$ can be written as $v = \sum_{a=1}^{2m} \lambda_a e_a$ with $\lambda_a \in \mathbb{R}$. Cl(V, g) is then generated by $\{e_a\}_{1 \leq a \leq 2m}$ and $\mathbb{1}_{2^m}$, so that a basis of $Cl(V, g) \simeq M_{2^m}(\mathbb{C})$ is given by the set

$$\mathcal{I} := \{\mathbb{1}_{2^m}, \{e_a\}_{1 \le a \le 2m}, \{e_{a_1}e_{a_2}\}_{1 \le a_1 < a_2 \le 2m}, \{e_{a_1}e_{a_2}e_{a_3}\}_{1 \le a_1 < a_2 < a_3 \le 2m}, \dots, e_1e_2 \dots e_{2m}\}.$$
(3.4)

Let define $g_v := g(v, v)$ and $g_a := g(e_a, e_a)$. The metric g induces the splitting $V = V^+ \oplus V^-$, where the set $\{e_a\}_{1 \le a \le n}$ is a basis of V^+ and the set $\{e_a\}_{n+1 \le a \le 2m}$ is a basis of V^- , with $g_{v^+} \ge 0$ and $g_{v^-} \le 0$ for $v^{\pm} \in V^{\pm}$. An irreducible representation of $Cl(V, g) \simeq M_{2^m}(\mathbb{C})$ is given by the module $M = \mathbb{C}^{2^m}$. A Hermitian inner product $\langle ., . \rangle$ can be associated with M, with adjoint denoted by \dagger . We define the Hilbert space $\mathcal{H}_m := (M, \langle ., . \rangle)$.

The commutant V' of V in Cl(V, g) is defined by

$$V' := \{a \in Cl(V,g) \mid \forall v \in V, \quad va = av\}.$$

$$(3.5)$$

Proposition 3.1 We have that $V' = Z(Cl(V, g)) = \mathbb{C}\mathbb{1}_{2^m}$.

PROOF Since $Cl(V, g) \simeq M_{2^m}(\mathbb{C})$, its center is $\mathbb{C}\mathbb{1}_{2^m}$. Now, taking any element $a \in Cl(V, g)$, it is a linear combination of elements in \mathcal{I} . The requirement [a, v] = 0 for any $v \in V$ implies that $a \in \mathbb{C}\mathbb{1}_{2^m}$, since for any element $e_{a_1}e_{a_2} \dots e_{a_k} \in \mathcal{I}$ with $1 \leq a_1 < a_2 < \dots < a_k \leq 2m$, there exists an element $e_a \in V$ that anticommutes with it, meaning that $V' = \mathbb{C}\mathbb{1}_{2^m}$.

The complex conjugation operator is defined on any operator $O \in \mathcal{B}(\mathcal{H})$ by

$$\overline{O} = cc \circ O \circ cc, \tag{3.6}$$

where $cc = cc^{-1}$ is the complex conjugation over \mathcal{H} .

Definition 3.2 The structural operators ρ , χ , κ , called twist, grading, and charge conjugation, are defined to be elements in Aut(Cl(V, g)) so that $\forall v \in V$ we have

$$\rho(v) := v^{\dagger}, \qquad \chi(v) := -v, \qquad \kappa(v) := -\bar{v}.$$
(3.7)

Remark 3.3 If we do not restrict to even-dimensional vector spaces *V*, the relation for κ is more generally given by

$$\kappa(v) := -\epsilon_1^K \bar{v},\tag{3.8}$$

where $\epsilon_1^K = \pm 1$. As shown in the KO-dimension table 1, a particular feature of the evendimensional case is that $\epsilon_1^K = 1$ in all cases, so the definition without this factor.

Proposition 3.4 The structural automorphisms ρ , χ , and κ are involutions on Cl(V, g).

PROOF Any element of Cl(V, g) can be expressed as a linear combination of elements in \mathcal{I} . We can directly check that

$$\rho^{2}(\alpha \mathbb{1}_{2^{m}}) = \chi^{2}(\alpha \mathbb{1}_{2^{m}}) = \kappa^{2}(\alpha \mathbb{1}_{2^{m}}) = \alpha \mathbb{1}_{2^{m}}, \quad \forall \alpha \in \mathbb{C},$$

$$(3.9)$$

and that for any element $\alpha e_{a_1} e_{a_2} \dots e_{a_k}$ with $k \in \mathbb{N}$, the same relation holds:

$$\rho^{2}(\alpha e_{a_{1}}e_{a_{2}}\dots e_{a_{k}}) = \alpha \rho(e_{a_{1}}^{\dagger}e_{a_{2}}^{\dagger}\dots e_{a_{k}}^{\dagger}) = \alpha e_{a_{1}}e_{a_{2}}\dots e_{a_{k}},$$
(3.10)

$$\chi^{2}(\alpha e_{a_{1}}e_{a_{2}}\dots e_{a_{k}}) = (-1)^{2k}\alpha e_{a_{1}}e_{a_{2}}\dots e_{a_{k}} = \alpha e_{a_{1}}e_{a_{2}}\dots e_{a_{k}},$$
(3.11)

$$\kappa^{2}(\alpha e_{a_{1}}e_{a_{2}}\dots e_{a_{k}}) = (-1)^{2k}\alpha e_{a_{1}}e_{a_{2}}\dots e_{a_{k}} = \alpha e_{a_{1}}e_{a_{2}}\dots e_{a_{k}}.$$
(3.12)

This implies that $\rho^2 = \chi^2 = \kappa^2 = \mathbb{1}_{2^m}$ on any element of \mathcal{I} , and thus on all of Cl(V, g).

This implies the existence of positive and negative eigenspaces in each case

$$Cl(V,g) = Cl^{+}(V,g) \oplus Cl^{-}(V,g), \quad \text{with} \quad \rho(a^{\pm}) = \pm a^{\pm}, \quad a^{\pm} \in Cl^{\pm}(V,g), \quad (3.13)$$

$$Cl(V,g) = Cl^{0}(V,g) \oplus Cl^{1}(V,g), \quad \text{with} \quad \gamma(a^{i}) = (-1)^{i}a^{i}, \quad a^{i} \in Cl^{i}(V,g), \quad (3.14)$$

$$Cl(V, g) = Cl^{\bar{0}}(V, g) \oplus Cl^{\bar{1}}(V, g), \quad \text{with} \quad \chi(a) = (-1)^{i}a^{i}, \quad a^{i} \in Cl^{\bar{i}}(V, g), \quad (3.14)$$
$$Cl(V, g) = Cl^{\bar{0}}(V, g) \oplus Cl^{\bar{1}}(V, g), \quad \text{with} \quad \kappa(a^{i}) = (-1)^{i}a^{i}, \quad a^{i} \in Cl^{\bar{i}}(V, g). \quad (3.15)$$

$$Cl(V,g) = Cl^{0}(V,g) \oplus Cl^{1}(V,g), \text{ with } \kappa(a^{i}) = (-1)^{i}a^{i}, a^{i} \in Cl^{i}(V,g).$$
 (3.15)

As a consequence, the structural automorphisms ρ, χ , and κ are also elements of Aut(V), as their action is linear (multiplication by ± 1 according to the corresponding eigenspace) on V in the corresponding decompositions of Cl(V, g). This notably implies that \bar{v}, v^{\dagger} , and $v^{\top} = \bar{v}^{\dagger}$ are elements of *V* for any $v \in V$, and consequently that

$$\rho(\bar{v}) = v^{\top} = \overline{\rho(v)}. \tag{3.16}$$

We then have the equivalent definition $\kappa(v) = \chi(\bar{v})$.

Proposition 3.5 We have $\rho \circ \chi = \chi \circ \rho$, $\rho \circ \kappa = \kappa \circ \rho$, and $\kappa \circ \chi = \chi \circ \kappa$ on Cl(V, g).

PROOF These relations are easily verified on the first element $\alpha \mathbb{1}_{2^m}$ of \mathcal{I} and on any element $v \in V$, since

$$\rho(\chi(v)) = -v^{\dagger} = \chi(\rho(v)), \tag{3.17}$$

$$\rho(\kappa(v)) = -v^{\top} = \kappa(\rho(v)), \tag{3.18}$$

$$\kappa(\chi(v)) = \bar{v} = \chi(\kappa(v)). \tag{3.19}$$

Taking any element $a = \alpha e_{a_1} e_{a_2} \dots e_{a_k}$ with $k \in \mathbb{N}$, we obtain the same equalities, as the operators commute with each $e_{a_i} \in V$, ensuring that the relations hold on \mathcal{I} , then extend to all of Cl(V, g).

The following proposition justifies the choice of notation ρ by establishing that it fulfills the defining properties of a twist i.e., satisfying the regularity condition.

Proposition 3.6 The automorphism ρ is $\mathcal{B}(\mathcal{H})$ -regular.

PROOF Since $\rho^2 = \mathbb{1}_{2^m}$ on all of Cl(V, g), we have $\rho^{-1} = \rho$. From the definition of ρ , taking any element $e_a \in V$, we obtain

$$\rho(e_a^{\dagger}) = e_a = \rho(e_a)^{\dagger}, \tag{3.20}$$

which implies that ρ satisfies

.

$$\rho(e_a^{\dagger}) = \rho^{-1}(e_a)^{\dagger}. \tag{3.21}$$

Thus, ρ is regular on V.

We recover this relation for any element of \mathcal{I} , since $\rho(\mathbb{1}_{2^m}^{\dagger}) = \rho^{-1}(\mathbb{1}_{2^m})^{\dagger}$, and for any $\alpha \in \mathbb{C}$ and $k \in \mathbb{N}$, we have

$$\rho((\alpha e_{a_1} e_{a_2} \dots e_{a_k})^{\dagger}) = \alpha^* \rho(e_{a_k}^{\dagger}) \dots \rho(e_{a_1}^{\dagger})$$

= $\alpha^* \rho^{-1}(e_{a_k})^{\dagger} \dots \rho^{-1}(e_{a_1})^{\dagger}$
= $\alpha^* \rho^{-1}(e_{a_1} \dots e_{a_k})^{\dagger} = \rho^{-1}(\alpha e_{a_1} e_{a_2} \dots e_{a_k})^{\dagger}$

By linearity, this extends to all of $Cl(V, g) \simeq M_{2^m}(\mathbb{C}) \simeq \mathcal{B}(\mathcal{H})$, proving that ρ is regular.

We thus recover an automorphism with the properties of a twist, satisfying the regularity condition defined in section 2, so the name twist for this operator.

Since Aut(Cl(V, g)) = Inn(Cl(V, g)), there exist operators $K, \Gamma, C \in Cl(V, g)$ such that

$$\rho(.) = K(.)K^{-1}, \qquad \chi(.) = \Gamma(.)\Gamma^{-1}, \qquad \kappa(.) = C(.)C^{-1}.$$
(3.22)

We require these operators to be unitary, ensuring that the structural automorphisms are *-automorphisms.

The real operator is defined from *C* by

$$J := C \circ cc, \tag{3.23}$$

so that the charge conjugation transformation is given by $\psi \rightarrow \psi_c := J\psi$ with

$$Jv = -vJ \qquad \forall v \in V. \tag{3.24}$$

From Proposition 3.6, we retrieve the result of equation (2.16), namely $K = \exp(i\theta)K^{\dagger}$, since ρ is $\mathcal{B}(\mathcal{H})$ -regular. Additionally, we have:

Proposition 3.7 We have that $\overline{K} = \exp(i\theta')K$ with $\theta' \in \mathbb{R}$.

PROOF The relation $\rho(\bar{v}) = \overline{\rho(v)}$ implies that

$$K\bar{v}K^{\dagger} = \overline{K}\bar{v}K^{\top}.$$
(3.25)

Equivalently, we write

. —

$$K^{\mathsf{T}}K\bar{v}K^{\mathsf{T}}K = \bar{v}, \quad \forall \bar{v} \in V.$$
(3.26)

This implies that $K^{\dagger}\overline{K}$ belongs to the commutant V' and is therefore of the form $\alpha \mathbb{1}_{2^m}$ with $\alpha \in \mathbb{C}$. Multiplying on the left by K, we obtain

$$\overline{K} = \alpha K. \tag{3.27}$$

Substituting into the previous expression, we find

$$K\bar{v}K^{\dagger} = \overline{K}\bar{v}K^{\dagger} = \alpha\alpha^*K\bar{v}K^{\dagger}, \qquad (3.28)$$

which requires $\alpha \alpha^* = 1$, meaning that $\alpha = \exp(i\theta')$ with $\theta' \in \mathbb{R}$.

From now on, we impose $\overline{K} = \pm K$.

Proposition 3.8 We have that

$$K\Gamma = \exp(i\theta_1)\Gamma K, \quad KC = \exp(i\theta_2)CK, \quad and \quad \Gamma C = \exp(i\theta_3)C\Gamma,$$
 (3.29)

where $\theta_i \in \mathbb{R}$ for $i \in \{1, 2, 3\}$.

PROOF From Proposition 3.5, taking any element $a \in Cl(V, g)$, we have

$$K\Gamma a(K\Gamma)^{\dagger} = \Gamma K a(\Gamma K)^{\dagger}, \quad KCa(KC)^{\dagger} = CKa(CK)^{\dagger}, \quad \Gamma Ca(\Gamma C)^{\dagger} = C\Gamma a(C\Gamma)^{\dagger}.$$
 (3.30)

This implies that $K^{-1}\Gamma^{-1}K\Gamma$, $K^{-1}C^{-1}KC$, and $\Gamma^{-1}C^{-1}\Gamma C$ belong to the center Z(Cl(V, g)), and are therefore of the form $\alpha \mathbb{1}_{2^m}$ with $\alpha \in \mathbb{C}$. As a consequence, we obtain

$$K\Gamma = \alpha_1 \Gamma K, \quad KC = \alpha_2 CK, \quad \text{and} \quad \Gamma C = \alpha_3 C\Gamma,$$
(3.31)

where $\alpha_i \in \mathbb{C}$ for $i \in \{1, 2, 3\}$. Substituting these expressions back into equation (3.30), we find that $\alpha_i \alpha_i^* = 1$, hence the result.

To prevent unnecessary complex factors, we impose $\exp(i\theta_i) = \pm 1$ for $i \in \{1, 2, 3\}$.

From now on, we assume the chosen representation of e_a to be unitary² with respect to the inner product $\langle ., . \rangle$. This, in particular implies that $e_a^{\dagger} = g_a e_a$. Consequently, the vector spaces $V^{\pm} \subset Cl^{\pm}(V, g)$ induced by ρ are also those induced by the metric, as

$$\rho(e_a) = e_a^{\dagger} = g_a e_a = \pm e_a \qquad \text{for} \qquad e_a \in V^{\pm}, \tag{3.32}$$

hence the notation $Cl^{\pm}(V, g)$ for the splitting induced by ρ . The twist then reverses the components associated with negative metrics. The automorphism ρ thus corresponds to the parity operator, a point that will be further emphasized in Subsection 4.2.

Remark 3.9 In the case of a Riemannian metric, i.e., n = 2m, we have $e_a^{\dagger} = g_a e_a = e_a$, so that $v = v^{\dagger}$, meaning that we can take $K = \mathbb{1}_{2^m}$.

Let \hat{V} be a vector space corresponding to a positive definite metric, such that the orthonormal basis $\{\hat{e}_a\}_{1 \le a \le 2m}$ for \hat{V} is connected to the one of V by a Wick rotation:

$$\hat{e}_a = e_a \qquad \qquad \text{for } 1 \le a \le n, \\ \hat{e}_a = ie_a \qquad \qquad \text{for } n+1 \le a \le 2m.$$

The unitarity of the e_a 's implies the unitarity of the \hat{e}_a 's. The corresponding real operator is denoted by $\hat{J} = \hat{C} \circ cc^3$, with $\hat{\kappa}(\hat{v}) = -\bar{v}$ for any $\hat{v} \in \hat{V}$.

Proposition 3.10 We have $\kappa = \hat{\kappa} \circ \rho = \rho \circ \hat{\kappa}$ on Cl(V, g).

PROOF Taking any element e_a of the basis of *V* we have

$$\begin{aligned} \hat{\kappa}(\rho(e_a)) &= \hat{\kappa}(e_a) = \hat{\kappa}(\hat{e}_a) = -\bar{e}_a & \text{for } 1 \le a \le n, \\ \hat{\kappa}(\rho(e_a)) &= -\hat{\kappa}(e_a) = i\hat{\kappa}(\hat{e}_a) = -\bar{e}_a & \text{for } n+1 \le a \le 2m. \end{aligned}$$

Similarly, for $\rho \circ \hat{\kappa}$, using $\rho(\bar{v}) = \overline{\rho(v)}$ we get

$$\rho(\hat{\kappa}(e_a)) = -\rho(\bar{e}_a) = -\overline{\rho(e_a)} = -\bar{e}_a \qquad \text{for } 1 \le a \le n,$$

$$\rho(\hat{\kappa}(e_a)) = i\rho(\bar{\hat{e}}_a) = \rho(\bar{e}_a) = \overline{\rho(e_a)} = -\bar{e}_a \qquad \text{for } n+1 \le a \le 2m.$$

Thus, we obtain $\kappa = \hat{\kappa} \circ \rho = \rho \circ \hat{\kappa}$ on all *V*. Then, for any element $a = \alpha e_{a_1} e_{a_2} \dots e_{a_k}$ with $\alpha \in \mathbb{C}$ and $k \in \mathbb{N}$, we have:

$$\kappa(a) = \rho(\hat{\kappa}(a)) = \hat{\kappa}(\rho(a)).$$

By linearity, this relation extends to all of Cl(V, g), concluding the proof.

We can then define $C = K\hat{C}$ with $K\hat{C} = \pm\hat{C}K$ as $KC = \pm CK$. We recover the relation

$$J = K\hat{J} \tag{3.33}$$

obtained in the context of twisted spectral triples (see equation (2.21)).

²This choice is natural and essential for the upcoming definition of the Dirac operator.

³Similarly to subsection 2.3, as this corresponds to the spectral triple case.

Remark 3.11 In full generality, if we do not restrict to the even-dimensional case, the defining relation for κ is given by $\kappa(v) := -\epsilon_1^K \bar{v}$ (see relation (3.95) for the definition of ϵ_1^K). By defining $\hat{\kappa}(\hat{v}) = -\hat{\epsilon}_1 \bar{\hat{v}}$ and $\delta = \epsilon_1^K / \hat{\epsilon}_1 = \pm 1$, the result of Proposition 3.10 becomes

$$\kappa = \delta \hat{\kappa} \circ \rho = \delta \rho \circ \hat{\kappa}.$$

The result of equation (3.33) is then a specificity of the case $\delta = 1$, which always holds in the even-dimensional case (see Table 1).

In addition, we have the relation

$$KJ = KC \circ cc = \pm CK \circ cc = \pm J\overline{K} = (\pm)JK, \tag{3.34}$$

where the last \pm sign is in parentheses to signify that it is not necessarily the same \pm sign as the ones without parentheses. This notation will not be retained in what follows. We recover relation (2.4), i.e., $\rho(J) = \epsilon J$, given in the context of twisted spectral triples.

It is interesting to note that most of these results would have been different in the odd-dimensional case, as the Clifford algebra is not simple in this case. This implies, in particular, that not all automorphisms are inner, and the commutant V' is no longer proportional to \mathbb{C} , properties that have been essential in many demonstrations.

3.2 From automorphisms of Clifford algebras to indefinite inner products

This section aims at showing how the twist previously defined plays a fundamental role in describing the automorphisms of the Clifford algebra Cl(V, g) which are also automorphisms of V and the induced automorphism invariant inner product in the case of the orthochronous spin group.

The following proposition establishes the Aut(V)-invariance of the structural automorphisms' definitions, justifying the term "structural".

Proposition 3.12 The relations given in (3.7) are preserved under Aut(V), i.e.,

$$\rho(\phi(v)) = \phi(v)^{\dagger}, \qquad \chi(\phi(v)) = -\phi(v), \qquad \kappa(\phi(v)) = -\overline{\phi(v)}$$
(3.35)

for all $v \in V$ and all $\phi \in Aut(V)$.

PROOF Taking $\phi \in \text{Aut}(V)$, ϕ acts as a linear transformation on V, so that $\phi(e_a) = \sum_{b=1}^{2m} \lambda_{ab} e_b$ for some $\lambda_{ab} \in \mathbb{R}$ (since V is a real vector space). Then,

$$\rho(\phi(e_a)) = \sum_{b=1}^{2m} \lambda_{ab} \rho(e_b) = \sum_{b=1}^{2m} \lambda_{ab} e_b^{\dagger} = (\sum_{b=1}^{2m} \lambda_{ab} e_b)^{\dagger} = \phi(e_a)^{\dagger},$$
(3.36)

$$\chi(\phi(e_a)) = \sum_{b=1}^{2m} \lambda_{ab} \chi(e_b) = -\sum_{b=1}^{2m} \lambda_{ab} e_b = -\phi(e_a),$$
(3.37)

$$\kappa(\phi(e_a)) = \sum_{b=1}^{2m} \lambda_{ab} \kappa(e_b) = -\sum_{b=1}^{2m} \lambda_{ab} \bar{e}_b = -\overline{\phi(e_a)}.$$
(3.38)

We then recover the result of relations (3.35) for any $v \in V$.

Notably, if $\phi \in Aut(V)$, we have $\phi \circ \chi = \chi \circ \phi$.

We will only consider automorphisms of *V* in what follows. More precisely, we are interested in the automorphisms of Cl(V, g) being also automorphisms of *V*, denoted by

$$Q(V,g) := \operatorname{Aut}(V) \cap \operatorname{Aut}(Cl(V,g)).$$
(3.39)

Proposition 3.13 Given $\phi \in Q(V, g)$, defined via $x \in Cl(V, g)^{\times}$ by $\phi_x(y) = ad_x(y)$ for all $y \in Cl(V, g)$, we have

$$x^{-1} = \alpha \rho(x^{\dagger}), \tag{3.40}$$

where α is a real number.

PROOF Since $\phi \in \text{Aut}(V)$, the preservation of the defining relation $\rho(v) = v^{\dagger}$ implies that $\phi_x(v)^{\dagger} = \rho(\phi_x(v))$ for $v \in V$, so that

$$(x^{-1})^{\dagger}v^{\dagger}x^{\dagger} = Kxvx^{-1}K^{\dagger}.$$
(3.41)

Using the fact that $v^{\dagger} = KvK^{\dagger}$, equation (3.41) becomes

$$x^{-1}K^{\dagger}(x^{-1})^{\dagger}KvK^{\dagger}x^{\dagger}Kx = v.$$
(3.42)

This rewrite $A^{-1}vA = v$, where $A = K^{\dagger}x^{\dagger}Kx$. This implies that A belongs to the commutant of V, and hence takes the form $\beta \mathbb{1}_{2^m}$ from Proposition 3.1, with $\beta \in \mathbb{C}$. Consequently, we obtain $K^{\dagger}x^{\dagger}Kx = \beta \mathbb{1}_{2^m}$ and then $x^{-1} = \frac{1}{\beta}K^{\dagger}x^{\dagger}K$. Multiplying $K^{\dagger}x^{\dagger}Kx = \beta \mathbb{1}_{2^m}$ on the left by K, this relation becomes $x^{\dagger}Kx = \beta K$. Taking the adjoint, we obtain $x^{\dagger}K^{\dagger}x = \beta^*K^{\dagger}$, which is equivalent to $x^{\dagger}Kx = \beta^*K$ since $K = \exp(i\theta)K^{\dagger}$. This implies that β is real. Identifying α with $1/\beta$, we obtain $x^{-1} = \alpha K^{\dagger}x^{\dagger}K = \alpha Kx^{\dagger}K^{\dagger}$, hence the result.

We define $V^{\times} = V \cap Cl^{\times}(V, g)$ to be the subset of invertible elements in V or equivalently⁴

$$V^{\times} = \{ v \in V \mid g(v, v) \neq 0 \}.$$
(3.43)

Taking $u, v \in V$ and observing that $ad_v(u) \in V$, the group $P(V, g) \subset Q(V, g)$ is defined as the group generated by the elements of V^{\times} :

$$P(V,g) = \{v_1 \dots v_k / v_1, \dots, v_k \in V^{\times} \text{ and } k \in \mathbb{N}^* \text{ is finite}\}.$$
(3.44)

Taking $x = v_1 \dots v_k \in P(V, g)$, we define $q(x) := \operatorname{card}\{v_i \mid g_{v_i} < 0\}$. We then have that

$$x^{-1} = v_k^{-1} \dots v_1^{-1} = v_k g_{v_k}^{-1} \dots v_1 g_{v_1}^{-1}$$
(3.45)

and that $\rho(x^{\dagger}) = v_k \dots v_1$, implying that equation (3.40) becomes

$$x^{-1} = \alpha \rho(x^{\dagger}) = g_{v_1}^{-1} \dots g_{v_k}^{-1} \rho(x^{\dagger}).$$
(3.46)

Now if we define the (twisted) adjoint action $\tilde{\phi}_x(y) := \widetilde{ad}_x(y) = \chi(x)yx^{-1}$, taking $u, v \in V$, we have that

$$\widetilde{\mathrm{ad}}_{v}(u) = u - 2\frac{g(v, u)}{g_{v}}v$$
(3.47)

⁴The relation $vv = g_v \mathbb{1}_{2^m}$ directly implies the existence of v^{-1} if $g_v \neq 0$. If $g_v = 0$ we have that vv = 0, so that the existence of v^{-1} leads to the contradiction $v = v(vv^{-1}) = (vv)v^{-1} = 0$, meaning that v^{-1} does not exist in this case. Hence, the equivalent definitions for V^{\times} .

is a reflection across the orthogonal complement v^{\perp} . Moreover, we have

$$\widetilde{\operatorname{ad}}_x = \widetilde{\operatorname{ad}}_{v_1...v_k} = \widetilde{\operatorname{ad}}_{v_1} \circ \cdots \circ \widetilde{\operatorname{ad}}_{v_k}$$
 and $\operatorname{ad}_{v_1...v_k} = \operatorname{ad}_{v_1} \circ \cdots \circ \operatorname{ad}_{v_k}$

since $\widetilde{ad}_v = -ad_v$. Therefore, the image of P(V, g) under \widetilde{ad} is the group generated by reflections, which is the full orthogonal group; see [20] for more details.

The unit sphere *S* for g is defined as

$$S \mathrel{\mathop:}= \{ v \in V / g_v = \pm 1 \}.$$

Taking $v \in V \cap S$ and $u \in V$, we obtain

$$\widetilde{\operatorname{ad}}_{v}(u) = u - 2 \frac{g(v, u)}{g_{v}} v = \begin{cases} u & \text{for } u \in v^{\perp}, \\ -u & \text{for } u = v. \end{cases}$$
(3.48)

Such a transformation corresponds precisely to an orthogonal reflection. The induced subgroup $Pin(V, g) \subset P(V, g)$ is defined as

$$Pin(V, g) = \{v_1 \dots v_k / v_1, \dots, v_k \in S \text{ and } k \in \mathbb{N}^*\}.$$
(3.49)

Taking any $x \in Pin(V, g)$, from equation (3.46) we have that

$$x^{-1} = \pm \rho(x^{\dagger})$$

The associated spin group is defined as

$$Spin(V,g) = Pin(V,g) \cap Cl^0(V,g)$$

or equivalently, as the subset corresponding to even values of k. The representations of Spin(V, g) are called spinor representations and will be denoted by S. For any $x \in Spin(V, g)$, the image under the adjoint and twisted adjoint actions coincide i.e., $\widetilde{ad}_x = ad_x$.

We define the "fixed" g-orthonormal basis

$$e_{(i)} := e_a|_{a=i}$$

which is, by definition, invariant under Aut(V). The corresponding "fixed" vector space is denoted as V_f . The subgroup $Pin(V_f, g)$, i.e., the fixed version of Pin(V, g), is generated by the elements of V_f in the unit sphere.

Proposition 3.14 The structural automorphisms ρ , χ , κ are elements of $Pin(V_f, g)$ represented by the adjoint action.

PROOF Obviously, we have that ρ , χ , κ are elements of Q(V, g). Taking the general notation ϕ for any of these automorphisms, from Proposition 3.4, we have that ϕ is an involution on Cl(V, g). As a consequence, we can define the eigenspaces $W^{\pm} \subset V$ so that

$$V = W^+ \oplus W^-$$
, with $\phi(w^{\pm}) = \pm w^{\pm}$ for $w^{\pm} \in W^{\pm}$. (3.50)

This implies that ϕ acts as an orthogonal reflection across W^+ . From Proposition 3.12, the definition of ϕ is Aut(*V*)-invariant.

By defining $\{f_{(i)}\}_{1 \le i \le l}$ and $\{f_{(i)}\}_{l+1 \le i \le 2m}$ as fixed *g*-orthonormal bases of W^+ and W^- respectively, then for all $u \in V$, we have

$$\phi(u) = \begin{cases} ad_{f_{(1)}} \circ \cdots \circ ad_{f_{(l)}}(u) = ad_{f_{(1)} \cdots f_{(l)}}(u) & \text{for } l \text{ odd,} \\ (\mathbb{1}_{2^m}) ad_{f_{(l+1)}} \circ \cdots \circ ad_{f_{(2m)}}(u) = ad_{f_{(l+1)} \cdots f_{(2m)}}(u) & \text{for } l \text{ even.} \end{cases}$$
(3.51)

where $f_{(1)} \dots f_{(l)}$ and $f_{(l+1)} \dots f_{(2m)}$ are elements of $Pin(V_f, g)$, thus providing the desired Aut(*V*)-invariant automorphism ϕ , hence the result.

For ρ , noting that $W^{\pm} = V^{\pm}$, the unitary operator *K* is then given by

$$K = \begin{cases} \alpha_1 \prod_{i=1}^n e_{(i)} & \text{for } n \text{ odd,} \\ \alpha_1 (\mathbb{1}_{2^m}) \prod_{i=n+1}^{2m} e_{(i)} & \text{for } n \text{ even.} \end{cases}$$
(3.52)

with $\alpha_1^{-1} = \alpha_1^*$ a complex number, where the fixed basis $e_{(i)}$ guarantees that *K* remains Aut(*V*)-invariant.

For Γ , we have that $V = W^-$, since $\chi(v) = -v$ for any $v \in V$, so that

$$\Gamma = \alpha_2 \prod_{i=1}^{2m} e_{(i)} \tag{3.53}$$

with $\alpha_2^{-1} = \alpha_2^*$ a complex number. We now fix $\alpha_2 = i^{-m(2m-1)-n}$ so that $\Gamma^2 = \mathbb{1}_{2^m}$ and $\Gamma^{\dagger} = \Gamma$.

For *C*, a convenient choice is to require that the basis $\{e_a\}_{1 \le a \le 2m}$ (which is then the fixed one) satisfies $\bar{e}_a = \pm e_a$. By defining the index set $I := \{i / \kappa(e_{(i)}) = -\bar{e}_{(i)} = e_{(i)}\}$ corresponding to the elements in W^+ , and $\bar{I} := \{1, ..., 2m\} \setminus I$ for W^- , we obtain

$$C = \begin{cases} \alpha_3 \prod_{i \in I} e_{(i)} & \text{for } \operatorname{card}(I) \text{ odd,} \\ \alpha_3 (\mathbb{1}_{2^m}) \prod_{i \in \overline{I}} e_{(i)} & \text{for } \operatorname{card}(I) \text{ even.} \end{cases}$$
(3.54)

with $\alpha_3^{-1} = \alpha_3^*$ a complex number.

The invariance of these three operators under Aut(V) follows from their definition based on the fixed basis, in accordance with Proposition 3.12.

The subgroup $Spin(V, g)^+ \subset Spin(V, g)$ called orthochronous spin group is defined as the set of elements of Spin(V, g) that preserve the orientation of V^{\pm} independently (spatial and temporal). Taking $v^{\pm} \in V^{\pm} \cap S$ and $u \in V$, we have that

$$\widetilde{\mathrm{ad}}_{v^{\pm}}(u) = u - 2\frac{g(v^{\pm}, u)}{g_{v^{\pm}}}v^{\pm} = \begin{cases} u & \text{for } u \in (v^{\pm})^{\perp}, \\ -u & \text{for } u = v^{\pm}. \end{cases}$$
(3.55)

meaning that $\widetilde{ad}_{v^{\pm}}$ reverses the orientation of V^{\pm} while leaving the one of V^{\mp} unchanged. Taking $x = v_1 \dots v_k \in Spin(V, g)$, the transformation \widetilde{ad}_x preserves the orientation of V^{\pm} only if q(x) is even, since

$$\widetilde{\mathrm{ad}}_{v_1\ldots v_k} = \widetilde{\mathrm{ad}}_{v_1} \circ \cdots \circ \widetilde{\mathrm{ad}}_{v_k}.$$

As a consequence, the orthochronous spin group corresponds to the subgroup of elements $x \in Spin(V, g)$ such that q(x) is even:

$$Spin(V,g)^{+} = \{x = v_1 \dots v_k / v_1, \dots, v_k \in S, q(x) \in 2\mathbb{N}, \text{ and } k \in 2\mathbb{N}^*\}.$$
 (3.56)

For any $x \in Spin(V, g)^+$, we directly obtain

$$x^{-1} = \rho(x^{\dagger}) \tag{3.57}$$

from equation (3.46), since q(x) is even.

Consequently, a natural $Spin(V, g)^+$ -invariant inner product is given by the *K*-inner product $\langle ., . \rangle_K$ defined in equation (2.14) from $\langle ., . \rangle$. The associated adjoint operator \dagger_K is unique, as ρ is $\mathcal{B}(\mathcal{H})$ -regular. equation (3.57) then rewrites as $x^{-1} = x^{\dagger_K}$, implying that any $x \in Spin(V, g)^+$ is a *K*-unitary operator. The relation $\rho(v) = v^{\dagger}$ then becomes

$$v = v^{\dagger_K} \qquad \forall v \in V, \tag{3.58}$$

which is preserved under Aut(V) and consequently under $Spin(V, g)^+$.

The inner product $\langle ., . \rangle_K$ allows the definition of the dual/adjoint spinor of $\psi \in S$ as

$$\overline{\psi} = \psi^{\dagger} K \tag{3.59}$$

through the relation $\overline{\psi}(\phi) = \langle \psi, \phi \rangle_K$.

Remark 3.15 The specific notation $\langle ., . \rangle_K$ emphasizes the fact that the operator K, being Aut(V)-invariant, is embedded within the algebraic structure. In the case n = 2m, i.e., a positive definite metric, we can take $K = \mathbb{1}_{2^m}$, as mentioned in Remark 3.9. In this case, we recover the usual Hermitian inner product $\langle ., . \rangle_K = \langle ., . \rangle$.

Definition 3.16 Given a vector $\psi \in S$ and the *K*-inner product $\langle ., . \rangle_K$, we define the function $\|.\|_K$ by

$$\|\psi\|_{K} := \sqrt{\langle\psi,\psi\rangle_{K}}.$$
(3.60)

This generalizes the usual notion of norm, which is retrieved when $K = \mathbb{1}_{2^m}$.

As stated in Subsection 2.2, an indefinite (or definite) and Hermitian product $\langle ., . \rangle_K$ can only be obtained by eliminating the phase shift in equation (2.16), enforcing $K = K^{\dagger}$ so that ρ acts as a fundamental twist. Such a product is an example of a Krein product (defined more abstractly in Subsection 3.4). In what follows, we will only consider such fundamental twists, ensuring that any product $\langle ., . \rangle_K$ is indefinite and Hermitian, with

$$K = K^{\dagger} = (\overline{K})^{\top} = \pm K^{\top} = \pm \overline{K}.$$
(3.61)

3.3 Even dimensional pseudo-Riemannian manifolds

Let's consider a 2m-dimensional smooth compact manifold \mathcal{M} equipped with a nondegenerate metric g of signature (n, 2m - n), defined on the tangent space $T\mathcal{M}$. A local oriented g-orthonormal basis of $T\mathcal{M}$ is given by $\{E_1, \ldots, E_{2m}\}$, with components satisfying $g_{ab} = g(E_a, E_b)$. The dual basis is given by $\{\theta^1, \ldots, \theta^{2m}\}$, defined by the relation $g(\theta^b, E_a) = \delta_a^b$. The metric g induces a splitting $T\mathcal{M} = T\mathcal{M}^+ \oplus T\mathcal{M}^-$, where a local oriented orthonormal basis is given by $\{E_1, \ldots, E_n\}$ for $T\mathcal{M}^+$ and $\{E_{n+1}, \ldots, E_{2m}\}$ for $T\mathcal{M}^-$. Consequently, any element $v \in T\mathcal{M}$ can be decomposed as $v = v^+ \oplus v^-$, with $v^{\pm} \in T\mathcal{M}^{\pm}$, such that $g(v^+, v^+) \ge 0$ and $g(v^-, v^-) \le 0$. Thus, the signature of g consists of n positive and 2m - n negative signs in this basis.

At each point $x \in \mathcal{M}$, we associate the local metric g_x and the tangent space $T\mathcal{M}_x$. The Clifford bundle $Cl_{n,2m-n}$ is the bundle of Clifford algebras, defined pointwise as

$$Cl_{n,2m-n} = \bigcup_{x\in\mathcal{M}} Cl(T\mathcal{M}_x, g_x),$$

where $Cl(T\mathcal{M}_x, g_x)$ is the Clifford algebra generated by the elements c(v) and $\mathbb{1}_{2^m}$ for $v \in T\mathcal{M}_x$, satisfying the relation

$$c(u)c(v) + c(v)c(u) = 2g_x(u,v)\mathbb{1}_{2^m}, \qquad \forall u, v \in T\mathcal{M}_x.$$
(3.62)

We define the vector space $V_x \subset Cl(T\mathcal{M}_x, g_x)$ associated with $T\mathcal{M}_x$ by

$$V_x := \{c(v) \mid v \in T\mathcal{M}_x\}$$

$$(3.63)$$

so that $Cl(T\mathcal{M}_x, g_x)$ is generated by V_x and $\mathbb{1}_{2^m}$. The corresponding vector bundle is then given by $V = \bigcup_{x \in \mathcal{M}} V_x$. As previously, unless explicitly mentioned otherwise, we replace the notation c(v) by v for any element $v \in V_x$ and for any $x \in \mathcal{M}$. Automorphisms of the vector bundle V are defined as

$$\operatorname{Aut}(V) = \bigcup_{x \in \mathcal{M}} \operatorname{Aut}(V_x).$$
(3.64)

The spinor bundle S over \mathcal{M} is a left-module for $Cl_{n, 2m-n}$. A section $\psi : \mathcal{M} \to S$ of S is a map that associates a fiber S_x to each point $x \in \mathcal{M}$. These sections, known as spinor fields, constitute the vector space denoted $\Gamma(S)$. In this framework, the Spin group is more commonly denoted as Spin(n, 2m - n).

We define the Hermitian inner product

$$\langle ., . \rangle : \Gamma(S) \times \Gamma(S) \to C^{\infty}(\mathcal{M}, \mathbb{C})$$

with adjoint \dagger . If \mathcal{M} is compact, then any vector bundle carries a continuously varying fiber-wise inner product $\langle . , . \rangle_x$ so that for $\psi, \psi' \in \Gamma(\mathcal{S})$, we have $\langle \psi, \psi' \rangle_x = \langle \psi(x), \psi'(x) \rangle_x$.

As before, the twist is (locally) defined on V_x by the relation

$$\rho_x(v) := \rho(v) = K_x v K_x^{\dagger} = v^{\dagger}, \qquad \forall v \in V_x, \qquad (3.65)$$

with $\rho_x(w) = 0$ for any $w \in V_{y \neq x}$.

From this, we define the fiber-wise indefinite inner product

$$\langle \, . \, , \, . \, \rangle_{K_x} = \langle \, . \, , K_x \, . \, \rangle_x, \tag{3.66}$$

which is invariant under $Spin(V_x, g_x)^+$.

Using the fiber-wise extended norm $\|\psi\|_{K_x}^2 = \langle \psi(x), \psi(x) \rangle_{K_x}$ from equation (3.60), we define its bundle counterpart as

$$\|\psi\|_{K}^{2} := \int_{\mathcal{M}} \|\psi\|_{K_{x}}^{2} \sqrt{|g|} dx, \qquad (3.67)$$

where $\sqrt{|g|}$ is the volume form.

The space of square-integrable sections of the spinor bundle over \mathcal{M} is defined by

$$L^{2}(\mathcal{M}, S) := \{ \psi \in \mathcal{H}_{m} / \|\psi\|_{K}^{2} < \infty \}.$$
(3.68)

Defining $g^{ab} = g(\theta^a, \theta^b)$ in the *g*-orthonormal basis, the gamma matrices are defined as the unitary operators $\gamma^a := c(\theta^a)$ for all $a \in \{1, ..., 2m\}$, satisfying

$$\{\gamma^a, \gamma^b\} = 2g^{ab}\mathbb{1}_{2^m}.$$
(3.69)

The relation to general coordinates $dx^{\mu 5}$ is provided by the vielbein:

$$c(dx^{\mu}) = \gamma^{\mu} = c(e^{\mu}_{a}dx^{a}) = e^{\mu}_{a}\gamma^{a}.$$
(3.70)

The relation between the twist and the adjoint remains true in general coordinates

$$(\gamma^{\mu})^{\dagger} = e_{a}^{\mu}(\gamma^{a})^{\dagger} = e_{a}^{\mu}\rho(\gamma^{a}) = \rho(\gamma^{\mu}).$$
(3.71)

⁵denoted using Greek indices

This implies that the operator K is independent of the choosen coordinate system. Consequently, the relation in equation (3.58) became

$$(\gamma^{\mu})^{\dagger_{K}} = \gamma^{\mu}, \tag{3.72}$$

which is preserved under the action of Aut(V).

The Dirac operator is then given by the \dagger_K -selfadjoint operator

$$D^{K} = i\gamma^{\mu}\nabla^{s}_{\mu}, \tag{3.73}$$

where the spin connection $\nabla_a^s = \partial_a + \frac{1}{4} \Gamma_{ac}^b \gamma^c \gamma_b$ is the lift of the Levi-Civita connection to the spinor bundle.

Using the Aut(*V*)-invariant matrices $\gamma^{(i)} := \gamma^a|_{a=i}$, from equation (3.52) the operator $K = K^{\dagger}$ takes the explicit form

$$K = \begin{cases} i^{-n(n-1)/2} \prod_{i=1}^{n} \gamma^{(i)} & \text{for } n \text{ odd,} \\ (\mathbb{1}_{2^m}) \prod_{i=n+1}^{2m} \gamma^{(i)} & \text{for } n \text{ even.} \end{cases}$$
(3.74)

Similarly, a grading operator satisfying $\Gamma^{\dagger} = \Gamma$ is defined from equation (3.53) as

$$\Gamma = i^{-m(2m-1)-n} \prod_{i=1}^{2m} \gamma^{(i)}.$$
(3.75)

The real structure operator *J* is defined from *K* and \hat{J} following equation (2.21). From equation (3.24), we obtain the fundamental relation

$$J\gamma^{\mu} = -\gamma^{\mu}J. \tag{3.76}$$

Consequently, the operators K, Γ , and J remain invariant under Aut(V), encompassing coordinate transformations and the Spin group action.

Proposition 3.17 In the case where n is odd, we have

$$K\Gamma = -\Gamma K$$
 and $KJ = (-1)^{n(3n-1)/2} JK.$ (3.77)

In the case where n is even, we obtain

 $K\Gamma = \Gamma K$ and KJ = JK. (3.78)

PROOF In the even case, we have

$$K\Gamma = (-1)^n \Gamma K = -\Gamma K. \tag{3.79}$$

Noting that $i^{-n(n-1)/2} = i^{3n(n-1)/2}$ and that Ji = -iJ, we obtain

$$JK = (-1)^{3n(n-1)/2} i^{3n(n-1)/2} J \prod_{i=1}^{n} \gamma^{(i)} = (-1)^{3n(n-1)/2} (-1)^n K J = (-1)^{n(3n-1)/2} K J.$$
(3.80)

Similarly, in the odd case, we obtain $K\Gamma = (-1)^{2m-n}\Gamma K = \Gamma K$ and $JK = (-1)^{2m-n}KJ = KJ$.

From equation (2.4), we specify the relation between J and K using the parameter ϵ :

$$KJ = \epsilon JK, \tag{3.81}$$

and the relation between *K* and Γ using the parameter $\epsilon' = \pm 1$:

$$K\Gamma = \epsilon' \Gamma K. \tag{3.82}$$

Note that ϵ and ϵ' are functions of *n* in all cases.

Remark 3.18 The odd case, where $\epsilon' = -1$, corresponds to a twist by the grading, since it implies $\rho(\Gamma) = -\Gamma$ so that ρ exchanges the representations π_+ and π_- in equation (2.11).

Now, let us focus on the four-dimensional Lorentzian case. The four gamma matrices (in the chiral representation) are given by γ^a with $a \in \{0, 1, 2, 3\}$,⁶ defined as follows:

$$\gamma^a = \begin{pmatrix} 0 & \sigma^a \\ \tilde{\sigma}^a & 0 \end{pmatrix}. \tag{3.83}$$

where, for a = 0 and $j \in \{1, 2, 3\}$, we define $\sigma^a := \{\mathbb{1}_2, -\sigma_j\}$ and $\tilde{\sigma}^a := \{\mathbb{1}_2, \sigma_j\}$, with the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \qquad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \qquad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$
(3.84)

They satisfy the anticommutation relation

$$\gamma^{a}\gamma^{b} + \gamma^{b}\gamma^{a} = 2\eta^{ab}\mathbb{1}_{4}, \quad \forall a, b = 0, ..., 3,$$
(3.85)

where η^{ab} is the Minkowski metric. The grading operator is given by

$$\Gamma = -i\gamma^{(0)}\gamma^{(1)}\gamma^{(2)}\gamma^{(3)}.$$

Noting that γ^2 is the only gamma matrix such that $\bar{\gamma}^2 = -\gamma^2$, from equation (3.54), the real structure is defined as

$$J = -i\gamma^{(2)} \circ cc.$$

Taking $K = \gamma^{(0)}$, the indefinite inner product is given by $\langle . , . \rangle_K$, with the adjoint

$$(.)^{\dagger_{K}} := \gamma^{(0)} (.)^{\dagger} \gamma^{(0)}, \tag{3.86}$$

so that the Lorentzian Dirac operator D^K is *K*-selfadjoint.

If $S \in Spin(1,3)^+$ is an element of the orthochronous Lorentz group, then S is unitary with respect to this inner product, i.e.,

$$SS^{\dagger_K} = S^{\dagger_K}S = \mathbb{1}_4. \tag{3.87}$$

These transformations naturally preserve the Dirac Lagrangian

$$\mathcal{L}_f = \overline{\psi} D^K \psi + m \overline{\psi} \psi. \tag{3.88}$$

⁶The index a now starts at 0 instead of 1, aligning with the standard physics notation.

Lorentz transformations act on coordinates and partial derivatives as

$$(x^{a})' = \Lambda^{a}_{b} x^{b}, \quad \text{and} \quad \partial_{a} \to \partial'_{a} = (\Lambda^{-1})^{b}_{a} \partial_{b}.$$
 (3.89)

Taking $S(\Lambda)$ as the bispinor representation of an arbitrary Lorentz transformation Λ , we have that $\psi'(x') = S(\Lambda)\psi(x)$, with the relation

$$S\gamma^{a}S^{-1} = (\Lambda^{-1})^{a}_{\ b}\gamma^{b}.$$
(3.90)

For the Dirac operator $D^K = i\gamma^a \partial_a$, Lorentz transformations act as

$$D^K \rightarrow SD^K S^{-1}$$

This transformation can be viewed in two equivalent ways:

1. As a transformation of the partial derivatives:

$$\partial_a \to \partial'_a = (\Lambda^{-1})^b_{\ a} \partial_b, \qquad \gamma^a \to \gamma^a, \qquad \gamma^{(0)} \to \gamma^{(0)}.$$
 (3.91)

2. As a transformation of the Clifford algebra:

$$\partial_a \to \partial'_a = \partial_a, \qquad \gamma^a \to \gamma'^a = S\gamma^a S^{-1}, \qquad \gamma^{(0)} \to \gamma^{(0)}.$$
 (3.92)

A distinction between the transformation behaviors of the matrices γ^0 and $\gamma^{(0)}$ arises only in the second case, which justifies the need to distinguish these two matrices.

The notation $\overline{\psi} = \psi^{\dagger} \gamma^{0}$, commonly used in quantum field theory textbooks instead of $\overline{\psi} = \psi^{\dagger} \gamma^{(0)}$, is therefore misleading. It suggests that $K = \gamma^{0}$ is a true gamma matrix, subject to coordinate changes and varying under Lorentz transformations when applied to the Clifford algebra. However, this does not pose an issue in standard quantum field theory textbooks, as they typically restrict themselves to Minkowski spacetime, where the Clifford algebra is fixed in the background and Lorentz transformations act only on coordinates and partial derivatives.

However, the approach that considers Lorentz transformations at the level of the gamma matrices becomes more appropriate in the generalized context of curved spacetimes. Indeed, the definition and properties of the adjoint spinor $\overline{\psi} = \psi^{\dagger} \gamma^{(0)}$ are crucial for maintaining consistency across curved geometries. The matrix $\gamma^{(0)}$ is fixed in the local tangent space as the canonical temporal gamma matrix in the Minkowski frame and does not transform as a vector, unlike the γ^{μ} matrices. The invariance of $\gamma^{(0)}$ is essential for constructing the adjoint spinor and preserving the indefinite inner product across all coordinate systems. More information can be found in [27].

3.4 Indefinite inner products and pseudo-Riemannian spectral triples

In the previous subsection, new notions of inner products and Dirac operator selfadjointness naturally emerged in the context of pseudo-Riemannian geometries. We now propose to present the consequent noncommutative generalization of these structures.

We begin with the unitary operator $K^{\dagger} = K$, known as fundamental symmetry. The associated indefinite inner product is given by

$$\langle \, . \, , \, . \, \rangle_K \mathrel{\mathop:}= \langle \, . \, , K \, . \, \rangle, \tag{3.93}$$

where $\langle ., . \rangle$ is a (positive definite) inner product associated with the Hilbert space $\mathcal{H} := (V, \langle ., . \rangle)$. Similarly, we define the associated Krein space as

$$\mathcal{K} := (V, \langle ., . \rangle_K).$$

Considering an involutive unital algebra \mathcal{A} represented by bounded operators on \mathcal{H} , the *K* should not be regarded as an element of \mathcal{A} . Consequently, the corresponding *K*-unitary actions on the algebra and the Hilbert space do not act on it:

$$\langle \psi, a\psi \rangle_K \to \langle U_K \psi, U_K a U_K^{\dagger_K} U_K \psi \rangle_K = \langle \psi, U_K^{\dagger_K} U_K a U_K^{\dagger_K} U_K \psi \rangle_K = \langle \psi, a\psi \rangle_K$$

where $U_K \in \mathcal{U}_K(\mathcal{B}(\mathcal{H}))$. This justifies the notation introduced in equation (3.93).

Conversely, fixing *K* as part of the structure constrains the transformations $\psi \to U\psi$ and $\psi \to U^{\dagger}\psi$ that preserve $\langle \psi, \psi \rangle_{K}$ to be *K*-unitaries, since they must satisfy

 $U^{\dagger}KU = UKU^{\dagger} = K, \tag{3.94}$

which implies that

 $UU^{\dagger_K} = U^{\dagger_K}U = \mathbb{1}.$

We can then generalize the notion of spectral triples by defining a spectral triple based on these Krein space structures.

Definition 3.19 (Pseudo-Riemannian spectral triple) A pseudo-Riemannian spectral triple $(\mathcal{A}, \mathcal{K}, D^K)$ is a spectral triple in which the Hilbert space is replaced by a Krein space \mathcal{K} , and the Dirac operator is a Krein self-adjoint operator D^K , with the requirement that $[D^K, a]$ is bounded for any $a \in \mathcal{A}$.

If the pseudo-Riemannian spectral triple admits a real structure, the operator J is derived from \hat{J} (defined in Subsection 2.3) and K via the relation (2.21).

We recover the case of the pseudo-Riemannian manifold described in Subsection 3.3 by considering the triple

$$(C^{\infty}(\mathcal{M}), (L^{2}(\mathcal{M}, \mathcal{S}), \langle . , . \rangle_{K}), D^{K}, J, \Gamma),$$

where D^{K} , K, $L^{2}(\mathcal{M}, S)$, and Γ are defined in Equations (3.73), (3.74), (3.68), and (3.75).

The algebraic constraints of a pseudo-Riemannian spectral triple are specified by:

$$J^{2} = \epsilon_{0}^{K}, \qquad JD^{K} = \epsilon_{1}^{K}D^{K}J, \qquad J\Gamma = \epsilon_{2}^{K}\Gamma J, \qquad \epsilon_{0}^{K}, \epsilon_{1}^{K}, \epsilon_{2}^{K} \in \{\pm 1\}.$$
(3.95)

The following table gives the relation between the KO-dimension of pseudo-Riemannian spectral triples and the numbers ϵ_i^K for $i \in \{0, 1, 2\}$:

$2(n-m) \mod 8$	0	2	4	6
ϵ_0^K	1	-1	-1	1
ϵ_1^K	1	1	1	1
ϵ_2^K	1	-1	1	-1

Table 1: KO-dimensions for even-dimensional pseudo-Riemannian spectral triples.

As for twisted spectral triples, standard spectral triples are retrieved in the special case K = 1, which corresponds to n = 0 in the KO-dimension table. More information on such spectral triples can be found in [28].

4 Isomorphism between pseudo-Riemannian and twisted spectral triples

In the previous sections, an interesting link between twisted spectral triples and pseudo-Riemannian spectral triples has been revealed. This arises through the twist and the associated indefinite inner product, the role played by the corresponding unitaries in each context, and the real structure. This suggests the existence of a connection between these two extended types of spectral triples. This section presents some important notions for what will follow and summarizes key results from [22] concerning this connection.

4.1 *K*-morphism of spectral triples

In this subsection, we present part of the results from [22], specifically focusing on the definition of the *K*-morphism between the (even real) twisted spectral triple ($\mathcal{A}, \mathcal{H}, D, J, \Gamma, K$) and the pseudo-Riemannian spectral triples ($\mathcal{A}, \mathcal{K}, D^K, J, \Gamma$).

Starting with the fundamental symmetry *K*, we construct the corresponding twist as $\rho(.) = K(.)K$. From the compatibility of the real structure with the twist and the result in equation (3.82) we require

$$JK = \epsilon KJ, \qquad K\Gamma = \epsilon' \Gamma K.$$
 (4.1)

The *K*-morphism, denoted Φ^{K} , is a bijective involutive morphism parametrized by *K*. It permits the transition from the twisted spectral triple $(\mathcal{A}, \mathcal{H}, D, J, \Gamma, K)$ to a pseudo-Riemannian spectral triple $(\mathcal{A}, \mathcal{K}, D^{K}, J, \Gamma)$ by constructing D^{K} and \mathcal{K} from *D* and \mathcal{H} through the following relations:

$$D \xrightarrow{\Phi^{K}} D^{K} := KD,$$

$$\mathcal{H} := (V, \langle ., . \rangle) \xrightarrow{\Phi^{K}} \mathcal{K} := (V, \langle ., . \rangle_{K} = \langle ., K. \rangle),$$

where *V* is a vector space. The *K*-morphism does not act on *J* and Γ .

As previously, the characteristic numbers of $(\mathcal{A}, \mathcal{H}, D, J, \Gamma, K)$ are given by $\epsilon_0, \epsilon_1, \epsilon_2, \epsilon$, and ϵ' . Similarly, the characteristic numbers of $(\mathcal{A}, \mathcal{K}, D^K, J, \Gamma)$ are given by $\epsilon_0^K, \epsilon_1^K, \epsilon_2^K, \epsilon$, and ϵ' , which lead to the following relations: $\epsilon_0 = \epsilon_0^K, \epsilon_1^K = \epsilon \epsilon_1$, and $\epsilon_2^K = \epsilon_2$. By doing this it was chacked in [22] that

By doing this it was checked in [22] that

- The assertions $D^{\dagger} = D$ and $(D^K)^{\dagger_K} = D^K$ implies each other.
- The derivations in each context are related by

$$[D,a]_{\rho} = K[D^K,a] \tag{4.2}$$

for any $a \in A$.

• The first order conditions in each contexts implies each other through the relation

$$[[D, a]_{\rho}, b^{\circ}]_{\rho^{\circ}} = K[[D^{K}, a], b^{\circ}].$$
(4.3)

• The fluctuations in each context are related by the relation $D_{A_{\rho}} = K D_{A^{K}}^{K}$ where

$$D_{A^{K}}^{K} = D^{K} + A^{K} + \epsilon_{1}^{K} J A^{K} J^{-1},$$
(4.4)

$$D_{A_{\rho}} = D + A_{\rho} + \epsilon_1 J A_{\rho} J^{-1}, \qquad (4.5)$$

where A^{K} is a connection of the form $A^{K} = \sum_{i} a_{i}[D^{K}, b_{i}]$ and $A_{\rho} = \sum_{i} \rho(a_{i})[D, b_{i}]\rho$, a twisted connection such that $A^{K} = KA_{\rho}$.

• In particular, fluctuations that preserve the \dagger -selfadjointness of D and the \dagger_{K} -selfadjointness of D^{K} are related by the K-unitary $U_{K} = u_{K}Ju_{K}J^{-1}$, where the K-unitary $u_{K} \in \mathcal{A}$

$$D_{A^K}^K := U_K D^K U_K^{\dagger_K} = K V_K D V_K^{\dagger} := K D_{A_\rho}, \tag{4.6}$$

where $V_K = \rho(U_K) = \rho(u_K) J \rho(u_K) J^{-1}$ is also a *K*-unitary operator. The computational details can be found in [22].

Note that all these results depend on the connection between the twist and the Krein product, by requiring the twist to be built from the fundamental symmetry that defines the Krein space structure.

The commutation relation between the grading and the Dirac operator cannot be the same for the two spectral triples. This is due to the presence of *K*, which connects *D* and D^{K} . Specifically, defining $\epsilon_{3}^{K} = \pm 1$ such that $D^{K}\Gamma = \epsilon_{3}^{K}\Gamma D^{K}$, we have $D\Gamma := \epsilon_{3}\Gamma D$ with $\epsilon_{3} = \epsilon'\epsilon_{3}^{K}$. In the case where $\epsilon' = -1$, we obtain a different relation between Γ and the respective Dirac operators.

If we require that $\epsilon_3^K = -1$, i.e., $\{D^K, \Gamma\} = 0$ (the usual case for spectral triples), then we find that D and Γ satisfy a twisted version of the usual anti-commutator relation:

$$\{D,\Gamma\}_{\rho} := D\Gamma + \rho(\Gamma)D = D\Gamma + \epsilon'\Gamma D = 0.$$
(4.7)

If $\epsilon_3 = -1$, then we can work with the twist by grading on the algebra $C^{\infty}(\mathcal{M}) \otimes \mathbb{C}^2$. However, if $\epsilon_3 = 1$, the boundedness condition on the twisted commutator will imply that we must work on the sub-algebra $C^{\infty}(\mathcal{M})$, i.e., the sub-algebra corresponding to elements $a \in C^{\infty}(\mathcal{M}) \otimes \mathbb{C}^2$ on which the twist is trivial, i.e., $\rho(a) = a$ (see [22] for the proof). In this case, the *K*-morphism connects untwisted spectral triples.

We will focus on the case $\epsilon_3^K = -1$ (context 2 in [22]). This choice is motivated by the upcoming results and the fact that physics provides us with the Lorentzian D^K (a special case of D^K here), which anti-commutes with the grading.

As for the grading, the real structure J = KJ is preserved by the *K*-morphism. This is an important property of the *K*-morphism, as we prove that such a real structure is relevant both in the context of twisted spectral triples (see subsection 2.3) and in the one of pseudo-Riemannian spectral triples (see subsection 3.1).

Now, let's see how this *K*-morphism is realized in the case of the spectral triple of even-dimensional manifolds.

4.2 Signature change for even-dimensional manifolds

Let's consider the 2m-dimensional pseudo-Riemannian manifold (\mathcal{M} , g) defined in subsection 3.3. The following subsection shows how the presented *K*-morphism acts in this context, changing the signature of the metric.

Definition 4.1 (Reflection) Taking (\mathcal{M}, g) with g being any metric, a reflection r is an isometric automorphism on $T\mathcal{M}$ such that $r^2 = 1$.

Definition 4.2 (Spacelike reflection) Taking (\mathcal{M}, g) to be a pseudo-Riemannian manifold, a spacelike reflection r is a reflection for the metric g so that

$$g_{R}(.,.) := g(.,r.)$$
 (4.8)

is a positive definite metric on $T\mathcal{M}$.

The fundamental symmetry *K* was defined in (3.74) with the twist $\rho(.) = K(.)K$. We have that ρ implements the spacelike reflection *r* at the level of the Clifford algebra

$$\rho(c(v)) = c(rv) \tag{4.9}$$

for any vector $v \in T\mathcal{M}$ (see [22]), i.e., ρ is the parity operator as it reverses the spacelike coordinates, i.e., those corresponding to negative metric components.

The corresponding spectral triple is given by $(C^{\infty}(\mathcal{M}), \mathcal{K}, D^{K}, J, \Gamma)$ with $\mathcal{K} = (L^{2}(\mathcal{M}, S), \langle ., . \rangle_{K})$ and D^{K}, J , and Γ as defined in subsection 3.3, so that $\epsilon_{3}^{K} = -1$ i.e.,

$$\{D^K, \Gamma\} = 0. \tag{4.10}$$

This pseudo-Riemannian spectral triple will be taken as the basis to construct a twisted spectral triple through the *K*-morphism. This justifies the use of the space $L^2(\mathcal{M}, S)$ associated with the metric *g*, with $L^2(\mathcal{M}, S)$ being preserved by the *K*-morphism.

Using equation (4.9), one can express relation (4.8) at the Clifford algebra level

$$2g_{R}(v,w)\mathbb{1}_{2^{m}} = 2g(v,rw)\mathbb{1}_{2^{m}} = \{c(v),c(rw)\} = \{c(v),\rho(c(w))\}.$$
(4.11)

We can then define $D = KD^K$ within the corresponding twisted spectral triple $(C^{\infty}(\mathcal{M}), \mathcal{H}, D, J, \Gamma, K)$ which satisfies $\epsilon_3 = -\epsilon_3^K = 1$ i.e.,

$$\{D,\Gamma\}_{\rho} = 0. \tag{4.12}$$

We define $\tilde{\gamma}^a = K\gamma^a$ so that $D = -i\tilde{\gamma}^a \nabla_a^{R,S}$ where $\nabla_a^{R,S} = \partial_a + \frac{1}{4}\tilde{\Gamma}^b_{\ ac}\tilde{\gamma}^c\tilde{\gamma}_b$ is the associated spin connection, defined from ∇_a^s by the relation $\Gamma^b_{\ ac} = g_c\tilde{\Gamma}^b_{\ ac}$.

For standard (untwisted) spectral triples, the metric is obtained by a relation between the norm induced by the inner product and the commutator of the Dirac operator with elements of the algebra as the local evaluation of $||[D, a]||^2$ for $a \in C^{\infty}(\mathcal{M})$, see [26, 8] for more information. A generalized formula to extract g from $(C^{\infty}(\mathcal{M}), \mathcal{K}, D^K, J, \Gamma)$ was given in [22] by extending the notion of norm of vectors and operators through the modification of the inner product into a Krein product in the definition, as follow.

In subsection 3.3 we have seen how the extension of the norm on S defined by $\|.\|_{K}$ was used to define $L^{2}(\mathcal{M}, S)$. The following definition follows this line for operators.

Definition 4.3 Given an operator $O \in \mathcal{B}(\mathcal{H})$ we define $||O||_K$ by

$$\|O\|_{K} := \sup_{\substack{\psi \in \mathcal{H} \\ \|\psi\|_{K} = 1}} \{ \|O\psi\|_{K} \}.$$
(4.13)

We recover the usual notions of operator norm in the case $K = \mathbb{1}_{2^m}$.

Using this, one can show that for any $a \in C^{\infty}(\mathcal{M})$, one obtains

$$\frac{1}{2} \left(\| [D^K, a] \|_K^2 + \| [D^K, a]^{\dagger_K} \|_K^2 \right) = g(\operatorname{grad} a, \operatorname{grad} \bar{a}).$$
(4.14)

In the Riemannian case, having $K = \mathbb{1}_{2^m}$ so that $\|(.)\|_K = \|(.)\|$ and $\dagger_K = \dagger$, we recover the usual formula $\|[D, a]\|^2$ for the extraction of the metric in the distance formula.

The *K*-morphism acts on formula (4.14) by changing the underlying inner product and turning the commutator $[D^K, a]$ into $[D, a]_{\rho} = K[D^K, a]$ so that equation (4.14) becomes

$$\frac{1}{2}\left(\|[D,a]_{\rho}\|^{2}+\|[D,a]_{\rho}^{\dagger_{K}}\|^{2}\right)=g(grad\,a,r\,grad\,\bar{a})=g_{R}(grad\,a,grad\,\bar{a}).$$
(4.15)

The metric g_R can then be associated to the twisted spectral triple ($C^{\infty}(\mathcal{M}), \mathcal{H}, D, J, \Gamma, K$). The *K*-morphism then acts on the metric as the transformation

$$g(.,.) \xrightarrow{\Phi^{K}} g_{R}(.,.) = g(.,r.)$$

$$(4.16)$$

hence by changing its signature through the spacelike reflection r associated with the fundamental symmetry K.

As shown in [22], this leads to a generalization of the concept of Clifford algebras. Indeed, the so-called twisted Clifford algebras are the finite-dimensional algebras generated by operators which are linear representations of the cotangent space $T\mathcal{M}^*$ together with a modified relation to the metric $g_{\mathbb{R}}$. From equation (4.15), two kinds of twisted Clifford algebras can be defined

$$c'_{\rho}(\theta^{a}) := \gamma^{a} \qquad \text{with} \qquad \gamma^{a} \rho(\gamma^{b}) + \rho(\gamma^{b}) \gamma^{a} = 2g_{\scriptscriptstyle R}^{ab} \mathbb{1}_{2^{m}}, \tag{4.17}$$

$$c_{\rho}(\theta^a) := \tilde{\gamma}^a \qquad \text{with} \qquad \tilde{\gamma}^a \tilde{\gamma}^b + \rho(\tilde{\gamma}^b \tilde{\gamma}^a) = 2g_{\scriptscriptstyle R}^{ab} \mathbb{1}_{2^m}.$$
 (4.18)

Using this last representation, equation (4.11) takes the equivalent form

$$2g_{R}(v,w)\mathbb{1}_{2^{m}} = c_{\rho}(v)c_{\rho}(w) + \rho(c_{\rho}(w)c_{\rho}(v)).$$
(4.19)

The relation to general coordinate μ is the same as in equation (3.70) since twisted Clifford representations are linear representations of the coordinates on which the Vielbeins act. Note that the operations producing the signature change through the *K*-morphism are defined locally, meaning that the signature change is also performed locally, in the same way as for Wick rotation in quantum field theory.

5 K-morphism from an almost-commutative twisted spectral triple

The previous section concludes with the introduction of the twisted spectral triple $(C^{\infty}(\mathcal{M}), \mathcal{H}, D, J, \Gamma, K)$ derived from the application of a *K*-morphism on a pseudo-Riemannian spectral triple.

We will focus on the case of a pseudo-Riemannian spectral triple of KO dimension 6. This choice is motivated by physical considerations, as it encompasses the 4-dimensional Lorentzian spectral triple case, which will be derived from the corresponding algebraic constraints in subsection 5.3.

The algebraic constraints of $(C^{\infty}(\mathcal{M}), \mathcal{K}, D^{K}, J, \Gamma)$ are then given by

$$\epsilon_0^K = 1, \qquad \epsilon_1^K = 1, \qquad \epsilon_2^K = -1, \qquad \epsilon_3^K = -1.$$
 (5.1)

The numbers ϵ, ϵ' which depend on *n* are left unfixed for the moment.

The twisted spectral triple $(C^{\infty}(\mathcal{M}), \mathcal{H}, D, J, \Gamma, K)$ obtained by the action of the *K*-morphism Φ^{K} is then specified by the four numbers

$$\epsilon_0 = \epsilon_0^K, \qquad \epsilon_1 = \epsilon \epsilon_1^K = \epsilon, \qquad \epsilon_2 = \epsilon_2^K, \qquad \epsilon_3 = \epsilon' \epsilon_3^K = -\epsilon'.$$
 (5.2)

We now turn our attention to the almost-commutative manifold constructed on $C^{\infty}(\mathcal{M}) \otimes \mathcal{A}_F$, where \mathcal{A}_F is a finite-dimensional algebra associated with the spectral triple $(\mathcal{A}_F, D_F, \mathcal{H}_F, J_F, \Gamma_F)$, also of KO dimension 6, whose manifold part is given by $(C^{\infty}(\mathcal{M}), \mathcal{H}, D, J, \Gamma, K)$. The motivations to do this are the following:

• Explicit appearance of D in the Lorentzian Dirac Lagrangian in equation (3.88) as

$$\overline{\psi}D^{K}\psi = \psi^{\dagger}D\psi. \tag{5.3}$$

- The Noncommutative standard model is based on the product $C^{\infty}(\mathcal{M}) \otimes \mathcal{A}_F$ with KO dimension 6 for the finite part. A justification for this choice can be found in [1], ensuring the elimination of the fermion doubling problem i.e. permitting to obtain the physically good number of degrees of freedom for the fermionic fields. Masses are given by D_F , corresponding to the second term of the Dirac Lagrangian. In the Lorentzian case, this mass is explicitly given as a Lorentz scalar $m\psi^{\dagger}\gamma^{(0)}\psi$.
- In quantum field theory and non-commutative geometry, modular flow arises from the Tomita-Takesaki theory of von Neumann algebras. Given a state and an associated von Neumann algebra, the modular operator and the anti-linear modular conjugation define a time evolution through the modular group, parametrized by *t*, which acts on elements of the algebra. If the algebra is commutative, the flow is trivial. This provides a geometric interpretation of time, linking the flow of observables to the algebra's internal symmetries.

On another side, twisted spectral triples have been introduced in the context of von Neumann algebras where the twist itself is connected with the modular flow [12]. In our approach, the twist is connected with the Krein structure of the dual pseudo-Riemannian spectral triple. The intuition is then that there may be a connection between the twist and a finite noncommutative part, hence the proposition to study the presented product structure with a noncommutative finite space. This section proposes a top-down reversed approach, where the starting point is now the presented twisted spectral triple within a product spectral triple. The aim will be to show how this very product structure harmonizes with the twisted spectral triple ones, to show how this implies the emergence of the dual pseudo-Riemannian spectral triple and to characterize this last one.

5.1 Product of twisted spectral triples and spectral triple

We start from the twisted spectral triple $(C^{\infty}(\mathcal{M}), \mathcal{H}, D, J, \Gamma, K)$ with algebraic constraints given by $\epsilon_i = 1$ for $i \in \{0, 1, 2, 3\}$ defined in (5.2) and the numbers ϵ, ϵ' . The finite spectral triple $(\mathcal{A}_F, D_F, \mathcal{H}_F, J_F, \Gamma_F)$ of KO dimension 6 is specified by the numbers $\epsilon_i^F = \epsilon_i^K$ for $i \in \{0, 1, 2, 3\}$ given in (5.1). The inner products on \mathcal{H} and \mathcal{H}_F are given by $\langle ., . \rangle$ and $\langle ., . \rangle_F$.

The product spectral, is denoted $(\mathcal{A}_p, \mathcal{H}_p, D_p, J_p, \Gamma_p, K_p)$ where the algebra is given by $\mathcal{A}_p := C^{\infty}(\mathcal{M}) \otimes \mathcal{A}_F$ with Hilbert space $\mathcal{H}_p := \mathcal{H} \otimes \mathcal{H}_F$ and adjoint is $(.)^{\dagger_p} = (.)^{\dagger} \otimes (.)^{\dagger_F}$ where \dagger_F is the adjoint associated with \mathcal{H}_F . The full real structures, grading, and twist are given by

 $J_p := J \otimes J_F \qquad \qquad \Gamma_p := \Gamma \otimes \Gamma_F \qquad \qquad \rho_p := \rho \otimes \mathbb{1}_F$

and the Dirac operator by

$$D_{p} := D \otimes \mathbb{1}_{F} + \mathcal{O} \otimes D_{F}$$

where \mathcal{O} is an operator that will be specified by the algebraic constraints connecting J_p , D_p , and Γ_p in the upcoming proposition. The operator's algebraic constraints are given by

$$J_p^2 = \epsilon_0^p \qquad J_p D_p = \epsilon_1^p D_p J_p \qquad J_p \Gamma_p = \epsilon_2^p \Gamma_p J_p \qquad \Gamma_p D_p = \epsilon_3^p D_p \Gamma_p.$$
(5.4)

The following proposition is an important result showing how the operator K may appear naturally in D_p .

Proposition 5.1 The spectral triple algebraic constraints (see (5.4)) imply

$$\mathcal{O} = \mathcal{O}^{\dagger} \qquad \qquad J\mathcal{O} = \epsilon \mathcal{O}J \qquad \qquad \Gamma \mathcal{O} = \epsilon' \mathcal{O}\Gamma. \tag{5.5}$$

PROOF The requirement $D_p^{\dagger_p} = D_p$ implies directly $\mathcal{O} = \mathcal{O}^{\dagger}$.

The "existence" of ϵ_1^p implies that

$$J_p D_p = JD \otimes J_F + J\mathcal{O} \otimes J_F D_F = \epsilon_1 DJ \otimes J_F + \epsilon_1^F J\mathcal{O} \otimes D_F J_F$$

so that $\epsilon_1^p = \epsilon_1 = \epsilon$ and then that $J\mathcal{O} = \epsilon \epsilon_1^F \mathcal{O}J = \epsilon \mathcal{O}J$ since $\epsilon_1^F = 1$. Then, the "existence" of ϵ_3^p implies that

$$\Gamma_{p}D_{p} = \Gamma D \otimes \Gamma_{F} + \Gamma \mathcal{O} \otimes \Gamma_{F}D_{F} = \epsilon_{3}\Gamma D \otimes \Gamma_{F} + \epsilon_{3}^{F}\Gamma \mathcal{O} \otimes D_{F}\Gamma_{F}$$

so that $\epsilon_3^p = \epsilon_3 = -\epsilon'$ and $\Gamma \mathcal{O} = -\epsilon' \epsilon_3^F \mathcal{O} \Gamma = \epsilon' \mathcal{O} \Gamma$ since $\epsilon_3^F = -1$.

In addition, we have $\epsilon_0^p = \epsilon_0 \epsilon_0^F = 1$ and $\epsilon_2^p = \epsilon_2 \epsilon_2^F = 1$.

In Proposition 5.1, we see that \mathcal{O} satisfies all the properties required for the unitary operator *K* (for compatibility with the real structure and the grading), while additionally imposing $\mathcal{O} = \mathcal{O}^{\dagger}$, i.e., exhibiting the property of a fundamental symmetry.

Proposition 5.2 The choice of operator $\mathcal{O} = K'$ such that $\rho(.) = K'(.)K'$ preserves the differential structure at the level of $(\mathcal{A}_F, D_F, \mathcal{H}_F, J_F, \Gamma_F)$.

PROOF Let us suppose that $\mathcal{O} = K'$ is a fundamental symmetry. Then, taking any $a = a_1 \otimes a_2 \in \mathcal{A}_F$, the full differential is obtained by computing the composed twisted commutator

$$[D_p,a]_{\rho_p} = [D,a_1]_\rho \otimes a_2 + K'a_1 \otimes D_F a_2 - \rho(a_1)K' \otimes a_2 D_F$$

which implies the requirement that $K'a_1 = \rho(a_1)K'$ in order to obtain the term $[D_F, a_2]$ in the full derivation

$$[D_p, a]_{\rho_p} = [D, a_1]_{\rho} \otimes a_2 + K' a_1 \otimes [D_F, a_2].$$
(5.6)

This requirement is equivalent to $\rho(a_1) = K'a_1K'$.

We therefore propose to replace \mathcal{O} by $K = K^{\dagger}$ so that we now consider the Dirac

$$D_p = D \otimes \mathbb{1}_F + K \otimes D_F. \tag{5.7}$$

This may appear surprising since it is now the algebraic structure of the almostcommutative manifold itself that invites us to introduce K, at the level of the finite Dirac operator. The conditions connecting K to J and Γ through ϵ and ϵ' are then induced by the spectral triple structure of the almost-commutative manifold. Conversely, starting from the unitary operator K deduced in subsections 2.2 and 3.1, the condition $K = K^{\dagger}$ ensuring the hermiticity of the product is now deduced from the requirement $D_p^{\dagger p} = D_p$.

This makes an interesting link between the product spectral triple structure and the twisted spectral triples, where the twist naturally emerges as a mechanism to decouple the differential structures. In addition, the term $K \otimes D_F$ is exactly the expected term; having the form of a mass term.

Remark 5.3 Although the twist ρ is trivial on $C^{\infty}(\mathcal{M})$ (as any element in $C^{\infty}(\mathcal{M})$ is proportional to identity), we keep the notation $\rho(a_1)$ to highlight the role played by the twist in full generality, i.e., for noncommutative generalizations of the twisted spectral triple $(C^{\infty}(\mathcal{M}), \mathcal{H}, D, J, \Gamma, K)$ for which the twist is not necessarily trivial. In the case $\epsilon' = -1$, as mentioned in Remark 3.18, the twist is the twist by the grading. Note that in this case, *K*-unitaries in $C^{\infty}(\mathcal{M})$ are also unitary operators.

We recover the twisted anticommutator relation for the product spectral triple

$$\{D_p, \Gamma_p\}_{\rho_p} = 0 \tag{5.8}$$

so that $\epsilon_3^p = -\epsilon'$.

Remark 5.4 In the case $\epsilon = -1$, we have that ϵ_i corresponds to the inverse sign of the usual KO dimension table in KO dimension 4, and that ϵ_i^p corresponds to the inverse sign of the KO dimension 2 for usual spectral triples. This may indicate that we can extend the KO dimension table by permitting a more general relation between the grading and the Dirac with ϵ_3^p , where the signs ϵ_i are reversed, as illustrated in the following table:

2m	0	2	4	6	(2)	(4)
ϵ_0	1	-1	-1	1	1	1
ϵ_1	1	1	1	1	-1	-1
ϵ_2	1	-1	1	-1	1	-1
ϵ_3	-1	-1	-1	-1	1	1

Table 2: Parameters $\epsilon_0, \epsilon_1, \epsilon_2, \epsilon_3$ for KO-dimensions (mod 8).

The last two columns correspond to the two twisted spectral triples obtained in this section. The notation (2) and (4) is used to underline the fact that the KO dimension interpretation is not clear in this twisted context. This point will be explored in future research. The apparent logic may then be the same as for the almost-commutative manifold of the noncommutative standard model of particle physics, i.e., we obtain a spectral triple of KO dimension (4 + 6) mod 8 = 2.

5.2 *K*-morphism from the spectral triple product structure

The following subsection aims at showing how the very presence of K in equation (5.7) induces the emergence of a Krein structure together with the pseudo-Riemannian spectral triple induced by the K-morphism.

Thanks to this, by defining the *K*-selfadjoint Dirac operator $D^K = KD$, we can rewrite equation (5.7) as

$$D_p = K(D^K \otimes \mathbb{1}_F + \mathbb{1}_{2^m} \otimes D_F) \tag{5.9}$$

and (5.6) as

$$[D_p, a]_{\rho_p} = [D, a_1]_{\rho} \otimes a_2 + Ka_1 \otimes [D_F, a_2]$$

= $K([D^K, a_1] \otimes a_2 + a_1 \otimes [D_F, a_2]).$

We recognize the first elements of transformation of a *K*-morphism for the Dirac and the corresponding derivation at the level of the first spectral triple.

The adjoint \dagger_K is defined from \dagger by the relation (.)^{\dagger_K} = ρ (.)^{\dagger_K} . Let's now examine how *K*-unitaries $U_K \otimes \mathbb{1}_F$ arise as the natural operator to preserve the fundamental symmetry *K* in D_p , thus the term $K \otimes D_F$ of the noncommutative part.

General transformations that preserve the \dagger -selfadjoint property of D are given by the transformations $D \rightarrow \tilde{U}D\tilde{U}^{\dagger}$ where \tilde{U} is not necessarily a unitary operator.

Natural unitary transformations for the first twisted spectral triple must not affect the finite part term, namely *K*. The following proposition shows that they must therefore be implemented by *K*-unitary operators.

Proposition 5.5 The transformations of D_p by operators $\tilde{U} \otimes \mathbb{1}_F$ (at the level of the first spectral triple) given by

$$D_p \to (\tilde{U} \otimes \mathbb{1}_F) D_p(\tilde{U}^{\dagger} \otimes \mathbb{1}_F) \quad and \quad D_p \to (\tilde{U}^{\dagger} \otimes \mathbb{1}_F) D_p(\tilde{U} \otimes \mathbb{1}_F)$$
(5.10)

that preserve $K \otimes D_F$ (corresponding to the second spectral triple) are K-unitary operators.

PROOF The transformation acts respectively on $K \otimes D_F$ as

 $\tilde{U}K\tilde{U}^{\dagger} \otimes D_F$ and $\tilde{U}^{\dagger}K\tilde{U} \otimes D_F$. (5.11)

The term $K \otimes D_F$ is then preserved if $\tilde{U}K\tilde{U}^{\dagger} = \tilde{U}^{\dagger}K\tilde{U} = K$ which implies $\tilde{U}\tilde{U}^{\dagger_{\kappa}} = \tilde{U}^{\dagger_{\kappa}}\tilde{U} = \mathbb{1}_{2^m}$, hence that \tilde{U} is *K*-unitary.

The *K*-unitary operators (for the first spectral triple) then arise naturally from the requirement that they must not affect the term $K \otimes D_F$. This choice is reinforced by the following proposition, where this appears as a necessity to make the fluctuations of *D* and D_F independent.

The following proposition shows how the fluctuations of the respective spectral triple are generated by generalized unitaries.

Proposition 5.6 Taking the operators $U_K \otimes U$ where $U_K = u_K J u_K (J)^{-1}$ with u_K a K-unitary operator in $C^{\infty}(\mathcal{M})$ and $U = u J_F u (J_F)^{-1}$ with u a unitary operator in \mathcal{A}_F , the corresponding fluctuations of D_p are given by

$$(U_K \otimes U)D_p(U_K^{\dagger} \otimes U^{\dagger_F}) = K(D_{A^K}^K \otimes \mathbb{1}_F + \mathbb{1}_{2^m} \otimes D_{A_F})$$
(5.12)

where

$$D_{A^{K}}^{K} := V_{K} D^{K} (V_{K})^{\dagger_{K}} = D^{K} + A^{K} + \epsilon_{1}^{K} J A^{K} J^{-1}$$
$$D_{A_{F}} := U D_{F} U^{\dagger_{F}} = D_{F} + A_{F} + \epsilon_{1}^{F} J_{F} A_{F} J_{F}^{-1}$$

with $V_K = \rho(U_K)$ being also a K-unitary operator and the respective one forms

$$A^{K} = v_{K}[D^{K}, v_{K}^{\dagger_{K}}]$$

 $A_{F} = u[D_{F}, u^{\dagger_{F}}]$

with $v_K = \rho(u_K)$.

PROOF For the first term of equation (5.9), we have that

$$(U_K \otimes U)K(D^K \otimes \mathbb{1}_F)(U_K^{\dagger} \otimes U^{\dagger_F}) = K(\rho(U_K)D^K U_K^{\dagger} \otimes \mathbb{1}_F)$$
$$= K(V_K D^K V_K^{\dagger_K} \otimes \mathbb{1}_F)$$

For the second term of equation (5.9), we have

$$(U_K \otimes U)(K \otimes D_F)(U_K^{\dagger} \otimes U^{\dagger_F}) = K \otimes UD_F U^{\dagger_F}$$

where the equality with the fluctuated Dirac operator D_{A_F} is a well known result.

Taking any $\psi = \psi_1 \otimes \psi_2$ and $\psi' = \psi'_1 \otimes \psi'_2$ in \mathcal{H}_p , the inner product $\langle ., . \rangle_p$ on \mathcal{H}_p is defined by the relation

$$\langle \psi, \psi' \rangle_p = \langle \psi_1, \psi'_1 \rangle \langle \psi_2, \psi'_2 \rangle_F.$$
(5.13)

Using the indefinite inner product $\langle ., . \rangle_K = \langle ., K . \rangle$, the induced product for \mathcal{H}_p is then

$$\langle .\,,\,.\rangle_{K\otimes\mathbb{1}_F} := \langle .\,,K\otimes\mathbb{1}_F\,.\rangle_p = \langle .\,,\,.\rangle_K\langle .\,,\,.\rangle_F. \tag{5.14}$$

Then, defining $D_p^K := KD_p = D^K \otimes \mathbb{1}_F + \mathbb{1}_{2^m} \otimes D_F$, the evaluation of D_p gives

$$\langle \psi, D_p \psi' \rangle_p = \langle \psi, D_p^K \psi' \rangle_{K \otimes \mathbb{1}_F} = \langle \psi_1, D^K \psi_1' \rangle_K \langle \psi_2, \psi_2' \rangle_F + \langle \psi_1, \psi_1' \rangle_K \langle \psi_2, D_F \psi_2' \rangle_F$$
(5.15)

as *K* is a fixed structure for the twisted spectral triple ($C^{\infty}(\mathcal{M}), \mathcal{H}, D, J, \Gamma, K$). This relation also holds true for the fluctuated Dirac operators.

We recognize the Krein structure at the level of the first spectral triple. By considering only the transformation at the level of the first spectral triple, this induces that only the general *K*-unitary operators $\hat{U}_K \otimes \mathbb{1}_F$ (not necessarily of the form $u_K J u_K (J)^{-1} \otimes \mathbb{1}_F$) acting on the structures of the first spectral triple as $\psi_1 \rightarrow \hat{U}_K \psi_1$, $\psi'_1 \rightarrow \hat{U}_K \psi'_1$ and $D^K \rightarrow \hat{U}_K D^K \hat{U}_K^{\dagger \kappa}$ will preserve the evaluation (5.15). This evaluation is more generally preserved by the $K \otimes \mathbb{1}_F$ -unitary operators' actions at the level of the full spectral triple, with the generation of fluctuations in equation (5.12) as a special case.

In this way, the very structure of the product spectral triple singularizes a "sub-spectral triple" ($C^{\infty}(\mathcal{M}), \mathcal{K}, D^{K}, J, \Gamma$) where \mathcal{K} is obtained from \mathcal{H} by replacing its inner product with $\langle ., K . \rangle$. This pseudo-Riemannian spectral triple is nothing else that the *K*-morphism Φ^{K} applied on ($C^{\infty}(\mathcal{M}), D, \mathcal{H}, J, \Gamma, K$). This therefore induces that this product spectral triple structure contains and brings to the forefront a pseudo-Riemannian spectral triple. Note that these conclusions also hold if the twisted spectral triple on $C^{\infty}(\mathcal{M})$ is replaced with a noncommutative generalization.

5.3 Structure of the induced pseudo-Riemannian spectral triple

This subsection aims to present the method for determining the properties (particularly the signature) of the emerged pseudo-Riemannian spectral triple, with special attention to the 4-dimensional case.

We start from the previous twisted spectral triple $(C^{\infty}(\mathcal{M}), D, \mathcal{H}, J, \Gamma, K)$ for the 2*m*dimensional manifold \mathcal{M} . The Dirac operator D takes the form $D = -i\tilde{\gamma}^a \nabla_a^{R,S}$, where the $\tilde{\gamma}^a$ matrices are self-adjoint unitary elements in $M_{2^m}(\mathbb{C})$ such that $\forall a, [\Gamma, \tilde{\gamma}^a] = 0$. We additionally ask them to be unitaries in the orthonormal basis. A general relation to the riemaniann metric g_R is given by the following:

$$\operatorname{tr}(\tilde{\gamma}^{\mu}\tilde{\gamma}^{\nu}) = 2^{m}g_{R}^{\mu\nu} \tag{5.16}$$

For the moment, we keep this general relation with the metric. A twisted Clifford-like relation will be provided at a later stage.

We can then define $\gamma_K^{\mu} := K \tilde{\gamma}^{\mu}$ such that we have $D^K = KD = -i\gamma_K^a \nabla_a^{R,S}$. The subscript K in γ_K^{μ} emphasizes the dependence of γ_K^{μ} on $\tilde{\gamma}^{\mu}$ through the choice of K in this top-down approach. The purpose will then be to understand how the operator K is constrained by the algebraic structure of the almost-commutative spectral triple and to characterize the resulting effects on the $\tilde{\gamma}^{\mu}$'s matrices and the associated metric.

Proposition 5.7 The γ_K^a 's are unitary operators, in general coordinate basis we have that

$$(\gamma_K^{\mu})^{\dagger_K} = \gamma_K^{\mu}.$$
(5.17)

PROOF Since $\tilde{\gamma}^a$ is unitary we have $K\gamma_K^a(\gamma_K^a)^{\dagger}K = \mathbb{1}_{2^m}$ and then by multiplying by K on the left and the right that $\gamma_K^a(\gamma_K^a)^{\dagger} = \mathbb{1}_{2^m}$. In the same way, $(\gamma_K^a)^{\dagger}KK\gamma_K^a = \mathbb{1}_{2^m}$ implies directly $(\gamma_K^a)^{\dagger}\gamma_K^a = \mathbb{1}_{2^m}$ hence the result. Since $\tilde{\gamma}^{\mu}$ is self-adjoint, it follows that $K\gamma_K^{\mu} = (\gamma_K^{\mu})^{\dagger}K$, leading to $\gamma_K^{\mu} = K(\gamma_K^{\mu})^{\dagger}K \equiv (\gamma_K^{\mu})^{\dagger_K}$.

In the same way, the operator D^K can be associated to a metric $g_{K}^{\mu\nu}$ defined by

$$tr(\gamma_{K}^{\mu}\gamma_{K}^{\nu}) = 2^{m}g_{K}^{\mu\nu}.$$
(5.18)

Since $K^2 = \mathbb{1}_{2^m}$, it follows that ρ is an involution. Consequently, we obtain the splitting $M_{2^m}(\mathbb{C}) = M_{2^m}(\mathbb{C})^+ \oplus M_{2^m}(\mathbb{C})^-$, with $\rho(a^{\pm}) = \pm a^{\pm}$ for any $a^{\pm} \in M_{2^m}(\mathbb{C})^{\pm}$.

The $\tilde{\gamma}^{a}$'s (local oriented orthonormal basis) are related to the $\tilde{\gamma}^{\mu}$'s through the relation $\tilde{\gamma}^{a} = e^{a}_{R,\mu} \tilde{\gamma}^{\mu}$ as they form linear representations of the tangent space on which the vielbeins $e^{a}_{R,\mu} \in C^{\infty}(\mathcal{M})$ operate, with the subscript R denoting Riemannian. Since the $\tilde{\gamma}^{a}$'s are mutually orthogonal, i.e. $\operatorname{tr}(\tilde{\gamma}^{a}\tilde{\gamma}^{b}) = 0$ for $a \neq b$, we choose the representation of the $\tilde{\gamma}^{a}$'s such that each $\tilde{\gamma}^{a}$ resides in either $M_{2^{m}}(\mathbb{C})^{+}$ or $M_{2^{m}}(\mathbb{C})^{-}$

$$\rho(\tilde{\gamma}^a) := s(a)\tilde{\gamma}^a \quad \text{with} \quad s(a) = \pm 1 \quad \text{for} \quad \tilde{\gamma}^a \in M_{2^m}(\mathbb{C})^{\pm}.$$
(5.19)

This will be a very convenient choice.

The following proposition shows the connection of the signature of g_R and g_K in the local orthonormal basis.

Proposition 5.8 The metric g_{κ}^{ab} is 0 if $a \neq b$ with signature $g_{\kappa}^{a} = s(a)$ or equivalently

$$g_{\kappa}^{ab} = g_{\kappa}^a g_{\kappa}^{ab}. \tag{5.20}$$

PROOF equation (5.19) is equivalent to

$$\rho(\gamma_K^a) = s(a)\gamma_K^a. \tag{5.21}$$

Then we have

$$2^{m}g_{R}^{ab} = \operatorname{tr}(\tilde{\gamma}^{a}\tilde{\gamma}^{b}) = \operatorname{tr}(\rho(\gamma_{K}^{a})\gamma_{K}^{b}) = s(a)\operatorname{tr}(\gamma_{K}^{a}\gamma_{K}^{b}) = s(a)2^{m}g_{K}^{ab}$$

and then $g_R^{ab} = s(a)g_K^{ab}$ which implies that $g_K^{ab} = 0$ for $a \neq b$ and $g_R^a = s(a)g_K^a$ so that $s(a) = g_K^a$ and $g_R^{ab} = g_K^a g_K^{ab}$.

We recover the result from relation (3.32), i.e., $\rho(\gamma_K^a) = g_{\kappa}^a \gamma_K^a$. The number of "temporal" directions *n* is then defined as card($a / g_{\kappa}^a = 1$), and the grading Γ is obtained from the γ_K^a 's through equation (3.75).

The relation $tr(\tilde{\gamma}^{\mu}\tilde{\gamma}^{\nu}) = tr(\rho(\gamma_{K}^{\mu})\gamma_{K}^{\nu})$ does not permit in itself to obtain a concrete link between g_{R} and g_{K} in general coordinates, as $\tilde{\gamma}^{\mu}$ and γ_{K}^{μ} are not necessarily unitaries. This is obtained by passing through the local orthonormal basis, see the following proposition.

Proposition 5.9 $g_{k}^{\mu\nu}$ and $g_{k}^{\mu\nu}$ are connected by the relation

$$g_{\scriptscriptstyle R}^{\mu\nu}=g_{\scriptscriptstyle K}^a e_{{\scriptscriptstyle R},a}^\mu e_{{\scriptscriptstyle R},b}^\nu e_{{\scriptscriptstyle K},\lambda}^a e_{{\scriptscriptstyle K},\kappa}^b g_{\scriptscriptstyle K}^{\lambda\kappa}$$

where $e_{K,\lambda}^a$'s are the vielbeins for the metric g_K .

PROOF We have that

$$2^{m}g_{R}^{\mu\nu} = e_{R,a}^{\mu}e_{R,b}^{\nu}\operatorname{tr}(\tilde{\gamma}^{a}\tilde{\gamma}^{b}) = g_{K}^{a}e_{R,a}^{\mu}e_{R,b}^{\nu}\operatorname{tr}(\gamma_{K}^{a}\gamma_{K}^{b}) = 2^{m}g_{K}^{a}e_{R,a}^{\mu}e_{K,b}^{\nu}e_{K,\lambda}^{a}e_{K,k}^{b}g_{K}^{\lambda\kappa}$$

We now require the \tilde{y}^{a} 's to satisfy the twisted Clifford relation of equation (4.18)

$$2g_{R}^{ab}\mathbb{1}_{2^{m}} = \tilde{\gamma}^{a}\tilde{\gamma}^{b} + \rho(\tilde{\gamma}^{b}\tilde{\gamma}^{a}).$$
(5.22)

This can be interpreted as a compatibility condition between *K* and the $\tilde{\gamma}^a$ matrices, which becomes necessary to obtain the following Clifford algebraic relation

$$2g_{\kappa}^{ab}\mathbb{1}_{2^{m}} = \gamma_{K}^{a}\gamma_{K}^{b} + \gamma_{K}^{b}\gamma_{K}^{a}, \tag{5.23}$$

to connect the dual γ_K^a 's with the metric g_K .

The relation between g_R and g_K at the level of the Clifford algebraic structures in equations (5.22) and (5.23) is given by observing that equation (5.22) is equivalent to $\rho(\gamma_K^a)\gamma_K^b + \gamma_K^b\rho(\gamma_K^a)$ so that

$$2g_{R}^{ab}\mathbb{1}_{2^{m}}=\rho(\gamma_{K}^{a})\gamma_{K}^{b}+\gamma_{K}^{b}\rho(\gamma_{K}^{a})=g_{K}^{a}(\gamma_{K}^{a}\gamma_{K}^{b}+\gamma_{K}^{b}\gamma_{K}^{a})=g_{K}^{a}2g_{K}^{ab}\mathbb{1}_{2^{m}},$$

using equation (5.20).

From the fact that

$$\rho(\tilde{\gamma}^a \tilde{\gamma}^b) = \rho(\tilde{\gamma}^a) \rho(\tilde{\gamma}^b) = g^a_{\kappa} g^b_{\kappa} \tilde{\gamma}^a \tilde{\gamma}^b, \tag{5.24}$$

relation (5.22) can be rewritten as

$$2g_{\scriptscriptstyle R}^{ab}\mathbb{1}_{2^m} = \tilde{\gamma}^a \tilde{\gamma}^b + s_{ab} \tilde{\gamma}^b \tilde{\gamma}^a, \tag{5.25}$$

where $s_{ab} = g^a_{\kappa} g^b_{\kappa}$ is equal to 1 if a = b and can take the values ± 1 if $a \neq b$.

This can be seen as a direct generalization of the usual Clifford relations, where Clifford algebras correspond to the special case when $s_{ab} = 1$ for all *a* and *b*.

We have in KO dimension 6 that 2(n - m) = 8N + 6 where $N \in \mathbb{Z}$, implying that n = 4N + m + 3. This shows that *n* is odd if *m* is even, which further implies that $\epsilon' = -1$ (from proposition 3.17), and that *n* is even if *m* is odd, leading to $\epsilon' = 1$.

We now focus on the four-dimensional case. Since *m* is even, it follows that $\epsilon' = -1$ and *n* is odd. The following proposition presents a significant result of this article, demonstrating how the algebraic constraints of the spectral triple structure of the almost-commutative Manifold are sufficiently strong to explicitly determine *K* and the signature of the induced metric.

Proposition 5.10 In dimension 4, the fundamental symmetry K induces the transformation from the metric g_R with signature (+, +, +, +) to the metric g_K with signature (+, -, -, -)when $\epsilon = -1$, and to (+, +, +, -) when $\epsilon = 1$.

PROOF We aim to demonstrate that the conditions

$$K^{\dagger} = K, \qquad K\Gamma = -\Gamma K, \qquad KJ = \epsilon JK$$
 (5.26)

which arise from the structure of the spectral triple of the almost-commutative manifold, together with the relation $\rho(\gamma_K^a) = g_{\kappa}^a \gamma_K^a$, lead to the conclusion that there is only one index *a* for which $g_{\kappa}^a = 1$ when $\epsilon = -1$, and exactly three such indices when $\epsilon = 1$.

a for which $g_K^a = 1$ when $\epsilon = -1$, and exactly three such indices when $\epsilon = 1$. We have that the matrices $\Gamma^a := \alpha_a \gamma_K^{(a)}$ where α_a is a complex number such that $(\Gamma^a)^{\dagger} = \Gamma^a$ and $(\Gamma^a)^2 = \mathbb{1}_4$ constitute a free family due to the metric relation, see equation (5.23). This free family can be extended by incorporating the elements $\mathbb{1}_4$ and the grading operator Γ (see equation (3.75)), together with the elements

$$\begin{split} \Gamma^{ab} &:= \alpha_{ab} \gamma_K^{(a)} \gamma_K^{(b)} & \text{with} & a < b \\ \Gamma^{abc} &:= \alpha_{abc} \gamma_K^{(a)} \gamma_K^{(b)} \gamma_K^{(c)} & \text{with} & a < b < c \end{split}$$

with α_{ab} and α_{abc} being complex numbers such that $(\Gamma^{ab})^{\dagger} = \Gamma^{ab}$ and $(\Gamma^{ab})^2 = \mathbb{1}_4$, and similarly $(\Gamma^{abc})^{\dagger} = \Gamma^{abc}$ and $(\Gamma^{abc})^2 = \mathbb{1}_4$.

There are 6 elements Γ^{ab} and 4 elements Γ^{abc} , such that the set $\mathcal{I} := \{\mathbb{1}_4, \{\Gamma^a\}_a, \{\Gamma^{ab}\}_{a < b}, \{\Gamma^{abc}\}_{a < b < c}, \Gamma\}$ forms a free family of 16 orthogonal elements⁷.

Since \mathcal{I} is a free family of 16 elements, it is therefore also a basis of $M_4(\mathbb{C})$ so that we can write $K \in M_4(\mathbb{C})$ according to this basis

$$K = \lambda \mathbb{1}_4 + e_a \Gamma^a + e_{bc} \Gamma^{bc} + e_{def} \Gamma^{def} + \epsilon \Gamma$$
(5.27)

where λ , e_a , e_{bc} , e_{def} , ϵ are complex numbers.

The condition $K\Gamma = -\Gamma K$ then implies that λ , e_{ab} and ϵ must be zero because $\Gamma \gamma_K^{(a)} = -\gamma_K^{(a)}\Gamma$ for any *a*. Consequently, *K* can be expressed as follows

$$K = e_a \Gamma^a + e_{bcd} \Gamma^{bcd}. \tag{5.28}$$

Now using the fact that $K\gamma_K^e = g_K^e \gamma_K^e K$, we have that

$$(e_a\Gamma^a + e_{bcd}\Gamma^{bcd})\gamma_K^e = g_\kappa^e\gamma_K^e(e_a\Gamma^a + e_{bcd}\Gamma^{bcd}).$$

We are left with two possibilities

• If $g_{\kappa}^{e} = 1$ for a given (fixed) index *e*, we then have that $e_{a} = 0$ if $a \neq e$ and $e_{bcd} = 0$ if $e \notin \{b, c, d\}$. As *e* is a fixed index, they remain 3 fixed different indices a_{1}, a_{2}, a_{3} so that $a_{i} \neq e$ for i = 1, 2, 3. In this way, *K* takes the form

$$K = e_e \Gamma^e + e_{(ea_1 a_2)} \Gamma^{(ea_1 a_2)} + e_{(ea_1 a_3)} \Gamma^{(ea_1 a_3)} + e_{(ea_2 a_3)} \Gamma^{(ea_2 a_3)}$$
(5.29)

where the notation (*abc*) in $e_{(abc)}\Gamma^{(abc)}$ indicates that the indices *a*, *b*, *c* are ordered in increasing magnitude, e.g., if b < c < a then $e_{(abc)}\Gamma^{(abc)} = e_{bca}\Gamma^{bca}$.

Now if we consider the relation $K\gamma_K^{a_1} = g_K^{a_1}\gamma_K^{a_1}K$, this very relation implies that either $e_{(ea_2a_3)} = 0$ or $e_{(ea_1a_2)} = e_{(ea_1a_3)} = 0$ according to the sign of $g_K^{a_1}$. In the case $e_{(ea_1a_2)} \neq 0$ and $e_{(ea_1a_3)} \neq 0$ we can now consider the relation $K\gamma_K^{a_2} = g_K^{a_2}\gamma_K^{a_2}K$, this implies that either $e_{(ea_1a_2)} = 0$ or $e_{(ea_1a_3)} = 0$ so that in any case, only one of the 3 coefficients $e_{(ea_1a_2)}, e_{(ea_1a_3)}, e_{(ea_2a_3)}$ will be non-zero. Thus, K ultimately takes the form

$$K = e_e \Gamma^e + e_{(ea_1 a_2)} \Gamma^{(ea_1 a_2)}$$
(5.30)

• If $g_{\kappa}^{e} = -1$ for a given (fixed) e, we then have that $e_{a} = 0$ if a = e and $e_{bcd} = 0$ if $e \in \{b, c, d\}$. We then have

$$K = e_{a_1} \Gamma^{a_1} + e_{a_2} \Gamma^{a_2} + e_{a_3} \Gamma^{a_3} + e_{(a_1 a_2 a_3)} \Gamma^{(a_1 a_2 a_3)}$$
(5.31)

⁷The metric properties ensure the freeness of the family, implying that the trace of any product of two distinct elements in \mathcal{I} vanishes, thus guaranteeing orthogonality.

where $a_i \neq e$.

Now if we consider the relation $K\gamma_K^{a_1} = g_K^{a_1}\gamma_K^{a_1}K$, then either $e_{a_1} = 0$ or $e_{a_2} = e_{a_3} = 0$. In the case $e_{a_2} \neq 0$ and $e_{a_3} \neq 0$, considering the relation $K\gamma_K^{a_2} = g_K^{a_2}\gamma_K^{a_2}K$ leads to the fact that either $e_{a_2} = 0$ or $e_{a_3} = 0$ so that in any case, only one of the indices e_{a_i} is non-zero. Then K takes the form

$$K = e_{a_1} \Gamma^{a_1} + e_{(a_1 a_2 a_3)} \Gamma^{(a_1 a_2 a_3)}$$
(5.32)

which is equivalent to (5.30).

Then, in any case, *K* is now given by (5.32). Now let's consider again $K\gamma_K^{a_2} = g_K^{a_2}\gamma_K^{a_2}K$ on this last expression, this clearly implies that either $e_{a_1} = 0$ or $e_{(a_1a_2a_3)} = 0$. In all cases, the condition $K^{\dagger} = K$ leads to either $e_{a_1} = 1$ or $e_{(a_1a_2a_3)} = 1$.

- In the case $K = \Gamma^{a_1}$, we have that $\rho(\gamma_K^{a_1}) = \gamma_K^{a_1}$. Then knowing that $\rho(\gamma_K^a) = (\gamma_K^a)^{\dagger}$, we get that $(\gamma_K^{a_1})^{\dagger} = \gamma_K^{a_1}$ inducing that $K = \Gamma^{a_1} = \gamma_K^{(a_1)}$. This is only consistent with the case $\epsilon = -1$ because $J\gamma_K^{(a)} = -\epsilon_1^K \gamma_K^{(a)} J = -\gamma_K^{(a)} J$. Consequently, $\rho(\gamma_K^a) = g_K^a \gamma_K^a = \gamma_K^a$ holds exclusively for $a = a_1$, leading to the signature (+, -, -, -) for g_K .
- Now if $K = \Gamma^{a_1 a_2 a_3}$ with $a_1 < a_2 < a_3$, for the same reason we have that $\gamma_K^{a_1}, \gamma_K^{a_2}, \gamma_K^{a_3}$ are selfadjoint. Then the requirement $K = K^{\dagger}$ becomes

$$\alpha_{a_1a_2a_3}\gamma_K^{(a_1)}\gamma_K^{(a_2)}\gamma_K^{(a_3)} = \alpha_{a_1a_2a_3}^*\gamma_K^{(a_3)}\gamma_K^{(a_2)}\gamma_K^{(a_1)} = -\alpha_{a_1a_2a_3}^*\gamma_K^{(a_1)}\gamma_K^{(a_2)}\gamma_K^{(a_3)}$$

inducing that $\alpha_{a_1a_2a_3} = i$. Then the condition $J\gamma_K^{(a)} = -\gamma_K^{(a)}J$ implies that

$$JK = J(i\gamma_K^{(a_1)}\gamma_K^{(a_2)}\gamma_K^{(a_3)}) = -i(-1)^3\gamma_K^{(a_1)}\gamma_K^{(a_2)}\gamma_K^{(a_3)}J = KJ.$$

This is consistent only with the case $\epsilon = 1$. Consequently, $\rho(\gamma_K^a) = g_K^a \gamma_K^a = \gamma_K^a$ holds exclusively for $a \in \{a_1, a_2, a_3\}$, leading to the signature (+, +, +, -) for g_K .

This structure consequently leads to the uniqueness of the time dimension in four dimensions, as the two cases $\epsilon = \pm 1$ are merely different conventions when time is associated with the negative metric component in the case $\epsilon = 1$. This is a very specific and welcoming result, revealing the connection between the signature and the algebraic constraints of the almost-commutative spectral triple.

In the case $\epsilon = -1$, which corresponds to the standard Lorentzian case described at the end of subsection 3.3, renaming a_1 as 0 leads to $\tilde{\gamma}^{(0)} = \mathbb{1}_4$ and $K = \gamma_K^{(0)}$, as we obtained in subsection 3.3. In this case, the evaluation of D_p in equation (5.15) is nothing else than the Dirac Lagrangian. This is a scalar (invariant) under Lorentz transformations, being Kunitary operators. We retrieves all the necessary components for a Lorentzian version of the Standard Model (excluding the spectral action), notably the correct form for the mass term given by $K \otimes D_F$. Note that the twisted spectral triple used to deduce the Lorentzian one has also been used in the context of the noncommutative standard model of particle physics in [4], recalling that the twist acts trivially on $C^{\infty}(\mathcal{M})$.

6 Conclusion

The role played by K is fundamental in most of the presented structures. Twisted spectral triples and pseudo-Riemannian spectral triples appear as generalizations of the usual spectral triple, recovered in the special case K = 1. This is precisely at this level that the K-morphism allows the desired signature change. The remaining question was why this may occur. This very question motivates the study of the so-called top-down approach presented in section 5, which places the twisted spectral triple within a product spectral triple, following the spirit of the noncommutative standard model of particle physics.

The main assertion of this article was the proposition that pseudo-Riemannian spectral triples may be seen as emergences from twisted spectral triples within an Almost-commutative spectral triple structure. The KO dimension of the emerged pseudo-Riemannian spectral triple and that of the finite noncommutative one are the same. In the end, the Almost-commutative spectral triple can be seen as a first given object, in which the twisted structure may be considered as a consequence of the "entanglement" between the two spectral triples of the product through the operator *K*. The special twisted Clifford algebraic structure then appears as a modified way to algebraically represent the tangent space $T\mathcal{M}$ and retrieve the metric within the algebraic relations.

In the four-dimensional case, we recover $K_x = \gamma^{(0)}(x) := \gamma^a|_{a=0}(x)$, the fundamental symmetry which appears in Lorentzian quantum field theory, specific to the signature (+, -, -, -). This very operator is linked to the finite noncommutative geometry of the Almost-commutative spectral triple and acts as a local fixed point in the theory, providing the mass term and enabling the definition of the desired coordinate-independent Krein structure, unsensitive to Lorentz transformations.

This article focuses on special cases of physical interest. We leave the generalization of these results for future study. This will, in particular, concern the study of what happens in the odd-dimensional manifold case, the exploration of more general KO dimensions (not only the KO dimension 6), and the characterization of the emerged signature in these more general contexts. An exploration of the potential connection with Tomita-Takesaki theory is also envisioned.

We hope that the structure presented will open concrete paths for studying General Relativity within the framework of spectral triples. The specific algebraic constraints arising from the noncommutative geometry of Yang-Mills-Higgs theory are algebraically linked to the causal structure of spacetime through the fundamental symmetry operator K. This conceptual connection may have significant physical implications that I intend to investigate in future works.

Acknowledgements

I would like to express my sincere gratitude to T. Masson, S. Lazzarini, and A. Sitarz for the insightful discussions and valuable exchanges I had with them throughout the development of this work.

References

- [1] John W Barrett. Lorentzian version of the noncommutative geometry of the standard model of particle physics. *Journal of mathematical physics*, 48(1), 2007.
- [2] Nadir Bizi, Christian Brouder, and Fabien Besnard. Space and time dimensions of algebras with application to lorentzian noncommutative geometry and quantum electrodynamics. *Journal of Mathematical Physics*, 59(6), 2018.
- [3] Arkadiusz Bochniak and Andrzej Sitarz. Finite pseudo-riemannian spectral triples and the standard model. *Physical Review D*, 97(11):115029, 2018.
- [4] Arkadiusz Bochniak and Andrzej Sitarz. Spectral geometry for the standard model without fermion doubling. *Physical Review D*, 101(7):075038, 2020.
- [5] Ali H Chamseddine and Alain Connes. The spectral action principle. *Communications in Mathematical Physics*, 186(3):731–750, 1997.
- [6] Ali H Chamseddine, Alain Connes, and Matilde Marcolli. Gravity and the standard model with neutrino mixing. *Advances in Theoretical and Mathematical Physics*, 11 (6):991–1089, 2007.
- [7] A. Connes and J. Lott. Particle models and noncommutative geometry. *Nuclear Phys. B Proc. Suppl.*, 18B:29–47, 1990.
- [8] Alain Connes. Gravity coupled with matter and the foundation of non-commutative geometry. *Communications in Mathematical Physics*, 182:155–176, 1996.
- [9] Alain Connes. On the spectral characterization of manifolds. *Journal of Noncommutative Geometry*, 7(1):1–82, 2013.
- [10] Alain Connes and John Lott. Particle models and noncommutative geometry. Nucl. Phys. B, 18:29–47, 1989.
- [11] Alain Connes and Matilde Marcolli. *Quantum fields and motives*, volume 56. Elsevier, 2006.
- [12] Alain Connes and Henri Moscovici. Type iii and spectral triples. *arXiv preprint math/0609703*, 2006.
- [13] Agostino Devastato and Pierre Martinetti. Twisted spectral triple for the standard model and spontaneous breaking of the grand symmetry. *Mathematical Physics, Analysis and Geometry*, 20:1–43, 2017.
- [14] Agostino Devastato, Shane Farnsworth, Fedele Lizzi, and Pierre Martinetti. Lorentz signature and twisted spectral triples. *Journal of High Energy Physics*, 2018(3):1–21, 2018.
- [15] M. Dubois-Violette, R. Kerner, and J. Madore. Noncommutative differential geometry of matrix algebras. *J. Math. Phys.*, 31:316, 1990.
- [16] Francesco D'Andrea, Maxim A Kurkov, and Fedele Lizzi. Wick rotation and fermion doubling in noncommutative geometry. *Physical Review D*, 94(2):025030, 2016.

- [17] Nicolas Franco. Temporal lorentzian spectral triples. *Reviews in Mathematical Physics*, 26(08):1430007, 2014.
- [18] Giovanni Landi and Pierre Martinetti. On twisting real spectral triples by algebra automorphisms. *Letters in Mathematical Physics*, 106(11):1499–1530, 2016.
- [19] Giovanni Landi and Pierre Martinetti. Gauge transformations for twisted spectral triples. *Letters in Mathematical Physics*, 108:2589–2626, 2018.
- [20] H Blaine Lawson and Marie-Louise Michelsohn. *Spin Geometry (PMS-38), Volume 38,* volume 20. Princeton university press, 2016.
- [21] Pierre Martinetti, Gaston Nieuviarts, and Ruben Zeitoun. Torsion and lorentz symmetry from twisted spectral triples. *preprint arXiv:2401.07848*, 2024.
- [22] G Nieuviarts. Signature change by an isomorphism of spectral triples. *arXiv preprint arXiv:2402.05839*.
- [23] Mario Paschke and Andrzej Sitarz. Equivariant lorentzian spectral triples. *arXiv* preprint math-ph/0611029, 2006.
- [24] Raphaël Ponge and Hang Wang. Index map, σ -connections, and connes-chern character in the setting of twisted spectral triples. 2016.
- [25] Raphaël Ponge and Hang Wang. Noncommutative geometry and conformal geometry. i. local index formula and conformal invariants. *Journal of Noncommutative Geometry*, 12(4):1573–1639, 2018.
- [26] Vincent Rivasseau and Alain Connes. Non-commutative geometry and the spectral model of space-time. In *Quantum Spaces: Poincaré Seminar 2007*, pages 203–227. Springer, 2007.
- [27] AA Saharian. Quantum field theory in curved spacetime, 2020.
- [28] Alexander Strohmaier. On noncommutative and pseudo-riemannian geometry. *Journal of Geometry and Physics*, 56(2):175–195, 2006.
- [29] Koen Van Den Dungen. Krein spectral triples and the fermionic action. *Mathematical Physics, Analysis and Geometry*, 19:1–22, 2016.