

Uniform stability for the inverse Sturm-Liouville problem with eigenparameter-dependent boundary conditions

Natalia P. Bondarenko

Abstract. We consider a class of self-adjoint Sturm-Liouville problems with rational functions of the spectral parameter in the boundary conditions. The uniform stability for direct and inverse spectral problems is proved for the first time for Sturm-Liouville operator pencils with boundary conditions depending on the eigenparameter. Furthermore, we obtain stability estimates for finite data approximations, which are important from the practical viewpoint. Our method is based on Darboux-type transforms and proving of their Lipschitz continuity.

Keywords: Sturm-Liouville equation; Herglotz-Nevanlinna functions; inverse spectral problems; uniform stability; Darboux-type transforms.

AMS Mathematics Subject Classification (2020): 34A55 34B07 34B09 34L40

1 Introduction

Consider the eigenvalue problem $\mathcal{L}(\sigma, f, F)$:

$$-(y_\sigma^{[1]})' - \sigma(x)y_\sigma^{[1]} - \sigma^2(x)y = \lambda y, \quad x \in (0, \pi), \quad (1.1)$$

$$\frac{y_\sigma^{[1]}(0)}{y(0)} = -f(\lambda), \quad \frac{y_\sigma^{[1]}(\pi)}{y(\pi)} = F(\lambda), \quad (1.2)$$

where σ is a real-valued function of $L_2[0, \pi]$, $y_\sigma^{[1]} := y' - \sigma y$ is the so-called quasi-derivative, λ is the spectral parameter, $f(\lambda)$ and $F(\lambda)$ are rational Herglotz-Nevanlinna functions (see Section 2 for details). Solutions of equation (1.1) are considered in the domain

$$\mathcal{D}_\sigma := \{y \in W_1^1[0, \pi] : y_\sigma^{[1]} \in W_1^1[0, \pi]\}. \quad (1.3)$$

Equation (1.1) is equivalent to the Sturm-Liouville equation

$$-y'' + q(x)y = \lambda y, \quad x \in (0, \pi), \quad (1.4)$$

with the potential $q(x) = \sigma'(x)$ of the distribution space $W_2^{-1}[0, \pi]$.

This paper is concerned with the theory of inverse spectral problems, which consist in the reconstruction of differential operators from their spectral data. Such kind of problems arise in various physical and engineering applications, when one needs to find unknown medium properties from some measurable data or to design a device with desired characteristics. The most complete results in inverse spectral theory have been obtained for the Sturm-Liouville operators with constant coefficients in the boundary conditions (see the monographs [1–4] and references therein). Eigenvalue problems with the spectral parameter in boundary conditions naturally arise in acoustics [5], quantum mechanics [6], fluid dynamics [7], and other physical applications [8]. Inverse spectral theory for self-adjoint Sturm-Liouville problems with rational Herglotz-Nevanlinna functions in the boundary conditions was studied in [9–15]. In particular,

Guliyev [11] has obtained the spectral data characterization for the problem (1.1)–(1.2) with $\sigma \in L_2[0, \pi]$. Furthermore, Freiling and Yurko [16, 17] proposed a method for solving non-self-adjoint inverse Sturm-Liouville problems with arbitrary polynomials of the spectral parameter in the boundary conditions. In recent years, the approach of [16] was developed for the case of distribution potential $q \in W_2^{-1}[0, \pi]$ in [18–20]. Some other issues of inverse spectral theory for differential and intergo-differential operators with polynomials in the boundary conditions were considered in [21, 22] and [23], respectively.

In recent years, a significant progress has been achieved in investigation of uniform stability for inverse spectral problems. In [25, 26], Savchuk and Shkalikov have proved the unconditional uniform stability for inverse Sturm-Liouville problems with potentials in the Sobolev spaces W_2^θ , $\theta \geq -1$. The case $\theta = 0$ was studied by Hryniv [27], who also obtained the results of this kind for the Dirac system [28]. Furthermore, the uniform stability for inverse problems has been proved for several classes of non-local operators, in particular, for intergo-differential operators [29], functional-differential operators with constant delay [30] and with frozen argument [31]. The uniform stability of the non-self-adjoint Sturm-Liouville problem was investigated in [32]. However, to the best of the author’s knowledge, there are no uniform stability results for the Sturm-Liouville inverse problems with eigenparameter-dependent boundary conditions. Local stability of such problems was studied in [19, 20, 24], but therein only small perturbations of spectral data were considered and the constants in the obtained stability estimates depend on a fixed potential. The investigation of uniform stability requires a different approach. One has to find large sets of spectral data or of problem parameters, on which stability estimates are uniform.

In this paper, we prove the uniform stability of direct and inverse spectral problems for the Sturm-Liouville problem (1.1)–(1.2) with Herglotz-Nevalinna functions in the boundary conditions and with σ in the scale of Sobolev spaces $W_2^\alpha[0, \pi]$. As spectral data, we use the eigenvalues $\{\lambda_n\}_{n \geq 1}$ and the norming constants $\{\gamma_n\}_{n \geq 1}$ according to the problem statement in [11]. The most challenging issue in studying the uniform stability is to describe such sets, on which stability estimates hold uniformly. We establish the correspondence between the sets $\mathcal{P}_{Q, \delta}$ of the problem parameters (σ, f, F) and the sets $\mathcal{B}_{R, \varepsilon}$ of the spectral data (see Section 3 for details) and obtain the uniform stability on these sets. Analogously to the previous works by Savchuk and Shkalikov [25, 26] and by Hryniv [27], the set $\mathcal{B}_{R, \varepsilon}$ of spectral data is bounded by the upper constraint R , as well as by the lower constraint $\varepsilon > 0$ on the “gap” between neighboring eigenvalues λ_n and λ_{n+1} and on the norming constant γ_n . A feature of our study is that the set $\mathcal{P}_{Q, \delta}$ of problem parameters is similarly bounded not only by an upper constraint Q but also by a lower constraint $\delta > 0$ related to the Herglotz-Nevalinna functions $f(\lambda)$ and $F(\lambda)$. For the Dirichlet or the Robin boundary conditions, this lower constraint is unnecessary, but it is crucial for forming a closed set of rational functions. Our proof method is based on the Darboux-type transforms that were constructed by Guliyev [11]. Those transforms allow us, by using finite changes of spectral data, reduce the boundary value problem (1.1)–(1.2) to the similar problem with the Dirichlet boundary conditions. We prove the Lipschitz continuity of Guliyev’s transforms and then transfer the results by Savchuk and Shkalikov [25] to the problem with eigenparameter-dependent boundary conditions.

Applying our main results, we obtain the uniform stability estimates for the solution of the inverse problem by the finite spectral data $\{\lambda_n, \gamma_n\}_{n=1}^m$. This kind of estimates is important from the practical viewpoint, since in applications only a finite amount of data is usually available. Therefore, a number of research works are focused on stability of finite data approximations for the Sturm-Liouville operators on a finite interval [33–37], on the half-line [3, Section 5], and on the line [38, 39]. Furthermore, this aspect caused interest for the transmission inverse eigenvalue

problem [40,41] and for inverse resonance problems [42,43]. In this paper, we consider two cases for the problem (1.1)–(1.2), when finite spectral data are known (i) precisely, (ii) with error at most ε . For the both cases, estimates for finite data approximations are readily deduced from our main theorems on the uniform stability.

The paper is organized as follows. Section 2 contains necessary preliminaries about the Sobolev spaces W_2^α and rational Herglotz-Nevalinna functions. In Section 3, we introduce the spectral data and present the main results. In Section 4, auxiliary lemmas about the Lipschitz continuity are proved for some characteristics of the problem $\mathcal{L}(\sigma, f, F)$. In Section 5, we consider the Darboux-type transforms from [11] and study their Lipschitz continuity. Section 6 contains the proofs of our main theorems.

2 Preliminaries

2.1 Spaces W_2^α and l_2^α

Throughout this paper, we denote by $L_p[0, \pi]$ and $W_p^k[0, \pi]$ for $p \geq 1$ and integer $k \geq 0$ the real Lebesgue and Sobolev spaces, respectively, with the corresponding norms:

$$\|u\|_{L_p[0,\pi]} = \left(\int_0^\pi |u(x)|^p dx \right)^{1/p}, \quad \|u\|_{W_p^k[0,\pi]} = \left(\sum_{j=0}^k \left\| \frac{d^j u}{dx^j} \right\|_{L_p[0,\pi]}^p \right)^{1/p}.$$

In particular, $W_2^k[0, \pi]$ ($k = 0, 1, 2, \dots$) are Hilbert spaces and $W_2^0[0, \pi] = L_2[0, \pi]$.

For $\alpha \geq 0$, denote by l_2^α the space of real infinite sequences $v = \{v_n\}_{n \geq 1}$ such that

$$\|v\|_{l_2^\alpha} := \left(\sum_{n=1}^{\infty} n^{2\alpha} v_n^2 \right)^{1/2} < \infty.$$

We define $W_2^\alpha[0, \pi]$ for $\alpha \in (0, \frac{1}{2})$ as the space of functions $u \in L_2[0, \pi]$, whose Fourier coefficients $\hat{u}_k := \frac{2}{\pi} \int_0^\pi u(x) \sin kx dx$ form a sequence of l_2^α , and $\|u\|_{W_2^\alpha[0,\pi]} := \|\{\hat{u}_k\}\|_{l_2^\alpha}$. For brevity, we use the notation $\|\cdot\|_\alpha = \|\cdot\|_{W_2^\alpha[0,\pi]}$ for the norms of functions and $\|\cdot\|_\alpha = \|\cdot\|_{l_2^\alpha}$ for the norms of sequences. Alternatively, the space $W_2^\alpha[0, \pi]$ can be defined by real interpolation $W_2^\alpha[0, \pi] := (L_2[0, \pi], W_2^1[0, \pi])_{\alpha,2}$ or by complex interpolation $W_2^\alpha[0, \pi] := [L_2[0, \pi], W_2^1[0, \pi]]_\alpha$ of Banach spaces for $\alpha \in (0, 1)$ (see [45, 46]).

The following proposition easily follows from the above definitions or can be deduced from the results of [46].

Proposition 2.1. *Let $\alpha \in (0, \frac{1}{2})$. Then, the following assertions hold.*

- (i) *The spaces $W_2^\alpha[0, \pi]$ and l_2^α are compactly embedded in $L_2[0, \pi]$ and l_2 , respectively.*
- (ii) *The space $W_1^1[0, \pi]$ is continuously embedded in $W_2^\alpha[0, \pi]$.*

Denote by $\mathring{L}_2[0, \pi]$ and $\mathring{W}_2^\alpha[0, \pi]$ the corresponding subspaces of functions with the zero mean value $\int_0^\pi u(x) dx = 0$. Note that the antiderivative $\sigma(x)$ of the potential $q(x)$ from (1.4) can be chosen up to an additive constant. For definiteness, we assume that $\sigma \in \mathring{L}_2[0, \pi]$.

2.2 Rational Herglotz-Nevalinna functions

A rational Herglotz-Nevalinna function has the form

$$f(\lambda) = h_0 \lambda + h + \sum_{j=1}^d \frac{\delta_j}{h_j - \lambda}, \tag{2.1}$$

where

$$h_0 \geq 0, \quad h \in \mathbb{R}, \quad \delta_j > 0, \quad j = \overline{1, d}, \quad h_1 < h_2 < \cdots < h_d. \quad (2.2)$$

Define the index of $f(\lambda)$ as

$$\text{ind } f := \begin{cases} 2d + 1, & h_0 > 0, \\ 2d, & h_0 = 0. \end{cases}$$

Additionally, we consider $f = \infty$ with the index $\text{ind } f = -1$. For $M \geq 0$, denote by \mathcal{R}_M the set of rational functions of form (2.1) with $\text{ind } f = M$. Put $\mathcal{R}_{-1} := \{\infty\}$ and

$$\mathcal{R} := \bigcup_{M=-1}^{\infty} \mathcal{R}_M.$$

A function of form (2.1) can be represented as a fraction of polynomials:

$$f(\lambda) = \frac{f_{\uparrow}(\lambda)}{f_{\downarrow}(\lambda)}, \quad f_{\downarrow}(\lambda) := h'_0 \prod_{j=1}^d (h_j - \lambda), \quad h'_0 = \begin{cases} h_0^{-1}, & h_0 > 0, \\ 1, & h_0 = 0. \end{cases} \quad (2.3)$$

If $\text{ind } f \geq 2$, then $f(\lambda)$ has singular points $\{h_j\}_{j=1}^d$ and strictly increases on the intervals $(-\infty, h_1)$, (h_1, h_2) , \dots , (h_{d-1}, h_d) , $(h_d, +\infty)$, since $f'(\lambda) > 0$. Denote

$$\dot{\pi}(f) := \begin{cases} h_1, & \text{ind } f \geq 2, \\ \infty, & \text{ind } f < 2. \end{cases}$$

Consider two cases separately.

Case $\text{ind } f = 2d$: $\lim_{\lambda \rightarrow \pm\infty} f(\lambda) = h$, so the polynomial $f_{\uparrow}(\lambda)$ has exactly d zeros for $h \neq 0$ and $(d-1)$ zeros for $h = 0$. Then, the polynomials $f_{\downarrow}(\lambda)$ and $f_{\uparrow}(\lambda)$ can be represented in terms of their coefficients as follows:

$$f_{\downarrow}(\lambda) = \sum_{n=0}^d a_n \lambda^n, \quad f_{\uparrow}(\lambda) = \sum_{n=0}^d b_n \lambda^n, \quad a_d = (-1)^d, \quad \deg f_{\uparrow} \in \{d-1, d\}.$$

Define

$$c(f) := (a_0, a_1, \dots, a_{d-1}, b_0, b_1, \dots, b_d).$$

Case $\text{ind } f = 2d + 1$: $\lim_{\lambda \rightarrow \pm\infty} f(\lambda) = \pm\infty$, so the polynomial $f_{\uparrow}(\lambda)$ has exactly $d+1$ zeros.

Taking (2.1) and (2.3) into account, we conclude that

$$f_{\downarrow}(\lambda) = \sum_{n=0}^d a_n \lambda^n, \quad f_{\uparrow}(\lambda) = \sum_{n=0}^{d+1} b_n \lambda^n, \quad b_{d+1} = (-1)^d, \quad \deg f_{\downarrow} = d.$$

Define

$$c(f) := (a_0, a_1, \dots, a_d, b_0, b_1, \dots, b_d).$$

In the both cases, we associate a rational function $f \in \mathcal{R}_M$, $M \geq 0$, with the vector $c(f)$ of \mathbb{R}^{M+1} and use the Euclidean norm $\|\cdot\| := \|\cdot\|_{\mathbb{R}^{M+1}}$ for this vector.

For $\text{ind } f = -1$, we set $f_{\downarrow}(\lambda) := -1$, $f_{\uparrow}(\lambda) := 0$, so $c(f)$ is the empty vector and we assume that $\|c(f)\| = 0$.

Consider some examples:

$$\begin{aligned} \text{ind } f = 0: \quad & f(\lambda) = h, \quad f_{\uparrow}(\lambda) = h, \quad f_{\downarrow}(\lambda) = 1, \quad \|c(f)\| = |h|; \\ \text{ind } f = 1: \quad & f(\lambda) = h_0\lambda + h, \quad f_{\uparrow}(\lambda) = \lambda + \frac{h}{h_0}, \quad f_{\downarrow}(\lambda) = \frac{1}{h_0}, \quad c(f) = \left(\frac{1}{h_0}, \frac{h}{h_0} \right). \end{aligned}$$

Note that the set $\{c(f): f \in \mathcal{R}_M\}$ is not closed with respect to the norm $\|\cdot\|_{\mathbb{R}^{M+1}}$ for $M \geq 1$. For an integer $M \geq 0$ and reals $Q > 0$, $\delta > 0$, define the set $\mathcal{R}_{M,Q,\delta} \subset \mathcal{R}_M$ of the functions $f(\lambda)$ of form (2.1) satisfying the conditions:

$$|h| \leq Q, \quad \delta \leq \delta_j \leq Q, \quad j = \overline{1, d}, \quad h_1 \geq 1, \quad h_j + \delta \leq h_{j+1}, \quad j = \overline{1, d-1}, \quad h_d \leq Q, \quad (2.4)$$

$$\begin{cases} \delta \leq h_0 \leq Q, & M = 2d + 1, \\ h_0 = 0, & M = 2d. \end{cases} \quad (2.5)$$

For $M = -1$, set $\mathcal{R}_{M,Q,\delta} = \mathcal{R}_M$.

Lemma 2.2. *Let $M \geq 0$, $Q > 0$, and $\delta > 0$. Then the set $\{c(f): f \in \mathcal{R}_{M,Q,\delta}\}$ is compact in \mathbb{R}^{M+1} .*

Proof. By construction, the coefficients $\{a_n\}$ and $\{b_n\}$ of the polynomials $f_{\uparrow}(\lambda)$ and $f_{\downarrow}(\lambda)$ depend continuously on the constants h, h_0 (if $h_0 \neq 0$), h_j and δ_j for $j = \overline{1, d}$. Furthermore, the set of vectors $(h, h_0, h_1, \dots, h_d, \delta_1, \dots, \delta_d)$ described by the conditions (2.4) and (2.5) is compact in \mathbb{R}^{2d+2} . Hence, its image under the continuous mapping is also compact. \square

Lemma 2.3. *Suppose that $M \geq 0$ and A is such a subset of \mathcal{R}_M that $\{c(f): f \in A\}$ is compact in \mathbb{R}^{M+1} and $\hat{\pi}(f) \geq 1$ for $f \in A$. Then, there exist $Q > 0$ and $\delta > 0$ such that $A \subset \mathcal{R}_{M,Q,\delta}$.*

Proof. For $f \in A$, the coefficients $\{a_n\}$ are uniformly bounded. In the case $M = 2d + 1$, we additionally have $a_d \neq 0$, so $a_d \geq \varepsilon > 0$. This implies (2.5) with $Q = \varepsilon^{-1}$ and some $\delta > 0$ for $f \in A$. Consequently, for both even and odd M , the roots $\{h_j\}_{j=1}^d$ of the polynomial $f_{\downarrow}(\lambda)$ depend continuously on its coefficients. Furthermore, (2.1) implies

$$\delta_j = -\text{Res}_{\lambda=h_j} f(\lambda) = -\frac{f_{\uparrow}(h_j)}{f'_{\downarrow}(h_j)}, \quad h = f(0) - \sum_{j=1}^d \frac{\delta_j}{h_j}.$$

Thus, the map $c(f) \mapsto (h, h_0, h_1, \dots, h_d, \delta_1, \dots, \delta_d)$ is continuous. In view of (2.2), this yields (2.4) with some positive Q and δ for all $f \in A$. \square

Throughout this paper, we use the following notations:

- When two problems $\mathcal{L}(\sigma_i, f_i, F_i)$, $i = 1, 2$, of the same class are considered, the lower index i denotes an object related to the corresponding problem.
- In estimates, the notation $C(A_1, A_2, \dots)$ is used for various positive constants depending on parameters A_1, A_2 , etc. (e.g., $C(Q, \delta)$, $C(R, \varepsilon)$).

3 Main results

Consider the problem $\mathcal{L}(\sigma, f, F)$ of form (1.1)–(1.2) and introduce the spectral data in accordance with [11].

Denote by $\varphi(x, \lambda)$ the solution of equation (1.1) under the initial conditions

$$\varphi(0, \lambda) = f_{\downarrow}(\lambda), \quad \varphi_{\sigma}^{[1]}(0, \lambda) = f_{\uparrow}(\lambda). \quad (3.1)$$

The boundary value problem (1.1)–(1.2) is self-adjoint. It has the countable set of real and simple eigenvalues $\lambda_1 < \lambda_2 < \dots < \lambda_n < \lambda_{n+1} < \dots$, which coincide with the zeros of the characteristic function

$$\chi(\lambda) := F_{\uparrow}(\lambda)\varphi(\pi, \lambda) - F_{\downarrow}(\lambda)\varphi_{\sigma}^{[1]}(\pi, \lambda). \quad (3.2)$$

In addition, define the norming constants

$$\gamma_n := \int_0^{\pi} \varphi^2(x, \lambda_n) dx + f'(\lambda_n)\varphi^2(0, \lambda_n) + F'(\lambda_n)\varphi^2(\pi, \lambda_n), \quad n \geq 1. \quad (3.3)$$

The numbers $\{\lambda_n, \gamma_n\}_{n \geq 1}$ are called the spectral data of the problem (1.1)–(1.2).

This paper is focused on the two spectral problems:

Direct Problem. Given (σ, f, F) , find $\{\lambda_n, \gamma_n\}_{n \geq 1}$.

Inverse Problem. Given $\{\lambda_n, \gamma_n\}_{n \geq 1}$, find (σ, f, F) .

The uniqueness of the inverse problem solution for $\sigma \in \dot{L}_2[0, \pi]$ was established in [11]. Furthermore, for $\sigma \in \dot{W}_2^{\alpha}[0, \pi]$, we get the following result.

Theorem 3.1 (Spectral data characterization). *Let $\alpha \in [0, \frac{1}{2})$ be real and $M, N \geq -1$ be integers. For real numbers $\{\lambda_n, \gamma_n\}_{n \geq 1}$ to be the spectral data of a problem $\mathcal{L}(\sigma, f, F)$ with $\sigma \in \dot{W}_2^{\alpha}[0, \pi]$, $f \in \mathcal{R}_M$, and $F \in \mathcal{R}_N$, the following conditions are necessary and sufficient:*

$$\lambda_n < \lambda_{n+1}, \quad \gamma_n > 0, \quad n \geq 1, \quad (3.4)$$

$$\sqrt{\lambda_n} = n - \frac{M+N}{2} - 1 + \kappa_n, \quad \gamma_n = \frac{\pi}{2} n^{2M} (1 + \beta_n), \quad n \geq 1, \quad (3.5)$$

where $\{\kappa_n\}$ and $\{\beta_n\}$ belong to l_2^{α} .

Note that, if the sequences $\{\lambda_n\}$ and $\{\gamma_n\}$ satisfy the asymptotics (3.5), then one can uniquely determine M and N , and then find $\{\kappa_n\}$ and $\{\beta_n\}$.

For $\alpha = 0$, Theorem 3.1 was proved in [11]. For $\alpha \in (0, \frac{1}{2})$, the proof is analogous. We only need to establish the correspondence between the spaces $W_2^{\alpha}[0, \pi]$ for σ and l_2^{α} for the remainder sequences $\{\kappa_n\}$ and $\{\beta_n\}$. It is worth mentioning that, for the cases of the Dirichlet and the Neumann-Dirichlet boundary conditions ($M \in \{-1, 0\}$, $N = -1$) and $\alpha > 0$, the spectral data characterization has been obtained in [25, 47]. Theorem 3.1 is consistent with those results for $\alpha \in (0, \frac{1}{2})$ and generalizes them to the case of rational Herglotz-Nevanlinna functions in the boundary conditions.

In order to formulate our main results on the uniform stability, we need to introduce metric spaces of problem parameters and of spectral data. Consider the metric space

$$\mathcal{P}^{\alpha, M, N} := \dot{W}_2^{\alpha}[0, \pi] \times \mathcal{R}_M \times \mathcal{R}_N$$

of triples $P = (\sigma, f, F)$ with the distance

$$\mathbf{d}_\alpha(P_1, P_2) = \|\sigma_1 - \sigma_2\|_\alpha + \|c(f_1) - c(f_2)\|_{\mathbb{R}^{M+1}} + \|c(F_1) - c(F_2)\|_{\mathbb{R}^{N+1}}, \quad (3.6)$$

where $P_i = (\sigma_i, f_i, F_i)$, $i = 1, 2$. Furthermore, introduce the metric space of real-valued sequences $\mathfrak{S} = \{\lambda_n, \gamma_n\}_{n \geq 1}$ satisfying the asymptotics (3.5) for fixed $\alpha \in [0, \frac{1}{2})$, $M, N \geq -1$, with the distance

$$\varrho_\alpha(\mathfrak{S}_1, \mathfrak{S}_2) = \|\{\kappa_{n,1} - \kappa_{n,2}\}\|_\alpha + \|\{\beta_{n,1} - \beta_{n,2}\}\|_\alpha, \quad (3.7)$$

where $\{\kappa_{n,i}\}$ and $\{\beta_{n,i}\}$ are the remainders from the asymptotics (3.5) for $\mathfrak{S}_i = \{\lambda_{n,i}, \gamma_{n,i}\}$, $i = 1, 2$.

Next, let us define sets in the introduced metric spaces, on which the uniform stability of direct and inverse spectral mappings will be established. Fix $\alpha \in (0, \frac{1}{2})$ and $M, N \geq -1$ such that $M + N$ is even. For $Q > 0$ and $\delta > 0$, introduce the set

$$\mathcal{P}_{Q,\delta}^{\alpha,M,N} := \{P = (\sigma, f, F) : \sigma \in \mathring{W}_2^\alpha[0, \pi], \|\sigma\|_\alpha \leq Q, f \in \mathcal{R}_{M,Q,\delta}, F \in \mathcal{R}_{N,Q,\delta}, \lambda_1(P) \geq 1\}. \quad (3.8)$$

We impose the technical restriction $\lambda_1(P) \geq 1$ according to the previous studies [25, 27], since it simplifies formulations. In general, the spectrum of the boundary value problem $\mathcal{L}(P)$ is bounded from below.

For $R > 0$ and $\varepsilon > 0$, denote by $\mathcal{B}_{R,\varepsilon}^{\alpha,M,N}$ the set of sequences $\mathfrak{S} = \{\lambda_n, \gamma_n\}_{n \geq 1}$ satisfying the asymptotics (3.5) and the additional requirements:

$$\lambda_1 \geq 1, \quad \sqrt{\lambda_{n+1}} - \sqrt{\lambda_n} \geq \varepsilon, \quad n \geq 1, \quad \|\{\kappa_n\}_{n \geq 1}\|_\alpha \leq R, \quad (3.9)$$

$$1 + \beta_n \geq \varepsilon, \quad n \geq 1, \quad \|\{\beta_n\}_{n \geq 1}\|_\alpha \leq R. \quad (3.10)$$

For brevity, we sometimes omit the indices α, M, N : $\mathcal{P}_{Q,\delta} := \mathcal{P}_{Q,\delta}^{\alpha,M,N}$, $\mathcal{B}_{R,\varepsilon} := \mathcal{B}_{R,\varepsilon}^{\alpha,M,N}$.

Denote by \mathcal{S} the spectral transform that maps $P = (\sigma, f, F)$ to the corresponding spectral data $\mathfrak{S} = \{\lambda_n, \gamma_n\}_{n \geq 1}$. Our main results on the uniform stability are formulated as follows.

Theorem 3.2 (Uniform stability for the direct problem). *Let $Q > 0$ and $\delta > 0$. Then \mathcal{S} maps $\mathcal{P}_{Q,\delta}$ into $\mathcal{B}_{R,\varepsilon}$, where $R > 0$ and $\varepsilon > 0$ depend on Q and δ . Moreover, the direct spectral transform \mathcal{S} is Lipschitz continuous on $\mathcal{P}_{Q,\delta}$, that is,*

$$\varrho_\alpha(\mathfrak{S}_1, \mathfrak{S}_2) \leq C(Q, \delta) \mathbf{d}_\alpha(P_1, P_2) \quad (3.11)$$

for any $P_i \in \mathcal{P}_{Q,\delta}$, $\mathfrak{S}_i := \mathcal{S}(P_i)$, $i = 1, 2$.

Theorem 3.3 (Uniform stability for the inverse problem). *Let $R > 0$ and $\varepsilon > 0$. Then \mathcal{S}^{-1} maps $\mathcal{B}_{R,\varepsilon}$ into $\mathcal{P}_{Q,\delta}$, where $Q > 0$ and $\delta > 0$ depend on R and ε . Moreover, the inverse spectral transform \mathcal{S}^{-1} is Lipschitz continuous on $\mathcal{B}_{R,\varepsilon}$, that is,*

$$\mathbf{d}_\alpha(P_1, P_2) \leq C(R, \varepsilon) \varrho_\alpha(\mathfrak{S}_1, \mathfrak{S}_2) \quad (3.12)$$

for any $\mathfrak{S}_i \in \mathcal{B}_{R,\varepsilon}$, $P_i := \mathcal{S}^{-1}(\mathfrak{S}_i)$, $i = 1, 2$.

Theorems 3.2 and 3.3 generalize the results of [25] to the problem (1.1)–(1.2) with rational Herglotz-Nevanlinna functions in the boundary conditions. Their proofs rely on the Lipschitz continuity of the Darboux-type transforms constructed in [11]. Note that the parameter $\delta > 0$ is unimportant for $M, N \in \{-1, 0\}$. Therefore, this constraint does not appear in the previous studies [25–27] for constant coefficients.

Theorems 3.2 and 3.3 establish the unconditional uniform stability for the direct and the inverse problems, respectively. This means that the constants in the estimates (3.11) and (3.12) depend on the constraints for the initial data of the corresponding problem (i.e. on P in (3.11) and on \mathfrak{S} in (3.12)). Using Theorems 3.2 and 3.3 together, we get the following corollary on the conditional uniform stability for the direct problem and the inverse problem. Therein, the constants in the estimates depend on the constraints for the solution.

Corollary 3.4. (i) For any $P_i \in \mathcal{P}_{Q,\delta}$ and $\mathfrak{S}_i := \mathcal{S}(P_i)$, $i = 1, 2$, there holds

$$\mathbf{d}_\alpha(P_1, P_2) \leq C(Q, \delta) \varrho_\alpha(\mathfrak{S}_1, \mathfrak{S}_2).$$

(ii) For any $\mathfrak{S}_i \in \mathcal{B}_{R,\varepsilon}$ and $P_i := \mathcal{S}^{-1}(\mathfrak{S}_i)$, there holds

$$\varrho_\alpha(\mathfrak{S}_1, \mathfrak{S}_2) \leq C(R, \varepsilon) \mathbf{d}_\alpha(P_1, P_2).$$

Suppose that we know a finite number of spectral data $\{\lambda_n, \gamma_n\}_{n=1}^m$ for the problem $\mathcal{L}(P)$, $P = (\sigma, f, F) \in \mathcal{P}^{\alpha, M, N}$. Then, we can complete them as follows:

$$\lambda_{n,m} := \begin{cases} \lambda_n, & n \leq m, \\ (n - \frac{M+N}{2} - 1)^2, & n > m, \end{cases} \quad \gamma_{n,m} := \begin{cases} \gamma_n, & n \leq m, \\ \frac{\pi}{2} n^{2M}, & n > m, \end{cases} \quad \mathfrak{S}_m := \{\lambda_{n,m}, \gamma_{n,m}\}_{n \geq 1}, \quad (3.13)$$

and obtain the finite data approximation $P_m = (\sigma_m, f_m, F_m) := \mathcal{S}^{-1}(\mathfrak{S}_m)$ of P .

Theorem 3.5. Let $0 < \alpha_1 < \alpha_2 < \frac{1}{2}$, $M, N \geq -1$ be fixed, and $Q, \delta > 0$. Then, for every $P \in \mathcal{P}_{Q,\delta}^{\alpha_2, M, N}$, the finite data approximation satisfies the uniform stability estimate

$$\mathbf{d}_{\alpha_1}(P, P_m) \leq C(Q, \delta) m^{\alpha_1 - \alpha_2} \quad (3.14)$$

for all sufficiently large m .

Note that the right-hand side of (3.14) tends to zero as $m \rightarrow \infty$ uniformly by $P \in \mathcal{P}_{Q,\delta}$.

Finally, consider the case when the finite spectral data are known with an error at most $\varepsilon > 0$:

$$|\tilde{\lambda}_n - \lambda_n| \leq \varepsilon, \quad |\tilde{\gamma}_n - \gamma_n| \leq \varepsilon, \quad n = \overline{1, m}. \quad (3.15)$$

Then, find the finite data approximation $\tilde{P}_m := \mathcal{S}^{-1}(\tilde{\mathfrak{S}}_m)$ by using the completion $\tilde{\mathfrak{S}}_m = \{\tilde{\lambda}_{n,m}, \tilde{\gamma}_{n,m}\}_{n \geq 1}$:

$$\tilde{\lambda}_{n,m} := \begin{cases} \tilde{\lambda}_n, & n \leq m, \\ (n - \frac{M+N}{2} - 1)^2, & n > m, \end{cases} \quad \tilde{\gamma}_{n,m} := \begin{cases} \tilde{\gamma}_n, & n \leq m, \\ \frac{\pi}{2} n^{2M}, & n > m. \end{cases}$$

We obtain the following theorem about the uniform stability of the approximation \tilde{P}_m .

Theorem 3.6. Let $0 < \alpha_1 < \alpha_2 < \frac{1}{2}$, $M, N \geq -1$ be fixed, and $Q, \delta > 0$. Then, there exists $\varepsilon_0 > 0$ such that, for every $P \in \mathcal{P}_{Q,\delta}^{\alpha_2, M, N}$, $\varepsilon \in (0, \varepsilon_0]$, and all sufficiently large m , the following estimate holds:

$$\mathbf{d}_{\alpha_1}(P, \tilde{P}_m) \leq C(Q, \delta) \left(\varepsilon m^{\alpha_1 + \frac{1}{2} - \min\{1, 2M\}} + m^{\alpha_1 - \alpha_2} \right). \quad (3.16)$$

4 Auxiliary lemmas

In this section, we prove several lemmas on the Lipschitz continuity by $P \in \mathcal{P}_{Q,\delta}$ for some auxiliary values.

Lemma 4.1. *Let $y(x, \lambda)$ be the solution of equation (1.1) satisfying the initial conditions*

$$y(x_0) = a, \quad y_\sigma^{[1]}(x_0) = b, \quad x_0 \in [0, \pi]. \quad (4.1)$$

Then, the maps $(\sigma, a, b, \lambda) \mapsto y$ and $(\sigma, a, b, \lambda) \mapsto y_\sigma^{[1]}$ from $L_2[0, \pi] \times \mathbb{R}^3$ to $C[0, \pi]$ are analytic and Lipschitz continuous on bounded sets.

Proof. The initial value problem (1.1), (4.1) is equivalent to the first-order system

$$\frac{d}{dx} \begin{bmatrix} y \\ y_\sigma^{[1]} \end{bmatrix} = \begin{bmatrix} \sigma & 1 \\ -(\sigma^2 + \lambda) & -\sigma \end{bmatrix} \begin{bmatrix} y \\ y_\sigma^{[1]} \end{bmatrix}, \quad \begin{bmatrix} y(x_0) \\ y_\sigma^{[1]}(x_0) \end{bmatrix} = \begin{bmatrix} a \\ b \end{bmatrix}.$$

Reducing this system to an integral equation and solving it by iterations yield the claim. Note that the continuity for solutions of equation (1.1) with respect to σ has been proved in [44]. \square

Lemma 4.2. *The set $\mathcal{P}_{Q,\delta}^{\alpha,M,N}$ for $\alpha > 0$ is compact in $\mathcal{P}^{0,M,N}$.*

Proof. By virtue of part (i) of Proposition 2.1, the ball $\{\sigma \in \dot{W}_2^\alpha[0, \pi]: \|\sigma\|_\alpha \leq Q\}$ is compact in $L_2[0, \pi]$. This fact together with Lemma 2.2 and the definitions (3.6) and (3.8) yield the claim. \square

Note that $\mathcal{P}_{Q,\delta}^{\alpha,M,N}$ is not compact in $\mathcal{P}^{\alpha,M,N}$. The compact embedding $W_2^\alpha[0, \pi] \subset L_2[0, \pi]$ is crucial.

Lemma 4.3. *Let $Q > 0$ and $\delta > 0$ be fixed. For any $P_i \in \mathcal{P}_{Q,\delta}$, $i = 1, 2$, there holds*

$$|\lambda_1(P_1) - \lambda_1(P_2)| + |\gamma_1(P_1) - \gamma_1(P_2)| \leq C(Q, \delta) \mathbf{d}_0(P_1, P_2). \quad (4.2)$$

Thus, the maps $\lambda_1(P)$ and $\gamma_1(P)$ are Lipschitz continuous on $\mathcal{P}_{Q,\delta}^{\alpha,M,N}$ in the metric space $\mathcal{P}^{0,M,N}$.

Proof. According to Lemma 4.1 and the initial conditions (3.1), the maps $(\sigma, c(f), \lambda) \mapsto \varphi(\cdot, \lambda)$ and $(\sigma, c(f), \lambda) \mapsto \varphi_\sigma^{[1]}(\cdot, \lambda)$ are analytic. Hence, the characteristic function $\chi(\lambda)$ given by (3.2) is analytic in σ , $c(f)$, $c(F)$, and λ . Therefore, $\chi(\lambda)$ is Lipschitz continuous for $P = (\sigma, f, F) \in \mathcal{P}_{Q,\delta}$, and λ on bounded sets. Obviously, the first zero $\lambda_1(P)$ of $\chi(\lambda)$ is continuous by P .

Let us show the Lipschitz continuity of λ_1 using the simplicity of this eigenvalue. Consider a small perturbation $\tilde{P} = (\tilde{\sigma}, \tilde{f}, \tilde{F})$ of $P = (\sigma, f, F)$ in $\mathcal{P}_{Q,\delta}$. The Taylor formula implies

$$\chi(\tilde{\lambda}_1) = \chi(\lambda_1) + \chi'(\lambda_1)(\tilde{\lambda}_1 - \lambda_1) + \frac{1}{2}\chi''(\xi)(\tilde{\lambda}_1 - \lambda_1)^2,$$

where $\xi = (1 - \theta)\lambda_1 + \theta\tilde{\lambda}_1$, $\theta \in (0, 1)$. Taking $\chi(\lambda_1) = \tilde{\chi}(\tilde{\lambda}_1) = 0$ into account, we derive

$$(\tilde{\lambda}_1 - \lambda_1) \left(\chi'(\lambda_1) + \frac{1}{2}\chi''(\xi)(\tilde{\lambda}_1 - \lambda_1) \right) = \chi(\tilde{\lambda}_1) - \tilde{\chi}(\tilde{\lambda}_1). \quad (4.3)$$

The Lipschitz continuity of $\chi(\lambda)$ implies

$$|\chi(\tilde{\lambda}_1) - \tilde{\chi}(\tilde{\lambda}_1)| \leq C(Q, \delta) \mathbf{d}_0(P, \tilde{P}). \quad (4.4)$$

Since $\chi(\lambda)$ is analytic and $\lambda_1(P)$ is continuous, then $\chi'(\lambda_1)$ is a continuous function on the set $\mathcal{P}_{Q,\delta}$, which is compact according to Lemma 4.2. Due to the simplicity of λ_1 , we have $\chi'(\lambda_1) \neq 0$, so

$$|\chi'(\lambda_1)| \geq \varepsilon > 0, \quad \varepsilon = \varepsilon(Q, \delta). \quad (4.5)$$

Analogously, we get

$$|\chi''(\xi)| \leq C(Q, \delta), \quad \text{for } |\xi| \leq R(Q, \delta). \quad (4.6)$$

Combining (4.3), (4.4), (4.5), and (4.6), we arrive at the estimate

$$|\lambda_1 - \tilde{\lambda}_1| \leq C(Q, \delta) \mathbf{d}_0(P, \tilde{P}). \quad (4.7)$$

Remark 2.2 in [11] implies that $\lambda_1 < \min\{\hat{\pi}(f), \hat{\pi}(F)\}$. This together with (2.3) yield $f_\downarrow(\lambda_1) > 0$ and $F_\downarrow(\lambda_1) > 0$. Since $f_\downarrow(\lambda_1(P))$ and $F_\downarrow(\lambda_1(P))$ are continuous on $\mathcal{P}_{Q,\delta}$, then $f_\downarrow(\lambda_1) \geq \varepsilon$ and $F_\downarrow(\lambda_1) \geq \varepsilon$, where $\varepsilon = \varepsilon(Q, \delta)$. The latter estimates together with (3.3), (4.7), and the Lipschitz continuity for $\varphi(\cdot, \lambda)$ imply the Lipschitz continuity for $\gamma_1(P)$, which concludes the proof. \square

Remark 4.4. The assertion similar to Lemma 4.3 holds for $\lambda_n(P)$ and $\gamma_n(P)$ for each fixed index $n \geq 2$.

For reals μ, τ, ρ and a rational function $f \in \mathcal{R}$, consider the transform $\Theta: (\mu, \tau, \rho, f) \mapsto \hat{f}$, which was defined in [11] by the formula

$$\hat{f}(\lambda) := \frac{\mu - \lambda}{f(\lambda) - \tau} + \rho.$$

The properties of this transform are described by the following proposition from [11].

Proposition 4.5 ([11]). *Suppose that $\mu < \hat{\pi}(f)$ and $\tau \geq f(\mu)$ if $\text{ind } f \geq 0$. Then $\hat{f} \in \mathcal{R}$. Moreover, if $\tau = f(\mu)$, then $\text{ind } \hat{f} = \text{ind } f - 1$ and*

$$\hat{f}_\uparrow(\lambda) = \frac{\rho f_\uparrow(\lambda) - (\lambda - \mu + \tau\rho)f_\downarrow(\lambda)}{\lambda - \mu}, \quad \hat{f}_\downarrow(\lambda) = \frac{f_\uparrow(\lambda) - \tau f_\downarrow(\lambda)}{\lambda - \mu}; \quad (4.8)$$

if $\tau > f(\mu)$, then $\text{ind } \hat{f} = \text{ind } f + 1$ and

$$\hat{f}_\uparrow(\lambda) = -\rho f_\downarrow(\lambda) + (\lambda - \mu + \tau\rho)f_\downarrow(\lambda), \quad \hat{f}_\downarrow(\lambda) = -f_\uparrow(\lambda) + \tau f_\downarrow(\lambda). \quad (4.9)$$

Lemma 4.6. *Suppose that $\hat{f} = \Theta(\mu, \tau, \rho, f)$, where $f \in \mathcal{R}_{M,Q,\delta}$ for fixed M, Q , and δ , parameters (μ, τ, ρ) belong to a compact set satisfying the hypothesis of Proposition 4.5 and either $\tau = f(\mu)$ or $\tau > f(\mu)$. Then, the coefficients $c(\hat{f})$ are Lipschitz continuous with respect to μ, τ, ρ , and $c(f)$.*

Proof. For the case $\tau > f(\mu)$, the assertion of the lemma trivially follows from (4.9). For $\tau = f(\mu)$, the Lipschitz continuity is proved by interpolation argument. The interpolation nodes are chosen to be separated from μ and from each other. \square

5 Transforms

In this section, we introduce the Darboux-type transforms that have been constructed in [11] and prove their Lipschitz continuity.

Define the domain

$$\mathcal{P} := \mathring{L}_2(0, \pi) \times \mathcal{R} \times \mathcal{R}.$$

For $P = (\sigma, f, F) \in \mathcal{P}$, $\Lambda \in \mathbb{R}$, $v \in \mathcal{D}_\sigma$ (the domain \mathcal{D}_σ is defined by (1.3)), $v(x) \neq 0$ for $x \in [0, \pi]$, consider the transform

$$\mathcal{T}: (P, \Lambda, v) \mapsto \widehat{P} = (\widehat{\sigma}, \widehat{f}, \widehat{F})$$

defined by the following rules:

$$\widehat{\sigma} := \sigma - \frac{2v'}{v} + \frac{2}{\pi} \ln \frac{v(\pi)}{v(0)}, \quad (5.1)$$

$$\widehat{f} := \Theta \left(\Lambda, -\frac{v_\sigma^{[1]}(0)}{v(0)}, -\frac{v_\sigma^{[1]}(0)}{v(0)} + \frac{2}{\pi} \ln \frac{v(\pi)}{v(0)}, f \right), \quad (5.2)$$

$$\widehat{F} := \Theta \left(\Lambda, \frac{v_\sigma^{[1]}(\pi)}{v(\pi)}, \frac{v_\sigma^{[1]}(\pi)}{v(\pi)} - \frac{2}{\pi} \ln \frac{v(\pi)}{v(0)}, F \right). \quad (5.3)$$

Denote by $\psi(x, \lambda)$ and $z(x, \lambda, \rho)$ the solutions of equation (1.1) satisfying the initial conditions

$$\psi(\pi, \lambda) = F_\downarrow(\lambda), \quad \psi_\sigma^{[1]}(\pi, \lambda) = F_\uparrow(\lambda), \quad z(0, \lambda, \rho) = 1, \quad z_\sigma^{[1]}(0, \lambda, \rho) = -\rho. \quad (5.4)$$

Introduce the transforms

$$\begin{aligned} \mathbf{T}_-(P) &:= \mathcal{T}(P, \lambda_1, \varphi(\cdot, \lambda_1)), & \text{dom}(\mathbf{T}_-) &= \{P \in \mathcal{P}: f \neq \infty, F \neq \infty\}, \\ \mathbf{T}_{-+}(P) &:= \mathcal{T}(P, \lambda_1 - 2, \varphi(\cdot, \lambda_1 - 2)), & \text{dom}(\mathbf{T}_{-+}) &= \{P \in \mathcal{P}: f \neq \infty\}, \\ \mathbf{T}_{+-}(P) &:= \mathcal{T}(P, \lambda_1 - 2, \psi(\cdot, \lambda_1 - 2)), & \text{dom}(\mathbf{T}_{+-}) &= \{P \in \mathcal{P}: F \neq \infty\}, \\ \mathbf{T}_+(\mu, \nu, P) &:= \mathcal{T}(P, \mu, u), & \text{dom}(\mathbf{T}_+) &= \{(\mu, \nu, P): P \in \mathcal{P}, \mu < \lambda_1, \nu > 0\}, \end{aligned}$$

where $\lambda_1 = \lambda_1(P)$ and

$$u(x) := z(x, \lambda_1, \rho), \quad \rho := \frac{\nu \varkappa + f_\uparrow(\mu)(\varkappa f_\downarrow(\mu) - f_\uparrow(\mu))}{\nu + f_\downarrow(\mu)(\varkappa f_\downarrow(\mu) - f_\uparrow(\mu))}, \quad \varkappa := -\frac{\psi_\sigma^{[1]}(0, \mu)}{\psi(0, \mu)}. \quad (5.5)$$

Our notations for the defined transforms have the following meaning:

- \mathbf{T}_- removes the first eigenvalue λ_1 , decreases $\text{ind } f$ and $\text{ind } F$.
- \mathbf{T}_{-+} does not change the spectrum, decreases $\text{ind } f$, and increases $\text{ind } F$.
- \mathbf{T}_{+-} does not change the spectrum, decreases $\text{ind } f$, and increases $\text{ind } F$.
- \mathbf{T}_+ adds the eigenvalue μ with the corresponding norming constant ν , increases $\text{ind } f$ and $\text{ind } F$.

More formally, the above properties are described in the following proposition.

Proposition 5.1 ([11]). *The transforms \mathbf{T}_- , \mathbf{T}_{-+} , \mathbf{T}_{+-} , and \mathbf{T}_+ are well-defined on their domains as mappings to \mathcal{P} . They change the indices of rational functions in the following way:*

$$\begin{aligned}\mathbf{T}_-: & \quad \text{ind } \widehat{f} = \text{ind } f - 1, & \text{ind } \widehat{F} = \text{ind } F - 1, \\ \mathbf{T}_{-+}: & \quad \text{ind } \widehat{f} = \text{ind } f - 1, & \text{ind } \widehat{F} = \text{ind } F + 1, \\ \mathbf{T}_{+-}: & \quad \text{ind } \widehat{f} = \text{ind } f + 1, & \text{ind } \widehat{F} = \text{ind } F - 1, \\ \mathbf{T}_+: & \quad \text{ind } \widehat{f} = \text{ind } f + 1, & \text{ind } \widehat{F} = \text{ind } F + 1.\end{aligned}$$

Moreover, \mathbf{T}_- and \mathbf{T}_+ are inverses of each other in the following sense: if $P \in \text{dom } \mathbf{T}_-$ and $\widehat{P} = \mathbf{T}_-(P)$, then

$$\mathbf{T}_+(\lambda_1(P), \gamma_1(P), \widehat{P}) = P, \quad (5.6)$$

and conversely, if $(\mu, \nu, P) \in \text{dom } \mathbf{T}_+$, then $\mathbf{T}_-\mathbf{T}_+(\mu, \nu, P) = P$. Furthermore, $\mathbf{T}_{-+} = \mathbf{T}_{+-}^{-1}$.

Let the problem \mathcal{L} has the spectral data $\mathcal{S}(P) = \{\lambda_n, \gamma_n\}_{n \geq 1}$. Then

$$\begin{aligned}\mathcal{S}(\mathbf{T}_-(P)) &= \{\lambda_n, \gamma_n(\lambda_n - \lambda_1)^{-1}\}_{n \geq 2}, \\ \mathcal{S}(\mathbf{T}_{-+}(P)) &= \{\lambda_n, \gamma_n(\lambda_n - \lambda_1 + 2)^{-1}\}_{n \geq 1}, \\ \mathcal{S}(\mathbf{T}_{+-}(P)) &= \{\lambda_n, \gamma_n(\lambda_n - \lambda_1 + 2)\}_{n \geq 1}, \\ \mathcal{S}(\mathbf{T}_+(\mu, \nu, P)) &= \{\mu, \nu\} \cup \{\lambda_n, \gamma_n(\lambda_n - \lambda_1)\}_{n \geq 1}.\end{aligned} \quad (5.7)$$

Proceed to studying the Lipschitz continuity of the introduced transforms.

Lemma 5.2. *Suppose that $\mathbf{T} \in \{\mathbf{T}_-, \mathbf{T}_{-+}, \mathbf{T}_{+-}\}$, fixed integers M, N are such that $\mathring{L}_2(0, \pi) \times \mathcal{R}_M \times \mathcal{R}_N \subset \text{dom } \mathbf{T}$, and $\alpha \in (0, \frac{1}{2})$ is fixed. Then \mathbf{T} maps $\mathcal{P}_{Q, \delta}^{\alpha, M, N}$ into $\mathcal{P}_{\widehat{Q}, \widehat{\delta}}^{\alpha, \widehat{M}, \widehat{N}}$, where $\widehat{Q} > 0$ and $\widehat{\delta} > 0$ depend on Q and δ . Moreover,*

$$\mathbf{d}_\alpha(\widehat{P}_1, \widehat{P}_2) \leq C(Q, \delta) \mathbf{d}_\alpha(P_1, P_2), \quad (5.8)$$

for any $P_i \in \mathcal{P}_{Q, \delta}$, $\widehat{P}_i = \mathbf{T}(P_i)$, $i = 1, 2$. Thus, the maps \mathbf{T}_- , \mathbf{T}_{-+} , and \mathbf{T}_{+-} are Lipschitz continuous on $\mathcal{P}_{Q, \delta}$.

Proof. For definiteness, let us prove the lemma for $\mathbf{T} = \mathbf{T}_-$. The proof for \mathbf{T}_{-+} and \mathbf{T}_{+-} is analogous and even simpler. Suppose that $P \in \mathcal{P}_{Q, \delta} \subset \text{dom } \mathbf{T}_-$. By Lemma 4.3, the first eigenvalue $\lambda_1(P)$ is Lipschitz continuous on $\mathcal{P}_{Q, \delta}$. Taking (3.1) and Lemma 4.1 into account, we conclude that $v(\cdot) = \varphi(\cdot, \lambda_1)$ and $v_\sigma^{[1]}(\cdot) = \varphi_\sigma^{[1]}(\cdot, \lambda_1)$ are Lipschitz continuous on $\mathcal{P}_{Q, \delta}$ as maps from $L_2[0, \pi] \times \mathbb{R}^{M+1} \times \mathbb{R}^{N+1}$ to $C[0, \pi]$. This implies

$$|v(x)| + |v_\sigma^{[1]}(x)| \leq C(Q, \delta), \quad \text{for all } P \in \mathcal{P}_{Q, \delta}, x \in [0, \pi]. \quad (5.9)$$

The previous argument yields that the map $(x, P) \mapsto v(x)$ is continuous on the set $[0, \pi] \times \mathcal{P}_{Q, \delta}$, which is compact in the sense of Lemma 4.2. Therefore, the image of this map is compact in \mathbb{R} .

Note that $v(x)$ is the eigenfunction of the problem $\mathcal{L}(\sigma, f, F)$ corresponding to the first eigenvalue λ_1 . Since $f, F \neq \infty$, then the oscillation theory implies $v(x) \neq 0$ for $x \in [0, \pi]$ (see, e.g., [11, Lemma 2.3]). Due to the compactness of the set $\{v(x)\}$, we obtain the estimate

$$|v(x)| \geq \varepsilon > 0, \quad \text{for all } P \in \mathcal{P}_{Q, \delta}, x \in [0, \pi], \quad (5.10)$$

where $\varepsilon = \varepsilon(Q, \delta)$.

Next, put $w := v_\sigma^{[1]}$ and note that

$$v' = w + \sigma v. \quad (5.11)$$

Substituting (5.11) into (5.1), we get

$$\widehat{\sigma} = -\sigma - \frac{2w}{v} + \frac{2}{\pi} \ln \frac{v(\pi)}{v(0)}. \quad (5.12)$$

Consider two triples $P_i = (\sigma_i, f_i, F_i) \in \mathcal{P}_{Q,\delta}$, $i = 1, 2$. Since $v_i \in \mathcal{D}_{\sigma_i}$ (1.3), then v_i and $w_i := (v_i)_{\sigma_i}^{[1]}$ belong to $W_1^1[0, \pi]$. Let us show that

$$\|v_1 - v_2\|_{W_1^1[0,\pi]} \leq C(Q, \delta) \mathbf{d}_\alpha(P_1, P_2), \quad \|w_1 - w_2\|_{W_1^1[0,\pi]} \leq C(Q, \delta) \mathbf{d}_\alpha(P_1, P_2). \quad (5.13)$$

Recall that

$$\|u\|_{W_1^1[0,\pi]} = \|u\|_{L_1[0,\pi]} + \|u'\|_{L_1[0,\pi]}, \quad \|u\|_{C[0,\pi]} = \max_{x \in [0,\pi]} |u(x)|.$$

Using (5.9), (5.11), the Lipschitz continuity of $v(\cdot)$ and $w(\cdot) = v^{[1]}(\cdot)$ by $P \in \mathcal{P}_{Q,\delta}$, and the continuous embeddings $W_2^\alpha[0, \pi] \subset L_2[0, \pi] \subset L_1[0, \pi]$, we get

$$\begin{aligned} \|v_1 - v_2\|_{L_1[0,\pi]} &\leq \pi \|v_1 - v_2\|_{C[0,\pi]} \leq C(Q, \delta) \mathbf{d}_\alpha(P_1, P_2), \\ \|v_1' - v_2'\|_{L_1[0,\pi]} &\leq \|w_1 - w_2\|_{L_1[0,\pi]} + \|\sigma_1 - \sigma_2\|_\alpha \|v_1\|_{C[0,\pi]} + \|\sigma_2\|_\alpha \|v_1 - v_2\|_{C[0,\pi]} \\ &\leq C(Q, \delta) \mathbf{d}_\alpha(P_1, P_2). \end{aligned}$$

Analogously, we obtain $\|w_1 - w_2\|_{L_1[0,\pi]} \leq C(Q, \delta) \mathbf{d}_\alpha(P_1, P_2)$. Using equation (1.1), we get

$$w_i' = -\sigma_i w_i - \sigma_i^2 w_i - \lambda_1(P_i) v_i, \quad i = 1, 2. \quad (5.14)$$

Using (5.14), the Lipschitz continuity of $v(\cdot)$, $w(\cdot)$, and λ_1 by $P \in \mathcal{P}_{Q,\delta}$, and the estimate $\|\sigma_i\|_0 \leq \|\sigma_i\|_\alpha \leq Q$, we derive

$$\begin{aligned} \|w_1' - w_2'\|_{L_1[0,\pi]} &\leq C(Q, \delta) (\|\sigma_1 - \sigma_2\|_0 + \|w_1 - w_2\|_{C[0,\pi]} + \|v_1 - v_2\|_{C[0,\pi]} + |\lambda_1(P_1) - \lambda_2(P_2)|) \\ &\leq C(Q, \delta) \mathbf{d}_\alpha(P_1, P_2). \end{aligned}$$

Summarizing the above arguments, we arrive at (5.13). Thus, $P \mapsto v$ and $P \mapsto w$ are Lipschitz continuous as maps of $\mathcal{P}_{Q,\delta}$ to $W_1^1[0, \pi]$. Recall that $W_1^1[0, \pi]$ is compactly embedded in $W_2^\alpha[0, \pi]$ due to part (ii) of Proposition 2.1. Therefore, using (5.9), (5.10), (5.12), and (5.13), we conclude that

$$\widehat{\sigma} \in W_2^\alpha[0, \pi], \quad \|\widehat{\sigma}\|_\alpha \leq C(Q, \delta), \quad (5.15)$$

$$\|\widehat{\sigma}_1 - \widehat{\sigma}_2\|_\alpha \leq C(Q, \delta) \mathbf{d}_\alpha(P_1, P_2). \quad (5.16)$$

In other words, the map $P \mapsto \widehat{\sigma}$ is Lipschitz continuous from $\mathcal{P}_{Q,\delta}$ to $W_2^\alpha[0, \pi]$. The equality $\int_0^\pi \widehat{\sigma}(x) dx = 0$ follows from Proposition 5.1.

Proceed to \widehat{f} and \widehat{F} . By virtue of Proposition 5.1, we have $\widehat{f} \in \mathcal{R}_{\widehat{M}}$ and $\widehat{F} \in \mathcal{R}_{\widehat{N}}$, where $\widehat{M} := M - 1$ and $\widehat{N} := N - 1$ for $\mathbf{T} = \mathbf{T}_-$. Furthermore, due to [11, Remark 2.2] and (5.7), we have

$$\min\{\widehat{\pi}(\widehat{f}), \widehat{\pi}(\widehat{F})\} > \lambda_2 > \lambda_1 \geq 1.$$

Note that the coefficients $\Lambda = \lambda_1, \frac{w(0)}{v(0)}, \frac{w(\pi)}{v(\pi)}$, and $\ln \frac{v(\pi)}{v(0)}$, which participate in formulas (5.2) and (5.3), are Lipschitz continuous by $P \in \mathcal{P}_{Q,\delta}$ according to the above arguments. Applying Lemma 4.6, we conclude that the maps $P \mapsto c(\widehat{f})$ and $P \mapsto c(\widehat{F})$ are Lipschitz continuous from $\mathcal{P}_{Q,\delta}$ to $\mathbb{R}^{\widehat{M}+1}$ and $\mathbb{R}^{\widehat{N}+1}$, respectively. This together with (5.16) and (3.6) imply the estimate (5.8). Furthermore, the images of these maps $\{c(\widehat{f}): P \in \mathcal{P}_{Q,\delta}\}$ and $\{c(\widehat{F}): P \in \mathcal{P}_{Q,\delta}\}$ are compact. Therefore, by Lemma 2.3 they are subsets of $\mathcal{R}_{\widehat{M},\widehat{Q},\widehat{\delta}}$ and $\mathcal{R}_{\widehat{N},\widehat{Q},\widehat{\delta}}$, respectively, where $\widehat{Q} > 0$ and $\widehat{\delta} > 0$ depend on Q and δ . This together with (5.15) and (3.8) imply $\widehat{P} = \mathbf{T}(P) \in \mathcal{P}_{\widehat{Q},\widehat{\delta}}^{\alpha,\widehat{M},\widehat{N}}$ and so conclude the proof. \square

The following lemma presents the result analogous to Lemma 5.2 for \mathbf{T}_+ .

Lemma 5.3. *Let $\alpha \in (0, \frac{1}{2})$ and integers $M, N \geq -1$ be fixed. Then \mathbf{T}_+ maps the set*

$$\mathcal{P}_{Q,\delta}^+ := \{(\mu, \nu, P) \in \text{dom } \mathbf{T}_+ : P \in \mathcal{P}_{Q,\delta}^{\alpha,M,N}, 1 \leq \mu \leq \lambda_1(P) - \delta, \delta \leq \nu \leq Q\}, \quad Q > 0, \delta > 0,$$

into $\mathcal{P}_{\widehat{Q},\widehat{\delta}}^{\alpha,\widehat{M},\widehat{N}}$, where \widehat{Q} and $\widehat{\delta}$ depend on Q and δ . Moreover, the map \mathbf{T}_+ is Lipschitz continuous on $\mathcal{P}_{Q,\delta}^+$:

$$\mathbf{d}_\alpha(\widehat{P}_1, \widehat{P}_2) \leq C(Q, \delta) (|\mu_1 - \mu_2| + |\nu_1 - \nu_2| + \mathbf{d}_\alpha(P_1, P_2))$$

for any $(\mu_i, \nu_i, P_i) \in \mathcal{P}_{Q,\delta}^+, \widehat{P}_i := \mathbf{T}_+(\mu_i, \nu_i, P_i), i = 1, 2$.

Proof. It follows from Lemmas 4.1 and 4.3 together with the initial conditions (5.4) that the maps $(\mu, P) \mapsto \psi(0, \mu)$ and $(\mu, P) \mapsto \psi_\sigma^{[1]}(0, \mu)$ are Lipschitz continuous on the compact set

$$\mathcal{G}_{Q,\delta} := \{(\mu, P) : P \in \mathcal{P}_{Q,\delta}, 1 \leq \mu \leq \lambda_1(P) - \delta\}$$

w.r.t. the metric $\mathbf{d}_0(\cdot, \cdot)$ for P . In addition, it has been shown in [11] that $\psi(0, \mu) \neq 0$. Consequently, there holds $|\psi(0, \mu)| \geq \varepsilon > 0$ for $(\mu, P) \in \mathcal{G}_{Q,\delta}$, where $\varepsilon = \varepsilon(Q, \delta)$. Consequently,

the map $(\mu, P) \mapsto \varkappa = -\frac{\psi_\sigma^{[1]}(0, \mu)}{\psi(0, \mu)}$ is Lipschitz continuous on $\mathcal{G}_{Q,\delta}$. In particular, $|\varkappa| \leq C(Q, \delta)$.

Analogously, we show that $\rho = \rho(\mu, \nu, P)$ defined by (5.5) is Lipschitz continuous on $\mathcal{P}_{Q,\delta}^+$. Note that the denominator of ρ is non-zero according to the construction in [11]. Next, we consider the function $u(x) = z(x, \mu, \rho)$ and conclude that the transforms $(\mu, \nu, P) \mapsto u(\cdot)$ and $(\mu, \nu, P) \mapsto u_\sigma^{[1]}(\cdot)$ are Lipschitz continuous as maps from $\mathbb{R} \times \mathbb{R} \times \mathcal{P}^{0,M,N}$ to $C[0, \pi]$. Furthermore, one can show that the function $u(x)$ possesses the same properties on $\mathcal{P}_{Q,\delta}^+$ as the function $v(x)$ on $\mathcal{P}_{Q,\delta}$ in the proof of Lemma 5.2 and analogously complete the proof. \square

Now, let us study transforms $\mathcal{S}\mathbf{T}\mathcal{S}^{-1}$ which map the spectral data $\mathfrak{S} \mapsto \widehat{\mathfrak{S}}$ while $\mathbf{T}: P \mapsto \widehat{P}$.

Lemma 5.4. *Suppose that $\mathbf{T} \in \{\mathbf{T}_-, \mathbf{T}_{-+}, \mathbf{T}_{+-}\}$, fixed integers M, N are such that $\mathcal{P}^{0,M,N} \subset \text{dom } \mathbf{T}$, and $\alpha \in (0, \frac{1}{2})$ is fixed. Then $\mathcal{S}\mathbf{T}\mathcal{S}^{-1}$ maps $\mathcal{B}_{R,\varepsilon}^{\alpha,M,N}$ into $\mathcal{B}_{\widehat{R},\widehat{\varepsilon}}^{\alpha,\widehat{M},\widehat{N}}$, where $\widehat{R} > 0$ and $\widehat{\varepsilon} > 0$ depend on R and ε . Moreover, the transform $\mathcal{S}\mathbf{T}\mathcal{S}^{-1}$ is Lipschitz continuous on $\mathcal{B}_{R,\varepsilon}$:*

$$\varrho_\alpha(\widehat{\mathfrak{S}}_1, \widehat{\mathfrak{S}}_2) \leq C(R, \varepsilon) \varrho_\alpha(\mathfrak{S}_1, \mathfrak{S}_2) \tag{5.17}$$

for any $\mathfrak{S}_i \in \mathcal{B}_{R,\varepsilon}, \widehat{\mathfrak{S}}_i = \mathcal{S}\mathbf{T}\mathcal{S}^{-1}(\mathfrak{S}_i), i = 1, 2$.

Proof. For definiteness, consider $\mathbf{T} = \mathbf{T}_-$. The proof for the other cases is analogous. Let $\mathfrak{S} = \{\lambda_n, \gamma_n\}_{n \geq 1} \in \mathcal{B}_{R, \varepsilon}$. By virtue of (3.9), (3.10), and Theorem 3.1 for $\alpha = 0$, which has been proved by Guliyev [11], we conclude that \mathfrak{S} are the spectral data of some problem $\mathcal{L}(P)$ with $P = (\sigma, f, F) \in \mathcal{P}^{0, M, N}$. According to Proposition 5.1, there hold $\widehat{P} = \mathbf{T}(P) \in \mathcal{P}^{0, \widehat{M}, \widehat{N}}$, where $\widehat{M} = M - 1$, $\widehat{N} = N - 1$. Moreover, the corresponding spectral data $\mathfrak{S} = \mathcal{S}(\widehat{P})$ have the form

$$\widehat{\lambda}_n = \lambda_n, \quad \widehat{\gamma}_n = \frac{\gamma_n}{\lambda_n - \lambda_1}, \quad n \geq 2. \quad (5.18)$$

Let us show that $\widehat{\mathfrak{S}} \in \mathcal{B}_{\widehat{R}, \widehat{\varepsilon}}^{\alpha, \widehat{M}, \widehat{N}}$. By virtue of the direct part of Theorem 3.1 for $\alpha = 0$, there hold the asymptotics

$$\sqrt{\widehat{\lambda}_n} = n - \frac{\widehat{M} + \widehat{N}}{2} - 2 + \widehat{\kappa}_n, \quad \widehat{\gamma}_n = \frac{\pi}{2}(n-1)^{2\widehat{M}}(1 + \widehat{\beta}_n), \quad n \geq 2. \quad (5.19)$$

Here, we take the index shift into account. Since $\widehat{\lambda}_n = \lambda_n$, we have $\widehat{\kappa}_n = \kappa_n$ and so $\{\widehat{\lambda}_n\}$ and $\{\widehat{\kappa}_n\}$ fulfill the conditions (3.9). Using (3.5), (5.18), and (5.19), we derive

$$\lambda_n - \lambda_1 = n^2(1 + O(n^{-1})), \quad 1 + \widehat{\beta}_n = (1 + \beta_n)(1 + O(n^{-1})), \quad (5.20)$$

where the estimates $O(n^{-1})$ are uniform by $\mathfrak{S} \in \mathcal{B}_{R, \varepsilon}$. Since $\lambda_n - \lambda_1 \geq \varepsilon$ and $1 + \beta_n \geq \varepsilon$, then $1 + \widehat{\beta}_n \geq \widehat{\varepsilon}$. Note that $\{\frac{1}{n}\} \in l_2^\alpha$ for $\alpha \in (0, \frac{1}{2})$. Therefore, (5.20) implies $\{\widehat{\beta}_n\} \in l_2^\alpha$ and $\|\{\widehat{\beta}_n\}\|_\alpha \leq C(R, \varepsilon)$. Thus, we have proved all the requirements (3.9) and (3.10) for $\widehat{\mathfrak{S}}$ with some positive constants $\widehat{R}, \widehat{\varepsilon}$ instead of R, ε .

It remains to prove the estimate (5.17). Consider the corresponding collections of the spectral data \mathfrak{S}_i and $\widehat{\mathfrak{S}}_i$ for $i = 1, 2$. It follows from (3.5) that

$$|\lambda_{n,1} - \lambda_{n,2}| \leq C(R)n|\kappa_{n,1} - \kappa_{n,2}|, \quad n \geq 1. \quad (5.21)$$

Using (3.5), (5.18), (5.19), and (5.21), we derive

$$|\widehat{\beta}_{n,1} - \widehat{\beta}_{n,2}| \leq C(R, \varepsilon) \left(|\beta_{n,1} - \beta_{n,2}| + \frac{|\kappa_{n,1} - \kappa_{n,2}|}{n} + \frac{|\kappa_{1,1} - \kappa_{1,2}|}{n^2} \right), \quad n \geq 2.$$

Hence

$$\|\{\widehat{\beta}_{n,1} - \widehat{\beta}_{n,2}\}\|_\alpha \leq C(R, \varepsilon) (\|\{\beta_{n,1} - \beta_{n,2}\}\|_\alpha + \|\{\kappa_{n,1} - \kappa_{n,2}\}\|_\alpha).$$

The similar estimate for $\{\widehat{\kappa}_{n,1} - \widehat{\kappa}_{n,2}\}$ is obtained trivially. Thus, in view of (3.7), we arrive at (5.17). \square

Lemma 5.5. *Let $\alpha \in (0, \frac{1}{2})$ and the integers $M, N \geq -1$ be fixed. Then, for any $\mathfrak{S} \in \mathcal{B}_{R, \varepsilon}^{\alpha, M, N}$, $1 \leq \mu \leq \lambda_1 - \varepsilon$, and $\varepsilon \leq \nu \leq R$, the spectral data $\mathcal{S}\mathbf{T}_+(\mu, \nu, \mathcal{S}^{-1}(\mathfrak{S}))$ belong to $\mathcal{B}_{\widehat{R}, \widehat{\varepsilon}}^{\alpha, \widehat{M}, \widehat{N}}$, where $\widehat{R} > 0$ and $\widehat{\varepsilon} > 0$ depend on R and ε . Furthermore, the Lipschitz continuity holds:*

$$\varrho_\alpha(\widehat{\mathfrak{S}}_1, \widehat{\mathfrak{S}}_2) \leq C(R, \varepsilon) (|\mu_1 - \mu_2| + |\nu_1 - \nu_2| + \varrho_\alpha(\mathfrak{S}_1, \mathfrak{S}_2))$$

for any $\mathfrak{S}_i \in \mathcal{B}_{R, \varepsilon}$, $1 \leq \mu_i \leq \lambda_{1,i} - \varepsilon$, $\varepsilon \leq \nu_i \leq R$, $\widehat{\mathfrak{S}}_i := \mathcal{S}\mathbf{T}_+(\mu_i, \nu_i, \mathcal{S}^{-1}(\mathfrak{S}_i))$, $i = 1, 2$.

The proof of Lemma 5.5 is analogous to Lemma 5.4.

6 Proofs of the main results

In this section, we present the proofs of the main theorems formulated in Section 3. Theorems 3.2 and 3.3 are proved by induction on $M + N = 2p$. The induction base follows from the results of [25]. At each step, we use the Lipschitz continuity of the Darboux-type transforms discussed in Section 5. Next, we consider the finite data approximation and prove Theorems 3.5 and 3.6.

Proof of Theorem 3.2. In the case of the Dirichlet boundary conditions $M = N = -1$, Theorem 3.2 follows from [25, Theorem 3.8] (see also [26, Theorem 3.15]).

Suppose that Theorem 3.2 is valid for $M + N = 2p$. Let us prove it for $M + N = 2p + 2$. Consider the case $M \geq 0$, $N \geq 0$. Fix $\alpha \in (0, \frac{1}{2})$, $Q > 0$, and $\delta > 0$. Let $P \in \mathcal{P}_{Q,\delta}^{\alpha,M,N}$. By Lemma 5.2, $\widehat{P} := \mathbf{T}_-(P)$ belongs to $\mathcal{P}_{\widehat{Q},\widehat{\delta}}^{\alpha,\widehat{M},\widehat{N}}$, where $\widehat{M} = M - 1$, $\widehat{N} = N - 1$, \widehat{Q} and $\widehat{\delta}$ depend on Q and δ . Moreover, the Lipschitz estimate (5.8) is fulfilled. For $\widehat{M} + \widehat{N} = 2p$, the assertion of Theorem 3.2 holds by virtue of the induction hypothesis. Therefore, $\widehat{\mathfrak{S}} := \mathcal{S}(\widehat{P})$ lies in $\mathcal{B}_{\widehat{R},\widehat{\varepsilon}}^{\alpha,\widehat{M},\widehat{N}}$, where \widehat{R} and $\widehat{\varepsilon}$ depend on \widehat{Q} and $\widehat{\delta}$, and

$$\varrho_\alpha(\widehat{\mathfrak{S}}_1, \widehat{\mathfrak{S}}_2) \leq C(\widehat{Q}, \widehat{\delta}) \mathbf{d}_\alpha(\widehat{P}_1, \widehat{P}_2), \quad \widehat{P}_i \in \mathcal{P}_{\widehat{Q},\widehat{\delta}}. \quad (6.1)$$

Next, in view of (5.6), the spectral data $\mathfrak{S} := \mathcal{S}(P)$ equal $\mathcal{S}\mathbf{T}_+(\lambda_1(P), \gamma_1(P), \mathcal{S}^{-1}(\widehat{\mathfrak{S}}))$. For applying Lemma 5.5, we have to show that

$$1 \leq \lambda_1(P) \leq \lambda_2(P) - \eta, \quad \eta \leq \gamma_1(P) \leq \Omega, \quad (6.2)$$

for any $P \in \mathcal{P}_{Q,\delta}$, where positive constants η and Ω depend in Q and δ . The inequality $\lambda_1(P) \geq 1$ follows from (3.8). According to Lemmas 4.2 and 4.3 and Remark 4.4, the functions $\lambda_2(P) - \lambda_1(P)$ and $\gamma_1(P)$ are continuous on the compact set $\mathcal{P}_{Q,\delta}$ in $\mathcal{P}^{0,M,N}$. This together with the inequalities $\lambda_1 < \lambda_2$ and $\gamma_1 > 0$ yield (6.2). Consequently, Lemma 5.5 implies $\mathfrak{S} \in \mathcal{B}_{R,\varepsilon}^{\alpha,M,N}$ and

$$\varrho_\alpha(\mathfrak{S}_1, \mathfrak{S}_2) \leq C(|\lambda_1(P_1) - \lambda_1(P_2)| + |\gamma_1(P_1) - \gamma_1(P_2)| + \varrho_\alpha(\widehat{\mathfrak{S}}_1, \widehat{\mathfrak{S}}_2)), \quad (6.3)$$

where the constants $R > 0$, $\varepsilon > 0$, and C depend on \widehat{R} , $\widehat{\varepsilon}$, η , and Ω . From the above arguments, one can easily see that these constants depend on Q and δ .

Thus, we have shown that, for any $P \in \mathcal{P}_{Q,\delta}$, the spectral data $\mathcal{S}(P)$ belong to $\mathcal{B}_{R,\varepsilon}$. Combining the Lipschitz estimates (5.8), (6.1), (6.3), and (4.2), we arrive at (3.11). So, the assertion of Theorem 3.2 is proved for $M + N = 2p + 2$, $M \geq 0$, $N \geq 0$. This proof can be presented by the following scheme:

$$P \xrightarrow[\text{Lemma 5.2}]{\mathbf{T}_-} \widehat{P} \xrightarrow[\text{induction hypothesis for } \widehat{M} + \widehat{N} = M + N - 2]{\mathcal{S}} \widehat{\mathfrak{S}} \xrightarrow[\text{Lemma 5.5}]{\mathbf{T}_+} \mathfrak{S}$$

The cases $M = -1$ and $N = -1$ can be analogously reduced to the previously studied case $\widehat{M} \geq 0$, $\widehat{N} \geq 0$, $\widehat{M} + \widehat{N} = M + N$ by using the transforms \mathbf{T}_{-+} and \mathbf{T}_{+-} :

$$\begin{aligned} M = -1: \quad & P \xrightarrow[\text{Lemma 5.2}]{\mathbf{T}_{+-}} \widehat{P} \xrightarrow[\text{induction hypothesis for } \widehat{M} + \widehat{N} = M + N, \widehat{M}, \widehat{N} \geq 0]{\mathcal{S}} \widehat{\mathfrak{S}} \xrightarrow[\text{Lemma 5.4}]{\mathbf{T}_{-+}} \mathfrak{S}, \\ N = -1: \quad & P \xrightarrow[\text{Lemma 5.2}]{\mathbf{T}_{-+}} \widehat{P} \xrightarrow[\text{induction hypothesis for } \widehat{M} + \widehat{N} = M + N, \widehat{M}, \widehat{N} \geq 0]{\mathcal{S}} \widehat{\mathfrak{S}} \xrightarrow[\text{Lemma 5.4}]{\mathbf{T}_{+-}} \mathfrak{S}. \end{aligned}$$

This concludes the proof for $M + N = 2p + 2$. The induction yields the assertion of Theorem 3.2 for any even $M + N \geq -2$. \square

Proof of Theorem 3.3. Let us use induction on $M + N = 2p$. For the case of the Dirichlet boundary conditions ($p = -1$), Theorem 3.3 follows from [25, Theorem 3.8] (see also [26, Theorem 3.15]).

Suppose that the assertion of Theorem 3.3 holds for $M + N = 2p$. Let us prove it for $M + N = 2p + 2$. Schematically, the proof for different cases can be presented as follows:

$$\begin{aligned}
M, N \geq 0: \quad \mathfrak{S} &\xrightarrow[\text{Lemma 5.4}]{\mathbf{T}_-} \widehat{\mathfrak{S}} \xrightarrow[\text{induction hypothesis for } \widehat{M} + \widehat{N} = M + N - 2]{\mathcal{S}^{-1}} \widehat{P} \xrightarrow[\text{Lemma 5.3}]{\mathbf{T}_+} P, \\
M = -1: \quad \mathfrak{S} &\xrightarrow[\text{Lemma 5.4}]{\mathbf{T}_{+-}} \widehat{\mathfrak{S}} \xrightarrow[\text{induction hypothesis for } \widehat{M} + \widehat{N} = M + N, \widehat{M}, \widehat{N} \geq 0]{\mathcal{S}^{-1}} \widehat{P} \xrightarrow[\text{Lemma 5.2}]{\mathbf{T}_{-+}} P, \\
N = -1: \quad \mathfrak{S} &\xrightarrow[\text{Lemma 5.4}]{\mathbf{T}_{-+}} \widehat{\mathfrak{S}} \xrightarrow[\text{induction hypothesis for } \widehat{M} + \widehat{N} = M + N, \widehat{M}, \widehat{N} \geq 0]{\mathcal{S}^{-1}} \widehat{P} \xrightarrow[\text{Lemma 5.2}]{\mathbf{T}_{+-}} P.
\end{aligned}$$

Consider the case $M, N \geq 0$ in more detail. Let $\alpha \in (0, \frac{1}{2})$, M , and N be fixed. Suppose that $\mathfrak{S} \in \mathcal{B}_{R,\varepsilon}^{\alpha,M,N}$ for some $R > 0$ and $\varepsilon > 0$. Put $\widehat{\mathfrak{S}} := \mathcal{S}\mathbf{T}_-\mathcal{S}^{-1}\mathfrak{S}$. By Lemma 5.4, we have $\widehat{\mathfrak{S}} \in \mathcal{B}_{\widehat{R},\widehat{\varepsilon}}^{\alpha,\widehat{M},\widehat{N}}$, where $\widehat{R} > 0$ and $\widehat{\varepsilon} > 0$ depend on R and ε , $\widehat{M} = M - 1$, $\widehat{N} = N - 1$. Moreover, the estimate (5.17) is valid. Since $\widehat{M} + \widehat{N} = M + N - 2$, the map $\mathcal{S}^{-1}: \widehat{\mathfrak{S}} \mapsto \widehat{P}$ satisfies the assertion of Theorem 3.3 by the induction hypothesis. Hence, $\widehat{P} \in \mathcal{P}_{\widehat{Q},\widehat{\delta}}^{\alpha,\widehat{M},\widehat{N}}$ with some \widehat{Q} and $\widehat{\delta}$ depending on \widehat{R} and $\widehat{\varepsilon}$. Additionally, the Lipschitz estimate holds:

$$\mathbf{d}_\alpha(\widehat{P}_1, \widehat{P}_2) \leq C(\widehat{R}, \widehat{\varepsilon}) \varrho_\alpha(\widehat{\mathfrak{S}}_1, \widehat{\mathfrak{S}}_2), \quad \widehat{\mathfrak{S}}_i \in \mathcal{B}_{\widehat{R},\widehat{\varepsilon}}. \quad (6.4)$$

Next, note that $P := \mathcal{S}^{-1}(\widehat{\mathfrak{S}})$ equals $\mathbf{T}_+(\lambda_1, \gamma_1, \widehat{P})$, where λ_1 and γ_1 are from $\widehat{\mathfrak{S}}$. In order to apply Lemma 5.3, we have to show that

$$1 \leq \lambda_1 \leq \widehat{\lambda}_2 - \eta, \quad \eta \leq \gamma_1 \leq \Omega, \quad (6.5)$$

for any $\widehat{\mathfrak{S}} \in \mathcal{B}_{R,\varepsilon}$, where η and Ω are positive constants depending on R and ε . The estimates (6.5) readily follow from $\widehat{\mathfrak{S}} \in \mathcal{B}_{R,\varepsilon}$ and the equality $\widehat{\lambda}_2 = \lambda_2$. Then, by Lemma 5.3, we get $P \in \mathcal{P}_{Q,\delta}^{\alpha,M,N}$, where $Q > 0$ and $\delta > 0$ depend on R and ε , and

$$\mathbf{d}_\alpha(P_1, P_2) \leq C(R, \varepsilon) (|\lambda_{1,1} - \lambda_{1,2}| + |\gamma_{1,1} - \gamma_{1,2}| + \mathbf{d}_\alpha(\widehat{P}_1, \widehat{P}_2)). \quad (6.6)$$

Combining the Lipschitz estimates (5.17), (6.4), and (6.6), we arrive at (3.12) and so conclude the proof for $M + N = 2p$, $M \geq 0$, $N \geq 0$. The cases $M = -1$ and $N = -1$ are studied analogously. Induction completes the proof. \square

Remark 6.1. The proof of Theorem 3.2, in particular, shows by induction that, for any $\sigma \in \mathring{W}_2^\alpha[0, \pi]$, $f \in \mathcal{R}_M$, and $F \in \mathcal{R}_N$, the sequences of the remainders $\{\kappa_n\}$ and $\{\beta_n\}$ from the spectral data asymptotics (3.5) belong to l_2^α . This proves the direct part of Theorem 3.1. Analogously, the inverse part follows from the proof of Theorem 3.3.

Remark 6.2. Note that the transforms \mathbf{T}_- , \mathbf{T}_{-+} , \mathbf{T}_{+-} , and \mathbf{T}_+ do not change the parity of $M + N$. Therefore, we have proved Theorems 3.2 and 3.3 only for even $M + N$, since they are based on the results of [25], which were obtained only for the Dirichlet boundary conditions ($M + N = -2$). Nevertheless, the spectral data characterization has been proved in [47] for the Dirichlet-Dirichlet and the Neumann-Dirichlet boundary conditions, so we get Theorem 3.1 for both even and odd $M + N$.

Proceed to considering the finite data approximation P_m of P .

Proof of Theorem 3.5. Let P satisfy the hypothesis of the theorem. In view of the continuous embedding $W_2^{\alpha_2}[0, \pi] \subset W_2^{\alpha_1}[0, \pi]$ for $\alpha_1 < \alpha_2$, we have $P \in \mathcal{P}_{Q, \delta}^{\alpha_1, M, N}$. By Theorem 3.1, the numbers $\mathfrak{S}_m = \{\lambda_{n,m}, \gamma_{n,m}\}_{n \geq 1}$ for each sufficiently large m are the spectral data of some problem $\mathcal{L}(P_m)$, where $P_m \in \mathcal{P}^{\alpha, M, N}$ for any $\alpha \in [0, \frac{1}{2})$. Applying Theorem 3.2 with $\alpha = \alpha_1$, we conclude that $\mathfrak{S} \in \mathcal{B}_{R, \varepsilon}$, where R and ε depend on Q and δ . Furthermore, we have $\mathfrak{S}_m \in \mathcal{B}_{R, \varepsilon}$ for sufficiently large m in view of (3.9), (3.10), and (3.13). Theorem 3.3 implies $P_m \in \mathcal{P}_{\widehat{Q}, \widehat{\delta}}$, where \widehat{Q} and $\widehat{\delta}$ depend on Q and δ . Therefore, part (i) of Corollary 3.4 yields

$$\mathbf{d}_{\alpha_1}(P, P_m) \leq C(Q, \delta) \varrho_{\alpha_1}(\mathfrak{S}, \mathfrak{S}_m). \quad (6.7)$$

According to (3.7), we have

$$\varrho_{\alpha_1}(\mathfrak{S}, \mathfrak{S}_m) = \|\{\kappa_n - \kappa_{n,m}\}\|_{\alpha_1} + \|\{\beta_n - \beta_{n,m}\}\|_{\alpha_1}.$$

Using (3.13), we get

$$\|\{\kappa_n - \kappa_{n,m}\}\|_{\alpha_1} = \sqrt{\sum_{n=m+1}^{\infty} n^{2\alpha_1} \kappa_n^2} \leq m^{\alpha_1 - \alpha_2} \|\{\kappa_n\}\|_{\alpha_2}.$$

Using the similar estimate for $\{\beta_n\}$ together with (3.9) and (3.10), we obtain

$$\varrho_{\alpha_1}(\mathfrak{S}, \mathfrak{S}_m) \leq m^{\alpha_1 - \alpha_2} (\|\{\kappa_n\}\|_{\alpha_2} + \|\{\beta_n\}\|_{\alpha_2}) \leq C(Q, \delta) m^{\alpha_1 - \alpha_2}. \quad (6.8)$$

Combining (6.7) and (6.8) yields the claim. \square

Proof of Theorem 3.6. Analogously to the previous proof, we get the estimate

$$\mathbf{d}_{\alpha_1}(P, \tilde{P}_m) \leq C(Q, \delta) \varrho_{\alpha_1}(\mathfrak{S}, \tilde{\mathfrak{S}}_m) \quad (6.9)$$

for all sufficiently small $\varepsilon > 0$ and sufficiently large m . It follows from (3.5) and (3.15) that

$$|\kappa_n - \tilde{\kappa}_n| \leq C(Q, \delta) \varepsilon n^{-1}, \quad |\beta_n - \tilde{\beta}_n| \leq C(Q, \delta) \varepsilon n^{-2M}, \quad n = \overline{1, m}.$$

Using the latter estimates together with (6.8) and (6.9), we arrive at (3.16). \square

Remark 6.3. In this paper, we confine ourselves by $\alpha < \frac{1}{2}$ for technical simplicity. Our methods also work for $\alpha \geq \frac{1}{2}$, but then spectral data asymptotics contain more terms:

$$\sqrt{\lambda_n} = s(n) + \frac{\omega_1}{s(n)} + \frac{\omega_2}{s^3(n)} + \cdots + \frac{\omega_k}{s^{2k-1}(n)} + o(n^{-(2k-1)}), \quad s(n) = n - \frac{M+N}{2} - 1.$$

The constants $\{\omega_j\}_{j=1}^k$ are related to σ and to the coefficients of the rational functions $f(\lambda)$ and $F(\lambda)$, which complicates the technique.

Acknowledgment. The author is grateful to Professor Maria A. Kuznetsova for reading the manuscript and for her valuable remarks.

Funding. This work was supported by Grant 24-71-10003 of the Russian Science Foundation, <https://rscf.ru/en/project/24-71-10003/>.

References

- [1] Levitan, B.M. Inverse Sturm-Liouville Problems, VNU Sci. Press, Utrecht (1987).
- [2] Freiling, G.; Yurko, V. Inverse Sturm-Liouville Problems and Their Applications, Nova Science Publishers, Huntington, NY, 2001.
- [3] Marchenko, V.A. Sturm-Liouville Operators and Applications. Revised edition, AMS, Providence, 2011.
- [4] Kravchenko, V.V. Direct and Inverse Sturm-Liouville Problems, Birkhäuser, Cham (2020).
- [5] Kraft, R.E.; Wells, W.R. Adjointness properties for differential systems with eigenvalue-dependent boundary conditions, with application to flow-duct acoustics, *J. Acoust. Soc. Am.* 61 (1977), 913–922.
- [6] Granet, E. Regularization of energy-dependent pointlike interactions in 1D quantum mechanics, *J. Phys. A* 55 (2022), no. 42, 425308.
- [7] Kaoullas, G.; Georgiou, G.C. Start-up and cessation Newtonian Poiseuille and Couette flows with dynamic wall slip, *Meccanica* 50 (2015), no. 7, 1747–1760.
- [8] Fulton, C.T. Singular eigenvalue problems with eigenvalue parameter contained in the boundary conditions, *Proc. R. Soc. Edinburgh, Sect. A.* 87 (1980), no. 1–2, 1–34.
- [9] Binding, P.A.; Browne, P.J.; Watson, B.A. Sturm-Liouville problems with boundary conditions rationally dependent on the eigenparameter. II, *J. Comput. Appl. Math.* 148 (2002), no. 1, 147–168.
- [10] Binding, P.A.; Browne, P.J.; Watson, B.A. Equivalence of inverse Sturm-Liouville problems with boundary conditions rationally dependent on the eigenparameter, *J. Math. Anal. Appl.* 291 (2004), 246–261.
- [11] Guliyev, N.J. Schrödinger operators with distributional potentials and boundary conditions dependent on the eigenvalue parameter, *J. Math. Phys.* 60 (2019), 063501.
- [12] Guliyev, N.J. Essentially isospectral transformations and their applications, *Annali di Matematica Pura ed Applicata*, 199 (2020), no. 4, 1621–1648.
- [13] Guliyev, N.J. On two-spectra inverse problems, *Proc. AMS.* 148 (2020), 4491–4502.
- [14] Guliyev, N.J. Inverse square singularities and eigenparameter-dependent boundary conditions are two sides of the same coin, *Quart. J. Math.* 74 (2023), no. 3, 889–910.
- [15] Yang, Y.; Wei, G. Inverse scattering problems for Sturm-Liouville operators with spectral parameter dependent on boundary conditions, *Math. Notes* 103 (2018), no. 1–2, 59–66.
- [16] Freiling, G.; Yurko V. Inverse problems for Sturm-Liouville equations with boundary conditions polynomially dependent on the spectral parameter, *Inverse Problems* 26 (2010), no. 5, 055003.
- [17] Freiling, G.; Yurko, V. Determination of singular differential pencils from the Weyl function, *Adv. Dyn. Sys. Appl.* 7 (2012), no. 2, 171–193.

- [18] Chitorkin, E.E.; Bondarenko, N.P. Solving the inverse Sturm-Liouville problem with singular potential and with polynomials in the boundary conditions, *Anal. Math. Phys.* 13 (2023), Article number: 79.
- [19] Chitorkin, E.E.; Bondarenko, N.P. Local solvability and stability for the inverse Sturm-Liouville problem with polynomials in the boundary conditions, *Math. Meth. Appl. Sci.* 47 (2024), no. 11, 8881–8903.
- [20] Chitorkin, E.E.; Bondarenko, N.P. Inverse Sturm-Liouville problem with polynomials in the boundary condition and multiple eigenvalues, *Journal of Inverse and Ill-Posed Problems* (2025), published online. <https://doi.org/10.1515/jiip-2024-0011>
- [21] Wang, Y.P. Uniqueness theorems for Sturm–Liouville operators with boundary conditions polynomially dependent on the eigenparameter from spectral data, *Results Math.* 63 (2013), 1131–1144.
- [22] Yang, C.-F. Uniqueness theorems for differential pencils with eigenparameter boundary conditions and transmission conditions, *J. Diff. Eqns.* 255 (2013), no. 9, 2615–2635.
- [23] Bondarenko, N.P. An inverse problem for an integro-differential pencil with polynomial eigenparameter-dependence in the boundary condition, *Anal. Math. Phys.* 9 (2019), no. 4, 2227–2236.
- [24] Chitorkin, E.E.; Bondarenko, N.P. Inverse Sturm-Liouville problem with singular potential and spectral parameter in the boundary conditions, *J. Diff. Eqns.* 421 (2025), 495–523.
- [25] Savchuk, A.M.; Shkalikov, A.A. Inverse problems for Sturm-Liouville operators with potentials in Sobolev spaces: Uniform stability, *Funct. Anal. Appl.* 44 (2010), no. 4, 270–285.
- [26] Savchuk, A.M.; Shkalikov, A.A. Uniform stability of the inverse Sturm-Liouville problem with respect to the spectral function in the scale of Sobolev spaces, *Proc. Steklov Inst. Math.* 283 (2013), 181–196.
- [27] Hryniv, R.O. Analyticity and uniform stability in the inverse singular Sturm-Liouville spectral problem, *Inverse Problems* 27 (2011), no. 6, 065011.
- [28] Hryniv, R.O. Analyticity and uniform stability in the inverse spectral problem for Dirac operators, *J. Math. Phys.* 52 (2011), 063513.
- [29] Buterin, S. Uniform full stability of recovering convolutional perturbation of the Sturm-Liouville operator from the spectrum, *J. Diff. Eqns.* 282 (2021), 67–103.
- [30] Buterin, S.; Djuric, N. Inverse problems for Dirac operators with constant delay: uniqueness, characterization, uniform stability, *Lobachevskii J. Math.* 43 (2022), no. 6, 1492–1501.
- [31] Kuznetsova, M. Uniform stability of recovering Sturm-Liouville-type operators with frozen argument, *Results Math.* 78 (2023), no. 5, Article ID 169.
- [32] Bondarenko, N.P. Uniform stability of the inverse problem for the non-self-adjoint Sturm-Liouville operator, [arXiv:2409.16175](https://arxiv.org/abs/2409.16175).
- [33] Ryabushko, T.I. Stability of the reconstruction of a Sturm-Liouville operator from two spectra, *Teor. Funkts., Funkts. Anal. Prilozh.* 18 (1973), 176–185 [in Russian].

- [34] Marletta, M.; Weikard, R. Weak stability for an inverse Sturm-Liouville problem with finite spectral data and complex potential, *Inverse Problems* 21 (2005), 1275–1290.
- [35] Savchuk, A.M.; Shkalikov, A.A. Recovering of a potential of the Sturm-Liouville problem from finite sets of spectral data, *Spectral Theory and Differential Equations*, Amer. Math. Soc. Transl. Ser. 2, 233, Amer. Math. Soc., Providence, RI, 2014, pp. 211–224.
- [36] Savchuk, A.M. Reconstruction of the potential of the Sturm-Liouville operator from a finite set of eigenvalues and normalizing constants, *Math. Notes* 99 (2016), no. 5, 715–728.
- [37] Guo, Y.; Ma, L.-J.; Xu, X.-C.; An, Q. Weak and strong stability of the inverse Sturm-Liouville problem, *Math. Meth. Appl. Sci.* 46 (2023), no. 14, 15684–15705.
- [38] Aktosun, T. Stability of the Marchenko inversion, *Inverse Problems*, 3 (1987), 555–563.
- [39] Hitrik, M. Stability of an inverse problem in potential scattering on the real line, *Commun. in Partial Differential Equations* 25 (2000), no.5-6, 925–955.
- [40] Xu, X.-C.; Yang, C.-F. On the inverse spectral stability for the transmission eigenvalue problem with finite data, *Inverse Problems* 36 (2020), no. 8, 085006.
- [41] Xu, X.-C.; Guo, Y.; Yang, C.-F. Stability of the inverse transmission eigenvalue problem for the Schrödinger operator with a radial potential, *Lett. Math. Phys.* 112 (2022), no. 4, 82.
- [42] Marletta, M.; Shterenberg, R.; Weikard, R. On the inverse resonance problem for Schrödinger operators, *Comm. Math. Phys.* 295 (2009), no. 2, 465–484.
- [43] Bledsoe, M. Stability of the inverse resonance problem on the line, *Inverse Problems* 28 (2012), no. 10, 105003.
- [44] Savchuk, A.M.; Shkalikov, A.A. Sturm-Liouville operators with singular potentials, *Math. Notes* 66 (1999), no. 6, 741–753.
- [45] Savchuk, A.M.; Shkalikov, A.A. Inverse problem for Sturm-Liouville operators with distribution potentials: reconstruction from two spectra, *Russ. J. Math. Phys.* 12 (2005), no. 4, 507–514.
- [46] Triebel, H. *Interpolation Theory, Function Spaces, Differential Operators*, VEB Deutscher Verlag der Wissenschaften, Berlin, 1978.
- [47] Hryniv, R.O.; Mykytyuk, Y.V. Inverse spectral problems for Sturm-Liouville operators with singular potentials. IV. Potentials in the Sobolev space scale, *Proc. Edinb. Math. Soc.* 49 (2006), no. 2, 309–329.

Natalia Pavlovna Bondarenko

1. Department of Mechanics and Mathematics, Saratov State University, Astrakhanskaya 83, Saratov 410012, Russia,

2. Department of Applied Mathematics, Samara National Research University, Moskovskoye Shosse 34, Samara 443086, Russia,

3. S.M. Nikolskii Mathematical Institute, Peoples' Friendship University of Russia (RUDN University), 6 Miklukho-Maklaya Street, Moscow, 117198, Russia,

4. Moscow Center of Fundamental and Applied Mathematics, Lomonosov Moscow State University, Moscow 119991, Russia.

e-mail: *bondarenkonp@sgu.ru*