# Mixing Any Cocktail with Limited Ingredients: On the Structure of Payoff Sets in Multi-Objective MDPs and its Impact on Randomised Strategies\*

James C. A. Main Mickael Randour

#### Abstract

We consider multi-dimensional payoff functions in Markov decision processes, and ask whether a given expected payoff vector can be achieved or not. In general, pure strategies (i.e., not resorting to randomisation) do not suffice for this problem.

We study the structure of the set of expected payoff vectors of all strategies given a multi-dimensional payoff function and its consequences regarding randomisation requirements for strategies. In particular, we prove that for any payoff for which the expectation is well-defined under all strategies, it is sufficient to mix (i.e., randomly select a pure strategy at the start of a play and committing to it for the rest of the play) finitely many pure strategies to approximate any expected payoff vector up to any precision. Furthermore, for any payoff for which the expected payoff is finite under all strategies, any expected payoff can be obtained exactly by mixing finitely many strategies.

## 1 Introduction

Markov decision processes (MDPs) are a classical framework for decision making in uncertain environments. MDPs are notably used in the fields of formal methods (e.g., [BK08, FBB+23]) and reinforcement learning (e.g., [SB18]). An MDP models the interaction of a controllable system with a stochastic environment: at each step of the process, the system selects an action and the state of the process is updated following a probability distribution that depends only the current state and chosen action. This interaction goes on forever and yields a play (i.e., an execution) of the process. The quality of a play is quantified by a payoff function, which assigns a numeric value to all plays. The goal is then to compute an optimal strategy, i.e., a decision-making procedure for the system that maximises the expected payoff. Such a strategy constitutes a formal blueprint for a controller of the modelled system [Ran13].

Multi-objective MDPs. More complex specifications, such as simultaneous constraints on the response time and energy consumption of a system, are usually modelled using multi-dimensional payoff functions. In this setting, some expected payoff vectors may be incomparable; an analysis of trade-offs between the different dimensions may be necessary.

The goal is generally to determine, given a vector, whether it is *achievable*, i.e., whether there exists a strategy whose expected payoff is greater than or equal to the vector in

<sup>\*</sup>James C. A. Main is an F.R.S.-FNRS Research Fellow and Mickael Randour is an F.R.S.-FNRS Research Associate. Both authors are members of the TRAIL institute. This work has been supported by the Fonds de la Recherche Scientifique – FNRS under Grant n° T.0188.23 (PDR ControlleRS).

all components (e.g., [EKVY08, CFW13, RRS17]). A related problem is to compute or approximate the *Pareto curve* of the set of expected payoffs, i.e., the expected payoffs that are *Pareto-optimal* (e.g., [FKP12, CKK17, QK21]). Intuitively, an expected payoff is Pareto-optimal if there is no strategy whose expected payoff is as good on all dimensions and strictly better on one dimension. Alternatively, one can look for strategies with expected payoffs that are optimal for the *lexicographic* order over vectors (e.g., [HPS<sup>+</sup>21, CKM<sup>+</sup>23, BCM<sup>+</sup>23]). For instance, in a two-dimensional setting, this equates to finding strategies that maximise the expected payoff on the second dimension among the strategies that maximise the expected payoff on the first dimension.

Strategy complexity. For controller design, simpler strategies, i.e., using limited memory and randomisation, are preferable. In the case of many classical (one-dimensional) specifications, there exist optimal strategies that use no memory and no randomisation. For instance, such strategies suffice to maximise the probability of a parity objective [CJH04] or the expectation of a mean [Bie87] or discounted-sum payoff [Sha53].

In contrast to the one-dimensional case, Pareto-optimal strategies in multi-objective MDPs may require both *memory* and *randomisation* (see, e.g., [RRS17, BGMR23]). For this reason, a major problem is understanding what are the simplest relevant strategies, e.g., whether randomisation is necessary, or what is the minimum amount of memory required for the application at hand. A related approach consists in directly searching for simple strategies that satisfy the desired constraints (e.g., [DKQR20]).


There is an extensive body of work aimed at understanding memory requirements for strategies in various game-based models. These works provide sufficient conditions on payoffs for which memoryless or finite-memory strategies suffice, or even complete characterisations of such payoffs. For instance, see [LPR18, BLO<sup>+</sup>22, BRV23, CO23] for games on deterministic graphs, [BORV23, GK23] for stochastic games and [Gim07] for MDPs. Most of these works focus on pure strategies.


In this work, our focus is on randomisation requirements in multi-objective MDPs. Instead of identifying bounds on memory, we study whether we can achieve vectors while using strategies with limited randomisation power. We remark that in general, it is not possible to jointly minimise memory and randomisation requirements. Indeed, there can be a trade-off between memory and randomisation, even in the one-dimensional case [CdH04, Hor09, CRR14, MPR20].

Randomised strategies. A strategy that does not use randomisation is called pure. Formally, a pure strategy is a function that assigns actions to histories (i.e., sequences of observations). Randomised strategies can be defined in two ways. On the one hand, a behavioural strategy is a function that assigns distributions over actions to histories. In other words, when following a behavioural strategy, the random choices of actions are made at each step of the process. On the other hand, a mixed strategy is a distribution over the set of pure strategies. When playing according to a mixed strategy, a pure strategy is randomly chosen at the start of the process and is followed for the entire play.

In MDPs, behavioural and mixed strategies are equivalent; any strategy from one class can be emulated by a strategy of the other. This result is known as *Kuhn's theo-* rem [Aum64], and it also shows that mixed and behavioural strategies are the most general type of randomised strategies.

When limiting ourselves to *finite-memory strategies*, the natural analogues of mixed strategies are strictly weaker than the analogue of behavioural strategies [MR24]. In fact, the emulation of a behavioural strategy that flips a coin at each round requires a





- (a) Transitions are labelled by actions and probabilities. Weights represent the time taken by each action. The target is doubly circled.
- (b) A set of expected payoffs. The probability of reaching work before 40 time units is given on the first dimension. The other dimension represents the expected time to reach work.

Figure 1.1: An MDP with two payoffs and its associated set of expected payoffs.

mixed strategy that randomises over *uncountably* many pure strategies, infinitely many of which require infinite memory to account for all patterns. It follows that mixed strategies with a finite support, i.e., that mix finitely many pure strategies, are a *strict sub-class* of the set of randomised strategies (regardless of memory). Arguably, this is one of the simplest ways of integrating randomness in strategies. Considering such strategies instead of general behavioural (or mixed) strategies amounts to *limiting the randomisation power* of strategies.

A simple example. For the sake of illustration, let us consider the MDP depicted in Figure 1.1a. This MDP models a situation where a person wants to go to work. They must choose between riding their bicycle or taking the train. However, the train may be delayed with high probability due to an ongoing strike. The goal of the commuter is twofold: maximise the likelihood of reaching work within 40 time units (to reach work on time) and do so as fast as possible on average.

We illustrate the set of expected payoffs for this situation in Figure 1.1b. We label expected payoffs with their corresponding strategies. On the one hand,  $\sigma_{\text{train}}$  and  $\sigma_{\text{bike}}$  denote the pure strategies that always choose the action in the subscript. On the other hand, we denote by  $\sigma_{\ell t+b}$  the strategy that attempts to take the train  $\ell$  times before choosing the bicycle.

This simple example highlights the need for randomisation and memory in multiobjective MDPs. When limited to pure strategies, for instance, it is not possible to reach work on time with probability exceeding 90% while guaranteeing an expected commute time lower than 27: any pure strategy whose payoff is not represented will reach work on time with a smaller probability than  $\sigma_{\text{train}}$ , and thus will not be satisfactory. However, as suggested by the point labelled by  $\sigma_{\text{mix}}$  on the illustration, these constraints can be satisfied by mixing  $\sigma_{\text{train}}$  with  $\sigma_{2\text{t+b}}$ . In fact, here, all expected payoffs can be obtained by finite-support mixed strategies. We explain below how this property generalises. We remark however that, in general, the set of expected payoffs need not be a convex polytope like in this example (see Section 4).

Our contributions. We study the structure of sets of expected payoffs in multiobjective MDPs, focusing on the relationship between what can be obtained with pure and with randomised strategies. Our goal is to provide general results, i.e., that apply to a broad class of payoffs. We only consider universally unambiguously integrable payoffs (Section 3), i.e., payoffs that have a well-defined (possibly infinite) expectation no matter the strategy, a natural requirement when aiming to reason about expectations. We obtain finer results for universally integrable payoffs, i.e., payoffs whose expectation is finite under all strategies.

First, we prove that for universally integrable (multi-dimensional) payoffs, for all strategies, there exists a *pure strategy* with a greater or equal expected payoff in the lexicographic sense (Theorem 5.1).

This first result serves as a building block to one of our main results: any expected payoff vector of a universally integrable (multi-dimensional) payoff is a convex combination of expected payoffs of pure strategies (Theorem 6.2). From a strategic perspective, this means that in multi-objective MDPs, finite-support mixed strategies suffice to exactly obtain any (Pareto-optimal) expected payoff vector. As a corollary, we obtain that any extreme point of a set of expected payoffs of a universally integrable payoff can be obtained by a pure strategy (Corollary 6.3). These results generalise known properties for classical combinations of objectives for which the set of expected payoffs is a convex polytope (e.g., combinations of  $\omega$ -regular specifications [EKVY08] and universally integrable total-reward payoffs [FKP12]). While none of the previous properties generalise to the whole class of universally unambiguously integrable payoffs (Example 5.1 and 6.3), we prove that, for such payoffs, convex combinations of pure strategies can be used to approximate any expected payoff (Theorem 6.5).

In both cases, we can bound the number of strategies that are mixed depending on the number of dimensions d. Depending on the setting, we can match or approximate expected payoffs by mixing no more than d+1 strategies (Theorem 6.6).

We also provide a sufficient condition to guarantee that the set of expected payoffs is closed. We show that for continuous universally *square-integrable* payoffs (which generalise real-valued continuous payoffs), the set of expected payoffs is closed (Theorem 7.6).

To prove our results, we borrow techniques from several fields. We mainly rely on topology, properties of convex sets (separating and supporting hyperplanes) and the theory of the Lebesgue integral. We assume familiarity with basic properties of the Lebesgue integral, and otherwise endeavour to keep the text self-contained by recalling most required notions, notably through an extensive appendix.

Applicability. The class of universally integrable payoffs is large: it contains most classical payoffs. Indeed, all bounded payoffs are defacto in this class. For example, all indicators of objectives are in it. This means that all settings where one considers the probabilities of sets of plays (i.e., either an inherently qualitative objective or one arising from fixing a threshold for a quantitative payoff) are in it, for any definition of such sets of plays. Classical examples include  $\omega$ -regular objectives [EKVY08], window objectives [BDOR20] or percentiles queries [RRS17]. Bounded payoffs also encompass discounted-sum [Sha53] and mean-payoff [BBC+14, CKK17] functions. Heterogeneous combinations, such as, e.g., combinations of energy and mean-payoff [BHRR19] also fit under this umbrella.

Payoffs that are completely out of the scope of our results—i.e., not even universally unambiguously integrable—include total payoff [FBB<sup>+</sup>23], average-energy [BMR<sup>+</sup>18] and shortest path [RRS17], all with both positive and negative weights: the expectation is ill-defined when plays of positive and negative infinite payoffs have non-zero measure. Let us focus on shortest-path objectives. When only positive weights are used, a classical

restriction [RRS17], the associated payoff is universally unambiguously integrable (but not universally integrable as plays of infinite payoffs may exist). Finally, when targets are almost-surely visited (as in the above example), the payoff is continuous and universally square integrable (Appendix C).

For many payoffs that are at least universally unambiguously integrable, extreme points of the set of expected payoffs can be obtained via pure finite-memory strategies (e.g.,  $\omega$ -regular objectives [EKVY08] or mean-payoff [BBC<sup>+</sup>14]). It follows that for these objectives, mixing over finitely many pure finite-memory strategies is sufficient to fulfil any achievability requirement. In other words, one of the least expressive models of randomised finite-memory strategies (see the classification of [MR24]) suffices. Furthermore, the blow-up in memory from this mixing argument is small: it suffices to mix, at most, one more strategy than the number of payoffs.

Our results also highlight the role of randomisation in strategies for multi-objective MDPs: it is useful *only* to balance the payoffs on different dimensions in many cases.

**Related work.** We provide a few references, in addition to the main references cited previously. In our proof of Theorem 6.2, we invoke the separating hyperplane theorem. This theorem also plays a role in approximation schemes of the set of achievable vectors: see, e.g., [FKP12, QK21]; a unifying approach is presented in [Qua23].

Specifications with multiple objectives have also been considered in the context of two-player (non-stochastic) games on graphs (e.g., [FH13, CRR14]) and two-player stochastic games (e.g., [CFK<sup>+</sup>13, ACK<sup>+</sup>20]). Closely related to multi-objective specifications are approaches that provide guarantees simultaneously in the worst case and the expected case [BFRR17].

Regarding randomisation in strategies, [CDGH15] studies when randomisation is helpful in strategies or in transitions of games. In particular, the authors show that if there exists an optimal strategy to maximise the probability of an event in an MDP, then there exists a pure optimal strategy. This property is generalised by our result on lexicographic MDPs.

Outline. Section 2 overviews notation and preliminary notions. We introduce payoff functions in Section 3 and establish relationships between expected payoffs of general strategies and expected payoffs of pure strategies. In Section 4, we provide an example highlighting the potential complexity of sets of expected payoffs in multi-objective MDPs. We study randomisation requirements for the lexicographic optimisation of several objectives in MDPs in Section 5. Section 6 presents our results regarding expected payoff sets and achievable sets. Finally, we study continuous payoff functions in Section 7.

Appendices A and B complement Sections 2 and 3 respectively. Appendix C shows that the results of Section 7 apply to universally integrable continuous shortest-path payoffs.

## 2 Preliminaries

This section defines the objects occurring in the main part of this paper. Additional definitions and results have been deferred to Appendix A.

**Set-theoretic notation.** We let  $\mathbb{N}$  and  $\mathbb{R}$  respectively denote the sets of natural and real numbers. We let  $\mathbb{N}_0 = \mathbb{N} \setminus \{0\}$ . We denote the extended real line by  $\mathbb{R} = \mathbb{R} \cup \{-\infty, +\infty\}$ .

Let  $A' \subseteq A$  and  $B' \subseteq B$  be sets and  $f: A \to B$  be a function. We let  $\mathbb{1}_{A'}: A \to \{0,1\}$  denote the indicator function of A'. We let  $\mathsf{Im}(f)$  denote the image of f. We let  $f^{-1}(B') = \{a \in A \mid f(a) \in B'\}$  denote the inverse image of B' by f. For any  $b \in B$ , we write  $f^{-1}(b)$  instead of  $f^{-1}(\{b\})$  to lighten notation. We let |A| denote the cardinal of A.

**Probability.** Let A be a countable set. We write  $\mathcal{D}(A)$  for the set of distributions over A, i.e., the set of functions  $\mu \colon A \to [0,1]$  such that  $\sum_{a \in A} \mu(a) = 1$ . The support of a distribution  $\mu \in \mathcal{D}(A)$  is  $\mathsf{supp}(\mu) = \{a \in A \mid \mu(a) > 0\}$ .

Given a set B and a  $\sigma$ -algebra  $\mathcal{A}$  over B, we denote by  $\mathcal{D}(B,\mathcal{A})$  the set of probability distributions over the measurable space  $(B,\mathcal{A})$ . Let  $\mu \in \mathcal{D}(B,\mathcal{A})$  and  $f \colon B \to \overline{\mathbb{R}}$  be a measurable function. We say that f is  $\mu$ -integrable it is integrable with respect to  $\mu$ , i.e., if  $\int_B |f| \mathrm{d}\mu \in \mathbb{R}$ . We extend the Lebesgue integral to non-positive functions in the following way: if f is non-positive, we let  $\int_B f \mathrm{d}\mu = -\int_B -f \mathrm{d}\mu$ . If f is non-negative, non-positive or  $\mu$ -integrable, we say that  $\int_B f \mathrm{d}\mu$  is the  $\mu$ -integral of f.

**Vector spaces.** Throughout this text, vectors are written in boldface to distinguish them from scalars. Let  $d \in \mathbb{N}_0$ . We let  $\mathbf{0}_d$  and  $\mathbf{1}_d \in \mathbb{R}^d$  respectively be the vectors of  $\mathbb{R}^d$  where all components are zero and one respectively. We omit the dimension subscript or superscript whenever there is no ambiguity on the dimension of the space.

Given  $\mathbf{v} = (v_j)_{1 \leq j \leq d}$ ,  $\mathbf{w} = (w_j)_{1 \leq j \leq d} \in \mathbb{R}^d$ , we let  $\langle \mathbf{v}, \mathbf{w} \rangle = \sum_{j=1}^d v_j w_j$  denote the scalar product of  $\mathbf{v}$  and  $\mathbf{w}$ . We let  $\| \cdot \|_2$  denote the Euclidean norm on  $\mathbb{R}^d$ , defined by  $\| \mathbf{v} \|_2 = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$  for all  $\mathbf{v} \in \mathbb{R}^d$ .

The affine span of a set  $D \subseteq \mathbb{R}^d$ , which we denote by  $\mathsf{aff}(D)$ , is the smallest affine set (i.e., translation of a vector subspace of  $\mathbb{R}^d$ ) in which D is included.

Given a linear map  $L: \mathbb{R}^d \to \mathbb{R}^{d'}$  (where  $d' \in \mathbb{N}_0$ ), we let  $\ker(L)$  denote the kernel of L, i.e., the set  $\{\mathbf{v} \in \mathbb{R}^d \mid L(\mathbf{v}) = \mathbf{0}_{d'}\}$ . A linear form is a linear map whose co-domain is  $\mathbb{R}$ .


**Convexity.** A convex combination of vectors  $\mathbf{v}_1, \ldots, \mathbf{v}_n \in \mathbb{R}^d$  is a linear combination  $\sum_{m=1}^n \alpha_m \cdot \mathbf{v}_m$  such that  $\alpha_1, \ldots, \alpha_n \in [0,1]$  and  $\sum_{m=1}^n \alpha_m = 1$ . We refer to a sequence of coefficients  $\alpha_1, \ldots, \alpha_n \in [0,1]$  such that  $\sum_{m=1}^n \alpha_m = 1$  as convex combination coefficients. Given  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^d$ , we let  $[\mathbf{v}, \mathbf{w}] = \{\alpha \cdot \mathbf{v} + (1 - \alpha)\mathbf{w} \mid \alpha \in [0,1]\}$  denote the (closed) segment from  $\mathbf{v}$  to  $\mathbf{w}$ ; it is the set of convex combinations of  $\mathbf{v}$  and  $\mathbf{w}$ . Open and half-open segments are defined analogously.


Let  $D \subseteq \mathbb{R}^d$ . The convex hull of D, denoted by  $\mathsf{conv}(D)$ , is the set of all convex combinations of elements of D. The set D is convex if for all  $\mathbf{v}$ ,  $\mathbf{w} \in D$ ,  $[\mathbf{v}, \mathbf{w}] \subseteq D$ , or, equivalently, if  $D = \mathsf{conv}(D)$ . If D is convex, we say that  $\mathbf{q} \in D$  is an extreme point of D if  $\mathbf{q} \notin \mathsf{conv}(D \setminus \{\mathbf{q}\})$ , i.e., if  $\mathbf{q}$  is not a convex combination of elements of D other than  $\mathbf{q}$  and we let  $\mathsf{extr}(D)$  denote the set of extreme points of D. Extreme points generalise the notion of vertices of polytopes.

The definition of a convex combination does not bound the number of involved vectors. However, in  $\mathbb{R}^d$ , it is sufficient to only consider convex combinations involving no more than d+1 vectors. This is formalised by the following theorem.

**Theorem 2.1** (Carathéodory's theorem for convex hulls (e.g., [Roc70, Thm. 17.1])). Let  $D \subseteq \mathbb{R}^d$  and  $\mathbf{q} \in \mathsf{conv}(D)$ . There exists  $D' \subseteq D$  such that  $|D'| \le d+1$  and  $\mathbf{q} \in \mathsf{conv}(D')$ .

Convexity does not generalise to  $\mathbb{R}^d$ . Most notably, convex combinations of vectors of  $\mathbb{R}^d$  (defined in the same way as above) may be ill-defined. Although we consider convex





(a) The dashed blue line  $(x + y = \frac{7}{2})$  strongly separates  $D_1$  and  $D_2$ : they lie on a different side of the line and there is a strip around the line that is disjoint from  $D_1$  and  $D_2$ .

(b) The dashed blue line  $(x+y=\sqrt{2})$  supports the unit ball for the Euclidean norm in  $\mathbb{R}^2$  at  $\mathbf{q} = \frac{\sqrt{2}}{2}\mathbf{1}$ .

Figure 2.1: Illustration of separating and supporting hyperplanes.

combinations of elements of  $\mathbb{R}^d$  in the sequel, these are always guaranteed to be well-defined, for reasons explained below.

**Topology.** We only provide notation in this section. For convenience, we recall some definitions in Appendix A.1, including the definitions of the product topology, continuity, compactness and the usual topology of  $\bar{\mathbb{R}}$ .

Let  $(X, \mathcal{T})$  be a Hausdorff topological space. For all  $D \subseteq X$ , we let  $\mathsf{cl}(D)$  and  $\mathsf{int}(D)$  denote the closure and interior of D. The border of  $D \subseteq X$  is the set  $\mathsf{bd}(D) = \mathsf{cl}(D) \setminus \mathsf{int}(D)$ .

In addition to the three notions above, we introduce an additional one for subsets of  $\mathbb{R}^d$  (with its usual topology). The *relative interior* of a set  $D \subseteq \mathbb{R}^d$ , denoted by ri(D), is the interior of D as a subset of aff(D) (with the induced topology). The relative interior of a subset of  $\mathbb{R}^d$  includes its interior and these sets may differ. For instance, the segment  $[\mathbf{0}_2, \mathbf{1}_2] \subseteq \mathbb{R}^2$  has empty interior. However, we have  $ri([\mathbf{0}_2, \mathbf{1}_2]) = ]\mathbf{0}_2, \mathbf{1}_2[$  (because  $aff([\mathbf{0}_2, \mathbf{1}_2])$  is the line of equation x = y).

**Hyperplane separation.** A hyperplane H of  $\mathbb{R}^d$  is a set of the form  $\{\mathbf{v} \in \mathbb{R}^d \mid x^*(\mathbf{v}) = \alpha\}$  for some non-zero linear form  $x^*$  and  $\alpha \in \mathbb{R}$ . Let  $D_1$  and  $D_2 \subseteq \mathbb{R}^d$ . The sets  $D_1$  and  $D_2$  are strongly separated by a hyperplane if there exists a non-zero linear form  $x^*$  such that  $\inf_{\mathbf{q} \in D_1} x^*(\mathbf{q}) > \sup_{\mathbf{p} \in D_2} x^*(\mathbf{p})$ . A convex set  $D \subseteq \mathbb{R}^d$  is supported by a hyperplane at  $\mathbf{q} \in D$  if there exists a non-zero linear form  $x^*$  such that, for all  $\mathbf{p} \in D$ ,  $x^*(\mathbf{p}) \leq x^*(\mathbf{q})$  (a supporting hyperplane in this case is  $H = \{\mathbf{v} \in \mathbb{R}^d \mid x^*(\mathbf{v}) = x^*(\mathbf{q})\}$ ). We provide an illustration of the notions of separating and supporting hyperplanes in Figure 2.1.

We recall a variant of the hyperplane separation theorem and the supporting hyperplane theorem. We first outline a sufficient condition such that two disjoint convex sets can be strongly separated.

**Theorem 2.2** (Hyperplane separation theorem [Roc70, Cor. 11.4.2]). Let  $D_1$  and  $D_2$  be two convex subsets of  $\mathbb{R}^d$ . If  $cl(D_1) \cap cl(D_2) = \emptyset$  and  $D_1$  or  $D_2$  is bounded, then there exists a hyperplane strongly separating  $D_1$  and  $D_2$ .

Figure 2.1a illustrates a setup in which we can apply the theorem. We often apply Theorem 2.2 in the case where  $D_1$  is a singleton set  $\{\mathbf{q}\}$  (it is bounded) and  $\mathbf{q} \notin \mathsf{cl}(D_2)$ . The

next theorem provides a sufficient condition for the existence of a supporting hyperplane at a given point of a convex set.

**Theorem 2.3** (Supporting hyperplane theorem [Roc70, Thm. 11.6]). Let  $D \subseteq \mathbb{R}^d$  be convex and  $\mathbf{q} \in D$ . If  $\mathbf{q} \notin \mathsf{ri}(D)$ , then there exists a hyperplane H supporting D at  $\mathbf{q}$  such that  $D \not\subseteq H$ .

Ordering vectors. We consider two order relations on  $\mathbb{R}^d$ : the component-wise order and the lexicographic order. Let  $\mathbf{q} = (q_j)_{1 \leq j \leq d}$  and  $\mathbf{p} = (p_j)_{1 \leq j \leq d} \in \mathbb{R}^d$ . For the component-wise order, we write  $\mathbf{q} \leq \mathbf{p}$  if and only if  $q_j \leq p_j$  for all  $1 \leq j \leq d$ . For the lexicographic ordering over  $\mathbb{R}^d$ , we write  $\mathbf{q} \leq_{\mathsf{lex}} \mathbf{p}$  if and only if  $\mathbf{q} = \mathbf{p}$  or  $q_j \leq p_j$  where  $j = \min\{j' \leq d \mid q_{j'} \neq p_{j'}\}$ . We write  $\mathbf{q} <_{\mathsf{lex}} \mathbf{p}$  if  $\mathbf{q} \leq_{\mathsf{lex}} \mathbf{p}$  and  $\mathbf{q} \neq \mathbf{p}$ . We recall that the component-wise order is partial, whereas the lexicographic order is a total order.

Let  $D \subseteq \mathbb{R}^d$ . We say that  $\mathbf{q} \in D$  is a Pareto-optimal element of D if it is maximal for the component-wise order, i.e., if there does not exist  $\mathbf{p} \in D$  such that  $\mathbf{q} \leq \mathbf{p}$  and  $\mathbf{q} \neq \mathbf{p}$ . We say that D is downward-closed if for all  $\mathbf{q} \in D$  and  $\mathbf{p} \in \mathbb{R}^d$ ,  $\mathbf{p} \leq \mathbf{q}$  implies  $\mathbf{p} \in D$ . We let down(D) denote the downward closure of D, which is defined as the smallest (with respect to set inclusion) downward-closed set in which D is included. A set and its downward closure have the same set of Pareto-optimal elements.

Markov decision processes. A Markov decision process models the interaction of a system with a stochastic environment. At each step of the process, an action is taken by the system and the state of the process is updated in a stochastic fashion, where the relevant distribution depends only on the current state and chosen action.

Formally, a Markov decision process (MDP) is a tuple  $\mathfrak{M} = (S, A, \delta)$  where S is a finite non-empty set of states, A is a finite set of actions and  $\delta \colon S \times A \to \mathcal{D}(S)$  is a partial probabilistic transition function. For  $s \in S$ , we let A(s) denote the set of actions  $a \in A$  such that  $\delta(s, a)$  is defined. If  $a \in A(s)$ , we say that a is enabled in s. We assume that MDPs have no deadlocks, i.e., there is an action enabled in each state.

A play of  $\mathfrak{M}$  is an infinite sequence  $\pi = s_0 a_0 s_1 a_1 \ldots \in (SA)^{\omega}$  such that  $a_{\ell} \in A(s_{\ell})$  and  $s_{\ell+1} \in \operatorname{supp}(\delta(s_{\ell}, a_{\ell}))$  for all  $\ell \in \mathbb{N}$ . A history is a finite prefix of a play ending in a state. Let  $\pi = s_0 a_0 s_1 a_1 \ldots$  be a play. We write  $\pi_{\leq r}$  for the history  $s_0 a_0 s_1 \ldots a_{r-1} s_r$ . We also use this notation for prefixes of histories. We denote the suffix  $s_r a_r s_{r+1} a_{r+1} \ldots$  of  $\pi$  by  $\pi_{\geq r}$ . For any history  $h = s_0 a_0 s_1 \ldots s_r$ , we let  $\operatorname{last}(h) = s_r$ . We write  $\operatorname{Plays}(\mathfrak{M})$  and  $\operatorname{Hist}(\mathfrak{M})$  for the sets of plays and histories of  $\mathfrak{M}$ .

We fix an MDP  $\mathfrak{M} = (S, A, \delta)$  for the rest of the section.

Strategies. A strategy is a function that describes how to select actions based on the history of the ongoing play. A strategy may resort to randomisation. Formally, a (behavioural) strategy of  $\mathfrak{M}$  is a function  $\sigma \colon \mathsf{Hist}(\mathfrak{M}) \to \mathcal{D}(A)$  such that for all  $h \in \mathsf{Hist}(\mathfrak{M})$ ,  $\mathsf{supp}(\sigma(h)) \subseteq A(\mathsf{last}(h))$ . A strategy is pure if it uses no randomisation, i.e., if it maps all histories to Dirac distributions. We view pure strategies as functions  $\sigma \colon \mathsf{Hist}(\mathfrak{M}) \to A$ . A strategy  $\sigma$  is memoryless if the distribution it assigns to a history depends only on its last state, i.e., if for all histories  $h, h' \in \mathsf{Hist}(\mathfrak{M})$ ,  $\mathsf{last}(h) = \mathsf{last}(h')$  implies  $\sigma(h) = \sigma(h')$ . Memoryless (resp. pure memoryless) strategies can be viewed as functions  $S \to \mathcal{D}(A)$  (resp.  $S \to A$ ). We let  $\Sigma(\mathfrak{M})$  and  $\Sigma_{\mathsf{pure}}(\mathfrak{M})$  respectively denote the set of all strategies of  $\mathfrak{M}$  and the set of pure strategies of  $\mathfrak{M}$ .

A play  $\pi = s_0 a_0 s_1 \dots$  is *consistent* with a strategy  $\sigma$  if for all  $\ell \in \mathbb{N}$ , we have  $a_{\ell} \in \mathsf{supp}(\sigma(\pi_{\leq \ell}))$ . An *outcome* of a strategy is any play consistent with the strategy. Consistency of a history with a strategy is defined similarly.

We now define the distribution over plays induced by a strategy from an initial state. First, we specify the relevant  $\sigma$ -algebra. A cylinder set is a set of continuations of a history; given  $h = s_0 a_0 s_1 \dots s_r \in \mathsf{Hist}(\mathfrak{M})$ , we let  $\mathsf{Cyl}(h) = \{\pi \in \mathsf{Plays}(\mathfrak{M}) \mid \pi_{\leq r} = h\}$ . We let  $\mathcal{A}$  denote the  $\sigma$ -algebra generated by the cylinder sets. Let  $\sigma \in \Sigma(\mathfrak{M})$  and  $s \in S$ . We define  $\mathbb{P}_s^{\sigma} \in \mathcal{D}(\mathsf{Plays}(\mathfrak{M}), \mathcal{A})$  over cylinder sets as follows. For  $h = s_0 a_0 s_1 \dots s_r \in \mathsf{Hist}(\mathfrak{M})$ , if  $s_0 = s$ , we let

$$\mathbb{P}^{\sigma}_s(\operatorname{Cyl}(h)) = \prod_{\ell=0}^{r-1} \sigma(h_{\leq \ell})(a_\ell) \cdot \delta(s_\ell, a_\ell)(s_{\ell+1}),$$

and otherwise, if  $s_0 \neq s$ , we let  $\mathbb{P}_s^{\sigma}(\mathsf{Cyl}(h)) = 0$ . This distribution extends in a unique fashion to  $\mathcal{A}$  by Carathéodory's extension theorem [Dur19, Theorem A.1.3.].

There exists an alternative definition for randomised strategies: a mixed strategy is a distribution over pure strategies. Intuitively, when playing following a mixed strategy, a pure strategy is selected at random at the start of the process and then the play proceeds according to this pure strategy. Behavioural and mixed strategies are equivalent in terms of the distributions over plays that they can induce in MDPs; this is a special case of Kuhn's theorem [Aum64]. However, the derivation of mixed strategies from behavioural strategies is non-trivial: mixed strategies that are equivalent to some behavioural strategies may have to randomise over an uncountable set of pure strategies. Furthermore, even when the behavioural strategy uses finite memory (i.e., has a finite automata-based representation), an equivalent mixed strategy may have to randomise over infinite-memory strategies [MR24]. In particular, the class of finite-support mixed strategies, which is at the core of our results, is a strict subclass of the class of mixed strategies.

In the sequel, unless otherwise stated, all strategies are assumed to be behavioural strategies due to the equivalence stated above. Some of our arguments explicitly refer to mixed strategies that randomise over countable sets of pure strategies. Distributions over plays for such strategies are defined as follows: for any countable class of pure strategies  $\Sigma \subseteq \Sigma_{\mathsf{pure}}(\mathfrak{M})$ , mixed strategy  $\mu \in \mathcal{D}(\Sigma)$  and state  $s \in S$ , the distribution defined by  $\mu$  from s is defined by  $\mathbb{P}^{\mu}_{s}(\Omega) = \sum_{\sigma \in \Sigma} \mu(\sigma) \mathbb{P}^{\sigma}_{s}(\Omega)$  for all  $\Omega \in \mathcal{A}$ .

## 3 Payoffs and multi-objective Markov decision processes

Payoffs are functions that quantify the quality of plays. Definitions are given in Section 3.1. Section 3.2 introduces a key technical property. Some classical payoffs are provided in Sect. 3.3. Continuous payoffs, which are the main object of study of Section 7, are introduced in Section 3.4. Section 3.5 presents notation for MDPs with multiple payoffs.

We fix an MDP  $\mathfrak{M} = (S, A, \delta)$  for the whole section.

## 3.1 Payoff functions and objectives

A payoff function (or payoff for short) is a measurable function  $f: \mathsf{Plays}(\mathfrak{M}) \to \overline{\mathbb{R}}$ . An objective is a measurable subset of the set of plays. An objective represents a qualitative specification, e.g., reaching a target set of states. To each objective  $\Omega$ , we associate the payoff  $\mathbb{1}_{\Omega}$ .

Let f be a payoff,  $\sigma \in \Sigma(\mathfrak{M})$  be a strategy and  $s \in S$  be an initial state. The  $\mathbb{P}_s^{\sigma}$ -integral of f is only formally defined whenever f is non-negative, non-positive or  $\mathbb{P}_s^{\sigma}$ -integrable. If f is such a payoff, we let  $\mathbb{E}_s^{\sigma}(f) = \int_{\pi \in \mathsf{Plays}(\mathfrak{M})} f(\pi) d\mathbb{P}_s^{\sigma}(\pi)$ ;  $\mathbb{E}_s^{\sigma}(f)$  is the expected payoff of the strategy  $\sigma$  from s (for f). We also generalise the notion of expected payoff to a broader class of payoff functions as follows. Let  $f^+ = \max(f, 0)$  and  $f^- = \max(-f, 0)$ 

denote the non-negative and non-positive parts of f. We say that f has an unambiguous  $\mathbb{P}_s^{\sigma}$ -integral if  $\mathbb{E}_s^{\sigma}(f^+) \in \mathbb{R}$  or  $\mathbb{E}_s^{\sigma}(f^-) \in \mathbb{R}$ . If f has an unambiguous  $\mathbb{P}_s^{\sigma}$ -integral, we abuse notation and let  $\mathbb{E}_s^{\sigma}(f) = \mathbb{E}_s^{\sigma}(f^+) - \mathbb{E}_s^{\sigma}(f^-)$ . We reserve the notation  $\mathbb{E}$  for the expectation of payoffs, and use integrals for other probability spaces.

In the sequel, we only consider payoffs for which the expected payoff is (unambiguously) defined for all strategies from all initial states. We say that a payoff f is universally unambiguously integrable if f has an unambiguous  $\mathbb{P}_s^{\sigma}$ -integral for all  $\sigma \in \Sigma(\mathfrak{M})$  and  $s \in S$ . In particular, all non-negative and non-positive payoffs are universally unambiguously integrable. In this work, we focus on universally unambiguously integrable payoffs: all payoffs considered from this point on are assumed to be universally unambiguously integrable.

Payoffs that are integrable no matter the strategy and initial state are of particular interest in the sequel. We say that a payoff f is universally integrable if it is  $\mathbb{P}_s^{\sigma}$ -integrable for all  $\sigma \in \Sigma(\mathfrak{M})$  and  $s \in S$ . All bounded functions (in particular indicators) are universally integrable.

Given a payoff function f, we say that a strategy  $\sigma$  is optimal from a state  $s \in S$  if for all strategies  $\tau$ , we have  $\mathbb{E}_s^{\sigma}(f) \geq \mathbb{E}_s^{\tau}(f)$ . Playing optimally with respect to an indicator thus equates to maximising the probability of the underlying objective. In spite of this maximisation-oriented viewpoint for optimisation, we can capture minimisation by considering -f in lieu of f: f is universally unambiguously integrable if and only if -f is, and minimising a payoff f is equivalent to maximising the payoff -f.

## 3.2 Relating payoffs of randomised and pure strategies

Our main goal is to show that a simpler class of randomised strategies, i.e., finite-support mixed strategies, suffices when considering MDPs with multiple payoffs. To prove our results, we rely on a result that relates the expectation of any strategy with the expectation of pure strategies. More precisely, our proofs rely on the following result: the expected payoff of a strategy from a given state can be reformulated as an integral of expected payoffs of pure strategies (under certain technical assumptions). This result is a generalisation of a part of the proof of [CDGH15, Lem. 1], which introduces the same property for probabilities of events instead of expected values of payoff.

Let  $\sigma \in \Sigma(\mathfrak{M})$ . Instead of introducing a measure on  $\Sigma_{\mathsf{pure}}(\mathfrak{M})$ , following [CDGH15], we consider a distribution over  $[0,1]^{\omega}$  and a mapping  $[0,1]^{\omega} \to \Sigma_{\mathsf{pure}}(\mathfrak{M})$ . Intuitively, each element of  $[0,1]^{\omega}$  can be seen as an a priori resolution of all random choices of  $\sigma$  since the uniform distribution of [0,1] can be used to simulate any distribution of a finite set. Thus, the pure strategies obtained through the aforementioned mapping assign, to all histories, actions that may occur with positive probability under  $\sigma$ .

To formalise the lemma, we introduce a  $\sigma$ -algebra on  $[0,1]^{\omega}$ . We consider the  $\sigma$ -algebra  $\mathcal{A}_{[0,1]^{\omega}}$  generated by the set of sequence-cylinders. A sequence-cylinder is a set of the form  $[0,x_1]\times\cdots\times[0,x_n]\times[0,1]^{\omega}$  for some  $x_1,\ldots,x_n\in[0,1]$ . We obtain the following relation between expected payoffs of randomised and pure strategies.

**Lemma 3.1.** Let  $\sigma \in \Sigma(\mathfrak{M})$ . There exists a distribution  $\nu_{\sigma}$  over  $([0,1]^{\omega}, \mathcal{A}_{[0,1]^{\omega}})$  and a mapping  $[0,1]^{\omega} \to \Sigma_{\mathsf{pure}}(\mathfrak{M}) \colon x \mapsto \tau_x$  such that, for all universally unambiguously integrable payoffs f and all  $s \in S$ , the mapping  $[0,1]^{\omega} \to \mathbb{R} \colon x \mapsto \mathbb{E}_s^{\tau_x}(f)$  is measurable and

$$\mathbb{E}_{s}^{\sigma}(f) = \int_{x \in [0,1]^{\omega}} \mathbb{E}_{s}^{\tau_{x}}(f) d\nu_{\sigma}(x)$$

if  $\inf_{\tau} \mathbb{E}_{s}^{\tau}(f) \geq 0$  or  $\sup_{\tau} \mathbb{E}_{s}^{\tau}(f) \leq 0$  or f is  $\mathbb{P}_{s}^{\sigma}$ -integrable.

Proof. Fix a state  $s \in S$  and let f be a universally unambiguously integrable payoff. We show the result for payoffs increasing complexity, similarly to how the Lebesgue integral is constructed. First, we prove it for indicators of objectives. Second, we show that it also holds for non-negative simple functions (i.e., linear combinations of indicators) by linearity of the integral. Third, we deal with non-negative payoffs with the monotone convergence theorem. Fourth, we consider  $\mathbb{P}_s^{\sigma}$ -integrable payoffs. Finally, we close the proof by considering universally unambiguously integrable payoffs such that  $\inf_{\tau} \mathbb{E}_s^{\tau}(f) \geq 0$  or  $\sup_{\tau} \mathbb{E}_s^{\tau}(f) \leq 0$ .

We consider the distribution  $\nu_{\sigma}$  and mapping  $x \mapsto \tau_x$  defined in the proof of [CDGH15, Lemma 1] such that, for all measurable  $\Omega \subseteq \mathsf{Plays}(\mathfrak{M}), x \mapsto \mathbb{P}_s^{\tau_x}(\Omega)$  is measurable and

$$\mathbb{P}_s^{\sigma}(\Omega) = \int_{x \in [0,1]^{\omega}} \mathbb{P}_s^{\tau_x}(\Omega) d\nu_{\sigma}(x). \tag{3.1}$$

This directly entails the claim of the lemma when f is an indicator of an objective.

For the second step of the argument, we assume that f is a non-negative simple function. Let  $\alpha_1, \ldots, \alpha_n \geq 0$  and  $\Omega_1, \ldots, \Omega_n \subseteq \mathsf{Plays}(\mathfrak{M})$  be objectives such that  $f = \sum_{j=1}^n \alpha_j \mathbb{1}_{\Omega_j}$ . The function  $x \mapsto \mathbb{E}_s^{\tau_x}(f)$  is measurable: it is a non-negative linear combination of measurable functions by the above. It follows from Equation (3.1) applied for all  $1 \leq j \leq n$  and the linearity of the Lebesgue integral that

$$\mathbb{E}_s^{\sigma}(f) = \sum_{j=1}^n \alpha_j \mathbb{P}_s^{\sigma}(\Omega_j) = \sum_{j=1}^n \alpha_j \int_{x \in [0,1]^{\omega}} \mathbb{P}_s^{\tau_x}(\Omega_j) = \int_{x \in [0,1]^{\omega}} \mathbb{E}_s^{\tau_x}(f).$$

Third, we assume that f is a non-negative measurable function. Let  $(f_n)_{n\in\mathbb{N}}$  be a sequence of measurable simple functions increasing to f (i.e., for all plays  $\pi \in \mathsf{Plays}(\mathfrak{M})$ ,  $f_n(\pi) \leq f_{n+1}(\pi)$  and  $\lim_{n\to\infty} f_n(\pi) = f(\pi)$ ). By the monotone convergence theorem and the previous point on simple functions, we have

$$\mathbb{E}_s^{\sigma}(f) = \lim_{n \to \infty} \mathbb{E}_s^{\sigma}(f_n) = \lim_{n \to \infty} \int_{x \in [0,1]^{\omega}} \mathbb{E}_s^{\tau_x}(f_n) d\nu_{\sigma}(x). \tag{3.2}$$

For all  $x \in [0,1]^{\omega}$ , by the monotone convergence theorem,  $\lim_{n\to\infty} \mathbb{E}_s^{\tau_x}(f_n) = \mathbb{E}_s^{\tau_x}(f)$ Therefore, the sequence of functions  $(x \mapsto \mathbb{E}_s^{\tau_x}(f_n))_{n\in\mathbb{N}}$  increases to  $x \mapsto \mathbb{E}_s^{\tau_x}(f)$ , implying that this function is measurable. The monotone convergence theorem allows us to exchange the limit and integral in the rightmost term of Equation (3.2), and implies that:

$$\mathbb{E}_s^{\sigma}(f) = \int_{x \in [0,1]^{\omega}} \lim_{n \to \infty} \mathbb{E}_s^{\tau_x}(f_n) d\nu_{\sigma}(x) = \int_{x \in [0,1]^{\omega}} \mathbb{E}_s^{\tau_x}(f) d\nu_{\sigma}(x).$$

We introduce some notation for the two last cases. We let  $f^+ = \max(f,0)$  and  $f^- = \max(-f,0)$  denote the non-negative and non-positive parts of f respectively; we have  $f = f^+ - f^-$ . From the above, we obtain that for all universally unambiguously integrable payoffs, the function  $x \mapsto \mathbb{E}_s^{\tau_x}(f)$  is measurable; it is the difference of the measurable non-negative functions  $x \mapsto \mathbb{E}_s^{\tau_x}(f^+)$  and  $x \mapsto \mathbb{E}_s^{\tau_x}(f^-)$ .

For the second-to-last case, we assume that f is  $\mathbb{P}_s^{\sigma}$ -integrable. We prove that the mappings  $x \mapsto \mathbb{E}_s^{\tau_x}(f^+)$  and  $x \mapsto \mathbb{E}_s^{\tau_x}(f^-)$  are  $\nu_{\sigma}$ -integrable. We proceed by bounding these functions by a  $\nu_{\sigma}$ -integrable function. For all  $x \in [0,1]^{\omega}$ , we have  $\mathbb{E}_s^{\tau_x}(f^+), \mathbb{E}_s^{\tau_x}(f^-) \leq \mathbb{E}_s^{\tau_x}(|f|)$ . By the above, we have  $\int_{x \in [0,1]^{\omega}} \mathbb{E}_s^{\tau_x}(|f|) \mathrm{d}\nu_{\sigma}(x) = \mathbb{E}_s^{\sigma}(|f|)$ , which is a real number since f is  $\mathbb{P}_s^{\sigma}$ -integrable. We have shown that  $x \mapsto \mathbb{E}_s^{\tau_x}(|f|)$  is  $\nu_{\sigma}$ -integrable, which implies that  $x \mapsto \mathbb{E}_s^{\tau_x}(f^+)$  and  $x \mapsto \mathbb{E}_s^{\tau_x}(f^-)$  also are. It follows that  $x \mapsto \mathbb{E}_s^{\tau_x}(f)$  is  $\nu_{\sigma}$ -integrable.

By definition, we have  $\mathbb{E}_s^{\sigma'}(f) = \mathbb{E}_s^{\sigma'}(f^+) - \mathbb{E}_s^{\sigma'}(f^-)$  for all strategies  $\sigma' \in \Sigma(\mathfrak{M})$ . Combining this with the linearity of the Lebesgue integral and the result for non-negative payoffs yields the following sequence of equalities:

$$\mathbb{E}_{s}^{\sigma}(f) = \mathbb{E}_{s}^{\sigma}(f^{+}) - \mathbb{E}_{s}^{\sigma}(f^{-}) 
= \int_{x \in [0,1]^{\omega}} \mathbb{E}_{s}^{\tau_{x}}(f^{+}) d\nu_{\sigma}(x) - \int_{x \in [0,1]^{\omega}} \mathbb{E}_{s}^{\tau_{x}}(f^{-}) d\nu_{\sigma}(x) 
= \int_{x \in [0,1]^{\omega}} \mathbb{E}_{s}^{\tau_{x}}(f^{+}) - \mathbb{E}_{s}^{\tau_{x}}(f^{-}) d\nu_{\sigma}(x) 
= \int_{x \in [0,1]^{\omega}} \mathbb{E}_{s}^{\tau_{x}}(f) d\nu_{\sigma}(x).$$

To deal with the last case, we assume that  $\inf_{\tau} \mathbb{E}_s^{\tau}(f) \geq 0$ . The analogous case  $\sup_{\tau} \mathbb{E}_s^{\tau}(f) \leq 0$  can be recovered from the case  $\inf_{\tau} \mathbb{E}_s^{\tau}(f) \geq 0$  by considering -f as the payoff function. We assume that f is not  $\mathbb{P}_s^{\sigma}$ -integrable, as this case has been examined above, i.e., that  $\mathbb{E}_s^{\sigma}(f) = +\infty$ . The integral  $\int_{x \in [0,1]^{\omega}} \mathbb{E}_s^{\tau_x}(f) d\nu_{\sigma}(x)$  is formally well-defined by the assumption that  $\inf_{\tau} \mathbb{E}_s^{\tau}(f) \geq 0$ . To end the argument, we must show that this integral is  $+\infty$ . Assume towards a contradiction that this is not the case. This implies that  $x \mapsto \mathbb{E}_s^{\tau_x}(f)$  is  $\nu_{\sigma}$ -integrable. From the result for non-negative payoffs, we obtain that  $\int_{x \in [0,1]^{\omega}} \mathbb{E}_s^{\tau_x}(f^+) d\nu_{\sigma}(x) = \mathbb{E}_s^{\sigma}(f^+) = +\infty$  and  $\int_{x \in [0,1]^{\omega}} \mathbb{E}_s^{\tau_x}(f^-) d\nu_{\sigma}(x) = \mathbb{E}_s^{\sigma}(f^-) \in \mathbb{R}$ . By linearity of the Lebesgue integral (for  $\nu_{\sigma}$ -integrable payoffs), we obtain that

$$\mathbb{E}_{s}^{\sigma}(f^{+}) = \int_{x \in [0,1]^{\omega}} \mathbb{E}_{s}^{\tau_{x}}(f^{+}) d\nu_{\sigma}(x) 
= \int_{x \in [0,1]^{\omega}} (\mathbb{E}_{s}^{\tau_{x}}(f) + \mathbb{E}_{s}^{\tau_{x}}(f^{-})) d\nu_{\sigma}(x) 
= \int_{x \in [0,1]^{\omega}} \mathbb{E}_{s}^{\tau_{x}}(f) d\nu_{\sigma}(x) + \int_{x \in [0,1]^{\omega}} \mathbb{E}_{s}^{\tau_{x}}(f^{-}) d\nu_{\sigma}(x).$$

This is a contradiction: on the one hand, we have  $\mathbb{E}_s^{\sigma}(f^+) = +\infty$  and, on the other hand, the sum in the last term is a real number. This ends the argument for the case  $\inf_{\tau} \mathbb{E}_s^{\tau}(f) \geq 0$ .

We highlight two major consequences of Lemma 3.1. On the one hand, for all strategies whose expected payoff is real, there exists a pure strategy with a greater expected payoff. On the other hand, if there exists a strategy with an infinite expected payoff, then there are pure strategies with arbitrarily large expected payoffs in absolute value. We note that even if a randomised strategy has an infinite expectation, there need not exist a pure strategy with infinite expectation. This is analogous to the fact that real-valued random variables can have an infinite expectation.

Using Lemma 3.1, we can obtain a characterisation of universally integrable payoffs. Let  $\Sigma \in \{\Sigma(\mathfrak{M}), \Sigma_{\mathsf{pure}}(\mathfrak{M})\}$ . A payoff f is universally integrable if and only if for all  $s \in S$ ,  $\sup_{\sigma \in \Sigma} \mathbb{E}^s_s(|f|)$  is real. The non-trivial part of the proof is showing that the definition of universally integrable and the property when the supremum ranges over pure strategies both imply the property with the supremum ranging over all strategies. We show the contrapositive of both implications. We assume that for some  $s \in S$ ,  $\sup_{\sigma \in \Sigma} \mathbb{E}^\sigma_s(|f|) = +\infty$ . Lemma 3.1 then implies that there are pure strategies  $\tau$  with arbitrarily large  $\mathbb{E}^\tau_s(|f|)$ , which implies the validity of one of the implications. For the other, we mix these pure strategies to construct a strategy  $\sigma$  such that  $\mathbb{E}^\sigma_s(|f|) = +\infty$ .

**Lemma 3.2.** Let f be a payoff. Let  $s \in S$ . The following assertions are equivalent.

- 1. f is  $\mathbb{P}_s^{\sigma}$ -integrable for all  $\sigma \in \Sigma(\mathfrak{M})$ .
- 2. We have  $\sup \{ \mathbb{E}_s^{\sigma}(|f|) \mid \sigma \in \Sigma(\mathfrak{M}) \} \in \mathbb{R}$ .
- 3. We have  $\sup \{ \mathbb{E}_s^{\sigma}(|f|) \mid \sigma \in \Sigma_{\mathsf{pure}}(\mathfrak{M}) \} \in \mathbb{R}$ .

In particular, f is universally integrable if and only if Item 2 (resp. 3) holds for all  $s \in S$ .

*Proof.* Item 2 directly implies the other two items. We now show that the other two items imply Item 2 via the contrapositive of these implications.

Assume that Item 2 does not hold, i.e., that  $\sup\{\mathbb{E}_s^{\sigma}(|f|) \mid \sigma \in \Sigma(\mathfrak{M})\} = +\infty$ . If there exists a pure strategy  $\sigma$  such that  $\mathbb{E}_s^{\sigma}(|f|) = +\infty$ , the negations of Items 1 and 3 follow directly. In the remainder of the proof, we assume that this is not the case.

First, we show that Item 3 does not hold. By Lemma 3.1, for all strategies  $\sigma$ , if  $\mathbb{E}_s^{\sigma}(|f|) \in \mathbb{R}$ , there exists a pure strategy  $\tau$  such that  $\mathbb{E}_s^{\tau}(|f|) \geq \mathbb{E}_s^{\sigma}(|f|)$  and, otherwise, if  $\mathbb{E}_s^{\sigma}(|f|) = +\infty$ , then for all  $M \in \mathbb{R}$ , there exists a pure strategy  $\tau$  such that  $\mathbb{E}_s^{\tau}(|f|) \geq M$ . In particular, Item 3 does not hold.

We now construct a strategy  $\sigma$  such that  $\mathbb{E}_s^{\sigma}(|f|) = +\infty$  from the pure strategies above. For all  $r \in \mathbb{N}$ , let  $\tau_r$  be a pure strategy such that  $\mathbb{E}_s^{\tau_r}(|f|) \geq 2^r$ . Let  $\sigma$  be a strategy with the same outcome as the mixed strategy that randomises over the set of strategies  $\{\tau_r \mid r \in \mathbb{N}\}$  and selects strategy  $\tau_r$  with probability  $\frac{1}{2^{r+1}}$ . Kuhn's theorem [Aum64], which implies the equivalence of mixed and behavioural strategies in MDPs, guarantees the existence of such a strategy. We directly obtain  $\mathbb{E}_s^{\sigma}(|f|) = +\infty$ . This ends the proof that Item 1 does not hold.

Let f be a universally unambiguously integrable payoff. We now use Lemma 3.2 to prove a useful property of universally unambiguously integrable payoffs. We show that, for all initial states s, there either exists a lower or upper bound on the expectation of strategies from s. In practice, this property guarantees that we can shift payoffs so Lemma 3.1 can be applied to the shifted payoff.

**Lemma 3.3.** Let f be a universally unambiguously integrable payoff function. For all  $s \in S$ , we have  $\inf_{\sigma \in \Sigma(\mathfrak{M})} \mathbb{E}^{\sigma}_{s}(f) \in \mathbb{R}$  or  $\sup_{\sigma \in \Sigma(\mathfrak{M})} \mathbb{E}^{\sigma}_{s}(f) \in \mathbb{R}$ .

*Proof.* Let  $s \in S$ . Let  $f^+ = \max(f,0)$  and  $f^- = \max(0,-f)$ . Assume towards a contradiction that  $\inf_{\sigma \in \Sigma(\mathfrak{M})} \mathbb{E}^{\sigma}_{s}(f) \notin \mathbb{R}$  and  $\sup_{\sigma \in \Sigma(\mathfrak{M})} \mathbb{E}^{\sigma}_{s}(f) \notin \mathbb{R}$ . Because  $\Sigma(\mathfrak{M})$  is non-empty, we therefore have  $\inf_{\sigma \in \Sigma(\mathfrak{M})} \mathbb{E}^{\sigma}_{s}(f) = -\infty$  and  $\sup_{\sigma \in \Sigma(\mathfrak{M})} \mathbb{E}^{\sigma}_{s}(f) = +\infty$ . We show that this implies that f is not universally unambiguously integrable.

We observe that for all  $\sigma \in \Sigma(\mathfrak{M})$ , we have  $\mathbb{E}_s^{\sigma}(f) \leq \mathbb{E}_s^{\sigma}(f^+)$  and  $\mathbb{E}_s^{\sigma}(-f) \leq \mathbb{E}_s^{\sigma}(f^-)$ . It follows from Lemma 3.2 that there exists a strategy  $\sigma_+$  (resp.  $\sigma_-$ ) such that  $f^+$  (resp.  $f^-$ ) is not  $\mathbb{P}_s^{\sigma_+}$ -integrable (resp.  $\mathbb{P}_s^{\sigma_-}$ -integrable). In particular, we obtain that  $\mathbb{E}_s^{\sigma_+}(f^+) = \mathbb{E}_s^{\sigma_-}(f^-) = +\infty$ .

By Kuhn's theorem [Aum64] (mixed and behavioural strategies are equally powerful in this context), there exist mixed strategies that are equivalent to  $\sigma_+$  and  $\sigma_-$ . The mixed strategy  $\sigma$  that averages these two mixed strategies can be shown to satisfy  $\mathbb{E}_s^{\sigma}(f^+) = \mathbb{E}_s^{\sigma}(f^-) = +\infty$ . This shows that f is not universally unambiguously integrable: f does not have an unambiguous  $\mathbb{P}_s^{\sigma}$ -integral.

## 3.3 Some classical objectives and payoffs

We now present some classical objectives and payoffs functions. First, we define reachability objectives. A reachability objective requires that a set of target states be visited. Formally, given a set of target states (target for short)  $T \subseteq S$ , we define the reachability objective Reach(T) as the set  $\{s_0a_0s_1a_1 \ldots \in \mathsf{Plays}(\mathfrak{M}) \mid \exists \ell \in \mathbb{N}, s_\ell \in T\}$ .

The other payoff functions we introduce are defined in a context where transitions of the MDP have a numerical weight assigned to them. Formally, a weight function is a function  $w \colon S \times A \to \mathbb{R}$ . We let w denote a weight function. A discounted-sum payoff is defined as an accumulated sum of weights multiplied by powers of a discount factor in [0,1[. Formally, given a discount factor  $\lambda \in [0,1[$ , we let  $\mathsf{DSum}_w^{\lambda} \colon \mathsf{Plays}(\mathfrak{M}) \to \mathbb{R}$  be the payoff function defined by  $\mathsf{DSum}_w^{\lambda}(\pi) = \sum_{\ell=0}^{\infty} \lambda^{\ell} w(s_{\ell}, a_{\ell})$  for all plays  $\pi = s_0 a_0 s_1 a_1 \ldots \in \mathsf{Plays}(\mathfrak{M})$ . A discounted-sum payoff is bounded and thus universally integrable.

A total-reward payoff corresponds to the accumulated weights along a play with no discounting. Formally, we let  $\mathsf{TRew}_w\colon\mathsf{Plays}(\mathfrak{M})\to\bar{\mathbb{R}}$  be defined, for all plays  $\pi=s_0a_0s_1a_1\ldots$ , by  $\liminf_{r\to\infty}\sum_{\ell=0}^r w(s_\ell,a_\ell)$ . We use a limit-inferior to ensure that this payoff is well-defined for all plays, as the series of weights along a play need not converge.

A shortest-path payoff can be seen as a quantitative variant of reachability and is defined as the accumulated sum of weights up to the first visit of a set of targets. Formally, given a target T, we let  $\mathsf{SPath}_w^T\colon\mathsf{Plays}(\mathfrak{M})\to\bar{\mathbb{R}}$  be defined by, for all plays  $\pi=s_0a_0s_1a_1\ldots$ ,  $\mathsf{SPath}_w^T(\pi)=+\infty$  if  $\pi\notin\mathsf{Reach}(T)$  and, otherwise,  $\mathsf{SPath}_w^T(\pi)=\sum_{\ell=0}^{r-1}w(s_\ell,a_\ell)$  where  $r=\min\{\ell\in\mathbb{N}\mid s_\ell\in T\}$ . Traditionally, the goal is to minimise the shortest-path payoff (i.e., it is a cost function rather than a payoff), hence the infinite payoff whenever the target is not visited.

All total-reward and shortest-path payoffs we consider are built on a non-negative weight function and are thus universally unambiguously integrable.

### 3.4 Continuous payoffs

The set of plays of  $\mathfrak{M}$  can be equipped with a natural topology that is metrisable and compact. This allows us to define continuous payoffs. We defer a formal presentation of the topology of  $\mathsf{Plays}(\mathfrak{M})$  to Section B.1, and instead provide a direct definition of continuous payoffs here.

Intuitively, a payoff is continuous if plays with a long common prefix have a close payoff. Let  $f \colon \mathsf{Plays}(\mathfrak{M}) \to \overline{\mathbb{R}}$  be a payoff and let  $\pi \in \mathsf{Plays}(\mathfrak{M})$ . If  $f(\pi) \in \mathbb{R}$ , then f is continuous at  $\pi$  if and only if for all  $\varepsilon > 0$ , there exists  $\ell \in \mathbb{N}$  such that for all  $\pi' \in \mathsf{Cyl}(\pi_{\leq \ell})$ ,  $|f(\pi) - f(\pi')| < \varepsilon$ . If  $f(\pi) = +\infty$  (resp.  $-\infty$ ), then f is continuous at  $\pi$  if and only if for all  $M \in \mathbb{R}$ , there exists  $\ell \in \mathbb{N}$  such that for all plays  $\pi' \in \mathsf{Cyl}(\pi_{\leq \ell})$ ,  $f(\pi') \geq M$  (resp.  $f(\pi') \leq -M$ ). The payoff f is continuous if it is continuous at all plays. By compactness of  $\mathsf{Plays}(\mathfrak{M})$ , continuous real-valued payoffs are uniformly continuous. Formally, if  $f \colon \mathsf{Plays}(\mathfrak{M}) \to \mathbb{R}$  is continuous, then for all  $\varepsilon > 0$ , there exists  $\ell \in \mathbb{N}$  such that for all plays  $\pi, \pi' \in \mathsf{Plays}(\mathfrak{M})$ ,  $\pi_{<\ell} = \pi'_{<\ell}$  implies that  $|f(\pi) - f(\pi')| < \varepsilon$ .

Discounted-sum payoffs are continuous (and thus uniformly continuous) no matter the discount factor and weight function. Given a positive weight function, any shortest-path function built on this weight function is also continuous. We prove slight generalisations of these claims and provide characterisation of continuous indicators and continuous prefix-independent payoffs in Appendix B.2.

## 3.5 Multiple payoffs

We now introduce terminology and notation for MDPs with multiple payoff functions. Let  $d \in \mathbb{N}_0$ . We summarise d payoffs  $f_1, \ldots, f_d$ : Plays $(\mathfrak{M}) \to \overline{\mathbb{R}}$  as a multi-dimensional payoff  $\overline{f}$ : Plays $(\mathfrak{M}) \to \mathbb{R}^d$ , and write  $\overline{f} = (f_j)_{1 \le j \le d}$ .

Let  $\bar{f} = (f_j)_{1 \leq j \leq d}$  and let  $s \in S$ . We say that  $\bar{f}$  is universally (resp. unambiguously) integrable whenever  $f_j$  is universally (resp. unambiguously) integrable for all  $1 \leq j \leq d$ . We now assume that  $\bar{f}$  is universally unambiguously integrable (recall that we focus on such payoffs). Given a set of strategies  $\Sigma \subseteq \Sigma(\mathfrak{M})$ , we let  $\mathsf{Pay}_s^{\Sigma}(\bar{f}) = \{\mathbb{E}_s^{\sigma}(\bar{f}) \mid \sigma \in \Sigma\}$  denote the set of expected payoffs of strategies in  $\Sigma$  from s. We let  $\mathsf{Pay}_s(\bar{f})$  and  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$  be shorthand for  $\mathsf{Pay}_s^{\Sigma(\mathfrak{M})}(\bar{f})$  and  $\mathsf{Pay}_s^{\Sigma(\mathfrak{M})}(\bar{f})$  respectively. We refer to elements of  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$  as pure expected payoffs. A set of expected payoffs need not have a maximum for the component-wise order, e.g., there can be several Pareto-optimal payoffs.

We now claim that convex combinations of elements of  $\mathsf{Pay}_s(\bar{f})$  are well-defined under the convention that  $0\cdot (+\infty) = 0\cdot (-\infty) = 0$ . This convention essentially boils down to ignoring terms of convex combinations that have a zero coefficient. These convex combinations are well-defined because  $\mathsf{Pay}_s(\bar{f}) \subseteq \prod_{j=1}^d \left[\inf_{\sigma \in \Sigma(\mathfrak{M})} \mathbb{E}^\sigma_s(f_j), \sup_{\sigma \in \Sigma(\mathfrak{M})} \mathbb{E}^\sigma_s(f_j)\right]$ . By Lemma 3.3, each interval in this product has a finite (upper or lower) bound and thus no terms of the form  $+\infty -\infty$  occur in any component of convex combinations of expected payoffs. It follows that any convex combination of elements of  $\mathsf{Pay}_s(\bar{f})$  is well-defined and is an element of  $\mathsf{Pay}_s(\bar{f})$ . Furthermore,  $\mathsf{Pay}_s(\bar{f}) \cap \mathbb{R}^d$  is a convex set. The last two properties can be established directly using mixed strategies.

In multi-objective optimisation, the goal is to ensure a given threshold on each dimension. This is formalised by the notion of achievable vectors. A vector  $\mathbf{q} \in \mathbb{R}^d$  is achievable (from s) if there exists a strategy  $\sigma$  such that  $\mathbf{q} \leq \mathbb{E}_s^{\sigma}(\bar{f})$ . In this case, we say that  $\sigma$  witnesses that  $\mathbf{q}$  is achievable. For any class of strategies  $\Sigma \subseteq \Sigma(\mathfrak{M})$ , we let  $\mathsf{Ach}_s^{\Sigma}(\mathbf{q}) = \mathsf{down}(\mathsf{Pay}_s^{\Sigma}(\bar{f}))$  denote the set of vectors such that a strategy of  $\Sigma$  witnesses that they are achievable. We define  $\mathsf{Ach}_s(\bar{f})$  and  $\mathsf{Ach}_s^{\mathsf{pure}}(\bar{f})$  as above.

We also consider the lexicographic optimisation of multiple objectives. For this case, we define an analogue of optimal strategies from the one-dimensional setting. A strategy  $\sigma$  is lexicographically optimal if  $\mathbb{E}_s^{\sigma}(\bar{f})$  is the lexicographic maximum of  $\mathsf{Pay}_s(\bar{f})$ .


## 4 An introductory example


The goal of this section is to illustrate that expected payoff sets in multi-objective MDPs may be complex. We provide a two-dimensional example illustrating that sets of expected payoffs are not necessarily polytopes, even when the payoffs are universally integrable. We introduce our example and comment on several of its properties in Section 4.1. We formally prove that these properties hold in Section 4.2.

### 4.1 Example overview

We consider the MDP  $\mathfrak{M}$  depicted in Figure 4.1a and let w denote the two-dimensional weight function from the illustration. On this MDP, we consider the two-dimensional payoff  $\bar{f} = (f_1, f_2)$  given by the discounted-sum payoffs  $f_1 = \mathsf{DSum}_{w_1}^{3/4}$  and  $f_2 = \mathsf{DSum}_{w_2}^{1/2}$ . MDPs with several discounted-sum payoffs with different discount factors have previously been studied in [CFW13].

Due to the absence of randomness in transitions, the expected payoff of any pure





(a) An MDP with deterministic transitions. Pairs next to actions represent two-dimensional weights.

(b) The set of expected payoffs for the MDP of Figure 4.1a for the payoff  $f_1 = \mathsf{DSum}_{w_1}^{3/4}$  and  $f_2 = \mathsf{DSum}_{w_2}^{1/2}$ .

Figure 4.1: An MDP with a two-dimensional discounted-sum payoff  $\bar{f}$  such that  $\mathsf{extr}(\mathsf{Pay}_{so}(\bar{f}))$  is infinite.

strategy from  $s_0$  is the payoff of a play from  $s_0$ . Therefore, we obtain that

$$\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) = \{(0,2), (1,2)\} \cup \left\{ \left(1 + \frac{3^r}{4^{r-1}}, 2 - \frac{1}{2^{r-1}}\right) \mid r \in \mathbb{N} \right\}.$$

On the one hand, the payoffs (0,2) and (1,2) are obtained by moving from  $s_0$  to  $s_1$  and  $s_2$  respectively and looping in these states forever. On the other hand, for all  $r \in \mathbb{N}$ , the payoff  $\left(1 + \frac{3^r}{4^{r-1}}, 2 - \frac{1}{2^{r-1}}\right)$  is obtained by spending r rounds in  $s_2$  then moving to  $s_3$ ; for r = 0, we move from  $s_0$  to  $s_3$  directly.

We approximately illustrate  $\mathsf{Pay}_{s_0}(\bar{f})$  in Figure 4.1b. This illustration is based on the equality  $\mathsf{Pay}_{s_0}(\bar{f}) = \mathsf{conv}(\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}))$ . This equality follows from Thm. 6.2, which states that if  $\bar{f}$  is universally integrable, then this equality holds. In the interest of keeping this section self-contained, we outline a direct argument for this simple example that does not rely on Theorem 6.2. Because  $\mathfrak{M}$  has countably many plays, the expected payoff of any (randomised) strategy can be seen as an infinite convex combination of payoffs of plays of  $\mathfrak{M}$ . Any such infinite convex combination can be shown to be in the closure of  $\mathsf{conv}(\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}))$ . Furthermore,  $\mathsf{conv}(\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}))$  is closed because the convex hull of a compact set is compact and  $\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f})$  is compact. This shows that  $\mathsf{Pay}_{s_0}(\bar{f}) = \mathsf{conv}(\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}))$  and justifies the illustration.

Since  $\mathsf{Pay}_{s_0}(\bar{f}) = \mathsf{conv}(\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}))$ , we conclude that all extreme points of  $\mathsf{Pay}_{s_0}(\bar{f})$  can be obtained by using pure strategies. Indeed, any vector of  $\mathsf{Pay}_{s_0}(\bar{f})$  that cannot be obtained by a pure strategy is a convex combination of the expected payoffs of pure strategies, and thus is not extreme. There exists another means of showing that a point can be obtained by a pure strategy. This technique involves reducing to a single-dimensional setting by composing a linear form with  $\bar{f}$ . However, this approach does not work for  $(1,2) \in \mathsf{Pay}_{s_0}(\bar{f})$ .

First, let us explain how to establish that a point can be obtained by a pure strategy via a reduction to a single-dimensional payoff. Let  $\mathbf{q} \in \mathsf{Pay}_{s_0}(\bar{f})$ . To prove that  $\mathbf{q} \in \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$ , one considers a linear form  $x^*$  such that  $x^*(\mathbf{q}) > x^*(\mathbf{p})$  for all  $\mathbf{p} \in \mathsf{Pay}_{s_0}(\bar{f})$ . The existence of such a form is equivalent to the existence of a hyperplane H supporting  $\mathsf{Pay}_{s_0}(\bar{f})$  at  $\mathbf{q}$  such that  $\mathsf{Pay}_{s_0}(\bar{f}) \cap H = \{\mathbf{q}\}$ . By linearity of the expectation,  $x^* \circ \bar{f}$  is universally integrable and any strategy  $\sigma$  that maximises the expectation of  $x^* \circ \bar{f}$  from  $s_0$  must satisfy  $\mathbb{E}^{\sigma}_{s_0}(\bar{f}) = \mathbf{q}$  (such a strategy exists because  $\mathbf{q} \in \mathsf{Pay}_{s_0}(\bar{f})$ ). Lemma 3.1 then implies that  $\mathbf{q}$  can be obtained with a pure strategy. This argument can be used whenever the set of

expected payoffs is a compact polytope to prove that all vertices can be attained by pure strategies.

We now demonstrate that this argument is not applicable to show that  $(1,2) \in \mathsf{extr}(\mathsf{Pay}_{s_0}(\bar{f}))$  is the payoff of a pure strategy. We observe that the only hyperplane that supports  $\mathsf{Pay}_{s_0}(\bar{f})$  at (1,2) is the line H carrying the segment [(0,2),(1,2)]. This is highlighted by the orange dotted lines in Figure 4.1b: angling H in any way yields a line that is not a supporting hyperplane. Furthermore, this hyperplane is not suitable for the above argument because  $H \cap \mathsf{Pay}_{s_0}(\bar{f}) = [(0,2),(1,2)]$ . This implies that there is no hyperplane supporting  $\mathsf{Pay}_{s_0}(\bar{f})$  at (1,2) that does not contain other points of  $\mathsf{Pay}_{s_0}(\bar{f})$ . Therefore, we cannot show that  $(1,2) \in \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$  by following the argument outlined above. In Section 6, we reason on MDPs with lexicographic optimisation to prove Theorem 6.2, and as a corollary, obtain that all extreme points of a set of expected payoffs can be attained by a pure strategy.

We conclude from the above that  $\mathsf{Pay}_{s_0}(\bar{f})$  is not a polytope. Indeed, the vertices of a polytope can be isolated with supporting hyperplanes. Thus, if  $\mathsf{Pay}_{s_0}(\bar{f})$  were a polytope, we would have been able to conclude that (1,2) is the payoff of a pure strategy via the described reduction to a single dimension. This implies that  $\mathsf{Pay}_{s_0}(\bar{f})$  has infinitely many extreme points. In fact, we can show that  $\mathsf{extr}(\mathsf{Pay}_{s_0}(\bar{f})) = \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$  and that all of these payoffs aside from (0,2) are Pareto-optimal. This illustrates that even for classical payoffs, the expected payoff and achievable sets can be quite complex.

Finally, we comment on the strategy complexity required to obtain certain expected payoffs. More precisely, we briefly discuss memory requirements. Memory requirements in games and MDPs are thoroughly studied (see Sect. 1) and constitute a central measure of complexity for strategies in the literature. All but three extreme points of  $\mathsf{Pay}_{s_0}(\bar{f})$  are obtained by moving from  $s_0$  to  $s_2$  and looping there finitely many times before moving to  $s_3$ . In other words, these extreme points require pure strategies that count up to some arbitrarily large number. In fact, in this case, we can only obtain these payoffs by using these specific pure strategies. Intuitively, the expected payoff of a randomised strategy that induces more than one play is a non-trivial convex combination of payoffs of several plays, and therefore not in  $\mathsf{extr}(\mathsf{Pay}_{s_0}(\bar{f}))$ . This implies that some expected payoffs need strategies with arbitrarily large albeit finite memory to be obtained in this instance.

### 4.2 Proofs and details

We now prove all statements that were made in Section 4.1. First, we prove that the description of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$  given above is accurate. Second, we show that  $\mathsf{Pay}_{s_0}(\bar{f}) = \mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}))$  in the context of our example. When then prove that we cannot conclude that  $(1,2) \in \mathsf{extr}(\mathsf{Pay}_{s_0}(\bar{f}))$  by reducing to a single-dimensional payoff via a linear form. Next, we establish that all pure payoffs are extreme points and that all of these points except (0,2) are Pareto-optimal. Finally, we close the section by proving that all payoffs of pure strategies can *only* be obtained by playing without randomisation, and comment on the consequences in terms of strategy complexity.

Throughout this section,  $\mathfrak{M}$  refers to the MDP of Figure 4.1a.

#### 4.2.1 Determining the set of payoffs of pure strategies

We prove that the description of  $\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f})$  given in Section 4.1 is correct. The argument is based on the fact that there are no randomised transitions in the MDP we consider. Therefore, the payoff of a pure strategy from  $s_0$  is the payoff of a single play. In the following proof, we directly compute the payoff of each play from  $s_0$ .

**Lemma 4.1.** We have 
$$\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}) = \{(0,2),(1,2)\} \cup \{(1 + \frac{3^r}{4^{r-1}}, 2 - \frac{1}{2^{r-1}}) \mid r \in \mathbb{N}\}$$
.

*Proof.* There are no randomised transitions in  $\mathfrak{M}$ . For this reason, any pure strategy induces a single play in  $\mathfrak{M}$  from any starting state. We compute the payoff of all plays of  $\mathfrak{M}$  from  $s_0$  to obtain the desired result.

We first consider the three plays  $s_0c(s_1a)^{\omega}$ ,  $s_0a(s_2a)^{\omega}$  and  $s_0b(s_3a)^{\omega}$  that never leave their second state once it is reached. By definition of discounted-sum payoff functions, we have  $\bar{f}(s_0c(s_1a)^{\omega}) = (0, \sum_{\ell=0}^{\infty} \frac{1}{2^{\ell}}) = (0, 2)$ ,  $\bar{f}(s_0a(s_2a)^{\omega}) = (1, \sum_{\ell=0}^{\infty} \frac{1}{2^{\ell}}) = (1, 2)$  and  $\bar{f}(s_0b(s_3a)^{\omega}) = (1 + \sum_{\ell=0}^{\infty} (\frac{3}{4})^{\ell}, 0) = (5, 0) = (1 + \frac{3^0}{4^{0-1}}, 2 - \frac{1}{2^{0-1}})$ .

It remains to deal with the plays that move from  $s_0$  to  $s_2$  and then eventually move to  $s_3$ . It suffices to show that for all  $r \geq 1$ , we have  $\bar{f}(s_0(as_2)^rb(s_3a)^\omega) = (1 + \frac{3^r}{4^{r-1}}, 2 - \frac{1}{2^{r-1}})$ . Let  $r \geq 1$ . We obtain, by definition of discounted-sum payoff functions, that

$$f_1(s_0(as_2)^r b(s_3 a)^\omega) = 1 + \sum_{\ell=r}^\infty \frac{3^\ell}{4^\ell} = 1 + \frac{3^r}{4^r} \cdot \sum_{\ell=0}^\infty \frac{3^\ell}{4^\ell} = 1 + \frac{3^r}{4^{r-1}},$$

and

$$f_2(s_0(as_2)^r b(s_3 a)^\omega) = \sum_{\ell=0}^{r-1} \frac{1}{2^\ell} = \frac{1 - \frac{1}{2^r}}{1 - \frac{1}{2}} = 2 - \frac{1}{2^{r-1}}.$$

This proves the required equality to end the proof.

## 4.2.2 Convex combinations of pure payoffs suffice

Our goal is to prove that  $\mathsf{Pay}_{s_0}(\bar{f}) = \mathsf{conv}(\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}))$ . The inclusion  $\mathsf{conv}(\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f})) \subseteq \mathsf{Pay}_{s_0}(\bar{f})$  is direct by convexity of sets of expected payoffs (see Section 3.5). For the other inclusion, we break the proof down into three arguments like in the sketch above. First, we show that  $\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f})$  is closed.

**Lemma 4.2.** The set  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \subseteq \mathbb{R}^2$  is closed.

*Proof.* Let  $\mathbf{q} = (q_1, q_2) \in \mathbb{R}^2$  such that there exists a sequence  $(\mathbf{p}^{(n)})_{n \in \mathbb{N}} \subseteq \mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$  such that  $\lim_{n \to \infty} \mathbf{p}^{(n)} = \mathbf{q}$ . We show that  $\mathbf{q} \in \mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$ . We observe that necessarily,  $\mathbf{q} \in [0, 5] \times [0, 2]$  because  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}) \subseteq [0, 5] \times [0, 2]$  (this inclusion follows from Lemma 4.1).

We distinguish three cases. First, we assume that  $\mathbf{p}^{(n)} = (0,2)$  infinitely often. In this case, the constant sequence  $((0,2))_{n\in\mathbb{N}}$  is a subsequence of  $(\mathbf{p}^{(n)})_{n\in\mathbb{N}}$ . Since a convergent sequence has the same limit as its subsequences, this implies that  $\mathbf{q} = (0,2)$ . We assume, for the remainder of the proof, that (0,2) does not occur in the sequence  $(\mathbf{p}^{(n)})_{n\in\mathbb{N}}$ .

Second, we assume that  $q_2 \neq 2$ . In this case, there exists  $r \in \mathbb{N}$  such that  $q_2 \leq 2 - \frac{1}{2^{r-1}}$ . This, along with Lemma 4.1, implies that there is a suffix of  $(\mathbf{p}^{(n)})_{n \in \mathbb{N}}$  that is included in the finite subset  $\left\{\left(1 + \frac{3^{\ell}}{4^{\ell-1}}, 2 - \frac{1}{2^{\ell-1}}\right) \mid \ell \leq r\right\}$  of  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$ . We conclude that  $\mathbf{q} \in \mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$  as all finite sets are closed.

Finally, we assume that  $q_2=2$  and show that  $\mathbf{q}=(1,2)$ . If there are infinitely many  $n\in\mathbb{N}$  such that  $\mathbf{p}^{(n)}=(1,2)$ , then  $\mathbf{q}=(1,2)$  (refer to the first case). We therefore assume that (1,2) does not occur in  $(\mathbf{p}^{(n)})_{n\in\mathbb{N}}$ . For all  $n\in\mathbb{N}$ , we let  $\ell_n\in\mathbb{N}$  such that  $\mathbf{p}^{(n)}=\left(1+\frac{3^{\ell_n}}{4^{\ell_n-1}},2-\frac{1}{2^{\ell_n-1}}\right)$ . From  $q_2=2$ , we obtain that  $\lim_{n\to\infty}\frac{1}{2^{\ell_n-1}}=0$ . It follows that

$$q_1 = \lim_{n \to \infty} 3 \cdot \left(\frac{3}{4}\right)^{\ell_n - 1} = \lim_{n \to \infty} 3 \cdot \left(\frac{1}{2}\right)^{(\ell_n - 1) \cdot \log_2(4/3)} = 0.$$

This implies that  $\mathbf{q}=(1,2)\in\mathsf{Pay}^{\mathsf{pure}}_s(\bar{f})$ . This ends the proof that  $\mathsf{Pay}^{\mathsf{pure}}_s(\bar{f})$  is closed.  $\square$ 

We now prove a general result that implies that  $\mathsf{conv}(\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}))$  is closed. We show that the convex hull of any compact subset of  $\mathbb{R}^d$  is also compact. Lemma 4.2 allows us to apply it to  $\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f})$ . Recall that a subset of  $\mathbb{R}^d$  is compact if and only if it is closed and bounded.

**Lemma 4.3.** Let  $d \ge 1$ . Let  $D \subseteq \mathbb{R}^d$ . If D is compact, then conv(D) is also compact.

*Proof.* Assume that D is compact. We assume that D is non-empty, as otherwise the result is direct.

We first show that conv(D) is bounded. Let  $\mathbf{q} \in conv(D)$ . Let  $\alpha_1, \ldots, \alpha_n$  be convex combination coefficients and let  $\mathbf{p}_1, \ldots, \mathbf{p}_n \in D$  such that  $\mathbf{q} = \sum_{m=1}^n \alpha_m \mathbf{p}_m$ . By triangulation, we obtain that  $\|\mathbf{q}\|_2 \leq \sum_{m=1}^n \alpha_m \|\mathbf{p}_m\|_2 \leq \sup\{\|\mathbf{p}\|_2 \mid \mathbf{p} \in D\} \in \mathbb{R}$ , where the second inequality is a consequence of  $\sum_{m=1}^n \alpha_m = 1$ . It follows that conv(D) is bounded. We now show that conv(D) is closed. Let  $\mathbf{q} \in \mathbb{R}^d$  such that there exists a sequence

We now show that  $\operatorname{conv}(D)$  is closed. Let  $\mathbf{q} \in \mathbb{R}^d$  such that there exists a sequence  $(\mathbf{q}^{(n)})_{n \in \mathbb{N}} \subseteq \operatorname{conv}(D)$  such that  $\lim_{n \to \infty} \mathbf{q}^{(n)} = \mathbf{q}$ . By Carathéodory's theorem for convex hulls (Theorem 2.1), all elements of the sequence  $(\mathbf{q}^{(n)})_{n \in \mathbb{N}}$  are a convex combination of no more than d+1 elements of D. For all  $n \in \mathbb{N}$ , let  $\alpha_1^{(n)}, \ldots, \alpha_{d+1}^{(n)}$  be convex combination coefficients and let  $\mathbf{p}_1^{(n)}, \ldots, \mathbf{p}_{d+1}^{(n)}$  such that  $\mathbf{q}^{(n)} = \sum_{j=1}^{d+1} \alpha_j^{(n)} \mathbf{p}_j^{(n)}$ . By compactness of [0,1] and D, we obtain an increasing sequence of natural numbers  $(n_m)_{m \in \mathbb{N}}$ , convex combination coefficients  $\alpha_1, \ldots, \alpha_{d+1}$  and  $\mathbf{p}_1, \ldots, \mathbf{p}_{d+1} \in D$  such that for all  $1 \le j \le d+1$ ,  $\lim_{m \to \infty} \alpha_j^{(n_m)} = \alpha_j$  and  $\lim_{m \to \infty} \mathbf{p}_j^{(n_m)} = \mathbf{p}_j$ . It follows (from the uniqueness of the limit) that  $\mathbf{q} = \sum_{j=1}^{d+1} \alpha_j \mathbf{p}_j$ . This shows that  $\mathbf{q} \in \operatorname{conv}(D)$  and ends the proof that  $\operatorname{conv}(D)$  is closed.

Finally, we show that  $\mathsf{Pay}_{s_0}(\bar{f}) \subseteq \mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}))$  by establishing that all elements of  $\mathsf{Pay}_{s_0}(\bar{f})$  are included in the closure of  $\mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}))$ . We use the argument based on infinite convex combinations that was mentioned in Section 4.1.

 $\mathbf{Lemma\ 4.4.\ We\ have\ Pay}_{s_0}(\bar{f}) = \mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})).$ 

*Proof.* By Lemma 4.2 and Lemma 4.3,  $\operatorname{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}))$  is closed. To end the proof, it thus suffices to show that  $\mathsf{Pay}_{s_0}(\bar{f}) \subseteq \mathsf{cl}(\mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})))$ . We recall that there are countably many plays in  $\mathfrak{M}$ . Let  $(\pi_\ell)_{\ell \in \mathbb{N}}$  be an enumeration of the set of plays of  $\mathfrak{M}$  from  $s_0$ . Let  $\sigma \in \Sigma(\mathfrak{M})$  and let  $\mathbf{q} = \mathbb{E}^{\sigma}_{s_0}(\bar{f})$ . We can write

$$\mathbf{q} = \sum_{\ell=0}^{\infty} \mathbb{P}_{s_0}^{\sigma}(\{\pi_{\ell}\}) \cdot \bar{f}(\pi_{\ell}). \tag{4.1}$$

Next, we observe that  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$  is the set of payoffs of plays from  $s_0$  in  $\mathfrak{M}$  due to the absence of random transitions. Therefore, informally, Equation (4.1) implies that  $\mathbf{q}$  is an infinite convex combination of elements of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$ . To end the proof, it suffices to show that we can approach this infinite convex combination with finite convex combinations of elements of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$ . For all  $r \in \mathbb{N}$ , let

$$\mathbf{p}^{(r)} = \left( \mathbb{P}_{s_0}^{\sigma}(\{\pi_0\}) + \sum_{\ell=r+1}^{\infty} \mathbb{P}_{s_0}^{\sigma}(\{\pi_\ell\}) \right) \cdot \bar{f}(\pi_0) + \sum_{\ell=1}^{r} \mathbb{P}_{s_0}^{\sigma}(\{\pi_\ell\}) \cdot \bar{f}(\pi_\ell).$$

For all  $r \in \mathbb{N}$ ,  $\mathbf{p}^{(r)}$  is derived from the partial sum of the series in Equation (4.1) up to index r by increasing the coefficient of  $\bar{f}(\pi_0)$  enough to obtain a convex combination of pure payoffs. For all  $r \in \mathbb{N}$ , we have  $\mathbf{p}^{(r)} \in \mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}))$  and  $\mathbf{q} = \lim_{r \to \infty} \mathbf{p}^{(r)}$ . This shows that  $\mathbf{q} \in \mathsf{cl}(\mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})))$  and ends the proof.

### 4.2.3 One dimension is not enough to show that extreme points are pure

We formalise the intuition given on Figure 4.1b that the only hyperplane supporting  $\mathsf{Pay}_{s_0}(\bar{f})$  at (1,2) is the line carrying the segment [(0,2),(1,2)]. We reach a contradiction by assuming that there is a linear form  $x^*$  such that  $x^*((1,2)) > x^*((0,2))$  and  $x^*((1,2)) \ge x^*(\mathbf{p})$  for all  $\mathbf{p} \in \mathsf{Pay}_{s_0}(\bar{f})$ .

**Lemma 4.5.** Let  $\mathbf{q} = (1,2)$ . The unique supporting hyperplane of  $\mathsf{Pay}_{s_0}(\bar{f})$  containing  $\mathbf{q}$  is the line defined by the equation y = 2.

*Proof.* We assume towards a contradiction that there exists a linear form  $x^* \colon \mathbb{R}^2 \to \mathbb{R}$  such that for all  $\mathbf{p} \in \mathsf{Pay}_{s_0}(\bar{f})$ ,  $x^*(\mathbf{q}) \geq x^*(\mathbf{p})$  and  $x^*(\mathbf{q}) \neq x^*((0,2))$ . Let  $\alpha, \beta \in \mathbb{R}$  such that for all  $\mathbf{v} = (v_1, v_2) \in \mathbb{R}^2$ ,  $x^*(\mathbf{v}) = \alpha v_1 + \beta v_2$ .

We observe that  $x^*(\mathbf{q}) > x^*((0,2))$  is equivalent to  $\alpha > 0$ . We also have, for all  $\ell \in \mathbb{N}$ ,  $x^*(\mathbf{q}) \geq x^*((1+\frac{3^\ell}{4^{\ell-1}},2-\frac{1}{2^{\ell-1}}))$ , i.e.,  $\beta \geq \frac{3^\ell}{2^{\ell-1}} \cdot \alpha$ . Together, these statements imply that  $\beta$  must be greater than all real numbers, which is a contradiction.

We have shown that we cannot necessarily isolate extreme points of sets of expected payoffs by intersecting them with a single hyperplane.

## 4.2.4 Extreme points and Pareto-optimality

We now show that  $\operatorname{extr}(\mathsf{Pay}_{s_0}(\bar{f})) = \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$  in the context of this example (this property does not hold in full generality). It follows from  $\mathsf{Pay}_{s_0}(\bar{f}) = \mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}))$  that all extreme points of  $\mathsf{Pay}_{s_0}(\bar{f})$  are the payoff of a pure strategy. It remains to show that  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \subseteq \mathsf{extr}(\mathsf{Pay}_{s_0}(\bar{f}))$ . We first observe that  $(0,2) \in \mathsf{extr}(\mathsf{Pay}_{s_0}(\bar{f}))$  because it is with the least first component among all elements of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$ . The main difficulty lies with the elements of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{(0,2)\}$ .

We handle the remaining points with a two-part argument. First, we show that any non-extreme element of the border of  $\mathsf{Pay}_{s_0}(\bar{f})$  is a convex combination of two extreme points. This implies that all Pareto-optimal elements of  $\mathsf{Pay}_{s_0}(\bar{f})$  are convex combinations of no more than two vectors. Second, we prove that for all vectors  $\mathbf{q} \in \mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}) \setminus \{(0,2)\}$ ,  $\mathbf{q}$  is either incomparable or strictly greater (with respect to the component-wise ordering) to convex combinations of any two vectors of  $\mathsf{Pay}_{s_0}(\bar{f}) \setminus \{\mathbf{q}\}$ . We prove this by reasoning on a strictly concave real function whose graph includes  $\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}) \setminus \{(0,2)\}$ . The graphical intuition is as follows: any segment joining two points of the graph of a strictly concave function is beneath the curve, so any points on the segment that are comparable to  $\mathbf{q}$  must be smaller. This approach also yields that all vectors of  $\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}) \setminus \{(0,2)\}$  are Pareto-optimal elements of  $\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f})$ .

We now prove a generalisation of the first property formulated above: given a compact set  $D \subseteq \mathbb{R}^2$ , any vector in the border of  $\mathsf{conv}(D)$  is a convex combination of no more than two elements of D. This can be seen as a refinement of Carathéodory's theorem for convex hulls (Theorem 2.1) when considering the border of the convex hull of a compact set in a two-dimensional setting.

**Lemma 4.6.** Let  $D \subseteq \mathbb{R}^2$  be compact. For all  $\mathbf{q} \in \mathsf{bd}(\mathsf{conv}(D))$ , either  $\mathbf{q} \in \mathsf{extr}(\mathsf{conv}(D))$  or  $\mathbf{q}$  is a convex combination of two vectors of  $D \setminus \{\mathbf{q}\}$ .

*Proof.* If D is empty or a singleton set, the result is direct. We thus assume that D has at least two elements. Let  $\mathbf{q} \in \mathsf{bd}(\mathsf{conv}(D))$ . Because D is compact,  $\mathsf{conv}(D)$  is closed (Lemma 4.3) and thus  $\mathsf{bd}(\mathsf{conv}(D)) \subseteq \mathsf{conv}(D)$ . We let  $\mathbf{p}^{(1)}, \ldots, \mathbf{p}^{(n)} \in D$  and  $\alpha_1, \ldots, \alpha_n \in ]0,1]$  be non-zero convex combination coefficients such that  $\mathbf{q} = \sum_{m=1}^n \alpha_n \mathbf{p}^{(m)}$ .

We first show that the vectors  $\mathbf{q}$ ,  $\mathbf{p}^{(1)}$ , ...,  $\mathbf{p}^{(n)}$  lie on a single line. If  $\mathsf{aff}(D)$  is a line, then this is direct. We thus assume that  $\mathsf{aff}(D)$  is not a line. We obtain that  $\mathsf{aff}(D) = \mathbb{R}^2$ : it contains a line because D has at least two elements, and therefore its dimension must be two. This implies that  $\mathsf{ri}(D) = \mathsf{int}(D)$ , and thus that  $\mathbf{q} \notin \mathsf{ri}(D)$ . By the supporting hyperplane theorem (Theorem 2.3), there exists a non-zero linear form  $x^*$  such that for all  $\mathbf{p} \in \mathsf{conv}(D)$ ,  $x^*(\mathbf{q}) \geq x^*(\mathbf{p})$ . It follows that for all  $1 \leq m \leq n$ , we have  $x^*(\mathbf{p}^{(m)}) = x^*(\mathbf{q})$  (by linearity, because  $\alpha^{(m)} \neq 0$ ). In other words,  $\mathbf{q}$  and the  $\mathbf{p}^{(m)}$  lie on the line  $(x^*)^{-1}(x^*(\mathbf{q}))$ .

We have shown that  $\mathbf{q}$  and the  $\mathbf{p}^{(m)}$  lie on a single line. There exists a non-zero vector  $\mathbf{v} \in \mathbb{R}^2$  such that for all  $1 \leq m \leq n$ , there exists  $\beta_m \in \mathbb{R}$  such that  $\mathbf{p}^{(m)} = \mathbf{q} + \beta_m \cdot \mathbf{v}$ . There must exist  $1 \leq m, m' \leq n$  such that  $\beta_m \geq 0$  and  $\beta_{m'} \leq 0$ . We conclude that  $\mathbf{q} \in \left[\mathbf{p}^{(m)}, \mathbf{p}^{(m')}\right]$ . We have shown that  $\mathbf{q}$  is a convex combination of at most two elements of D.

We now introduce the strictly concave function  $\mathcal{F}: [1,5] \to \mathbb{R}$  used in the remainder of our argument. For all  $x \in [1,5]$ , we let

$$\mathcal{F}(x) = 2 - \left(\frac{x-1}{3}\right)^{\log_{4/3}(2)}$$

We observe that  $\mathcal{F}$  is well-defined in 0 because  $\log_{4/3}(2) > 0$ . We recall that  $\mathcal{F}$  is strictly concave if and only if for all  $x, x' \in [1, 5]$  such that x < x' and all  $\beta \in ]0, 1[$ , we have  $\beta \mathcal{F}(x) + (1-\beta)\mathcal{F}(x') < \mathcal{F}(\beta x + (1-\beta)x')$ . We now show that  $\mathcal{F}$  is decreasing and strictly concave.

**Lemma 4.7.** The function  $\mathcal{F}$  is decreasing and strictly concave.

*Proof.* To prove that  $\mathcal{F}$  is decreasing (resp. strictly concave), it suffices to show that it is differentiable over ]1,5[ and its derivative  $\mathcal{F}'$  is strictly negative (resp. decreasing). For the strict concavity of  $\mathcal{F}$ , in practice, we show that the second derivative  $\mathcal{F}''$  of  $\mathcal{F}$  (defined over ]1,5[) is negative. For all  $x \in ]1,5[$ , we have

$$\mathcal{F}'(x) = -\log_{4/3}(2) \cdot \left(\frac{x-1}{3}\right)^{\log_{4/3}(2)-1}$$

and

$$\mathcal{F}''(x) = -\log_{4/3}(2) \cdot \left(\log_{4/3}(2) - 1\right) \cdot \left(\frac{x-1}{3}\right)^{\log_{4/3}(2)-2}.$$

To prove that  $\mathcal{F}'$  and  $\mathcal{F}''$  are negative over ]1,5[, it suffices to show that  $\log_{4/3}(2) > 0$  and  $\log_{4/3}(2) - 1 > 0$ . The second inequality implies the first and can be shown to be equivalent to  $2 > \frac{4}{3}$ . This ends the proof that  $\mathcal{F}$  is decreasing and strictly concave.

Next, we prove that  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{(0,2)\}$  is included in the graph of  $\mathcal{F}$ .

**Lemma 4.8.** The graph of  $\mathcal F$  includes  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar f)\setminus\{(0,2)\}$ , i.e., for all  $(x,y)\in\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar f)$ ,  $\mathcal F(x)=y$ .

*Proof.* First, we observe that  $\mathcal{F}(1) = 2$ , i.e., (1,2) is in the graph of  $\mathcal{F}$ . It remains to show that for all  $\ell \in \mathbb{N}$ , we have

$$\mathcal{F}\left(1 + \frac{3^{\ell}}{4^{\ell - 1}}\right) = 2 - \frac{1}{2^{\ell - 1}}.$$

We let  $\ell \in \mathbb{N}$ . We obtain the following equalities:

$$\mathcal{F}\left(1 + \frac{3^{\ell}}{4^{\ell-1}}\right) = 2 - \left(\frac{3^{\ell-1}}{4^{\ell-1}}\right)^{\log_{4/3}(2)}$$

$$= 2 - \left(\frac{3}{4}\right)^{(\ell-1)\cdot\log_{4/3}(2)}$$

$$= 2 - \left(\left(\frac{4}{3}\right)^{\log_{4/3}(2)}\right)^{-(\ell-1)}$$

$$= 2 - \frac{1}{2^{\ell-1}}.$$

We have shown that all elements of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{(0,2)\}$  are in the graph of  $\mathcal{F}$ .

We now prove that all vectors  $\mathbf{q} \in \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{(0,2)\}$  are not smaller than convex combinations of two elements of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{\mathbf{q}\}$ .

**Lemma 4.9.** Let  $\mathbf{q} \in \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{(0,2)\}$ . For all  $\mathbf{p}_1, \mathbf{p}_2 \in \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{\mathbf{q}\}$  and all  $\alpha \in [0,1]$ ,  $\mathbf{q}$  is incomparable to or strictly greater than  $\alpha \cdot \mathbf{p}_1 + (1-\alpha) \cdot \mathbf{p}_2$ .

*Proof.* Let  $\mathbf{q} = (q_1, q_2)$ . Let  $\mathbf{p}_1, \mathbf{p}_2 \in \mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}) \setminus \{\mathbf{q}\}$ . We fix  $\alpha \in ]0,1[$ ; the above statement for  $\alpha \in \{0,1\}$  is covered by the cases  $\mathbf{p}_1 = \mathbf{p}_2$ . We let  $\mathbf{p} = \alpha \cdot \mathbf{p}_1 + (1-\alpha) \cdot \mathbf{p}_2$ . In the first part of this proof, we assume that  $\mathbf{p}_1 \neq (0,2)$  and  $\mathbf{p}_2 \neq (0,2)$  and discuss the case when this assumption is lifted at the end of the proof.

By Lemma 4.8, we have  $q_2 = \mathcal{F}(q_1)$  and we can write  $\mathbf{p}_1 = (x_1, \mathcal{F}(x_1))$  and  $\mathbf{p}_2 = (x_2, \mathcal{F}(x_2))$  where  $x_1$  and  $x_2$  are the first components of  $\mathbf{p}_1$  and  $\mathbf{p}_2$  respectively. We assume without loss of generality that  $x_1 \leq x_2$ . All elements of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$  have different x-components, hence  $x_1 \neq q_1$  and  $x_2 \neq q_1$ .

We first assume that  $x_1 \leq x_2 < q_1$ . In this case, by Lemma 4.7 ( $\mathcal{F}$  is strictly decreasing), we have  $\mathcal{F}(x_1) \geq \mathcal{F}(x_2) > q_1$ . It follows that  $\mathbf{q}$  and  $\mathbf{p}$  are incomparable: the first component of  $\mathbf{q}$  is greater than that of  $\mathbf{p}$  but the second component of  $\mathbf{q}$  is smaller than that of  $\mathbf{p}$ . In the case that  $q_1 < x_1 \leq x_2$ , we obtain in a similar fashion that  $\mathbf{q}$  and  $\mathbf{p}$  are incomparable.

We now assume that  $x_1 < q_1 < x_2$ . We let  $\beta \in ]0,1[$  such that  $q_1 = \beta \cdot x_1 + (1-\beta)x_2$ , which exists because  $q_1 \in ]x_1,x_2[$ . If **p** is not comparable to **q**, there is nothing to show. We assume that these two vectors are comparable and show that  $\mathbf{p} < \mathbf{q}$ . We proceed by contradiction and assume that  $\mathbf{p} \geq \mathbf{q}$ .

We consider the linear form  $x^* \colon \mathbb{R}^2 \to \mathbb{R}$  defined by  $x^*(\mathbf{v}) = -\frac{\mathcal{F}(x_2) - \mathcal{F}(x_1)}{x_2 - x_1} \cdot v_1 + v_2$  for all  $\mathbf{v} = (v_1, v_2) \in \mathbb{R}^2$ . We have  $x^*(\mathbf{p}_1) = x^*(\mathbf{p}_2)$  and thus  $(x^*)^{-1}(x^*(\mathbf{p}_1))$  is the line carrying the segment  $[\mathbf{p}_1, \mathbf{p}_2]$ . Because  $\mathcal{F}$  is decreasing (Lemma 4.7), we have  $\frac{\mathcal{F}(x_2) - \mathcal{F}(x_1)}{x_2 - x_1} < 0$ . This implies that  $x^*$  is increasing in the sense that for all  $\mathbf{v}_1, \mathbf{v}_2 \in \mathbb{R}^2$ ,  $\mathbf{v}_1 < \mathbf{v}_2$  implies that  $x^*(\mathbf{v}_1) < x^*(\mathbf{v}_2)$ .

Let  $\mathbf{v} = (q_1, \beta \cdot \mathcal{F}(x_1) + (1 - \beta)\mathcal{F}(x_2))$ . We have  $\mathbf{v} < \mathbf{q}$  by strict concavity of  $\mathcal{F}$  (Lemma 4.7). Furthermore, we have  $x^*(\mathbf{v}) = x^*(\mathbf{p})$  because both of these vectors are in the segment  $[\mathbf{p}_1, \mathbf{p}_2]$ . We obtain, because  $x^*$  is increasing and  $\mathbf{v} < \mathbf{q} < \mathbf{p}$ , that  $x^*(\mathbf{v}) < x^*(\mathbf{q}) < x^*(\mathbf{p}) = x^*(\mathbf{v})$ . This is a contradiction. This ends the proof in the case that  $\mathbf{p}_1 \neq (0, 2)$  and  $\mathbf{p}_2 \neq (0, 2)$ .

Next, we assume that  $\mathbf{p}_1 = \mathbf{p}_2 = (0,2)$ . We obtain that  $\mathbf{p} = (0,2)$ , which is smaller than (1,2) and incomparable to all elements of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{(1,2),(0,2)\}$ . Finally, we assume that only one of  $\mathbf{p}_1$  or  $\mathbf{p}_2$  are (0,2). We assume without loss of generality that

 $\mathbf{p}_1 = (0,2)$ . If  $\mathbf{p} > \mathbf{q}$ , then we would have  $\alpha \cdot (1,2) + (1-\alpha) \cdot \mathbf{p}_2 > \mathbf{p} > \mathbf{q}$ , which would contradict the first part of the proof.

With Lemma 4.6 and Lemma 4.9, we directly obtain that all elements of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{(0,2)\}$  are Pareto-optimal elements of  $\mathsf{Pay}_{s_0}(\bar{f})$ : if they were not Pareto-optimal, then they would be dominated by an element of the border of  $\mathsf{Pay}_{s_0}(\bar{f})$  which would be the convex combination of two vectors in  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$ .

**Lemma 4.10.** Let  $\mathbf{q} \in \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{(0,2)\}$ . Then  $\mathbf{q}$  is a Pareto-optimal element of  $\mathsf{Pay}_{s_0}(\bar{f})$ .

*Proof.* Assume towards a contradiction that there exists  $\mathbf{p} \in \mathsf{Pay}_{s_0}(\bar{f})$  such that  $\mathbf{q} < \mathbf{p}$ . We show that there exists  $\mathbf{p}' \in \mathsf{bd}(\mathsf{Pay}_{s_0}(\bar{f}))$  such that  $\mathbf{p}' > \mathbf{q}$ . This yields a contradiction with Lemma 4.6 and Lemma 4.9.

If  $\mathbf{p} \in \mathsf{bd}(\mathsf{Pay}_{s_0}(\bar{f}))$ , then we let  $\mathbf{p}' = \mathbf{p}$ . We thus assume that  $\mathbf{p} \in \mathsf{int}(\mathsf{Pay}_{s_0}(\bar{f}))$ . Let  $\alpha = \sup\{\beta \geq 0 \mid \mathbf{p} + \beta \mathbf{1} \in \mathsf{Pay}_{s_0}(\bar{f})\}$ . We have  $\alpha \in \mathbb{R}$  because  $\mathbf{p} \in \mathsf{Pay}_{s_0}(\bar{f})$  (i.e.,  $0 \in \{\beta \geq 0 \mid \mathbf{p} + \beta \mathbf{1} \in \mathsf{Pay}_{s_0}(\bar{f})\}$  thus  $\alpha > -\infty$ ) and  $\mathsf{Pay}_{s_0}(\bar{f})$  is bounded (thus  $\alpha < +\infty$ ). Furthermore, because  $\mathsf{Pay}_{s_0}(\bar{f})$  is closed, we have  $\mathbf{p} + \alpha \mathbf{1} \in \mathsf{bd}(\mathsf{Pay}_{s_0}(\bar{f}))$ . We obtain the announced contradiction by letting  $\mathbf{p}' = \mathbf{p} + \alpha \mathbf{1} > \mathbf{q}$ .

We conclude this section by showing that the set of extreme points of  $\mathsf{Pay}_{s_0}(\bar{f})$  is the set of pure payoffs.

**Lemma 4.11.** We have  $\mathsf{extr}(\mathsf{Pay}_{s_0}(\bar{f})) = \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$ 

Proof. First, we show that  $\operatorname{extr}(\mathsf{Pay}_{s_0}(\bar{f})) \subseteq \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$ . It suffices to show that all vectors  $\mathbf{q} \in \mathsf{Pay}_{s_0}(\bar{f}) \setminus \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$  are not extreme points of  $\mathsf{Pay}_{s_0}(\bar{f})$ . Let  $\mathbf{q} \in \mathsf{Pay}_{s_0}(\bar{f}) \setminus \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$ . By Lemma 4.4, we have  $\mathbf{q} \in \mathsf{Pay}_{s_0}(\bar{f}) = \operatorname{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}))$ . This implies that  $\mathbf{q} \in \operatorname{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{\mathbf{q}\})$  because  $\mathbf{q} \notin \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$ , and thus  $\mathbf{q} \notin \operatorname{extr}(\mathsf{Pay}_{s_0}(\bar{f}))$ .

conv(Pay<sub>s0</sub>( $\bar{f}$ ) \ {**q**}) because  $\mathbf{q} \notin \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$ , and thus  $\mathbf{q} \notin \mathsf{extr}(\mathsf{Pay}_{s_0}(\bar{f}))$ . Conversely, let  $\mathbf{q} \in \mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$ . First, we assume that  $\mathbf{q} = (0,2)$ . The first coordinate of all other vectors of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$  is greater than or equal to 1. This therefore also applies to any convex combination of vectors in  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{(0,2)\}$ . It follows that  $\mathbf{q} \in \mathsf{extr}(\mathsf{Pay}_{s_0}(\bar{f}))$ . Second, we assume that  $\mathbf{q} \neq (0,2)$ . Assume towards a contradiction that  $\mathbf{q} \notin \mathsf{extr}(\mathsf{Pay}_{s_0}(\bar{f}))$ . By Lemma 4.10,  $\mathbf{q}$  is a Pareto-optimal element of  $\mathsf{Pay}_{s_0}(\bar{f})$ , and thus lies on the border of this set. We obtain that  $\mathbf{q}$  is the convex combination of two elements of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{\mathbf{q}\}$  by Lemma 4.6 (which is applicable here by Lemmas 4.4 and 4.2) and the assumption that  $\mathbf{q} \notin \mathsf{extr}(\mathsf{Pay}_{s_0}(\bar{f}))$ . This yields a contradiction with Lemma 4.9, which states that there is no convex combination of two vectors of  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{\mathbf{q}\}$  that is greater than or equal to  $\mathbf{q}$ .

## 4.2.5 Memory cannot be traded for randomness

We prove that the only way to obtain an expected payoff in  $\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f})$  is through a strategy that induces a single play from  $s_0$  (i.e., intuitively, a strategy that is pure in practice).

**Lemma 4.12.** Let  $\sigma \in \Sigma(\mathfrak{M})$  be a strategy such that at least two plays starting in  $s_0$  are consistent with  $\sigma$ . Then  $\mathbb{E}_{s_0}^{\sigma}(\bar{f}) \notin \mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$ .

Proof. We consider the following enumeration of the set of plays of  $\mathfrak{M}$  that start in  $s_0$ . We let  $\pi_{-2} = s_0 c(s_1 a)^{\omega}$ ,  $\pi_{-1} = s_0 a(s_2 a)^{\omega}$ , and, for all  $r \in \mathbb{N}$ , we let  $\pi_r = s_0 (as_2)^r b(s_3 a)^{\omega}$ . We have  $\bar{f}(\pi_{-2}) = (0,2)$ ,  $\bar{f}(\pi_{-1}) = (1,2)$  and, for all  $r \in \mathbb{N}$ ,  $\bar{f}(\pi_r) = \left(1 + \frac{3^r}{4^{r-1}}, 2 - \frac{1}{2^{r-1}}\right)$  (refer to the proof of Lemma 4.1 for the relevant computations). Due to the absence of

randomised transitions in  $\mathfrak{M}$ ,  $\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f})$  is the set of payoffs of the plays starting in  $s_0$ . Therefore, to end the proof, we must show that for all  $r \geq -2$ ,  $\mathbb{E}^{\sigma}_{s_0}(\bar{f}) \neq \bar{f}(\pi_r)$ .

For all  $r \in \{-1, -2\} \cup \mathbb{N}$ , let  $\alpha_r = \mathbb{P}^{\sigma}_{s_0}(\{\pi_r\})$ . Through this notation, we obtain that

$$\mathbb{E}_{s_0}^{\sigma}(\bar{f}) = \sum_{r > -2} \alpha_r \bar{f}(\pi_r). \tag{4.2}$$

First, we show that  $\mathbb{E}^{\sigma}_{s_0}(\bar{f}) \notin \{(0,2),(1,2)\}$ . If there exists  $r \in \mathbb{N}$  such that  $\alpha_r > 0$ , then we have  $\mathbb{E}^{\sigma}_{s_0}(f_2) < 2$ , which implies that  $\mathbb{E}^{\sigma}_{s_0}(\bar{f}) \notin \{(0,2),(1,2)\}$ . Next, assume that  $\alpha_r = 0$  for all  $r \in \mathbb{N}$ . Then  $\alpha_{-2}$  and  $\alpha_{-1}$  must sum to one. Furthermore, since  $\sigma$  has at least two outcomes, both  $\alpha_{-2}$  and  $\alpha_{-1}$  must be non-zero. It follows that  $\mathbb{E}^{\sigma}_{s_0}(f_1) \in ]0,1[$ . We conclude that  $\mathbb{E}^{\sigma}_{s_0}(\bar{f}) \notin \{(0,2),(1,2)\}$  in this case as well.

We now fix  $r \in \mathbb{N}$  and show that  $\mathbb{E}^{\sigma}_{s_0}(\bar{f}) \neq \bar{f}(\pi_r)$ . We proceed by contradiction. Assume towards a contradiction that  $\mathbb{E}^{\sigma}_{s_0}(\bar{f}) = \bar{f}(\pi_r)$ . Our goal is to contradict Lemma 4.11, i.e., to show that  $\bar{f}(\pi_r)$  is not an extreme point of  $\mathsf{Pay}_{s_0}(\bar{f})$ . We do so in two steps. First, we show that  $\mathbb{E}^{\sigma}_{s_0} \in \mathsf{cl}(\mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{\bar{f}(\pi_r)\}))$ . Next, we prove that  $\mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{\bar{f}(\pi_r)\})$  is closed. Together, these statements imply that  $\bar{f}(\pi_r) \in \mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s_0}(\bar{f}) \setminus \{\bar{f}(\pi_r)\})$ , which is the sought contradiction.

We now establish that  $\mathbb{E}_{s_0}^{\sigma} \in \text{cl}(\text{conv}(\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}) \setminus \{\bar{f}(\pi_r)\}))$ . It follows from  $\mathbb{E}_{s_0}^{\sigma}(\bar{f}) = \bar{f}(\pi_r)$  and Equation (4.2) that  $\mathbb{E}_{s_0}^{\sigma}(\bar{f}) = \sum_{\ell \neq r} \frac{\alpha_\ell}{1 - \alpha_r} \bar{f}(\pi_\ell)$ . By an argument analogous to the one in the proof of Lemma 4.4, we conclude from this last equality that  $\mathbb{E}_{s_0}^{\sigma}(\bar{f}) \in \text{cl}(\text{conv}(\mathsf{Pay}_{s_0}^{\mathsf{pure}}(\bar{f}) \setminus \{\bar{f}(\pi_r)\}))$ .

It remains to show that  $\operatorname{conv}(\operatorname{Pay}_{s_0}^{\operatorname{pure}}(\bar{f})\setminus\{\bar{f}(\pi_r)\})$  is closed. The vector  $\bar{f}(\pi_r)$  is an isolated point of  $\operatorname{Pay}_{s_0}^{\operatorname{pure}}(\bar{f})$ : all other elements of  $\operatorname{Pay}_{s_0}^{\operatorname{pure}}(\bar{f})$  are at distance at least  $2^{-r}$  of  $\bar{f}(\pi_r)$ . Therefore,  $\operatorname{Pay}_{s_0}^{\operatorname{pure}}(\bar{f})\setminus\{\bar{f}(\pi_r)\}$  is closed (when removing an isolated point from a closed set, the resulting set is still closed), and thus  $\operatorname{conv}(\operatorname{Pay}_{s_0}^{\operatorname{pure}}(\bar{f})\setminus\{\bar{f}(\pi_r)\})$  is closed by Lemma 4.3.

We close this section by commenting on the significance of Lemma 4.12 in terms of strategy complexity with respect to memory requirements. When we only consider pure strategies, some Pareto-optimal payoffs may require strategies with an arbitrarily large memory, in the sense that we may need to count to some high (but finite) counter value to enact a given number of loops in  $s_2$  to obtain certain expected payoffs. Lemma 4.12 implies that we cannot substitute this counting by randomisation, even when only considering Pareto-optimal payoffs. In particular, for this example, we obtain that although all expected payoffs can be obtained with strategies that only count up to a finite number of steps (i.e., to count loops in  $s_2$ ), we cannot bound this number of steps uniformly for all expected payoff vectors.

## 5 Lexicographic Markov decision processes

In this section, we consider lexicographic optimisation of multiple payoff functions in MDPs. In Section 5.1, we assert that for universally integrable payoffs, randomness does not provide additional power for lexicographic optimisation. We show that this property does not generalise to the whole class of universally unambiguously integrable payoffs in Section 5.2, i.e., randomness may be required for lexicographically optimal strategies in general.

For this section, we fix an MDP  $\mathfrak{M}=(S,A,\delta)$  and a d-dimensional payoff function  $\bar{f}=(f_j)_{1\leq j\leq d}$ .

## 5.1 Universally integrable payoffs

Assume that  $\bar{f}$  is universally integrable. We show that for all initial states  $s \in S$  and all strategies  $\sigma$ , there exists a pure strategy  $\tau$  such that  $\mathbb{E}^{\sigma}_{s}(\bar{f}) \leq_{\mathsf{lex}} \mathbb{E}^{\tau}_{s}(\bar{f})$ . The crux of the proof is showing that the Lebesgue integral is compatible with the lexicographic order over  $\bar{\mathbb{R}}^d$ . Once this is shown, we assume towards a contradiction that there is no suitable pure strategy  $\tau$ . By Lemma 3.1, we can write  $\mathbb{E}^{\sigma}_{s}(\bar{f})$  as an integral over pure expected payoffs. We then reach the contradiction that  $\mathbb{E}^{\sigma}_{s}(\bar{f}) <_{\mathsf{lex}} \mathbb{E}^{\sigma}_{s}(\bar{f})$ . We formalise this argument below.

**Theorem 5.1.** Assume that  $\bar{f}$  is universally integrable. Let  $\sigma$  be a strategy and  $s \in S$ . There exists a pure strategy  $\tau$  such that  $\mathbb{E}_s^{\sigma}(\bar{f}) \leq_{\mathsf{lex}} \mathbb{E}_s^{\tau}(\bar{f})$ .

Proof. To prove the theorem, we reason on random variables of  $([0,1]^{\omega}, \mathcal{A}_{[0,1]^{\omega}})$ . We consider the distribution  $\nu_{\sigma} \in \mathcal{D}([0,1]^{\omega}, \mathcal{A}_{[0,1]^{\omega}})$  and the mapping  $[0,1]^{\omega} \to \Sigma_{\mathsf{pure}}(\mathfrak{M}) \colon x \mapsto \tau_x$  such that  $\mathbb{E}_s^{\sigma}(\bar{f}) = \int_{x \in [0,1]^{\omega}} \mathbb{E}_s^{\tau_x}(\bar{f}) \mathrm{d}\nu_{\sigma}(x)$  given by Lemma 3.1. For all  $1 \leq j \leq d$ , we consider the real random variable  $X_j \colon x \mapsto \mathbb{E}_s^{\tau_x}(f_j)$  over  $[0,1]^{\omega}$ . We let  $\mathcal{X} = (X_1, \dots, X_d)$ . By Lemma 3.2,  $\mathcal{X}$  is bounded and thus integrable. For any real or multivariate random variable Y over  $[0,1]^{\omega}$ , we write  $\int Y \mathrm{d}\nu_{\sigma}$  for  $\int_{x \in [0,1]^{\omega}} Y(x) \mathrm{d}\nu_{\sigma}(x)$  to lighten notation.

Let  $\mathcal{Y} = (Y_1, \dots, Y_d)$  be an integrable multi-variate real random variable over  $[0, 1]^{\omega}$ . We first show that if  $\mathcal{X} <_{\text{lex}} \mathcal{Y}$ , then  $\mathbb{E}_s^{\sigma}(\bar{f}) = \int \mathcal{X} d\nu_{\sigma} <_{\text{lex}} \int \mathcal{Y} d\nu_{\sigma}$ . We use this claim below to prove the theorem.

Assume that  $\mathcal{X} <_{\mathsf{lex}} \mathcal{Y}$ . We partition  $[0,1]^\omega$  as follows. For all  $1 \leq j \leq d$ , we let  $E_j = \{x \in [0,1]^\omega \mid X_j(x) < Y_j(x) \text{ and } \forall j' < j, X_{j'}(x) = Y_{j'}(x)\}$ . Intuitively,  $E_j$  is the set of elements such that the strict lexicographic ordering of their respective images by  $\mathcal{X}$  and  $\mathcal{Y}$  is witnessed in component j. The sets  $E_1, \ldots, E_d$  partition  $[0,1]^\omega$  because  $\mathcal{X} <_{\mathsf{lex}} \mathcal{Y}$ . It follows that, for  $\mathcal{Z} \in \{\mathcal{X}, \mathcal{Y}\}$ , we have  $\int \mathcal{Z} d\nu_\sigma = \sum_{j=1}^d \int \mathcal{Z} \cdot \mathbbm{1}_{E_j} d\nu_\sigma$ .

Let  $j^* = \min\{1 \leq j \leq d \mid \nu_{\sigma}(E_j) > 0\}$ . To obtain that  $\int \mathcal{X} d\nu_{\sigma} <_{\text{lex}} \int \mathcal{Y} d\nu_{\sigma}$ , we show that, for  $j < j^*$ , we have  $\int X_j d\nu_{\sigma} = \int Y_j d\nu_{\sigma}$  and  $\int X_{j^*} d\nu_{\sigma} < \int Y_{j^*} d\nu_{\sigma}$ . To prove these relations, we formulate two observations. First, we observe that for all  $1 \leq j < j' \leq d$ ,  $X_j$  and  $Y_j$  agree over  $E_{j'}$  (by definition), and thus we have

$$\int X_j \cdot \mathbb{1}_{E_{j'}} d\nu_{\sigma} = \int Y_j \cdot \mathbb{1}_{E_{j'}} d\nu_{\sigma}.$$
 (5.1)

Second, since  $\nu_{\sigma}(E_{j'}) = 0$  for all  $j' < j^*$ , it follows that for all  $1 \leq j \leq d$  and all  $Z_j \in \{X_j, Y_j\}$  that

$$\int Z_j d\nu_{\sigma} = \sum_{j'=j^*}^d \int Z_j \cdot \mathbb{1}_{E_{j'}} d\nu_{\sigma}.$$
(5.2)

Let  $1 \le j < j^*$ . By combining Equations (5.1) and (5.2), we obtain that

$$\int X_j d\nu_{\sigma} = \sum_{j'=j^*}^d \int X_j \cdot \mathbb{1}_{E_{j'}} d\nu_{\sigma} = \sum_{j'=j^*}^d \int Y_j \cdot \mathbb{1}_{E_{j'}} d\nu_{\sigma} = \int Y_j d\nu_{\sigma}.$$

To end the proof that  $\int \mathcal{X} d\nu_{\sigma} <_{\text{lex}} \int \mathcal{Y} d\nu_{\sigma}$ , it remains to show that  $\int X_{j^{\star}} d\nu_{\sigma} < \int Y_{j^{\star}} d\nu_{\sigma}$ . This inequality is equivalent to  $\int X_{j^{\star}} \cdot \mathbb{1}_{E_{j^{\star}}} d\nu_{\sigma} < \int Y_{j^{\star}} \cdot \mathbb{1}_{E_{j^{\star}}} d\nu_{\sigma}$  by Equations (5.1) and (5.2). This inequality holds by compatibility of the Lebesgue integral with the order of  $\mathbb{R}$  (recall that  $X_{j^{\star}}(x) < Y_{j^{\star}}(x)$  for all  $x \in E_{j^{\star}}$ ) and because  $\nu_{\sigma}(E_{j^{\star}}) > 0$ . We have shown that  $\int \mathcal{X} d\nu_{\sigma} <_{\text{lex}} \int \mathcal{Y} d\nu_{\sigma}$  whenever  $\mathcal{X} <_{\text{lex}} \mathcal{Y}$  holds.




Figure 5.1: An MDP with deterministic transitions. The doubly circled state denotes a target for a reachability objective.

Now assume that for all pure strategies  $\tau$ , we have  $\mathbb{E}_s^{\tau}(\bar{f}) <_{\mathsf{lex}} \mathbb{E}_s^{\sigma}(\bar{f})$  towards a contradiction. It follows that  $\mathcal{X}(x) <_{\mathsf{lex}} \mathbb{E}_s^{\sigma}(\bar{f})$  for all  $x \in [0,1]^{\omega}$ . From the above, we obtain that  $\mathbb{E}_s^{\sigma}(\bar{f}) = \int \mathcal{X} d\nu_{\sigma} <_{\mathsf{lex}} \mathbb{E}_s^{\sigma}(\bar{f})$ , which is a contradiction.

A direct corollary of Theorem 5.1 is the following.

Corollary 5.2. Assume that  $\bar{f}$  is universally integrable. For all  $s \in S$ , if there exists a lexicographically optimal strategy from s, then there exists a pure lexicographically optimal strategy.

## 5.2 General payoffs

In this section, we present an example that shows that the conclusion of Theorem 5.1 does not hold for all universally unambiguously integrable payoffs.

**Example 5.1.** We consider the MDP  $\mathfrak{M}$  depicted in Figure 5.1. Let w denote the illustrated weight function. We consider the two-dimensional payoff function  $\bar{f} = (f_1, f_2)$  such that  $f_1 = \mathbb{1}_{\mathsf{Reach}(\{t\})}$  and  $f_2 = \mathsf{TRew}_w$ . We prove that randomisation is necessary to play lexicographically optimally from s.

Since transitions of  $\mathfrak{M}$  are deterministic,  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$  is the set of payoffs of plays from s. We introduce notation for these plays: for all  $r \in \mathbb{N}$ , let  $\pi_r = (sa)^r s(bt)^\omega$  and let  $\pi_\infty = (sa)^\omega$ . It follows that  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}) = \{\bar{f}(\pi_r) \mid r \in \mathbb{N} \cup \{\infty\}\} = \{(1,r) \mid r \in \mathbb{N}\} \cup \{(0,+\infty)\}$ . In particular, no pure strategy has an expected payoff of  $(1,+\infty)$ , which is the greatest that could occur with  $\bar{f}$ .

However, there exists a randomised strategy whose expected payoff from s is  $(1, +\infty)$ . To show this, we construct a strategy  $\sigma$  such that, for all  $r \in \mathbb{N}$ , the play  $\pi_{2^r}$ , whose payoff is  $\bar{f}(\pi_r) = (1, 2^r)$ , occurs with probability  $\frac{1}{2^{r+1}}$ . We set  $\sigma(h)(a) = 1$  for all histories such that  $h \notin \{(sa)^{2^r}s \mid r \in \mathbb{N}\}$  and, for all  $r \in \mathbb{N}$ , we set  $\sigma((sa)^{2^r}s)(a) = \sigma((sa)^{2^r}s)(b) = \frac{1}{2}$ . For all  $r \in \mathbb{N}$ , we obtain that  $\mathbb{P}_s^{\sigma}(\{\pi_{2^r}\}) = \mathbb{P}_s^{\sigma}(\mathsf{Cyl}((sa)^{2^r}sbt)) = \frac{1}{2^{r+1}}$  due to the structure of  $\mathfrak{M}$ . It follows that  $\mathbb{E}_s^{\sigma}(\bar{f}) = \sum_{r=0}^{\infty} \mathbb{P}_s^{\sigma}(\{\pi_{2^r}\}) \cdot \bar{f}(\pi_{2^r}) = (1,\infty)$ . This shows that  $\sigma$  is lexicographically optimal, and ends the proof that randomisation is required to play lexicographically optimally from s.

## 6 The structure of expected payoff sets

In this section, we study the relationship between the set of expected payoffs of pure strategies and of general strategies. In Section 6.1, we prove that for universally integrable payoffs, the set of expected payoffs (for general strategies) is the convex hull of the set of expected payoffs of pure strategies. In other words, mixed strategies with a finite support can match the expected payoff of any strategy when considering universally integrable payoffs. We provide a variant of this result in Section 6.2 for the more general universally unambiguously integrable payoffs: we show that any expected payoff can be approximated (in the sense of the topology of  $\mathbb{R}$ ) by the expected payoff of a finite-support mixed

strategy. Finally, in Section 6.3, we provide bounds on support sizes of finite-support mixed strategies analogous to Carathéodory's theorem for convex hulls (Theorem 2.1).

We fix, for this whole section, an MDP  $\mathfrak{M}=(S,A,\delta),\ d\in\mathbb{N}_0$  and a d-dimensional payoff function  $\bar{f}=(f_j)_{1\leq j\leq d}$ .

## 6.1 Universally integrable payoffs

Throughout this section, we assume that  $\bar{f}$  is universally integrable. Our goal is to show that, for all  $s \in S$ ,  $\mathsf{Pay}_s(\bar{f}) = \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}))$ . We first provide an overview of the proof, and then provide a formal proof.

**Proof overview.** Let  $s \in S$ . By convexity of  $\mathsf{Pay}_s(\bar{f})$ , the inclusion  $\mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})) \subseteq \mathsf{Pay}_s(\bar{f})$  holds. Therefore, the main difficulty of the proof is to prove the other inclusion. Let  $\mathbf{q} \in \mathsf{Pay}_s(\bar{f})$ . The first step of the proof is to construct a linear map  $L_{\mathbf{q}} \colon \mathbb{R}^d \to \mathbb{R}^{d'}$  with  $d' \leq d$  such that (i)  $L_{\mathbf{q}}(\mathbf{q})$  is the lexicographic maximum of  $L_{\mathbf{q}}(\mathsf{Pay}_s(\bar{f}))$  and (ii)  $\mathbf{q} \in \mathsf{ri}(\mathsf{Pay}_s(\bar{f}) \cap V)$  where  $V = L_{\mathbf{q}}^{-1}(L_{\mathbf{q}}(\mathbf{q}))$  denotes the set of vectors that share their image by  $L_{\mathbf{q}}$  with  $\mathbf{q}$ .

The mapping  $L_{\mathbf{q}}$  is constructed as follows (Examples 6.1 and 6.2 below illustrate this construction). If  $\mathbf{q}$  is in the relative interior of  $\mathsf{Pay}_s(\bar{f})$ , we let  $L_{\mathbf{q}}$  be the zero-valued linear form. Otherwise, by the supporting hyperplane theorem (Theorem 2.3), there exists a linear form  $x_1^*$  such that  $x_1^*(\mathbf{q}) \geq x_1^*(\mathbf{p})$  for all  $\mathbf{p} \in \mathsf{Pay}_s(\bar{f})$ . Let  $H_1 = (x_1^*)^{-1}(x_1^*(\mathbf{q}))$  denote the supporting hyperplane given by  $x_1^*$ . We check whether  $\mathbf{q}$  is in the relative interior of  $\mathsf{Pay}_s(\mathbf{q}) \cap H_1$ . If it is the case, we obtain the desired linear mapping by letting  $L_{\mathbf{q}} = x_1^*$ . Otherwise, we continue: there is a linear form  $x_2^*$  describing a hyperplane  $H_2$  that supports  $\mathsf{Pay}_s(\mathbf{q}) \cap H_1$  at  $\mathbf{q}$ . We choose  $x_2^*$  such that  $x_1^*$  and  $x_2^*$  have distinct kernels, i.e.,  $H_1 \neq H_2$ . To ensure that the kernels are distinct, we construct  $x_2^*$  as an extension to  $\mathbb{R}^d$  of a non-zero linear form over  $\mathsf{ker}(x_1^*)$ . By induction, we continue constructing linear forms with pairwise distinct kernels (i.e., defining pairwise distinct supporting hyperplanes) until  $\mathbf{q} \in \mathsf{ri}(\mathsf{Pay}_s(\bar{f}) \cap \bigcap_{1 \leq j \leq d'} H_j)$ . When this condition is satisfied, we define, for all  $\mathbf{v} \in \mathbb{R}^d$ ,  $L_{\mathbf{q}}(\mathbf{v}) = (x_1^*(\mathbf{v}), \dots, x_{d'}^*(\mathbf{v}))$ . The invocation of the supporting hyperplane theorem at each iteration guarantees that  $L_{\mathbf{q}}(\mathbf{q})$  is a lexicographic maximum of  $L_{\mathbf{q}}(\mathsf{Pay}_s(\bar{f}))$ . We remark that, when the stopping condition is fulfilled, the supporting hyperplane theorem is no longer applicable.

Let  $V=L_{\mathbf{q}}^{-1}(L_{\mathbf{q}}(\mathbf{q}))$ . By construction,  $\mathbf{q}\in \mathrm{ri}(V\cap \mathsf{Pay}_s(\bar{f}))$ . To conclude, we establish that  $\mathrm{ri}(V\cap \mathsf{Pay}_s(\bar{f}))=\mathrm{ri}(V\cap \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})))$ . This suffices because the latter set is a subset of  $\mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}))$ . This is equivalent to showing that  $\mathsf{cl}(V\cap \mathsf{Pay}_s(\bar{f}))=\mathsf{cl}(V\cap \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})))$ : convex subsets of  $\mathbb{R}^d$  have the same relative interior as their closure [Roc70, Thm. 6.3].

The inclusion  $\operatorname{conv}(\operatorname{Pay}_s^{\operatorname{pure}}(\bar{f})) \subseteq \operatorname{Pay}_s(\bar{f})$  implies that  $\operatorname{cl}(V \cap \operatorname{conv}(\operatorname{Pay}_s^{\operatorname{pure}}(\bar{f}))) \subseteq \operatorname{cl}(V \cap \operatorname{Pay}_s(\bar{f}))$ . For the other inclusion, we need only show that  $V \cap \operatorname{Pay}_s(\bar{f}) \subseteq \operatorname{cl}(V \cap \operatorname{conv}(\operatorname{Pay}_s^{\operatorname{pure}}(\bar{f})))$ . We assume towards a contradiction that this is not the case and let  $\mathbf{p} \in V \cap \operatorname{Pay}_s(\bar{f}) \setminus \operatorname{cl}(V \cap \operatorname{conv}(\operatorname{Pay}_s^{\operatorname{pure}}(\bar{f})))$ . Using the hyperplane separation theorem (Theorem 2.2), we obtain  $x_\star^*$  such that  $x_\star^*(\mathbf{p}) > x_\star^*(\mathbf{v})$  for all  $\mathbf{v} \in \operatorname{cl}(V \cap \operatorname{conv}(\operatorname{Pay}_s^{\operatorname{pure}}(\bar{f})))$ . Because  $\mathbf{p} \in \operatorname{Pay}_s(\bar{f})$ , Theorem 5.1 implies that there exists a pure strategy  $\sigma$  such that  $(L_{\mathbf{q}}(\mathbf{p}), x_\star^*(\mathbf{p})) \leq_{\operatorname{lex}} (L_{\mathbf{q}}(\mathbb{E}_s^{\sigma}(\bar{f})), x_\star^*(\mathbb{E}_s^{\sigma}(\bar{f})))$ . Furthermore,  $\mathbf{p} \in V$  implies that  $L_{\mathbf{q}}(\mathbf{p})$  is the lexicographic maximum of  $L_{\mathbf{q}}(\operatorname{Pay}_s(\bar{f}))$ , and thus so is  $L_{\mathbf{q}}(\mathbb{E}_s^{\sigma}(\bar{f}))$ , i.e.,  $\mathbb{E}_s^{\sigma}(\bar{f}) \in V \cap \operatorname{Pay}_s^{\operatorname{pure}}(\bar{f})$ . It follows that  $x_\star^*(\mathbf{p}) \leq x_\star^*(\mathbb{E}_s^{\sigma}(\bar{f}))$ , which is contradictory. This closes the argument that  $\operatorname{cl}(V \cap \operatorname{Pay}_s(\bar{f})) = \operatorname{cl}(V \cap \operatorname{conv}(\operatorname{Pay}_s^{\operatorname{pure}}(\bar{f}))$ , which implies that  $\mathbf{q} \in \operatorname{conv}(\operatorname{Pay}_s^{\operatorname{pure}}(\bar{f}))$ , ending the sketch.







Figure 6.1: An MDP. The doubly circled and filled states respectively highlight the targets of the reachability objectives  $Reach(\{s_1, s_4\})$  and  $Reach(\{s_2, s_4\})$ .







(b) The image of the set on the left by the linear mapping  $L_{\mathbf{q}}$ :  $(v_1, v_2) \mapsto (v_1 + 3v_2, 6v_1 - 2v_2)$  obtained through the equations of the hyperplanes on the left. Remark that the image of  $\mathbf{q}$  is lexicographically optimal.

Figure 6.2: The set of expected payoffs for Example 6.1 and its image by a linear mapping.

We complement the sketch above with two examples that illustrate the construction of the mapping  $L_{\bf q}$ . In the first example, we select  ${\bf q}$  as an extreme point of the set of expected payoffs. In this case, the constructed linear mapping  $L_{\bf q}$  is such that  ${\bf q}$  is the unique vector  ${\bf p} \in \mathbb{R}^d$  such that  $L_{\bf q}({\bf p})$  is the lexicographic maximum of  $L_{\bf q}(\mathsf{Pay}_s(\bar f))$ . In our second example,  $\mathsf{Pay}_s(\bar f)$  is not closed and we choose  ${\bf q}$  as a non-extreme point such that no pure strategy has expected payoff  ${\bf q}$ . In this case, we observe that the uniqueness property of the first example cannot be obtained no matter the constructed  $L_{\bf q}$ .

**Example 6.1** (Extreme point). We consider the MDP depicted in Figure 6.1. Throughout this example, we (implicitly) consider  $s_0$  as the initial state. We study indicators of reachability objectives: we let  $\bar{f} = (\mathbbm{1}_{\mathsf{Reach}(T_1)}, \mathbbm{1}_{\mathsf{Reach}(T_2)})$  where  $T_1 = \{s_1, s_4\}$  and  $T_2 = \{s_2, s_4\}$ . The set  $\mathsf{Pay}_{s_0}(\bar{f})$  is depicted in Figure 6.2: it is the convex hull of the expected payoffs of the pure strategies that play one of the actions a, b or c in  $s_0$ . We focus on the extreme point  $\mathbf{q} = (\frac{3}{4}, \frac{3}{4})$  in the remainder of this example.

The construction of  $L_{\mathbf{q}}$  is in two steps. First, we consider the linear form  $x_1^*$  defined by, for all  $\mathbf{v} \in \mathbb{R}^d$ ,  $x_1^*(\mathbf{v}) = v_1 + 3v_2$ . The hyperplane  $H = (x_1^*)^{-1}(3)$  supports  $\mathsf{Pay}_{s_0}(\bar{f})$  at  $\mathbf{q}$  and is depicted by the blue dashed line in Figure 6.2a. It is not satisfactory to set  $L_{\mathbf{q}} = x_1^*$ :  $\mathbf{q}$  is an endpoint of the segment  $[(0,1),\mathbf{q}] = \mathsf{Pay}_{s_0}(\bar{f}) \cap H$ , and is therefore not in its relative interior.

We recall that for any linear form  $y^*$  of  $\ker(x_1^*)$  (i.e., the vector space corresponding to H), there exists  $\mathbf{v} \in \ker(x_1^*)$  such that  $y^*(\mathbf{w}) = \langle \mathbf{w}, \mathbf{v} \rangle$  for all  $\mathbf{w} \in \ker(x_1^*)$ . Since any non-zero linear form of  $\ker(x_1^*)$  is bijective, all of them induce a hyperplane of H




Figure 6.3: An MDP with deterministic transitions. The pairs beneath actions represent a two-dimensional weight function. State t is doubly circled because it is a target.

supporting  $\mathsf{Pay}_s(\bar{f}) \cap H$  at  $\mathbf{q}$ . We proceed with the linear form  $x_2^* \colon \mathbb{R}^2 \to \mathbb{R}$  defined by  $x_2^*(\mathbf{v}) = 6v_1 - 2v_2$  for all  $\mathbf{v} = (v_1, v_2) \in \mathbb{R}^2$  (derived from the vector  $\mathbf{w} = (6, -2) \in \mathsf{ker}(x_1^*)$ ). Observe that  $\mathsf{ker}(x_1^*) \cap \mathsf{ker}(x_2^*) = \{\mathbf{0}\}$ .

We define  $L_{\mathbf{q}}(\mathbf{v})=(x_1^*(\bar{\mathbf{v}}),x_2^*(\bar{\mathbf{v}}))$ . Since  $L_{\mathbf{q}}$  is bijective,  $L_{\mathbf{q}}^{-1}(L_{\mathbf{q}}(\mathbf{q}))$  is a singleton set. Therefore,  $\mathbf{q}$  is in the relative interior of  $L_{\mathbf{q}}^{-1}(L_{\mathbf{q}}(\mathbf{q}))\cap \mathsf{Pay}_{s_0}(\bar{f})$ . By linearity of the expectation, it holds that  $\mathsf{Pay}_{s_0}(L_{\mathbf{q}}\circ\bar{f})=L_{\mathbf{q}}(\mathsf{Pay}_{s_0}(\bar{f}))$ . This set is illustrated in Figure 6.2b; it is easy to check that  $L_{\mathbf{q}}(\mathbf{q})$  is the lexicographic maximum of this set.

In this case,  $L_{\mathbf{q}}$  allows us to deduce that there exists a pure strategy  $\sigma$  such that  $\mathbb{E}^{\sigma}_{s_0}(\bar{f}) = \mathbf{q}$ . On the one hand, by Theorem 5.1, there exists a pure strategy  $\sigma$  that is lexicographically optimal from  $s_0$  for  $L_{\mathbf{q}} \circ \bar{f}$ . On the other hand, the only payoff vector  $\mathbf{p} \in \mathsf{Pay}_{s_0}(\bar{f})$  such that  $L_{\mathbf{q}}(\mathbf{p})$  is the lexicographic maximum of  $L_{\mathbf{q}}(\mathsf{Pay}_{s_0}(\bar{f}))$  is  $\mathbf{q}$ . This implies that  $\mathbf{q}$  is the payoff of the pure strategy  $\sigma$ , and thus  $\mathbf{q} \in \mathsf{Pay}^\mathsf{pure}_{s_0}(\bar{f})$ .


**Remark 6.1** (The necessity of induction). In Ex. 6.1, it is possible to isolate  $\mathbf{q}$  by a supporting hyperplane of  $\mathsf{Pay}_{s_0}(\bar{f})$ . We could thus choose  $L_{\mathbf{q}}$  as a linear form and bypass the induction step of the construction of  $L_{\mathbf{q}}$  here. We present it for the sake of illustration, as we cannot use linear forms to isolate extreme points in general (see Section 4).


The construction outlined in Example 6.1 can be generalised to show that all extreme points of the set of expected payoffs of  $\bar{f}$  from a given state are the expected payoff of a pure strategy. However, this property is not sufficient to show that all expected payoffs are convex combinations of pure expected payoffs. In particular, the following example highlights that some expected payoffs are not convex combinations of extreme points of the set of expected payoffs.


**Example 6.2** (Non-extreme point). We consider the MDP depicted in Figure 6.3. We assume that state s is the initial state throughout this example. Let  $w=(w_1,w_2)$  denote the two-dimensional weight function given on the illustration. We consider a two-dimensional payoff function  $\bar{f}$ . The payoff of a play, for each dimension, is zero if t is not visited and, otherwise, its payoff is given by a discounted-sum payoff. We formalise this as the product of a discounted-sum payoff and an indicator. Therefore, formally,  $\bar{f}=(f_1,f_2)$  is such that, for  $j\in\{1,2\}$ ,  $f_j=\mathbbm{1}_{\mathsf{Reach}(\{t\})}\cdot\mathsf{DSum}_{w_j}^{\frac{3}{4}}$ . We observe that, by definition of w,  $f_2=\mathsf{DSum}_{w_2}^{\frac{3}{4}}$ .

The set  $\mathsf{Pay}_s(\bar{f})$  is illustrated in Figure 6.4a. Any vector in  $\mathsf{Pay}_s(\bar{f})$  is a convex combination of  $\mathbf{0}$  and a vector in the segment [(0,4),(4,0)[. In particular, no strategy has an expected payoff of (4,0) from s. We can derive  $\mathsf{Pay}_s(\bar{f})$  from  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}) = \{\mathbf{0}\} \cup \{(4-\frac{3^\ell}{4^\ell-1},\frac{3^\ell}{4^\ell-1}) \mid \ell \in \mathbb{N}\}$ . To obtain  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$ , we note that any pure strategy in this MDP induces a single play from s, because all transitions are deterministic. On the one hand, we can obtain the payoff  $\mathbf{0}$  with the play  $(sb)^\omega$  (the payoff is zero on the first dimension because t is not visited). On the other hand, for all  $\ell \in \mathbb{N}$ , we have  $\bar{f}((sb)^\ell s(at)^\omega) = \left(4-\frac{3^\ell}{4^\ell-1},\frac{3^\ell}{4^{\ell-1}}\right)$ .

We consider the payoff vector  $\mathbf{q}=(2,2)$  and construct  $L_{\mathbf{q}}$ . We remark that the vector  $\mathbf{q}$  is not a convex combination of extreme points of  $\mathsf{Pay}_s(\bar{f})$ . Therefore, is not possible to







(a) The blue dashed line (x + y = 4)is a hyperplane supporting  $Pay_s(f)$  at in Figure 6.4a by the lining Figure 6.4a by the lining **q**. The orange dotted line (x-y=0)is an orthogonal hyperplane included  $(v_1, v_2) \mapsto (v_1 + v_2, v_1 - v_2)$ .  $(v_1, v_2) \mapsto (v_1 + v_2, v_2 - v_1)$ . for reference for the adjacent figures.

(b) Image of the payoff set (c) Image of the payoff set

ear mapping  $L_1$  such that ear mapping  $L_2$  such that

Figure 6.4: The set of expected payoffs for Example 6.2 and its image by two (related) linear functions. The segment (0,0),(4,0) in grey does not intersect  $\mathsf{Pay}_s(f)$ . Its image is similarly coloured in the two other illustrations.

conclude that  $\mathbf{q} \in \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}))$  by adapting the argument of Example 6.1 to deal with all extreme points. The only hyperplane H that support  $\mathsf{Pay}_s(\bar{f})$  at  $\mathbf{q}$  is the line depicted in blue in Figure 6.4a. We let  $x_1^* : \mathbb{R}^2 \to \mathbb{R}$  be the linear form defined by  $x_2^*(\mathbf{v}) = v_1 + v_2$ for all  $\mathbf{v} = (v_1, v_2) \in \mathbb{R}^2$ . We have  $H = (x_1^*)^{-1}(4)$ . We observe (via Figure 6.4a) that  $\mathbf{q}$  is in the relative interior of  $\mathsf{Pay}_s(f) \cap H$ . We define  $L_{\mathbf{q}} = x_1^*$ .

To close this example, we provide an argument based on  $L_{\mathbf{q}}^{-1}(L_{\mathbf{q}}(\mathbf{q}))$  being a line to show that  $\mathbf{q}$  is a convex combination of expected payoffs of pure strategies. While this argument differs from the general proof provided below, it can be generalised to show that  $\mathbf{q} \in \mathsf{conv}(\mathsf{Pay}^\mathsf{pure}_s(\bar{f}))$  whenever  $L^{-1}_{\mathbf{q}}(L_{\mathbf{q}}(\mathbf{q}))$  is a line. This argument consists in showing that there are payoffs of pure strategies on either side of  $\mathbf{q}$  on the line segment  $\mathsf{Pay}_s(f) \cap H = [(0,4),(4,0)[$ . This implies that  $\mathbf{q} \in \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})).$ 

We fix a direction vector  $\mathbf{v}_H = (1, -1)$  of H. For all vectors  $\mathbf{p}$  of  $\mathsf{Pay}_s(\bar{f})$  in the segment  $[\mathbf{q}, (4,0)[ \text{ (resp. } [\mathbf{q}, (0,4)]), \text{ we have } \langle \mathbf{p}, \mathbf{v}_H \rangle \geq \langle \mathbf{q}, \mathbf{v}_H \rangle \text{ (resp. } \langle \mathbf{p}, -\mathbf{v}_H \rangle \geq \langle \mathbf{q}, -\mathbf{v}_H \rangle).$ Consider the linear mappings  $L_1: \mathbf{w} \mapsto (x_1^*, \langle \mathbf{w}, \mathbf{v}_H \rangle)$  and  $L_2: \mathbf{w} \mapsto (x_1^*, \langle \mathbf{w}, -\mathbf{v}_H \rangle)$  over  $\mathbb{R}^2$ . We illustrate the image of  $\mathsf{Pay}_s(\bar{f})$  by  $L_1$  and  $L_2$  respectively in Figure 6.4b and Figure 6.4c.

Theorem 5.1 implies that, for i = 1, 2, there exists a pure strategy  $\sigma_i$  such that  $L_i(\mathbf{p}_i) \geq_{\mathsf{lex}} L_i(\mathbf{q})$  where  $\mathbf{p}_i = \mathbb{E}_s^{\sigma_i}(\bar{f})$ . We obtain, by definition of  $L_i$ , that  $x_1^*(\mathbf{p}_i) =$  $x_1^*(\mathbf{q})$  for i=1,2, as  $x_1^*$  supports  $\mathsf{Pay}_s(\bar{f})$  at  $\mathbf{q}$ . Therefore,  $\alpha_1 \coloneqq \langle \mathbf{p}_1, \mathbf{v}_H \rangle \geq 0$  and  $\alpha_2 := \langle \mathbf{p}_2, \mathbf{v}_H \rangle \leq 0$ . Furthermore, for i = 1, 2, it holds that  $\mathbf{p}_i = \mathbf{q} + \frac{\alpha_i}{\|\mathbf{v}_H\|_2} \mathbf{v}_H$  because  $(\frac{1}{\|\mathbf{q}\|_2} \mathbf{q}, \frac{1}{\|\mathbf{v}_H\|_2} \mathbf{v}_H)$  is an orthonormal basis of  $\mathbb{R}^2$ ,  $\langle \mathbf{p}_i, \frac{1}{\|\mathbf{q}\|_2} \mathbf{q} \rangle = \frac{2}{\|\mathbf{q}\|_2} \cdot x_1^*(\mathbf{p}_i) = 2\sqrt{2} = \|\mathbf{q}\|_2$  (as  $x_1^*(\mathbf{p}_i) = x_1^*(\mathbf{q})$ ) and  $\langle \mathbf{p}_i, \frac{1}{\|\mathbf{v}_H\|_2} \mathbf{v}_H \rangle = \frac{\alpha_i}{\|\mathbf{v}_H\|_2}$  (by definition of  $\alpha_i$ ). Thus,  $\mathbf{q} \in [\mathbf{p}_1, \mathbf{p}_2] \subseteq (\mathbf{p}_1, \mathbf{p}_2)$  $conv(Pay_s^{pure}(\bar{f})).$ 

Formal statement and proof. We formally state the main theorem of this section and prove it.

**Theorem 6.2.** Assume that  $\bar{f}$  is universally integrable. For all  $s \in S$ , we have  $\mathsf{Pay}_s(\bar{f}) =$  $conv(Pay_s^{pure}(\bar{f}))$ . In other words, the expected payoff of any strategy is also the expected payoff of a finite-support mixed strategy.

*Proof.* Throughout this proof, we assume that for all  $1 \leq j \leq d$ ,  $f_j$  is a real-valued payoff. This is without loss of generality: these payoffs are universally integrable, and thus are  $\mathbb{P}_s^{\sigma}$ -almost-surely real-valued for all  $\sigma \in \Sigma(\mathfrak{M})$  and  $s \in S$ .

It is sufficient to show that  $\mathsf{Pay}_s(\bar{f}) \subseteq \mathsf{conv}(\mathsf{Pay}_s^\mathsf{pure}(\bar{f}))$ . Let  $\mathbf{q} \in \mathsf{Pay}_s(\bar{f})$ . We construct the linear mapping  $L_{\mathbf{q}}$  as explained in the sketch, i.e., such that  $L_{\mathbf{q}}(\mathbf{q})$  is the lexicographic maximum of  $L_{\mathbf{q}}(\mathsf{Pay}_s(\bar{f}))$  and  $\mathbf{q} \in \mathsf{ri}(L_{\mathbf{q}}^{-1}(L_{\mathbf{q}}(\mathbf{q})) \cap \mathsf{Pay}_s(\bar{f}))$ .

Let  $D = \mathsf{Pay}_s(\bar{f}) - \mathbf{q}$ . We observe that  $L_{\mathbf{q}}$  satisfies the conditions above if and only if  $L_{\mathbf{q}}(\mathbf{0})$  is the lexicographic maximum of  $L_{\mathbf{q}}(D)$  and  $\mathbf{0}$  is in the relative interior of  $D \cap \ker(L_{\mathbf{q}})$ . We construct  $L_{\mathbf{q}}$  by working with D and  $\mathbf{0}$  instead of  $\mathsf{Pay}_s(\bar{f})$  and  $\mathbf{q}$ . This allows us to work with vector sub-spaces instead of affine spaces, overall simplifying the presentation.

We let  $y_0^* \colon \mathbb{R}^d \to \mathbb{R}$  be the constant zero function. If  $\mathbf{0} \in \mathsf{ri}(D)$ , we let  $L_{\mathbf{q}} = y_0^*$ . This function satisfies the desired properties. We now assume that  $\mathbf{0} \notin \mathsf{ri}(D)$ . We inductively define a sequence of non-zero linear forms  $y_1^*, \ldots, y_{d'}^*$  such that  $y_j^* \colon \mathsf{ker}(y_{j-1}^*) \to \mathbb{R}$  for all  $1 \le j \le d'$ . Next, for all  $1 \le j \le d'$ , we extend  $y_j^*$  to a form  $x_j^* \colon \mathbb{R}^d \to \mathbb{R}$ . Finally, we define the mapping  $L_{\mathbf{q}}$  as  $L_{\mathbf{q}}(\mathbf{v}) = (x_1^*(\mathbf{v}), \ldots, x_{d'}^*(\mathbf{v}))$  for all  $\mathbf{v} \in \mathbb{R}^d$  and show that it satisfies the desired properties.

Let  $j \geq 1$ . By induction, assume that  $y_{j-1}^*$  is defined (this is the case even for j=1). We distinguish two cases. If  $\mathbf{0} \in \mathsf{ri}(\mathsf{ker}(y_{j-1}^*) \cap D)$ , we stop the construction. We remark that if j=d+1, then  $\mathsf{ker}(y_{j-1}^*)$  is a singleton set and we are necessarily in this case (i.e.,  $d' \leq d$ ). Now, assume that  $\mathbf{0} \notin \mathsf{ri}(\mathsf{ker}(y_{j-1}^*) \cap D)$ . The supporting hyperplane theorem (Theorem 2.3) implies that there exists a linear form  $y_j^* \colon \mathsf{ker}(y_{j-1}^*) \to \mathbb{R}$  such that for all  $\mathbf{p} \in \mathsf{Pay}_s(\bar{f}) \cap \mathsf{ker}(y_{j-1}^*)$ , we have  $y^*(\mathbf{p}) \leq 0 = y^*(\mathbf{0})$ . This allows us to continue with the induction.

Assume that the procedure above has provided linear forms  $y_1^*, \ldots, y_{d'}^*$ . We now extend them to  $\mathbb{R}^d$ . Let  $1 \leq j \leq d'$ . There exists  $\mathbf{w}_j \in \ker(y_{j-1}^*) \subseteq \mathbb{R}^d$  such that for all  $\mathbf{v} \in \ker(y_{j-1}^*)$ , we have  $y_j^*(\mathbf{v}) = \langle \mathbf{v}, \mathbf{w}_j \rangle$ . We define  $x_j^* \colon \mathbb{R}^d \to \mathbb{R}$  by, for all  $\mathbf{v} \in \mathbb{R}^d$ ,  $x_j^*(\mathbf{v}) = \langle \mathbf{v}, \mathbf{w}_j \rangle$ . We let  $L_{\mathbf{q}} \colon \mathbb{R}^d \to \mathbb{R}^{d'}$  be such that  $L_{\mathbf{q}}(\mathbf{v}) = (x_1^*(\mathbf{v}), \ldots, x_{d'}^*(\mathbf{v}))$  for all  $\mathbf{v} \in \mathbb{R}^d$ .

We now show that  $L_{\mathbf{q}}$  satisfies the required properties. By construction, for all  $\mathbf{p} \in D$  and all  $1 \leq j \leq d'$ , if  $x_{j'}^*(\mathbf{p}) = 0$  for all  $1 \leq j' < j$ , then necessarily  $\mathbf{p} \in \ker(y_{j-1}^*)$ , and thus  $x_j^*(\mathbf{p}) \leq 0$ . This implies that for all  $\mathbf{p} \in D$ ,  $L_{\mathbf{q}}(\mathbf{p}) \leq_{\mathsf{lex}} L_{\mathbf{q}}(\mathbf{0})$ . This shows that  $L_{\mathbf{q}}(\mathbf{0})$  is the lexicographic maximum of  $L_{\mathbf{q}}(D)$ . Next, we show that  $\mathbf{0} \in \operatorname{ri}(D \cap \ker(L_{\mathbf{q}}))$ . This follows from the stopping condition in the construction of  $L_{\mathbf{q}}$  and the equality  $\ker(L_{\mathbf{q}}) = \bigcap_{1 \leq j \leq d'} \ker(x_j^*) = \ker(y_{d'}^*)$  (the second equality follows from  $x_1^* = y_1^*$  and  $\ker(y_1^*) \supsetneq \ldots \supsetneq \ker(y_{d'}^*)$ ).

Let  $V=L_{\mathbf{q}}^{-1}(L_{\mathbf{q}}(\mathbf{q}))$ . We have shown that  $\mathbf{q}\in \mathrm{ri}(V\cap \mathsf{Pay}_s(\bar{f}))$ . It suffices to show that  $\mathrm{ri}(V\cap \mathsf{Pay}_s(\bar{f}))=\mathrm{ri}(V\cap \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})))$  to conclude that  $\mathbf{q}\in \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}))$ . Since all convex subsets of  $\mathbb{R}^d$  have the same relative interior as their closure [Roc70, Theorem 6.3], the equality of the relative interiors stated before is implied by the relation  $\mathrm{cl}(V\cap \mathsf{Pay}_s(\bar{f}))=\mathrm{cl}(V\cap \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})))$ . To end the proof, we show that  $\mathrm{cl}(V\cap \mathsf{Pay}_s(\bar{f}))=\mathrm{cl}(V\cap \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})))$ . The inclusion  $\mathrm{cl}(V\cap \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})))\subseteq \mathrm{cl}(V\cap \mathsf{Pay}_s(\bar{f}))$  is direct.

For the other inclusion, it suffices to show that  $V \cap \mathsf{Pay}_s(\bar{f}) \subseteq \mathsf{cl}(V \cap \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})))$ . Let  $\mathbf{p} \in V \cap \mathsf{Pay}_s(\bar{f})$ . Assume, by contradiction, that  $\mathbf{p} \notin \mathsf{cl}(V \cap \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})))$ . By the hyperplane separation theorem (Theorem 2.2), there exists a linear form  $x^*$  over  $\mathbb{R}^d$  such that for all  $\mathbf{p}' \in V \cap \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}))$ , we have  $x^*(\mathbf{p}) > x^*(\mathbf{p}')$ . Let  $\mathcal{L}_{\mathbf{q}} \colon \mathbb{R}^d \to \mathbb{R}^{d'+1}$  such that for all  $\mathbf{v} \in \mathbb{R}^d$ , we have  $\mathcal{L}_{\mathbf{q}}(\mathbf{v}) = (L_{\mathbf{q}}(\mathbf{v}), x^*(\mathbf{v}))$ .

Let  $\sigma$  such that  $\mathbf{p} = \mathbb{E}_s^{\sigma}(\bar{f})$ . By Theorem 5.1, there exists a pure strategy  $\tau$  such

that  $\mathbb{E}_{s}^{\tau}(\mathcal{L}_{\mathbf{q}} \circ \bar{f}) \geq_{\mathsf{lex}} \mathbb{E}_{s}^{\sigma}(\mathcal{L}_{\mathbf{q}} \circ \bar{f}) = \mathcal{L}_{\mathbf{q}}(\mathbf{p})$ . We have  $\mathbb{E}_{s}^{\tau}(\bar{f}) \in V$  because  $L_{\mathbf{q}}(\mathbb{E}_{s}^{\tau}(\bar{f})) = \mathbb{E}_{s}^{\tau}(L_{\mathbf{q}} \circ \bar{f}) \geq_{\mathsf{lex}} L_{\mathbf{q}}(\mathbf{p}) = L_{\mathbf{q}}(\mathbf{q})$  and  $L_{\mathbf{q}}(\mathbf{q})$  is lexicographically optimal in  $L_{\mathbf{q}}(\mathsf{Pay}_{s}(\bar{f}))$ . It follows that  $\mathbb{E}_{s}^{\tau}(x^{*} \circ \bar{f}) = x^{*}(\mathbb{E}_{s}^{\tau}(\bar{f})) \geq x^{*}(\mathbf{p})$ . This is a contradiction with  $x^{*}$  defining a strongly separating hyperplane.

We now formulate two corollaries of Theorem 6.2. The first one relates to extreme points of payoffs sets.

Corollary 6.3. Assume that  $\bar{f}$  is universally integrable. For all  $s \in S$ ,  $\text{extr}(\mathsf{Pay}_s(\bar{f})) \subseteq \mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$ , i.e., all extreme points of  $\mathsf{Pay}_s(\bar{f})$  are payoffs of pure strategies.

*Proof.* By Theorem 6.2, we have  $\mathsf{Pay}_s(\bar{f}) = \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}))$ . All extreme points of  $\mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}))$  must be in  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$  by definition of the convex hull.

Second, we establish that, for all  $s \in S$ ,  $\mathsf{Pay}_s(\bar{f})$  is closed whenever  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$  is closed. We note that, in general, the convex hull of a closed set need not be closed. However, the convex hull of a compact subset of  $\mathbb{R}^d$  is closed (Lemma 4.3). Since the set of expected payoffs of a universally integrable payoff function is bounded, it is thus compact, implying the claimed property.

Corollary 6.4. Assume that  $\bar{f}$  is universally integrable. For all  $s \in S$ , if  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$  is closed, then  $\mathsf{Pay}_s(\bar{f})$  is compact.

*Proof.* Let  $s \in S$  such that  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$  is closed. By Lemma 3.2,  $\bar{f}$  is universally integrable if and only if  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$  is bounded. It follows that  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$  is compact. Theorem 6.2 ensures that  $\mathsf{Pay}_s(\bar{f}) = \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}))$ , and thus  $\mathsf{Pay}_s(\bar{f})$  is compact by Lemma 4.3.  $\square$ 

We close this section by showing that Theorem 6.2 does not generalise to universally unambiguously integrable payoffs.

**Example 6.3** (Example 5.1 continued). We consider the MDP  $\mathfrak{M}$  depicted in Figure 5.1 and the payoff function  $\bar{f}=(\mathbb{1}_{\mathsf{Reach}(\{t\})},\mathsf{TRew}_w)$  where w is the weight function of Figure 5.1. In Example 5.1, we have shown that there exists a randomised strategy  $\sigma$  such that  $\mathbb{E}_s^{\sigma}(\bar{f})=(1,+\infty)$  and that  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})=\{(0,+\infty)\}\cup\{(1,\ell)\mid \ell\in\mathbb{N}\}.$ 

We show that  $(1, +\infty) \notin \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}))$ . On the one hand, convex combinations of these vectors that give a non-zero coefficient to  $(0, +\infty)$  have a first component is not equal to 1. On the other hand, convex combinations that assign a zero coefficient to  $(0, +\infty)$  have a finite second component. We obtain that  $(1, +\infty) \notin \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}))$ .

Although we cannot have a payoff of  $(1, +\infty)$  with finite-support mixed strategies, we can approximate it with such strategies. We generalise this observation in the next section.

## 6.2 Universally unambiguously integrable payoffs

We now relax the assumption that  $\bar{f}$  is universally integrable and assume that  $\bar{f}$  is universally unambiguously integrable. We formulate an approximate variant of Theorem 6.2: from a given state, any expected payoff of a strategy can be approached by convex combinations of expected payoffs of pure strategies (in the sense of limits in  $\mathbb{R}^d$ ).

We provide a proof sketch, followed by the formal statement of the result and its proof.




Figure 6.5: Illustration of the approach used to construct the approximation  $\mathcal{Y}$  of  $\mathcal{X}$  adapted to a function over [0,1]. We round the blue function down to the closest multiple of  $\frac{1}{4}$  to obtain the red function. This yields a linear combination of indicators that is  $\frac{1}{4}$ -close in all points to the function in blue.

**Proof overview.** Fix  $s \in S$  and a strategy  $\sigma$ . The goal is to show that all neighbourhoods of  $\mathbb{E}^{\sigma}_{s}(\bar{f})$  intersect  $\mathsf{conv}(\mathsf{Pay}^{\mathsf{pure}}_{s}(\bar{f}))$ . In other words, we must prove that for all  $\varepsilon > 0$  and all  $M \in \mathbb{R}$ , there exist finitely many pure strategies  $\tau_{1}, \ldots, \tau_{n}$  and convex combination coefficients  $\alpha_{1}, \ldots, \alpha_{n} \in [0,1]$  and, for all  $1 \leq j \leq d$ , if  $\mathbb{E}^{\sigma}_{s}(f_{j}) = +\infty$ , then  $\sum_{m=1}^{n} \alpha_{m} \mathbb{E}^{\tau_{m}}_{s}(f_{j}) \geq M$ , if  $\mathbb{E}^{\sigma}_{s}(f_{j}) = -\infty$ , then  $\sum_{m=1}^{n} \alpha_{m} \mathbb{E}^{\tau_{m}}_{s}(f_{j}) \leq -M$  and, otherwise,  $\sum_{m=1}^{n} \alpha_{m} \mathbb{E}^{\tau_{m}}_{s}(f_{j}) \geq \mathbb{E}^{\sigma}_{s}(f_{j}) - \varepsilon$ . We fix  $\varepsilon > 0$  and  $M \in \mathbb{R}$ .

The proof is based on the reformulation of  $\mathbb{E}_{s}^{\sigma}(\bar{f})$  as an integral (Lemma 3.1) and the manipulation of random variables. While Lemma 3.1 is not applicable to all payoffs, Lemma 3.3 allows us to invoke it on a shifted payoff. We can then recover the result for the original payoff using the linearity of the expectation. We thus assume without loss of generality that Lemma 3.1 applies to all payoffs  $f_1, \ldots, f_d$ . We consider the distribution  $\nu_{\sigma} \in \mathcal{D}([0,1]^{\omega}, \mathcal{A}_{[0,1]^{\omega}})$  and the mapping  $[0,1]^{\omega} \to \Sigma_{\mathsf{pure}}(\mathfrak{M}) \colon x \mapsto \tau_x$  such that  $\mathbb{E}_{s}^{\sigma}(\bar{f}) = \int_{x \in [0,1]^{\omega}} \mathbb{E}_{s}^{\tau_x}(\bar{f}) d\nu_{\sigma}(x)$  that are given by Lemma 3.1. For all  $1 \leq j \leq d$ , we let  $X_j \colon [0,1]^{\omega} \to \mathbb{R} \colon x \mapsto \mathbb{E}_{s}^{\tau_x}(f_j)$  denote the random variable appearing in the previous integral. We let  $\mathcal{X} = (X_1, \ldots, X_d)$ . We sketch the proof when the  $X_j$  are  $(\nu_{\sigma}$ -almost-surely) real-valued functions. We comment on the generalisation at the end of the sketch.

The broad idea is as follows. First, we approximate  $\mathcal{X}$  with a multivariate random variable  $\mathcal{Y} = (Y_1, \dots, Y_d)$  over  $[0,1]^{\omega}$ . We then approximate the integral of  $\mathcal{Y}$  with a convex combination  $\sum_{m=0}^{n} \alpha_m \mathcal{Y}(x_m) \in \mathsf{conv}(\mathsf{Im}(\mathcal{Y}))$ . Finally, we derive the convex combination  $\sum_{m=0}^{n} \alpha_m \mathcal{X}(x_m)$  from the previous one. The successive approximations above ensure that the last convex combination respects the claims of the theorem.

We now expand on the broad idea above. First, we construct  $\mathcal{Y}$  such that  $\mathcal{X} - \frac{\varepsilon}{3}\mathbf{1} \leq \mathcal{Y} \leq \mathcal{X} + \frac{\varepsilon}{3}\mathbf{1}$ . It follows that, for all  $1 \leq j \leq d$ ,  $\int_{x \in [0,1]^{\omega}} Y_j d\nu_{\sigma}(x)$  is equal to  $\mathbb{E}_s^{\sigma}(f_j)$  whenever  $\mathbb{E}_s^{\sigma}(f_j) \in \{-\infty, +\infty\}$  and otherwise is  $\frac{\varepsilon}{3}$ -close. Intuitively, we construct  $\mathcal{Y}$  as an infinite linear combination of indicators, following the rounding idea illustrated in Figure 6.5 (where the rounding precision depends on  $\varepsilon$ ). Its integral is thus (informally) an infinite convex combination of images of  $\mathcal{Y}$ : there are sequences  $(\beta_m)_{m \in \mathbb{N}}$  and  $(x_m)_{m \in \mathbb{N}}$  respectively of coefficients and elements of  $[0,1]^{\omega}$  such that  $\sum_{m=0}^{\infty} \beta_m = 1$  and  $\int_{x \in [0,1]^{\omega}} \mathcal{Y} d\nu_{\sigma}(x) = \sum_{m \in \mathbb{N}} \beta_m \mathcal{Y}(x_m)$ . We derive a sequence  $(\mathbf{p}^{(n)})_{n \in \mathbb{N}}$  in  $\mathsf{conv}(\mathsf{Im}(\mathcal{Y}(x_m)))$  that converges to  $\int_{x \in [0,1]^{\omega}} \mathcal{Y} d\nu_{\sigma}(x)$  from this series: we let  $\mathbf{p}^{(n)} = \sum_{m=0}^{n} \beta_m \mathcal{Y}(x_m) + (1 - \sum_{m=0}^{n} \beta_m) \mathcal{Y}(x_0)$  for all  $n \in \mathbb{N}$ .

Fix  $n \in \mathbb{N}$  large enough such that, for all  $1 \leq j \leq d$ , component j of  $\mathbf{p}^{(n)}$  is  $\frac{\varepsilon}{3}$ -close to  $\int_{x \in [0,1]^{\omega}} Y_j d\nu_{\sigma}(x)$  if it is a real number or greater than  $M + \varepsilon$  in absolute value otherwise. The convex combination  $\mathbf{q} = \sum_{m=0}^n \beta_m \mathbb{E}_s^{\tau_{x_m}}(\bar{f}) + (1 - \sum_{m=0}^n \beta_m) \mathbb{E}_s^{\tau_{x_0}}(\bar{f})$  is a satisfactory convex combination with respect to the claim of the theorem. Let  $1 \leq j \leq d$ . If  $\mathbb{E}_s^{\sigma}(f_j) = +\infty$ , we obtain that  $q_j \geq M$ . Similarly, if  $\mathbb{E}_s^{\sigma}(f_j) = -\infty$ , we obtain that  $q_j \leq -M$ . Otherwise, we have that  $q_j$  is  $\varepsilon$ -close to  $\mathbb{E}_s^{\sigma}(f_j)$ .

We now briefly discuss the case where some  $X_j$  is not  $\nu_{\sigma}$ -almost-surely real-valued. For the sake of illustration, we assume that this only applies to j=d and that  $X_d \geq 0$ . Therefore, we have  $\mathbb{E}_s^{\sigma}(f_d) = +\infty$  and there exists some  $x \in [0,1]^{\omega}$  such that  $\mathbb{E}_s^{\tau_x}(f_d) = +\infty$  and  $\mathbb{E}_s^{\tau_x}(f_j) \in \mathbb{R}$  for all  $1 \leq j \leq d-1$ . Let  $\tau_1, \ldots, \tau_n$  and  $\alpha_1, \ldots, \alpha_n$  given by the theorem for  $(f_1, \ldots, f_{d-1})$ ,  $\frac{\varepsilon}{2}$  and  $M + \varepsilon$ . Let  $\mathbf{q} = \sum_{m=1}^n \alpha_m \mathbb{E}_s^{\tau_m}(\bar{f})$ . By choosing  $\eta \in ]0, 1[$  such that all components of  $\mathbb{E}_s^{\tau_x}(\bar{f})$  other than the last have absolute value no more than  $\frac{\varepsilon}{2}$  and the finite components of  $(1-\eta)\mathbf{q}$  are  $\frac{\varepsilon}{2}$ -close to the corresponding components of  $\mathbf{q}$ , we obtain a suitable convex combination in the form of  $\eta \mathbb{E}_s^{\tau_x}(\bar{f}) + (1-\eta)\mathbf{q}$ .

Formal statement and proof. We state our theorem below and prove it.

**Theorem 6.5.** Assume that  $\bar{f}$  is universally unambiguously integrable. Let  $s \in S$ . We have  $\operatorname{cl}(\mathsf{Pay}_s(\bar{f})) = \operatorname{cl}(\mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})))$ . In particular, for all strategies  $\sigma$ , all  $\varepsilon > 0$  and all  $M \in \mathbb{R}$ , there exist finitely many pure strategies  $\tau_1, \ldots, \tau_n$  and convex combination coefficients  $\alpha_1, \ldots, \alpha_n \in [0, 1]$  such that for all  $1 \leq j \leq d$ :

- if  $\mathbb{E}_s^{\sigma}(f_i) = +\infty$ , then  $\sum_{m=1}^n \alpha_m \mathbb{E}_s^{\tau_m}(f_i) \geq M$ ,
- if  $\mathbb{E}_s^{\sigma}(f_j) = -\infty$ , then  $\sum_{m=1}^n \alpha_m \mathbb{E}_s^{\tau_m}(f_j) \leq -M$ , and,
- otherwise, if  $\mathbb{E}_s^{\sigma}(f_j) \in \mathbb{R}$ ,  $\mathbb{E}_s^{\sigma}(f_j) \varepsilon \leq \sum_{m=1}^n \alpha_m \mathbb{E}_s^{\tau_m}(f_j) \leq \mathbb{E}_s^{\sigma}(f_j) + \varepsilon$ .

*Proof.* The inclusion  $\operatorname{cl}(\operatorname{conv}(\operatorname{\mathsf{Pay}}^{\mathsf{pure}}_s(\bar{f}))) \subseteq \operatorname{cl}(\operatorname{\mathsf{Pay}}_s(\bar{f}))$  follows from the convexity of  $\operatorname{\mathsf{Pay}}_s(\bar{f})$ . For other inclusion, it suffices to show that  $\operatorname{\mathsf{Pay}}_s(\bar{f}) \subseteq \operatorname{cl}(\operatorname{\mathsf{conv}}(\operatorname{\mathsf{Pay}}^{\mathsf{pure}}_s(\bar{f})))$ . This inclusion is equivalent to the last property of the theorem statement.

Let  $\sigma$  be a strategy,  $\varepsilon > 0$  and  $M \in \mathbb{R}$ . To prove the theorem, we reason on random variables of  $([0,1]^{\omega}, \mathcal{A}_{[0,1]^{\omega}})$ . We consider the distribution  $\nu_{\sigma} \in \mathcal{D}([0,1]^{\omega}, \mathcal{A}_{[0,1]^{\omega}})$  and the mapping  $[0,1]^{\omega} \to \Sigma_{\mathsf{pure}}(\mathfrak{M}) \colon x \mapsto \tau_x$  given by Lemma 3.1.

We make two assumptions without loss of generality. We defer the proof that these assumptions are without loss of generality to the end of the proof. First, we assume that for all  $1 \leq j \leq d$ , either  $\mathbb{E}_s^{\tau}(f_j) \geq 0$  for all strategies  $\tau$  or  $\mathbb{E}_s^{\tau}(f_j) \leq 0$  for all strategies  $\tau$ . This assumption guarantees that  $\mathbb{E}_s^{\sigma}(\bar{f}) = \int_{x \in [0,1]^{\omega}} \mathbb{E}_s^{\tau_x}(\bar{f}) d\nu_{\sigma}(x)$  by Lemma 3.1. Second, we assume that for all  $1 \leq j \leq d$ , the set of  $x \in [0,1]^{\omega}$  such that  $\mathbb{E}_s^{\tau_x}(f_j) \notin \mathbb{R}$  has  $\nu_{\sigma}$ -measure zero.

For all  $1 \leq j \leq d$ , we consider the random variable  $X_j : x \mapsto \mathbb{E}_s^{\tau_x}(f_j)$  over  $[0,1]^{\omega}$ . We let  $\mathcal{X} = (X_1, \dots, X_d)$ . The first assumption above implies that for all  $1 \leq j \leq d$ ,  $X_j$  is a non-negative or non-positive random variable. The second assumption implies that  $\mathcal{X}$  is almost-surely  $\mathbb{R}^d$ -valued. We thus interpret the random variable  $\mathcal{X}$  as a function  $[0,1]^{\omega} \to \mathbb{R}^d$ .

We now prove the result under the assumptions above. The first step of the proof is to construct a random variable  $\mathcal{Y} = (Y_1, \dots, Y_d) \colon [0,1]^{\omega} \to \mathbb{R}^d$  which approximates  $\mathcal{X}$ . To ensure that the integral of  $\mathcal{Y}$  is an infinite convex combination of images of  $\mathcal{Y}$  (in the sense of the proof overview), we define  $\mathcal{Y}$  as a function similar to a simple function. More precisely, we define  $\mathcal{Y}$  as a series of indicators multiplied by coefficients.

We fix  $k \in \mathbb{N}$  such that  $\frac{1}{2^k} \leq \frac{\varepsilon}{3}$ . We define  $\mathcal{Y}$  component by component first, and introduce its series form later. Let  $1 \leq j \leq d$ . We generalise the construction illustrated in Figure 6.5. For all  $k \in \mathbb{N}$ , we let  $Y_j$  be the random variable over  $[0,1]^{\omega}$  such that

$$Y_j = \sum_{\ell=0}^{\infty} \sum_{i=0}^{2^k - 1} \left( \ell + \frac{i}{2^k} \right) \mathbb{1}_{\left[\ell + \frac{i}{2^k}, \ell + \frac{i+1}{2^k}\right]} (X_j)$$

if  $X_j$  is non-negative and, otherwise,

$$Y_j = \sum_{\ell=0}^{\infty} \sum_{i=0}^{2^k - 1} - \left(\ell + \frac{i}{2^k}\right) \mathbb{1}_{\left] - \ell - \frac{i+1}{2^k}, -\ell - \frac{i}{2^k}\right]}(X_j).$$

We have  $X_j - \frac{1}{2^k} \le Y_j \le X_j + \frac{1}{2^k}$ . This implies that  $X_j - \frac{\varepsilon}{3} \le Y_j^{(k)} \le X_j + \frac{\varepsilon}{3}$ . In particular,  $Y_j$  is integrable if and only if  $X_j$  is and, and, if both are integrable:

$$\mathbb{E}_{s}^{\sigma}(f_{j}) - \frac{\varepsilon}{3} \leq \int_{x \in [0,1]^{\omega}} Y_{j}(x) d\nu_{\sigma}(x) \leq \mathbb{E}_{s}^{\sigma}(f_{j}) + \frac{\varepsilon}{3}.$$

$$(6.1)$$

We also have  $\mathbb{E}_s^{\sigma}(f_j) = +\infty$  (resp.  $-\infty$ ) if and only if  $\int_{x \in [0,1]^{\omega}} Y_j(x) d\nu_{\sigma}(x) = +\infty$  (resp.  $-\infty$ ).

Now that we have shown that  $\mathcal Y$  approximates  $\mathcal X$ , we prove that the integral of  $\mathcal Y$  can be written as an infinite convex combination of elements of  $\operatorname{Im}(\mathcal Y)$ . To this end, we rewrite  $\mathcal Y$  in the series form mentioned above. Let  $1 \leq j \leq d$ . For all  $(\ell,i) \in \mathbb N \times \{0,1,\ldots,2^k-1\}$ , we define  $I_j(\ell,i) = \left[\ell + \frac{i}{2^k}, \ell + \frac{i+1}{2^k}\right]$  and  $v_j = \ell + \frac{i}{2^k}$  if  $X_j \geq 0$ , and  $I_j(\ell,i) = \left] - \ell - \frac{i+1}{2^k}, -\ell - \frac{i}{2^k}\right]$  and  $v_j = -\ell - \frac{i}{2^k}$  if  $X_j \leq 0$ . We fix an enumeration  $(\bar\ell^{(m)},\bar\imath^{(m)})_{m\in\mathbb N}$  of  $\mathbb N^d \times \{0,\ldots,2^k-1\}^d$ . For all  $m\in\mathbb N$ , let  $\bar\ell^{(m)} = (\ell_1^{(m)},\ldots,\ell_d^{(m)}) \in \mathbb N^d$  and  $\bar\imath^{(m)} = (i_1^{(m)},\ldots,i_d^{(m)}) \in \{0,\ldots,2^k-1\}^d$ . We let  $B_m = \prod_{j=1}^d I_j(\ell_j^{(m)},i_j^{(m)})$  and  $\mathbf v_m = (v_j(\ell_j^{(m)},i_j^{(m)}))_{1\leq j\leq d}$ . We can rewrite  $\mathcal Y$  as follows, using this notation:

$$\mathcal{Y} = \sum_{m \in \mathbb{N}} \mathbf{v}_m \mathbb{1}_{B_m}(\mathcal{X}).$$

Through this, we obtain that the integral of  $\mathcal{Y}$  is an infinite convex combination: the monotone convergence theorem ensures that

$$\int_{x\in[0,1]^{\omega}} \mathcal{Y}(x) d\nu_{\sigma}(x) = \sum_{m\in\mathbb{N}} \mathbf{v}_{m} \nu_{\sigma} \left( \left\{ x \in [0,1]^{\omega} \mid \mathcal{X}(x) \in B_{m} \right\} \right).$$

Next, we define a sequence in  $\mathsf{conv}(\mathsf{Im}(\mathcal{Y}))$  converging to  $\int_{x\in[0,1]^{\omega}} \mathcal{Y}(x) \mathrm{d}\nu_{\sigma}(x)$ . It is used to determine the coefficients and pure strategies we seek: strategies are obtained via the mapping  $x\mapsto \tau_x$ . For all  $m\in\mathbb{N}$ , we let  $\beta_m=\nu_{\sigma}\left(\{x\in[0,1]^{\omega}\mid\mathcal{X}(x)\in B_m\}\right)$  and let  $x_m\in[0,1]^{\omega}$  such that  $\mathbf{v}_m=\mathcal{Y}(x_m)$  if  $\beta_m\neq 0$ , and  $x_m$  is left arbitrary otherwise. We consider the sequence  $(\mathbf{p}^{(n)})_{n\in\mathbb{N}}$  defined by, for all  $n\in\mathbb{N}$ ,

$$\mathbf{p}^{(n)} = \sum_{m=1}^{n} \beta_m \mathcal{Y}(x_m) + \left(1 - \sum_{m=1}^{n} \beta_m\right) \mathcal{Y}(x_0).$$

We have  $\lim_{n\to\infty} \mathbf{p}^{(n)} = \int_{x\in[0,1]^{\omega}} \mathcal{Y}(x) d\nu_{\sigma}(x)$  and for all  $n\in\mathbb{N}$ ,  $\mathbf{p}^{(n)}\in\mathsf{conv}(\mathsf{Im}(\mathcal{Y}))$ . For all  $n\in\mathbb{N}$ , we let  $\mathbf{p}^{(n)}=(p_1^{(n)},\ldots,p_d^{(n)})$ .

We now fix n such that, for all  $1 \leq j \leq d$ ,  $\int_{x \in [0,1]^{\omega}} Y_j(x) d\nu_{\sigma}(x) \in \mathbb{R}$  implies that

$$-\frac{\varepsilon}{3} \le p_j^{(n)} - \int_{x \in [0,1]^\omega} Y_j(x) d\nu_\sigma(x) \le \frac{\varepsilon}{3}, \tag{6.2}$$

 $\int_{x\in[0,1]^{\omega}}Y_j(x)\mathrm{d}\nu_{\sigma}(x) = +\infty \text{ implies that } p_j^{(n)} \geq M + \varepsilon \text{ and } \int_{x\in[0,1]^{\omega}}Y_j(x)\mathrm{d}\nu_{\sigma}(x) = -\infty \text{ implies that } p_j^{(n)} \leq -M - \varepsilon. \text{ We set } \alpha_0 = 1 - \sum_{m=1}^n \beta_m \text{ and for } 1 \leq m \leq n, \ \alpha_m = \beta_m. \text{ We remark that } \mathbf{p}^{(n)} = \sum_{m=0}^n \alpha_m \mathcal{Y}(x_m). \text{ For all } 1 \leq m \leq n, \text{ we set } \tau_m = \tau_{x_m}. \text{ We show that the convex combination } \mathbf{q} = \sum_{m=0}^n \alpha_m \mathcal{X}(x_m) = \sum_{m=0}^n \alpha_m \mathbb{E}_s^{\tau_m}(\bar{f}) \text{ satisfies the claim of the theorem. We write } \mathbf{q} = (q_1, \dots, q_d). \text{ It follows from the inequalities } Y_j - \frac{\varepsilon}{3} \leq X_j \leq Y_j + \frac{\varepsilon}{3} \text{ for all } 1 \leq j \leq d \text{ and the definitions of } \mathbf{q} \text{ and } \mathbf{p}^{(n)} \text{ that}$ 

$$\mathbf{p}^{(n)} - \frac{\varepsilon}{3} \mathbf{1} \le \mathbf{q} \le \mathbf{p}^{(n)} + \frac{\varepsilon}{3} \mathbf{1}. \tag{6.3}$$

Let  $1 \leq j \leq d$ . First, assume that  $\mathbb{E}_s^{\sigma}(f_j) = +\infty$  (i.e.,  $\int_{x \in [0,1]^{\omega}} Y_j(x) d\nu_{\sigma}(x) = +\infty$ ). In this case, it follows from Equation (6.3) and  $p_j^{(n)} \geq M + \varepsilon$  that

$$q_j = \sum_{m=0}^n \alpha_m \mathbb{E}_s^{\tau_{x_n}}(f_j) \ge p_j^{(n)} - \frac{\varepsilon}{3} \ge M + \frac{2\varepsilon}{3} \ge M.$$

The argument for the case  $\mathbb{E}_s^{\sigma}(f_j) = -\infty$  follows an analogous reasoning and is omitted. We now assume that  $\mathbb{E}_s^{\sigma}(f_j) \in \mathbb{R}$  (i.e.,  $\int_{x \in [0,1]^{\omega}} Y_j(x) d\nu_{\sigma}(x) \in \mathbb{R}$ ). By applying Equations (6.3), (6.2) and (6.1) in succession, we obtain that

$$q_{j} \leq p_{j}^{(n)} + \frac{\varepsilon}{3}$$

$$\leq \int_{x \in [0,1]^{\omega}} Y_{j}(x) d\nu_{\sigma}(x) + \frac{2\varepsilon}{3}$$

$$\leq \mathbb{E}_{s}^{\sigma}(f_{j}) + \varepsilon.$$

A similar succession of inequalities (referring to the same equations), yields  $q_j \geq \mathbb{E}_s^{\sigma}(f_j) - \varepsilon$ . This ends the argument that **q** satisfies the conditions outlined in the statement of the theorem.

To end the proof, it remains to show that the assumptions made above are without loss of generality. We recall them first:

- 1. for all  $1 \leq j \leq d$ ,  $\mathbb{E}_s^{\tau}(f_i) \geq 0$  for all strategies  $\tau$  or  $\mathbb{E}_s^{\tau}(f_i) \leq 0$  for all strategies  $\tau$ ;
- 2. for all  $1 \leq j \leq d$ , the set of  $x \in [0,1]^{\omega}$  such that  $\mathbb{E}_s^{\tau_x}(f_j) \notin \mathbb{R}$  has  $\nu_{\sigma}$ -measure zero.

In the above, we have shown that the claim of the theorem holds with Assumptions 1 and 2. In the following, we first show that the theorem with both Assumptions 1 and 2 implies the theorem with only Assumption 2. After this, we show that the theorem with Assumption 2 implies the theorem with neither additional assumption.

Assume that Assumption 2 holds and let us show that the claim of the theorem holds. To obtain the result for  $\bar{f}$ , we derive a payoff  $\bar{g} = (g_1, \ldots, g_d)$  from  $\bar{f}$  such that  $\bar{g}$  satisfies the conditions outlined in Assumptions 1 and 2, so that we can apply the variant of the theorem with Assumptions 1 and 2 to  $\bar{g}$ .

Let  $1 \leq j \leq d$ . If  $\inf_{\tau \in \Sigma(\mathfrak{M})} \mathbb{E}_s^{\tau}(f_j) \in \mathbb{R}$ , we let  $g_j = f_j - \inf_{\tau \in \Sigma(\mathfrak{M})} \mathbb{E}_s^{\tau}(f_j)$ . We obtain that  $\mathbb{E}_s^{\tau}(g_j) \geq 0$  for all strategies  $\tau$ . Indeed, for all strategies  $\tau$ , this follows by linearity of  $\mathbb{E}$ 

if  $f_j$  is  $\mathbb{P}_s^{\tau}$ -integrable (which is equivalent to  $g_j$  being  $\mathbb{P}_s^{\tau}$ -integrable) and otherwise the non-negative parts of  $f_j$  and  $g_j$  are  $|\inf_{\tau'\in\Sigma(\mathfrak{M})}\mathbb{E}_s^{\tau'}(f_j)|$ -close to one another, thus share their infinite integral and we obtain  $\mathbb{E}_s^{\tau}(g_j)=\mathbb{E}_s^{\tau}(f_j)=+\infty\geq 0$ . Otherwise, by Lemma 3.3, we have  $\sup_{\tau\in\Sigma(\mathfrak{M})}\mathbb{E}_s^{\tau}(f_j)\in\mathbb{R}$  and we let  $g_j=f_j-\sup_{\tau\in\Sigma(\mathfrak{M})}\mathbb{E}_s^{\tau}(f_j)$ . By adapting the argument of the previous case, we obtain that for all strategies  $\tau$ , we have  $\mathbb{E}_s^{\tau}(g_j)\leq 0$  and the equivalence  $\mathbb{E}_s^{\tau}(g_j)=-\infty$  if and only if  $\mathbb{E}_s^{\tau}(f_j)=-\infty$ .

Let  $M'=M+\max_{1\leq j\leq d}|\gamma_j|$  where  $\gamma_j$  is the constant such that  $g_j=f_j-\gamma_j$  for all  $1\leq j\leq d$ . We let  $\tau_1,\ldots,\tau_n$  be the strategies and  $\alpha_1,\ldots,\alpha_n\in[0,1]$  be the coefficients given by the restricted theorem for  $\bar{g},\,\sigma,\,\varepsilon$  and M'. Let  $1\leq j\leq d$ . For all  $1\leq m\leq n,\,f_j$  is  $\mathbb{P}^{\tau_m}_s$ -integrable (by the remaining additional assumption). We thus have  $\sum_{m=1}^n\alpha_m\mathbb{E}^{\tau_m}(f_j)=\gamma_j+\sum_{m=1}^n\alpha_m\mathbb{E}^{\tau_m}(g_j)$ . If  $\mathbb{E}^{\sigma}_s(f_j)\in\mathbb{R}$ , i.e.,  $f_j$  is  $\mathbb{P}^{\sigma}_s$ -integrable, then so is  $g_j$  and we directly obtain  $\mathbb{E}^{\sigma}_s(f_j)-\varepsilon\leq\sum_{m=1}^n\alpha_m\mathbb{E}^{\tau_m}_s(f_j)\leq\mathbb{E}^{\sigma}_s(f_j)+\varepsilon$  from the similar inequality for  $g_j$ . Next, assume that  $\mathbb{E}^{\sigma}_s(f_j)=+\infty$ . By the above, this implies that  $\mathbb{E}^{\sigma}_s(g_j)=+\infty$ . We obtain (from the application of the theorem to  $g_j$ ) that  $\sum_{m=1}^n\alpha_m\mathbb{E}^{\tau_m}_s(f_j)\geq\gamma_j+M'\geq M$ . Finally, if  $\mathbb{E}^{\sigma}_s(g_j)=-\infty$ , we obtain that  $\sum_{m=1}^n\alpha_m\mathbb{E}^{\tau_m}_s(f_j)\leq\gamma_j-M'\leq -M$  in the same way as the previous case.

It remains to show that the claim of the theorem with Assumption 2 implies the theorem with no assumption. For convenience of notation, we assume that there exists  $1 \leq d' \leq d$  such that  $x \mapsto \mathbb{E}_s^{\tau_x}(f_j)$  is not  $\nu_{\sigma}$ -almost-surely real-valued for  $1 \leq j \leq d'$  and is  $\nu_{\sigma}$ -almost-surely real-valued is for  $d' + 1 \leq j \leq d$ .

For all dimensions  $1 \leq j \leq d'$ ,  $\mathbb{E}_s^{\sigma}(f_j)$  is infinite and there exists  $x_j \in [0,1]^{\omega}$  such that the strategy  $\tau_j = \tau_{x_j}$  satisfies  $\mathbb{E}_s^{\tau_j}(f_j) = \mathbb{E}_s^{\sigma}(f_j)$  (because  $\nu_{\sigma}(\{x \in [0,1]^{\omega} \mid \mathbb{E}_s^{\tau_x}(f_j) \notin \mathbb{R}\}) \neq 0$ ) and, for all  $d'+1 \leq j' \leq d$ ,  $\mathbb{E}_s^{\tau_{x_j}}(f_j) \in \mathbb{R}$  (because  $\nu_{\sigma}(\{x \in [0,1]^{\omega} \mid \mathbb{E}_s^{\tau_x}(f_j) \in \mathbb{R}\}) = 1$ ). If d'=d, we conclude using the convex combination  $\sum_{j=1}^d \frac{1}{d} \mathbb{E}_s^{\tau_j}(\bar{f}) = \mathbb{E}_s^{\sigma}(\bar{f})$ . Therefore, we assume that d' < d. The payoff  $(f_{d'+1}, \ldots, f_d)$  satisfies Assumption 2. We apply the theorem with Assumption 2 to  $(f_{d'+1}, \ldots, f_d)$ , strategy  $\sigma$ ,  $\frac{\varepsilon}{3}$  and  $M + \varepsilon$  to obtain pure strategies  $\tau_{d'+1}, \ldots, \tau_n$  and convex combination coefficients  $\beta_{d'+1}, \ldots, \beta_n$  that satisfy the implications given in the last property of the statement of the theorem.

We now define convex combination coefficients  $\alpha_1, \ldots, \alpha_n$  to obtain the claim of the theorem for  $\bar{f}$ . Fix  $\eta \in ]0,1[$  such that (i) for all  $1 \leq j \leq d'$ , the real components of  $\eta \mathbb{E}_s^{\tau_j}(\bar{f})$  are no more than  $\frac{\varepsilon}{3}$  in absolute value and (ii)  $(1-\eta)\sum_{m=d'+1}^n \beta_m \mathbb{E}_s^{\tau_m}((f_{d'+1},\ldots,f_d))$  is  $\frac{\varepsilon}{3}$ -close to  $\sum_{m=d'+1}^n \beta_m \mathbb{E}_s^{\tau_m}((f_{d'+1},\ldots,f_d))$ . For all  $1 \leq m \leq d'$ , we set  $\alpha_m = \frac{\eta}{d'}$  and for  $d'+1 \leq m \leq n$ , we set  $\alpha_m = (1-\eta)\beta_m$ . It follows from  $\eta \in ]0,1[$  that  $1-\eta>0$  and  $\sum_{m=1}^n \alpha_m = 1$ .

We show that the pure strategies  $\tau_1, \ldots, \tau_m$  and coefficients  $\alpha_1, \ldots, \alpha_n$  are witnesses to the implications in the theorem statement. Let  $1 \leq j \leq d$ . If  $j \leq d'$ , it follows from  $\eta > 0$  that  $\sum_{m=1}^{n} \alpha_m \mathbb{E}_s^{\tau_m}(f_j) = \eta \mathbb{E}_s^{\tau_j}(f_j) = \mathbb{E}_s^{\sigma}(f_j) \in \{-\infty, +\infty\}$ , and the required inequality is trivially satisfied. We assume from here that  $j \geq d' + 1$ . It follows from properties (i) and (ii) above that  $\sum_{m=1}^{n} \alpha_m \mathbb{E}_s^{\tau_m}(f_j)$  is  $\frac{2\cdot\varepsilon}{3}$ -close to  $\sum_{m=d'+1}^{n} \beta_m \mathbb{E}_s^{\tau_m}(f_j)$ . Assume that  $\mathbb{E}_s^{\sigma}(f_j) = +\infty$ . Then, we have

$$\sum_{m=1}^{n} \alpha_m \mathbb{E}_s^{\tau_m}(f_j) \ge \sum_{m=d'+1}^{n} \beta_m \mathbb{E}_s^{\tau_m}(f_j) - \frac{2 \cdot \varepsilon}{3} \ge M + \varepsilon - \frac{2 \cdot \varepsilon}{3} \ge M.$$

We obtain that  $\mathbb{E}_s^{\sigma}(f_j) = -\infty$  implies  $\sum_{m=1}^n \alpha_m \mathbb{E}_s^{\tau_m}(f_j) \leq -M$  similarly. Finally, assume that  $\mathbb{E}_s^{\sigma}(f_j) \in \mathbb{R}$ . We recall that  $|\sum_{m=d'+1}^n \beta_m \mathbb{E}_s^{\tau_m}(f_j) - \mathbb{E}_s^{\sigma}(f_j)| \leq \frac{\varepsilon}{3}$  by definition of the

strategies  $\tau_{d'+1}, \ldots, \tau_n$  and coefficients  $\beta_{d'+1}, \ldots, \beta_n$ . We obtain, by triangulation, that

$$\left| \sum_{m=1}^{n} \alpha_m \mathbb{E}_s^{\tau_m}(f_j) - \mathbb{E}_s^{\sigma}(f_j) \right| \leq \left| \sum_{m=1}^{n} \alpha_m \mathbb{E}_s^{\tau_m}(f_j) - \sum_{m=d'+1}^{n} \beta_m \mathbb{E}_s^{\tau_m}(f_j) \right| + \left| \sum_{m=d'+1}^{n} \beta_m \mathbb{E}_s^{\tau_m}(f_j) - \mathbb{E}_s^{\sigma}(f_j) \right| \leq \frac{2 \cdot \varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon.$$

This ends the proof that theorem with Assumption 2 implies the theorem without any additional assumption.  $\Box$ 

### 6.3 Bounding the support of mixed strategies

Theorems 6.2 and 6.5 state that it suffices to mix finitely many pure strategies to respectively match or approximate the payoff of any strategy. We provide bounds on the number of pure strategies to mix in this section, in the same vein as Carathéodory's theorem for convex hulls (Theorem 2.1). We show that the expected payoff of a finite-support mixed strategy can be obtained exactly by mixing no more than d+1 strategies that are in its support and that a greater or equal payoff can be obtained by mixing no more than d of these strategies.

Let  $s \in S$ ,  $\sigma_1, \ldots, \sigma_n$  be pure strategies and  $\alpha_1, \ldots, \alpha_n \in [0,1]$  be convex combination coefficients. Let  $\mathbf{q} = (q_j)_{1 \leq j \leq d} = \sum_{m=1}^n \alpha_m \mathbb{E}_s^{\sigma_m}(\bar{f})$ . First, let us discuss the case when  $\mathbf{q} \in \mathbb{R}^d$ , i.e., when all considered pure strategies have a finite expected payoff on all dimensions. In this case, the first bound is direct by Carathéodory's theorem for convex hulls.

For the second bound, we observe that  $\mathbf{q}$  is an element of the compact convex polytope  $D = \mathsf{conv}(\{\mathbb{E}_s^{\sigma_m}(\bar{f}) \mid 1 \leq m \leq n\})$ . In particular,  $\mathbf{q}$  is dominated by an element  $\mathbf{p}$  lying on a proper face of D (i.e., the intersection of D and a hyperplane). For instance, we can consider  $\mathbf{q} + \beta \cdot \mathbf{1}$  for  $\beta = \max\{\gamma \geq 0 \mid \mathbf{q} + \gamma \mathbf{1} \in D\}$ . Because proper faces have dimension no more than d-1, Carathéodory's theorem implies that  $\mathbf{p}$  is a convex combination of no more than d vectors of the form  $\mathbb{E}_s^{\sigma_m}(\bar{f})$ , taken among those lying on the considered proper face.

Assume now that  $\mathbf{q} \notin \mathbb{R}^d$ . In this case, Carathéodory's theorem does not apply directly. The idea is to reduce ourselves to the previous case. For each  $1 \leq j \leq d$ , if  $q_j$  is infinite, there is  $1 \leq m \leq n$  such that  $\mathbb{E}_s^{\sigma_m}(f_m) = q_j$ . For each infinite component of  $\mathbf{q}$ , we fix  $\alpha_m \mathbb{E}_s^{\sigma_m}(\bar{f})$  in the convex combination for one such m. We then obtain the sought bounds from the above; we reason on the real-valued components of  $\mathbf{q}$  in the non-fixed part of the convex combination (after normalising its coefficients to sum to one).

We formalise our theorem and the previous proof sketch below.

**Theorem 6.6.** Assume that  $\bar{f}$  is universally unambiguously integrable. Let  $s \in S$ ,  $\sigma_1$ , ...,  $\sigma_n$  be pure strategies and  $\alpha_1, \ldots, \alpha_n \in [0,1]$  be convex combination coefficients. There exist convex combination coefficients  $\beta_1, \ldots, \beta_n \in [0,1]$  with  $|\{1 \le m \le n \mid \beta_m \ne 0\}| \le d+1$  and convex combination coefficients  $\gamma_1, \ldots, \gamma_n \in [0,1]$  with  $|\{1 \le m \le n \mid \gamma_m \ne 0\}| \le d$  such that

$$\sum_{m=1}^{n} \alpha_m \mathbb{E}_s^{\sigma_m}(\bar{f}) = \sum_{m=1}^{n} \beta_m \mathbb{E}_s^{\sigma_m}(\bar{f}) \le \sum_{m=1}^{n} \gamma_m \mathbb{E}_s^{\sigma_m}(\bar{f}).$$

Proof. For all  $1 \leq m \leq n$ , let  $\mathbf{p}^{(m)} = (p_j^{(m)})_{1 \leq j \leq d} = \mathbb{E}_s^{\sigma_m}(\bar{f})$  and let  $\mathbf{q} = (q_j)_{1 \leq j \leq d} = \sum_{m=1}^n \alpha_m \mathbf{p}^{(m)}$ . We assume that for all  $1 \leq m \leq n$ ,  $\alpha_m \neq 0$ .

First, we assume that  $\mathbf{q} \in \mathbb{R}^d$ . The existence of coefficients  $\beta_1, \ldots, \beta_n$  obeying the required conditions is direct by Carathéodory's theorem for convex hulls (Theorem 2.1). We let  $D = \mathsf{conv}(\{\mathbf{p}^{(m)} \mid 1 \le m \le n\})$ , which is a compact set (see Lemma 4.3).

We define  $\mathbf{p} = \mathbf{q} + \beta \cdot \mathbf{1}$  for  $\beta = \sup\{\gamma \geq 0 \mid \mathbf{q} + \gamma \cdot \mathbf{1} \in D\}$ . We remark that  $\beta$  is a real number. On the one hand,  $0 \in \{\gamma \geq 0 \mid \mathbf{q} + \gamma \cdot \mathbf{1} \in D\}$ , and thus  $\beta \neq -\infty$ . On the other hand, D is bounded, therefore  $\beta \neq +\infty$ . Furthermore,  $\mathbf{p} \in D$ . By convexity of D,  $\mathbf{q} + \gamma \cdot \mathbf{1} \in D$  for all  $0 \leq \gamma < \beta$ . It follows that  $\mathbf{p} \in \mathsf{cl}(D) = D$ .

We have  $\mathbf{q} \leq \mathbf{p}$ . To end the case  $\mathbf{q} \in \mathbb{R}^d$ , we show that there exists a hyperplane H such that  $\mathbf{p}$  is a convex combination of the vectors  $\mathbf{p}^{(m)}$  that lie in H. This suffices, because Carathéodory's theorem then ensures that  $\mathbf{p}$  is a convex combination of no more than d vectors among the  $\mathbf{p}^{(m)}$ .

If D is included in a hyperplane, there is nothing to show. We thus assume that D is not included in a hyperplane and obtain a hyperplane using the supporting hyperplane theorem (Theorem 2.3). By construction,  $\mathbf{p} \notin \operatorname{int}(D) = \operatorname{ri}(D)$ : for all  $\gamma > 0$ ,  $\mathbf{p} + \gamma \cdot \mathbf{1} \notin D$ . Therefore, there exists a linear form  $x^* \colon \mathbb{R}^d \to \mathbb{R}$  such that for all  $\mathbf{v} \in D$ ,  $x^*(\mathbf{v}) \le x^*(\mathbf{p})$ . We claim that  $\mathbf{p}$  is a convex combination of the  $\mathbf{p}^{(m)}$  that lie in the hyperplane  $(x^*)^{-1}(x^*(\mathbf{p}))$ . Write  $\mathbf{p}$  as a convex combination  $\sum_{m=1}^n \alpha_m' \mathbf{p}^{(m)}$ . We observe that for all  $1 \le m \le n$ ,  $\alpha_m' \ne 0$  implies  $x^*(\mathbf{p}^{(m)}) = x^*(\mathbf{p})$ , as otherwise we would obtain that  $x^*(\mathbf{p}) < x^*(\mathbf{p})$ . This ends the proof in the case  $\mathbf{q} \in \mathbb{R}^d$ .

We now assume that some components of  $\mathbf{q}$  are infinite. For convenience of notation, we assume that there is  $1 \leq d' \leq d$  such that, for all  $1 \leq j \leq d'$ ,  $q_j \in \{-\infty, +\infty\}$  and, for all  $d'+1 \leq j \leq d$ ,  $q_j \in \mathbb{R}$ . For all  $1 \leq j \leq d'$ , let  $1 \leq m_j \leq n$  such that  $p_j^{(m_j)} = q_j$ . If d'=d, we have  $\mathbf{q} = \frac{1}{d} \sum_{j=1}^{d} \mathbf{p}^{(m_j)}$ . We now assume that d' < d.

d'=d, we have  $\mathbf{q}=\frac{1}{d}\sum_{j=1}^{d}\mathbf{p}^{(m_{j})}$ . We now assume that d'< d. Let  $I_{\infty}=\{m_{j}\mid 1\leq j\leq d'\}$ ,  $I_{\mathbb{R}}=\{1,\ldots,n\}\setminus I_{\infty}$ . We have  $|I_{\infty}|\leq d'\leq d$ . In particular, if  $I_{\mathbb{R}}$  is empty, the sought result is direct. We assume that  $I_{\mathbb{R}}\neq\emptyset$ . Let  $\alpha_{\mathbb{R}}=\sum_{m\in I_{\mathbb{R}}}\alpha_{m}>0$ , and let  $\operatorname{proj}_{>d'}\colon (\mathbb{R})^{d}\to (\mathbb{R})^{d-d'}$  denote the projection of a vector onto its d-d' last components. We apply the result for vectors in  $\mathbb{R}^{d-d'}$  to  $\operatorname{proj}_{>d'}(\sum_{m\in I_{\mathbb{R}}}\frac{\alpha_{m}}{\alpha_{\mathbb{R}}}\mathbf{p}^{(m)})$  with respect to the payoff function  $(f_{d'+1},\ldots,f_{d})$ . It follows that there exist convex combination coefficients  $(\beta'_{m})_{m\in I_{\mathbb{R}}}$  of which at most d-d'+1 are positive and convex combination coefficients  $(\gamma'_{m})_{m\in I_{\mathbb{R}}}$  of which at most d-d' are positive such that

$$\operatorname{proj}_{>d'}\left(\sum_{m\in I_{\mathbb{R}}}\frac{\alpha_m}{\alpha_{\mathbb{R}}}\mathbf{p}^{(m)}\right) = \sum_{m\in I_{\mathbb{R}}}\beta'_m\operatorname{proj}_{>d'}(\mathbf{p}^{(m)}) \leq \sum_{m\in I_{\mathbb{R}}}\gamma'_m\operatorname{proj}_{>d'}(\mathbf{p}^{(m)}).$$

We conclude by observing that

$$\mathbf{q} = \sum_{m \in I_{\infty}} \alpha_m \mathbf{p}^{(m)} + \alpha_{\mathbb{R}} \cdot \sum_{m \in I_{\mathbb{R}}} \frac{\alpha_m}{\alpha_{\mathbb{R}}} \mathbf{p}^{(m)}$$
$$= \sum_{m \in I_{\infty}} \alpha_m \mathbf{p}^{(m)} + \sum_{m \in I_{\mathbb{R}}} \alpha_{\mathbb{R}} \beta'_m \mathbf{p}^{(m)}$$
$$\leq \sum_{m \in I_{\infty}} \alpha_m \mathbf{p}^{(m)} + \sum_{m \in I_{\mathbb{R}}} \alpha_{\mathbb{R}} \gamma'_m \mathbf{p}^{(m)},$$

i.e., for all  $m \in I_{\infty}$ , we let  $\beta_m = \gamma_m = \alpha_m$ , and for all  $m \in I_{\mathbb{R}}$ , we let  $\beta_m = \alpha_{\mathbb{R}} \cdot \beta_m'$  and  $\gamma_m = \alpha_{\mathbb{R}} \cdot \gamma_m'$ .

We now highlight two corollaries of Theorem 6.6. First, we obtain bounds when dealing with universally integrable payoffs that follow from Theorem 6.2.

Corollary 6.7. Assume that  $\bar{f}$  is universally integrable. Let  $s \in S$ .

- For all  $\mathbf{q} \in \mathsf{Pay}_s(\bar{f})$ ,  $\mathbf{q}$  is a convex combination of at most d+1 elements of  $\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f})$ .
- For all  $\mathbf{q} \in \mathsf{Ach}_s(\bar{f})$ , there exists a convex combination  $\mathbf{p}$  of at most d elements of  $\mathsf{Pay}^{\mathsf{pure}}_s(\bar{f})$  such that  $\mathbf{q} \leq \mathbf{p}$ .

*Proof.* By Theorem 6.2,  $\mathsf{Pay}_s(\bar{f}) = \mathsf{conv}(\mathsf{Pay}_s^{\mathsf{pure}}(\bar{f}))$ . Furthermore, we recall that  $\mathbf{q} \in \mathsf{Ach}_s(\bar{f})$  if and only if there exists a strategy  $\sigma$  such that  $\mathbf{q} \leq \mathbb{E}_s^{\sigma}(\bar{f})$ . We obtain both claims of the corollary directly by Thm. 6.6.

Example 6.3 implies that Corollary 6.7 does not directly extend to universally unambiguously integrable payoffs. Nonetheless, we can identify a class of vectors that can be achieved by mixing no more than d strategies: the vectors in  $\operatorname{int}(\operatorname{Ach}_s(\bar{f}) \cap \mathbb{R}^d)$ . These are vectors  $\mathbf{q} \in \operatorname{Ach}_s(\bar{f}) \cap \mathbb{R}^d$  that can be improved in all dimensions simultaneously, i.e., such that  $\mathbf{q} + \varepsilon \mathbf{1} \in \operatorname{Ach}_s(\bar{f})$  for some  $\varepsilon > 0$ . To prove our result, intuitively, we approximate, via Theorem 6.5, the payoff of a strategy achieving  $\mathbf{q} + \varepsilon \mathbf{1}$  with a finite-support mixed strategy then invoke Theorem 6.6.

Corollary 6.8. Assume that  $\bar{f}$  is universally unambiguously integrable. Let  $s \in S$ . For  $\mathbf{q} \in \operatorname{int}(\operatorname{Ach}_s(\bar{f}) \cap \mathbb{R}^d)$ , there exists a convex combination  $\mathbf{p}$  of at most d elements of  $\operatorname{Pay}_s^{\operatorname{pure}}(\bar{f})$  such that  $\mathbf{q} \leq \mathbf{p}$ .

Proof. Let  $\mathbf{q} = (q_j)_{1 \leq j \leq d} \in \operatorname{int}(\operatorname{Ach}_s(\bar{f}) \cap \mathbb{R}^d)$ . By definition of the interior of a subset of  $\mathbb{R}^d$ , there exists  $\varepsilon > 0$  such that  $\mathbf{q} + \varepsilon \mathbf{1} \in \operatorname{Ach}_s(\bar{f})$ . Therefore, there exists a strategy  $\sigma$  such that, for all  $1 \leq j \leq d$ ,  $q_j < \mathbb{E}^{\sigma}_s(f_j)$ . For all  $1 \leq j \leq d$ , we have  $\mathbb{E}^{\sigma}_s(f_j) \neq -\infty$  because  $\mathbb{E}^{\sigma}_s(f_j) > q_j \in \mathbb{R}$ . Let  $\eta = \min(\{1\} \cup \{\mathbb{E}^{\sigma}_s(f_j) - q_j \mid \mathbb{E}^{\sigma}_s(f_j) \in \mathbb{R}, 1 \leq j \leq d\})$  and  $M = \max(\{1\} \cup \{q_j + 1 \mid \mathbb{E}^{\sigma}_s(f_j) = +\infty, 1 \leq j \leq d\})$ .

Theorem 6.5 implies that there exist pure strategies  $\tau_1, \ldots, \tau_n$  and convex combination coefficients  $\alpha_1, \ldots, \alpha_n \in [0,1]$  such that if  $\mathbb{E}_s^{\sigma}(f_j) = +\infty$ , then  $\sum_{m=1}^n \alpha_m \mathbb{E}_s^{\tau_m}(f_j) \geq M$  and, otherwise,  $\sum_{m=1}^n \alpha_m \mathbb{E}_s^{\tau_m}(f_j) - \mathbb{E}_s^{\sigma}(f_j) \geq -\eta$ . By Theorem 6.6, we can assume that  $n \leq d$ . Let  $\mathbf{p} = (p_j)_{1 \leq j \leq d} = \sum_{m=1}^n \alpha_m \mathbb{E}_s^{\tau_m}(\bar{f})$ .

We show that  $\mathbf{q} \leq \mathbf{p}$ . Let  $1 \leq j \leq d$ . First, assume that  $\mathbb{E}_s^{\sigma}(f_j) = +\infty$ . In this case,  $M \geq q_j$ , and we obtain  $q_j \leq M \leq p_j$ . Second, assume that  $\mathbb{E}_s^{\sigma}(f_j) \in \mathbb{R}$ . In that case, we have  $\eta \leq \mathbb{E}_s^{\sigma}(f_j) - q_j$ , and therefore  $q_j \leq \mathbb{E}_s^{\sigma}(f_j) - \eta \leq p_j$ .

## 7 Continuous payoffs

In this section, we study the topological properties of sets of expected payoff when all payoffs are continuous. The main result of this section applies to payoffs whose square is universally integrable, or *universally square integrable* for short. We prove that the set of expected payoffs and of achievable vectors are closed for such payoff functions. In general, the set of expected payoffs need not be closed when considering (universally unambiguously integrable) continuous multi-dimensional payoffs. Our result for universally square integrable continuous payoffs is a generalisation of [CFW13, Lem. 2] for the special case where all payoffs are discounted-sum payoffs (as defined in Section 2).

The Cauchy-Schwarz inequality (e.g., [Dur19, Thm. 1.5.2.]) guarantees that any universally square integrable payoff is universally integrable. This class subsumes real-valued continuous payoffs because such payoffs are bounded (since the set of plays is compact) and universally integrable continuous shortest-path payoffs (see Appendix C).

We divide this section in three parts. In Section 7.1, we introduce a topology on the space of randomised strategies such that the resulting topological space is metrisable and

compact to formalise a notion of convergence of strategies. In Section 7.2, we formulate our result for continuous universally square integrable payoffs. Section 7.3 provides examples illustrating that the results of Section 7.2 do not hold for continuous payoffs that are not universally integrable. We leave open whether our results extend to all universally integrable continuous payoffs. For this section, we fix an MDP  $\mathfrak{M} = (S, A, \delta)$  and a d-dimensional continuous payoff  $\bar{f} = (f_i)_{1 \le i \le d}$ .

## 7.1 A topology on the space of strategies

In this section, we define a topology on the space of strategies to formalise the convergence of sequences of strategies.

First, we define a topology over  $\mathcal{D}(A(s))$  for all  $s \in S$ . Let  $s \in S$ . The set  $\mathcal{D}(A(s))$  is a compact subset of  $\mathbb{R}^{A(s)}$ . In the sequel, we consider the metric (induced by the Euclidean norm)  $\operatorname{dist_{proba}}(\mu, \mu') = \sqrt{\sum_{a \in A(s)} |\mu(a) - \mu'(a)|^2}$  for all  $\mu, \mu' \in \mathcal{D}(A(s))$  on distributions over actions

We endow the set  $\Sigma(\mathfrak{M}) = \prod_{h \in \mathsf{Hist}(\mathfrak{M})} \mathcal{D}(A(\mathsf{last}(h)))$  of all strategies with the product topology. We obtain that  $\Sigma(\mathfrak{M})$  is a compact metrisable topological space. We do not define a metric over strategies, as it is not necessary in the sequel. Instead, we recall that a sequence of strategies  $(\sigma^{(n)})_{n \in \mathbb{N}}$  converges to a strategy  $\sigma$  if and only if, for all  $h \in \mathsf{Hist}(\mathfrak{M})$ ,  $(\sigma^{(n)}(h))_{n \in \mathbb{N}}$  converges to  $\sigma(h)$ .

We now formulate a result intuitively stating that close strategies induce distributions that assign similar probabilities to cylinders of histories of bounded length. We first require the following technical lemma.

**Lemma 7.1.** Let  $\alpha_1, \ldots, \alpha_n, \beta_1, \ldots, \beta_n \in [0, 1]$ . Then

$$\left| \prod_{m=1}^{n} \alpha_m - \prod_{m=1}^{n} \beta_m \right| \leq \sum_{m=1}^{n} |\alpha_m - \beta_m|.$$

*Proof.* We proceed by induction. The claim trivially holds for n = 1. To lighten notation, we prove the case n = 2 separately first. We perform the general induction step below by building on this simpler case. We have:

$$|\alpha_{1}\alpha_{2} - \beta_{1}\beta_{2}| = \left|\frac{1}{2}(\alpha_{1} - \beta_{1})(\alpha_{2} + \beta_{2}) + \frac{1}{2}(\alpha_{1} + \beta_{1})(\alpha_{2} - \beta_{2})\right|$$

$$\leq |\alpha_{1} - \beta_{1}|\frac{|\alpha_{2} + \beta_{2}|}{2} + |\alpha_{2} - \beta_{2}|\frac{|\alpha_{1} + \beta_{1}|}{2}$$

$$\leq |\alpha_{1} - \beta_{1}| + |\alpha_{2} - \beta_{2}|.$$

The second line is obtained by triangulation and the last line is obtained by using the fact that  $\alpha_1, \alpha_2, \beta_1, \beta_2 \in [0, 1]$ .

We now perform the general induction step. We assume that the result holds for some  $n \in \mathbb{N}_0$  and prove that it holds for n + 1. By applying the simpler case proven above and then the induction hypothesis, we obtain that

$$\left| \prod_{m=1}^{n+1} \alpha_m - \prod_{m=1}^{n+1} \beta_m \right| \le \left| \prod_{m=1}^n \alpha_m - \prod_{m=1}^n \beta_m \right| + |\alpha_{n+1} - \beta_{n+1}|$$

$$\le \sum_{m=1}^{n+1} |\alpha_m - \beta_m|.$$

We now state the lemma regarding induced distributions.

**Lemma 7.2.** Let  $\sigma, \tau$  be two strategies,  $k \in \mathbb{N}_0$  and  $\eta > 0$ . Assume that, for all histories h that are at most k states long,  $\mathsf{dist}_{\mathsf{proba}}(\sigma(h), \tau(h)) \leq \frac{\eta}{k}$ . Then, for all histories h that are at most k+1 states long and all  $s \in S$ , we have  $|\mathbb{P}_s^{\sigma}(\mathsf{Cyl}(h)) - \mathbb{P}_s^{\tau}(\mathsf{Cyl}(h))| \leq \eta$ .

*Proof.* Let  $h = s_0 a_0 s_1 \dots s_r$  with  $r \leq k$ . We only prove the claim for  $s = s_0$ . The other case is direct because, for  $s \in S \setminus \{s_0\}$ , we have  $\mathbb{P}_s^{\sigma}(\mathsf{Cyl}(h)) = \mathbb{P}_s^{\tau}(\mathsf{Cyl}(h)) = 0$ .

The proof follows from the following sequence of inequations.

$$\begin{split} |\mathbb{P}^{\sigma}_{s_0}(\mathsf{Cyl}(h)) - \mathbb{P}^{\tau}_{s_0}(\mathsf{Cyl}(h))| &= \prod_{\ell=0}^{r-1} \delta(s_\ell, a_\ell)(s_{\ell+1}) \cdot \left| \prod_{\ell=0}^{r-1} \sigma(h_{\leq \ell}) - \prod_{\ell=0}^{r-1} \tau(h_{\leq \ell}) \right| \\ &\leq \left| \prod_{\ell=0}^{r-1} \sigma(h_{\leq \ell}) - \prod_{\ell=0}^{r-1} \tau(h_{\leq \ell}) \right| \\ &\leq \sum_{\ell=0}^{r-1} |\sigma(h_{\leq \ell}) - \tau(h_{\leq \ell})| \\ &\leq k \cdot \frac{\eta}{k} = \eta. \end{split}$$

The first line is by definition of probability measures induced by the strategies from  $s_0$ . The second line uses the fact that transition probabilities are at most 1. The third line is obtained by Lemma 7.1. The last line follows from the assumption of the lemma and  $r \leq k$ .

We close this section by showing that  $\Sigma_{\mathsf{pure}}(\mathfrak{M})$  is a closed subset of  $\Sigma(\mathfrak{M})$ . In the following section, we prove that if  $\bar{f}$  is universally square integrable and  $\Sigma \subseteq \Sigma(\mathfrak{M})$  is closed, then  $\mathsf{Pay}_s^{\Sigma}(\bar{f})$  is compact. Therefore, establishing that  $\Sigma_{\mathsf{pure}}(\mathfrak{M})$  is closed implies that the result is also applicable to  $\Sigma_{\mathsf{pure}}(\mathfrak{M})$ . The proof boils down to showing that the limit of a converging sequence of Dirac distributions is a Dirac distribution. This follows from such sequences being ultimately constant.

**Lemma 7.3.** The set  $\Sigma_{\mathsf{pure}}(\mathfrak{M})$  is a closed subset of  $\Sigma(\mathfrak{M})$ .

*Proof.* Let  $(\sigma^{(n)})_{n\in\mathbb{N}}$  be a sequence of pure strategies that converges to a strategy  $\sigma$ , i.e., for all  $h \in \mathsf{Hist}(\mathfrak{M})$ , the sequence  $(\sigma^{(n)}(h))_{n\in\mathbb{N}}$  converges to  $\sigma(h)$ . We must show that, for all  $h \in \mathsf{Hist}(\mathfrak{M})$ ,  $\sigma(h)$  is a Dirac distribution.

Let  $h \in \mathsf{Hist}(\mathfrak{M})$ . Because  $(\sigma^{(n)}(h))_{n \in \mathbb{N}}$  is convergent, it is a Cauchy sequence. Thus, there exists some  $n_0 \in \mathbb{N}$  such that for all  $n, m \geq n_0$ ,  $\mathsf{dist}_{\mathsf{proba}}(\sigma^{(n)}(h), \sigma^{(m)}(h)) < 1$ . It follows from  $(\sigma^{(n)}(h))_{n \in \mathbb{N}}$  being a sequence of a Dirac distributions and the definition of  $\mathsf{dist}_{\mathsf{proba}}$  that the sequence  $(\sigma^{(n)}(h))_{n \geq n_0}$  is constant. It follows that  $\sigma(h) = \sigma^{(n_0)}(h)$ , and thus  $\sigma(h)$  is a Dirac distribution.

#### 7.2 Universally square integrable continuous payoffs

We prove that if  $\bar{f}$  is universally square integrable, then for all  $s \in S$ ,  $\operatorname{Pay}_s(\bar{f})$  and  $\operatorname{Ach}_s(\bar{f})$  are closed. Our proof relies on the following property: if  $\bar{f}$  is universally square integrable, then for all sequences  $(\sigma^{(n)})_{n \in \mathbb{N}}$  of strategies converging to a strategy  $\sigma$ , and for all  $s \in S$ ,  $(\mathbb{E}_s^{\sigma^{(n)}}(\bar{f}))_{n \in \mathbb{N}}$  converges to  $\mathbb{E}_s^{\sigma}(\bar{f})$ . It suffices to prove the convergence on each dimension to obtain this property. Therefore, we need only consider one-dimensional payoffs for now.

We split the proof into two parts: first, we consider the particular case of real-valued continuous payoffs and then generalise to universally square integrable payoffs. We remark

that we may not assume that a universally integrable continuous payoff is real-valued without loss of generality: changing the payoff of plays that have an infinite payoff to a real number will violate the continuity property. Therefore, we do not generalise the property for real-valued continuous payoffs through such an assumption.

Let  $f \colon \mathsf{Plays}(\mathfrak{M}) \to \mathbb{R}$  be a continuous real-valued payoff. The proof in this case relies on the uniform continuity of f. Recall that f is uniformly continuous if for all  $\varepsilon > 0$ , there exists  $\ell \in \mathbb{N}$  such that for all plays  $\pi$ ,  $\pi'$ ,  $\pi_{\leq \ell} = \pi'_{\leq \ell}$  implies that  $|f(\pi) - f(\pi')| < \varepsilon$ . In particular, for all  $\varepsilon > 0$ , f can be  $\varepsilon$ -approximated by a linear combination of indicator functions of cylinders of histories of a fixed length. This provides a means to  $\varepsilon$ -approximate  $\mathbb{E}^{\tau}_{s}(f)$  for any strategy  $\tau$  as a linear combination of probabilities of cylinders of histories of bounded length. Since Lemma 7.2 guarantees that the distributions over plays induced by strategies that are intuitively close assign similar probabilities to such cylinders, through the approximations above, we can obtain that  $\mathbb{E}^{\sigma^{(n)}}_{s}(\bar{f})$  is  $3\varepsilon$ -close  $\mathbb{E}^{\sigma}_{s}(\bar{f})$  for all  $s \in S$  for large values of n. We formalise this argument below.

**Theorem 7.4.** Let  $s \in S$ . Assume that  $\bar{f} : \mathsf{Plays}(\mathfrak{M}) \to \mathbb{R}^d$  and that  $\bar{f}$  is continuous. Then the function  $\Sigma(\mathfrak{M}) \to \mathbb{R}^d : \sigma \mapsto \mathbb{E}^{\sigma}_s(\bar{f})$  is continuous. In other words, for all sequences  $(\sigma^{(n)})_{n \in \mathbb{N}}$  of strategies that converge to a strategy  $\sigma$ ,  $\lim_{n \to \infty} \mathbb{E}^{\sigma^{(n)}}_s(\bar{f}) = \mathbb{E}^{\sigma}_s(\bar{f})$ .

*Proof.* It suffices to prove the theorem in the case d=1 to obtain the general case. For this reason, we consider a one-dimensional real-valued continuous payoff f below. Let  $(\sigma^{(n)})_{n\in\mathbb{N}}$  be a sequence of strategies converging to a strategy  $\sigma$ . We start with some notation. By continuity of f, f is bounded. We let  $||f||_{\infty} = \sup_{\pi \in \mathsf{Plays}(\mathfrak{M})} |f(\pi)|$ . For any history  $h = s_0 a_0 \dots s_r$ , we let |h| = r denote the index of the last state of the history. We also fix, for all histories  $h \in \mathsf{Hist}(\mathfrak{M})$ , a play  $\pi(h) \in \mathsf{Cyl}(h)$  which is a continuation of h.

We must prove that  $(\mathbb{E}_s^{\sigma^{(n)}}(f))_{n\in\mathbb{N}}$  converges to  $\mathbb{E}_s^{\sigma}(f)$ . Let  $\varepsilon > 0$ . We assume that  $||f||_{\infty} > 0$ , as otherwise the result is direct: if  $||f||_{\infty} = 0$ , then  $\mathbb{E}_s^{\sigma^{(n)}}(f) = \mathbb{E}_s^{\sigma}(f) = 0$  for all  $n \in \mathbb{N}$ .

We start by constructing a simple function which  $\frac{\varepsilon}{3}$ -approximates f by exploiting the uniform continuity of f. By uniform continuity of f, there exists some  $k \in \mathbb{N}_0$  such that, for any two plays  $\pi, \pi' \in \mathsf{Plays}(\mathfrak{M})$ , if  $\pi_{\leq k} = \pi'_{\leq k}$ , then  $|f(\pi) - f(\pi')| \leq \frac{\varepsilon}{3}$ . It follows that

$$\left| f - \sum_{|h| = k} f(\pi(h)) \cdot \mathbb{1}_{\operatorname{Cyl}(h)} \right| \leq \frac{\varepsilon}{3}.$$

Since f is universally integrable, it follows that, for all  $\tau \in \Sigma(\mathfrak{M})$ ,

$$\left| \mathbb{E}_{s}^{\tau}(f) - \sum_{|h|=k} f(\pi(h)) \cdot \mathbb{P}_{s}^{\tau}(\mathsf{Cyl}(h)) \right| \leq \frac{\varepsilon}{3}. \tag{7.1}$$

To end the argument, we now determine  $n_0$  such that, for all  $n \geq n_0$ , we have

$$\left| \sum_{|h|=k} f(\pi(h)) \cdot \left( \mathbb{P}_s^{\sigma}(\mathsf{Cyl}(h)) - \mathbb{P}_s^{\sigma^{(n)}}(\mathsf{Cyl}(h)) \right) \right| \le \frac{\varepsilon}{3}. \tag{7.2}$$

Let M denote the number of histories h such that |h|=k. Since  $\lim_{n\to\infty}\sigma^{(n)}=\sigma$ , there exists  $n_0\in\mathbb{N}$  such that for all histories h such that  $|h|\leq k$  (i.e., with at most k+1 states), we have  $\operatorname{dist}_{\mathsf{proba}}(\sigma^{(n)}(h),\sigma(h))\leq \frac{\varepsilon}{3\cdot M\cdot\|f\|_{\infty}\cdot k}$  (here, we use the fact that there are

finitely many such histories). Equation (7.2) follows from the triangular inequality and Lemma 7.2: we have, for all  $n \ge n_0$ ,

$$\begin{split} \left| \sum_{|h|=k} f(\pi(h)) \cdot \left( \mathbb{P}^{\sigma}_{s}(\mathsf{Cyl}(h)) - \mathbb{P}^{\sigma^{(n)}}_{s}(\mathsf{Cyl}(h)) \right) \right| &\leq \|f\|_{\infty} \cdot \sum_{|h|=k} \left| \mathbb{P}^{\sigma}_{s}(\mathsf{Cyl}(h)) - \mathbb{P}^{\sigma^{(n)}}_{s}(\mathsf{Cyl}(h)) \right| \\ &\leq \|f\|_{\infty} \cdot \sum_{|h|=k} \frac{\varepsilon}{3 \cdot M \cdot \|f\|_{\infty}} \\ &\leq \frac{\varepsilon}{3}. \end{split}$$

Let  $n \geq n_0$ . We now show that  $|\mathbb{E}_s^{\sigma}(f) - \mathbb{E}_s^{\sigma_n}(f)| \leq \varepsilon$ . From the triangular inequality, Equation (7.1) and Equation (7.2), we obtain

$$\begin{split} \left| \mathbb{E}_{s}^{\sigma}(f) - \mathbb{E}_{s}^{\sigma^{(n)}}(f) \right| &\leq \left| \mathbb{E}_{s}^{\sigma}(f) - \sum_{|h| = k} f(\pi(h)) \cdot \mathbb{P}_{s}^{\sigma}(\mathsf{Cyl}(h)) \right| \\ &+ \left| \sum_{|h| = k} f(\pi(h)) \cdot \left( \mathbb{P}_{s}^{\sigma}(\mathsf{Cyl}(h)) - \mathbb{P}_{s}^{\sigma^{(n)}}(\mathsf{Cyl}(h)) \right) \right| \\ &+ \left| \sum_{|h| = k} f(\pi(h)) \cdot \mathbb{P}_{s}^{\sigma^{(n)}}(\mathsf{Cyl}(h)) - \mathbb{E}_{s}^{\sigma^{(n)}}(f) \right| \\ &\leq \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon. \end{split}$$

We have thus shown that  $(\mathbb{E}_s^{\sigma^{(n)}}(f))_{n\in\mathbb{N}}$  converges to  $\mathbb{E}_s^{\sigma}(f)$ .

We now generalise Theorem 7.4 to universally square integrable payoffs. We note that the previous proof relies on the boundedness of continuous payoffs and their uniform continuity, and thus is not valid in this more general case.

Similarly to above, it suffices to consider one-dimensional non-negative payoffs; the general case can be recovered by writing payoffs on each dimension as the difference of their non-negative and non-positive parts. The non-negative and non-positive parts of a continuous payoff are also continuous (see Lemma A.1 of Appendix A.1). We let  $f: \mathsf{Plays}(\mathfrak{M}) \to \bar{\mathbb{R}}$  be a non-negative continuous square integrable payoff,  $s \in S$ , let  $(\sigma^{(n)})_{n \in \mathbb{N}}$  be a sequence of strategies that converges to a strategy  $\sigma$  and  $\varepsilon > 0$ . Given  $M \in \mathbb{R}$ , we abbreviate  $\{f(\pi) \geq M \mid \pi \in \mathsf{Plays}(\mathfrak{M})\}$  by  $\{f \geq M\}$ , for all strategies  $\tau$ , we write  $\mathbb{P}^{\tau}_{s}(f \geq M)$  instead of  $\mathbb{P}^{\tau}_{s}(\{f \geq M\})$  and we let  $\min(f, M)$  denote the payoff  $\pi \mapsto \min\{f(\pi), M\}$ .

The goal is to show that  $|\mathbb{E}_s^{\sigma^{(n)}}(f) - \mathbb{E}_s^{\sigma}(f))| \leq \varepsilon$  for all large enough values of n. We show that there exists a constant  $M \geq 0$  (dependent on  $\varepsilon$ ) such that the real-valued continuous payoff  $\min(f, M)$  (whose continuity follows from Lemma A.1) satisfies  $0 \leq \mathbb{E}_s^{\tau}(f) - \mathbb{E}_s^{\tau}(\min(f, M)) \leq \frac{\varepsilon}{3}$  for all strategies  $\tau \in \Sigma(\mathfrak{M})$ . By the triangular inequality,  $|\mathbb{E}_s^{\sigma^{(n)}}(f) - \mathbb{E}_s^{\sigma}(f))|$  is no more than

$$|\mathbb{E}_{s}^{\sigma^{(n)}}(f) - \mathbb{E}_{s}^{\sigma^{(n)}}(\min(f, M))| + |\mathbb{E}_{s}^{\sigma^{(n)}}(\min(f, M)) - \mathbb{E}_{s}^{\sigma}(\min(f, M))| + |\mathbb{E}_{s}^{\sigma}(\min(f, M)) - \mathbb{E}_{s}^{\sigma}(f)|.$$

The first and last terms are no more than  $\frac{\varepsilon}{3}$  by choice of M, and the second term is smaller than  $\frac{\varepsilon}{3}$  for large values of n by Theorem 7.4.

Thus, the main hurdle of the proof is establishing the existence of a suitable M. The first step consists in showing that the probability of f being large can be made arbitrarily small, in the sense that for all  $\eta > 0$ , there exists  $M(\eta)$  such that  $\mathbb{P}_s^{\tau}(f \geq M(\eta)) \leq \eta$  for all strategies  $\tau$ . The negation of this property would allow the existence of strategies with an arbitrarily large expected payoff from s, contradicting Lemma 3.2. It then follows from the Cauchy-Schwarz inequality that, for all strategies  $\tau$ ,  $\mathbb{E}_s^{\tau}(f - \min(f, M)) \leq \sqrt{\mathbb{E}_s^{\tau}(f^2) \cdot \eta}$  because  $f - \min(f, M) \leq f \cdot \mathbb{1}_{f \geq M}$ . Because  $f^2$  is universally integrable, Lemma 3.2 guarantees that  $\sup_{\tau} \mathbb{E}_s^{\tau}(f^2)$  is finite, i.e., we obtain the desired inequality by choosing  $\eta$  small enough.

We provide the details of the above sketch in the following proof.

**Theorem 7.5.** Let  $s \in S$ . Assume that  $\bar{f}$  is continuous and universally square integrable. Then the function  $\Sigma(\mathfrak{M}) \to \mathbb{R}^d \colon \sigma \mapsto \mathbb{E}^{\sigma}_s(\bar{f})$  is continuous. In other words, for all sequences  $(\sigma^{(n)})_{n \in \mathbb{N}}$  of strategies that converge to a strategy  $\sigma$ ,  $\lim_{n \to \infty} \mathbb{E}^{\sigma^{(n)}}_s(\bar{f}) = \mathbb{E}^{\sigma}_s(\bar{f})$ .

*Proof.* It suffices to prove the theorem in the case d=1 to obtain the general case. For this reason, we consider a one-dimensional real-valued continuous payoff f. Let  $(\sigma^{(n)})_{n\in\mathbb{N}}$  be a sequence of strategies that converges to a strategy  $\sigma$ . We assume that f is non-negative. We show that this implies the general case at the end of the proof.

We show that  $\lim_{n\to\infty}(\mathbb{E}_s^{\sigma(n)}(\bar{f}))_{n\in\mathbb{N}}=\mathbb{E}_s^{\sigma}(\bar{f})$  by the standard definition of convergence. Let  $\varepsilon>0$ . Our goal is to determine some  $M\geq 0$  and some  $n_0$  such that for all  $n\geq n_0$ , we have

$$|\mathbb{E}_{s}^{\sigma^{(n)}}(f) - \mathbb{E}_{s}^{\sigma^{(n)}}(\min(f, M))| + |\mathbb{E}_{s}^{\sigma^{(n)}}(\min(f, M)) - \mathbb{E}_{s}^{\sigma}(\min(f, M))| + |\mathbb{E}_{s}^{\sigma}(\min(f, M)) - \mathbb{E}_{s}^{\sigma}(f)| \le \varepsilon$$

This is sufficient: the sum highlighted above is greater or equal to  $|\mathbb{E}_s^{\sigma^{(n)}}(f) - \mathbb{E}_s^{\sigma}(f)|$  by the triangular inequality. We bound each term of the above sum by  $\frac{\varepsilon}{3}$  in the following.

The crux of the proof is determining a bound  $M \geq 0$  such that for all strategies  $\tau$ ,  $\mathbb{E}_s^{\tau}(f) - \mathbb{E}_s^{\tau}(\min(f,M)) < \frac{\varepsilon}{3}$ . To establish this, we show the following property: for all  $\eta > 0$ , there exists a bound  $M(\eta) \geq 0$  such that, for all strategies  $\tau$ ,  $\mathbb{P}_s^{\tau}(f \geq M(\eta)) \leq \eta$ . This last property is shown by contradiction. Assume that there exists some  $\eta > 0$  such that for all  $M \geq 0$ , there exists a strategy  $\tau_M$  such that  $\mathbb{P}_s^{\tau_M}(f \geq M) > \eta$ . Then, we obtain that for all  $M \geq 0$ ,  $\mathbb{E}_s^{\tau_M}(f) \geq \mathbb{E}_s^{\tau_M}(M \cdot \mathbb{1}_{\{f \geq M\}}) = M \cdot \mathbb{P}_s^{\tau_M}(f \geq M) \geq \eta \cdot M$ . This contradicts the fact that f is universally integrable (Lemma 3.2).

Let  $\alpha = 1 + \sup_{\tau \in \Sigma(\mathfrak{M})} \mathbb{E}_s^{\tau}(f^2) > 0$ . The value  $\alpha$  is real by universal square integrability of f (Lemma 3.2). For the remainder of the proof, we fix  $M \geq 0$  such that for all strategies  $\tau$ , we have  $\mathbb{P}_s^{\tau}(f \geq M) \leq \frac{\varepsilon^2}{9 \cdot \alpha}$ . We prove that M is the bound sought above. First, we observe that  $f - \min(f, M) = (f - M) \cdot \mathbb{1}_{\{f \geq M\}} \leq f \cdot \mathbb{1}_{\{f \geq M\}}$ . Because indicators are equal to their square, they are universally square integrable. By applying the Cauchy-Schwarz inequality, we obtain that, for all strategies  $\tau$ ,

$$\mathbb{E}_{s}^{\tau}(f - \min(f, M)) \leq \mathbb{E}_{s}^{\tau}(f \cdot \mathbb{1}_{\{f \geq M\}}) \leq \sqrt{\mathbb{E}_{s}^{\tau}(f^{2}) \cdot \mathbb{E}_{s}^{\tau}(\mathbb{1}_{\{f \geq M\}})} \leq \frac{\varepsilon}{3}.$$

To conclude (for non-negative payoffs), it remains to show that there exists  $n_0$  such that for all  $n \geq n_0$ , we have  $|\mathbb{E}_s^{\sigma^{(n)}}(\min(f,M)) - \mathbb{E}_s^{\sigma}(\min(f,M))| \leq \frac{\varepsilon}{3}$ . To this end, we observe that the payoff  $\min(f,M)$  is continuous (Lemma A.1). Theorem 7.4 then implies that a suitable  $n_0$  exists  $(\min(f,M)$  is real-valued).

We have shown that the theorem holds for non-negative continuous universally square integrable payoffs. We now generalise to arbitrary continuous universally square integrable

payoffs. Let  $f^+ = \max(f,0)$  and  $f^- = \max(-f,0)$  denote the non-negative and non-positive parts of f. These payoffs are continuous by Lemma A.1. We observe that  $f^2 = (f^+)^2 + (f^-)^2$ , and, in particular,  $f^2 \ge (f^+)^2 + (f^-)^2$ . It follows that  $(f^+)^2$  and  $(f^-)^2$  are universally integrable. We obtain the claim of the theorem from the above and the following sequence of equations:

$$\lim_{n \to \infty} \mathbb{E}_s^{\sigma^{(n)}}(f) = \lim_{n \to \infty} \mathbb{E}_s^{\sigma^{(n)}}(f^+) - \lim_{n \to \infty} \mathbb{E}_s^{\sigma^{(n)}}(f^-)$$
$$= \mathbb{E}_s^{\sigma}(f^+) - \mathbb{E}_s^{\sigma}(f^-)$$
$$= \mathbb{E}_s^{\sigma}(f).$$

We now prove that for multi-dimensional universally square integrable payoffs that are continuous, given a closed set of strategies, its set of expected payoffs and achievable set from a state are compact. For sets of expected payoffs, this follows from Theorem 7.5: since the image of compact set by a continuous function is compact, the result is direct. For achievable sets, it follows from the property that the downward-closure of any compact set is compact.

**Theorem 7.6.** Let  $s \in S$  and  $\Sigma \subseteq \Sigma(\mathfrak{M})$  be a closed set of strategies. Assume that  $\bar{f}$  is a continuous universally square integrable payoff. Then  $\mathsf{Pay}^\Sigma_s(\bar{f})$  and  $\mathsf{Ach}^\Sigma_s(\bar{f})$  are compact subsets of  $\mathbb{R}^d$  and  $\bar{\mathbb{R}}^d$  respectively. In particular,  $\mathsf{Pay}_s(\bar{f})$ ,  $\mathsf{Ach}_s(\bar{f})$ ,  $\mathsf{Pay}^\mathsf{pure}_s(\bar{f})$  and  $\mathsf{Ach}^\mathsf{pure}_s(\bar{f})$  are compact.

*Proof.* Since  $\Sigma$  is closed and  $\Sigma(\mathfrak{M})$  is compact, we obtain that  $\Sigma$  is compact. It follows from Theorem 7.4 that  $\mathsf{Pay}_s^\Sigma(\bar{f})$  is the image of  $\Sigma$  by a continuous function, and thus is compact. The claim regarding  $\mathsf{Pay}_s^\mathsf{pure}(\bar{f})$  follows from Lemma 7.3. The claims related to achievable sets follow from the property that for all compact  $D \subseteq \bar{\mathbb{R}}^d$ ,  $\mathsf{down}(D)$  is compact (see Lemma A.4 of Appendix A.2 for a proof) and the fact that  $\mathsf{Ach}_s^\Sigma(\bar{f}) = \mathsf{down}(\mathsf{Pay}_s^\Sigma(\bar{f}))$  by definition.


Theorem 7.6 provides a sufficient condition on payoffs which guarantees that we can dominate any expected payoff by a Pareto-optimal expected payoff. This property does not hold for all universally integrable payoffs in full generality, e.g., in the one-dimensional setting, optimal strategies need not exist.


### 7.3 Non-universally integrable continuous payoffs

In this section, we present counterexamples to Theorems 7.5 and 7.6 when the assumption of universal square integrability is relaxed. The MDPs used in these counterexamples are depicted in Figure 7.1.

For the first example, we provide an MDP with a shortest-path payoff witnessing that Theorem 7.5 does not generalise to shortest-path payoffs that are not universally (square) integrable, even when only considering pure strategies and when  $\mathsf{Pay}_s(\bar{f})$  is closed.

**Example 7.1.** We consider the MDP  $\mathfrak{M}$  depicted in Figure 7.1a, the constant weight function w=1 and the shortest-path function  $\mathsf{SPath}_w^{\{t\}}$ . We provide a sequence of pure strategies  $(\sigma^{(n)})_{n\in\mathbb{N}}$  converging to a pure strategy  $\sigma$  such that  $\lim_{n\to\infty} \mathbb{E}_s^{\sigma^{(n)}}(\mathsf{SPath}_w^{\{t\}}) \neq \mathbb{E}_s^{\sigma}(\mathsf{SPath}_w^{\{t\}})$ . For all  $n\in\mathbb{N}$ , we define  $\sigma^{(n)}$  as the pure strategy such that, for all  $h\in\mathsf{Hist}(\mathfrak{M})$ ,  $\sigma^{(n)}(h)=b$  if  $\mathsf{last}(h)=s$  and there are at least n actions in h, and otherwise,  $\sigma^{(n)}(h)=a$ . Intuitively, when starting in s,  $\sigma^{(n)}$  uses action a for the first n rounds of the





(a) An MDP with one randomised transition. Weights are omitted and are all 1.

(b) An MDP with two-dimensional weights.

Figure 7.1: MDPs for counter-examples to Theorems 7.5 (left) and 7.6 (right) without the universally square integrable assumption.

play and then uses b for all subsequent rounds. The pure memoryless strategy  $\sigma$  assigning action a to all states is easily checked to be  $\lim_{n\to\infty} \sigma^{(n)}$ .

Let  $n \in \mathbb{N}$ . Let  $\pi_n$  denote the play  $(sa)^n(sb)^\omega$ . We have  $\mathbb{P}_s^{\sigma^{(n)}}(\{\pi_n\}) = \frac{1}{2^n}$ . Indeed, for all  $r \in \mathbb{N}$ , the definition of the probability distribution over plays under a strategy implies that

$$\mathbb{P}_s^{\sigma^{(n)}}(\mathsf{Cyl}((sa)^n(sb)^rs)) = \mathbb{P}_s^{\sigma^{(n)}}(\mathsf{Cyl}((sa)^ns)) = \frac{1}{2^n}.$$

It follows from  $\mathsf{SPath}_w^{\{t\}}(\pi_n) = +\infty$  that  $\mathbb{E}_s^{\sigma^{(n)}}(\mathsf{SPath}_w^{\{t\}}) = +\infty$ . We conclude that  $\lim_{n\to\infty} \mathbb{E}_s^{\sigma^{(n)}}(\mathsf{SPath}_w^{\{t\}}) = +\infty$ .

We now show that  $\mathbb{E}_s^{\sigma}(\mathsf{SPath}_w^{\{t\}}) \in \mathbb{R}$ . First, we observe that  $\mathbb{P}_s^{\sigma}(\mathsf{Reach}(\{t\})) = 1$ . Therefore,  $\mathbb{E}_s^{\sigma}(\mathsf{SPath}_w^{\{t\}})$  is the unique solution of the equation  $x = 1 + \frac{1}{2}x$  (see, e.g., [BK08]), i.e.,  $\mathbb{E}_s^{\sigma}(\mathsf{SPath}_w^{\{t\}}) = 2 \in \mathbb{R}$ . We have shown that  $\lim_{n \to \infty} \mathbb{E}_s^{\sigma^{(n)}}(\mathsf{SPath}_w^{\{t\}}) \neq \mathbb{E}_s^{\sigma}(\mathsf{SPath}_w^{\{t\}})$ .

We now show that  $\mathsf{Pay}_s(\mathsf{SPath}_w^{\{t\}})$  is closed. The memoryless strategy playing a is optimal when adopting a minimisation point of view. Furthermore, there exist strategies with arbitrarily large but finite expected payoffs from s. For instance, for all  $n \in \mathbb{N}$ , the strategy that plays b for the first n rounds in s and then only uses a after ensures a finite payoff greater than n. By convexity of  $\mathsf{Pay}_s(\mathsf{SPath}_w^{\{t\}}) \cap \mathbb{R}$ , it follows that  $\mathsf{Pay}_s(\mathsf{SPath}_w^{\{t\}}) = [2, +\infty]$  and thus is closed.

We now present a counter-example to Theorem 7.6 when the considered payoffs are not universally integrable: we provide a continuous payoff such that its set of expected payoffs and achievable sets from a given state are not closed.

**Example 7.2.** We consider the MDP  $\mathfrak{M}$  and the weight function  $w=(w_1,w_2)$  depicted in Figure 7.1b. Let  $\bar{f}=(f_1,f_2)$  be the two-dimensional payoff such that  $f_1=\mathsf{DSum}_{w_1}^{1/2}$  and  $f_2=\mathsf{SPath}_{w_2}^{\{t\}}$ . To show that  $\mathsf{Pay}_s(\bar{f})$  is not closed, we show that  $(2,+\infty)\in\mathsf{cl}(\mathsf{Pay}_s(\bar{f}))\setminus\mathsf{Pay}_s(\bar{f})$ .

First, we show that  $(2, +\infty) \in \mathsf{cl}(\mathsf{Pay}_s(\bar{f}))$ . We have  $\bar{f}((sa)^\omega) = (0, \infty)$  and  $\bar{f}(s(bt)^\omega) = (2, 1)$ . By convexity of  $\mathsf{Pay}_s(\bar{f})$ , we obtain that for all  $\eta \in ]0, 1[, (2\eta, +\infty) \in \mathsf{Pay}_s(\bar{f})$ . It follows that  $(2, +\infty) \in \mathsf{cl}(\mathsf{Pay}_s(\bar{f}))$ .

Next, we argue that  $(2,+\infty) \notin \mathsf{Pay}_s(\bar{f})$ . We observe that the only play  $\pi$  from s such that  $f_1(\pi) = 2$  is the play  $s(bt)^\omega$ . All other plays  $\pi'$  starting in s are such that  $f_1(\pi') < 2$ . Indeed, for all  $\ell \in \mathbb{N}$ , we have  $f_1((sa)^\ell s(bt)^\omega) = \sum_{\ell' \geq \ell} \frac{1}{2^{\ell'}} = \frac{1}{2^{\ell-1}} < 2$ . It follows that, to obtain a payoff of 2 on the first dimension from s, we require a strategy whose only outcome is  $s(bt)^\omega$ . This implies that  $(2,+\infty) \notin \mathsf{Pay}_s(\bar{f})$ . We have shown that  $\mathsf{Pay}_s(\bar{f})$  is not closed.

## References

- [ACK<sup>+</sup>20] Pranav Ashok, Krishnendu Chatterjee, Jan Kretínský, Maximilian Weininger, and Tobias Winkler. Approximating values of generalized-reachability stochastic games. In Holger Hermanns, Lijun Zhang, Naoki Kobayashi, and Dale Miller, editors, *Proceedings of the 35th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2020, Saarbrücken, Germany, July 8–11, 2020*, pages 102–115. ACM, 2020.
- [Aum64] Robert J. Aumann. Mixed and behavior strategies in infinite extensive games. In Melvin Dresher, Lloyd S. Shapley, and Albert William Tucker, editors, Advances in Game Theory. (AM-52), Volume 52, pages 627–650. Princeton University Press, 1964.
- [BBC<sup>+</sup>14] Tomáš Brázdil, Václav Brožek, Krishnendu Chatterjee, Vojtěch Forejt, and Antonín Kučera. Markov decision processes with multiple long-run average objectives. *Logical Methods in Computer Science*, Volume 10, Issue 1, Feb 2014.
- [BCM<sup>+</sup>23] Damien Busatto-Gaston, Debraj Chakraborty, Anirban Majumdar, Sayan Mukherjee, Guillermo A. Pérez, and Jean-François Raskin. Bi-objective lexicographic optimization in Markov decision processes with related objectives. In Étienne André and Jun Sun, editors, Proceedings (Part I) of the 21st Interval Symposium on Automated Technology for Verification and Analysis, ATVA 2023, Singapore, October 24–27, 2023, volume 14215 of Lecture Notes in Computer Science, pages 203–223. Springer, 2023.
- [BDOR20] Thomas Brihaye, Florent Delgrange, Youssouf Oualhadj, and Mickael Randour. Life is random, time is not: Markov decision processes with window objectives. *Logical Methods in Computer Science*, 16(4), 2020.
- [BFRR17] Véronique Bruyère, Emmanuel Filiot, Mickael Randour, and Jean-François Raskin. Meet your expectations with guarantees: Beyond worst-case synthesis in quantitative games. *Information and Computation*, 254:259–295, 2017.
- [BGMR23] Thomas Brihaye, Aline Goeminne, James C. A. Main, and Mickael Randour. Reachability games and friends: A journey through the lens of memory and complexity (invited talk). In Patricia Bouyer and Srikanth Srinivasan, editors, Proceedings of the 43rd IARCS Annual Conference on Foundations of Software Technology and Theoretical Computer Science, FSTTCS 2023, IIIT Hyderabad, Telangana, India, December 18–20, 2023, volume 284 of LIPIcs, pages 1:1–1:26. Schloss Dagstuhl Leibniz-Zentrum für Informatik, 2023.
- [BHRR19] Véronique Bruyère, Quentin Hautem, Mickael Randour, and Jean-François Raskin. Energy mean-payoff games. In Wan J. Fokkink and Rob van Glabbeek, editors, *Proceedings of the 30th International Conference on Concurrency Theory, CONCUR 2019, Amsterdam, the Netherlands, August 27–30, 2019*, volume 140 of *LIPIcs*, pages 21:1–21:17. Schloss Dagstuhl Leibniz-Zentrum für Informatik, 2019.
- [Bie87] Kark-Josef Bierth. An expected average reward criterion. Stochastic processes and their applications, 26:123–140, 1987.

- [BK08] Christel Baier and Joost-Pieter Katoen. *Principles of model checking*. MIT Press, 2008.
- [BLO<sup>+</sup>22] Patricia Bouyer, Stéphane Le Roux, Youssouf Oualhadj, Mickael Randour, and Pierre Vandenhove. Games where you can play optimally with arenaindependent finite memory. *Logical Methods in Computer Science*, 18(1), 2022.
- [BMR<sup>+</sup>18] Patricia Bouyer, Nicolas Markey, Mickael Randour, Kim G. Larsen, and Simon Laursen. Average-energy games. *Acta Informatica*, 55(2):91–127, 2018.
- [BORV23] Patricia Bouyer, Youssouf Oualhadj, Mickael Randour, and Pierre Vandenhove. Arena-independent finite-memory determinacy in stochastic games. *Log. Methods Comput. Sci.*, 19(4), 2023.
- [BRV23] Patricia Bouyer, Mickael Randour, and Pierre Vandenhove. Characterizing omega-regularity through finite-memory determinacy of games on infinite graphs. *TheoretiCS*, 2, 2023.
- [CDGH15] Krishnendu Chatterjee, Laurent Doyen, Hugo Gimbert, and Thomas A. Henzinger. Randomness for free. *Information and Computation*, 245:3–16, 2015.
- [CdH04] Krishnendu Chatterjee, Luca de Alfaro, and Thomas A. Henzinger. Trading memory for randomness. In *Proceedings of the 1st International Conference on Quantitative Evaluation of Systems, QEST 2004, Enschede, The Netherlands, 27–30 September 2004*, pages 206–217. IEEE Computer Society, 2004.
- [CFK<sup>+</sup>13] Taolue Chen, Vojtech Forejt, Marta Z. Kwiatkowska, Aistis Simaitis, and Clemens Wiltsche. On stochastic games with multiple objectives. In Krishnendu Chatterjee and Jirí Sgall, editors, *Proceedings of the 38th International Symposium on Mathematical Foundations of Computer Science, MFCS 2013, Klosterneuburg, Austria, August 26–30, 2013*, volume 8087 of *Lecture Notes in Computer Science*, pages 266–277. Springer, 2013.
- [CFW13] Krishnendu Chatterjee, Vojtech Forejt, and Dominik Wojtczak. Multiobjective discounted reward verification in graphs and MDPs. In Kenneth L. McMillan, Aart Middeldorp, and Andrei Voronkov, editors, Proceedings of the 19th International Conference on Logic for Programming, Artificial Intelligence, and Reasoning, LPAR-19, Stellenbosch, South Africa, December 14– 19, 2013, volume 8312 of Lecture Notes in Computer Science, pages 228–242. Springer, 2013.
- [CJH04] Krishnendu Chatterjee, Marcin Jurdzinski, and Thomas A. Henzinger. Quantitative stochastic parity games. In J. Ian Munro, editor, *Proceedings of the Fifteenth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2004, New Orleans, Louisiana, USA, January 11–14, 2004*, pages 121–130. SIAM, 2004.
- [CKK17] Krishnendu Chatterjee, Zuzana Kretínská, and Jan Kretínský. Unifying two views on multiple mean-payoff objectives in Markov decision processes. *Logical Methods in Computer Science*, 13(2), 2017.
- [CKM<sup>+</sup>23] Krishnendu Chatterjee, Joost-Pieter Katoen, Stefanie Mohr, Maximilian Weininger, and Tobias Winkler. Stochastic games with lexicographic objectives. Formal methods in system design, pages 1–41, 2023.

- [CO23] Antonio Casares and Pierre Ohlmann. Characterising memory in infinite games. In Kousha Etessami, Uriel Feige, and Gabriele Puppis, editors, Proceedings of the 50th International Colloquium on Automata, Languages, and Programming, ICALP 2023, July 10–14, 2023, Paderborn, Germany, volume 261 of LIPIcs, pages 122:1–122:18. Schloss Dagstuhl Leibniz-Zentrum für Informatik, 2023.
- [CRR14] Krishnendu Chatterjee, Mickael Randour, and Jean-François Raskin. Strategy synthesis for multi-dimensional quantitative objectives. *Acta Informatica*, 51(3-4):129–163, 2014.
- [DKQR20] Florent Delgrange, Joost-Pieter Katoen, Tim Quatmann, and Mickael Randour. Simple strategies in multi-objective MDPs. In Armin Biere and David Parker, editors, Proceedings (Part I) of the 26th International Conference on Tools and Algorithms for the Construction and Analysis of Systems, TACAS 2020, Held as Part of ETAPS 2020, Dublin, Ireland, April 25–30, 2020, volume 12078 of Lecture Notes in Computer Science, pages 346–364. Springer, 2020.
- [Dur19] Rick Durrett. *Probability: Theory and Examples*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 5th edition, 2019.
- [EKVY08] Kousha Etessami, Marta Z. Kwiatkowska, Moshe Y. Vardi, and Mihalis Yannakakis. Multi-objective model checking of Markov decision processes. Logical Methods in Computer Science, 4(4), 2008.
- [FBB<sup>+</sup>23] Nathanaël Fijalkow, Nathalie Bertrand, Patricia Bouyer-Decitre, Romain Brenguier, Arnaud Carayol, John Fearnley, Hugo Gimbert, Florian Horn, Rasmus Ibsen-Jensen, Nicolas Markey, Benjamin Monmege, Petr Novotný, Mickael Randour, Ocan Sankur, Sylvain Schmitz, Olivier Serre, and Mateusz Skomra. Games on graphs. *CoRR*, abs/2305.10546, 2023.
- [FH13] Nathanaël Fijalkow and Florian Horn. Les jeux d'accessibilité généralisée. Technique et Science Informatiques, 32(9-10):931–949, 2013.
- [FKP12] Vojtech Forejt, Marta Z. Kwiatkowska, and David Parker. Pareto curves for probabilistic model checking. In Supratik Chakraborty and Madhavan Mukund, editors, Proceedings of the 10th International Symposium on Automated Technology for Verification and Analysis, ATVA 2014, Thiruvananthapuram, India, October 3–6, 2012, volume 7561 of Lecture Notes in Computer Science, pages 317–332. Springer, 2012.
- [Gim07] Hugo Gimbert. Pure stationary optimal strategies in Markov decision processes. In Wolfgang Thomas and Pascal Weil, editors, *Proceedings of the 24th Annual Symposium on Theoretical Aspects of Computer Science, STACS 2007, Aachen, Germany, February 22–24, 2007*, volume 4393, pages 200–211. Springer, 2007.
- [GK23] Hugo Gimbert and Edon Kelmendi. Submixing and shift-invariant stochastic games. *International Journal of Game Theory*, 52(4):1179–1214, 2023.

- [HM18] Serge Haddad and Benjamin Monmege. Interval iteration algorithm for MDPs and IMDPs. *Theoretical Computer Science*, 735:111–131, 2018.
- [Hor09] Florian Horn. Random fruits on the Zielonka tree. In Susanne Albers and Jean-Yves Marion, editors, Proceedings of the 26th International Symposium on Theoretical Aspects of Computer Science, STACS 2009, Freiburg, Germany, February 26–28, 2009, volume 3 of LIPIcs, pages 541–552. Schloss Dagstuhl—Leibniz-Zentrum für Informatik, Germany, 2009.
- [HPS<sup>+</sup>21] Ernst Moritz Hahn, Mateo Perez, Sven Schewe, Fabio Somenzi, Ashutosh Trivedi, and Dominik Wojtczak. Model-free reinforcement learning for lexicographic omega-regular objectives. In Marieke Huisman, Corina S. Pasareanu, and Naijun Zhan, editors, Proceedings of the 24th International Symposium on Formal Methods, FM 2021, Virtual Event, November 20–26, 2021, volume 13047 of Lecture Notes in Computer Science, pages 142–159. Springer, 2021.
- [LPR18] Stéphane Le Roux, Arno Pauly, and Mickael Randour. Extending finite-memory determinacy by Boolean combination of winning conditions. In Sumit Ganguly and Paritosh K. Pandya, editors, Proceedings of the 38th IARCS Annual Conference on Foundations of Software Technology and Theoretical Computer Science, FSTTCS 2018, Ahmedabad, India, December 11–13, 2018, volume 122 of LIPIcs, pages 38:1–38:20. Schloss Dagstuhl Leibniz-Zentrum fuer Informatik, 2018.
- [MPR20] Benjamin Monmege, Julie Parreaux, and Pierre-Alain Reynier. Reaching your goal optimally by playing at random with no memory. In Igor Konnov and Laura Kovács, editors, Proceedings of the 31st International Conference on Concurrency Theory, CONCUR 2020, Vienna, Austria, September 1–4, 2020, volume 171 of LIPIcs, pages 26:1–26:21. Schloss Dagstuhl –Leibniz-Zentrum für Informatik, 2020.
- [MR24] James C. A. Main and Mickael Randour. Different strokes in randomised strategies: Revisiting Kuhn's theorem under finite-memory assumptions. *Information and Computation*, 301:105229, 2024.
- [Mun97] James R. Munkres. *Topology: A First Course*. Prentice Hall International, 1997.
- [QK21] Tim Quatmann and Joost-Pieter Katoen. Multi-objective optimization of long-run average and total rewards. In Jan Friso Groote and Kim Guldstrand Larsen, editors, Proceedings (Part I) of the 27th International Conference on Tools and Algorithms for the Construction and Analysis of Systems, TACAS 2021, Held as Part of ETAPS 2021, Luxemburg City, Luxemburg, March 27–April 1, 2021, volume 12651 of Lecture Notes in Computer Science, pages 230–249. Springer, 2021.
- [Qua23] Tim Quatmann. Verification of multi-objective Markov models. PhD thesis, RWTH Aachen University, Germany, 2023.
- [Ran13] Mickael Randour. Automated synthesis of reliable and efficient systems through game theory: A case study. In *Proc. of ECCS 2012*, Springer Proceedings in Complexity XVII, pages 731–738. Springer, 2013.

- [Roc70] R. Tyrrell Rockafellar. *Convex Analysis*. Princeton Landmarks in Mathematics and Physics. Princeton University Press, 1970.
- [RRS17] Mickael Randour, Jean-François Raskin, and Ocan Sankur. Percentile queries in multi-dimensional Markov decision processes. Formal methods in system design, 50(2-3):207–248, 2017.
- [SB18] Richard S. Sutton and Andrew G. Barto. Reinforcement Learning: An Introduction. MIT Press, 2018.
- [Sha53] Lloyd S. Shapley. Stochastic games. *Proceedings of the National Academy of Sciences*, 39(10):1095–1100, 1953.

## A Additional preliminaries

This section complements Section 2.

## A.1 General notions of topology

The goal of this first section is to recall topological definitions and results mentioned in the body of the paper. Some definitions provided in this section are relevant for the later Section B.1. We refer the reader to [Mun97] for a reference on topology.

**Topology.** Let X be a non-empty set. A topology over X is a set  $\mathcal{T} \subseteq 2^X$  of subsets of X such that (i)  $\emptyset$ ,  $X \in \mathcal{T}$ , (ii) for any family  $(U_i)_{i \in I}$  such that  $U_i \in \mathcal{T}$  for all  $i \in I$ ,  $\bigcup_{i \in I} U_i \in \mathcal{T}$  and (iii) if  $U, U' \in \mathcal{T}$ , then  $U \cap U' \in \mathcal{T}$ . The pair  $(X, \mathcal{T})$  is called a topological space. Elements of  $\mathcal{T}$  are open sets. A set  $F \subseteq X$  is closed if it is the complement of an open set, i.e., if there exists  $U \in \mathcal{T}$  such that  $F = X \setminus U$ .

We say that  $(X, \mathcal{T})$  is a *Hausdorff space* when for any two distinct elements x and  $y \in X$ , there exists disjoint open sets  $U_x$  and  $U_y$  such that  $x \in U_x$  and  $y \in U_y$ . We assume that all topological spaces below are Hausdorff.

Simple examples of topologies include the discrete topology  $\mathcal{T}_{\mathsf{dis}} = 2^X$  and the trivial topology  $\{\emptyset, X\}$ . The discrete topology is Hausdorff. The trivial topology is not Hausdorff whenever X has at least two elements.

A set  $N \subseteq X$  is a neighbourhood of  $x \in X$  if there exists an open set  $U_x \in \mathcal{T}$  such that  $x \in U_x \subseteq N$ . A set is open if and only if it is a neighbourhood of all of its elements. A point  $x \in X$  is an isolated point if  $\{x\}$  is a neighbourhood of x.

Let  $Y \subseteq X$ . The *closure*  $\operatorname{cl}(Y)$  of Y is the smallest closed set in which Y is included. An element  $x \in X$  is in  $\operatorname{cl}(Y)$  if and only if all (open) neighbourhoods of x intersect Y. The *interior*  $\operatorname{int}(Y)$  of Y is the greatest open set that is included in Y. An element  $x \in X$  is in  $\operatorname{int}(Y)$  if and only if there exists an open neighbourhood  $N_x$  of x such that  $N_x \subseteq Y$ . A set is closed (resp. open) if and only if it is equal to its closure (resp. interior).

A base of  $\mathcal{T}$  is a set  $\mathcal{B} \subseteq \mathcal{T}$  such that all elements of  $\mathcal{T}$  are (arbitrary) unions of elements of  $\mathcal{B}$ . For instance, a topology is a base of itself. A base of the discrete topology is the set of all singleton sets. Another example is the usual topology of the extended real line  $\mathbb{R}$ ; a base of this topology is given by the set of intervals

$$\{ |\alpha, \beta[, [-\infty, \alpha[, ]\alpha, +\infty] \mid \alpha, \beta \in \mathbb{R}, \alpha < \beta \}.$$

This topological space is Hausdorff.

Given a non-empty set  $Y \subseteq X$ , we define the *induced* (or subspace) topology  $\mathcal{T}_{\mathsf{ind}}$  on Y as the topology defined by  $\mathcal{T}_{\mathsf{ind}} = \{U \cap Y \mid U \in \mathcal{T}\}$ . An element  $x \in Y$  is an *isolated* point of Y if it is an isolated point in  $(Y, \mathcal{T}_{\mathsf{ind}})$ .

Metric and normed spaces. Let X be a non-empty set. A metric over X is a function  $\mathsf{dist} \colon X \times X \to [0, +\infty[$  such that, for all  $x, y, z \in X$ , (i)  $\mathsf{dist}(x,y) = 0$  if and only if x = y, (ii)  $\mathsf{dist}(x,y) = \mathsf{dist}(y,x)$  and (iii)  $\mathsf{dist}(x,z) \le \mathsf{dist}(x,y) + \mathsf{dist}(y,z)$ . The last condition is called the triangle inequality. An open ball centred in  $x \in X$  of radius  $\varepsilon > 0$  is the set  $B(x,\varepsilon) = \{y \in X \mid \mathsf{dist}(x,y) < \varepsilon\}$ . A base of the topology induced by a metric is the set of open balls. A topological space  $(X,\mathcal{T})$  is metrisable if there exists a metric that induces  $\mathcal{T}$ . We remark that all metrisable spaces are necessarily Hausdorff.

For instance, the usual topology of  $\mathbb{R}$  is metrisable and is induced by the metric dist defined by  $\operatorname{dist}(x,y) = |x-y|$  for all  $x, y \in \mathbb{R}$ . The extended real line  $\mathbb{R}$  with its usual

topology is also metrisable; it is homeomorphic (i.e., topologically isomorphic) to [0,1] with the induced topology inherited from  $\mathbb{R}$ . Topological spaces X with the discrete topology are also metrisable; the *discrete metric*  $\mathsf{dist}_{\mathsf{disc}}$  defined by  $\mathsf{dist}_{\mathsf{disc}}(x,y) = 1$  whenever  $x \neq y$  induces the discrete topology (observe that all singleton sets are open balls).

Let  $d \in \mathbb{N}_0$ . Any norm  $\|\cdot\|$  on  $\mathbb{R}^d$  induces a topology via the metric  $\operatorname{dist}(\mathbf{v}, \mathbf{w}) = \|\mathbf{v} - \mathbf{w}\|$  for all  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^d$ . All norms of  $\mathbb{R}^d$  are *equivalent*, i.e., induce the same topology, which is the usual topology of  $\mathbb{R}^d$ . For infinite-dimensional spaces, which we do not consider here, some norms may not be equivalent, and can induce different topologies.

**Convergence.** Let  $(X, \mathcal{T})$  be a Hausdorff topological space. A sequence  $(x_n)_{n \in \mathbb{N}}$  of elements of X is said to converge to  $x \in X$  if for all open neighbourhoods  $U_x$  of x, there exists  $n_0 \in \mathbb{N}$  such that for all  $n \geq n_0$ ,  $x_n \in U_x$ . It is equivalent to universally quantify only over open neighbourhoods of x in a fixed base of  $\mathcal{T}$  instead of all open neighbourhoods. The uniqueness of the limit is guaranteed by the Hausdorff assumption. We note that in general spaces, sequences can have several limits: e.g., for the trivial topology  $\{\emptyset, X\}$ , all sequences converge to all elements of the set.

In a metric space  $(X, \mathsf{dist})$ , this definition of convergence is equivalent to the usual definition recalled hereafter: for all  $\varepsilon > 0$ , there exists some  $n_0 \in \mathbb{N}$  such that for all  $n > n_0$ ,  $\mathsf{dist}(x_n, x) < \varepsilon$  (i.e.,  $x_n \in B(x, \varepsilon)$ ).

Convergence of (real) sequences in  $\mathbb{R}$  in the above sense is equivalent to the classical definitions for convergence to a real limit,  $+\infty$  or  $-\infty$ .

Continuity. A function  $f:(X,\mathcal{T})\to (Y,\mathcal{T}')$  is continuous at  $x\in X$  if for all neighbourhoods  $N_{f(x)}\subseteq Y$  of f(x),  $f^{-1}(N_{f(x)})$  is a neighbourhood of x. If  $\mathcal{B}_Y$  is a basis of  $(Y,\mathcal{T}')$ , continuity can be checked by looking only at elements of the basis in the following sense: f is continuous at x if and only if for all  $U_{f(x)}\in \mathcal{B}_Y$  such that  $f(x)\in U_{f(x)}$ ,  $f^{-1}(U_{f(x)})$  is a neighbourhood of x. The function f is said to be continuous if it is continuous at x for all  $x\in X$ .

If  $f:(X,\mathsf{dist}_X) \to (Y,\mathsf{dist}_Y)$  is a function between metric spaces, the definition above is directly equivalent to the usual  $\varepsilon$ - $\delta$  definition of continuity: f is continuous at  $x \in X$  if for all  $\varepsilon > 0$ , there exists  $\delta > 0$  such that for all  $x' \in X$ ,  $\mathsf{dist}_X(x,x') < \delta$  implies that  $\mathsf{dist}_Y(f(x),f(x')) < \varepsilon$ .

For functions between metric spaces, there exists a stronger variant of continuity, called uniform continuity. A function  $f:(X,\mathsf{dist}_X)\to (Y,\mathsf{dist}_Y)$  is uniformly continuous if for all  $\varepsilon>0$ , there exists some  $\delta>0$  such that for all  $x,x'\in X$ ,  $\mathsf{dist}_X(x,x')<\delta$  implies that  $\mathsf{dist}_Y(f(x),f(x'))<\varepsilon$ . The difference with continuity is the quantification order. For continuity,  $\delta$  may depend on both  $\varepsilon$  and the point at which we check continuity, whereas for uniform continuity,  $\delta$  may only depend on  $\varepsilon$  and must work for all points.

For instance, the function  $[0,+\infty[ \to [0,+\infty[ : x \mapsto \sqrt{x} \text{ is uniformly continuous. This can be shown via the observation that errors at a neighbourhood of some <math>x \ge 0$  can be bounded independently of x, i.e., we have that for all  $x \ge 0$  and  $|h| \le x$  (this ensures that  $\sqrt{x+h}$  is well-defined), we have  $|\sqrt{x+h}-\sqrt{x}| \le \sqrt{|h|}$ . On the other hand, the function  $\mathbb{R} \to \mathbb{R}$ :  $x \mapsto x^2$  is not uniformly continuous. For any  $x \in \mathbb{R}$  and  $h \in \mathbb{R}$ , we have  $|(x+h)^2-x^2|=|h^2-2xh|$ , and thus, intuitively, we cannot choose  $\delta$  independently of x in the definition of continuity because errors depend on x (which cannot be bounded).

For functions from a metric space to another, continuity at x is equivalent to sequential continuity at x. A function  $f: (X, \mathcal{T}) \to (Y, \mathcal{T}')$  is sequentially continuous at  $x \in X$  if for all sequences  $(x_n)_{n \in \mathbb{N}}$  that converge to x, the sequence  $(f(x_n))_{n \in \mathbb{N}}$  converges to f(x).

We now prove a result implying that the non-negative and non-positive parts of a continuous function are continuous, as this is used in the main text.

**Lemma A.1.** Let  $(X, \mathcal{T})$  be a topological space,  $x \in X$ ,  $f: X \to \mathbb{R}$  be a function that is continuous at x and  $M \in \mathbb{R}$ . Then the functions  $\min(f, M): y \mapsto \min\{f(y), M\}$  and  $\max(f, M): y \mapsto \max\{f(y), M\}$  are continuous at x.

*Proof.* We provide a proof only for  $\min(f, M)$  as the argument is analogous for  $\max(f, M)$ . We distinguish three cases.

First, assume that f(x) < M. Let  $N_{f(x)}$  be a neighbourhood of  $f(x) = \min(f, M)(x)$ . We must show that  $\min(f, M)^{-1}(N_{f(x)})$  is a neighbourhood of x. By continuity of f at x, since  $N_{f(x)} \cap [-\infty, M[$  is a neighbourhood of f(x),  $f^{-1}(N_{f(x)} \cap [-\infty, M[)$  is a neighbourhood of x. To close the first case, it suffices to establish that  $f^{-1}(N_{f(x)} \cap [-\infty, M[) \subseteq \min(f, M)^{-1}(N_{f(x)})$ . This inclusion follows from the fact that for all  $y \in X$ , if f(y) < M, then  $f(y) = \min(f, M)(y)$ . This ends the proof of the first case.

Second, assume that f(x) > M. Let  $N_M$  be a neighbourhood of  $M = \min(f, M)(x)$ . By definition of  $\min(f, M)$ , we obtain that for all  $y \in f^{-1}(]M, +\infty]$ ,  $\min(f, M)(y) = M$ . It follows that  $f^{-1}(]M, +\infty]$ )  $\subseteq \min(f, M)^{-1}(N_M)$ . By continuity of f at  $x, f^{-1}(]M, +\infty]$ ) is a neighbourhood of x. We conclude from the above that  $\min(f, M)^{-1}(N_M)$  is a neighbourhood of x.

Finally, assume that f(x) = M and let  $N_M$  be a neighbourhood of  $M = \min(f, M)(x)$ . It suffices to show that  $f^{-1}(N_M) \subseteq \min(f, M)^{-1}(N_M)$  by continuity of f at x. Let  $y \in X$  such that  $f(y) \in N_M$ . If  $f(y) \leq M$ , then  $\min(f, M)(y) = f(y) \in N_M$ . Otherwise,  $\min(f, M)(y) = M \in N_M$ . This shows the required inclusion and ends the proof.

**Compactness.** A topological space  $(X, \mathcal{T})$  is *compact* if for any open cover  $(U_i)_{i \in I}$  of X (i.e., for all  $i \in I$ ,  $U_i$  is open and  $\bigcup_{i \in I} U_i = X$ ), one can extract a finite open cover of X, i.e., there exists  $I' \subseteq I$  finite such that  $\bigcup_{i \in I'} U_i = X$ . A subset Y of a topological space  $(X, \mathcal{T})$  is compact if  $(Y, \mathcal{T}_{ind})$  is compact, where  $\mathcal{T}_{ind}$  is the induced topology. A compact subset of a Hausdorff topological space is closed. For instance, any finite set with the discrete topology is compact. Any closed bounded interval of  $\mathbb{R}$  is also compact. This can be used to show that the extended real line  $\mathbb{R}$  is compact (without relying on the fact that  $\mathbb{R}$  and [0,1] are homeomorphic).

In metrisable spaces, there is an equivalent characterisation of compactness based on sequences. A topological space  $(X, \mathcal{T})$  is sequentially compact if for all sequences  $(x_n)_{n\in\mathbb{N}}$  of X, there exists a convergent subsequence. A metrisable space is compact if and only if it is sequentially compact (e.g., [Mun97, Chap. 3, Thm. 7.4]).

For subsets of  $\mathbb{R}^d$  (and more generally, of finite-dimensional normed vector spaces), there is yet another equivalent formulation of compactness. A set  $D \subseteq \mathbb{R}^d$  is compact if and only if it is closed and bounded (e.g., [Mun97, Chap. 3, Thm. 6.3]).

Let  $(X, \mathcal{T})$  and  $(Y, \mathcal{T}')$  be topological spaces and let  $f: X \to Y$ . If  $(X, \mathcal{T})$  is compact and f is continuous, then f(X) is compact. Furthermore, if  $(X, \mathsf{dist}_X)$  is a compact metric space and  $(Y, \mathsf{dist}_Y)$  is a metric space, f is continuous if and only if f is uniformly continuous (e.g., [Mun97, Chap. 3, Thm. 7.3]).

**Product topology.** The *product topology* is a topology defined over Cartesian products of topological spaces. Let I be an arbitrary non-empty set. For all  $i \in I$ , let  $(X_i, \mathcal{T}_i)$  be a topological space. A base of open sets for the product topology over  $\prod_{i \in I} X_i$  consists of the open sets  $\prod_{i \in I} U_i$  where for all  $i \in I$ ,  $U_i \in \mathcal{T}_i$  and  $U_i = X_i$  for all but finitely

many  $i \in I$ . The product topology is the coarsest topology for which the projections  $\prod_{i' \in I} X_{i'} \to X_i : (x_{i'})_{i' \in I} \mapsto x_i$  are continuous for all  $i \in I$ .

Ā simple example of the product topology is  $\mathbb{R}^d$ ; the usual topology of  $\mathbb{R}^d$  corresponds to the product topology of d copies of  $\mathbb{R}$  with its usual topology.

A sequence in a product of topological spaces converges with respect to the product topology if and only if it converges component-wise (see, e.g., [Mun97, Chap. 2, Sect. 8, Exercise 6]). This is formalised in the following result.

**Lemma A.2.** Let I be a non-empty set. For all  $i \in I$ , let  $(X_i, \mathcal{T}_i)$  be a topological space. Let  $\mathcal{T}_{\Pi}$  denote the product topology on  $\prod_{i \in I} X_i$ . Let  $(\mathbf{x}^{(n)})_{n \in \mathbb{N}}$  be a sequence of elements of  $\prod_{i \in I} X_i$  and  $\mathbf{x} = (x_i)_{i \in \mathbb{N}} \in \prod_{i \in I} X_i$ . Then  $(\mathbf{x}^{(n)})_{n \in \mathbb{N}}$  converges to  $\mathbf{x}$  if and only if for all  $i \in I$ ,  $(x_i^{(n)})_{n \in \mathbb{N}}$  converges to  $x_i$ .

A product of metrisable spaces need not be metrisable in general. However, countable products of metrisable spaces are metrisable. This can be shown in the same way that [Mun97, Chap 2, Thm. 9.5] proves that  $\mathbb{R}^{\omega}$  with the product topology is metrisable (where the usual topology is assumed on  $\mathbb{R}$ ).

Finally, we show that countable products of compact metrisable spaces are compact. In full generality, arbitrary products of compact topological spaces are compact; this result is known as Tychonoff's theorem (e.g., [Mun97, Chap. 5, Thm. 1.1]). Tychonoff's theorem is equivalent to the axiom of choice. Because of this, we provide an alternative proof below for the case of countable products.

**Theorem A.3.** For all  $i \in \mathbb{N}$ , let  $(X_i, \mathcal{T}_i)$  be a sequentially compact topological space. Then  $(\prod_{i \in \mathbb{N}} X_i, \mathcal{T}_{\Pi})$  is sequentially compact, where  $\mathcal{T}_{\Pi}$  denotes the product topology. In particular, countable products of compact metrisable spaces are compact.

*Proof.* The second claim of the theorem follows from the first and the fact that, in metrisable spaces, compactness and sequential compactness are equivalent. We thus focus on the first claim of the theorem. Let  $(\mathbf{x}^{(n)})_{n\in\mathbb{N}}$  be a sequence of elements of  $\prod_{i\in\mathbb{N}} X_i$ . Our goal is to show that  $(\mathbf{x}^{(n)})_{n\in\mathbb{N}}$  has a convergent subsequence.

First, for all  $i \in \mathbb{N}$ , we construct a subsequence  $(\mathbf{x}^{(n)})_{n \in I_i}$  of  $(\mathbf{x}^{(n)})_{n \in \mathbb{N}}$  such that for all j < i,  $(x_j^{(n)})_{n \in I_i}$  converges in  $X_j$ . We proceed by induction. We let  $I_0 = \mathbb{N}$  for the base case. For the induction step, we assume that  $I_i$  is defined. By compactness of  $X_{i+1}$ , there exists  $I_{i+1} \subseteq I_i$  such that  $(x_i^{(n)})_{n \in I_{i+1}}$  converges. The induction hypothesis holds by construction because  $(\mathbf{x}^{(n)})_{n \in I_{i+1}}$  is a subsequence of  $(\mathbf{x}^{(n)})_{n \in I_i}$ .

We now construct a convergent subsequence of  $(\mathbf{x}^{(n)})_{n\in\mathbb{N}}$ . Let  $n_0 = \min I_1$  and, for all i > 0, let  $n_i$  be the least element of  $I_{i+1}$  strictly greater than  $n_{i-1}$ . It is easy to check that  $(\mathbf{x}^{(n_i)})_{i\in\mathbb{N}}$  converges via Lemma A.2.

### A.2 Properties of the downward closure

Let  $d \in \mathbb{N}_0$ . We show that the downward closure of a closed subset of  $\mathbb{R}^d$  is closed. We remark below that the statement, in general, does not apply to closed sets of  $\mathbb{R}^d$ .

**Lemma A.4.** Let  $D \subseteq \mathbb{R}^d$  be closed (i.e., compact) with respect to the topology of  $\mathbb{R}^d$ . Then down(D) is compact with respect to the topology of  $\mathbb{R}^d$  and  $down(D) \cap \mathbb{R}^d$  is closed with respect to the topology of  $\mathbb{R}^d$ .

*Proof.* Recall that  $\bar{\mathbb{R}}^d$  is a compact metrisable space. Thus, it suffices to show that down(D) is closed with respect to the topology of  $\bar{\mathbb{R}}^d$  to end the proof.

Let  $\mathbf{q} \in \mathbb{R}^d$  and  $(\mathbf{q}_n)_{n \in \mathbb{N}}$  a sequence of elements of  $\mathsf{down}(D)$  such that  $\mathbf{q}_n \to \mathbf{q}$  when  $n \to \infty$ . We must show that  $\mathbf{q} \in \mathsf{down}(D)$ . The idea of the proof is to bound  $\mathbf{q}$  from above by a vector that is the limit of some sequence of elements of D.

For all  $n \in \mathbb{N}$ , we let  $\mathbf{p}_n \in D$  such that  $\mathbf{p}_n \geq \mathbf{q}_n$ . By (sequential) compactness of D,  $(\mathbf{p}_n)_{n \in \mathbb{N}}$  has a convergent subsequence. Let  $\mathbf{p} \in D$  denote the limit of one such subsequence. It follows that  $\mathbf{q} \leq \mathbf{p}$ . This shows that  $\mathbf{q} \in \mathsf{down}(D)$ , and thus  $\mathsf{down}(D)$  is a closed subset of  $\mathbb{R}^d$ . Since the topology of  $\mathbb{R}^d$  can be seen as the topology induced on  $\mathbb{R}^d$  by that of  $\mathbb{R}^d$ , it follows that  $\mathsf{down}(D) \cap \mathbb{R}^d$  is a closed subset of  $\mathbb{R}^d$ .

**Remark A.5** (Assumption of Lemma A.4). The subsets of  $\mathbb{R}^d$  that are closed with respect to the topology of  $\mathbb{R}^d$  are the compact subsets of  $\mathbb{R}^d$ . Therefore, the assumption of Lemma A.4 does not apply to all closed subsets of  $\mathbb{R}^d$ .

We provide closed subset of  $\mathbb{R}^2$  the downward-closure of which is not a closed subset of  $\mathbb{R}^2$ . We consider  $D = \{(-\frac{1}{n}, n) \mid n \in \mathbb{N}_0\}$ . To see that D is  $\mathbb{R}^2$ -closed, consider a convergent sequence of elements of D. As it is a Cauchy sequence, from some point on, all subsequent elements of the sequence are at a distance of at most  $\frac{1}{2}$  from one another. Because the distance between two different elements of D is at least 1, it follows the sequence that is considered is ultimately constant, thus its limit lies in D. This shows that D is  $\mathbb{R}^2$ -closed.

We now argue that  $\mathsf{down}(D)$  and  $\mathsf{down}(D) \cap \mathbb{R}^2$  are not closed: (0,0) is in the closure of these sets, but not in them. To show that  $(0,0) \in \mathsf{cl}(\mathsf{down}(D))$ , we observe that the sequence  $((-\frac{1}{n},0))_{n\in\mathbb{N}_0}$  is a sequence of elements of  $\mathsf{down}(D)$  that converges to (0,0). On the other hand, we see that for all  $n \in \mathbb{N}_0$ ,  $(0,0) \leq (-\frac{1}{n},n)$  does not hold, i.e.,  $(0,0) \notin \mathsf{down}(D)$ .

## B Complement to Section 3

### B.1 Topology on the set of plays

Let  $\mathfrak{M} = (S, A, \delta)$  be an MDP. In this section, we introduce the usual topology over  $\mathsf{Plays}(\mathfrak{M})$ . At the end of this section, we prove that the definition of a continuous payoff given in Section 3 is coherent with the usual one.

We consider  $(SA)^{\omega}$  equipped with the product topology, where S and A are both equipped with the discrete topology. It follows that  $(SA)^{\omega}$  is both metrisable (as a countable product of metrisable spaces) and compact (by Theorem A.3). The topology of  $\mathsf{Plays}(\mathfrak{M})$  is the topology induced on  $\mathsf{Plays}(\mathfrak{M})$  by the topology of  $(SA)^{\omega}$ . A base of the topology of  $\mathsf{Plays}(\mathfrak{M})$  is the set of cylinder sets. We show below that  $\mathsf{Plays}(\mathfrak{M})$  is a closed subset of  $(SA)^{\omega}$ , and is therefore compact.

**Lemma B.1.** For any MDP  $\mathfrak{M} = (S, A, \delta)$ , Plays $(\mathfrak{M})$  is a closed subset of  $(SA)^{\omega}$ . In particular, Plays $(\mathfrak{M})$  is a compact space.

*Proof.* The first part of the statement implies the second, because all closed subsets of compact spaces are themselves compact [Mun97, Chap. 3, Thm. 5.2].

For all  $w \in (SA)^*$ , the subset of  $(SA)^{\omega}$  that contains all continuations of w is an open set (in fact, this set is in the base of the product topology). Furthermore, any  $u \in (SA)^{\omega} \setminus \mathsf{Plays}(\mathfrak{M})$  has a prefix  $w_u \in (SA)^*S$  that is not a history. We obtain that, for all  $u \in (SA)^{\omega} \setminus \mathsf{Plays}(\mathfrak{M})$ , the set of continuations of  $w_u$  does not intersect  $\mathsf{Plays}(\mathfrak{M})$ . Therefore,  $(SA)^{\omega} \setminus \mathsf{Plays}(\mathfrak{M})$  can be written as the union of the sets of continuations of each  $w_u$ , thus is an open set. We have shown that  $\mathsf{Plays}(\mathfrak{M})$  is closed.

We now introduce a metric on  $\mathsf{Plays}(\mathfrak{M})$  to justify the definition of a continuous payoff provided in the main text. We consider the metric  $\mathsf{dist}_{\mathsf{play}}$  over  $\mathsf{Plays}(\mathfrak{M})$  defined by, for all  $\pi, \pi' \in \mathsf{Plays}(\mathfrak{M})$ ,  $\mathsf{dist}_{\mathsf{play}}(\pi, \pi') = 2^{-r}$ , where  $r \in \mathbb{N} \cup \{+\infty\}$  is the number of elements in the longest common prefix of  $\pi$  and  $\pi'$  (states and actions are counted separately). This distance is derived from the discrete metric and a construction that shows that countable products of metric spaces are metrisable.

We now show that the definition of continuity and uniform continuity for payoffs given in the main text is indeed equivalent to the definition of the original topological definitions. The crux of the argument for both cases is that cylinder sets are open balls with respect to  $\operatorname{dist}_{\mathsf{play}}$ : for all  $h = s_0 a_0 s_1 \dots a_{r-1} s_r \in \operatorname{Hist}(\mathfrak{M})$ , by definition of  $\operatorname{dist}_{\mathsf{play}}$ ,  $\operatorname{Cyl}(h) = B(\pi, 2^{-(2r+1)})$  for all  $\pi \in \operatorname{Cyl}(h)$ . Furthermore, any open neighbourhood of a play contains a cylinder, as all balls include balls with smaller radii. We provide separate statements for continuity and uniform continuity. We start with continuity.

**Lemma B.2.** Let  $f: \mathsf{Plays}(\mathfrak{M}) \to \overline{\mathbb{R}}$  be a payoff function and let  $\pi \in \mathsf{Plays}(\mathfrak{M})$ .

- If  $f(\pi) \in \mathbb{R}$ , then f is continuous at  $\pi$  if and only if for all  $\varepsilon > 0$ , there exists  $\ell \in \mathbb{N}$  such that for all plays  $\pi' \in \mathsf{Cyl}(\pi_{<\ell}), |f(\pi) f(\pi')| < \varepsilon$ .
- If  $f(\pi) = +\infty$  (resp.  $-\infty$ ), then f is continuous at  $\pi$  if and only if for all  $M \in \mathbb{R}$ , there exists  $\ell \in \mathbb{N}$  such that for all plays  $\pi' \in \mathsf{Cyl}(\pi_{<\ell})$ ,  $f(\pi') \geq M$  (resp.  $f(\pi') \leq -M$ ).

Proof. We proceed by equivalences. We first open with general equivalences without assuming anything on  $f(\pi)$ . By definition, f is continuous at  $\pi$  if and only if, for all open neighbourhoods  $N_{f(\pi)}$  of  $f(\pi)$ , there exists an open neighbourhood  $N_{\pi}$  of  $\pi$  such that  $f(N_{\pi}) \subseteq N_{f(\pi)}$ . All open neighbourhoods of  $\pi$  contain an open ball centred in  $\pi$  with respect to  $\operatorname{dist}_{\mathsf{play}}$ , and thus a cylinder set of a prefix of  $\pi$ . Thus f is continuous at  $\pi$  if and only if, for all open neighbourhoods  $N_{f(\pi)}$  of  $f(\pi)$ , there exists  $\ell \in \mathbb{N}$  such that for all  $\pi' \in \mathsf{Cyl}(\pi_{\leq \ell})$ ,  $f(\pi') \in N_{f(\pi)}$ .

We now distinguish cases following whether  $f(\pi) \in \mathbb{R}$ . First, assume that  $f(\pi) \in \mathbb{R}$ . Then, all neighbourhoods of  $f(\pi)$  contain an interval  $]f(\pi) - \varepsilon, f(\pi) + \varepsilon[$  for some  $\varepsilon > 0$ . The equivalence in the statement of the lemma follows. Now, assume that  $f(\pi) = +\infty$ . By definition of the topology of  $\mathbb{R}$  (see the base specified previously), all neighbourhoods of  $+\infty$  contain an interval  $]M,\infty]$  for some  $M \in \mathbb{R}$ . The equivalence in the statement of the lemma follows. The argument is similar in the case  $f(\pi) = -\infty$ .

We now continue with uniform continuity.

**Lemma B.3.** Let  $f: \mathsf{Plays}(\mathfrak{M}) \to \mathbb{R}$  be a real-valued payoff. The payoff f is continuous if and only if for all  $\varepsilon > 0$ , there exists  $\ell \in \mathbb{N}$  such that for all plays  $\pi$ ,  $\pi' \in \mathsf{Plays}(\mathfrak{M})$ ,  $\pi_{\leq \ell} = \pi'_{\leq \ell}$  implies that  $|f(\pi) - f(\pi')| < \varepsilon$ .

Proof. Since (Plays( $\mathfrak{M}$ ), dist<sub>play</sub>) is a compact metric space (Theorem A.3), f is continuous if and only if f is uniformly continuous, i.e., if for all  $\varepsilon > 0$ , there exists some  $\eta > 0$  such that for all  $\pi$ ,  $\pi' \in \mathsf{Plays}(\mathfrak{M})$ , dist<sub>play</sub> $(\pi, \pi') < \eta$  implies that  $|f(\pi) - f(\pi')| < \varepsilon$ . The desired equivalence can be obtained by exploiting the fact that for all  $\ell \in \mathbb{N}$  and  $\pi$ ,  $\pi' \in \mathsf{Plays}(\mathfrak{M})$ , dist<sub>play</sub> $(\pi, \pi') \leq 2^{-(2\ell+1)}$  if and only if  $\pi \leq \ell = \pi' \leq \ell$ .

#### B.2 Examples of continuous payoffs

We present examples of continuous payoffs in this section. First, we show that (a generalisation of) the discounted-sum payoff is continuous. Second, we show that the shortest-path payoff is continuous whenever the considered weight function is positive. Finally,

we provide characterisations of the objectives whose indicator is continuous and of prefixindependent payoffs that are continuous. We fix an MDP  $\mathfrak{M} = (S, A, \delta)$  for the remainder of the section.

### B.2.1 Discounted-sum payoff

In the main text, we defined the discounted-sum payoff with a fixed discount factor. Certain authors study a variant of this payoff where the discount factor changes at each step of the play depending on the current state. We define a generalisation of the discounted-sum payoff in which both the discount factor and the weights depend not only on the current state-action pair, but on the history and current action at each step. We provide sufficient conditions ensuring that this generalisation is well-defined and continuous.

We consider a history-dependent weight function  $w: (SA)^+ \to \mathbb{R}$  and a history-dependent discount factor function  $\lambda: (SA)^+ \to [0,1[$ . We assume that w is no more than  $W \in \mathbb{R}$  in absolute value and that  $\lambda$  is bounded away from 1, i.e., there exists  $\lambda_{\star} \in [0,1[$  such that  $\lambda(u) \leq \lambda_{\star}$  for all  $u \in (SA)^+$ . We define the generalised discounted-sum payoff as the function  $\mathsf{GDSum}_w^{\lambda}$  defined, for all  $\pi = s_0 a_0 s_1 \ldots \in \mathsf{Plays}(\mathfrak{M})$ , by

$$\mathsf{GDSum}_w^\lambda(\pi) = \sum_{r=0}^\infty \left( \prod_{\ell=0}^{r-1} \lambda(s_0 a_0 \dots s_\ell a_\ell) \right) w(s_0 a_0 \dots s_r a_r).$$

This function is well-defined for all plays: the defining series is absolutely convergent by the assumptions on w and  $\lambda$ . We now prove that  $\mathsf{GDSum}_w^{\lambda}$  is continuous.

**Lemma B.4.** Let  $w: (SA)^+ \to \mathbb{R}$  be a history-dependent weight function such that w is no more than  $W \in \mathbb{R}$  in absolute value and  $\lambda: (SA)^+ \to [0,1[$  be a history-dependent discount factor function such that there exists  $\lambda_{\star} \in [0,1[$  such that  $\lambda(u) \leq \lambda_{\star}$  for all  $u \in (SA)^+$ . Then  $\mathsf{GDSum}_w^{\lambda}$  is a continuous payoff.

*Proof.* Let  $\pi = s_0 a_1 s_1 \ldots \in \mathsf{Plays}(\mathfrak{M})$  and  $\varepsilon > 0$ . Let  $\ell \in \mathbb{N}$  such that  $\frac{2 \cdot W \cdot \lambda_x^{\ell}}{1 - \lambda_x} < \varepsilon$  (whose existence is guaranteed by  $\lambda_x \in [0,1[)$ . Let  $\pi' = t_0 b_0 t_0 \ldots \in \mathsf{Cyl}(\pi_{\leq \ell})$ . For all  $r \in \mathbb{N}$ , let  $u_r = s_0 a_0 \ldots s_r a_r$  and  $u'_r = t_0 b_0 \ldots t_r b_r$ . By definition of  $\mathsf{GDSum}_w^{\lambda}$ , we obtain that

$$\begin{split} \left| \mathsf{GDSum}_w^\lambda(\pi) - \mathsf{GDSum}_w^\lambda(\pi') \right| \\ &= \left| \sum_{r=\ell}^\infty \left( \left( \prod_{n=0}^{r-1} \lambda(u_n) \right) w(u_r) - \left( \prod_{n=0}^{r-1} \lambda(u_n') \right) w(u_r') \right) \right| \\ &\leq 2 \cdot \sum_{r=\ell}^\infty \lambda_\star^r \cdot W \\ &= \frac{2 \cdot W \cdot \lambda_\star^\ell}{1 - \lambda_\star} \\ &< \varepsilon. \end{split}$$

We have shown that  $\mathsf{GDSum}_w^{\lambda}$  is continuous in  $\pi$ .

#### B.2.2 Shortest-path payoff

We consider a weight function  $w \colon S \times A \to \mathbb{R}$  and a target  $T \subseteq S$ . The shortest-path payoff  $\mathsf{SPath}_w^T$  is continuous over  $\mathsf{Reach}(T)$ , since the payoff of a play depends only on its prefix prior to the first visit to T. However, without imposing any conditions on w, the shortest-path payoff  $\mathsf{SPath}_w^T$  is not necessarily continuous everywhere.

**Example B.1.** Consider a two-state MDP  $\mathfrak{M}=(\{s,t\},\{a\},\delta)$  such that  $\delta(s,a)$  is the uniform distribution on  $\{s,t\}$  and  $\delta(t,a)(t)=1$ . Let w be a the constant zero weight function and consider the target  $T=\{t\}$ . The payoff  $\mathsf{SPath}_w^T$  is not continuous at  $\pi=(sa)^\omega$ . Indeed, for all  $\ell\in\mathbb{N}$ , the play  $\pi'=(sa)^{\ell+1}(ta)^\omega$  is such that  $\pi_{\leq \ell}=\pi'_{\leq \ell}$  and  $\mathsf{SPath}_w^T(\pi')=0$ . Therefore, for  $\mathsf{SPath}_w^T$  to be continuous, 0 would have to be in all neighbourhoods of  $\mathsf{SPath}_w^T(\pi)=+\infty$ , which is not the case.

The same example can be used to show that  $\mathbb{1}_{\mathsf{Reach}(T)}$  is not continuous in general.  $\triangleleft$ 

The problem occurring in the previous example is that there is a play with an infinite payoff that can be approached (in the sense of convergence) by plays that have a bounded payoff. A sufficient condition to avoid this phenomenon is to require that all weights are positive. In that case, the payoff of a play is no less than the smallest weight multiplied by the number of actions occurring before the first visit to T. We show that this condition implies the continuity of  $\mathsf{SPath}^T_w$ .

**Lemma B.5.** Let  $w: S \times A \to \mathbb{R}$  be a weight function and  $T \subseteq S$  be a target. Assume that there exists  $\eta > 0$  such that  $w(s, a) \ge \eta$  for all  $s \in S$  and all  $a \in A$ . Then  $\mathsf{SPath}_w^T$  is continuous.

Proof. Let  $\pi = s_0 a_0 s_1 \ldots \in \mathsf{Plays}(\mathfrak{M})$ . First, assume that  $\mathsf{SPath}_w^T(\pi) \in \mathbb{R}$ , i.e.,  $\pi \in \mathsf{Reach}(T)$ . Let  $r \in \mathbb{N}$  such that  $s_r \in T$ . By definition of  $\mathsf{SPath}_w^T$ , for all  $\pi' \in \mathsf{Cyl}(\pi_{\leq r})$ , we have  $\mathsf{SPath}_w^T(\pi) = \mathsf{SPath}_w^T(\pi')$ . This implies that  $\mathsf{SPath}_w^T$  is continuous in  $\pi$ .

Now, assume that  $\mathsf{SPath}_w^T(\pi) = +\infty$ . Let  $M \in \mathbb{R}$ . Let  $r \in \mathbb{N}$  such that  $M \leq r \cdot \eta$ . We claim that for all  $\pi' \in \mathsf{Cyl}(\pi_{\leq r})$ ,  $\mathsf{SPath}_w^T(\pi') \geq M$ . Let  $\pi' \in \mathsf{Cyl}(\pi_{\leq r})$ . If  $\pi' \notin \mathsf{Reach}(T)$ , the sought inequality is direct. We now assume that  $\pi' \in \mathsf{Reach}(T)$ . Because no state of T occurs in  $\pi_{\leq r}$  and all weights are non-negative, we obtain that

$$\mathsf{SPath}_w^T(\pi') \geq \sum_{\ell=0}^r w(s_\ell, a_\ell) \geq r \cdot \eta \geq M.$$

We have shown that  $\mathsf{SPath}_w^T$  is continuous.

### B.2.3 Objectives and indicators.

We now characterise objectives that have a continuous indicator function. Let  $\Omega$  be an objective. The co-domain of an indicator function is  $\{0,1\}$ . This implies that  $\mathbbm{1}_{\Omega}$  is continuous if and only if, for all plays  $\pi$ , there exist  $\ell \in \mathbb{N}$  such that, for all plays  $\pi' \in \mathsf{Cyl}(\pi_{\leq \ell})$ , we have  $\pi' \in \Omega$  if and only if  $\pi \in \Omega$ . Furthermore it follows from uniform continuity that  $\ell$  can be chosen independently of the play in the previous statement. Therefore,  $\mathbbm{1}_{\Omega}$  is continuous if and only if membership in  $\Omega$  depends only on a bounded prefix of plays. We show below that this is also equivalent to  $\Omega$  being both open and closed.

**Lemma B.6.** Let  $\Omega$  be an objective. The three following statements are equivalent:

- 1.  $\mathbb{1}_{\Omega}$  is continuous;
- 2. there exists  $\ell \in \mathbb{N}$  such that for all  $\pi \in \mathsf{Plays}(\mathfrak{M})$ ,  $\mathsf{Cyl}(\pi_{\leq \ell})$  is either included in or disjoint from  $\Omega$ ;
- 3.  $\Omega$  is open and closed, i.e., there exist finitely many histories  $h_1, \ldots, h_n$  such that  $\Omega = \bigcup_{m=1}^n \mathsf{Cyl}(h_m)$ .

*Proof.* We show that Items 1 and 2 are equivalent, then that Items 2 and 3 are equivalent. In the interest of this proof being self-contained, we also show that the two properties in Item 3 are equivalent at the end of the proof.

We first show that Items 1 and 2 are equivalent. Since  $\mathbb{1}_{\Omega}$  is real-valued, we have  $\mathbb{1}_{\Omega}$  continuous if and only if, for all  $\varepsilon > 0$  there exists  $\ell \in \mathbb{N}$  such that for all plays  $\pi, \pi' \in \mathsf{Plays}(\mathfrak{M})$ , if  $\pi_{\leq \ell} = \pi'_{\leq \ell}$ , then  $|\mathbb{1}_{\Omega}(\pi) - \mathbb{1}_{\Omega}(\pi')| < \varepsilon$ . It follows from the co-domain of  $\mathbb{1}_{\Omega}$  being  $\{0,1\}$  that  $\mathbb{1}_{\Omega}$  is continuous if and only if there exists  $\ell \in \mathbb{N}$  such that for all plays  $\pi, \pi' \in \mathsf{Plays}(\mathfrak{M})$ , if  $\pi' \in \mathsf{Cyl}(\pi_{\leq \ell})$ , then  $\mathbb{1}_{\Omega}(\pi) = \mathbb{1}_{\Omega}(\pi')$  (the non-trivial direction follows by choosing  $\varepsilon = \frac{1}{2}$ ), i.e.,  $\mathsf{Cyl}(\pi_{\leq \ell})$  is included in or disjoint from  $\Omega$ . This establishes the equivalence of Items 1 and 2.

Next, we establish that Items 2 and 3 are equivalent. First, assume that Item 2 holds and let  $\ell \in \mathbb{N}$  be given by this property. We obtain that  $\Omega = \bigcup_{\pi \in \Omega} \mathsf{Cyl}(\pi_{\leq \ell})$ . There are finitely many histories of the form  $\pi_{\leq \ell}$  ( $\pi \in \mathsf{Plays}(\mathfrak{M})$ ) because S and A are finite. This shows that Item 3 holds. Conversely, assume that Item 3 holds and let  $h_1, \ldots, h_n$  such that  $\Omega = \bigcup_{m=1}^n \mathsf{Cyl}(h_m)$ . Item 2 follows by letting  $\ell$  be the greatest number of states in the histories  $h_1, \ldots, h_n$  or zero if there are no histories.

For the sake of completeness, we close the proof by showing that the two properties given in Item 3 are equivalent. First, assume that  $\Omega$  is open and closed. A base of the topology of  $\mathsf{Plays}(\mathfrak{M})$  is the set of cylinder sets, therefore  $\Omega$  is a union of cylinder sets. Since  $\mathsf{Plays}(\mathfrak{M})$  is compact and  $\Omega$  is closed, it follows that  $\Omega$  is compact. We conclude that  $\Omega$  can be written as a finite union of cylinder sets by compactness.

Conversely, assume that  $\Omega = \bigcup_{m=1}^n \mathsf{Cyl}(h_m)$  for some histories  $h_1, \ldots, h_n$ . It suffices to show that  $\mathsf{Cyl}(h)$  is open and closed for all  $h \in \mathsf{Hist}(\mathfrak{M})$ . Let  $h = s_0 a_0 \ldots a_{r-1} s_r \in \mathsf{Hist}(\mathfrak{M})$ . By definition of the product topology,  $\mathsf{Cyl}(h)$  is open. To show that  $\mathsf{Cyl}(h)$  is closed, we consider  $\pi \in \mathsf{cl}(\mathsf{Cyl}(h))$  and show that  $\pi \in \mathsf{Cyl}(h)$ . By definition of closure, the cylinder  $\mathsf{Cyl}(\pi_{\leq r})$  intersects  $\mathsf{Cyl}(h)$ , i.e., there exists a play with the prefixes  $\pi_{\leq r}$  and h. It follows that  $\pi_{\leq r} = h$ , and thus  $\pi \in \mathsf{Cyl}(h)$ . We have shown that  $\mathsf{Cyl}(h)$  is both open and closed, ending the proof.

## B.2.4 Prefix-independent payoffs.

Prefix-independent functions assign the same payoff to any two plays that share a common suffix. Formally, a payoff  $f cdots Plays(\mathfrak{M}) \to \overline{\mathbb{R}}$  is prefix-independent if for any play  $\pi \in \text{Plays}(\mathfrak{M})$  and any  $r \in \mathbb{N}$ ,  $f(\pi) = f(\pi_{\geq r})$ . We open the section by introducing some useful notation. Given a history  $h = s_0 a_0 s_1 \dots a_{\ell-1} s_\ell$  and a play  $\pi = s_\ell a_\ell l s_{\ell+1}$  such that the last state of h and the first state of  $\pi$  coincide, we let  $h \cdot \pi = s_0 a_0 s_1 \dots a_{\ell-1} s_\ell a_\ell s_{\ell+1} \dots$  denote the play obtained by concatenating h and  $\pi$  without a repetition of  $s_\ell$ . For any play  $\pi = s_0 a_0 s_1 \dots \in \text{Plays}(\mathfrak{M})$ , we let  $\inf(\pi) = \{s \in S \mid \forall r \in \mathbb{N}, \exists \ell \geq r, s_\ell = s\}$  denote the set of states that occur infinitely often in  $\pi$ .

Let  $f \colon \mathsf{Plays}(\mathfrak{M}) \to \mathbb{R}$  be a prefix-independent payoff. Assume that f is continuous. We claim that the payoff of a play  $\pi$  is uniquely determined by  $\mathsf{inf}(\pi)$ . Let  $\pi$  and  $\pi'$  be two plays such that  $\mathsf{inf}(\pi) = \mathsf{inf}(\pi')$ . By continuity of f, to prove that  $f(\pi) = f(\pi')$ , it suffices to show that in all neighbourhoods of  $\pi$ , there is a play whose payoff is  $f(\pi')$ . By prefix-independence of f, we need only establish that in all neighbourhoods of  $\pi$ , there is a play that shares a common suffix with  $\pi'$ . Let  $\ell \in \mathbb{N}$  such that  $\mathsf{last}(\pi_{\leq \ell}) \in \mathsf{inf}(\pi)$ . We construct a play  $\pi''$  in  $\mathsf{Cyl}(\pi_{\leq \ell})$  such that  $f(\pi'') = f(\pi)$  as follows. Since  $\mathsf{inf}(\pi) = \mathsf{inf}(\pi')$ , there exists  $r \in \mathbb{N}$  such that the first state of  $\pi'_{\geq r}$  is  $\mathsf{last}(\pi_{\leq \ell})$ . We thus define  $\pi'' = \pi_{\leq \ell} \cdot \pi'_{\geq r}$ . By prefix-independence of f, we obtain that  $f(\pi'') = f(\pi'_{\geq r}) = f(\pi')$ . This shows that all neighbourhoods of  $\pi$  contain a play whose payoff is  $f(\pi')$  and this closes our argument.

This argument above can be adapted to provide a more general property of continuous prefix-independent payoffs: for any two plays  $\pi$  and  $\pi'$ , if  $\inf(\pi)$  is reachable from  $\inf(\pi')$ , then both plays have the same payoff. We show that this property characterises continuous prefix-independent payoffs.

To formalise our characterisation, we introduce some additional terminology and notation. First, we define Büchi objectives. Given a set  $T \subseteq S$  of target states, the Büchi objective for T requires that states of T be visited infinitely often. Formally, given  $T \subseteq S$ , we define Büchi $T = \{\pi \in \mathsf{Plays}(\mathfrak{M}) \mid \mathsf{inf}(\pi) \cap T \neq \emptyset\}$ . Second, we consider the strongly connected components (SCCs) of (the graph induced by)  $\mathfrak{M}$ . An SCC is a maximal set of states  $C \subseteq S$  such that, for all  $s, t \in C$  there exists a history with at least one action starting in S and ending in S. An SCC S is reachable from an SCC S if there exists a history starting in S and ending in S.

We obtain the following characterisation. For the sake of conciseness, we apply the convention  $0 \cdot (+\infty) = 0 \cdot (-\infty) = 0$  below.

**Lemma B.7.** Let f be a prefix-independent payoff function. Let  $C_1, \ldots, C_k$  be the SCCs of  $\mathfrak{M}$ . Then f is continuous if and only if there exist constants  $\alpha_1, \ldots, \alpha_k \in \overline{\mathbb{R}}$  such that  $f = \sum_{i=1}^k \alpha_i \cdot \mathbb{1}_{\mathsf{B\"{u}chi}(C_i)}$  and, for all  $1 \leq i, i' \leq k$ ,  $\alpha_i = \alpha_{i'}$  whenever  $C_{i'}$  is reachable from  $C_i$ .

*Proof.* We first observe that the set of states visited infinitely often along a play is a subset of an SCC. In other words, for all plays  $\pi \in \mathsf{Plays}(\mathfrak{M})$ , there exists a unique SCC C such that  $\pi \in \mathsf{B\ddot{u}chi}(C)$ ; we say that  $\pi$  stabilises in C for short.

We first assume that f is continuous. Let  $\pi$ ,  $\pi' \in \mathsf{Plays}(\mathfrak{M})$ . We show that if  $\pi$  stabilises in an SCC C and  $\pi'$  stabilises in an SCC C' such that C is reachable from C', then  $f(\pi) = f(\pi')$ . Let  $r \in \mathbb{N}$  such that  $\pi_{\geq r}$  starts in a state of C. For all  $\ell \in \mathbb{N}$ , since C is reachable from C', there exists a play  $\pi^{(\ell)} \in \mathsf{Cyl}(\pi'_{\leq \ell})$  such that  $\pi_{\geq r}$  is a suffix of  $\pi^{(\ell)}$ . By prefix-independence of f, for all  $\ell \in \mathbb{N}$ , we have  $f(\pi^{(\ell)}) = f(\pi)$ . Furthermore, all neighbourhoods of  $\pi'$  contain at least one play  $\pi^{(\ell)}$  by construction. It follows from the continuity of f that  $f(\pi)$  is in all neighbourhoods of  $f(\pi')$ , i.e.,  $f(\pi) = f(\pi')$ .

The previous argument implies that the payoff of a play depends only on the SCC in which the play stabilises. For all  $1 \le i \le k$ , let  $\alpha_i$  be the payoff of any play that stabilises in  $C_i$ . From the above, we obtain that  $f = \sum_{i=1}^k \alpha_i \cdot \mathbb{1}_{\mathsf{B\"{u}chi}(C_i)}$  and that the coefficients  $\alpha_i$  satisfy the required conditions. This ends the proof of the first implication.

We now let  $\alpha_1, \ldots, \alpha_k \in \mathbb{R}$  such that  $f = \sum_{i=1}^k \alpha_i \cdot \mathbb{1}_{\mathsf{B\"{u}chi}(C_i)}$  and, for all  $1 \leq i, i' \leq k$ ,  $\alpha_i = \alpha_{i'}$  whenever  $C_{i'}$  is reachable from  $C_i$ . We show that f is continuous. Let  $\pi \in \mathsf{Plays}(\mathfrak{M})$ . It suffices to show that f is constant over a neighbourhood of  $\pi$ . Let  $1 \leq i \leq k$  such that  $\pi$  stabilises in  $C_i$  and let  $r \in \mathbb{N}$  such that all states of  $\pi_{\geq r}$  are in  $C_i$ . Let  $\pi' \in \mathsf{Cyl}(\pi_{\geq r})$ . Then  $\pi'$  remains in an SCC that is reachable from  $C_i$ , thus  $f(\pi') = \alpha_i = f(\pi)$ . We have shown that f is constant over  $\mathsf{Cyl}(\pi_{\geq r})$ , ending the proof of the second implication.

# C Universally square integrable shortest-path payoffs

In this appendix, our goal is to prove that all universally integrable shortest-path payoffs are universally square integrable. In particular, this implies that Theorem 7.5 is applicable to universally integrable continuous shortest-path payoffs. We fix an MDP  $\mathfrak{M} = (S, A, \delta)$  for this entire section.

Let  $T \subseteq S$  be a set of vertices. We show that the payoff  $\mathsf{SPath}_w^T$  is universally square integrable for all weight functions  $w \colon S \times A \to \mathbb{R}$  if and only if, for all strategies  $\tau$  and

all initial states  $s \in S$ , we have  $\mathbb{P}_s^{\tau}(\mathsf{Reach}(T)) = 1$ . For any weight function w, if  $\mathsf{SPath}_w^T$  is universally (square) integrable, then the set of plays with payoff  $+\infty$  has zero  $\mathbb{P}_s^{\sigma}$ -probability for all  $\sigma$  and  $s \in S$ , i.e., a target state is reached almost surely no matter the strategy and initial state.

To show the converse, it suffices to show that  $\mathsf{SPath}_1^T$ , i.e., the shortest-path payoff where all weights are 1, is universally square integrable, because all shortest-path payoffs are bounded in absolute value by a multiple of  $\mathsf{SPath}_1^T$  (as there are finitely many weights). Because all weights are 1, the expectation of  $(\mathsf{SPath}_1^T)^2$  from  $s \in S$  under a strategy  $\sigma$  can be written as a series over  $r \in \mathbb{N}$  of  $r^2$  multiplied by the  $\mathbb{P}_r^\sigma$ -probability of reaching T after exactly r actions are used. Convergence of this series is guaranteed by the fact that the sequence of probabilities of reaching the target after exactly r actions is in  $\mathcal{O}(\alpha^r)$  for some  $\alpha \in [0,1[$ . This asymptotic bound on these probabilities follows from the linear convergence of value iteration for reachability in MDPs of a specific form [HM18].

The formal proof below requires the notion of end-components of MDPs. An end-component of  $\mathfrak{M}$  is a pair  $\mathcal{E} = (E, A_{\mathcal{E}})$  such that (i)  $E \subseteq S$  is a non-empty set of states, (ii)  $A_{\mathcal{E}} \colon E \to 2^A \setminus \{\emptyset\}$  is a mapping assigning, to all states  $s \in E$ , a non-empty set of actions such that for all  $a \in A_{\mathcal{E}}(s)$ ,  $\operatorname{supp}(\delta(s, a)) \subseteq E$  and (iii) from any  $s, s' \in E$ , there exists a history  $s_0 a_0 s_1 \dots a_{r-1} s_r$  with  $r \geq 1$  such that  $s_0 = s$ ,  $s_r = s'$  and  $a_\ell \in A_{\mathcal{E}}(s_\ell)$  for all  $0 \leq \ell \leq r-1$ . Condition (iii) means that an end-component is strongly connected.

**Lemma C.1.** Let  $T \subseteq S$  be a set of targets and  $s_{\mathsf{init}} \in S$  be an initial state. We have  $\mathbb{P}^{\sigma}_{s_{\mathsf{init}}}(\mathsf{Reach}(T)) = 1$  for all strategies  $\sigma$  if and only if, for all strategies  $\sigma$  and all weight functions  $w \colon S \times A \to \mathbb{R}$ ,  $\mathsf{SPath}^T_w$  is  $\mathbb{P}^{\sigma}_{s_{\mathsf{init}}}$ -square integrable. In particular, for any weight function w, if  $\mathsf{SPath}^T_w$  is universally integrable, then  $\mathsf{SPath}^T_w$  is universally square integrable.

*Proof.* If all shortest-path payoffs with target T are  $\mathbb{P}^{\sigma}_{s_{\text{init}}}$ -integrable for all strategies  $\sigma$ , then under all strategies and from all initial states, the set of plays with payoff  $+\infty$  (which is independent of the weight function) must have probability zero, i.e., the set of targets is reached almost-surely. This implies that the equivalence stated in the lemma entails the claim made in particular. In the remainder of the proof, we assume that for all strategies  $\sigma$ , we have  $\mathbb{P}^{\sigma}_{s_{\text{init}}}(\mathsf{Reach}(T)) = 1$  and show that for all weight functions w and strategies  $\sigma$ ,  $\mathsf{SPath}^T_w$  is  $\mathbb{P}^{\sigma}_{s_{\text{init}}}$ -square integrable.

If  $s_{\mathsf{init}} \in T$ , then for all plays  $\pi \in \mathsf{Cyl}(s_{\mathsf{init}})$ ,  $\mathsf{SPath}_w^T(\pi) = 0$ . Therefore, we directly obtain that  $\mathsf{SPath}_w^T$  is  $\mathbb{P}^{\sigma}_{s_{\mathsf{init}}}$ -square integrable for all strategies  $\sigma$ . We thus assume that  $s_{\mathsf{init}} \notin T$ .

We assume without loss of generality that all states of  $\mathfrak{M}$  are reachable from  $s_{\mathsf{init}}$  (i.e., for all states  $s \in S$ , there exists a history starting in  $s_{\mathsf{init}}$  and ending in s). Furthermore, we also assume that there is a unique absorbing target state, i.e., that  $T = \{t\}$  such that for all  $a \in A(t)$ ,  $\delta(t,a)(t) = 1$ . We can always reduce to this case by considering the MDP  $\mathfrak{M}'$  obtained from  $\mathfrak{M}$  by merging all target states into a single absorbing state in which all actions are enabled. For any strategy  $\sigma$  and all weight functions w,  $\mathsf{SPath}_w^T$  is  $\mathbb{P}_{s_{\mathsf{init}}}^\sigma$ -square integrable if and only if its counterpart in  $\mathfrak{M}'$  is  $\mathbb{P}_{s_{\mathsf{init}}}^{\sigma'}$ -square integrable, where  $\sigma' \in \Sigma(\mathfrak{M}')$  agrees with  $\sigma$  over  $\mathsf{Hist}(\mathfrak{M}) \cap \mathsf{Hist}(\mathfrak{M}')$ .

We now fix a strategy  $\sigma$ . First, we consider the constant weight function w=1 and show that  $\mathsf{SPath}_1^T$  is  $\mathbb{P}^\sigma_{s_{\mathsf{init}}}$ -square integrable. The general case follows from this one.

We preface the argument with some notation. For all  $r \in \mathbb{N}$ , we let  $\mathsf{Reach}^{\leq r}(T) = \{s_0a_0s_1\ldots\in\mathsf{Plays}(\mathfrak{M})\mid \exists\,\ell\leq r,\,s_\ell\in T\}$  denote the set of plays in which a target is reached in at most r transitions. For all  $r\in\mathbb{N}_0$ , we let  $\mathsf{Reach}^{=r}(T)=\mathsf{Reach}^{\leq r}(T)\setminus\mathsf{Reach}^{\leq r}(T)$  and  $\mathsf{Reach}^{=0}(T)=\mathsf{Reach}^{\leq 0}(T)$ . Given  $\Omega\in\{\mathsf{Reach}(T)\}\cup\{\mathsf{Reach}^{\leq r}(T)\mid r\in T\}$ 

 $\mathbb{N}$ }, we let,  $\mathbb{P}^{\min}_{s_{\mathsf{init}}}(\Omega) = \min_{\tau \in \Sigma(\mathfrak{M})} \mathbb{P}^{\tau}_{s_{\mathsf{init}}}(\Omega)$  denote the least probabilities that strategies can obtain for  $\Omega$  from s. These minima are well-defined and we have  $(\mathbb{P}^{\min}_{s_{\mathsf{init}}}(\mathsf{Reach}^{\leq r}(T)))_{r \in \mathbb{N}}$  is an increasing sequence that converges to  $\mathbb{P}^{\min}_{s_{\mathsf{init}}}(\mathsf{Reach}(T)) = 1$  (see, e.g., [BK08]).

The equality  $\mathbb{P}_{s_{\text{init}}}^{\sigma}(\mathsf{Reach}(T)) = 1$  implies that

$$\mathbb{E}^{\sigma}_{s_{\mathrm{init}}}((\mathsf{SPath}_1^T)^2) = \sum_{r \in \mathbb{N}} r^2 \cdot \mathbb{P}^{\sigma}_{s_{\mathrm{init}}}(\mathsf{Reach}^{=r}(T)).$$

It suffices to show that there exists constants  $\alpha \in ]0,1[$  and  $\beta \in [0,+\infty[$  such that for all  $r \in \mathbb{N}$ ,  $\mathbb{P}^{\sigma}_{s_{\text{init}}}(\mathsf{Reach}^{=r}(T)) \leq \beta \cdot \alpha^r$  to obtain convergence. Indeed, the convergence of the above series is implied by that of the series  $\sum_{r \in \mathbb{N}} \beta \cdot r^2 \cdot \alpha^r$ , whose convergence is guaranteed by Cauchy's convergence test for non-negative series (because  $\lim_{r \to \infty} \sqrt[r]{\beta \cdot r^2 \cdot \alpha^r} = \alpha < 1$ ).

We now seek the constants  $\alpha$  and  $\beta$  satisfying the above condition. We obtain from  $\mathbb{P}^{\sigma}_{s_{\text{init}}}(\mathsf{Reach}(T)) = \mathbb{P}^{\min}_{s_{\text{init}}}(\mathsf{Reach}(T)) = 1$  that, for all  $r \in \mathbb{N}$ , we have

$$\begin{split} \mathbb{P}^{\sigma}_{s_{\mathsf{init}}}(\mathsf{Reach}^{=r}(T)) &\leq \mathbb{P}^{\sigma}_{s_{\mathsf{init}}}(\mathsf{Reach}(T) \setminus \mathsf{Reach}^{\leq r-1}(T)) \\ &= \mathbb{P}^{\sigma}_{s_{\mathsf{init}}}(\mathsf{Reach}(T)) - \mathbb{P}^{\sigma}_{s_{\mathsf{init}}}(\mathsf{Reach}^{\leq r-1}(T)) \\ &\leq \mathbb{P}^{\min}_{s_{\mathsf{init}}}(\mathsf{Reach}(T)) - \mathbb{P}^{\min}_{s_{\mathsf{init}}}(\mathsf{Reach}^{\leq r-1}(T)). \end{split}$$

To bound the last term, we prove that  $\mathfrak{M}$  is min-reduced in the sense of [HM18]. An MDP is min-reduced if there are two states  $s^+$  and  $s^-$  such that  $T = \{s^+\}$ ,  $s^- \neq s^+$  and all end-components have a singleton  $\{s^+\}$  or  $\{s^-\}$  as their set of states. In our case, the only end-components of  $\mathfrak{M}$  are of the form  $(\{t\}, A')$  where  $A' \subseteq A$  is non-empty. By contradiction, assume that there is an end-component  $\mathcal{E} = (E, A_{\mathcal{E}})$  such that  $E \neq \{t\}$ . Then  $t \notin E$  because t is absorbing and an end-component is strongly connected. Because E is be reachable from  $s_{\text{init}}$ , there exists a strategy reaching E with positive probability and then surely remaining in E from there (by only using actions authorised by  $A_{\mathcal{E}}$ ). This contradicts the fact that for all strategies  $\tau$ ,  $\mathbb{P}^{\tau}_{s_{\text{init}}}$  (Reach(T)) = 1.

contradicts the fact that for all strategies  $\tau$ ,  $\mathbb{P}^{\tau}_{s_{\text{init}}}(\mathsf{Reach}(T)) = 1$ . It follows from [HM18, Proof of Thm. 2] that there exists  $I \in \mathbb{N}_0$  such that, for all  $\ell \in \mathbb{N}$ ,  $\mathbb{P}^{\min}_{s_{\text{init}}}(\mathsf{Reach}(T)) - \mathbb{P}^{\min}_{s_{\text{init}}}(\mathsf{Reach}^{\leq \ell I}(T)) \leq (1 - \eta^I)^{\ell}$  where  $\eta$  is the smallest positive transition probability in  $\mathfrak{M}$ . Let  $r \in \mathbb{N}$ ,  $\ell \in \mathbb{N}$  and  $0 \leq p < I$  such that  $r = \ell I + p$ . We obtain, using the fact that the sequence  $(\mathbb{P}^{\min}_{s_{\text{init}}}(\mathsf{Reach}^{\leq r'}(T)))_{r' \in \mathbb{N}}$  is non-decreasing, that

$$\begin{split} \mathbb{P}^{\min}_{s_{\mathsf{init}}}(\mathsf{Reach}(T)) - \mathbb{P}^{\min}_{s_{\mathsf{init}}}(\mathsf{Reach}^{\leq r}(T)) &\leq \mathbb{P}^{\min}_{s_{\mathsf{init}}}(\mathsf{Reach}(T)) - \mathbb{P}^{\min}_{s_{\mathsf{init}}}(\mathsf{Reach}^{\leq \ell I}(T)) \\ &\leq (1 - \eta^I)^{\ell} \\ &= \left(\sqrt[I]{1 - \eta^I}\right)^{r - p} \\ &\leq \left(\sqrt[I]{1 - \eta^I}\right)^{r - (I - 1)}. \end{split}$$

We conclude that it suffices to set  $\alpha = \sqrt[I]{1 - \eta^I}$  and  $\beta = \left(\sqrt[I]{1 - \eta^I}\right)^{-I}$  to conclude the proof that  $\mathsf{SPath}_1^T$  is  $\mathbb{P}_{s_{\mathsf{init}}}^\sigma$ -square integrable.

We end the proof by establishing that for all weight functions w, the shortest-path payoff  $\mathsf{SPath}_w^T$  is  $\mathbb{P}_{s_{\mathsf{init}}}^\sigma$ -square integrable. Let  $W = \max_{(s,a) \in S \times A} |w(s,a)|$ . By the triangular inequality, we obtain that  $|\mathsf{SPath}_w^T| \leq W \cdot \mathsf{SPath}_1^T$ . We have thus bounded  $\mathsf{SPath}_w^T$  in absolute value by a multiple of a  $\mathbb{P}_{s_{\mathsf{init}}}^\sigma$ -square integrable function, thus implying that  $\mathsf{SPath}_w^T$  is also  $\mathbb{P}_{s_{\mathsf{init}}}^\sigma$ -square integrable.