

Characterisation of circular light rays in a plasma

Volker Perlick

Faculty 1, University of Bremen
28359 Bremen, Germany
Email: perlick@uni-bremen.de

Abstract. It is the purpose of this paper to give a characterisation of circular light rays in a plasma on an axially symmetric and stationary spacetime. We restrict to the case of an unmagnetised, pressure-free electron-ion plasma and we assume that the plasma shares the symmetry of the spacetime. As a main tool we use two potentials, one for prograde and one for retrograde light rays, whose critical points are exactly the circular light rays in the plasma. In the case that the plasma density vanishes, the corresponding equipotential surfaces reduce to the relativistic Von Zeipel cylinders which have been discussed in many papers since the 1970s. In a plasma, the gradients of the potentials give the centrifugal and the Coriolis forces experienced by a light ray, where the plasma has an influence only on the centrifugal force. The introduction of these potentials allows us to generalise topological methods that have been successfully used for proving the existence or non-existence of circular vacuum light rays to the plasma case. The general results are illustrated with examples on Minkowski, Schwarzschild, Kerr and NUT spacetimes.

1. Introduction

If gravitating objects are sufficiently compact, the light-bending effect of their gravitational fields may be so strong that some light rays are circular. Examples for such objects, which are sometimes called *ultracompact*, are black holes and wormholes and there are also speculations that ultracompact stars might exist. The existence of circular light rays has important consequences for the light-bending effects in general. In particular, circular light rays that are unstable against perturbations in the radial direction are associated with light rays that asymptotically spiral towards them, giving rise to infinite bending angles and to the formation of a *shadow*. For reviews on the latter we refer to Cunha and Herdeiro [1] and to Perlick and Tsupko [2].

When discussing the light bending of gravitating objects, it is justified in most cases to treat the light rays as lightlike geodesics of the spacetime metric, i.e., to disregard any influence of a medium. For some applications, however, this influence might be non-negligible. This is true, in particular, in the radio regime if the medium is a sufficiently dense plasma. E.g., for a light ray that crosses the ionosphere of our Earth or the Solar corona, it is known that there is a measurable influence of the medium on the travel time and on the spatial path of a light ray unless the frequency is much bigger than 10

MHz or 100 MHz, respectively. It is not unplausible to assume that near a black hole the plasma density might be even higher, i.e., to assume that near a black hole light rays of even higher frequencies could be influenced by a plasma in a non-negligible way. This gives a good motivation for investigating the influence of a plasma on the lensing features of an ultracompact object.

Gravitational light deflection on a general-relativistic spacetime in the presence of a plasma has been investigated by many authors. The first papers on this subject date back to the 1960s when the Shapiro time delay of light rays (in the radio regime) that pass through the Solar corona was at the centre of interest. Muhleman et al. [3, 4] calculated, on the basis of the linearised Schwarzschild metric, the time delay and the bending angle of such light rays. Light bending in a plasma, without the weak-field approximation, on the Schwarzschild spacetime and in the equatorial plane of a Kerr metric, was first calculated by Perlick [5]. This was followed by many other papers on light bending by a gravitational field in the presence of a plasma. Here, in particular the work by Bisnovatyi-Kogan and Tsupko [6, 7] has to be mentioned, which considerably pushed the subject, and two papers by Perlick and Tsupko [8, 9] where the influence of a plasma on lensing by a Kerr black hole was discussed in detail. In all these papers, the simplest model of a plasma was used, namely a non-magnetised pressure-less electron-ion plasma.

It is the purpose of the present paper to present a mathematical formalism that allows to investigate the existence of circular light rays in the presence of a plasma. As outlined in the preceding paragraphs, this is motivated by the two facts that the existence of circular light rays is of crucial relevance for the light bending and that the influence of a plasma might be non-negligible in some cases of interest. To that end, we restrict to axially symmetric and stationary spacetimes and we assume that the plasma shares these symmetries. Moreover, as in the works quoted above, we restrict to the simple case of a non-magnetised pressure-less electron-ion plasma. As a main tool, we define two potentials, one for prograde and one for retrograde light rays, in such a way that the critical points of these potentials give the locations of circular light rays. If there is no plasma, the corresponding equipotential surfaces reduce to the *relativistic Von Zeipel cylinders* which were introduced by Marek Abramowicz, originally for axially symmetric and static spacetimes, already in the 1970s and have been discussed in numerous papers, see in particular Abramowicz [10, 11, 12]. These equipotential surfaces have many interesting properties. What is most important for our purpose is the fact that they give us the circular light rays which are located exactly where the potentials have critical points. The generalised potentials that will be introduced in this paper have the same crucial property in the plasma case.

With the help of these potentials we can visualise the behaviour of light rays in an instructive way: If a light ray is sent tangent to a circle, the gradient of the corresponding potential points in the direction in which this light ray is deflected from this circle. The corresponding “force” acting on the light ray can be interpreted as the sum of a “centrifugal force” and a “Coriolis force”. Interestingly, we will see that only the

centrifugal force is influenced by the plasma. However, as already mentioned it is most important that the location of circular light rays is given by the critical points of the corresponding potential. In addition to just visualising the light-deflection properties in a suggestive way, this mathematical construction allows us to generalise theorems on the existence of circular light rays that have been established with topological methods, using the Brouwer degree of appropriate maps, by Cunha et al. [13, 14, 15], to the plasma case.

It should be emphasised that throughout the paper Einstein's field equation is not used, so the general results apply to the case that the plasma is self-gravitating and equally well to the case that the gravitational field of the plasma is neglected. The latter case is, probably, more interesting in view of applications to astrophysics, so in our examples we consider only vacuum solutions to Einstein's field equation, namely Minkowski, Schwarzschild, Kerr and NUT spacetimes.

In Section 2 we briefly review the Hamiltonian formalism for light rays in a non-magnetised pressure-less electron-ion plasma on an unspecified general-relativistic spacetime. In Section 3 we specialise to the axially symmetric and stationary case. In particular, we characterise, in Proposition 1, light rays that are tangent to a circle at one point. This proposition is crucial for the rest of the paper. On the basis of this proposition, we introduce in Section 4 two potentials \mathcal{R}_\pm . Proposition 1 makes sure that these potentials determine the centrifugal and the Coriolis force acting on light rays in the plasma and that, in particular, the critical points determine the locations of circular light rays. In Section 5 we discuss the properties of these two potentials, in particular their limiting behaviour if a horizon, a regular axis or an asymptotic flat end is approached. In Section 6 we point out how the preceding results allow to generalise the above-quoted work of Cunha et al. to the plasma case. Finally, in Sections 7 and 8 we illustrate the general results with several examples.

Throughout, we use Einstein's summation convention for greek indices μ, ν, \dots that take values 0, 1, 2, 3. When specialising to axially symmetric and static spacetimes, we will occasionally use Einstein's summation convention for upper case latin indices A, B, \dots that take values 0, 1 and for lower case latin indices i, j, \dots that take values 2, 3. We use units making the vacuum speed of light c equal to 1 and we assume that the spacetime has signature $(-+++)$.

2. Light rays in a plasma

Light rays in a non-magnetised and pressure-less electron-ion plasma on a general-relativistic spacetime with metric $g_{\mu\nu}$ are the solutions of Hamilton's equations

$$\dot{x}^\mu = \frac{\partial \mathcal{H}}{\partial p_\mu}, \quad \dot{p}^\mu = -\frac{\partial \mathcal{H}}{\partial x_\mu}, \quad \mathcal{H} = 0 \quad (1)$$

with the Hamiltonian

$$\mathcal{H}(x, p) = \frac{1}{2} \left(g^{\rho\sigma}(x) p_\rho p_\sigma + \omega_p(x)^2 \right). \quad (2)$$

Here ω_p is a non-negative function of the spacetime coordinates with the dimension of a frequency. It is known as the *plasma frequency*. Its square ω_p^2 equals, up to a constant factor that depends on the system of units used, the number density of electrons. We will refer to ω_p^2 as to the *plasma density* in the following.

The overdot denotes the derivative with respect to a curve parameter s . This parameter has no particular physical meaning. It is *not* an affine parameter, i.e., the equations are *not* invariant under affine changes of this parameter, unless $\omega_p = 0$.

For a derivation from Maxwell's equations of this Hamiltonian formalism for light rays in a plasma on a curved spacetime we refer to Breuer and Ehlers [16] who even treated the case of a magnetised plasma. For a non-magnetised plasma a derivation can also be found in a book by Perlick [5]. A non-magnetised pressure-less plasma belongs to the class of isotropic dispersive media which have been treated first in a book by Synge [17].

Written out explicitly, the equations (1) with the Hamiltonian (2) read

$$\dot{x}^\mu = g^{\mu\sigma} p_\sigma, \quad (3)$$

$$\dot{p}_\mu = -\frac{1}{2} \left(\frac{\partial g^{\rho\sigma}}{\partial x^\mu} p_\rho p_\sigma + \frac{\partial \omega_p^2}{\partial x^\mu} \right), \quad (4)$$

$$\mathcal{H} = \frac{1}{2} \left(g^{\rho\sigma} p_\rho p_\sigma + \omega_p^2 \right) = 0. \quad (5)$$

Eq. (3) is equivalent to

$$p_\nu = g_{\nu\rho} \dot{x}^\rho, \quad (6)$$

so (5) can be rewritten as

$$\mathcal{H} = \frac{1}{2} \left(p_\rho \dot{x}^\rho + \omega_p^2 \right) = 0 \quad (7)$$

or

$$\mathcal{H} = \frac{1}{2} \left(g_{\rho\sigma} \dot{x}^\rho \dot{x}^\sigma + \omega_p^2 \right) = 0. \quad (8)$$

From (8) we read that light rays in a plasma are *timelike* curves. More precisely, they are *timelike geodesics* of the conformally rescaled metric $\omega_p^2 g_{\mu\nu}$, as can be easily verified. The parameter s is proper time with respect to this (unphysical) metric.

If we choose an observer field, i.e., a timelike vector field U^μ with $g_{\mu\nu} U^\mu U^\nu = -1$, we can assign a frequency

$$\omega(s) = -p_\mu(s) U^\mu(x(s)) \quad (9)$$

to every light ray. The minus sign makes sure that ω is positive for a light ray that is future-oriented with respect to U^μ . In vacuum, the frequency of a light ray has no influence on its path. In a plasma, however, the dispersion relation (5) is inhomogeneous (i.e., not invariant under multiplication of p_μ with a non-zero factor), which implies that light rays that start in the same spatial direction with different frequencies have different trajectories. This inhomogeneity is the defining property of a *dispersive medium*, see e.g. Perlick [5] for a detailed discussion.

3. The axially symmetric and stationary case

We now specify to the case that the metric is axially symmetric and stationary and that the plasma density shares this symmetry. This means that the metric can be written in coordinates $(x^0 = t, x^1 = \varphi, x^2, x^3)$ such that

$$\frac{\partial g_{\mu\nu}}{\partial x^A} = 0, \quad \frac{\partial \omega_p^2}{\partial x^A} = 0. \quad (10)$$

Here and in the following we use the summation convention for upper-case latin indices $A, B, \dots = 0, 1$ and for lower-case latin indices $i, j, \dots = 2, 3$. The x^i could be, e.g., spherical polar coordinates (r, ϑ) or cylindrical polar coordinates (ρ, z) . As before, greek indices take values $0, 1, 2, 3$. We assume that the metric coefficients g_{AB} and the plasma density ω_p^2 are smooth (C^2 would do) on the domain under consideration.

We will assume in the following that the coordinate function φ is 2π -periodic,

$$(t, \varphi, x^2, x^3) \equiv (t, \varphi + 2\pi, x^2, x^3), \quad (11)$$

and that the two-dimensional surfaces $x^i = \text{constant}$ are timelike, i.e.

$$g_{\varphi\varphi}g_{tt} - g_{t\varphi}^2 < 0. \quad (12)$$

The first condition means that φ can be interpreted as an angular coordinate, the second condition means that there is a linear combination of the Killing vector fields ∂_t and ∂_φ that is timelike. Both assumptions are necessary to justify calling the spacetime ‘‘axially symmetric and stationary’’. Note that in a black-hole spacetime the condition (12) is satisfied in the domain of outer communication, i.e., outside the horizon, while the left-hand side of (12) is equal to zero on the horizon and positive inside.

As the spacetime metric ($g_{\mu\nu}$) has signature $(-+++)$, (12) implies that the (2×2) -matrix (g^{ij}) is positive definite. This will be relevant for the following considerations.

Note that we do *not* require the mixed metric components g_{iA} to be zero. If the metric is invariant under a transformation $(t, \varphi) \mapsto (-t, -\varphi)$, the g_{iA} must be zero. It is indeed true that such a choice of coordinates is possible for most axially symmetric and stationary spacetimes with relevance to physics. However, even in these cases it is nonetheless meaningful to allow for another choice of coordinates where $g_{iA} \neq 0$. An example is the Schwarzschild spacetime in Eddington-Finkelstein or Painlevé-Gullstrand coordinates.

Also note that we do *not* require the spacetime to admit a horizontal plane, i.e., we do *not* require that it is possible to choose cylindrical polar coordinates (ρ, z) such that the metric coefficients are invariant under a transformation $z \mapsto -z$.

Moreover, we will allow the metric functions g_{tt} and $g_{\varphi\varphi}$ to have either sign, i.e., we will allow for the presence of an ergoregion, where $g_{tt} > 0$, and of a causality-violating region, where $g_{\varphi\varphi} < 0$. An ergoregion occurs, e.g., in the domain of outer communication of a Kerr black hole while a causality-violating region occurs, e.g., in the domain of outer communication of a NUT black hole.

The following proposition, which is at the basis of all that follows, characterises light rays that are tangent, at one point, to a surface $x^i = \text{constant}$. Obviously, this

applies in particular to circular light rays which, by definition, are light rays with $\dot{\varphi} \neq 0$ that are completely contained in a surface $x^i = \text{constant}$.

Proposition 1. Let $x^\mu(s)$ be a solution to Hamilton's equations in an axially symmetric and stationary spacetime. Then, at points where $\dot{x}^i = 0$, the following equations are true.

$$p_\varphi = \frac{g_{t\varphi}}{g_{tt}} p_t \mp \frac{1}{g_{tt}} \sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}} \sqrt{p_t^2 + \omega_p^2 g_{tt}} \quad (13)$$

$$\dot{\varphi} = \pm \frac{\sqrt{p_t^2 + \omega_p^2 g_{tt}}}{\sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}}} \quad (14)$$

$$\dot{t} = \frac{p_t}{g_{tt}} \mp \frac{g_{t\varphi}}{g_{tt}} \frac{\sqrt{p_t^2 + \omega_p^2 g_{tt}}}{\sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}}} \quad (15)$$

$$\ddot{x}^\mu = - \frac{\sqrt{p_t^2 + \omega_p^2 g_{tt}}}{\sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}}} g^{\mu i} \frac{\partial}{\partial x^i} \left(\mp \frac{g_{t\varphi}}{g_{tt}} p_t + \frac{1}{g_{tt}} \sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}} \sqrt{p_t^2 + \omega_p^2 g_{tt}} \right) \quad (16)$$

The upper sign is for prograde light rays ($\dot{\varphi} > 0$) and the lower sign for retrograde ones ($\dot{\varphi} < 0$).

The proof of this proposition is given in the appendix. It is a straight-forward generalisation from the case $\omega_p = 0$. For the latter, cf. e.g. Cunha et al. [13].

Proposition 1 gives us all relevant information where in phase space light rays with $\dot{x}^i = 0$ can exist. Firstly, we see that (12) must be satisfied. We have already mentioned that this condition assures that the two-space $x^i = \text{constant}$ contains a timelike vector. In a plasma, where light rays are timelike, this condition is obviously necessary for the existence of a light ray with $\dot{x}^i = 0$, and this is confirmed by the occurrence of the $\sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}}$ terms in (13), (14), (15) and (16). One might argue that in the vacuum case, where the light rays are lightlike, we should also allow the limiting case that the surface $x^i = \text{constant}$ is lightlike, i.e., $g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi} = 0$. It is indeed true that vacuum light rays can be contained in such a surface, but this would be the generators of a horizon and we do not want them to be considered as circular light rays. So we require, also in vacuum, the strict inequality (12).

Secondly, we read from Proposition 1 that we should restrict to the region in phase space where

$$p_t^2 + \omega_p^2 g_{tt} > 0. \quad (17)$$

Again, one might argue about the limiting case that the left-hand side of (17) is equal to zero, but then (14) can hold only with $\dot{\varphi} = 0$ and we do not consider such light rays

as circular. To be sure, it is obvious and well known that light rays with this property do exist in a plasma: In a region with $\omega_p \neq 0$ and $g_{tt} < 0$ there is always a value of $|p_t|$ such that the left-hand side of (17) is zero. Light rays with this frequency are tangent to a t -line, i.e., they “stand still”. This happens, e.g., at the boundary of the Earth’s ionosphere for $|p_t| \approx 10$ MHz. A light ray with a lower frequency is reflected from the boundary of the ionosphere. As we do not consider a light ray as “circular” if it stands still, we require that (17) holds with the strict inequality sign.

Thirdly, the occurrence of factors of g_{tt} in the denominators of (13), (15) and (16) requires to investigate what happens if $g_{tt} = 0$. If we multiply in (13) and (15) both sides with g_{tt} and use the fact that p_φ and \dot{t} must be finite, we see that in the limit $g_{tt} \rightarrow 0$ the condition $g_{t\varphi}p_t = \pm|g_{t\varphi}p_t|$ must hold. As $g_{tt} = 0$ requires, by (12) and (17), that $g_{t\varphi} \neq 0$ and $p_t \neq 0$, this condition holds for only one of the two signs, i.e., at points where $g_{tt} = 0$ only a prograde or a retrograde circular light ray can exist, but not both. This gives us the following third restriction on the phase space:

$$g_{tt}^2 + p_t^2 \left(\pm g_{t\varphi} + |g_{t\varphi}| \right)^2 > 0, \quad (18)$$

where the upper sign gives the restriction for prograde and the lower sign for retrograde light rays.

4. Definition of the potentials R_\pm

As a plasma is a dispersive medium, the propagation of light rays depends on their frequencies. In a stationary spacetime, where p_t is a constant of motion, it is convenient to consider all light rays with fixed p_t . Henceforth, we write

$$p_t = -\omega_0 \quad (19)$$

and it is our goal to characterise all circular light rays with fixed ω_0 . Note that in an asymptotically flat spacetime, where

$$U^\mu = \frac{1}{\sqrt{-g_{tt}}} \delta_t^\mu \longrightarrow \delta_t^\mu \quad (20)$$

at infinity, ω_0 is the frequency (9) with respect to this observer field U^μ at infinity (provided that the light ray under consideration reaches infinity). We have chosen the sign in (19) such that in regions where ∂_t is timelike ω_0 is positive for light rays that are future-oriented with respect to ∂_t .

For any fixed $\omega_0 \geq 0$, we now introduce the following potentials \mathcal{R}_+ and \mathcal{R}_- , which is motivated by Proposition 1.

$$\mathcal{R}_\pm = \pm \frac{g_{t\varphi}}{g_{tt}} \omega_0 + \frac{1}{g_{tt}} \sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}} \sqrt{\omega_0^2 + \omega_p^2 g_{tt}}. \quad (21)$$

Restricting to values $\omega_0 \geq 0$ is justified because the potentials for $-\omega_0$ are the same as for ω_0 , just interchanged. As we will outline in Section 5.3 below, this convention has the consequence that inside an ergoregion the upper sign in our equations corresponds to a future-oriented parametrisation of the light rays and the lower sign to a past-oriented

one, or vice versa. Note that in a plasma it is indeed possible that a circular light ray with $\omega_0 = 0$ exists: In vacuum, (17) excludes the case $p_t = 0$; in a plasma, however, $\omega_p^2 g_{tt}$ is positive in an ergoregion, so the case $p_t = 0$ is not forbidden by (17). Strictly speaking, \mathcal{R}_\pm should carry an index ω_0 . We do not do this, to ease notation, so we have to keep in mind that \mathcal{R}_\pm is to be considered for a fixed $\omega_0 \geq 0$.

With the help of these potentials \mathcal{R}_\pm , (13) can be rewritten as

$$p_\varphi = \mp \mathcal{R}_\pm \quad (22)$$

and (16) can be rewritten, for $\mu = j$, as

$$\ddot{x}^j = - \frac{\sqrt{\omega_0^2 + \omega_p^2 g_{tt}}}{\sqrt{g_{t\varphi}^2 - g_{tt} g_{\varphi\varphi}}} g^{ji} \frac{\partial \mathcal{R}_\pm}{\partial x^i}. \quad (23)$$

As the 2×2 matrix (g^{ij}) is invertible (and even positive definite), this equation shows that a light ray that is sent tangential to a circle $x^i = \text{constant}$ is experiencing a “force” that is a negative multiple of the gradient of \mathcal{R}_\pm . In particular, it implies that a circular light ray exists at $x^i = x_0^i$ if and only if \mathcal{R}_\pm has a critical point at $x^i = x_0^i$. Defining the potential such that the direction of the force is opposite to the gradient of the potential is the usual convention in mechanics which implies that a stable critical point corresponds to a local minimum of the potential.

Drawing the surfaces $\mathcal{R}_\pm = \text{constant}$ illustrates the way in which light rays are deflected and, in particular, where circular light rays exist. These equipotential surfaces generalise the *relativistic Von Zeipel cylinders* which were introduced by Abramowicz [10, 11] already in the 1970s and have been discussed in many papers. To make the connection, we specify \mathcal{R}_\pm to the case that $g_{t\varphi} = 0$ and $\omega_p = 0$. By (12), $g_{t\varphi} = 0$ implies $g_{tt} g_{\varphi\varphi} < 0$ and, if we exclude causality violation, $g_{tt} < 0$, hence

$$\mathcal{R}_+ = \mathcal{R}_- = - \sqrt{- \frac{g_{\varphi\varphi}}{g_{tt}}} \omega_0 \quad (24)$$

which is exactly the potential whose equipotential surfaces have been called the *relativistic Von Zeipel cylinders* in axisymmetric and static spacetimes, see e.g. Abramowicz [12]. (For vacuum light rays, these equipotential surfaces are, of course, independent of the frequency constant ω_0 .) In flat spacetime, where we can choose cylindrical polar coordinates such that $g_{tt} = -1$ and $g_{\varphi\varphi} = \rho^2$, these equipotential surfaces are indeed straight cylinders in three-dimensional space, i.e., they are represented by vertical lines in the $(x^2, x^3) = (\rho, z)$ half-plane. In curved spacetimes, however, they do not always have the topology of a cylinder. Therefore, the name should be taken with a grain of salt.

As shown by Abramowicz [12], particles in *timelike* motion along a circle $x^i = \text{constant}$ experience a *centrifugal force* that points in the direction of the negative gradient of the potential (24), with a prefactor that depends on the velocity. The potential itself is actually independent of the velocity, so the interpretation as a potential for the centrifugal force is also valid for particles that move at the speed of light (although

in this case the prefactor becomes infinite). In the general case, allowing for a non-zero $g_{t\varphi}$ and a non-zero ω_p , the first term on the right-hand side of (21) can be viewed as a potential for the Coriolis force and the second one as a potential for the centrifugal force. Note that the plasma only influences the centrifugal force. For a discussion of Von Zeipel cylinders in the case that $\omega_p = 0$ but $g_{t\varphi} \neq 0$ we refer to Jefremov and Perlick [18] where two possible definitions are compared. One of them corresponds to our potential (21) specified to $\omega_p = 0$, the other one, denoted $\tilde{\mathcal{R}}$ in [18], is of no relevance for the present paper.

5. Properties of the potentials \mathcal{R}_{\pm}

5.1. Domain of definition of the potentials \mathcal{R}_{\pm}

From the discussion at the end of Section 3 we know that circular light rays can exist only where (12), (17) and (18) hold. Correspondingly, we consider the potential \mathcal{R}_{\pm} , for fixed $\omega_0 \geq 0$, on the open subset of the (x^2, x^3) -plane where

$$g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi} > 0, \quad (25)$$

$$g_{tt}^2 + \omega_0^2 \left(\pm g_{t\varphi} + |g_{t\varphi}| \right)^2 > 0, \quad (26)$$

$$\omega_0^2 + \omega_p^2 g_{tt} > 0. \quad (27)$$

If this set is not connected, we have to consider the potential \mathcal{R}_{\pm} on each connected component separately. Let us choose one of these connected components, denoted \mathcal{V}_{\pm} . In the following we will discuss the various possibilities for \mathcal{V}_{\pm} . We first observe that the open set where (25) holds need not be connected. We denote the connected component that contains our chosen \mathcal{V}_{\pm} by the letter \mathcal{U} . We call the open subset of \mathcal{U} where (26) holds \mathcal{U}_{\pm} .

If $g_{tt} < 0$ on all of \mathcal{U} , (26) gives no further restriction, hence $\mathcal{U}_+ = \mathcal{U}_- = \mathcal{U}$. If $-\omega_p^2 g_{tt}$ is bounded on \mathcal{U} , i.e. if

$$\sup(-\omega_p^2 g_{tt}) < \infty, \quad (28)$$

\mathcal{R}_+ and \mathcal{R}_- are defined on all of \mathcal{U} if the frequency constant ω_0 has been chosen such that $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$, i.e. $\mathcal{V}_+ = \mathcal{V}_- = \mathcal{U}$. For smaller values of ω_0 , the sets \mathcal{V}_+ and \mathcal{V}_- are further restricted; on the boundary of these restricted domains light rays are tangent to a t -line, i.e., they “stand still”.

If $g_{tt} > 0$ on all of \mathcal{U} , it is again true that (26) gives no further restriction, hence $\mathcal{U}_+ = \mathcal{U}_- = \mathcal{U}$, hence $\mathcal{V}_+ = \mathcal{V}_- = \mathcal{U}$ for all $\omega_0 > 0$ and in the case that ω_p is bounded away from zero on \mathcal{U} even for $\omega_0 = 0$.

The situation is more complicated if g_{tt} takes positive and negative values on \mathcal{U} . In this case we decompose \mathcal{U} into the sets

$$\mathcal{U}^{\text{out}} = \{(x^2, x^3) \in \mathcal{U} \mid g_{tt}(x^2, x^3) < 0\}, \quad (29)$$

$$\mathcal{U}^{\text{erg}} = \{(x^2, x^3) \in \mathcal{U} \mid g_{tt}(x^2, x^3) > 0\}, \quad (30)$$

$$\mathcal{S} = \{(x^2, x^3) \in \mathcal{U} \mid g_{tt}(x^2, x^3) = 0\}. \quad (31)$$

\mathcal{U}^{out} and \mathcal{U}^{erg} are open while \mathcal{S} is closed in \mathcal{U} . If \mathcal{U} is the domain of outer communication of a Kerr black hole, \mathcal{U}^{erg} is the ergoregion, \mathcal{U}^{out} is the region outside the ergoregion, and \mathcal{S} is the boundary of the ergoregion in \mathcal{U} .

Because of (25) $g_{t\varphi}$ must be non-zero on \mathcal{S} . Let us assume that $g_{t\varphi} < 0$ on \mathcal{S} . Then, by (26), $\mathcal{U}_+ = \mathcal{U}$ and $\mathcal{U}_- = \mathcal{U}^{\text{out}} \cup \mathcal{U}^{\text{erg}}$. If (28) holds on \mathcal{U} the potential \mathcal{R}_+ is defined on the domain $\mathcal{V}_+ = \mathcal{U}$ for all $\omega_0 > \sup(-\omega_p^2 g_{tt})$. For \mathcal{R}_- we have to consider the domains \mathcal{U}^{out} and \mathcal{U}^{erg} separately. If (28) holds on \mathcal{U}^{out} , then \mathcal{R}_- is defined on all of \mathcal{U}^{out} for all $\omega_0 > \sup(-\omega_p^2 g_{tt})$, so for these values of ω_0 one possibility for our chosen domain of definition is $\mathcal{V}_- = \mathcal{U}^{\text{out}}$. On \mathcal{U}^{erg} the potential \mathcal{R}_- is defined for all $\omega_0 > 0$ and in the case that ω_p is bounded away from zero on this domain even for $\omega_0 = 0$, so for these values of ω_0 our chosen domain of definition can be $\mathcal{V}_- = \mathcal{U}^{\text{erg}}$. If \mathcal{S} is approached from \mathcal{U}_{out} , the potential \mathcal{R}_- goes to $-\infty$; the same is true for $\partial\mathcal{R}_-/\partial r$ unless $\partial g_{tt}/\partial r$ goes to zero. Correspondingly, if \mathcal{S} is approached from \mathcal{U}_{erg} , the potential \mathcal{R}_- goes to $+\infty$; the same is true for $\partial\mathcal{R}_-/\partial r$ unless $\partial g_{tt}/\partial r$ goes to zero. The potential \mathcal{R}_+ and its derivative are finite and continuous if \mathcal{S} is crossed.

The statements of the previous paragraph are also true in the case that $g_{t\varphi} > 0$ on \mathcal{S} , just with the potentials \mathcal{R}_+ and \mathcal{R}_- interchanged.

A further decomposition of \mathcal{U} is necessary if $g_{t\varphi}$ takes positive and negative values on \mathcal{S} , which is possible only if \mathcal{S} is disconnected. We will not work out such cases here.

5.2. Coordinate transformations

We are free to make coordinate transformations of the form

$$t \mapsto \tilde{t} = t + h(x^2, x^3), \quad \varphi \mapsto \tilde{\varphi} = \pm\varphi, \quad x^i \mapsto \tilde{x}^i = f^i(x^2, x^3) \quad (32)$$

where h , f^2 and f^3 are functions of (x^2, x^3) that are arbitrary except for the condition that they define an allowed coordinate transformation, i.e., that the determinant of the (2×2) -matrix $(\partial f^i/\partial x^j)$ is non-zero.

Under such a transformaton,

$$d\tilde{t} = dt + \frac{\partial h(x^2, x^3)}{\partial x^j} dx^j, \quad d\tilde{\varphi} = \pm d\varphi, \quad d\tilde{x}^i = \frac{\partial f^i(x^2, x^3)}{\partial x^j} dx^j, \quad (33)$$

hence

$$\frac{\partial}{\partial \tilde{t}} = \frac{\partial}{\partial t}, \quad \frac{\partial}{\partial \tilde{\varphi}} = \pm \frac{\partial}{\partial \varphi}, \quad \frac{\partial f^j(x^2, x^3)}{\partial x^i} \frac{\partial}{\partial \tilde{x}^j} = \frac{\partial}{\partial x^i} - \frac{\partial h(x^2, x^3)}{\partial x^i} \frac{\partial}{\partial t}. \quad (34)$$

This implies that

$$g_{tt} \mapsto \tilde{g}_{tt} = g_{tt}, \quad g_{t\varphi} \mapsto \tilde{g}_{t\varphi} = \pm g_{t\varphi}, \quad g_{\varphi\varphi} \mapsto \tilde{g}_{\varphi\varphi} = g_{\varphi\varphi} \quad (35)$$

and, thus,

$$g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi} \mapsto \tilde{g}_{t\varphi}^2 - \tilde{g}_{tt}\tilde{g}_{\varphi\varphi} = g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}. \quad (36)$$

As also

$$p_t \mapsto \tilde{p}_t = p_t, \quad (37)$$

the potentials \mathcal{R}_+ and \mathcal{R}_- remain unchanged. Of course, after the transformation we have to express them in terms of the new coordinates $(f^2(x^2, x^3), f^3(x^2, x^3))$.

The transformations (32) allow, e.g., in the Schwarzschild spacetime to switch from standard Schwarzschild coordinates to Eddington-Finkelstein or Painlevé-Gullstrand coordinates.

Note that, apart from a trivial rescaling of the time coordinate, $t \mapsto kt$ with a constant k , the transformations (32) are indeed the most general ones that preserve all our assumptions. A transformation of the form $t \mapsto t + \Omega_0^{-1}\varphi$ with a non-zero constant Ω_0 would leave each of the two potentials \mathcal{R}_+ and \mathcal{R}_- invariant up to an additive constant, but it would violate the periodicity condition (11); the φ -lines would no longer be closed.

5.3. Sign conventions

There are several conventions that go into our definition of \mathcal{R}_\pm . Firstly, it is not necessary, though convenient, to require that the force points into the direction of the *negative* gradient of the potential. This means that one could put a minus sign in front of one potential, or both.

Secondly, it is arbitrary which potential we label with a plus sign and which with a minus sign. We have chosen the signs such that $\dot{\varphi}$ is positive for the upper sign and negative for the lower sign, see (14). This, however, does not in general determine the sign of the angular velocity

$$\Omega = \frac{\dot{\varphi}}{\dot{t}} \tag{38}$$

because \dot{t} , which is given according to (15) by

$$\dot{t} = -\frac{\omega_0}{g_{tt}} \mp \frac{g_{t\varphi} \sqrt{\omega_0^2 + \omega_p^2 g_{tt}}}{g_{tt} \sqrt{g_{t\varphi}^2 - g_{tt} g_{\varphi\varphi}}}, \tag{39}$$

may have either sign. This is related to the question of whether the chosen parametrisation is future-oriented or past-oriented with respect to a time orientation.

If $g_{\varphi\varphi} > 0$ (i.e., if there is no causality violation), the hypersurfaces $t = \text{constant}$ are spacelike, i.e., t is a time function which defines a time orientation. Then, if $g_{tt} < 0$ (i.e., outside of an ergoregion), the first term in (39) dominates the second one, so our convention $\omega_0 \geq 0$ implies that \dot{t} is positive for both signs, so all light rays are future-oriented and Ω has the same sign as $\dot{\varphi}$. (Recall that, actually, the case $\omega_0 = 0$ is forbidden if $g_{tt} < 0$.) If, on the other hand, $g_{tt} > 0$ (i.e., inside an ergoregion), (25) requires $g_{t\varphi} \neq 0$, so the second term in (39) dominates the first one. Our convention $\omega_0 \geq 0$ implies that $\dot{t} > 0$ if $g_{t\varphi} < 0$ and $\dot{t} < 0$ if $g_{t\varphi} > 0$. (Here the case $\omega_0 = 0$ is possible provided that $\omega_p \neq 0$.) So we have to keep in mind that inside an ergoregion our light rays are parametrised in a future-oriented way for one sign and in a past-oriented way for the other.

If $g_{\varphi\varphi} < 0$ (i.e. in a causality-violating region), it depends on ω_p which term in (39) dominates, so no general statements are possible about the sign of the angular velocity Ω .

Thirdly, we could use, instead of \mathcal{R}_{\pm} , a function of \mathcal{R}_{\pm} that has the same critical points, e.g. the negative inverse whose gradient is given by

$$\frac{\partial}{\partial x^i} \left(\frac{-1}{\mathcal{R}_{\pm}} \right) = \frac{1}{\mathcal{R}_{\pm}^2} \frac{\partial \mathcal{R}_{\pm}}{\partial x^i}. \quad (40)$$

The relation between the potentials and their inverses can be read from the identity

$$g_{tt} \mathcal{R}_+ \mathcal{R}_- = (g_{t\varphi}^2 - g_{tt} g_{\varphi\varphi}) \omega_p^2 - g_{\varphi\varphi} \omega_0^2. \quad (41)$$

Note, however, that in our equation (23) the prefactor of the gradient on the right-hand side is strictly negative and finite at all points where (25) and (27) hold, i.e., at all points where circular light rays may exist. This is no longer true if we replace \mathcal{R}_{\pm} by $-1/\mathcal{R}_{\pm}$: The new prefactor is still non-negative, but it is infinite at points where \mathcal{R}_{\pm} has a zero. This unwanted feature is avoided if we use \mathcal{R}_{\pm} rather than its negative inverse. We also mention that in the vacuum case, but not in a plasma, the inverse potentials give directly the angular velocity (38),

$$\Omega = \frac{g_{tt} \sqrt{\omega_0^2 + \frac{\omega_p^2 g_{tt}}{\omega_0^2 c^2}}}{\mp \omega_0 \sqrt{g_{t\varphi}^2 - g_{\varphi\varphi} g_{tt}} - g_{t\varphi} \sqrt{\omega_0^2 + \omega_p^2 g_{tt}}} \xrightarrow{\omega_p \rightarrow 0} \frac{\omega_0}{\mp \mathcal{R}_{\pm}} \quad (42)$$

In the case $\omega_p = 0$, the inverse potentials $1/\mathcal{R}_{\pm}$ have been used for studying gravitational lensing in the Kerr-Newman spacetime by Hasse and Perlick [19]. Using a different sign convention, Cunha et al [13, 14, 15] have utilised these potentials in their work on the existence of circular vacuum light rays in axially symmetric and stationary spacetimes. We will generalise their results to the plasma case in the next section. The above arguments demonstrate that, in particular in the presence of a plasma, it is more convenient and more natural to choose \mathcal{R}_{\pm} rather than their inverses.

5.4. Properties of the potentials \mathcal{R}_{\pm} near a horizon

For analysing the properties of the potentials when a horizon is approached, we choose spherical polar coordinates $(t, \varphi, r, \vartheta)$ such that the horizon is represented as the hypersurface $r = r_h$. Then the metric coefficients g_{AB} and g^{ij} and their derivatives are finite on the horizon. As we are free to change to horizon-penetrating coordinates, by a transformation (32), we may even assume that *all* metric coefficients $g_{\mu\nu}$ and $g^{\mu\nu}$ and their derivatives are finite on the horizon.

A horizon is a lightlike hypersurface with finite circumference, so we must have

$$g_{t\varphi}^2 - g_{tt} g_{\varphi\varphi} = 0, \quad g_{\varphi\varphi} > 0 \quad (43)$$

at all points on the horizon with $0 < \vartheta < \pi$. We denote the domain of outer communication of the black hole by \mathcal{U} . By definition, this is the open set adjacent to the horizon where (25) holds. We choose the coordinate r such that $r > r_h$ on \mathcal{U} .

In order to derive the limit behaviour of the potentials and their derivatives at the horizon, we have to make sure that both potentials are defined on a subset of \mathcal{U} which is adjacent to the horizon. This is achieved by requiring that $-\omega_p^2 g_{tt}$ is bounded on a neighbourhood of the horizon and that we have chosen a frequency constant ω_0 such that $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$. In addition we also have to assume that the condition

$$\sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}} \frac{\partial\omega_p^2}{\partial r} \rightarrow 0 \quad (44)$$

holds for $r \rightarrow r_h$. This is certainly true if the gradient of ω_p^2 is bounded near the horizon.

For deriving the limit of \mathcal{R}_{\pm} for $r \rightarrow r_h$ we first observe that

$$\frac{1}{\mathcal{R}_{\pm}} = \frac{\mathcal{R}_{\mp}}{\mathcal{R}_{\pm}\mathcal{R}_{\mp}} = \frac{\mp g_{t\varphi}\omega_0 + \sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}}\sqrt{\omega_0^2 + \omega_p^2 g_{tt}}}{-\omega_0^2 g_{\varphi\varphi} + \omega_p^2 (g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi})}. \quad (45)$$

If we exclude the case $\omega_0 = 0$, which will be considered at the end of this section, we find with the help of our assumption that $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$ that

$$\frac{1}{\mathcal{R}_{\pm}} \mp \frac{g_{t\varphi}}{g_{\varphi\varphi}\omega_0} \rightarrow -0 \quad (46)$$

for $r \rightarrow r_h$. Here we write -0 to indicate that the limit is approached from below.

To evaluate this result further, we first consider the case of a non-rotating horizon, i.e., we assume that $g_{t\varphi} = 0$ and thus $g_{tt} = 0$ on the horizon. From (46) we read that then

$$\mathcal{R}_{\pm} \rightarrow -\infty \quad (47)$$

for $r \rightarrow r_h$. This case covers all spherically symmetric and static black holes and also, e.g., the NUT metric. In the latter case $g_{t\varphi}$ is non-zero in general, but it is zero on the horizon.

The situation is different if the horizon rotates, i.e., if $g_{t\varphi} \neq 0$ and hence $g_{tt} > 0$ on the horizon. From (46) we read that then

$$\mathcal{R}_{\pm} \rightarrow \pm \left. \frac{g_{t\varphi}}{g_{\varphi\varphi}\omega_0} \right|_{r=r_h}, \quad (48)$$

i.e., the potentials approach finite and non-zero values that are equal in magnitude and opposite in sign. This case covers all metrics that describe stationarily rotating black holes such as the Kerr metric.

Knowing the limit behaviour of the potentials, we now derive the limit behaviour of their derivatives. We will see that in this case it makes a difference whether the horizon is degenerate or non-degenerate. We start out from the equations

$$\frac{1}{\mathcal{R}_-} + \frac{1}{\mathcal{R}_+} = \frac{\mathcal{R}_+ + \mathcal{R}_-}{\mathcal{R}_+\mathcal{R}_-} = \frac{2\sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}}\sqrt{\omega_0^2 + \omega_p^2 g_{tt}}}{-g_{\varphi\varphi}\omega_0^2 + \omega_p^2 (g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi})} \quad (49)$$

and

$$\frac{1}{\mathcal{R}_-} - \frac{1}{\mathcal{R}_+} = \frac{\mathcal{R}_+ - \mathcal{R}_-}{\mathcal{R}_+\mathcal{R}_-} = \frac{\pm 2 g_{t\varphi}\omega_0}{-g_{\varphi\varphi}\omega_0^2 + \omega_p^2 (g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi})} \quad (50)$$

which follow directly from (21). Differentiating these equations, and excluding again the case $\omega_0 = 0$, shows that

$$\frac{1}{\mathcal{R}_+^2} \frac{\partial \mathcal{R}_+}{\partial r} + \frac{1}{\mathcal{R}_-^2} \frac{\partial \mathcal{R}_-}{\partial r} + \frac{2 \sqrt{\omega_0^2 + \omega_p^2 g_{tt}}}{g_{\varphi\varphi} \omega_0^2} \frac{\partial \sqrt{g_{t\varphi}^2 - g_{tt} g_{\varphi\varphi}}}{\partial r} \rightarrow 0 \quad (51)$$

and

$$\frac{1}{\mathcal{R}_+^2} \frac{\partial \mathcal{R}_+}{\partial r} - \frac{1}{\mathcal{R}_-^2} \frac{\partial \mathcal{R}_-}{\partial r} \quad \text{stays finite} \quad (52)$$

if the horizon is approached. Here we have used our assumptions that $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$ and that (44) holds.

If the horizon is non-degenerate, $\partial \sqrt{g_{t\varphi}^2 - g_{tt} g_{\varphi\varphi}} / \partial r$ goes to $+\infty$ for $r \rightarrow r_h$. (Recall that we approach the horizon from the side where $r > r_h$.) As \mathcal{R}_\pm does not go to 0, (51) and (52) imply that

$$\frac{\partial \mathcal{R}_\pm}{\partial r} \rightarrow -\infty \quad (53)$$

for $r \rightarrow r_h$.

If the horizon is degenerate, $\partial \sqrt{g_{t\varphi}^2 - g_{tt} g_{\varphi\varphi}} / \partial r$ approaches a finite and positive value for $r \rightarrow r_h$. In the non-rotating case, where $\mathcal{R}_\pm^2 \rightarrow \infty$, (51) and (52) imply that again (53) holds. In the rotating case, however, where (48) holds, (51) and (52) imply that the derivatives of the potentials approach finite values.

Finally, we consider the case $\omega_0 = 0$ which was left out so far. In this case the potentials, which are defined near the horizon only if $g_{tt} > 0$ and $\omega_p^2 > 0$, reduce to

$$\mathcal{R}_+ = \mathcal{R}_- = \frac{\omega_p}{\sqrt{g_{tt}}} \sqrt{g_{t\varphi}^2 - g_{tt} g_{\varphi\varphi}}. \quad (54)$$

From this expression and from its r -derivative we read that no general statements about the limit behaviour of the potentials and their derivatives are possible: Apart from the obvious fact that the limit of $\mathcal{R}_+ = \mathcal{R}_-$ cannot be negative, everything is possible, depending on the limit behaviour of g_{tt} , ω_p and $\partial \omega_p / \partial r$.

So we see that the case that $\omega_0 = 0$ and the case that the horizon is rotating and degenerate are special. In all other cases $\partial \mathcal{R}_\pm / \partial r$ goes to $-\infty$ if a horizon is approached. This implies that there is a positive $r_0(\vartheta)$ such that neither the potential \mathcal{R}_\pm itself nor any small perturbation thereof has a critical point in the domain

$$\mathcal{U}_{r_0} = \{ (r \sin \vartheta, r \cos \vartheta) \mid r_h < r < r_h + \delta(\vartheta), 0 < \vartheta < \pi \}, \quad (55)$$

or on its boundary. Here the case that $\delta(\vartheta) \rightarrow 0$ for $\sin \vartheta \rightarrow 0$ cannot be excluded. We keep in mind that this is true not only for vacuum light rays but also in a plasma, provided that $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$ and that (44) holds.

5.5. Properties of the potentials \mathcal{R}_\pm on an axis

In order to analyse the properties of the potentials near a regular axis of symmetry, we use cylindrical polar coordinates (t, φ, ρ, z) such that the axis is represented as the boundary $\rho = 0$ of the half-plane $\rho > 0$. The axis is called regular if the metric satisfies the condition of “elementary flatness” there. This means that the coordinates can be chosen such that on the interval $z_1 < z < z_2$ where the axis is regular there is a strictly positive function $f(z)$ such that

$$g_{tt} \rightarrow -f(z)^2, \quad \frac{g_{t\varphi}}{\rho} \rightarrow 0, \quad \frac{g_{\varphi\varphi}}{\rho^2} \rightarrow 1, \quad (56)$$

$$\frac{\partial g_{tt}}{\partial \rho} \rightarrow 0, \quad \frac{\partial g_{t\varphi}}{\partial \rho} \rightarrow 0, \quad \frac{1}{\rho} \frac{\partial g_{\varphi\varphi}}{\partial \rho} \rightarrow 2, \quad (57)$$

for $\rho \rightarrow 0$. This implies that

$$\frac{1}{\rho} \sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}} \rightarrow f(z), \quad \frac{\partial \sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}}}{\partial \rho} \rightarrow f(z), \quad (58)$$

for $\rho \rightarrow 0$.

We have to make sure that the domain of definition of \mathcal{R}_\pm extends to the axis. This requires that we have to assume that $-\omega_p^2 g_{tt}$ is bounded near the axis, and we have to choose a frequency constant ω_0 such that $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$. Moreover, we have to require that

$$\rho \frac{\partial \omega_p^2}{\partial \rho} \rightarrow 0 \quad (59)$$

for $\rho \rightarrow 0$. This condition, which is analogous to (44), is certainly true if the gradient of ω_p^2 is bounded near the axis.

Then the axis is on the boundary of the domain where the potential \mathcal{R}_\pm is defined, for both signs, and we find

$$\mathcal{R}_\pm \rightarrow 0, \quad \frac{\partial \mathcal{R}_\pm}{\partial \rho} + \frac{1}{f(z)} \sqrt{\omega_0^2 + \omega_p^2 g_{tt}} \rightarrow 0 \quad (60)$$

for $\rho \rightarrow 0$. This is true for all values of z in the interval $z_1 < z < z_2$ where the axis is regular.

This observation has important consequences for spacetimes that are asymptotically flat, see Section 5.6 below. If this is true, and if the axis is regular on an interval $z_1 < z < \infty$, we have $f(z) \rightarrow 1$ for $z \rightarrow \infty$. As the square-root in (60) is strictly positive, the gradient of \mathcal{R}_\pm is bounded away from zero near the axis, i.e., there is a constant ρ_0 such that neither the potential \mathcal{R}_\pm itself nor any small perturbation thereof has a critical point in the domain $0 < \rho < 2\rho_0$, $z_1 < z < \infty$. Similarly, if the axis is regular on the interval $-\infty < z < z_2$, asymptotic flatness implies that $f(z) \rightarrow 1$ for $z \rightarrow -\infty$, so there is a constant ρ_0 such that neither the potential itself nor any small perturbation thereof can have a critical point on the domain $0 < \rho < 2\rho_0$, $-\infty < z < z_2$. We will later apply this result to asymptotically flat spacetimes where the entire axis is regular, i.e. $z_1 = -\infty$ and $z_2 = \infty$, and to asymptotically flat black-hole spacetimes

with horizon at $\sqrt{\rho^2 + z^2} = r_h$ where the axis is regular between the horizon and infinity, i.e. the result is true for any $z_1 \geq r_h$ and any $z_2 \leq -r_h$. Note that the limiting values $z_1 = r_h$ and $z_2 = -r_h$ are indeed included because $f(z)$ does not go to ∞ for $z \rightarrow \pm r_h$: It goes to 0 for a non-rotating horizon and to a non-zero finite value for a rotating horizon.

5.6. Properties of the potentials \mathcal{R}_\pm at infinity

We now turn to an investigation of the behaviour of the potentials at infinity. We assume that the spacetime is asymptotically flat in the sense that we may choose spherical polar coordinates $(t, \varphi, r, \vartheta)$ such that

$$g_{tt} \rightarrow -1, \quad \frac{g_{t\varphi}}{r \sin \vartheta} \rightarrow 0, \quad \frac{g_{\varphi\varphi}}{r^2 \sin^2 \vartheta} \rightarrow 1, \quad (61)$$

$$r \frac{\partial g_{tt}}{\partial r} \rightarrow 0, \quad \frac{\partial g_{t\varphi}}{\partial r} \rightarrow 0, \quad \frac{1}{r \sin^2 \vartheta} \frac{\partial g_{\varphi\varphi}}{\partial r} \rightarrow 2 \quad (62)$$

for $r \rightarrow \infty$ and $0 < \vartheta < \pi$. This implies that

$$\frac{\sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}}}{r \sin \vartheta} \rightarrow 1, \quad \frac{\partial \sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}}}{\partial r} \rightarrow \sin \vartheta. \quad (63)$$

We have to assume that $-\omega_p^2 g_{tt}$ is bounded at infinity, and we have to choose a frequency constant ω_0 with $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$. Moreover, we have to require that

$$\sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}} \frac{\partial \omega_p^2}{\partial r} \rightarrow 0 \quad (64)$$

for $r \rightarrow \infty$ which is certainly true if the gradient of ω_p^2 falls off stronger than $1/r$.

Then it is straight-forward to verify that

$$\mathcal{R}_\pm(r, \vartheta) \rightarrow -\infty, \quad \frac{\partial \mathcal{R}_\pm(r, \vartheta)}{\partial r} + \sin \vartheta \sqrt{\omega_0^2 + \omega_p(r, \vartheta)^2 g_{tt}(r, \vartheta)} \rightarrow 0 \quad (65)$$

for $r \rightarrow \infty$. As the square-root is bounded away from zero this guarantees that there is an $r_0(\vartheta)$ such that neither the potential \mathcal{R}_\pm itself nor any small perturbation thereof has a critical point in the domain $r_0(\vartheta)/2 < r < \infty$, $0 < \vartheta < \pi$. We cannot exclude the possibility that $r_0(\vartheta) \rightarrow \infty$ for $\sin \vartheta \rightarrow 0$.

6. Existence theorems for circular light rays

Cunha et al. [13, 14, 15] have proven existence theorems for circular light rays in vacuum (i.e. for circular lightlike geodesics) in axially symmetric and stationary spacetimes. To that end they have used two potentials H_\pm that are related to our potentials \mathcal{R}_\pm by the equation $H_\pm = \mp 1/\mathcal{R}_\pm$ if the latter are restricted to the case that $\omega_p = 0$. The reasons why we use \mathcal{R}_\pm , rather than H_\pm , have been outlined in Section 5.3. The fact that \mathcal{R}_\pm is well defined also for $\omega_p \neq 0$ allows us to generalize the results of Cunha et al. to the plasma case.

Following Cunha et al. closely, we utilise the Brouwer degree, which is a standard concept in differential topology, and the fact that it is a homotopy invariant. The Brouwer degree is defined for a smooth (C^2 would do) function f from a compact manifold \mathcal{K} to a manifold \mathcal{N} of the same dimension. In our case f is the gradient of the potential \mathcal{R}_\pm , \mathcal{K} is the closure of an open and bounded subset of the connected domain \mathcal{V}_\pm where the potential \mathcal{R}_\pm is defined, and \mathcal{N} is \mathbb{R}^2 . One has to assume that 0 is a regular value of f . In our case, this means that \mathcal{R}_\pm is a Morse function, i.e. that at all critical points of the potential its Hessian is non-degenerate. As we have the positive definite metric g^{ij} at our disposal, this assumption is tantamount to the condition that the (2×2) -matrix $(g^{ik} \partial^2 \mathcal{R}_\pm / \partial x^k \partial x^j)$ has two non-zero eigenvalues, so its determinant is non-zero. Following the terminology of Cunha et al., we assign to each critical point a “topological charge” which, by definition, is the sign of this determinant. In other words, the topological charge is +1 if the critical point is an extremum (i.e., a local minimum or a local maximum), while it is -1 if the critical point is a saddle. The Brouwer degree of the map f is defined as the sum of the topological charges over all critical points. If the potential has no critical points, it is a Morse function and one assigns the Brouwer degree 0 to its gradient. Now the homotopy invariance of the degree says that two maps $f^0 : \mathcal{K} \rightarrow \mathcal{N}$ and $f^1 : \mathcal{K} \rightarrow \mathcal{N}$ have the same degree if they can be deformed into each other by a one-parameter family of maps $f^\varepsilon : \mathcal{K} \rightarrow \mathcal{N}$, continuously depending on $\varepsilon \in [0, 1]$, such that f^ε does not take the value 0 on the boundary of \mathcal{K} , for all $\varepsilon \in [0, 1]$. While every textbook on differential topology discusses the homotopy invariance of the Brouwer degree, some of them restrict to the case that \mathcal{K} is closed, i.e., compact without boundary. A complete proof for the case that \mathcal{K} is compact with boundary, which is the case of relevance to us, can be found e.g. in the book by Dinca and Mawhin [20].

If applied to our potentials \mathcal{R}_+ and \mathcal{R}_- , the homotopy invariance of the Brouwer degree gives us the following result.

Proposition 2. Consider a one-parameter family of axially symmetric and stationary spacetimes with metrics $g_{\mu\nu}^\varepsilon$ and axially symmetric and stationary plasma frequencies ω_p^ε , depending continuously on a parameter $\varepsilon \in [0, 1]$. Let $\mathcal{R}_\pm^\varepsilon$ be the corresponding potentials, defined on the open and connected subsets $\mathcal{V}_\pm^\varepsilon$. Assume that there is a family of compact subsets $\mathcal{K}_\pm^\varepsilon$ and a family of diffeomorphisms $\mathcal{K}_\pm^\varepsilon \rightarrow \mathcal{K}_\pm^0$, depending continuously on $\varepsilon \in [0, 1]$, such that all critical points of $\mathcal{R}_\pm^\varepsilon$ are contained in the interior of $\mathcal{K}_\pm^\varepsilon$. If \mathcal{R}_\pm^0 and \mathcal{R}_\pm^1 are Morse functions, then the number of critical points of \mathcal{R}_\pm^1 differs from the number of critical points of \mathcal{R}_\pm^0 by an even number $2n$, where n of the additional critical points are saddles and the other n are extrema.

When applying this proposition one has to make sure that $\mathcal{R}_\pm^\varepsilon$ satisfies all the assumptions. The proposition is not applicable if \mathcal{R}_\pm^0 or \mathcal{R}_\pm^1 fails to be a Morse function. Then its critical points need not be isolated, i.e., there may be a continuum of critical points which means that the number of critical points is infinite and not even countable. Morse functions are generic in the sense that every non-Morse function becomes a Morse

function under an appropriate small perturbation. Also, the proposition is not applicable if both functions \mathcal{R}_\pm^0 and \mathcal{R}_\pm^1 are Morse functions but if one of them has infinitely many critical points. As critical points are isolated if they are non-degenerate, it is not possible that infinitely many of them are contained in a compact set. Finally, it is possible that \mathcal{R}_\pm^1 and \mathcal{R}_\pm^0 are Morse functions all of whose critical points lie in the interior of a compact set \mathcal{K}_\pm^1 , but that it is impossible to find a deformation $\mathcal{R}_\pm^\varepsilon$ and compact sets $\mathcal{K}_\pm^\varepsilon$, both depending continuously on ε , such that the critical points of $\mathcal{R}_\pm^\varepsilon$ are in the interior of $\mathcal{K}_\pm^\varepsilon$ for all $\varepsilon \in [0, 1]$; this situation happens if, intuitively speaking, \mathcal{R}_\pm^1 has a critical point at infinity. Examples for all three cases will be given in Section 7.1 below.

In their first paper Cunha et al. [13] considered the case that $\omega_p = 0$ throughout, and they applied what we have formulated as Proposition 2 to the case that $g_{\mu\nu}^0$ is the metric of a spacetime without circular light rays and $g_{\mu\nu}^1$ is the metric of a spacetime with circular light rays. The interpretation is that $g_{\mu\nu}^0$ describes the spacetime of a regular star before it undergoes gravitational collapse, while $g_{\mu\nu}^1$ describes the corresponding spacetime after the star has collapsed to an ultracompact object that is not a black hole. The crucial observation is that, in our notation, circular light rays of the metric $g_{\mu\nu}^1$ come in pairs, always a saddle together with an extremum. In their follow-up papers, Cunha et al. [14, 15] investigated the Brouwer degree of the gradients of their potentials for black-hole spacetimes.

Our generalisation to the plasma case allows for applications where both the spacetime metric and the plasma density are deformed. We will concentrate, however, in the following on one-parameter families of the form

$$\mathcal{R}_\pm^\varepsilon = \frac{g_{t\varphi}}{g_{tt}}\omega_0 + \frac{1}{g_{tt}}\sqrt{g_{t\varphi}^2 - g_{tt}g_{\varphi\varphi}}\sqrt{\omega_0^2 + \varepsilon\omega_p^2g_{tt}} \quad (66)$$

where the spacetime metric is kept fixed. $\mathcal{R}_\pm^\varepsilon$ is defined on the domain \mathcal{V}_\pm where \mathcal{R}_\pm^1 is defined, for all $\varepsilon \in [0, 1]$, if $-\omega_p^2g_{tt}$ is bounded on \mathcal{V}_\pm and $\omega_0^2 > \sup(-\omega_p^2g_{tt})$. In all applications we will assume that this is the case. For other values of ω_0 one could apply Proposition 2 as well, but one would have to consider domains $\mathcal{V}_\pm^\varepsilon$ that depend on ε and possibly also frequency constants ω_0 that depend on ε .

Moreover, we have to make sure that \mathcal{R}_\pm^1 and \mathcal{R}_\pm^0 are Morse functions and that all critical points of $\mathcal{R}_\pm^\varepsilon$ are contained in the interior of a compact set \mathcal{K} . (If all $\mathcal{R}_\pm^\varepsilon$ are defined on the same domain \mathcal{V}_\pm , there is no need to consider ε -dependent compact sets.) Then Proposition 2 guarantees that the circular light rays in the plasma density ω_p^2 differ from the circular vacuum light rays on the same spacetime by n saddles and n extrema. This implies, in particular, that there is at least one circular light ray in the plasma if there is an odd number of circular light rays in vacuum. In other words, by introducing a plasma we cannot destroy all circular light rays if their number is odd.

Note that Proposition 2 has to be applied to \mathcal{R}_+ and \mathcal{R}_- separately and that these two potentials may have different domains of definition.

In the following sections we will consider several specific axially symmetric and stationary spacetimes. We will discuss the general results for circular light rays that

can be concluded from Proposition 2 and we will illustrate this with particular plasma densities.

7. Regular spacetimes

In this section we exemplify Proposition 2 with the case that the axially symmetric and stationary spacetime and the axially symmetric and stationary plasma density under consideration are regular. By that we mean that the metric and the plasma density are defined on the entire half-plane

$$\mathcal{U} = \{(\rho, z) \mid \rho > 0, -\infty < z < \infty\} = \{(r \sin \vartheta, r \cos \vartheta,) \mid r > 0, 0 < \vartheta < \pi\}, \quad (67)$$

that (25) holds on \mathcal{U} , that $\rho = 0$ is a regular axis (recall Section 5.5), that the spacetime is asymptotically flat (recall Section 5.6), and that the plasma density is bounded on the half-plane \mathcal{U} . These assumptions allow us to choose a frequency constant ω_0 such that $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$. If we connect with the vacuum case on the same spacetime by defining the one-parameter family of potentials (66), we find that $\mathcal{R}_\pm^\varepsilon$ is defined on the entire halfplane \mathcal{U} for all $\varepsilon \in [0, 1]$.

If in addition the conditions (59) and (64) hold for $r \sin \vartheta \rightarrow 0$ and $r \rightarrow \infty$, respectively, the results of Sections 5.5 and 5.6 guarantee that there is a compact subset \mathcal{K} of \mathcal{U} such that all critical points of $\mathcal{R}_\pm^\varepsilon$ lie in the interior of \mathcal{K} , for all $\varepsilon \in [0, 1]$. In the notation of Sections 5.5 and 5.6, this compact set is of the form

$$\mathcal{K} = \{(r \sin \vartheta, r \cos \vartheta,) \mid r \sin \vartheta \geq \rho_0, r \leq r_0(\vartheta)\}. \quad (68)$$

If \mathcal{R}_\pm^0 and \mathcal{R}_\pm^1 are Morse functions, which is generically true, Proposition 2 guarantees that the circular light rays in the plasma density ω_p^2 differ from the circular light rays in vacuum by n saddles and n extrema. This result can be applied, e.g., to the spacetime of a star that is not ultracompact or to Minkowski spacetime. In both cases \mathcal{R}_\pm^0 has no critical points, so in the plasma the number of circular light rays must be even.

In the following we illustrate this situation with plasma densities on Minkowski spacetime. We choose three examples where the assumptions of Proposition 2 are *not* satisfied, just to demonstrate in which situations this is the case.

7.1. Minkowski spacetime

In this section we give three examples for plasma densities on Minkowski spacetime. Note that then the light rays in the plasma are timelike geodesics of the conformally rescaled metric $\omega_p^2 g_{\mu\nu}$, i.e., they mimic the motion of freely falling massive particles in this rescaled metric. In this sense, light propagation in a plasma on Minkowski spacetime may be viewed as an example of *analogue gravity*.

In each of the three following examples we choose a frequency constant such that the potentials (66) are defined on the entire half-plane (67), but that the assumptions of Proposition 2 are violated in three different ways: In the first example $\mathcal{R}_\pm = \mathcal{R}_\pm^1$ fails to be a Morse function, in the second it is a Morse function but it has infinitely many

critical points (which is possible because in this example the gradient of the potential is unbounded) and in the third we choose ω_0^2 equal to the supremum of $-\omega_p^2 g_{tt}$, rather than strictly bigger.

In the first example we consider Minkowski spacetime in cylindrical polar coordinates (t, φ, ρ, z) ,

$$g_{tt} = -1, \quad g_{\varphi\varphi} = \rho^2, \quad g_{\rho\rho} = 1 \quad g_{zz} = 1. \quad (69)$$

All other metric coefficients are zero. Then the two potentials coincide,

$$\mathcal{R}_+(\rho, z) = \mathcal{R}_-(\rho, z) = -\rho \sqrt{\omega_0^2 - \omega_p(\rho, z)^2}. \quad (70)$$

We choose the plasma density

$$\omega_p(\rho, z)^2 = \omega_c^2 \sin^2 \frac{\rho}{\rho_0} \quad (71)$$

where ω_c is a constant with the dimension of a frequency and ρ_0 is a constant with the dimension of a length. Then $-\omega_p^2 g_{tt}$ is bounded on the half-plane \mathcal{U} , with $\sup(-\omega_p^2 g_{tt}) = \omega_c^2$. We choose a frequency constant $\omega_0 > \omega_c$ which guarantees that

$$\mathcal{R}_+(\rho, z) = \mathcal{R}_-(\rho, z) = -\rho \sqrt{\omega_0^2 - \omega_c^2 \sin^2 \frac{\rho}{\rho_0}}, \quad (72)$$

is defined on the entire half-plane \mathcal{U} . However, there is a continuum of critical points, located on infinitely many vertical lines in \mathcal{U} which are given by the equation

$$\omega_c^2 \sin^2 \frac{\rho}{\rho_0} \left(\sin \frac{\rho}{\rho_0} + \frac{\rho}{\rho_0} \cos \frac{\rho}{\rho_0} \right) = \omega_0^2, \quad (73)$$

so the potential fails to be a Morse function and Proposition 2 is not applicable.

For the second (equally contrived) example we consider again Minkowski spacetime, this time in spherical polar coordinates $(t, \varphi, r, \vartheta)$, with the non-zero metric coefficients

$$g_{tt} = -1, \quad g_{\varphi\varphi} = r^2 \sin^2 \vartheta, \quad g_{rr} = 1, \quad g_{\vartheta\vartheta} = r^2. \quad (74)$$

We choose the plasma density

$$\omega_p(r, \vartheta)^2 = \omega_c^2 \sin^2 \frac{r}{r_0}, \quad (75)$$

where ω_c is a constant with the dimension of a frequency and r_0 is a constant with the dimension of a length. Again, $-\omega_p^2 g_{tt}$ is bounded on the half-plane \mathcal{U} with $\sup(-\omega_p^2 g_{tt}) = \omega_c^2$. We choose an $\omega_0 > \omega_c$ which guarantees that

$$\mathcal{R}_+^\varepsilon(r, \vartheta) = \mathcal{R}_-^\varepsilon(r, \vartheta) = -r \sin \vartheta \sqrt{\omega_0^2 - \varepsilon \omega_c^2 \sin^2 \frac{r}{r_0}} \quad (76)$$

is defined on the entire half-plane $r \sin \vartheta > 0$, for all $\varepsilon \in [0, 1]$. In this case, \mathcal{R}_+^1 and \mathcal{R}_+^0 are, indeed, Morse functions. However, as the gradient of \mathcal{R}_+^1 is

$$\frac{\partial \mathcal{R}_+^1(r, \vartheta)}{\partial r} = - \frac{\sin \vartheta \left(\omega_0^2 - \omega_c^2 \sin^2 \frac{r}{r_0} \left(\sin \frac{r}{r_0} + \frac{r}{r_0} \cos \frac{r}{r_0} \right) \right)}{\sqrt{\omega_0^2 - \omega_c^2 \cos^2 \frac{r}{r_0}}}, \quad (77)$$




Figure 1. This picture shows the equipotential lines $\mathcal{R}_+ = \mathcal{R}_- = \text{constant}$ for the plasma density (75) on Minkowski spacetime for $\omega_0 = \sqrt{4/3}\omega_c$ in the $(r \sin \vartheta, r \cos \vartheta)$ -halfplane \mathcal{U} . r_0 is chosen as the unit on the axes. The circular light rays are marked by black dots. There are infinitely many saddles with local maxima in between. The saddles are minima (stable) in r direction and maxima (unstable) in ϑ direction. The potential goes, in an oscillatory fashion, to $-\infty$ for $r \sin \vartheta \rightarrow \infty$.

$$\frac{\partial \mathcal{R}_{\pm}^1(r, \vartheta)}{\partial \vartheta} = -r \cos \vartheta \sqrt{\omega_0^2 - \omega_c^2 \cos^2 \frac{r}{r_0}}, \quad (78)$$

\mathcal{R}_{\pm}^1 has infinitely many isolated critical points, located at

$$\omega_0^2 = \omega_c^2 \sin \frac{r}{r_0} \left(\sin \frac{r}{r_0} + \frac{r}{r_0} \cos \frac{r}{r_0} \right), \quad \vartheta = \frac{\pi}{2}, \quad (79)$$

so Proposition 2 is again not applicable; we cannot find a compact subset of \mathcal{U} which contains all critical points of \mathcal{R}_{\pm}^1 . This can happen only because (64) is violated for $r \rightarrow \infty$. Fig. 1 shows the potential (76) for $\omega_0 = \sqrt{4/3}\omega_c$.

For the final example on Minkowski spacetime we work again in cylindrical polar coordinates (69). We choose the plasma density

$$\omega_p(\rho, z)^2 = \frac{\omega_c^2(\rho_0^2 + (\rho - \rho_0)^2)}{2r_0^2 + (\rho - \rho_0)^2 + z^2} \quad (80)$$

where again ω_c is a constant with the dimension of a frequency and ρ_0 is a constant with the dimension of a length. This time we choose $\omega_0 = \omega_c$ for which the potentials

$$\mathcal{R}_+^{\varepsilon}(\rho, z) = \mathcal{R}_-^{\varepsilon}(\rho, z) = \rho \sqrt{\omega_0^2 - \frac{\varepsilon \omega_c^2(r_0^2 + (\rho - r_0)^2)}{2r_0^2 + (\rho - r_0)^2 + z^2}} \quad (81)$$




Figure 2. This picture shows the equipotential lines $\mathcal{R}_+ = \mathcal{R}_- = \text{constant}$ for the plasma density (80) on Minkowski spacetime in the (ρ, z) -half-plane \mathcal{U} , with ρ_0 used as the units on the axes. We have chosen ω_0 equal to ω_c which implies that the strict inequality $\omega_0 > \omega_p(\rho, z)$ holds for all $0 < \rho < \infty$ but not in the limit $\rho \rightarrow \infty$. The picture shows that there is exactly one circular light ray, namely a saddle, marked by a black dot. In this case the assumptions of Proposition 2 are violated because it is impossible to find a one-parameter family of compact sets such that the critical points of $\mathcal{R}_\pm^\varepsilon$ are in their interior for all $\varepsilon \in [0, 1]$.

are indeed defined on the entire half-plane $\{(\rho, z) \mid \rho > 0, -\infty < z < \infty\}$. In this case \mathcal{R}_\pm^1 and \mathcal{R}_\pm^0 are Morse functions. \mathcal{R}_\pm^0 has no critical points whereas \mathcal{R}_\pm^1 has exactly one critical point, namely a saddle, see Fig. 2. Clearly, this can be reconciled with Proposition 2 only if one of the assumptions of this Proposition is violated. This is indeed the case: For $0.992 \lesssim \varepsilon < 1$ the potential $\mathcal{R}_\pm^\varepsilon$ has *two* critical points, a saddle and a local maximum. With $\varepsilon \rightarrow 1$ the maximum moves to infinity while the saddle approaches the saddle of the potential \mathcal{R}_\pm^1 which lies in the interior of the half-plane \mathcal{U} . Therefore it is impossible to include all critical points in a compact set, or in a family of compact sets that depend continuously on ε . This situation cannot occur if ω_0^2 is *strictly* bigger than the supremum of $-\omega_p^2 g_{tt}$.

8. Black-hole spacetimes

In this section we apply Proposition 2 to the domain of outer communication \mathcal{U} of a black hole. We assume that $g_{\varphi\varphi} > 0$, i.e. that there is no causality violation, on \mathcal{U}

which guarantees that t is a time function. Moreover, we restrict to the case that the spacetime is asymptotically flat and that the axis is regular outside the horizon, recall Sections 5.5 and 5.6. Note that these assumptions are *not* satisfied, e.g., in the NUT spacetime, which features a causality-violating region.

The domain of outer communication is the set

$$\begin{aligned} \mathcal{U} &= \{(\rho, z) \mid r_h^2 < \rho^2 + z^2, \rho > 0, -\infty < z < \infty\} \\ &= \{(r \sin \vartheta, r \cos \vartheta) \mid r_h < r < \infty, 0 < \vartheta < \pi\} \end{aligned} \quad (82)$$

where $r = r_h$ is the horizon.

We first consider black holes without an ergoregion, i.e., we assume that $g_{tt} < 0$ on all of \mathcal{U} . If we connect with the vacuum case by the one-parameter family of potentials (66), our assumptions guarantee that $\mathcal{R}_\pm^\varepsilon$ is defined on $\mathcal{V}_\pm = \mathcal{U}$, for all $\varepsilon \in [0, 1]$, provided that $-\omega_p^2 g_{tt}$ is bounded on \mathcal{U} and that we choose a frequency constant ω_0 with $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$ where the supremum is to be taken over \mathcal{U} . Moreover, if the plasma density satisfies conditions (44), (59) and (64) at the horizon, at the regular axis and at infinity, respectively, then the results of Sections 5.4, 5.5 and 5.6 guarantee that there is a compact subset \mathcal{K} of \mathcal{U} such that the critical points of $\mathcal{R}_\pm^\varepsilon$ lie in the interior of \mathcal{K} , for all $\varepsilon \in [0, 1]$. This compact set is of the form

$$\mathcal{K} = \{(r \sin \vartheta, r \cos \vartheta) \mid r \sin \vartheta \geq \rho_0, \quad r_h + \delta \leq r \leq r_0\} \quad (83)$$

with some positive ρ_0 , δ and r_0 . Then the one-parameter family (66) satisfies the assumptions of Proposition 2 for all $\varepsilon \in [0, 1]$, provided that $\mathcal{R}_\pm^1 = \mathcal{R}_\pm$ and \mathcal{R}_\pm^0 are Morse functions. In this case, the number of critical points of \mathcal{R}_\pm^1 differs from the number of critical points of \mathcal{R}_\pm^0 by an even number. As $\mathcal{R}_\pm^\varepsilon \rightarrow -\infty$ at the horizon and at infinity, each $\mathcal{R}_\pm^\varepsilon$ must have at least one critical point. Actually, for \mathcal{R}_\pm^0 (i.e., vacuum light rays) in spherically symmetric and static black-hole spacetimes the latter observation is rather trivial and well known, see e.g. Hasse and Perlick [21], Section 6.1, or Perlick [22], Section 4.3.

We now turn to the case that there is an ergoregion. Disregarding more complicated situations, we will assume in the following that \mathcal{U}^{out} , \mathcal{U}^{erg} and \mathcal{S} are connected, where we use the notation of Section 5.1. To fix the sign ambiguity of the potentials inside an ergoregion (recall Section 5.3), we assume that there $g_{t\varphi} < 0$. This is no restriction of generality, because we are free to transform $\varphi \mapsto -\varphi$ and $g_{t\varphi}$ cannot have zeros inside an ergoregion. If the ergoregion extends to the horizon, this sign convention means that prograde ($\dot{\varphi} > 0$) light rays are co-rotating with the horizon.

The potential \mathcal{R}_+ , which describes future-oriented ($\dot{t} > 0$) prograde ($\dot{\varphi} > 0$) light rays, is defined on $\mathcal{V}_+ = \mathcal{U}$. In contrast to the case without an ergoregion, it is not guaranteed that there is at least one critical point, because \mathcal{R}_+ does not go to $-\infty$ at the horizon. If the horizon is non-degenerate, our assumptions on the plasma density imply that the critical points of \mathcal{R}_+ lie inside a compact set. If \mathcal{R}_+^1 and \mathcal{R}_+^0 are Morse functions, Proposition 2 guarantees that the number of critical points of \mathcal{R}_+^1 differs from the number of critical points of \mathcal{R}_+^0 by an even number.

The potential \mathcal{R}_- has to be considered on \mathcal{U}^{out} and \mathcal{U}^{erg} separately. If $-\omega_p^2 g_{tt}$ is bounded on \mathcal{U}^{out} , we can choose a frequency constant ω_0 with $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$, where the supremum is to be taken over \mathcal{U}^{out} . Then \mathcal{R}_- is defined on $\mathcal{V}_- = \mathcal{U}^{\text{out}}$. On this domain \mathcal{R}_- describes future-oriented ($\dot{t} > 0$) retrograde ($\dot{\varphi} < 0$) light rays. $\mathcal{V}_- = \mathcal{U}^{\text{out}}$ extends to infinity and its boundary may contain points on the regular axis, on the set \mathcal{S} where $g_{tt} = 0$ and on the horizon. In order to make sure that all critical points of \mathcal{R}_- in \mathcal{U}_{out} lie inside a compact set, we need the additional assumptions that the horizon is non-degenerate (if \mathcal{U}_{out} extends to the horizon) and that $\partial g_{tt}/\partial r$ does not go to zero if a point on \mathcal{S} is approached from \mathcal{U}^{out} . If \mathcal{R}_-^1 and \mathcal{R}_-^0 are Morse functions, Proposition 2 guarantees that the number of critical points of \mathcal{R}_-^1 differ from the number of critical points of \mathcal{R}_-^0 by an even number. As $\mathcal{R}_-^\varepsilon$ goes to $-\infty$ for $r \rightarrow \infty$ and if a point on \mathcal{S} is approached, this potential must have at least one critical point, for all $\varepsilon \in [0, 1]$.

On the other hand, we can also consider \mathcal{R}_- on the domain $\mathcal{V}_- = \mathcal{U}^{\text{erg}}$. There the potential \mathcal{R}_- is defined for all $\omega_0 > 0$ and if the supremum of $-\omega_p^2 g_{tt}$ is strictly negative on \mathcal{U}^{erg} also for $\omega_0 = 0$. On this domain \mathcal{R}_- describes past-oriented ($\dot{t} < 0$) retrograde ($\dot{\varphi} < 0$) light rays, i.e, if we reparametrise the light rays in the future-oriented sense, then they are prograde. This reflects the known fact that inside an ergoregion timelike or lightlike curves cannot be retrograde. (Recall that we have fixed the sign ambiguity by requiring $g_{t\varphi} < 0$ in the ergoregion.) The boundary of \mathcal{U}^{erg} may consist of points on the set \mathcal{S} where $g_{tt} = 0$ and of points on the horizon, including possibly the points where the horizon meets the axis. If $\partial g_{tt}/\partial r$ does not go to zero if a point on \mathcal{S} is approached from \mathcal{U}^{erg} , and if the horizon is non-degenerate, our assumptions guarantee that the critical points of \mathcal{R}_- in \mathcal{U}^{erg} lie inside a compact set. Hence, if \mathcal{R}_-^1 and \mathcal{R}_-^0 are Morse functions, the number of critical points of \mathcal{R}_-^1 differs from the number of critical points of \mathcal{R}_-^0 by an even number. In this case the existence of at least one critical point is not guaranteed because \mathcal{R}_- approaches a finite value at the horizon.

An important conclusion from this section is that in a black-hole spacetime with an ergoregion that extends to the horizon the existence of a future-oriented co-rotating circular light ray is not guaranteed, while there is always at least one counter-rotating circular light ray, necessarily outside the ergoregion. So it is possible that in the vacuum case there is an even number of co-rotating circular light rays which are all destroyed if a plasma is introduced.

8.1. Schwarzschild spacetime

In spherical polar coordinates $(t, \varphi, r, \vartheta)$ the Schwarzschild spacetime has the following non-zero metric coefficients:

$$g_{tt}(r, \vartheta) = -\left(1 - \frac{2m}{r}\right), \quad g_{\varphi\varphi}(r, \vartheta) = r^2 \sin^2 \vartheta, \quad (84)$$

$$g_{rr}(r, \vartheta) = \left(1 - \frac{2m}{r}\right)^{-1}, \quad g_{\vartheta\vartheta}(r, \vartheta) = r^2 \quad (85)$$

with a positive mass parameter m , hence

$$\mathcal{R}_+(r, \vartheta) = \mathcal{R}_-(r, \vartheta) = -\frac{r^2 \sin \vartheta}{r - 2m} \sqrt{\omega_0^2 - \omega_p(r, \vartheta)^2 \left(1 - \frac{2m}{r}\right)}. \quad (86)$$

(25) is valid on the domain of outer communication

$$\mathcal{U} = \{(r \sin \vartheta, r \cos \vartheta) \mid r > 2m, 0 < \vartheta < \pi\}. \quad (87)$$

If the plasma density is a function of r only, we may restrict to the equatorial plane and discuss the properties of light rays in terms of a one-dimensional effective potential $V_{\text{eff}}(r)$. So if we want to demonstrate the merits of the formalism developed here we have to consider plasma densities that depend on r and ϑ . We will assume that $-\omega_p^2 g_{tt}$ is bounded on \mathcal{U} . Then the potential (86) is defined on all of \mathcal{U} for $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$.

In vacuum, calculating the gradient of $\mathcal{R}_+ = \mathcal{R}_-$ and equating it to zero produces the well-known result that there is one circular light ray, located at $r = 3m$ and $\vartheta = \pi/2$. Calculating the second derivatives shows that this critical point is a maximum (unstable) in the r -direction and a minimum (stable) in the ϑ direction, so it is a saddle.

In the plasma case, with $\omega_0^2 > \sup(-\omega_p^2 g_{tt})$, we read from (86) that $\mathcal{R}_+ = \mathcal{R}_-$ goes to $-\infty$ at infinity and at the horizon, which exemplifies our general results. This observation already demonstrates that there must be at least one circular light ray in \mathcal{U} .

Stronger statements are possible if we use Proposition 2. If (44), (59) and (64) are satisfied if the horizon, the regular axis or infinity is approached, the critical points of the potential $\mathcal{R}_\pm^\varepsilon$ are confined to a compact set. If \mathcal{R}_\pm^1 is a Morse function, Proposition 2 shows that in comparison to the vacuum case (one saddle) there can be $2n$ additional circular light rays, n extrema and n saddles, so there are n extrema and $n + 1$ saddles.

If the plasma density is symmetric with respect to the equatorial plane, then each extremum in the northern hemisphere must be accompanied by a similar extremum in the southern hemisphere, and the same is true for saddles. Therefore, at least one extremum lies in the equatorial plane if n is odd and at least one saddle lies in the equatorial plane if n is even.

As a specific example, we consider the plasma density.

$$\omega_p^2 = \frac{\omega_c^2 (r^2 + 16 m^2 \sin^2 \vartheta)}{r^2} \quad (88)$$

where ω_c is a constant with the dimension of a frequency. It was already briefly mentioned in Perlick [23] that in this case there are three circular light rays, one in the equatorial plane and two off the equatorial plane. For off-equatorial circular light rays in a plasma on Schwarzschild or Kerr spacetime we also refer to Perlick and Tsupko [9].




Figure 3. The picture on the left shows the equipotential lines $\mathcal{R}_+ = \mathcal{R}_- = \text{constant}$ for the plasma density (88) on Schwarzschild spacetime with $\omega_0 = \sqrt{1.001} \omega_c$. We plot $\rho = r \sin \vartheta$ on the horizontal axis and $z = r \cos \vartheta$ on the vertical axis, choosing m as the unit on both axes. The region inside the horizon is shown as a black disc. The potential goes to $-\infty$ at the horizon and at infinity, while it goes to zero on the axis; this confirms our general results of Sections 5.4, 5.5 and 5.6. There are three circular light rays, marked by black dots: a local maximum of the potential in the equatorial plane and two saddles off the equatorial plane. They all lie on the circle $r = 4m$ which is marked by a dashed line in the picture on the left. In the picture on the right, this sphere with its three circular light rays is shown in three-space. The two off-equatorial light rays are unstable in r -direction and stable in ϑ -direction, while the light ray in the equatorial plane is unstable in both directions. Light rays from the asymptotic region can spiral asymptotically towards these three circular light rays. This is relevant for the construction of the shadow in this spacetime which turns out to be a circular disc, cf. Perlick and Tsupko [8], Section VII.

8.2. Kerr spacetime

In the Kerr spacetime with mass parameter m and spin parameter s , the non-zero metric coefficients in Boyer-Lindquist coordinates are

$$g_{tt} = -\frac{r^2 + a^2 \cos^2 \vartheta - 2mr}{r^2 + a^2 \cos^2 \vartheta}, \quad g_{t\varphi} = -\frac{2ma r \sin^2 \vartheta}{r^2 + a^2 \cos^2 \vartheta},$$

$$g_{\varphi\varphi} = \sin^2 \vartheta \left(r^2 + a^2 + \frac{2ma r \sin^2 \vartheta}{r^2 + a^2 \cos^2 \vartheta} \right) \quad (89)$$

$$g_{rr} = \frac{r^2 + a^2 \cos^2 \vartheta}{r^2 - 2mr + a^2}, \quad g_{\vartheta\vartheta} = r^2 + a^2 \cos^2 \vartheta, \quad (90)$$

hence

$$\mathcal{R}_{\pm} = \pm \frac{2\omega_0 m a r \sin^2 \vartheta}{r^2 + a^2 \cos^2 \vartheta - 2mr}$$

$$- \frac{r^2 + a^2 \cos^2 \vartheta}{r^2 + a^2 \cos^2 \vartheta - 2mr} \sqrt{r^2 - 2mr + a^2} \sin \vartheta \sqrt{\omega_0^2 - \omega_p^2 \frac{(r^2 + a^2 \cos^2 \vartheta - 2mr)}{(r^2 + a^2 \cos^2 \vartheta)}}. \quad (91)$$




Figure 4. These pictures show the equipotential lines $\mathcal{R}_+ = \text{constant}$ (on the left) and $\mathcal{R}_- = \text{constant}$ (on the right) for the plasma density (93) on Kerr spacetime with $a = 3m/4$ and $\omega_0 = \sqrt{1.001} \omega_c$. On the horizontal axis we plot $\rho = r \sin \vartheta$ and on the vertical axis we plot $z = r \cos \vartheta$, using m as the unit on both axes. The region inside the horizon is shown as a black disc, in the picture on the right the ergoregion is shown in grey. There are three corotating light rays in the domain of outer communication, marked by black dots in the picture on the left, a local maximum of the potential in the equatorial plane and two saddles off the equatorial plane. The saddles are maxima (unstable) in the r direction and minima (stable) in the ϑ direction. This is similar to our Schwarzschild example shown in Fig. 3. For the potential \mathcal{R}_- we show in the picture on the right the equipotential lines only on \mathcal{U}^{out} . There are no circular light rays for this particular plasma density in the ergoregion, so we left \mathcal{U}^{erg} in the picture grey. Outside of the ergoregion, we read from the picture that there is exactly one circular light ray, and that it is a saddle, very similar to the vacuum case.

We consider a Kerr black hole with $0 < a < m$, but we will later also give an example for the extreme case $a = m$. We concentrate on the domain of outer communication

$$\mathcal{U} = \{(r \sin \vartheta, r \cos \vartheta) \mid r > m + \sqrt{m^2 - a^2}, 0 < \vartheta < \pi\}. \quad (92)$$

Our general results imply that, if \mathcal{R}_+ and \mathcal{R}_- are Morse functions and if ω_p^2 satisfies the required boundedness conditions, then there is an odd number of co-rotating light rays in \mathcal{U} and an odd number of counter-rotating light rays in \mathcal{U}^{out} , i.e. outside the ergoregion.

As an example we consider on a Kerr spacetime with $a = 3m/4$ the plasma density

$$\omega_p^2 = \frac{\omega_c^2 m^2 (\sin^2(50 r/m) + 2 \sin^2 \vartheta)}{r^2 + a^2 \cos^2 \vartheta} \quad (93)$$

where ω_c is a constant with the dimension of a frequency. For $\omega_0 > \omega_c$, \mathcal{R}_+ is defined on \mathcal{U} while \mathcal{R}_- is defined on the disjoint open sets \mathcal{U}^{out} and \mathcal{U}^{erg} . Then all assumptions of Proposition 2 are satisfied, so we know that there must be an odd number of co-rotating circular light rays in \mathcal{U} and an odd number of counter-rotating circular light rays in \mathcal{U}^{out} . This is indeed the case, as Fig. 4 demonstrates.




Figure 5. This picture shows the equipotential lines $\mathcal{R}_- = \text{constant}$ for the plasma density (94) on Kerr spacetime with $a = m$ and $\omega_0 = 0$. Again, we plot $\rho = r \sin \vartheta$ on the horizontal axis and $z = r \cos \vartheta$ on the vertical axis, choosing m as the unit on both axes. The regions below the (degenerate) horizon and outside of the ergoregion are shown in black, i.e., only the ergoregion is displayed. There are two circular light rays in the ergoregion, marked by black dots in the picture. The inner one is a local maximum, the outer one is a saddle which is a minimum (stable) in r direction and a maximum (unstable) in the ϑ direction. Note that in comparison to the vacuum light rays in (Schwarzschild or) Kerr spacetime the stability properties of the saddle are reversed. In this case the assumptions of Proposition 2 are indeed satisfied, but this was not implied by our general results because the horizon is degenerate.

We have mentioned several times that in a plasma circular light rays with $\omega_0 = 0$ are possible in an ergoregion. Here is an example that illustrates this fact. We choose an extremal Kerr black hole, $a = m$, and a plasma density

$$\omega_p^2 = \omega_c^2 \left(\frac{3}{2} + \sin^2 \left(\frac{2\pi r}{m} \right) \right) \quad (94)$$

where ω_c is again a constant with the dimension of a frequency. Inside the ergoregion, the potential \mathcal{R}_- is defined for $\omega_0 = 0$. There are two counter-rotating circular light rays in this domain, see Fig. 5.

8.3. NUT spacetime

The NUT metric, which was found by Newman, Unti and Tamburino [24] in 1963 as a solution to Einstein's vacuum field equation, depends on two parameters which have the dimension of a length, a mass parameter m and a NUT parameter, also known as

gravitomagnetic charge, n . For a detailed discussion we refer to Griffiths and Podolský [25].

The non-zero metric coefficients are

$$g_{tt} = -\frac{r^2 - 2mr - n^2}{r^2 + n^2}, \quad g_{t\varphi} = \frac{(r^2 - 2mr - n^2) 2n \cos \vartheta}{r^2 + n^2},$$

$$g_{\varphi\varphi} = (r^2 + n^2) \sin^2 \vartheta - \frac{(r^2 - 2mr - n^2) 4n^2 \cos^2 \vartheta}{r^2 + n^2} \quad (95)$$

$$g_{rr} = \frac{r^2 + n^2}{r^2 - 2mr - n^2}, \quad g_{\vartheta\vartheta} = r^2 + n^2, \quad (96)$$

hence

$$\mathcal{R}_{\pm} = \pm \frac{r^2 - 2mr - n^2}{r^2 + n^2} 2n \cos \vartheta \omega_0$$

$$+ \frac{\sqrt{r^2 - 2mr - n^2}}{\sqrt{r^2 + n^2}} \sin \vartheta \sqrt{\omega_0^2 (r^2 - 2mr - n^2) - \omega_p^2 (r^2 + n^2)}. \quad (97)$$

The NUT metric features a black-hole horizon at $r_h = m + \sqrt{m^2 + n^2}$. In the following we concentrate on the domain of outer communication

$$\mathcal{U} = \{(r \sin \vartheta, r \cos \vartheta) \mid r > m + \sqrt{m^2 + n^2}, 0 < \vartheta < \pi\}. \quad (98)$$

Manko and Ruiz [26] have introduced an additional parameter, C , into the NUT metric. Here we have chosen $C = 0$. With this choice, the NUT metric is singular on both axes, $\vartheta = 0$ and $\vartheta = \pi$, outside the horizon and there is a causality-violating region near these axes. There is no ergoregion, and (12) is satisfied on all of \mathcal{U} . Following Bonnor [27], one may interpret the singular axes as spinning rods.

In vacuum, there are two circular light rays, one prograde and one retrograde, off the equatorial plane and symmetric with respect to it, see Jefremov and Perlick [18]. More generally, if $\omega_p(r, \vartheta) = \omega_p(r, \pi - \vartheta)$, one potential is the mirror image of the other, $\mathcal{R}_+(r, \vartheta) = \mathcal{R}_-(r, \pi - \vartheta)$, i.e., for every prograde circular light ray in the northern hemisphere there is a retrograde one in the southern hemisphere and vice versa.

As a specific example we choose $n = 0.635 m$ and

$$\omega_p(r, \vartheta)^2 = \omega_c^2 \left(1 + \frac{16m^2}{r^2} \sin^2 \vartheta \right) \quad (99)$$

where ω_c is a constant with the dimension of a frequency. If $\omega_0 > \omega_c$, the one-parameter family (66) satisfies all assumptions of Proposition 2, for both signs, so there must be an odd number of retrograde and an odd number of prograde circular light rays in the plasma. This is indeed true, as is illustrated by Fig. 6.




Figure 6. This picture shows the equipotential lines $\mathcal{R}_+ = \text{constant}$ for the plasma density (99) on the NUT spacetime with $n = 0.635 m$ and $\omega_0 = \sqrt{1.001} \omega_c$. Again, we plot $\rho = r \sin \vartheta$ on the horizontal axis and $z = r \cos \vartheta$ on the vertical axis, with m as the unit on both axes. The region inside the horizon is shown as a black disc. There are three circular light rays, marked by black dots, a local maximum of the potential and two saddles. The saddles are maxima (unstable) in r direction and minima (stable) in ϑ direction. As in the Schwarzschild example shown in Fig. 3, the three circular light rays lie on a sphere, which is marked by a dashed line. Since $\mathcal{R}_+(r, \vartheta) = \mathcal{R}_-(r, \pi - \vartheta)$, the equipotential lines of \mathcal{R}_- are given by the same picture turned upside down.

9. Conclusions

The potentials \mathcal{R}_+ and \mathcal{R}_- which are at the centre of this paper are useful for several reasons. Firstly, they illustrate the influence of a plasma on the lensing features in axially symmetric and stationary spacetimes in a suggestive way. In particular, plotting the corresponding equipotential surfaces (i.e., the generalised Von Zeipel cylinders) immediately locates the circular light rays and allows to read the direction of the centrifugal-plus-Coriolis force experienced by light rays. Secondly, and maybe even more importantly, these potentials allow to determine in a mathematically precise way how the light rays in a plasma differ from the light rays in vacuum on the same spacetime. This is highly relevant, in particular, in the spacetime of a black hole or some other ultracompact object: With the help of Proposition 2 one can find out if the circular vacuum light rays in the spacetime persist if a plasma is taken into account. Among other things, the answer to this question gives important information on the shadow of such an object in the presence of a plasma.

Roughly speaking, Proposition 2 guarantees that generically the number of circular

light rays in a plasma differs from the number of circular vacuum light rays on the same spacetime by an even number, provided that the plasma density satisfies certain boundedness conditions. As a consequence, if there is an odd number of circular light rays in vacuum, it is impossible that all of them are destroyed when a plasma is introduced. More specifically, we have seen that on the Kerr spacetime Proposition 2, if applicable, implies that in a plasma there is at least one co-rotating and at least one counter-rotating circular light ray in the domain of outer communication, where one counter-rotating circular light ray must be outside the ergoregion. Again, this is of relevance for the shadow.

It is to be emphasised that throughout the present study Einstein's field equation was not used. Therefore, the general results may be applied to the case of a self-gravitating plasma and also to the case that the gravitational field of the plasma is ignored. In the examples we have restricted ourselves to the latter case.

It is certainly desirable to generalise the techniques used in this paper beyond the case of circular light rays, namely to spatially bounded light rays. Potentials very similar to the ones used in this paper have been utilised by Hasse and Perlick [19] and by Halla and Perlick [28] for characterising those regions in axially symmetric and stationary spacetimes where vacuum light rays can have turning points, i.e., where the radius coordinate can have a local maximum or a local minimum, respectively, along a vacuum light ray. However, it is hard to see if and how for that purpose the Brouwer degree of the gradient of the potentials can be helpful.

Acknowledgement

The idea for this paper grew out of a discussion with Pedro Cunha during the Workshop on “Lensing and wave optics in strong gravity” in December 2024 at the Erwin Schrödinger Institute in Vienna, Austria. Quite generally, this workshop proved exceptionally fruitful for me.

Appendix: Proof of Proposition 1

To prove (13), we start out from (6),

$$p_A = g_{AB}\dot{x}^B + g_{Ai}\dot{x}^i, \quad (100)$$

Our assumption (12) guarantees that the matrix

$$(g_{AB}) = \begin{pmatrix} g_{tt} & g_{t\varphi} \\ g_{t\varphi} & g_{\varphi\varphi} \end{pmatrix} \quad (101)$$

is invertible. We denote the inverse

$$(\gamma^{AB}) = \frac{1}{g_{tt}g_{\varphi\varphi} - g_{t\varphi}^2} \begin{pmatrix} g_{\varphi\varphi} & -g_{t\varphi} \\ -g_{t\varphi} & g_{tt} \end{pmatrix}. \quad (102)$$

This allows to solve (100) for \dot{x}^C ,

$$\dot{x}^C = \gamma^{CB} (p_B - g_{bj} \dot{x}^j). \quad (103)$$

Inserting this expression into (7) results in

$$\begin{aligned} 2\mathcal{H} &= g_{ij} \dot{x}^i \dot{x}^j + 2g_{iA} \dot{x}^i \gamma^{AB} (p_B - g_{Bj} \dot{x}^j) \\ &+ g_{AB} \gamma^{AC} (p_B - g_{Bj} \dot{x}^j) \gamma^{BD} (p_D - g_{Dj} \dot{x}^j) + \omega_p^2. \end{aligned} \quad (104)$$

With $g_{AB} \gamma^{BC} = \delta_A^C$ this results in

$$\mathcal{H} = \frac{1}{2} \left(F + \gamma^{BD} p_B p_D + \omega_p^2 \right) \quad (105)$$

with

$$F = \left(g_{ij} - g_{iA} \gamma^{AB} g_{Bj} \right) \dot{x}^i \dot{x}^j. \quad (106)$$

We rewrite this equation with the explicit form of the matrix (γ^{AB}) from (102):

$$2\mathcal{H} - F = \frac{g_{tt} p_\varphi^2 - 2g_{t\varphi} p_t p_\varphi + g_{\varphi\varphi} p_t^2}{g_{tt} g_{\varphi\varphi} - g_{t\varphi}^2} + \omega_p^2, \quad (107)$$

hence

$$\frac{1}{g_{tt}} (g_{tt} g_{\varphi\varphi} - g_{t\varphi}^2) (2\mathcal{H} - F) = p_\varphi^2 - 2 \frac{g_{t\varphi}}{g_{tt}} p_\varphi p_t + \frac{g_{\varphi\varphi}}{g_{tt}} p_t^2 + (g_{tt} g_{\varphi\varphi} - g_{t\varphi}^2) \frac{\omega_p^2}{g_{tt}}. \quad (108)$$

Factorising the second-order polynomial on the right-hand side yields

$$\frac{1}{g_{tt}} (g_{tt} g_{\varphi\varphi} - g_{t\varphi}^2) (2\mathcal{H} - F) = \left(p_\varphi - \frac{g_{t\varphi}}{g_{tt}} p_t - Q \right) \left(p_\varphi - \frac{g_{t\varphi}}{g_{tt}} p_t + Q \right) \quad (109)$$

where

$$Q = \frac{1}{g_{tt}} \sqrt{g_{t\varphi}^2 - g_{tt} g_{\varphi\varphi}} \sqrt{p_t^2 + \omega_p^2 g_{tt}} \quad (110)$$

With $\mathcal{H} = 0$ and the fact that F vanishes at points where $\dot{x}^i = 0$, this proves (13). To prove (14) and (15), we write (103) explicitly at points where $\dot{x}^i = 0$:

$$\dot{\varphi} = \frac{g_{tt} p_\varphi - g_{t\varphi} p_t}{g_{\varphi\varphi} g_{tt} - g_{t\varphi}^2}, \quad \dot{t} = \frac{g_{\varphi\varphi} p_t - g_{t\varphi} p_\varphi}{g_{\varphi\varphi} g_{tt} - g_{t\varphi}^2}. \quad (111)$$

Inserting (13) proves (14) and (15). Finally, we prove (16). To that end we differentiate (109) with respect to x^i and set \mathcal{H} and the \dot{x}^k equal to zero afterwards. This results in

$$\frac{1}{g_{tt}} (g_{tt} g_{\varphi\varphi} - g_{t\varphi}^2) \frac{\partial \mathcal{H}}{\partial x^i} = \mp Q \frac{\partial}{\partial x^i} \left(- \frac{g_{t\varphi}}{g_{tt}} p_t \pm Q \right). \quad (112)$$

Here we have used the fact that $\partial F / \partial x^i$ vanishes at points where $\dot{x}^i = 0$. This is not quite trivial because in the Hamiltonian formalism a partial derivative $\partial / \partial x^\mu$ means that the p_ρ are kept fixed, i.e., we have to express the \dot{x}^i in (106) in terms of p_ρ and x^μ before we can calculate the derivative $\partial F / \partial x^i$. The result gives indeed zero at points where $\dot{x}^i = 0$, because the \dot{x}^j occur quadratic in (106). With (112) in our hands, we now differentiate (3) which yields

$$\ddot{x}^\mu = \frac{\partial g^{\mu\sigma}}{\partial x^i} \dot{x}^i p_\sigma + g^{\mu i} \dot{p}_i. \quad (113)$$

Here we have used that $\dot{p}^A = 0$. With the help of (4) this can be rewritten at points where $\dot{x}^i = 0$ as

$$\ddot{x}^\mu = -g^{\mu i} \frac{\partial \mathcal{H}}{\partial x^i}. \quad (114)$$

Inserting (112) into this expression proves (16). \square

References

- [1] Cunha P V P and Herdeiro C A R 2018 *Gen. Relativ. Gravit.* **50** 42
- [2] Perlick V and Tsupko O Y 2022 *Phys. Rep.* **947** 1–39
- [3] Muhleman D O and Johnston I D 1966 *Phys. Rev. Lett.* **17** 455–458
- [4] Muhleman D O, Ekers R D and Fomalont E 1970 *Phys. Rev. Lett.* **24** 1377–1380
- [5] Perlick V 2000 *Ray Optics, Fermat's Principle, and Applications to General Relativity (Lecture Notes in Physics. Monographs vol m61)* (Berlin: Springer)
- [6] Bisnovaty-Kogan G S and Tsupko O Y 2009 *Gravitation and Cosmology* **15** 20–27
- [7] Bisnovaty-Kogan G S and Tsupko O Y 2010 *Mon. Not. Roy. Astron. Soc.* **404** 1790–1800
- [8] Perlick V and Tsupko O Y 2017 *Phys. Rev.* **D 95** 104003
- [9] Perlick V and Tsupko O Y 2024 *Phys. Rev.* **D 109** 064063
- [10] Abramowicz M A 1971 *Acta Astron.* **21** 81
- [11] Abramowicz M A 1974 *Acta Astron.* **46** 81
- [12] Abramowicz M A 1990 *Mon. Not. Roy. Astron. Soc.* **245** 733–746
- [13] Cunha P V P, Berti E and Herdeiro C A R 2017 *Phys. Rev. Lett.* **119** 251102
- [14] Cunha P V P and Herdeiro C A R 2020 *Phys. Rev. Lett.* **124** 181101
- [15] Cunha P V P, Herdeiro C A R and Novo J P A 2020 *Phys. Rev.* **D 109** 064050
- [16] Breuer R A and Ehlers J 1980 *Proc. Roy. Soc. London A* **370** 389–406
- [17] Synge J L 1960 *Relativity. The general theory* (Amsterdam: North-Holland)
- [18] Jefremov P and Perlick V 2016 *Class. Quant. Grav.* **33** 245014
- [19] Hasse W and Perlick V 2006 *J. Math. Phys.* **47** 042503
- [20] Dinca G and Mawhin J 2000 *Brouwer Degree: The Core of Nonlinear Analysis* (Berlin: Springer)
- [21] Hasse W and Perlick V 2002 *Gen. Relativ. Gravit.* **34** 415–433
- [22] Perlick V 2004 *Liv. Rev. Relativity* **7(9)** <http://www.livingreviews.org/lrr-2004-9>
- [23] Perlick V 2023 *Astron. Rep.* **67, Suppl. 2** S102–S107
- [24] Newman E T, Tamburino L and Unti T W J 1963 *J. Math. Phys.* **4** 915–923
- [25] Griffiths J B and Podolský J 2009 *Exact Space-Times in Einstein's General Relativity* (Cambridge, UK: Cambridge University Press)
- [26] Manko V S and Ruiz E 2005 *Class. Quant. Grav.* **22** 3555–3560
- [27] Bonnor W B 1969 *Math. Proc. Cambr. Philos. Soc.* **66** 145–151
- [28] Halla M and Perlick V 2022 *Phys. Rev.* **D 105** 024027