AROUND THE MERINO–WELSH CONJECTURE: IMPROVING JACKSON'S INEQUALITY

PÉTER CSIKVÁRI

ABSTRACT. The Merino-Welsh conjecture states that for a graph G without loops and bridges we have

 $\max(T_G(2,0), T_G(0,2)) \ge T_G(1,1).$

Later Jackson proved that for any matroid M without loop and coloop we have

$$T_M(3,0)T_M(0,3) \ge T_M(1,1)^2.$$

The value 3 in this statement was improved to 2.9242 by Beke, Csáji, Csikvári and Pituk. In this paper, we further improve on this result by showing that

 $T_M(2.355, 0)T_M(0, 2.355) \ge T_M(1, 1)^2.$

We also prove that the Merino–Welsh conjecture is true for matroids M, where all circuits of M and its dual M^* have length between ℓ and $(\ell - 2)^4$ for some $\ell \ge 6$.

1. INTRODUCTION

Let G be a connected graph without loops and bridges. Merino and Welsh [18] conjectured that

$$\max\left(\alpha(G), \alpha^*(G)\right) \ge \tau(G),$$

where $\alpha(G), \alpha^*(G), \tau(G)$ denote the number of acyclic orientations, strongly connected orientations, and spanning trees of G, respectively. These quantities are evaluations of the Tutte polynomial $T_G(x, y)$, namely $T_G(2, 0) = \alpha(G), T_G(0, 2) = \alpha^*(G)$ and $T_G(1, 1) = \tau(G)$.

Conde and Merino [5] proposed "additive" and "multiplicative" versions of this conjecture:

$$T_G(2,0) + T_G(0,2) \ge 2T_G(1,1)$$

 $T_G(2,0)T_G(0,2) > T_G(1,1)^2$,

respectively. The multiplicative version implies the additive version, which in turn implies the original conjecture. These conjectures also naturally extend to the Tutte polynomial of a matroid without loops and coloops.

While the conjecture holds for certain classes of graphs [16, 19, 21] and matroids [4, 10, 14, 15, 17] it fails for general matroids. Beke, Csáji, Csikvári and Pituk [2] showed that there exist infinitely many matroids without loops and coloops violating the multiplicative version.

Theorem 1.1 (Beke, Csáji, Csikvári and Pituk [2]). There are infinitely many matroids *M* without loops and coloops for which

$$T_M(2,0)T_M(0,2) < T_M(1,1)^2.$$

The research was supported by the MTA-Rényi Counting in Sparse Graphs "Momentum" Research Group, and by Dynasnet European Research Council Synergy project – grant number ERC-2018-SYG 810115.

Furthermore, let x_0 be the largest root of the polynomial $x^3 - 9(x - 1)$. $(x_0 \approx 2.22668...)$ Then for $0 \leq a < x_0$ there are infinitely many matroids M without loops and coloops for which

$$T_M(a,0)T_M(0,a) < T_M(1,1)^2.$$

This paper investigates the following question: for what values of a does the inequality

$$T_M(a,0)T_M(0,a) \ge T_M(1,1)^2$$

hold for all loopless and coloopless matroids M? In this direction the first major result is due to Jackson.

Theorem 1.2 (Jackson [13]). For any matroid M without loops and coloops,

$$T_M(3,0)T_M(0,3) \ge T_M(1,1)^2.$$

Jackson's result was improved by Beke, Csáji, Csikvári and Pituk in the paper [1]. They showed that one can write 2.9243 instead of 3. In this paper, we further improve on this inequality.

Theorem 1.3. For any matroid M without loops and coloops and $a \ge 2.355$,

 $T_M(a,0)T_M(0,a) \ge T_M(1,1)^2.$

We may also study which matroid classes satisfy the product version of the Merino–Welsh conjecture. The following theorem is motivated by the fact that paving matroids satisfy the Merino–Welsh conjecture. A matroid of rank r is a paving matroid if all circuits have length r or r + 1.

Theorem 1.4. Suppose that there exists an $\ell \geq 6$ such that all circuits of the matroid M and its dual M^* have length between ℓ and $(\ell - 2)^4$. Then

$$T_M(2,0)T_M(0,2) \ge T_M(1,1)^2.$$

The proof of Theorem 1.3 and Theorem 1.4 are based on the theory of the permutation Tutte polynomial developed in the paper [1]. While Theorem 1.4 does not imply that paving matroids satisfy the Merino–Welsh conjecture, one can prove this fact by modifying the proof of Theorem 1.4.

Notations. Throughout the paper G = (V, E) is an arbitrary graph and H = (A, B, E) is a bipartite graph. $K_{a,b}$ denotes the complete bipartite graph with parts of size a and b. S_k denotes the star graph on k vertices. For a vertex v the degree of v is denoted by d_v . $N_H(v)$ denotes the set of neighbors of v. If H is clear from the context, then we simply write N(v).

This paper is organized as follows.

- In the next section we introduce the basic concepts from matroid theory that we will use, and revisit the theory of permutation Tutte polynomial $\tilde{T}_H(x, y)$ developed in the paper [1].
- In Section 3 we compute the growth constant $\lim_{n\to\infty} \widetilde{T}_{H_n}(x,0)^{1/n}$ for some family of bipartite graphs. While this section is not necessary for the proofs of Theorem 1.3 and 1.4 it provides an important intuition to study the permutation Tutte polynomial.
- In Section 4 we prove a technical, but very important lemma that provides the basis of the proof of Theorems 1.3 and 1.4.
- In Section 5 we give the proof of Theorem 1.3.
- In Section 6 we prove Theorem 1.4.

- In Section 7 we end the paper some conjectures.
- In the Appendix one can find some tables that are used in the proof of Theorem 1.3.

2. Preliminaries

This section collects the necessary tools from matroid theory together, and recall some of the basic facts from the theory of permutation Tutte polynomial.

2.1. Tutte polynomial and matroids. The Tutte polynomial $T_G(x, y)$ of a graph G is defined as

$$T_G(x,y) = \sum_{A \subseteq E} (x-1)^{k(A)-k(E)} (y-1)^{k(A)+|A|-v(G)},$$

with k(A) denoting the number of connected components of the graph (V, A), see [22]. There is a vast amount of literature on the properties of the Tutte polynomial and its applications, see for instance, [3, 6, 8, 23], or the book [9].

The Tutte polynomial naturally extends to matroids. Recall that a matroid M is a pair (E, \mathcal{I}) such that $\mathcal{I} \subseteq 2^E$, called independent sets, satisfies the axioms (i) $\emptyset \in \mathcal{I}$, (ii) if $A' \subseteq A \in \mathcal{I}$, then $A' \in \mathcal{I}$, and (iii) if $A, B \in \mathcal{I}$ such that |B| < |A|, then there exists an $x \in A \setminus B$ such that $B \cup \{x\} \in \mathcal{I}$. Given a set $S \subseteq E$, the maximal independent subsets of S all have the same cardinality, and this cardinality is called the rank of the matroid, denoted by $\mathcal{P}(S)$. The maximum size independent sets of M are called bases, and their set is denoted by $\mathcal{B}(M)$. The dual of a matroid M is the matroid M^* whose bases are $\{E \setminus B \mid B \in \mathcal{B}(M)\}$. For further details on matroids, see for instance [20]

Given a graph G = (V, E), the edge sets of the spanning forests of G form the independent sets of a matroid M_G called the cycle matroid of G. If G is connected, then the basis of M_G are the spanning trees of G. One can define the Tutte polynomial of a matroid as

$$T_M(x,y) = \sum_{S \subseteq E} (x-1)^{r(E)-r(S)} (y-1)^{|S|-r(S)},$$

where r(S) is the rank of a set $S \subseteq E$. When $M = M_G$, then $T_{M_G}(x, y) = T_G(x, y)$. A loop in a matroid M is an element $x \in E$ such that $r(\{x\}) = 0$, that is, $\{x\} \notin \mathcal{I}$, and a coloop is an element that is a loop in the dual M^* of the matroid M. Equivalently, a coloop is an element that is in every base of M. For a cycle matroid M_G , loops correspond to loop edges and coloops correspond to bridges in the graph G.

Hence it was suggested that the inequalities

$$\max(T_M(2,0), T_M(0,2) \ge T_M(1,1),$$
$$T_M(2,0) + T_M(0,2) \ge 2T_M(1,1),$$
$$T_M(2,0)T_M(0,2) \ge T_M(1,1)^2$$

may hold true for all matroids M without loops and coloops. (These versions appear explicitly in [10], but were treated much earlier without explicitly calling them conjectures.) Note that for general matroids, all these versions are equivalent in the following sense: if one of them is true for all matroids, then the others are also true for all matroids. Applying the maximum version to $M \oplus M^*$ with M^* being the dual of M leads to the multiplicative version of the conjecture. (Here $M \oplus N$ denotes the direct sum of the matroids M and N.) 2.2. **Permutation Tutte polynomial.** The proof of Theorem 1.3 heavily relies on the theory of permutation Tutte polynomial. The idea is that Tutte polynomial $T_G(x, y)$ can be written as a sum of permutation Tutte polynomials $\tilde{T}_{H_j}(x, y)$ for certain bipartite graphs H_j . As a consequence certain inequalities valid for the permutation Tutte polynomial transfers to the Tutte polynomial.

Definition 2.1 ([1]). Let H = (A, B, E) be a bipartite graph. Suppose that V(H) = [m]. For a permutation $\pi : [m] \to [m]$, we say that a vertex $i \in A$ is internally active if

$$\pi(i) > \max_{j \in N_H(i)} \pi(j),$$

where the maximum over an empty set is set to be $-\infty$. Similarly, we say that vertex $j \in B$ is externally active if

$$\pi(j) > \max_{i \in N_H(j)} \pi(i).$$

Let $ia(\pi)$ and $ea(\pi)$ be the number of internally and externally active vertices in A and B, respectively. Let

$$\widetilde{T}_H(x,y) = \frac{1}{m!} \sum_{\pi \in S_m} x^{\mathrm{ia}(\pi)} y^{\mathrm{ea}(\pi)}.$$

We will call $\widetilde{T}_H(x, y)$ the permutation Tutte polynomial of H.

The above definition is motivated by the following theorem of Tutte.

Theorem 2.2 (Tutte [22]). Let G be a connected graph with m edges. Label the edges with 1, 2, ..., m arbitrarily. In the case of a spanning tree T of G, let us call an edge $e \in E(T)$ internally active if e has the largest label among the edges in the cut determined by T and e by removing e from T. Let us call an edge $e \notin E(T)$ externally active if e has the largest label among the edges in the cycle determined by T and e by adding e to T. Let ia(T) and ea(T) be the number of internally and externally active edges, respectively. Then

$$T_G(x,y) = \sum_{T \in \mathcal{T}(G)} x^{\mathrm{ia}(T)} y^{\mathrm{ea}(T)},$$

where the summation goes for all spanning trees of G.

Theorem 2.2 was originally a definition for the Tutte polynomial [22]. This characterization of the Tutte polynomial immediately shows that the coefficients of the Tutte polynomial are non-negative. In this theorem, we are restricted to the same labelling of the edges for all spanning trees. For those who have never seen this definition before, it might be very surprising that the Tutte polynomial is independent of the actual choice of the labelling.

To explain the connection between $T_G(x, y)$ and $\widetilde{T}_H(x, y)$, we need the concept of the local basis exchange graph.

Definition 2.3. The local basis exchange graph H[T] of a graph G = (V, E) with respect to a spanning tree T is defined as follows. The graph H[T] is a bipartite graph whose vertices are the edges of G. One bipartite class consists of the edges of T, the other consists of the edges of $E \setminus T$, and we connect a spanning tree edge e with a non-edge f if f is in the cut determined by e and T, equivalently, e is in the cycle determined by f and T.

Clearly, this definition works for general matroids and their basis: if A is a basis, then $e \in A$ and $f \in E \setminus A$ are adjacent in the bipartite graph H[A] if A - e + f is again a basis.

Figure 1 depicts a graph G with a spanning tree T and the bipartite graph H[T] obtained from T.

For a fixed labelling of the edges of G, we get a labelling of the vertices of H[T], and the internally (externally) active edges of G correspond to internally (externally) active vertices of H[T], so the two definitions of internal and external activity are compatible. The following lemma is crucial for us, so we even included its proof.



FIGURE 1. Example for a graph G and the local basis exchange graph H[T] obtained from a spanning tree T.

Lemma 2.4 (Beke, Csáji, Csikvári, Pituk [1]). Let G be a graph. For each spanning tree T of G, let H[T] be the local basis exchange graph with respect to T. Then

$$T_G(x,y) = \sum_{T \in \mathcal{T}(G)} \widetilde{T}_{H[T]}(x,y),$$

where the sum is over the set of spanning trees $\mathcal{T}(G)$ of G.

Proof. For a fixed spanning tree T and a permutation π of the edges, the internally and externally active edges correspond to the internally and externally active vertices of H[T]. Hence

$$T_G(x,y) = \sum_{T \in \mathcal{T}(G)} x^{\mathrm{ia}_{H[T]}(\pi)} y^{\mathrm{ea}_{H[T]}(\pi)}.$$

Now averaging it for all permutations $\pi \in S_m$ we get that

$$T_{G}(x,y) = \frac{1}{m!} \sum_{\pi \in S_{m}} T_{G}(x,y)$$

= $\frac{1}{m!} \sum_{\pi \in S_{m}} \sum_{T \in \mathcal{T}(G)} x^{\mathrm{ia}_{H[T]}(\pi)} y^{\mathrm{ea}_{H[T]}(\pi)}$
= $\sum_{T \in \mathcal{T}(G)} \frac{1}{m!} \sum_{\pi \in S_{m}} x^{\mathrm{ia}_{H[T]}(\pi)} y^{\mathrm{ea}_{H[T]}(\pi)}$
= $\sum_{T \in \mathcal{T}(G)} \widetilde{T}_{H[T]}(x,y).$

Remark 2.5. The local basis exchange graph H[T] has an isolated vertex if and only if G contains a bridge or a loop. Furthermore, H[T] is connected if and only G is 2-connected.

The following lemma enables us to study quadratic inequalities of the Tutte polynomial.

Lemma 2.6 (Transfer lemma [1]). Let $x_0, x_1, x_2, y_0, y_1, y_2 \ge 0$. Suppose that for any bipartite graph H, we have

$$\widetilde{T}_H(x_1, y_1)\widetilde{T}_H(x_2, y_2) \ge \widetilde{T}_H(x_0, y_0)^2.$$

Then for any graph G, we have

 $T_G(x_1, y_1)T_G(x_2, y_2) \ge T_G(x_0, y_0)^2.$

Remark 2.7. For us the following version of the transfer lemma will be a bit more convenient. If for any basis A of a matroid M, the local basis exchange graph H[A] satisfies that

$$\widetilde{T}_{H[A]}(x_1, y_1)\widetilde{T}_{H[A]}(x_2, y_2) \ge \widetilde{T}_{H[A]}(x_0, y_0)^2$$

then

$$T_M(x_1, y_1)T_M(x_2, y_2) \ge T_M(x_0, y_0)^2$$

The proof of this claim is exactly the same as of the transfer lemma.

A key example for bounding the permutation Tutte polynomial is the following theorem proved in [1].

Theorem 2.8 ([1]). Let H be an arbitrary bipartite graph, and let d_i be the degree of vertex i. Suppose that $0 \le x \le 1$ and $y \ge 1$ or $0 \le y \le 1$ and $x \ge 1$. Then

$$\widetilde{T}_H(x,y) \ge \prod_{i \in A} \left(1 + \frac{x-1}{d_i+1} \right) \cdot \prod_{j \in B} \left(1 + \frac{y-1}{d_j+1} \right).$$

The proof of Lemma 2.8 relies on following inequality of Harris that is also a special case of the FKG-inequality [11].

Lemma 2.9 (Harris [12], Fortuin, Kasteleyn, Ginibre [11]). Suppose that $\mu = \mu_1 \otimes \cdots \otimes \mu_N$ is a product measure on a set $S_1 \times S_2 \times \cdots \times S_N$, and X_1, \ldots, X_t are non-negative monotone increasing functions in the sense that if $x_i \geq x'_i$ for $i = 1, \ldots, N$, then for $1 \leq j \leq t$ we have

Then

$$\mathbb{E}\left[\frac{t}{\mathbf{\Pi}}\mathbf{V}\right] > \frac{t}{\mathbf{\Pi}}\mathbb{E}\left[\mathbf{V}\right]$$

 $X_i(x_1,\ldots,x_N) > X_i(x'_1,\ldots,x'_N).$

$$\mathbb{E}_{\mu}\left[\prod_{j=1}^{j} X_{j}\right] \geq \prod_{j=1}^{j} \mathbb{E}_{\mu}[X_{j}].$$

Furthermore, if X is monotone increasing and Y is monotone decreasing, then

$$\mathbb{E}[XY] \le \mathbb{E}[X]\mathbb{E}[Y].$$

In this paper, we will use the Harris inequality for both $[0,1]^N$ and the set $\{0,1\}^A$ for some set A.

In what follows, we repeatedly use the following crucial idea to express $T_H(x, y)$. We can create a random ordering of the vertices of H as follows: for each vertex i we choose a uniform random number x_i from the interval [0, 1]. The numbers x_i then determine an ordering of the edges. The probability that two numbers are equal is 0.

Lemma 2.10 ([1]). Let H be a bipartite graph and let $\widetilde{T}_H(x, y) = \sum t_{i,j}(H)x^iy^j$. Let v(H) = m and let $x_1, x_2, \ldots x_m$ be i.i.d. random variables with distribution $x_i \sim U(0, 1)$. Let $I(A) = |\{v \in A | x_v \ge x_{v'} \text{ for } v' \in N_H(v)\}|$ and $I(B) = |\{v \in B | x_v \ge x_{v'} \text{ for } v' \in N_H(v)\}|$. Then

$$\mathbb{P}\left(I(A) = i, I(B) = j\right) = t_{i,j}(H).$$

In what follows we do a little trick. For $i \in A$ we generate $x_i \sim U(0, 1)$ as before, but for $j \in B$ we actually first generate a uniformly random number y_j from [0, 1] and let $x_j = 1 - y_j$. The role of this trick will be apparent soon.

For $i \in A$, let us introduce the random variable

$$X_i(x_i, \{y_j\}_{j \in B}) = \begin{cases} x & \text{if } \max_{j \in N_H(i)} (1 - y_j) \le x_i, \\ 1 & \text{if } \max_{j \in N_H(i)} (1 - y_j) > x_i. \end{cases}$$

and for $j \in B$, let

$$Y_j(\{x_i\}_{i \in A}, y_j) = \begin{cases} y & \text{if } \max_{i \in N_H(j)} x_i \le 1 - y_j, \\ 1 & \text{if } \max_{i \in N_H(j)} x_i \ge 1 - y_i. \end{cases}$$

Lemma 2.11. (a) We have

$$\widetilde{T}_H(x,y) = \mathbb{E}\left[\prod_{i \in A} X_i \cdot \prod_{j \in B} Y_j\right].$$

(b1) If $x \ge 1$, then $X_i(x_i, \{y_j\}_{j \in B})$ is a monotone increasing function for each $i \in A$. (b2) If $0 \le x \le 1$, then $X_i(x_i, \{y_j\}_{j \in B})$ is a monotone decreasing function for each $i \in A$. (b3) For $0 \le y \le 1$ the function $Y_j(\{x_i\}_{i \in A}, y_j)$ is monotone increasing for each $j \in B$. (b4) Finally, for $0 \le y \le 1$ the function $Y_j(\{x_i\}_{i \in A}, y_j)$ is monotone decreasing for each $j \in B$.

Remark 2.12. This lemma is the reason why we generated x_j by $x_j = 1 - y_j$ for $j \in B$. If we consider the function

$$X'_{i}(x_{i}, \{x_{j}\}_{j \in B}) = \begin{cases} x & \text{if } \max_{j \in N_{H}(i)} x_{j} \le x_{i}, \\ 1 & \text{if } \max_{j \in N_{H}(i)} x_{j} > x_{i}. \end{cases}$$

instead of $X_i(x_i, \{y_j\}_{j \in B})$, this would be neither increasing, nor decreasing.

We will also need a very little extension of the above ideas, where each vertex gets its own activity: let For $i \in A$, let us introduce the random variable

$$\widehat{X}_{i}(x_{i}, \{y_{j}\}_{j \in B}) = \begin{cases} x^{(i)} & \text{if } \max_{j \in N_{H}(i)}(1-y_{j}) \leq x_{i}, \\ 1 & \text{if } \max_{j \in N_{H}(i)}(1-y_{j}) > x_{i}. \end{cases}$$

and for $j \in B$, let

$$\widehat{Y}_{j}(\{x_{i}\}_{i\in A}, y_{j}) = \begin{cases} y^{(j)} & \text{if } \max_{i\in N_{H}(j)} x_{i} \leq 1 - y_{j}, \\ 1 & \text{if } \max_{i\in N_{H}(j)} x_{i} \geq 1 - y_{i}. \end{cases}$$

The following lemma is just a trivial extension of the previous lemma together with Harris' inequality.

Lemma 2.13. (a1) If $i \in A$ and $x^{(i)} \ge 1$, then $\widehat{X}_i(x_i, \{y_j\}_{j \in B})$ is a monotone increasing function.

(a2) If $i \in A$ and $0 \leq x^{(i)} \leq 1$, then $\widehat{X}_i(x_i, \{y_j\}_{j \in B})$ is a monotone decreasing function. (a3) For $j \in B$ and $0 \leq y^{(j)} \leq 1$ the function $\widehat{Y}_j(\{x_i\}_{i \in A}, y_j)$ is monotone increasing for each $j \in B$. (a4) Finally, if $j \in B$ and $0 \leq y^{(j)} \leq 1$ the function $Y_j(\{x_i\}_{i \in A}, y_j)$ is monotone decreas-

(a4) Finally, if $j \in B$ and $0 \le y^{(j)} \le 1$ the function $Y_j(\{x_i\}_{i \in A}, y_j)$ is monotone decreasing.

(:)

(b) If
$$x^{(i)} \ge 1$$
 for all $i \in A$ and $0 \le y^{(j)} \le 1$, then

$$\mathbb{E}\left[\prod_{i \in A} \widehat{X}_i \cdot \prod_{j \in B} \widehat{Y}_j\right] \ge \prod_{i \in A} \mathbb{E}[\widehat{X}_i] \cdot \prod_{j \in B} \mathbb{E}[\widehat{Y}_j] = \prod_{i \in A} \left(\frac{x^{(i)}}{d_i + 1} + \frac{d_i}{d_i + 1}\right) \cdot \prod_{j \in B} \left(\frac{y^{(j)}}{d_j + 1} + \frac{d_j}{d_j + 1}\right)$$

We will use one more lemma from the paper [1], namely the gluing lemma. Originally, this lemma used the condition that H_1 and H_2 are trees, but the proof never used this condition.

Lemma 2.14 (Gluing lemma [1]). Let $x \ge 1$ and $0 \le y \le 1$. Let H_1 be a rooted graph with root vertex v_1 . Let H_2 be another rooted graph with root vertex v_2 . Let H be obtained from H_1 and H_2 by identifying v_1 and v_2 in the union of H_1 and H_2 . Let v be the vertex obtained from identifying v_1 and v_2 . Assume that the bipartite parts of H determines the bipartite parts of H_1 and H_2 , that is, if $v \in A(H)$, then $v_1 \in A(H_1)$ and $v_2 \in A(H_2)$, and if $v \in B(H)$, then $v_1 \in B(H_1)$ and $v_2 \in B(H_2)$.

(a) If $v \in A$, then

8

$$x\widetilde{T}_H(x,y) \ge \widetilde{T}_{H_1}(x,y)\widetilde{T}_{H_2}(x,y).$$

(b) If $v \in B$, then

$$\widetilde{T}_H(x,y) \ge \widetilde{T}_{H_1}(x,y)\widetilde{T}_{H_2}(x,y).$$

(c) In particular,

$$\widetilde{T}_{H}(x,0)\widetilde{T}_{H}(0,x) \ge \frac{1}{x}(\widetilde{T}_{H_{1}}(x,0)\widetilde{T}_{H_{1}}(0,x))(\widetilde{T}_{H_{2}}(x,0)\widetilde{T}_{H_{2}}(0,x))$$

3. WARM-UP: ASYMPTOTIC COMPUTATION OF SOME PERMUTATION TUTTE POLYNOMIAL

In this section we motivate an important technique of this paper by computing the asymptotic value of the permutation Tutte polynomial of certain bipartite graphs.

Let us start with a very simple example, the complete bipartite graph.

Theorem 3.1. We have

$$\widetilde{T}_{K_{a,b}}(x,0) = ab \int_0^1 \int_s^1 s^{b-1} (s+x(t-s))^{a-1} dt \, ds.$$

Proof. We generate the random permutation on $V(H) = A \cup B$ by first generating an $x_v \in (0, 1)$ uniformly at random for all $v \in V$, and then we take the relative order of x_v 's. Let $t = \max_{v \in A} x_v$ and $s = \max_{v \in B} x_v$. We have *ab* choices for the vertices that take the value t and s, let these vertices be v_A and v_B . If s > t, then the permutation has weight 0 as v_B is an active vertex. If t > s, then no vertex can be active in B. For each $w \in A$ that is not equal to v_A two things can happen: if $x_w < s$, then w is not active, so $X_w = 1$, or $s < x_w < t$ and then w is active so $X_w = x$. For a vertex $w \in B$ we simply need to have $x_w < s$ and w will not be active. The integral formula then follows. \Box

Remark 3.2. Suppose that $a = \alpha m$ and $b = \beta m$, where α, β are fixed such that $\alpha + \beta = 1$ and $m \to \infty$. Then the exponential growth constant of $\widetilde{T}_{K_{a,b}}(x,0)$ is simply

$$\lim_{m \to \infty} \widetilde{T}_{K_{a,b}}(x,0)^{1/m} = \max_{s,t} s^{\beta} (s+x(t-s))^{\alpha}.$$

Clearly, at the maximum we have t = 1 and we simply need to maximize $s^{\beta}(s+x(1-s))^{\alpha}$. This turns out to be at $s = \min\left(1, \frac{\beta x}{\beta-1}\right)$. If $\beta < \frac{x-1}{x}$, then the growth constant is

$$\left(\frac{\beta x}{\beta - 1}\right)^{\beta} \left(x + (1 - x)\frac{\beta x}{x - 1}\right)^{\alpha} = \alpha^{\alpha} (1 - \alpha)^{1 - \alpha} \frac{x}{(x - 1)^{1 - \alpha}}.$$

If $\beta \geq \frac{x-1}{x}$, then the exponential growth constant is simply 1.

Let us consider the graph $H_{a,b,c}$ introduced in [1]: we start with a complete bipartite graph $K_{a,b}$ with vertex set $A \cup B$, and then attach c pendant leaves to c distinct vertices of B, let C be the set of these leaf vertices. So the resulting bipartite graph has a+c vertices on one side and b vertices on the other side. The graphs $H_{n,n,n}$ played an important role in the refutation of the matroidal version of the Merino–Welsh conjecture.



FIGURE 2. The graph $H_{6,6,6}$.

It turns out that for even n there are matroids for which all local basis exchange graphs are isomorphic to $H_{n,n,n}$. Indeed, all local basis exchange graphs of the matroid $U_{\frac{3}{2}n,n}^{(2)}$ are isomorphic to $H_{n,n,n}$, where $U_{\frac{3}{2}n,n}$ is the uniform matroid of rank n on $\frac{3}{2}n$ elements, and the matroid $M^{(2)}$ is obtained from M by replacing each element of M with two parallel elements. Furthermore, we have

$$\widetilde{T}_{H_{n,n,n}}(2,0)\widetilde{T}_{H_{n,n,n}}(2,0) < \widetilde{T}_{H_{n,n,n}}(1,1)^2$$

for large enough n.

We can now give a simple proof of this fact.

Theorem 3.3. We have

$$\lim_{n \to \infty} \widetilde{T}_{H_{n,n,n}}(x,0)^{1/n} = \max_{s \in [0,1]} \left(s + x(1-s) \right) \left(xs + (1-x)\frac{s^2}{2} \right) = \begin{cases} \frac{1}{3\sqrt{3}}\frac{x^3}{x-1} & \text{if } x \ge \sqrt{3}, \\ \frac{1}{2}(x+1) & \text{if } 1 < x \le \sqrt{3} \end{cases}$$

and

$$\lim_{n \to \infty} \widetilde{T}_{H_{n,n,n}}(0,x)^{1/n} = \max_{s \in [0,1]} t\left(\frac{t^2}{2} + \left(\frac{1}{2} - \frac{t^2}{2}\right)x\right) = \begin{cases} \frac{1}{3\sqrt{3}} \frac{x^{3/2}}{(x-1)^{1/2}} & \text{if } x \ge \frac{3}{2}, \\ \frac{1}{2} & \text{if } 1 < x \le \frac{3}{2}. \end{cases}$$

In particular, if $x \ge \sqrt{3}$ we have

$$\lim_{n \to \infty} \left(\widetilde{T}_{H_{n,n,n}}(x,0) \widetilde{T}_{H_{n,n,n}}(0,x) \right)^{1/n} = \left(\frac{x^3}{9(x-1)} \right)^{3/2}.$$

Sketch of the proof. As before we generate the random permutation on V(H) by first generating an $x_v \in (0,1)$ uniformly at random for all $v \in V$, and then we take the relative order of x_v 's. Let

$$t = \max_{v \in A} x_v$$
 and $s = \max_{v \in B} x_v$.

Let v_A and v_B be the vertices, where these maximum are achieved. (Note that t is only maximum on A and not on $A \cup C$.) If s < t, then the contribution of this case to

 $\widetilde{T}_{H_{n,n,n}}(x,0)$ is

(3.1)
$$I_1 := n^2 \int_0^1 \int_s^1 (s + x(t - s))^{n-1} \left(\int_0^s (s_i + x(1 - s_i)) \, ds_i \right)^{n-1} (s + x(1 - s)) \, dt ds.$$

Here s + x(t - s) is the contribution of a vertex $w \in A$ that is not v_A . If $x_w = s_i < s$ for some $w \in B$ that is not v_B , then the attached leaf in C has contribution $s_i + x(1 - s_i)$. (Note that attached leaf vertex u can have a value bigger than t as t was only a maximum among the vertices of A.) The last term s + x(1 - s) is simply the contribution of the leaf attached to v_B .

If t < s, then the contribution of this case to $\widetilde{T}_{H_{n,n,n}}(x,0)$ is

(3.2)
$$I_2 := n^2 \int_0^1 \int_s^1 t^{n-1} \left(\int_0^t (s_i + x(1-s_i)) \, ds_i + \int_t^s x(1-s_i) \, ds_i \right)^{n-1} x(1-s) \, dt ds.$$

Here the contribution of a vertex $w \in A$ is simply t as it can never be active since s < t. If we have a vertex $w \in B$ that is not v_B with $x_w = s_i < s$, then we need to distinguish two cases. If $s_i < t$, then w is not active, and so the attached leaf has a contribution $s_i + x(1 - s_i)$. If $s_i > t$ then, in order to make w inactive the attached leaf u should have a value $x_u > s_i$, but then it is automatically active. The contribution of the leaf attached to v_B is x(1 - s) as it has to be active to make v_B inactive.

We have

$$\lim_{k \to \infty} I_1^{1/n} = \max_{0 \le s \le t \le 1} (s + x(t - s)) \left(xs + (1 - x) \frac{s^2}{2} \right)$$
$$= \max_{0 \le s \le 1} (s + x(1 - s)) \left(xs + (1 - x) \frac{s^2}{2} \right)$$

The function $(s+x(1-s))\left(xs+(1-x)\frac{s^2}{2}\right)$ has either maximum at $\left(1-\frac{1}{\sqrt{3}}\right)\frac{x}{x-1}$ if this is less than 1, or at s = 1. In the first case, its value is $\frac{1}{3\sqrt{3}}\frac{x^3}{x-1}$. In the second case, its value is $\frac{1}{2}(x+1)$.

On the other hand,

r

$$\lim_{n \to \infty} I_2^{1/n} = \max_{0 \le t \le s \le 1} t \left((1-x)\frac{t^2}{2} + xt + x(s-t) - x \left(\frac{s^2}{2} - \frac{t^2}{2}\right) \right)$$
$$= \max_{0 \le t \le s \le 1} t \left(xs - x\frac{s^2}{2} + \frac{t^2}{2} \right)$$
$$= \max_{0 \le s \le 1} \left(x \left(s - \frac{s^2}{2} \right) + \frac{1}{2} \right)$$
$$= \frac{1}{2}(x+1),$$

so this is always at most $\lim_{n\to\infty} I_1^{1/n}$.

Then we have

$$\lim_{n \to \infty} \widetilde{T}_{H_{n,n,n}}(x,0)^{1/n} = \lim_{n \to \infty} (I_1 + I_2)^{1/n}$$
$$= \max\left(\lim_{n \to \infty} I_1^{1/n}, \lim_{n \to \infty} I_2^{1/n}\right)$$
$$= \lim_{n \to \infty} I_1^{1/n}.$$

This proves the first statement. The second statement is actually simpler. We have

$$\widetilde{T}_{H_{n,n,n}}(0,x) = n^2 \int_0^1 \int_0^s t^{n-1} \left(\int_0^t s_i \, ds_i + x \int_t^s s_i \, ds_i \right)^{n-1} s \, dt ds$$
$$= n^2 \int_0^1 \int_0^s t^{n-1} \left(\frac{t^2}{2} + x \left(\frac{s^2}{2} - \frac{t^2}{2} \right) \right)^{n-1} s \, dt ds.$$

From this we get that

$$\lim_{n \to \infty} \widetilde{T}_{H_{n,n,n}}(0,x)^{1/n} = \max_{0 \le t \le s \le 1} t \left(\frac{t^2}{2} + x \left(\frac{s^2}{2} - \frac{t^2}{2} \right) \right)$$
$$= \max_{0 \le t \le 1} t \left(\frac{t^2}{2} + x \left(\frac{1}{2} - \frac{t^2}{2} \right) \right)$$

This expression has either maximum at $x = \left(\frac{x}{3(x-1)}\right)^{1/2}$ if $x \ge \frac{3}{2}$ or at 1 if $x \le \frac{3}{2}$. In the first case its value is $\frac{1}{3\sqrt{3}} \frac{x^{3/2}}{(x-1)^{1/2}}$. In the second case, its value is $\frac{1}{2}$. This completes the sketch of the proof.

Remark 3.4. It is clear from the above examples that even though the permutation Tutte polynomial might be complicated still it is possible to compute the exponential growth of it for many families of bipartite graphs. Another important take-away that it is often possible to understand those permutations that contributes the most weight to the permutation Tutte polynomial, and concentrating for these permutations might give a better lower or upper bound for the permutation Tutte polynomial.

4. The main lemma

In this section we give the main ingredient of the proofs of Theorems 1.3 and 1.4.

Theorem 4.1. Let H = (A, B, E) be a bipartite graph. For $x \ge 2$ and a fixed $0 \le s \le 1$ let us introduce the function

$$\gamma_{x,s}(d) = \frac{(d+x)s}{(d+x)s + (d+1)x(1-s)}$$

Then

$$\widetilde{T}_H(x,0) \ge \prod_{v \in A} \left(\frac{(d_v + x)s}{d_v + 1} + x(1-s) \right) \cdot \prod_{u \in B} \left(s - \frac{s}{d_u + 1} \prod_{v \in N(u)} \gamma_{x,s}(d_v) \right).$$

Proof. Recall that

$$\widetilde{T}_{H}(x,0) = \int_{[0,1]^{A}} \int_{[0,1]^{B}} \prod_{v \in A} X_{v}(\underline{x},\underline{y}) \cdot \prod_{u \in B} Y_{u}(\underline{x},\underline{y}) d\underline{x} d\underline{y}.$$

The first idea is that we only take those values for which $x_u = 1 - y_u \leq s$ for $u \in B$. Hence

$$\widetilde{T}_{H}(x,0) \ge \int_{[0,1]^{A}} \int_{[1-s,1]^{B}} \prod_{v \in A} X_{v}(\underline{x},\underline{y}) \cdot \prod_{u \in B} Y_{u}(\underline{x},\underline{y}) d\underline{x} d\underline{y}.$$

For each $v \in A$ we decompose the integral over [0, 1] to [0, s] and [s, 1], this way we decomposed the integral $[0, 1]^A$ to the sum of $2^{|A|}$ integrals. Since $x_u \leq s$ for $u \in B$ we

immediately get that if $x_v \ge s$, then $X_v(\underline{x}, \underline{y}) = x$. Let $T \subseteq A$ be the set of those vertices $v \in A$ for which $x_v \ge s$. Then

$$\int_{[0,1]^A} \int_{[1-s,1]^B} \prod_{v \in A} X_v(\underline{x},\underline{y}) \cdot \prod_{u \in B} Y_u(\underline{x},\underline{y}) d\underline{x} d\underline{y}$$
$$= \sum_{T \subseteq A} (x(1-s))^{|T|} \int_{[0,s]^{A \setminus T}} \int_{[1-s,1]^B} \prod_{v \in A} X_v(\underline{x},\underline{y}) \cdot \prod_{u \in B} Y_u(\underline{x},\underline{y}) d\underline{x} d\underline{y}.$$

Let $N(T) \subseteq B$ be the neighbors of T. If $u \in N(T)$, then $\widehat{Y}_u(\underline{x}, \underline{y}) := Y_u(\underline{x}, \underline{y}) = 1$, because the vertex u cannot be active anymore. Hence

$$I_T := \int_{[0,s]^{A\setminus T}} \int_{[1-s,1]^B} \prod_{v \in A} X_v(\underline{x},\underline{y}) \cdot \prod_{u \in B} Y_u(\underline{x},\underline{y}) d\underline{x} d\underline{y}$$

corresponds to $s^{|A \setminus T| + |B|}$ times a permutation Tutte polynomial, where an active vertex $i \in A \setminus T$ gets a weight $x^{(i)} = x$ while in B an active vertex j gets a weight $y^{(j)} = 1$ if it is N(T), and $y^{(j)} = 0$ if it is $B \setminus N(T)$, so for this term we have a lower bound from Lemma 2.13:

$$I_T = s^{|A \setminus T| + |B|} \mathbb{E} \left[\prod_{i \in A \setminus T} \widehat{X}_i \cdot \prod_{j \in B} \widehat{Y}_j \right]$$

$$\geq s^{|A \setminus T| + |B|} \prod_{i \in A} \mathbb{E}[\widehat{X}_i] \cdot \prod_{j \in B} \mathbb{E}[\widehat{Y}_j]$$

$$= s^{|A \setminus T| + |B|} \prod_{v \in A \setminus T} \left(1 + \frac{x - 1}{d_v + 1} \right) \prod_{u \in B \setminus N(T)} \left(1 - \frac{1}{d_u + 1} \right).$$

So far we got that

$$\widetilde{T}_{H}(x,0) = \sum_{T \subseteq A} (x(1-s))^{|T|} I_{T}$$

$$\geq \sum_{T \subseteq A} (x(1-s))^{|T|} \cdot s^{|A \setminus T| + |B|} \prod_{v \in A \setminus T} \left(1 + \frac{x-1}{d_{v}+1} \right) \prod_{u \in B \setminus N(T)} \left(1 - \frac{1}{d_{u}+1} \right).$$

$$(4.1) \qquad = \sum_{T \subseteq A} (x(1-s))^{|T|} \cdot \prod_{v \in A \setminus T} \left(\frac{d_{v}}{d_{v}+1} s + \frac{x}{d_{v}+1} s \right) \cdot \prod_{u \in B \setminus N(T)} \frac{d_{u}s}{d_{u}+1} \cdot \prod_{u \in N(T)} s$$

Next let

$$Z = \prod_{v \in A} \left(\frac{d_v}{d_v + 1} s + \frac{x}{d_v + 1} s + x(1 - s) \right),$$

and let us consider the measure μ on $\{0,1\}^A$ for which

$$\mu(T) = \frac{1}{Z} (x(1-s))^{|T|} \prod_{A \setminus T} \left(\frac{d_v}{d_v + 1} s + \frac{x}{d_v + 1} s \right).$$

Note that μ is a product measure $\mu = \bigoplus_{v \in A} \mu_v$, where

$$\mu_v(0) = \frac{\frac{d_v}{d_v+1}s + \frac{x}{d_v+1}s}{\frac{d_v}{d_v+1}s + \frac{x}{d_v+1}s + x(1-s)} \quad \text{and} \quad \mu_v(1) = \frac{x(1-s)}{\frac{d_v}{d_v+1}s + \frac{x}{d_v+1}s + x(1-s)}$$

For each $u \in B$ consider the function

$$f_u(T) = \begin{cases} s & \text{if } u \in N(T), \\ \frac{d_u}{d_u + 1}s & \text{if } u \notin N(T). \end{cases}$$

Then we can rewrite inequality (4.1) as follows:

$$\widetilde{T}_{H}(x,0) \geq \sum_{T \subseteq A} (x(1-s))^{|T|} \cdot \prod_{v \in A \setminus T} \left(\frac{d_v}{d_v+1} s + \frac{x}{d_v+1} s \right) \cdot \prod_{u \in B \setminus N(T)} \frac{d_u s}{d_u+1} \cdot \prod_{u \in N(T)} s$$
$$= Z \sum_{T \subseteq A} \mu(T) \prod_{u \in B} f_u(T).$$

All functions f_u are monotone increasing on the space $\{0,1\}^A$ identified with the subsets T of A, since adding vertices to T can only increase N(T), so by Harris' inequality we have

$$Z\sum_{T\subseteq A}\mu(T)\prod_{u\in B}f_u(T)\geq Z\prod_{u\in B}\left(\sum_{T\subseteq A}\mu(T)f_u(T)\right).$$

Here

$$\sum_{T \subseteq A} \mu(T) f_u(T) = \sum_{\substack{T \subseteq A \\ u \notin N(T)}} \mu(T) s + \sum_{\substack{T \subseteq A \\ u \notin N(T)}} \mu(T) s \left(1 - \frac{1}{d_u + 1}\right)$$
$$= s - \frac{s}{d_u + 1} \sum_{\substack{T \subseteq A \\ u \notin N(T)}} \mu(T) = s - \frac{s}{d_u + 1} \sum_{\substack{T \subseteq A \\ T \cap N(u) = \emptyset}} \mu(T)$$
$$= s - \frac{s}{d_u + 1} \prod_{v \in N(u)} \frac{\frac{d_v + x}{d_v + 1}s}{x(1 - s) + \frac{d_v + x}{d_v + 1}s} = s \left(1 - \frac{1}{d_u + 1} \prod_{v \in N(u)} \gamma_{x,s}(d_v)\right).$$

Hence

$$\widetilde{T}_H(x,0) \ge \prod_{v \in A} \left(\frac{(d_v + x)s}{d_v + 1} + x(1-s) \right) \cdot \prod_{u \in B} \left(s - \frac{s}{d_u + 1} \prod_{v \in N(u)} \gamma_{x,s}(d_v) \right).$$

By applying Theorem 4.1 for both $\widetilde{T}_H(x,0)$ and $\widetilde{T}_H(0,x)$ we get the following corollary. Corollary 4.2. We have

$$\widetilde{T}_H(x,0)\widetilde{T}_H(0,x) \ge \prod_{v \in V} \left[\left(\frac{(d_v + x)s}{d_v + 1} + x(1-s) \right) \left(s - \frac{s}{d_v + 1} \prod_{u \in N(v)} \gamma_{x,s}(d_u) \right) \right].$$

Lemma 4.3. If $x \ge 1$ and $0 \le s \le 1$, then the function $\gamma_{x,s}(d)$ is monotone decreasing in d. Furthermore, $\gamma_{x,s}(d) \le s$.

Proof. The function $\gamma_{x,s}(d)$ is a rational function in d:

$$\gamma_{x,s}(d) = \frac{(d+x)s}{(d+x)s + (d+1)x(1-s)} = \frac{sd+sx}{(x(1-s)+s)d+x}$$

In general, the derivative of a function $f(t) = \frac{at+b}{ct+d}$ is $\frac{ad-bc}{(ct+d)^2}$, so it is enough to check that

$$sx - sx(x(1-s) + s) = sx(1-x)(1-s) \le 0$$

which is clearly true. The second statement follows from $d + x \leq (d + 1)x$ as follows:

$$\gamma_{x,s}(d) = \frac{(d+x)s}{(d+x)s + (d+1)x(1-s)} \le \frac{(d+x)s}{(d+x)s + (d+x)(1-s)} = s.$$

Definition 4.4. Let

$$G(d, x, s, \gamma) := \left(\frac{(d+x)s}{d+1} + x(1-s)\right) \left(s - \frac{s}{d+1}\gamma^d\right).$$

For any $0 \leq \gamma < 1$ we have

$$G(\infty, x, s) := \lim_{d \to \infty} G(d, x, s, \gamma) = (s + x(1 - s))s.$$

By combining Corollary 4.2 with Lemma 4.6 we immediately get the following statement.

Corollary 4.5. Let H be a bipartite graph with minimum degree at least δ . Then for $x \geq 2$ and $0 \leq s \leq 1$ we have

$$\widetilde{T}_H(x,0)\widetilde{T}_H(0,x) \ge \prod_{v \in V} G(d_v, x, s, \gamma_{x,s}(\delta)) \ge \prod_{v \in V} G(d_v, x, s, s).$$

Proof. We have

$$\begin{split} \widetilde{T}_{H}(x,0)\widetilde{T}_{H}(0,x) &\geq \prod_{v \in V} \left[\left(\frac{(d_{v}+x)s}{d_{v}+1} + x(1-s) \right) \left(s - \frac{s}{d_{v}+1} \prod_{u \in N(v)} \gamma_{x,s}(d_{u}) \right) \right] \\ &\geq \prod_{v \in V} \left[\left(\frac{(d_{v}+x)s}{d_{v}+1} + x(1-s) \right) \left(s - \frac{s}{d_{v}+1} \prod_{u \in N(v)} \gamma_{x,s}(\delta) \right) \right] \\ &= \prod_{v \in V} G(d_{v},x,s,\gamma_{x,s}(\delta)) \\ &\geq \prod_{v \in V} G(d_{v},x,s,s). \end{split}$$

Lemma 4.6. Let $x \ge 2$ and $\gamma \le s < 1$. A sufficient condition for

$$G(d, x, s, \gamma) \ge G(d+1, x, s, \gamma)$$

 $is \left(\frac{x-1}{x(d+2)}\right)^{1/(d-1)} \ge \gamma.$ Furthermore, the function $\left(\frac{x-1}{x(d+2)}\right)^{1/(d-1)}$ is monotone increasing in d for $x \ge 1$ and $d \ge 2$, so it is enough to check the above inequality for some fixed d_0 to conclude that the sequence $G(d, x, s, \gamma)$ is decreasing for $d \ge d_0$.

$$\begin{aligned} Proof. \text{ We have} \\ G(d, x, s, \gamma) &- G(d+1, x, s, \gamma) \\ &= \left(\frac{(d+x)s}{d+1} + x(1-s)\right) \left(s - \frac{s}{d+1}\gamma^d\right) - \left(\frac{(d+1+x)s}{d+2} + x(1-s)\right) \left(s - \frac{s}{d+2}\gamma^{d+1}\right) \\ &= s^2 \left(\frac{d+x}{d+1} - \frac{d+1+x}{d+2}\right) - \frac{s}{d+1}\gamma^d \left(\frac{(d+x)s}{d+1} + x(1-s)\right) + \frac{s}{d+2}\gamma^{d+1} \left(\frac{(d+1+x)s}{d+2} + x(1-s)\right) \\ &\geq \frac{(x-1)s^2}{(d+1)(d+2)} - \frac{s}{d+1}\gamma^d \left(\frac{(d+x)s}{d+1} + x(1-s)\right) \\ &\geq \frac{(x-1)s^2}{(d+1)(d+2)} - \frac{sx}{d+1}\gamma^d \end{aligned}$$

which is non-negative if $\frac{x-1}{x(d+2)} \ge \gamma^{d-1}$. Finally, the derivative of $h(t) := \frac{1}{t-1} \ln \left(\frac{x-1}{x(t+2)} \right)$ is

$$h'(t) = \frac{1}{(t-1)^2} \ln\left(\frac{x(t+2)}{x-1}\right) - \frac{1}{(t-1)(t+2)}.$$

Then

if t > 1.

$$h'(t) \ge \frac{\ln(t+2)}{(t-1)^2} - \frac{1}{(t-1)(t+2)} \ge \frac{\ln(3)}{(t-1)^2} - \frac{1}{(t-1)(t+2)} > 0$$

5. Proof of Theorem 1.3.

In this section we prove that

$$T_M(x,0)T_M(0,x) \ge T_M(1,1)^2$$

whenever $x \ge 2.355$. By the transfer lemma it is enough to prove that

$$\widetilde{T}_H(x,0)\widetilde{T}_H(0,x) \ge \widetilde{T}_H(1,1)^2$$

for any bipartite graph H with minimum degree at least 1. Note that $\widetilde{T}_H(1,1) = 1$.

We do not immediately prove the best value 2.355, but give various improvements to the value 2.9243 step by step to digest one idea at a time. All computations are carried out in the jupyter notebook [7] and we also give the tables in the Appendix.

Idea 1. By Corollary 4.5 we have

$$\widetilde{T}_H(x,0)\widetilde{T}_H(0,x) \ge \prod_{v \in V} G(d_v, x, s, \gamma_{x,s}(1)).$$

Since $\widetilde{T}_H(1,1)^2 = 1$ it is enough if $G(d, x, s, \gamma_{x,s}(1)) \ge 1$ for all $d \ge 1$. Let

$$x = \frac{3 + \sqrt{5}}{2} \approx 2.6180.$$
 and $s = \frac{6 + 2\sqrt{5}}{7 + 3\sqrt{5}} \approx 0.7639.$

These numbers are chosen in such a way that $G(1, x, s, \gamma_{x,s}(1)) = 1$. It is easy to check that for d = 9 we have $\left(\frac{x-1}{x(d+2)}\right)^{1/(d-1)} \ge s \ge \gamma_{x,s}(1)$. By Lemma 4.6 this implies that for $d \geq 9$ we have $G(d, x, s, \gamma_{x,s}(1)) \geq G(\infty, x, s) > 1$. It is also easy to check (see the jupyter notebook or Table 1 in the Appendix) that $G(d, x, s, \gamma_{x,s}(1)) \ge 1$ for $d \in \{2, \ldots, 10\}$. So with $x = \frac{3+\sqrt{5}}{2}$ we have

$$\widetilde{T}_H(x,0)\widetilde{T}_H(0,x) \ge \widetilde{T}_H(1,1)^2$$
$$T_H(x,0)T_H(0,x) \ge T_H(1,1)^2$$

implying that

$$T_M(x,0)T_M(0,x) \ge T_M(1,1)^2$$

for all matroids without loops and coloops, in particular for all graphs G without loops and bridges.

Idea 2. Observe that we can assume that H is connected since $\widetilde{T}_H(x, y) = \prod_{i=1}^k \widetilde{T}_{H_i}(x, y)$ if H has connected components H_1, \ldots, H_k . The statement $\widetilde{T}_H(2, 0)\widetilde{T}(0, 2) \geq 1$ is trivially true for $H = K_2$, the complete graph on 2 vertices. So we can assume that H has at least 3 vertices. On the other hand, if H is connected and has at least 3 vertices, then a degree one vertex cannot be adjacent to another degree one vertex. Thus it is enough if

$$G(1, x, s, \gamma_{x,s}(2)) \ge 1$$
 and $G(d, x, s, \gamma_{x,s}(1)) \ge 1$

for all $d \ge 2$. This is satisfied if x = 2.54 and s = 0.76. (See Table 2.)

Idea 3. Observe that it is not really necessary that $G(1, x, s, \gamma) \ge 1$, it is quite enough if for a non-leaf vertex the product with its leaf neighbors is at least 1. If $x \ge 2.2$, then any vertex can have at most 2 leaf neighbors, since for the star S_4 on 4 vertices we have

$$\frac{1}{x}\widetilde{T}_{S_4}(x,0)\widetilde{T}_{S_4}(0,x) = \frac{1}{x} \cdot \frac{x^3 + x^2 + x}{4} \cdot \frac{x}{4} = \frac{x^3 + x^2 + x}{16} > 1$$

so by the gluing lemma (Lemma 2.14) we know that a graph containing a pending S_4 cannot be a minimal counter example. So it is enough to check if for every $d \ge 2$ we have $G(d, x, s, \gamma_{x,s}(1))G(1, x, s, \gamma_{x,s}(d))^{\min(2,d-1)} \ge 1$

and

$$G(d, x, s, \gamma_{x,s}(1)) \ge 1.$$

In order to check the first inequality for large d it is enough if for some d_0 we have

$$G(\infty, x, s)G(1, x, s, \gamma_{x,s}(d_0))^2 \ge 1,$$

then we get that

$$G(d, x, s, \gamma_{x,s}(1))G(1, x, s, \gamma_{x,s}(d))^{\min(2,d-1)} \ge 1$$

for $d \ge d_0$. So we only need to check the first inequality for $d = 2, \ldots, d_0$ and the inequality $G(\infty, x, s)G(1, x, s, \gamma_{x,s}(d_0))^2 \ge 1$. These inequalities are satisfied for x = 2.36, s = 0.78 and $d_0 = 44$. Again see the jupyter notebook [7] or alternatively, Table 3.

Idea 4. We can gain a very small further improvement as follows. For $d \ge 2$ we can use that

$$\left(\frac{(d_v+x)s}{d_v+1} + x(1-s)\right) \left(s - \frac{s}{d_v+1} \prod_{u \in N(v)} \gamma_{x,s}(d_u)\right)$$

$$\geq \left(\frac{(d_v+x)s}{d_v+1} + x(1-s)\right) \left(s - \frac{s}{d_v+1} \gamma_{x,s}(1)^{\min(2,d_v-1)} \gamma_{x,s}(2)^{d_v-\min(2,d_v-1)}\right),$$

Let

$$G_2(d, x, s, \gamma_1, \gamma_2) := \left(\frac{(d+x)s}{d+1} + x(1-s)\right) \left(s - \frac{s}{d+1}\gamma_1^{\min(2,d-1)}\gamma_2^{d-\min(2,d-1)}\right).$$

Then it is enough to check that

$$G_2(d, x, s, \gamma_{x,s}(1), \gamma_{x,s}(2)) \ge 1$$

for $d \geq 2$ and

$$G_2(d, x, s, \gamma_{x,s}(1), \gamma_{x,s}(2))G(d, x, s, \gamma_{x,s}(1))^{\min(2,d-1)} \ge 1$$

for all $d \ge 2$. This leads to the tiny improvement x = 2.355 (s = 0.78 and d = 100). See [7] or Table 4 in the Appendix for further details.

6. MATROIDS WITH PRESCRIBED CIRCUIT LENGTHS

In this section, we prove Theorem 1.4. The heart of the argument is the following lemma.

Lemma 6.1. Let $k \ge 4$ be a real number. Let H be a bipartite graph such that all degrees of H lie in the interval $[k + 1, k^4 - 2k^2 - 1]$. Then

$$\widetilde{T}_H(2,0)\widetilde{T}(0,2) \ge \widetilde{T}(1,1)^2.$$

Proof. We will use that $\widetilde{T}_H(1,1) = 1$, and Corollary 4.5:

$$\widetilde{T}_H(2,0)\widetilde{T}(0,2) \ge \prod_{v \in V} G(d_v,2,s,s).$$

Note that

$$G(d, 2, s, s) = \left(\frac{d+2}{d+1} \cdot s + 2(1-s)\right) \left(s - \frac{s^{d+1}}{d+1}\right) = \left(\frac{d+2}{d+1} + \frac{d}{d+1}(1-s)\right) \left(s - \frac{s^{d+1}}{d+1}\right)$$

We will show that if $s = 1 - \frac{1}{k^2}$, then for all $d \in [k+1, k^4 - 2k^2 - 1]$ we have $G(d, 2, s, s) \ge 1$. First observe that G(d, 2, s, s) is a concave function of s on the interval (0, 1) as its second derivative is

$$-\frac{(2-s)(d^2+d)s^d+2ds}{(d+1)s},$$

which is clearly negative. This means that $I_d := \{s \mid G(d, 2, s, s) \ge 1\}$ is an interval for each d. We will show that $1 - \frac{1}{k^2} \in I_{k+1} \cap I_{k^4 - 2k^2 - 1}$. By first choosing k = d - 1 we get that $1 - \frac{1}{(d-1)^2} \in I_d$. If we choose $k^2 = 1 + \sqrt{d+2}$, then $k^4 - 2k^2 - 1 = d$ and so $1 - \frac{1}{1 + \sqrt{d+2}} \in I_d$. Hence $\left[1 - \frac{1}{1 + \sqrt{d+2}}, 1 - \frac{1}{(d-1)^2}\right] \subseteq I_d$. This means that if $k + 1 \le d \le k^4 - 2k^2 - 1$, then $1 - \frac{1}{k^2} \in I_d$.

First we show that $1 - \frac{1}{k^2} \in I_{k+1}$, that is, $G(k+1, 2, 1 - \frac{1}{k^2}, 1 - \frac{1}{k^2}) \ge 1$. We will use that

$$(1-t)^r \le 1 - \binom{r}{1}t + \binom{r}{2}t^2$$

for $0 \leq t \leq 1$ by Bonferroni's inequality applied to independent events with probability t. Hence

$$G\left(d,2,1-\frac{1}{k^2},1-\frac{1}{k^2}\right) = \left(\frac{d+2}{d+1} + \frac{d}{d+1} \cdot \frac{1}{k^2}\right) \left(1 - \frac{1}{k^2} - \frac{1}{d+1}\left(1 - \frac{1}{k^2}\right)^{d+1}\right)$$
$$\geq \left(\frac{d+2}{d+1} + \frac{d}{d+1} \cdot \frac{1}{k^2}\right) \left(1 - \frac{1}{k^2} - \frac{1}{d+1}\left(1 - \frac{d+1}{k^2} + \frac{(d+1)d}{2k^2}\right)\right)$$

For d = k + 1 we get that

$$G\left(k+1,2,1-\frac{1}{k^2},1-\frac{1}{k^2}\right) \ge \left(\frac{k+3}{k+2} + \frac{k+1}{k+2} \cdot \frac{1}{k^2}\right) \left(1 - \frac{1}{k^2} - \frac{1}{k+2} \left(1 - \frac{k+2}{k^2} + \frac{(k+2)(k+1)}{2k^2}\right)\right)$$
$$= 1 + \frac{1}{2} \frac{3k^5 - 4k^4 - 12k^3 - 10k^2 - 5k - 2}{k^8 + 4k^7 + 4k^6}$$

For $k \ge 4$ we have $3k^5 - 4k^4 - 12k^3 - 10k^2 - 5k - 2 > 0$ showing that $G\left(k + 1, 2, 1 - \frac{1}{k^2}, 1 - \frac{1}{k^2}\right) > 1$.

Next we show that $1 - \frac{1}{k^2} \in I_{k^4 - 2k^2 - 1}$. For $s = 1 - \frac{1}{k^2}$ and $d = k^4 - 2k^2 - 1$ we have

$$\left(\frac{d+2}{d+1} + \frac{d}{d+1}(1-s)\right)s = 1 + \frac{1}{k^8 - 2k^6}$$

and

$$\left(\frac{d+2}{d+1} + \frac{d}{d+1}(1-s)\right)\frac{s^{d+1}}{d+1} = \frac{k^6 - k^4 - k^2 - 1}{k^6 - 2k^4} \cdot \frac{1}{k^4 - 2k^2} \left(1 - \frac{1}{k^2}\right)^{k^4 - 2k^2}.$$

So we need to prove that

$$\frac{1}{k^8 - 2k^6} > \frac{k^6 - k^4 - k^2 - 1}{k^6 - 2k^4} \cdot \frac{1}{k^4 - 2k^2} \left(1 - \frac{1}{k^2}\right)^{k^4 - 2k^2}$$

which is equivalent with

$$\frac{k^2 - 2}{k^6 - k^4 - k^2 - 1} > \left(1 - \frac{1}{k^2}\right)^{k^4 - 2k^2}$$

By $1 - t < e^{-t}$ we have

$$\left(1 - \frac{1}{k^2}\right)^{k^4 - 2k^2} < e^{-\frac{1}{k^2}(k^4 - 2k^2)} = e^{-k^2 - 2}.$$

Let us introduce $y = k^2 - 2$. Then $k^6 - k^4 - k^2 - 1 = y^3 + 5y^2 + 7y + 1$, so we need that

$$\frac{y}{y^3 + 5y^2 + 7y + 1} < e^{-1}$$

or equivalently

$$y^2 + 5y + 7 + \frac{1}{y} < e^y.$$

Since $k \ge 3$ we have $y = k^2 - 2 \ge 7$, but $e^y > y^2 + 5y + 7 + \frac{1}{y}$ already true for $y \ge 4$.

Remark 6.2. There was no attempt to optimize Lemma 6.1, not even concerning the minimum degree. It turns out that if s = 0.9226, then

$$G(d, 2, s, \gamma_{2,s}(3)) > 1$$

for $3 \le d \le 141$. So if all degrees of a bipartite graph H lies in the interval [3, 141], then

$$\widetilde{T}_H(2,0)\widetilde{T}(0,2) \ge \widetilde{T}(1,1)^2.$$

Similarly, for s = 0.9622 we have

$$G(d, 2, s, \gamma_{2,s}(4)) > 1$$

for $4 \le d \le 646$. So if all degrees of a bipartite graph H lies in the interval [4, 646], then

$$\widetilde{T}_H(2,0)\widetilde{T}(0,2) \ge \widetilde{T}(1,1)^2.$$

This shows that in Theorem 1.4 we could have written $\ell \geq 4$ and probably a faster growing function than $(\ell - 2)^4$.

Now we are ready to finish the proof of Theorem 1.4.

Proof of Theorem 1.4. Let A be a basis of the matroid M on ground set E. Let $B = E \setminus A$. Let us consider the local basis exchange graph H[A] on $A \cup B$. Observe that $v \in B$ are adjacent with exactly those elements that are in his fundamental circuit. So $\ell \leq d_v + 1 \leq (\ell - 2)^4$. Similarly, if $u \in A$, then it is adjacent with those elements of B that are in its fundamental circuits with respect to B in M^* . So $\ell \leq d_u + 1 \leq (\ell - 2)^4$.

By choosing $k = \ell - 1$ we get that $k^4 - 2k^2 - 1 = (\ell - 1)^4 - 2(\ell - 1)^2 - 1 \ge (\ell - 2)^4$ if $\ell \ge 3$. So by Lemma 1.4 we have

$$\widetilde{T}_{H[A]}(2,0)\widetilde{T}_{H[A]}(0,2) \ge \widetilde{T}_{H[A]}(1,1)^2.$$

Since this is true for all basis A of the matroid M, we get by the Transfer lemma (Lemma 2.6 and Remark 2.7) that

$$T_M(2,0)T_M(0,2) \ge T_M(1,1)^2.$$

7. Concluding Remarks

We end this paper with some conjectures.

Conjecture 7.1. If the bipartite graph H has minimum degree at least 2, then

$$\widetilde{T}_H(2,0)\widetilde{T}_H(0,2) \ge \widetilde{T}_H(1,1)^2.$$

If this conjecture is true, then the following conjecture would be an immediate corollary.

Conjecture 7.2. If M is a simple co-simple matroid, then

$$T_M(2,0)T_M(0,2) \ge T_M(1,1)^2.$$

Acknowledgment. The author is very grateful to Csongor Beke, Gergely Kál Csáji and Sára Pituk for discussions on the topic of this paper.

References

- Csongor Beke, Gergely Kál Csáji, Péter Csikvári, and Sára Pituk. Permutation Tutte polynomial. European Journal of Combinatorics, 120:104003, 2024.
- [2] Csongor Beke, Gergely Kál Csáji, Péter Csikvári, and Sára Pituk. The Merino–Welsh conjecture is false for matroids. Advances in Mathematics, 446:109674, 2024.
- [3] Thomas Brylawski and James Oxley. The Tutte polynomial and its applications. Matroid applications, 40:123–225, 1992.
- [4] Laura E. Chávez-Lomelí, Criel Merino, Steven D. Noble, and Marcelino Ramírez-Ibáñez. Some inequalities for the Tutte polynomial. European Journal of Combinatorics, 32(3):422–433, 2011.
- [5] Rodolfo Conde and Criel Merino. Comparing the number of acyclic and totally cyclic orientations with that of spanning trees of a graph. Int. J. Math. Com, 2:79–89, 2009.
- [6] Henry H. Crapo. The Tutte polynomial. Aequationes Mathematicae, 3(3):211–229, 1969.
- [7] Péter Csikvári. Jupyter Notebook 'Merino-Welsh optimization'. https://github.com/Merino-Welsh-conjecture/blob/main/Merino-Welsh_optimization.ipynb, 2025.
- [8] Joanna A. Ellis-Monaghan and Criel Merino. Graph polynomials and their applications i: The Tutte polynomial. In *Structural analysis of complex networks*, pages 219–255. Springer, 2011.
- Joanna A. Ellis-Monaghan and Iain Moffatt. Handbook of the Tutte polynomial and related topics. CRC Press, 2022.
- [10] Luis Ferroni and Benjamin Schröter. The Merino–Welsh conjecture for split matroids. Annals of Combinatorics, 27(3):737–748, 2023.
- [11] Cees M. Fortuin, Pieter W. Kasteleyn, and Jean Ginibre. Correlation inequalities on some partially ordered sets. *Communications in Mathematical Physics*, 22(2):89–103, 1971.

P. CSIKVÁRI

- [12] Theodore E Harris. A lower bound for the critical probability in a certain percolation process. In Mathematical Proceedings of the Cambridge Philosophical Society, volume 56, pages 13–20. Cambridge University Press, 1960.
- [13] Bill Jackson. An inequality for Tutte polynomials. Combinatorica, 30:69-81, 2010.
- [14] Kolja Knauer, Leonardo Martínez-Sandoval, and Jorge Luis Ramírez Alfonsín. A Tutte polynomial inequality for lattice path matroids. Advances in Applied Mathematics, 94:23–38, 2018.
- [15] Joseph P. S. Kung. Inconsequential results on the Merino-Welsh conjecture for Tutte polynomials. arXiv preprint arXiv:2105.01825, 2021.
- [16] Fenggen Lin. A note on spanning trees and totally cyclic orientations of 3-connected graphs. Journal of Combinatorics, 4(1):95–104, 2013.
- [17] Criel Merino, Marcelino Ibañez, and M Guadalupe Rodríguez. A note on some inequalities for the Tutte polynomial of a matroid. *Electronic Notes in Discrete Mathematics*, 34:603–607, 2009.
- [18] Criel Merino and Dominic Welsh. Forests, colorings and acyclic orientations of the square lattice. Annals of Combinatorics, 3(2-4):417–429, 1999.
- [19] Steven D. Noble and Gordon F. Royle. The Merino–Welsh conjecture holds for series–parallel graphs. European Journal of Combinatorics, 38:24–35, 2014.
- [20] James Oxley. Matroid theory. Oxford University Press, 1992.
- [21] Carsten Thomassen. Spanning trees and orientations of graphs. Journal of Combinatorics, 1(2):101– 111, 2010.
- [22] William Thomas Tutte. A contribution to the theory of chromatic polynomials. Canadian Journal of mathematics, 6:80–91, 1954.
- [23] Dominic Welsh. The Tutte polynomial. Random Structures & Algorithms, 15(3-4):210–228, 1999.

8. Appendix

Below, one can find the tables belonging to the various ideas.

8.1. Table for Idea 1. Let $x := \frac{3+\sqrt{5}}{2} \approx 2.6180$ and $s := \frac{6+2\sqrt{5}}{7+3\sqrt{5}} \approx 0.7639$. Below we give the values of $G(d, x, s, \gamma_{x,s}(1))$.

d	$G(d, x, s, \gamma_{x,s}(1))$	
1	1.0000000000000000000000000000000000000	
2	1.15236921034711	
3	1.18525033351524	
4	1.18783805465536	
5	1.18125856327950	
6	1.17211090590244	
7	1.16272735027462	
8	1.15394531114347	
9	1.14602516001225	
10	1.13899595137302	
11	1.13279660374225	
∞	1.05572809000084	
TABLE 1.		

For d = 9 we have $\left(\frac{x-1}{x(d+2)}\right)^{1/(d-1)} \approx 0.6977$. This is larger than $\gamma_{x,s}(1) \approx 0.6909$. This means that the sequence $G(d, x, s, \gamma_{x,s}(1))$ is decreasing from d = 9, so the smallest value of $G(d, x, s, \gamma_{x,s}(1))$ from that point on is $G(\infty, x, s)$. (As we can see it is actually a decreasing sequence already from d = 4.)

8.2. Table for Idea 2. Let x = 2.54 and s = 0.76.

d	$G(d, x, s, \gamma_{x,s}(1))$
1*	1.00015021063798
2	1.12628760116317
3	1.16035420716839
4	1.16413305093218
5	1.15856178434317
6	1.15024896467994
7	1.14155987842924
8	1.13336130037553
9	1.12593636935915
10	1.11933141925454
11	1.11349857234605
∞	1.04089600000000

TABLE 2. The value at d = 1 is not $G(1, x, s, \gamma_{x,s}(1))$ but $G(1, x, s, \gamma_{x,s}(2))$.

Again we can check that

$$\left(\frac{x-1}{x(d+2)}\right)^{1/(d-1)} > \gamma_{x,s}(1)$$

for $d \ge 9$ so the sequence $G(d, x, s, \gamma_{x,s}(1))$ is monotone decreasing from that point on. This gives that

$$G(d, x, s, \gamma_{x,s}(1)) \ge G(\infty, x, s) > 1$$

for all $d \ge 9$. Since $G(d, x, s, \gamma_{x,s}(1)) > 1$ for $2 \le d \le 8$ we get this inequality for all $d \ge 2$.

8.3. Table for Idea 3. Let x = 2.36 and s = 0.78.

d	$G(d, x, s, \gamma_{x,s}(1))$	$G(d, x, s, \gamma_{x,s}(1))G(1, x, s, \gamma_{x,s}(d))^{\min(2,d-1)}$
2	1.06874465202436	1.00215345922882
3	1.10815264651986	1.00086197145259
4	1.11681913369317	1.02640833194772
5	1.11511667337790	1.03743759976094
6	1.10975849131510	1.04182882517377
7	1.10330750172561	1.04300865188036
8	1.09680798068072	1.04261001534334
9	1.09068230467836	1.04145583550358
10	1.08508115163529	1.03997408296620
11	1.08003342261565	1.03838980790695
12	1.07551442455619	1.03682008545765
43	1.03218107718904	1.01881210598816
44	1.03176319039565	1.01863210924050
∞	1.01337600000000	1.00047892960579

TABLE 3.

8.4. Table for Idea 4. Let $x = 2.355$ and $s = 0.78$	3.
---	----

d	$G_2(d, x, s, \gamma_{x,s}(1), \gamma_{x,s}(2))$	$G_2(d, x, s, \gamma_{x,s}(1), \gamma_{x,s}(2))G(1, x, s, \gamma_{x,s}(d))^{\min(2,d-1)}$
2	1.07641984643180	1.00750701821492
3	1.11135556808369	1.00001551323253
4	1.12051595966294	1.02591566665079
5	1.11815485337910	1.03629473774471
6	1.11192862150501	1.03985270835852
7	1.10468853332594	1.04027529602599
8	1.09755066213067	1.03926180701977
9	1.09093594198563	1.03763439690775
10	1.08497152313990	1.03579937829855
11	1.07965934020252	1.03395657259571
12	1.07495066897719	1.03220035567236
98	1.02084509090909	1.00934521974652
99	1.02076182000000	1.00930958644862
∞	1.01251800000000	1.00115825634209

Table	4.
-------	----

HUN-REN Alfréd Rényi Institute of Mathematics, H-1053 Budapest Reáltanoda utca 13-15 and ELTE: Eötvös Loránd University Mathematics Institute, Department of Computer Science H-1117 Budapest, Pázmány Péter sétány $1/{\rm C}$

 $Email \ address: \ {\tt peter.csikvari@gmail.com}$