

Wormhole geometries in $f(R, T^2)$ gravity satisfying the energy conditions

Nailya Ganiyeva

*Instituto de Astrofísica e Ciências do Espaço, Faculdade de Ciências da Universidade de Lisboa,
Edifício C8, Campo Grande, P-1749-016 Lisbon, Portugal*
*E-mail: fc57452@alunos.fc.ul.pt

João Luís Rosa

University of Gdańsk, Jana Bażyńskiego 8, 80-309 Gdańsk, Poland
Institute of Physics, University of Tartu, W. Ostwaldi 1, 50411 Tartu, Estonia
*E-mail: joaoluis92@gmail.com

Francisco S. N. Lobo

*Instituto de Astrofísica e Ciências do Espaço, Faculdade de Ciências da Universidade de Lisboa,
Edifício C8, Campo Grande, P-1749-016 Lisbon, Portugal*
*Departamento de Física, Faculdade de Ciências da Universidade de Lisboa, Edifício C8, Campo
Grande, P-1749-016 Lisbon, Portugal*
*E-mail: fslobo@fc.ul.pt

We explore the properties of traversable wormhole spacetimes within the framework of energy-momentum squared gravity, also known as $f(R, T^2)$ gravity, where R represents the Ricci scalar, T_{ab} is the energy-momentum tensor, and $T^2 = T_{ab}T^{ab}$. Adopting a linear functional form $f(R, T^2) = R + \gamma T^2$, we demonstrate the existence of a wide range of wormhole solutions that satisfy all of the energy conditions without requiring fine-tuning of the model parameters. Due to the inherent complexity of the field equations, these solutions are constructed through an analytical recursive method. However, they lack a natural localization, requiring a junction with an external vacuum region. To address this, we derive the corresponding junction conditions and establish that the matching must always be smooth, precluding the formation of thin shells at the interface. Using these conditions, we match the interior wormhole geometry to an exterior Schwarzschild solution, yielding localized, static, and spherically symmetric wormholes that satisfy all the energy conditions throughout the entire spacetime. Finally, we extend our analysis to more intricate dependencies on T^2 , demonstrating that the methodology remains applicable as long as no mixed terms between R and T^2 are introduced.

Keywords: Wormholes, modified theories of gravity, junction conditions.

1. Introduction

Wormholes are theoretical structures that connect two separate spacetime manifolds or distinct regions within the same manifold. These objects have been extensively studied in the context of General Relativity (GR)¹⁻⁶, where it is found that traversable wormholes must satisfy the flaring-out condition¹. Combining the requirement with the Einstein field equations results in the violation of the null energy condition (NEC) and, consequently, all other energy conditions^{3,7,8}. Matter violating the NEC, termed as *exotic* matter, has limited physical relevance since it

lacks experimental evidence for its existence.

One way to avoid the use of exotic matter to sustain wormhole geometries is to examine them within the framework of modified gravity theories^{9–23}. In these theories, additional curvature terms contribute to sustain the geometry of the wormhole throat, ensuring its traversability, while allowing the matter content to remain non-exotic. This result can be achieved across various extensions of GR, ranging from $f(R)$ gravity and its generalizations^{24–30}, to models involving curvature-matter couplings^{31,32}, theories incorporating additional fundamental fields^{33,34}, Einstein–Cartan gravity³⁵, Gauss–Bonnet gravity^{36–38}, and brane-world scenarios^{39,40}. Regarding $f(R, T^2)$ gravity, specific solutions were first obtained using a Noether symmetry approach. However, these solutions lacked physical significance as they violated the NEC^{9,10}.

More recently, physically viable traversable wormhole spacetimes that satisfy all energy conditions have been found¹¹, and these are the solutions presented in this manuscript. More specifically, we analyze traversable wormholes in $f(R, T^2)$ gravity with $f(R, T^2) = R + \gamma T^2$. A broad class of solutions satisfying all energy conditions is found via an analytical recursive method. To ensure localization, we derive junction conditions, proving smooth matching to an external Schwarzschild solution. The approach extends to more general T^2 dependencies, provided no mixed terms with R appear.

2. Theory and framework

2.1. Action and field equation of $f(R, T^2)$

In this work, we investigate wormholes within the framework of the $f(R, T^2)$ theory of gravity. The action S governing this theory is given by

$$S = \frac{1}{2\kappa^2} \int_{\Omega} \sqrt{-g} f(R, T^2) d^4x + \int_{\Omega} \sqrt{-g} \mathcal{L}_m d^4x, \quad (1)$$

where $\kappa^2 = 8\pi$, where we adopt a geometrized unit system with $c = G = 1$. Here, Ω represents a spacetime manifold, and g is the determinant of the metric tensor g_{ab} . The function $f(R, T^2)$ is an arbitrary function of the Ricci scalar $R = g^{ab} R_{ab}$, where R_{ab} is the Ricci tensor, and the scalar $T^2 = T_{ab} T^{ab}$, where T_{ab} denotes the energy-momentum tensor. The term \mathcal{L}_m corresponds to the matter Lagrangian, and the energy-momentum tensor is defined as usual by

$$T_{ab} = -\frac{2}{\sqrt{-g}} \frac{\delta(\sqrt{-g} \mathcal{L}_m)}{\delta g^{ab}}. \quad (2)$$

Varying the action (1) with respect to the metric g_{ab} , we obtain the modified gravitational field equations

$$f_R R_{ab} - \frac{1}{2} g_{ab} f - (\nabla_a \nabla_b - g_{ab} \square) f_R = 8\pi T_{ab} - f_{T^2} \Theta_{ab}, \quad (3)$$

where we use the shorthand notation $f_R \equiv \partial f / \partial R$ and $f_{T^2} = \partial f / \partial T^2$. Here, ∇_a denotes the covariant derivative, and $\square = g^{ab} \nabla_a \nabla_b$ is the d'Alembert operator, both

defined with respect to the metric g_{ab} . The term Θ_{ab} arises from the variation of T_{ab} given by

$$\Theta_{ab} = \frac{\delta T^2}{\delta g^{ab}}. \quad (4)$$

Choosing a specific matter Lagrangian \mathcal{L}_m or, equivalently, a particular form of the energy-momentum tensor T_{ab} , the explicit expression for the auxiliary tensor Θ_{ab} is determined. Taking the covariant derivative of Eq. (3) then yields the conservation equation

$$8\pi\nabla_b T^{ab} = \nabla_b (f_{T^2}\Theta^{ab}) + f_R\nabla_b R^{ab} - \frac{1}{2}g^{ab}\nabla_b f. \quad (5)$$

This equation reveals that, in this theory, the energy-momentum tensor T_{ab} is not necessarily conserved, meaning that in general, $\nabla_b T^{ab} \neq 0$. This feature distinguishes it from GR.

In this work, we assume that the function $f(R, T^2)$ is both separable and linear in terms of R and T^2 , that is

$$f(R, T^2) = R + \gamma T^2, \quad (6)$$

where γ represents a coupling constant. Thus, the field equations (3) and the conservation equation (5) take the following form:

$$G_{ab} = 8\pi T_{ab} - \gamma \left(\Theta_{ab} - \frac{1}{2}g_{ab}T^2 \right), \quad (7)$$

$$8\pi\nabla_b T^{ab} = \gamma\nabla_b \left(\Theta^{ab} - \frac{1}{2}g^{ab}T^2 \right), \quad (8)$$

where we have introduced the Einstein tensor $G_{ab} = R_{ab} - \frac{1}{2}Rg_{ab}$. Possible generalizations of these assumptions are discussed later in this manuscript.

2.2. Wormhole geometry and matter distribution

We begin by considering a static, spherically symmetric traversable wormhole metric in spherical coordinates (t, r, θ, φ) , expressed as

$$ds^2 = -e^{\zeta(r)}dt^2 + \left(1 - \frac{b(r)}{r}\right)^{-1}dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2), \quad (9)$$

where $\zeta(r)$ is the redshift function, $b(r)$ is the shape function. To ensure that the wormhole is traversable, the redshift function must be finite throughout the entire spacetime, i.e., $|\zeta(r)| < \infty$, to prevent event horizons. Additionally, the flaring-out condition at the wormhole throat $r = r_0$ is required^a, resulting in the following two boundary conditions: $b(r_0) = r_0$, and $b'(r_0) < 1$.

^aThe flaring-out condition close to the throat takes the form of $(b - b')/b^2 > 0^1$.

Upon these requirements, we consider two general families of solutions for the functions $\zeta(r)$ and $b(r)$, given by

$$\zeta(r) = \zeta_0 \left(\frac{r_0}{r}\right)^\alpha, \quad b(r) = r_0 \left(\frac{r_0}{r}\right)^\beta, \quad (10)$$

where ζ_0 is an arbitrary constant to be specified, and α and β are arbitrary positive exponents.

For the matter sector, we assume that the matter distribution is described by an anisotropic perfect fluid. Therefore, the energy-momentum tensor T_{ab} takes the form

$$T_a^b = \text{diag}(-\rho, p_r, p_t, p_t), \quad (11)$$

where $\rho \equiv \rho(r)$ is the energy density, $p_r \equiv p_r(r)$ represents the radial pressure, and $p_t \equiv p_t(r)$ denotes the tangential pressure.

Based on these assumptions, and with the choice in Eq.(10), the field equation (7) leads to three independent components, which are given by the following expressions

$$8\pi\rho = \frac{\gamma}{6} (p_r^2 - 2p_t^2 - 3\rho^2 - 8p_r p_t - 8p_r \rho - 16p_t \rho) - \frac{\beta}{r^2} \left(\frac{r_0}{r}\right)^{\beta+1}, \quad (12)$$

$$8\pi p_r = \frac{\gamma}{6} (p_r^2 + 2p_t^2 - 3\rho^2 - 12p_r p_t + 4p_r \rho - 4p_t \rho) - \frac{1}{r^2} \left(\frac{r_0}{r}\right)^{\beta+1} - \frac{\alpha\zeta_0}{r^2} \left(\frac{r_0}{r}\right)^\alpha \left[1 - \left(\frac{r_0}{r}\right)^{\beta+1}\right], \quad (13)$$

$$8\pi p_t = -\frac{\gamma}{6} (p_r^2 + 6p_t^2 + 3\rho^2 + 2p_r p_t + 2p_r \rho - 2p_t \rho) + \frac{1+\beta}{2r^2} \left(\frac{r_0}{r}\right)^{\beta+1} + \frac{\alpha^2\zeta_0^2}{4r^2} \left(\frac{r_0}{r}\right)^{2\alpha} \left[1 - \left(\frac{r_0}{r}\right)^{\beta+1}\right] + \frac{\alpha\zeta_0}{4r^2} \left(\frac{r_0}{r}\right)^\alpha \left[2\alpha - (1+2\alpha+\beta) \left(\frac{r_0}{r}\right)^{\beta+1}\right]. \quad (14)$$

The Eqs. (12)–(14) constitute a system of three equations with three unknowns: ρ , p_r , and p_t . Each of these equations is quadratic in its respective unknown, which suggests that the system could have up to eight independent solutions. However, the nature of these solutions may vary, with some solutions potentially being complex, depending on the particular values of the involved parameters.

2.3. Wormhole solutions

Due to the complexity of the system of Eqs.(12)–(14), obtaining explicit analytical solutions for ρ , p_r , and p_t is not feasible. However, using a recursive approach, we can determine these quantities by starting with specific values for the free parameters and solving the system at the initial radius r_0 . From there, we incrementally calculate ρ , p_r , and p_t at larger radii, up to a sufficient large radius r . This method also works

for any $f(R, T^2)$ function with higher powers of T^2 and no cross terms of RT^2 , although the set of solutions is larger.

Among the solutions obtained, we focus on those that are astrophysically relevant, meaning their matter components satisfy the energy conditions. For a diagonal energy-momentum tensor T_{ab} as described in Eq. (11), these energy conditions are expressed as follows

$$\rho + p_r \geq 0, \quad \rho + p_t \geq 0, \quad \rho \geq 0, \quad \rho + p_r + 2p_t \geq 0, \quad \rho \geq |p_r|, \quad \rho \geq |p_t|. \quad (15)$$

From the set of eight solutions for the matter quantities, any solutions that violate any of these energy conditions are excluded, and only those that satisfy all the energy conditions are studied.

As a specific example, we consider the parameter combination $\alpha = \beta = -\gamma = 1$, $r_0 = 3M$, and $\zeta_0 = -\frac{6}{5}$.^b The matter density ρ , along with the combinations $\rho + p_r$, $\rho + p_t$, $\rho + p_r + 2p_t$, $\rho - |p_r|$, and $\rho - |p_t|$, are plotted in Figure 1 for the solution that satisfies all the energy conditions discussed above.

As in linear $f(R, T)$ gravity³⁰, our analysis of solutions satisfying all the energy conditions in linear $f(R, T^2)$ confirms that they only exist for negative values of γ . However, for higher-order terms, $f(R, T^2) = R + \gamma T^2 + \sigma(T^2)^n$, such solutions can exist for positive values of γ , provided that σ remains negative. In both cases, the matter components remain nonzero across all r , despite asymptotic flatness, as seen in Figure 1, thus necessitating a matching with an exterior vacuum spacetime at a finite radius. This issue is addressed in the following section.

3. Junction Conditions and Matching

To obtain physically meaningful spacetime solutions describing localized objects, junction conditions must be applied to match the interior and exterior spacetimes at a finite radius. First derived in GR⁴¹, these conditions have been widely used in studies of traversable wormholes⁴²⁻⁴⁷, fluid stars⁴⁸⁻⁵⁰, and gravitational collapse^{51,52}. Since these conditions depend on the underlying theory, various works have explored them in modified gravity (see⁵³ for a review), including $f(R)$ gravity⁵⁴⁻⁶⁰, theories with additional fundamental fields⁶¹⁻⁶⁴, and metric-affine gravity⁶⁵⁻⁶⁷. More recently, junction conditions have been studied in $f(R, T^2)$ gravity¹¹, which we focus on here by deriving the conditions and performing the matching between the interior wormhole with an exterior vacuum solution.

3.1. Junction conditions of linear $f(R, T^2)$ gravity

We first derive the junction conditions for the linear case, $f(R, T^2) = R + \gamma T^2$, then briefly extend the analysis to higher-order terms, $f(R, T^2) = R + \gamma T^2 + \sigma(T^2)^n$.

^bWhile the choice of ζ_0 is somewhat arbitrary at this stage, this particular value is selected for reasons explained in the following section. Other values of ζ_0 , including positive ones, would lead to qualitatively similar solutions.




Fig. 1. Matter density ρ , along with the expressions $\rho + p_r$, $\rho + p_t$, $\rho + p_r + 2p_t$, $\rho - |p_r|$, and $\rho - |p_t|$, are shown for the parameter set $\alpha = \beta = -\gamma = 1$, $r_0 = 3M$, and $\zeta_0 = -\frac{6}{5}$.

Let us consider a spacetime manifold Ω divided into two regions Ω^\pm , each with metric tensors g_{ab}^\pm in coordinates x_\pm^a . The interior and exterior spacetimes are Ω^- and Ω^+ , respectively, separated by a three-dimensional hypersurface Σ with induced metric $h_{\alpha\beta}$, expressed in coordinates y^α , where greek indices exclude the directional normal to Σ . The projection tensors mapping Ω onto Σ are $e_\alpha^a = \partial x^a / \partial y^\alpha$. The normal vector to Σ is $n_a = \epsilon \partial_a l$, where l is an affine parameter along geodesics normal to Σ , and $\epsilon = 1, -1, 0$ for space-like, time-like, and null geodesic congruences, respectively. By construction, the condition $n^a e_a^\alpha = 0$ is always satisfied. With this notation, the induced metric $h_{\alpha\beta}$ and extrinsic curvature $K_{\alpha\beta}$ of Σ are given by

$$h_{\alpha\beta} = e_\alpha^a e_\beta^b g_{ab}, \quad K_{\alpha\beta} = e_\alpha^a e_\beta^b \nabla_a n_b. \quad (16)$$

The junction conditions are obtained using the distribution formalism, where any quantity X and its derivative $\nabla_a X$ are expressed through distribution functions as

$$X = X^+ \theta(l) + X^- \theta(-l), \quad (17)$$

$$\nabla_a X = \nabla_a X^+ \theta(l) + \nabla_a X^- \theta(-l) + \epsilon n_a [X] \delta(l), \quad (18)$$

where X^\pm denotes X in the respective spacetimes Ω^\pm ; $\theta(l)$ is the Heaviside distribution function, defined as $\theta(l) = 0$ for $l < 0$, $\theta(l) = 1$ for $l > 0$, and $\theta(l) = \frac{1}{2}$ for $l = 0$; and $\delta(l) = \partial_l \theta(l)$ is the Dirac delta distribution. The discontinuity of X across Σ is denoted by $[X]$, defined as

$$[X] = X^+|_\Sigma - X^-|_\Sigma. \quad (19)$$

If the quantity X is continuous across Σ , then $[X] = 0$. By definition, we have $[n^a] = [e_\alpha^a] = 0$.

To derive the junction conditions, we reformulate the field equations (7) within the distribution formalism. Starting with the metric tensor g_{ab} , it is expressed in this framework as

$$g_{ab} = g_{ab}^+ \theta(l) + g_{ab}^- \theta(-l). \quad (20)$$

From the metric expression in Eq.(20), we now compute the Christoffel symbols Γ_{ab}^c associated with the metric g_{ab} . This involves computing the derivatives $\partial_c g_{ab}$, which, using Eq.(18), take the form

$$\partial_c g_{ab} = \partial_c g_{ab}^+ \theta(l) + \partial_c g_{ab}^- \theta(-l) + \epsilon n_c [g_{ab}] \delta(l). \quad (21)$$

The term proportional to $\delta(l)$ poses a challenge in defining the Riemann tensor R_{bcd}^a within the distributional formalism. Since the Riemann tensor R_{bcd}^a involves products of Christoffel symbols Γ_{ab}^c , it results in terms proportional to $\delta^2(l)$, which are singular in this formalism. The junction conditions are introduced precisely to eliminate such singularities from the field equations. To ensure the removal of these singularities in the Riemann tensor R_{bcd}^a , we impose the continuity of the metric g_{ab} across Σ , requiring $[g_{ab}] = 0$. Given that $[e_\alpha^a] = 0$, this condition can be expressed in a coordinate-independent form by projecting both indices onto Σ , leading to

$$[h_{\alpha\beta}] = 0. \quad (22)$$

This equation gives the first junction condition, ensuring the induced metric on Σ is continuous. Consequently, the derivatives of g_{ab} are expressed as

$$\partial_c g_{ab} = \partial_c g_{ab}^+ \theta(l) + \partial_c g_{ab}^- \theta(-l). \quad (23)$$

With the metric derivatives, we can now compute the Christoffel symbols, followed by the Riemann tensor and its contractions, specifically the Ricci tensor R_{ab} and the Ricci scalar R , which remain well-defined. These are given by

$$R_{abcd} = R_{abcd}^+ \theta(l) + R_{abcd}^- \theta(-l) + \bar{R}_{abcd} \delta(l), \quad (24)$$

$$R_{ab} = R_{ab}^+ \theta(l) + R_{ab}^- \theta(-l) + \bar{R}_{ab} \delta(l), \quad (25)$$

$$R = R^+ \theta(l) + R^- \theta(-l) + \bar{R} \delta(l), \quad (26)$$

where \bar{R}_{abcd} , \bar{R}_{ab} , and \bar{R} represent the $\delta(l)$ -proportional terms, expressed in terms of geometric quantities as

$$\bar{R}_{abcd} = 4 [K_{\alpha\beta}] e_{[a}^\alpha n_b] e_{[d}^\beta n_c], \quad (27)$$

$$\bar{R}_{ab} = - \left(\epsilon [K_{\alpha\beta}] e_a^\alpha e_b^\beta + n_a n_b [K] \right), \quad (28)$$

$$\bar{R} = -2\epsilon [K], \quad (29)$$

where $X_{[ab]} \equiv \frac{1}{2}(X_{ab} - X_{ba})$ defines index anti-symmetrization, and $K = h^{\alpha\beta} K_{\alpha\beta}$ is the trace of the extrinsic curvature.

Regarding the matter sector, it is helpful to associate the presence of a thin shell at the hypersurface Σ with terms that are proportional to $\delta(l)$ in the gravitational sector of the modified field equations. The energy-momentum tensor is then written as

$$T_{ab} = T_{ab}^+ \theta(l) + T_{ab}^- \theta(-l) + S_{ab} \delta(l), \quad (30)$$

where $S_{ab} = S_{\alpha\beta} e_a^\alpha e_b^\beta$, with $S_{\alpha\beta}$ being the three-dimensional energy-momentum tensor of the thin shell. To derive T^2 in the distributional formalism, we contract T_{ab} with itself using Eq.(30), resulting in

$$T^2 = (T^2)^+ \theta(l) + (T^2)^- \theta(-l) + (\bar{T}^2) \delta(l) + (\hat{T}^2) \delta^2(l), \quad (31)$$

Note that the term proportional to $\delta^2(l)$ in Eq.(31) is singular and must be eliminated. This can only be achieved by imposing that the energy-momentum tensor of the thin shell vanishes, that is,

$$S_{ab} = 0. \quad (32)$$

When Eq.(32) is satisfied, the matching is called a smooth matching. In $f(R, T^2)$ gravity, this is the only method that preserves the regularity of the action, unlike in other gravity theories where it is a special case of a more general thin-shell matching.

Under the constraint from Eq.(32), we project the field equations for $f(R, T^2) = R + \gamma T^2$ onto the hypersurface Σ using $e_a^\alpha e_b^\beta$, leading to $[K_{\alpha\beta}] - [K] h_{\alpha\beta} = 0$. Taking the trace of this equation with $h^{\alpha\beta}$ gives $[K] = 0$, which, when substituted back into the original equation, results in

$$[K_{\alpha\beta}] = 0. \quad (33)$$

Thus, the second junction condition requires that the extrinsic curvature $K_{\alpha\beta}$ remains continuous across Σ .

In conclusion, the matching in linear $f(R, T^2)$ gravity is always smooth, meaning there is no thin shell involved. The junction conditions are the same as in GR: both the induced metric $h_{\alpha\beta}$ and the extrinsic curvature $K_{\alpha\beta}$ must be continuous across Σ , i.e.,

$$[h_{\alpha\beta}] = 0, \quad [K_{\alpha\beta}] = 0. \quad (34)$$

When considering higher-order powers of T^2 in the function, the field equations involve terms like $(T^2)^{n-1} \Theta_{ab}$ and $(T^2)^n$. In the linear case, we showed that for T^2 to be well-defined in the distributional formalism, the matching must be smooth, implying $S_{ab} = 0$. In fact, this condition ensures that both T^2 and the auxiliary tensor Θ_{ab} remain regular, with terms proportional to $\theta(l)$ and excluding $\delta(l)$. Consequently, the products of T^2 and Θ_{ab} , as well as the powers of $(T^2)^n$, preserve this regularity. As a result, higher powers of T^2 do not introduce additional junction conditions, provided that mixed RT^2 terms are absent.

3.2. Matching with an exterior vacuum spacetime

We now apply the junction conditions to match the interior wormhole spacetime with an exterior spherically symmetric and static vacuum solution. The metric expressions for the interior and exterior spacetimes are

$$ds_-^2 = -C e^{\zeta_0 \left(\frac{r_0}{r}\right)^\alpha} dt^2 + \left[1 - \left(\frac{r_0}{r}\right)^{\beta+1}\right]^{-1} dr^2 + r^2 d\Omega^2, \quad (35)$$

$$ds_+^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2, \quad (36)$$

respectively. The metric in Eq.(35) is derived from Eq.(9) using the ansatz for the redshift and shape functions in Eq.(10). The constant C is introduced for convenience to ensure that the time coordinates of the interior and exterior metrics are continuous. Meanwhile, the metric in Eq.(36) corresponds to the Schwarzschild solution with mass M ⁶⁸.

The analysis is simplified by first considering the second junction condition (34). Due to the spherical symmetry of the metrics under consideration, the extrinsic curvatures $K_{\alpha\beta}^\pm$ have two independent components: K_{00} and $K_{\theta\theta} = K_{\phi\phi} \sin^2 \theta$. This leads to the constraints $[K_{00}] = 0$ and $[K_{\theta\theta}] = 0$, expressed as

$$\frac{\alpha\zeta_0}{2} \left(\frac{r_0}{r}\right)^\alpha \sqrt{1 - \left(\frac{r_0}{r}\right)^{\beta+1}} + \frac{M}{r} \sqrt{\frac{r}{r-2M}} = 0, \quad (37)$$

$$\sqrt{1 - \left(\frac{r_0}{r}\right)^{\beta+1}} = \sqrt{1 - \frac{2M}{r}}, \quad (38)$$

respectively. Solving the second equation for the radius r using Eq.(38), we obtain unique real solutions for $M > 0$ and $r_0 > 0$, given by

$$r_\Sigma = (2M)^{-\frac{1}{\beta}} (r_0)^{1+\frac{1}{\beta}}, \quad (39)$$

This determines the matching radius r_Σ , which must satisfy $r_\Sigma > 2M$ to avoid event horizons. This implies $r_0 > 2M$ for any $\beta \geq 1$. Substituting the solutions for r_Σ into Eq.(37), we solve for ζ_0 , resulting in the expression

$$\zeta_0 = \frac{(2M)^{\frac{1-\alpha+\beta}{\beta}} (r_0)^{\frac{\alpha}{\beta}}}{\alpha \left[(2M)^{1+\frac{1}{\beta}} - (r_0)^{1+\frac{1}{\beta}} \right]}. \quad (40)$$

Since $r_0 > 2M$, for $\alpha \geq 1$ and $\beta \geq 1$, the condition $\zeta_0 < 0$ holds. This result is expected, as negative values of ζ_0 guarantee that the derivative of g_{00} has consistent signs in both the interior and exterior metrics, which is essential for achieving a smooth matching.

Next, we examine the first junction condition stated in Eq.(34). Given that the angular components of the metrics in Eqs.(35) and (36) are identical, the angular

parts of the induced metric $h_{\alpha\beta}$ are straightforwardly continuous. Consequently, the continuity condition $[h_{00}] = 0$ can be analyzed separately and is expressed as follows

$$C e^{\zeta_0 \left(\frac{r_0}{r}\right)^\alpha} = \left(1 - \frac{2M}{r}\right). \quad (41)$$

Using the results from the second junction condition, specifically the radius r_Σ from Eq.(39) and the corresponding ζ_0 from Eq.(40), we substitute these into Eq.(41) to solve for the constant C , obtaining

$$C = \left[1 - \left(\frac{2M}{r_0}\right)^{1+\frac{1}{\beta}}\right] e^{-\alpha \left[\left(\frac{r_0}{2M}\right)^{1+\frac{1}{\beta}} - 1\right]}. \quad (42)$$

Since $r_0 > 2M$, the constant C remains positive for all $\alpha \geq 1$ and $\beta \geq 1$, ensuring the correct metric signature.

In summary, for $r_0 > 2M$, $\alpha \geq 1$, and $\beta \geq 1$, the second junction condition $[K_{\alpha\beta}] = 0$ determines r_Σ (Eq.(39)) and ζ_0 (Eq.(40)), while the first condition $[h_{\alpha\beta}] = 0$ fixes C (Eq.(42)), ensuring the continuity of the full spacetime metric.

As an example, let us consider $r_0 = 3M$ and $\alpha = \beta = 1$, for which Eq.(39) gives $r_\Sigma = \frac{9}{2}M$, Eq.(40) yields $\zeta_0 = -\frac{6}{5}$, and Eq.(42) gives $C = \frac{5}{9}e^{\frac{4}{5}}$. The left panel of Figure 2 shows the g_{00} components of the interior, exterior, and matched metrics, illustrating the smooth transition of g_{00} , thus ensuring the continuity of both the induced metric and extrinsic curvature.

Examining the radial metric component g_{rr} , right panel of Figure 2, we observe that although g_{rr} is not explicitly constrained by the junction conditions, since both $h_{\alpha\beta}$ and $K_{\alpha\beta}$ are three-dimensional tensors on Σ , it remains continuous but not differentiable at $r = r_\Sigma$. This continuity of g_{rr} is expected due to its dependence on the mass function within a spherical hypersurface of radius r , given by $g_{rr} = \left(1 - \frac{2m(r)}{r}\right)^{-1}$, from which we obtain $m(r) = \frac{r_0}{2} \left(\frac{r_0}{r}\right)^\beta$ (see Eqs. (10) and (35)). Since the transition between the interior and exterior spacetimes is smooth, the mass function $m(r)$ remains continuous at r_Σ , therefore ensuring the continuity of the g_{rr} component of the metric.

4. Conclusion

In conclusion, this analysis of traversable wormholes in $f(R, T^2)$ gravity with a linear dependence on R and T^2 reveals numerous solutions satisfying all energy conditions, emphasizing their physical relevance. While the spacetime metrics of these solutions are asymptotically flat, the theory allows for non-localized matter distributions, which can be localized via junction conditions. These conditions, derived here, permit only a smooth matching, since the scalar T^2 becomes singular in the presence of a thin shell. As a result, they reduce to the junction conditions of GR, requiring continuity of the induced metric and extrinsic curvature at the hypersurface separating the interior and exterior spacetime regions. The matching procedure then yields localized wormhole solutions satisfying all energy conditions,




Fig. 2. The interior wormhole spacetime from Eq.(35) (red dashed curve) and the exterior Schwarzschild spacetime from Eq.(36) (blue dotted curve) are shown for $\beta = 1$, $r_0 = 3M$, and $\alpha = 1$. The thin black line represents the matched solution at $r = r_\Sigma$, illustrating g_{00}^{tot} (left panel) and g_{rr}^{tot} (right panel).

enhancing their astrophysical significance. Moreover, the methods presented can be extended to more general $f(R, T^2)$ models, provided no crossed terms between R and T^2 are present. Furthermore, the smooth matching requirement ensures that, in the absence of crossed terms, no additional junction conditions arise, allowing for the effective localization of solutions as in the linear T^2 case.

5. Acknowledgments

JLR acknowledges the European Regional Development Fund and the programme Mobilias Plus for financial support through Project No. MOBJD647, and project No. 2021/43/P/ST2/02141 co-funded by the Polish National Science Centre and the European Union Framework Programme for Research and Innovation Horizon 2020 under the Marie Skłodowska-Curie grant agreement No. 94533. N.G. and F.S.N.L. acknowledge funding from the Fundação para a Ciência e a Tecnologia (FCT) research grants UIDB/04434/2020 and UIDP/04434/2020. F.S.N.L. acknowledges support from the FCT Scientific Employment Stimulus contract with reference CEECINST/00032/2018, and funding from the FCT research grant PTDC/FIS-AST/0054/2021.

References

1. Morris, M.S.; Thorne, K.S. Wormholes in spacetime and their use for interstellar travel: A tool for teaching general relativity. *Am. J. Phys.* **1988**, *56*, 395.
2. Morris, M.S.; Thorne, K.S.; Yurtsever, U. Wormholes, Time Machines, and the Weak Energy Condition. *Phys. Rev. Lett.* **1988**, *61*, 1446–1449.
3. Visser, M. *Lorentzian Wormholes: From Einstein to Hawking*; Springer: New York, NY, USA, 1996.
4. Lemos, J.P.S.; Lobo, F.S.N.; Quinet de Oliveira, S. Morris-Thorne wormholes with a cosmological constant. *Phys. Rev. D* **2003**, *68*, 064004.
5. Visser, M.; Kar, S.; Dadhich, N. Traversable wormholes with arbitrarily small energy condition violations. *Phys. Rev. Lett.* **2003**, *90*, 201102.
6. Kar, S.; Sahdev, D. Evolving Lorentzian wormholes. *Phys. Rev. D* **1996**, *53*, 722–730.

7. Hawking, S.W.; Ellis, G.F.R. *The Large Scale Structure of Space-Time*; Cambridge University Press: Cambridge, UK, 1973. ISBN 978-1-00-925316-1.
8. Sajadi, S.N.; Riazi, N. Gravitational lensing by multi-polytropic wormholes. *Can. J. Phys.* **2020**, *98*, 1046–1054.
9. Sharif, M.; Zeeshan Gul, M. Viable wormhole solutions in energy-momentum squared gravity. *Eur. Phys. J. Plus* **2021**, *136*, 503.
10. Zeeshan Gul, M.; Sharif, M. Traversable Wormhole Solutions Admitting Noether Symmetry in Theory. *Symmetry* **2023**, *15*, 684.
11. Rosa, J.L.; Ganiyeva, N.; Lobo, F.S.N. Non-exotic traversable wormholes in $f(R, T_{ab}T^{ab})$ gravity. *Eur. Phys. J. C* **2023**, *83*, 1040.
12. Agnese, A.G.; La Camera, M. Wormholes in the Brans-Dicke theory of gravitation. *Phys. Rev. D* **1995**, *51*, 2011–2013.
13. Nandi, K.K.; Bhattacharjee, B.; Alam, S.M.K.; Evans, J. Brans-Dicke wormholes in the Jordan and Einstein frames. *Phys. Rev. D* **1998**, *57*, 823–828.
14. La Camera, M. Wormhole solutions in the Randall-Sundrum scenario. *Phys. Lett. B* **2003**, *573*, 27–32.
15. Lobo, F.S.N. Exotic solutions in general relativity: Traversable wormholes and ‘warp drive’ spacetimes. In *Classical and Quantum Gravity Research*; Christiansen, M.N., Rasmussen, T.K., Eds.; Nova Science Publishers: Hauppauge, NY, USA, 2008; p. 1.
16. Garattini, R.; Lobo, F.S.N. Self sustained phantom wormholes in semi-classical gravity. *Class. Quantum Gravity* **2007**, *24*, 2401–2413.
17. Lobo, F.S.N. General class of wormhole geometries in conformal Weyl gravity. *Class. Quantum Gravity* **2008**, *25*, 175006.
18. Garattini, R.; Lobo, F.S.N. Self-sustained traversable wormholes in noncommutative geometry. *Phys. Lett. B* **2009**, *671*, 146–152.
19. Lobo, F.S.N.; Oliveira, M.A. General class of vacuum Brans-Dicke wormholes. *Phys. Rev. D* **2010**, *81*, 067501.
20. Montelongo Garcia, N.; Lobo, F.S.N. Exact solutions of Brans-Dicke wormholes in the presence of matter. *Mod. Phys. Lett. A* **2011**, *40*, 3067–3076.
21. Garattini, R.; Lobo, F.S.N. Self-sustained wormholes in modified dispersion relations. *Phys. Rev. D* **2012**, *85*, 024043.
22. Myrzakulov, R.; Sebastiani, L.; Vagnozzi, S.; Zerbini, S. Static spherically symmetric solutions in mimetic gravity: Rotation curves and wormholes. *Class. Quantum Gravity* **2016**, *33*, 125005.
23. Lobo, F.S.N. (Ed.) *Wormholes, Warp Drives and Energy Conditions*; Fundamental Theories of Physics; Springer: New York, NY, USA, 2017; Volume 189.
24. Lobo, F.S.N.; Oliveira, M.A. Wormhole geometries in $f(R)$ modified theories of gravity. *Phys. Rev. D* **2009**, *80*, 104012.
25. Capozziello, S.; Harko, T.; Koivisto, T.S.; Lobo, F.S.N.; Olmo, G.J. Wormholes supported by hybrid metric-Palatini gravity. *Phys. Rev. D* **2012**, *86*, 127504.
26. Rosa, J.L.; Lemos, J.P.S.; Lobo, F.S.N. Wormholes in generalized hybrid metric-Palatini gravity obeying the matter null energy condition everywhere. *Phys. Rev. D* **2018**, *98*, 064054.
27. Rosa, J.L. Double gravitational layer traversable wormholes in hybrid metric-Palatini gravity. *Phys. Rev. D* **2021**, *104*, 064002.
28. Rosa, J.L.; Lemos, J.P.S. Junction conditions for generalized hybrid metric-Palatini gravity with applications. *Phys. Rev. D* **2021**, *104*, 124076.
29. Rosa, J.L.; André, R.; Lemos, J.P.S. Traversable wormholes with double layer thin shells in quadratic gravity. *Gen. Relativ. Gravit.* **2023**, *55*, 65.
30. Rosa, J.L.; Kull, P.M. Non-exotic traversable wormhole solutions in linear $f(R, T)$

- gravity. *Eur. Phys. J. C* **2022**, *82*, 1154.
31. Garcia, N.M.; Lobo, F.S.N. Wormhole geometries supported by a nonminimal curvature-matter coupling. *Phys. Rev. D* **2010**, *82*, 104018.
 32. Montelongo Garcia, N.; Lobo, F.S.N. Nonminimal curvature-matter coupled wormholes with matter satisfying the null energy condition. *Class. Quantum Gravity* **2011**, *28*, 085018.
 33. Harko, T.; Lobo, S.N.; Mak, M.K.; Sushkov, S.V. Modified-gravity wormholes without exotic matter. *Phys. Rev. D* **2013**, *87*, 067504.
 34. Anchordoqui, L.A.; Perez Bergliaffa, S.E.; Torres, D.F. Brans-Dicke wormholes in nonvacuum space-time. *Phys. Rev. D* **1997**, *55*, 5226–5229.
 35. Di Grezia, E.; Battista, E.; Manfredonia, M.; Miele, G. Spin, torsion and violation of null energy condition in traversable wormholes. *Eur. Phys. J. Plus* **2017**, *132*, 537.
 36. Bhawal, B.; Kar, S. Lorentzian wormholes in Einstein–Gauss-Bonnet theory. *Phys. Rev. D* **1992**, *46*, 2464.
 37. Dotti, G.; Oliva, J.; Troncoso, R. Exact solutions for the Einstein-Gauss-Bonnet theory in five dimensions: Black holes, wormholes and spacetime horns. *Phys. Rev. D* **2007**, *76*, 064038.
 38. Mehdizadeh, M.R.; Kord Zangeneh, M.; Lobo, F.S.N. Einstein-Gauss-Bonnet traversable wormholes satisfying the weak energy condition. *Phys. Rev. D* **2015**, *91*, 084004.
 39. Bronnikov, K.A.; Kim, S.W. Possible wormholes in a brane world. *Phys. Rev. D* **2003**, *67*, 064027.
 40. Lobo, F.S.N. A General class of braneworld wormholes. *Phys. Rev. D* **2007**, *75*, 064027.
 41. Israel, W. Singular hypersurfaces and thin shells in general relativity. *Nuovo Cimento B* **44**, 1 (1966).
 42. Visser, M. Traversable wormholes: Some simple examples. *Phys. Rev. D* **1989**, *39*, 3182.
 43. Visser, M. Traversable wormholes from surgically modified Schwarzschild spacetimes. *Nucl. Phys. B* **1989**, *328*, 203.
 44. Lobo, F.S.N. Energy conditions, traversable wormholes and dust shells. *Gen. Relativ. Gravit.* **2005**, *37*, 2023–2038.
 45. Lobo, F.S.N. Surface stresses on a thin shell surrounding a traversable wormhole. *Gen. Relativ. Gravit.* **2004**, *21*, 4811–4832.
 46. Lobo, F.S.N. Phantom energy traversable wormholes. *Phys. Rev. D* **2005**, *71*, 084011.
 47. Lobo, F.S.N. Stability of phantom wormholes. *Phys. Rev. D* **2005**, *71*, 124022.
 48. Schwarzschild, K. *Über das Gravitationsfeld Einer Kugel aus Inkompressibler Flüssigkeit Nach der Einsteinschen Theorie*; Sitzungsberichte der Königlich-Preussischen Akademie der Wissenschaften Berlin: Berlin, Germany, 1916; pp. 424–434.
 49. Rosa, J.L.; Piçarra, P. Existence and stability of relativistic fluid spheres supported by thin-shells. *Phys. Rev. D* **2020**, *102*, 6.
 50. Rosa, J.L. Observational properties of relativistic fluid spheres with thin accretion disks. *arXiv* **2023**, arXiv:2302.11915.
 51. Oppenheimer, J.R.; Snyder, H. On Continued Gravitational Contraction. *Phys. Rev.* **1939**, *56*, 455.
 52. Rosa, J.L.; Carloni, S. Junction conditions for general LRS spacetimes in the $1+1+2$ covariant formalism. *arXiv* **2023**, arXiv:2303.12457.
 53. Rosa, J.L. Junction conditions in gravity theories with extra scalar degrees of freedom. *Phys. Rev. D* **2024**, *109*, 064018.

54. Senovilla, J.M.M. Junction conditions for $F(R)$ -gravity and their consequences. *Phys. Rev. D* **2013**, *88*, 064015.
55. Vignolo, S.; Cianci, R.; Carloni, S. On the junction conditions in $f(R)$ -gravity with torsion. *Class. Quantum Gravity* **2018**, *35*, 095014.
56. Reina, B.; Senovilla, J.M.M.; Vera, R. Junction conditions in quadratic gravity: Thin shells and double layers. *Class. Quantum Gravity* **2016**, *33*, 105008.
57. Deruelle, N.; Sasaki, M.; Sendouda, Y. Junction conditions in $f(R)$ theories of gravity. *Prog. Theor. Exp. Phys.* **2008**, *119*, 237–251.
58. Olmo, G.J.; Rubiera-Garcia, D. Junction conditions in Palatini $f(R)$ gravity. *Class. Quantum Gravity* **2020**, *37*, 215002.
59. Rosa, J.L. Junction conditions and thin shells in perfect-fluid $f(R, T)$ gravity. *Phys. Rev. D* **2021**, *103*, 104069.
60. Rosa, J.L.; Rubiera-Garcia, D. Junction conditions of Palatini $f(R, T)$ gravity. *Phys. Rev. D* **2022**, *106*, 064007.
61. Suffern, K.G. Singular hypersurfaces in the Brans-Dicke theory of gravity. *J. Phys. A Math. Gen.* **1982**, *15*, 1599.
62. Barrabes, C.; Bressange, G.F. Singular hypersurfaces in scalar—Tensor theories of gravity. *Class. Quantum Gravity* **1997**, *14*, 805–824.
63. Padilla, A.; Sivanesan, V. Boundary Terms and Junction Conditions for Generalized Scalar-Tensor Theories. *J. High Energy Phys.* **2012**, *08*, 122.
64. Casado-Turrión, A.; de la Cruz-Dombriz, Á.; Jiménez-Cano, A.; Maldonado Torralba, F.J. Junction conditions in bi-scalar Poincaré gauge gravity. *J. Cosmol. Astropart. Phys.* **2023**, *07*, 023.
65. de la Cruz-Dombriz, Á.; Dunsby, P.K.S.; Saez-Gomez, D. Junction conditions in extended Teleparallel gravities. *J. Cosmol. Astropart. Phys.* **2014**, *12*, 048.
66. Arkuszewski, W.; Kopczynski, W.; Ponomarev, V.N. Matching Conditions in the Einstein-Cartan Theory of Gravitation. *Commun. Math. Phys.* **1975**, *45*, 183–190.
67. Macias, A.; Lammerzahl, C.; Pimentel, L.O. Matching conditions in metric affine gravity. *Phys. Rev. D* **2002**, *66*, 104013.
68. Schwarzschild, K. On the gravitational field of a mass point according to Einstein's theory. *Sitzungsber. Preuss. Akad. Wiss. Berlin Math. Phys.* **1916**, *1916*, 189–196.