

Quantum-Corrected Gravitational Interaction Incorporated with a Noninertial Cosmic String Spacetime

M. Baradaran^{1*}, L.M. Nieto^{2†}, and S. Zarrinkamar^{2,3‡}

¹ Department of Physics, Faculty of Science, University of Hradec Králové,
Rokitanského 62, 500 03 Hradec Králové, Czechia

² Departamento de Física Teórica, Atómica y Óptica, and Laboratory for Disruptive
Interdisciplinary Science (LaDIS), Universidad de Valladolid, 47011 Valladolid, Spain

³ Department of Basic Sciences, Garmsar Branch, Islamic Azad University, Garmsar, Iran

February 27, 2025

Abstract

Motivated by the great interest in studying quantum and gravitational phenomena in a unified way, scalar bosons are considered in a cosmic-string spacetime and in a non-inertial framework, with a generalized gravitational interaction containing both relativistic and quantum corrections. Solutions for the quantum-corrected gravitational interaction for an arbitrary state are reported in a quasi-exact manner and a discussion of the structure of the problem, whose special case appears in double confluent Heun form, is given.

Keywords: cosmic string, non-inertial effect, scalar field, quasi-exact solution, double-confluent Heun functions.

1 Introduction

The existence of cosmic strings, which are a kind of topological defect, was originally proposed by Kibble, whose ideas have greatly affected several fields of physics, including cosmology, quantum field theory, high-energy physics, and, currently, quantum optics [1]. Following the idea of Kibble and others, several studies have been carried out on the formulation of cosmic strings [2–4]. A direct consequence of working on cosmic string formalism is the modification of the structure of the equations. In this way, various problems have been raised. The modified equations of motion in the case of a massless scalar field for realistic cosmic string models in two- and four-dimensional cases was studied analytically using the Green’s function approach by Allen and Ottewill [5]. The Klein-Gordon equation in cosmic string spacetime was investigated within a non-inertial reference frame by Motta and Bakke [6]. The Klein-Gordon equation in a rotating cosmic string spacetime was solved in [7] for a combination of linear and Coulomb interaction terms. Analytical solutions of the linear and Coulomb interactions within a non-inertial cosmic string spacetime within a generalized metric were obtained by Santos and Basros [8, 9]. The cosmic string formulation in the so-called q -deformed approach was studied in [10]. The spin-zero version of the Duffin-Kemmer-Petiau equation with a linear interaction, i.e. the so-called Dirac oscillator, was solved in [11].

*marzieh.baradaran@uhk.cz, ORCID: 0000-0002-8455-9973

†luismiguel.nieto.calzada@uva.es, ORCID: 0000-0002-2849-2647

‡saber.zarrinkamar@uva.es, ORCID: 0000-0001-9128-4624

In [12] the cosmic string spacetime of the Klein-Gordon equation with several dislocations was considered and the associated Aharonov-Bohm and Sagnac effects were discussed. The so-called Dirac oscillator was solved exactly for a certain form of cosmic string spacetime by Hosseinpour and collaborators [13]. Some interaction terms, including linear, Yukawa and Cornell-type terms, were solved in this framework in [14]. The Fisher entropy within the Klein-Gordon framework of cosmic strings was derived by Huang et al. [15].

On the other hand, He et al. studied the problem of a two-level system with cosmic string spacetime and extensively analyzed the corresponding entanglement problem [16]. The Klein-Gordon oscillator in connection to the Kaluza-Klein theory was discussed by Ahmed [17]. The quantum coherence of a weakly coupled two-level atom to a bath of fluctuating electromagnetic field within cosmic string spacetime was comprehensively analyzed by [18]. The effect of non-commutativity in momentum was studied for the case of the Klein-Gordon oscillator by Cuzinatto and collaborators [19]. In a pair of papers, the Feshbach-Villars representation of the Klein-Gordon equation was analyzed within rotating and non-inertial cosmic string frameworks [20, 21]. The theory of gravitational lensing in the framework of cosmic strings was reviewed in [22]. Vacuum polarization effects due to a cosmic string in an anti-de Sitter ($D+1$)-dimensional spacetime were studied analytically in [23, 24]. Interestingly, the concept of non-inertial frame has also been investigated in relation to modern applications of quantum mechanics and its interface with other concepts and fields, including electrodynamics [25], teleportation [26–28], entanglement transfer [29–32], quantum encryption protocol [33], Fisher information [34], quantum metrology [35], and Airy beam propagation [36].

Here, motivated by the importance of cosmic string space, non-inertial formalism and the long-standing desire to consider quantum effects in connection with gravitation, we consider the very recently proposed generalized gravitational interaction [37–39] within the framework of the Klein-Gordon equation when a non-inertial term is present. The generalized interaction, in addition to the usual Coulomb term, includes inverse quadratic and inverse cubic terms, which correspond to post-Newtonian (relativistic) and quantum corrections, respectively. This is perhaps the simplest way to incorporate the three important concepts of physics mentioned above in a single study. The structure of the present study, which is also of pedagogical interest, is as follows. In Section 2 we first review the Klein-Gordon equation in a cosmic string spacetime when a non-inertial term is present. In Section 3 we will review the mathematical structure of the problem, which, as will be shown, does not appear, as far as we know, in the form of a special function known for mathematical physics. Rather, it appears as a generalization of the double-confluent Heun equation, which has so far not been addressed in the literature in closed series form. To tackle with the equation, we propose the quasi-exact Bethe-ansatz approach. Next, the special case of the Coulomb interaction is recovered. The paper ends with some concluding remarks.

2 Scalar field in cosmic string spacetime in the presence of a non-inertial term

The cosmic string is usually considered to be cylindrically symmetric and static with the metric

$$ds^2 = -dt'^2 + dr'^2 + \alpha^2 r'^2 d\phi'^2 + dz'^2, \quad (2.1)$$

where $\alpha = 1 - 4G\mu$ (where μ is the mass density of the cosmic string) denotes the form of curvature and obviously $\alpha = 1$ results in the flat spacetime case. Generalization to a non-inertial frame that is

rotating with constant velocity is possible by the transformation

$$t' = t, \quad r' = r, \quad \phi' = \phi + \omega t, \quad z' = z, \quad (2.2)$$

where ω is the angular velocity of the rotating frame. Therefore, the line element is

$$ds^2 = -(1 - \alpha^2 r^2 \omega^2) dt^2 + 2\alpha^2 r^2 \omega dt d\phi + dr^2 + \alpha^2 r^2 d\phi^2 + dz^2, \quad (2.3)$$

where $0 < r < \frac{1}{\alpha\omega}$ to assure that the velocity of the particle is not greater than that of light. Considering the gauge $m \rightarrow m + V(r)$, the scalar Klein-Gordon equation has the form [2]

$$-\frac{1}{\sqrt{-g}} \partial_\mu (g^{\mu\nu} \sqrt{-g} \partial_\nu \psi) + (m + V)^2 \psi = 0, \quad (2.4)$$

where m is the mass of the particle and g denotes the determinant of the metric tensor. We will look for solutions of the form

$$\psi(t, r, z, \phi) = e^{-i\epsilon t} e^{-i\ell\phi} e^{ip_z z} R(r), \quad (2.5)$$

where $\ell = 0, \pm 1, \pm 2, \pm 3, \dots$, p_z stands for the momentum components in the z -direction and ϵ can be regarded as the energy of the particle. Some simple algebra gives [8]

$$\left\{ \frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{\ell^2/\alpha^2}{r^2} - (m + V(r))^2 + (\epsilon + \omega\ell)^2 - p_z^2 \right\} R(r) = 0. \quad (2.6)$$

Performing the transformation $R(r) = r^{-1/2} u(r)$, we arrive at a Schrödinger-type equation

$$\left\{ \frac{d^2}{dr^2} + \frac{\alpha^2 - 4\ell^2}{4\alpha^2 r^2} - (m + V(r))^2 + (\epsilon + \omega\ell)^2 - p_z^2 \right\} u(r) = 0. \quad (2.7)$$

Let us now introduce the following generalized Coulomb/gravitational interaction:

$$V(r) = \frac{a_c}{r} + \frac{a_r}{r^2} + \frac{a_q}{r^3} \quad \text{with} \quad a_c, a_r, a_q < 0. \quad (2.8)$$

As the indices represent, the first term is the ordinary Coulomb term, and the second and third terms, respectively, denote the relativistic and quantum corrections [37–39].

3 Bethe-ansatz approach for the generalized Coulomb interaction

Before addressing the essential part of this work, let us first discuss the special cases of the generalized gravitational interaction. Obviously, when only the ordinary Coulomb term is present, the problem resembles the case of the non-relativistic Coulomb problem already addressed in [8] and finally recovered here as a special case. The problem when both relativistic and Coulomb corrections are present resembles the so-called double-confluent Heun function addressed by several authors through different techniques in the framework of various wave equations [40–47]. We now start the analysis of the equation structure through the Bethe-ansatz approach.

3.1 General solution

Substituting (2.8) into (2.7), we have the Schrödinger-like equation

$$\left\{ \frac{d^2}{dr^2} + V_{eff}(r) \right\} u_n(r) = 0, \quad (3.1)$$

with an effective potential

$$V_{eff}(r) = (\omega\ell + \epsilon_n)^2 - \left(m^2 + p_z^2 \right) - \frac{2a_c m}{r} - \frac{\alpha^2 (4a_c^2 + 8a_r m - 1) + 4\ell^2}{4\alpha^2 r^2} - \frac{2(a_c a_r + a_q m)}{r^3} - \frac{2a_c a_q + a_r^2}{r^4} - \frac{2a_q a_r}{r^5} - \frac{a_q^2}{r^6}. \quad (3.2)$$

To distinguish between the various quantum states, we have added the index n to ϵ and the reduced radial function $u(r)$. Next, to ensure the appropriate asymptotic behavior of the wave function at the origin and at infinity, we propose the solution to be of the form

$$u_n(r) = \exp \left[Ar + \frac{B}{r} + \frac{\Gamma}{r^2} + \Delta \ln r \right] \psi_n(r), \quad (3.3)$$

where the new radial function $\psi_n(r)$ must be a polynomial function of order $n = 0, 1, 2, \dots$, and $A, B, \Gamma, < 0$ and $\Delta > 0$ are parameters to be determined. Substituting (3.3) into (3.1) yields the following ordinary differential equation for $\psi_n(r)$:

$$\left\{ r^3 \frac{d^2}{dr^2} + \left(2A r^3 + 2\Delta r^2 - 2B r - 4\Gamma \right) \frac{d}{dr} + \left(\Lambda_2 r^2 + \Lambda_1 r + \Lambda_0 \right) \right\} \psi_n(r) = 0, \quad (3.4)$$

where

$$\Lambda_2 = 2A\Delta - 2a_c m, \quad (3.5a)$$

$$\Lambda_1 = -2AB - a_c^2 - 2a_r m + \Delta^2 - \Delta - \frac{\ell^2}{\alpha^2} + \frac{1}{4}, \quad (3.5b)$$

$$\Lambda_0 = -2(2A\Gamma + a_c a_r + a_q m + B(\Delta - 1)), \quad (3.5c)$$

and the parameters A, B, Γ and Δ are given by

$$A = -\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_n)^2}, \quad B = a_r, \quad \Gamma = \frac{a_q}{2}, \quad \Delta = \frac{3}{2} - a_c > 0. \quad (3.6)$$

Now, following the Bethe-ansatz method [48], we assume that (3.4) has polynomial solutions of degree n :

$$\psi_n(r) = \begin{cases} 1, & n = 0, \\ \prod_{i=1}^n (r - r_i), & n = 1, 2, \dots, \end{cases} \quad (3.7)$$

where r_i are unknown roots to be determined. Then, according to Theorem 1.1 of [48], the values of the coefficients Λ_j , $j = 0, 1, 2$ in (3.4) are given by

$$\Lambda_0 = -2 \left(-nB + (\Delta + n - 1) \sum_{i=1}^n r_i + A \sum_{i=1}^n r_i^2 \right), \quad (3.8a)$$

$$\Lambda_1 = -n(2\Delta + n - 1) - 2A \sum_{i=1}^n r_i, \quad \Lambda_2 = -2nA, \quad (3.8b)$$

in which the roots r_1, r_2, \dots, r_n are determined by the Bethe ansatz equations

$$\sum_{j \neq i}^n \frac{1}{r_i - r_j} + \frac{A r_i^3 + \Delta r_i^2 - B r_i - 2\Gamma}{r_i^3} = 0, \quad i = 1, 2, \dots, n. \quad (3.9)$$

Let us now obtain the general solutions of our equation (3.1). Substituting Λ_2 and A from (3.5a) and (3.6) into the second equation of (3.8b), after some manipulations, the energy ϵ_n is found to be

$$\epsilon_n(a_c; m, \ell, \omega, p_z) = -\ell\omega \pm \sqrt{p_z^2 + m^2 \left[1 - \frac{a_c^2}{\left(\frac{3}{2} + n - a_c\right)^2} \right]}, \quad (3.10)$$

whenever the expression under the square root is positive. Note that although the energy expression (3.10) depends only on the Coulomb term a_c of the potential (2.8), the other two potential parameters a_r and a_q also play an important role. In other words, (3.8a) and the first relation in (3.8b) result in severe restrictions on the potential parameters given explicitly by the following closed-form expressions:

$$a_r = \frac{2(n+1)a_c + 2\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_n)^2} \sum_{i=1}^n r_i - (n+1)^2 + \frac{\ell^2}{\alpha^2}}{2 \left(\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_n)^2} - m \right)}, \quad (3.11a)$$

$$a_q = \frac{-(2n+1)a_r - 2\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_n)^2} \sum_{i=1}^n r_i^2 + (2n+1 - 2a_c) \sum_{i=1}^n r_i}{2 \left(\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_n)^2} - m \right)}. \quad (3.11b)$$

On the other hand, the closed form expression for the wave function associated with the energy (3.10), from (3.3) together with (3.6) and (3.7), is given explicitly by

$$u_n(r) = \exp \left[-\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_n)^2} r + \frac{a_r}{r} + \frac{a_q}{2r^2} + \left(\frac{3}{2} - a_c \right) \ln r \right] \psi_n(r). \quad (3.12)$$

In the Eqs.(3.11) and (3.12), the roots r_i are determined by the Bethe ansatz equations

$$\sum_{j \neq i}^n \frac{1}{r_i - r_j} + \left(\frac{3}{2} - a_c \right) \frac{1}{r_i} - \frac{a_r}{r_i^2} - \frac{a_q}{r_i^3} - \sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_n)^2} = 0, \quad i = 1, 2, \dots, n. \quad (3.13)$$

In summary, for a given n , the ‘general solutions’ of our equation (3.1) are given explicitly through the Eqs.(3.10)-(3.13). Below, we explicitly report the exact solutions of the ground state and the first excited state.

3.2 Ground state solution

For $n = 0$, from (3.10) and (3.12), the ground state energy and the corresponding wave function are given respectively by

$$\epsilon_0(a_c; m, \ell, \omega, p_z) = -\ell\omega \pm \sqrt{p_z^2 + m^2 \left[1 - \frac{a_c^2}{\left(\frac{3}{2} - a_c\right)^2} \right]}, \quad (3.14)$$

and

$$u_0(r) = r^{\left(\frac{3}{2} - a_c\right)} \exp \left[-\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_0)^2} r + \frac{a_r}{r} + \frac{a_q}{2r^2} \right]. \quad (3.15)$$

Furthermore, from (3.11), the allowed values of the potential parameters, in terms of the Coulomb term a_c , are given by

$$a_r = \frac{1}{2} \frac{2a_c - 1 + \frac{\ell^2}{\alpha^2}}{\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_0)^2} - m}, \quad a_q = -\frac{1}{2} \frac{a_r}{\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_n)^2} - m}. \quad (3.16)$$

Note that the Bethe ansatz equation (3.13) plays no role for the ground state case. As stated in (3.7), $\psi_0(r) \equiv 1$.

3.3 First excited state solution

Similarly, for $n = 1$, from (3.10) and (3.12), the energy of the first excited state and its wave function are given, respectively, by

$$\epsilon_1(a_c; m, \ell, \omega, p_z) = -\ell\omega \pm \sqrt{p_z^2 + m^2 \left[1 - \frac{a_c^2}{\left(\frac{5}{2} - a_c\right)^2} \right]}, \quad (3.17)$$

and

$$u_1(r) = (r - r_1)r^{\left(\frac{3}{2} - a_c\right)} \exp \left[-\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_1)^2} r + \frac{a_r}{r} + \frac{a_q}{2r^2} \right]. \quad (3.18)$$

In this case, starting from (3.11), the potential parameters a_r and a_q , again in terms of the Coulomb term a_c as well as r_1 , are given by the following expressions

$$a_r = \frac{1}{2} \frac{4(a_c - 1) + 2\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_1)^2} r_1 + \frac{\ell^2}{\alpha^2}}{\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_1)^2} - m}, \quad (3.19a)$$

$$a_q = -\frac{1}{2} \frac{3a_r + 2\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_1)^2} r_1^2 + (2a_c - 3)r_1}{\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_1)^2} - m}. \quad (3.19b)$$

The unknown parameter r_1 appearing in (3.18) and (3.19) is determined, from (3.13), from the following Bethe ansatz equation

$$\sqrt{m^2 + p_z^2 - (\omega\ell + \epsilon_1)^2} r_1^3 + \left(a_c - \frac{3}{2}\right) r_1^2 + a_r r_1 + a_q = 0. \quad (3.20)$$

3.4 The special case of Coulomb interaction

Now, let us check the special case of Coulomb interaction, i.e. when the potential parameters a_r and a_q in (2.8) vanish, by comparing the results with Ref. [8]. Note that due to the cumbersome structure of the potential constraints (3.11) and the roots of the Bethe ansatz equations (3.13) appearing within them, there is probably no hope of inspecting the limit for general n . However, we can check this explicitly for the ground state. To this purpose, let us rewrite the ground state energy (3.14) as

$$\epsilon_0 = -\ell\omega \pm \sqrt{p_z^2 + m^2 \left[1 - \frac{a_c^2}{\left(\frac{1}{2} + \sqrt{a_c^2 - 2a_c + 1}\right)^2} \right]}, \quad (3.21)$$

where we have replaced a_c in the denominator of the fraction in (3.14) by $1 - \sqrt{(a_c - 1)^2}$ since a_c and hence $a_c - 1$ are negative. On the other hand, from (3.16), we see that the potential parameters a_r and a_q vanish whenever $a_c = \frac{1}{2} - \frac{\ell^2}{2\alpha^2}$. Then, replacing a_c in (3.21) from this condition (only the first power term) we get

$$\epsilon_0 = -\ell\omega \pm \sqrt{p_z^2 + m^2 \left[1 - \frac{a_c^2}{\left(\frac{1}{2} + \sqrt{a_c^2 + \frac{\ell^2}{\alpha^2}}\right)^2} \right]},$$

which is the same as Eq. (27) in Ref. [8] for $\tilde{N} = 0$ (note that η appearing there is equivalent to our potential parameter a_c).

4 Conclusion

Relativistic scalar particles were investigated within a cosmic string spacetime exhibiting non-inertial effects. Based on very recent results, the generalized gravitational potential, which includes post-Newtonian (relativistic) and quantum corrections, was considered as the interaction case. It was briefly discussed that the resulting equation appears in a form that can be regarded as a generalization of the double-confluent Heun equation for which no closed form solutions are available. It was commented that the equation can not be solved, as far as we know, by other analytical (non-perturbative) techniques including the general form of Lie algebra approach, integral transforms frequently used in quantum mechanics, etc. As a result and based on a Bethe ansatz approach, arbitrary state solutions were reported. Having obtained the energy spectrum, the energy shift due to the quantum and relativistic terms was given in (3.10) in a non-explicit way. The special case of the ordinary gravitational potential was recovered [8]. The results obtained are important because it is a simultaneous study of gravitational and quantum effects in

a non-inertial cosmic string framework. We hope the present work, in connection with other theoretical and experimental studies in this field, will lead to a better understanding of the interface of quantum mechanics and gravity. As the final point, it should be mentioned that a particular difficulty of this way of approaching the problem at hand is that the values of the potential parameters are quite diverse and this motivates even more studies in order to have a more deterministic numerical statement on the modification of ordinary case.

Acknowledgments

The work of MB was supported by the Czech Science Foundation, project 22-18739S. The research of LMN and SZ was supported by the Q-CAYLE project, funded by the European Union-Next Generation UE/MICIU/Plan de Recuperacion, Transformacion y Resiliencia/Junta de Castilla y Leon (PRTRC17.11), and also by project PID2023-148409NB-I00, funded by MICIU/AEI/10.13039/501100011033. Financial support of the Department of Education of the Junta de Castilla y Leon and FEDER Funds is also gratefully acknowledged (Reference: CLU-2023-1-05). The authors are also grateful to M. J. Lake for his kind comments and critics on this manuscript.

References

- [1] T. W. K.ibble, Topology of cosmic domains and strings, *J. Phys. A: Math. Gen.* **9** (1976) 1387.
- [2] N. Birrell and P. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, 1984).
- [3] T. Harko and M. J. Lake, Cosmic strings in $f(R, L_m)$ gravity, *Eur. Phys. J. C* **75** (2015) 60.
- [4] A. Vilenkin, Cosmic strings and domain walls, *Phys. Rep.* **121** (1985) 263.
- [5] A. Allen and A. C. Ottewill, Effects of curvature couplings for quantum fields on cosmic-string spacetimes, *Phys. Rev. D* **42** (1990) 2669.
- [6] H. F. Motta and K. Bakke, Noninertial effects on the ground state energy of a massive scalar field in the cosmic string spacetime, *Phys. Rev. D* **89** (2014) 027702.
- [7] M. S. Cunha, C. R. Muniz, H. R. Christiansen, V. B. Bezerra, Relativistic Landau levels in the rotating cosmic string spacetime, *Eur. Phys. J. C* **76** (2016) 512.
- [8] L. C. N. Santos and C. C. Barros Jr., Scalar bosons under the influence of noninertial effects in the cosmic string spacetime, *Eur. Phys. J. C* **77** (2017) 186.
- [9] L. C. N. Santos and C. C. Barros Jr., Relativistic quantum motion of spin-0 particles under the influence of noninertial effects in the cosmic string spacetime, *Eur. Phys. J. C* **78** (2018) 13.
- [10] H. Sobhani, H. Hassanabadi and W. S. Chung, Effects of cosmic-string framework on the thermodynamical properties of anharmonic oscillator using the ordinary statistics and the q-deformed superstatistics approaches, *Eur. Phys. J. C* **78** (2018) 106.

- [11] M. Hosseinpour, H. Hassanabadi and F. M. Andrade, The DKP oscillator with a linear interaction in the cosmic string space-time, *Eur. Phys. J. C* **78** (2018) 93.
- [12] R. L. L. Vitória and K. Bakke, Rotating effects on the scalar field in the cosmic string spacetime, in the spacetime with space-like dislocation and in the spacetime with a spiral dislocation, *Eur. Phys. J. C* **78** (2018) 175.
- [13] M. Hosseinpour, H. Hassanabadi and M. de Montigny, The Dirac oscillator in a spinning cosmic string spacetime, *Eur. Phys. J. C* **79** (2019) 311.
- [14] L.-F. Deng, C.-Y. Longa, Z.-W. Long, and T. Xu, The generalized K-G oscillator in the cosmic string space-time, *Eur. Phys. J. Plus* **134** (2019) 355.
- [15] Z. Huang, H. Situ and Z. He, Quantum Fisher information in the cosmic string spacetime, *Class. Quant. Grav.* **37** (2020) 175002.
- [16] P. He, F. Yu and J. Hu, Entanglement dynamics for static two-level atoms in cosmic string spacetime, *Eur. Phys. J. C* **80** (2020) 134.
- [17] F. Ahmed, The generalized Klein–Gordon oscillator in the background of cosmic string space-time with a linear potential in the Kaluza–Klein theory, *Eur. Phys. J. C* **80** (2020) 211.
- [18] Z. Huang, Quantum coherence for an atom interacting with an electromagnetic field in the background of cosmic string spacetime, *Quantum Inf. Process.* **19** (2020) 370.
- [19] R. R. Cuzinatto, M. de Montigny and P. J. Pompeia, Non-commutativity and non-inertial effects on a scalar field in a cosmic string space-time: I. Klein–Gordon oscillator, *Class. Quant. Grav.* **39** (2022) 075006.
- [20] A. Bouzenada and A. Boumali, Statistical properties of the two dimensional Feshbach–Villars oscillator (FVO) in the rotating cosmic string space–time, *Ann. Phys.* **452** (2023) 169302.
- [21] A. Bouzenada, A. Boumali and E. O. Silva, Applications of the Klein–Gordon equation in the Feshbach–Villars representation in the non-inertial cosmic string space–time, *Ann. Phys.* **458** (2023) 169479.
- [22] I. I. Bulygin, M. V. Sazhin and O. S. Sazhina, Theory of gravitational lensing on a curved cosmic string, *Eur. Phys. J. C* **83** (2023) 844.
- [23] W. O. dos Santos and E. R. B. de Mello, Vacuum polarization induced by a cosmic string and a brane in AdS spacetime, *Eur. Phys. J. C* **83** (2023) 726.
- [24] W. O. dos Santos and E. R. B. de Mello, Induced current by a cosmic string and a brane in high-dimensional AdS spacetime, *Eur. Phys. J. C* **84** (2024) 224.
- [25] Y. N. Obukhov, Electrodynamics in noninertial frames, *Eur. Phys. J. C* **81** (2021) 919.
- [26] P. M. Alsing and G. J. Milburn, Teleportation with a Uniformly Accelerated Partner, *Phys. Rev. Lett.* **91** (2003) 180404.

- [27] P. M. Alsing, D. McMahon and G. J. Milburn, Teleportation in a non-inertial frame, *J. Opt. B* **6** (2004) S834.
- [28] X. Chen and K. W. C. Chan, Quantum teleportation of Dirac fields in noninertial frames with amplitude damping, *Phys. Rev. A* **99** (2019) 022334.
- [29] I. Fuentes-Schuller and R. B. Mann, Alice Falls into a Black Hole: Entanglement in Noninertial Frames, *Phys. Rev. Lett.* **95** (2005) 120404.
- [30] P. M. Alsing, I. Fuentes-Schuller, R. B. Mann, and T. E. Tessier, Entanglement of Dirac fields in noninertial frames, *Phys. Rev. A* **74** (2006) 3232.
- [31] M. Fink, A. Rodriguez-Aramendia, J. Handsteiner, A. Ziarkash, F. Steinlechner, T. Scheidl, I. Fuentes, J. Pienaar, T.C. Ralph, R. Ursin, Experimental test of photonic entanglement in accelerated reference frames, *Nat. Comm.* **8** (2017) 15304 .
- [32] A. J. Torres-Arenas, Q. Dong, G.-H. Sun, W.-C. Qiang, S.-H. Dong, Entanglement measures of W-state in noninertial frames, *Phys. Lett. B* **789** (2019) 93.
- [33] K. Brádler, Eavesdropping of quantum communication from a noninertial frame, *Phys. Rev. A* **75** (2007) 022311.
- [34] Y. Yao, X. Xiao, L. Ge, X. Wang, and C. Sun, Quantum Fisher information in noninertial frames, *Phys. Rev. A* **89** (2014) 042336.
- [35] M. Ahmadi, D. Edward Bruschi and I. Fuentes, Quantum metrology for relativistic quantum fields, *Phys. Rev. D* **89** (2014) 065028.
- [36] F. Wang, D. Liu and L. Wang, Propagation of Airy beams in uniformly accelerated space, *Opt. Comm.* **537** (2023) 129445.
- [37] T. de Paula Netto, L. Modesto and I. L. Shapiro, Universal leading quantum correction to the Newton potential, *Eur. Phys. J. C* **82** (2022) 160.
- [38] M. B. Fröb, C. Rein and R. Verch, Graviton corrections to the Newtonian potential using invariant observables, *JHEP* **01** (2022)180.
- [39] G. Lambiase and F. Scargildi, Generalized uncertainty principle and asymptotically safe gravity, *Phys. Rev. D* **105** (2022) 124054.
- [40] A. Ronveaux, Heun's differential equations (Oxford University Press, New York, 1995).
- [41] L. El-Jaick and B. D. B. Figueiredo, Solutions for confluent and double-confluent Heun equations, *J. Math. Phys.* **49** (2008) 083508.
- [42] A. M. Ishkhanyan, Schrödinger potentials solvable in terms of the confluent Heun function, *Theor. Math. Phys.* **188**(1) (2016) 980.
- [43] A.V. Turbiner, One-dimensional quasi-exactly solvable Schrödinger equations, *Phys. Rep.* **642** (2016) 1.

- [44] A. Turbiner, Quasi-exactly-solvable problems and $sl(2)$ algebra, *Commun. Math. Phys.* **118**, (1988) 467.
- [45] A. Gonzalez-Lopez, N. Kamran, and P.J. Olver, Quasi-exact solvability, *Contemp. Math.* **160** (1994) 113.
- [46] M. Baradaran, and H. Panahi, Perturbed Coulomb potentials in the Klein–Gordon equation: quasi-exact solution, *Few-Body Syst.* **59** (2018) 42.
- [47] M. Baradaran, L. M. Nieto and S. Zarrinkamar, On some quantum correction to the Coulomb potential in generalized uncertainty principle approach, *Phys. Lett. B* **852** (2024) 138603.
- [48] Y.Z. Zhang, Exact polynomial solutions of second order differential equations and their applications, *J. Phys. A: Math. Theor.* **45**, (2012) 065206.
- [49] D. Agboola, and Y.Z. Zhang, Novel quasi-exactly solvable models with anharmonic singular potentials, *Ann. Phys.* **330** (2013) 246.