

FROBENIUS SUBALGEBRA LATTICES IN TENSOR CATEGORIES

MAINAK GHOSH AND SEBASTIEN PALCOUX

ABSTRACT. This paper investigates Frobenius subalgebra lattices within tensor categories and extends Watatani's theorem on the finiteness of intermediate subfactors to this setting, under certain positivity conditions encompassing the pseudo-unitary fusion categories. The primary tools employed in this study are semisimplification and a concept of formal angle. Additionally, we have broadened several intermediate results, such as the exchange relation theorem and Landau's theorem, to apply to abelian monoidal categories.

Key applications of our findings include a stronger version of the Ino-Watatani result: we show that the finiteness of intermediate C^* -algebras holds in a finite index unital irreducible inclusion of C^* -algebras without requiring the simple assumption. Moreover, for a finite-dimensional semisimple Hopf algebra H , we demonstrate that H^* contains a finite number of Frobenius subalgebra objects in $\text{Rep}(H)$. Finally, we explore a range of applications, including abstract spin chains, vertex operator algebras, and speculations on quantum arithmetic involving the generalization of Ore's theorem, Euler's totient and sigma functions, and RH.

1. INTRODUCTION

A *subfactor* is a unital inclusion of factors. The modern theory of subfactors (type II_1) was initiated by V. Jones [26]. In [25], it was shown that for any finite group G , there exists an outer action of G on the hyperfinite II_1 factor R . This outer action allows the formation of a crossed-product $R \rtimes G$. The lattice of intermediate subfactors of $(R \subseteq R \rtimes G)$ is isomorphic to the subgroup lattice of G . Since G is finite, this lattice is also finite. Generalizing this, Watatani [63] proved that any irreducible finite-index subfactor has a finite lattice of intermediate subfactors (henceforth referred to as *Watatani's theorem*). This paper aims to extend Watatani's theorem to the setting of rigid abelian monoidal categories and explore its applications.

The original proof in [63] relies on functional analysis. An alternative proof using planar algebra and angles between biprojections was later provided in [1]. Dave Penneys observed that Watatani's theorem can be reformulated in the framework of unitary tensor categories using the notion of Frobenius algebras in monoidal categories [34, 35, 58, 18, 7, 27]. A *Frobenius algebra* (in Vec) is a finite-dimensional unital algebra A that is isomorphic to its dual A^* as an A -module. Alternatively, it can be characterized by the existence of a nondegenerate associative bilinear form $\langle -, - \rangle$, as described in [56, Chapter IV, §1]. If this bilinear form is symmetric, the Frobenius algebra is said to be *symmetric*. Notably, any complexified fusion ring forms a symmetric Frobenius algebra, where $\langle a, b \rangle$ is the coefficient of the unit summand in ab . The standard invariant of subfactors, as discussed in [29], can be axiomatized as a Frobenius algebra. For further details, refer to [35] and see Examples 2.23.

A crucial first step toward our goal was establishing a tensor-categorical version of the *exchange relation* from planar algebras [30, 32]. This is achieved in Theorem 3.7. Building on this, we extended *Landau's theorem* (Theorem 3.15) to connected Frobenius algebras in any k -linear abelian monoidal category. The connectedness of Frobenius algebras, which extends the concept of subfactor irreducibility, plays a crucial role in the proof.

A major challenge in extending Watatani's theorem was defining a *lattice structure* for Frobenius subalgebras in a tensor category. For subgroups and intermediate subfactors, the lattice structure is inherent, but this is not the case for Frobenius subalgebras. By using the concept of pullback of monomorphisms along with additional assumptions, we proved the following theorem:

Theorem 1.1. *The poset of Frobenius subalgebras of a Frobenius algebra X , which is of finite length and semisimple as an object in an abelian rigid monoidal category, forms a lattice.*

Semisimplicity of the Frobenius algebra object X is a sufficient condition. A counterexample in [51] shows that the poset of Frobenius subalgebras may fail to be a lattice when X is not semisimple.

The notion of an *equivalence class* (Definition 4.14) is not the same as an isomorphism class; rather, it captures substructures in a categorical way. A Frobenius algebra generalizes the concept of a subfactor, while the equivalence class of its Frobenius subalgebras corresponds to intermediate subfactors. For more on its relation to biprojections, see Definition 3.5 and Proposition 4.15.

To prove finiteness results, we typically rely on comparison arguments, which in turn require some form of *positivity*. In unitary tensor categories, positivity is intrinsic. However, since we are working with more general

tensor categories, we introduce the notion of *total positivity* and focus on Frobenius algebras that satisfy this property (Definition 5.2). This is the best relaxation of unitarity that we have identified as sufficient for our proofs.

In §7, we introduce the concept of a *formal angle* between two biprojections arising from Frobenius subalgebras of a connected Frobenius algebra, building on ideas from the arXiv version of [1], see Remark 7.10. This allows us to prove the following result:

Theorem 1.2. *Let X be a totally positive, connected Frobenius algebra of finite length that is semisimple as an object in a \mathbb{C} -linear abelian rigid monoidal category \mathcal{C} , where $\mathbf{1}$ is linear-simple. Then the Frobenius subalgebra lattice of X is finite.*

The notions of *finite length*, *connected*, and *Frobenius algebra* within a monoidal category extend the notions of *finite index*, *irreducible*, and *subfactor*, respectively. The connected assumption cannot be avoided because there are non-irreducible finite index subfactors with infinite intermediate subfactor lattice.

After proving Theorem 1.2, we sought to relax the assumption of semisimplicity for X . However, as previously noted, without this assumption, the Frobenius subalgebra poset may not form a lattice. Additionally, defining a formal angle between two biprojections becomes challenging unless X is semisimple. This challenge is addressed in §6 through the concept of semisimplification [13]. For a spherical tensor category \mathcal{C} , its semisimplification is a new tensor category $\overline{\mathcal{C}}$, where the morphism compositions and tensor products remain unchanged, while the simple objects consist of the indecomposable objects of \mathcal{C} that have a nonzero dimension. Moreover, by utilizing total positivity, we have embedded the Frobenius subalgebra poset of X into a poset within $\overline{\mathcal{C}}$. By integrating all these concepts, we establish a non-semisimple version of Watatani's theorem:

Theorem 1.3. *Let X be a totally positive connected Frobenius algebra in a spherical tensor category \mathcal{C} . Then its Frobenius subalgebra poset is finite.*

As a corollary, we obtain the following result.

Corollary 1.4. *For any connected Frobenius algebra in a pseudo-unitary fusion category, the lattice of Frobenius subalgebras is finite.*

We are interested in exploring how much the pseudo-unitary assumption in Corollary 1.4 can be relaxed (see Conjecture 8.10).

In §9, §10, and §11, we explore the applications of Theorem 1.2 in various contexts. In [23], the authors extended Watatani's theorem for subfactors to irreducible unital inclusions of finite index simple C^* -algebras. In §9, we build upon this result by removing the assumption of simplicity for the C^* -algebras. We establish the following theorem:

Theorem 1.5. *A finite index unital irreducible inclusion of C^* -algebras contains a finite number of intermediate C^* -subalgebras.*

In §10, we offer a comprehensive proof of Theorem 10.7, which states that for every finite-dimensional Hopf algebra H , its dual H^* can be structured as a connected Frobenius algebra object within $\text{Rep}(H)$. This structure features a different comultiplication and counit, derived from the Fourier transform. As a consequence, we find that H^* contains a finite number of Frobenius subalgebra objects in $\text{Rep}(H)$ when H is semisimple (see Corollary 10.14).

Furthermore, as indicated by [24], the left coideal subalgebras of H^* are Frobenius subalgebras in $\text{Rep}(H)$ in the unitary case. In general, there are Frobenius algebras in Vec , and are connected unital algebra objects in $\text{Rep}(H)$. However, we encountered issues regarding the Frobenius subalgebra structure in $\text{Rep}(H)$, even within the semisimple context, where finiteness is already established by [14, Theorem 3.6]. Moreover, it is well-known that the semisimple condition is essential for establishing finiteness, even for Hopf subalgebras, as illustrated in Example 10.19.

In §11, we introduce additional examples of connected Frobenius algebra objects. In §11.1, we first demonstrate that the canonical Frobenius algebra object in a unimodular multitensor category \mathcal{C} [12, §7.20] is connected if and only if \mathcal{C} is tensor (Proposition 11.1). This result allows us to provide a connected Frobenius algebra structure on H^* as an object in $\text{Rep}(H \otimes H^{\text{cop}})$, where H is a finite-dimensional unimodular Hopf algebra. Finally, we note that in a pivotal unimodular non-semisimple finite tensor category, the canonical Frobenius algebra object cannot be totally positive. In §11.2, we discuss inclusions of abstract spin chains. In [28], an algebraic model of categorical inclusions was employed to examine the extensions of bounded-spread isomorphisms of symmetric local algebras to quantum cellular automata (QCA), defined on either the full or edge-restricted local operator algebras. We demonstrate that the lattice of categorical inclusions is finite. In §11.3, we include a brief discussion with Shimizu about the challenge of establishing a Frobenius algebra structure on a vertex operator algebra (VOA) V as an object within $\text{Rep}(V)$.

In §12, we present a comprehensive list of open problems and speculations that have emerged from our current research. This list has been made possible by the finiteness of the Frobenius subalgebra lattice. We propose a tensor categorical generalization of Ore’s theorem. Additionally, we introduce a concept of Euler’s totient and sigma functions for a connected Frobenius algebra object in a pseudo-unitary fusion category. We have also expanded the idea of subfactor depth to Frobenius algebras. Finally, we propose a categorical generalization of the Riemann hypothesis. We plan to address the questions and speculations outlined in §12 in our future research.

The primary audience for this paper is tensor categorists who may not have a background in subfactor theory. However, it can also serve as an entry point for subfactor theorists interested in exploring tensor category theory. For an introduction to tensor category theory, we refer readers to the book [12]. This paper presents proofs of several intermediate results without assuming semisimplicity. These findings could be valuable for future research that adopts a direct non-semisimple approach (i.e., without relying on semisimplification, as discussed in §6), potentially broadening the scope of the main results in this paper.

CONTENTS

1. Introduction	1
2. Frobenius algebra	3
3. Frobenius subalgebras	11
4. Lattice structure	15
5. Karoubian categories	20
6. Semisimplification	20
7. Formal angle	22
8. Watatani’s theorem	24
9. Inclusion of C^* -algebras	25
10. Hopf algebras	27
11. Other examples	29
11.1. Connected canonical Frobenius algebra	30
11.2. Abstract spin chains	30
11.3. Vertex operator algebras	31
12. Quantum arithmetic	31
12.1. Ore’s theorem	31
12.2. Euler’s totient	32
12.3. Depth	32
12.4. Riemann hypothesis	33
Acknowledgments	33
References	34


2. FROBENIUS ALGEBRA


Let us review some definitions and fundamental results. For the basic definition of monoidal and tensor categories, we refer to [12]. According to Mac Lane’s strictness theorem, we can assume that monoidal categories are strict without any loss of generality. We will use graphical calculus, interpreting diagrams from top to bottom.

Definition 2.1. *An algebra in a monoidal category \mathcal{C} consists of a triple (X, m, e) , where X is an object in \mathcal{C} , m is a multiplication morphism in $\text{Hom}_{\mathcal{C}}(X \otimes X, X)$, and e is a unit morphism in $\text{Hom}_{\mathcal{C}}(\mathbf{1}, X)$. These elements must satisfy the following axioms:*

- (Associativity) $m \circ (m \otimes \text{id}_X) = m \circ (\text{id}_X \otimes m)$,
- (Unitality) $m \circ (e \otimes \text{id}_X) = \text{id}_X = m \circ (\text{id}_X \otimes e)$.

These relations are typically represented as follows:







Definition 2.2. A coalgebra in a monoidal category \mathcal{C} is defined as a triple (X, δ, ϵ) , where X is an object in \mathcal{C} , δ is a comultiplication morphism in $\text{Hom}_{\mathcal{C}}(X, X \otimes X)$, and ϵ is a counit morphism in $\text{Hom}_{\mathcal{C}}(X, \mathbf{1})$. These components must satisfy the following axioms:

- (Coassociativity) $(\delta \otimes \text{id}_X) \circ \delta = (\text{id}_X \otimes \delta) \circ \delta$,
- (Counitality) $(\epsilon \otimes \text{id}_X) \circ \delta = \text{id}_X = (\text{id}_X \otimes \epsilon) \circ \delta$.


These relations are typically represented as follows:




Definition 2.3. A Frobenius algebra in a monoidal category \mathcal{C} is a quintuple $(X, m, e, \delta, \epsilon)$, where (X, m, e) is an algebra and (X, δ, ϵ) is a coalgebra. These must satisfy the following axiom:

- (Frobenius) $(\text{id}_X \otimes m) \circ (\delta \otimes \text{id}_X) = \delta \circ m = (m \otimes \text{id}_X) \circ (\text{id}_X \otimes \delta)$.

This relation is typically illustrated as follows:




If the Frobenius condition is relaxed, we define a weak Frobenius algebra as follows:




We will demonstrate (Lemma 2.7) that an algebra is weak Frobenius if and only if it is Frobenius.

Lemma 2.4. Let $(X, m, \delta, e, \epsilon)$ be a weak Frobenius algebra. Then X is selfdual (i.e., $X^* = X$), with the evaluation morphism defined as $\text{ev}_X := \epsilon \circ m$ and the coevaluation morphism defined as $\text{coev}_X := \delta \circ e$, as illustrated below:



Proof. The following diagram shows that $\epsilon \circ m$ and $\delta \circ e$ satisfy the zigzag relation:



The result follows from [12, Definition 2.10.1]. □

Remark 2.5. *The definitions $\text{ev}_X := \epsilon \circ m$ and $\text{coev}_X := \delta \circ e$ may not align perfectly with a given rigid structure in the monoidal category. However, it is possible to adjust the rigid structure to achieve a perfect match, as noted in [12, Definition 2.10.11]. Additionally, the rigid structure on an object is unique up to a unique isomorphism, as stated in [12, Proposition 2.10.5]. In this paper, we will assume that this alignment is always satisfied.*


Lemma 2.6. *An algebra-coalgebra $(X, m, \delta, e, \epsilon)$ is a weak Frobenius algebra if and only if the following equalities hold:*

$$(m \otimes \text{id}_X) \circ (\text{id}_X \otimes \text{coev}_X) = \delta = (m \otimes \text{id}_X) \circ (\text{coev}_X \otimes \text{id}_X)$$


and

$$(\text{ev}_X \otimes \text{id}_X) \circ (\text{id}_X \otimes \delta) = m = (\text{id}_X \otimes \text{ev}_X) \circ (\text{id}_X \otimes \delta),$$


as illustrated below:



Proof. We will demonstrate the first equality using the following diagram (where *self-dual* refers to Lemma 2.4):




The other equalities follow similarly. Conversely, if these equalities hold, let us show the weak Frobenius condition. First, note that Lemma 2.4 can be established as well. Next, we have:



Thus, the result follows. □


Lemma 2.7. *An algebra-coalgebra is weak Frobenius if and only if it is Frobenius.*

Proof. The proof is illustrated by the following diagram (where *weak Frobenius* refers to Lemma 2.6):




The reverse direction is immediate. □

Lemma 2.8. *Let $(X, m, \delta, e, \epsilon)$ be a Frobenius algebra. Then $m^* = \delta$ and $\delta^* = m$, as shown below:*




Proof. To prove the first equality, we apply Lemma 2.6 three times:




The second equality follows similarly. □

Lemma 2.9. *Let X and X' be two selfdual objects, and let $\alpha \in \text{Hom}_{\mathcal{C}}(X', X)$.*


- *If $(X, m, \delta, e, \epsilon)$ is a Frobenius algebra, then*



- *If $(X', m', \delta', e', \epsilon')$ is a Frobenius algebra, then*




Proof. We can prove the first equality using the following diagram:




The proof of the other equalities follows a similar approach. □

In the rest of the paper, we will refer to the application of Lemma 2.9 as *weak Frobenius*.

Lemma 2.10. *Let $(X, m, \delta, e, \epsilon)$ be a Frobenius algebra. Then $e^* = \epsilon$ and $\epsilon^* = e$, as depicted below:*




Proof. We can demonstrate the first equality using the following diagram:




The second equality follows similarly. □

Lemma 2.11. *The following equalities hold in a Frobenius algebra:*




Proof. This can be proven using the following diagram:




Lemma 2.12. *Let $(X, m, \delta, e, \epsilon)$ be a Frobenius algebra, and let α be a morphism in $\text{Hom}_{\mathcal{C}}(X, X)$. Then:*

$$m \circ (\alpha \otimes \text{id}_X) \circ \delta = ((e \circ m \circ (\alpha \otimes \text{id}_X) \circ \delta) \otimes \text{id}_X) \circ m,$$


which can be depicted as follows:



Proof. Here is a graphical proof:



Definition 2.13. *An algebra-coalgebra $(X, m, \delta, e, \epsilon)$ in a \mathbb{k} -linear abelian monoidal category is called connected (or haploid) if $\text{Hom}_{\mathcal{C}}(\mathbf{1}, X)$ is one-dimensional, specifically equal to $\mathbb{k}e$. It is termed separable if there exists $\lambda \in \mathbb{k}$ such that $m \circ \delta = \lambda \text{id}_X$, as illustrated below:*



Lemma 2.14. *A connected Frobenius algebra is separable.*

Proof. Note that $(m \circ \delta) \circ e \in \text{Hom}_{\mathcal{C}}(\mathbf{1}, X)$. By the connectedness condition, there exists $\lambda \in \mathbb{k}$ such that $(m \circ \delta) \circ e = \lambda e$, as shown below:

$$\text{Circle with dot on top and line down} = \lambda \text{ Vertical line with dot on top}$$

Applying Lemma 2.12 with $\alpha = \text{id}_X$, along with the previous equality and the unitality property, we obtain:

$$\text{Circle with dot on top and line down} = \text{Circle with dot on top and line down} = \lambda \text{ Vertical line with dot on top} = \lambda \text{ Cup shape with dot on top and line down}$$

Thus, the result follows. \square

If there is no possibility for confusion, a Frobenius algebra $(X, m, \delta, e, \epsilon)$ may simply be referred to as X .

Definition 2.15. *Let X be an object in a monoidal category \mathcal{C} . Assume that X has a left dual and a double dual, denoted X^* and X^{**} , respectively. The trace of a morphism $\alpha \in \text{Hom}_{\mathcal{C}}(X, X^{**})$, denoted $\text{tr}(\alpha)$, is defined as*

$$\text{tr}(\alpha) = \text{ev}_{X^*} \circ (\alpha \otimes \text{id}_{X^*}) \circ \text{coev}_X \in \text{Hom}_{\mathcal{C}}(\mathbf{1}, \mathbf{1}),$$

as illustrated below:

$$\text{tr}(\alpha) := \text{Box } \alpha \text{ (X above, X** below)} \text{ connected to } X^*$$

If X^{**} also has a left dual, then the following holds:

$$\text{Loop with } \alpha^* \text{ and } X^* \text{ boxes} = \text{Loop with } \alpha \text{ and } X^{**} \text{ boxes} = \text{Loop with } \alpha \text{ and } X^{**} \text{ boxes} = \text{tr}(\alpha)$$

If \mathcal{C} is a \mathbb{k} -linear abelian category and the unit object $\mathbf{1}$ is linear-simple (i.e., $\text{Hom}_{\mathcal{C}}(\mathbf{1}, \mathbf{1}) = \mathbb{k}$, as in a tensor category), then the elements of $\text{Hom}_{\mathcal{C}}(\mathbf{1}, \mathbf{1})$ can be interpreted as scalars. By Schur's lemma, if $\mathbf{1}$ is simple, then $\text{Hom}_{\mathbb{k}}(\mathbf{1}, \mathbf{1})$ forms a division algebra. Therefore, if \mathbb{k} is algebraically closed, then $\mathbf{1}$ is linear-simple.

Lemma 2.16. *Let A and B be two objects in a monoidal category \mathcal{C} such that $A^{**} = A$ and $B^{**} = B$. Consider two morphisms $f : A \rightarrow B$ and $g : B \rightarrow A$. Then, it holds that*

$$\text{tr}(g \circ f) = \text{tr}(f^{**} \circ g).$$

Proof. This can be proven pictorially:

$$\text{tr}(f^{**} \circ g) = \text{Loop with } g \text{ and } f^{**} \text{ boxes} = \text{Loop with } g \text{ and } f \text{ boxes} = \text{Loop with } g \text{ and } f \text{ boxes} = \text{Loop with } f \text{ and } g \text{ boxes} = \text{tr}(g \circ f)$$

\square

Let a be a pivotal structure on \mathcal{C} rigid monoidal, i.e. a collection of isomorphisms $a_X : X \rightarrow X^{**}$ such that $a_{X \otimes Y} = a_X \otimes a_Y$, and for all f in $\text{Hom}_{\mathcal{C}}(X, Y)$ then $f = a_Y^{-1} \circ f^{**} \circ a_X$. If $X = Y$, let us define $\text{tr}_a(f) := \text{tr}(a_X \circ f)$.

Lemma 2.17. *Let X and Y be two objects in a rigid monoidal category \mathcal{C} with pivotal structure a . Consider two morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Then, it holds that*

$$\text{tr}_a(g \circ f) = \text{tr}_a(f \circ g).$$

Proof. By Lemma 2.16 and the pivotal structure:

$$\text{tr}_a(g \circ f) = \text{tr}(a_X \circ g \circ f) = \text{tr}(f^{**} \circ a_X \circ g) = \text{tr}(a_Y \circ f \circ g) = \text{tr}_a(f \circ g). \quad \square$$


According to [36, Theorem 2.2], every pivotal monoidal category is equivalent, as a pivotal monoidal category, to a strictly pivotal monoidal category. In particular, we can assume, without loss of generality, that the pivotal structure (i.e., $a : ()^{**} \rightarrow \text{id}_{\mathcal{C}}$) is trivial (i.e., $a_X = \text{id}_X$).

Lemma 2.18. *Let $(X, m, \delta, e, \epsilon)$ be a Frobenius algebra. Then,*

$$\text{tr}(\text{id}_X) = \text{ev}_X \circ \text{coev}_X = \epsilon \circ m \circ \delta \circ e.$$

In the separable case, where $m \circ \delta = \lambda_X \text{id}_X$, and assuming that the unit object $\mathbf{1}$ is linear-simple, let μ_X denote the scalar defined by $\epsilon \circ e$. Then, we have $\text{tr}(\text{id}_X) = \lambda_X \mu_X$.


Proof. This follows immediately from Lemma 2.4. The relation is illustrated below:




Lemma 2.19. *Using the notation from Lemma 2.18, let $\alpha \in \text{Hom}_{\mathcal{C}}(X, X)$ and assume the connectedness condition. Then,*

$$\mu_X m \circ (\alpha \otimes \text{id}_X) \circ \delta = \text{tr}(\alpha) \text{id}_X,$$


as depicted below:



Proof. Here is a graphical proof:



Thus, we have $c_X \mu_X = \text{tr}(\alpha)$. Next, by applying Lemma 2.12 along with the preceding equalities, we obtain:



Proposition 2.20. *Let Y be a simple object with left dual Y^* and double dual Y^{**} in a \mathbb{k} -linear abelian monoidal category. Assume that $Y^{**} = Y$. Then, the object $X := Y \otimes Y^*$ can be equipped with a structure of connected Frobenius algebra, defined by $m = \text{id}_Y \otimes \text{ev}_Y \otimes \text{id}_{Y^*}$, $\delta = \text{id}_Y \otimes \text{coev}_{Y^*}$, $e = \text{coev}_Y$, and $\epsilon = \text{ev}_{Y^*}$, as depicted below:*

$$\begin{aligned}
m &= \begin{array}{c} X \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ X \end{array} := \begin{array}{c} Y \quad Y^* \quad Y \quad Y^* \\ | \quad | \quad | \quad | \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ | \quad | \quad | \quad | \\ Y \quad Y^* \quad Y \quad Y^* \end{array} = \text{id}_Y \otimes \text{ev}_Y \otimes \text{id}_{Y^*} \\
\delta &= \begin{array}{c} X \\ | \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ X \end{array} := \begin{array}{c} | \quad | \quad | \quad | \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ | \quad | \quad | \quad | \\ Y \quad Y^* \quad Y \quad Y^* \end{array} = \text{id}_Y \otimes \text{coev}_{Y^*} \otimes \text{id}_{Y^*} \\
e &= \begin{array}{c} 1 \\ \bullet \\ | \\ X \end{array} := \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ Y \quad Y^* \end{array} = \text{coev}_Y \\
\epsilon &= \begin{array}{c} X \\ | \\ \bullet \\ 1 \end{array} := \begin{array}{c} Y \quad Y^* \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} = \text{ev}_{Y^*}
\end{aligned}$$

Proof. The connectedness follows from the fact that Y is a simple object. We can verify all other properties graphically:

The same applies for coassociativity and counitality. □

Proposition 2.20 can be extended to 2-categories in the following manner:

Proposition 2.21. *Consider a 2-category \mathfrak{A} , with two objects, denoted as C and D (i.e., $\mathfrak{A}_0 = \{C, D\}$). Define \mathcal{C} and \mathcal{D} as the monoidal categories $\text{End}_{\mathfrak{A}}(C)$ and $\text{End}_{\mathfrak{A}}(D)$, respectively. Let \mathcal{M} represent the $(\mathcal{C}, \mathcal{D})$ -bimodule category, specifically $\text{Hom}_{\mathfrak{A}}(C, D)$. Suppose Y is a simple object in \mathcal{M} , and let Y^\vee in $\text{Hom}_{\mathfrak{A}}(D, C)$ be a two-sided dual of Y . Then the object $X := Y \circ Y^\vee$ in \mathcal{C} can be structured as a connected Frobenius algebra.*

Proof. The proof uses a pictorial approach similar to that in Proposition 2.20 (refer to [35, Lemma 3.4] for details), incorporating the natural adjunction isomorphism (Frobenius reciprocity) which also has a standard pictorial proof. □

Remark 2.22. *Note that the converse is also true by [35, Proposition 3.8].*

Example 2.23. *Let $(N \subset M)$ be a finite index inclusion of II_1 factors. Then $X = {}_N M_N$ is a Frobenius algebra in the tensor category \mathcal{C} of N - N -bimodules. This follows the structure outlined in Proposition 2.21, where \mathcal{D} is the tensor category of M - M -bimodules, $Y = {}_N M_M$, and $Y^\vee = {}_M M_N$, such that $X = Y \otimes_M Y^\vee$. Moreover, the subfactor $(N \subset M)$ is irreducible (i.e., $N' \cap M = \mathbb{C}$) if and only if the Frobenius algebra X is connected.*


3. FROBENIUS SUBALGEBRAS

Example 2.23 demonstrates how Theorem 1.2 extends Watatani’s theorem in subfactor theory. To generalize the concept of intermediate subfactors, it is essential to clarify what we mean by Frobenius subalgebras. First, let us recall why a morphism of Frobenius algebras is an isomorphism:

Definition 3.1. A morphism f between Frobenius algebras $(X, m, \delta, e, \epsilon)$ and $(X', m', \delta', e', \epsilon')$ in a monoidal category \mathcal{C} is called a Frobenius algebra morphism if it satisfies the following conditions:


- (Algebra morphism) $f \circ m' = m \circ (f \otimes f)$ and $e = e' \circ f$,
- (Coalgebra morphism) $\delta' \circ f = (f \otimes f) \circ \delta$ and $\epsilon = f \circ \epsilon'$.

These conditions can be illustrated as follows:



Lemma 3.2. Let (X, m, e) be an algebra in a monoidal category. If (X', m', e') is a triple along with a monomorphism $f : X' \rightarrow X$ that satisfies the conditions of an algebra morphism from Definition 3.1, then (X', m', e') is also an algebra.

Proof. To prove unitality, we begin with the following illustration:



Since the morphism f is left-cancellative (by the definition of a monomorphism), the result follows. The proof for associativity is analogous. \square


Lemma 3.3. Let (X, δ, ϵ) be a coalgebra in a monoidal category. If (X', δ', ϵ') is a triple along with an epimorphism $f : X \rightarrow X'$ that satisfies the conditions of a coalgebra morphism from Definition 3.1, then (X', δ', ϵ') is also a coalgebra.

Proof. The proof is similar to that of Lemma 3.2, as an epimorphism is defined to be right-cancellative. \square

The following is a well-known result:

Lemma 3.4. A Frobenius algebra morphism is an isomorphism.

Proof. Let f be a morphism as defined in Definition 3.1. The following illustration demonstrates that $f^* \circ f = \text{id}_X$:



Similarly, we can show that $f \circ f^* = \text{id}_{X'}$. Thus, the result follows. \square

Lemma 3.4 indicates that the concept of a Frobenius algebra morphism is too restrictive for defining the notion of a Frobenius subalgebra.


Definition 3.5. A Frobenius algebra $(X', m', \delta', e', \epsilon')$ is termed a Frobenius subalgebra of $(X, m, \delta, e, \epsilon)$ if there exists an algebra monomorphism $i : X' \rightarrow X$ such that $i^* \circ i = \text{id}_{X'}$ and $i^{**} = i$. We define $b_{X'} := i \circ i^* \in \text{Hom}_{\mathcal{C}}(X, X)$.

The morphism $b_{X'}$, as defined in Definition 3.5, generalizes the notion of a biprojection from the theory of subfactor planar algebras (see Proposition 4.15 and [32, §4]). In Definition 3.5, since $i^* \circ i = \text{id}_{X'}$, it follows that $i : X' \rightarrow X$ is a split algebra monomorphism, and $i^* : X \rightarrow X'$ is a split coalgebra epimorphism. Recall from Lemma 2.4 that both X and X' are selfdual.


Remark 3.6. Let A and B be two selfdual objects in an abelian monoidal category \mathcal{C} . Consider the monomorphism $i = \text{id}_A \oplus 0 \in \text{Hom}_{\mathcal{C}}(A \oplus 0, A \oplus B)$. In this case, i^* corresponds to the epimorphism $\text{id}_A \oplus 0 \in \text{Hom}_{\mathcal{C}}(A \oplus B, A \oplus 0)$, and we have $i^* \circ i = \text{id}_A$.

Note that X is a Frobenius subalgebra of itself with $i = i^* = \text{id}_X$. Furthermore, if the category is \mathbb{k} -linear and the unit object $\mathbf{1}$ is linear-simple, and if $\lambda_X := \epsilon \circ e$ is nonzero, then $\mathbf{1}$ is a Frobenius subalgebra of X with $i = \lambda_X^{-1/2} e$. This is because $i^* = \lambda_X^{-1/2} \epsilon$ as shown in Lemma 2.10. In this case, we have $b_X = \text{id}_X$ and $b_{\mathbf{1}} = \lambda_X^{-1} e \circ \epsilon$.

Theorem 3.7 (Exchange Relations). Let X and X' be two Frobenius algebras in a monoidal category \mathcal{C} . Let $f \in \text{Hom}_{\mathcal{C}}(X', X)$ be an algebra morphism such that $f^{**} = f$, and define $g := f \circ f^*$. The following relation holds:



Proof. First, we show that $f^* \in \text{Hom}_{\mathcal{C}}(X, X')$ is a coalgebra morphism. Since f is an algebra morphism, meaning $f \circ m = m \circ (f \otimes f)$, we have $(f \circ m)^* = (m \circ (f \otimes f))^*$, which can be reformulated as $\delta \circ f^* = (f^* \otimes f^*) \circ \delta$ (noting that $m^* = \delta$ by Lemma 2.8). Thus, f^* is confirmed as a coalgebra morphism. Next, we note that $g^* = g$ because $(f \circ f^*)^* = f^{**} \circ f^*$ and $f^{**} = f$. The rest of the proof follows from the illustration:




The second equality follows a similar reasoning. □

Corollary 3.8. Let X, X_1 , and X_2 be three Frobenius algebras in a \mathbb{k} -linear abelian monoidal category \mathcal{C} , where $\mathbf{1}$ is linear-simple. Let $f_i \in \text{Hom}_{\mathcal{C}}(X_i, X)$ be an algebra morphism such that $f_i^{**} = f_i$, and define $g_i := f_i \circ f_i^*$. If X is connected, then the following relation holds:

$$\mu_X(g_1 * g_2) \circ g_i = \text{tr}(g_1 \circ g_2) g_i = \mu_X g_i \circ (g_1 * g_2).$$

Proof. We first prove the left-hand side for $i = 2$ using the following illustration, applying the exchange relation (Theorem 3.7) and the connectedness condition (Lemma 2.19):



For the case when $i = 1$, the reasoning is similar. By Definition 2.15, we conclude with $\text{tr}((g_2 \circ g_1)^*) = \text{tr}(g_1 \circ g_2)$ since $g_i^* = g_i$. The second equality follows a similar pattern, resulting in $\text{tr}(g_2 \circ g_1) = \text{tr}(g_1 \circ g_2)$ by Lemma 2.16, given that $g_i^{**} = g_i$. □

Corollary 3.9. *Following the notations of Definition 3.5, the morphism $b_{X'}$ satisfies the exchange relations. Moreover, if X_1 and X_2 are Frobenius subalgebras of X in a \mathbb{k} -linear abelian monoidal category \mathcal{C} where $\mathbf{1}$ is linear-simple, then:*

$$\mu_X(b_{X_1} * b_{X_2}) \circ b_{X_i} = \text{tr}(b_{X_1} \circ b_{X_2}) b_{X_i}.$$

Proof. The first part is immediate from Theorem 3.7 with $f = i$. For the second part, we can apply Corollary 3.8 with $g_i = b_{X_i}$. \square

We need to expand the concept of the convolution product.

Definition 3.10 (Convolution Product). *Let $(X, m, \delta, e, \epsilon)$ be an algebra-coalgebra in a monoidal category \mathcal{C} . For two morphisms $a, b \in \text{Hom}_{\mathcal{C}}(X, X)$, we define the convolution product as $a * b := \delta \circ (a \otimes b) \circ m$, illustrated as follows:*

$$a * b := \delta \circ (a \otimes b) \circ m = \begin{array}{c} \text{---} \\ | \\ \text{---} \circlearrowleft \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} \begin{array}{c} \boxed{a} \\ \boxed{b} \end{array}$$

Note that Lemma 2.19 can be restated as $\mu_X(\alpha * \text{id}_X) = \text{tr}(\alpha) \text{id}_X$.

Remark 3.11 (Fourier Transform). *Although it may not be utilized in this paper, it is noteworthy that the convolution product defined in Definition 3.10 can be derived from the Fourier transform:*

$$\begin{array}{ccc} \mathcal{F}: \text{Hom}_{\mathcal{C}}(X, X) & \rightarrow & \text{Hom}_{\mathcal{C}}(X \otimes X, X \otimes X) \\ a & \mapsto & (\text{id}_X \otimes m) \circ (\text{id}_X \otimes a \otimes \text{id}_X) \circ (\delta \otimes \text{id}_X) \end{array}$$

with a left inverse given by $\mathcal{F}^{-1}(x) = (\epsilon \otimes \text{id}_X) \circ x \circ (\text{id}_X \otimes \epsilon)$. These transformations are depicted below:

$$\mathcal{F}(a) = \begin{array}{c} \text{---} \\ | \\ \text{---} \circlearrowleft \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} \boxed{a} \quad \mathcal{F}^{-1}(x) = \begin{array}{c} \bullet \\ | \\ \boxed{x} \\ | \\ \bullet \end{array}$$

It can be shown pictorially that $a * b = \mathcal{F}^{-1}(\mathcal{F}(a) \circ \mathcal{F}(b))$; refer to [6, Proposition 1] for further details.

Proposition 3.12. *Let X and X' be two algebras-coalgebras in a \mathbb{k} -linear abelian monoidal category \mathcal{C} , with the assumption that X' is separable. Let $f \in \text{Hom}_{\mathcal{C}}(X', X)$ be an algebra morphism and $h \in \text{Hom}_{\mathcal{C}}(X, X')$ a coalgebra morphism. Define $g := f \circ h$. Then, it holds that*

$$g * g = \lambda_{X'} g,$$

where $\lambda_{X'}$ is the constant associated with the separability of X' (see Definition 2.13).

Moreover, if both X and X' are Frobenius, X is connected, and $\mathbf{1}$ is linear-simple, with $h = f^*$, $h^* = f$, and $g = f \circ f^*$ being a nonzero idempotent (e.g., $f^* \circ f = \text{id}_{X'}$), then

$$\text{tr}(g) = \mu_X \lambda_{X'},$$


where μ_X is the scalar defined by $\epsilon \circ e$. Consequently, we have

$$\mu_X(g * g) = \text{tr}(g)g.$$


Proof. The first part is illustrated by the following diagram:

$$g * g = \begin{array}{c} \text{---} \\ | \\ \text{---} \circlearrowleft \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} \begin{array}{c} \boxed{g} \\ \boxed{g} \end{array} = \begin{array}{c} \text{---} \\ | \\ \text{---} \circlearrowleft \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} \begin{array}{c} \boxed{h} \\ \boxed{h} \\ \boxed{f} \\ \boxed{f} \end{array} \begin{array}{c} \text{coalgebra map} \\ \downarrow \\ \text{---} \\ | \\ \text{---} \circlearrowleft \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} \begin{array}{c} \boxed{h} \\ \boxed{f} \end{array} \begin{array}{c} \text{separability} \\ \downarrow \\ \text{---} \\ | \\ \text{---} \circlearrowleft \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} \begin{array}{c} \boxed{h} \\ \boxed{f} \end{array} = \lambda_{X'} f \circ h = \lambda_{X'} g$$

For the second part, define $A \in \text{Hom}_{\mathcal{C}}(X, X \otimes X)$ as follows:



We observe that



By the first part and Lemma 2.19, we obtain

$$\mu_X A^*A = 2(\mu_X \lambda_{X'} - \text{tr}(g))g,$$

but $A = 0$ by Theorem 3.7. The result follows. \square

Corollary 3.13. *Using the notation from Definition 3.5, the following holds in the connected case if $\mathbf{1}$ is linear-simple:*

$$\text{tr}(b_{X'}) = \mu_X \lambda_{X'} \quad \text{and} \quad \mu_X(b_{X'} * b_{X'}) = \text{tr}(b_{X'})b_{X'}.$$

Remark 3.14 ([5, 30, 32]). *Theorem 3.7 and Proposition 3.12 align with the theory of irreducible finite index subfactors ($N \subset P \subset M$), where $X = {}_N M_N$, $X' = {}_N P_N$, $g = b_{X'} = e_P^M$, and $\lambda_X = \mu_X = |M : N|^{1/2}$. It is important to note that the trace tr here is not normalized, in contrast to its typical normalization within the subfactor framework.*

The following result generalizes Landau's theorem (e.g., see [19, Theorem 3.10]):

Theorem 3.15 (Landau's theorem). *Let X , X_1 , and X_2 be three Frobenius algebras in a \mathbb{k} -linear abelian monoidal category \mathcal{C} with $\mathbf{1}$ being linear-simple. Assume that X is connected. For $i = 1, 2$, let $f_i \in \text{Hom}_{\mathcal{C}}(X_i, X)$ be an algebra morphism such that $f_i^{**} = f_i$, and let $g_i := f_i \circ f_i^*$. Then*

$$\mu_X(g_1 * g_2) \circ (g_1 * g_2) = \text{tr}(g_2 \circ g_1)(g_1 * g_2).$$

Furthermore,

$$\mu_X \text{tr}(g_1 * g_2) = \text{tr}(g_1)\text{tr}(g_2).$$

As a result, if $\text{tr}(g_2 \circ g_1)$ is nonzero, then


$$g_{12} := \frac{\mu_X}{\text{tr}(g_2 \circ g_1)} g_1 * g_2$$

is an idempotent, and we have

$$\text{tr}(g_{12}) = \frac{\text{tr}(g_1)\text{tr}(g_2)}{\text{tr}(g_2 \circ g_1)}.$$

Finally, $g_{12} \circ g_i = g_i = g_i \circ g_{12}$ for $i \in \{1, 2\}$.


Proof. The proof begins with the diagram below, utilizing Lemmas 2.9, 2.11, and Theorem 3.7:



Next, we apply Lemma 2.19 to conclude that

$$\mu_X(g_1 * g_2) \circ (g_1 * g_2) = \text{tr}(g_2 \circ g_1)(g_1 * g_2).$$

The following diagram illustrates the proof of the second equality, where *connectedness* is invoked according to Lemma 2.19:



The final equality follows from Corollary 3.8. \square

Corollary 3.16. *Let X be a connected Frobenius algebra in a \mathbb{k} -linear abelian monoidal category \mathcal{C} with $\mathbf{1}$ being linear-simple. Let X' and X'' be two Frobenius subalgebras of X . Using the notation from Definition 3.5, we have*

$$\mu_X(b_{X'} * b_{X''}) \circ (b_{X'} * b_{X''}) = \text{tr}(b_{X''} \circ b_{X'})(b_{X'} * b_{X''}).$$

Additionally,

$$\mu_X \text{tr}(b_{X'} * b_{X''}) = \text{tr}(b_{X'}) \text{tr}(b_{X''}).$$

Consequently, if $\text{tr}(b_{X''} \circ b_{X'})$ is nonzero, then

$$b_{X'X''} := \frac{\mu_X}{\text{tr}(b_{X''} \circ b_{X'})} b_{X'} * b_{X''}$$

is an idempotent, and therefore

$$\text{tr}(b_{X'X''}) = \frac{\text{tr}(b_{X'}) \text{tr}(b_{X''})}{\text{tr}(b_{X''} \circ b_{X'})}.$$

Finally, $b_{X'X''} \circ b_Y = b_Y = b_Y \circ b_{X'X''}$ for all $Y \in \{X', X''\}$.

4. LATTICE STRUCTURE

We will define a lattice structure on the set of Frobenius subalgebras (up to equivalence) of a Frobenius algebra, semisimple as an object in an abelian monoidal category. However, without the semisimplicity assumption, we will establish only a poset structure. We begin by introducing the equivalence relation and the partial order, following [15, §1.5]. Two monomorphisms $i_1 : A_1 \rightarrow C$ and $i_2 : A_2 \rightarrow C$ are deemed *equivalent* if there exist morphisms $i_{1,2} : A_1 \rightarrow A_2$ and $i_{2,1} : A_2 \rightarrow A_1$ such that

$$i_1 = i_2 \circ i_{1,2} \quad \text{and} \quad i_2 = i_1 \circ i_{2,1}.$$

A *subobject* of C is an equivalence class of monomorphisms into C . We define the subobject represented by $j_1 : B_1 \rightarrow C$ to be *contained* in the subobject represented by $j_2 : B_2 \rightarrow C$ if there exists a morphism $j_{1,2} : B_1 \rightarrow B_2$ such that $j_1 = j_2 \circ j_{1,2}$. This containment relation defines a partial order on subobjects. When no confusion arises, we will simply write $B_1 \subseteq B_2$.

Next, following [15, §2.1], the *intersection* of two subobjects of C is defined as their greatest lower bound in the family of subobjects of C , which always exists according to [15, Theorem 2.13]. This intersection is represented by the pullback of their representatives $i_A : A \rightarrow C$ and $i_B : B \rightarrow C$.

$$\begin{array}{ccc} D & \xrightarrow{j_A} & A \\ j_B \downarrow & & \downarrow i_A \\ B & \xrightarrow{i_B} & C \end{array}$$

We denote a pullback generically as $A \times_C B$, but the pullback of monomorphisms can simply be denoted as $A \cap B$, with the monomorphism $i_A \circ j_A = i_B \circ j_B$ represented as $i_{A \cap B}$ (noting that a pullback of monomorphisms is also a monomorphism). The question arises as to whether the structure of a Frobenius subalgebra is preserved under intersection (we will later show that it is not always the case).

Lemma 4.1. *Let (A, i_A) and (B, i_B) be two subobject representatives of an object C in an abelian category. Consider another object C' with monomorphisms $i : C \rightarrow C'$ and $i'_Z : Z \rightarrow C'$ such that $i'_Z = i \circ i_Z$ for all $Z \in \{A, B\}$. Then, (A, i'_A) and (B, i'_B) are subobject representatives of C' , and the pullback of (i_A, i_B) can be taken to be the same as that of (i'_A, i'_B) .*

$$\begin{array}{ccccc}
 A \cap B & \xrightarrow{j_A} & A & & \\
 j_B \downarrow & & \downarrow i_A & \searrow i'_A & \\
 B & \xrightarrow{i_B} & C & \xrightarrow{i} & C' \\
 & \searrow i'_B & & & \\
 & & & &
 \end{array}$$

Proof. This follows directly from [15], which states that the pullback of monomorphisms corresponds to the intersection defined as the greatest lower bound in the subobject poset of C or C' . The relationship $Z \subseteq C$ as subobjects of C' holds as $i'_Z = i \circ i_Z$ for all $Z \in \{A, B\}$. \square

Remark 4.2. *According to [15], the union of two subobjects A and B of C is defined as their least upper bound in the subobject poset of C , which always exists and is represented by the pushout of the morphisms j_A and j_B .*

$$\begin{array}{ccccc}
 A \cap B & \xrightarrow{j_A} & A & & \\
 j_B \downarrow & & \downarrow k_A & \searrow i_A & \\
 B & \xrightarrow{k_B} & A \cup B & & C \\
 & \searrow i_B & & &
 \end{array}$$

Furthermore, the subobject poset of C has a lattice structure with respect to \cap and \cup . In an abelian category, the union $A \cup B$ is more appropriately denoted as $A + B$.

Roughly speaking, the following theorem states that the intersection of subalgebras is a subalgebra. The proof in a general abelian monoidal category is partly attributed to Brian Shin [54].

Proposition 4.3. *Let $L \in \{A, B, C\}$, and let (L, m_L, e_L) be algebras in an abelian monoidal category, along with algebra monomorphisms $i_L : L \rightarrow C$ for $L \in \{A, B\}$. Then there exists a unique algebra structure on the pullback $A \cap B$ such that the monomorphisms $j_L : A \cap B \rightarrow L$ are algebra morphisms.*

Proof. For $L \in \{A, B\}$, consider the morphisms

$$f_L := i_L \circ m_L \circ (j_L \otimes j_L) : (A \cap B) \otimes (A \cap B) \rightarrow C.$$

Since i_L is an algebra morphism, we have $i_L \circ m_L = m \circ (i_L \otimes i_L)$. Therefore,

$$f_L = m \circ (i_L \otimes i_L) \circ (j_L \otimes j_L) = m \circ ((i_L \circ j_L) \otimes (i_L \circ j_L)).$$

However, $i_A \circ j_A = i_B \circ j_B$ by the definition of the pullback, so $f_A = f_B$. By universality, there exists a unique map

$$m_{A \cap B} : (A \cap B) \otimes (A \cap B) \rightarrow A \cap B$$

such that

$$j_L \circ m_{A \cap B} = m_L \circ (j_L \otimes j_L).$$

Here is the corresponding diagram:

$$\begin{array}{ccccc}
 (A \cap B) \otimes (A \cap B) & \xrightarrow{j_A \otimes j_A} & A \otimes A & & \\
 \downarrow j_B \otimes j_B & \searrow \exists! m_{A \cap B} & \downarrow m_A & & \\
 & & A \cap B & \xrightarrow{j_A} & A \\
 & & \downarrow j_B & & \downarrow i_A \\
 B \otimes B & \xrightarrow{m_B} & B & \xrightarrow{i_B} & C
 \end{array}$$

Similarly, for $L \in \{A, B\}$, since i_L is an algebra morphism, we have $i_A \circ e_A = e_C = i_B \circ e_B$. By universality, there exists a unique $e_{A \cap B}$ such that $j_L \circ e_{A \cap B} = e_L$. The corresponding diagram is as follows:

$$\begin{array}{ccccc}
 \mathbb{1} & & & & \\
 \downarrow e_B & \searrow \exists! e_{A \cap B} & & \xrightarrow{e_A} & \\
 & & A \cap B & \xrightarrow{j_A} & A \\
 & & \downarrow j_B & & \downarrow i_A \\
 & & B & \xrightarrow{i_B} & C
 \end{array}$$

The result follows from Lemma 3.2. \square

The following lemma and the proof of Lemma 4.5 are partly due to Maxime Ramzi [47].

Lemma 4.4. *A square $A_0 \rightarrow (A_1, A_2) \rightarrow A_3$ with $A_0 \rightarrow A_1$ and $A_2 \rightarrow A_3$ being monomorphisms is a pullback square if and only if the induced morphism $A_1/A_0 \rightarrow A_3/A_2$ is a monomorphism.*

Proof. The map $A_1 \times_{A_3} A_2 \rightarrow A_1$ is a monomorphism because it is pulled back from a monomorphism, and the inclusion $A_0 \rightarrow A_1$ factors through it. Therefore, the map $A_1/A_0 \rightarrow A_3/A_2$ factors through $A_1/A_0 \rightarrow A_1/(A_1 \times_{A_3} A_2)$. For the total composite to be a monomorphism, this must also be a monomorphism, thus proving the claim. \square

The following lemma extends [50, Chapter 14, Exercises 10-12]:

Lemma 4.5. *Let A, B, C, D, E, F be objects in an abelian rigid monoidal category.*

- (1) *If $A, B \subseteq C$, then $(A \otimes D) \cap (B \otimes D) \simeq (A \cap B) \otimes D$,*
- (2) *If $A \subseteq C$ and $B \subseteq D$, then $(A \otimes D) \cap (C \otimes B) \simeq (A \otimes B)$,*
- (3) *If $A, B \subseteq C$ and $D, E \subseteq F$, then $(A \otimes D) \cap (B \otimes E) \simeq (A \cap B) \otimes (D \cap E)$.*

Proof. By [12, Exercise 1.6.4 and Proposition 2.10.8], the functor $- \otimes D$ is exact in an abelian rigid monoidal category. The intersection is defined as a pullback of monomorphisms. Since a pullback is a finite limit and an exact functor preserves finite limits, the result in (1) follows.

For (2), $X \otimes -$ preserves monomorphisms for all X , hence the square $A \otimes B \rightarrow (A \otimes D, C \otimes B) \rightarrow C \otimes D$ satisfies Lemma 4.4. In particular, a tensor product of monomorphisms is a monomorphism (i.e., $A \otimes B \subseteq C \otimes D$).

For (3), applying (2) yields that

$$(A \otimes D) \cap (B \otimes E) = (A \otimes F) \cap (C \otimes D) \cap (B \otimes F) \cap (C \otimes E).$$

Next, applying (1) with $- \otimes F$ and $C \otimes -$, along with the commutativity of intersections, gives us $((A \cap B) \otimes F) \cap (C \otimes (D \cap E))$. We then apply (2) again to obtain $(A \cap B) \otimes (D \cap E)$. \square

Remark 4.6. *We deduce from the proof of Lemma 4.5 that in an abelian rigid monoidal category, the tensor product of two monomorphisms is a monomorphism.*

The following result does not hold without the semisimple assumption. Dave Benson provided a counterexample in Vec , available at [3].

Proposition 4.7. *Let A, B, C be objects in an abelian rigid monoidal category with $A, B \subseteq C$ and C semisimple. If A and B are selfdual, then so is $A \cap B$.*

Proof. Let (X_i) be the simple objects up to isomorphism. Without loss of generality, we can take $C = \bigoplus_i C_i \otimes X_i$, $A = \bigoplus_i A_i \otimes X_i$, and $B = \bigoplus_i B_i \otimes X_i$, where A_i and B_i are subspaces of the multiplicity space C_i for all i . We can also take i_A and i_B induced by the inclusions $A_i, B_i \subseteq C_i$. Then, $A \cap B = \bigoplus_i (A_i \cap B_i) \otimes X_i$. Since A, B , and C are selfdual, we have $C_{i^*} = C_i$, $A_{i^*} = A_i$, and $B_{i^*} = B_i$. Thus, for all i ,

$$(A \cap B)^* = \bigoplus_i (A_i \cap B_i) \otimes X_i^* = \bigoplus_i (A_{i^*} \cap B_{i^*}) \otimes X_i = \bigoplus_i (A_i \cap B_i) \otimes X_i = (A \cap B). \quad \square$$


Remark 4.8. *Following the proof of Proposition 4.7, j_A is simply the monomorphism induced by the inclusions $(A_i \cap B_i) \subseteq A_i$. Thus, $j_A^* \circ j_A = \text{id}_{A \cap B}$ and $j_A^{**} = j_A$.*

Corollary 4.9. *Following Proposition 4.3 in the rigid case, for all $K \in \{A, B, A \cap B\}$, if K is selfdual, then $\delta_K = m_K^*$ and $\epsilon_K = e_K^*$ define a coalgebra structure such that j_L^* is a coalgebra epimorphism for all $L \in \{A, B\}$.*


Proof. This follows immediately from Proposition 4.3, since the dual of a pullback of monomorphisms is a pushout of epimorphisms. \square

Proposition 4.10. *In an abelian rigid monoidal category, let A and B be two Frobenius subalgebras of C such that $A \cap B$ is self-dual and $j_A^* \circ j_A = \text{id}_{A \cap B}$. Then $A \cap B$ admits a structure of Frobenius subalgebra of C .*


Proof. The results regarding the algebra and coalgebra structure on $A \cap B$ follow from Lemma 2.4, Proposition 4.3, and Corollary 4.9. The Frobenius structure is illustrated by the following diagrams. By assumption, $j_A^* \circ j_A = \text{id}_{A \cap B}$.



Since (j_A, j_B) is a pullback of (i_A, i_B) , it follows that (j_A^{**}, j_B^{**}) is a pullback of (i_A^{**}, i_B^{**}) . By Definition 3.5, we have $(i_A^{**}, i_B^{**}) = (i_A, i_B)$, so by universality, there is an isomorphism $\phi \in \text{Hom}_C(A \cap B, A \cap B)$ such that $j_A = j_A^{**} \circ \phi$. We conclude that



Consequently, we obtain



This implies $\phi = \text{id}_{A \cap B}$, demonstrating that the Frobenius property holds for $A \cap B$ (recall Lemma 2.7), and also that $j_A^{**} = j_A$. We have established that $j_A^* \circ j_A = \text{id}_{A \cap B}$. By Definition 3.5, we find $i_A^{**} = i_A$ and $i_A^* \circ i_A = \text{id}_A$. Thus, $i_{A \cap B}^{**} = i_{A \cap B}$ and $i_{A \cap B}^* \circ i_{A \cap B} = \text{id}_{A \cap B}$, where $i_{A \cap B} := i_A \circ j_A : A \cap B \rightarrow C$ is an algebra monomorphism, since both i_A and j_A are so (a pullback of a monomorphism is a monomorphism). \square

The following result does not hold without the semisimple assumption. Dave Benson provided a counterexample in Vec , available at [4].

Proposition 4.11. *In an abelian rigid monoidal category, let A and B be two Frobenius subalgebras of C , which is semisimple as an object. Then $A \cap B$ admits a structure of Frobenius subalgebra of C .*

Proof. Immediate by Proposition 4.7 and Proposition 4.10. Recall from Remark 4.8 that $j_A^* \circ j_A = \text{id}_{A \cap B}$. \square

Let us recall the definition of a lattice:

Definition 4.12. A meet-semilattice (L, \wedge) is a partially ordered set (poset) L in which every pair of elements a and b has a unique infimum (or meet) denoted by $a \wedge b$. A join-semilattice (L, \vee) is a poset L in which every pair of elements a and b has a unique supremum (or join) denoted by $a \vee b$. A lattice (L, \wedge, \vee) is a poset that is both a meet-semilattice and a join-semilattice.

Recall that the *height* of a poset is defined as the length of its longest chain, denoted as $\cdots < a_1 < a_2 < \cdots$.

Lemma 4.13. A meet-semilattice with a top element and finite height is a lattice.

Proof. Given the existence of a top element and finite height, any two elements a and b have a minimal upper bound. If there were two distinct minimal upper bounds c and d , then $c \wedge d$ would also serve as a minimal upper bound, leading to a contradiction. \square

The equivalence classes (and the poset structure) will be defined similarly to the subobjects described in [15, §1.5], as reviewed at the beginning of §4.

Definition 4.14. Following the notation from Definition 3.5, two Frobenius subalgebras X_1 and X_2 of X are considered equivalent if there exists an isomorphism $i_{1,2} : X_1 \rightarrow X_2$ such that $i_1 = i_2 \circ i_{1,2}$ and $i_2 = i_1 \circ i_{1,2}^*$, along with $i_{1,2}^* \circ i_{1,2} = \text{id}_{X_1}$ and $i_{1,2} \circ i_{1,2}^* = \text{id}_{X_2}$. We denote the equivalence class by $[X_1]$.

We can alternatively define a subalgebra as an equivalence class, similar to the concept of a subobject, to avoid confusion with isomorphism classes. The equivalence classes defined in Definition 4.14 are more restrictive than typical isomorphism classes. In particular, these equivalence classes correspond to generalized biprojections, as demonstrated in the following proposition:

Proposition 4.15. Using the notation from Definitions 3.5 and 4.14, the equivalence classes satisfy $[X_1] = [X_2]$ if and only if $b_{X_1} = b_{X_2}$.

Proof. Recall that $b_{X_k} = i_k \circ i_k^*$. If X_1 and X_2 belong to the same class, then:

$$b_{X_1} = i_1 \circ i_1^* = (i_2 \circ i_{1,2}) \circ (i_{1,2}^* \circ i_2^*) = i_2 \circ (i_{1,2} \circ i_{1,2}^*) \circ i_2^* = i_2 \circ i_2^* = b_{X_2}.$$

Conversely, if $b_{X_1} = b_{X_2}$, set $i_{1,2} = i_2^* \circ i_1$. Then:

$$i_2 \circ i_{1,2} = i_2 \circ i_2^* \circ i_1 = b_{X_2} \circ i_1 = b_{X_1} \circ i_1 = i_1 \circ i_1^* \circ i_1 = i_1,$$

and similarly $i_1 \circ i_{1,2}^* = i_2$. Finally,

$$i_{1,2} \circ i_{1,2}^* = i_2^* \circ i_1 \circ i_1^* \circ i_2 = i_2^* \circ b_{X_1} \circ i_2 = i_2^* \circ b_{X_2} \circ i_2 = \text{id}_{X_2},$$

and likewise $i_{1,2}^* \circ i_{1,2} = \text{id}_{X_1}$. \square

Definition 4.16. An equivalence class $[X_1]$ is said to be contained in $[X_2]$ if there exists a monomorphism $i_{1,2} : X_1 \rightarrow X_2$ such that $i_1 = i_2 \circ i_{1,2}$ and $i_{1,2}^* \circ i_{1,2} = \text{id}_{X_1}$.

Lemma 4.17. The containment relation defined in Definition 4.16 forms a partial order.

Proof. Assume that $[X_1]$ is contained in $[X_2]$ and $[X_2]$ is contained in $[X_1]$. Then $i_{1,2}$ must be an isomorphism with inverse $i_{1,2}^*$, leading to the conclusion that $[X_1] = [X_2]$. The remaining properties of a partial order are straightforward to verify. \square

Definition 4.18. The poset established in Lemma 4.17 will be referred to as the Frobenius subalgebra poset of the Frobenius algebra X .

The following result fails without the assumption of semisimplicity. A counterexample provided by Will Sawin is available in Vec at [51].

Theorem 4.19. The Frobenius subalgebra poset of a Frobenius algebra C of finite length and semisimple as an object in an abelian rigid monoidal category is a lattice. It can be identified with the interval $[b_1, b_C]$.

Proof. Lemma 4.17 establishes the poset structure. By Proposition 4.11, this poset forms a meet-semilattice with $[X_1] \wedge [X_2] := [X_1 \cap X_2]$, where the Frobenius algebra acts as the top element. The finite length of the Frobenius algebra ensures that the poset has finite height. Thus, the result follows from Lemma 4.13. The final assertion follows from Proposition 4.15. \square

In the case of connected Frobenius algebras, the lattice in Theorem 4.19 is finite, as stated in Theorem 1.2 and proved in §8.

5. KAROUBIAN CATEGORIES

A preadditive category \mathcal{C} is called *Karoubian* (or pseudo-abelian) if it is idempotent-complete. This means that for every object X in \mathcal{C} and every morphism b in $\text{Hom}_{\mathcal{C}}(X, X)$ that satisfies $b \circ b = b$ (i.e., b is idempotent), there exists an object Y in \mathcal{C} , a monomorphism $i : Y \rightarrow X$, and an epimorphism $p : X \rightarrow Y$ such that $i \circ p = b$ and $p \circ i = \text{id}_Y$. Notably, every abelian category is Karoubian. Typically, the idempotents in $\text{Hom}_{\mathcal{C}}(X, X)$ are partially ordered, defined by $b \leq b'$ if and only if $b \circ b' = b = b' \circ b$.

Lemma 5.1. *In a Karoubian category, using the notations described above, $b \leq b'$ if and only if there exists a monomorphism $i_{Y, Y'} : Y \rightarrow Y'$ and an epimorphism $p_{Y', Y} : Y' \rightarrow Y$ such that $i = i' \circ i_{Y, Y'}$ and $p_{Y', Y} \circ p' = p$, satisfying $p_{Y', Y} \circ i_{Y, Y'} = \text{id}_Y$. Consequently, $Y' \cong Y \oplus Z$ for some object Z .*

Proof. By definition, $b \leq b'$ if and only if $b \circ b' = b = b' \circ b$. This is equivalent to $i \circ p \circ i' \circ p' = i \circ p = i' \circ p' \circ i \circ p$. Hence, we have $p \circ i' \circ p' = p$ and $i = i' \circ p' \circ i$, where i is left-cancellative and p is right-cancellative. Therefore, this holds if and only if $p_{Y', Y} \circ p' = p$ and $i = i' \circ i_{Y, Y'}$, since by applying $_ \circ i'$ or $p' \circ _$, we find $p_{Y', Y} = p \circ i'$ and $i_{Y, Y'} = p' \circ i$. Consequently, $p \circ i' \circ p' = p$ implies $p \circ i' \circ p' \circ i = \text{id}_Y$, and we have $p \circ i' \circ p' \circ i = p_{Y', Y} \circ i_{Y, Y'}$. The last sentence follows from the splitting lemma. \square

Definition 5.2. *An object X in a \mathbb{C} -linear abelian monoidal category \mathcal{C} with a linear-simple unit $\mathbf{1}$ is called totally positive (or totally nonzero) if there exists an isomorphism $\phi \in \text{Hom}_{\mathcal{C}}(X, X^{**})$ such that for all nonzero idempotent morphisms $b \in \text{Hom}_{\mathcal{C}}(X, X)$, the trace $\text{tr}_{\phi}(b) := \text{tr}(\phi \circ b)$, as defined in Definition 2.15, is positive (or nonzero). If ϕ is not specified, it implies that $X = X^{**}$ and $\phi = \text{id}_X$.*

Lemma 5.3. *Let X be a totally positive object with isomorphism ϕ . For all nonzero idempotents $f, g \in \text{Hom}_{\mathcal{C}}(X, X)$ such that $g \leq f$, it follows that $\text{tr}_{\phi}(g) \leq \text{tr}_{\phi}(f)$. Moreover, if $f \neq g$, then $\text{tr}_{\phi}(g) < \text{tr}_{\phi}(f)$.*

Proof. Recall that $g \leq f$ if and only if $g \circ f = g = g \circ f$. By the linearity of the trace, we have:

$$\text{tr}_{\phi}(f) = \text{tr}_{\phi}(g) + \text{tr}_{\phi}(f - g).$$

But $f - g$ is an idempotent because:

$$(f - g) \circ (f - g) = (f \circ f) - (f \circ g) - (g \circ f) + (g \circ g) = f - g - g + g = f - g.$$

Thus, the result follows from the assumption that X is totally positive. \square

Definition 5.4. *Recall the notion of pivotal structure a and trace tr_a above Lemma 2.17. A finite tensor category is called positive if it has a pivotal structure a such that $\text{tr}_a(\text{id}_X) = \text{FPdim}(X)$.*

Proposition 5.5. *If \mathcal{C} is a positive finite tensor category, then every object X in \mathcal{C} is totally positive.*

Proof. Let (a_X) be the pivotal structure that ensures positivity. For any nonzero idempotent $b \in \text{Hom}_{\mathcal{C}}(X, X)$, consider $b = i \circ p$ and $p \circ i = \text{id}_Y$ from the Karoubian structure. By Lemma 2.17 and positivity, we have:

$$\text{tr}_a(b) = \text{tr}_a(i \circ p) = \text{tr}_a(p \circ i) = \text{tr}_a(\text{id}_Y) = \text{FPdim}(Y) > 0. \quad \square$$

Proposition 5.6. *A finite tensor category is positive if and only if it is a pseudo-unitary fusion category.*

Proof. According to [12, Proposition 9.5.1], a pseudo-unitary fusion category is positive. Conversely, if \mathcal{C} is a pivotal non-semisimple finite tensor category, then as shown in the proof of [12, Theorem 6.6.1], if P is projective in \mathcal{C} , then $\text{tr}_a(\text{id}_P) = 0$. Hence, \mathcal{C} cannot be positive. Furthermore, a positive fusion category is intrinsically pseudo-unitary. \square

While every unitary fusion category is inherently pseudo-unitary, it remains an open question whether every pseudo-unitary fusion category can be endowed with a unitary structure (see [12, §9.4]).

6. SEMISIMPLIFICATION

The concept of semisimplification for spherical tensor categories was introduced in [2]. In this subsection, we refer to [13]. Recall that a morphism $f : X \rightarrow Y$ in a spherical tensor category \mathcal{C} over a field \mathbb{k} is termed *negligible* if, for any morphism $g : Y \rightarrow X$, the trace $\text{tr}(f \circ g) = 0$. The collection \mathcal{N} of negligible morphisms forms a tensor ideal, allowing us to define $\bar{\mathcal{C}} = \mathcal{C}/\mathcal{N}$ as a semisimple tensor category. In this new category, the morphism compositions and tensor products remain unchanged from those in \mathcal{C} , while the simple objects consist of the indecomposable objects of \mathcal{C} that have nonzero dimension. Objects with zero dimension map to the zero object. This process is referred to as the *semisimplification* of \mathcal{C} .

Remark 6.1. *This result can be extended to \mathbb{k} -linear Karoubian pivotal rigid monoidal categories, where all morphism spaces are finite-dimensional, and with a pivotal structure such that the trace of a nilpotent endomorphism is zero, and the left and right dimensions of indecomposable objects vanish simultaneously (see [13, Theorem 2.6]).*

We refer to [16, Definition 3.4] for the following definition.

Definition 6.2. *Let $(X, m, \delta, e, \epsilon)$ be an algebra-coalgebra in a \mathbb{k} -linear Karoubian monoidal category \mathcal{C} . It is called special if $\epsilon \circ e = \alpha_X \text{id}_{\mathbf{1}}$ and $m \circ \delta = \beta_X \text{id}_X$, where α_X and β_X are nonzero scalars. This is illustrated as follows:*

$$\begin{array}{c} \bullet \\ | \\ X \\ | \\ \bullet \end{array} = \alpha_X \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \quad \text{and} \quad \begin{array}{c} | \\ \bigcirc \\ | \\ X \end{array} = \beta_X \begin{array}{c} | \\ | \\ | \end{array}$$

Note that a special algebra-coalgebra is inherently separable (see Definition 2.13). The term *connected* in Definition 2.13 is referred to as *haploid* in [16]. According to [16, Corollary 3.10 and (3.50)], for any connected special Frobenius algebra, it holds that $\alpha_X \beta_X = \dim(X)$ is nonzero. The image below provides a proof that does not rely on the connected assumption, but the alignment $\text{ev}_X = \epsilon \circ m$ and $\text{coev}_X = \delta \circ e$, see Remark 2.5.

$$\dim(X) = X \begin{array}{c} \bigcirc \\ | \\ \bullet \end{array} = X \begin{array}{c} \bullet \\ | \\ \bigcirc \\ | \\ \bullet \end{array} = \beta_X \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ X \end{array} = \alpha_X \beta_X$$

Furthermore, we can, without loss of generality, assume that $\alpha_X = \dim(X)$ and $\beta_X = 1$.

Lemma 6.3. *A separable Frobenius algebra X of nonzero dimension within a monoidal category \mathcal{C} , where the unit object $\mathbf{1}$ is linear-simple, is special.*

Proof. Since $\mathbf{1}$ is linear-simple, $\text{Hom}_{\mathcal{C}}(\mathbf{1}, \mathbf{1})$ is one-dimensional. Thus, from the above discussion, we have $\dim(X) = \alpha_X \beta_X$, which is nonzero by assumption. Therefore, both α_X and β_X must be nonzero, indicating that X is special. \square

Lemma 6.4. *A totally nonzero Frobenius algebra X has nonzero dimension.*

Proof. Recall that $\dim(X) = \text{tr}(\text{id}_X)$. Since id_X is an idempotent morphism, its trace is nonzero by assumption. \square

Lemma 6.5. *Let $(X, m, \delta, e, \epsilon)$ be a special Frobenius algebra within a \mathbb{k} -linear Karoubian monoidal category \mathcal{C} . Then, none of the morphisms m, δ, e, ϵ are negligible.*

Proof. Given the special condition and applying Lemmas 2.4, 2.8, 2.10, and 2.16, we find that

$$\text{tr}(\delta \circ m) = \text{tr}(m^{**} \circ \delta) = \text{tr}(m \circ \delta) = \beta_X \dim(X)$$

and

$$\text{tr}(e \circ \epsilon) = \text{tr}(\epsilon^{**} \circ e) = \text{tr}(\epsilon \circ e) = \alpha_X$$

are both non-zero. This confirms that none of the morphisms are negligible. \square

Proposition 6.6. *Let $(X, m, \delta, e, \epsilon)$ be a special Frobenius algebra in a monoidal category \mathcal{C} that satisfies the conditions in Remark 6.1 (e.g., a spherical tensor category). Then, it is also a Frobenius algebra \overline{X} in the semisimplification $\overline{\mathcal{C}}$. Furthermore, if X is totally nonzero (see Definition 5.2), the Frobenius subalgebra poset (see Definition 4.18) of X in \mathcal{C} embeds into the corresponding poset in $\overline{\mathcal{C}}$.*

Proof. A special Frobenius algebra in \mathcal{C} remains a Frobenius algebra in $\overline{\mathcal{C}}$ due to Lemma 6.5 and the preservation of morphism compositions and tensor products from \mathcal{C} , which ensures that the axioms are satisfied. For any indecomposable component Y of X , there exists a monomorphism $i : Y \rightarrow X$ and an epimorphism $p : X \rightarrow Y$ such that $i \circ p = b$ and $p \circ i = \text{id}_Y$, indicating that b is a nonzero idempotent. Given that X is totally nonzero, we also have $\text{tr}(b) \neq 0$, which implies i is not negligible, leading to $\dim(Y) \neq 0$ by [13, Lemma 2.2]. Therefore, X has no indecomposable components of zero dimension, resulting in the poset embedding. \square

The primary application of Proposition 6.6 is to simplify the proof of the finiteness of the Frobenius subalgebra poset by reducing it to the semisimple case.

7. FORMAL ANGLE

We extend the concept of the angle between two intermediate subfactors, as defined in [1], to introduce a formal angle between two Frobenius subalgebras. We will employ the idempotent $b_{X'}$ from Definition 3.5, which generalizes the biprojection in subfactor planar algebra theory.

Assume C is a totally positive connected Frobenius algebra within a \mathbb{C} -linear abelian rigid monoidal category \mathcal{C} , where the unit object $\mathbf{1}$ is linear-simple. Recall the partial order on the idempotents in $\text{Hom}_{\mathcal{C}}(C, C)$ as defined in §5. Let A and B be two Frobenius subalgebras of C . We further assume that $\text{tr}(b_A \circ b_B)$ is nonzero (the zero case will be addressed later, see Remark 7.9). By applying Lemma 2.16 and Corollary 3.16, we can define the idempotent

$$b_{AB} := \frac{\mu_C}{\text{tr}(b_A \circ b_B)} b_A * b_B,$$

which satisfies $\text{tr}(b_{AB}) = \frac{\text{tr}(b_A)\text{tr}(b_B)}{\text{tr}(b_A \circ b_B)}$, and $b_Y \leq b_{AB}$ for all $Y \in \{A, B\}$.

Remark 7.1. *The notation b_{AB} is inspired by [19, Theorem 3.10]. However, it is used here solely as a notation; there is no requirement to show that b_{AB} corresponds to the idempotent associated with the image of $m \circ (i_A \otimes i_B)$.*

Remark 7.2. *Note that $\text{tr}(b_A \circ b_B)$ is positive, as it equals $\frac{\text{tr}(b_A)\text{tr}(b_B)}{\text{tr}(b_{AB})}$, which is positive due to the total positivity assumption and the fact that b_A , b_B , and b_{AB} are idempotent.*

Following the notation from §4, we define $b_{A \cap B}$ as $i_{A \cap B} \circ i_{A \cap B}^*$, where $i_{A \cap B} = i_Y \circ j_Y$ for $Y \in \{A, B\}$.

Lemma 7.3. *For all $Y \in \{A, B\}$, we have $b_{A \cap B} \leq b_Y$. Consequently, $\text{tr}(b_{A \cap B}) \leq \text{tr}(b_Y)$.*

Proof. By definition, and the equality $i_Y^* \circ i_Y = \text{id}_Y$ (from Definition 3.5):

$$\begin{aligned} b_Y \circ b_{A \cap B} &= i_Y \circ i_Y^* \circ (i_Y \circ j_Y) \circ (i_Y \circ j_Y)^* \\ &= i_Y \circ (i_Y^* \circ i_Y) \circ j_Y \circ (i_Y \circ j_Y)^* \\ &= (i_Y \circ j_Y) \circ (i_Y \circ j_Y)^* = b_{A \cap B}. \end{aligned}$$

Similarly, we find $b_{A \cap B} \circ b_Y = b_{A \cap B}$. The last statement follows from Lemma 5.3. \square

Lemma 7.4. *Let A, B, C be objects in an abelian rigid monoidal category with $A, B \subseteq C$ and C being semisimple. If A and B are selfdual, then so is $A + B$. Furthermore, i_{A+B} satisfies $i_{A+B}^{**} = i_{A+B}$ and $i_{A+B}^* \circ i_{A+B} = \text{id}_{A+B}$.*

Proof. Following the argument from Proposition 4.7, we have $A + B = \bigoplus_i (A_i + B_i) \otimes X_i$, leading to

$$(A + B)^* = \bigoplus_i (A_i + B_i) \otimes X_i^* = \bigoplus_i (A_i^* + B_i^*) \otimes X_i = \bigoplus_i (A_i + B_i) \otimes X_i = A + B.$$

Moreover, i_{A+B} is induced by the inclusion $(A_i + B_i) \subset C_i$. The rest of the proof is straightforward. \square

We wonder whether Lemma 7.4 can be proven without the assumption that C is semisimple, by defining $A + B$ using the pushout $(k_Y : Y \rightarrow A + B)_{Y \in \{A, B\}}$ of the pullback $(j_Y : A \cap B \rightarrow Y)$ of $(i_Y : Y \rightarrow C)$. By universality, there exists a monomorphism $i_{A+B} : A + B \rightarrow C$ such that $i_{A+B} \circ k_Y = i_Y$.

$$\begin{array}{ccc} A \cap B & \xrightarrow{j_A} & A \\ j_B \downarrow & & \downarrow k_A \\ B & \xrightarrow{k_B} & A + B \\ & \searrow i_B & \downarrow i_{A+B} \\ & & C \end{array}$$

(Note: In the original image, there is a curved arrow from A to C labeled i_A, and a curved arrow from B to C labeled i_B. The arrow from A+B to C is labeled i_{A+B}.)

Recall that C is totally positive. We now assume it is also semisimple as an object. Let b_{A+B} denote the idempotent $i_{A+B} \circ i_{A+B}^*$.

Lemma 7.5. *For all $Y \in \{A, B\}$, it holds that $b_Y \leq b_{A+B}$. Thus, $\text{tr}(b_Y) \leq \text{tr}(b_{A+B})$.*

Proof. By definition, the equality $i_{A+B}^* \circ i_{A+B} = \text{id}_{A+B}$ and above discussion,

$$\begin{aligned} b_{A+B} \circ b_Y &= (i_{A+B} \circ i_{A+B}^*) \circ (i_Y \circ i_Y^*) \\ &= i_{A+B} \circ i_{A+B}^* \circ (i_{A+B} \circ k_Y) \circ (i_{A+B} \circ k_Y)^* \\ &= i_{A+B} \circ (i_{A+B}^* \circ i_{A+B}) \circ k_Y \circ (i_{A+B} \circ k_Y)^* \\ &= (i_{A+B} \circ k_Y) \circ (i_{A+B} \circ k_Y)^* = b_Y. \end{aligned}$$

Similarly, we also find $b_Y \circ b_{A+B} = b_Y$. The last statement follows from Lemma 5.3. \square

Lemma 7.6. *The equality $\text{tr}(b_{A+B}) = \text{tr}(b_A) + \text{tr}(b_B) - \text{tr}(b_{A \cap B})$ holds.*

Proof. Using Lemma 2.16 and the property $i_Z^{**} = i_Z$, we get $\text{tr}(b_Z) = \text{tr}(i_Z \circ i_Z^*) = \text{tr}(i_Z^* \circ i_Z) = \text{tr}(\text{id}_Z)$. Therefore,

$$\begin{aligned} \text{tr}(b_{A+B}) &= \text{tr}(\text{id}_{A+B}) = \sum_i \dim(A_i + B_i) \text{tr}(X_i) \\ &= \sum_i (\dim(A_i) + \dim(B_i) - \dim(A_i \cap B_i)) \text{tr}(X_i) \\ &= \text{tr}(\text{id}_A) + \text{tr}(\text{id}_B) - \text{tr}(\text{id}_{A \cap B}) = \text{tr}(b_A) + \text{tr}(b_B) - \text{tr}(b_{A \cap B}). \end{aligned} \quad \square$$

Lemma 7.7. *Let P be a poset and let p, q_1, q_2 belong to P such that $q_i \leq p$ for $i \in \{1, 2\}$. If q_1 and q_2 have a unique supremum $q_1 \vee q_2$ (called the join), then $q_1 \vee q_2 \leq p$.*

Proof. This statement is immediate. \square

Lemma 7.8. *The inequality $b_{A+B} \leq b_{AB}$ holds. Consequently, $\text{tr}(b_{A+B}) \leq \text{tr}(b_{AB})$.*

Proof. This follows directly from the inequality $b_Y \leq b_{AB}$ and Lemma 7.7, since $b_{A+B} = b_A \vee b_B$. The last statement follows from Lemma 5.3. \square

By combining Lemmas 7.3, 7.6, 7.8, and Corollary 3.16, for all $Y \in \{A, B\}$ we have:

$$0 < \text{tr}(b_Y) \leq \text{tr}(b_A) + \text{tr}(b_B) - \text{tr}(b_{A \cap B}) = \text{tr}(b_{A+B}) \leq \text{tr}(b_{AB}) = \frac{\text{tr}(b_A) \text{tr}(b_B)}{\text{tr}(b_A \circ b_B)},$$

where positivity holds because b_Y is a nonzero idempotent and C is totally positive. Therefore, we derive:

$$\begin{aligned} \text{tr}(b_A \circ b_B) &\leq \frac{\text{tr}(b_A) \text{tr}(b_B)}{\text{tr}(b_A) + \text{tr}(b_B) - \text{tr}(b_{A \cap B})}, \\ \text{tr}(b_A \circ b_B) - \text{tr}(b_{A \cap B}) &\leq \frac{\text{tr}(b_A) \text{tr}(b_B)}{\text{tr}(b_A) + \text{tr}(b_B) - \text{tr}(b_{A \cap B})} - \text{tr}(b_{A \cap B}), \\ &= \frac{\text{tr}(b_A) \text{tr}(b_B) - \text{tr}(b_{A \cap B})(\text{tr}(b_A) + \text{tr}(b_B)) + \text{tr}(b_{A \cap B})^2}{\text{tr}(b_A) + \text{tr}(b_B) - \text{tr}(b_{A \cap B})}, \\ &= \frac{(\text{tr}(b_A) - \text{tr}(b_{A \cap B}))(\text{tr}(b_B) - \text{tr}(b_{A \cap B}))}{\text{tr}(b_A) + \text{tr}(b_B) - \text{tr}(b_{A \cap B})}. \end{aligned}$$

By Lemmas 7.3 and 5.3, $(b_A - b_{A \cap B}) \circ (b_B - b_{A \cap B}) = b_A \circ b_B - b_{A \cap B}$, and $\text{tr}(b_Y - b_{A \cap B}) > 0$ if $b_Y \neq b_{A \cap B}$. Thus,

$$\begin{aligned} \frac{\text{tr}((b_A - b_{A \cap B}) \circ (b_B - b_{A \cap B}))}{\sqrt{\text{tr}(b_A - b_{A \cap B}) \text{tr}(b_B - b_{A \cap B})}} &\leq \frac{\sqrt{\text{tr}(b_A - b_{A \cap B}) \text{tr}(b_B - b_{A \cap B})}}{\text{tr}(b_A) + \text{tr}(b_B) - \text{tr}(b_{A \cap B})}, \\ &< \frac{\sqrt{\text{tr}(b_A - b_{A \cap B}) \text{tr}(b_B - b_{A \cap B})}}{\text{tr}(b_A - b_{A \cap B}) + \text{tr}(b_B - b_{A \cap B})}, \\ &\leq \frac{1}{2}. \end{aligned}$$

The final inequality follows from the AM–GM inequality, which states that for two non-negative numbers x and y , we have $\frac{x+y}{2} \geq \sqrt{xy}$.

Remark 7.9. *We initially assumed $\text{tr}(b_A \circ b_B)$ to be nonzero. However, if it is zero, the above inequality trivially holds.*

Let us define the *formal angle* between b_A and b_B with respect to $b_A \wedge b_B = b_{A \cap B}$ as follows:

$$\theta(b_A, b_B) := \begin{cases} \arccos \left(\frac{\text{tr}((b_A - b_{A \cap B}) \circ (b_B - b_{A \cap B}))}{\sqrt{\text{tr}(b_A - b_{A \cap B}) \text{tr}(b_B - b_{A \cap B})}} \right), & \text{if } b_A \neq b_B \\ 0, & \text{if } b_A = b_B. \end{cases}$$

Remark 7.10. *The term angle draws inspiration from the work in [1], while the descriptor formal indicates that the trace tr may not define an inner product space on $\text{Hom}_C(C, C)$, as the tensor category is not assumed to be unitary, and thus it may not yield a genuine angle.*

8. WATATANI'S THEOREM

This section is dedicated to proving Theorem 1.2.

Definition 8.1. Let L be a lattice and $a \in L$. We define the subset $m_a \subset L$ as the collection of elements $b \in L$ such that if $a < c \leq b$, then $b = c$. In simpler terms, m_a is the set of minimal elements in L that are greater than a .

Theorem 8.2. Let Z be a Frobenius subalgebra of a totally positive connected Frobenius algebra X of finite length and semisimple as an object in a \mathbb{C} -linear abelian rigid monoidal category \mathcal{C} with $\mathbf{1}$ being linear-simple. Consider the subset m_{b_Z} , as defined in Definition 8.1, of the interval lattice $[b_{\mathbf{1}}, b_X]$ from Theorem 4.19. Then m_{b_Z} is a finite set.

Proof. Let the elements of m_{b_Z} be denoted by b_{A_i} for each i in some index set I . Define the vectors

$$v_i := \frac{b_{A_i} - b_Z}{\sqrt{\operatorname{tr}(b_{A_i} - b_Z)}}$$

for each $i \in I$. Let V_Z be the \mathbb{R} -subspace of $\operatorname{Hom}_{\mathcal{C}}(X, X)$ spanned by the vectors $(v_i)_{i \in I}$. Since $\operatorname{Hom}_{\mathcal{C}}(X, X)$ is isomorphic to $\operatorname{Hom}_{\mathcal{C}}(\mathbf{1}, X \otimes X^*)$, it is finite-dimensional, as X is semisimple and $\mathbf{1}$ is linear-simple. Therefore, there exists a finite subset $J \subseteq I$ such that $(v_i)_{i \in J}$ forms a basis for V_Z .

We define a real inner product on V_Z as follows:

$$\left\langle \sum_{j \in J} \lambda_j v_j, \sum_{j \in J} \mu_j v_j \right\rangle := \sum_{j \in J} \lambda_j \mu_j.$$

Let M_Z be the matrix defined by $(\operatorname{tr}(v_i \circ v_j))_{i, j \in J}$. For all $i, j \in I$,

$$\begin{aligned} \langle v_i, M_Z v_j \rangle &= \left\langle \sum_{k \in J} \lambda_k v_k, M_Z \sum_{l \in J} \mu_l v_l \right\rangle \\ &= \sum_{k, l \in J} \lambda_k \mu_l \langle v_k, M_Z v_l \rangle \\ &= \sum_{k, l \in J} \lambda_k \mu_l \operatorname{tr}(v_k \circ v_l) \\ &= \operatorname{tr}\left(\left(\sum_{k \in J} \lambda_k v_k\right) \circ \left(\sum_{l \in J} \mu_l v_l\right)\right) \\ &= \operatorname{tr}(v_i \circ v_j). \end{aligned}$$

If $i \neq j$, then by minimality, we have $b_{A_i \cap A_j} = b_Z$. Therefore, based on the discussion at the end of §7,

$$\langle v_i, M_Z v_j \rangle = \operatorname{tr}(v_i \circ v_j) = \cos(\theta(b_{A_i}, b_{A_j})) < \frac{1}{2}.$$

Assume there exists $i \in I$ and a sequence $(i_n)_{n \in \mathbb{N}}$ such that $i \neq i_n$ for all n , and the angle $\angle(v_i, v_{i_n})$ tends to zero in the real inner product space V_Z . As a result, there exists a sequence of positive real numbers (α_n) such that $\lim_{n \rightarrow \infty} (\alpha_n v_{i_n}) = v_i$. For all $j \in I$, we have

$$\langle v_j, M_Z v_j \rangle = \operatorname{tr}(v_j \circ v_j) = \operatorname{tr}\left(\left(\frac{b_{A_j} - b_Z}{\sqrt{\operatorname{tr}(b_{A_j} - b_Z)}}\right) \circ \left(\frac{b_{A_j} - b_Z}{\sqrt{\operatorname{tr}(b_{A_j} - b_Z)}}\right)\right) = \operatorname{tr}\left(\frac{b_{A_j} - b_Z}{\operatorname{tr}(b_{A_j} - b_Z)}\right) = 1.$$

By continuity, we find that

$$\lim_{n \rightarrow \infty} \langle \alpha_n v_{i_n}, M_Z \alpha_n v_{i_n} \rangle = \langle v_i, M_Z v_i \rangle = 1.$$

However, since

$$\langle \alpha_n v_{i_n}, M_Z \alpha_n v_{i_n} \rangle = \alpha_n^2 \langle v_{i_n}, M_Z v_{i_n} \rangle = \alpha_n^2,$$

it follows that

$$\lim_{n \rightarrow \infty} \alpha_n = 1.$$

Again, by continuity, we have

$$\lim_{n \rightarrow \infty} \langle v_i, M_Z v_{i_n} \rangle = \lim_{n \rightarrow \infty} \alpha_n \langle v_i, M_Z v_{i_n} \rangle = \lim_{n \rightarrow \infty} \langle v_i, M_Z \alpha_n v_{i_n} \rangle = \langle v_i, M_Z v_i \rangle = 1.$$

This contradicts the inequality $\langle v_i, M_Z v_{i_n} \rangle < \frac{1}{2}$, which holds because we assumed that $i \neq i_n$ for all n . Thus, our initial assumption must be incorrect. This means that there exists a constant $\beta > 0$ such that for all $i, j \in I$ with $i \neq j$, the angle $\angle(v_i, v_j) \geq \beta$. Since V_Z is finite-dimensional, the index set I must also be finite. \square

An estimate of β , possibly involving $\|M_Z\|$, should provide an upper bound for the size of the finite lattice, similarly to [1].

Definition 8.3. Let L be a lattice and $a \in L$. Define the subset $m_a^s \subset L$ as follows: $m_a^0 = \{a\}$ if $s = 0$, and $m_a^s = \bigcup_{b \in m_a^{s-1}} m_b$ if $s \geq 1$. In particular, $m_a^1 = m_a$.

Remark 8.4. If L has finite height h and a bottom element 1 , then $L = \bigcup_{s=0}^h m_1^s$.

Theorem 8.5. Let X be a totally positive connected Frobenius algebra of finite length and semisimple as an object in a \mathbb{C} -linear abelian rigid monoidal category \mathcal{C} with $\mathbf{1}$ being linear-simple. Then its Frobenius subalgebra lattice is finite.

Proof. By Theorem 4.19, the Frobenius subalgebra lattice of X is isomorphic to the interval $[b_1, b_X]$. The bottom element is b_1 and it has finite height, say h , since X has finite length. By Remark 8.4, we have $[b_1, b_X] = \bigcup_{s=0}^h m_{b_1}^s$. According to Theorem 8.2, Definition 8.3, and by induction, each set $m_{b_1}^s$ is finite. Therefore, $[b_1, b_X]$ is finite. \square

Corollary 8.6. Let X be a totally positive connected Frobenius algebra in a spherical tensor category \mathcal{C} . Then its Frobenius subalgebra poset is finite.

Proof. By Lemmas 2.14, 6.3, and 6.4, X is determined to be special. The conclusion follows from Proposition 6.6 and Theorem 8.5. Note that as an object in a tensor category, X has finite length and can be uniquely decomposed (up to isomorphism) into a direct sum of finitely many indecomposable components (Krull-Schmidt theorem). Due to total positivity, all these indecomposables have nonzero dimensions. Consequently, \overline{X} in the semisimplification $\overline{\mathcal{C}}$ also admits a unique decomposition (up to isomorphism) as a direct sum of finitely many simple objects corresponding to the aforementioned indecomposables. \square

We can obtain a more general statement by considering the conditions in Remark 6.1.

Corollary 8.7. Let X be a totally positive connected Frobenius algebra in a semisimple tensor category. Then its Frobenius subalgebra lattice is finite.

Proof. This follows immediately from Theorem 8.5. \square

Corollary 8.8. Let X be a connected Frobenius algebra in a pseudo-unitary fusion category \mathcal{C} . Then its Frobenius subalgebra lattice is finite.

Proof. This follows immediately from Proposition 5.6 and Corollary 8.7. \square

The structure of a (connected) Frobenius algebra object relies solely on the fusion data of the Grothendieck ring. In other words, determining all possible Frobenius algebra structures on an object reduces to solving polynomial equations that depend only on the object's isomorphism class and the Grothendieck ring. Therefore:

Corollary 8.9. Let X be a connected Frobenius algebra in a fusion category \mathcal{C} that is Grothendieck equivalent to a fusion category \mathcal{D} in which the corresponding Frobenius subalgebra lattice is finite. Then it is also finite in \mathcal{C} .

Conjecture 8.10. Let X be a connected Frobenius algebra in a fusion category. Then its Frobenius subalgebra lattice is finite.

According to Corollaries 8.8 and 8.9, to find a counterexample to Conjecture 8.10, we need a fusion category that is not Grothendieck equivalent to a pseudo-unitary one. Such fusion categories are quite rare in the literature. The first known example was identified in this paper [55], specifically the rank 6 modular fusion category $\mathcal{C}(\mathfrak{so}_5, 3/2)_{ad}$. It would be intriguing to classify the connected Frobenius algebras within this category. Note that some of its connected Frobenius algebras are already totally positive (e.g., $X \otimes X^*$).

The remaining section presents families of examples of connected Frobenius algebras and discusses the applicability of Watatani's theorem.

9. INCLUSION OF C^* -ALGEBRAS

Let $A \subset B$ be a unital inclusion of \mathbb{k} -algebras. Define \mathcal{C} as the \mathbb{k} -linear abelian monoidal category of A -bimodules, denoted $\text{Bimod}(A)$. More generally for rings, we refer the reader to [45, Example 1.6]. It is convenient to use the (abuse of) notation $A' \cap B$ to denote the commutant of A within B , defined as $\{b \in B \mid ab = ba \ \forall a \in A\}$, even though the term A' on its own may not make sense in general.

Lemma 9.1. The \mathbb{k} -vector spaces $\text{Hom}_{\mathcal{C}}({}_A A_A, {}_A B_A)$ and $A' \cap B$ are isomorphic.

Proof. Consider the linear map $T : A' \cap B \rightarrow \text{Hom}_{\mathcal{C}}({}_A A_A, {}_A B_A)$ defined by $T(x) = \hat{x} : a \mapsto xa$. The map \hat{x} is a morphism in \mathcal{C} because $\hat{x}(a_1 a a_2) = x a_1 a a_2 = a_1 x a a_2 = a_1 \hat{x}(a) a_2$, where $a_1 \in A$ commutes with $x \in A' \cap B$.

Next, define the linear map $S : \text{Hom}_{\mathcal{C}}({}_A A_A, {}_A B_A) \rightarrow A' \cap B$ by $S(f) = f(1)$. We will show that S and T are inverses of each other. First, we have:

$$T(S(f)) = T(f(1)) = \widehat{f(1)}.$$

Since $\widehat{f(1)}(a) = f(1)a = f(a)$ (because f is a bimodule morphism), we find $\widehat{f(1)} = f$, implying $T \circ S = \text{id}$.

Next, for $x \in A' \cap B$:

$$T(S(x)) = T(\hat{x}) = \hat{x}(1) = x,$$

which shows that $S \circ T = \text{id}$. Thus, S and T are indeed inverses. \square

Definition 9.2. A unital inclusion of algebras $A \subset B$ is called irreducible if $A' \cap B$ is one-dimensional.

Lemma 9.3. A unital inclusion of algebras $A \subset B$ is irreducible if and only if ${}_A B_A$ is connected (and consequently, A is centerless and $\mathbf{1}$ is linear-simple).

Proof. Since ${}_A A_A = \mathbf{1}$ is the unit in \mathcal{C} , the result follows directly from Definitions 2.13 and 9.2, along with Lemma 9.1. Furthermore, since $(A' \cap A) \subset (A' \cap B)$, if ${}_A B_A$ is connected, it follows that both $A' \cap A$ and $\text{Hom}_{\mathcal{C}}(\mathbf{1}, \mathbf{1})$ are one-dimensional. This means that A is centerless and that $\mathbf{1}$ is linear-simple. \square

Theorem 9.4. A finite index unital irreducible inclusion of C^* -algebras has a finite number of intermediate C^* -algebras.

Proof. According to [62], a finite index unital inclusion of C^* -algebras is defined through a faithful conditional expectation $E_A : B \rightarrow A$. As a result, based on [9, Lemma 3.11], the object ${}_A B_A$ is a Frobenius algebra object within the C^* -tensor category $\text{Bimod}(A)$, and it is connected by Lemma 9.3. To proceed with applying Corollary 8.8, observe that under the finite index assumption, the semisimple object ${}_A B_A$ generates a unitary fusion subcategory of $\text{Bimod}(A)$. Additionally, every intermediate C^* -algebra corresponds to a Frobenius subalgebra of ${}_A B_A$. \square

The theorem presented above extends [23, Corollary 3.9] by removing the requirement for simplicity. However, as indicated in Lemma 9.3, the algebra A is centerless when the inclusion is irreducible. Therefore, it is crucial to identify non-simple, centerless C^* -algebras (see Proposition 9.8). First, let us review some basic definitions and results.

Definition 9.5. A C^* -algebra (or von Neumann algebra) is called simple if it has no nonzero proper ideals that are closed under the operator norm topology (or the weak operator topology, respectively).

Proposition 9.6. A von Neumann algebra M is simple if and only if it is centerless (i.e., a factor).

Proof. The simplicity of a factor is specifically addressed in [29, Proposition A.3.1]. Conversely, if M is not a factor, it contains a nontrivial central projection $p \in M \cap M'$. Thus, pMp forms a nontrivial closed ideal in M , demonstrating that M is not simple. \square

Proposition 9.7. Let H be a separable Hilbert space, and let $A \subset B(H)$ be a unital C^* -algebra. Let $M := A''$ be the von Neumann algebra generated by A . If M is a factor then A is centerless.

Proof. This follows immediately since $(A \cap A'')' = \overline{A \cap A'}^{\text{wot}} \subseteq (\overline{A}^{\text{wot}} \cap \overline{A'}^{\text{wot}}) = M \cap M'$. \square

Recall that the von Neumann algebra $L(\Gamma)$ of a countable group Γ is a factor if and only if Γ is ICC (infinite conjugacy classes).

Proposition 9.8. Let Γ be a countable ICC group with a non-trivial amenable normal subgroup $N \subseteq \Gamma$. Then the reduced C^* -algebra $C_r^*(\Gamma)$ is centerless but non-simple.

Proof. Since Γ is a countable ICC group, $L(\Gamma)$ is a factor. Therefore, by Proposition 9.7, $C_r^*(\Gamma)$ is centerless because $C_r^*(\Gamma)'' = L(\Gamma)$. The non-simplicity follows from [22, Proposition 3]. \square

In particular, for all non-trivial ICC groups G and Γ , with G being amenable (e.g., S_∞), $C_r^*(G)$ and $C_r^*(G \times \Gamma)$ are centerless, non-simple C^* -algebras. In contrast, as noted in [22], for every torsion-free non-elementary hyperbolic group Γ , the C^* -algebra $C_r^*(\Gamma)$ is simple. This implies that Γ has no non-trivial amenable normal subgroups (i.e., it has a trivial amenable radical) and is ICC.

10. HOPF ALGEBRAS

As we will elaborate in this subsection (but, see Remark 10.6), the dual Hopf algebra H^* serves as a connected Frobenius algebra object in $\text{Rep}(H)$. One might consider examining their Frobenius subalgebra objects, specifically focusing on whether their right (or left) coideal subalgebras qualify as such. This is supported by [24, Theorem 4.4] in the unitary case, but it remains unclear in general, even under the semisimple assumption. This exploration is motivated by the desire to determine if we can recover [14, Theorem 3.6] as an application of Corollary 8.6, while potentially relaxing the assumption of semisimplicity. However, this investigation will not be addressed in the current paper.

Definition 10.1 (Bialgebra). *A bialgebra is a quintuple $(H, m, e, \Delta, \epsilon)$ consisting of:*

- vector space H over a field \mathbb{k} ,
- multiplication map $m : H \otimes H \rightarrow H$,
- unit map $e : \mathbb{k} \rightarrow H$,
- comultiplication map $\Delta : H \rightarrow H \otimes H$,
- counit map $\epsilon : H \rightarrow \mathbb{k}$,

that satisfies the following axioms:

- (1) Associativity: $m \circ (m \otimes \text{id}) = m \circ (\text{id} \otimes m)$,
- (2) Unit axiom: $m \circ (e \otimes \text{id}) = \text{id} = m \circ (\text{id} \otimes e)$,
- (3) Coassociativity: $(\Delta \otimes \text{id}) \circ \Delta = (\text{id} \otimes \Delta) \circ \Delta$,
- (4) Counit axiom: $(\epsilon \otimes \text{id}) \circ \Delta = \text{id} = (\text{id} \otimes \epsilon) \circ \Delta$,
- (5) Compatibility axioms: Δ and ϵ are unital algebra morphisms, i.e., for $a, b \in H$,

$$\Delta(ab) = \Delta(a)\Delta(b), \quad \Delta(1_H) = 1_H \otimes 1_H, \quad \epsilon(ab) = \epsilon(a)\epsilon(b), \quad \epsilon(1_H) = 1,$$

where $e(1) = 1_H$. It is equivalent to say that m and e are counital coalgebra morphisms.

Definition 10.2 (Antipode). *An antipode is a linear map $S : H \rightarrow H$ that satisfies the antipode axiom:*

$$m \circ (\text{id} \otimes S) \circ \Delta = e \circ \epsilon = m \circ (S \otimes \text{id}) \circ \Delta.$$

Equivalently, this can be expressed as:

$$a_{(1)}S(a_{(2)}) = \epsilon(a)1_H = S(a_{(1)})a_{(2)}$$

for all $a \in H$, using Sweedler notation $\Delta(a) = a_{(1)} \otimes a_{(2)}$ for comultiplication.

Recall that S is an antihomomorphism of (co)unital (co)algebras (see e.g., [60, Proposition 1.3.12]), which means:

$$S(1_H) = 1_H, \quad S(ab) = S(b)S(a), \quad \epsilon(S(h)) = \epsilon(h), \quad (S(h))_{(1)} \otimes (S(h))_{(2)} = S(h_{(2)}) \otimes S(h_{(1)}).$$

Definition 10.3 (Hopf Algebra). *A Hopf algebra is a bialgebra $(H, m, e, \Delta, \epsilon)$ that is equipped with an invertible antipode S .*

Definition 10.4. *Let H be a Hopf algebra. A subalgebra $K \subseteq H$ is termed a right (resp. left) coideal subalgebra if $\Delta(K) \subset K \otimes H$ (resp. $\Delta(K) \subset H \otimes K$), where Δ represents the comultiplication of H .*

Theorem 10.5 ([31, 44]). *A finite-dimensional Hopf algebra is a Frobenius algebra (in Vec).*

Remark 10.6. *The application of Watatani's theorem necessitates a connected Frobenius algebra object within a tensor category. In the category Vec , an object is connected if and only if it is trivial. This is why Theorem 10.5 needs to be revised to Theorem 10.7. The proof of this well-known result is presented for readers who may not be experts in Hopf algebra theory. It is included here because the authors feel that the existing literature lacks a self-contained and detailed proof.*

Theorem 10.7. *Let H be a finite-dimensional Hopf algebra. The dual H^* can be endowed with the structure of a connected Frobenius algebra object in $\text{Rep}(H)$, but it uses a different comultiplication and counit.*

Proof. The proof will be divided into several lemmas. For $K = H$ or H^* , let $m_K, e_K, \Delta_K, \epsilon_K$, and S_K denote the multiplication, unit, comultiplication, counit, and antipode of K . If there is no risk of confusion, the subscript K will be omitted. Let $\text{Rep}(H)$ be the tensor category of left H -modules. For all $h, k \in H$ and $f, g \in H^*$, recall that $(fg)(h) = f(h_{(1)})g(h_{(2)})$, $\Delta(f)(h \otimes k) = f(hk)$, $1_{H^*} = e_{H^*}(1) = \epsilon_H$, and $\epsilon_{H^*}(f) = f(1_H)$.

Lemma 10.8. *The triple (H^*, m_{H^*}, e_{H^*}) is a unital algebra object in $\text{Rep}(H)$.*

Proof. First, H^* is an object in $\text{Rep}(H)$ because for all $h, k, x \in H$ and for all $f \in H^*$, the expression $(h \cdot f)(x) := f(xh)$ defines a left H -module structure on H^* since:

$$(h \cdot (k \cdot f))(x) = (k \cdot f)(xh) = f(xhk) = ((hk) \cdot f)(x).$$

Next, we need to check that m_{H^*} and e_{H^*} are morphisms in $\text{Rep}(H)$. Let $h, k \in H$ and $f, g \in H^*$, then:

$$\begin{aligned} (m_{H^*}(h \cdot (f \otimes g)))(k) &= (m_{H^*}((h_{(1)} \cdot f) \otimes (h_{(2)} \cdot g)))(k) = (h_{(1)} \cdot f)(k_{(1)})(h_{(2)} \cdot g)(k_{(2)}) \\ &= f(k_{(1)}h_{(1)})g(k_{(2)}h_{(2)}) = f((kh)_{(1)})g((kh)_{(2)}) = (m_{H^*}(f \otimes g))(kh) = (h \cdot (m_{H^*}(f \otimes g)))(k), \\ (e_{H^*}(h \cdot 1))(k) &= (e_{H^*}(\epsilon_H(h)))(k) = (e_{H^*}(1))(k)\epsilon_H(h) = \epsilon_H(k)\epsilon_H(h) = \epsilon_H(kh) = (h \cdot (e_{H^*}(1)))(k). \quad \square \end{aligned}$$

Lemma 10.9. *The triple $(H, \Delta_H, \epsilon_H)$ is a counital coalgebra object in $\text{Rep}(H)$.*

Proof. Consider the left action of H on itself defined by $h \cdot k = hk$. We need to verify that Δ_H and ϵ_H are morphisms in $\text{Rep}(H)$:

$$\begin{aligned} \Delta_H(h \cdot k) &= \Delta_H(hk) = \Delta_H(h)\Delta_H(k) = (h_{(1)}k_{(1)}) \otimes (h_{(2)}k_{(2)}) = h \cdot (k_{(1)} \otimes k_{(2)}) = h \cdot \Delta_H(k), \\ \epsilon_H(h \cdot k) &= \epsilon_H(hk) = \epsilon_H(h)\epsilon_H(k) = h \cdot (\epsilon_H(k)). \quad \square \end{aligned}$$

Lemma 10.10. *Let $f \in H^*$. Then $\phi_f : H \rightarrow H^*$ defined by $\phi_f(h) = h \cdot f$ is a morphism in $\text{Rep}(H)$.*

Proof. This reduces to the following computation:

$$(\phi_f(h \cdot k))(x) = (\phi_f(hk))(x) = ((hk) \cdot f)(x) = f(xhk) = (k \cdot f)(xh) = (\phi_f(k))(xh) = (h \cdot (\phi_f(k)))(x). \quad \square$$

By [31, Theorem on page 79], H^* admits a non-degenerate right integral λ . Let $\phi := \phi_\lambda$ from Lemma 10.10. It should be regarded as a Fourier transform, see for example [61, Definition 1.3].

Lemma 10.11. *The map ϕ is an isomorphism in $\text{Rep}(H)$.*

Proof. We already know that it is a morphism in $\text{Rep}(H)$ by Lemma 10.10. By finite-dimensionality, we only need to show that ϕ is injective. Let $h \in H$ such that $\phi(h) = 0$. Then we have $\lambda(kh) = (\phi(h))(k) = 0$ for all $k \in H$. Thus, $h = 0$ since λ is non-degenerate. \square

Regarding connectedness, we need to prove that:

Lemma 10.12. *The space $\text{Hom}_{\text{Rep}(H)}(\mathbf{1}, H^*)$ is one-dimensional.*

Proof. Let α be a morphism in this space. Then $\alpha(h \cdot 1) = h \cdot \alpha(1)$, but

$$\alpha(h \cdot 1)(k) = \alpha(\epsilon_H(h))(k) = \epsilon_H(h)\alpha(1)(k) \text{ and } (h \cdot \alpha(1))(k) = (\alpha(1))(kh).$$

Thus, we have $\epsilon_H(h)\alpha(1)(k) = (\alpha(1))(kh)$. Setting $k = 1_H$ gives us $\alpha(1) = \alpha(1)(1_H)\epsilon_H \in k\epsilon_H$. The result follows. \square

Finally, we define new comultiplication and counit on H^* as follows:

$$\Delta'_{H^*} := (\phi \otimes \phi) \circ \Delta_H \circ \phi^{-1} \quad \text{and} \quad \epsilon'_{H^*} = \epsilon_H \circ \phi^{-1}.$$

These are morphisms in $\text{Rep}(H)$, as they are compositions of morphisms in $\text{Rep}(H)$. We still need to verify that the morphisms $m_{H^*}, e_{H^*}, \Delta'_{H^*}, \epsilon'_{H^*}$ satisfy all the identities for a Frobenius algebra (see §2), but they hold in Vec by Theorem 10.5, and thus automatically hold in $\text{Rep}(H)$. \square

Remark 10.13 ([43]). *One Frobenius identity can be reformulated as follows:*

$$\lambda(hx_{(1)})\lambda(k_{(1)}x_{(2)})\lambda(k_{(2)}y) = \lambda(h_{(1)}x)\lambda(h_{(2)}y_{(1)})\lambda(ky_{(2)}),$$

for all elements x, y, h , and k in H . However, it is unclear how to prove this directly.

Let us apply Corollary 8.8 together with Theorem 10.7.

Corollary 10.14. *Let H be a finite-dimensional semisimple Hopf algebra. Then the dual Hopf algebra H^* admits a finite number of Frobenius subalgebra objects in $\text{Rep}(H)$.*

The assertion of Corollary 10.14 should allow for a reformulation purely in terms of Hopf algebra, incorporating identities similar to those mentioned in Remark 10.13. Based on Corollary 11.5, it may be challenging to find a non-semisimple finite-dimensional Hopf algebra that satisfies the totally positive assumption. However, if one exists, we can apply Corollary 8.6 as well. Now, let us turn our attention to the left coideal subalgebras.

Theorem 10.15 ([57]). *A left coideal subalgebra of a finite-dimensional Hopf algebra is a Frobenius algebra (in Vec).*

The following question is supported by [24, Theorem 4.4] in the unitary case.

Question 10.16. *Is it true that every left coideal subalgebra K of H^* can be given the structure of a connected Frobenius subalgebra object of H^* in $\text{Rep}(H)$, where H is a finite-dimensional Hopf algebra?*

From Definition 10.4, we have that $\Delta_{H^*}(K) \subseteq H^* \otimes K$. Let $i_K : K \rightarrow H^*$ be the embedding induced by this inclusion. We define the multiplication and unit on K as follows: $m_K := m_{H^*} \circ i_K$ and $e_K(1) = 1_H$.

Lemma 10.17. *A left coideal subalgebra K of H^* is a unital algebra object in $\text{Rep}(H)$.*

Proof. First, K is an object in $\text{Rep}(H)$. For any $f \in K$ and $h, k \in H$, we have:

$$(h \cdot f)(k) = f(kh) = (\Delta_{H^*}(f))(h \otimes k) = f_{(1)}(h)f_{(2)}(k),$$

which shows that $(h \cdot f)$ is in K since it is a linear combination of elements $f_{(2)}$ in K with coefficients $f_{(1)}(h)$. Specifically, i_K is a morphism in $\text{Rep}(H)$, since $i_K(h \cdot f) = h \cdot f = h \cdot i_K(f)$. Together with Lemma 10.8, this implies that m_K and e_K are morphisms in $\text{Rep}(H)$. \square

Since $\text{Hom}_{\text{Rep}(H)}(\mathbf{1}, K) \subset \text{Hom}_{\text{Rep}(H)}(\mathbf{1}, H^*)$, we conclude that K is connected, by Lemma 10.12 and the fact that $1_H \in K$. However, we *cannot* define the comultiplication and counit on K as $\Delta'_{H^*} \circ i_K$ and $\epsilon'_{H^*} \circ i_K$ to create a Frobenius algebra structure. If we were to do so, i_K would be an isomorphism according to Lemma 3.4. Alternatively, if we define the comultiplication and counit on K (similar to the approach for H^*) using an isomorphism of K -modules $\phi : K \rightarrow K^*$ as provided by Theorem 10.15, then ϕ may not be a morphism in $\text{Rep}(H)$.

Our motivation for Question 10.16 has already been addressed in the semisimple case by the following result.

Theorem 10.18 ([14], Theorem 3.6). *A semisimple (thus finite-dimensional) Hopf algebra over an algebraically closed field of characteristic zero has a finite number of left coideal subalgebras.*

We are still uncertain about how much the semisimple assumption can be relaxed, but we do know that it cannot be completely removed, even when limited to Hopf subalgebras.

Example 10.19. *Nichols' Hopf algebras H_{2^n} for $n \geq 1$, introduced in [37] and further explored in [12, Example 5.5.8], serve as a family of counterexamples for $n \geq 3$. The Hopf algebra H_{2^n} (for $n \geq 1$) is 2^n -dimensional and generated by elements g, x_1, \dots, x_{n-1} subject to the relations:*

$$g^2 = 1, \quad x_i g = -g x_i, \quad x_i^2 = 0, \quad x_i x_j = -x_j x_i \quad (i \neq j),$$

$$\Delta(g) = g \otimes g, \quad \Delta(x_i) = g \otimes x_i + x_i \otimes 1, \quad \epsilon(g) = 1, \quad \epsilon(x_i) = 0, \quad S(g) = g, \quad S(x_i) = -g x_i.$$

For $n = 1$, this corresponds to the group algebra of the cyclic group of order 2, and for $n = 2$, it represents Sweedler's 4-dimensional Hopf algebra. When $n \geq 3$, it contains infinitely many Hopf subalgebras. Specifically, for any subspace $E \subset \bigoplus_i \mathbb{C}x_i$, the subalgebra $H_E = \langle E, g \rangle$ is a Hopf subalgebra, since for any $x \in E$, we have $\Delta(x) = g \otimes x + x \otimes 1$ and $S(x) = -gx$. The dimension of H_E is $2^{\dim(E)+1}$ and is completely determined by E . Nichols' Hopf algebra H_{2^n} is unimodular if and only if n is odd. For $n = 2, 3$, refer to [46, Proposition 7 (e) and Proposition 10 (d)], where they are denoted $H_{(2,1,-1)}$ and $U_{(2,1,-1)}$. For general cases, see the correction in [11, Exercise 6.5.10(i)].

Every Hopf subalgebra is a left coideal subalgebra, but not vice versa. For example, Sweedler's 4-dimensional Hopf algebra has finitely many Hopf subalgebras but infinitely many left coideal subalgebras [14, Example 3.5].

Example 10.20. *The quantum double $D(H)$ of a finite-dimensional Hopf algebra H is a bicrossed product of H with H^{op*} . Hence, H is a Hopf subalgebra of $D(H)$. Moreover, $D(H)$ is factorizable and therefore unimodular ([12, Definition 6.5.7, Exercise 8.6.4(i), Propositions 7.14.6, 8.6.3, and 8.10.10]). Consequently, if H has infinitely many Hopf subalgebras (as is the case with Nichols' Hopf algebra H_8), then $D(H)$ is a factorizable finite-dimensional Hopf algebra that also has infinitely many Hopf subalgebras.*

11. OTHER EXAMPLES

This section presents additional examples—connected canonical Frobenius algebras (11.1), abstract spin chains (11.2), and vertex operator algebras (11.3)—that may fall within the scope of Watatani's theorem in §8.

11.1. Connected canonical Frobenius algebra. In [12, Corollary 7.20.4], the canonical Frobenius algebra $\underline{\text{Hom}}(\mathbf{1}, \mathbf{1})$ in $\mathcal{C} \boxtimes \mathcal{C}^{\text{op}}$ is examined for a unimodular *multitensor* category \mathcal{C} . However,

Proposition 11.1. *The Frobenius algebra $\underline{\text{Hom}}(\mathbf{1}, \mathbf{1})$ is connected if and only if \mathcal{C} is a tensor category.*

Proof. Let \mathcal{D} be a multitensor category and \mathcal{M} a \mathcal{D} -module category with objects M_1 and M_2 in \mathcal{M} . According to [12, (7.20)], the space $\underline{\text{Hom}}(M_1, M_2)$ is defined via the natural isomorphism:

$$\text{Hom}_{\mathcal{M}}(X \otimes M_1, M_2) \simeq \text{Hom}_{\mathcal{D}}(X, \underline{\text{Hom}}(M_1, M_2)).$$

Assuming \mathcal{M} is also monoidal with unit $\mathbf{1}_{\mathcal{M}}$, and letting $\mathbf{1}_{\mathcal{D}}$ be the unit of \mathcal{D} , we set $M_1 = M_2 = \mathbf{1}_{\mathcal{M}}$ and $X = \mathbf{1}_{\mathcal{D}}$. This yields:

$$\text{Hom}_{\mathcal{M}}(\mathbf{1}_{\mathcal{M}}, \mathbf{1}_{\mathcal{M}}) \simeq \text{Hom}_{\mathcal{D}}(\mathbf{1}_{\mathcal{D}}, \underline{\text{Hom}}(\mathbf{1}_{\mathcal{M}}, \mathbf{1}_{\mathcal{M}})).$$

Therefore, $\underline{\text{Hom}}(\mathbf{1}_{\mathcal{M}}, \mathbf{1}_{\mathcal{M}})$ is connected if and only if $\mathbf{1}_{\mathcal{M}}$ is linear-simple, meaning $\text{Hom}_{\mathcal{M}}(\mathbf{1}_{\mathcal{M}}, \mathbf{1}_{\mathcal{M}})$ is one-dimensional. For a multitensor category \mathcal{M} , this indicates that $\underline{\text{Hom}}(\mathbf{1}_{\mathcal{M}}, \mathbf{1}_{\mathcal{M}})$ is connected if and only if \mathcal{M} is a tensor category. The result follows by taking $\mathcal{D} = \mathcal{C} \boxtimes \mathcal{C}^{\text{op}}$ and $\mathcal{M} = \mathcal{C}$. \square

Corollary 11.2. *Let H be a finite-dimensional unimodular Hopf algebra. Then H^* serves as the connected canonical Frobenius algebra in $\text{Rep}(H \otimes H^{\text{cop}})$, albeit with a different comultiplication and counit.*

Proof. This characterization of the canonical Frobenius algebra is noted at the end of [12, §7.20]. The connectedness follows from Theorem 11.1. \square

Proposition 11.3 ([12, Proposition 7.18.9]). *Let \mathcal{C} be a unimodular non-semisimple finite tensor category. Let $f : \underline{\text{Hom}}(\mathbf{1}, \mathbf{1}) \rightarrow \underline{\text{Hom}}(\mathbf{1}, \mathbf{1})^{**}$ be a morphism in $\mathcal{C} \boxtimes \mathcal{C}^{\text{op}}$. Then $\text{tr}(f) = 0$.*

Referring to Lemma 2.4, we understand that a Frobenius algebra object is self-dual. Therefore, in Proposition 11.3, the morphism f can be viewed as an isomorphism, which implies that the categorical dimension of $\underline{\text{Hom}}(\mathbf{1}, \mathbf{1})$ with respect to f must be zero.

Corollary 11.4. *Let \mathcal{C} be a unimodular non-semisimple finite tensor category. Assume that $\mathcal{C} \boxtimes \mathcal{C}^{\text{op}}$ has a pivotal structure a . Then $\dim_a(\underline{\text{Hom}}(\mathbf{1}, \mathbf{1})) = 0$.*

Proof. This follows directly from Proposition 11.3. \square

Corollary 11.5. *Let H be a finite-dimensional unimodular non-semisimple Hopf algebra. Assume that $\text{Rep}(H \otimes H^{\text{cop}})$ has a pivotal structure a . Then $\dim_a(H^*) = 0$.*

Proof. This follows immediately from Corollaries 11.2 and 11.4. \square

In Corollary 11.5, we note that due to the fiber functor, FPdim is equal to $\dim_{\mathbb{k}}$. However, in characteristic zero, $\dim_{\mathbb{k}}(H^*)$ is non-zero. Thus, \dim_a must differ from FPdim in this case.

Remark 11.6. *In the pivotal non-semisimple case, the connected Frobenius algebras discussed in this subsection cannot be totally positive, which means that Corollary 8.6 is not applicable.*

11.2. Abstract spin chains. Quantum cellular automata (QCA) are quantum operations that reflect the core principles of unitarity and locality [64]. In [28], the authors explore extensions of bounded-spread isomorphisms of symmetric local algebras to QCA defined on complete (or edge-restricted) local operator algebras. They introduce the notions of abstract spin chains and categorical inclusions, offering an algebraic framework to study these inclusions. In this section, we aim to demonstrate that the lattice of categorical inclusions (as discussed in [28, §4]) corresponding to a connected, commutative Frobenius algebra (Q-system) is finite.

As detailed in [28, §3], let \mathcal{C} be an indecomposable unitary multi-fusion category, and let X be an object in \mathcal{C} . For any finite interval $I \subset \mathbb{Z}$, define $A(\mathcal{C}, X)_I$ as the finite-dimensional $*$ -algebra $\text{End}_{\mathcal{C}}(X^{\otimes |I|})$. The colimit $A(\mathcal{C}, X)$ of $A(\mathcal{C}, X)_I$, taken in the category of $*$ -algebras, is called an *abstract spin chain*.

Let \mathcal{C} be an indecomposable multi-fusion category. Following [28, Definition 3.2], a *quotient* of \mathcal{C} is defined as an indecomposable multi-fusion category \mathcal{D} , equipped with a dominant¹ tensor functor $F : \mathcal{C} \rightarrow \mathcal{D}$.

As noted in [28], a dominant tensor functor $F : \mathcal{C} \rightarrow \mathcal{D}$ induces an inclusion of C^* -algebras:

$$i_F : A(\mathcal{C}, X) \hookrightarrow A(\mathcal{D}, F(X)).$$

Given two quotients, $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{E}$, the composition $G \circ F : \mathcal{C} \rightarrow \mathcal{E}$ is also a quotient, and the inclusion satisfies $i_{G \circ F} = i_G \circ i_F$. These inclusions are known as *categorical inclusions*.

¹That is, *surjective*, as defined in [12, Definition 1.8.3].

According to [28, §4], citing a result from [8], for a quotient $F : \mathcal{C} \rightarrow \mathcal{D}$, there exists a connected commutative Frobenius algebra object L in $\mathcal{Z}(\mathcal{C})$ such that \mathcal{D} is equivalent to the category \mathcal{C}_L of right L -modules. Moreover, F is equivalent to the functor $F_L : \mathcal{C} \rightarrow \mathcal{C}_L$, where $x \mapsto x \otimes L$, and L is identified with its image under the forgetful functor. Using this, [28, §4] demonstrates an equivalence between the lattice of intermediate categorical inclusions between $A(\mathcal{C}, X)$ and $A(\mathcal{C}_L, X)$, and the lattice of Frobenius subalgebras of L . Since the Drinfeld center of a multi-fusion category is a fusion category [12, Exercise 7.13.7], we can apply the result from §8 to get the following:

Corollary 11.7. *The lattice of intermediate categorical inclusions between $A(\mathcal{C}, X)$ and $A(\mathcal{C}_L, X)$ is finite.*

This provides a rigidity-type result for abstract spin chains.

11.3. Vertex operator algebras. We posed the following question to Kenichi Shimizu in [52]:

Question 11.8. *Under what conditions is a vertex operator algebra (VOA) V a connected Frobenius algebra object in $\text{Rep}(V)$?*

Shimizu’s response can be understood through the categorical perspective on VOA extensions, as explored in [10]. This work highlights that a VOA extension A of V naturally forms a commutative algebra in the representation category $\text{Rep}(V)$. This algebra is characterized by two key properties: it is connected (or haploid; see Definition 2.13) and has a trivial twist, for the ribbon structure of $\text{Rep}(V)$.

Although it is not fully established that every VOA extension gives rise to a Frobenius algebra, a related partial result is known. Specifically, if \mathcal{C} is a modular tensor category and A is a connected, commutative, and exact algebra in \mathcal{C} , and if the category of local A -modules, $\mathcal{C}_A^{\text{loc}}$, forms a ribbon category with the same twist as \mathcal{C} , then A is a symmetric Frobenius algebra [53, Proposition 5.19]. However, this assumption may be too restrictive when applied to VOA extensions.

12. QUANTUM ARITHMETIC

In this section, we will explore additional applications of Watatani’s theorem from §8. Our focus will be on generalizing Ore’s theorem [40] and Euler’s totient theorem [39] in the context of fusion categories, as well as Robin’s reformulation of the Riemann hypothesis [49], which involves the sigma function. Each of these results requires a finite lattice, making Watatani’s theorem relevant.

For the first two, we will present the statements without proofs. While they may currently be considered conjectures, we believe their proofs could follow approaches similar to those used in [40] and [39]. It would be valuable to formalize these proofs in the future. Regarding the Riemann hypothesis, we do not claim that our generalization is relevant; rather, we suggest that it is a possibility.

Remark 12.1. *We caution the reader that this section is largely speculative. Speculation involves imagining possibilities or exploring ideas without a solid foundation in evidence or structure. In contrast, a conjecture is a more formalized hypothesis, often based on observed patterns or logical reasoning, and is typically intended to be proven or disproven.*

12.1. Ore’s theorem. Øystein Ore demonstrated in [38] in 1938 that a finite group is cyclic if and only if its subgroup lattice is distributive. He also extended this result in one direction as follows:

Theorem 12.2 ([38], Theorem 7). *Let $[H, G]$ be a distributive interval of finite groups. Then there exists an element $g \in G$ such that $\langle Hg \rangle = G$.*

The paper [40] generalizes Ore’s Theorem 12.2 to planar algebras as follows, and applies it to establish a connection between combinatorics and representation theory.

Theorem 12.3 ([40]). *Let \mathcal{P} be an irreducible subfactor planar algebra with a distributive biprojection lattice. Then there exists a minimal 2-box projection that generates the identity biprojection.*

We propose the following generalization to pseudo-unitary fusion categories, supported by Corollary 8.8.

Statement 12.4. *Let X be a connected Frobenius algebra object in a pseudo-unitary fusion category \mathcal{C} with a distributive Frobenius subalgebra lattice. Then there is a minimal idempotent in $\text{Hom}_{\mathcal{C}}(X, X)$ generating id_X .*

The elimination of the pseudo-unitary assumption is contingent upon Conjecture 8.10. Following the ordering described in §5, an idempotent p is termed *minimal* if, for any nonzero idempotent q such that $p \circ q = q = q \circ p$, it follows that $p = q$. Additionally, based on Proposition 4.15, a *biprojection* refers to the idempotent b_Y associated with a Frobenius subalgebra Y of X . The biprojection *generated* by an idempotent p is defined as the minimal biprojection b such that $b \geq p$. It would be advantageous to obtain a tensor category extension of [32, Theorem 4.12] in the following way, thereby offering a clearer reformulation of (generated) biprojections.

Statement 12.5. *A selfdual idempotent b is a biprojection if and only if $b * b = \lambda b$, for some nonzero scalar λ .*

12.2. Euler's totient. The traditional Euler's totient function, denoted as $\varphi(n)$, counts the number of positive integers up to n that are relatively prime to n . Let G be a finite group and μ the Möbius function associated with its subgroup lattice $\mathcal{L}(G)$. In 1936, Philip Hall established in [20] that the Euler totient of a group G , as defined below, corresponds to the cardinality of the set $\{g \in G \mid \langle g \rangle = G\}$:

$$\varphi(G) := \sum_{H \in \mathcal{L}(G)} \mu(H, G) |H|.$$

Consequently, if $\varphi(G)$ is nonzero, then G must be cyclic, and it follows that $\varphi(C_n) = \varphi(n)$. This result has been generalized to planar algebras as follows.

Theorem 12.6 ([39]). *Let \mathcal{P} be an irreducible subfactor planar algebra, and let μ denote the Möbius function of its biprojection lattice $[e_1, \text{id}]$. We introduce the Euler totient of the planar algebra \mathcal{P} as follows:*

$$\varphi(\mathcal{P}) := \sum_{b \in [e_1, \text{id}]} \mu(b, \text{id}) |b : e_1|.$$

If $\varphi(\mathcal{P})$ is nonzero, then there exists a minimal 2-box projection that generates the identity biprojection.

We propose the following generalization, which is subject to the same considerations outlined in §12.1, where $|Y| := \text{FPdim}(\underline{\text{Hom}}(Y, Y))$ generalizes the concept of the subfactor index, as discussed in [12, §7.25.2].

Statement 12.7. *Let X be a connected Frobenius algebra object in a pseudo-unitary fusion category \mathcal{C} , and let μ denote the Möbius function of its Frobenius subalgebra lattice, isomorphic to $[b_1, \text{id}_X]$. We introduce the Euler totient of X as follows:*

$$\varphi(X) := \sum_{b_Y \in [b_1, \text{id}_X]} \mu(b_Y, \text{id}_X) |Y|.$$

If $\varphi(X)$ is nonzero, then there is a minimal idempotent in $\text{Hom}_{\mathcal{C}}(X, X)$ generating id_X .

Question 12.8. *When $\varphi(X)$ is nonzero, can we also deduce that there exists a faithful simple component S of X , meaning that X is a subobject of $S^{\otimes n}$ for sufficiently large n ?*

The fact that $\varphi(n)$ is nonzero for any positive integer n prompts the following speculation:

Speculation 12.9. *If the interval $[b_1, \text{id}_X]$ is distributive, then $\varphi(X)$ is nonzero.*

Here is an application of Statement 12.7:

Speculation 12.10. *The minimum number of minimal idempotents that generate id_X is at most the minimal length ℓ of a chain given by*

$$b_1 < b_2 < \dots < b_{\ell+1} = \text{id}_X,$$

where $\varphi(b_i, b_{i+1})$, defined similarly to [39, Definition 4.3], is nonzero for each i .

12.3. Depth. We aim to broaden the concept of subfactor depth. Consider a Frobenius algebra object X within a monoidal category \mathcal{C} . We can assume, without loss of generality, that X generates \mathcal{C} . According to Remark 2.22, $X = Y \circ Y^\vee$ for some Y as described in Proposition 2.21. The *depth* of X is defined as the largest number $n = 2k$ or $2k + 1$ such that $X^{\otimes k}$ or $X^{\otimes k} \otimes Y$ includes a new simple subquotient, up to isomorphism.

Speculation 12.11. *The concept of depth, as defined above, is independent of the choice of Y .*

Speculation 12.12. *Let X_i be a Frobenius algebra in \mathcal{C}_i , with $i = 1, 2$. Then, $X_1 \boxtimes X_2$ is a Frobenius algebra in the Deligne tensor product $\mathcal{C}_1 \boxtimes \mathcal{C}_2$, and its depth is the maximum of the depths of X_1 and X_2 .*

According to [59], a finite index irreducible subfactor is a Hopf C^* -algebra subfactor if and only if it has depth 2. Drawing inspiration from this and Morita contexts of depth 2 in [35], we propose the following speculation:

Speculation 12.13. *A tensor category admits a fiber functor if and only if it is generated by a connected Frobenius algebra of depth 2.*

Recall by [12, Theorem 5.3.12] that a finite tensor category admits a fiber functor if and only if it is equivalent to $\text{Rep}(H)$ for some finite dimensional Hopf algebra H (and see [12, Theorem 5.4.1] for the infinite case).

12.4. Riemann hypothesis. The sigma function $\sigma(n) := \sum_{d|n} d$ is defined as the sum of the positive divisors of n . Let γ denote the Euler–Mascheroni constant. Then, we have [21, Theorem 323]:

$$\limsup_{n \rightarrow \infty} \frac{\sigma(n)}{n \log \log n} = e^\gamma.$$

In 1984, Guy Robin proved in [49] that the Riemann Hypothesis (RH) is true if and only if, for sufficiently large n ,

$$\sigma(n) < e^\gamma n \log \log n.$$

Let X be a connected Frobenius algebra object in a pseudo-unitary fusion category \mathcal{C} . We define the sigma function of X as:

$$\sigma(X) := \sum_{d \in \mathcal{D}(X)} d,$$

where $\mathcal{D}(X)$ is the divisor set of X , defined as $\mathcal{D}(X) := \{|Y| \mid b_Y \in [b_1, \text{id}_X]\}$. The set $\mathcal{D}(X)$ is finite, as shown in Corollary 8.8. We aim to extend (RH) by employing Robin’s reformulation for the class \mathfrak{C}_d of pairs (X, \mathcal{C}) , with X connected Frobenius algebra object of depth d (as defined in §12.3) in a pseudo-unitary \mathcal{C} it generates. It generalizes the quantum Riemann hypothesis proposed in [42], which serves as a revised version of the earlier, disproven version presented in [41]. Let us speculate a (RH) of depth d , denoted (RH_d) .

Speculation 12.14. *For all $d \geq 2$, there is a constant γ_d such that:*

$$\limsup_{(X, \mathcal{C}) \in \mathfrak{C}_d, |X| \rightarrow \infty} \frac{\sigma(X)}{|X| \log \log |X|} = e^{\gamma_d},$$

Furthermore, for all $(X, \mathcal{C}) \in \mathfrak{C}_d$ with $|X|$ large enough,

$$\sigma(X) < e^{\gamma_d} |X| \log \log |X|.$$

Let \mathcal{I}_d be the set $\{|X| \mid (X, \mathcal{C}) \in \mathfrak{C}_d\}$. Using Speculation 12.12:

Speculation 12.15. *The (RH_d) as stated in Speculation 12.14 implies that \mathcal{I}_d countable without accumulation point, and that γ_d is strictly increasing in d .*

Question 12.16. *Can we further infer from (RH_d) that there is a minimum gap length in \mathcal{I}_d ?*

The following holds true for irreducible depth 2 subfactors, by Ore’s theorem and the proof of Proposition 12.19.

Speculation 12.17. *Speculation 12.14 can be simplified by focusing on the subclass \mathfrak{D}_d which consists of $(X, \mathcal{C}) \in \mathfrak{C}_d$ where the Frobenius subalgebra lattice of X is distributive.*

Given an intermediate subfactor $N \subset P \subset M$, we derive two subfactors: $N \subset P$ and $P \subset M$. However, the Frobenius subalgebra associated with the intermediate subfactor pertains solely to $N \subset P$. It is necessary to explore the categorical generalization of $P \subset M$ and the algebraic integer $\frac{|M:N|}{|M:P|} = |P:N|$, leading to:

Speculation 12.18. *Let Y be a Frobenius subalgebra of a Frobenius algebra X in \mathcal{C} . Then $\frac{|X|}{|Y|}$ is an algebraic integer.*

Proposition 12.19. *Assuming Speculations 12.13 and 12.18, (RH_2) is equivalent to (RH).*

Proof. In this case, $\mathcal{D}(X)$ is a subset of $\mathcal{D}(|X|)$, the set of divisors of the integer $|X|$. Consequently, $\sigma(X) \leq \sigma(|X|)$. However, equality is attained when X is the connected Frobenius algebra associated with the irreducible cyclic group subfactor of index $n = |X|$. This is due to the one-to-one correspondence among the divisors m of n , the subgroups C_m of C_n , and the Frobenius subalgebras. \square

ACKNOWLEDGMENTS

We extend our gratitude to Keshab Chandra Bakshi, Dave Benson, Pavel Etingof, Jürgen Fuchs, Shamindra Ghosh, Dave Penneys, Maxime Ramzi, Ingo Runkel, Will Sawin, Brian Shin, and Serge Skryabin for insightful discussions. The names are listed in alphabetical order by first name. Mainak Ghosh’s work is supported by the BJNSF (Grant No. 1S24063). Sebastien Palcoux’s work is supported by the NSFC (Grant No. 12471031).

REFERENCES

- [1] K. S. BAKSHI, S. DAS, Z. LIU, Y. REN, *An angle between intermediate subfactors and its rigidity*. Trans. Amer. Math. Soc. 371 (2019), no. 8, 5973–5991, and arXiv:1710.00285.
- [2] J.W. BARRETT, B.W. WESTBURY, *Spherical categories*. Adv. Math. 143 (1999), no. 2, 357–375.
- [3] D. BENSON, *Pullback of monomorphisms between selfdual objects in a tensor category*, <https://mathoverflow.net/a/478229/34538>
- [4] D. BENSON, *Intersection of Frobenius subalgebra objects*, <https://mathoverflow.net/a/478374/34538>
- [5] D. BISCH, *A note on intermediate subfactors*. Pacific J. Math. 163 (1994), no. 2, 201–216.
- [6] M. BISCHOFF, A. DAVYDOV, *Hopf algebra actions in tensor categories*. Transform. Groups 26 (2021), no. 1, 69–80.
- [7] M. BISCHOFF, Y. KAWAHIGASHI, R. LONGO, R.-H. REHREN, *Tensor categories and endomorphisms of von Neumann algebras—with applications to quantum field theory*. Springer Briefs in Mathematical Physics, 3. Springer, Cham, 2015. x+94 pp.
- [8] A. BRUGUIÈRES, S. NATALE, *Exact sequences of tensor categories*. International Mathematics Research Notices, 2011:5644–5705, 6 2010.
- [9] Q. CHEN, R. HERNÁNDEZ PALOMARES, C. JONES, D. PENNEYS, *Q-system completion for C^* 2-categories*. J. Funct. Anal. 283 (2022), no. 3, Paper No. 109524, 59 pp.
- [10] T. CREUTZIG, S. KANADE, R. MCRAE, *Tensor categories for vertex operator superalgebra extensions*. Mem. Amer. Math. Soc. 295 (2024), no. 1472, vi+181 pp.
- [11] P. ETINGOF, *Corrections to the book “Tensor categories”, 2023*, <https://math.mit.edu/~etingof/tensbookcor.pdf>
- [12] P. ETINGOF, S. GELAKI, D. NIKSHYCH, AND V. OSTRIK, *Tensor Categories*, American Mathematical Society, (2015). Mathematical Surveys and Monographs Volume 205.
- [13] P. ETINGOF, V. OSTRIK, *On semisimplification of tensor categories*. Representation theory and algebraic geometry, 3–35, Trends Math., Birkhäuser/Springer, Cham, 2022
- [14] P. ETINGOF, C. WALTON, *Semisimple Hopf actions on commutative domains*. Adv. Math. 251 (2014), 47–61.
- [15] P. FREYD, *Abelian categories. An introduction to the theory of functors*. Harper’s Series in Modern Mathematics. Harper & Row, Publishers, New York, 1964. xi+164 pp.
- [16] J. FUCHS, I. RUNKEL, C. SCHWEIGERT, *TFT construction of RCFT correlators. I. Partition functions*. Nuclear Phys. B 646 (2002), no. 3, 353–497.
- [17] J. FUCHS, C. SCHWEIGERT, C. STIGNER, *Modular invariant Frobenius algebras from ribbon Hopf algebra automorphisms*. J. Algebra 363 (2012), 29–72.
- [18] J. Fuchs, C. Stigner, *On Frobenius algebras in rigid monoidal categories*. Arab. J. Sci. Eng. Sect. C Theme Issues 33 (2008), no. 2, 175–191.
- [19] P. GROSSMAN, V.F.R. JONES, *Intermediate subfactors with no extra structure* J. Amer. Math. Soc. 20 (2007), no. 1, 219–265.
- [20] P. HALL, *The Eulerian functions of a group*. The Quarterly Journal of Mathematics, Volume os-7, Issue 1, 1936, Pages 134–151.
- [21] G.H. HARDY, E.M. WRIGHT, *An introduction to the theory of numbers*. Sixth edition. Revised by D. R. Heath-Brown and J. H. Silverman. With a foreword by Andrew Wiles. Oxford University Press, Oxford, 2008. xxii+621 pp.
- [22] P. DE LA HARPE, *On simplicity of reduced C^* -algebras of groups*. Bull. Lond. Math. Soc. 39 (2007), no. 1, 1–26.
- [23] S. INO, Y. WATATANI, *Perturbations of intermediate C^* -subalgebras for simple C^* -algebras*. Bull. Lond. Math. Soc. 46 (2014), no. 3, 469–480.
- [24] M. IZUMI, R. LONGO, S. POPA, SORIN, *A Galois correspondence for compact groups of automorphisms of von Neumann algebras with a generalization to Kac algebras*. J. Funct. Anal. 155 (1998), no. 1, 25–63.
- [25] V.F.R. JONES, *Actions of finite groups on the hyperfinite type II_1 factor*. Mem. Amer. Math. Soc. 28 (1980), no. 237, v+70 pp.
- [26] V.F.R. JONES, *Index for subfactors*. Invent. Math. 72 (1983), no. 1, 1–25.
- [27] C. JONES, D. PENNEYS, *Q-systems and compact W^* -algebra objects*. Topological phases of matter and quantum computation, 63–88, Contemp. Math., 747, Amer. Math. Soc. (2020)
- [28] C. JONES, K. SCHATZ, D. J. WILLIAMSON, *Quantum cellular automata and categorical duality of spin chains*, arXiv:2410.08884
- [29] V. JONES, V.S. SUNDER, *Introduction to subfactors*. London Mathematical Society Lecture Note Series, 234. Cambridge University Press, Cambridge, 1997. xii+162 pp.
- [30] Z. LANDAU, *Exchange relation planar algebras*. Proceedings of the Conference on Geometric and Combinatorial Group Theory, Part II (Haifa, 2000). Geom. Dedicata 95 (2002), 183–214.
- [31] R.G. LARSON, M.E. SWEEDLER, *An associative orthogonal bilinear form for Hopf algebras*. Amer. J. Math. 91 (1969), 75–94.
- [32] Z. LIU, *Exchange relation planar algebras of small rank*. Trans. Amer. Math. Soc. 368 (2016), no. 12, 8303–8348.
- [33] Z. LIU, S. PALCOUX, Y. REN, *Triangular prism equations and categorification*, arXiv:2203.06522
- [34] R. LONGO, *A duality for Hopf algebras and for subfactors. I*. Comm. Math. Phys. 159 (1994), no. 1, 133–150.
- [35] M. MÜGER, *From subfactors to categories and topology. I. Frobenius algebras in and Morita equivalence of tensor categories*. J. Pure Appl. Algebra 180 (2003), no. 1-2, 81–157.
- [36] S.H. NG, P. SCHAUBURG, *Higher Frobenius-Schur indicators for pivotal categories*. Hopf algebras and generalizations, 63–90, Contemp. Math., 441, Amer. Math. Soc., 2007.
- [37] W.D. NICHOLS, *Bialgebras of type one*. Comm. Algebra 6 (1978), no. 15, 1521–1552.
- [38] O. ORE, *Structures and group theory. II*. Duke Math. J. 4 (1938), no. 2, 247–269.
- [39] S. PALCOUX, *Euler totient of subfactor planar algebras*. Proc. Amer. Math. Soc. 146 (2018), no. 11, 4775–4786.
- [40] S. PALCOUX, *Ore’s theorem on subfactor planar algebras*. Quantum Topol. 11 (2020), no. 3, 525–543.
- [41] S. PALCOUX, *On a quantum Riemann Hypothesis*, URL (version: 2021-02-28): <https://mathoverflow.net/q/308300/34538>
- [42] S. PALCOUX, *On a revised quantum Riemann hypothesis*, URL (version: 2021-02-28): <https://mathoverflow.net/q/364311/34538>
- [43] S. PALCOUX, *Frobenius identity on finite-dimensional Hopf algebras*, URL (version: 2024-12-25): <https://mathoverflow.net/q/484771>
- [44] B. PAREIGIS, *When Hopf algebras are Frobenius algebras*. J. Algebra 18 (1971), 588–596.
- [45] H.H. PHÙNG, *An embedding theorem for abelian monoidal categories*. Compositio Math. 132 (2002), no. 1, 27–48.

- [46] D.E. RADFORD, *On Kauffman's knot invariants arising from finite-dimensional Hopf algebras*. Advances in Hopf algebras, 205–266, Lecture Notes in Pure and Appl. Math., 158, Dekker, New York, 1994.
- [47] M. RAMZI, *Tensor product of intersections in an abelian rigid monoidal category*, <https://mathoverflow.net/a/478863/34538>, 2024-09-14.
- [48] J. RICKARD, *Tensor product of monomorphisms is a monomorphism?*, <https://mathoverflow.net/a/232343/34538>
- [49] G. ROBIN, *Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann*. (French) [[Large values of the sum-of-divisors function and the Riemann hypothesis]] J. Math. Pures Appl. (9) 63 (1984), no. 2, 187–213.
- [50] S. ROMAN, *Advanced linear algebra*. Third edition. Graduate Texts in Mathematics, 135. Springer, New York, 2008. xviii+522 pp.
- [51] W. SAWIN, *Do Frobenius subalgebras form a lattice?*, <https://mathoverflow.net/a/481533/34538>
- [52] K. SHIMIZU, *Rigidity of the category of local modules*, BIMS-Tsinghua Quantum Symmetry Seminar, 2024-12-19, <https://youtu.be/-T6UmspdIUM>
- [53] K. SHIMIZU, H. YADAV, *ommutative exact algebras and modular tensor categories*, <https://arxiv.org/pdf/2408.06314>
- [54] B. SHIN, *Intersection of subalgebras in an abelian monoidal category*, <https://mathoverflow.net/a/478656/34538>, 2024-09-11
- [55] A. SCHOPIERAY, *Non-pseudounitary fusion*. J. Pure Appl. Algebra 226 (2022), no. 5, Paper No. 106927, 19 pp.
- [56] A. SKOWROŃSKI, K. YAMAGATA, *Frobenius algebras. I. Basic representation theory*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2011. xii+650 pp.
- [57] S. SKRYABIN, *Projectivity and freeness over comodule algebras*. Trans. Amer. Math. Soc. 359 (2007), no. 6, 2597–2623.
- [58] R. STREET, *Frobenius monads and pseudomonoids*. J. Math. Phys. 45 (2004), no. 10, 3930–3948.
- [59] W. SZYMAŃSKI, *Finite index subfactors and Hopf algebra crossed products*. Proc. Amer. Math. Soc. 120 (1994), no. 2, 519–528.
- [60] T. TIMMERMANN, *An invitation to quantum groups and duality. From Hopf algebras to multiplicative unitaries and beyond*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2008. xx+407 pp
- [61] A. VAN DAELE, *The Fourier transform in quantum group theory*. New techniques in Hopf algebras and graded ring theory, 187–196, K. Vlaam. Acad. Belgie Wet. Kunsten (KVAB), Brussels, 2007.
- [62] Y. WATATANI, *Index for C^* -subalgebras*. Mem. Amer. Math. Soc. 83 (1990), no. 424, vi+117 pp.
- [63] Y. WATATANI, *Lattices of intermediate subfactors*. J. Funct. Anal. 140 (1996), no. 2, 312–334.
- [64] J. WATROUS, *On one-dimensional quantum cellular automata*. 36th Annual Symposium on Foundations of Computer Science (1995), 528–537, IEEE Comput. Soc. Press.

M. GHOSH, BEIJING INSTITUTE OF MATHEMATICAL SCIENCES AND APPLICATIONS, HUAIROU DISTRICT, BEIJING, CHINA
Email address: mainakghosh@bimsa.cn

S. PALCOUX, BEIJING INSTITUTE OF MATHEMATICAL SCIENCES AND APPLICATIONS, HUAIROU DISTRICT, BEIJING, CHINA
Email address: sebastien.palcoux@gmail.com
URL: <https://sites.google.com/view/sebastienpalcoux>