

ALGEBRAIC STUDY ON PERMUTATION GRAPHS

ANTONINO FICARRA, SOMAYEH MORADI

ABSTRACT. Let G be a permutation graph. We show that G is Cohen-Macaulay if and only if G is unmixed and vertex decomposable. When this is the case, we obtain a combinatorial description for the a -invariant of G . Moreover, we characterize the Gorenstein permutation graphs.

INTRODUCTION

Permutation graphs arise naturally in combinatorics and graph theory. They are characterized as those graphs which are both comparability and co-comparability graphs of posets [25], making them an intriguing subject of algebraic and combinatorial investigation. For a finite simple graph G on n vertices and with the edge set $E(G)$, the *edge ideal* of G , introduced by Villarreal [29], is the ideal of the polynomial ring $S = K[x_i : i \in V(G)]$ over a field K defined as $I(G) = (x_i x_j : \{i, j\} \in E(G))$. The main theme in the study of edge ideals is to translate the algebraic properties of the ring $S/I(G)$ to the combinatorics of the underlying graph G and vice versa. The study of the Cohen-Macaulay property of graphs has been well-established for various classes of graphs such as bipartite graphs [10], very well-covered graphs [5, 19], chordal graphs [15], Cameron-Walker graphs [17], fully-whiskered graphs [4] and graphs of girth at least five [1]. For the aforementioned classes of graphs it is shown that G is Cohen-Macaulay if and only if G is unmixed and vertex decomposable.

Vertex decomposable simplicial complexes were introduced by Provan and Billera in [26]. Their recursive definition allows to determine algebraic invariants of their Stanley-Reisner rings inductively, see [22]. A graph G is called *vertex decomposable* if the independence complex of G is vertex decomposable. These graphs were first considered by Dochtermann-Engström [6] and Woodroffe [30]. Any unmixed, vertex decomposable graph is Cohen-Macaulay, but the converse does not hold in general.

In this work, we study the Cohen-Macaulay and the Gorenstein properties for permutation graphs. These graphs were first introduced in [7] and [25]. They form a subclass of weakly chordal graphs, as was shown in [8]. Different characterizations of permutation graphs are given in [8, 18, 25]. In Theorem 2.3 we show that a permutation graph is Cohen-Macaulay if and only if it is unmixed and vertex decomposable. To this aim, we use a characterization of Cohen-Macaulay permutation graphs given in [3, Theorem 1.1] in terms of the maximal cliques of the graph. This implies that the cover ideal $J(G)$ of a Cohen-Macaulay permutation graph is vertex splittable (Corollary 2.5) and that the Rees algebra $\mathcal{R}(J(G))$ and the toric algebra $K[J(G)]$ are normal Cohen-Macaulay domains (Corollary 2.6).

2020 *Mathematics Subject Classification.* Primary 13C05, 13C14; Secondary 05E40.

Key words and phrases. Cohen-Macaulay, Gorenstein, permutation graph, edge ideal.

In Theorem 2.7 we characterize the Gorenstein permutation graphs. To prove it, we use a result by Oboudi and Nikseresht [24] regarding the Gorenstein graphs. In Proposition 2.8 we obtain the a -invariant of a Cohen-Macaulay permutation graph in terms of the induced matching number and the vertex cover number of G . As a consequence we determine when $I(G)$ is Hilbertian, that is the Hilbert function and the Hilbert polynomial of $S/I(G)$ coincide. We conclude the paper with Proposition 2.9 which gives a combinatorial description for bi-Cohen-Macaulay graphs.

1. PRELIMINARIES

In this section, we recall some concepts and introduce some notation which are needed in the sequel. Throughout, G is a finite simple graph with the vertex set $V(G) = [n] = \{1, 2, \dots, n\}$ and the edge set $E(G)$, and $S = K[x_1, \dots, x_n]$ is the polynomial ring over a field K . The *edge ideal* of G is defined as the ideal of S ,

$$I(G) = (x_i x_j : \{i, j\} \in E(G)).$$

The graph G is called *Cohen-Macaulay*, respectively *Gorenstein*, if $S/I(G)$ is a Cohen-Macaulay, respectively Gorenstein ring. A subset $F \subseteq V(G)$ is called an *independent set* of G , if it contains no edge of G . The maximal cardinality of independent sets of G is denoted by $\alpha(G)$.

A *vertex cover* of G is a subset $C \subseteq V(G)$ which intersects all the edges of G and a vertex cover which is minimal with respect to inclusion is called a *minimal vertex cover* of G . The graph G is called *unmixed* if all the minimal vertex covers of G have the same cardinality. The *vertex cover number* of G is defined as the minimum cardinality of the vertex covers of G and is denoted by $\tau(G)$. A subset $A \subseteq V(G)$ is called a *clique* of G if any two vertices in A are adjacent in G . A *maximal clique* is a clique of G which is not contained in any other clique of G .

A *matching* of G is a subset of $E(G)$ consisting of pairwise disjoint edges of G . The maximum size of matchings of G is denoted by $m(G)$. We say that the edges e and e' form a *gap* in G , if they are disjoint and no vertex in e is adjacent to a vertex in e' . A subset E of edges *forms a gap* in G , when each two elements in E form a gap in G . The maximum cardinality of a set $E \subseteq E(G)$ which forms a gap is called the *induced matching number* of G and is denoted by $\text{im}(G)$.

For a graph G , the *complementary graph* of G is the graph G^c with the same vertex set as G whose edges are the non-edges of G .

For a simplicial complex Δ and a face $F \in \Delta$, the *link* of F in Δ is defined as

$$\text{lk}_\Delta(F) = \{G \in \Delta : G \cap F = \emptyset, G \cup F \in \Delta\},$$

and the *deletion* of F is the simplicial complex

$$\text{del}_\Delta(F) = \{G \in \Delta : G \cap F = \emptyset\}.$$

A simplicial complex Δ is called *vertex decomposable* if either Δ is a simplex, or Δ contains a vertex x such that

- (i) both $\text{del}_\Delta(x)$ and $\text{lk}_\Delta(x)$ are vertex decomposable, and
- (ii) any facet of $\text{del}_\Delta(x)$ is a facet of Δ .

A vertex x which satisfies condition (ii) is called a *shedding vertex* of Δ .

The *independence complex* of a graph G is defined as the simplicial complex

$$\Delta_G = \{F \subseteq V(G) : F \text{ is an independent set of } G\}.$$

The graph G is called *vertex decomposable* if Δ_G is vertex decomposable.

Vertex decomposability has a nice translation to independence complexes of graphs. For a vertex $i \in V(G)$, let $N_G(i)$ be the set of all vertices of G adjacent to i and let $N_G[i] = N_G(i) \cup \{i\}$. Translating the definition of vertex decomposable to independence complexes of graphs we have that:

A graph G is vertex decomposable, if either G consists of isolated vertices or it has a vertex i such that

- (i) $G \setminus \{i\}$ and $G \setminus N_G[i]$ are vertex decomposable.
- (ii) Any maximal independent set of $G \setminus \{i\}$ is a maximal independent set of G .

It can be easily seen that (ii) is equivalent to say that no independent set of $G \setminus N_G[i]$ is a maximal independent set of $G \setminus \{i\}$. Such a vertex i is called a *shedding vertex* of G .

Let $\sigma = (k_1, \dots, k_n)$ be a permutation of $[n]$ that is $\sigma(i) = k_i$ for all i . The *permutation graph* $G(\sigma)$ corresponding to σ is the graph on the vertex set $[n]$ for which $\{i, j\} \in E(G(\sigma))$ if and only if $i < j$ and j appears before i in the list k_1, \dots, k_n . For example if $\sigma = (2, 4, 5, 1, 3)$, then $G(\sigma)$ is the graph on $[5]$ with the edge set $\{\{1, 2\}, \{1, 4\}, \{3, 4\}, \{1, 5\}, \{3, 5\}\}$.

For a poset (P, \prec) with the vertex set $V = \{v_1, \dots, v_n\}$, the *comparability graph* $\text{comp}(P)$ of P is defined to be the graph on V with $\{v_i, v_j\} \in E(\text{comp}(P))$ if and only if v_i and v_j are comparable in P and $v_i \neq v_j$. Whereas, the *co-comparability graph* $\text{co-comp}(P)$ of P is defined to be the graph on V with $\{v_i, v_j\} \in E(\text{co-comp}(P))$ if and only if v_i and v_j are not comparable in P . In other words, $\text{co-comp}(P)$ is the complementary graph of $\text{comp}(P)$. By [25, Theorem 3], a graph G is a permutation graph if and only if G is both a comparability and a co-comparability graph.

For two elements x, y in a poset (P, \prec) , we write $x \prec y$, whenever $x \prec y$ and if z is an element in P with $x \preceq z \preceq y$, then $z = x$ or $z = y$.

A graph G is called *weakly chordal* if G and G^c have no induced cycles of length $m \geq 5$. A cycle in G is called an *induced cycle* if no two non-consecutive vertices in the cycle are adjacent in G .

We denote by K_n the complete graph on n vertices and by P_n the path graph on n vertices. Notice that K_n and P_n^c are permutation graphs.

2. COHEN-MACAULAY AND GORENSTEIN PERMUTATION GRAPHS

Permutation graphs are characterized in terms of the existence of a so-called cohesive order on their vertex sets. A graph G is said to have a *cohesive order* if there is a labeling $[n]$ on $V(G)$ such that

- (i) If $i < j < k$ and $\{i, j\} \in E(G)$, $\{j, k\} \in E(G)$, then $\{i, k\} \in E(G)$.
- (ii) If $i < j < k$ and $\{i, k\} \in E(G)$, then $\{i, j\} \in E(G)$ or $\{j, k\} \in E(G)$.

Theorem 2.1. [9, Theorem 2.3] *A graph G is a permutation graph if and only if it has a cohesive order.*

The following characterization of Cohen-Macaulay permutation graphs by Cheri, et al. [3] will be extensively used in the proof of Theorem 2.3.

Theorem 2.2. [3, Theorem 1.1] *Let G be a permutation graph. Then the following statements are equivalent.*

- (i) G is Cohen-Macaulay.
- (ii) G is unmixed and there exists a unique way of partitioning $V(G)$ into r disjoint maximal cliques, where r is the cardinality of a maximal independent set of G .

Now, we are in the position to prove

Theorem 2.3. *Let G be a permutation graph. The following statements are equivalent.*

- (i) G is Cohen-Macaulay.
- (ii) G is unmixed and vertex decomposable.
- (iii) G is unmixed and shellable.

Proof. (i) \Rightarrow (ii): Without loss of generality we assume that G has no isolated vertices. By Theorem 2.1, we may assume that $V(G) = [n]$ is a cohesive order of G in the given labeling. For $i, j \in [n]$, we set $i \prec j$, if $i < j$ and $\{i, j\} \in E(G)$. It follows from the property (i) of cohesive order that $(V(G), \prec)$ is a poset. We denote this poset by P . Then it is clear that $G = \text{comp}(P)$, and the maximal cliques of G are just the maximal chains in P .

Since G is Cohen-Macaulay, it is unmixed. We may assume that the cardinality of any maximal independent set of G is r . By Theorem 2.2, $V(G)$ is partitioned in a unique way into r disjoint maximal chains in P . Let $V(G) = A_1 \cup \dots \cup A_r$ be this unique partition. Let $j_k = \max A_k$ and $i_k \in A_k$ be the unique element in A_k with $i_k \prec j_k$ for $1 \leq k \leq r$.

Claim. There exists an integer $1 \leq t \leq r$, such that

$$\{s \in [n] : i_t \prec s\} = \{j_t\}.$$

Proof of the claim. Suppose on the contrary that this is not the case. Then for any $1 \leq k \leq r$, there exists $1 \leq \ell_k \leq r$ with $\ell_k \neq k$ such that $i_k \prec j_{\ell_k}$. Without loss of generality we may assume that $i_1 \prec j_2$.

First suppose that $i_2 \prec j_1$. We show that $B_1 = (A_1 \setminus \{j_1\}) \cup \{j_2\}$ and $B_2 = (A_2 \setminus \{j_2\}) \cup \{j_1\}$ are maximal chains of P . Since $i_1 \prec j_2$ and $i_2 \prec j_1$, B_1 and B_2 are chains of P . Suppose that B_1 is not a maximal chain of P . This would mean that there exists $s \in [n]$ with $i_1 \prec s \prec j_2$. Let F be a maximal independent set of G which contains s . Since $|F| = r$, F contains precisely one element from each A_k . Since $i_1 \prec s$ and $s \in F$, it follows that $F \cap A_1 = \{j_1\}$. Now, from $j_1 \in F$ and $i_2 \prec j_1$, it follows that $F \cap A_2 = \{j_2\}$. Thus $s, j_2 \in F$. This contradicts to $s \prec j_2$. Therefore, B_1 is a maximal chain of P . The same argument shows that B_2 is a maximal chain of P . Then $V(G) = B_1 \cup B_2 \cup A_3 \cup \dots \cup A_r$ is another partition of $V(G)$ into maximal chains of P and this contradicts to Theorem 2.2.

So we have $i_2 \not\prec j_1$. Then by our assumption we may assume that $i_2 \prec j_3$. Similar to the argument in the previous paragraph, if $i_3 \prec j_1$, then $V(G) = B_1 \cup B_2 \cup$

$B_3 \cup A_4 \cup \cdots \cup A_r$ is another partition of $V(G)$ into maximal chains of P , where $B_1 = (A_1 \setminus \{j_1\}) \cup \{j_2\}$, $B_2 = (A_2 \setminus \{j_2\}) \cup \{j_3\}$ and $B_3 = (A_3 \setminus \{j_3\}) \cup \{j_1\}$, and this again contradicts to Theorem 2.2. Moreover, if $i_3 \prec j_2$, then $V(G) = A_1 \cup B_2 \cup B_3 \cup A_4 \cup \cdots \cup A_r$ is another partition of $V(G)$ into maximal chains of P , where $B_2 = (A_2 \setminus \{j_2\}) \cup \{j_3\}$ and $B_3 = (A_3 \setminus \{j_3\}) \cup \{j_2\}$, which is absurd. Hence, we may assume $i_3 \prec j_4$.

Proceeding with the same argument, and after relabeling we obtain that $i_k \prec j_{k+1}$ for $1 \leq k \leq r-1$. Our assumption implies that $i_r \prec j_s$ for some $s < r$. We set $B_k = (A_k \setminus \{j_k\}) \cup \{j_{k+1}\}$ for $s \leq k \leq r-1$ and $B_r = (A_r \setminus \{j_r\}) \cup \{j_s\}$. We show that each B_k is a maximal chain of P . To simplify the notation, we set $j_{r+1} = j_s$. Fix an integer $s \leq q \leq r$ and suppose that B_q is not a maximal chain of P . Then $i_q \prec m \prec j_{q+1}$ for some m . We let F be a maximal independent set of G which contains m . Since F contains precisely one element from each of A_1, \dots, A_r , from $i_q \prec m$, we obtain $F \cap A_q = \{j_q\}$. Since $i_{q-1} \prec j_q$ and $j_q \in F$, we get $F \cap A_{q-1} = \{j_{q-1}\}$. Similarly, the relations $i_k \prec j_{k+1}$, imply that $F \cap A_k = \{j_k\}$ for all $s \leq k \leq r$. Therefore, $m, j_{q+1} \in F$. This contradicts to $m \prec j_{q+1}$. Thus we have proved that each B_k is a maximal chain of P . Having this, we obtain the partition $(\bigcup_{i=1}^{s-1} A_i) \cup (\bigcup_{i=s}^r B_i)$ of $V(G)$ into maximal chains of P , which is different from $\bigcup_{i=1}^r A_i$. This contradicts to Theorem 2.2. So our claim is proved. \square

Let t be an integer satisfying the claim. Without loss of generality we let $t = 1$. We prove that j_1 is a shedding vertex of G so that $G' = G \setminus \{j_1\}$ and $G'' = G \setminus N_G[j_1]$ are vertex decomposable. This will show that G is vertex decomposable. Since any induced subgraph of a permutation graph is a permutation graph, G' and G'' are permutation graphs. We show that they are Cohen-Macaulay. Then by induction on the number of vertices of the graph, it follows that they are vertex decomposable. By [29, Proposition 4.3], G'' is Cohen-Macaulay.

To show that G' is Cohen-Macaulay, we set P' to be the poset obtaining from P by removing j_1 . Then $P' = \text{comp}(G')$. By our assumption on i_1 , the chain $A'_1 = A_1 \setminus \{j_1\}$ is a maximal chain of P' . Thus $V(G') = A'_1 \cup A_2 \cup \cdots \cup A_r$ is a partition of $V(G')$ into maximal chains of P' . We show that $V(G')$ is uniquely partitioned into maximal chains of P' . Let $V(G') = C_1 \cup C_2 \cup \cdots \cup C_p$ be an arbitrary partition of $V(G')$ into maximal chains C_k of P' . First notice that the set consisting of the maximal elements of C_1, \dots, C_p is an independent set of G' and hence an independent set of G of cardinality p . Therefore, $p \leq r$. Moreover, by the assumption on i_1 , we know that i_1 is a maximal element of P' . So it is the maximal element of some maximal chain, let say C_1 . Then $C_1 \cup \{j_1\}$ is a maximal chain of P , since $i_1 \prec j_1$. We show that each C_k for $2 \leq k \leq p$ is a maximal chain of P . Suppose this is not the case. Then $C_h \cup \{j_1\}$ is a chain in P for some $2 \leq h \leq p$. Now, consider a maximal independent set F of G with $j_1 \in F$. Since F has at most one element from each chain of P and $C_1 \cup \{j_1\}$ and $C_h \cup \{j_1\}$ are chains in P , we conclude that $F \cap C_1 = F \cap C_h = \emptyset$ and $|F \cap C_\ell| \leq 1$ for all ℓ . Thus

$$r-1 = |F \setminus \{j_1\}| = \sum_{\ell=1}^p |F \cap C_\ell| \leq p-2.$$


This contradicts to $p \leq r$. So C_2, \dots, C_p are maximal chains of P , and hence $V(G) = (C_1 \cup \{j_1\}) \cup C_2 \cup \dots \cup C_p$ is a partition of $V(G)$ into maximal chains of P . Since such a partition is unique, we obtain $p = r$, $A_1 = C_1 \cup \{j_1\}$ and, after a suitable relabeling, $A_k = C_k$ for $2 \leq k \leq r$. So $V(G')$ is uniquely partitioned into maximal chains as $V(G') = A'_1 \cup A_2 \cup \dots \cup A_r$.

Next, we show that any maximal independent set of G' is a maximal independent set of G . This will show that j_1 is a shedding vertex of G and that G' is unmixed. Consider a maximal independent set F of G' . If $i_1 \in F$, then $F \cup \{j_1\}$ is not an independent set of G . In other words, F is a maximal independent set of G .

Now, assume that $i_1 \notin F$. Since F is maximal, this means that $b \in F$ for some $b \in N_{G'}(i_1)$. Otherwise, $F \cup \{i_1\}$ would be an independent set of G' which strictly contains F . By our assumption on i_1 , we have $N_G[i_1] \subseteq N_G[j_1]$. Indeed, it follows from the equality $\{s \in [n] : i_1 \prec s\} = \{j_1\}$ that if $\{s, i_1\} \in E(G)$ with $s \neq j_1$, then we have $s \prec i_1$. So $s \prec i_1 \prec j_1$, and hence $\{s, j_1\} \in E(G)$. So $N_G[i_1] \subseteq N_G[j_1]$. Therefore, $b \in N_G[j_1]$. Since $b \neq j_1$ we have $b \in N_G(j_1) \cap F$. This shows that $F \cup \{j_1\}$ is not an independent set of G . So F is a maximal independent set of G . This implies that G' is unmixed and j_1 is a shedding vertex of G . Now, by Theorem 2.2, we conclude that G' is Cohen-Macaulay. So by induction, G' is vertex decomposable. The proof is complete.

(ii) \Rightarrow (iii) and (iii) \Rightarrow (i) follow from [2, Theorem 11.3] and [11, Theorem 8.2.6], respectively. \square

Remark 2.4. *A permutation graph is not necessarily vertex decomposable. Indeed, the graph G depicted below is an unmixed permutation graph, which is not vertex decomposable. It follows from Theorem 2.3 that G is not Cohen-Macaulay.*



Vertex splittable ideals were defined in [22]. They appear as the Alexander duals of the Stanley-Reisner ideals of vertex decomposable simplicial complexes. A monomial ideal $I \subset S$ is called *vertex splittable* if it can be obtained by the following recursive procedure.

- (i) If u is a monomial and $I = (u)$, $I = (0)$ or $I = S$, then I is vertex splittable.
- (ii) If there is a variable x_i and vertex splittable ideals I_1 and I_2 of $K[X \setminus \{x_i\}]$ so that $I = x_i I_1 + I_2$, $I_2 \subseteq I_1$ and $\mathcal{G}(I)$ is the disjoint union of $\mathcal{G}(x_i I_1)$ and $\mathcal{G}(I_2)$, then I is vertex splittable.

The *cover ideal* $J(G)$ of a graph G is defined as the monomial ideal generated by those monomials whose support is a vertex cover of G .

The following corollary is obtained from Theorem 2.3 and [22, Theorem 2.2].

Corollary 2.5. *Let G be an unmixed permutation graph, and let $J(G)$ be the cover ideal of G . The following are equivalent:*

- (i) $J(G)$ is vertex splittable.

- (ii) $J(G)$ has linear quotients.
- (iii) G is Cohen-Macaulay.

We expect that for a Cohen-Macaulay permutation graph G , all powers of the cover ideal $J(G)$ have linear resolution.

Corollary 2.6. *Let G be a Cohen-Macaulay permutation graph. Then*

- (a) *The Rees algebra $\mathcal{R}(J(G))$ and the toric algebra $K[J(G)]$ are normal Cohen-Macaulay domains*
- (b) *$J(G)$ satisfies the strong persistence property.*
- (c) $\lim_{k \rightarrow \infty} \text{depth } S/J(G)^k = n - \ell(J(G))$.
- (d) $\text{reg}(K[I(G)]) \leq m(G)$.
- (e) $m(G) \leq \text{reg}(\mathcal{R}(I(G))) \leq m(G) + 1$.

Proof. (a), (b) and (c) follow from Theorem 2.3 and [21, Theorem 3.1]. Whereas, (d) and (e) follow from Theorem 2.3, [12, Theorem 1] and [13, Theorem 2.2]. \square

Let (R, \mathfrak{m}, K) be either a local ring or a standard graded K -algebra, with (graded) maximal ideal \mathfrak{m} , which is Cohen-Macaulay and admits a canonical module ω_R . The *canonical trace* of R is defined as the ideal

$$\text{tr}(\omega_R) = \sum_{\varphi \in \text{Hom}_R(\omega_R, R)} \varphi(\omega_R).$$

Following [14], we say that R is *nearly Gorenstein* if $\mathfrak{m} \subseteq \text{tr}(\omega_R)$. It is clear from the definition that any Gorenstein ring is nearly Gorenstein.

We say that a graph G is *nearly Gorenstein* if $S/I(G)$ is a nearly Gorenstein ring.

Next we characterize Gorenstein and nearly Gorenstein permutation graphs. To this aim we use the properties of the poset P associated to a Cohen-Macaulay permutation graph G , employed in the proof of Theorem 2.3.

Theorem 2.7. *Let G be a permutation graph without isolated vertices. Then*

- (a) *G is Gorenstein if and only if G is the disjoint union of edges.*
- (b) *G is nearly Gorenstein but not Gorenstein if and only if G is either K_n or P_n^c for some $n \geq 3$.*

Proof. (a) If G is the disjoint union of edges, then $I(G)$ is a complete intersection, and so G is Gorenstein. Conversely, suppose that G is Gorenstein. Since G is Cohen-Macaulay, we will adopt the notation and results shown in the proof of Theorem 2.3. The set $L = \{j_1, \dots, j_r\}$ is a maximal independent set of G . Since G is unmixed, this means that $\alpha(G) = r$. If $\alpha(G) = 1$, then P is a chain, which means that G is a complete graph. On the other hand, the only complete graph which is Gorenstein is K_2 . Hence G is just an edge. Now, let $\alpha(G) \geq 2$. Set $e_\ell = \{i_\ell, j_\ell\}$ for $1 \leq \ell \leq r$. We show that G is the disjoint union of the edges e_1, \dots, e_r .

For any $F \subseteq [n]$, we set $G_F = G \setminus N_G[F]$. Then by [24, Theorem 2.3], for any independent set F of G with $|F| = r - 2$, we have $G_F = C_m^c$, where C_m denotes the cycle graph on $m \geq 4$ vertices. Since any permutation graph is weakly chordal, we obtain $m = 4$. Consider a subset $F \subset L$ with $|F| = r - 2$. Without loss of generality assume that $F = L \setminus \{j_1, j_2\}$. Since $G_F = C_4^c$, we get $E(G_F) = \{e_1, e_2\}$.

This means that e_1 and e_2 form a gap in G . By choosing the set F as $L \setminus \{j_s, j_t\}$ for any $s \neq t$, with the same argument we conclude that any two edges e_s, e_t form a gap. This means that e_1, \dots, e_r form a gap in G . If $|V(G)| = 2r$, it follows that $E(G) = \{e_1, \dots, e_r\}$, as desired.

Now, by contradiction assume that $|V(G)| > 2r$. Since G has no isolated vertices, we have $|A_\ell| \geq 2$ for $1 \leq \ell \leq r$. Then by the assumption that $|V(G)| > 2r$, we may assume that $|A_\ell| > 2$ for $1 \leq \ell \leq s$ and $|A_\ell| = 2$ for $\ell > s$, where s is an integer with $1 \leq s \leq r$. For any maximal chain A_ℓ with $|A_\ell| > 2$, let $t_\ell \in A_\ell$ be the element with $t_\ell \prec i_\ell$. Since e_1, \dots, e_r form a gap, the set $L' = \{i_1, \dots, i_r\}$ is an independent set of G . For any $1 \leq \ell \leq s$, choose $F_\ell \subset L'$ such that $|F_\ell| = r - 2$ and $i_\ell \notin F_\ell$. Then $G_{F_\ell} = C_4^c$. Moreover, $e_\ell = \{i_\ell, j_\ell\}$ is an edge of G_{F_ℓ} . This together with $t_\ell \in A_\ell$ implies that $t_\ell \in N_G(F_\ell)$. In other words, for any $1 \leq \ell \leq s$ there exists $h_\ell \neq \ell$ such that $\{t_\ell, i_{h_\ell}\} \in E(G)$. This in fact means that for any $1 \leq \ell \leq s$, $t_\ell \prec i_{h_\ell}$. Hence, $\{i_{h_\ell}, i_\ell\} \subseteq \{k : t_\ell \prec k\}$. We show that this is not possible.

We set $G_1 = G \setminus \{j_1\}$ and $G_p = G_{p-1} \setminus \{j_p\}$ for $2 \leq p \leq r$. Since e_1, \dots, e_r form a gap in G , for any integer p we have $\{k \in [n] : i_p \prec k\} = \{j_p\}$. Then as is shown in the proof of Theorem 2.3, each j_p is a shedding vertex of G_p and each G_p is a Cohen-Macaulay permutation graph. In particular $G_r = G \setminus \{j_1, \dots, j_r\}$ is Cohen-Macaulay. Let $P' = P \setminus \{j_1, \dots, j_r\}$. Then clearly $G_r = \text{comp}(P')$ and P' is the disjoint union of the maximal chains $A'_\ell = A_\ell \setminus \{j_\ell\}$. Moreover, i_ℓ is the maximal element of A'_ℓ for all ℓ . As the proof of Theorem 2.3(Claim) shows, there should exist an integer $1 \leq \ell \leq s$ such that $\{k \in V(P') : t_\ell \prec k\} = \{i_\ell\}$. This contradicts to $\{i_{h_\ell}, i_\ell\} \subseteq \{k : t_\ell \prec k\}$ for all $1 \leq \ell \leq s$. So we have $|V(G)| = 2r$, and this concludes the proof of (a).

(b) By [20, Theorem A(Y)], G is nearly Gorenstein but not Gorenstein if and only if Δ_G is isomorphic either to the disjoint union of n vertices or to a path on n vertices. This implies that either $G = K_n$ or $G = P_n^c$. Since both K_n and P_n^c are Cohen-Macaulay permutation graphs, the result follows. \square

In the next proposition we give a combinatorial description for the a -invariant $a(S/I(G))$ of the ring $S/I(G)$, when G is a Cohen-Macaulay permutation graph. A graded ideal $I \subset S$ is called *Hilbertian* if $P_{S/I}(t) = H(S/I, t)$ for all $t \geq 0$, where $P_{S/I}(t)$ and $H(S/I, t)$ denote the Hilbert polynomial and the Hilbert function of S/I , respectively.

Proposition 2.8. *Let G be a Cohen-Macaulay permutation graph on n vertices. Then*

- (a) $\text{reg}(S/I(G)) = \text{im}(G)$.
- (b) $a(S/I(G)) = \text{im}(G) + \tau(G) - n$.
- (c) $I(G)$ is Hilbertian if and only if $\tau(G) + \text{im}(G) < n$.

Proof. (a) follows from the fact that any permutation graph is weakly chordal, together with [31, Theorem 14], which shows that if G is a weakly chordal graph, then $\text{reg}(S/I(G)) = \text{im}(G)$.

(b) Since G is Cohen-Macaulay, we know that the degree of the h -polynomial $h(t)$ in the Hilbert series of $S/I(G)$ is equal to $\text{reg}(S/I(G))$, see [28, Corollary B.28].

Hence, $a(S/I(G)) = \text{reg}(S/I(G)) - d$, where $d = \dim(S/I(G))$. Using (a) and the equality $d = n - \tau(G)$, we obtain $a(S/I(G)) = \text{im}(G) + \tau(G) - n$.

(c) By [23, Lemma 5.3], the ideal $I(G)$ is Hilbertian if and only if $a(S/I(G)) < 0$. By (ii) we have $a(S/I(G)) < 0$ if and only if $\tau(G) + \text{im}(G) < n$. \square

A graph G is called *bi-Cohen-Macaulay* if $S/I(G)$ and $S/J(G)$ are Cohen-Macaulay rings. Combining Theorem 2.3 and Proposition 2.8(a) we obtain

Proposition 2.9. *Let G be a permutation graph on n vertices. Then G is bi-Cohen-Macaulay if and only if G is an unmixed vertex decomposable graph and $\text{im}(G) = 1$.*

Proof. Having Theorem 2.3, it is enough to show that $S/J(G)$ is Cohen-Macaulay if and only if $\text{im}(G) = 1$. By [27, Theorem 2.1] and Proposition 2.8(a) we have $\text{proj dim}(S/J(G)) = \text{reg}(I(G)) = \text{im}(G) + 1$. So $\text{depth}(S/J(G)) = n - \text{im}(G) - 1$. Moreover, since $J(G)$ is unmixed of height two, we have $\dim(S/J(G)) = n - 2$. So $S/J(G)$ is Cohen-Macaulay if and only if $\text{im}(G) = 1$. \square

Acknowledgment. A. Ficarra was partly supported by INDAM (Istituto Nazionale di Alta Matematica), and also by the Grant JDC2023-051705-I funded by MI-CIU/AEI/10.13039/501100011033 and by the FSE+. S. Moradi is supported by the Alexander von Humboldt Foundation.

REFERENCES

- [1] T. Biyikouğlu, Y. Civan, Vertex-decomposable graphs, codismantlability Cohen-Macaulayness, and Castelnuovo-Mumford regularity, *Electron. J. Combin.* **21** (2014), no. 1, Paper 1.1, 17 pp.
- [2] A. Björner, M. Wachs, Shellable nonpure complexes and posets. II, *Trans. Amer. Math. Soc.* **349** (1997), no. 10, 3945–3975.
- [3] P. V. Cheri, D. Dey, A. K., N. Kotal, D. Veer, Cohen-Macaulay permutation graphs, *Math. Scand.* **130** (2024), no. 3, 419–431.
- [4] D. Cook and U. Nagel, Cohen-Macaulay graphs and face vectors of flag complexes, *SIAM J. Discrete Math.* **26** (2012), no. 1, 89–101.
- [5] M. Crupi, G. Rinaldo, N. Terai, Cohen–Macaulay edge ideals whose height is half of the number of vertices, *Nagoya Math. J.* **201** (2011), 116–130.
- [6] A. Dochtermann, A. Engström, Algebraic properties of edge ideals via combinatorial topology, *Electron. J. Combin.* **16** (2009), no. 2.
- [7] S. Even, A. Pnueli, A. Lempel, Permutation graphs and transitive graphs, *Journal of the Association for Computing Machinery*, **19** (1972), 400–410 .
- [8] T. Gallai, Transitiv orientierbare Graphen, *Acta Math. Acad. Sci. Hungar.*, **18** (1967), 25–66.
- [9] S. V. Gervacio, T.A. Rapanut, P.F. Ramos, Characterization and construction of permutation graphs, *Open Journal of Discrete Mathematics* **3** (2013) 33–38.
- [10] J. Herzog, T. Hibi, Distributive lattices, bipartite graphs and Alexander duality, *J. Algebraic Combin.* **22** (2005) 289–302.
- [11] J. Herzog, T. Hibi, *Monomial ideals*, Graduate texts in Mathematics **260**, Springer, 2011.
- [12] J. Herzog, T. Hibi, The Regularity of Edge Rings and Matching Numbers, *Mathematics*, (2020), **8**(1), 39.
- [13] J. Herzog, T. Hibi, Matching numbers and the regularity of the Rees algebra of an edge ideal, *Ann. Comb.* **24**, (2020), 577–586.
- [14] J. Herzog, T. Hibi, D. I. Stamate, The trace of the canonical module, *Israel Journal of Mathematics* **233** (2019), 133–165.

- [15] J. Herzog, T. Hibi, X. Zheng, Cohen-Macaulay chordal graphs, *J. Combin. Theory Ser. A* **113** (2006), no. 5, 911–916.
- [16] J. Herzog, A.A. Qureshi, Persistence and stability properties of powers of ideals, *J. Pure Appl. Algebra*, **219** (2015), 530–542.
- [17] T. Hibi, A. Higashitani, K. Kimura, A. O’Keefe, Algebraic study on Cameron-Walker graphs, *J. Algebra* **422** (2015), 257–269.
- [18] V. Limouzy, Seidel Minor, Permutation Graphs and Combinatorial Properties, In: *Lecture Notes in Computer Science Volume 6506*, Springer, Berlin, 2010, pp. 194-205.
- [19] M. Mahmoudi, A. Mousivand, M. Crupi, G. Rinaldo, N. Terai, S. Yassemi, Vertex decomposability and regularity of very well-covered graphs, *J. Pure Appl. Algebra* **215** (2011), 2473-2480.
- [20] S. Miyashita, M. Varbaro, The canonical trace of Stanley-Reisner rings that are Gorenstein on the punctured spectrum, arXiv preprint arXiv:2412.12860 (2024).
- [21] S. Moradi, Normal Rees algebras arising from vertex decomposable simplicial complexes, arXiv preprint arXiv:2311.15135 (2023).
- [22] S. Moradi, F. Khosh-Ahang, On vertex decomposable simplicial complexes and their Alexander duals. *Math. Scand.* **118** (2016), no. 1, 43–56.
- [23] T. Nguyen, J. Rajchgot, A. Van Tuyl, Three invariants of geometrically vertex decomposable ideals, *Pacific Journal of Mathematics*, Vol. **333**(2024), No. 2, 357–390.
- [24] M. R. Oboudi, A. Nikseresht, Some combinatorial characterizations of Gorenstein graphs with independence number less than four, *Iran. J. Sci. Technol. Trans. A Sci.* **44** (2020), no. 6, 1667–1671.
- [25] A. Pnueli, A. Lempel, S. Even, Transitive orientation of graphs and identification of permutation graphs, *Canadian J. Math.*, **23** (1971), 160–175.
- [26] J. S. Provan, L. J. Billera, Decompositions of simplicial complexes related to diameters of convex polyhedra, *Math. Oper. Res.* **5** (1980), no. 4, 576–594. MR 593648 (82c:52010)
- [27] N. Terai, Alexander duality theorem and Stanley-Reisner rings. Free resolutions of coordinate rings of projective varieties and related topics, (Japanese) (Kyoto, 1998). *Sūrikaiseikikenkyūsho Kōkyūroku* no. 1078 (1999), 174–184.
- [28] W. V. Vasconcelos, Computational methods in commutative algebra and algebraic geometry, volume 2 of *Algorithms and Computation in Mathematics*. Springer-Verlag, Berlin, 1998. With chapters by David Eisenbud, Daniel R. Grayson, Jürgen Herzog and Michael Stillman.
- [29] R. H. Villarreal, Cohen-Macaulay graphs, *Manuscripta Math.* **66** (1990), no. 3, 277–293.
- [30] R. Woodroffe, Vertex decomposable graphs and obstructions to shellability, *Proc. Amer. Math. Soc.* **137** (2009), no. 10, 3235–3246.
- [31] R. Woodroffe, Matchings, coverings, and Castelnuovo-Mumford regularity, *J. Commut. Algebra* **6** (2014), no. 2, 287–304.

ANTONINO FICARRA, DEPARTAMENTO DE MATEMÁTICA, ESCOLA DE CIÊNCIAS E TECNOLOGIA, CENTRO DE INVESTIGAÇÃO, MATEMÁTICA E APLICAÇÕES, INSTITUTO DE INVESTIGAÇÃO E FORMAÇÃO AVANÇADA, UNIVERSIDADE DE ÉVORA, RUA ROMÃO RAMALHO, 59, P-7000-671 ÉVORA, PORTUGAL

Email address: antonino.ficarra@uevora.pt

Email address: antficarra@unime.it

SOMAYEH MORADI, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, ILAM UNIVERSITY, P.O.BOX 69315-516, ILAM, IRAN

Email address: so.moradi@ilam.ac.ir