

New q -identities Via q -Derivative of Basic Hypergeometric Series with Respect to Parameters

Ronald Orozco López

February 28, 2025

Abstract

In this paper, we use the effect of the q -differential and deformed q -exponential operators on basic hypergeometric series to find new q -identities from the q -Gauss sum, the q -Chu-Vandermonde's sum, and Jackson's transformation formula.

2020 *Mathematics Subject Classification*: Primary 05A30. Secondary 33D15.

Keywords: Deformed basic hypergeometric series, q -Gauss sum, q -Chu-Vandermonde's sum, deformed q -exponential operator, q -differential operator.

1 Introduction and Motivation

Throughout this paper, we take $0 < |q| < 1$. The q -differential operator D_q is defined by:

$$D_q f(x) = \frac{f(x) - f(qx)}{x}$$

and the Leibniz rule for D_q

$$D_q^n \{f(x)g(x)\} = \sum_{k=0}^n q^{k(k-n)} \begin{bmatrix} n \\ k \end{bmatrix}_q D_q^k \{f(x)\} D_q^{n-k} \{g(q^k x)\}, \quad (1)$$

where the q -binomial coefficient is defined by

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}},$$

and the q -shifted factorial of a is given by $(a; q)_0 = 1$ and

$$(a; q)_n = \prod_{k=0}^{n-1} (1 - q^k a), \quad \text{for } n = 1, 2, \dots,$$

$$(a; q)_\infty = \lim_{n \rightarrow \infty} (a; q)_n = \prod_{k=0}^{\infty} (1 - aq^k).$$

Following Gasper and Rahman [4], the ${}_r\phi_s$ basic hypergeometric series is defined by

$${}_r\phi_s \left(\begin{matrix} a_1, a_2, \dots, a_r \\ b_1, \dots, b_s \end{matrix} ; q, z \right) = \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_r; q)_n}{(q; q)_n (b_1, b_2, \dots, b_s; q)_n} \left[(-1)^n q^{\binom{n}{2}} \right]^{1+s-r} z^n$$

where

$$(a_1, a_2, \dots, a_m; q)_n = (a_1; q)_n (a_2; q)_n \cdots (a_m; q)_n.$$

Equivalently,

$$(a_1, a_2, \dots, a_m; q)_{\infty} = (a_1; q)_{\infty} (a_2; q)_{\infty} \cdots (a_m; q)_{\infty}.$$

In [1], we define the deformed basic hypergeometric series ${}_r\Phi_s$ as

$${}_r\Phi_s \left(\begin{matrix} a_1, a_2, \dots, a_r \\ b_1, \dots, b_s \end{matrix} ; q, u, z \right) = \sum_{n=0}^{\infty} u^{\binom{n}{2}} \frac{(a_1, a_2, \dots, a_r; q)_n}{(q, b_1, b_2, \dots, b_s; q)_n} \left[(-1)^n q^{\binom{n}{2}} \right]^{1+s-r} z^n. \quad (2)$$

If $u = 1$, then ${}_r\Phi_s = {}_r\phi_s$. If $u = q$,

$${}_{r+1}\Phi_r \left(\begin{matrix} a_1, a_2, \dots, a_r, 0 \\ b_1, \dots, b_r \end{matrix} ; q, q, z \right) = {}_r\phi_r \left(\begin{matrix} a_1, a_2, \dots, a_r \\ b_1, b_2, \dots, b_r \end{matrix} ; q, -z \right).$$

In this paper, we will frequently use the following identities:

$$(a; q)_n = \frac{(a; q)_{\infty}}{(aq^n; q)_{\infty}}, \quad (3)$$

$$(a; q)_{n+k} = (a; q)_n (aq^n; q)_k, \quad (4)$$

$$(aq^k; q)_{n-k} = \frac{(a; q)_k}{(a; q)_i}, \quad (5)$$

$$(aq^n; q)_k = \frac{(a; q)_k (aq^k; q)_n}{(a; q)_n}. \quad (6)$$

In addition, we will use the identities for binomial coefficients:

$$\binom{n+k}{2} = \binom{n}{2} + \binom{k}{2} + nk,$$

$$\binom{n-k}{2} = \binom{n}{2} + \binom{k}{2} + k(1-n).$$

This paper provides new q -identities by applying the q -differential and deformed q -exponential operators to the q -Gauss sum, the q -Chu-Vandermonde's sum, and Jackson's transformation formula. These q -identities have the following quotient forms of the q -shifted factorials

$$\frac{1}{(c; q)_n}, \quad \frac{x^n}{(c; q)_n}, \quad \frac{(ac; q)_n}{(bc; q)_n}, \quad \text{and} \quad \frac{(ac, bc; q)_n}{(dc, ec; q)_n}.$$

The method will consist of applying the operators D_q and $T(yD_q|u)$, with respect to the parameter c , to both sides of the above q -identities and then using the identities Eqs.(3)-(6). Something similar was done in [7].

2 Some identities of the q -differential operator

For all $n \in \mathbb{N}$, we have the identities:

$$D_q^k x^n = \begin{cases} \frac{(q; q)_n}{(q; q)_{n-k}} x^{n-k}, & \text{if } k \leq n; \\ 0, & \text{if } k > n. \end{cases} \quad (7)$$

$$D_q^k (ax; q)_n = \begin{cases} (-a)^k q^{\binom{k}{2}} \frac{(q; q)_n (ax; q)_n}{(q; q)_{n-k} (ax; q)_k}, & \text{if } k \leq n; \\ 0, & \text{if } k > n. \end{cases} \quad (8)$$

Theorem 1. For all $n \geq 1$,

$$D_q^n \left(\frac{1}{1-ax} \right) = \frac{(q; q)_n a^n}{(ax; q)_{n+1}}. \quad (9)$$

Proof. Note that

$$D_q \left(\frac{1}{1-ax} \right) = \frac{(1-q)a}{(1-ax)(1-axq)} = \frac{(q; q)_1 a}{(ax; q)_2}. \quad (10)$$

Suppose that Eq.(9) is true for n and let us prove by induction for $n+1$. As

$$D_q (ax; q)_{n+1} = -(1-q^{n+1})a(ax; q)_n, \quad (11)$$

then

$$\begin{aligned} D_q^{n+1} \left(\frac{1}{1-ax} \right) &= D_q D_q^n \left(\frac{1}{1-ax} \right) \\ &= D_q \left(\frac{(q; q)_n a^n}{(ax; q)_{n+1}} \right) \\ &= \frac{-(q; q)_n a^n D_q (ax; q)_{n+1}}{(ax; q)_{n+1} (ax; q)_{n+1}} \\ &= \frac{(q; q)_{n+1} (ax; q)_n a^{n+1}}{(ax; q)_{n+1} (ax; q)_{n+1}} \\ &= \frac{(q; q)_{n+1} a^{n+1}}{(1-q^{n+1}xa)(ax; q)_{n+1}} \\ &= \frac{(q; q)_{n+1} a^{n+1}}{(ax; q)_{n+2}}. \end{aligned}$$

The proof is reached. □

Theorem 2.

$$D_q^m \left\{ \frac{1}{(ax; q)_n} \right\} = \frac{(q^n; q)_m a^m}{(ax; q)_{m+n}}.$$

Proof.

$$\begin{aligned} D_q^m \left\{ \frac{1}{(ax; q)_n} \right\} &= \frac{1}{a^{n-1}(q; q)_{n-1}} D_q^{m+n-1} \left\{ \frac{1}{1-ax} \right\} \\ &= \frac{1}{a^{n-1}(q; q)_{n-1}} \frac{(q; q)_{m+n-1} a^{m+n-1}}{(ax; q)_{m+n}} \\ &= \frac{(q^n; q)_m a^m}{(ax; q)_{m+n}}. \end{aligned}$$

□

Theorem 3.

$$D_q^k \left\{ \frac{x^n}{(ax; q)_n} \right\} = \frac{(q; q)_n}{(ax; q)_n} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{(q^n; q)_{k-i} (ax; q)_i a^{k-i}}{(q; q)_{n-i} (aq^n x; q)_k} x^{n-i}.$$

Proof.

$$\begin{aligned} D_q^k \left\{ \frac{x^n}{(ax; q)_n} \right\} &= \sum_{i=0}^k q^{i(i-k)} \begin{bmatrix} k \\ i \end{bmatrix}_q D_q^i \{x^n\} D_q^{k-i} \left\{ \frac{1}{(aq^i x; q)_n} \right\} \\ &= \sum_{i=0}^k q^{i(i-k)} \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{(q; q)_n}{(q; q)_{n-i}} x^{n-i} \frac{(q^n; q)_{k-i} (aq^i)^{k-i}}{(aq^i x; q)_{n+k-i}} \\ &= \frac{(q; q)_n}{(ax; q)_n} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{(q^n; q)_{k-i} (ax; q)_i a^{k-i}}{(q; q)_{n-i} (aq^n x; q)_k} x^{n-i}. \end{aligned}$$

□

Theorem 4. For $k \leq n$,

$$D_q^k \left\{ \frac{(ax; q)_n}{(bx; q)_n} \right\} = \frac{(ax, q; q)_n}{(bx; q)_n (bq^n x; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{\binom{i}{2}} (-a)^i b^{k-i} \frac{(bx; q)_i (q^n; q)_{k-i}}{(q; q)_{n-i} (ax; q)_i}.$$

Proof. From Eq.(8),

$$\begin{aligned} D_q^k \left\{ \frac{(ax; q)_n}{(bx; q)_n} \right\} &= \sum_{i=0}^k q^{i(i-k)} \begin{bmatrix} k \\ i \end{bmatrix}_q D_q^i \{(ax; q)_n\} D_q^{k-i} \left\{ \frac{1}{(bq^i x; q)_n} \right\} \\ &= \sum_{i=0}^k q^{i(i-k)} \begin{bmatrix} k \\ i \end{bmatrix}_q (-a)^i q^{\binom{i}{2}} \frac{(q; q)_n}{(q; q)_{n-i}} \frac{(ax; q)_n}{(ax; q)_i} \frac{(bx; q)_i (q^n; q)_{k-i} (bq^i)^{k-i}}{(bx; q)_{n+k}} \\ &= \frac{(ax, q; q)_n}{(bx; q)_n (bq^n x; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{\binom{i}{2}} (-a)^i b^{k-i} \frac{(bx; q)_i (q^n; q)_{k-i}}{(q; q)_{n-i} (ax; q)_i}. \end{aligned}$$

□

If $a = b$, then for all $n \geq 0$

$$\sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{\binom{i}{2}} (-1)^i \frac{(q^n; q)_{k-i}}{(q; q)_{n-i}} = 0.$$

Theorem 5.

$$\begin{aligned} D_q^k \left\{ \frac{(ax, bx; q)_n}{(cx, dx; q)_n} \right\} &= (q; q)_n^2 (ax, bx; q)_n \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{q^{\binom{i}{2}}}{(bx; q)_i} \sum_{j=0}^i \begin{bmatrix} i \\ j \end{bmatrix}_q q^{j(j-i)} \frac{(-a)^j}{(q; q)_{n-j}} \frac{(-b)^{i-j}}{(q; q)_{n-i+j}} \frac{(bq^n x; q)_j}{(ax; q)_j} \\ &\times \frac{1}{(cq^i x, dq^i x; q)_n (dq^{n+i} x; q)_{k-i}} \sum_{l=0}^{k-i} \begin{bmatrix} k-i \\ l \end{bmatrix}_q \frac{(q^n; q)_l (q^n; q)_{k-i-l} (dq^i x; q)_l}{(cq^{n+i} x; q)_l} c^l d^{k-i-l}. \end{aligned}$$

Proof.

$$\begin{aligned}
& D_q^k \left\{ \frac{(ax, bx; q)_n}{(cx, dx; q)_n} \right\} \\
&= \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{i(i-k)} D_q^i \{(ax, bx; q)_n\} D_q^{k-i} \left\{ \frac{1}{(cq^i x, dq^i x; q)_n} \right\} \\
&= (q; q)_n^2 (ax, bx; q)_n \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{q^{\binom{i}{2}}}{(bx; q)_i} \sum_{j=0}^i \begin{bmatrix} i \\ j \end{bmatrix}_q q^{j(j-i)} \frac{(-a)^j}{(q; q)_{n-j}} \frac{(-b)^{i-j}}{(q; q)_{n-i+j}} \frac{(bq^n x; q)_j}{(ax; q)_j} \\
&\times \frac{1}{(cq^i x, dq^i x; q)_n (dq^{n+i} x; q)_{k-i}} \sum_{l=0}^{k-i} \begin{bmatrix} k-i \\ l \end{bmatrix}_q \frac{(q^n; q)_l (q^n; q)_{k-i-l} (dq^i x; q)_l}{(cq^{n+i} x; q)_l} c^l d^{k-i-l}.
\end{aligned}$$

□

Por hacer $n \rightarrow \infty$

$$\begin{aligned}
& D_q^k \left\{ \frac{(ax, bx; q)_\infty}{(cx, dx; q)_\infty} \right\} \\
&= \frac{(ax, bx; q)_\infty}{(cx, dx; q)_\infty} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{q^{\binom{i}{2}} (cx, dx; q)_i}{(bx; q)_i} \sum_{j=0}^i \begin{bmatrix} i \\ j \end{bmatrix}_q q^{j(j-i)} \frac{(-a)^j (-b)^{i-j}}{(ax; q)_j} \\
&\quad \times \sum_{l=0}^{k-i} \begin{bmatrix} k-i \\ l \end{bmatrix}_q (dq^i x; q)_l c^l d^{k-i-l}.
\end{aligned}$$

3 q -derivative of ${}_r\Phi_s$ with respect to parameters

Theorem 6. For all $n \in \mathbb{N}$,

$$\begin{aligned}
& D_{c_1, q}^k \left\{ {}_r\Phi_s \left(\begin{matrix} a_1, \dots, a_r \\ c_1, c_2, \dots, c_s \end{matrix}; q, u, z \right) \right\} \\
&= \frac{(-1)^{1+s-r} \prod_{i=1}^r (1-a_i) (q; q)_k z}{(c_1; q)_{k+1} (1-q) \prod_{i=2}^s (1-c_i)} \\
&\quad {}_{r+1}\Phi_{s+1} \left(\begin{matrix} a_1 q, \dots, a_r q, q^{k+1} \\ c_1 q^{k+1}, c_2, \dots, c_s, q^2 \end{matrix}; q, u, q^{1+s-r} u z \right).
\end{aligned}$$

Proof. From Eq.(2) and Theorem 2,

$$\begin{aligned}
& D_{c_1, q}^k \left\{ {}_r\Phi_s \left(\begin{matrix} a_1, \dots, a_r \\ c_1, c_2, \dots, c_s \end{matrix}; q, u, z \right) \right\} \\
&= \sum_{n=0}^{\infty} u^{\binom{n}{2}} \frac{(a_1; q)_n \cdots (a_r; q)_n}{(c_2; q)_n \cdots (c_s; q)_n (q; q)_n} \left((-1)^n q^{\binom{n}{2}} \right)^{1+s-r} z^n D_{c_1, q}^k \left\{ \frac{1}{(c_1; q)_n} \right\} \\
&= c_1^k \sum_{n=0}^{\infty} u^{\binom{n}{2}} \frac{(a_1; q)_n \cdots (a_r; q)_n (q^n; q)_k}{(c_1; q)_{k+n} (c_2; q)_n \cdots (c_s; q)_n (q; q)_n} \left((-1)^n q^{\binom{n}{2}} \right)^{1+s-r} z^n.
\end{aligned}$$

Taking into account that $(q^0; q)_k = 0$, we shift the index n by one unit and apply the identity Eq.(4),

$$\begin{aligned}
& D_{c_1, q}^k \left\{ {}_r\Phi_s \left(\begin{matrix} a_1, \dots, a_r \\ c_1, c_2, \dots, c_s \end{matrix} ; q, u, z \right) \right\} \\
&= \frac{z}{(c_1; q)_k} \sum_{n=0}^{\infty} u^{\binom{n+1}{2}} \frac{(a_1; q)_{n+1} \cdots (a_r; q)_{n+1} (q^{n+1}; q)_k}{(c_1 q^k; q)_{n+1} (c_2; q)_{n+1} \cdots (c_s; q)_{n+1} (q; q)_{n+1}} \\
&\quad \left((-1)^{n+1} q^{\binom{n+1}{2}} \right)^{1+s-r} z^n \\
&= \frac{(-1)^{1+s-r} \prod_{i=1}^r (1-a_i) z}{(cx; q)_{k+1} (1-q) \prod_{i=2}^s (1-c_i)} \sum_{n=0}^{\infty} \frac{u^{\binom{n}{2}} (a_1 q; q)_n \cdots (a_r q; q)_n (q^{n+1}; q)_k}{(c_1 q^{k+1}; q)_n (c_2 q; q)_n \cdots (c_s q; q)_n (q^2; q)_n} \\
&\quad \times \left((-1)^n q^{\binom{n}{2}} \right)^{1+s-r} (q^{1+s-r} uz)^n \\
&= \frac{(-1)^{1+s-r} \prod_{i=1}^r (1-a_i) (q; q)_k z}{(cx; q)_{k+1} (1-q) \prod_{i=2}^s (1-c_i)} \\
&\quad \times \sum_{n=0}^{\infty} \frac{u^{\binom{n}{2}} (a_1 q; q)_n \cdots (a_r q; q)_n (q^{k+1}; q)_n}{(c_1 q^{k+1}; q)_n (c_2; q)_n \cdots (c_s; q)_n (q^2; q)_n (q; q)_n} \left((-1)^n q^{\binom{n}{2}} \right)^{1+s-r} (q^{1+s-r} uz)^n \\
&= \frac{(-1)^{1+s-r} \prod_{i=1}^r (1-a_i) (q; q)_k z}{(c_1; q)_{k+1} (1-q) \prod_{i=2}^s (1-c_i)} \\
&\quad {}_{r+1}\Phi_{s+1} \left(\begin{matrix} a_1 q, \dots, a_r q, q^{k+1} \\ c_1 q^{k+1}, c_2, \dots, c_s, q^2 \end{matrix} ; q, u, q^{1+s-r} uz \right).
\end{aligned}$$

□

Theorem 7. For all $n \in \mathbb{N}$,

$$\begin{aligned}
& D_{c, q}^k \left\{ {}_r\Phi_1 \left(\begin{matrix} a_1, \dots, a_r \\ c \end{matrix} ; q, u, z \right) \right\} \\
&= \frac{(-1)^{2-r} \prod_{i=1}^r (1-a_i) (q; q)_k z}{(c; q)_{k+1} (1-q)} \\
&\quad {}_{r+1}\Phi_2 \left(\begin{matrix} a_1 q, \dots, a_r q, q^{k+1} \\ c q^{k+1}, q^2 \end{matrix} ; q, u, q^{1+s-r} uz \right).
\end{aligned}$$

4 New identities of the deformed q -exponential operator

Set $u \in \mathbb{C}$. In [1] the u -deformed q -exponential operator $\mathbb{T}(yD_q|u)$ is defined by letting

$$\mathbb{T}(yD_q|u) = \sum_{n=0}^{\infty} u^{\binom{n}{2}} \frac{(yD_q)^n}{(q; q)_n}.$$

The u -deformed q -exponential operator $\mathbb{T}(yD_q|u)$ generalizes the operators given by Chen and Liu [2, 3]:

$$\mathbb{T}(bD_q) = \sum_{n=0}^{\infty} \frac{(bD_q)^n}{(q; q)_n} = \mathbb{T}(bD_q|1),$$

and by Saad et al. [5, 6]:

$$R(bD_q) = \sum_{n=0}^{\infty} (-1)^n \frac{(bD_q)^n q^{\binom{n}{2}}}{(q; q)_n} = T(-bD_q|q).$$

Some q -operator identities:

Theorem 8. For all $n \geq 0$

$$T(yD_q|u) \left\{ \frac{1}{(ax; q)_n} \right\} = \frac{1}{(ax; q)_n} {}_1\Phi_1 \left(\begin{matrix} q^n \\ aq^n x \end{matrix}; q, uq^{-1}, -ay \right). \quad (12)$$

Proof. We have

$$\begin{aligned} T(yD_q|u) \left\{ \frac{1}{(ax; q)_n} \right\} &= \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{y^k}{(q; q)_k} D_q^k \left\{ \frac{1}{(ax; q)_n} \right\} \\ &= \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{y^k}{(q; q)_k} \frac{(q^n; q)_k a^k}{(ax; q)_{k+n}} \\ &= \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{(ay)^k}{(q; q)_k} \frac{(q^n; q)_k}{(ax; q)_n (aq^n x; q)_k} \\ &= \frac{1}{(ax; q)_n} {}_1\Phi_1 \left(\begin{matrix} q^n \\ aq^n x \end{matrix}; q, uq^{-1}, -ay \right) \end{aligned}$$

as claimed. \square

Theorem 9. For all $n \geq 0$

$$\begin{aligned} T(yD_q|u) \left\{ \frac{(ax; q)_n}{(bx; q)_n} \right\} \\ = \frac{(ax; q)_n}{(bx; q)_n} \sum_{i=0}^n (uq)^{\binom{i}{2}} \begin{bmatrix} n \\ i \end{bmatrix}_q \frac{(bx; q)_i (-ay)^i}{(ax; q)_i (bq^n x; q)_i} {}_1\Phi_1 \left(\begin{matrix} q^n \\ bq^{n+i} x \end{matrix}; q, uq^{-1}, -u^i by \right). \end{aligned} \quad (13)$$

Proof.

$$\begin{aligned} T(yD_q|u) \left\{ \frac{(ax; q)_n}{(bx; q)_n} \right\} \\ = \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{y^k}{(q; q)_k} D_q^k \left\{ \frac{(ax; q)_n}{(bx; q)_n} \right\} \\ = \frac{(ax; q; q)_n}{(bx; q)_n} \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{y^k}{(q; q)_k} \frac{1}{(bq^n x; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{\binom{i}{2}} (-a)^i b^{k-i} \frac{(bx; q)_i (q^n; q)_{k-i}}{(q; q)_{n-i} (ax; q)_i} \\ = \frac{(ax; q)_n}{(bx; q)_n} \sum_{i=0}^n (uq)^{\binom{i}{2}} \begin{bmatrix} n \\ i \end{bmatrix}_q \frac{(bx; q)_i (-ay)^i}{(ax; q)_i (bq^n x; q)_i} \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{(q^n; q)_k}{(q; q)_k (bq^{n+i} x; q)_k} (u^i y b)^k \\ = \frac{(ax; q)_n}{(bx; q)_n} \sum_{i=0}^n (uq)^{\binom{i}{2}} \begin{bmatrix} n \\ i \end{bmatrix}_q \frac{(bx; q)_i (-ay)^i}{(ax; q)_i (bq^n x; q)_i} {}_1\Phi_1 \left(\begin{matrix} q^n \\ bq^{n+i} x \end{matrix}; q, uq^{-1}, -u^i by \right). \end{aligned}$$

\square

If $a = b$, then for all $n \geq 0$

$$\sum_{i=0}^n (uq)^{\binom{i}{2}} \begin{bmatrix} n \\ i \end{bmatrix}_q \frac{(-ay)^i}{(aq^n x; q)_i} {}_1\Phi_1 \left(\begin{matrix} q^n \\ aq^{n+i} x \end{matrix}; q, uq^{-1}, -u^i ay \right) = 1$$

Theorem 10.

$$\begin{aligned} & \mathbb{T}(yD_q|u) \left\{ \frac{x^n}{(ax; q)_n} \right\} \\ &= \frac{1}{(ax; q)_n} \sum_{i=0}^n \begin{bmatrix} n \\ i \end{bmatrix}_q u^{\binom{i}{2}} \frac{(ax; q)_i}{(aq^n x; q)_i} x^{n-i} y^i {}_1\Phi_1 \left(\begin{matrix} q^n \\ aq^{n+i} x \end{matrix}; q, uq^{-1}, -u^i ay \right). \end{aligned}$$

Proof.

$$\begin{aligned} & \mathbb{T}(yD_q|u) \left\{ \frac{x^n}{(ax; q)_n} \right\} \\ &= \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{y^k}{(q; q)_k} D_q^k \left\{ \frac{x^n}{(ax; q)_n} \right\} \\ &= \frac{(q; q)_n}{(ax; q)_n} \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{y^k}{(q; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{(q^n; q)_{k-i} (ax; q)_i a^{k-i}}{(q; q)_{n-i} (aq^n x; q)_k} x^{n-i} \\ &= \frac{1}{(ax; q)_n} \sum_{i=0}^n \begin{bmatrix} n \\ i \end{bmatrix}_q u^{\binom{i}{2}} \frac{(ax; q)_i}{(aq^n x; q)_i} x^{n-i} y^i \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{(q^n; q)_k (u^i ay)^k}{(q; q)_k (aq^{n+i} x; q)_k} \\ &= \frac{1}{(ax; q)_n} \sum_{i=0}^n \begin{bmatrix} n \\ i \end{bmatrix}_q u^{\binom{i}{2}} \frac{(ax; q)_i}{(aq^n x; q)_i} x^{n-i} y^i {}_1\Phi_1 \left(\begin{matrix} q^n \\ aq^{n+i} x \end{matrix}; q, uq^{-1}, -u^i ay \right). \end{aligned}$$

□

Theorem 11. For all $n \geq 0$

$$\begin{aligned} & \mathbb{T}(zD_q|v) \left\{ \frac{1}{(ax; q)_n} {}_1\Phi_1 \left(\begin{matrix} q^n \\ aq^n x \end{matrix}; q, uq^{-1}, -ay \right) \right\} \\ &= \frac{1}{(ax; q)_n} \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{(q^n; q)_k (ay)^k}{(q; q)_k (axq^n; q)_k} {}_1\Phi_1 \left(\begin{matrix} q^{n+k} \\ aq^{n+k} x \end{matrix}; q, vq^{-1}, -az \right). \quad (14) \end{aligned}$$

Proof. From Theorem 8, we have

$$\begin{aligned} & \mathbb{T}(zD_q|v) \left\{ \frac{1}{(ax; q)_n} {}_1\Phi_1 \left(\begin{matrix} q^n \\ aq^n x \end{matrix}; q, uq^{-1}, -ay \right) \right\} \\ &= \mathbb{T}(zD_q|v) \left\{ \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{(q^n; q)_k}{(q; q)_k (ax; q)_{n+k}} (-ay)^k \right\} \\ &= \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{(q^n; q)_k}{(q; q)_k} (-ay)^k \mathbb{T}(zD_q|v) \left\{ \frac{1}{(ax; q)_{n+k}} \right\} \\ &= \frac{1}{(ax; q)_n} \sum_{k=0}^{\infty} u^{\binom{k}{2}} \frac{(q^n; q)_k (ay)^k}{(q; q)_k (axq^n; q)_k} {}_1\Phi_1 \left(\begin{matrix} q^{n+k} \\ aq^{n+k} x \end{matrix}; q, vq^{-1}, -az \right) \end{aligned}$$

as claimed. □

Theorem 12.

$$\begin{aligned} & \mathbb{T}(yD_q|v) \left\{ {}_r\Phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1x, b_2, \dots, b_s \end{matrix} ; q, u, z \right) \right\} \\ &= \sum_{n=0}^{\infty} u^{\binom{n}{2}} \frac{(a_1, a_2, \dots, a_r; q)_n}{(b_1x, b_2, \dots, b_s; q)_n (q; q)_n} \\ & \quad {}_1\Phi_1 \left(\begin{matrix} q^n \\ q^n b_1x \end{matrix} ; q, vq^{-1}, -b_1y \right) \left((-1)^n q^{\binom{n}{2}} \right)^{1+s-r} z^n. \end{aligned}$$

Proof.

$$\begin{aligned} & \mathbb{T}(yD_q|v) \left\{ {}_r\Phi_s \left(\begin{matrix} a_1, a_2, \dots, a_r \\ b_1x, b_2, \dots, b_s \end{matrix} ; q, u, z \right) \right\} \\ &= \mathbb{T}(yD_q|v) \left\{ \sum_{n=0}^{\infty} u^{\binom{n}{2}} \frac{(a_1, a_2, \dots, a_r; q)_n}{(b_1x, b_2, \dots, b_s; q)_n (q; q)_n} \left((-1)^n q^{\binom{n}{2}} \right)^{1+s-r} z^n \right\} \\ &= \sum_{n=0}^{\infty} u^{\binom{n}{2}} \mathbb{T}(yD_q|v) \left\{ \frac{1}{(b_1x; q)_n} \right\} \frac{(a_1, a_2, \dots, a_r; q)_n}{(b_2, \dots, b_s; q)_n (q; q)_n} \left((-1)^n q^{\binom{n}{2}} \right)^{1+s-r} z^n \\ &= \sum_{n=0}^{\infty} u^{\binom{n}{2}} \frac{(a_1, a_2, \dots, a_r; q)_n}{(b_1x, b_2, \dots, b_s; q)_n (q; q)_n} \\ & \quad {}_1\Phi_1 \left(\begin{matrix} q^n \\ q^n b_1x \end{matrix} ; q, vq^{-1}, -b_1y \right) \left((-1)^n q^{\binom{n}{2}} \right)^{1+s-r} z^n. \end{aligned}$$

□

5 New q -identities from old one

5.1 q -identity from the q -Gauss sum

The q -Gauss sum is given by [Gasper]:

$${}_2\phi_1 \left(\begin{matrix} a, b \\ c \end{matrix} ; q, c/ab \right) = \frac{(c/a, c/b; q)_{\infty}}{(c, c/ab; q)_{\infty}}. \quad (15)$$

Theorem 13.

$$\begin{aligned} & \frac{(q; q)_k}{(1-q)(c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{(1-aq^i)(1-bq^i)(c; q)_i (a; q)_i (b; q)_i}{(1-cq^{k+1})(cq^k; q)_i (q; q)_i} \left(\frac{1}{ab} \right)^n \\ & \quad \times {}_4\phi_3 \left(\begin{matrix} aq^{i+1}, bq^{i+1}, q^{k+1}, q \\ cq^{k+i+1}, q^{i+1}, q^2 \end{matrix} ; q, c/ab \right) \\ &= \frac{(ax, bx; q)_{\infty}}{(cx, dx; q)_{\infty}} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{q^{\binom{i}{2}} (cx, dx; q)_i}{(bx; q)_i} \sum_{j=0}^i \begin{bmatrix} i \\ j \end{bmatrix}_q q^{j(j-i)} \frac{(-a)^j (-b)^{i-j}}{(ax; q)_j} \\ & \quad \times \sum_{l=0}^{k-i} \begin{bmatrix} k-i \\ l \end{bmatrix}_q (dq^i x; q)_l c^l d^{k-i-l}. \end{aligned}$$

Proof. On the one hand,

$$\begin{aligned}
& D_{c,q}^k \left\{ {}_2\phi_1 \left(\begin{matrix} a, b \\ c \end{matrix}; q, c/ab \right) \right\} \\
&= \sum_{n=0}^{\infty} \frac{(a; q)_n (b; q)_n}{(q; q)_n} \left(\frac{1}{ab} \right)^n D_q^k \left\{ \frac{c^n}{(c; q)_n} \right\} \\
&= \sum_{n=0}^{\infty} \frac{(a; q)_n (b; q)_n}{(q; q)_n} \left(\frac{1}{ab} \right)^n \frac{(q; q)_n}{(c; q)_k (cq^k; q)_n} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{(q^n; q)_{k-i} (c; q)_i}{(q; q)_{n-i}} c^{n-i} \\
&= \frac{1}{(c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q (c; q)_i c^{-i} \sum_{n=i}^{\infty} \frac{(a; q)_n (b; q)_n (q^n; q)_{k-i}}{(cq^k; q)_n (q; q)_{n-i}} \left(\frac{c}{ab} \right)^n \\
&= \frac{1}{(c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q (c; q)_i c^{-i} \sum_{n=0}^{\infty} \frac{(a; q)_{n+i} (b; q)_{n+i} (q^{n+i}; q)_{k-i}}{(cq^k; q)_{n+i} (q; q)_n} \left(\frac{c}{ab} \right)^{n+i} \\
&= \frac{1}{(c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{(c; q)_i (a; q)_i (b; q)_i}{(cq^k; q)_i} \left(\frac{1}{ab} \right)^n \sum_{n=0}^{\infty} \frac{(aq^i; q)_n (bq^i; q)_n (q^{n+i}; q)_{k-i}}{(cq^{k+i}; q)_n (q; q)_n} \left(\frac{c}{ab} \right)^n.
\end{aligned}$$

As $(q^{n+i}; q)_{k-i} = \frac{(q^n; q)_k}{(q^n; q)_i}$ and $(q^0; q)_m = 0$, then using Eq.(5) we have

$$\begin{aligned}
& D_{c,q}^k \left\{ {}_2\phi_1 \left(\begin{matrix} a, b \\ c \end{matrix}; q, c/ab \right) \right\} \\
&= \frac{1}{(c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{(c; q)_i (a; q)_i (b; q)_i}{(cq^k; q)_i} \left(\frac{1}{ab} \right)^n \sum_{n=0}^{\infty} \frac{(aq^i; q)_{n+1} (bq^i; q)_{n+1} (q^{n+1}; q)_k}{(cq^{k+i}; q)_{n+1} (q; q)_{n+1} (q^{n+1}; q)_i} \left(\frac{c}{ab} \right)^n \\
&= \frac{(q; q)_k}{(1-q)(c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{(1-aq^i)(1-bq^i)(c; q)_i (a; q)_i (b; q)_i}{(1-cq^{k+1})(cq^k; q)_i (q; q)_i} \left(\frac{1}{ab} \right)^n \\
&\quad \times \sum_{n=0}^{\infty} \frac{(aq^{i+1}; q)_n (bq^{i+1}; q)_n (q^{k+1}; q)_n}{(cq^{k+i+1}; q)_n (q^2; q)_n (q^{i+1}; q)_n} \left(\frac{c}{ab} \right)^n \\
&= \frac{(q; q)_k}{(1-q)(c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{(1-aq^i)(1-bq^i)(c; q)_i (a; q)_i (b; q)_i}{(1-cq^{k+1})(cq^k; q)_i (q; q)_i} \left(\frac{1}{ab} \right)^n \\
&\quad \times {}_4\phi_3 \left(\begin{matrix} aq^{i+1}, bq^{i+1}, q^{k+1}, q \\ cq^{k+i+1}, q^{i+1}, q^2 \end{matrix}; q, c/ab \right).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
& D_q^k \left\{ \frac{(ax, bx; q)_{\infty}}{(cx, dx; q)_{\infty}} \right\} \\
&= \frac{(ax, bx; q)_{\infty}}{(cx, dx; q)_{\infty}} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q \frac{q^{\binom{i}{2}} (cx, dx; q)_i}{(bx; q)_i} \sum_{j=0}^i \begin{bmatrix} i \\ j \end{bmatrix}_q q^{j(j-i)} \frac{(-a)^j (-b)^{i-j}}{(ax; q)_j} \\
&\quad \times \sum_{l=0}^{k-i} \begin{bmatrix} k-i \\ l \end{bmatrix}_q (dq^i x; q)_l c^l d^{k-i-l}.
\end{aligned}$$

□

5.2 q -identities from q -Chu-Vandermonde's sum

The q -Chu-Vandermonde's sum [4] is given by:

$${}_2\phi_1(q^{-n}, a; c; q, q) = \frac{(c/a; q)_n}{(c; q)_n} a^n. \quad (16)$$

Theorem 14. For $k \leq n$,

$$\begin{aligned} & {}_3\phi_2 \left(\begin{matrix} q^{-n+1}, aq, q^{k+1} \\ cq^{k+1}, q^2 \end{matrix}; q, q \right) \\ &= a^n \frac{(1-q)(q; q)_n (c; q)_{k+1} (c/a; q)_n}{q(q; q)_k (c; q)_n (q^n c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{\binom{i}{2}} (-1/a)^i \frac{(c; q)_i (q^n; q)_{k-i}}{(q; q)_{n-i} (c/a; q)_i}. \end{aligned}$$

Proof. On the one hand,

$$D_{c,q}^k \left\{ {}_2\phi_1 \left(\begin{matrix} q^{-n}, a \\ c \end{matrix}; q, q \right) \right\} = \frac{(q; q)_k q}{(c; q)_{k+1} (1-q)} {}_3\phi_2 \left(\begin{matrix} q^{-n+1}, aq, q^{k+1} \\ cq^{k+1}, q^2 \end{matrix}; q, q \right).$$

On the other hand,

$$D_{c,q}^k \left\{ \frac{(c/a; q)_n}{(c; q)_n} \right\} = \frac{(c/a, q; q)_n}{(c; q)_n (q^n c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{\binom{i}{2}} (-1/a)^i \frac{(c; q)_i (q^n; q)_{k-i}}{(q; q)_{n-i} (c/a; q)_i}.$$

□

Set $a = 1$ in the above theorem. Then for all $c \in \mathbb{R}$,

$${}_3\phi_2 \left(\begin{matrix} q^{-n+1}, q, q^{k+1} \\ cq^{k+1}, q^2 \end{matrix}; q, q \right) = 0$$

Theorem 15.

$$\begin{aligned} & \sum_{k=0}^n \frac{(q^{-n}, a; q)_k}{(c; q)_k (q; q)_k} {}_1\Phi_1 \left(\begin{matrix} q^k \\ q^k c \end{matrix}; q, uq^{-1}, -y \right) q^k \\ &= \frac{(c/a; q)_n}{(c; q)_n} \sum_{i=0}^n (uq)^{\binom{i}{2}} \begin{bmatrix} n \\ i \end{bmatrix}_q \frac{(c; q)_i (-a^{-1}y)^i}{(c/a; q)_i (q^n c; q)_i} {}_1\Phi_1 \left(\begin{matrix} q^n \\ q^{n+i} c \end{matrix}; q, uq^{-1}, -u^i y \right). \end{aligned}$$

Proof. On the one hand,

$$\begin{aligned} & \Gamma(yD_{c,q}|u) \left\{ {}_2\phi_1 \left(\begin{matrix} q^{-n}, a \\ c \end{matrix}; q, q \right) \right\} \\ &= \sum_{k=0}^n \frac{(q^{-n}, a; q)_k}{(q; q)_k} \Gamma(yD_{c,q}|u) \left\{ \frac{1}{(c; q)_k} \right\} q^k \\ &= \sum_{k=0}^n \frac{(q^{-n}, a; q)_k}{(c; q)_k (q; q)_k} {}_1\Phi_1 \left(\begin{matrix} q^k \\ q^k c \end{matrix}; q, uq^{-1}, -y \right) q^k \end{aligned}$$

On the other hand,

$$\begin{aligned} & \Gamma(yD_{c,q}|u) \left\{ \frac{(c/a; q)_n}{(c; q)_n} \right\} \\ &= \frac{(c/a; q)_n}{(c; q)_n} \sum_{i=0}^n (uq)^{\binom{i}{2}} \begin{bmatrix} n \\ i \end{bmatrix}_q \frac{(c; q)_i (-a^{-1}y)^i}{(c/a; q)_i (q^n c; q)_i} {}_1\Phi_1 \left(\begin{matrix} q^n \\ q^{n+i} c \end{matrix}; q, uq^{-1}, -u^i y \right). \quad (17) \end{aligned}$$

□

5.3 q -identity from Jackson's transformation formula

Jackson's transformation of ${}_2\phi_1$ -series is

$${}_2\phi_1 \left(\begin{matrix} a, b \\ c \end{matrix}; q, z \right) = \frac{(az; q)_\infty}{(z; q)_\infty} {}_2\phi_2 \left(\begin{matrix} a, c/b \\ c, az \end{matrix}; q, bz \right). \quad (18)$$

Theorem 16.

$$\begin{aligned} & {}_3\phi_2 \left(\begin{matrix} aq, bq, q^{k+1} \\ cq^{k+1}, q^2 \end{matrix}; q, z \right) \\ &= -\frac{b(1-cq^k)(az; q)_\infty}{(1-a)(1-b)(z; q)_\infty} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{2\binom{i}{2}} \frac{(1-aq^i)(1-cq^i/b)(1-q^{i+1})(a; q)_i (c; q)_i}{(1-azq^i)(1-cq^{k+i})(az; q)_i (cq^k; q)_i} (qz)^i \\ & \quad \times {}_4\phi_4 \left(\begin{matrix} aq^{i+1}, cq^{i+1}/b, q^{i+2}, q^{k+1} \\ azq^{i+1}, cq^{k+i+1}, q^2, q^{i+1} \end{matrix}; q, q^{i+1}bz \right). \end{aligned}$$

Proof. On the one hand,

$$D_{c,q}^k \left\{ {}_2\phi_1 \left(\begin{matrix} a, b \\ c \end{matrix}; q, z \right) \right\} = \frac{(1-a)(1-b)(q; q)_{kz}}{(c; q)_{k+1}(1-q)} {}_3\phi_2 \left(\begin{matrix} aq, bq, q^{k+1} \\ cq^{k+1}, q^2 \end{matrix}; q, z \right).$$

On the other hand,

$$\begin{aligned} & D_{c,q}^k \left\{ {}_2\phi_2 \left(\begin{matrix} a, c/b \\ c, az \end{matrix}; q, bz \right) \right\} \\ &= \sum_{n=0}^{\infty} (-1)^n q^{\binom{n}{2}} \frac{(a; q)_n}{(az; q)_n} (bz)^n D_{c,q}^k \left\{ \frac{(c/b; q)_n}{(c; q)_n} \right\} \\ &= \sum_{n=0}^{\infty} (-1)^n q^{\binom{n}{2}} \frac{(a; q)_n}{(az; q)_n} (bz)^n \frac{(c/b; q)_n (q; q)_n}{(c; q)_n (cq^n; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{\binom{i}{2}} (-1/b)^i \frac{(c; q)_i (q^n; q)_{k-i}}{(q; q)_{n-i} (c/b; q)_i} \\ &= \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{\binom{i}{2}} (-1/b)^i \frac{(c; q)_i}{(c/b; q)_i} \sum_{n=i}^{\infty} (-1)^n q^{\binom{n}{2}} \frac{(a; q)_n (c/b; q)_n (q; q)_n (q^n; q)_{k-i}}{(az; q)_n (c; q)_n (cq^n; q)_k (q; q)_{n-i}} (bz)^n. \end{aligned}$$

From Identities Eq.(4) and Eq.(5) and simplifying it, we have

$$\begin{aligned} & D_{c,q}^k \left\{ {}_2\phi_2 \left(\begin{matrix} a, c/b \\ c, az \end{matrix}; q, bz \right) \right\} \\ &= \frac{1}{(c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{2\binom{i}{2}} \frac{(a; q)_i (c; q)_i (q; q)_i}{(az; q)_i (cq^k; q)_i} z^i \\ & \quad \times \sum_{n=0}^{\infty} (-1)^n q^{\binom{n}{2}} \frac{(aq^i; q)_n (cq^i/b; q)_n (q^{i+1}; q)_n (q^n; q)_k}{(azq^i; q)_n (q; q)_n (cq^{k+i}; q)_n (q^n; q)_i} (q^i bz)^n. \end{aligned}$$

As $(q^0; q)_n = 0$, then using Eq.(4) we have

$$\begin{aligned}
& D_{c,q}^k \left\{ {}_2\phi_2 \left(\begin{matrix} a, c/b \\ c, az \end{matrix}; q, bz \right) \right\} \\
&= \frac{1}{(c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{2\binom{i}{2}} \frac{(a; q)_i (c; q)_i (q; q)_i}{(az; q)_i (cq^k; q)_i} z^i \\
&\quad \times \sum_{n=0}^{\infty} (-1)^{n+1} q^{\binom{n+1}{2}} \frac{(aq^i; q)_{n+1} (cq^i/b; q)_{n+1} (q^{i+1}; q)_{n+1} (q^{n+1}; q)_k}{(azq^i; q)_{n+1} (q; q)_{n+1} (cq^{k+i}; q)_{n+1} (q^{n+1}; q)_i} (q^i bz)^{n+1} \\
&= -\frac{bz(q; q)_k}{(1-q)(c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{2\binom{i}{2}} \frac{(1-aq^i)(1-cq^i/b)(1-q^{i+1})(a; q)_i (c; q)_i}{(1-azq^i)(1-cq^{k+i})(az; q)_i (cq^k; q)_i} (qz)^i \\
&\quad \times \sum_{n=0}^{\infty} (-1)^n q^{\binom{n}{2}} \frac{(aq^{i+1}; q)_n (cq^{i+1}/b; q)_n (q^{i+2}; q)_n (q^{k+1}; q)_n}{(azq^{i+1}; q)_n (q^2; q)_n (cq^{k+i+1}; q)_n (q^{i+1}; q)_n} (q^{i+1}bz)^n \\
&= -\frac{bz(q; q)_k}{(1-q)(c; q)_k} \sum_{i=0}^k \begin{bmatrix} k \\ i \end{bmatrix}_q q^{2\binom{i}{2}} \frac{(1-aq^i)(1-cq^i/b)(1-q^{i+1})(a; q)_i (c; q)_i}{(1-azq^i)(1-cq^{k+i})(az; q)_i (cq^k; q)_i} (qz)^i \\
&\quad \times {}_4\phi_4 \left(\begin{matrix} aq^{i+1}, cq^{i+1}/b, q^{i+2}, q^{k+1} \\ azq^{i+1}, cq^{k+i+1}, q^2, q^{i+1} \end{matrix}; q, q^{i+1}bz \right).
\end{aligned}$$

□

References

- [1] R. Orozco, Deformed Rogers-Szegö Polynomials and the Generalized q -Exponential Operator, arXiv:2306.07431v4, (2024).
- [2] W.Y.C. Chen, Z.G. Liu, Parameter augmenting for basic hypergeometric series, I, Mathematical Essays in Honor of Gian-Carlo Rota, Eds., B. E. Sagan and R.P. Stanley, Birkhäuser, Boston, 1998, pp. 111–129.
- [3] W.Y.C. Chen, Z.G. Liu, Parameter augmenting for basic hypergeometric series, II, J. Combin. Theory, Ser. A **80** (1997) 175–195.
- [4] G. Gasper, M. Rahman, Basic Hypergeometric Series, 2nd ed., Cambridge University Press, Cambridge, MA, 1990.
- [5] H. L. Saad and M.A. Abdul, The q -Exponential Operator and Generalized Rogers-Szego Polynomials, Journal of Advances in Mathematics, 8(1), (2014) 1440–1455. <https://doi.org/10.24297/jam.v8i1.6912>
- [6] H. L. Saad, and A. A. Sukhi, The q -exponential operator, Applied Mathematical Sciences, **7**, 6369–6380, (2013) <http://dx.doi.org/10.12988/ams.2013.24287>.
- [7] H. A. Ghany, q -derivative of basic hypergeometric series with respect to parameters, Int. J. Math. Anal. **3** 33 (2009).