Efficient Risk-sensitive Planning via Entropic Risk Measures

Alexandre Marthe¹ Samuel Bounan¹ Aurélien Garivier¹ Claire Vernade²

Abstract

Risk-sensitive planning aims to identify policies maximizing some tail-focused metrics in Markov Decision Processes (MDPs). Such an optimization task can be very costly for the most widely used and interpretable metrics such as threshold probabilities or (Conditional) Values at Risk. Indeed, previous work showed that only Entropic Risk Measures (EntRM) can be efficiently optimized through dynamic programming, leaving a hard-to-interpret parameter to choose.

We show that the computation of the *full set of optimal policies* for EntRM across parameter values leads to tight approximations for the metrics of interest. We prove that this *optimality front* can be computed effectively thanks to a novel structural analysis and smoothness properties of entropic risks. Empirical results demonstrate that our approach achieves strong performance in a variety of decision-making scenarios.

1. Introduction

Markov Decision Processes (MDPs) (Puterman, 2014), the central models of modern Reinforcement Learning (RL). capture sequential decision making in domains as diverse as robotics, finance, healthcare, and operations research (Sutton & Barto, 2018; Silver et al., 2017; Charpentier et al., 2021; Polydoros & Nalpantidis, 2017). At their core, MDPs allow agents to optimize actions so as to maximize the expected cumulative reward via dynamic programming or other policy optimization methods. However, in many highstakes applications, such as healthcare, average performance alone is not sufficient. In fact, it may be critical to limit the probability of catastrophic outcomes or to ensure that returns remain above certain thresholds with high confidence. This need has driven research on risk-sensitive RL, which incorporates measures of uncertainty and tail behavior into the control objective (Bäuerle & Rieder, 2014; Tamar et al.,

2015).

One of the central challenges in risk-sensitive RL is to identify risk criteria that are both (1) *meaningful* for real-world decision making and (2) *tractable* in an MDP context. Popular approaches often revolve around the quantile-based Value at Risk (VaR) and Conditional Value at Risk (CVaR) (Artzner et al., 1999; Rockafellar et al., 2000), which are widely used in finance and operations research for bounding tail risk. Another common objective relies on controlling a Threshold Probability (White, 1993), that is the probability that total returns fall below a specified level.

Despite their practical interest and interpretability properties, these common risk metrics cannot be directly and efficiently optimized in MDPs (Marthe et al., 2024; Rowland et al., 2019). Our work addresses precisely this gap by connecting the Threshold Probability and the (C)VaR metrics to the moment-generating function. In fact, in the context of MDPs, we show that these two optimization problems can be carefully approximated by the Entropic (Exponential) Risk Measure (EntRM) (Howard & Matheson, 1972).

The work of (Föllmer & Schied, 2011; Marthe et al., 2024) shows that EntRM is unique among non-linear transformations of the return in admitting a dynamic programming decomposition, making it arguably the "best possible" extension of the risk-neutral Bellman recursion to a risk-aware MDP setting. Yet, despite its computational appeal, practical usage of EntRM can be hindered by interpretability concerns, especially around selecting the risk tolerance parameter β .

We propose a unifying framework for risk-sensitive planning in MDPs through the study of the EntRM. We prove that optimal policies evolve in a structured manner as the risk parameter changes, which allows us to derive an efficient algorithm to compute all the optimal policies for EntRM, a set that we call *the optimality front*. We demonstrate the consequence of this new method on the optimization of the Threshold Probability, the VaR and CVaR via dynamic programming. We provide both theoretical guarantees on the computational complexity and on the approximation error, and an empirical study on an inventory management problem.

¹UMPA, ENS de Lyon, Lyon, France ²University of Tübingen, Tübingen, Germany. Correspondence to: Alexandre Marthe <alexandre.marthe@ens-lyon.fr>.

2. Risk Sensitive Planning

2.1. Risk-sensitivity in MDPs

We consider a finite-horizon MDP $\mathcal{M} = (\mathcal{X}, \mathcal{A}, p, r, H, p_1)$, where \mathcal{X} is a finite set of states, \mathcal{A} is a finite set of actions, $p(x' \mid x, a)$ is the probability of transitioning to state x' from state x when action ais chosen, r(x, a) is the reward function specifying the immediate reward for taking action a in state x (bounded in [-1,1]), H is the finite horizon, i.e., the number of decision steps, and p_1 is the distribution of initial states $x_1 \sim p_1(\cdot)$. To simplify notation, we assume all rewards and transitions are stationary, i.e. $p_t = p$ and $r_t = r \forall t$, and the rewards are deterministic, but our results naturally extend to the non-stationary setting.

A policy $\pi = (\pi_1, \ldots, \pi_H)$ is a sequence of decision rules $\pi_t : \mathcal{X} \to \mathcal{A}$ that, at each step $t \in \{1, 2, \ldots, H\}$, selects an action based on the current state. For simplicity, we assume policies to be deterministic, but not necessarily stationary (a stationary policy verifies $\forall t, \pi_1 = \pi_t$). We denote $\tau = (x_1, a_1, \ldots, x_H, a_H)$ a trajectory induced by π . The corresponding *cumulative reward* (or *return*) is

$$R^{\pi} = \sum_{t=1}^{H} r(x_t, a_t)$$

where $x_0 \sim p_1(\cdot)$, $a_t = \pi_t(x_t)$ and $x_{t+1} \sim p(\cdot | x_t, a_t)$. We also denote $R_h^{\pi}(x, a) = \sum_{t=h}^{H} r(x_t, a_t)$ with $x_h = x, a_h = a$, and $R_h^{\pi}(x) = R_h^{\pi}(x, \pi(x))$.

The classical objective in MDPs is to find a policy π^* that maximizes the expected return, $\mathbb{E}_{\pi}[R^{\pi}]$, which is inadequate for risk-sensitive applications. In turn, risk-sensitive RL studies generic optimization problems $\max_{\pi} \rho(R^{\pi})$ that extend the standard MDP framework by incorporating a functional ρ that captures *tail events* or *uncertainty aversion*.

2.2. Risk metrics and computational limitations in MDPs

The theory of risk measures is too rich for us to provide more than a few insights below on the most widely used examples. However, it has been shown (Kupper & Schachermayer, 2009; Föllmer & Schied, 2011; Rowland et al., 2019; Marthe et al., 2024) that only the *Entropic Risk Measure* satisfies a recursive Dynamic Programming equation akin to the Bellman Equation (Sutton & Barto, 2018), which is the cornerstone of RL algorithms and what makes the optimization problem tractable.

Entropic Risk Measure (EntRM). Originally introduced by Howard & Matheson (1972) for MDPs, the EntRM for a

random variable X and parameter $\beta \in \mathbb{R}$ is defined as

$$\operatorname{EntRM}_{\beta}[X] = \begin{cases} \frac{1}{\beta} \log(\mathbb{E}[e^{\beta X}]), & \text{if } \beta \neq 0, \\ \mathbb{E}[X], & \text{if } \beta = 0. \end{cases}$$
(1)

In the MDP context, maximizing $\text{EntRM}_{\beta}[R^{\pi}]$ is also equivalent to maximizing $\text{sign}(\beta) \mathbb{E}[e^{\beta R^{\pi}}]$, by composition with a monotonous function. We refer to this as the *exponential form* of EntRM, and will use both forms throughout the paper, noting that the core optimization problem remains the same.

The parameter β encodes risk tolerance: large positive β values emphasize a risk-seeking attitude (amplifying high returns), while negative values induce conservativeness, diminishing the impact of large potential losses. Despite its interpretive appeal in terms of tail emphasis, β can still be difficult to set in practice, and the measure is not *positively homogeneous*, i.e. $EntRM_{\beta}[cX] \neq c EntRM_{\beta}[X]$ for $c \in \mathbb{R}$. In finance, this is problematic when scaling outcomes by different currencies or units. However, MDPs typically normalize rewards, mitigating some of these issues. Practitioners nonetheless often prefer quantile-based risk measures like VaR or CVaR. See Appendix A for more details on the EntRM. In the following, the risk parameter is always assumed to be non-positive.

Value at Risk (VaR). VaR represents the quantile of the distribution. At level $\alpha \in (0, 1)$:

$$\operatorname{VaR}_{\alpha}[R^{\pi}] = \inf \left\{ x \mid \Pr(R^{\pi} \le x) \ge \alpha \right\}.$$
 (2)

Although VaR is widely used, it suffers from a lack of *sub-additivity*: it can be manipulated by adding or removing redundant risks. In the context of MDPs, VaR is particularly challenging to optimize due to its non-linearity and non-monotonicity (Chow et al., 2018).

Conditional Value at Risk (CVaR). To address some limitations of VaR, the *Conditional Value at Risk* (Rockafellar et al., 2000; Föllmer & Schied, 2011; Bäuerle & Ott, 2011; Chow & Ghavamzadeh, 2014) at level α is given by

$$\operatorname{CVaR}_{\alpha}[R^{\pi}] = \frac{1}{\alpha} \int_{0}^{\alpha} \operatorname{VaR}_{\gamma}[R^{\pi}] d\gamma.$$
 (3)

CVaR represents the expected return in the α -worst fraction of outcomes, providing a more comprehensive view of the risk than VaR. What makes it particularly interesting is that it is a *coherent risk measure*, meaning that it satisfies both sub-additivity and positive homogeneity, compared to the risk measure mentioned previously.

In this article, we chose the convention of VaR and CVaR representing the low tail of the distributions. In the literature,

it is often defined using the high tail. This has no impact on the theory as a change of variable $X \leftarrow -X$ will transpose the results. More details about the different notations can be found in Appendix B.

Threshold Probability. Certainly the most intuitive measure of risk is the probability of falling below a userspecified threshold level T. Solving $\min_{\pi} \Pr(R^{\pi} \leq T)$, means seeking a policy whose probability of yielding a return below T is minimal. The VaR and Threshold optimization are **dual** problems. A farmer who worries about the possibility of a very poor harvest in the coming year will either ask, "What is the chance that my yield will be below 2 tons?" (Threshold Probability viewpoint), or "With 90% confidence, how large will my yield be?" (VaR perspective). Lowering the probability of dropping below a certain threshold (Threshold Probability) is directly tied to choosing a quantile-based cutoff for outcomes, as $\operatorname{VaR}_{\alpha}[X] = T$ just means that $P(X \leq T) = \alpha$.

Computational limitations. The EntRM verifies a Bellman Equation similar to the one of the expected return, making it computationally efficient (See Appendix A for details). Yet, optimization is much more complex for the other risk measures. Optimization of VaR and especially CVaR in MDPs have been studied extensively (Chow & Ghavamzadeh, 2014; Chow et al., 2015; 2018; Achab & Neu, 2021; Bäuerle & Ott, 2011), but remains a challenging task. Optimal policies are usually non-Markovian, meaning they depend on the accumulated reward and not only the current state. This makes it harder to compute and implement in practice. Common Dynamic Programming schemes require augmenting the state space with a continuous variable, which is computationally expensive and makes the problem intractable except for very simple MDPs.

Approximation schemes have been proposed, such as Dynamic Risk Measures (Bäuerle & Glauner, 2022) (also called Nested or Recursive Risk Measures). Where the goal is to optimize recursively $V_h(x) = \max_a \rho[r(x, a) + V_{h+1}(X')],$ where ρ is the risk measure, such as Var or CVaR. This problem has a natural Dynamic Programming formula that make computationnaly simple, but it does not optimize precisely the risk measure of choice. Also, the approximation is usually quite poor and the optimized objective is not law invariant, which makes it harder to interpret. Discretizing the augmented-state is another common approach, but the results are not always satisfying (Hau et al., 2024). Optimizing the Threshold Probability raises the same challenges as VaR and CVaR (White, 1993; Wu & Lin, 1999; Kira et al., 2012) but seems to have been less studied and the literature lacks approximation schemes.

In short, while the EntRM is the only risk measure that can be efficiently optimized in MDPs, it is not what people use in practice. Practitioners prefer using more interpretable measures of risk, which are usually not tractable in MDPs. How can we leverage the computational properties of EntRM to optimize those more preferred measures of risk ? In the sequel, we show that optimizing the EntRM over a range of risk parameters can lead to good approximation of optimal policies for several problems.

3. A unified framework for Threshold Probability and VaR/CVaR optimization.

The Threshold Probability and VaR/CVaR objectives are all related to the tail probabilities of the return distribution. These tail probabilities can be approximated with the help of exponential moments of the distribution by Chernoff's bound:

$$\Pr(X \le T) \le \inf_{\beta \le 0} \exp(-\beta T) \mathbb{E}[e^{\beta X}].$$
(4)

Exponential moments are the core of Entropic Risk Measure and we explain in this section how Inequality (4) leads to proxies for the risk metrics introduced above.

A proxy for the Threshold Probability. In the case of the Threshold Probability, the related proxy is pretty direct:

$$\min_{\pi} \Pr\left(R^{\pi} \leq T\right) \leq \min_{\beta < 0} \min_{\pi} e^{-\beta T} \mathbb{E}\left[e^{\beta R^{\pi}}\right].$$
(5)

Some insights are given in Appendix C.1.

From (C)VaR to EVaR. Solving for the rhs of (4) to equal α , Ahmadi-Javid (2012) introduced the *Entropic Value at Risk (EVaR)* as a proxy for the VaR defined by

$$\operatorname{EVaR}_{\alpha}[X] = \sup_{\beta < 0} \operatorname{EntRM}_{\beta}[X] - \frac{1}{\beta} \log(\alpha).$$

It is the best approximation of the VaR based on exponential moments, and one proves that $\operatorname{VaR}_{\alpha}[X] \geq \operatorname{CVaR}_{\alpha}[X] \geq \operatorname{EVaR}_{\alpha}[X]$. Hence, the EVaR indeed appears as an even better approximation for the CVaR. EVaR is a *coherent* risk measure on its own, and its use for approximating VaR and CVaR has been of growing interest recently in MDPs (Ni & Lai, 2022; Hau et al., 2023; Su et al., 2024). The related Proxy for VaR and CVaR is

$$\max_{\pi} \operatorname{VaR}_{\alpha} \left[R^{\pi} \right] \geq \sup_{\beta < 0} \max_{\pi} \operatorname{EntRM}_{\beta} \left[R^{\pi} \right] - \frac{\log(\alpha)}{\beta}.$$
(6)

Relaxed optimization problems. These proxies directly result in new optimization problems for the respective risk metrics. We note π_{β}^{*} the optimal policy for the EntRM with

parameter β . For the Threshold Probability, the optimal policy is π_{β}^* where the risk parameter β corresponds to

$$\operatorname*{argmin}_{\beta < 0} e^{-\beta T} \mathbb{E} \left[e^{\beta R^{\pi \hat{\beta}}} \right], \tag{7}$$

Similarly, for the Var/CVaR, we have

$$\arg \sup_{\beta < 0} \operatorname{EntRM}_{\beta} \left[R^{\pi_{\beta}^{*}} \right] - \frac{1}{\beta} \log(\alpha) \tag{8}$$

(which is effectively an *EVaR optimization problem*).

Those two problems are reduced to optimizing over the EntRM. There are some direct implications of this transformation in terms of the properties of the optimal policies.

Proposition 3.1 (Hau et al. (2023)). The optimal policies for the previous problems (7)-(8) are deterministic and Markovian. For each problem, there exists $\beta < 0$ such that the optimal policy is also optimal for the EntRM with parameter β .

Compared to the initial problems where optimal policies are usually not markovian, the transformation to the EntRM allows finding optimal policies that are easier to compute and implement.

Yet, in those new optimization problems, optimizing the EntRM is not done only for a specific parameter, but for a whole range of values. Current methods for optimizing the EntRM in MDPs are not adapted to this problem, and works for a single parameter at a time. A natural way of getting around this issue is by discretizing over the range of values of β .

Quality of the approximations. The bounds above depend on the tail of the distributions which are known to be more accurate for distributions with light tails (Vershynin, 2018). In MDPs, it means that they are tighter for rich reward signals.

On the other hand, existing methods to optimize VaR, CVaR and Threshold Probability rely on dynamic programming on extended MDPs, where the state space is augmented with a continuous variable that correspond to the achievable values of the return (Chow & Ghavamzadeh, 2014; White, 1993). Thus, even for small MDPs, rich reward signals may quickly increase the dimension to an extent that renders the optimization intractable. Conveniently, this is when our approximation method is relevant.

Moreover, while there is work on CVaR/VaR, this approximation scheme for the Threshold Probability problem remains unstudied so far in the literature.

3.1. Grid-Based Optimization of the Risk Parameter

A natural first idea to optimize the EVaR in MDPs is to discretize β on a grid. By considering well-chosen values of β ,

Hau et al. (2023) show that they can get an ε -approximation of the optimal policy for the EVaR problem with a complexity of $O(|\mathcal{S}|^2 |\mathcal{A}| H \frac{\log(1/\varepsilon)}{\varepsilon^2})$.

We prove a similar result for the Chernoff approximation of the Threshold Probability problem.

Proposition 3.2. Let R be the return of the MDP such that there exists a < 0 and p > 0 with the property that for any policy π , $\Pr(R \le a) \ge p$. Then solving Equation (7) with accuracy $2\log(1 + \varepsilon)/H$ on β and $\beta_{\min} = \ln(p)/a$, finds a policy π that satisfies

$$\mathbb{P}(R^{\pi} \leq 0) \leq B$$
 and $B \leq B \leq (1+\varepsilon)B$,

where *B* is the true optimal value.

Hence, one needs to compute

$$\frac{H\,\beta_{\min}}{2\,\log(1+\varepsilon)}$$

policies to obtain a value within a factor $(1 + \varepsilon)$ of the optimum. Given that computing an optimal policy for EntRM involves a complexity of $O(|\mathcal{S}|^2 |\mathcal{A}| H)$, the overall complexity to achieve an approximation ratio of $(1 + \varepsilon)$ is

$$O\left(\frac{H^2 |\mathcal{S}|^2 |\mathcal{A}| \beta_{\min}}{\log(1+\varepsilon)}\right)$$

As expected, those bounds show that the complexity explodes as the grid is refined to obtain more accurate approximations. However, intuition suggests that computing very close values of risk parameters should lead to the same optimal policy via similar computations. Can there be an efficient method to optimize over a continuous range of risk parameters for the EntRM problem, avoiding redundant computation? Besides, the grid-based optimization computes several EntRM-optimal policies in order to optimize the proxy. Yet, the obtained result might not be the best EntRM-optimal policy for the initial problem (Threshold Probability or (C)VaR). The next section provides answers to these points by a careful study of the structure of optimal policies for the entropic risk over all values of the risk parameter.

4. Structural Insights into Entropic Risk Measures

Optimizing EntRM for an entire range of risk parameters requires understanding the structure of EntRM optimal policies. We now show that exploiting the regularity of the EntRM_{β} function leads to an efficient algorithm that computes *all the optimal policies* along $\beta \in \mathbb{R}$ much more efficiently than using a grid. We also show that this proves useful for surprisingly many risk-sensitive objectives. In passing, understanding how a small perturbation of the risk parameter can influence the optimal policy is also helpful from the point of view of interpretability and robustness (Bäuerle et al., 2024). All the proofs are given in the appendix.

4.1. Structure Analysis

Definition 4.1 (Optimality Front). For any MDP, the *optimality front* is defined as $\Gamma = (\pi_k, I_k)_k$, where $(I_k)_k$ is a partition of the risk tolerance set and where π_k is the optimal policy of EntRM for all risk tolerance parameters $\beta \in I_k$.

This definition is justified by the following property, formalizing the intuition that a small perturbation of the risk parameter typically does not change the optimal policy: the mapping $\beta \mapsto \pi_{\beta}^{*}$ is piecewise constant.

Proposition 4.2. The Optimality Front Π^* is a finite set of policies. Each policy in Π^* is optimal on a finite union of closed intervals.

Crucially, we are able to prove lower bounds on the length of these intervals for a specific risk parameter β , knowing the Advantage function at this specific point.

Theorem 4.3. Let $\beta \in \mathbb{R}$ be such that there is a unique deterministic policy π_{β}^* optimizing EntRM_{β}. Define the Generalized Advantage function:

$$A_{h,\beta}^{\pi}(x,a) = \operatorname{EntRM}_{\beta}[R_{h}^{\pi}(x)] - \operatorname{EntRM}_{\beta}[R_{h}^{\pi}(x,a)].$$

Then, define optimality gaps as the smallest differences over the entire MDP:

•
$$\Delta = \frac{|\beta|}{2} \min_{h,x} \min_{a \neq \pi^*_{\beta,h}(x)} \frac{1}{h} A_{h,\beta}^{\pi^*_{\beta}}(x,a) \text{ if } \beta \neq 0,$$

•
$$\Delta = 2 \min_{h,x} \min_{a \neq \pi^*_{\beta,h}(x)} \frac{1}{h^2} A_{h,\beta}^{\pi^*_{\beta}}(x,a) \text{ if } \beta = 0.$$

Then, for all $\beta' \in [\beta - \Delta, \beta + \Delta]$, the optimal policy for EntRM_{β'} remains π^*_{β} .

The first bound is usually quite good when the risk parameter is not too small, as it scales with β . For large values of β , it balances the small optimality gap (remember that EntRM_{β}[R^{π}] \rightarrow ess inf R^{π} when $\beta \rightarrow -\infty$, so the gaps tend to 0). The degeneracy at $\beta = 0$ is circumvented by the second bound.

Knowing this structure of intervals, the only information we need to determine the optimality front is the location of these subinterval boundaries. We call *breakpoints* these risk-parameter values at which the optimal policy changes, and we show that the resulting change is generally only local. **Proposition 4.4.** Consider a random MDP with reward functions and transitions $(r_t(x, a))_{t,x,a}$ and $(p_t(x))_{t,x}$ generated from, say, independent uniform distributions. With probability 1, for each breakpoint $\beta \in \mathcal{B}$ there is a single state-horizon pair for which the optimal action changes: if π^1 is optimal for $\beta \in [\beta_1, \beta_2]$ and π^2 is optimal for $\beta \in [\beta_2, \beta_3]$, then there exists a unique state x and time step t such that $\pi_t^1(x) \neq \pi_t^2(x)$.

See Appendix D.3 for more details.

4.2. Application to Risk-Sensitive Objectives

The structure of the optimality front allows to rewrite the proxy optimization problems as minimization over a finite set of optimal policies.

Proposition 4.5. Consider $\Gamma = (\pi_k, I_k)_k$ the optimality front.

$$Eq. (7) = \min_{k} \inf_{\beta \in I_{k}} \mathbb{E}\left[e^{\beta(R^{\pi_{k}} - T)}\right]$$
(9)

$$Eq. (8) = \min_{k} \inf_{\beta \in I_{k}} \operatorname{EntRM}_{\beta} \left[R^{\pi_{k}} \right] - \frac{1}{\beta} \log(\alpha) \quad (10)$$

Both these problems are reduced to more simple optimization problems on small intervals that can be solved using gradient methods. Indeed, the first one, for the Threshold Probability, is concave (Boyd & Vandenberghe, 2004), and the second one is quasiconcave¹ (Hau et al., 2023).

While this method can be better if one wishes to truly optimize the EVaR, remember that the initial problem was to find "good" policies for the Threshold Probability, the VaR and the CVaR. Instead of finding the best EntRM optimal policy for the proxy, one can directly evaluate the metric of choice for each policy in the optimality front.

The previous optimization problem is thus reduced to the single and more general:

$$\max_{k} \rho \left(R^{\pi_k} \right) \tag{11}$$

where ρ is the metric of choice and $(\pi_k)_k$ the policies of the optimality front.

4.3. Computing the Optimality Front

The first step towards computing the Optimality Front is to *find the breakpoints*. We first show that a direct approach is not feasible but instead we can exploit Theorem 4.3. Then, we present our algorithm, *Distributional Optimality Front Iteration* (DOLFIN), based on efficient (distributional) value iteration (Bellemare et al., 2023).

¹It becomes a concave problem after using the change of variable $\beta \leftarrow \frac{1}{\beta}$ and thus can still be optimized efficiently

Finding the Breakpoints. Breakpoints mark the transition between two different intervals of optimality. According to Proposition 4.4 there must be at least two optimal policies for those particular parameter values. This yields a system of equations characterizing the breakpoints.

Proposition 4.6. Assume π^1 and π^2 are such that π^1 is optimal for $\beta \in [\beta_1, \beta_b]$ and π^2 is optimal for $\beta \in [\beta_b, \beta_2]$. Then β_b satisfies:

$$\forall h, x, \quad \operatorname{EntRM}_{\beta_2}[R_h^{\pi^1}(x)] = \operatorname{EntRM}_{\beta_2}[R_h^{\pi^2}(x)]$$

Note that since the policies only differ in one state-horizon pair (Prop. 4.4), most of these equations are trivial. Nevertheless, one could attempt to use them to directly compute the breakpoints by resolving a system of equations. Unfortunately, we argue in Appendix E that this is unfeasible due to the lack of regularity of the EntRM functions and to the computational complexity of the problem. In general, we show that the lack of "regularity" in these functions makes it impossible in practice to know in advance how many breakpoints, or optimal policies, might appear between two given points.

However, the advantage of using the EntRM in MDPs is that the optimal policies can be computed recursively by Dynamic Programming and we show that this implies a structure on the breakpoints.

Proposition 4.7. (Informal) Let \mathcal{B}^h be the set of breakpoints at time h and $\mathcal{B}^h(x)$ the corresponding set of breakpoints at state x. Then, the sets verify the recursive equation

$$\mathcal{B}^{h} = \mathcal{B}^{h+1} \cup \left(\bigcup_{x \in \mathcal{X}} \mathcal{B}^{h}(x)\right)$$

A formal statement and the proof can be found in Appendix D.6. This result shows that the set of breakpoints is simply a union over the per-state breakpoints, and they can be computed via a backward recursion.

We now have all the tools to build an incremental approach that we call *FindBreaks*. Theorem 4.3 tells us that knowing the Entropic risk at a given point allows us to identify an interval of β values over which the optimal policy does not change. This fact can be utilized to 'jump' over β values. The process is the following. At a given state, assume the distribution of the return for each action is known $(\eta(x, a))_{a \in \mathcal{A}}$. Start with $\beta = 0$ and iterate the following steps:

- 1. Evaluate the Generalized Advantage function and the optimality gaps (see Theorem 4.3),
- 2. Use the optimality gaps to get a lower bound $\beta \Delta$ on the next breakpoint, and 'jump': $\beta \leftarrow \beta \Delta$

At some point, when getting close to a breakpoint, the increments Δ will get closer to 0. Then use a minimal increment ϵ until the optimal action changes. This can be done in parallel over states thanks to Proposition 4.7. Details and pseudo-code are in Appendix G.

This may not be the optimal way to compute the breakpoints but it exploits all the structure of the problem: both the regularity of the exponential functions and the recursive properties of the MDP optimization allow to speed up the process. The general question of characterizing optimality for this problem is a challenging open problem we leave for future work. Our final risk-sensitive optimization algorithm below is fully modular and could integrate any other breakpoint-search algorithm.

Distributional Optimality Front Iteration. Combining both insights from Dynamic Programming and the approximation of Optimality Intervals, we derive an algorithm to compute the Optimality Front up to a desired accuracy.

This algorithm keeps in memory the distribution of the return recursively. While not compulsory, it accelerates the computation of several values of the EntRM with same reward distribution. For more details on the Dynamic Programming computation of return distributions, see Bellemare et al. (2023).

Algorithm 1 DOLFIN - Distributional Optimality Front Iteration

Require: Precision $\varepsilon \in (0,1)$; MDP $\mathcal{M}(\mathcal{X}, \mathcal{A}, p, R, H)$ parameters. 1: Select lower bound β_{\min} {Computed or handpicked} 2: $\mathcal{I}_H \leftarrow [\beta_{\min}, 0]$ {Starting interval} 3: $\nu_H(x) \leftarrow \delta_0$ {Optimal return distribution at timestep *H*} 4: for h = H to 1 do 5: for $x \in \mathcal{X}$ do 6: for $I \in \mathcal{I}_h$ do $\eta_h^I(x,a) \leftarrow \varrho(x,a) * \sum_{x'} p(x' \mid x,a) \nu_{h+1}^I(x')$ 7: {Return distributions} $\{\mathcal{J}, (a_i^*)_{i \in \mathcal{J}}\} \leftarrow \text{FINDBREAKS}(\epsilon, (\eta_h^I(x, a))_a, I)$ 8: {Apply Algorithm 2 on $(\eta_h^I(x, a))_a$ as the reward distribution for each action} 9: for $j \in \mathcal{J}$ do 10: Add *j* to \mathcal{I}_{h-1} {Update intervals for next timestep} $\nu_h^j(x) \leftarrow \eta_h^I(x, a_i^*)$ {Store optimal return distribu-11: tion} 12: end for 13: end for 14: end for 15: end for **output** $\Gamma = (\pi_k, I_k)_k, (\eta_0^k)^k$ {Optimality Front, distributions}

DOLFIN returns the Optimality Front Γ and it remains to solve (11): $\min_k \rho(\pi^k)$. In the experimental section below, we simply call this the *Optimality Front* method.

About the computational cost. Calling B the number of breakpoints in the optimality front of the MDP, the number of calls to FindBreaks is bounded by $\Theta(|\mathcal{X}|HB)$ and thus heavily depend on the number of optimal policies. For a low number, only a few calls will be made and only a few O-value evaluations will have to be computed. Using the empirical observations on the number of breakpoints, the number of calls to can be estimated to be in the order of $(|\mathcal{X}|H)^2$. The total complexity ultimately depends on the complexity of FindBreaks. An empirical study of our implementation can be found in Appendix G. In practice we observe that FindBreaks runs in $O(|A|f(1/\varepsilon))$ time, with some sublinear f. When the support of the accumulated reward is too large, the implementation of the distributional induction can benefit from the approximation schemes described in (Bellemare et al., 2023).

5. Numerical Experiments

This section evaluates the effectiveness and efficiency of the Optimality Front approach. As this is the first study to explore policies for Threshold Probability optimization in MDPs, we aim to assess how well these policies perform, compared to naive approximation methods. We then compare the performance of the Optimality Front on the VaR and CVaR metrics against the method introduced by Hau et al. (2023). Lastly, we analyze the efficiency of our algorithm, focusing on the proposed approximation method. We quantify the improvements due to *FindBreaks* compared to a naive grid-based approach, highlighting its practical advantages in reducing redundant computations.

Environment We implemented *Optimality Front* on the Inventory Management MDP (Bellman et al., 1955; Scarf et al., 1960). In this example, the goal is to maximize the profit of a store selling one extensive good. The store has a strict maximal capacity of M = 10. At each time step, the state of the store is its number of available goods, $x_t \in [M]$, and it can buy (action) a quantity $a_t \in [M]$ of new goods. The reward obtained is $r_t = [f(D_t, x_t, a_t) - C_m(x_t) - C_c(a_t)]/4M$ with D_t is the random demand modeled by a binomial $D_t \sim B(0.5, M)$ and $f(D_t, x_t, a_t) = 4 \min(D_t, x_t + a_t)$ is the sales profit, $C_m(x_t) = 1x_t$ is the maintenance cost, and $C_c(a_t) =$ $3 + 2a_t$ is the order cost. We considered a horizon H = 10with $s_0 = 0$. Optimal policies in the Inventory Management MDP can be expressed in the form (s_t, S_t) : at time t, if the stock is less than s_t , then agent should buy so that they have a stock of exactly S_t . More precisely: if $x_t \leq s_t$, then $a_t^* = S_t - x_t$, else $a_t^* = 0$.

Another environment, Cliff, can be found in Appendix H.

Optimality Front For our method, *Optimality Front*, we executed DOLFIN only once, with accuracy $\varepsilon = 10^{-3}$ on the chosen environment. We then computed all the metrics following Equation (11) by applying the functional of choice on all returned optimal distributions and selecting the optimal value. The Optimality Front contained 20 different optimal policies.

5.1. Threshold Probability

We compare the *Optimality Front* against several methods to compute a policy: *Proxy Optimization* is the value obtained by optimizing Equation (7). The *Risk neutral optimal* policy (i.e. the policy optimizing $\mathbb{E}[R^{\pi}]$) serves as a baseline. *Nested Probability Threshold* corresponds to the metric of the policy obtained following the Nested Risk Measure method (Bäuerle & Glauner, 2022). We also add, for reference, the optimal value obtained using augmented MDPs dynamic programming, resulting in non-markovian policies.

The value of the thresholds T are selected as ratios of the optimal mean μ^* . We observe in Table 1 that using EntRM

Table 1. Evaluation of $P(R^{\pi} \leq T)$.

T/μ^*	0.25	0.33	0.5
Optimality Front	$1.26\mathrm{e}^{-5}$	$8.40\mathrm{e}^{-5}$	$3.26\mathrm{e}^{-3}$
Proxy Optimization	$2.33e^{-5}$	$1.18e^{-4}$	$3.28e^{-3}$
Risk neutral optimal	$1.11e^{-4}$	$4.24e^{-4}$	$5.73e^{-3}$
Nested Prob. Thresh.	$1.54e^{-3}$	$8.37e^{-3}$	1
Optimal value	$6.29e^{-7}$	$8.22e^{-6}$	$7.85e^{-4}$

optimal policies outperforms the risk-neutral optimal policy by up to a factor 10 for a threshold ratio of 0.25. On the opposite, the Nested Probability Threshold fails to find a good policy and performs worse than the risk-neutral one. The experiment also highlights the benefit of choosing the best policy from the Optimality Front instead of simply optimizing on the proxy. Not only is the optimization problem simpler, but the performances are also better.

The true optimal value here is significantly better than what any markovian policy can achieve, especially for low thresholds. This gap is intensified by the nature of the MDP: in Inventory Management, the agent receives some reward with high variance at every timestep. For a specific state and timestep, the accumulated reward (and so, the distance to the threshold) can vary a lot, and so does the action. In some more goal-oriented MDPs, with very scarce reward, this gap can become insignificant (ex. see Cliff in Appendix H).

5.2. Value at Risk family

Hau et al. (2023) already showed that optimizing over the EVaR performed better than previous methods when it

comes to optimizing both VaR and CVaR, including using augmented MDPs. We compare our method to theirs (here, *Proxy Optimization*) and two other baselines for better illustration: the *risk-neutral optimal* policy and the *Nested Risk Measure* method. We compute both the VaR and CVaR with two different risk levels : 0.05 and 0.1.

	Table 2.	Evaluation	of (C)VaR	$[R^{\pi}]$	l
--	----------	------------	-------	------	-------------	---

Risk Measure	VaR			CVaR		
Risk parameter α	0.05	0.1	0.05	0.1		
Optimality Front	37	40	32.54	35.56		
Proxy Optimization	36	40	32.28	35.46		
Risk neutral optimal	36	40	31.45	34.68		
Nested Risk Measure	35	38	30.00	33.80		

We observe in Table 2 that for VaR and CVaR also, our algorithm outperforms pre-existing methods.

5.3. Optimality Front

We first give a simple illustration of the optimality front on the Cliff environment. The agent starts in the blue cell and is encouraged to reach the green cell as fast as possible, while absolutely avoiding the cliff depicted in red. She may choose her moves, but with probability 0.3 each choice is replaced by a random direction. Her risk tolerance is easily interpreted as her acceptance to walk close to the edge. Figure 1 shows the variety of optimal policies as the risk parameter β varies from -10 (very risk averse) to +10 (risk-prone). Non-violet arrows show the action that is preferred as soon as β is larger than a breakpoint whose value is given by the arrow's color.




Figure 1. Illustration of all cliffs β -optimal policies. Arrows are of the color of the smallest β value corresponding to the optimal action $[\pi_{\beta}^*]$.

We now illustrate the computational efficiency of our algorithms. DOLFIN computes intervals on which the current policy stays optimal, in order to avoid computing unnecessary values. But as the values get close to the breakpoint, using a regular grid becomes necessary as the computed values become irrelevant. We show however in Figure 2 that the number of evaluations grows sublinearly with the increasing accuracy, highlighting the gain of using such method.




Figure 2. Number of evaluation of EntRM required to obtain an accuracy of ε on the breakpoints using our method (blue) com a naive grid (red). The efficiency ratio goes up to 15 in this example.

In practice, this algorithm already performs better than the grid-based methods when comparing the number of evaluations. Yet, this algorithm can also be used as a one-shot preprocessing step to consider several risk-sensitive problems (such as different quantile levels), each of which can then be solved efficiently. For instance, DOLFIN was only executed once for the evaluation of all previous metrics, while all other methods have to be executed separately for each value.

6. Conclusion

We propose a unified framework for optimizing risksensitive objectives in Markov Decision Processes. Leveraging the computational advantages of the Entropic Risk Measure (EntRM), we provide an efficient algorithm for computing the optimality front, a family of policies which are optimal on a range of risk tolerance values. This also allows us to approximate key metrics such as Threshold Probabilities, Values at Risk and Conditional Values at Risk. Our algorithm demonstrates significant practical benefits in both efficiency and policy quality. This approach not only enhances risk-sensitive planning but also provides a versatile tool for tackling a variety of decision-making problems under uncertainty. It remains to be extended beyond planning in the context of learning unknown transition probabilities and reward distributions.

Acknowledgements

C. Vernade is funded by the Deutsche Forschungsgemeinschaft (DFG) under both the project 468806714 of the Emmy Noether Program and under Germany's Excellence Strategy – EXC number 2064/1 – Project number 390727645. CV also thanks the international Max Planck Research School for Intelligent Systems (IMPRS-IS). Aurélien Garivier and Alexandre Marthe thank the Chaire SeqALO (ANR-20-CHIA-0020-01) and PEPR IA project FOUNDRY (ANR-23-PEIA-0003).

Impact Statement

This paper presents work whose goal is to advance the field of Machine Learning. There are many potential societal consequences of our work, none which we feel must be specifically highlighted here.

References

- Achab, M. and Neu, G. Robustness and risk management via distributional dynamic programming. *arXiv preprint arXiv:2112.15430*, 2021.
- Ahmadi-Javid, A. Entropic Value-at-Risk: A New Coherent Risk Measure. *Journal of Optimization Theory and Applications*, 155(3):1105–1123, December 2012. ISSN 1573-2878. doi: 10.1007/s10957-011-9968-2.
- Akritas, A., Strzebonski, A., and Vigklas, P. Improving the performance of the continued fractions method using new bounds of positive roots. *Nonlinear Analysis: Modelling* and Control, 13(3):265–279, 2008.
- Artzner, P., Delbaen, F., Eber, J.-M., and Heath, D. Coherent measures of risk. *Mathematical finance*, 9(3):203–228, 1999.
- Atallah, M. J. Some dynamic computational geometry problems. *Computers & Mathematics with Applications*, 11 (12):1171–1181, 1985.
- Bäuerle, N. and Glauner, A. Markov decision processes with recursive risk measures. *European Journal of Operational Research*, 296(3):953–966, 2022.
- Bäuerle, N. and Ott, J. Markov decision processes with average-value-at-risk criteria. *Mathematical Methods of Operations Research*, 74:361–379, 2011.
- Bäuerle, N., Pitera, M., and Stettner, Ł. Blackwell optimality and policy stability for long-run risk-sensitive stochastic control. *SIAM Journal on Control and Optimization*, 62(6):3172–3194, 2024.
- Bellemare, M. G., Dabney, W., and Rowland, M. Distributional reinforcement learning. MIT Press, 2023.
- Bellman, R., Glicksberg, I., and Gross, O. On the optimal inventory equation. *Management Science*, 2(1):83–104, 1955.

- Boyd, S. P. and Vandenberghe, L. Convex optimization. Cambridge University Press, Cambridge, UK ; New York, 2004. ISBN 978-0-521-83378-3.
- Bäuerle, N. and Rieder, U. More Risk-Sensitive Markov Decision Processes. *Mathematics of Operations Research*, 39(1):105–120, February 2014. ISSN 0364-765X. doi: 10.1287/moor.2013.0601. Publisher: INFORMS.
- Cauchy, A. L. B. *Exercices de mathématiques*, volume 3. De Bure Frères, 1828.
- Charpentier, A., Elie, R., and Remlinger, C. Reinforcement learning in economics and finance. *Computational Economics*, pp. 1–38, 2021.
- Chow, Y. and Ghavamzadeh, M. Algorithms for CVaR Optimization in MDPs. In *Advances in Neural Information Processing Systems*, volume 27. Curran Associates, Inc., 2014.
- Chow, Y., Tamar, A., Mannor, S., and Pavone, M. Risksensitive and robust decision-making: a cvar optimization approach. *Advances in neural information processing systems*, 28, 2015.
- Chow, Y., Ghavamzadeh, M., Janson, L., and Pavone, M. Risk-constrained reinforcement learning with percentile risk criteria. *Journal of Machine Learning Research*, 18 (167):1–51, 2018.
- Collins, G. E. Polynomial minimum root separation. *Journal of Symbolic Computation*, 32(5):467–473, 2001.
- Durrett, R. *Probability: theory and examples*, volume 49. Cambridge university press, 2019.
- Föllmer, H. and Schied, A. *Stochastic finance: an introduction in discrete time.* Walter de Gruyter, 2011.
- Hau, J. L., Petrik, M., and Ghavamzadeh, M. Entropic risk optimization in discounted mdps. In *International Conference on Artificial Intelligence and Statistics*, pp. 47–76. PMLR, 2023.
- Hau, J. L., Delage, E., Ghavamzadeh, M., and Petrik, M. On dynamic programming decompositions of static risk measures in markov decision processes. *Advances in Neural Information Processing Systems*, 36, 2024.
- Howard, R. A. and Matheson, J. E. Risk-sensitive markov decision processes. *Management science*, 18(7):356–369, 1972.
- Kira, A., Ueno, T., and Fujita, T. Threshold probability of non-terminal type in finite horizon markov decision processes. *Journal of Mathematical Analysis and Applications*, 386(1):461–472, 2012.

- Kupper, M. and Schachermayer, W. Representation results for law invariant time consistent functions. *Mathematics and Financial Economics*, 2(3):189–210, September 2009. ISSN 1862-9660. doi: 10.1007/ s11579-009-0019-9.
- Marthe, A., Garivier, A., and Vernade, C. Beyond average return in Markov Decision Processes. Advances in Neural Information Processing Systems, 36, 2024.
- Massart, P. Concentration inequalities and model selection: Ecole d'Eté de Probabilités de Saint-Flour XXXIII-2003. Springer, 2007.
- Ni, X. and Lai, L. Evar optimization for risk-sensitive reinforcement learning. *IEEE Transactions on Information Theory*, 2022.
- Polydoros, A. S. and Nalpantidis, L. Survey of modelbased reinforcement learning: Applications on robotics. *Journal of Intelligent & Robotic Systems*, 86(2):153–173, 2017.
- Puterman, M. L. Markov decision processes: discrete stochastic dynamic programming. John Wiley & Sons, 2014.
- Rockafellar, R. T., Uryasev, S., et al. Optimization of conditional value-at-risk. *Journal of risk*, 2:21–42, 2000.
- Rowland, M., Dadashi, R., Kumar, S., Munos, R., Bellemare, M. G., and Dabney, W. Statistics and samples in distributional reinforcement learning. In *International Conference on Machine Learning*, pp. 5528–5536. PMLR, 2019.
- Scarf, H., Arrow, K., Karlin, S., and Suppes, P. The optimality of (s, s) policies in the dynamic inventory problem. *Optimal pricing, inflation, and the cost of price adjustment*, pp. 49–56, 1960.
- Silver, D., Schrittwieser, J., Simonyan, K., Antonoglou, I., Huang, A., Guez, A., Hubert, T., Baker, L., Lai, M., Bolton, A., et al. Mastering the game of go without human knowledge. *nature*, 550(7676):354–359, 2017.
- Su, X., Petrik, M., and Grand-Clément, J. Evar optimization in mdps with total reward criterion. In Seventeenth European Workshop on Reinforcement Learning, 2024.
- Sutton, R. S. and Barto, A. G. *Reinforcement learning: An introduction*. MIT press, 2018.
- Tamar, A., Chow, Y., Ghavamzadeh, M., and Mannor, S. Policy gradient for coherent risk measures. *Advances in neural information processing systems*, 28, 2015.
- Tossavainen, T. The lost cousin of the fundamental theorem of algebra. *Mathematics Magazine*, 80(4):290–294, 2007.

- Vershynin, R. High-dimensional probability: An introduction with applications in data science, volume 47. Cambridge university press, 2018.
- White, D. J. Minimizing a threshold probability in discounted markov decision processes. *Journal of mathematical analysis and applications*, 173(2):634–646, 1993.
- Wu, C. and Lin, Y. Minimizing risk models in markov decision processes with policies depending on target values. *Journal of mathematical analysis and applications*, 231 (1):47–67, 1999.

A. EntRM supplements

In this section, we give more insights about the Entropic Risk Measure. We first recall the definition of the EntRM.

Let X be a random variable. The Entropic Risk $\operatorname{EntRM}_{\beta}[X]$ of parameter $\beta \in \mathbb{R}$ is defined as

EntRM_{$$\beta$$}[X] =

$$\begin{cases}
\frac{1}{\beta} \log(\mathbb{E}[e^{\beta X}]), & \text{if } \beta \neq 0, \\
\mathbb{E}[X], & \text{if } \beta = 0.
\end{cases}$$

Risk parameter interpretation. The parameter β of EntRM measures risk tolerance. Large (positive) values of β encourage a riskier behavior, as only the largest values taken by X contribute significantly to the expectation. Conversely, negative values of β promote a conservative behavior that minimizes potential losses, irrespective of the maximum reward. This behavior is illustrated by the values of the EntRM when the risk parameter approaches $\pm \infty$.

If the support of X is (R_{\min}, R_{\max}) , then

$$\lim_{\beta \to -\infty} \mathrm{EntRM}_{\beta}[X] = R_{\min} \text{ and } \lim_{\beta \to +\infty} \mathrm{EntRM}_{\beta}[X] = R_{\max}$$

When β approaches zero, the EntRM converges to the expected value, leading to risk-neutral behavior:

EntRM_{$$\beta$$}(X) $\underset{\beta \to 0}{=} \mathbb{E}[X] + \frac{\beta}{2} \mathbb{V}[X] + o(\beta).$

The risk neutral behavior obtained for $\beta = 0$ is just the expectation, and the variance appears as the sensitivity of EntRM. For a Gaussian random variable $X \sim \mathcal{N}(\mu, \sigma^2)$, it holds exactly that $\text{EntRM}_{\beta}(X) = \mu + \frac{\beta}{2}\sigma^2$.

Dynamic programming. Q-value function in the case of Entropic Risk Measure is defined as follows:

$$Q_{h,\beta}^{\pi}(x,a) = \frac{1}{\beta} \log \left(\mathbb{E} \left[\exp \left(\beta R_h^{\pi}(x,a) \right) \right] \right).$$
(12)

As shown in Howard & Matheson (1972), this Q-value also satisfies a Bellman equation and thus it can be optimized efficiently by value iteration, and can result in a deterministic optimal policy.

The Q-value function for the entropic risk of parameter β satisfies the following optimal Bellman equations:

$$Q_{h,\beta}^{\pi}(x,a) = \operatorname{EntRM}_{\beta}[r_h(x,a)] + \operatorname{EntRM}_{\beta}\left[Q_{h+1,\beta}^{\pi}(X',\pi(X'))\right]$$
(13)

$$Q_{h,\beta}^*(x,a) = \operatorname{Ent} \operatorname{RM}_{\beta}[r_h(x,a)] + \operatorname{Ent} \operatorname{RM}_{\beta}\left[\max_{a'} Q_{h+1,\beta}^*(X',a')\right]$$
(14)

Where $X' \sim p(\cdot | x, a)$.

Principle of Optimality. Another implication of the Bellman equation is that the Entropic Risk Measure verifies the principle of Optimality, meaning there exists an optimal policy that is optimal for any subproblem:

$$\exists \pi^*, \, \forall x, h, \quad \pi^*_{h:H} \in \operatorname{argmax} \operatorname{Ent} \operatorname{RM}_\beta[R^\pi(x)] \tag{15}$$

with $\pi_{h:H} = (\pi_h, \dots, \pi_H)$. This is in general not true for other risk measures, where a policy may only be optimal for a specific timestep and state, and can be suboptimal at intermediate timestep.

Exponential form While the EntRM is often defined as in Equation (1), the $\frac{1}{\beta}$ ln serves solely as a renormalization factor. It does not influence the preference ordering between returns, and thus has no impact on the optimal policy in an MDP. For this reason, we also consider the exponential form $\operatorname{sign}(\beta)\mathbb{E}_{\pi}[\exp(\beta X)]$ (also called *exponential utility*), which can appear in some applications, like for the Threshold Probability objective. If a policy π is optimal for EntRM_{β}, it is also optimal for E_{β} , and vice versa. In general:

$$\operatorname{EntRM}_{\beta}(X_1) > \operatorname{EntRM}_{\beta}(X_2) \iff \operatorname{sign}(\beta) \cdot \mathbb{E}[\exp(\beta X_1)] > \operatorname{sign}(\beta) \cdot \mathbb{E}[\exp(\beta X_2)]$$

and thus,

$$\operatorname*{argmax}_{\pi} \operatorname{EntRM}_{\beta}[R^{\pi}] = \operatorname*{argmax}_{\pi} \operatorname{sign} \mathbb{E}_{\pi}[\exp(\beta R^{\pi})].$$

This equivalence is used at several occasions below.

B. More on the Value at Risk family of risk measures

In this section, we provide additional details on the *Value at Risk* (VaR), *Conditional Value at Risk* (CVaR), and *Entropic Value at Risk* (EVaR) measures discussed in the main text. Three main points are covered: (1) a clarification of conventions and notation, (2) key properties of VaR/CVaR, and (3) the derivation and theoretical advantages of EVaR.

B.1. Conventions and Notation

In the finance literature, VaR is generally introduced as the $(1 - \alpha)$ -quantile of a *loss* distribution, referring to a worst-case tail. It can also be defined via an α -quantile for *gains* (or returns), leading to slightly different formulas or sign conventions. For instance, one would usually write

$$\Pr(X \ge \operatorname{VaR}_{1-\alpha}[X]) = \alpha$$
. while others use $\Pr(X \le \operatorname{VaR}_{\alpha}[X]) = \alpha$,

Both definitions capture the idea of a critical quantile but from opposite sides of the distribution (gains vs. losses).

Throughout this work, we consistently adopt the low-tail perspective, defining

$$\operatorname{VaR}_{\alpha}[X] = \inf\{ x \mid \Pr(X \le x) \ge \alpha \},\$$

so that VaR at level α is simply the α -quantile of X from below. We focus on this definition to remain consistent with our usage of "negative outcomes" or "low returns" as the main source of risk. Should the convention change, one can substitute $X \leftarrow -X$ to convert between definitions without altering the mathematical properties.

To follow the change of convention, we also define the Conditional Value at Risk (CVaR) as

$$\operatorname{CVaR}_{\alpha}[X] = \frac{1}{\alpha} \int_{0}^{\alpha} \operatorname{VaR}_{\gamma}[X] \, d\gamma. \quad \text{instead of} \quad \operatorname{CVaR}_{1-\alpha}[X] = \frac{1}{\alpha} \int_{0}^{\alpha} \operatorname{VaR}_{1-\gamma}[X] \, d\gamma.$$

and the Entropic Value at Risk (EVaR) as

$$\operatorname{EVaR}_{\alpha}[X] = \sup_{\beta < 0} \left\{ \frac{1}{\beta} \ln \mathbb{E}[e^{\beta X}] - \frac{1}{\beta} \ln(\alpha) \right\} \quad \text{instead of} \quad \inf_{\beta > 0} \left\{ \frac{1}{\beta} \ln \mathbb{E}[e^{\beta X}] - \frac{1}{\beta} \ln(1 - \alpha) \right\}.$$

B.2. Entropic Value at Risk (EVaR)

Entropic Value at Risk (EVaR) was proposed as a tighter exponential-based bound on VaR and CVaR (Ahmadi-Javid, 2012). Its derivation stems from the fact that exponential moment bounding can yield a close approximation to quantile-based measures.

Proof. By Chernoff's inequality, for any $\beta < 0$ we have:

$$\Pr(X \le \ell) \le \mathbb{E}[e^{\beta X}] \exp(-\beta \ell).$$

Solving the equation $\mathbb{E}[e^{\beta X}] \exp(-\beta \ell) = \alpha$ for ℓ , we obtain

$$\ell = a_X(\beta, \alpha) = \frac{1}{\beta} \ln(\mathbb{E}[e^{\beta X}]) - \frac{1}{\beta} \ln(\alpha).$$

Hence, for any $\beta < 0$ and $\alpha \in (0, 1]$, the inequality

$$\Pr(X \le a_X(\beta, \alpha)) \le \alpha$$

implies

$$a_X(\beta, \alpha) \leq \operatorname{VaR}_{\alpha}(X).$$

Hence,

$$\operatorname{EVaR}_{\alpha}(X) = \sup_{\beta < 0} a_X(\beta, \alpha) \leq \operatorname{VaR}_{\alpha}(X).$$

C. Details on Section 3

C.1. Derivation of Proxy Problems

We here detail how the proxy problems are derived from the original problem.

Probability Threshold Problem. The initial problem is

$$\min_{\pi} \Pr(R^{\pi} \leq T),$$

Using the Chernoff bound, and a few manipulations.

$$\min_{\pi} \Pr(R^{\pi} \le T) \le \min_{\pi} \min_{\beta < 0} \mathbb{E}\left[e^{\beta(R^{\pi} - T)}\right]$$
$$= \min_{\beta < 0} \min_{\pi} \mathbb{E}\left[e^{\beta(R^{\pi} - T)}\right]$$
$$= \min_{\beta < 0} \mathbb{E}\left[e^{\beta\left(R^{\pi^{*}_{\beta}} - T\right)}\right].$$

Where the inversion between the minimum on the policy and the minimum on β is justified by the fact that there is only a finite number of policies.

Using results from Section 4, we can further simplify the problem. Consider the EntRM optimal policies π_1, \ldots, π_K and the corresponding optimality intervals I_1, \ldots, I_K . We have

$$\min_{\pi} \Pr(R^{\pi} \le T) \le \dots$$
$$= \min_{\beta < 0} \mathbb{E} \left[e^{\beta(R^{\pi^*_{\beta}} - T)} \right]$$
$$= \min_{k} \min_{\beta \in I_k} \mathbb{E} \left[e^{\beta(R^{\pi_k} - T)} \right].$$

Value at Risk Problem. The initial problem is

$$\max_{\pi} (\mathbf{C}) \operatorname{VaR}_{\alpha} [R^{\pi}].$$

The similar manipulations lead to

$$\max_{\pi} (C) \operatorname{VaR}_{\alpha} [R^{\pi}] \geq \max_{\pi} \operatorname{EVaR}_{\alpha} [R^{\pi}]$$
$$= \max_{\pi} \sup_{\beta < 0} \operatorname{EntRM}_{\beta} [R^{\pi}] + \frac{1}{\beta} \ln(\alpha)$$
$$= \sup_{\beta < 0} \max_{\pi} \operatorname{EntRM}_{\beta} [R^{\pi}] + \frac{1}{\beta} \ln(\alpha)$$
$$= \sup_{\beta < 0} \operatorname{EntRM}_{\beta} [R^{\pi^{*}}_{\beta}] + \frac{1}{\beta} \ln(\alpha)$$
$$= \max_{k} \sup_{\beta \in I_{k}} \operatorname{EntRM}_{\beta} [R^{\pi_{k}}] + \frac{1}{\beta} \ln(\alpha).$$

C.2. Proof of Proposition 3.2

We write $M_R(\beta) = \mathbb{E}[\exp(\beta R)]$ the moment generating function of the random variable R. Lemma C.1. Assume that R is bounded between -H and H, then for a fixed stepsize $\epsilon > 0$, for $k \in \mathbb{N}$ and $\beta \in [\epsilon k, \epsilon(k+1)]$, noting $b_k = M_R(-\epsilon k)$, we have

$$M_R(-\beta) \ge \exp(-\epsilon H)b_k$$
 and $M_R(-\beta) \ge \exp(\epsilon H)b_{k+1}$,

and,

$$\frac{M_R(-\beta)}{\min\{b_k, b_{k+1}\}} \ge \exp\left(\epsilon \frac{-H}{2}\right) \left(\frac{b_k}{b_{k+1}}\right)^{\frac{1}{2H}} \ge \exp\left(\epsilon \frac{-H}{2}\right),$$

otherwise the ratio is greater or equal to 1.

Proof. First we have

$$M_{R}(-\beta) = \mathbb{E}[\exp(-(\beta - \epsilon k)R)\exp(-\epsilon kR)]$$

$$\geq \exp(-(\beta - \epsilon k)H)\mathbb{E}[\exp(-\epsilon kR)] \geq \exp(-\epsilon H)b_{k},$$

$$M_{R}(-\beta) = \mathbb{E}[\exp((-\beta + \epsilon(k+1))R)\exp(-\epsilon(k+1)R)]$$

$$\geq \exp(-(\epsilon(k+1) - \beta)H)\mathbb{E}[\exp(-\epsilon(k+1)R)] \geq \exp(-\epsilon H)b_{k+1}.$$

To obtain the following inequality it is enough to consider the intersection of the functions $t \mapsto \exp(-tH)b_k$ and $t \mapsto \exp((-\epsilon - t)H)b_{k+1}$ on $t \in [0, \epsilon]$: as $M_R(-\beta)$ follows both left decreasing and right increasing constraint, the minimal value possible is on the intersection.

We can now give a proof of Proposition 3.2.

Proof. Let π_c and β_c be the policy and the β optimizing the Chernoff bound *B*. First, we can consider that $\beta_c < \ln(1/p)/a$. Indeed if not we can choose $\beta_c = 0$ instead:

$$\begin{split} M_R(\beta_c) &= \mathbb{E}[\exp(-\beta R)\mathbb{1}\{R \le a\}] + \mathbb{E}[\exp(-\beta R)\mathbb{1}\{R > a\}] \\ &\geq \exp(-\beta_c a)\mathbb{P}[R \le a] + \exp(-\beta M)\mathbb{P}[R \ge a] \\ &\geq \exp(-\beta_c a)p \ge 1 = M_R(0) \quad \text{using the hypothesis on } \beta_c \;. \end{split}$$

Let then $k \in \mathbb{N}$ be such that $\epsilon k \leq \beta_c < \epsilon(k+1)$, and suppose $M_{R^{\pi_c}}(-\epsilon k) \leq M_{R^{\pi_c}}(-\epsilon(k+1))$ without loss of generality. We have

$$\tilde{B} \le \min_{\pi} M_{R^{\pi}}(-\epsilon k) \le M_{R^{\pi_c}}(-\epsilon k) \le \exp\left(\epsilon \frac{H}{2}\right) M_{R^{\pi_c}}(\beta_c) \le \exp\left(\epsilon \frac{H}{2}\right) B,$$

using Lemma C.1.

For $\epsilon \leftarrow 2\log(1\varepsilon)/H$, we have that $\tilde{B} \leq (1+\varepsilon)B$.

D. Proof of Section 4

D.1. Proof of Proposition 4.2.

Proof of Proposition 4.2. By Theorem D.5, there exist values β_{\min} and β_{\max} such that for all $\beta < \beta_{\min}$ (respectively, $\beta > \beta_{\max}$), a single action remains optimal. Consequently, any "breakpoint" (a value of β at which the optimal action changes) must lie in the finite interval $[\beta_{\min}, \beta_{\max}]$.

Let a_i and a_j be two different actions. Suppose at some β , both a_i and a_j are optimal for EntRM_{β}($R(\cdot)$). That implies:

$$\sum_{i} \mu_i e^{\beta x_i} = \sum_{j} \mu_j e^{\beta x_j},$$

where $\{\mu_i, x_i\}$ and $\{\mu_j, x_j\}$ refer to the reward distributions (or points in the returns) associated with actions a_i and a_j . The map $\beta \mapsto \sum_k \mu_k e^{\beta x_k}$ is holomorphic (analytic) as it is a finite sum of exponentials. Two distinct holomorphic functions on a finite interval can only coincide at a finite number of points, unless they coincide everywhere.

Therefore, there are only finitely many β values where

$$\operatorname{EntRM}_{\beta}(R(a_i)) = \operatorname{EntRM}_{\beta}(R(a_j)).$$

Because the optimal policy can only change when two action-value curves intersect (thus altering the argmax set), a finite number of pairwise intersections implies a finite number of breakpoints in $[\beta_{\min}, \beta_{\max}]$.

Assume an action a is the unique optimum at some β_0 . Formally,

$$\operatorname{EntRM}_{\beta_0}(R(a)) > \operatorname{EntRM}_{\beta_0}(R(a'))$$
 for all $a' \neq a$.

By continuity of $\beta \mapsto \text{EntRM}(R(a))$, there exists $\varepsilon > 0$ such that for all $\beta' \in [\beta_0 - \varepsilon, \beta_0 + \varepsilon]$, the same strict inequality holds. Hence, *a* remains the unique optimal action throughout this interval around β_0 .

D.2. Proof of Theorem 4.3.

We first start by deriving bounds on the growth of the EntRM when the risk parameter is changed slightly.

Proposition D.1. Let X be a random variable, $\beta \in \mathbb{R}$, and $0 < \varepsilon < |\beta|$. Then:

If
$$\beta > 0$$
:

$$\frac{\beta}{\beta+\varepsilon}\operatorname{EntRM}_{\beta}[X] + \frac{\varepsilon}{\beta+\varepsilon}r_{\min} \le \operatorname{EntRM}_{\beta+\varepsilon}[X] \le \frac{\beta}{\beta+\varepsilon}\operatorname{EntRM}_{\beta}[X] + \frac{\varepsilon}{\beta+\varepsilon}r_{\max}, \tag{16}$$

$$\frac{\beta}{\beta - \varepsilon} \operatorname{EntRM}_{\beta}[X] - \frac{\varepsilon}{\beta - \varepsilon} r_{\max} \le \operatorname{EntRM}_{\beta - \varepsilon}[X] \le \frac{\beta}{\beta - \varepsilon} \operatorname{EntRM}_{\beta}[X] - \frac{\varepsilon}{\beta - \varepsilon} r_{\min}.$$
(17)

If $\beta < 0$:

$$\frac{\beta}{\beta + \varepsilon} \operatorname{EntRM}_{\beta}[X] + \frac{\varepsilon}{\beta + \varepsilon} r_{\max} \le \operatorname{EntRM}_{\beta + \varepsilon}[X] \le \frac{\beta}{\beta + \varepsilon} \operatorname{EntRM}_{\beta}[X] + \frac{\varepsilon}{\beta + \varepsilon} r_{\min},$$
(18)

$$\frac{\beta}{\beta - \varepsilon} \operatorname{EntRM}_{\beta}[X] - \frac{\varepsilon}{\beta - \varepsilon} r_{\min} \le \operatorname{EntRM}_{\beta - \varepsilon}[X] \le \frac{\beta}{\beta - \varepsilon} \operatorname{EntRM}_{\beta}[X] - \frac{\varepsilon}{\beta - \varepsilon} r_{\max}.$$
(19)

Proof. In the following, $0 < \varepsilon < |\beta|$. We write $X = \sum \mu_i \delta_{x_i}$, with $r_{\min} = \min_i x_i$ and $r_{\max} = \max_i x_i$. For $\beta > 0$:

$$\operatorname{Ent}\operatorname{RM}_{\beta+\varepsilon}[X] = \frac{1}{\beta+\varepsilon} \ln\left(\sum \mu_i e^{(\beta+\varepsilon)x_i}\right)$$
$$= \frac{1}{\beta+\varepsilon} \ln\left(\sum \mu_i e^{\beta x_i} e^{\varepsilon x_i}\right)$$
$$\leq \frac{1}{\beta+\varepsilon} \ln\left(e^{\varepsilon r_{\max}} \sum \mu_i e^{\beta x_i}\right) \quad (\text{since } \frac{1}{\beta+\varepsilon} > 0)$$
$$= \frac{1}{\beta+\varepsilon} \ln\left(\sum \mu_i e^{\beta x_i}\right) + \frac{\varepsilon}{\beta+\varepsilon} r_{\max}$$
$$= \frac{\beta}{\beta+\varepsilon} \operatorname{Ent}\operatorname{RM}_{\beta}[X] + \frac{\varepsilon}{\beta+\varepsilon} r_{\max},$$

and

$$\operatorname{EntRM}_{\beta-\varepsilon}[X] = \frac{1}{\beta-\varepsilon} \ln\left(\sum \mu_i e^{(\beta-\varepsilon)x_i}\right)$$
$$= \frac{1}{\beta-\varepsilon} \ln\left(\sum \mu_i e^{\beta x_i} e^{-\varepsilon x_i}\right)$$
$$\geq \frac{1}{\beta-\varepsilon} \ln\left(e^{-\varepsilon r_{\max}} \sum \mu_i e^{\beta x_i}\right) \quad (\text{since } \frac{1}{\beta-\varepsilon} > 0)$$
$$= \frac{1}{\beta-\varepsilon} \ln\left(\sum \mu_i e^{\beta x_i}\right) - \frac{\varepsilon}{\beta-\varepsilon} r_{\max}$$
$$= \frac{\beta}{\beta-\varepsilon} \operatorname{EntRM}_{\beta}[X] - \frac{\varepsilon}{\beta-\varepsilon} r_{\max}.$$

For $\beta < 0$:

$$\operatorname{EntRM}_{\beta+\varepsilon}[X] = \frac{1}{\beta+\varepsilon} \ln\left(\sum \mu_i e^{(\beta+\varepsilon)x_i}\right)$$
$$= \frac{1}{\beta+\varepsilon} \ln\left(\sum \mu_i e^{\beta x_i} e^{\varepsilon x_i}\right)$$
$$\leq \frac{1}{\beta+\varepsilon} \ln\left(e^{\varepsilon r_{\min}} \sum \mu_i e^{\beta x_i}\right) \quad (\text{since } \frac{1}{\beta+\varepsilon} < 0)$$
$$= \frac{1}{\beta+\varepsilon} \ln\left(\sum \mu_i e^{\beta x_i}\right) + \frac{\varepsilon}{\beta+\varepsilon} r_{\min}$$
$$= \frac{\beta}{\beta+\varepsilon} \operatorname{EntRM}_{\beta}[X] + \frac{\varepsilon}{\beta+\varepsilon} r_{\min},$$

and

$$\operatorname{EntRM}_{\beta-\varepsilon}[X] = \frac{1}{\beta-\varepsilon} \ln\left(\sum \mu_i e^{(\beta-\varepsilon)x_i}\right)$$
$$= \frac{1}{\beta-\varepsilon} \ln\left(\sum \mu_i e^{\beta x_i} e^{-\varepsilon x_i}\right)$$
$$\geq \frac{1}{\beta-\varepsilon} \ln\left(e^{-\varepsilon r_{\min}} \sum \mu_i e^{\beta x_i}\right) \quad (\text{since } \frac{1}{\beta-\varepsilon} < 0)$$
$$= \frac{1}{\beta-\varepsilon} \ln\left(\sum \mu_i e^{\beta x_i}\right) - \frac{\varepsilon}{\beta-\varepsilon} r_{\min}$$
$$= \frac{\beta}{\beta-\varepsilon} \operatorname{EntRM}_{\beta}[X] - \frac{\varepsilon}{\beta-\varepsilon} r_{\min}.$$

The other sides of the inequalities are obtained in a similar manner.

We then consider the easier case of a single state MDP, which is what will be used in practice for the algorithm. There is n actions a_1, \ldots, a_n with reward distribution $R(a_1), \ldots, R(a_n)$.

Theorem D.2 (Interval of Action Optimality). Let r_{\min} (resp. r_{\max}) be the minimum (resp. maximum) achievable reward, and $\Delta R = r_{\max} - r_{\min}$. Suppose we have actions $(a_{(i)})_i$ ordered so that

$$\operatorname{EntRM}_{\beta}[R(a_{(1)})] > \operatorname{EntRM}_{\beta}[R(a_{(2)})] \geq \ldots \geq \operatorname{EntRM}_{\beta}[R(a_{(n)})].$$

In particular, $a_{(1)}$ is the unique optimal action and $a_{(2)}$ is the second-best. Define

$$\Delta U = \operatorname{EntRM}_{\beta} \left[R(a_{(1)}) \right] - \operatorname{EntRM}_{\beta} \left[R(a_{(2)}) \right].$$

Then:

• If
$$\beta \neq 0$$
, for all $\beta' \in \left[\beta\left(1 - \frac{\Delta U}{\Delta R}\right), \ \beta\left(1 + \frac{\Delta U}{\Delta R}\right)\right]$ and for all $i \geq 2$, we have
 $\operatorname{EntRM}_{\beta'}\left[R(a_{(1)})\right] > \operatorname{EntRM}_{\beta'}\left[R(a_{(i)})\right]$.

• If
$$\beta = 0$$
, then for all $\beta' \in \left[-\frac{8 \Delta U}{\Delta R^2}, \frac{8 \Delta U}{\Delta R^2}\right]$ and all $i \ge 2$, we have
 $\operatorname{EntRM}_{\beta'}\left[R(a_{(1)})\right] > \operatorname{EntRM}_{\beta'}\left[R(a_{(i)})\right]$

Proof of Theorem D.2 for the case $\beta \neq 0$ *.* Assume $\beta \neq 0$ *.* Let

$$U_{\beta}^{1} = \operatorname{EntRM}_{\beta} [R(a_{(1)})], \quad U_{\beta}^{2} = \operatorname{EntRM}_{\beta} [R(a_{(2)})].$$

By hypothesis, $U_{\beta}^1 > U_{\beta}^2$. Define $\Delta U = U_{\beta}^1 - U_{\beta}^2$ and $\Delta R = r_{\text{max}} - r_{\text{min}}$. We aim to show that if β' remains within the specified range around β , action $a_{(1)}$ remains strictly optimal for $\text{EntRM}_{\beta'}$.

Case $\beta' > \beta$ and $\beta > 0$.

Write $\beta' = \beta + \varepsilon$ with $\varepsilon > 0$. Assume

$$\varepsilon < \beta \cdot \frac{\Delta U}{\Delta R}$$

From the previously established bounds (see Proposition D.1), we know

$$\frac{\beta}{\beta+\varepsilon}U_{\beta}^{1} + \frac{\varepsilon}{\beta+\varepsilon}r_{\min} \leq \operatorname{EntRM}_{\beta+\varepsilon}[R(a_{(1)})] = U_{\beta'}^{1},$$

and

$$\operatorname{EntRM}_{\beta+\varepsilon} \left[R(a_{(2)}) \right] = U_{\beta'}^2 \leq \frac{\beta}{\beta+\varepsilon} U_{\beta}^2 + \frac{\varepsilon}{\beta+\varepsilon} r_{\max}$$

Thus, showing

$$\frac{\beta}{\beta+\varepsilon}U_{\beta}^{1} + \frac{\varepsilon}{\beta+\varepsilon}r_{\min} > \frac{\beta}{\beta+\varepsilon}U_{\beta}^{2} + \frac{\varepsilon}{\beta+\varepsilon}r_{\max}$$

implies $U_{\beta'}^1 > U_{\beta'}^2$. Rewriting, we get

$$\frac{\beta}{\beta+\varepsilon} \left(U_{\beta}^{1} - U_{\beta}^{2} \right) > \frac{\varepsilon}{\beta+\varepsilon} \left(r_{\max} - r_{\min} \right),$$

i.e.

$$\beta \, \frac{\Delta U}{\Delta R} \, > \, \varepsilon.$$

But this is exactly our assumption on ε . Hence $a_{(1)}$ remains strictly better than $a_{(2)}$ at $\beta' = \beta + \varepsilon$, and by extension, better than all other actions.

Case
$$\beta' < \beta$$
 and $\beta > 0$.
Now let $\beta' = \beta - \varepsilon$ with $\varepsilon > 0$. We assume

$$\varepsilon < \beta \, \frac{\Delta U}{\Delta R}.$$

An analogous argument shows

$$rac{eta}{eta-arepsilon} \left(U_eta^1 - U_eta^2
ight) \ > \ rac{arepsilon}{eta-arepsilon} \left(r_{
m max} - r_{
m min}
ight).$$

This inequality, combined with the upper and lower bounds for $U^1_{\beta-\varepsilon}$ and $U^2_{\beta-\varepsilon}$, yields $U^1_{\beta'} > U^2_{\beta'}$. Consequently, $a_{(1)}$ remains the strictly optimal action at $\beta' = \beta - \varepsilon$.

The case $\beta < 0$ is similar, using the associated inequalities from Proposition D.1.

Hence, for all parameter shifts $\beta' \in [\beta(1 - \Delta U/\Delta R), \ \beta(1 + \Delta U/\Delta R)]$, the action $a_{(1)}$ remains strictly optimal. This completes the proof for $\beta \neq 0$.

The proof of the second part of Theorem D.2, when $\beta = 0$, uses *Hoeffding's lemma* (see e.g. Massart (2007)):

$$\forall \lambda \in \mathbb{R}, \quad \mathbb{E}[\exp(\lambda X)] \le \exp\left(\lambda \mathbb{E}[X] + \frac{\lambda^2 \Delta R^2}{8}\right)$$
 (20)

Proof. (second part of Theorem D.2) Hoeffding's lemma gives

$$\begin{cases} \operatorname{EntRM}_{\beta}[X] \leq \mathbb{E}[X] + \beta \frac{\Delta R^2}{8} & \text{if } \beta > 0 \text{, and} \\ \operatorname{EntRM}_{\beta}[X] \geq \mathbb{E}[X] - \beta \frac{\Delta R^2}{8} & \text{if } \beta < 0 \text{.} \end{cases}$$
(21)

We can then proceed similarly as the previous proof. Consider $0 < \beta < \frac{8\Delta\mu}{\Delta R^2}$. Using the monotonicity of $\beta \mapsto U_{\beta}$, $U_{\beta}^1 \ge E[R(a_1)]$ and

$$U_{\beta}^{1} - U_{\beta}^{2} \ge \mathbb{E}[R(a_{1})] - \mathbb{E}[R(a_{2})] - \beta \frac{\Delta R^{2}}{8} \ge \Delta \mu - \frac{8\Delta \mu}{\Delta R^{2}} \frac{\Delta R^{2}}{8} = 0 ,$$

and $U_{\beta}^1 > U_{\beta}^2$. The case $\beta < 0$ is similar.

Better bounds can also be derived, yet they are not symmetrical. These better bounds will be those used in practice in Algorithm 2.

Proposition D.3. Better bounds Let $\beta \in \mathbb{R} \setminus \{0\}$. Let $a_1, \ldots, a_n \in \mathcal{A}$ and $R(a_i)$ be the associated rewards. Without loss of generality, assume that $U_{\beta}(R(a_1)) > U_{\beta}(R(a_2)) \ge \cdots \ge U_{\beta}(R(a_n))$ (i.e., a_1 is the unique optimal action). Define $\Delta U = U_{\beta}(R(a_1)) - U_{\beta}(R(a_2))$.

If
$$\beta > 0$$
, for all $\beta' \in \left[\beta\left(1 - \frac{\Delta U}{r_{\max} - U_2}\right), \beta\left(1 + \frac{\Delta U}{r_{\max} - U_1}\right)\right]$, the optimal action for $U_{\beta'}$ is a_1 .
If $\beta < 0$, for all $\beta' \in \left[\beta\left(1 - \frac{\Delta U}{U_2 - r_{\min}}\right), \beta\left(1 + \frac{\Delta U}{U_1 - r_{\min}}\right)\right]$, the optimal action for $U_{\beta'}$ is a_1 .

We can verify that these intervals are better than those given by Theorem D.2 by noting that $r_{\min} \leq U_{\beta}(R) \leq r_{\max}$. Therefore, $\frac{\Delta U}{r_{\max}-U} > \frac{\Delta U}{\Delta R}$ and $\frac{\Delta U}{U-r_{\min}} > \frac{\Delta U}{\Delta R}$.

Proof. The proof is the same as in Proposition 4.2, but using the monotonicity bounds $U_{\beta}(\mu) \leq U_{\beta+\varepsilon}(\mu)$ and $U_{\beta}(\mu) \geq U_{\beta-\varepsilon}(\mu)$.

Finally, we can prove Theorem 4.3. We use the same principle as in the proof of Theorem D.2 and the principle of optimality (i.e. the optimal policy is always greedy with respect to itself) to show that the optimal policy remains the same. The inequalities from Proposition D.1 are adapted by replacing r_{max} by $h \cdot r_{\text{max}}$ and r_{min} by $h \cdot r_{\text{min}}$ (for timestep h, the return is in $[h \cdot r_{\text{min}}, h \cdot r_{\text{max}}]$).

Proof. (of Theorem 4.3) Let us address the case where $\beta > 0$. By the principle of optimality, we have:

$$\forall h, x, \quad \pi_{h,\beta}^*(x) = \arg\max_a \operatorname{EntRM}_{\beta}[R_h^{\pi^*}(x,a)] .$$

Let $\varepsilon < \Delta$ (with $\varepsilon < \beta$). We then obtain the inequalities:

$$\forall h, x, a, \quad \frac{\beta}{\beta + \varepsilon} \operatorname{EntRM}_{\beta}[R_{h}^{\pi^{*}}(x, a)] + \frac{\varepsilon}{\beta + \varepsilon} hr_{\min} \leq \operatorname{EntRM}_{\beta + \varepsilon}[R_{h}^{\pi^{*}}(x, a)] \leq \frac{\beta}{\beta + \varepsilon} \operatorname{EntRM}_{\beta}[R_{h}^{\pi^{*}}(x, a)] + \frac{\varepsilon}{\beta + \varepsilon} hr_{\max}$$

By the definition of ε , we get:

$$\forall h, x, \forall a \neq \pi_h^*(x), \quad \frac{\beta}{\beta + \varepsilon} \mathrm{EntRM}_\beta[R_h^{\pi^*}(x, a)] + \frac{\varepsilon}{\beta + \varepsilon} hr_{\max} \leq \frac{\beta}{\beta + \varepsilon} \mathrm{EntRM}_\beta[R_h^{\pi^*}(x, \pi_h^*(x))] + \frac{\varepsilon}{\beta + \varepsilon} hr_{\min}.$$

Therefore,

$$\operatorname{EntRM}_{\beta'}[R_{h}^{\pi^{*}}(x,a)] \leq \frac{\beta}{\beta+\varepsilon} \operatorname{EntRM}_{\beta}[R_{h}^{\pi^{*}}(x,a)] + \frac{\varepsilon}{\beta+\varepsilon} hr_{\max}$$
$$\leq \frac{\beta}{\beta+\varepsilon} \operatorname{EntRM}_{\beta}[R_{h}^{\pi^{*}}(x,\pi_{h}^{*}(x))] + \frac{\varepsilon}{\beta+\varepsilon} hr_{\min}$$
$$\leq \operatorname{EntRM}_{\beta'}[R_{h}^{\pi^{*}}(x,\pi_{h}^{*}(x))] .$$

Thus, π^* remains greedy with respect to itself and continues to be the optimal policy.

The cases $\beta = 0$ and $\beta < 0$ follow similarly, using the associated inequalities.

D.3. About Proposition 4.4.

We first start with a lemma.

Lemma D.4. Let Y be a random variable with continuous law. Let X_1, \ldots, X_n be random variables with Y independent from (X_1, \ldots, X_n) . Then,

$$\Pr(Y = f(X_1, \dots, X_n) \mid X_1, \dots, X_n) = 0$$

Proof. This is a special case of Exercise 2.1.5 in Durrett (2019). It comes down to writing the definition of conditional probabilities and computing the integrals with Fubini's theorem. \Box

Proof. (of Proposition 4.4)

Let π be a policy. Assume the probability transitions are fixed and only the rewards are random. We consider, for x, h, a, a' the set $\mathcal{B}_{a,a'}^{x,h}$ of parameters such that $\operatorname{EntRM}_{\beta}[R_h^{\pi}(x,a)] = \operatorname{EntRM}_{\beta}[R_h^{\pi}(x,a')]$. We aim to show that $(\mathcal{B}_{a,a'}^{x,h})_{x,h,a,a'}$ have no element in common pairwise, with probability one.

Consider orders on both \mathcal{X} and \mathcal{A} , and consider the associated lexigocraphic order of $[H, 1] \times \mathcal{X} \times \mathcal{A}$. We proceed by induction on this order.

Case t = H.

Let $x_1, x_2 \in \mathcal{X}$, $a_1, a'_1, a_2, a'_2 \in \mathcal{A}$, with $(x_1, a_1, a'_1) \neq (x_2, a_2, a'_2)$ (the triplets are different, but some elements of the triplets can be equal). Consider known $\mathcal{B}^{x_1,H}_{a_1,a'_1}$.

$$\forall \beta \in \mathbb{R}, \quad \Pr\left(\operatorname{EntRM}_{\beta}[R_{H}^{\pi}(x_{2}, a_{2})] = \operatorname{EntRM}_{\beta}[R_{H}^{\pi}(x_{2}, a_{2}')] \middle| r_{H}(x_{1}, a_{1}), r_{H}(x_{1}, a_{1}')\right)$$
$$= \Pr\left(r_{H}(x_{2}, a_{2}) = r_{H}(x_{2}, a_{2}') \middle| r_{H}(x_{1}, a_{1}), r_{H}(x_{1}, a_{1}')\right)$$
$$= 0.$$

Because $r_H(x_2, a_2)$ and $r_H(x_2, a'_2)$ are continuous random variables and at least one of the two is not conditioned on, the probability that they are equal is zero according to Lemma D.4.

It is in particular true for all $\beta \in \mathcal{B}_{a_1,a'_1}^{x_1,H}$. Using the union bound, we get that $\mathcal{B}_{a_1,a'_1}^{x_1,H}$ and $\mathcal{B}_{a_2,a'_2}^{x_2,H}$ have no element in common with probability one.

By considering elements in order and conditioning on the previously *observed* rewards, the induction is verified for t = H. Case t < H.

Let $u = h_0, x_0, a_0$. Consider all breakpoints observed before

$$\mathcal{B} = \bigcup_{\substack{(h,x,a) < u \\ (h,x,a') \le u}} \mathcal{B}^{x,h}_{a,a'}.$$

By induction, all of them are disjoint pairwise with probability one.

$$\begin{aligned} \forall \beta \in \mathcal{B}, \quad \Pr\left(\operatorname{EntRM}_{\beta}[R_{h_{0}}^{\pi}(x_{0}, a_{0})] = \operatorname{EntRM}_{\beta}[R_{h_{0}}^{\pi}(x_{0}, a'_{0})] \left| \left(r_{h}(x, a)\right)_{(h, x, a) \leq u} \right) \right. \\ &= \Pr\left(r_{h_{0}}(x_{0}, a_{0}) + \operatorname{EntRM}_{\beta}[R_{h_{0}+1}^{\pi}(X_{0})] = r_{h_{0}}(x_{0}, a'_{0}) + \operatorname{EntRM}_{\beta}[R_{h_{0}+1}^{\pi}(X'_{0})] \left| \left(r_{h}(x, a)\right)_{(h, x, a) \leq u} \right) \right. \\ &= P\left(r_{h_{0}}(x_{0}, a_{0}) = f\left(\left(r_{h}(x, a)\right)_{(h, x, a) \leq u}\right) \left| \left(r_{h}(x, a)\right)_{(h, x, a) \leq u} \right) \right. \\ &= 0. \end{aligned}$$

Where $f\left((r_h(x,a))_{(h,x,a)\leq u}\right) = r_{h_0}(x_0,a'_0) + \text{EntRM}_{\beta}[R^{\pi}_{h_0+1}(X'_0)] - \text{EntRM}_{\beta}[R^{\pi}_{h_0+1}(X_0)]$ and using Lemma D.4.

The induction is then proved.

As there is a finite number of policy, this result remains when considering the set of all policies.

To conclude, notice that a breakpoint β_0 is a value of risk parameter for which two different action a_1, a_2 have the same expected return for some state x and timestep t. Hence, $\beta_0 \in \mathcal{B}^{x,h}_{a_1,a_2}$. With probability one, those set are pairwise disjoint, meaning a single action changes in β_0 .

As this proof works for any value of the probability transitions, the result also remains for p random.

The result is what we observe for all the MDPs used in practice. Yet it is simple to construct an MDP where several actions change at the same time, it is enough to have two states x_1, x_2 have the same transitions function and rewards : $\forall a, x', p(x'|x_1, a) = p(x'|x_2, a)$ and $r(x_1, a) = r(x_2, a)$.

D.4. Proof of Proposition 4.6.

Proof. (of Proposition 4.6) It all relies on the continuity of the function $\beta \to \text{EntRM}_{\beta}[R^{\pi}]$.

$$\forall h, x, \quad \operatorname{EntRM}_{\beta_1}[R_h^{\pi^1}(x)] \ge \operatorname{EntRM}_{\beta_1}[R_h^{\pi^2}(x)]$$
$$\operatorname{EntRM}_{\beta_3}[R_h^{\pi^1}(x)] \le \operatorname{EntRM}_{\beta_3}[R_h^{\pi^2}(x)]$$

By continuity, there exists β_2 such that there is an equality, and, since we assume that there are no other optimal policy in $[\beta_1, \beta_3]$, the equality is verified for all x, h.

D.5. On the lowest breakpoint.

Theorem D.5. Consider $\pi_{\inf} = \lim_{\beta \to -\infty} \pi_{\beta}^*$. Consider the return R taking value in $x_1 \leq \ldots \leq x_n$. We write $a_i(\pi) = P(R^{\pi_{\inf}} = x_i) - \Pr(R^{\pi} = x_i)$. Then, the lowest breakpoint β_{\inf} verifies

$$\beta_{\inf} \geq \frac{-\log\left(1 + \max_{\pi \neq \pi_{\inf}} \max_{i>0} \frac{|a_i(\pi)|}{|a_0(\pi)|}\right)}{\min_{i \neq j} |x_i - x_j|} .$$

Proof. (of Theorem D.5)

We write $\Delta R = R_{\text{max}} - R_{\text{min}}$. We assume here for simplicity that all the rewards are even spaced on a grid. This mean that the values of the return are of the form $R_i = R_{\min} + i \frac{\Delta R}{n}$ for $i \in \{0, \dots, n\}$.

With this natural assumption, the problem of finding β such that,

$$\operatorname{EntRM}_{\beta}[R^{\pi}] = \operatorname{EntRM}_{\beta}[R^{\pi'}]$$

can be transformed into a problem of finding the roots of a polynomial.

$$\operatorname{Ent}\operatorname{RM}_{\beta}[R^{\pi}] = \operatorname{Ent}\operatorname{RM}_{\beta}[R^{\pi'}] \Longrightarrow \mathbb{E}[\exp(\beta R^{\pi})] = \mathbb{E}[\exp(\beta R^{\pi'})]$$
$$\iff \sum_{i=0}^{n} \mu_{i} \exp(\beta R_{i}) = \sum_{i=0}^{n} \mu_{i}' \exp(\beta R_{i}')$$
$$\iff \sum_{i=0}^{n} a_{i} \exp(\beta R_{i}) = 0$$
$$\iff \sum_{i=0}^{n} a_{i} \exp(\beta i \frac{\Delta R}{n}) = 0$$
$$\iff \sum_{i=0}^{n} a_{i} X^{-i} = 0.$$
$$\iff \sum_{i=0}^{n} a_{n-i} X^{i} = 0.$$

Where $X = \exp\left(-\beta \frac{\Delta R}{n}\right)$. The first implication is not an equivalence because of the case $\beta = 0$, where the exponential form (rhs) is always equal to 1 and thus the equality is always verified. The last implication is verified by multiplying by X^n because of that same fact that 0 is already a root of the equation.

Hence, if X is a solution, β verifies $\beta = -\frac{\log(X)}{\Delta R}$.

Cauchy's bound on the size of the largest polynomial root (Cauchy, 1828) claims that the largest root verifies

$$1 + \max_{i>0} \frac{|a_i|}{|a_0|}.$$

The lowest breakpoints corresponds to the largest breakpoint solution between the π_{inf} and any other policy. Hence, the lowest breakpoint verifies

$$\beta_{\inf} \geq \frac{-\log\left(1 + \max_{\pi \neq \pi_{\inf}} \max_{i} \frac{|a_{i}(\pi)|}{|a_{0}(\pi)|}\right)}{\frac{\Delta R}{n}}.$$

This proof uses Cauchy's bound on the size of the largest polynomial root as it is simple to write and an efficient bound. Tighter bound have been developed in the litterature that could also be used here. See Akritas et al. (2008) for example.

Using this polynomial formulation, it is also possible to derive a theoretical bound on the smallest distance between breakpoints. Mignotte's separation bound (Collins, 2001) gives a lower bound on the distance between two roots of the polynomial, as a function of the coefficient of such polynomial. This bound can then be transfered to a bound on the distance between breakpoints. This kind of bound could be useful to choose the necessary precision for the computation of the breakpoints, but are intractable to compute and the obtained values are too small to be relevant in practice.

D.6. About Proposition 4.7.

Proposition D.6 (Formal). Let \mathcal{B}^h and Γ^h respectively be the set of breakpoints and the optimality front when the MDP starts at timestep h. Let $\mathcal{B}((X_i)_i, I)$ the set of breakpoints for the MDP with a single state and reward distributions $(X_i)_i$ (not assumed deterministic).

$$\mathcal{B}^{h} = \mathcal{B}^{h+1} \cup \left(\bigcup_{x \in \mathcal{X}, (\pi_{k}^{h}, I_{k}^{h}) \in \Gamma^{h}} \mathcal{B}\left([R_{h}^{\pi_{k}^{h}}(x, a)]_{a}, I_{k}^{h} \right) \right)$$

Proof. Consider β_0 a breakpoint. There exists h such that $\beta_0 \in \mathcal{B}^h \setminus \mathcal{B}^{h+1}$. This h corresponds to the last timestep where the optimal policy changes when the risk parameter crosses β_0 .

In particular, let $(\pi^{I}, I) \in \Gamma^{h+1}$ such that $\beta_{0} \in I$. By definition,

$$\forall t \ge h+1, \forall \beta \in I, \quad [\pi_{\beta}^*]_t(x) = [\pi^I]_t(x).$$

Also, since β_0 is a breakpoint, there exists $x, a_1 \neq a_2, \pi_1 \neq \pi_2$ such that

$$\operatorname{EntRM}_{\beta_0}[R_h^{\pi^1}(x, a_1)] = \operatorname{EntRM}_{\beta_0}[R_h^{\pi^2}(x, a_2)],$$

By definition of π^I , it is also equal to

$$\operatorname{EntRM}_{\beta_0}[R_h^{\pi^I}(x,a_1)] = \operatorname{EntRM}_{\beta_0}[R_h^{\pi^I}(x,a_2)],$$

Hence, $\beta_0 \in \mathcal{B}\left([R_h^{\pi^I}(x,a)]_a, I\right)$.

E. Solving the exact breakpoints

Several issues rises when trying to computing the Optimality front with this method. The first one is that there is no easy way to compute the function $\beta \mapsto Q_{h,\beta}^{\pi}(x,\pi(x))$ without having access to the exact distribution of return for this policy, or to perform a Policy Evaluation step for each value of β , which quickly becomes computationally inefficient.

A second issue is that this system of equation is only valid if there is a single breakpoint (i.e., no other optimal policy) between those two policies. Assume that we know π_1 is optimal for β_1 and π_2 is optimal for π_2 , solving the equations gives value of the risk parameter for which one policy becomes better than the other. However, there could also be a third (or more) policy π_3 which is optimal for values of β between β_1 and β_2 . Computing the optimality front would require computing the breakpoints between π_1 and π_3 , and between π_3 and π_2 . As there are no simple conditions to verify the existence of another optimal policy between two known one, solving the exact breakpoint for a policy π_1 optimal for β_1 would require to solve the system of equation for all possible policy π_2 , and retrieving the lowest breakpoint obtained among them all. Selecting this lowest breakpoint β_2 ensures that π_1 is only optimal for the a risk parameter up to β_2 . Because of the exponential number of possible policies, such method is intractable in practice, and some approximation will be required.

Lastly, the equations themselves are non-trivial. Indeed, knowing the distribution of rewards, such an equation takes the form $\sum_{i=1}^{k} a_i e^{\alpha_i \beta} = 0$, where the coefficient a_i may be non-positive. The function in question can be expressed as the difference of two convex functions, but it lacks many of the regularity properties needed for simple optimization methods. To the best of our knowledge, the best algorithms for solving such problems reduce it to finding roots of polynomial and using efficient solvers. Yet, the non-linear transformation required (see proof of Theorem D.5) makes the approximation error become significant when transposed back to our initial problem.

F. Analysis of the number of breakpoints

F.1. Theoretical bound on the number of breakpoints

Proposition F.1. Let n the number of possible values of \mathbb{R}^{π} . The number of breakpoints B verifies

$$B < n \cdot |\mathcal{A}|^{2|X|H}$$

Proof. (of Proposition F.1)

We first consider this lemma, on the number of roots of an exponential sum.

Lemma F.2 ((Tossavainen, 2007)). Let $f_n(x) = \sum_{i=0}^{n} b_i k_i^x$, $b_i \in \mathbb{R}$, $k_i > 0$. Then f_n has at most n-1 roots.

This lemma allows for bounding the number of values of the risk parameter β for which two different policies have the same entropic risk.

Proposition F.3. Consider two distributions $\mu_1 \neq \mu_2$ with support on $X = \{x_1, \ldots, x_n\} \subset \mathbb{R}$ of size n. Consider $X_1 \sim \mu_1, X_2 \sim \mu_2$ random variables following those distributions. Consider $\mathcal{B} = \{\beta \in \mathbb{R} \mid \text{EntRM}_{\beta}(X_1) = \text{EntRM}_{\beta}(X_2)\}$. Then: $|\mathcal{B}| \leq n-1$

The proposition is straightforward by considering the exponential form of the EntRM, which is a sum of exponentials.

Finally we conclude by considering that there are $|\mathcal{A}|^{|X|H}$ deterministic markovian policies for a specific MDP, and thus $|\mathcal{A}|^{2|X|H}$ pairs of policies. The breakpoints are values for which the entropic risk of two policies is equal, and thus are included in the union of the "breakpoints" of all pairs of policies. By Proposition F.3, the number of breakpoints is at most $(n-1)|\mathcal{A}|^{2|X|H}$.

This proposition is not in general. In the conclusion we consider the number of point of equality between any two policies, but those points cannot all be breakpoints. Only the points where one of the two policy is optimal count as breakpoints. This problem is reduced to finding the number of components of the function $h(\beta) = \max\{f_{\pi}(\beta)\}_{\pi \in \Pi}$, where $f_{\pi}(\beta) = \text{EntRM}_{\beta}[R^{\pi}]$. This combinatorial problem has been studied before (see Lemma 2.4 in Atallah (1985)), but to the best of our knowledge, no better explicit formula has been found.

F.2. Evolution with the number of actions and atoms in the distributions

We conducted experiments in the single-state setting (the *FindBreaks* setting) to determine whether the theoretical bound on the number of breakpoints is reached in practice. We generated numerous random reward distributions and used our algorithm to find the number of breakpoints. We performed two experiments: one evaluating how the number of breakpoints evolves with respect to the number of actions, and the other with respect to the number of atoms in the reward distributions.

All the distributions considered have values in [0, 1] with atoms evenly spaced over this interval. To generate these distributions, we treat each as an element of $[0, 1]^n$, where the sum of the elements equals 1, representing a point on the *n*-dimensional simplex. The distributions are thus generated by uniformly sampling a point on this simplex.

For the first experiment, we fixed the number of atoms to 10 and varied the number of actions from 5 to 50. For the second experiment, we fixed the number of actions to 10 and varied the number of atoms from 5 to 50 as well. For each action, the reward distribution was generated randomly as previously described. In the solving algorithm, we searched for breakpoints for β values in the range [-15, 15] with a precision of 0.01. For each plot, we generated 100 independent problems and displayed a histogram of the number of breakpoints found across these 100 problems.




Figure 3. Illustration of the evolution of the number of breakpoints when the number of actions (Up) and atoms in the distributions (Down) is increasing. While the number of actions and atoms is multiplied by 10, the average number of breakpoints increases by less than a factor 2.

The results are shown in Figure 3. We observe that the number of breakpoints increases according to the studied parameters, with an average number of crossings of 1.25, 1.9, 2.19, and 2.36 for the first experiment, and 1.42, 1.66, 1.93, and 2.05 for the second. However, this increase is far from the theoretical bound established in Proposition F.1. Indeed, the number of breakpoints is much lower than the theoretical bound, showing sublinear growth. This experiment confirms the efficiency of our algorithm, whose complexity is strongly tied to the number of breakpoints.

G. FindBreaks Algorithm

Here *FindBreaks* is designed to be able to handle values of the risk parameter both positive and negative. In practice, using DOLFIN, all the values will be non-positive. The values of the increments Δ come from Theorem D.2.

Empirical Evaluation and Performance Analysis A simple simulation illustrates the behavior or Algorithm 2. We consider two actions, a_1 and a_2 , with reward distributions $\rho(a_1) = \frac{1}{2}\delta_0 + \frac{1}{2}\delta_1$ and $\rho(a_2) = \frac{99}{100}\delta_0 + \frac{1}{100}\delta_2$. Action a_1 is better in expectation (lower risk) but action a_2 can achieve a higher reward with small probability and should only

Algorithm 2 FindBreaks: Computing all optimal actions

Require: Precision $\varepsilon \in (0, 1)$, Random rewards $(R(a_i))_i$, interval *I*.

- 1: Compute $a^* = \arg \max_a \operatorname{EntRM}_0(R(a))$ {Initial optimial action}
- Compute A = min_{a2≠a*} EntRM₀(R(a*)) EntRM₀(R(a₂)), ΔR = r_{max} r_{min} {Advantage function and range of rewards}
- 3: Initialize $\beta = \frac{8\Delta\mu}{\Delta R^2}$, $\beta_{old} = 0$ {Initial parameter values}
- 4: Initialize $b_{\ell} = \overline{0}, a_{\beta_{\text{old}}} = a^*$
- 5: while $\beta_{\min} < \beta < \beta_{\max}$ do
- 6: Compute the β risks EntRM_{β}(R(a)) for each action a {Evaluate risks for all actions}
- 7: $a_{\beta} = \arg \max_{a} \operatorname{EntRM}_{\beta}(R(a))$ {Select the action with the highest utility}
- 8: **if** $a_{\beta} \neq a_{\beta_{\text{old}}}$ **then**
- 9: Add $[b_{\ell}, \beta]$ to the interval set \mathcal{I} and a_{β} to the optimality front Π^* {Update intervals and front}
- 10: $b_{\ell} \leftarrow \beta$ {Update the lower bound of the next interval}
- 11: end if

12: $a_{\beta_{\text{old}}} \leftarrow a_{\beta}$ {Update the last optimal action} $\Delta U = \min_{a \neq a_{\beta}} \left(\text{EntRM}_{\beta}(R(a_{\beta})) - \text{EntRM}_{\beta}(R(a)) \right)$ {Smallest optimality gap for non-optimal actions} 13: 14: if $\beta < 0$ then $\beta \leftarrow \beta - \max\{\beta \frac{\Delta U}{\Delta R}, \varepsilon\}$ {Decrease β for negative values} 15: 16: end if 17: if $\beta > 0$ then $\beta \leftarrow \beta + \max\{\beta \frac{\Delta U}{\Lambda R}, \varepsilon\}$ {Increase β for positive values} 18: 19: end if 20: end while 21: Add $[b_{\ell}, \beta_{\max}]$ to \mathcal{I} {Add the last interval to the set} **output** Optimality Front Γ^*

outperform action a_1 for large risk parameters. Figure 4 illustrates this: the functions EntRM(R(a)) are plotted for $\beta > 0$. As soon as the risk parameter β is large enough (specifically, $\beta > 3.9$), action a_2 becomes optimal.

Algorithm 2 was executed on this example for $\beta \in [0, 8]$, with a precision of $\varepsilon = 10^{-2}$, but our theoretical upper bound³ is $\beta_{\max} = \ln(100) = 4.6$. The green markers correspond to the values of β where the algorithm computed the EntRM, while the red markers represent the values that would be computed using a plain grid search with precision ε . As expected, we observe that the intervals shrink near the breakpoint, but grow significantly larger as we move away from these regions. Figure 4 (Right) zooms in on the interval $\beta \in [3.6, 4.2]$ to better visualize the concentration of intervals. Around $\beta = 3.9$, we observe that a few intervals are indeed capped by the maximal precision.

In this simple example, the naive grid uses 800 evaluations while Algorithm 2 only requires 22, with an *efficiency ratio* of 800/22 = 36. To better quantify this gain, we run another experiment on random problems with 8 actions and reward function supported on 20 atoms in [0, 1] (hence with possibly much more that 1 breakpoint). On Figure 5, for each level of precision $\varepsilon \in [10^{-3}, 10^{-1}]$, we compare the number of evaluations required by Algorithm 2 with the naive $1/\varepsilon$ obtained with a plain grid search. We report the gain on average over 20 random problem and notice efficiency ratios up to 35 for high-precision ε values.

The performance of this algorithm is also closely tied to the number of breakpoints. The more breakpoints there are, the smaller the intervals will tend to be, which in turn reduces the algorithm's efficiency compared to a plain grid search. Therefore, the number of breakpoints is a critical factor. Appendix F show that the number of breakpoints grows much slower than the theoretical bound. This allows our algorithm to be more efficient in practice than predicted by theory.

²This experiment is design to test the transition to a risky action, so it is only relevant to observe the optimality front for $\beta > 0$.

³We show a larger upper bound for visualisation purposes. Our algorithm only evaluates 3 values past this limit so it does not hurt performance significantly.




Figure 4. Utility functions on a single-state problem: a conservative (blue) and risky (red) actions, with respective ranges of optimality intersecting at $\beta_{bp} = 3.9 \pm 10^{-2}$. In red, we show the 800 values of β tested with a naive grid; in green the 22 values tested by Algorithm 2 to identify the breakpoint. Right-hand figure zooms around β_{bp} .




Figure 5. Performance gained using Algorithm 2. (Left) Number of evaluations, (Right) efficiency ratio (red over blue).

H. More Numerical Experiments

H.1. Inventory Management

We first consider some more values for the Inventory Management environments.

T/μ^*	0.25	0.33	0.5	0.66	0.75
Optimality Front	$1.26\mathrm{e}^{-5}$	$8.40\mathrm{e}^{-5}$	$3.26\mathrm{e}^{-3}$	$3.91e^{-2}$	$8.78e^{-2}$
Proxy Optimization	$2.33e^{-5}$	$1.18e^{-4}$	$3.28e^{-3}$	$3.91e^{-2}$	$8.78e^{-2}$
Risk neutral optimal	$1.11e^{-4}$	$4.24e^{-4}$	$5.73e^{-3}$	$4.62e^{-2}$	$9.77e^{-2}$
Nested Prob. Thresh.	$1.54e^{-3}$	$8.37e^{-3}$	1	1	1
Optimal value	$6.29e^{-7}$	$8.22e^{-6}$	$7.85e^{-4}$	$1.65e^{-2}$	$4.47e^{-2}$

Table 3. Evaluation of $P(R^{\pi} \leq T)$ for Inventory Management.

The values in Table 3 and Table 4 reveal that the improvements of our *Optimality Front* method over the *Proxy Optimization* becomes less significant as the level of risks decreases.

We also plot in Figure 6 the efficiency ratio of using the Findbreaks, similar to Figure 5. This figure highlights the polynomial gain in performances of using DOLFIN and Findbreaks when one wants to compute all the optimal policies up to a certain accuracy.

Risk Measure					CVaR				
Risk parameter α	0.05	0.1	0.2	0.5	0.05	0.1	0.2	0.5	
Optimality Front	37	40	44	53	32.54	35.56	39.03	45.03	
Proxy Optimization	36	40	44	53	32.28	35.46	39.00	45.03	
Risk neutral optimal	36	40	44	53	31.45	34.68	38.44	44.74	
Nested Risk Measure	35	38	42	51	30.00	33.80	37.98	44.89	

Table 4. Evaluation of (C)VaR $[R^{\pi}]$ for Inventory Management.




Figure 6. Performance gained using Algorithm 1 on Inventory Management.

H.2. Cliff Grid World

Here we consider the Cliff grid world (Sutton & Barto, 2018) illustrated in Figure 1. The agent starts in the blue state. At each step, she has a small probability (0.1 here) of moving to another random direction. Due to these random transitions, it is risky to walk too close to the cliff (bottom, in red, negative reward $-\frac{1}{2}$), and conservative policies will prefer to walk further away to reach the goal (in green). The horizon is fixed at H = 15, so in principle, the agent has enough time to reach the end using the safe path. The reward when the goal is reached at step h is $1 - \frac{h}{2H}$, which encourages the agent to reach it as fast as possible.

The agent thus faces a dilemma: she could either walk along the cliff, risking to fall but reaching the goal faster and consequently receiving a higher reward; or she could go all the way around, taking more time but with a lower probability to fall down. This can be observed in Figure 1, where the optimal policies for different values of β are shown. For high values of beta (e.g., $\beta = 10$), the agent takes the risky path, while for low values of beta (e.g., $\beta = -5$), the agent takes the safe path. One can even observe that for extremely negative values of β values (e.g., $\beta = -10$) the agent prefers to stay away from the cliff, not even trying to reach the goal (see purple arrow in top right corner pointing up).

The Cliff environment is very different from Inventory Management in its nature. The agent only receives rewards when a certain state is reached, making the reward scarce and the return distribution simpler. This implies that the optimal policy for different measures of risks, such as Probability Threshold, VaR and CVaR are markovian for this MDP.

For the Threshold Probability problem, we only consider 2 values of the threshold, -0.5 corresponding to falling into the cliff, and 0 corresponding to not reaching the goal. For the first threshold, the objective is to find the policy that is least likely to fall, while for the second it is to find the policy with the most chances of reaching the goal.

Table 1 confirms that our *Optimality Front* method performs better than other methods for the Threshold probability. An important remark is that, here, the real optimal value is reached by the optimality front. Similar performances are observed for the CVaR in Table 6. For the VaR, the gain in performance is limited, which is explained by the scarcity of rewards in the environment (the small changes in the return distribution does not change the value of the VaR).

Compared to the Inventory Management, the gain performance for computing the optimality front is much better, with a ratio up to a factor 100, as seen in Figure 7.

	$I(n \geq I)$	IOI CIIII.
Т	-0.5	0
Optimality Front	$3.72e^{-2}$	$4.65\mathrm{e}^{-2}$
Proxy Optimization	$3.38e^{-2}$	$4.67e^{-2}$
Risk neutral optimal	$4.50e^{-2}$	$4.84e^{-2}$
Nested Prob. Thresh.	1	1
Optimal value	$3.72e^{-2}$	$4.65\mathrm{e}^{-2}$

Table 5. Evaluation of $P(R^{\pi} \leq T)$ for Cliff.

Table 6. Evaluation of (C)VaR_{α}[R^{π}] for Cliff.

Risk Measure		Va	αR			CVa	R	
Risk parameter α	0.05	0.1	0.2	0.5	0.05	0.1	0.2	0.5
Optimality Front	0.53	0.63	0.70	0.76	-0.37	0.11	0.38	0.58
Proxy Optimization	0.00	0.6	0.70	0.76	-0.39	-0.08	0.38	0.58
Risk neutral optimal	0.53	0.63	0.70	0.76	-0.43	0.08	0.37	0.58
Nested Risk Measure	-0.5	-0.5	-0.5	-0.5	-0.5	-0.5	-0.5	-0.5




Figure 7. Performance gained using Algorithm 1 on Cliff. (Left) Number of evaluations, (Right) efficiency ratio (red over blue).