Quantum group deformation of the Kittel–Shore model

A. BALLESTEROS¹, I. GUTIÉRREZ-SAGREDO², V. MARISCAL¹ AND J.J. RELANCIO^{2,3}

¹Departamento de Física, Universidad de Burgos, 09001 Burgos, Spain

²Departamento de Matemáticas y Computación, Universidad de Burgos, 09001 Burgos, Spain

³Centro de Astropartículas y Física de Altas Energías (CAPA), Universidad de Zaragoza, Zaragoza 50009, Spain

e-mail: angelb@ubu.es, igsagredo@ubu.es, vmariscal@ubu.es, jjrelancio@ubu.es

Abstract

The Kittel–Shore (KS) Hamiltonian describes N spins with long-range interactions that are identically coupled. In this paper, the underlying $\mathfrak{su}(2)$ coalgebra symmetry of the KS model is demonstrated for arbitrary spins, and the quantum deformation of the KS Hamiltonian (q-KS model) is obtained using the corresponding $\mathfrak{su}_q(2)$ quantum group. By construction, the existence of such a symmetry guarantees that all integrability properties of the KS model are preserved under q-deformation. In particular, the q-KS model for spin-1/2 particles is analysed in both ferromagnetic and antiferromagnetic couplings, and the cases with N = 2, 3, and 4 spins are studied in detail. The higher-spin q-KS models are sketched.

PACS:

KEYWORDS: Kittel-Shore model; quantum algebras; coalgebras; integrability; spin systems

Contents

T	Intr	oduction	2	
2	The Kittel-Shore model			
	2.1	Coalgebra symmetry of the KS model	5	
	2.2	Eigenvectors and eigenvalues	7	
	2.3	The spin- $1/2$ model	8	
3 The <i>q</i> -deformed KS model		q-deformed KS model	13	
	3.1	The quantum algebra $su_q(2)$	14	
	3.2	The generic q -Kittel Shore model \ldots	15	

4	The	e spin-1/2 q -KS model	17
	4.1	Eigenvectors and eigenvalues	17
	4.2	The $N = 2, 3, 4$ cases \ldots	19
		4.2.1 The $N = 2$ case and the <i>q</i> -exchange operator $\ldots \ldots \ldots$	20
		4.2.2 The $N = 3$ case \ldots	22
		4.2.3 The $N = 4$ case	23
5	Hig	her spin q -KS models	24
	5.1	The q-KS model for $j = 1$	24
	5.2	The q-KS model for arbitrary spin	25
6	Con	ncluding remarks	25

1 Introduction

Let us consider the KS Hamiltonian [1] that describes a system of N long-range interacting spins with identical coupling among all of them, namely

$$H_{KS} = -I \sum_{i < j}^{N} \vec{J_i} \cdot \vec{J_j} , \qquad (1)$$

where $\vec{J}_i = (J_x^{(i)}, J_y^{(i)}, J_z^{(i)})$ are the (arbitrary) spin operators for the *i*-th particle. The coupling constant distinguishes between ferromagnetic (I > 0) and antiferromagnetic cases (I < 0), respectively. When a constant external magnetic field $\vec{h} = (0, 0, h)$ is considered, the Hamiltonian reads

$$\tilde{H}_{KS} = -I \sum_{i < j}^{N} \vec{J}_{i} \cdot \vec{J}_{j} - \gamma h \sum_{i=1}^{N} J_{z}^{(i)},$$
(2)

where $\gamma = g\mu_B$, in the usual notation.

The KS Hamiltonian (1) is simply the long-range generalisation for arbitrary spin of the Heisenberg XXX model with a fully symmetric constant coupling among all spins and presents a number of outstanding features. As Kittel and Shore emphasised in their original paper [1], it provides a 'rigorously solvable' many-body system, whose exact solvability stems from the well-known representation theory of the $\mathfrak{su}(2)$ Lie algebra, particularly because of the well-known decomposition of the tensor product of N irreducible representations of $\mathfrak{su}(2)$ into irreducible blocks. Consequently, there is no need to use the Bethe Ansatz technique to solve this problem. Moreover, this system provides an exactly solvable framework to study the theory of phase transitions for long-range interacting systems; in particular, it was found to present a ferromagnetic phase transition which develops very slowly as N increases [1].

Although the spin-1/2 case was studied in [1] and later in [2, 3, 4], the underlying integrability properties of the model are preserved for arbitrary spins. The study of the KS model was extended to the antiferromagnetic case in [2, 5]. In [3], group-theoretical properties were analysed, and an integral formula for the partition function in the ferromagnetic case was presented. Note also that H_{KS} can be viewed as a model of N spins located on the vertices of a (N-1)-dimensional simplex and, in this context, the model can be considered as infinite dimensional in the thermodynamic limit $N \to \infty$. Nevertheless, the cases with N = 2, 3, 4 (dimer, equilateral triangle, and regular tetrahedron) arrays of spins are realised in the context of ultra-small magnetic clusters (see [6] and references therein) or as Hamiltonians for quantum computation with quantum dots [7, 8].

The KS model for classical spins has also been studied systematically from different viewpoints as a long-range integrable generalisation of the Heisenberg model for classical spins, the dynamical behaviour of which, including the KS case, was analysed in [9]. The N independent integrals in involution of the KS model, as well as its action-angle variables were presented in [10], and the autocorrelation function of the so-called 'equivalent-neighbour Heisenberg model' (classical KS model) was calculated analytically in [11]. As another example of its applications, in [12, 13], the KS model was used to demonstrate the extension of Fisher's and Barber's finite-size scaling hypothesis [14] to the case of 'infinitely coordinated systems' (long-range interactions). Moreover, the correlation functions for the anisotropic KS model (the so-called 'equivalent-neighbour XYZ model') have also been considered [15, 16].

On the other hand, it is also well-known that quantum groups [17, 18] provide symmetries of certain integrable spin chains endowed with specific boundary conditions. This is the case for the XXZ Heisenberg model for N spin-1/2 particles, for which $\mathfrak{su}_q(2)$ turns out to be the symmetry algebra, provided that the quantum deformation parameter q and the anisotropy parameter δ of the model are related through $\delta = (q+q^{-1})/2$, and the extra boundary term is just $(q-q^{-1})(\sigma_z^1 - \sigma_z^N)/2$ (see [19, 20, 21] and references therein). In this sense, the spin-1/2 Heisenberg XXZ Hamiltonian can be understood as a $\mathfrak{su}_q(2)$ deformation of the isotropic XXX model, which is recovered in the limit $q \to 1$ because under this condition, both the anisotropy parameter δ and boundary term vanish.

The aim of this paper is to explicitly present and solve the $\mathfrak{su}_q(2)$ analogue of the KS model (2) for q not being a root of unity. The definition of this new quantum-group invariant model will be possible after unveiling the $\mathfrak{su}(2)$ coalgebra symmetry [22] of the undeformed KS Hamiltonian, which will be presented in the next Section. Thus, the KS model will also be revisited from a coalgebraic perspective: its eigenvalues and eigenvectors will be derived, thus allowing the exact derivation of its partition function. It is also worth noting that the Zeeman effect induced by the magnetic field is not affected by the q-deformation.

The q-deformation of the KS model is introduced in Section 3 by imposing its $\mathfrak{su}_q(2)$ coalgebra symmetry, thus obtaining a new exactly solvable N-dimensional quantum model with long-range interactions. In particular, the q-KS Hamiltonian for N spin-1/2 particles is thoroughly studied in Section 4. As essential results, we found that owing to the underlying coalgebra symmetry, the q-KS Hamiltonian preserves the complete integrability of the KS model and presents the same degeneracies of its spectrum, which nevertheless becomes nonlinearly deformed with respect to the KS model. These results are based on the fact that, where q is not a root of unity, the representation theory of $\mathfrak{su}_q(2)$ is structurally identical to that of $\mathfrak{su}(2)$ [23]. Moreover, to realise the $\mathfrak{su}_q(2)$ invariance of the Hamiltonian, the q-KS model is forced to break the characteristic KS symmetry between the interactions among all the N spins, which now become quite involved and site dependent, a fact that is induced by the non-local nature of the $\mathfrak{su}_q(2)$ coproduct maps. Nevertheless, the underlying quantum group symmetry allows the explicit knowledge of the full spectrum and eigenvectors, from which the partition function can be derived.

In particular, the density of states of the q-KS model will be compared with that of the KS model, and the smooth effect of the deformation can be appreciated. A deeper analysis of the q-KS Hamiltonian for N = 2, 3, and 4 spin-1/2 particles is provided, including the explicit form of the eigenvectors, which can be obtained through iterated application of the q-Clebsch–Gordan coefficients [23, 24]. In the ferromagnetic case, the ground state eigenvectors are just the so-called q-Dicke states [25, 26, 27], which are the most excited states in the antiferromagnetic model. The ground states for the antiferromagnetic q-KS model are also presented, which conversely provide the highest excited states for the ferromagnetic case. In Section 5, the q-deformed KS model for N spin-1 particles is explicitly presented, and the q-KS models with arbitrary spin are sketched using the so-called functional realisation approach to the $\mathfrak{su}_q(2)$ representations [28, 29, 30]. Finally, a Section that includes some remarks and open problems closes this paper.

2 The Kittel-Shore model

In this Section, we review the well-known KS model, unveiling its $\mathfrak{su}(2)$ coalgebra symmetry. This fact will be shown to have profound implications for the complete solvability of the model and for the possibility of defining an integrable q-deformation of the KS Hamiltonian, which will be presented in the next Section.

Let us start by considering the $\mathfrak{su}(2)$ Lie algebra, whose generators satisfy the following commutation relations

$$[J_x, J_y] = iJ_z, \qquad [J_y, J_z] = iJ_x, \qquad [J_z, J_x] = iJ_y.$$
 (3)

The Casimir operator for this Lie algebra is given by

$$C = J_x^2 + J_y^2 + J_z^2 \,. \tag{4}$$

If we perform the change of basis

$$J_{\pm} = J_x \pm i J_y, \qquad \qquad J_x = \frac{1}{2} (J_+ + J_-), \qquad J_y = -\frac{i}{2} (J_+ - J_-), \qquad (5)$$

the commutation rules become

$$[J_z, J_{\pm}] = \pm J_{\pm}, \qquad [J_+ J_-] = 2 J_z , \qquad (6)$$

and the Casimir operator now reads

$$C = J_{-}J_{+} + J_{z}^{2} + J_{z} = J_{+}J_{-} + J_{z}^{2} - J_{z}.$$
(7)

As it is well-known, the spin-j irreducible representation D^j of the $\mathfrak{su}(2)$ algebra is defined on a (2j+1)-dimensional space by the action on the $|j,m\rangle$ basis vectors given by

$$J_{\pm}|j,m\rangle = \sqrt{(j \mp m)(j \pm m + 1)}|j,m \pm 1\rangle, \qquad J_{z}|j,m\rangle = m|j,m\rangle, \tag{8}$$

where j = 0, 1/2, 1, 3/2, ... and m = -j, ..., +j. The common eigenvalue of the Casimir operator on all the D^j basis vectors is given by $C|j,m\rangle = j(j+1)|j,m\rangle$ and provides the expected value of the squared modulus of the spin.

The tensor product representation $D^{j_1} \otimes D^{j_2}$ turns out to be fully reducible, and it can be decomposed into irreducible components in the form

$$D^{j_1+j_2} \oplus D^{j_1+j_2-1} \oplus \dots \oplus D^{|j_1-j_2|}$$

$$\tag{9}$$

The Clebsch–Gordan coefficients provide the transformation matrix between the $((2j_1 + 1)(2j_2 + 1))$ dimensional basis $|j_1, m_1\rangle \otimes |j_2, m_2\rangle$ corresponding to the tensor product representation $D^{j_1} \otimes D^{j_2}$ and the block-diagonal one (9), whose basis will be given by the vectors

$$\{|j_1+j_2,m_{j_1+j_2}\rangle, |j_1+j_2-1,m_{j_1+j_2-1}\rangle, \dots, ||j_1-j_2|,m_{|j_1-j_2|}\rangle\}, \qquad m_{\alpha} = -j_{\alpha}, \dots, j_{\alpha}, \qquad (10)$$

where subscript α labels each irreducible subspace.

The iteration of this procedure allows the transformation of the tensor product representation of N arbitrary irreducible representations of the $\mathfrak{su}(2)$ algebra

$$D^{j_1} \otimes D^{j_2} \otimes \dots \otimes D^{j_N},$$
 (11)

into its block-diagonal form in terms of k irreducible representations

$$D^{j_{\alpha_1}} \oplus D^{j_{\alpha_2}} \oplus \dots \oplus D^{j_{\alpha_k}},$$
 (12)

where $j_{\alpha_1} = j_1 + j_2 + \cdots + j_N$ is the spin of the representation with the highest dimension and $(j_{\alpha_2}, j_{\alpha_3}, \ldots, j_{\alpha_k})$ are the spins of each of the lower-dimensional irreducible representations that arise in the decomposition. In general, some of these spins j_{α_i} can be the same, giving rise to the so-called multiplicities for a given irreducible representation with spin J (see [31] and references therein for the derivation of explicit formulas giving the multiplicities when the tensor product (11) of N representations with identical spin j are considered). The associated block-diagonal basis for the representation (12) will be given by the vectors

$$\{|j_{\alpha_1}, m_{\alpha_1}\rangle, |j_{\alpha_2}, m_{\alpha_2}\rangle, \dots, |j_{\alpha_k}, m_{\alpha_k}\rangle\}, \qquad m_{\alpha_i} = -j_{\alpha_i}, \dots, j_{\alpha_i},$$
(13)

that can be expressed in terms of the initial basis vectors corresponding to (11)

$$|j_1, m_1\rangle \otimes |j_2, m_2\rangle \otimes \cdots \otimes |j_N, m_N\rangle, \qquad m_i = -j_i, \dots, j_i, \qquad i = 1, \dots, N,$$
(14)

using the corresponding iteration of the Clebsch–Gordan transformations. Explicit examples of this will be provided in the following Sections.

2.1 Coalgebra symmetry of the KS model

A Hopf algebra (see, for instance [32]) is a (unital, associative) algebra A endowed with two homomorphisms called coproduct ($\Delta : A \longrightarrow A \otimes A$) and counit ($\epsilon : A \longrightarrow \mathbb{C}$), as well as an antihomomorphism (the antipode $\gamma : A \longrightarrow A$) such that, $\forall a \in A$:

$$(id \otimes \Delta)\Delta(a) = (\Delta \otimes id)\Delta(a), \tag{15}$$

$$(id \otimes \epsilon)\Delta(a) = (\epsilon \otimes id)\Delta(a) = a, \tag{16}$$

$$m((id \otimes \gamma)\Delta(a)) = m((\gamma \otimes id)\Delta(a)) = \epsilon(a)1,$$
(17)

where *m* is the usual multiplication mapping $m(a \otimes b) = a \cdot b$ and $id : A \to A$ is the identity map. The condition (15) is called the coassociativity property of Δ . When *A* is thought of as a module endowed with comultiplication and counit maps (Δ, ϵ) fulfilling (15) and (16), then *A* is said to be endowed with a coalgebra structure.

This means that the coproduct map Δ underlies the construction of the tensor product representations of two copies of the Hopf algebra A. Moreover, the coassociativity (15) of the coproduct map Δ provides its generalisation to any number of copies of A. If we denote $\Delta^{(2)} \equiv \Delta$, then the homomorphism $\Delta^{(3)}$: $A \to A \otimes A \otimes A$ can be defined through any of the two equivalent expressions:

$$\Delta^{(3)} := (id \otimes \Delta^{(2)}) \circ \Delta^{(2)} = (\Delta^{(2)} \otimes id) \circ \Delta^{(2)}, \qquad (18)$$

and this construction can be generalized to an arbitrary number N of tensor products of A either by the recurrence relation

$$\Delta^{(N)} := (id \otimes id \otimes \dots^{N-2)} \otimes id \otimes \Delta^{(2)}) \circ \Delta^{(N-1)}, \tag{19}$$

or by the equivalent one

$$\Delta^{(N)} := (\Delta^{(2)} \otimes id \otimes \dots^{N-2)} \otimes id \otimes id) \circ \Delta^{(N-1)}.$$
⁽²⁰⁾

In particular, the universal enveloping algebra $\mathcal{U}(\mathfrak{g})$ of any Lie algebra \mathfrak{g} can be endowed with a Hopf algebra structure by defining

$$\Delta_0(X) = \mathbf{I} \otimes X + X \otimes \mathbf{I}, \quad \forall X \in \mathfrak{g} \qquad \Delta_0(\mathbf{I}) = \mathbf{I} \otimes \mathbf{I}, \tag{21}$$

where **I** denotes the identity element in $\mathcal{U}(\mathfrak{g})$. The coproduct map can be extended to any monomial in $\mathcal{U}(\mathfrak{g})$ by means of the homomorphism condition for the coproduct $\Delta_0(X \cdot Y) = \Delta_0(X) \cdot \Delta_0(Y)$, which implies the compatibility of the coproduct Δ with the Lie bracket

$$[\Delta_0(X_i), \Delta_0(X_j)]_{\mathfrak{g}\otimes\mathfrak{g}} = \Delta_0([X_i, X_j]_{\mathfrak{g}}), \qquad \forall X_i, X_j \in \mathfrak{g}.$$
(22)

From (19) is straightforward to prove that the N-th coproduct of the Lie algebra generators is given by

$$\Delta_{0}^{(N)}(X_{i}) = X_{i} \otimes \mathbf{I} \otimes \mathbf{I} \otimes \dots^{N-1} \otimes \mathbf{I} + \mathbf{I} \otimes X_{i} \otimes \mathbf{I} \otimes \dots^{N-2} \otimes \mathbf{I} + \dots + \mathbf{I} \otimes \mathbf{I} \otimes \dots^{N-1} \otimes \mathbf{I} \otimes X_{i}.$$
(23)

By recalling the so-called "coalgebra symmetry" approach introduced in [33, 22], let us now consider N copies of the $\mathfrak{su}(2)$ Lie algebra (6) with generators $\{J_{\pm}^{(i)}, J_{z}^{(i)}\}$, where the N-th coproduct map (23) of the $\mathfrak{su}(2)$ generators provide the collective spin operators

$$\Delta_0^{(N)}(J_{\pm}) = \sum_{i=1}^N J_{\pm}^{(i)}, \qquad \Delta_0^{(N)}(J_z) = \sum_{i=1}^N J_z^{(i)}, \tag{24}$$

and superindex i represents the spin operator of the *i*-th particle. By making use of these expressions, it can be straightforwardly proven that the KS Hamiltonian (2) can be rewritten as

$$\tilde{H}_{KS} = -I \sum_{i < j}^{N} \vec{J}_{i} \cdot \vec{J}_{j} - \gamma h \sum_{i=1}^{N} J_{z}^{(i)} = -\frac{I}{2} \left(\Delta_{0}^{(N)}(C) - \sum_{i=1}^{N} C^{(i)} \right) - \gamma h \Delta_{0}^{(N)}(J_{z}),$$
(25)

where $C^{(i)} = J_{-}^{(i)} J_{+}^{(i)} + (J_z^{(i)})^2 + J_z^{(i)} = (J_x^{(i)})^2 + (J_y^{(i)})^2 + (J_z^{(i)})^2$ is the Casimir operator for the *i*-th $\mathfrak{su}(2)$ algebra. Equation (25) is proven by making use of the expressions (24) and by expanding the coproduct $\Delta_0^{(N)}(C)$ as $\Delta_0^{(N)}(J_{-}J_{+} + (J_z)^2 + J_z)$, namely

$$\tilde{H}_{KS} = -\frac{I}{2} \left(\Delta_0^{(N)}(J_-) \Delta_0^{(N)}(J_+) + (\Delta_0^{(N)}(J_z))^2 + \Delta_0^{(N)}(J_z) - \sum_{i=1}^N C^{(i)} \right) - \gamma h \Delta_0^{(N)}(J_z) , \qquad (26)$$

where, in general, different spins j_i can be considered for each site.

The expression (26) implies that the KS model is endowed with coalgebra symmetry [22] (see also [33, 34, 35], and references therein), as the Hamiltonian can be written as a function of the coproducts of the generators of a given Hopf algebra, in this case $\mathcal{U}(\mathfrak{su}(2))$ (note that the term $\sum_{i=1}^{N} C^{(i)}$ will be just a constant because the Casimir operators $C^{(i)}$ take constant values at each site). As it was demonstrated in [22], the complete integrability of any coalgebra-symmetric system is a consequence of the fact that

for any coalgebra-symmetric Hamiltonian H^N the coproducts $\Delta^{(m)}(C)$ with m = 2, ..., N of the Casimir operators provide (N-1) constants of the motion in involution, namely

$$\left[H^{(N)}, \Delta^{(m)}(C)\right] = 0, \qquad \left[\Delta^{(m)}(C), \Delta^{(k)}(C)\right] = 0, \qquad \forall m, k = 2, \dots, N.$$
(27)

Here the previous commutators are defined on $\mathcal{U}(\mathfrak{su}(2)) \otimes \ldots^{(N)} \otimes \mathcal{U}(\mathfrak{su}(2))$, and we also understand that $\Delta^{(m)}(C) \equiv \Delta^{(m)}(C) \otimes \mathbf{I} \otimes \ldots^{N-m} \otimes \mathbf{I}$ for any $m = 2, \ldots, N$.

Therefore, for any Hopf algebra A, the previous result enables the construction of a large class of integrable systems, which can be in principle diagonalised on a basis constructed from common eigenvectors of the $\Delta^{(m)}(C)$ operators (see, for instance, the case of different $\mathfrak{sl}(2)$ classical and quantum Calogero-Gaudin models and their q-deformations [22, 36, 37, 38]). In the case of the KS model presented here, the basis of $\Delta^{(m)}(C)$ eigenvectors will not be needed because the block-diagonal one (10) provides eigenvectors and eigenvalues in a straightforward manner, as we shall see below.

More importantly, a second essential consequence of the coalgebra symmetry of a given Hamiltonian H defined on a Hopf algebra A is the fact that any quantum deformation A_q of the underlying Hopf algebra structure automatically induces an integrable deformation H_q of the initial Hamiltonian [22]. Such a deformed Hamiltonian will be defined by substituting the coproduct of the generators with its quantum deformation, both in the definition of the Hamiltonian and in the *m*-th order coproducts of the q-deformed Casimir C_q . This is just the procedure that will be used in the next Section to construct a quantum deformation of the KS model by using the quantum deformation of $\mathcal{U}(\mathfrak{su}(2))$.

2.2 Eigenvectors and eigenvalues

Let us firstly consider the KS Hamiltonian (1) without external magnetic field (h = 0) in the coalgebrasymmetric form (25), namely

$$H_{KS} = -\frac{I}{2} \left(\Delta_0^{(N)}(C) - \sum_{i=1}^N C^{(i)} \right) , \qquad (28)$$

where we recall that C denotes the $\mathfrak{su}(2)$ Casimir operator (6). Since the coproduct map $\Delta_0^{(N)}$ is an algebra homomorphism, we have that

$$\left[\Delta_0^{(N)}(C), \Delta_0^{(N)}(X)\right] = \Delta_0^{(N)}([C, X]) = 0, \qquad \forall X \in \mathcal{U}(\mathfrak{su}(2)).$$
(29)

The previous expression implies, in particular, that the collective spin operators (24) commute with the KS Hamiltonian, which is essentially the *N*-th coproduct of the Casimir operator for $\mathfrak{su}(2)$. Moreover, the total spin component in the *z* direction $\Delta_0^{(N)}(J_z) = \sum_{i=1}^N J_z^{(i)}$ can be considered a good quantum number for the eigenstates of the model.

Therefore, we consider the block-diagonal basis (10) for the tensor product of N irreducible representations of $\mathfrak{su}(2)$. Thus, the $\Delta_0^{(N)}(C) \equiv C^{(N)}$ operator will be represented as a block-diagonal matrix. Within each of the irreducible blocks with basis vectors

$$|j_{\alpha_i}, m_{\alpha_i}\rangle, \qquad m_{\alpha_i} = -j_{\alpha_i}, \dots, j_{\alpha_i},$$
(30)

relations (29) imply that $C^{(N)}$ commutes with the representation of all $\mathfrak{su}(2)$ generators and, as a consequence of Schur's lemma, within each irreducible block the operator $C^{(N)}$ has to be proportional to the identity. In fact, the proportionality factor will be simply the $\mathfrak{su}(2)$ Casimir eigenvalue within such a block, that is, $j_{\alpha_i}(j_{\alpha_i} + 1)$. Therefore, we have proven that

$$\Delta_0^{(N)}(C) |j_{\alpha_i}, m_{\alpha_i}\rangle = j_{\alpha_i}(j_{\alpha_i} + 1) |j_{\alpha_i}, m_{\alpha_i}\rangle, \qquad \forall \alpha_i = 1, \dots, k, \qquad m_{\alpha_i} = -j_{\alpha_i}, \dots, j_{\alpha_i}.$$
(31)

Finally, since the term $\sum_{i=1}^{N} C^{(i)}$ acting on the initial tensor product basis

$$|j_1, m_1\rangle \otimes |j_2, m_2\rangle \otimes \cdots \otimes |j_N, m_N\rangle,$$
(32)

is just a constant, namely, $\sum_{i=1}^{N} j_i(j_i + 1)$, this term will not be affected by the consecutive Clebsch–Gordan transformations leading to the block-diagonal basis. Thus, we have proven that the eigenvectors of the KS Hamiltonian (28) are just the basis vectors (30) of the block-diagonal basis, and their eigenvalues are

$$H_{KS}|j_{\alpha_i}, m_{\alpha_i}\rangle = -\frac{I}{2} \left(\Delta_0^{(N)}(C) - \sum_{i=1}^N C^{(i)} \right) |j_{\alpha_i}, m_{\alpha_i}\rangle = -\frac{I}{2} \left(j_{\alpha_i}(j_{\alpha_i}+1) - \sum_{i=1}^N j_i(j_i+1) \right) |j_{\alpha_i}, m_{\alpha_i}\rangle.$$
(33)

Finally, when the magnetic field h is considered we have that, within each irreducible block and after the Clebsch–Gordan transformation is performed onto the $\Delta_0^{(N)}(J_z)$ operator

$$-\gamma h \Delta_0^{(N)}(J_z) |j_{\alpha_i}, m_{\alpha_i}\rangle = -\gamma h J_z^{(\alpha_i)} |j_{\alpha_i}, m_{\alpha_i}\rangle = -\gamma h m_{\alpha_i} |j_{\alpha_i}, m_{\alpha_i}\rangle.$$
(34)

Summarizing, the eigenvalues of the KS model with magnetic field for arbitrary spins are given by

$$\tilde{E}(j_{\alpha_i}, m_{\alpha_i}, K) = -\frac{I}{2} \left(j_{\alpha_i}(j_{\alpha_i} + 1) - K \right) - \gamma h \, m_{\alpha_i} \,, \tag{35}$$

where α_i (with i = 1, ..., k) labels the k irreducible blocks of the completely reduced form of the $D^{j_1} \otimes D^{j_2} \otimes \cdots \otimes D^{j_N}$ representation, where $m_{\alpha_i} = -j_{\alpha_i}, \ldots, +j_{\alpha_i}$ and $K = \sum_{i=1}^N j_i(j_i+1)$.

We stress that this result can be thought of as a consequence of (27) because H_{KS} with h = 0 is proportional to $\Delta^{(N)}(C)$ and its eigenvalues $E(j_{\alpha_i}, K)$ depend only on the modulus of the total spin for each lattice site. Moreover, $\Delta^{(N)}(C)$ commutes with the collective operators (24) that represent the three components of total spin, all of which are conserved when h = 0. When the magnetic field is turned on, the isotropy of the model is broken, and only $\Delta_0^{(N)}(J_z)$ remains as a constant of the motion, and therefore its eigenvalue m_{α_i} provides the second quantum number for the eigenstates $|j_{\alpha_i}, m_{\alpha_i}\rangle$ of the model.

2.3 The spin-1/2 model

When all the N particles carry the same j = 1/2 irreducible representation, the spin operators can be written as

$$J_x = \frac{1}{2}\sigma_x, \qquad J_y = \frac{1}{2}\sigma_y, \qquad J_z = \frac{1}{2}\sigma_z, \tag{36}$$

where σ_i are the Pauli matrices

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \qquad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \qquad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$
(37)

Therefore, the matrix form of the $\{J_+, J_-, J_z\}$ operators reads

$$J_{+} = J_{x} + i J_{y} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \sigma_{+}, \qquad J_{-} = J_{x} - i J_{y} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = \sigma_{-}, \qquad J_{z} = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (38)$$

and in this case the eigenvalues (35) read

$$\tilde{E}(j_{\alpha_i}, m_{\alpha_i}, 3N/4) = -\frac{I}{2} \left(j_{\alpha_i}(j_{\alpha_i} + 1) - 3N/4 \right) - \gamma h m_{\alpha_i}, \qquad i = 1, \dots, k,$$
(39)

where the irreducible blocks arising from the decomposition of the $D^{1/2} \otimes D^{1/2} \otimes \dots^{N} \otimes D^{1/2}$ representation are given by spins j_{α_i} that can take the values

$$J = N/2, N/2 - 1, \dots, \delta,$$
(40)

with $\delta = 0$ for N even and $\delta = 1/2$ for N odd. In general, the multiplicities d_{JNj} of the irreducible representation with spin J within the tensor product of N copies of spin j representations are explicitly given by [39]

$$d_{JNj} = \Omega(J, N, j) - \Omega(J+1, N, j), \quad \text{with} \quad \Omega(J, N, j) \equiv \text{coefficient of } x^J \text{ in } \left(x^j + x^{j-1} + \dots + x^{-j}\right)^N.$$
(41)

As mentioned in [31], this expression can be written in a simplified way

$$\Omega(J,N,j) = \sum_{k=0}^{\min\left(N, \lfloor \frac{Nj+J}{2j+1} \rfloor\right)} (-1)^k \binom{N}{k} \binom{Nj+J-(2j+1)k+N-1}{Nj+J-(2j+1)k},$$
(42)

where the floor function $\lfloor x \rfloor$ is the function that takes as input a real number x and gives as output the greatest integer less than or equal to x.

For the spin-1/2 case, these coefficients are explicitly given by [39]

$$d_{JN\frac{1}{2}} = \Omega(J, N, 1/2) - \Omega(J+1, N, 1/2) = \binom{N}{N/2 - J} - \binom{N}{N/2 - (J+1)},$$
(43)

where $0 \le J \le N/2$ and J = 0 is allowed only in the case with even N.

In the original KS paper [1] without an external magnetic field, the energy spectrum (39) is modified in the form

$$E'(j_{\alpha_i}, 3N/4) = -\frac{I}{2} \left(j_{\alpha_i}(j_{\alpha_i}+1) - 3N/4 \right) - E_0(N), \quad \text{with} \quad E_0(N) = \begin{cases} \frac{3IN}{8} & I < 0\\ -\frac{IN}{8}(N-1) & I > 0 \end{cases}$$
(44)

where the constant term $E_0(N)$ denotes the energy of the ground state. In this way, the corrected groundstate energy E' is always zero, independent of the number N of spins. In the ferromagnetic case, $E_0(N)$ corresponds to the eigenvalue of the block with the highest spin J = N/2, and in the antiferromagnetic case to the block with the lowest spin $J = \delta$. When the magnetic field is introduced, we will have

$$E'(j_{\alpha_i}, m_{\alpha_i}, 3N/4)) = -\frac{I}{2} \left(j_{\alpha_i}(j_{\alpha_i} + 1) - 3N/4 \right) - E_0(N) - \gamma h \, m_{\alpha_i} \,. \tag{45}$$

We will also adopt this approach because in this way we can compare the effect of the q-deformation in the model more easily. Also, in [1] the spectrum for the ferromagnetic case was written in terms of a quantum number p = 0, 1, 2, ..., p (where p = N/2 if N is even and p = (N - 1)/2 if N is odd). This quantum number is associated to the spin of each irreducible block and defined as

$$p = \frac{N}{2} - j_{\alpha_i} \,. \tag{46}$$

This means that p = 0 corresponds to the block with the highest spin J = N/2 (*i.e.* the one with the lowest energy), p = 1 corresponds to the first excited states corresponding to the next block(s), and so on. In terms of this quantum number and by correcting the eigenvalues in the form (44), the ferromagnetic energy spectrum without magnetic field is simply

$$E'(p,N) = \frac{I}{2}(-p^2 + (N+1)p).$$
(47)

The energy levels (44) for N = 2, 3, 4 in the antiferromagnetic and ferromagnetic cases, together with their degeneracies, are shown in Figs. 1 and 2, respectively. As expected, the presence of an external magnetic field partially breaks these degeneracies.

Therefore, the partition function for the spin-1/2 KS model is given by

$$Z_{N} = \sum_{i=1}^{k} \sum_{m_{\alpha_{i}}=-j_{\alpha_{i}}}^{m_{\alpha_{i}}=j_{\alpha_{i}}} \exp(-\beta E'(j_{\alpha_{i}}, m_{\alpha_{i}}, 3N/4)),$$
(48)

where $\beta = 1/k_B T$ and k_B is the Boltzmann constant. As we have mentioned, irreducible blocks with the same spin j_{α_i} are obtained and, by taking into account the corresponding multiplicities $d_{JN\frac{1}{2}}$ (43), the previous formula can be rewritten as

$$Z_N = \sum_J \sum_{m_J = -J}^{m_J = +J} d_{JN\frac{1}{2}} \exp(-\beta E'(J, m_J, 3N/4)), \tag{49}$$

where J denotes all the different spins obtained in the block-diagonal decomposition and the coefficients $d_{JN\frac{1}{2}}$ represent the total multiplicity of a given energy level with spin J.

The density of energy states can be obtained from the energy spectrum (35) and the explicit expression of the degeneracies (43). We present in Fig. 3 the density of states for N = 20, 100, 1000 spins in both the antiferromagnetic and ferromagnetic cases (always without an external magnetic field). As noted in [5], in the ferromagnetic case, the ground and first excited energy states are the most separated energy levels and exhibit the lowest degeneracies. In contrast, in the antiferromagnetic case, this situation is reversed, and the lowest energy levels are the closest ones and also have the highest degeneracies. Moreover, in [3] an integral formula for the partition function was obtained.

The partition function (49) can be now used to obtain the Helmholtz free energy as

$$F = -(k_B T/N) \log(Z_N), \qquad (50)$$

and all the thermodynamic functions can be obtained from F. In particular, the specific heat, magnetic susceptibility, and magnetization are given by

$$C_V = -T \left. \frac{\partial^2 F}{\partial T^2} \right|_{h=0}, \qquad \chi = - \left. \frac{\partial^2 F}{\partial h^2} \right|_{h=0}, \qquad M = -\frac{\partial F}{\partial h}.$$
(51)

In addition, as shown in [1], the Curie temperature corresponding to the ferromagnetic phase transition of the model can be estimated analytically, and it was proven that the phase transition develops very slowly as the number N of spins increases.



Figure 1: Energy levels (in the units |I|/4) and their degeneracies for N = 2, 3, 4, 5 spins with j = 1/2 coupled antiferromagnetically (a) without an external magnetic field (h = 0) and (b) with an external magnetic field (h = 0.5). Adapted from [2].



Figure 2: Energy levels (in the units |I|/4) and their degeneracies for N = 2, 3, 4, 5 spins j = 1/2 coupled ferromagnetically (a) without an external magnetic field (h = 0) and (b) with an external magnetic field (h = 0.5).



Figure 3: Density of energy levels for N = 20 (A) in the ferromagnetic case and (B) in the antiferromagnetic case.





Figure 4: Density of energy levels for N = 100, $I \rightarrow \frac{I}{N}$ (A) for the ferromagnetic case and (B) for the antiferromagnetic case.



Figure 5: Density of energy levels for N = 1000, $I \rightarrow \frac{I}{N}$ (A) for the ferromagnetic case and (B) for the antiferromagnetic case.

3 The q-deformed KS model

As anticipated, the coalgebra symmetry of the KS model allows the construction of its q-deformation by making use of the quantum universal enveloping algebra $\mathcal{U}_q(\mathfrak{su}(2))$, where q is a real parameter (hereafter referred to as quantum $\mathfrak{su}_q(2)$ algebra), because the structure of its representation theory is exactly the same as that for $\mathfrak{su}(2)$. In the following, we recall the basic notions of this quantum algebra and present the construction of the q-KS model by imposing its $\mathfrak{su}_q(2)$ coalgebra symmetry.

3.1 The quantum algebra $su_q(2)$

The three generators of the $\mathfrak{su}_q(2)$ algebra satisfy the following commutation relations, where $q \in \mathbb{R}^+$ (see [23] and references therein):

$$[L_z, L_{\pm}] = \pm L_{\pm}, \qquad [L_+, L_-] = [2L_z]_q = \frac{q^{L_z} - q^{-L_z}}{q^{1/2} - q^{-1/2}}, \tag{52}$$

where we have made use of the q-symbol

$$[n]_q \coloneqq \frac{q^{n/2} - q^{-n/2}}{q^{1/2} - q^{-1/2}}.$$
(53)

We will frequently write $q = e^{\eta}$, which implies that

$$[2L_z]_q = \frac{q^{L_z} - q^{-L_z}}{q^{1/2} - q^{-1/2}} = \frac{e^{\eta L_z} - e^{-\eta L_z}}{e^{\eta/2} - e^{-\eta/2}} = 2L_z + o[\eta^2],$$
(54)

and we recover the $\mathfrak{su}(2)$ Lie algebra in the $q \to 1 \ (\eta \to 0)$ limit. We remark that as a consequence of (52), the following relations hold:

$$q^{L_z} L_{\pm} q^{-L_z} = q^{\pm 1} L_{\pm}.$$
 (55)

The coalgebra structure for the $\mathfrak{su}_q(2)$ algebra is generated by the deformed coproduct map

$$\Delta(L_{\pm}) = q^{-L_z/2} \otimes L_{\pm} + L_{\pm} \otimes q^{L_z/2}, \qquad \Delta(L_z) = \mathbf{I} \otimes L_z + L_z \otimes \mathbf{I}, \tag{56}$$

which is an algebraic homomorphism with respect to (52) and leads to the undeformed coproduct Δ_0 (21) in the limit $q \to 1$ ($\eta \to 0$). The generalisation to an arbitrary number N of tensor products is given by

$$\Delta^{(N)}(L_{\pm}) = L_{\pm} \otimes q^{L_z/2} \otimes q^{L_z/2} \otimes \dots^{N-1} \otimes q^{L_z/2} + q^{-L_z/2} \otimes L_{\pm} \otimes q^{L_z/2} \otimes \dots^{N-2} \otimes q^{L_z/2} + \dots + q^{-L_z/2} \otimes q^{-L_z/2} \otimes \dots^{N-1} \otimes q^{-L_z/2} \otimes L_{\pm},$$
(57)

and for the L_z generator, the deformed coproduct map coincides with the undeformed one, $\Delta^{(N)}(L_z) \equiv \Delta_0^{(N)}(L_z)$ (23).

Irreducible representations for $\mathfrak{su}_q(2)$ are defined through the action of the algebra generators onto an state $|j,m\rangle$ in the form

$$L_{\pm}|j,m\rangle := \sqrt{[j \mp m]_q [j \pm m + 1]_q} |j,m \pm 1\rangle, \qquad L_z|j,m\rangle = m|j,m\rangle.$$
(58)

Moreover, the deformed Casimir operator for the algebra (52) reads

$$C_q = L_- L_+ + [L_z]_q [L_z + \mathbf{I}]_q = L_+ L_- + [L_z]_q [L_z - \mathbf{I}]_q,$$
(59)

and the eigenvalues of C_q in the representation states $|j,m\rangle$ are just $[j]_q[j+1]_q$. Therefore, if q is not a root of unity, the representation theory of $\mathfrak{su}_q(2)$ is just structurally the same as that of $\mathfrak{su}(2)$ (see [23]).

Moreover, the tensor product of two $\mathfrak{su}_q(2)$ irreducible representations j_1 and j_2 is constructed using the deformed coproduct Δ , and its decomposition into irreducible modules with spin j is the same as in the undeformed case, as we obtain invariant subspaces with $j = (j_1 + j_2), \ldots, |j_1 - j_2|$. This result can obviously be generalised to the direct sum of N irreducible representations of $\mathfrak{su}(2)$ and constitutes the keystone for the exact solvability of the q-deformed KS model introduced in the following.

3.2 The generic *q*-Kittel Shore model

The generic q-KS Hamiltonian with a external magnetic field is defined by substituting the undeformed coproducts and Casimir operators in (28) by their q-deformed counterparts, namely

$$\tilde{H}_{KS}^{q} = -\frac{I}{2} \left(\Delta^{(N)}(C_q) - \sum_{i=1}^{N} C_q^{(i)} \right) - \gamma h \Delta^{(N)}(J_z) , \qquad (60)$$

where the q-deformed Casimir operator C_q is the $\mathfrak{su}_q(2)$ Casimir operator given by (59). Again, the deformed coproduct map $\Delta^{(N)}$ is an algebra homomorphism with respect to the defining relations of the $\mathfrak{su}_q(2)$ algebra (52)

$$\left[\Delta^{(N)}(C_q), \Delta^{(N)}(X)\right] = \Delta^{(N)}([C_q, X]) = 0, \qquad \forall X \in \mathfrak{su}_q(2).$$
(61)

Therefore, by defining the q-KS Hamiltonian in the form (60), we have preserved, by construction, the $\mathfrak{su}_q(2)$ coalgebra symmetry of the model. This implies that the Hamiltonian (60) commutes with $\Delta^{(N)}(L_z)$, which is also the total z-component of the q-deformed spin operators. Moreover, in the h = 0 case, the collective raising and lowering q-spin operators (57) also provide the commuting operators with H_{KS}^q .

As a consequence, we can mimic the construction performed for the $\mathfrak{su}(2)$ algebra. We start from a block-diagonal basis (10) for the tensor product of N irreducible representations of $\mathfrak{su}_q(2)$, where the operator $C_q^{(N)} \equiv \Delta^{(N)}(C_q)$ arising from the q-deformed coproduct and Casimir will be again represented as a block-diagonal matrix. Within each irreducible block defined by a set of basis vectors (30), the relations (61) imply that $C_q^{(N)}$ has to be proportional to the identity operator with eigenvalues

$$\Delta^{(N)}(C) |j_{\alpha_i}, m_{\alpha_i}\rangle = [j_{\alpha_i}]_q [j_{\alpha_i} + 1]_q |j_{\alpha_i}, m_{\alpha_i}\rangle, \qquad \forall \alpha_i = 1, \dots, k, \qquad m_{\alpha_i} = -j_{\alpha_i}, \dots, j_{\alpha_i}.$$
(62)

Also, the operator $\sum_{i=1}^{N} C_q^{(i)}$ acting on the tensor product basis (32) is just the identity times the factor $K_q = \sum_{i=1}^{N} [j_i]_q [(j_i+1)]_q$, and the eigenvectors of the q-KS Hamiltonian (60) are again the block-diagonal basis vectors (30) with eigenvalues given by

$$H_{KS}^{q}|j_{\alpha_{i}},m_{\alpha_{i}}\rangle = -\frac{I}{2} \left(\Delta^{(N)}(C_{q}) - \sum_{i=1}^{N} C_{q}^{(i)} \right) |j_{\alpha_{i}},m_{\alpha_{i}}\rangle = -\frac{I}{2} \left([j_{\alpha_{i}}]_{q} [j_{\alpha_{i}} + 1]_{q} - K_{q} \right) |j_{\alpha_{i}},m_{\alpha_{i}}\rangle.$$
(63)

Finally, when $h \neq 0$, since the coproduct for the L_z generator is not deformed, by following the same reasoning as in the non-deformed case we have that

$$-\gamma h \Delta^{(N)}(L_z) |j_{\alpha_i}, m_{\alpha_i}\rangle = -\gamma h m_{\alpha_i} |j_{\alpha_i}, m_{\alpha_i}\rangle.$$
(64)

In conclusion, the eigenvalues of the generic q-KS model for N arbitrary spins are

$$\tilde{E}_{q}(j_{\alpha_{i}}, m_{\alpha_{i}}, K_{q}) = -\frac{I}{2} \left([j_{\alpha_{i}}]_{q} [j_{\alpha_{i}} + 1]_{q} - K_{q} \right) - \gamma h m_{\alpha_{i}}.$$
(65)

Note that when comparing this expression with (65), all numbers are substituted by q-numbers, except for the magnetic field term. In addition, the ground-state eigenvalue can be set to zero for any N by subtracting the ground-state energy from all the eigenvalues.

Let us make the abstract definition (60) of the q-KS model more explicit in terms of the $\mathfrak{su}_q(2)$ generators. A first step consists in rewriting it in the form (66), namely

$$\tilde{H}_{KS}^{q} = -\frac{I}{2} \left(\Delta^{(N)}(L_{-}) \Delta^{(N)}(L_{+}) + \left[\Delta^{(N)}(L_{z}) \right]_{q} \left[\Delta^{(N)}(L_{z}) + \Delta^{(N)}(\mathbf{I}) \right]_{q} - \sum_{i=1}^{N} C_{q}^{(i)} \right) - \gamma h \Delta^{(N)}(L_{z}).$$
(66)

Now we consider the explicit form of the N-th deformed coproducts (57)

$$\Delta^{(N)}(L_{\pm}) = \sum_{i=1}^{N} \exp\left[-\frac{\eta}{2} \sum_{j=1}^{i-1} L_{z}^{(j)}\right] L_{\pm}^{(i)} \exp\left[\frac{\eta}{2} \sum_{h=i+1}^{N} L_{z}^{(h)}\right], \qquad \Delta^{(N)}(L_{z}) = \sum_{i=1}^{N} L_{z}^{(i)}, \qquad (67)$$

where the superindex *i* represents the spin for the *i*-th particle and we have used that $q = e^{\eta}$ (which, as we will see, will be useful in the following). Expanding the Hamiltonian (66), we find the expression

$$\tilde{H}_{KS}^{q} = -\frac{I}{2} \left(\sum_{i=1}^{N} \exp\left[-\eta \sum_{j=1}^{i-1} L_{z}^{(j)} \right] \left(C_{q}^{(i)} - [L_{z}^{(i)}]_{q} \left[L_{z}^{(i)} + \mathbf{I}^{(i)} \right]_{q} \right) \exp\left[\eta \sum_{h=i+1}^{N} L_{z}^{(h)} \right] + \sum_{i=1}^{N-1} \sum_{k=i+1}^{N} \prod_{t=1}^{i-1} \exp\left[-\eta L_{z}^{(t)} \right] \right) \\
+ \exp\left[-\eta \frac{L_{z}^{(i)}}{2} \right] \left(L_{+}^{(i)} L_{-}^{(k)} + L_{-}^{(i)} L_{+}^{(k)} \right) \exp\left[\eta \frac{L_{z}^{(k)}}{2} \right] \prod_{r=k+1}^{N} \exp\left[\eta L_{z}^{(r)} \right] + \left[\sum_{i=1}^{N} L_{z}^{(i)} \right]_{q} \left[\sum_{s=1}^{N} L_{z}^{(s)} + \mathbf{I}^{(N)} \right]_{q} - \sum_{i=1}^{N} C_{q}^{(i)} \right) - \gamma h \sum_{i=1}^{N} L_{z}^{(i)}.$$
(68)

Now, by taking into account that $q^{L_z/2}L_{\pm}q^{-L_z/2} = q^{\pm 1/2}L_{\pm}$, we obtain the final expression

$$\tilde{H}_{KS}^{q} = -\frac{I}{2} \left(\sum_{i=1}^{N} \exp\left[-\eta \sum_{j=1}^{i-1} L_{z}^{(j)} \right] L_{-}^{(i)} L_{+}^{(i)} \exp\left[\eta \sum_{h=i+1}^{N} L_{z}^{(h)} \right] + \sum_{i=1}^{N-1} \sum_{r=i+1}^{N} \left(e^{\eta/2} L_{-}^{(i)} L_{+}^{(r)} + e^{-\eta/2} L_{+}^{(i)} L_{-}^{(r)} \right) \cdot \\
\cdot \exp\left[-\eta \frac{L_{z}^{(i)}}{2} \right] \cdot \exp\left[\eta \frac{L_{z}^{(r)}}{2} \right] \cdot \prod_{t=1}^{i-1} \exp\left[-\eta L_{z}^{(t)} \right] \prod_{k=r+1}^{N} \exp\left[\eta L_{z}^{(k)} \right] + \left[\sum_{i=1}^{N} L_{z}^{(i)} \right]_{q} \left[\sum_{s=1}^{N} L_{z}^{(s)} + \mathbf{I}^{(N)} \right]_{q} - \\
- \sum_{i=1}^{N} C_{q}^{(i)} \right) - \gamma h \sum_{i=1}^{N} L_{z}^{(i)},$$
(69)

which, for q = 1 (equivalently, $\eta = 0$), leads to the undeformed KS Hamiltonian (25).

Note that the q-KS Hamiltonian can be also written in terms of the L_x , L_y , and L_z spin operators, where we consider the same change of basis as in the non-deformed case, namely

$$L_x = \frac{1}{2}(L_+ + L_-), \qquad L_y = -\frac{i}{2}(L_+ - L_-).$$
 (70)

In this way, we have that

$$\tilde{H}_{KS}^{q} = -\frac{I}{2} \left(\Delta(L_{x})^{2} + \Delta(L_{y})^{2} - \frac{1}{2} [2\Delta(L_{z})]_{q} + [\Delta(L_{z})]_{q} [\Delta(L_{z}) + \Delta(\mathbf{I})]_{q} - \sum_{i=1}^{N} C_{q}^{(i)} \right) - \gamma h \Delta^{(N)}(L_{z}),$$
(71)

which leads to the final expression

$$\begin{split} \tilde{H}_{KS}^{q(N)} &= -\frac{I}{2} \left(\sum_{i=1}^{N} \exp\left[-\eta \sum_{j=1}^{i-1} L_{z}^{(j)} \right] \left((L_{x}^{(i)})^{2} + (L_{y}^{(i)})^{2} - \frac{1}{2} [2L_{z}^{(i)}]_{q} \right) \exp\left[\eta \sum_{h=i+1}^{N} L_{z}^{(h)} \right] + \sum_{i=1}^{N-1} \sum_{l=i+1}^{N} \left[\left(L_{x}^{(i)} L_{x}^{(l)} + L_{y}^{(l)} L_{y}^{(l)} \right) \left(e^{\eta/2} - e^{-\eta/2} \right) \right] \exp\left[-\eta \frac{L_{z}^{(i)}}{2} \right] \cdot \exp\left[\eta \frac{L_{z}^{(l)}}{2} \right] \cdot \\ & \cdot \prod_{t=1}^{i-1} \exp\left[-\eta L_{z}^{(t)} \right] \prod_{k=l+1}^{N} \exp\left[\eta L_{z}^{(k)} \right] + \left[\sum_{i=1}^{N} L_{z}^{(i)} \right]_{q} \left[\sum_{s=1}^{N} L_{z}^{(s)} + \mathbf{I}^{(N)} \right]_{q} - \sum_{i=1}^{N} C_{q}^{(i)} \right) - \gamma h \sum_{i=1}^{N} L_{z}^{(i)}. \end{split}$$

$$(72)$$

As can be seen from the previous expressions, the explicit form of the deformation in terms of the quantum algebra generators is far from simple. Nevertheless, the shape of the deformation is absolutely dictated by the $\mathfrak{su}_q(2)$ symmetry to preserve all the integrability properties of the KS model and, despite its complexity, leads to the simple form (65) for its eigenvalues, which preserves the same degeneracies as in the undeformed model but becomes non-linearly modified in terms of the quantum deformation parameter η .

4 The spin-1/2 q-KS model

4.1 Eigenvectors and eigenvalues

Taking into account that the 2-dimensional j = 1/2 fundamental irreducible representation (58) for $\mathfrak{su}_q(2)$ is given by the undeformed Pauli matrices (38), this means that the q-KS model for N copies of the $\mathfrak{su}_q(2)$ algebra gives rise to a spin Hamiltonian whose explicit form is obtained by substituting

$$L_{\pm}^{(i)} = \sigma_{\pm}^{(i)}, \qquad L_{z}^{(i)} = \frac{1}{2}\sigma_{z}^{(i)}, \qquad i = 1, \dots, N,$$
(73)

within (69), where $e^{\eta} = q$, namely

$$\widetilde{H}_{KS}^{q(N)} = -\frac{I}{2} \left(\sum_{i=1}^{N} \exp\left[-\eta \sum_{j=1}^{i-1} \frac{1}{2} \sigma_{z}^{(j)} \right] \sigma_{-}^{(i)} \sigma_{+}^{(i)} \exp\left[\eta \sum_{h=i+1}^{N} \frac{1}{2} \sigma_{z}^{(h)} \right] \right. \\
\left. + \sum_{i=1}^{N-1} \sum_{r=i+1}^{N} \left(e^{\eta/2} \sigma_{-}^{(i)} \sigma_{+}^{(r)} + e^{-\eta/2} \sigma_{+}^{(i)} \sigma_{-}^{(r)} \right) \cdot \\
\left. \cdot \exp\left[-\eta \frac{\sigma_{z}^{(i)}}{4} \right] \cdot \exp\left[\eta \frac{\sigma_{z}^{(r)}}{4} \right] \cdot \prod_{t=1}^{i-1} \exp\left[-\frac{\eta}{2} \sigma_{z}^{(t)} \right] \prod_{k=r+1}^{N} \exp\left[\frac{\eta}{2} \sigma_{z}^{(k)} \right] \\
\left. + \left[\sum_{i=1}^{N} \frac{1}{2} \sigma_{z}^{(i)} \right]_{q} \left[\sum_{s=1}^{N} \frac{1}{2} \sigma_{z}^{(s)} + \mathbf{I}^{(N)} \right]_{q} - N \left[1/2 \right]_{q} \left[3/2 \right]_{q} \right) - \gamma h \sum_{i=1}^{N} \frac{1}{2} \sigma_{z}^{(i)}.$$
(74)

We stress that this is a new integrable spin model with long-range interactions that, in contrast to the undeformed KS model, presents nonidentical couplings among different spins, which are determined by the deformed coproducts. Nevertheless, the underlying $\mathfrak{su}_q(2)$ coalgebra symmetry of the model makes

that these involved interactions are such that the eigenvalues of this model are can be straightforwardly deduced from (65) and read

$$\tilde{E}_{q}(j_{\alpha_{i}}, m_{\alpha_{i}}, K_{q}) = -\frac{1}{2} \left([j_{\alpha_{i}}]_{q} [j_{\alpha_{i}} + 1]_{q} - N [1/2]_{q} [3/2]_{q} \right) - \gamma h m_{\alpha_{i}},$$
(75)

which should be compared to (35). On the other hand, eigenvectors for N = 2, 3, 4, ... can be obtained by applying in a recurrent way the well-known expressions for the *q*-Clebsch–Gordan coefficients (see [24] and references therein)

$$CG[j_{1}, m_{1}, j_{2}, m_{2}, j, m] = \delta_{m, m_{1} + m_{2}} q^{\frac{1}{2}(j_{1}m_{2} - j_{2}m_{1}) - \frac{1}{4}(-j + j_{1} + j_{2})(j + j_{1} + j_{2} + 1)} \\ \sqrt{\frac{[2j + 1]_{q}[j + m]_{q}![j_{2} - m_{2}]_{q}![j + j_{1} - j_{2}]_{q}![-j + j_{1} + j_{2}]_{q}![j + j_{1} + j_{2} + 1]_{q}!}{[j - m]_{q}![j_{1} - m_{1}]_{q}![j_{1} + m_{1}]_{q}![j_{2} + m_{2}]_{q}![j - j_{1} + j_{2}]_{q}!}} \\ \sum_{n=0}^{\min(-j+j_{1}+j_{2},j_{2}-m_{2})} \frac{[2j_{2} - n]_{q}!(-1)^{-j+j_{1}+j_{2}+n}q^{\frac{1}{2}n(j_{1}+m_{1})}[j_{1} + j_{2} - m - n]_{q}!}{[n]_{q}![j_{2} - m_{2} - n]_{q}![-j + j_{1} + j_{2} - n]_{q}![j + j_{1} + j_{2} - n + 1]_{q}!}$$

$$(76)$$

As a consequence, the partition function Z_N^q of this model with h = 0 is directly obtained from (49), and reads

$$Z_N^q = \sum_J \sum_{m_J=-J}^{m_J=+J} d_{NJ\frac{1}{2}} \exp(-\beta E'_q(J, m_J, K_q)),$$
(77)

where $E'_q(J, m_J, K_q)$ is obtained from (75) with h = 0 and the degeneracies $d_{NJ\frac{1}{2}}$ are the same as those in the undeformed model (43) because of the identical structure of the representation theory of $\mathfrak{su}_q(2)$ and $\mathfrak{su}(2)$. From this expression, all the thermodynamic properties of the *q*-deformed model can be derived and studied. As in the undeformed case, the energy levels E'_q are obtained by subtracting the energy of the ground state.

Regarding the density of the (corrected) energy levels of the q-deformed model, we can easily see that the deformation enlarges the distance between energy states, as shown in Fig. 6, which plots the densities of the energy levels for the deformed model with N = 20,100,1000 and different values of η , because the larger N is, the smaller η must be considered in order to obtain reasonable values for the energies. We observe that, as expected, also in the deformed case the two distributions corresponding to the ferromagnetic and anti-ferromagnetic models are specular images one of the other.



Figure 6: Density of (corrected) energy levels for N = 20 as a function of the parameter $\eta = 0$ (red) and $\eta = 0.2$ (dashed blue) (A) for the ferromagnetic case and (B) for the antiferromagnetic case.



Figure 7: Density of (corrected) energy levels for N = 100 as a function of the parameter $\eta = 0$ (red) and $\eta = 0.02$ (dashed blue) (A) for the ferromagnetic case and (B) for the antiferromagnetic case.



Figure 8: Density of (corrected) energy levels for N = 1000 as a function of the parameter $\eta = 0$ (red) and $\eta = 0.005$ (blue) (A) for the ferromagnetic case and (B) for the antiferromagnetic case.

4.2 The N = 2, 3, 4 cases

To illustrate the main features of the spin-1/2 q-KS model, an explicit study of the N = 2, 3, 4 cases is required. The eigenvalues of the q-KS antiferromagnetic and ferromagnetic Hamiltonians without an external magnetic field (h = 0) are plotted in Figs. 9 and 10 together with their undeformed counterparts. Thus, the nonlinear effect of the deformation in the energy gaps between consecutive eigenvalues of the model can be clearly observed. The degeneracies for each energy level are not modified with respect to the undeformed KS Hamiltonian, and the addition of a constant external magnetic field would generate the same uniform Zeeman splittings as in the q = 1 case. Indeed, the antiferromagnetic and ferromagnetic spectra are inverted versions of the very same energy level structure.

The eigenvectors for the N = 2, 3, 4 q-KS model can be computed using the q-Clebsch–Gordan coefficients (76) and are presented in the following. In this way, the role that the q-Dicke states [25, 26] play in the model is shown, which provides a basis for the fundamental eigenspace in the ferromagnetic case and for the most excited state in the antiferromagnetic case. The remaining eigenstates are no longer q-Dicke eigenstates and are also given and analysed explicitly.

4.2.1 The N = 2 case and the *q*-exchange operator

In this case the q-KS Hamiltonian of two interacting spins without magnetic field is given, in the fundamental representation (73), by

$$H_{KS}^{q(2)} = -\frac{I}{2} \left(\sigma_{-}^{(1)} \sigma_{+}^{(1)} \exp\left[\eta \frac{1}{2} \sigma_{z}^{(2)}\right] + \exp\left[-\eta \frac{1}{2} \sigma_{z}^{(1)}\right] \sigma_{-}^{(2)} \sigma_{+}^{(2)} + \left(e^{\eta/2} \sigma_{-}^{(1)} \sigma_{+}^{(2)} + e^{-\eta/2} \sigma_{+}^{(1)} \sigma_{-}^{(2)}\right) \cdot \exp\left[-\eta \frac{\sigma_{z}^{(1)}}{4}\right] \cdot \exp\left[\eta \frac{\sigma_{z}^{(2)}}{4}\right] + \left[\sum_{i=1}^{2} \frac{1}{2} \sigma_{z}^{(i)}\right]_{q} \left[\sum_{s=1}^{2} \frac{1}{2} \sigma_{z}^{(s)} + \mathbf{I}^{(2)}\right]_{q} - 2\left[1/2\right]_{q}\left[3/2\right]_{q}\right).$$
(78)

Obviously, the $\eta \to 0$ limit of this expression is the N = 2 KS model rewritten in the form

$$H_{KS}^{(2)} = -\frac{I}{2} \left(\left(\sigma_{-}^{(1)} \sigma_{+}^{(1)} + \sigma_{-}^{(2)} \sigma_{+}^{(2)} + \sigma_{-}^{(1)} \sigma_{+}^{(2)} + \sigma_{+}^{(1)} \sigma_{-}^{(2)} \right) + \left(\sum_{i=1}^{2} \frac{1}{2} \sigma_{z}^{(i)} \right)^{2} + \left(\sum_{s=1}^{2} \frac{1}{2} \sigma_{z}^{(s)} \right) - 3/2 \right)$$

$$= -I \vec{\sigma}_{1} \cdot \vec{\sigma}_{2} .$$
(79)

The structural complexity of the q-deformation can be appreciated by comparing the latter expression with (78). In fact, the operator (78) can be taken as the definition of the q-exchange interaction induced by the deformed coalgebra structure of $\mathfrak{su}_q(2)$. In matrix form, the operator (78) reads

$$H_{KS}^{q(2)} = -\frac{I}{\left(\sqrt{q}+1\right)^2} \left(\begin{pmatrix} \frac{q^2+3}{2\sqrt{q}} & 0 & 0 & 0\\ 0 & \frac{1-q}{2\sqrt{q}} & \frac{1}{2} & 0\\ 0 & \frac{1}{2} & \frac{(q-1)\sqrt{q}}{2} & 0\\ 0 & 0 & 0 & \frac{q^2+3}{2\sqrt{q}} \end{pmatrix} - \begin{pmatrix} \frac{1}{\sqrt{q}} & 0 & 0 & 0\\ 0 & q & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & \frac{1}{\sqrt{q}} \end{pmatrix} \right),$$
(80)

which in the limit $q \to 1$ provides the usual form of the exchange operator (79):

$$H_{KS}^{q(2)} = -\frac{I}{2} \left(\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} - \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \right) .$$
(81)

In the ferromagnetic case, the ground state for (78) with h = 0 can be straightforwardly shown to be the J = 1 subspace, which is generated by the q-triplet

$$|1,-1\rangle = |\downarrow\downarrow\rangle, \qquad |1,0\rangle = \frac{1}{\sqrt{2q}} \left(q^{1/4} |\downarrow\uparrow\rangle + q^{-1/4} |\uparrow\downarrow\rangle \right), \qquad |1,1\rangle = |\uparrow\uparrow\rangle, \tag{82}$$

and the singlet with j = 0 is the only excited state

$$|0,0\rangle = \frac{1}{\sqrt{2q}} \left(-q^{-1/4} |\downarrow\uparrow\rangle + q^{1/4} |\uparrow\downarrow\rangle \right).$$
(83)

The situation is reversed in the antiferromagnetic case, and when the magnetic field $h \neq 0$ is included, the triplet becomes nondegenerate.

The states (82) are just the q-Dicke states for j = 1. The states $|1,1\rangle$ and $|1,-1\rangle$ are factorised, where $|1,0\rangle$ and $|0,0\rangle$ are q-Bell states whose entanglement is not maximal unless $q \to 1$ (the limit of the undeformed Bell states). However, the limit $q \to \infty$ gives rise to factorised states, thus showing that the q-deformation of the model (or, equivalently, the q-exchange operator (78)) induces a loss of entanglement governed by the deformation parameter q.



Figure 9: Energy levels (in the units |I|/4) for N = 2, 3, 4, 5 spins j = 1/2 coupled antiferromagnetically for values of the deformation parameter $\eta = 0, 0.5, 0.75$.



Figure 10: Energy levels (in the units |I|/4) for N = 2, 3, 4, 5 spins j = 1/2 coupled ferromagnetically for values of the deformation parameter $\eta = 0, 0.5, 0.75$.

4.2.2 The N = 3 case

When three spins are considered, the q-KS model in the fundamental representation (73) and with h = 0 becomes

$$\begin{aligned} H_{KS}^{q(3)} &= -\frac{I}{2} \Bigg(\sigma_{-}^{(1)} \sigma_{+}^{(1)} \exp\left[\eta \frac{1}{2} (\sigma_{z}^{(2)} + \sigma_{z}^{(3)}) \right] + \exp\left[-\eta \frac{1}{2} \sigma_{z}^{(1)} \right] \sigma_{-}^{(2)} \sigma_{+}^{(2)} \exp\left[\eta \frac{1}{2} \sigma_{z}^{(3)} \right] + \\ &+ \exp\left[-\eta \frac{1}{2} (\sigma_{z}^{(1)} + \sigma_{z}^{(2)}) \right] \sigma_{-}^{(3)} \sigma_{+}^{(3)} + \left(e^{\eta/2} \sigma_{-}^{(1)} \sigma_{+}^{(2)} + e^{-\eta/2} \sigma_{+}^{(1)} \sigma_{-}^{(2)} \right) \exp\left[-\eta \frac{\sigma_{z}^{(1)}}{4} \right] \cdot \\ &\cdot \exp\left[\eta \frac{\sigma_{z}^{(2)}}{4} \right] \cdot \exp\left[\frac{\eta}{2} \sigma_{z}^{(3)} \right] + \left(e^{\eta/2} \sigma_{-}^{(1)} \sigma_{+}^{(3)} + e^{-\eta/2} \sigma_{+}^{(1)} \sigma_{-}^{(3)} \right) \exp\left[-\eta \frac{\sigma_{z}^{(1)}}{4} \right] \cdot \\ &\cdot \exp\left[\eta \frac{\sigma_{z}^{(3)}}{4} \right] + \left(e^{\eta/2} \sigma_{-}^{(2)} \sigma_{+}^{(3)} + e^{-\eta/2} \sigma_{+}^{(2)} \sigma_{-}^{(3)} \right) \exp\left[-\eta \frac{\sigma_{z}^{(2)}}{4} \right] \exp\left[\eta \frac{\sigma_{z}^{(3)}}{4} \right] \cdot \\ &\cdot \exp\left[-\eta \frac{\sigma_{z}^{(1)}}{2} \right] + \left[\sum_{i=1}^{3} \frac{1}{2} \sigma_{z}^{(i)} \right]_{q} \left[\sum_{s=1}^{3} \frac{1}{2} \sigma_{z}^{(s)} + \mathbf{I}^{(3)} \right]_{q} - 3 \left[1/2 \right]_{q} \left[3/2 \right]_{q} \right), \end{aligned} \tag{84}$$

which is not a straightforward superposition of q-exchange operators (78), and is indeed much more involved than its underformed counterpart

$$H_{KS}^{(3)} = -I\left(\vec{\sigma}_1 \cdot \vec{\sigma}_2 + \vec{\sigma}_1 \cdot \vec{\sigma}_3 + \vec{\sigma}_2 \cdot \vec{\sigma}_3\right).$$
(85)

The eigenstates can be obtained using the q-Clebsch–Gordan coefficients (76) to compose the J = 1and J = 0 representations with an additional spin 1/2 representation. As a result, in the ferromagnetic case the ground states generate a J = 3/2 quadruplet, given by the q-Dicke states

$$|3/2, -3/2\rangle = |\downarrow\downarrow\downarrow\rangle, \qquad |3/2, -1/2\rangle = \frac{1}{\sqrt{3q}} \left(q^{1/2} |\downarrow\downarrow\uparrow\rangle + |\downarrow\uparrow\downarrow\rangle + q^{-1/2} |\uparrow\downarrow\downarrow\rangle \right),$$

$$|3/2, 1/2\rangle = \frac{1}{\sqrt{3q}} \left(q^{1/2} |\downarrow\uparrow\uparrow\rangle + |\uparrow\downarrow\uparrow\rangle + q^{-1/2} |\uparrow\uparrow\downarrow\rangle \right), \qquad |3/2, 3/2\rangle = |\uparrow\uparrow\uparrow\rangle,$$
(86)

in which again q governs the entanglement properties of the q-KS eigenstates. The four (degenerate) excited states are provided by two J = 1/2 doublets, given by

$$|1/2, -1/2\rangle_{1} = \frac{q^{-1/4}}{\sqrt{2q^{3}q}} \left(q^{1/2} |\uparrow\downarrow\downarrow\rangle + q |\downarrow\uparrow\downarrow\rangle - 2_{q} |\downarrow\downarrow\uparrow\rangle \right),$$

$$|1/2, 1/2\rangle_{1} = \frac{q^{1/4}}{\sqrt{2q^{3}q}} \left(2_{q} |\uparrow\uparrow\downarrow\rangle - q^{-1} |\uparrow\downarrow\uparrow\rangle - q^{-1/2} |\downarrow\uparrow\uparrow\rangle \right),$$
(87)

and

$$|1/2, -1/2\rangle_2 = \frac{1}{\sqrt{2q}} \left(q^{1/4} |\uparrow\downarrow\downarrow\rangle - q^{-1/4} |\downarrow\uparrow\downarrow\rangle \right),$$

$$|1/2, 1/2\rangle_2 = \frac{1}{\sqrt{2q}} \left(-q^{-1/4} |\downarrow\uparrow\uparrow\rangle + q^{1/4} |\uparrow\downarrow\uparrow\rangle \right).$$

(88)

These two doublets span the subspace of the ground states in the antiferromagnetic case.

4.2.3 The N = 4 case

Starting from the basis vectors of the J = 3/2 and J = 1/2 irreducible representations of quantum $\mathfrak{su}(2)$ obtained in the N = 3 case, and using the q-Clebsch–Gordan coefficients providing their composition with another spin 1/2 representation, we obtain the eigenvectors of the q-KS model in the N = 4 case.

The q-Dicke states are now given by a quintuplet with J = 2, namely

$$\begin{aligned} |2,-2\rangle &= |\downarrow\downarrow\downarrow\downarrow\rangle, \\ |2,-1\rangle &= \frac{1}{\sqrt{4_q}} \left(q^{3/4} |\downarrow\downarrow\downarrow\uparrow\uparrow\rangle + q^{1/4} |\downarrow\downarrow\uparrow\downarrow\rangle + q^{-1/4} |\downarrow\uparrow\downarrow\downarrow\rangle + q^{-3/4} |\uparrow\downarrow\downarrow\downarrow\rangle \right), \\ |2,0\rangle &= \frac{\sqrt{2_q}}{\sqrt{3_q 4_q}} \left(q |\downarrow\downarrow\uparrow\uparrow\uparrow\rangle + q^{1/2} |\downarrow\uparrow\downarrow\uparrow\rangle + |\downarrow\uparrow\uparrow\downarrow\rangle + |\uparrow\downarrow\downarrow\uparrow\rangle + q^{-1/2} |\uparrow\downarrow\uparrow\downarrow\rangle + q^{-1} |\uparrow\uparrow\downarrow\downarrow\rangle \right), \end{aligned}$$
(89)
$$|2,1\rangle &= \frac{1}{\sqrt{4_q}} \left(q^{3/4} |\downarrow\uparrow\uparrow\uparrow\uparrow\rangle + q^{1/4} |\uparrow\downarrow\uparrow\uparrow\rangle + q^{-1/4} |\uparrow\uparrow\downarrow\uparrow\rangle + q^{-3/4} |\uparrow\uparrow\uparrow\downarrow\rangle \right), \\ |2,2\rangle &= |\uparrow\uparrow\uparrow\uparrow\rangle. \end{aligned}$$

We also obtain three triplets with J = 1 given by

$$\begin{split} |1,-1\rangle_{1} &= \frac{1}{\sqrt{3}_{q}4_{q}} \left(-q^{-1/4}3_{q} |\downarrow\downarrow\downarrow\uparrow\rangle + q^{5/4} |\downarrow\downarrow\uparrow\downarrow\rangle + q^{3/4} |\downarrow\uparrow\downarrow\downarrow\rangle + q^{1/4} |\uparrow\downarrow\downarrow\downarrow\rangle \right), \tag{90} \\ |1,0\rangle_{1} &= \frac{\sqrt{2}_{q}}{\sqrt{3}_{q}4_{q}} \left(-|\downarrow\downarrow\uparrow\uparrow\uparrow\rangle - q^{-1/2} |\downarrow\uparrow\downarrow\uparrow\downarrow\rangle + q|\downarrow\uparrow\uparrow\downarrow\downarrow\rangle - q^{-1} |\uparrow\downarrow\downarrow\uparrow\rangle + q^{1/2} |\uparrow\downarrow\downarrow\downarrow\rangle + |\uparrow\uparrow\downarrow\downarrow\rangle \right), \tag{90} \\ |1,1\rangle_{1} &= \frac{1}{\sqrt{3}_{q}4_{q}} \left(-q^{-1/4} |\downarrow\uparrow\uparrow\uparrow\uparrow\rangle - q^{-3/4} |\uparrow\downarrow\uparrow\uparrow\uparrow\rangle - q^{-5/4} |\uparrow\uparrow\downarrow\downarrow\rangle + 3_{q}q^{1/4} |\uparrow\uparrow\uparrow\downarrow\rangle \right), \\ |1,-1\rangle_{2} &= \frac{q^{-1/4}}{\sqrt{2}q^{3}q} \left(-2_{q} |\downarrow\downarrow\uparrow\downarrow\downarrow\rangle + q|\downarrow\uparrow\downarrow\downarrow\downarrow\rangle + q^{1/2} |\uparrow\downarrow\downarrow\downarrow\rangle \right), \\ |1,0\rangle_{2} &= \frac{1}{2_{q}\sqrt{3}_{q}} \left(-2_{q} |\downarrow\downarrow\uparrow\uparrow\downarrow\rangle + q|\downarrow\uparrow\downarrow\downarrow\uparrow\rangle - q^{-1/2} |\downarrow\uparrow\uparrow\downarrow\downarrow\rangle + q^{1/2} |\uparrow\downarrow\downarrow\downarrow\uparrow\rangle - q^{-1} |\uparrow\downarrow\uparrow\downarrow\downarrow\rangle + 2_{q} |\uparrow\uparrow\downarrow\downarrow\rangle \right), \tag{91} \\ |1,1\rangle_{2} &= \frac{q^{1/4}}{\sqrt{2}q^{3}q} \left(-q^{-1/2} |\downarrow\uparrow\uparrow\uparrow\uparrow\rangle - q^{-1} |\uparrow\downarrow\uparrow\uparrow\downarrow\rangle + 2_{q} |\uparrow\uparrow\downarrow\downarrow\rangle \right), \\ |1,0\rangle_{3} &= \frac{1}{2_{q}} \left(-q^{-1/4} |\downarrow\uparrow\uparrow\downarrow\uparrow\rangle + q^{1/4} |\uparrow\downarrow\downarrow\downarrow\rangle \right), \\ |1,0\rangle_{3} &= \frac{1}{2_{q}} \left(-q^{-1/4} |\downarrow\uparrow\uparrow\uparrow\uparrow\rangle + q^{1/4} |\uparrow\downarrow\downarrow\downarrow\rangle \right), \tag{92} \\ |1,1\rangle_{3} &= \frac{1}{\sqrt{2q}} \left(-q^{-1/4} |\downarrow\uparrow\uparrow\uparrow\uparrow\rangle + q^{1/4} |\uparrow\downarrow\downarrow\uparrow\uparrow\rangle \right), \end{aligned}$$

and two singlets with J = 0

$$|0,0\rangle_{1} = \frac{1}{2q\sqrt{3q}} \left(q^{-1/2} 2_{q} |\downarrow\downarrow\uparrow\uparrow\rangle - q^{1/2} |\downarrow\uparrow\downarrow\uparrow\rangle - |\downarrow\uparrow\uparrow\downarrow\rangle - |\uparrow\downarrow\downarrow\uparrow\rangle - q^{-1/2} |\uparrow\downarrow\uparrow\downarrow\rangle + q^{1/2} 2_{q} |\uparrow\uparrow\downarrow\downarrow\rangle \right), \quad (93)$$

$$|0,0\rangle_2 = \frac{1}{2q} \left(q^{-1/2} |\downarrow\uparrow\downarrow\uparrow\rangle - |\downarrow\uparrow\uparrow\downarrow\rangle - |\uparrow\downarrow\downarrow\uparrow\rangle + q^{1/2} |\uparrow\downarrow\uparrow\downarrow\rangle \right).$$
(94)

The latter singlet states are the (degenerate) antiferromagnetic ground states, which contain the Neel states $|\uparrow\downarrow\uparrow\downarrow\rangle$ and $|\downarrow\uparrow\downarrow\uparrow\rangle$ when taking the limit $\eta \to \pm\infty$. Similarly, the ground states for the ferromagnetic

case contain the typical ferromagnetic configurations $|\uparrow\uparrow\uparrow\uparrow\rangle$ and $|\downarrow\downarrow\downarrow\downarrow\downarrow\rangle\rangle$ of the XXX Heisenberg model (the corresponding states do not depend on q), and when the magnetic field h > 0 is turned on, the state $|\psi_{2,-2}^4\rangle = |\uparrow\uparrow\uparrow\uparrow\rangle$ is the ground state. Indeed, the explicit eigenstates for the undeformed KS model are trivially obtained within the limit $q \to 1$.

5 Higher spin *q*-KS models

The generic q-KS model (60) is expressed in terms of $\mathfrak{su}_q(2)$ generators, which only coincide with $\mathfrak{su}(2)$ spin operators in the j = 1/2 representation. However, higher spin q-KS models can be obtained using the known relationships between both sets of generators for arbitrary j.

5.1 The q-KS model for j = 1

For j = 1, the fundamental representation of the undeformed angular momenta operators is obtained by applying (8), namely

$$J_{+} = \begin{pmatrix} 0 & \sqrt{2} & 0 \\ 0 & 0 & \sqrt{2} \\ 0 & 0 & 0 \end{pmatrix} \equiv \Sigma_{+}, \qquad J_{-} = \begin{pmatrix} 0 & 0 & 0 \\ \sqrt{2} & 0 & 0 \\ 0 & \sqrt{2} & 0 \end{pmatrix} \equiv \Sigma_{-}, \qquad J_{z} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} \equiv \Sigma_{z}. \tag{95}$$

In the q-deformed case, the j = 1 representation (58) of the $\mathfrak{su}_q(2)$ generators is given by

$$L_{+} = \begin{pmatrix} 0 & \sqrt{[2]_{q}} & 0\\ 0 & 0 & \sqrt{[2]_{q}} \\ 0 & 0 & 0 \end{pmatrix}, \qquad L_{-} = \begin{pmatrix} 0 & 0 & 0\\ \sqrt{[2]_{q}} & 0 & 0\\ 0 & \sqrt{[2]_{q}} & 0 \end{pmatrix}, \qquad L_{z} = \begin{pmatrix} 1 & 0 & 0\\ 0 & 0 & 0\\ 0 & 0 & -1 \end{pmatrix}, \tag{96}$$

and we realise that the quantum group generators are just proportional to the non-deformed j = 1 spin operators (95):

$$L_{\pm} = \sqrt{\frac{[2]_q}{2}} \Sigma_{\pm}, \qquad L_z = \Sigma_z \,. \tag{97}$$

Therefore, by substituting (97) in the generic expression of the q-KS model given in (69) we obtain the following Hamiltonian

$$\tilde{H}_{KS}^{q(N)} = -\frac{I}{2} \left(\sum_{i=1}^{N} \exp\left[-\eta \sum_{j=1}^{i-1} \Sigma_{z}^{(j)} \right] \frac{[2]_{q}}{2} \Sigma_{-}^{(i)} \Sigma_{+}^{(i)} \exp\left[\eta \sum_{h=i+1}^{N} \Sigma_{z}^{(h)} \right] \right. \\
\left. + \sum_{i=1}^{N-1} \sum_{r=i+1}^{N} \frac{[2]_{q}}{2} \left(e^{\eta/2} \Sigma_{-}^{(i)} \Sigma_{+}^{(r)} + e^{-\eta/2} \Sigma_{+}^{(i)} \Sigma_{-}^{(r)} \right) \cdot \\
\left. \cdot \exp\left[-\eta \frac{\Sigma_{z}^{(i)}}{2} \right] \cdot \exp\left[\eta \frac{\Sigma_{z}^{(r)}}{2} \right] \cdot \prod_{t=1}^{i-1} \exp\left[-\eta \Sigma_{z}^{(t)} \right] \prod_{k=r+1}^{N} \exp\left[\eta \Sigma_{z}^{(k)} \right] \\
\left. + \left[\sum_{i=1}^{N} \Sigma_{z}^{(i)} \right]_{q} \left[\sum_{s=1}^{N} \Sigma_{z}^{(s)} + \mathbf{I}^{(N)} \right]_{q} - N [2]_{q} \right) - \gamma h \sum_{i=1}^{N} \frac{1}{2} \Sigma_{z}^{(i)},$$
(98)

which defines a new exactly solvable model for N particles with a spin j = 1.

Eigenvalues of this model are given by

$$E_q(j_{\alpha_i}, m_{\alpha_i}) = -\frac{I}{2} \left([j_{\alpha_i}]_q \, [j_{\alpha_i} + 1]_q - N \, [2]_q \right) - \gamma \, h \, m_{\alpha_i}, \tag{99}$$

and the irreducible blocks for N particles with spin j = 1 will have a total spin given by $j_{\alpha_i} = N, N - 1, \ldots, 0$. The corresponding degeneracies d_{NS} can be found in [39, 31], and the partition function is given again by the formula (77).

5.2 The q-KS model for arbitrary spin

For any spin j > 1, the q-KS model can also be written in terms of non-deformed physical spin-j undeformed operators by making use of the so-called 'deforming functional' approach to the $\mathfrak{su}_q(2)$ representation theory [28, 29, 30]. In particular, if we impose L_{\pm} to be Hermitian conjugate, the $\mathfrak{su}_q(2)$ generators L can be written in terms of the $\mathfrak{su}(2)$ physical spin operators J as:

$$L_{3} = J_{3}, \qquad L_{+} = J_{+} \sqrt{\frac{[j - J_{3}]_{q}[j + J_{3} + 1]_{q}}{(j - J_{3})(j + J_{3} + 1)}}, \qquad L_{-} = \sqrt{\frac{[j - J_{3}]_{q}[j + J_{3} + 1]_{q}}{(j - J_{3})(j + J_{3} + 1)}} J_{-}, \qquad (100)$$

and the q-KS model for spin j is straightforwardly obtained by inserting (100) into (69). The resulting expression is quite involved, but it provides a family of exactly solvable models for any spin j whose spectra and eigenvectors would be deduced again using the representation theory of $\mathfrak{su}_q(2)$.

6 Concluding remarks

The new integrable spin Hamiltonians presented here were constructed as integrable deformations of the KS models by imposing *ab initio* the preservation of a *q*-deformation of the underlying coalgebra symmetry of the latter. The second essential component that leads to the exact solvability of the *q*-KS models is the full structural coincidence between the theory of irreducible representations of $\mathfrak{su}_q(2)$ and the one of $\mathfrak{su}(2)$ (when *q* is not a root of unity).

From a mathematical perspective, an interesting open problem would be the construction of q-KS models for q a root of unity. Moreover, KS models (both deformed and undeformed) in which each particle i has arbitrarily different spins j_i could be considered and, in particular, spin models with two alternating spin values would be interesting. In all these cases, explicit formulae for the eigenvalues of the model rely on the knowledge of the explicit expressions for the multiplicities and eigenvalues of the corresponding tensor product representations, which is by no means an immediate task that, to the best of our knowledge, has not been considered so far in the literature. In addition, the study of the dynamics of the q-KS model for classical spins would be interesting from an integrability point of view and could be performed by following the analysis given in [10] for the undeformed KS Hamiltonian and by applying the techniques developed in [34] for dynamical systems endowed with coalgebra symmetry.

Obviously, a second pending task consists of the study of the physical properties of the q-KS models, starting from the j = 1/2 and j = 1 cases. This requires the computation of their relevant thermodynamic quantities, including the derivation of their associated Curie temperatures and thermodynamic limits. All these properties are expected to behave as smooth deformations of those for the KS model, and a careful analysis will be presented in [40]. In addition, the quantum informational perspective for both the KS and q-KS models is worth developing: entanglement entropies should be analysed and, in particular, the role of q-Dicke states in this context could be relevant. Finally, possible applications of q-KS Hamiltonians for

small N as effective models for systems with few-body spin couplings should be addressed (for instance, in small magnetic clusters [6] and coupled quantum dots [7, 8]). All of these problems are currently under investigation and will be presented elsewhere.

Acknowledgements

The authors acknowledge partial support from the grant PID2023-148373NB-I00 funded by MCIN/AEI/ 10.13039/501100011033/FEDER – UE, and the Q-CAYLE Project funded by the Regional Government of Castilla y León (Junta de Castilla y León) and the Ministry of Science and Innovation MICIN through NextGenerationEU (PRTR C17.I1). I. Gutierrez-Sagredo thanks Universidad de La Laguna, where part of the work has been done, for the hospitality and support.

References

- C. Kittel and H. Shore. Development of a Phase Transition for a Rigorously Solvable Many-Body System. *Physical Review*, 138:A1165–A1169, 5 1965. doi:10.1103/PhysRev.138.A1165.
- H. Al-Wahsh, M. Urbán, and A. Czachor. Exact solutions for a model antiferromagnet with identical coupling between spins. *Journal of Magnetism and Magnetic Materials*, 185:L144–158, 6 1998. doi: 10.1016/S0304-8853(97)01147-5.
- [3] A. J. van der Sijs. Heisenberg models and a particular isotropic model. *Physical Review B*, 48:7125–7133, 9 1993. doi:10.1103/PhysRevB.48.7125.
- [4] A. Czachor. A verification of the random-phase-approximation using exact thermodynamic functions for the Kittel-Shore-Kac model magnet. *Physica B: Condensed Matter*, 311:56–60, 1 2002. doi: 10.1016/S0921-4526(01)01056-0.
- [5] A. Czachor. Energy Spectrum for the System of N Ising Spins with Identical Spin-Spin Coupling K/N Anatomy of Phase Transition. Acta Physica Polonica A, 113:1161–1169, 4 2008. doi: 10.12693/APhysPolA.113.1161.
- [6] O. Ciftja, M. Luban, M. Auslender, and J. H. Luscombe. Equation of state and spin-correlation functions of ultrasmall classical Heisenberg magnets. *Physical Review B*, 60:10122–10133, 10 1999. doi:10.1103/PhysRevB.60.10122.
- [7] D. Loss and D. P. DiVincenzo. Quantum computation with quantum dots. *Physical Review A*, 57:120–126, 1 1998. doi:10.1103/PhysRevA.57.120.
- [8] R. Woodworth, A. Mizel, and D. A. Lidar. Few-body spin couplings and their implications for universal quantum computation. *Journal of Physics: Condensed Matter*, 18:S721–S744, 5 2006. doi:10.1088/0953-8984/18/21/S02.
- [9] J.-M. Liu and G. Müller. Dynamics of an integrable two-sublattice spin model with long-range interaction. *Physical Review B*, 44:12020–12022, 12 1991. doi:10.1103/PhysRevB.44.12020.
- [10] E. Magyari, H. Thomas, R. Weber, C. Kaufman, and G. Muller. Integrable and nonintegrable classical spin clusters. Zeitschrift for Physik B Condensed Matter, 65:363–374, 9 1987. doi:10. 1007/BF01303725.

- [11] G. Müller. High-temperature spin dynamics of the classical Heisenberg magnet in one, two, three and infinite dimensions. Le Journal de Physique Colloques, 49:C8-1403-C8-1404, 12 1988. doi: 10.1051/jphyscol:19888644.
- [12] R. Botet, R. Jullien, and P. Pfeuty. Size Scaling for Infinitely Coordinated Systems. *Physical Review Letters*, 49:478–481, 8 1982. doi:10.1103/PhysRevLett.49.478.
- [13] R. Botet and R. Jullien. Large-size critical behavior of infinitely coordinated systems. *Physical Review B*, 28:3955–3967, 10 1983. doi:10.1103/PhysRevB.28.3955.
- [14] J. L. Cardy. Introduction to Theory of Finite-Size Scaling, volume 2, pages 1–7. 1988. doi:10.1016/ B978-0-444-87109-1.50006-6.
- [15] R. Dekeyser and M. H. Lee. Time-dependent correlations for spin Van der Waals systems. *Physical Review B*, 19:265–273, 1 1979. doi:10.1103/PhysRevB.19.265.
- [16] M. H. Lee, I. M. Kim, and R. Dekeyser. Time-Dependent Behavior of the Spin-¹/₂ Anisotropic Heisenberg Model in Infinite Lattice Dimensions. *Physical Review Letters*, 52:1579–1582, 4 1984. doi:10.1103/PhysRevLett.52.1579.
- [17] V. G. Drinfeld. Quantum Groups, in Proc. Int. Congr. Math. (Berkeley 1986), pages 798–820. American Mathematical Society, 1987.
- [18] M. Jimbo. A q-Difference Analogue of U(g) and the Yang-Baxter Equation. Letters in Mathematical Physics, 10:63–69, 7 1985. doi:10.1007/BF00704588.
- [19] E. K. Sklyanin. Boundary conditions for integrable quantum systems. Journal of Physics A: Mathematical and General, 21:2375–2389, 5 1988. doi:10.1088/0305-4470/21/10/015.
- [20] V. Pasquier and H. Saleur. Common structures between finite systems and conformal field theories through quantum groups. *Nuclear Physics B*, 330:523–556, 1 1990. doi:10.1016/0550-3213(90) 90122-T.
- [21] C. Gómez, M. Ruiz-Altaba, and G. Sierra. Quantum groups in two-dimensional physics. Quantum Groups in Two-Dimensional Physics, page 475, 1996.
- [22] A. Ballesteros and O. Ragnisco. A systematic construction of completely integrable Hamiltonians from coalgebras. Journal of Physics A: Mathematical and General, 31:3791–3813, 4 1998. doi: 10.1088/0305-4470/31/16/009.
- [23] L. C. Biedenharn and M. A. Lohe. Quantum Group Symmetry and Q-Tensor Algebras. World Scientific, 8 1995. doi:10.1142/2815.
- [24] R. Alvarez Nodarse and A. Arenas-Gómez. The q-Analog of the Quantum Theory of Angular Momentum: a Review from Special Functions. *Russian Journal of Mathematical Physics*, 31:24– 43, 3 2024. doi:10.1134/S106192084010023.
- [25] Z.-H. Li and A.-M. Wang. Entanglement entropy in quasi-symmetric multi-qubit states. International Journal of Quantum Information, 13:1550007, 3 2015. doi:10.1142/S0219749915500070.
- [26] D. Raveh and R. I. Nepomechie. q-analog qudit Dicke states. Journal of Physics A: Mathematical and Theoretical, 57:065302, 2 2024. doi:10.1088/1751-8121/adlea4.

- [27] A. Ballesteros, J. de Ramón, I. Gutierrez-Sagredo, and J. J. Relancio. Deformations of the symmetric subspace of qubit chains, in preparation.
- [28] T. Curtright and C. Zachos. Deforming maps for quantum algebras. *Physics Letters B*, 243:237–244, 6 1990. doi:10.1016/0370-2693(90)90845-W.
- [29] D. B. Fairlie. Quantum deformations of SU(2). Journal of Physics A: Mathematical and General, 23:L183–L187, 3 1990. doi:10.1088/0305-4470/23/5/001.
- [30] A. Ballesteros and J. Negro. A characterization of functional realizations of three-dimensional quantum groups. Journal of Physics A: Mathematical and General, 25:5945–5962, 11 1992. doi: 10.1088/0305-4470/25/22/021.
- [31] T. Curtright, T. V. Kortryk, and C. Zachos. Spin multiplicities. *Physics Letters A*, 381:422–427, 2 2017. doi:10.1016/j.physleta.2016.12.006.
- [32] V. Chari and A. Pressley. A guide to quantum groups. Cambridge University Press, 1995.
- [33] A. Ballesteros, M. Corsetti, and O. Ragnisco. N-dimensional classical integrable systems from Hopf algebras. Czechoslovak Journal of Physics, 46:1153–1163, 2017. doi:10.1007/BF01690329.
- [34] A. Ballesteros and O. Ragnisco. Classical dynamical systems from q-algebras: 'cluster' variables and explicit solutions. Journal of Physics A: Mathematical and General, 36:10505–10518, 10 2003. doi:10.1088/0305-4470/36/42/007.
- [35] A. Ballesteros, A. Blasco, F. J. Herranz, F. Musso, and O. Ragnisco. (Super)integrability from coalgebra symmetry: Formalism and applications. *Journal of Physics: Conference Series*, 175:012004, 6 2009. doi:10.1088/1742-6596/175/1/012004.
- [36] F. Musso and O. Ragnisco. Exact solution of the quantum Calogero-Gaudin system and of its q-deformation. Journal of Mathematical Physics, 41:7386-7401, 11 2000. doi:10.1063/1.1308508.
- [37] F. Musso and O. Ragnisco. The spin-¹/₂ Calogero-Gaudin system and its q-deformation. Journal of Physics A: Mathematical and General, 34:2625-2635, 3 2001. doi:10.1088/0305-4470/34/12/309.
- [38] F. Musso, M. Petrera, O. Ragnisco, and G. Satta. Gaudin models with $\mathcal{U}_q(\mathfrak{osp}(1|2))$ symmetry. *Nuclear Physics B*, 716:543–555, 6 2005. doi:10.1016/j.nuclphysb.2005.03.008.
- [39] J. H. van Vleck. The Theory of Electric and Magnetic Susceptibilities. Oxford University Press, 1932.
- [40] A. Ballesteros, I. Gutierrez-Sagredo, V. Mariscal, and J. J. Relancio. Thermodynamical properties of the *q*-deformed Kittel-Shore model, in preparation.