RESOLVING SUBCATEGORIES FOR GENTLE ALGEBRAS I: MONOGENEOUS RESOLVING SUBCATEGORIES FOR GENTLE TREES

BENJAMIN DEQUÊNE AND MICHAËL SCHOONHEERE

ABSTRACT. Let (Q, R) be a gentle quiver such that Q is a directed tree, and \mathbb{K} be an algebraically closed field. We study the resolving subcategories generated by a single nonprojective indecomposable module of $\mathbb{K}Q/\langle R \rangle$ -mod. We modify the geometric model of Baur–Coelho-Simões and Opper–Plamondon–Schroll to compute such subcategories. Then we prove that these subcategories are the join-irreducible of the poset of all the resolving subcategories ordered by inclusion.

Contents

1. Introduction	2
2. Resolving subcategories	4
2.1. General context and definition	4
2.2. Resolving closure	5
2.3. Algorithm for explicit calculations of resolving closures	6
2.4. Discussion on the usability of the algorithm	9
3. Gentle algebras	10
3.1. Gentle quivers and gentle algebras	10
3.2. Representations	12
3.3. String representations	12
3.4. Morphisms between indecomposable representations	13
3.5. Extensions between indecomposable representations	15
3.6. Kernels of epimorphisms	17
3.7. Projective covers and projective resolutions	19
4. Surface model for the representations of gentle algebras	19
4.1. Surfaces and dissections	20
4.2. The gentle quiver of a dissection	21
4.3. Accordions and indecomposable representations	22
4.4. Morphisms and extensions	23
4.5. Projective perspective	25
5. Resolving posets for gentle trees	29
5.1. Generalities on gentle trees	29
5.2. Vocabulary on posets	30
5.3. Resolving order on non-projective indecomposable representations	32
5.4. Resolving posets and posets of resolving subcategories	34
6. Monogeneous resolving subcategories for gentle trees	35

Date: March 3, 2025.

6.1.	Monogeneous geometric resolving sets	35
6.2.	Monogeneous resolving subcategories	38
Acknowledgements		41
References		42

1. INTRODUCTION

The notion of a resolving subcategory of a module category goes back at least to [AB69]. This notion is obtained by extracting some important properties of the subcategory of modules of Gorenstein dimension 0: this subcategory contains the projective modules, it is closed under direct sums, summands, extensions and kernels of epimorphisms. The dual notion of a coresolving subcategory generalizes the subcategory of Gorenstein injective objects. The biresolving i.e. resolving and coresolving subcategories are a generalization of the subcategory of projectiveinjective objects. Inspired from the case of Frobenius categories, [Rum21] states that, given a biresolving subcategory \mathscr{R} of an exact category \mathscr{E} , the localisation of the additive quotient $\mathscr{E}/(\mathscr{R})$ at its morphisms that are both monomorphisms and epimorphisms is triangulated.

Another natural occurrence of the contravariantly finite resolving subcategories is through hereditary cotorsion pairs [AR91, AT22]. Hereditary cotorsion pairs are in bijection with contravariantly finite resolving subcategories and this bijection preserve the natural order given by inclusion. This motivates the computation of all the contravariantly finite resolving subcategories. This question is adressed in [Tak09] where an algorithm is given to generate the smallest resolving subcategory \mathscr{R} containing the family of objects \mathscr{X} . The subcategory \mathscr{R} is called the *resolving closure* of \mathscr{X} and denoted by $\operatorname{Res}(\mathscr{X})$. This algorithm is built to work over every algebra of finite representation type and thus is not profiting of any potential insight on the combinatorics on the algebra.

Gentle algebras are algebras arising from quivers with relations and appeared for the first time in [AH81, AH82]. They have a good combinatorial behaviour: the extension of indecomposable objects have at most two indecomposable summands [BS80, WW85, GP68, SW83, DS87], the homspaces and extensions are endowed with a basis following from string and band combinatorics [CB89, Kra88, ÇPS21]. Moreover the module category over the path algebra, the derived category and the two term homotpy category of projectives are in bijection with dissected surfaces in several geometric models [BCS19, OPS18, APS23, PPP18, PPP17] that give a combinatorial way to compute in these categories.

Our aim is to compute the lattice of the resolving subcategories of mod A for A a gentle algebra of finite representation type. We first focus on the case of gentle algebras arising from an unpunctured disc, or equivalently, the quiver of the gentle algebra is a tree. We call *monogeneous* a resolving subcategory of the form Res(X) for X an non-projective indecomposable object. Our first result concerns the structure of the lattice of resolving subcategories.

Theorem 1.1 (5.29). The join-irreducible elements in the poset of resolving subcategories in $\operatorname{rep}(Q, R)$ ordered by inclusion are exactly given by the monogeneous resolving subcategories. To introduce the main result we have to look closer to the combinatorics of surfaces which helps the computation of extensions and kernels. The dissected disc is endowed with a so-called set of red points on the boundary : one in each cell of the dissected surface. These points are the endpoints of the accordions that are curves in bijection with indecomposable modules. To each accordion δ we associate the set $N_{\text{proj}}(\delta)$ (Definition 4.24) of accordions in bijection with projective modules appearing in the projective resolution of $M(\delta)$ or having a non trivial extension with $M(\delta)$. The set of endpoints of curves in $N_{\text{proj}}(\delta)$ are colored according to a combinatorial rule explained in Definition 6.1 as red orange or green. We can now state the main result :

Theorem 1.2. Let (Q, R) be a gentle tree, and $S(Q, R) = (\Sigma, \mathcal{M}, \Delta^{\circ})$ be the dissected surface associated with (Q, R). For any accordion δ the monogeneous resolving closure $\operatorname{Res}(M(\delta))$ is the union of the indecomposable projectives and the additive closure of $\{M(\varsigma) \mid \varsigma \in \mathscr{R}(\delta)\}$ where the set $\mathscr{R}(\delta)$ is the set of accordions whose sources are coloured either red or orange and targets are coloured either orange or green.

The paper is organized as follows. Section 2 contains generalities about resolving subcategories. In Section 2.3 we improve on control on the steps of the algorithm used to compute the resolving closure using the Krull-Schmidt property of \mathscr{C} . It is still possible to improve on control over the different steps this algorithm by adding a few hypotheses and we do so in Section 2.4. Section 3 recalls what is needed about gentle algebras arising from trees for the study of resolving subcategories. It goes from the definition in Section 3.1, its representations in Section 3.2 and the indecomposables in Section 3.3, the morphisms between indecomposables in Section 3.4, extensions between indecomposable modules in Section 3.5, kernels of epimorphisms in Section 3.6 and projective resolutions in Section 3.7. This proves that modules over such an algebra have the extra hypotheses that are required in the improved algorithm. Section 4 gives a translation of the previous results in the surface model of gentle algebras. Section 4.1 defines the combinatorial shell of dissected surfaces which is linked bijectively in Section 4.2 to gentle quivers. Modules are seen as accordions through Section 4.3, thus we can translate morphisms and extensions between indecomposables in Section 4.4. In order to compute the resolving subcategories we need to consider slight changes to the geometric model detailed in Section 4.5 and we define a first set of tools that gives us the combinatorial insight. With these tools we can start to study resolving closures and the resolving poset in Section 5 by studying properties of gentle trees in Section 5.1, Section 5.2 gives standard tools on posets and notations. Gentle trees enable us to create an order on non-projective indecomposable modules in Section 5.3 that conveys the data of the monogeneous resolving subcategories. In Section 5.4 we study the interaction between the poset we built and the poset of resolving subcategories ordered by inclusion. Then we compute monogeneous resolving subcategories in Section 6. The first step is to give a geometric construction in Section 6.1 and to check that it is a resolving subcategory and then in Section 6.2 we check that the construction is precisely the resolving closure.

The lattice of resolving subcategories is not semidistributive in general. However in [DSar] we introduce a specific decomposition of all resolving subcategories as a union of monogeneous resolving subcategories. This allows us to describe combinatorially all resolving closures and thus all resolving subcategories. We also consider the bijection between resolving subcategories and tilting modules from [AR91] and give a combinatorial construction of these tilting modules.

2. Resolving subcategories

2.1. General context and definition. Let \mathscr{C} be a Krull–Schmidt abelian category.

Convention 2.1. A subcategory is said to be *additive*, or *additively closed*, if it is closed under direct sums and summands

Given $A, B \in \mathscr{C}$ and an epimorphism $f: A \longrightarrow B$, we recall that f is *minimal* whenever there are no summand in Ker(f) that is isomorphic to one of A. Assume that \mathscr{C} has enough projective objects: for any object $M \in \mathscr{C}$, there exist a projective object $P \in \mathscr{C}$ and an epimorphism $P \longrightarrow M$. If such an epimorphism is minimal, we say that P is a projective cover of M. We recall that a projective cover of a given object of \mathscr{C} is unique up to isomorphism. A minimal projective resolution of M is a complex

$$(\mathbf{P}_M^*): \qquad \cdots \xrightarrow{h_3} \mathbf{P}_M^2 \xrightarrow{h_2} \mathbf{P}_M^1 \xrightarrow{h_1} \mathbf{P}_M^0 \xrightarrow{h_0} M$$

such that:

- P_M^0 is the projective cover of M and h_0 is an epimorphism; and,
- for all $i \in \mathbb{N}$, \mathbb{P}_M^{i+1} is the projective cover of $\operatorname{Ker}(h_i)$, and h_{i+1} is the composition of the inclusion map $\operatorname{Ker}(h_i) \longrightarrow \mathbb{P}_M^i$ followed by an epimorphism defining \mathbb{P}_M^{i+1} .

By assumption, any $M \in \mathscr{C}$ admits a unique minimal projective resolution, up to isomorphism of complexes. We recall pdim(M) the *projective dimension* of Mas the greatest integer $i \ge 0$ such that $P_M^i \ne 0$. We recall therefore the global dimension of \mathscr{C} , denoted by $pldim(\mathscr{C})$, as the maximum value of pdim on $M \in \mathscr{C}$. In the following, we denote by $proj(\mathscr{C})$ the subcategory of projective objects of \mathscr{C} .

For any subcategory $\mathscr{D} \subseteq \mathscr{C}$, we write $\operatorname{ind}(\mathscr{D})$ for the set of isomorphism classes of indecomposable objects in \mathscr{D} . Recall that, by the Krull–Schmidt properties, every additive subcategory of \mathscr{C} is characterized by its indecomposable objects. We set $\operatorname{ind} \operatorname{proj}(\mathscr{D}) = \operatorname{ind}(\mathscr{D}) \setminus \operatorname{proj}(\mathscr{C})$.

Let $\mathscr{G} \subseteq \mathscr{C}$ be a subcategory of \mathscr{C} . We say that an object $M \in \mathscr{C}$ admits a cover in \mathscr{G} if there exists an epimorphism $G \longrightarrow M$ with $G \in \mathscr{G}$. We say that the subcategory $\mathscr{G} \subseteq \mathscr{C}$ generates \mathscr{C} whenever every object in \mathscr{C} admits a cover in \mathscr{G} . For instance, $\operatorname{proj}(\mathscr{C})$ is generates \mathscr{C} by hypothesis.

Definition 2.2. A full subcategory $\mathscr{R} \subseteq \mathscr{C}$ is called *resolving* if it satisfies the following conditions:

- (R0) \mathscr{R} is additively closed,
- (R1) \mathscr{R} is generating \mathscr{C} ,
- (R2) \mathscr{R} is closed under extensions, and,
- $(\mathsf{R3})$ $\mathcal R$ is closed under kernels of epimorphisms

Remark 2.3. Note that the resolving subcategories are colsed under isomorphism

We can simplify some conditions to check that a subcategory is resolving under the hypotheses we enforce on \mathscr{C} .

Lemma 2.4. Let $\mathscr{R} \subseteq \mathscr{C}$ be a full additive subcategory generated. We have the following equivalences:

- (i) \mathscr{R} satisfies (R1) if, and only if, $\operatorname{proj}(\mathscr{C}) \subseteq \mathscr{R}$;
- (ii) R is resolving if, and only if, R satisfies (R0), (R1) (R2), and R is closed under syzygies.

Example 2.5.

- The largest resolving subcategory of \mathscr{C} is \mathscr{C} itself.
- The smallest resolving subcategory of $\mathscr C$ is $\operatorname{proj}(\mathscr C)$ by the fact that $\mathscr C$ has enough projectives.
- Given {𝔅_i}_{i∈I} an arbitrary family of resolving subcategories of 𝔅, the intersection ∩_{i∈I}𝔅_i is a resolving subcategory, using Lemma 2.4 (i).

2.2. **Resolving closure.** The last example allows us to introduce the following notion.

Definition 2.6. Let \mathcal{X} be a set of objects of \mathscr{C} . The *resolving closure* of \mathcal{X} , denoted by $\operatorname{Res}(\mathcal{X})$, is the smallest resolving subcategory of \mathscr{C} containing \mathcal{X} . If $\mathcal{X} = \{X\}$, we will write $\operatorname{Res}(\mathcal{X})$ for $\operatorname{Res}(\mathcal{X})$.

A resolving subcategory $\mathscr{R} \subseteq \mathscr{C}$ is said to be *monogeneous* if there exists a non-projective indecomposable object $X \in \mathscr{C}$ such that \mathscr{R} is equal to $\operatorname{Res}(X)$.

The following result is a direct consequence of Definition 2.2 (R0) and the Krull–Schmidt property we imposed to \mathscr{C} .

Corollary 2.7. Every resolving subcategory of C is the resolving closure of its non-projective indecomposable objects.

Therefore, from now on, we focus exclusively on resolving closure of subsets of $\operatorname{ind} \operatorname{proj}(\mathscr{C})$.

Lemma 2.8. Let \mathcal{X} and \mathcal{Y} be two collections of objects in \mathcal{C} . We have the following properties:

- (i) $\mathcal{X} \subseteq \operatorname{Res}(\mathcal{X}),$
- (*ii*) $\operatorname{Res}(\operatorname{Res}(\mathcal{X})) = \operatorname{Res}(\mathcal{X}), and,$
- (*iii*) if $\mathcal{X} \subseteq \mathcal{Y}$, then $\operatorname{Res}(\mathcal{X}) \subseteq \operatorname{Res}(\mathcal{Y})$.

This lemma highlights behaviours satisfied by the so-called *closure operator* whose abstract definition is given below.

Definition 2.9. Let \mathfrak{A} be a set. A *closure operator* on \mathfrak{A} is a map

$$\mathrm{cl}:\mathcal{P}(\mathfrak{A})\to\mathcal{P}(\mathfrak{A})$$

which satisfies the following properties:

- for all $\mathcal{X} \in \mathcal{P}(\mathfrak{A}), \ \mathcal{X} \subseteq \mathrm{cl}(\mathcal{X}),$
- for all $\mathcal{X} \in \mathcal{P}(\mathfrak{A})$, $cl(\mathcal{X}) = cl(cl(\mathcal{X}))$, and,
- for all $\mathcal{X}, \mathcal{Y} \in \mathcal{P}(\mathfrak{A})$, if $\mathcal{X} \subseteq \mathcal{Y}$, then $cl(\mathcal{X}) \subseteq cl(\mathcal{Y})$.

We focus on the operator $(-)^{\text{Res}}$ on subsets of **ind \ proj**(\mathscr{C}) defined by

$$\forall \mathcal{X} \subseteq \operatorname{ind} \backslash \operatorname{proj}(\mathscr{C}), \ \mathcal{X}^{\operatorname{Res}} = \operatorname{ind} \backslash \operatorname{proj}(\operatorname{Res}(\mathcal{X})).$$

The result below follows from Lemma 2.8 and Krull–Schmidt properties.

Lemma 2.10. The operator $(-)^{\text{Res}}$ is a closure operator on $\operatorname{ind} \operatorname{proj}(\mathscr{C})$.

2.3. Algorithm for explicit calculations of resolving closures. Given a commutative noetherian ring R, Takahashi introduces an inductive construction of the resolving closure of any collections of objects in $\mathscr{C} = \mod R$ [Tak09]. Actually, this construction can be used in our more general setting. Moreover, by Corollary 2.7, we give a new version of this algorithm which allows one to get the resolving closures using step-by step calculations by only focusing on collections of non-projective indecomposable objects, and by refining the extensions and kernels calculations.

Theorem 2.11. An additive subcategory $\mathscr{R} \subseteq \mathscr{C}$ is resolving if and only if all the following properties are satisfied:

- $(\mathsf{R}'1) \operatorname{proj}(\mathscr{C}) \subseteq \mathscr{R},$
- (R'2) For any short exact sequence

$$X \xrightarrow{k} E \xrightarrow{f} Y$$
,

if $X, Y \in ind(\mathscr{R})$, then $E \in \mathscr{R}$, and,

(R'3) For any short exact sequence

$$K \xrightarrow{k} M \xrightarrow{f} Y$$
,

if
$$M \in \mathscr{R}$$
 and $Y \in ind(\mathscr{R})$, then $K \in \mathscr{R}$.

Proposition 2.12. Let \mathscr{R} be an additive subcategory of \mathscr{C} . Then \mathscr{R} satisfies (R3) if, and only if, \mathscr{R} satisfies (R'3).

Proof. If \mathscr{R} satisfies (R3), then it is obvious that \mathscr{R} satisfies (R'3).

Assume that ${\mathscr R}$ satisfies (R'3). We will prove that if there is a short exact sequence

$$K \xrightarrow{k} M \xrightarrow{f} N$$

where $M, N \in \mathscr{R}$ and $K \in \mathscr{C}$, then $K \in \mathscr{R}$. Using the Krull–Schmidt property of \mathscr{C} , we can, and we will, proceed by induction on the number of indecomposable summands of N. The initialisation is obvious by ($\mathbb{R}'3$).

Now assume that N is not indecomposable. In this case, we write $N = N_1 \oplus N'$ with $N_1 \in \text{ind}(\mathscr{C})$. As \mathscr{R} is additive, we have that $N_1, N' \in \mathscr{R}$. Therefore we have the following short exact sequences.

$$K \xrightarrow{k} M \xrightarrow{f} N \text{ and } N_1 \xrightarrow{i} N \xrightarrow{p} N'$$

By taking the pull back of f along i, we obtain the following diagram of short exact sequences.

$$K \xrightarrow{} M' \xrightarrow{f_1} N_1$$

$$\| \qquad \downarrow \qquad \downarrow \qquad \downarrow$$

$$K \xrightarrow{k} M \xrightarrow{f'} N$$

$$f'_{\downarrow} \qquad \downarrow$$

$$N' = N'$$

$$6$$

We already know that $N_1 \in ind(\mathscr{R})$. Moreover

$$M' \longrightarrow M \xrightarrow{f'} N'$$

is a short exact sequence, where $M, N' \in \mathscr{R}$, and the number of indecomposable summands of N' is strictly smaller than the number of indecomposable summands of N. By induction hypothesis, we have that $M' \in \mathscr{R}$. therefore, by (R'3), we have that $K \in \mathscr{R}$ on the exact sequence

$$K \longrightarrow M' \xrightarrow{f_1} N_1$$

. This ends the induction.

Proposition 2.13. Let \mathscr{R} be an additive subcategory of \mathscr{C} satisfying (R3). Then \mathscr{R} satisfies (R2) if, and only if, \mathscr{R} satisfies (R'2).

Proof. It is clear that if \mathscr{R} satisfies (R2), then \mathscr{R} satisfies (R'2).

Assume that \mathscr{R} satisfies (R'2). We will show that if we have a short exact sequence

$$M \xrightarrow{k} E \xrightarrow{f} N$$

where $M, N \in \mathscr{R}$ and $E \in \mathscr{C}$, then $E \in \mathscr{R}$. We will proceed in two steps: we first study the case where $N \in ind(\mathscr{R})$, and then we study the general case.

First assume that $N \in ind(\mathscr{R})$. Consider the decomposition into indecomposable summands: $M = \bigoplus_{i=0}^{n} M_i$. By pushing out along the *i*th projection we obtain the family of short exact sequences:

$$M_i \longrightarrow E_i \longrightarrow N$$

As $M_i, N \in ind(\mathscr{R})$, we have $E_i \in \mathscr{R}$ by (R'2). Consider the family of the following diagrams:

$$\begin{array}{cccc} M \rightarrowtail & E & \longrightarrow & N \\ \pi_i & & \downarrow & & \parallel \\ M_i \rightarrowtail & E_i & \longrightarrow & N \end{array}$$

Summing the bottom rows we obtain:

By taking the mapping cone, we obtain the following short exact sequence.

$$E \longrightarrow \bigoplus_{i=0}^{n} (E_i \oplus N) \longrightarrow \bigoplus_{i=0}^{n} N$$

Thus E is in \mathscr{R} by (R3).

Now we prove the general result by induction on the number of indecomposable summands of N, as previously. The initialisation is done above. Asume that N is not indecomposable. We can write $N = N_1 \oplus N'$ with $N_1 \in ind(\mathscr{R})$ and $N' \in \mathscr{R}$

by additivity of \mathscr{R} . As in the proof of Proposition 2.12 we obtain the following diagram.



As we have the following short exact sequence

$$M \longrightarrow E' \longrightarrow N'$$

by induction hypothesis, we have that $E' \in \mathscr{R}$. Moreover the following short exact sequence

$$E' \longrightarrow E \longrightarrow N_1$$

allows us to conclude that $E \in \mathscr{R}$ by the first step of the proof.

Proof of Theorem 2.11. The result follows from Lemma 2.4 (i), Propositions 2.12 and 2.13.

Thanks to our previous result, we update the algorithm of Takahashi by only focusing on non projective indecomposable objects in \mathscr{C} .

Algorithm 2.14. Let \mathscr{C} be an abelian Krull–Schmidt category which has enough projective objects. We input a set $\mathcal{X} \subseteq \operatorname{ind} \operatorname{proj}(\mathscr{C})$.

- (1) Set $\mathcal{X}^0 = \mathcal{X} \cup (\operatorname{ind}(\mathscr{C}) \cap \operatorname{proj}(\mathscr{C})).$
- (2) At the *i*th iteration of the algorithm, we define \mathcal{X}^{i+1} as the union of \mathcal{X}^i and indecomposable summands of $Y \in \mathscr{C}$ appearing either: (2*a*) in a short exact sequence

$$A \rightarrowtail Y \longrightarrow B$$

with $A, B \in \mathcal{X}^i$, or,

(2b) in a short exact sequence

$$Y \longmapsto \bigoplus_{k=1}^p A_k \longrightarrow B$$

for some $p \in \mathbb{N}^*$ with $B, A_1, \ldots, A_p \in \mathcal{X}^i$;

- (3) If $\mathcal{X}^{i+1} \neq \mathcal{X}^i$, then go back to Step (2);
- (4) Otherwise, return \mathcal{X}^{i+1} .

Theorem 2.15. Let \mathscr{C} be an abelian Krull–Schmidt category with enough projective objects. For any set $\mathcal{X} \subseteq \operatorname{ind} \operatorname{proj}(\mathscr{C})$, we have

$$\operatorname{Res}(\mathcal{X}) = \operatorname{add}\left(\bigcup_{i \ge 0} \mathcal{X}^i\right),$$

and, equivalently, $\mathcal{X}^{\text{Res}} = \left(\bigcup_{i \geqslant 0} \mathcal{X}^i\right) \setminus \text{proj}(\mathscr{C})$

Proof. It follows from [Tak09, Remark 3.2] and Propositions 2.12 and 2.13.

2.4. Discussion on the usability of the algorithm. First, note that Algorithm 2.14 is less effective than the one given by Takahashi. Nevertheless, we support the fact that we earn in precision to construct step by step \mathcal{X}^{Res} for all collections $\mathcal{X} \subseteq \text{ind} \setminus \text{proj}(\mathcal{C})$. Recall that, by Corollary 2.7, it is enough to describe any resolving subcategory of \mathscr{C} .

Secondly, one can be disappointed by noticing that, in general, Algorithm 2.14 does not end. For instance, we could be in trouble if $\operatorname{ind}(\mathscr{C})$ is infinite. However, even though we assume that $\operatorname{ind}(\mathscr{C})$ is finite, another issue could come from the step (2b) as we have to go around all the objects of $\operatorname{add}(\mathcal{X}^i)$. To overcome this problem, we can restrict ourselves to categories \mathscr{C} which satisfy the following condition:

 $(\mathsf{Mid})_{p_0}$ There exists $p_0 \in \mathbb{N}^*$ such that, for any short exact sequence

$$X \longrightarrow Y \longrightarrow Z$$

in \mathscr{C} where $X, Z \in ind(\mathscr{C})$, the object Y admits at most p_0 indecomposable summands, up to isomorphism.

Under those hypotheses, we ensure that the algorithm ends.

Lemma 2.16. Let \mathscr{C} be an abelian Krull–Schmidt category with enough projective objects. Assume that \mathscr{C} satisfies $(\mathsf{Mid})_{p_0}$ for some $p_0 \in \mathbb{N}^*$. Let $\mathscr{R} \subseteq \mathscr{C}$ be an additive subcategory satisfying (R2). Then \mathscr{R} satisfies (R'3) if and only if \mathscr{R} satisfies the following:

 $(\mathsf{R}''3)_{p_0}$ for any short exact sequence

$$K \rightarrowtail \widetilde{M} \longrightarrow Y$$

with $Y \in ind(\mathscr{R})$ and $\widetilde{M} \in \mathscr{R}$ admitting at most p_0 indecomposable summands, then $K \in \mathscr{R}$.

Proof. It is clear that if \mathscr{R} satisfies $(\mathsf{R}'3)$ then \mathscr{R} satisfies $(\mathsf{R}''3)_{p_0}$.

Assume that \mathscr{R} satisfies $(\mathsf{R}''3)_{p_0}$. Consider a short exact sequence

$$K \longrightarrow M \longrightarrow Y$$

with $Y \in \operatorname{ind}(\mathscr{R})$ and $M \in \mathscr{R}$. Consider the decomposition of K into indecomposable summands : $K = \bigoplus_{i=0}^{n} K_i$. By pushing out along the *i*th projection, we get the family of short exact sequences.

$$K_i \longrightarrow M_i \longrightarrow Y$$

By $(\mathsf{Mid})_{p_0}$, for all $i \in \{0, \ldots, n\}$, we have that M_i admits at most p_0 indecomposable summands. Moreover, from these short exact sequence, by proceeding the same way that we did in the proof of Proposition 2.13, namely summing the family of short exact sequences, and taking some mapping cone, we get the following exact sequence.

$$M \rightarrowtail \left(\bigoplus_{i=0}^{n} M_{i} \right) \oplus Y \longrightarrow \bigoplus_{i=0}^{n} Y$$

By (R2) and additivity of \mathscr{R} , we have that $M_i \in \mathscr{R}$ for all $i \in \{0, ..., n\}$. By $(\mathbb{R}^{\prime\prime}3)_{p_0}$, we get that $K_i \in \mathscr{R}$ for all $i \in \{0, ..., n\}$ and, by additivity of \mathscr{R} , we have that $K \in \mathscr{R}$.

Let \mathbb{K} be a field. For the following statement, we will restrict ourself to \mathbb{K} -linear categories, meaning, in particular, that the homspaces between objects of \mathscr{C} are \mathbb{K} -vector spaces. This restriction is natural when we look at module categories over a \mathbb{K} -algebra. We will also impose that $\operatorname{ind}(\mathscr{C})$ is finite, and the homspaces between indecomposable representations of \mathscr{C} are at most one-dimensional. This is the kind of restriction we will handle with in our setting later on (see Section 5).

Proposition 2.17. Let \mathbb{K} be a field and \mathscr{C} be a \mathbb{K} -linear Krull–Schmidt category with enough projective objects. Assume that $\operatorname{ind}(\mathscr{C})$ is finite, \mathscr{C} satisfies $(\operatorname{Mid})_{p_0}$ for some $p_0 \in \mathbb{N}^*$, and

 $\forall X, Y \in \operatorname{ind}(\mathscr{C}), \dim_{\mathbb{K}}(\operatorname{Hom}(X, Y)) \leq 1.$

Then, for any $\mathcal{X} \subseteq \operatorname{ind} \backslash \operatorname{proj}(\mathscr{C})$, Algorithm 2.14 ends.

Proof. By the fact that, for any object $U, V \in \mathscr{C}$, we can describe the $\operatorname{Hom}(U, V)$ via the homspaces from indecomposable summands of U and the ones of V. By assumption, it implies that to describe a morphism of $\operatorname{Hom}(U, V)$, up to a change of bases, we only need to know if the induced morphism from each summand of U to each one of V is whether a zero map or not. It implies that, up to a change of bases, their is a finite number of morphisms from U to V.

Thanks to Lemma 2.16, as $\operatorname{ind}(\mathscr{C})$ is finite, and by previous arguments, we know that the step (2) of Algorithm 2.14 can be done with a finite number of calculations. Moreover $(\mathcal{X}^i)_{i\geq 0}$ is an increasing sequence of subsets of $\operatorname{ind}(\mathscr{C})$. Thus, as $\operatorname{ind}(\mathscr{C})$ is finite, there must be an integer $j \geq 0$ such that $\mathcal{X}^j = \mathcal{X}^{j+1}$. This is equivalent to say that the algorithm ends.

Remark 2.18. Let \mathbb{K} be a field, and A be a artinian \mathbb{K} -algebra. Butler and Ringel [BR87] showed that $\mathscr{C} = \mod A$ satisfies $(\mathsf{Mid})_{p_0}$ for a p_0 which depends of the maximal lengths of both indecomposable projective and injective A-modules.

However, one can still disappointed as we did not give an explicit recipe to get the objects appearing in step (2) to construct \mathcal{X}^{Res} from any $\mathcal{X} \subseteq \text{ind} \setminus \text{proj}(\mathscr{C})$.

In the following, we restrict ourselves on the study of module categories of some gentle algebras, and we refine Algorithm 2.14 in this setting until handling the explicit calculations of the objects appearing in step (2). Consequently, it will allow us to explicitly describe all the resolving subcategories of those categories.

3. Gentle Algebras

In this section, in addition to recalling some results about gentle quivers, we establish results on explicit combinatorial calculations of the kernel of any (minimal) epimorphism between representations of gentle quivers.

3.1. Gentle quivers and gentle algebras. A *quiver* is given by a quadruplet $Q = (Q_0, Q_1, s, t)$ where:

- Q_0 is the vertex set of Q;
- Q_1 is the arrow set of Q;
- $s, t; Q_1 \longrightarrow Q_0$ are respectively source and target function.

In the following, we assume that Q is *finite*; meaning that Q_0 and Q_1 are finite sets.

A *path* of Q is either a formal element e_q for some $q \in Q_0$ called the *lazy* path at q, or a finite sequence of arrows $(\alpha_1, \ldots, \alpha_k)$, for some $k \ge 1$, such that

 $s(\alpha_{i+1}) = t(\alpha_i)$ for all $1 \leq i < k$. From now on, we write $c = \alpha_k \cdots \alpha_1$ for the non-lazy path $(\alpha_1, \ldots, \alpha_k)$, and we set $s(c) = s(\alpha_1)$ and $t(c) = t(\alpha_k)$. We also set $s(e_q) = t(e_q) = q$ for all $q \in Q_0$. The *length* of a path c, denoted by $\ell(c)$, is the number of arrows whose composing c. A *cycle* is a non-lazy path whose source and target coincide.

For $\alpha \in Q_1$, write α^{-1} for the formal inverse of α . Set $s(\alpha^{-1}) = t(\alpha)$ and $t(\alpha^{-1}) = s(\alpha)$. A walk of Q is either a lazy path, or a finite sequence of arrows and inverse arrows $\rho = (\beta_1, \ldots, \beta_k)$, for some $k \ge 1$, such that $s(\beta_{i+1}) = t(\beta_i)$ for all $1 \le i < k$. From now on, we write $\rho = \alpha_k^{\varepsilon_k} \ldots \alpha_1^{\varepsilon_1}$ for the non-lazy walk $(\alpha_1^{\varepsilon_1}, \ldots, \alpha_k^{\varepsilon_k})$, where $\alpha_i \in Q_1$ and $\varepsilon_i \in \{\pm 1\}$. As previously for paths, we set $s(\rho) = s(\alpha_1^{\varepsilon_1})$ and $t(\rho) = t(\alpha_k^{\varepsilon_k})$.

In the following, we assume that Q is *connected*; meaning that for any pair $(v_1, v_2) \in Q_0$, there exists a walk ρ of Q such that $s(\rho) = v_1$ and $t(\rho) = v_2$.

Let \mathbb{K} be a field. To state results which are mainly based on algebraic geometry arguments, we assume that \mathbb{K} is algebraically closed. The *path algebra* of Q (over \mathbb{K}), denoted by $\mathbb{K}Q$, is the \mathbb{K} -vector space with a basis given by all the paths of Q endowed with a multiplication defined on paths as follows: for c_1 and c_2 two paths of Q, we set

$$c_2 \cdot c_1 = \begin{cases} c_2 c_1 & \text{if } s(c_2) = t(c_1), \\ 0 & \text{otherwise.} \end{cases}$$

A relation on Q is a K-linear combination involving a finite set of paths of length greater than or equal to 2 that have the same sources and the same sinks. Such a relation is said to be *quadratic monomial* if it involves only one path and that path is of length 2. For any set of relations R, write $\langle R \rangle$ for the (bi-sided) ideal in $\mathbb{K}Q$ generated by the elements of R.

For $\ell \in \mathbb{N}$, write $\mathbb{K}Q_{\geq \ell}$ for the ideal generated by all the paths c such that $\ell(c) \geq \ell$. An ideal $I \subseteq \mathbb{K}Q$ is said to be *admissible* if $\mathbb{K}Q_{\geq N} \subseteq I \subseteq \mathbb{K}Q_{\geq 2}$ for some $N \geq 2$. Recall that if I is an admissible ideal, the quotient algebra $\mathbb{K}Q/I$ is finite-dimensional.

Definition 3.1. A *gentle quiver* is a pair (Q, R) such that:

- Q is a quiver such that there are at most two incoming arrows and at most two outgoing arrows at each vertex of Q;
- R is a set of quadratic monomial relations such that:
 - for any arrow $\alpha \in Q_1$:
 - there is at most one $\beta \in Q_1$ such that $s(\beta) = t(\alpha)$ and $\beta \alpha \in R$;
 - there is at most one $\gamma \in Q_1$ such that $s(\gamma) = t(\alpha)$ and $\gamma \alpha \notin R$;
 - there is at most one $\beta' \in Q_1$ such that $t(\beta') = s(\alpha)$ and $\alpha\beta' \in R$;
 - there is at most one $\gamma' \in Q_1$ such that $t(\gamma') = s(\alpha)$ and $\alpha \gamma' \notin R$;
 - for any cycle $c = \alpha_k \cdots \alpha_1$, either there exists $i \in \{1, \ldots, k-1\}$ such that $\alpha_{i+1}\alpha_i \in R$, or $\alpha_1\alpha_k \in R$.

A \mathbb{K} -algebra Λ is said to be *gentle* if there exists a gentle quiver (Q, R) such that $\Lambda \cong \mathbb{K}Q/\langle R \rangle$.

Remark that if (Q, R) is a gentle quiver, then $\langle R \rangle$ is an admissible ideal in $\mathbb{K}Q$, and thus any gentle algebra is finite-dimensional.

3.2. Representations. Let (Q, R) be a gentle quiver.

Definition 3.2. A representation of (Q, R) is a pair $E = ((E_q)_{q \in Q_0}, (E_\alpha)_{\alpha \in Q_1})$ such that:

- for any $q \in Q_0$, E_q is a K-vector space;
- for any $\alpha \in Q_1, E_\alpha : E_{s(\alpha)} \longrightarrow E_{t(\alpha)}$ is a K-linear transformation;
- for any $(\alpha, \beta) \in (Q_1)^2$ such that $\beta \alpha \in R$, then $E_\beta E_\alpha = 0$.

We can understand the construction of a representation of (Q, R) as assigning a \mathbb{K} -vector space to each vertex of Q, and assigning a \mathbb{K} -linear map to each arrow of Q. A representation E of (Q, R) is *finite dimensional* whenever E_q is a finite-dimensional \mathbb{K} -vector space for all $q \in Q_0$.

Let *E* and *F* be two representations of (Q, R). Write $E \cong F$ whenever *E* and *F* are *isomorphic*. Write $E \oplus F$ for the *direct sum of representations E* and *F*. A representation *G* is said to be *indecomposable* if $E \cong 0$ or $F \cong 0$, whenever we have $G \cong E \oplus F$.

Denote by $\operatorname{rep}(Q, R)$ the category of finite-dimensional representations of (Q, R). Note that, even though we omitted it with this notation, this category depends on the field K. We highlight the structure of $\operatorname{rep}(Q, R)$ in the following well-known result.

Proposition 3.3. The category rep(Q, R) is a Krull–Schmidt K-linear category that has enough projective representations.

Write $\operatorname{proj}(Q, R)$ for the subcategory of projective representations in $\operatorname{rep}(Q, R)$

Another relevant point is that there is a correspondence between representations of (Q, R) and $(\mathbb{K}Q/\langle R \rangle)$ -modules.

Theorem 3.4 ([ASS06]). The category rep(Q, R) is equivalent to the category $mod(\mathbb{K}Q/\langle R \rangle)$ of finitely generated $(\mathbb{K}Q/\langle R \rangle)$ -modules.

Write $\operatorname{ind}(Q, R)$ for the set of isomorphism classes of indecomposable representations of (Q, R). We say that (Q, R) is *representation-finite* whenever we have $\# \operatorname{ind}(Q, R) < \infty$. In the following, and for our purpose, we focus on representation-finite gentle quivers.

3.3. String representations. In this subsection, we recall the string combinatorics which give a complete classification of indecomposable representations of any representation-finite gentle quiver (Q, R). Let us first begin with the following definition.

Definition 3.5. A *string* of (Q, R) is either a lazy path, or a non-lazy walk $\rho = \alpha_k^{\varepsilon_k} \cdots \alpha_1^{\varepsilon_1}$ of Q such that, for all $1 \leq i < k$:

- $\alpha_{i+1} \neq \alpha_i$ whenever $\varepsilon_{i+1} = -\varepsilon_i$: meaning that ρ is *reduced*;
- $\alpha_{i+1}\alpha_i \notin R$ whenever $\varepsilon_{i+1} = \varepsilon_i = 1$; and,
- $\alpha_i \alpha_{i+1} \notin R$ whenever $\varepsilon_{i+1} = \varepsilon_i = -1$.

Those strings allow us to construct a particular family of isomorphism classes of indecomposable representations of (Q, R).

Definition 3.6. Let (Q, R) be a gentle quiver. Consider $\rho = \alpha_k^{\varepsilon_k} \cdots \alpha_1^{\varepsilon_1}$ to be a string of (Q, R), the *standard string representation* $M(\rho)$ is a representation of Q obtained from ρ as follows:

- Let $v_0 = s(\alpha_1^{\varepsilon_1})$ and for all $i \in \{1, \ldots, k\}, v_i = t(\alpha_i^{\varepsilon_i});$
- For $q \in Q_0$, $M(\rho)_q$ is the vector space having as basis $\{x_i \mid v_i = q\}$, where the x_i are formal elements;
- For $\beta \in Q_1$, $M(\rho)_{\beta} : M(\rho)_{s(\beta)} \longrightarrow M(\rho)_{t(\beta)}$ is the linear transformation such that:

$$M(\rho)_{\beta}(x_i) = \begin{cases} x_{i-1} & \text{if } \alpha_i = \beta \text{ and } \varepsilon_i = -1\\ x_{i+1} & \text{if } \alpha_{i+1} = \beta \text{ and } \varepsilon_{i+1} = 1\\ 0 & \text{otherwise.} \end{cases}$$

A representation E of (Q, R) is a *string representation* whenever E is isomorphic to some standard string representation.

The following well-known result shows that the string representations describe $\operatorname{ind}(Q, R)$.

Theorem 3.7 ([BR87]). Let \mathbb{K} be an algebraically closed field. Let (Q, R) be a representation-finite gentle quiver. Any $E \in ind(Q, R)$ is isomorphic to a string representation. Moreover, given two strings of (Q, R), ρ and ρ' , $M(\rho)$ is isomorphic to $M(\rho')$ if and only if $\rho' = \rho^{\pm 1}$.

We identify ρ with its inverse ρ^{-1} as they give isomorphic representations.

Remark 3.8. The previous result can be stated for any gentle quiver by considering, in addition, the so-called *band representations*. They are parameterized by primitive cyclic strings of (Q, R), nonzero elements of \mathbb{K} , and positive integers. However, as we are restricting ourselves to representation-finite gentle quivers, we will not need to consider band representations in this paper.

3.4. Morphisms between indecomposable representations. Now we recall a combinatorial way to describe the homomorphism space between two fixed indecomposable representations of (Q, R).

Definition 3.9. Let $\rho = \alpha_k^{\varepsilon_k} \cdots \alpha_1^{\varepsilon_1}$ be a string of (Q, R). A *substring* of ρ is either a lazy path e_q for some $q \in \text{Supp}_0(\rho)$ or a string $\alpha_j^{\varepsilon_j} \cdots \alpha_i^{\varepsilon_i}$ for some $1 \leq i \leq j \leq k$ together with its position in the string. A substring $\sigma = \alpha_j^{\varepsilon_j} \cdots \alpha_i^{\varepsilon_i}$ of ρ is said to be *on top* of ρ if i = 1 or $\varepsilon_{i-1} = -1$, and j = k or $\varepsilon_{j+1} = 1$. In the same way σ is said to be *at the bottom* of ρ if i = 1 or $\varepsilon_{i-1} = 1$, and j = k or $\varepsilon_{j+1} = -1$. Note that any string is both on top and at the bottom of itself.

These combinatorial notions on strings are sufficient to describe the homomorphism space between any pair of indecomposable representations of (Q, R).

Theorem 3.10 ([CB89]). Let (Q, R) be a representation-finite gentle quiver. Consider ρ and ρ' two strings of (Q, R). Then $\operatorname{Hom}(\operatorname{M}(\rho), \operatorname{M}(\rho')) \cong \mathbb{K}^{\#[\rho, \rho']}$ where $[\rho, \rho']$ is the set of pairs (σ, σ') such that the substring σ is on top of ρ , the substring σ' is at the bottom of ρ' and $\sigma' = \sigma^{\pm 1}$.

More precisely, we can describe a basis $(\varphi_{(\sigma,\sigma')})_{(\sigma,\sigma')\in[\rho,\rho']}$ of $\mathsf{Hom}(\mathsf{M}(\rho),\mathsf{M}(\rho'))$ such that $\varphi_{(\sigma,\sigma')}$ is defined as follows:

Let (x₀,...,x_p) and (y₀,...,y_q) be the collections of formal variables linearly independent that define M(ρ) and M(ρ') as in Definition 3.6;



FIGURE 1. Figures showing a substring σ of ρ on top of ρ (above), and a substring σ of ρ at the bottom of ρ (below).

- The substring σ depends only of the variables x_i, \ldots, x_{j+1} , for some $0 \leq i \leq j+1 \leq p$, and the substring σ' depends only of the variables $y_k, \ldots, y_{\ell+1}$ for some $0 \leq k \leq \ell+1 \leq q$;
- If $\sigma = \sigma'$, we define $\varphi_{(\sigma,\sigma')}$ on the variable (x_0, \ldots, x_p) by:

$$\forall u \in \{0, \dots, p\}, \ \varphi_{(\sigma, \sigma')}(x_u) = \begin{cases} y_{k+u-i} & \text{if } i \leq u \leq j+1, \\ 0 & \text{otherwise.} \end{cases}$$

A similar definition can be given if $\sigma = (\sigma')^{-1}$.



FIGURE 2. Figure illustrating combinatorially the map $\varphi_{(\sigma,\sigma')}$ defined in Theorem 3.10, for $\sigma = \sigma'$ a substring on the top of ρ and at the bottom of ρ' .

Remark 3.11. In a more general setting, it is still possible to describe the homomorphism space between indecomposable representations, including band representations, for any gentle quiver (Q, R). We refer the reader to [Kra88] for more details. As we focus on representation-finite gentle quivers, Theorem 3.10 is enough for our purposes.

To compute rep(Q, R), its indecomposable objects and its morphisms, we often use the *Auslander–Reiten quiver* of (Q, R). This is a quiver, denoted by AR(Q, R), whose vertices are the (isomorphism classes of) indecomposable representations of (Q, R), and whose arrows correspond bijectively to the basis elements of the K-vector spaces of irreducibles morphisms between indecomposable representations (see [ASS06] for more details). In Section 5, we will state useful results on AR(Q, R) in the case where Q is a tree. 3.5. Extensions between indecomposable representations. Thanks to the description of all the homomorphism spaces, the extension between indecomposable representations can be explicited combinatorially.

Definition 3.12. Let ρ and μ be two strings of (Q, R).

• An arrow extension of ρ by μ is a string ν in (Q, R) such that there exists an arrow $a \in Q_1$ such that $\nu = \mu a \rho$. We denote by $\operatorname{ArExt}(\rho, \mu)$ the set of arrow extensions of ρ by μ .



FIGURE 3. Illustration of an arrow extension of ρ by μ .

- Assume that there exist five strings $\rho_{\ell}, \rho_r, \mu_{\ell}, \mu_r, \sigma$ such that $\rho = \rho_r \sigma \rho_{\ell}$ and $\mu = \mu_r \sigma \mu_{\ell}$ with the following conditions:
 - (i) If ρ_{ℓ} is lazy, then there exists $c \in Q_1$ and μ'_r a string of (Q, R) such that $\mu_r = c^{-1}\mu'_r$
 - (ii) If ρ_r is lazy, then there exist $d \in Q_1$ and μ'_r a string of (Q, R), such that $\mu_r = \mu'_r d$,
 - (*iii*) If σ is lazy, then:
 - (iii)(a) if $\rho_r = \rho'_r b^{-1}$ and $\mu_r = \mu'_r d$, for some $b, d \in Q_1$ and some strings ρ'_r and μ'_r of (Q, R), then $db \in R$, and
 - $\begin{array}{l} (iii)(b) \ \ \text{if} \ \rho_\ell = a\rho'_\ell \ \text{and} \ \mu_\ell = c^{-1}\mu'_\ell, \ \text{for some} \ a,c \in Q_1 \ \text{and some strings} \\ \rho'_r \ \text{and} \ \mu'_\ell \ \text{of} \ (Q,R) \ \text{then} \ ca \in R. \end{array}$

An overlap extension of ρ by μ is a string ν of (Q, R) such that either $\nu = \mu_r \sigma \rho_\ell$ or $\nu = \rho_r \sigma \mu_\ell$. We say that $\mu_r \sigma \rho_\ell$ and $\rho_r \sigma \mu_\ell$ are conjugated overlap extensions of ρ by μ . We denote by $\text{OvExt}(\rho, \mu)$ the set of overlap extensions of ρ by μ , by setting, if ρ and μ do not satisfy the above conditions, $\text{OvExt}(\rho, \mu) = \emptyset$.



FIGURE 4. Illustration of the general configuration of an overlap extension of ρ by μ . The dashed and dotted colored lines correspond to the overlap extensions of ρ by μ .

Proposition 3.13 ([QPS21]). Let ρ and μ be two string in (Q, R).

• Assume that there is an arrow extension of ρ by μ . Let $\nu = \mu a \rho$ in the notation of Definition 3.12. We have the following nonsplit short exact sequence.

$$M(\mu) \longrightarrow M(\nu) \longrightarrow M(\rho)$$

FIGURE 5. The nonsplit exact sequence corresponding to the arrow extension of ρ by μ pictured with the strings.

• Assume that there is an overlap extension of ρ by μ . Let $\nu_1 = \mu_r \sigma \rho_\ell$ and $\nu_2 = \rho_r \sigma \mu_\ell$ in the notation of Definition 3.12. We have the following nonsplit short exact sequence.

$$M(\mu) \longrightarrow M(\nu_1) \oplus M(\nu_2) \longrightarrow M(\rho)$$



FIGURE 6. The nonsplit exact sequence corresponding to the overlap extension of ρ by μ pictured with the strings.

The theorem below justifies the terminology of arrow and overlap extensions, and shows that they are enough to describe any extension space.

Theorem 3.14 ([BDM⁺20, QPS21]). Let (Q, R) be a representation-finite gentle quiver. Let ρ and μ be two strings of (Q, R). A basis of $Ext^1(M(\mu), M(\rho))$ is given by the nonsplit short exact sequences

$$\mathbf{M}(\mu) \rightarrowtail E \longrightarrow \mathbf{M}(\rho),$$

Such that, either:

- $E \cong M(\nu)$ for some $\nu \in ArExt(\rho, \mu)$; or,
- $E \cong M(\nu_1) \oplus M(\nu_2)$ with $\nu_1, \nu_2 \operatorname{OvExt}(\rho, \mu)$ that are conjugated.

Corollary 3.15. The category $\operatorname{rep}(Q, R)$ satisfies $(\operatorname{Mid})_{p_0}$ for $p_0 = 2$. More precisely, for any short exact sequence

$$X \rightarrowtail E \longrightarrow Y$$

where $X, Y \in ind(Q, R)$, the representation E admits at most two indecomposable summands.

Remark 3.16. Corollary 3.15 is a well-known result for string algebras (of which the gentle algebras are a part) proved by using calculations due to [BS80, WW85], with methods given in [GP68], and, independently by using covering theory (see [SW83, DS87]).

3.6. Kernels of epimorphisms. In this subsection, we compute the kernels of a certain family of minimal epimorphisms in $\operatorname{rep}(Q, R)$, which will be enough to study all the possible epimorphisms in $\operatorname{rep}(Q, R)$, under some more assumptions on (Q, R) (see Section 5).

Definition 3.17. Let (Q, R) be a gentle quiver. Let ρ be a string of (Q, R). Given some $p \in \mathbb{N}^*$, the *p*-tuple $(\sigma_1, \ldots, \sigma_p)$ of strings is said to be a *p*-cover of ρ whenever both of the following hold:

- For each $i \in \{1, \ldots, p\}$, σ_i is a substring of ρ ; and,
- There exists a *p*-tuple of strings $(\widetilde{\sigma_1}, \ldots, \widetilde{\sigma_p})$ such that:
 - for all $i \in \{1, \ldots, p\}$, $\tilde{\sigma}_i$ is a substring of σ_i ; and,
 - $\rho = \widetilde{\sigma_p} \cdots \widetilde{\sigma_1}$.

Such a *p*-cover of ρ is reduced if there are no $j \in \{1, \ldots, p-1\}$ such that for some $(i_1, \ldots, i_j) \in \{1, \ldots, p\}, (\sigma_{i_1}, \ldots, \sigma_{i_j})$ is a *j*-cover of ρ .

Example 3.18. In Figure 7, the strings $\sigma_1, \ldots, \sigma_7$ are substrings of the string μ in some ambient gentle quiver (Q, R).

- The 7-tuple $(\sigma_1, \sigma_2, \sigma_3, \sigma_4, \sigma_5, \sigma_6, \sigma_7)$ is a 7-cover of μ but it is not reduced.
- The 4 tuple $(\sigma_1, \sigma_4, \sigma_6, \sigma_7)$ is a reduced 4-cover of μ .



FIGURE 7. An example of a *p*-cover of a string μ

Lemma 3.19. Let (Q, R) be a gentle quiver, and let $p \ge 2$ be an integer. Let μ be a string of (Q, R). For any reduced p-cover $(\sigma_1, \ldots, \sigma_p)$, and for any $i \in \{1, \ldots, p-1\}$, there exists a unique triplet $(\sigma_i^{\ell}, \sigma_{i,i+1}, \sigma_{i+1}^{r})$ such that:

- $\sigma_i = \sigma_{i,i+1} \sigma_i^{\ell};$
- $\sigma_{i+1} = \sigma_{i+1}^r \sigma_{i,i+1}$; and,
- $\sigma_{i,i+1}$ is the maximal (for inclusion) common substring of σ_i and σ_{i+1} satisfying the equalities above.

Proof. As $(\sigma_1, \ldots, \sigma_p)$ is a *p*-cover of μ , there exists a substring $\widetilde{\sigma_i}$ of σ_i and a substring $\widetilde{\sigma_{i+1}}$ of σ_{i+1} , such that $\widetilde{\sigma_{i+1}}\widetilde{\sigma_i}$ is a string. Moreover, as $(\sigma_1, \ldots, \sigma_p)$ is reduced, σ_i cannot be a substring of σ_{i+1} and vice-versa. Therefore, there exists at least one triplet of strings $(\sigma_i^\ell, \sigma_{i,i+1}, \sigma_{i+1}^r)$ such that $\sigma_i = \sigma_{i,i+1}\sigma_i^\ell$ and $\sigma_{i+1} = \sigma_{i+1}^r \sigma_{i,i+1}$.

Note that if $(\nu_i^{\ell}, \nu_{i,i+1}, \nu_{i+1}^r)$ is another triplet such that $\sigma_i = \nu_{i,i+1}\nu_i^{\ell}$ and $\sigma_{i+1} = \nu_{i+1}^r \nu_{i,i+1}$, and either $\nu_{i,i+1}$ is a substring of $\sigma_{i,i+1}$ or $\sigma_{i,i+1}$ is a substring of $\nu_{i,i+1}$.

Therefore, we justified the existence of a unique triplet $(\sigma_i^{\ell}, \sigma_{i,i+1}, \sigma_{i+1}^r)$ satisfying the required properties.



FIGURE 8. Illustration of Lemma 3.19 on the reduced 4-cover seen in Example 3.18.

Proposition 3.20. Let (Q, R) be a gentle quiver. For some $p \in \mathbb{N}^*$, consider p+1 strings, ρ_1, \ldots, ρ_p , and μ , of (Q, R) such that there exists a minimal epimorphism

$$f: \bigoplus_{i=1}^{p} \mathcal{M}(\rho_{i}) \longrightarrow \mathcal{M}(\mu)$$

such that $f_{|M(\rho_i)} = k_i \varphi_{(\sigma_i, \sigma'_i)}$, for some $k_i \in \mathbb{K}^{\times}$ and $(\sigma_i, \sigma'_i) \in [\rho_i, \mu]$. Then there exists a reduced p-cover (v_1, \ldots, v_p) of μ and a permutation $\tau \in \mathfrak{S}_p$ such that, for all $i \in \{1, \ldots, p\}$, $\nu_i = \sigma_{\tau(i)}^{\pm 1}$.

Proof. The result holds by the fact that f is an epimorphism. Indeed, up to reordering the string ρ_1, \ldots, ρ_p , and up to reversing the strings σ_i , it implies that $(\sigma_1, \ldots, \sigma_p)$ is a *p*-cover of ρ . As f is a minimal epimorphism, then this *p*-cover is reduced.

Theorem 3.21. Let (Q, R) be a representation-finite gentle quiver. For some $p \in \mathbb{N}^*$, consider p + 1 strings, ρ_1, \ldots, ρ_p , and μ , of (Q, R) such that there exists a minimal epimorphism

$$f: \bigoplus_{i=1}^{p} \mathcal{M}(\rho_i) \longrightarrow \mathcal{M}(\mu) \ ,$$

such that $f_{|M(\rho_i)} = k_i \varphi_{(\sigma_i,\sigma'_i)}$ for some $k_i \in \mathbb{K}^{\times}$ and $(\sigma_i,\sigma'_i) \in [\rho_i,\mu]$. Then we have $\operatorname{Ker}(f) = \bigoplus_{i=0}^{p} \operatorname{M}(\nu_i)$ where ν_0, \ldots, ν_p are strings of (Q, R) constructed from ρ_1, \ldots, ρ_p , and μ as follows:

- consider (v₁,..., v_p) to be the reduced p-cover of μ obtained by Proposition 3.20, and order the strings ρ₁,..., ρ_p according to this p-cover;
- for $i \in \{1, \ldots, p\}$, decompose ρ_i with respect to v_i : write $\rho_i = \rho_i^r v_i \rho_i^\ell$;
- consider the unique triplet (v^ℓ_i, v_{i,i+1}, v^r_{i+1}) satisfying the conditions given in Lemma 3.19.
- Then we define ν_0, \ldots, ν_p as follows:
 - $\nu_0 = \rho_1^{\ell'}$, where $\rho_1^{\ell'}$ is the string obtained from ρ_1^{ℓ} by deleting its last arrow (if ρ_1^{ℓ} is lazy, then there is no string ν_0);
 - $\nu_p = \rho_p^{r'}$ where $\rho_p^{r'}$ is the string obtained from ρ_p^r by deleting its first arrow (if ρ_p^r is lazy, then there is no string ν_p); and,
 - for $i \in \{1, \dots, p-1\}$, $\nu_i = \rho_i^r \upsilon_{i,i+1} \rho_{i+1}^{\ell}$.



FIGURE 9. Illustration of the construction of the string ν_0 (left), and the string ν_i for $i \in \{1, \ldots, p-1\}$ (right), pictured as orange dotted lines, thanks to the strings ρ_1, \ldots, ρ_p (purple dashed lines), and μ (blue line). On the right, the gray part corresponds to the common substring $v_{i,i+1}$ of μ, ρ_i, ρ_{i+1} and ν_i .

Proof. It is clear that $M(\nu_0)$ and $M(\nu_p)$ are summands of Ker(f). For $i \in \{1, \ldots, p\}$, by construction, we have $\operatorname{OvExt}(v_{i+1}^r v_{i,i+1} v_i^\ell, \nu_i) = \{\rho_i, \rho_{i+1}\}$. In particular, we have the following short exact sequence:

$$\mathbf{M}(\nu_i) \xrightarrow{\psi_1} \mathbf{M}(\rho_{i+1}) \oplus \mathbf{M}(\rho_i) \xrightarrow{\psi_2} \mathbf{M}(v_{i+1}^r v_{i,i+1} v_i^\ell)$$

where $\psi_1 = \begin{bmatrix} k_i \varphi_{(\upsilon_{i,i+1},\rho_{i+1}^{\ell},\upsilon_{i,i+1},\rho_{i+1}^{\ell})} \\ -k_{i+1}\varphi_{(\rho_i^r\upsilon_{i,i+1},\rho_i^r\upsilon_{i,i+1})} \end{bmatrix}$ and $\psi_2 = \begin{bmatrix} k_{i+1}\varphi_{(\upsilon_{i+1},\upsilon_{i+1})} & k_i\varphi_{(\upsilon_i,\upsilon_i)} \end{bmatrix}$. By noting that $f_{|M(\rho_i)\oplus M(\rho_{i+1})} = \psi_2$, it follows that $M(\nu_i)$ is a summand of Ker(f).

As f is a minimal epimorphism, we get the desired result.

3.7. Projective covers and projective resolutions. In this subsection, we recall some useful notions and results about projective objects of rep(Q, R). For more details, we refer the reader to [ASS06].

Let (Q, R) be a representation-finite gentle quiver, and $M \in \operatorname{rep}(Q, R)$. By Proposition 3.3, we can consider minimal projective resolution of any object in $\operatorname{rep}(Q, R)$. We denote by $\operatorname{gldim}(Q, R)$ the global dimension of $\operatorname{rep}(Q, R)$.

Note that to determine $\operatorname{gldim}(Q, R)$, it is enough to compute $\operatorname{pdim}(Y)$ for $Y \in$ $\operatorname{ind}(Q, R).$

Proposition 3.22 ([ASS06]). Let (Q, R) be a representation-finite gentle quiver. Then $gldim(Q, R) = \infty$ if and only if there exists a finite sequence of arrow $\alpha_1, \ldots, \alpha_p$, for some $p \in \mathbb{N}^*$ such that:

- $s(\alpha_1) = t(\alpha_p)$, and, for all $i \in \{1, ..., p-1\}$, $s(\alpha_{i+1}) = t(\alpha_i)$; and,
- $\alpha_1 \alpha_p \in R$, and, for all $i \in \{1, \ldots, p-1\}$, $\alpha_{i+1} \alpha_i \in R$.

4. Surface model for the representations of gentle algebras

For our purpose, Theorem 3.14 is a key result to describe all the resolving subcategories of $\operatorname{rep}(Q, R)$.

In this section, we introduce a variant of the topological model developed in [BCS19, OPS18, PPP18], practical to study gentle quivers and their representations. This geometric model will make calculations of resolving closures of any subset of ind $\ \operatorname{proj}(Q, R)$ easier to handle.

4.1. **Surfaces and dissections.** We set some conventions and we give some definitions on essential geometric materials in the following.

Convention 4.1. Whenever we talk about *surfaces*, we will always mean an oriented compact surface Σ with a finite number (which may be zero) of open discs removed. Denote the boundary of Σ by $\partial \Sigma$. Such a surface is determined, up to homeomorphism, by its genus and by the number of connected components of $\partial \Sigma$; we will refer to these connected components as the *boundary components* of Σ .

Definition 4.2. A marked surface is a pair (Σ, \mathcal{M}) , where Σ is a surface and \mathcal{M} is a finite set of marked points of Σ such that \mathcal{M} admits a bipartition $\{\mathcal{M}_{\circ}, \mathcal{M}_{\bullet}\}$ such that, on each boundary component of Σ :

- there is at least one marked point of each colour;
- marked points in \mathcal{M}_{\circ} and \mathcal{M}_{\bullet} alternate.

Note that marked points are also allowed in the interior of the surface.

Definition 4.3. Let (Σ, \mathcal{M}) be a marked surface.

- A \circ -arc is a curve on (Σ, \mathcal{M}) joining two points in \mathcal{M}_{\circ} ; it is a continuous map from the closed interval [0, 1] to Σ with endpoints in \mathcal{M}_{\circ} , and such that the image of its interior is disjoint from \mathcal{M} .
- A o-arc is said to be *simple* if it does not intersect itself (except perhaps at its endpoints).
- A ◦-dissection of (Σ, M) is a collection Δ° of pairwise non-intersecting simple ◦-arcs which cut the surface into polygons (that is to say, into simply-connected regions), called the *cells* of the dissection, each of which contains exactly one marked point in M_•. The triplet (Σ, M, Δ°) is called a ◦-dissected marked surface.
- We define •-*arcs* and •-*dissections* the same way.
- Given a \circ -dissection or a •-dissection Δ of a marked surface (Σ, \mathcal{M}) , we denote by $\Gamma(\Delta)$ the set of cells of $(\Sigma, \mathcal{M}, \Delta)$.

Remark 4.4. Note that our definitions are the one of a *dualizable* dissection in [OPS18].

Example 4.5. Here are examples of \circ -dissections of an annulus (or a sphere with two boundary components), and a torus with one boundary component.



FIGURE 10. Examples of a o-dissection of an annulus (left), and another of a torus with one boundary component (right).

20

4.2. The gentle quiver of a dissection. Let (Σ, \mathcal{M}) be a marked surface.

Definition 4.6. Let Δ° be a \circ -dissection of (Σ, \mathcal{M}) . We define a quiver $\mathbf{Q}(\Delta^{\circ})$ and a set of relations $\mathbf{R}(\Delta^{\circ})$ as follows:

- (i) the vertices of $\mathbf{Q}(\Delta^{\circ})$ correspond bijectively to the arcs in Δ° ;
- (*ii*) for each configuration as drawn on the left in Figure 11 in Δ° (that is to say, for each common endpoint of a pair of arcs *i* and *j* around which *i* comes immediately after *j* in clockwise orientation), there is an arrow $j \to i$ in $\mathbf{Q}(\Delta^{\circ})$;
- (*iii*) for each configuration as drawn on the right in Figure 11 in Δ° with corresponding arrows $j \xrightarrow{\alpha} i$ and $k \xrightarrow{\beta} j$, the path $\alpha\beta$ is a relation in $\mathbf{R}(\Delta^{\circ})$.



FIGURE 11. Drawings representing the construction rules of $\mathbf{Q}(\Delta^{\circ})$ and $\mathbf{R}(\Delta^{\circ})$ described above; (*ii*) on the left and (*iii*) on the right.

The pair $(\mathbf{Q}(\Delta^{\circ}), \mathbf{R}(\Delta^{\circ}))$ is the bounded quiver associated to $(\Sigma, \mathcal{M}, \Delta^{\circ})$.

Convention 4.7. We choose to orient our marked surface clockwise around the green points.

Example 4.8. See in Figure 12 the quivers with relations associated with the dissection drawn in Figure 10.



FIGURE 12. Quivers with relations arising from the dissected surfaces of Figure 10

To construct the quiver with relations associated with the example on the right, we have to identify the two square points (blue) and the two cross points (orange).

Then, following the rules given previously, the quiver with relations $(\mathbf{Q}(\Delta^{\circ}), \mathbf{R}(\Delta^{\circ}))$ associated to this dissection is the one below.

$$1 \xrightarrow[]{\alpha_1} 2 \xrightarrow[]{\alpha_2} 3$$

The following result motivates our interest in the geometric model.

Theorem 4.9 ([BCS19, OPS18, PPP18]).

- (i) If Δ° is a \circ -dissection of the marked surface (Σ, \mathcal{M}) such that there are no \circ -marked points in the interior of Σ , then $(\mathbf{Q}(\Delta^{\circ}), \mathbf{R}(\Delta^{\circ}))$ is a gentle quiver.
- (ii) For any gentle quiver (Q, R), there exists a marked surface (Σ, M) without o-marked point in the interior of Σ and with a o-dissection Δ[°] such that (Q, R) is isomorphic to (Q(Δ[°]), R(Δ[°])). Moreover, (Σ, M) and Δ[°] are unique up to oriented homeomorphism of surfaces and homotopy of arcs.

An explicit recipe to construct $(\Sigma, \mathcal{M}, \Delta^{\circ})$ from the gentle quiver (Q, R) is given in [PPP18, Section 4.2]. In the following, we set $\mathcal{S}(Q, R) = (\Sigma, \mathcal{M}, \Delta^{\circ})$.

4.3. Accordions and indecomposable representations. The representations of gentle quivers are related to certain arcs on the associated o-dissected marked surface: these are the so-called *accordions*.

Let $(\Sigma, \mathcal{M}, \Delta^{\circ})$ be a \circ -dissected marked surface such that $\mathcal{M}_{\circ} \subset \partial \Sigma$.

Definition 4.10. A •-arc δ in $(\Sigma, \mathcal{M}, \Delta^{\circ})$ is an *accordion* if it satisfies the following conditions: Whenever δ enters a cell of Δ° by crossing an \circ -arc η ,

- (a) if it leaves the cell, it leaves it by crossing an \circ -arc ζ adjacent to η ;
- (b) the relevant segments of the arcs η , ζ and δ bound a disk that does not contain the unique marked point in \mathcal{M}_{\bullet} belonging to the cell.

See Figure 13 for a picture illustrating the rules (a) and (b).



FIGURE 13. Drawings representing the rules that an accordion must satisfy: on the left, δ satisfies the rule (a) and (b); on the right, δ does not satisfy the rule (b).

Example 4.11. See Figure 14 for an example of an accordion over an annulus, and an example of an accordion over a torus.



FIGURE 14. An example of an accordion over each \circ -dissection seen in Example 4.5.

Theorem 4.12 ([BCS19, OPS18, PPP17]). Consider $(\Sigma, \mathcal{M}, \Delta^{\circ})$ a \circ -dissected marked surface such that $\mathcal{M}_{\circ} \subset \partial \Sigma$, and let $(\mathbf{Q}(\Delta^{\circ}), \mathbf{R}(\Delta^{\circ}))$ be its associated gentle quiver. Then we have a one-to-one correspondence between accordions on $(\Sigma, \mathcal{M}, \Delta^{\circ})$ and string representations of $(\mathbf{Q}(\Delta^{\circ}), \mathbf{R}(\Delta^{\circ}))$.

Remark 4.13. We adapted Theorem 4.12 as we restrict ourselves to representationfinite gentle quivers. Indeed [BCS19, OPS18, PPP17] states a result which allows one to interpret geometrically all the elements of ind(Q, R).

Example 4.14. See Figure 15 for an illustration of the bijection from accordions in a \circ -dissected marked surface ($\Sigma, \mathcal{M}, \Delta^{\circ}$) to strings in the associated gentle quiver ($\mathbf{Q}(\Delta^{\circ}), \mathbf{R}(\Delta^{\circ})$).



FIGURE 15. Illustration of the one-to-one correspondence between accordions $\mathcal{S}(Q, R)$ and strings of (Q, R) via an example.

From now on, we denote by $\mathscr{A}(\Sigma, \mathcal{M}, \Delta^{\circ}) = \mathscr{A}$ the set of accordions on a given \circ -dissected marked surface $(\Sigma, \mathcal{M}, \Delta^{\circ})$. For any $\delta \in \mathscr{A}$, by abuse of notations with strings, we set $M(\delta)$ as its associated string representation of $(\mathbf{Q}(\Delta^{\circ}), \mathbf{R}(\Delta^{\circ}))$. Conversely, for any $M \in \operatorname{ind}(Q, R)$, we write $\gamma_{(M)}$ for its associated accordion on $\mathscr{S}(Q, R)$.

4.4. Morphisms and extensions. We recall some results allowing us to read morphisms and extensions between indecomposable representations of (Q, R) directly via the associated accordions on $\mathcal{S}(Q, R)$.

First we recall the result which allows us to describe the homomorphism space the accordions. The following proposition is the consequence of a transcription of Theorem 3.10 on \mathcal{A} .

Proposition 4.15. Let $(\Sigma, \mathcal{M}, \Delta^{\circ})$ be a \circ -dissected marked surface with $\mathcal{M}_{\circ} \subset \partial \Sigma$. For any pair $(\delta, \eta) \in \mathscr{A}$, a basis of Hom $(M(\delta), M(\eta))$ is given by crossings as depicted in Figure 16.



FIGURE 16. Illustration of a crossing corresponding to a basis element of Hom(M(δ), M(η)). The gray part is a part where all the segments of δ and all the ones of η , given by the cutting of Σ with $\Gamma(\Delta^{\circ})$, are homotopic.

Remark 4.16. Note that, given two accordions that are crossing, the orientation of the surface allows one to read the direction of the morphism between the two associated indecomposable representations.

For our purpose, the following proposition extracts a geometric interpretation of the kernel of any epimorphism $M \longrightarrow Y$ involving $M \in \operatorname{rep}(Q, R)$ and $Y \in \operatorname{ind}(Q, R)$.

Proposition 4.17. Let $(\Sigma, \mathcal{M}, \Delta^{\circ})$ be a \circ -dissected marked surface with $\mathcal{M}_{\circ} \subset \partial \Sigma$. Consider $p \in \mathbb{N}^*$ and $\delta_1, \ldots, \delta_p \in \mathscr{A}$. Let $\eta \in \mathscr{A}$ such that there exists a minimal epimorphism

$$f: \bigoplus_{i=1}^{p} \mathcal{M}(\delta_{i}) \longrightarrow \mathcal{M}(\eta)$$

such that $f_{|M(\delta_i)}$ is given by exactly one crossing between δ_i and η . Then $\text{Ker}(f) = \bigoplus_{i=0}^{p} M(\kappa_i)$ where $\kappa_0, \ldots, \kappa_p \in \mathscr{A}$ are constructed from $\delta_1, \ldots, \delta_p$, and η as it follows in Figure 17. Note that κ_0 is defined in a dual way to κ_p and thus the drawing of κ_p is dual to the one of κ_0 .



FIGURE 17. The two types of accordions κ_i representing the indecomposable summands of Ker(f).

Proof. This is direct transcription of Theorem 3.21 in the geometric model.

We also recall a result which allows us to read extensions between indecomposable representations in the geometric model.

Proposition 4.18. Let $(\Sigma, \mathcal{M}, \Delta^{\circ})$ be a \circ -dissected marked surface with $\mathcal{M}_{\circ} \subset \partial \Sigma$. Let $\delta, \eta \in \mathscr{A}$. We distinguish two types of extensions of $M(\delta)$ by $M(\eta)$ which are:

• Overlap extensions : whenever we have a nonsplit short exact sequence

 $\mathbf{M}(\eta) \longrightarrow E_1 \oplus E_2 \longrightarrow \mathbf{M}(\delta)$,

where $E_1, E_2 \in \operatorname{ind}(Q, R)$, then δ and η are intersecting each other, and $\gamma_{(E_1)}$ and $\gamma_{(E_2)}$ can be obtained from δ and η as pictured in Figure 18. By abuse of notation with strings of (Q, R), in such a case, we set $\operatorname{OvExt}(\delta, \eta) = \{\gamma_{(E_1)}, \gamma_{(E_2)}\}$; otherwise $\operatorname{OvExt}(\delta, \eta) = \emptyset$.



FIGURE 18. Illustration of an overlap extension.

• Arrow extension : whenever we have a nonsplit short exact sequence

$$\mathbf{M}(\eta) \longrightarrow E \longrightarrow \mathbf{M}(\delta)$$
,

where $E \in ind(Q, R)$, then we can construct $\gamma_{(E)}$ from δ and η as described in Figure 19. Similarly to overlap extensions, by abuse of notations with the strings of (Q, R), in such case, we set $\operatorname{ArExt}(\delta, \eta) = \{\gamma_{(E)}\}$; otherwise $\operatorname{ArExt}(\delta, \eta) = \emptyset$.



FIGURE 19. Illustration of an arrow extension. The gray part corresponds to one cell of $\Gamma(\Delta^{\circ})$.

Proof. This is a direct transcription of Theorem 3.14 in the geometric model.

4.5. **Projective perspective.** As we will use substantially the minimal projective resolution in the following, we first ask for a transcription of the fact that (Q, R) has a finite global dimension in the geometric model.

Proposition 4.19 ([OPS18],[LGH24]). Let (Q, R) be a gentle quiver. Consider $\mathcal{S}(Q, R) = (\Sigma, \mathcal{M}, \Delta^{\circ})$ its associated \circ -dissected marked surface. Then we have that $\operatorname{gldim}(Q, R) < \infty$ if and only if $\mathcal{M} \subset \partial \Sigma$.

Remark 4.20. In particular, the fact that $\mathcal{M}_{\bullet} \subset \partial \Sigma$ ensures that we avoid the pattern highlighted in Proposition 3.22.

Under those conditions, we introduce a geometric model based on the indecomposable projective objects of $\operatorname{rep}(Q, R)$.

Definition 4.21. Let $(\Sigma, \mathcal{M}, \Delta^{\circ})$ be a \circ -dssected marked surface. We define the *projective dissection* of $(\Sigma, \mathcal{M}, \Delta^{\circ})$ as the set $\operatorname{Prj}(\Delta^{\circ})$ of accordions corresponding to the indecomposable representations of $\operatorname{proj}(\mathbf{Q}(\Delta^{\circ}), \mathbf{R}(\Delta^{\circ}))$.

Proposition 4.22 ([BCS19, OPS18, PPP18]). For any \circ -dissected marked surface $(\Sigma, \mathcal{M}, \Delta^{\circ})$, the triplet $(\Sigma, \mathcal{M}, \operatorname{Prj}(\Delta^{\circ}))$ gives a \bullet -dissected marked surface. Moreover, if $\mathcal{M} \subset \partial \Sigma$, then there exists an oriented homeomorphism from $(\Sigma, \mathcal{M}, \Delta^{\circ})$ to $(\Sigma, \mathcal{M}, \operatorname{Prj}(\Delta^{\circ}))$.

Example 4.23. See Figure 20 for examples of $Prj(\Delta^{\circ})$ on our red thread examples.



FIGURE 20. Examples of $Prj(\Delta^{\circ})$ of the \circ -dissections seen in Example 4.5. For the torus example (right), we draw a part of its universal cover (obtained by pasting copies of the unit squares along sides that are identified), and the dashed red lines corresponds to the fundamental copy of all the projective accordions.

From now on, we consider \bullet -dissected marked surfaces by seeing them as the projective dissection of the associated \circ -dissected one.

In the remaining of this section, we introduce the notion of *neighboring projective* accordions of a given accordion γ in $(\Sigma, \mathcal{M}, \Delta^{\circ})$, which will reveal to be useful in the next sections.

Definition 4.24. Let (Q, R) be a representation-finite gentle quiver. Let δ be an accordion of $\mathcal{S}(Q, R)$. A *neighboring projective accordion* of δ is an accordion $\gamma_{(P)}$ in $\mathcal{S}(Q, R)$ associated to an indecomposable projective representation $P \in \operatorname{proj}_{\mathbb{K}}(Q, R)$ such that either :

- P and $M(\delta)$ have non trivial higher extension;
- *P* is a direct summand of some projective representations appearing in the minimal projective resolution $(P^*_{M(\delta)})$.

For any accordion γ of $\boldsymbol{S}(Q, R)$, write $N_{\text{proj}}(\delta)$ for the set of all the neighboring projective accordions of δ .

Proposition 4.25 ([OPS18],[LGH24]). Let (Q, R) be a representation-finite gentle quiver. Let δ be an accordion of $\mathcal{S}(Q, R)$. Then $N_{\text{proj}}(\delta)$ is the subset of $\text{Prj}(\Delta^{\circ})$ made of accordions η satisfying at least one of the following conditions:

- (i) the curve η crosses δ ;
- (ii) the curve η is part of the border of a cell crossed by δ containing one of its endpoint, and is a part of the path of projective curves that links the last projective crossed to the endpoint of δ
- (iii) the curve η and δ have a common endpoint v, and, at the vertex v, all the projective accordions are smaller than δ with respect to the counterclockwise order on v.

Remark 4.26. The projective accordions η from condition (*i*) are the ones such that $\operatorname{OvExt}(\delta, \eta)$, those from condition (*ii*) are coming from $\left(\mathbf{P}^*_{\mathrm{M}(\delta)}\right)$, and those from condition (*iii*) are the ones having non trivial higher extensions.

Example 4.27. In Figure 21, we represent a projective dissection of a \circ -dissection of a marked disc (drawn as thick red curves), and we consider an accordion δ (drawn as a dotted blue curve). The densely dotted thick red lines correspond to the accordions forming the set of its neighboring projective accordions N_{proj}(δ). We pictured the projective accordions appearing in (P^{*}_{M(δ)}) in Figure 22.



FIGURE 21. An example of $N_{proj}(\delta)$.

Corollary 4.28. Let $(\Sigma, \mathcal{M}, \Delta^{\circ})$ be a \circ -dissected marked surface with $\mathcal{M}_{\circ} \subset \partial \Sigma$. Consider $p \in \mathbb{N}^*$ and $\delta_1, \ldots, \delta_p \in \mathscr{A}$. Let $\eta \in \mathscr{A}$ such that the following epimorphism is the projective cover of $M(\eta)$:

$$f: \bigoplus_{i=1}^{p} \mathcal{M}(\delta_{i}) \longrightarrow \mathcal{M}(\eta)$$

Then the syzygy $\operatorname{Ker}(f) = \Omega(M(\eta)) = \bigoplus_{i=0}^{p} \operatorname{M}(\kappa_{i})$ where $\kappa_{0}, \ldots, \kappa_{p} \in \mathscr{A}$ are constructed from $\delta_{1}, \ldots, \delta_{p}$, and η as shown in Figure 17. The modules $M(\kappa_{0})$ and $M(\kappa_{p})$ are the only indecomposable summand of the syzygy that can be nonprojective. The nonprojective summands of the higher syzygies are forming the fan, depicted in Figure 23, in the large cells of $\operatorname{N}_{\operatorname{proj}}(\eta)$.

Proof. The first statement is a corollary of Proposition 4.17. The second part is obtained by iteration of the result using Proposition 4.22 which describes the projective dissection as obtained from the standard dissection through a counterclockwise



FIGURE 22. Explicit calculation of the projective accordions $\eta \in N_{proj}(\delta)$ of Figure 21 such that $M(\eta)$ is a summand of projective representations in $(P^*_{M(\delta)})$.



FIGURE 23. The nonprojective summands $M(\omega_i)$ of the higher syzygy $\Omega^i(M(\eta))$ of $M(\eta)$.

rotation and the accordions appearing in the projective cover of $M(\delta)$ are lying in $N_{proj}(\delta)$.

Remark 4.29. We recover the result of [BS21], but here we expressed it in our geometric model.

Notation 4.30. In the following, for any $\delta \in \mathscr{A}$, we denote by $N_{\text{proj}}(\delta)_0$ the subset of \mathcal{M}_{\bullet} made of the unions of the endpoints of accordions in $N_{\text{proj}}(\delta)$.

5. Resolving posets for gentle trees

From now on, we focus on a precise familly of gentle quivers, called *gentle trees*. We exhibit a poset structure using monogeneous resolving subcategories.

Convention 5.1. In this section all epimorphisms are assumed *minimal*. We note that an additively closed subcategory of rep(Q, R) is closed under kernels of epimorphisms if and only if it is closed under kernels of minimal epimorphisms.

5.1. Generalities on gentle trees. Let Q be a quiver. A reduced walk $\rho = \alpha_k^{\varepsilon_k} \cdots \alpha_1^{\varepsilon_1}$ is cyclic whenever $s(\rho) = t(\rho)$. We say that Q is a *tree* if Q does not admit a reduced cyclic walk.

Definition 5.2. A *gentle tree* is a gentle quiver (Q, R) such that Q is a tree.

Example 5.3. See in Figure 24 for an example of a gentle tree.



FIGURE 24. An example of a gentle tree. This is the gentle quiver corresponding to the •-dissected marked surface given in Figure 21.

The following theorem gives a complete description of the marked surface associated to a gentle tree.

Theorem 5.4 ([OPS18],[APS23]). Let (Q, R) be a gentle tree. Set $(\Sigma, \mathcal{M}, \Delta^{\circ}) = \mathcal{S}(Q, R)$. Then $\mathcal{M} \subset \partial \Sigma$ and Σ is homeomorphic to a disc.

The following two propositions can be seen as corollaries of Theorem 5.4, even though they are proved independently.

Proposition 5.5. Let (Q, R) be a gentle tree. For any $M \in ind(Q, R)$, and for any $q \in Q_0$, we have $dim(M_q) \leq 1$.

Remark 5.6. The previous proposition implies that, for a given gentle tree (Q, R):

- all the strings of (Q, R) are characterized by their vertex support; and,
- all the accordions of $\mathcal{S}(Q, R)$ are characterized by the o-arcs of the o-dissection they are crossing.

Proposition 5.7. Let (Q, R) be a gentle tree. For any $M, N \in ind(Q, R)$, then $dim(Hom(M, N)) \leq 1$.

Remark 5.8. We can infer from the previous result that, given a gentle tree (Q, R), the accordions of $\mathcal{S}(Q, R)$ are crossing at most once, up to homotopy.

The following notion will also be useful to introduce a poset structure on the set of non-projective indecomposable modules in the next section.

Definition 5.9. Let (Q, R) be a gentle algebra. A representation $X \in \operatorname{rep}(Q, R)$ is said to be *directing* whenever, for any $t \ge 1$, and for any sequence

$$X \xrightarrow{f_1} M_1 \xrightarrow{f_2} \cdots \xrightarrow{f_t} M_t,$$

where f_1, \ldots, f_t are nonzero nonisomorphisms between $M_1, \ldots, M_t \in \operatorname{ind}_{\mathbb{K}}(Q, R)$, we have that M_t and X are not isomorphic.

Theorem 5.10 ([ASS06, Chapiter IX]). Let (Q, R) be a gentle tree. Any $X \in \operatorname{rep}(Q, R)$ is directing, and $\operatorname{AR}(Q, R)$ is acyclic.

We also give a useful corollary of Proposition 5.7 which allows us to focus only on the basis elements of extensions between indecomposable representations.

Corollary 5.11. Let (Q, R) be a gentle tree. For any pair (ρ, μ) of strings of (Q, R), we have that dim $(\text{Ext}^1(M(\rho), M(\mu)))) \leq 1$

It implies the following result.

Corollary 5.12. Let (Q, R) be a gentle tree.

- (i) All the nonsplit exact sequences are the ones decribed in Proposition 3.13 using arrow and overlap extensions;
- (ii) For any pair (ρ, μ) of strings of (Q, R), $\# \operatorname{ArExt}(\rho, \mu) \in \{0, 1\}$ and $\# \operatorname{OvExt}(\rho, \mu) \in \{0, 2\}$; and,
- (iii) For any pair (ρ, μ) of strings of (Q, R), the sets $\operatorname{ArExt}(\rho, \mu)$ and $\operatorname{OvExt}(\rho, \mu)$, cannot be both nonempty.

5.2. Vocabulary on posets. In this subsection, we recall some notions on finite posets that will be discussed in the following. We refer the reader to [DP02, Sta11, CLM12] for further details.

Let (\mathfrak{A}, \leq) be a finite poset. A *cover relation* is a pair $(x, y) \in \mathfrak{A}^2$ such that $x \leq y, x \neq y$, and, for all $z \in \mathfrak{A}$, whenever $x \leq z \leq y$, we have x = z or y = z. The *Haase diagram* of (\mathfrak{A}, \leq) is a directed graph whose vertices are elements in \mathfrak{A} , and arrows $y \longrightarrow x$ are given by cover relations $x \leq y$.

Given $\mathfrak{B} \subset \mathfrak{A}$, we define a *maximal element* of \mathcal{B} as an element $x \in \mathfrak{B}$ such that, for any $y \in \mathfrak{B}$, if $y \ge x$ then y = x. We define dually *minimal elements* of \mathfrak{B} . Given $x, y \in \mathfrak{A}$, we say that x and y admit a *least upper bound* if there exists $z \in \mathfrak{A}$ such that

$$\forall t \in \mathfrak{A}, t \geq z \iff t \geq x \text{ and } t \geq y.$$

In such a case, it is called the *join* of x and y, and we denote it by $x \vee y$. We define dually the *greatest lower bound*, called the *meet* of x and y, and denoted by $x \wedge y$, whenever it exists. We say that (\mathfrak{A}, \leq) is a *lattice* whenever every pair $(x, y) \in \mathfrak{A}^2$ admits a join and a meet. Note that, as \mathfrak{A} is finite, if (\mathfrak{A}, \leq) is a lattice, then it is *complete*: (\mathfrak{A}, \leq) admits a unique minimal element, denoted by $\widehat{0}_{\mathfrak{A}}$, and a unique maximal element, denoted by $\widehat{1}_{\mathfrak{A}}$. We recall the following useful characterization of a lattice.

Proposition 5.13. Let $(\mathfrak{A}, \leqslant)$ be a finite poset. If:

(1) (\mathfrak{A},\leqslant) admits a unique maximal element $\widehat{1}_{\mathfrak{A}}$, and

(2) every pair $(x, y) \in \mathfrak{A}$ admits a greatest lower bound,

then $(\mathfrak{A}, \leqslant)$ is a (complete) lattice.

Remark 5.14. The dual esult also holds.

A lattice (\mathfrak{A}, \leq) is *distributive* whenever the meet and join operations are distributive; meaning that, for all $x, y, z \in \mathfrak{A}$:

- $x \land (y \lor z) = (x \land y) \lor (x \land z)$, and
- $(x \wedge y) \lor z = (x \lor z) \land (y \lor z).$

An order ideal of $(\mathfrak{A}, \leqslant)$ is a set I such that, for all $x \in I$, $\langle x \rangle := \{y \in \mathfrak{A} \mid y \leqslant x\} \subseteq I$. Note that \emptyset is an order ideal following our definition. We denote by $\mathscr{J}(\mathfrak{A}, \leqslant)$ the set of order ideals of $(\mathfrak{A}, \leqslant)$. We recall that $(\mathscr{J}(\mathfrak{A}, \leqslant), \subseteq)$ is a complete distributive lattice, where:

- the join operation is the union,
- the meet operation is the intersection,
- $\widehat{0}_{\mathscr{J}(\mathfrak{A},\leqslant)} = \emptyset$, and $\widehat{1}_{\mathscr{J}(\mathfrak{A},\leqslant)} = \mathcal{P}$.

We say that $I \in \mathscr{J}(\mathfrak{A}, \leq)$ is *generated by a subset* \mathfrak{B} if $I = \bigcup_{x \in \mathfrak{B}} \langle x \rangle$. Given a subset \mathfrak{B} of \mathfrak{A} , we set $\langle \mathfrak{B} \rangle = \bigcup_{x \in \mathfrak{B}} \langle x \rangle$ which is the order ideal generated by \mathfrak{B} .

Two elements $x, y \in \mathfrak{A}$ are said to be *comparable* whenever $x \leq y$ or $y \leq x$. Otherwise, we say that they are *non-comparable*. An *antichain* in (\mathfrak{A}, \leq) is a collection \mathcal{X} of elements in \mathfrak{A} that are pairwise non-comparable. We recall that any order ideal is generated by a unique antichain \mathcal{X}_I : if $I = \emptyset$, then $\mathcal{X}_I = \emptyset$; otherwise \mathcal{X}_I is the set of maximal elements in I.

A non minimal element $x \in \mathfrak{A}$ is said to be *join-irreducible* if, for all $(y, z) \in \mathfrak{A}^2$, whenever we have $x = y \lor z$, we get x = y or x = z. As (\mathfrak{A}, \leq) is a finite lattice, a join-irreducible element $x \in \mathfrak{A}$ is *completely join-irreducible*; meaning that, for all collection $(x_i)_{i \in I}$ of elements in \mathfrak{A} , if $x = \bigvee_{i \in I} x_i$ then there exists $i \in I$ such that $x_i = x$.

Proposition 5.15. Let (\mathfrak{A}, \leq) be a finite lattice, and $x \in \mathfrak{A}$. Then x is joinirreducible if and only if there exists exactly one element in \mathfrak{A} covered by x. 5.3. Resolving order on non-projective indecomposable representations. In this section, given a gentle tree (Q, R), we will construct a poset structure on the non-projective indecomposable representations of (Q, R) based on the monogeneous resolving subcategories they generate.

Definition 5.16. Let (Q, R) be a gentle quiver. We define the Hom-*relation*, denoted by \preccurlyeq , on ind(Q, R) as follows:

$$\forall X, Y \in \operatorname{ind}(Q, R), \ X \preccurlyeq Y \Longleftrightarrow \operatorname{Hom}(X, Y) \neq 0.$$

We write $\preccurlyeq_{\mathsf{Tr.}}$ for the transitive closure of \preccurlyeq . We also define the Res-*relation*, denoted by \triangleleft , on **ind \proj**(Q, R) as follows:

$$\forall X, Y \in \operatorname{ind} \operatorname{\mathsf{proj}}(Q, R), \ X \leqslant Y \Longleftrightarrow \operatorname{Res}(X) \subseteq \operatorname{Res}(Y).$$

We can easily check that the Hom-relation is reflexive on $\operatorname{ind}(Q, R)$, and the Resrelation is a pre-order on $\operatorname{ind} \operatorname{proj}(Q, R)$: it is reflexive and transitive. However, in general, neither of them is an order relation.

Example 5.17. Let (Q, R) be the following gentle quiver, known as the *Kronecker* quiver.

$$(Q,R) = 1 \xrightarrow{\alpha}_{\beta} 2$$

Fix $\lambda \in \mathbb{K}^{\times}$, and consider the following non-isomorphic non-projective indecomposable representations.

$$X = \mathbb{K} \xrightarrow{1} \mathbb{K} \quad \text{and} \quad Y = \mathbb{K}^2 \xrightarrow{\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}} \mathbb{K}^2$$

Then $X \leq Y$ by the fact that X appears as a summand of a representation in $\operatorname{Ext}^1(Y,Y)$. For an analogous reason, we also have $Y \leq X$.

One can notice that the gentle quiver given in the previous example is not a representation-finite gentle quiver.

Example 5.18. Let (Q, R) be the following gentle quiver.

$$(Q,R) = 1 \underbrace{\alpha}_{\beta} 2$$

Consider the following two non-isomorphic non-projective indecomposable representations.

$$X = \mathbb{K} \underbrace{\stackrel{0}{\swarrow}_{\swarrow}}_{0} 0 \qquad \qquad Y = 0 \underbrace{\stackrel{0}{\rightthreetimes}_{\rightthreetimes}_{\swarrow}}_{1} \mathbb{K}$$

Then $X \leq Y$, as $Y \cong \text{Ker}(\text{P}^0_X \twoheadrightarrow X)$, and $Y \leq X$ for a similar reason.

In the previous two examples, the Auslander–Reiten quiver of each gentle quiver is not acyclic. We will prove that, for gentle trees (Q, R), $\preccurlyeq_{\mathsf{Tr.}}$ and \preccurlyeq are order relations.

Lemma 5.19. Let (Q, R) be a gentle tree. Then $\preccurlyeq_{\mathsf{Tr.}}$ is an order on $\operatorname{ind}(Q, R)$. Moreover $\operatorname{AR}(Q, R)$ is the Hasse diagram of $(\operatorname{ind}(Q, R), \preccurlyeq_{\mathsf{Tr.}})$.

Proof. First of all, as \preccurlyeq is reflexive, we already know that $\preccurlyeq_{\text{Tr.}}$ is reflexive and transitive. Moreover $\preccurlyeq_{\text{Tr.}}$ is antisymmetric by Theorem 5.10.

By Proposition 5.7, for any $X, Y \in ind(Q, R)$, there exists at most one nonzero nonisomorphism, up to a multiplication by a scalar, in Hom(Y, X); meaning that, in AR(Q, R) there is at most one arrow from Y to X.

Assume that we have $X, Y \in ind(Q, R)$ such that $X \ncong Y$ and $X \preccurlyeq_{\mathsf{Tr.}} Y$ is a cover relation. Then $X \preccurlyeq Y$, and the nonzero nonisomorphism $f: Y \longrightarrow X$ must be irreducible. So we have an arrow from Y to X in the Auslander-Reiten quiver of (Q, R). Conversely, if we have an arrow from Y to X in the Auslander-Reiten quiver, it is obvious to check that $X \preccurlyeq_{\mathsf{Tr.}} Y$ and it is a cover relation. We get the desired result.

Lemma 5.20. Let (Q, R) be a gentle tree. Let $\mathcal{X}^0 = \mathcal{X} \subset \operatorname{ind} \operatorname{proj}(Q, R)$, and, for $i \in \mathbb{N}$, we denote by \mathcal{X}^i the subset of $\operatorname{ind} \operatorname{proj}(Q, R)$ obtained at the *i*th iteration of Algorithm 2.14. Then, for any $Y \in \mathcal{X}^{i+1} \setminus \operatorname{proj}(Q, R)$, there exists $X \in \mathcal{X}^i \setminus \operatorname{proj}(Q, R)$ such that $X \preccurlyeq_{\mathsf{Tr}} Y$.

Proof. Fix $i \in \mathbb{N}$. Consider $E \in \mathcal{X}^{i+1}$. If $E \in \mathcal{X}^i$, then we are done. Otherwise, we have that $E \in \mathcal{X}^{i+1} \setminus \mathcal{X}^i$. Following Algorithm 2.14, E can appear in at most two different cases.

(1) Assume that E is a summand of F which is in a short exact sequence

$$A \xrightarrow{f} F \xrightarrow{g} B ,$$

with $A, B \in \mathcal{X}^i$. If $B \in \operatorname{proj}(Q, R)$, then the short exact sequence splits and we get that $F \cong A \oplus B$. So $E \cong A \in \mathcal{X}^i$ or $E \cong B \in \mathcal{X}^i$. Otherwise, $B \in \mathcal{X}^i \setminus \operatorname{proj}(Q, R)$, and as g is a nonzero minimal epimorphism, we know that $E \preccurlyeq B$ and, so $E \preccurlyeq_{\mathsf{Tr}} B$;

(2) Assume that E is a summand of K in a short exact sequence

$$K \xrightarrow{f} \bigoplus_{\ell=1}^{p} A_{\ell} \xrightarrow{g} B$$
,

for some $p \in \mathbb{N}^*$, with $B, A_1, \ldots, A_p \in \mathcal{X}^i$. If $B \in \operatorname{proj}(Q, R)$, then the short exact sequence splits, and K is isomorphic to a summand of $\bigoplus_{i=1}^p A_p$. So $E \cong A_j \in \mathcal{X}^i$ for some $j \in \{1, \ldots, p\}$, which is a contradiction. Otherwise, as g is a minimal epimorphism and not an isomorphism, we have that $f_{|E}$ is a nonzero morphism. Therefore $E \preccurlyeq A_j$ for some $j \in \{1, \ldots, p\}$.

Corollary 5.21. Let (Q, R) be a gentle tree. Let $\mathcal{X} \subset \operatorname{ind} \backslash \operatorname{proj}(Q, R)$. Then, for any $Y \in \mathcal{X}^{\operatorname{Res}} \setminus \operatorname{proj}(Q, R)$, there exists $X \in \mathcal{X} \setminus \operatorname{proj}(Q, R)$ such that $Y \preccurlyeq_{\operatorname{Tr.}} X$.

Proof. By induction using Lemma 5.20 and by transitivity of $\preccurlyeq_{\mathsf{Tr.}}$.

We can state a more precise result which highlights a compatibility between \triangleleft and $\preccurlyeq_{\mathsf{Tr.}}$ on **ind** \ **proj**(Q, R).

Corollary 5.22. Let (Q, R) be a gentle tree. Let $X, Y \in \operatorname{ind} \operatorname{proj}(Q, R)$. If we have $X \leq Y$, then $X \leq_{\mathsf{Tr.}} Y$.

Proposition 5.23. Let (Q, R) be a gentle tree. Then the Res-relation is an order relation on ind $\operatorname{proj}(Q, R)$.

Proof. We only have to check that \leq is antisymmetric.

By Corollary 5.22, whenever we have $X \leq Y$ for some $X, Y \in \operatorname{ind} \backslash \operatorname{proj}(Q, R)$, we have $X \preccurlyeq_{\mathsf{Tr}} Y$. By Lemma 5.19, $\preccurlyeq_{\mathsf{Tr}}$ is an order on $\operatorname{ind}(Q, R)$, and so it is antisymmetric. Thus \leq is antisymmetric.

Remark 5.24. This is not a characterization of the gentle quivers (Q, R) such that \triangleleft is an order on **ind** \ **proj**(Q, R).

Definition 5.25. Let (Q, R) be a gentle quiver. Whenever $(ind \mid proj(Q, R), \triangleleft)$ is a poset, we call it the resolving poset of (Q, R).

We will see, in the following, and in futur works, that this poset is a crucial key in order to construct the lattice of resolving subcategories of a gentle tree (see [DSar]).

5.4. Resolving posets and posets of resolving subcategories. Given a gentle quiver (Q, R), we denote by **Res**(Q, R) the set of all the resolving subcategories of $\operatorname{rep}(Q, R).$

Proposition 5.26. For any representation-finite gentle quiver (Q, R), the pair $(\mathbf{Res}(Q, R), \subseteq)$ is a finite complete lattice.

Proof. It follows from Propositions 3.3 and 5.13.

Proposition 5.27. Let (Q, R) be a gentle tree. Then the map

$$\underline{\operatorname{Res}}: \begin{cases} (\operatorname{\textit{Res}}(Q,R),\subseteq) & \longrightarrow & (\mathscr{J}(\operatorname{ind} \backslash \operatorname{proj}(Q,R), \triangleleft),\subseteq) \\ \mathscr{R} & \longmapsto & \operatorname{ind} \backslash \operatorname{proj}(\mathscr{R}) \end{cases}$$

is an injective poset morphism.

Proof. It follows from Lemma 2.10, and by the fact that, for any $\mathscr{R} \in \mathbf{Res}(Q, R)$, we have that $(\operatorname{ind} \operatorname{proj}(\mathscr{R}))^{\operatorname{Res}} = \operatorname{ind} \operatorname{proj}(\mathscr{R}).$

Lemma 5.28. Let (Q, R) be a gentle tree, and $X \in \operatorname{ind} \operatorname{proj}(Q, R)$. Set $\mathcal{Y}_X =$ ind \ proj(Res(X)) \ {X}. Then add($\mathcal{Y}_X \cup \operatorname{proj}(Q, R)$) $\in \operatorname{Res}(Q, R)$.

Proof. The case $\mathcal{Y}_X = \emptyset$ is obvious. Assume that $\mathcal{Y}_X \neq \emptyset$. By Proposition 5.27, we get that X^{Res} is the ideal generated by X in $(\text{ind} \setminus \text{proj}(Q, R), \triangleleft)$. Therefore \mathcal{Y}_X is also an ideal of $(\operatorname{ind} \operatorname{proj}(Q, R), \triangleleft)$. We must show that $(\mathcal{Y}_X)^{\operatorname{Res}} = \mathcal{Y}_X$. Assume, by contradiction, that $(\mathcal{Y}_X)^{\operatorname{Res}} \supseteq \mathcal{Y}_X$. On one hand, there exists $Z \in (\mathcal{Y}_X)^{\operatorname{Res}} \setminus \mathcal{Y}_X$, and, by Lemma 2.10, we have that

 $\{X\} \cup \mathcal{Y}_X = X^{\operatorname{Res}} \supseteq (\mathcal{Y}_X)^{\operatorname{Res}}$. Therefore $Z \cong X$.

On the other hand, by using Corollary 5.21, As $Z \in (\mathcal{Y}_X)^{\text{Res}}$ there exists $Y \in \mathcal{Y}_X$ such that $Z \preccurlyeq_{\mathsf{Tr}} Y$. By Corollary 5.22 by the fact that $Y \preccurlyeq X$, we get that $X \cong Z \preccurlyeq_{\mathsf{Tr.}} Y \preccurlyeq_{\mathsf{Tr.}} X$. Therefore $X \cong Y \in \mathcal{Y}_X$, which raises a contradiction.

Theorem 5.29. Let (Q, R) be a gentle tree. Then the join-irreducible elements in $(\operatorname{Res}(Q, R), \subseteq)$ are exactly given by the monogeneous resolving subcategories, and $\operatorname{proj}(Q, R)$.

Proof. Let $\mathscr{R} \in \mathbf{Res}(Q, R)$ be a join-irreducible element. We consider \mathcal{X} a subset of non-projective indecomposable representations in \mathscr{R} . If $\mathcal{X} = \emptyset$, then $\mathscr{R} = \operatorname{proj}(Q, R)$ and we are done. Otherwise, we can write that $\mathscr{R} = \bigvee_{X \in \mathcal{X}} \operatorname{Res}(X)$ as $\mathscr{R} = \operatorname{Res}(\mathcal{X})$. By hypothesis, it implies that \mathscr{R} is monogeneous.

We still have to show that the monogeneous resolving subcategories are joinirreducible. Let $X \in \operatorname{ind} \operatorname{proj}(Q, R)$. Set $\mathcal{Y}_X = \operatorname{ind} \operatorname{proj}(\operatorname{Res}(X)) \setminus \{X\}$. By Lemma 5.28, we showed that $\operatorname{Res}(X)$ covers $\operatorname{Res}(\mathcal{Y}_X) = \operatorname{add}(\mathcal{Y}_X \cup \operatorname{proj}(Q, R))$ in $(\operatorname{Res}(Q, R), \subseteq)$. Indeed, this is the unique $\operatorname{add}(\mathcal{Y}_X \cup \operatorname{proj}(Q, R))$ resolving subcategory covered by $\operatorname{Res}(X)$ by construction. Thus, by Proposition 5.15, we get that $\operatorname{Res}(X)$ is join-irreducible in $(\operatorname{Res}(Q, R), \subseteq)$.

6. Monogeneous resolving subcategories for gentle trees

6.1. Monogeneous geometric resolving sets. Let (Q, R) be a gentle tree. In this section, we introduce monogeneous accordion sets on S(Q, R) which are closed under geometric operations coming from Propositions 4.17 and 4.18. In the next section, we show that they correspond bijectively to the sets of indecomposable representations appearing in monogeneous resolving subcategories of rep(Q, R).

Definition 6.1. Let (Q, R) be a gentle tree and $\mathcal{S}(Q, R) = (\Sigma, \mathcal{M}, \Delta^{\circ})$. Let $\delta \in \mathscr{A}$. We define a *coloration* of $N_{\text{proj}}(\delta)_0$ to be a partition of $N_{\text{proj}}(\delta)_0$ as follows :

- 1) As δ cuts the disc into two connected parts, we consider, on one side of the curve, points of $N_{\text{proj}}(\delta)_0$ are colored red \blacksquare , on the other side they are green \square . In the following, we choose arbitrarily to consider the red dots on the top of δ , and the green ones at the bottom.
- 2) The left endpoint of the curve is called the *source* and is colored red \blacksquare if all the accordions in $N_{proj}(\delta)$ sharing the same source extremity as δ admit a green endpoint \Box at the other end. It is colored orange \boxtimes otherwise.
- 3) The right endpoint of the curve is called the *target*, and is colored dually
- 4) In larger cells the coloration are changed : all the intermediate points except the second to last are colored orange ⊠

5) The last intermediate point in the large cells is finally colored in pink \times . We denote by:

- $N_{\text{proj}}(\delta)_0^{\blacksquare}$ the set of points in $N_{\text{proj}}(\delta)_0$ colored in red \blacksquare ;
- $N_{\text{proj}}(\delta)_0^{\boxtimes}$ the set of points in $N_{\text{proj}}(\delta)_0$ colored in orange \boxtimes ;
- $N_{\text{proj}}(\delta)_0^{\Box}$ the set of points in $N_{\text{proj}}(\delta)_0$ colored in green \Box ;
- $N_{\text{proj}}(\delta)_0^{\times}$ the set of points in $N_{\text{proj}}(\delta)_0$ colored in pink \times .

Remark 6.2. Note that the coloration of $N_{\text{proj}}(\delta)_0$ is unique up to exchanging the and \Box points. Moreover, it induces a laterality of the surface.

Convention 6.3. From now on, given such a coloration of $N_{\text{proj}}(\delta)_0$, for any nonprojective accordion whose endpoints are in $N_{\text{proj}}(\delta)_0$, we consider it oriented in the same way. That its source is always on the left of its target. **Example 6.4.** We consider Example 4.27. We give in Figure 25 a coloration of $N_{proj}(\delta)_0$. Note that the points • are points in $\mathcal{M}_{\bullet} \setminus N_{proj}(\delta)_0$.



FIGURE 25. An example of a coloration of $N_{proj}(\delta)_0$.

Definition 6.5. Let (Q, R) be a gentle tree, and $\mathcal{S}(Q, R) = (\Sigma, \mathcal{M}, \Delta^{\circ})$. For any $\delta \in \mathscr{A}$, we define the *monogeneous geometric resolving set of* δ as the set $\mathscr{R}(\delta) = \mathscr{R}'(\delta) \cup \operatorname{Prj}(\Delta^{\circ})$ where:

$$\mathscr{R}'(\delta) = \{\eta \in \mathscr{A} \mid s(\eta) \in \mathrm{N}_{\mathrm{proj}}(\delta)_0^{\blacksquare} \cup \mathrm{N}_{\mathrm{proj}}(\delta)_0^{\boxtimes}, t(\eta) \in \mathrm{N}_{\mathrm{proj}}(\delta)_0^{\square} \cup \mathrm{N}_{\mathrm{proj}}(\delta)_0^{\boxtimes} \}.$$

The monogeneous geometric resolving subcategory of δ , denoted by \mathscr{U}_{δ} , is the additive subcategory of rep(Q, R) generated by $\{M(\varsigma) \mid \varsigma \in \mathscr{R}(\delta)\}$.

Example 6.6. Let $(\Sigma, \mathcal{M}, \Delta^{\circ})$ be the \circ -dissection of the marked surface as seen in Example 4.27 and in Figure 25. Consider $\delta \in \mathscr{A}$ as previously. In Figure 26, we represent the accordions in $\mathscr{R}(\delta)$ with dotted and dashed lines. More precisely,

- the blue loosely-dotted line is the accordion δ ;
- the red densely dotted lines are the neighboring projectives of δ as seen in Example 4.27 and in Figure 25; and,
- the purple dash-dotted lines are accordions in $\mathscr{R}'(\delta)$.



FIGURE 26. An explicit calculation of $\mathscr{R}(\delta)$, done on Example 4.27. They are all the dashed lines drawn above.

Lemma 6.7. Let (Q, R) be a gentle tree, and $\mathcal{S}(Q, R) = (\Sigma, \mathcal{M}, \Delta^{\circ})$. Consider $\delta \in \mathscr{A}$. Then, for any pair $(\varsigma_1, \varsigma_2) \in \mathscr{R}(\delta)^2$, we have

$$\operatorname{OvExt}(\varsigma_1, \varsigma_2) \cup \operatorname{ArExt}(\varsigma_1, \varsigma_2) \subseteq \mathscr{R}(\delta).$$

Proof. Consider first the case of overlap extensions. Let $(\varsigma_1, \varsigma_2) \in \mathscr{R}(\delta)^2$ such that $\operatorname{OvExt}(\varsigma_1, \varsigma_2) \neq \emptyset$. By Proposition 4.18, the extension is involving at most two accordions v_1 and v_2 and as the four curves share a common part the curves are oriented following the orientation of δ thus their sources are either the source of ς_1 or the source of ς_2 and the targets either the target of ς_1 or the target of ς_2 . We schematize the situation in Figure 27. Up to exchanging the role of v_1 and v_2 ,



FIGURE 27. Accordions v_1 and v_2 involving in the overlap extension of ς_1 by ς_2 .

assume that $s(v_1) = s(\varsigma_2)$, $t(v_1) = t(\varsigma_1)$, $s(v_2) = s(\varsigma_1)$ and $t(v_2) = t(\varsigma_2)$. Then $v_1, v_2 \in \mathscr{A}$, and, as $s(\varsigma_i) \in \mathrm{N}_{\mathrm{proj}}(\delta)_0^{\square} \cup \mathrm{N}_{\mathrm{proj}}(\delta)_0^{\square}$ and $t(\varsigma_i) \in \mathrm{N}_{\mathrm{proj}}(\delta)_0^{\square} \cup \mathrm{N}_{\mathrm{proj}}(\delta)_0^{\square}$ for $i \in \{1, 2\}$, it occurs that $v_1, v_2 \in \mathscr{R}(\delta)$.

Consider the arrow extensions. Let $(\varsigma_1, \varsigma_2) \in \mathscr{R}(\delta)^2$ such that $\operatorname{ArExt}(\varsigma_1, \varsigma_2) \neq \emptyset$. The curve corresponding to the indecomposable summand of the arrow extension θ is such that it crosses all the projective curves crossed by ς_1 and ς_2 . Thus they have a common orientation induced by δ . The curve θ is such that $s(\theta) = s(\varsigma_1)$ and $t(\theta) = t(\varsigma_2)$ or the dual configuration. Up to renaming consider this configuration. As $\varsigma_1 \in \mathscr{R}(\delta), \ s(\varsigma_1) \in \operatorname{N}_{\operatorname{proj}}(\delta)_0^{\textcircled{o}} \cup \operatorname{N}_{\operatorname{proj}}(\delta)_0^{\textcircled{o}}$. Dually as $\varsigma_2 \in \mathscr{R}(\delta), \ t(\varsigma_2) \in \operatorname{N}_{\operatorname{proj}}(\delta)_0^{\textcircled{o}} \cup \operatorname{N}_{\operatorname{proj}}(\delta)_0^{\textcircled{o}}$.

Lemma 6.8. Let (Q, R) be a gentle tree, and $\mathcal{S}(Q, R) = (\Sigma, \mathcal{M}, \Delta^{\circ})$. Consider $\delta \in \mathscr{A}$. For any $\varsigma \in \mathscr{R}(\delta)$ a non projective summand of the syzygy, $\Omega(\varsigma)$ is such that $\Omega(\varsigma) \in \mathscr{R}(\delta)$

Proof. Let (Q, R) be a gentle tree, and $\mathcal{S}(Q, R) = (\Sigma, \mathcal{M}, \Delta^{\circ})$. Consider $\delta \in \mathscr{A}$. Let $\varsigma \in \mathscr{R}(\delta)$ and $\Omega(\varsigma)$ a non projective summand of the syzygy. It is computed through Corollary 4.28. It follows that $\Omega(\varsigma) \in \mathscr{R}(\delta)$

Proposition 6.9. Let (Q, R) be a gentle tree, and set $(\Sigma, \mathcal{M}, \Delta^{\circ}) = \mathcal{S}(Q, R)$. For any $\delta \in \mathscr{A}$ the monogeneous geometric resolving subcategory \mathscr{U}_{δ} is resolving.

Proof. By using Proposition 2.12 and 2.13, and by the fact that \mathscr{U}_{δ} is additive, It is enough to prove that $\operatorname{ind}(\mathscr{U}_{\delta})$ is closed under taking summands from non trivial extensions between indecomposable representations, and from kernels of epimorphisms whose image is an indecomposable representation.

By Proposition 4.18 and Corollary 4.28, we have the result from Lemma 6.7 and Lemma 6.8.

6.2. Monogeneous resolving subcategories. Let (Q, R) be a gentle quiver. We recall that a resolving subcategory $\mathscr{R} \subseteq \operatorname{rep}(Q, R)$ is monogeneous whenever there exists $X \in ind(Q, R)$ such that $Res(X) = \mathscr{R}$. In this section, we will prove that the monogeneous resolving subcategories of $\operatorname{rep}(Q, R)$ are exactly the monogeneous geometric resolving subcategories. Here is the more precise statement.

Theorem 6.10. Let (Q, R) be a gentle tree, and $\mathcal{S}(Q, R) = (\Sigma, \mathcal{M}, \Delta^{\circ})$. For any $\delta \in \mathscr{A}$, we have that $\mathscr{U}_{\delta} = \operatorname{Res}(\operatorname{M}(\delta))$.

Let us introduce some useful notations.

For any $\delta \in \mathscr{A}$ which is not contained in a cell of the \bullet -dissection ($\Sigma, \mathcal{M}, \operatorname{Prj}(\Delta^{\circ})$), define $t_{\text{cell}}(\delta)$ to be the pair (η, C) where $\eta \in \text{Prj}(\Delta^{\circ})$, and $C \in \Gamma(\text{Prj}(\Delta^{\circ}))$ admitting η as one of its edges, such that δ crosses η and $t(\delta)$ is a vertex of C. If $\delta \subset C$, then we set $t_{\text{cell}}(\delta) = (\emptyset, C)$. We define $s_{\text{cell}}(\delta)$ dually.

Given a $C \in \Gamma(\operatorname{Prj}(\Delta^{\circ}))$, we denote by $C(\delta)_{0}^{\boxtimes}$ the set of vertices of C in $\operatorname{N}_{\operatorname{proj}}(\delta)_{0}^{\boxtimes}$. We define similarly $C(\delta)_0^{\square}$, $C(\delta)_0^{\bullet}$ and $C(\delta)_0^{\times}$. Define $\vartheta_1^R, \vartheta_2^R \in \operatorname{Prj}(\Delta^\circ)$ such that:

- $t(\vartheta_1^R) = t(\delta)$ and ϑ_1^R covers δ with respect to the counterclockwise order around $t(\delta)$ on $Prj(\Delta^{\circ}) \cup \{\delta\}$; and,
- $t(\vartheta_2^R) = s(\vartheta_1^R)$ and ϑ_2^R covers ϑ_1^R with respect to the counterclockwise order around $s(\vartheta_1^R)$ on $\operatorname{Prj}(\Delta^\circ) \cup \{\delta\}$.

We define $w_R(\delta) = s(\vartheta_2^R)$. We define dually $w_L(\delta)$. See Figure 28 to visualize the different objects introduced.



FIGURE 28. Illustration of $s_{\text{cell}}(\delta) = (\eta_L, C_L)$ and $t_{\text{cell}}(\delta) =$ (η_R, C_R) given $\delta \in \mathscr{A}$.

To prove Theorem 6.10, as we already know that \mathscr{U}_{δ} is a resolving subcategory that contains $M(\delta)$, by additivity of both \mathscr{U}_{δ} and $\operatorname{Res}(M(\delta))$, it is sufficient to show that the indecomposable representations in \mathscr{U}_{δ} are in Res(M(δ)).

First of all, given $\delta \in \mathscr{A}$, we enumerate necessary conditions on \bullet -arcs of $\mathscr{S}(Q, R)$ to be accordions in $\mathscr{R}'(\delta)$

Lemma 6.11. Let $(\Sigma, \mathcal{M}, \Delta^{\circ})$ a \circ -dissected marked disc with $\mathcal{M} \subset \partial \Sigma$. Let $\delta \in$ \mathscr{A} . If $\delta \not\subseteq C$ for some $C \in \Gamma(\operatorname{Prj}(\Delta^{\circ}))$, then we set $s_{\operatorname{cell}}(\delta) = (\eta_L, C_L)$ and $t_{\text{cell}}(\delta) = (\eta_R, C_R);$ otherwise, we set $C = C_L = C_R$ the cell containing δ . The curve $\varsigma \in \mathscr{R}'(\delta)$ must satisfy all of the following assertions:

(i) if ς crosses η_L and $w_L(\delta) \neq t(\eta_L)$, then $s(\varsigma) = s(\delta)$, and, dually, if ς crosses η_R and $w_R(\delta) \neq s(\eta_R)$, then $t(\varsigma) = t(\delta)$;

- (ii) if $\varsigma \subset C_L$, then $s(\varsigma) = s(\delta)$, and, dually, if $\varsigma \subset C_R$, then $t(\varsigma) = t(\delta)$;
- (iii) if $s(\varsigma) \in N_{\text{proj}}(\delta)_0^{\blacksquare}$ and $t(\varsigma) \in C_R(\delta)_0$, then $t(\varsigma) = w_L(\delta)$, and, dually, if $t(\varsigma) \in N_{\text{proj}}(\delta)_0^{\square}$ and $s(\varsigma) \in C_R(\delta)_0$, then $s(\varsigma) = w_R(\delta)$;
- (iv) $s(\varsigma) \notin C_L(\delta)_0^{\boxtimes} \setminus \{s(\delta), t(\eta_L)\}, and t(\varsigma) \notin C_R(\delta)_0^{\boxtimes} \setminus \{t(\delta), s(\eta_R)\}.$

Proof. Recall that the green dissection is obtained through a clockwise rotation of the neighboring projectives. Thus Figure 28 gives that all the listed cases correspond to curves passing through a relation.

Now we will treat the case where δ is contained in a cell of $\Gamma(\Delta^{\circ})$.

Lemma 6.12. Let $(\Sigma, \mathcal{M}, \Delta^{\circ})$ a \circ -dissected marked disc with $\mathcal{M} \subset \partial \Sigma$. Let $\delta \in \mathscr{A} \setminus \operatorname{Prj}(\Delta^{\circ})$ such that $\delta \subset C$ for some $C \in \Gamma(\operatorname{Prj}(\Delta^{\circ}))$. Then

 $\mathscr{R}(\delta) \subseteq \{\gamma_{(E)} \mid E \in \operatorname{ind} (\operatorname{Res}(\mathcal{M}(\delta)))\}.$

Proof. We illustrate the situation in Figure 29. We can note that:



FIGURE 29. The case where δ is contained in a cell C of the •-dissected marked disc $(\Sigma, \mathcal{M}, \operatorname{Prj}(\Delta^{\circ}))$. All the dashed lines are •-arcs in $\mathscr{R}(\delta)$. The dual case is obtained by rotating the figure of 180°, by exchanging the role of $\operatorname{N}_{\operatorname{proj}}(\delta)_0^{\Box}$ and the one of $\operatorname{N}_{\operatorname{proj}}(\delta)_0^{\Box}$, and re-labelling $w_R(\delta)$ by $w_L(\delta)$.

- There is exactly one vertex in $N_{\text{proj}}(\delta)_0^{\times}$: up to duality, we assume that we can define $w_R(\delta)$;
- there is no $\varsigma \in \mathscr{A}$ such that ς crosses δ ;
- If $v \in N_{\text{proj}}(\delta)_0^{\bullet}$, then $v = s(\delta) = w_R(\delta)$: note that $N_{\text{proj}}(\delta)_0^{\bullet} \neq \emptyset$ if and only if $\#C(\delta)_0 = 3$.

By Lemma 6.11, if $\varsigma \in \mathscr{R}'(\delta)$ such that $\delta \neq \varsigma$, either:

- 1) if $s(\varsigma), t(\varsigma) \in N_{\text{proj}}(\delta)_0^{\boxtimes}$, then either $t(\varsigma) = t(\delta)$ or $s(\varsigma) = s(\delta)$: as we impose the existence of $w_R(\delta)$, we must have $t(\varsigma) = t(\delta)$; or,
- 2) if $t(\varsigma) \in N_{\text{proj}}(\delta)_0^{\square}$, then $s(\varsigma) = w_R(\delta)$.

If ς satisfies 1), as $\varsigma \notin \operatorname{Prj}(\Delta^{\circ})$ and $\delta \neq \varsigma$, then $s(\delta) \neq w_R(\delta)$. Consider the vertex $v_1 \in C(\delta)_0^{\boxtimes}$ define as the target of the greatest projective accordion that is smaller that δ with respect to the clockwise order on $s(\delta)$. Note that, in the projective resolution $(\operatorname{P}^*_{\mathrm{M}(\delta)})$ of $\mathrm{M}(\delta)$, $\operatorname{P}^0_{\mathrm{M}(\delta)} = \mathrm{M}(\nu)$ where $\nu \in \operatorname{Prj}(\Delta^{\circ})$ such that $s(\nu) = s(\delta)$ and $t(\nu) = v_1$. Then the accordion δ_1 such that $s(\delta_1) = v_1$ (and so $t(\delta_1) = t(\gamma_{(M)})$)

correspond to the indecomposable representation $M(\delta_1)$ which is isomorphic to the kernel of the unique (by Proposition 5.7) epimorphism $P_M^0 \to M$. As $\operatorname{Res}(M(\delta))$ is closed under taking kernels of epimorphisms, we have that $M(\delta_1) \in \operatorname{Res}(M(\delta))$. By induction, for any ς satisfying 1), we have $M(\varsigma) \in \operatorname{Res}(M(\delta))$.

If $\varsigma \notin \mathscr{R}'(\delta)$ satisfying 2), then $\varsigma \in \operatorname{ArExt}(\rho_{\varsigma}, \delta_R)$ where:

- $\delta_R \in \mathscr{A}$ such that $s(\delta_R) = w_R(\delta)$, and $t(\delta_R) = t(\delta)$; and,
- $\rho_{\varsigma} \in \mathscr{A}$ such that $s(\rho_{\varsigma}) = t(\delta)$ and $t(\rho_{\varsigma}) = t(\delta) \in \mathrm{N}_{\mathrm{proj}}(\gamma_{(M)})_0^{\Box}$.

As δ_R satisfies 1), then $M(\delta_R) \in \text{Res}(M(\delta))$. Moreover, $M(\rho_{\varsigma}) \in \text{proj}(Q, R)$. Therefore, by Theorem 3.14, and Definition 2.2, $M(\varsigma) \in \text{Res}(M(\delta))$.

Thus we get the wished result.

We now consider the case where δ is not contained in a cell of $\Gamma(\Pr(\Delta^{\circ}))$.

Lemma 6.13. Let $(\Sigma, \mathcal{M}, \Delta^{\circ})$ a \circ -dissected marked disc with $\mathcal{M} \subset \partial \Sigma$. Let $\delta \in \mathscr{A} \setminus \operatorname{Prj}(\Delta^{\circ})$ such that $\delta \nsubseteq C$ for all $C \in \Gamma(\operatorname{Prj}(\Delta^{\circ}))$. Then $\mathscr{R}(\delta) \subseteq \{\gamma_{(E)} \mid E \in \operatorname{ind}(\operatorname{Res}(\operatorname{M}(\delta)))\}.$

Proof. We illustrated a general situation, up to dualization, in Figure 28. As δ is not contained in a cell of $(\Sigma, \mathcal{M}, \operatorname{Prj}(\Delta^{\circ}))$, we can set $s_{\operatorname{cell}}(\delta) = (\eta_L, C_L)$ and $t_{\operatorname{cell}}(\delta) = (\eta_R, C_R)$.

Let us show first that any $\varsigma \in \mathscr{A}$ such that $s(\varsigma) = s(\delta)$ and $t(\varsigma) \in \mathrm{N}_{\mathrm{proj}}(\delta)_0^{\Box}$ is in $\{\gamma_{(E)} \mid E \in \mathrm{ind}(\mathrm{Res}(\mathrm{M}(\delta)))\}$. Consider ς as stated. There exists $\rho \in \mathrm{N}_{\mathrm{proj}}(\delta)$ and $t(\rho) = t(\varsigma)$. Different cases happen:

- if $s(\rho) \in C_R(\delta)_0$, then $t(\rho) = t(\delta)$: in this case, $\varsigma = \delta$ and we are done;
- else, if $s(\rho) = s(\delta)$, then $\rho = \varsigma$, and we are done;
- otherwise, ρ crosses δ , and $s(\rho) \in N_{\text{proj}}(\delta)_0^{\bullet}$: we have $\varsigma \in \text{OvExt}(\rho, \delta) \neq \emptyset$ (see Figure 30), and as $\text{Res}(M(\delta))$ is closed under extension, we also have $M(\varsigma) \in \text{Res}(M(\delta))$).

By an analogous argument, we can prove that any $\varsigma' \in \mathscr{A}$ such that $t(\varsigma') = t(\delta)$ and $s(\varsigma') \in \mathcal{N}_{\text{proj}}(\delta)^{\bullet}_{0}$ (denoted by w in Figure 30) is in $\{\gamma_{(E)} \mid E \in \operatorname{ind}(\operatorname{Res}(\mathcal{M}(\delta)))\}$. By



FIGURE 30. Accordions (in purple dashed lines) corresponding to accordions appearing in $\text{OvExt}(\rho, \delta)$.

repeating the procedure on the curves obtained, we have that $M(\varsigma'') \in \operatorname{Res}(M(\delta))$ for all $\varsigma'' \in \mathscr{A}$ such that $s(\varsigma'') \in \operatorname{N}_{\operatorname{proj}}(\delta)_0^{\square}$ and $t(\varsigma'') \in \operatorname{N}_{\operatorname{proj}}(\delta)_0^{\square}$.

Now we focus on $\varsigma \in \mathscr{R}'(\delta)$ such that $s(\delta) \neq s(\varsigma), t(\delta) \neq t(\varsigma)$ and either:

• $s(\varsigma), t(\varsigma) \in \mathcal{N}_{\text{proj}}(\gamma_{(M)})_0^{\boxtimes}$; or,

- $s(\varsigma) \in N_{\text{proj}}(\delta)_0^{\bullet}$ and $t(\varsigma) \in C_L$; or,
- $s(\varsigma) \in C_R$ and $t(\varsigma) \in N_{\text{proj}}(\delta)_0^{\square}$.

By defining $\delta_L^{\circ} \in \mathscr{A}$ such that $s(\delta_L^{\circ}) = s(\delta)$ and $t(\delta_L^{\circ}) = t(\eta_L)$, and by defining $\delta_R^{\circ} \in \mathscr{A}$ dually, then the previous nonprojective accordions ς appear either in $\mathscr{R}'(\delta_L^{\circ})$ or in $\mathscr{R}'(\delta_R^{\circ})$. As $\delta_L^{\circ} \subset C_L$ and $\delta_R^{\circ} \subset C_R$, by proving that $M(\delta_L^{\circ}), M(\delta_R^{\circ}) \in \text{Res}(M(\delta))$, the result follows by Lemma 6.12.

Consider ϑ the first accordion in $\operatorname{Prj}(\Delta^{\circ})$ such that both ϑ crosses δ , including the case where $s(\vartheta) = s(\delta)$, and $t(\vartheta) = t(\eta_R) \in C(\gamma_{(M)})_0^{\boxtimes}$ with respect to the clockwise order around $t(\eta_R)$ on $\operatorname{Prj}(\Delta^{\circ})$. We depict the situation in Figure 31. Then $\operatorname{M}(\vartheta)$



FIGURE 31. Reduction to the case where δ is contained in a cell of $\Gamma(\Pr(\Delta^{\circ}))$.

corresponds to one summand of $P^0_{M(\delta)}$ in the minimal projective resolution (P^{\bullet}_M) . In the kernel of the epimorphism $P^0_{M(\delta)} \twoheadrightarrow M(\delta)$, there is an indecomposable summand K such that $\gamma_{(K)} = \delta^{\circ}_R$, by Proposition 4.17. As $\text{Res}(M(\delta))$ is closed under kernels of epimorphisms, then $M(\delta^{\circ}_R) \in \text{Res}(M(\delta))$. By dual arguments, we also get that $M(\delta^{\circ}_L) \in \text{Res}(M(\delta))$.

Finally, we get the following result.

Proposition 6.14. Let (Q, R) be a gentle tree, and $\mathcal{S}(Q, R) = (\Sigma, \mathcal{M}, \Delta^{\circ})$. For any $\delta \in \mathscr{A}$, The category \mathscr{U}_{δ} is a subcategory of $\operatorname{Res}(\operatorname{M}(\delta))$.

Proof. To prove the result, as $\operatorname{Res}(M)$ and $\mathscr{U}_{\gamma_{(M)}}$ are additive, we only have to prove that indecomposable representations in \mathscr{U}_M are in $\operatorname{Res}(M)$. Therefore we will use the geometric model to do so.

Let $\mathcal{S}(Q,R) = (\Sigma, \mathcal{M}, \Delta^{\circ})$. As (Q,R) is a gentle tree, by Theorem 5.4, Σ is homeomorphic to a disc, and $\mathcal{M} \subset \partial \Sigma$.

If $M \in \operatorname{proj}(Q, R)$, then we have that $\mathscr{R}(\gamma_{(M)}) = \operatorname{Prj}(\Delta^{\circ})$ and we are done. In the remaining of the proof, we assume that $M \in \operatorname{ind} \operatorname{proj}(Q, R)$.

Either $\gamma_{(M)}$ is contained in a cell of $\Gamma(\operatorname{Prj}(\Delta^{\circ}))$ or not. By Lemmas 6.12 and 6.13, we get that $\mathscr{R}(\gamma_{(M)}) \subset \{\gamma_{(E)} \mid E \in \operatorname{ind}(\operatorname{Res}(M))\}$. Hence the result.

Proof of Theorem 6.10. This result follows from Propositions 6.9 and 6.14.

Acknowledgements

B.D. thanks the Institut des Sciences Mathematiques (UQAM) and the Engineering and Physical Sciences Research Council (EP/W007509/1)) for their partial funding support. The authors acknowledge the CHARMS program grant (ANR-19-CE40-0017-02) for their funding support.

The authors, together, thank Yann Palu and Baptiste Rognerud for the various discussions they had, and their advises, all along this work.

References

[AB69]	Maurice Auslander and Mark Bridger. Stable module theory. Memoirs of the American Mathematical Society, No. 04, American Mathematical Society, Providence, PL 1060.
[AH81]	Mathematical Society, No. 94. American Mathematical Society, Providence, R1, 1969. Ibrahim Assem and Dieter Happel. Generalized tilted algebras of type A_n . Comm. Algebra 9(20):2101–2125 1981
[AH82]	Ibrahim Assem and Dieter Happel. Erratum: "Generalized tilted algebras of type A_n '. Comm. Algebra. 10(13):1475, 1982.
[APS23]	Claire Amiot, Pierre-Guy Plamondon, and Sibylle Schroll. A complete derived invari- ant for gentle algebras via winding numbers and arf invariants. <i>Selecta Mathematica</i> , 29, 03 2023.
[AR91]	Maurice Auslander and Idun Reiten. Applications of contravariantly finite subcate- gories. Advances in Mathematics. 86(1):111–152, 1991
[ASS06]	Ibrahim Assem, Andrzej Skowroński, and Daniel Simson. Elements of the Representa- tion Theory of Associative Algebras: Techniques of Representation Theory, volume 1 of London Mathematical Society Student Texts. Cambridge University Press, 2006
[AT22]	Takahide Adachi and Mayu Tsukamoto. Hereditary cotorsion pairs and silting sub- categories in extriangulated categories. <i>Journal of Algebra</i> , 594:109–137, 2022.
[BCS19]	Karin Baur and Raquel Coelho Simões. A Geometric Model for the Module Category of a Gentle Algebra. <i>International Mathematics Research Notices</i> , 2021(15):11357–11392, 07 2019.
[BDM+20]	Thomas Brüstle, Guillaume Douville, Kaveh Mousavand, Hugh Thomas, and Emine Yıldırım. On the combinatorics of gentle algebras. <i>Canad. J. Math.</i> , 72(6):1551–1580, 2020.
[BR87]	Michael Charles Richard Butler and Claus Michael Ringel. Auslander-Reiten sequences with few middle terms and applications to string algebras. Communications in Algebra $15(1-2):145-179$ 1987
[BS80]	Michael Charles Richard Butler and Mostafa Shahzamanian. The construction of al- most split sequences, III: Modules over two classes of tame local algebras. <i>Mathema-</i> <i>tische Annalen</i> . 250:111–122, 1980.
[BS21]	Karin Baur and Sibylle Schroll. Higher extensions for gentle algebras. Bulletin des Sciences Mathématiques, 170:103010, 2021.
[CB89]	W.W Crawley-Boevey. Maps between representations of zero-relation algebras. <i>Journal of Algebra</i> , 126(2):259–263, 1989.
[CLM12]	Nathalie Caspard, Bruno Leclerc, and Bernard Monjardet. <i>Finite Ordered Sets: Concepts, Results and Uses.</i> Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2012.
[ÇPS21]	Ilke Çanakçı, David Pauksztello, and Sibylle Schroll. On extensions for gentle algebras. Canadian Journal of Mathematics, 73(1):249–292, 2021.
[DP02]	B. A. Davey and H. A. Priestley. <i>Introduction to Lattices and Order</i> . Cambridge University Press, 2 edition, 2002.
[DS87]	Piotr Dowbor and Andrzej Skowronski. Galois coverings of representation-infinite al- gebras. <i>Commentarii mathematici Helvetici</i> , 62:311–338, 1987.
[DSar]	Benjamin Dequêne and Michael Schoonheere. Resolving subcategories over gentle algebras ii: Monogenous resolving subcategories over gentle trees. ArXiv, 2025 (to appear).
[GP68]	Israel Moiseevich Gel'fand and V.A. Ponomarev. Indecomposable representations of the lorentz group. <i>Russian Mathematical Surveys</i> , 23(2):1, apr 1968.
[Kra88]	Henning Krause. Maps between tree and band modules. <i>Journal of Algebra</i> , 137(1):186–194, 1988.
[LGH24]	Yu-Zhe Liu, Hanpeng Gao, and Zhaoyong Huang. Homological dimensions of gentle algebras via geometric models. <i>Science China Mathematics</i> , 67(4):733–766, Apr 2024.
[OPS18]	Sebastian Opper, Pierre-Guy Plamondon, and Sibylle Schroll. A geometric model for the derived category of gentle algebras. <i>arXiv:1801.09659</i> , 2018.
	42

- [PPP17] Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon. Non-kissing complexes and tau-tilting for gentle algebras. *Memoirs of the American Mathematical Society*, 274, 07 2017.
- [PPP18] Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon. Non-kissing and non-crossing complexes for locally gentle algebras. *Journal of Combinatorial Algebra*, 2018.
- [Rum21] Wolfgang Rump. The acyclic closure of an exact category and its triangulation. Journal of Algebra, 565:402–440, 2021.
- [Sta11] Richard P. Stanley. Enumerative Combinatorics. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2 edition, 2011.
- [SW83] Andrzej Skowronski and Josef Waschbüsch. Representation-finite biserial algebras. Journal für die reine und angewandte Mathematik, 345:172–181, 1983.
- [Tak09] Ryo Takahashi. Modules in resolving subcategories which are free on the punctured spectrum. *Pacific Journal of Mathematics*, 241, 02 2009.
- [WW85] Burkhard Wald and Josef Waschbüsch. Tame biserial algebras. Journal of Algebra, 95(2):480–500, 1985.

(B. Dequêne) SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS *Email address*: B.D.Dequene@leeds.ac.uk

(M. Schoonheere) LAMFA, UNIVERSITÉ DE PICARDIE JULES VERNE Email address: michael.schoonheere@u-picardie.fr