

HANKEL DETERMINANTS FOR CONVOLUTION POWERS OF MOTZKIN NUMBERS

YING WANG¹ AND YINGRUI ZHANG^{2,*}

ABSTRACT. We evaluate the Hankel determinants of the convolution powers of Motzkin numbers for $r \leq 27$ by finding shifted periodic continued fractions, which arose in application of Sulanke and Xin's continued fraction method. We also conjecture some polynomial characterization of these determinants.

Mathematic subject classification: Primary 15A15; Secondary 05A15, 11B83.

Keywords: Hankel determinants; Continued fractions; Motzkin numbers.

1. INTRODUCTION

This paper is along the line of using generating functions to deal with Hankel determinants. The Hankel determinant of a generating function $A(x) = \sum_{n \geq 0} a_n x^n$ is defined by

$$H_n(A(x)) = \det(a_{i+j})_{0 \leq i, j \leq n-1}, \quad H_0(A(x)) = 1.$$

In recent years, a considerable amount of work has been devoted to Hankel determinant evaluations of various counting coefficients. Many of such Hankel determinants have attractive compact closed formulas, such as binomial coefficients, Catalan numbers [19], Motzkin numbers [1, 8], large and little Schröder numbers [7]. See [2–7, 10, 11, 14, 16–18, 20–23] for further references.

The classical method of continued fractions, either by J -fractions [17, 24], or by S -fractions [15], requires $H_n(A(x)) \neq 0$ for all n . There also have Gessel-Xin's [12], Han's [13] and Sulanke-Xin's [22] continued fraction method allow $H_n(A(x)) = 0$ for some values of n . Sulanke-Xin defined a quadratic transformation τ such that $H(F(x))$ and $H(\tau(F(x)))$ have simple connections. The shifted periodic continued fractions (of order q) of the form

$$F_0^{(p)}(x) \xrightarrow{\tau} F_1^{(p)}(x) \xrightarrow{\tau} \cdots \xrightarrow{\tau} F_q^{(p)}(x) = F_0^{(p+1)}(x)$$

were found in [25], which appear in Hankel determinants of many path counting numbers. Here p is an additional parameter. If one can guess an explicit formula for $F_0^{(p)}(x)$ by Maple, then their Hankel determinants can be easily computed. This is also the primary method of this paper.

We have know that the Hankel determinants for the convolution power of Catalan numbers exhibits some perfect results and Conjectures [26]. As we also know the close relationship between Motzkin numbers and Catalan numbers, it is necessary to investigate the Hankel

Date: March 3, 2025.

determinants for the convolution power of Motzkin numbers.

$$H_n(M(x)^r) = \det(M_n^{(r)})_{0 \leq i, j \leq n-1}$$

is the r -th convolution power of the well-known Motzkin numbers $M_n = \sum_{k \geq 0} \binom{n}{2k} C_k$. C_k is the Catalan numbers. The name comes after the generating function identity

$$\sum_{n \geq 0} M_n^{(r)} x^n = M(x)^r = \left(\sum_{n \geq 0} M_n x^n \right)^r. \quad (1)$$

In what follows, we will denote by $F(x, r) = M(x)^r$. The cases $r = 1$ are well-known [9].

$$H_n(F(x, 1)) = H_n(M(x)) = 1.$$

The cases $r = 2, 3$ are nice.

Theorem 1.

$$H_n(F(x, 2)) = \begin{cases} 1, & n \bmod 12 = 0, 1, 2, 3. \\ 0, & n \bmod 12 = 4, 5, 10, 11. \\ -1, & n \bmod 12 = 6, 7, 8, 9. \end{cases}$$

$$H_{3n}(F(x, 3)) = H_{3n+1}(F(x, 3)) = (-1)^n (n+1)^2, \quad H_{3n+2}(F(x, 3)) = 0.$$

When computing the $H_n(F(x, r))$ for $r \geq 3$, they exhibit distinct patterns. We need to discuss the cases where the $r \equiv 0 \pmod{3}$, or not.

For $r \equiv 0 \pmod{3}$, we have the following Theorem and Conjectures:

Theorem 2.

$$\begin{aligned} H_{6n}(F(x, 6)) &= H_{6n+1}(F(x, 6)) = (-1)^n (n+1)^5, \\ H_{6n+2}(F(x, 6)) &= (-1)^n \frac{1}{10} (4n+5)(2n+3)(4n-3)(n+2)(n+1)^4, \\ H_{6n+3}(F(x, 6)) &= (-1)^n \frac{1}{45} (144n^2 + 72n - 155)(n+2)^2(2n+3)^2(n+1)^3, \\ H_{6n+4}(F(x, 6)) &= (-1)^{n+1} \frac{1}{45} (n+2)^3(2n+3)^2(144n^2 + 792n + 925)(n+1)^2, \\ H_{6n+5}(F(x, 6)) &= (-1)^n \frac{1}{10} (4n+15)(4n+7)(n+2)^4(2n+3)(n+1). \end{aligned}$$

Conjecture 3. For $r \equiv 0 \pmod{3}$, we have

$$\begin{aligned} H_{rn}(F(x, r)) &= H_{rn+1}(F(x, r)) = (-1)^n (n+1)^{r-1}. \\ H_{rn+2}(F(x, r)) + H_{rn-1}(F(x, r)) &= 3^2 \alpha (n+1)^{r-1}. \end{aligned}$$

where $|\alpha| = \frac{1}{18} r(r-3)$.

For $r \equiv 1$ or $2 \pmod{3}$, the situation is more complex. We have developed software packages to compute $H_n(F(x, r))$ for $r \leq 27$ (this software package can also handle larger values of r , but there is not much necessity for it, as the method remains unchanged). Now we just present several results for $r = 4, 5, 7$, along with some Conjectures.

Theorem 4.

$$\begin{aligned}
H_{12n}(F(x, 4)) &= H_{12n+1}(F(x, 4)) = 1, \\
H_{12n+2}(F(x, 4)) &= 2(2n+1)(8n-1), \\
H_{12n+3}(F(x, 4)) &= -2(8n+9)(2n+1), \\
H_{12n+4}(F(x, 4)) &= H_{12n+5}(F(x, 4)) = -1, \\
H_{12n+6}(F(x, 4)) &= 4(n+1)(8n+5), \\
H_{12n+7}(F(x, 4)) &= H_{12n+10}(F(x, 4)) = -64(n+1)^2, \\
H_{12n+8}(F(x, 4)) &= H_{12n+9}(F(x, 4)) = 0, \\
H_{12n+11}(F(x, 4)) &= -4(n+1)(8n+11).
\end{aligned}$$

Theorem 5.

$$\begin{aligned}
H_{15n}(F(x, 5)) &= H_{15n+1}(F(x, 5)) = 1, \\
H_{15n+2}(F(x, 5)) &= 5(2n+1)(50n^2+15n-1), \\
H_{15n+3}(F(x, 5)) &= 25(25n^2+25n-2)(2n+1)^2, \\
H_{15n+4}(F(x, 5)) &= -5(50n^2+85n+34)(2n+1), \\
H_{15n+5}(F(x, 5)) &= H_{15n+6}(F(x, 5)) = 1, \\
H_{15n+7}(F(x, 5)) &= 5(n+1)(100n^2+120n+33), \\
H_{15n+8}(F(x, 5)) &= 25(n+1)^2(100n^2+100n+17), \\
H_{15n+9}(F(x, 5)) &= H_{15n+12}(F(x, 5)) = 1000(n+1)^3, \\
H_{15n+10}(F(x, 5)) &= H_{15n+11}(F(x, 5)) = 0, \\
H_{15n+13}(F(x, 5)) &= 25(n+1)^2(100n^2+300n+217), \\
H_{15n+14}(F(x, 5)) &= -5(n+1)(100n^2+280n+193).
\end{aligned}$$

Theorem 6.

$$\begin{aligned}
H_{21n}(F(x, 7)) &= H_{21n+1}(F(x, 7)) = (-1)^n, \\
H_{21n+2}(F(x, 7)) &= (-1)^{n+1} \frac{7}{30} (2n+1) (374556n^4 + 385532n^3 + 127449n^2 + 13363n + 60), \\
H_{21n+3}(F(x, 7)) &= (-1)^n \frac{49}{180} (29647548n^6 + 54252996n^5 + 27993259n^4 + 2344748n^3 \\
&\quad - 1393217n^2 - 227934n - 1080) (2n+1)^2, \\
H_{21n+4}(F(x, 7)) &= (-1)^{n+1} \frac{343}{180} (2n+1)^3 (4235364n^6 + 12706092n^5 + 12161065n^4 \\
&\quad + 3145310n^3 - 906059n^2 - 361032n + 3420), \\
H_{21n+5}(F(x, 7)) &= (-1)^{n+1} \frac{49}{180} (29647548n^6 + 123632292n^5 + 201441499n^4 + 160049288n^3 \\
&\quad + 61715353n^2 + 9000600n - 123300) (2n+1)^2, \\
H_{21n+6}(F(x, 7)) &= (-1)^{n+1} \frac{7}{30} (374556n^4 + 1112692n^3 + 1218189n^2 + 583163n + 103170) (2n+1),
\end{aligned}$$

$$\begin{aligned}
H_{21n+7}(F(x, 7)) &= H_{21n+8}(F(x, 7)) = (-1)^{n+1}, \\
H_{21n+9}(F(x, 7)) &= (-1)^n \frac{7}{30} (n+1) (749112n^4 + 1850828n^3 + 1698242n^2 + 686273n + 103230), \\
H_{21n+10}(F(x, 7)) &= (-1)^{n+1} \frac{49}{180} (n+1)^2 (118590192n^6 + 375938976n^5 + 459474568n^4 \\
&\quad + 261223312n^3 + 59779951n^2 - 1682814n - 1959840), \\
H_{21n+11}(F(x, 7)) &= (-1)^n \frac{343}{90} (n+1)^3 (16941456n^6 + 84707280n^5 + 148881208n^4 + 115228792n^3 \\
&\quad + 34672057n^2 - 1047963n - 1697400), \\
H_{21n+12}(F(x, 7)) &= (-1)^n \frac{2401}{15} (n+1)^4 (115248n^4 + 460992n^3 + 587608n^2 + 288512n + 43695), \\
H_{21n+13}(F(x, 7)) &= H_{21n+16}(F(x, 7)) = (-1)^{n+1} 537824 (n+1)^5, \\
H_{21n+14}(F(x, 7)) &= H_{21n+15}(F(x, 7)) = 0, \\
H_{21n+17}(F(x, 7)) &= (-1)^n \frac{2401}{15} (n+1)^4 (115248n^4 + 460992n^3 + 587608n^2 + 217952n - 26865), \\
H_{21n+18}(F(x, 7)) &= (-1)^{n+1} \frac{343}{90} (n+1)^3 (16941456n^6 + 118590192n^5 + 318295768n^4 \\
&\quad + 398162632n^3 + 205815337n^2 - 2633505n - 27024030), \\
H_{21n+19}(F(x, 7)) &= (-1)^{n+1} \frac{49}{180} (n+1)^2 (118590192n^6 + 1047143328n^5 + 3815496328n^4 \\
&\quad + 7351444912n^3 + 7906357711n^2 + 4503507834n + 1062057240), \\
H_{21n+20}(F(x, 7)) &= (-1)^{n+1} \frac{7}{30} (n+1) (749112n^4 + 4142068n^3 + 8571962n^2 + 7868343n + 2702820).
\end{aligned}$$

Conjecture 7. For $r \equiv 1$ or $2 \pmod{3}$, we have

$$\begin{aligned}
H_{3rn}(F(x, r)) &= H_{3rn+1}(F(x, r)) = H_{3rn+r}(F(x, r)) = H_{3rn+r+1}(F(x, r)) = \alpha. \\
H_{3rn+2r}(F(x, r)) &= H_{3rn+2r+1}(F(x, r)) = 0. \\
H_{3rn+2r-1}(F(x, r)) + H_{3rn+2r+2}(F(x, r)) &= \beta ((2r)(n+1))^{r-2}. \\
H_{3rn+2}(F(x, r)) + H_{3rn-1}(F(x, r)) &= \frac{1}{2} \gamma r(r-3).
\end{aligned}$$

where $|\alpha| = |\beta| = |\gamma| = 1$.

Conjecture 8. For $r \equiv 0 \pmod{3}$, the shifted periodic of $H_n(F(x, r))$ is r . For $r \equiv 1$ or $2 \pmod{3}$, the shifted periodic of $H_n(F(x, r))$ is $3r$.

Our main result is the following.

Theorem 9. The above Conjectures hold for $r \leq 27$.

The paper is organized as follows. Section 2.1 introduces Sulanke-Xin's continued fraction method, especially the quadratic transformation τ . Section 3 detailed description of the process for calculating $H_n(F(x, 3))$ and $H_n(F(x, 4))$ by using the Sulanke-Xin method. This is also the process in developing the software package.

2. MAIN TOOL

We will introduce the continued fraction method of Sulanke and Xin, especially their quadratic transformation τ in [22]. This is the main tool of this paper.

2.1. Sulanke-Xin's quadratic transformation τ . This subsection is copied from [25]. We include it here for the reader's convenience.

Suppose the generating function $F(x)$ is the unique solution of a quadratic functional equation which can be written as

$$F(x) = \frac{x^d}{u(x) + x^k v(x) F(x)}, \quad (2)$$

where $u(x)$ and $v(x)$ are rational power series with nonzero constants, d is a nonnegative integer, and k is a positive integer. We need the unique decomposition of $u(x)$ with respect to d : $u(x) = u_L(x) + x^{d+2}u_H(x)$ where $u_L(x)$ is a polynomial of degree at most $d+1$ and $u_H(x)$ is a power series. Then Propositions 4.1 and 4.2 of [22] can be summarized as follows.

Proposition 10. *Let $F(x)$ be determined by (2). Then the quadratic transformation $\tau(F)$ of F defined as follows gives close connections between $H(F)$ and $H(\tau(F))$.*

- i) *If $u(0) \neq 1$, then $\tau(F) = G = u(0)F$ is determined by $G(x) = \frac{x^d}{u(0)^{-1}u(x) + x^k u(0)^{-2}v(x)G(x)}$, and $H_n(\tau(F)) = u(0)^n H_n(F(x))$;*
- ii) *If $u(0) = 1$ and $k = 1$, then $\tau(F) = x^{-1}(G(x) - G(0))$, where $G(x)$ is determined by*

$$G(x) = \frac{-v(x) - xu_L(x)u_H(x)}{u_L(x) - x^{d+2}u_H(x) - x^{d+1}G(x)},$$

and we have

$$H_{n-d-1}(\tau(F)) = (-1)^{\binom{d+1}{2}} H_n(F(x));$$

- iii) *If $u(0) = 1$ and $k \geq 2$, then $\tau(F) = G$, where $G(x)$ is determined by*

$$G(x) = \frac{-x^{k-2}v(x) - u_L(x)u_H(x)}{u_L(x) - x^{d+2}u_H(x) - x^{d+2}G(x)},$$

and we have

$$H_{n-d-1}(\tau(F)) = (-1)^{\binom{d+1}{2}} H_n(F(x)).$$

 3. THE CALCULATIONS OF $H_n(F(x, 3))$ AND $H_n(F(x, 4))$

3.1. For $r = 3$, we have the functional equation

$$F(x, 3) = \frac{1}{1 - 3x + 2x^3 - x^6 F(x, 3)}.$$

Proof of Theorem 1. We apply Proposition 10 to $F_0 := F(x, 3)$ by repeatedly using the transformation τ . This results in a shifted periodic continued fractions of order 2:

$$F_0(x) \xrightarrow{\tau} F_1^{(1)}(x) \xrightarrow{\tau} F_2^{(1)}(x) \xrightarrow{\tau} F_3^{(1)}(x) = F_1^{(2)}(x) \cdots .$$

We obtain

$$H_k(F_0) = H_{k-1}(F_1^{(1)}). \quad (3)$$

For $p \geq 1$, computer experiments suggest us to define

$$F_1^{(p)} = -\frac{x(x^3 + 3p(p+1)x - p(p+1))}{p^2x^2F_1^{(p)} + 2px^3 + 3p^2x - p^2}.$$

Then the results can be summarized as follows:

$$\begin{aligned} H_{k-1}(F_1^p) &= -\left(\frac{p+1}{p}\right)^{k-1} H_{k-3}(F_2^p) \\ H_{k-3}(F_2^p) &= \left(\frac{p}{p+1}\right)^{k-3} H_{k-4}(F_1^{p+1}). \end{aligned}$$

Combination of the above formulas gives the recursion

$$H_{k-1}(F_1^p) = -\left(\frac{p+1}{p}\right)^2 H_{k-4}(F_1^{p+1}).$$

Let $k-1 = 3n+j$, where $0 \leq j < 3$. We then deduce that

$$H_{3n+j}(F_1^1) = (-1)^n(n+1)^2 H_j(F_1^{n+1}). \quad (4)$$

The initial values are

$$H_0(F_1^{n+1}) = 1; \quad H_1(F_1^{n+1}) = 0; \quad H_2(F_1^{n+1}) = \left(\frac{n+2}{n+1}\right)^2.$$

Then the theorem follows by the above initial values, (3) and (4). \square

3.2. For $r = 4$, we have the functional equation

$$F(x, 4) = -\frac{1}{F(x, 4)x^8 + x^4 - 4x^3 - 2x^2 + 4x - 1}.$$

Proof of Theorem 4. We apply Proposition 10 to $F_0 := F(x, 4)$ by repeatedly using the transformation τ . This results in a shifted periodic continued fractions of order 10:

$$\begin{aligned} F_0(x) &\xrightarrow{\tau} F_1^{(1)}(x) \xrightarrow{\tau} F_2^{(1)}(x) \xrightarrow{\tau} F_3^{(1)}(x) \xrightarrow{\tau} F_4^{(1)}(x) \xrightarrow{\tau} F_5^{(1)}(x) \xrightarrow{\tau} F_6^{(1)}(x) \xrightarrow{\tau} F_7^{(1)}(x) \\ &\xrightarrow{\tau} F_8^{(1)}(x) \xrightarrow{\tau} F_9^{(1)}(x) \xrightarrow{\tau} F_{10}^{(1)}(x) \xrightarrow{\tau} F_{11}^{(1)}(x) = F_1^{(2)}(x) \cdots \end{aligned}$$

We obtain

$$H_k(G_0) = H_{k-1}(G_1^{(1)}). \quad (5)$$

For $p \geq 1$, computer experiments suggest us to define

$$\begin{aligned} F_1^{(p)} &= -\frac{1}{x^2F_1^{(p)} - x^4 + (16p-12)x^3 + (-64p^2 + 104p - 38)x^2 + 4x - 1} \times \\ &x^6 + (-8p+8)x^5 + (96p^2 - 148p + 52)x^4 + (-512p^3 + 1216p^2 - 928p + 220)x^3 + \\ &(1024p^4 - 3328p^3 + 3920p^2 - 1944p + 345)x^2 + (-128p^2 + 200p - 68)x + 32p^2 - 52p + 18. \end{aligned}$$

Then the results can be summarized as follows:

$$\begin{aligned}
 H_{k-1}(F_1^p) &= (32p^2 - 52p + 18)^{k-1} H_{k-2}(F_2^p) \\
 H_{k-2}(F_2^p) &= \left(-\frac{8p+1}{256p^3 - 704p^2 + 612p - 162} \right)^{k-2} H_{k-3}(F_3^p) \\
 H_{k-3}(F_3^p) &= \left(-1/2 \frac{8p-9}{(2p-1)(8p+1)^2} \right)^{k-3} H_{k-4}(F_4^p) \\
 H_{k-4}(F_4^p) &= (2(8p+1)(2p-1))^{k-4} H_{k-5}(F_5^p) \\
 H_{k-5}(F_5^p) &= (-4p(8p-3))^{k-5} H_{k-6}(F_6^p) \\
 H_{k-6}(F_6^p) &= (-4(8p-3)^{-2})^{k-6} H_{k-1}(F_1^p) H_{k-7}(F_7^p) \\
 H_{k-7}(F_7^p) &= -\left(1/16 \frac{8p-3}{p} \right)^{k-7} H_{k-10}(F_8^p) \\
 H_{k-10}(F_8^p) &= \left(1/16 \frac{8p+3}{p} \right)^{k-10} H_{k-11}(F_9^p) \\
 H_{k-12}(F_9^p) &= (-4(8p+3)^{-2})^{k-11} H_{k-12}(F_{10}^p)
 \end{aligned}$$

Combination of the above formulas gives the recursion

$$H_{k-1}(F_1^{(p)}) = H_{k-13}(F_1^{(p+1)}).$$

Let $k-1 = 12n+j$, where $0 \leq j < 11$. We then deduce that

$$H_{12n+j+1}(F) = H_{12n+j}(F_1^{(1)}) = H_j(F_1^{(n+1)}). \quad (6)$$

The initial values are

$$\begin{aligned}
 H_0(F_1^{n+1}) &= 1, & H_1(F_1^{n+1}) &= 32(n+1)^2 - 52n - 34, \\
 H_2(F_1^{n+1}) &= -2(8n+9)(2n+1), & H_3(F_1^{n+1}) &= H_4(F_1^{n+1}) = -1, \\
 H_5(F_1^{n+1}) &= 4(n+1)(8n+5), & H_6(F_1^{n+1}) &= 64(n+1)^2, \\
 H_7(F_1^{n+1}) &= H_8(F_1^{n+1}) = 0, & H_9(F_1^{n+1}) &= -64(n+1)^2, \\
 H_{10}(F_1^{n+1}) &= -4(n+1)(8n+11), & H_{11}(F_1^{n+1}) &= H_{12}(F_1^{n+1}) = 1.
 \end{aligned}$$

Then the theorem follows by the above initial values, (5) and (6). \square

Declaration of Interest Statement The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data Availability Data availability is not applicable to this article as no new data were created or analyzed in this study.

Acknowledgments The authors would like to thank zihao Zhang for his careful reading and very useful comments.

REFERENCES

- [1] M. Aigner. Motzkin numbers. *European Journal of Combinatorics*, 19(6) (1998) 663–675.
- [2] M. Aigner. Catalan-like numbers and determinants, *J. Combin. Theory Ser. A* 87 (1999) 33–51.
- [3] M. Aigner. Catalan and other numbers: a recurrent theme, in: H. Crapo, D. Senato (Eds.), *Algebraic Combinatorics and Computer Science*, Springer, Berlin, 2001, pp. 347–390.
- [4] P. Barry. Some conjectures on the ratio of Hankel transforms for sequences and series reversion, 17 Jan. 2007. arxiv:math/0701483v1[math.CO].
- [5] P. Barry. Generalized Catalan numbers, Hankel transforms and Somos-4 sequences, *J. Integer Seq.* 13 (2010) Article 10.7.2.
- [6] D. M. Bressoud. *Proofs and Confirmations: The story of the alternating sign matrix conjecture*. Cambridge University Press Cambridge, England, 1999.
- [7] R. A. Brualdi and S. Kirkland. Aztec diamonds and digraphs, and Hankel determinants of schröder numbers. *J. Combin. Theory, Series B*, 94(2) (2005) 334–351.
- [8] J. Cigler. Some nice Hankel determinants. Arxiv preprint arXiv:1109.1449, 2011.
- [9] Hankel determinants of linear combinations of moments of orthogonal polynomials. Arxiv preprint arXiv:2003.01676, 2020.
- [10] M. Elouafi. A unified approach for the Hankel determinants of classical combinatorial numbers, *J. Math. Anal. Appl.* 431 (2015) 1253–1274.
- [11] I. Gessel and G. Viennot. Binomial determinants, paths, and hook length formulae. *Adv. in Math.*, 58(3) (1985) 300–321.
- [12] I. M. Gessel and G. Xin. The generating function of ternary trees and continued fractions. *Electronic J. Combin.* 13 (2008) # R53.
- [13] G.-N. Han. Hankel continued fraction and its applications. *Adv. in Math.*, 303 (2016) 295–321.
- [14] Q.-H. Hou, A. Lascoux, Y.-P. Mu. Evaluation of some Hankel determinants, *Adv. Appl. Math.* 34 (2005) 845–852.
- [15] W. B. Jones and W. J. Thron. *Continued Fractions: Analytic Theory and Applications*, *Encyclopedia of Mathematics and its Applications*. vol. 11, Addison-Wesley Publishing Co., Reading, Mass., 1980.
- [16] C. Krattenthaler. Advanced determinant calculus. *Séminaire Lotharingien Combin.*, 42 (1999) B42q.
- [17] C. Krattenthaler. Advanced determinant calculus: a complement. *Linear Algebra Appl.*, 411:68–166, 2005.
- [18] C. Krattenthaler. Determinants of (generalised) Catalan numbers, *J. Statist. Plann. Inference* 140 (2010) 2260–2270.
- [19] M. E. Mays and J. Wojciechowski. A determinant property of Catalan numbers. *Discrete Math.*, 211(1-3) (2000) 125–134.
- [20] L. Mu, Y. Wang and Y. Y. Nan. Hankel determinants of linear combinations of consecutive Catalan-like numbers. *Discrete Math.* 340 (2017), no. 12, 3097–3103.
- [21] L. Mu and Y. Wang. Hankel determinants of shifted Catalan-like numbers. *Discrete Math.* 340 (2017), no. 6, 1389–1396.
- [22] R. Sulanke and G. Xin. Hankel determinants for some common lattice paths. *Adv. Appl. Math.* 40 (2008) 1410–167.
- [23] U. Tamm. Some aspects of Hankel matrices in coding theory and combinatorics, *Electron. J. Combin.* 8 (1) (2001) 31.

- [24] H. S. Wall. Analytic Theory of Continued Fractions. Van Nostrand, New York, 1948.
- [25] Y. Wang, G. Xin and M. Zhai. Hankel determinants and shifted periodic continued fractions, Adv. Appl. Math. 102 (2019) 83–112.
- [26] Y. Wang and G. Xin. Hankel determinants for convolution powers of Catalan numbers. Discrete Math. , 2019,342(9):2694-2716.

¹SCHOOL OF MATHEMATICS AND STATISTICS, NORTH CHINA UNIVERSITY OF WATER RESOURCES AND ELECTRIC POWER, ZHENGZHOU 450045, PR CHINA

²SCHOOL OF STATISTICS AND MATHEMATICS, YUNNAN UNIVERSITY OF FINANCE AND ECONOMICS, KUNMING 650221, PR CHINA

Email address: ¹wangying2019@ncwu.edu.cn & ²zyrzuhe@126.com