

# BUBBLE SORT AND HOWE DUALITY FOR STAIRCASE MATRICES

ANTON KHOROSHKIN AND IEVGEN MAKEDONSKYI

ABSTRACT. In this paper, we present an independent proof of the Cauchy identities for staircase matrices, originally discovered in [FKhM3], using the combinatorics of the Bruhat poset and the bubble-sort procedure. Additionally, we derive new insights into certain coefficients appearing in one of these identities.

The first part of the paper focuses on combinatorial aspects. It is self-contained, of independent interest, and introduces a generalization of parabolic Bruhat graphs for monotone functions on an arborescent poset. The second part examines the intersections of Demazure modules within a given integrable representation. Finally, we propose a generalization of the classical Howe duality for staircase matrices in terms of the corresponding distributive lattice of Demazure submodules. Computing the associated character yields the desired Cauchy identities for staircase matrices.

## CONTENTS

0. Introduction	1
1. Recollection of (Bruhat) Posets and Distributive Lattices	6
2. Bubble-Sort Algorithm for Arborescent Posets with (Anti)Linearization	13
3. Distributive lattice of Demazure submodules	24
4. Staircase shapes, corners and $DL$ -dense arrays	29
5. Howe Duality for Staircase Matrices	38
References	42
Appendix A. Pictorial examples	44

## 0. INTRODUCTION

We focus in the introduction by explaining our main goal – “*Howe duality for staircase matrices*” – rather than detailing the combinatorial structures we discovered along the way. However, the key combinatorial ingredients must be introduced early on.

Let us fix the space of rectangular matrices  $\mathbf{Mat}_{n \times m}$  with  $n$  rows and  $m$  columns. The Lie algebras of matrices  $\mathfrak{gl}_n$  and  $\mathfrak{gl}_m$  act on the space  $\mathbf{Mat}_{n \times m}$  from the left and right, respectively. Classical “*Howe duality*” (discovered by R.Howe [Ho]) describes the decomposition of the symmetric algebra into a direct sum of irreducible  $\mathfrak{gl}_n$ - $\mathfrak{gl}_m$ -modules:

$$(A) \quad S^N(\mathbf{Mat}_{n \times m}) \simeq \bigoplus_{\lambda \vdash N} V_\lambda^{\mathfrak{gl}_n} \otimes (V_\lambda^{\mathfrak{gl}_m})^{\text{op}}, \text{ with } l(\lambda) \leq \min(m, n).$$

Here,  $V_\lambda$  is the irreducible left  $\mathfrak{gl}_n$ -module with highest weight  $\lambda$ , while  $(V_\lambda^{\mathfrak{gl}_m})^{\text{op}}$  denotes the opposite irreducible right  $\mathfrak{gl}_m$ -module with highest weight  $\lambda$ . Next, let us consider the



**Theorem D.** For each partition  $\lambda \vdash N$  with  $l(\lambda) \leq \#\mathfrak{S}_{\bar{n}}$ , the following isomorphism of  $\mathfrak{b}_n$ - $\mathfrak{b}_m$ -(bi)-submodules holds:

$$(E) \quad \text{gr}\mathcal{F}^\lambda(S^N(\text{Mat}_{\bar{n}})) \simeq \sum_{A \in \text{DL}_{\bar{n}}(\lambda)} D_{\text{hor}(A)} \otimes D_{\text{vrt}(A)}^{\text{op}} \subset V_\lambda^{\mathfrak{gl}_n} \otimes (V_\lambda^{\mathfrak{gl}_m})^{\text{op}}.$$

Here  $D_{\sigma\lambda}$  denotes the "Demazure  $\mathfrak{b}$ -submodule" of the  $\mathfrak{gl}$ -irreducible representation  $V_\lambda$ , generated by the extremal vector of weight  $\sigma\lambda \in \mathbb{S}_n\lambda$  (see [Dem1, Dem2]). Notations  $\text{hor}(A)$ ,  $\text{vrt}(A)$  stands for the horizontal and the vertical weights of the DL-dense array  $A$ . We suggest below pictorial illustration of  $\text{vrt}$  and  $\text{hor}$  and refer to the formal Definition 4.12 in the main body of the text:

$$\mathbb{Z}_{\geq 0}^5 \ni (1, 2, 3, 1, 4) = \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline 3 \\ \hline 1 \\ \hline 4 \\ \hline \end{array} \xleftarrow{\text{hor}} \begin{array}{|c|c|c|c|} \hline & & 1 & \\ \hline 2 & & & \\ \hline & 3 & & \\ \hline & & & 1 \\ \hline & & & 4 \\ \hline \end{array} \xrightarrow{\text{vrt}} \boxed{2\ 3\ 1\ 4\ 1} = (2, 3, 1, 0, 4, 1) \in \mathbb{Z}_{\geq 0}^6.$$

Second, to extract any character formula from the description (E), we need to analyze the intersection of Demazure modules. The following theorem appears to be known among specialists; however, we were unable to find it explicitly stated in the literature<sup>1</sup>:

**Theorem F.** (Theorem 3.10) The subspaces  $\{D_{w\lambda} : \omega \in W\}$  of the integrable representation  $V_\lambda$  form a distributive lattice, denoted by  $\mathcal{L}_D(V_\lambda)$ .

In other words, there exists a "common" basis of  $V_\lambda$  such that its intersection with any Demazure submodule forms a basis for that submodule. Consequently, any submodule obtained through an iterative process of sums and intersections of Demazure submodules is itself a sum of Demazure submodules. Furthermore, the character of any such  $\mathfrak{b}$ -submodule of  $V_\lambda$  is given by the sum of the characters of its minimal subquotients:

$$K_{w\lambda} := D_{w\lambda} / \sum_{\tau\lambda \prec w\lambda} D_{\tau\lambda},$$

which are known as "van der Kallen modules", following [vdK].

Note that the sets of  $\mathfrak{gl}_n$  and  $\mathfrak{gl}_m$  weights appearing in the summation range of the right-hand side of (E) are very special. Moreover, in Corollary 4.26, we show that the induced Bruhat partial orders  $\prec_{\text{Br}}$  on the horizontal and vertical weights of DL-dense arrays are equivalent:

$$(G) \quad \forall A, B \in \text{DL}_{\bar{n}}(\lambda), \quad (\text{hor}(A) \prec_{\text{Br}} \text{hor}(B)) \in \mathbb{S}_n\lambda \Leftrightarrow (\text{vrt}(A) \prec_{\text{Br}} \text{vrt}(B)) \in \mathbb{S}_m\lambda.$$

The most combinatorial part of this paper explores the Bruhat partial order on the set  $\text{DL}_{\bar{n}}(\lambda)$ , which is essential for understanding equivalence (G). In particular, we show that the poset  $\mathfrak{S}_{\bar{n}}$  is "arborescent" and that the maps  $\text{vrt}$  and  $\text{hor}$  define a "consistent (anti)linearization" (see Definition 2.1). The subset  $\text{vrt}(\text{DL}_{\bar{n}}(\lambda)) \subset \mathbb{S}_m\lambda$  consists of compositions that define a nondecreasing function on the poset  $\mathfrak{S}_{\bar{n}}$ . We refer to such compositions as "  $\mathfrak{S}_{\bar{n}}$ -dominant" (Definition 2.5) and establish the following properties of the poset  $\text{DL}_{\bar{n}}(\lambda)$ , which hold for any arborescent poset with consistent (anti)linearization:

<sup>1</sup>It is worth mentioning that, to the best of our knowledge, the statement of Theorem F does not follow from Crystal Basis theory, despite the fact that Demazure characters can be realized using crystals, as suggested by Kashiwara [Ka].

**Theorem H.**

- The set  $\text{DL}_{\bar{\pi}}(\lambda)$  is empty if  $l(\lambda) > \#\mathfrak{S}_{\bar{\pi}}$ .
- (Theorem 2.35) If  $\lambda = (\lambda_1 > \dots > \lambda_{\#\mathfrak{S}_{\bar{\pi}}})$  is a regular partition of length  $\#\mathfrak{S}_{\bar{\pi}}$ , then the poset  $\text{DL}_{\bar{\pi}}$  is independent of  $\lambda$  and is shown to be bounded, graded, and EL-shellable.
- If  $l(\lambda) \leq \#\mathfrak{S}_{\bar{\pi}}$ , then:
  - (Theorem 2.18) The covering relations for the Bruhat partial order on the set  $\text{DL}_{\bar{\pi}}(\lambda)$  of  $\mathfrak{S}_{\bar{\pi}}$ -dominant compositions are determined by transpositions in minimal disorders (see Definitions 2.13 and 4.20).
  - (§2.3, Corollary 2.31) A natural generalization of the bubble-sort algorithm for the arborescent poset  $\mathfrak{S}_{\bar{\pi}}$  defines a nonincreasing monotone idempotent on the corresponding interval in the Bruhat graph  $\mathbb{S}_m\lambda$ :

$$\text{bs}_{\bar{\pi}} : [\lambda_+^{\bar{\pi}}, \lambda_-]^{S_m\lambda} \rightarrow \text{DL}_{\bar{\pi}}(\lambda).$$

- There exists a monotone idempotent  $\pi_\lambda : \text{DL}_{\bar{\pi}} \rightarrow \text{DL}_{\bar{\pi}}(\lambda)$  that accounts for the difference between regular and non-regular  $\lambda$ .

It is worth noting that Theorem H can be viewed as a generalization of a well-known special case. Indeed, when  $n = m$  and  $\text{Mat}_{\bar{\pi}}$  is the parabolic Lie subalgebra of  $\mathfrak{gl}_n$  with parabolic Weyl subgroup  $\mathbb{S}_{\bar{\pi}}$ , the poset of staircase corners  $\mathfrak{S}_{\bar{\pi}}$  is a disjoint union of linearly ordered sets. In this case, all statements of Theorem H are well known for regular  $\lambda$ . In particular,  $\text{DL}_{\bar{\pi}}(\lambda)$  corresponds to the parabolic Bruhat graph, and the bubble-sort process maps a permutation  $w$  to the permutation of minimal length in the coset  $w\mathbb{S}_{\bar{\pi}} \subset \mathbb{S}_n$ . The monotonicity of the bubble-sort map  $\text{bs}_{\bar{\pi}}$  plays a crucial role in the following result:

**Corollary I.** (Corollary 3.13) *The following sets consist of all  $\vee$ -indecomposable elements in the distributive sublattice of vector subspaces they generate:*

- $\{D_{\text{hor}(A)} : A \in \text{DL}_{\bar{\pi}}(\lambda)\}$ , a sublattice of  $\mathcal{L}_D(V_\lambda)$ .
- $\{D_{\text{vrt}(A)}^{\text{op}} : A \in \text{DL}_{\bar{\pi}}(\lambda)\} \subset \mathcal{L}_{D^{\text{op}}}(V_\lambda^{\text{op}})$ .
- $\{D_{\text{hor}(A)} \otimes D_{\text{vrt}(B)}^{\text{op}} : A, B \in \text{DL}_{\bar{\pi}}(\lambda)\} \subset \mathcal{L}_{D \times D^{\text{op}}}(V_\lambda \otimes V_\lambda^{\text{op}})$ .

Here,  $\mathcal{L}_D(V_\lambda)$  (resp.  $\mathcal{L}_{D^{\text{op}}}(V_\lambda^{\text{op}})$ ) denotes the distributive lattice of left (resp. right) Demazure submodules as described in Theorem F. The corresponding minimal subquotients in these lattices are the generalized van der Kallen modules:

$$\mathbf{K}_{\mathfrak{S}_{\bar{\pi}}, \text{hor}(A)} := D_{\text{hor}(A)} / \sum_{C \prec_{\text{Br}} A} D_{\text{hor}(C)}, \quad \mathbf{K}_{\mathfrak{S}_{\bar{\pi}}, \text{vrt}(B)}^{\text{op}} := D_{\text{vrt}(B)}^{\text{op}} / \sum_{C \succ_{\text{Br}} B} D_{\text{hor}(C)}^{\text{op}}.$$

Finally, the desired "Cauchy identities for staircase matrices" (L) and (M) (first discovered in [FKhM3]) follows from a straightforward computation of the character of the isomorphism (E). The right-hand side of this isomorphism belongs to the aforementioned distributive lattice  $\{D_{\text{hor}(A)} \otimes D_{\text{vrt}(B)}^{\text{op}}\}$  and, consequently, is expressed as a sum of characters of its minimal subquotients:

$$\mathbf{K}_{A,B}^{\bar{\pi}} := \mathbf{K}_{\mathfrak{S}_{\bar{\pi}}, \text{hor}(A)} \otimes \mathbf{K}_{\mathfrak{S}_{\bar{\pi}}, \text{vrt}(B)}^{\text{op}}.$$

**Corollary J.** (Corollary 3.16 and Corollary 5.16) *The following character identities hold for any staircase matrix shape:*

$$\begin{aligned}
\text{(K)} \quad \text{ch}_{\mathfrak{b}_m}(\mathbf{K}_{\mathfrak{S}_{\bar{\pi}}, \text{vrt}(A)}^{\text{op}}) &= \sum_{\bar{d}: \text{bs}_{\bar{\pi}}(\bar{d})=\text{vrt}(A)} a^{\bar{d}}(y) = \sum_{B \succ_{\text{Br}A} A} \mu^{\text{DL}_{\bar{\pi}}(\lambda)}(A, B) \kappa^{\text{vrt}(B)}(y); \\
\text{(L)} \quad \prod_{(i,j) \in \mathbf{Y}_{\bar{\pi}}} \frac{1}{1 - x_i y_j} &= \sum_N \sum_{\substack{\lambda \vdash N \\ l(\lambda) \leq \#\mathfrak{S}_{\bar{\pi}}}} \sum_{A \in \text{DL}_{\bar{\pi}}(\lambda)} \kappa_{\text{hor}(A)}(x) \cdot \left( \sum_{\bar{d}: \text{bs}_{\bar{\pi}}(\bar{d})=\text{vrt}(A)} a^{\bar{d}}(y) \right), \\
\text{(M)} \quad \prod_{(i,j) \in \mathbf{Y}_{\bar{\pi}}} \frac{1}{1 - x_i y_j} &= \sum_N \sum_{\substack{\lambda \vdash N \\ l(\lambda) \leq \#\mathfrak{S}_{\bar{\pi}}}} \sum_{\substack{A \succ B \\ A, B \in \text{DL}_{\bar{\pi}}(\lambda)}} \mu^{\text{DL}_{\bar{\pi}}(\lambda)}(A, B) \kappa_{\text{hor}(A)}(x) \kappa^{\text{vrt}(B)}(y).
\end{aligned}$$

Here,  $\kappa_{\bar{d}}(x)$  and  $\kappa^{\bar{d}}(y)$  are the left and right "key polynomials", which compute the characters of left and right Demazure modules. Similarly,  $a_{\bar{d}}(x)$  and  $a^{\bar{d}}(y)$  are "Demazure atoms", which correspond to the characters of left and right van der Kallen modules (see, e.g., [Al, AGL, Mas] for combinatorial definitions of these polynomials). The numbers  $\mu^{\text{DL}_{\bar{\pi}}(\lambda)}(A, B)$  represent the "Möbius function" of the poset  $\text{DL}_{\bar{\pi}}(\lambda)$  of "DL-dense arrays". This function has detailed description for regular  $\lambda$  (Corollary 2.36), and it is conjectured to take values in  $\{0, \pm 1\}$  for all partitions  $\lambda$  (Conjectures 2.43 and 2.44).

It is worth mentioning that our original proof of the generalized Cauchy identities (L) and (M), presented in [FKhM3], was based on the Pieri rules for key polynomials discovered in [AQ1, AQ2]. In this paper, we provide an independent proof of these identities. Moreover, our results naturally lead to a rediscovery of the generalized Pieri rules, much like how the classical Pieri rules follow from the classical Cauchy identity.

The classical Cauchy identity can be interpreted as a character equality arising from Howe duality and has numerous applications and generalizations across various fields of mathematics, including combinatorics, representation theory, probability theory, and mathematical physics (see, e.g., [BC, BP, BW, Ok, OR, St1, CK, FL, Las]). Several attempts have been made to generalize the Cauchy identity for staircase matrices for combinatorial and probabilistic purposes ([Las, AE, AGL]). However, we believe that our result exhibits a particularly elegant internal structure, and the methods we propose may be applicable to other known instances of Howe duality.

**0.1. Structure of the paper.** Sections §1, §2, and §4 focus entirely on combinatorial aspects, while Sections §3 and §5 are dedicated to the representation-theoretic side of the paper.

In §1, we recall key definitions from the theory of posets (§1.1) and distributive lattices (§1.3) and review the combinatorial structure of the Bruhat graph in §1.2.

In §2, we introduce a generalization of the bubble-sort algorithm for arborescent posets:

- In §2.1, we define the main subposet of the Bruhat graph consisting of dominant compositions.
- In §2.2, we describe its Hasse diagram.
- In §2.3, we present the bubble-sort algorithm and analyze its key properties.
- The desired properties of the poset of dominant compositions are proved in §2.4.1 for regular  $\lambda$  and are conjectured in §2.4.2 for  $\lambda$  with repeating elements.

Section 3 focuses on the structure theory of Demazure modules. We recall their definition in §3.1 and describe their intersections in §3.2.

Starting from §4, we begin working with staircase shapes and develop their combinatorics:

- The poset  $\mathfrak{Sc}_{\overline{n}}$  of staircase corners is defined in §4.2.
- $DL$ -dense arrays are introduced in §4.3.
- The poset structure of these arrays is described in §4.4, relying on the combinatorial results obtained in §2.

The final section, §5, builds upon all the combinatorial techniques developed so far to establish the "*Howe duality for staircase matrices*".

Appendix A contains pictorial examples of the Hasse graph of the set  $DL_{\overline{n}}(\lambda)$ .

**Acknowledgements.** We are grateful to Michel Brion for useful correspondence that leads to the main idea of Theorem F and Evgeny Feigin, Sergei Ivanov, Igor Makhlin and Alek Vainstein for useful conversations. The work of Ie. M. is partially supported by Beijing NSF grant.

## 1. RECOLLECTION OF (BRUHAT) POSETS AND DISTRIBUTIVE LATTICES

1.1. **Recollection of poset terminology.** Let us recall some properties of partially ordered sets that are important for our purposes. We refer to [W] and references therein for the introduction to the theory of posets. All posets in this paper are finite and, in particular, all intervals

$$(x, y) := \{z : x \prec z \prec y\}$$

are finite.

**Definition 1.1.** A poset  $(X, \preceq)$  is called

- bounded if there exists an element  $\hat{1}$  greater or equal than every other element in  $X$ , and an element  $\hat{0}$  less or equal to every other element in  $X$ .
- graded (also called ranked or pure) if all chains between two comparable elements have the same length.

The length of the maximal chain from  $\hat{0}$  to  $x$  in a bounded graded poset is called the rank function and is denoted by  $\text{rk}(x)$ .

The Hasse graph or Hasse diagram  $\Gamma_X$  of the poset  $X$  is a directed graph whose vertices are elements of  $X$  and edges represent the covering relations in the poset  $X$ . For the pictorial description we orient the edges from the smaller element to the bigger one.

**Definition 1.2.** A graded poset  $(X, \preceq)$  is called

- thin if all open intervals of length 2 consist of two elements:

$$\forall x \prec y \ \& \ \text{rk}(y) - \text{rk}(x) = 2 \ \Rightarrow \ (x, y) := x \prec z_1, z_2 \prec y.$$

- subthin if all open intervals of length 2 consist of at most two elements.

**Definition 1.3.** An  $A$ -labelling of the poset  $(X, \prec)$  is the map  $\mathcal{E}$  from the set of edges of the Hasse diagram  $\Gamma_X$  to a given set  $A$ .

The  $A$ -labelling is called EL-labelling (*Edge-Lexicographical*) if  $A$  is a poset and for any pair of comparable elements  $x < y \in X$  there exists a unique maximal chain

$$\text{path}_{\mathcal{E}}(x \prec y) := (x \prec z_1 \prec \dots \prec z_k \prec y)$$

going from  $x$  to  $y$  which has increasing  $\mathcal{E}$  labels (when reading the covering relations from minima to maxima)

$$\mathcal{E}(x \prec z_1) < \mathcal{E}(z_1 \prec z_2) < \dots < \mathcal{E}(z_k \prec y)$$

and, moreover, this unique maximal chain is minimal for the lexicographic order on maximal chains (comparing the words given by the successive  $\mathcal{E}$  labels).

**Definition 1.4.** A poset  $X$  is called EL-shellable if there exists an EL-labelling of its Hasse diagram.

We refer to [W] and to [BB, BW1] for the details of aforementioned notions and applications to Coxeter groups.

**Definition 1.5.** A map  $F : X \rightarrow X$  on the poset  $(X, \leq)$  is called

- decreasing if  $F(x) \leq x$  (respectively increasing if  $F(x) \geq x$ );
- monotone if  $\forall x \leq y$  we have  $F(x) \leq F(y)$ ;
- idempotent if  $\forall x \in X$   $F(x) = F(F(x))$ .

**Lemma 1.6.** Suppose  $F : X \rightarrow X$  is a monotone idempotent. Then

- if  $F$  is decreasing then  $\forall x \in X$  the element  $F(x)$  is the supremum element among the elements  $y \in F(X)$  that are less or equal to  $x$ :

$$\forall x \in X \quad F(x) = \sup\{y \in F(X) : y \leq x\}.$$

- Similarly, if  $F$  is an increasing monotone idempotent then

$$\forall x \in X \quad F(x) = \inf\{y \in F(X) : y \geq x\}.$$

*Proof.* Suppose that  $F$  is decreasing and  $F(x') \leq x$ . Then since  $F$  is an order-preserving map we have:

$$F(x') = F(F(x')) \leq F(x).$$

□

In particular, we see that if  $x < x'$  and  $x < x''$  is a pair of different edges in the Hasse diagram of  $F(X)$  then the open intervals  $(x, x')$  and  $(x, x'')$  of the poset  $X$  do not intersect.

**Definition 1.7.** Let  $X$  be a locally finite poset. The Möbius function  $\mu^X : X \times X \rightarrow \mathbb{Z}$  is defined recursively as follows:

- For all  $x \in X$ ,  $\mu(x, x) = 1$ .
- For all  $x, y \in X$  such that  $x < y$ , the Möbius function is given by:

$$\mu(x, y) = - \sum_{z \in [x, y)} \mu(x, z),$$

where the sum is taken over all  $z \in X$  such that  $x \leq z < y$ .

- If  $x \not\leq y$ , then  $\mu(x, y) = 0$ .

The main application of the Möbius function is the following Möbius Inversion Formula

**Fact 1.8.** Let  $f, g : X \rightarrow \mathbb{C}$  be two functions on a locally finite poset  $X$  related by the following summation:

$$\forall x \in X \quad g(x) = \sum_{y \leq x} f(y).$$

Then,  $f$  can be expressed in terms of  $g$  using the Möbius function  $\mu^X$  of the poset  $X$ :

$$\forall y \in X \quad f(y) = \sum_{x \leq y} \mu^X(x, y)g(x).$$

**Definition 1.9.** The order complex  $\Delta(X)$  of a poset  $X$  is the simplicial complex whose vertices are indexed by elements of  $X$  and simplices  $\{x_0, \dots, x_k\}$  are in one-to-one correspondence with chains  $x_0 < \dots < x_k$  in the poset  $X$ .

**Fact 1.10.** The Möbius function  $\mu^X(x, y)$  is equal to the reduced Euler characteristic of the geometric realization of the order complex of the open interval  $(x, y)$ :

$$\mu^X(x, y) = \tilde{\chi}(|\Delta(x, y)|).$$

The following facts follow from the seminal Quillen's Theorem A ([Q], see also [Wl, Th.2.3] and [G, Corollary 10.12 on p.1853]):

**Fact 1.11.** If the poset  $X$  has a minimal element then  $\Delta(X)$  is contractible.

**Corollary 1.12.** Let  $F : X \rightarrow X$  be an order-preserving idempotent then

- $\forall y \in F(X)$  the simplicial complex  $\Delta(F^{-1}(y))$  is contractible;
- The order complexes  $\Delta(X)$  and of  $\Delta(F(X))$  are homotopy equivalent.

The subsequent fact was stated by Danaraj-Klee for simplicial complexes ([DK]) and later on was generalized to posets by Bjorner ([B1]). It has numerous applications for the parabolic BGG resolution:

**Fact 1.13.** If  $(X, \leq)$  is a shellable subthin graded poset then the geometric realization of an ordered complex  $\Delta(X)$  is homeomorphic to

- a sphere, if the poset  $X$  is thin;
- a ball, if the poset  $X$  is subthin but not thin.

**Corollary 1.14.** If  $X$  is a graded bounded subthin shellable poset then  $\forall x < y \in X$  we have

$$\mu^X(x, y) = \begin{cases} (-1)^{\text{rk}(y) - \text{rk}(x)}, & \text{if the interval } [x, y] \text{ is thin,} \\ 0, & \text{if the interval } [x, y] \text{ is subthin, but not thin.} \end{cases}$$

**1.2. Bruhat Order and Bruhat Graph.** Let  $W$  be the Weyl group associated with a semisimple Lie algebra  $\mathfrak{g}$  and the root system  $\Phi$ . We assume that the Cartan decomposition  $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$  is fixed, and that the set of simple reflections  $S \subset W$  corresponds to this decomposition. The group  $W$  is naturally equipped with the length function, which counts the number of simple reflections in a reduced decomposition.

Moreover, for any weight  $\lambda \in P$  in the weight lattice, we define:

- the dominant weight  $\lambda_+ := W\lambda \cap P_+$  in the  $W$ -orbit of  $\lambda$ ;
- the Weyl subgroup  $W_J := \text{Stab}(\lambda_+) \subset W$ , generated by reflections that fix  $\lambda_+$ ;
- the element  $\sigma_\lambda \in W$  of minimal length such that  $\lambda = \sigma_\lambda \lambda_+$ , which represents the left coset  $W/W_J$ .

**Definition 1.15.** The Bruhat partial order on the  $W$ -orbit  $W\lambda$  is defined as follows. For  $\lambda = \sigma_\lambda \lambda_+$  and  $\mu = \sigma_\mu \lambda_+$ , we set

$$\lambda \preceq \mu \quad \text{if and only if} \quad \Phi_- \cap \sigma_\lambda(\Phi_+) \subset \Phi_- \cap \sigma_\mu(\Phi_+).$$

The Hasse diagram of the poset  $W\lambda \simeq W/W_J$  is called the parabolic Bruhat graph. If  $W_J$  is trivial, it is called the Bruhat graph.

For further details, we refer to the classical textbooks on this subject [Hu, BB].



**Fact 1.16.** *The (parabolic) Bruhat graph  $W\lambda$  is bounded and graded, with*

$$\hat{0} := \lambda_+, \quad \hat{1} := \lambda_- = \omega_0 \lambda_+,$$

where  $\omega_0$  (respectively,  $\omega_0^J$ ) is the longest element of the Weyl group  $W$  (resp. the Weyl subgroup  $W_J$ ). The rank function coincides with the length function:

$$\text{rk}(\lambda) = l(\sigma_\lambda).$$

Moreover, the Bruhat graph is thin for regular  $\lambda$  (i.e., when  $W\lambda \simeq W$ ), while the parabolic Bruhat graph is subthin. Additionally,  $W\lambda$  admits an EL-labelling.

For this paper, we focus on the case  $\mathfrak{g} = \mathfrak{gl}_n$ . The corresponding Weyl group  $W$  is the symmetric group  $\mathbb{S}_n$ , and the set of weights we consider consists of compositions  $\mathbb{Z}_{\geq 0}^n$ .

When  $W = \mathbb{S}_n$ , the "reflections" are "transpositions", and the set of "simple reflections" consists of "consecutive" transpositions. This allows us to express the length function and the edges of the Bruhat graph in terms of compositions. In particular, if  $\lambda_+ = (\lambda_1 > \dots > \lambda_m)$  is a "strictly decreasing partition" also called regular, then:

$$(1.17) \quad \text{For } \nu \in \mathbb{S}_m \lambda_+, \quad \text{rk}(\nu) = l(\nu) := \#\{i < j : \nu_i < \nu_j\};$$

$$(1.18)$$

$$\nu \xrightarrow{\prec} (ij)\nu \quad \Leftrightarrow \quad \begin{cases} \nu_i > \nu_j, \\ \forall k = i+1, \dots, j-1, \quad \nu_i \notin [\nu_j, \nu_i]. \end{cases}$$

is an edge in the Bruhat graph  $\mathbb{S}_m \mu$

A detailed combinatorial description of the (strong) Bruhat order for  $\mathbb{S}_m$  can be found in [BB, Th.2.1.5].

**Fact 1.19.** *The labelling of the edges of the Bruhat graph  $\mathbb{S}_m$  by transpositions defines an EL-labelling if we consider the following total order on transpositions  $\{(ij) : 1 \leq i < j \leq n\}$ :*

$$(1.20) \quad (il) \triangleleft (jk) \quad \stackrel{\text{def}}{\Leftrightarrow} \quad \begin{cases} l < k, \\ l = k \text{ and } i > j. \end{cases}$$

It is well-known that for any given parabolic subgroup  $W_J$  of a given Weyl group  $W$  each coset  $wW_J$  is an interval in the Bruhat graph  $W$  with the unique element of minimal  $w^+$  and maximal  $w^-$  length correspondingly. Moreover, the assignments  $w \mapsto w^+$  (resp.  $w \mapsto w^-$ ) are monotone nonincreasing (respectively nondecreasing) idempotents acting on  $W$  (see e.g. [BB, §2.5]). Let us adopt this observation to our case of a symmetric group.

From now on, we suppose that  $\lambda$  is a partition with repeating elements. Then for  $\delta := (m-1, m-2, \dots, 1, 0)$  the partition  $\lambda + \delta$  has no equal elements. Consequently,  $\mathbb{S}_m(\lambda + \delta) \simeq \mathbb{S}_m$ . Let  $\mathbb{S}_\lambda$  be the subgroup that stabilizes partition  $\lambda$ . To each permutation  $\sigma$  we assign  $\sigma_\lambda^+$  and  $\sigma_\lambda^-$  that are elements of minimal and maximal length among those permutations whose action on  $\lambda$  is equal to  $\sigma\lambda$ . Then the assignments:

$$\psi_\lambda^+ : \mathbb{S}_m \xrightarrow{\sigma \mapsto \sigma_\lambda^+} \mathbb{S}_m, \quad \psi_\lambda^- : \mathbb{S}_m \xrightarrow{\sigma \mapsto \sigma_\lambda^-} \mathbb{S}_m$$

are known to be monotone idempotents (nonincreasing and nondecreasing correspondingly). The images of these idempotents are isomorphic to the poset  $\mathbb{S}_m \lambda$  and we end up with the order-preserving projection that forgets elements from  $\delta$ :

$$(1.21) \quad \pi_\lambda : \mathbb{S}_m(\lambda + \delta) \twoheadrightarrow \mathbb{S}_m \lambda,$$

whose left and right adjoints are called by the same letters  $\psi_\lambda^+$  and  $\psi_\lambda^-$  correspondingly.

### 1.3. Distributive Lattice of Vector Subspaces.

**Definition 1.22.** A lattice is a partially ordered set  $(\mathcal{L}, \leq)$  in which every pair of elements  $a, b \in \mathcal{L}$  has a unique supremum  $a \vee b$  (also called the least upper bound or join) and a unique infimum  $a \wedge b$  (also called the greatest lower bound or meet). Both operations are monotone with respect to the given order:

$$(a_1 \leq a_2) \ \& \ (b_1 \leq b_2) \ \Rightarrow \ (a_1 \vee b_1 \leq a_2 \vee b_2) \ \& \ (a_1 \wedge b_1 \leq a_2 \wedge b_2).$$

A lattice  $\mathcal{L}$  is called distributive if it satisfies the following identity:

$$\forall x, y, z \in \mathcal{L} \quad x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z).$$

Throughout this paper, we assume that all lattices under consideration are finite.

**Example 1.23.** (Boolean lattice) The power set  $2^M$  of a given set  $M$  forms a distributive lattice with the following operations:

$$\begin{aligned} S \leq T &\Leftrightarrow S \subset T, \\ S \wedge T &:= S \cap T, \quad S \vee T := S \cup T. \end{aligned}$$

Birkhoff's theorem ([Bi], see also [DP, §5]) states that any finite distributive lattice is a sublattice of a Boolean lattice. The proof of this theorem is straightforward and relies on the following notion:

**Definition 1.24.** An element  $V$  of a distributive lattice  $\mathcal{L}$  is called join-irreducible (or  $\vee$ -irreducible) if, whenever  $V = V_1 \vee V_2$ , we must have either  $V = V_1$  or  $V = V_2$ .

The set of all  $\vee$ -irreducible elements of  $\mathcal{L}$  is denoted by  $\mathcal{J}(\mathcal{L})$ .

In particular, for any two  $\vee$ -irreducible elements  $V_1, V_2$  in a distributive lattice  $\mathcal{L}$ , the following holds:

$$V_1 \leq V_2 \Leftrightarrow V_1 \vee V_2 = V_2.$$

**Example 1.25.** To each finite poset  $(X, \leq)$ , we associate a sublattice of the Boolean lattice  $2^X$  consisting of upper sets:

$$\mathcal{L}_X := \{S \subset X : x \in S \Rightarrow \forall y \geq x \ y \in S\}.$$

The set of  $\vee$ -irreducible elements of  $\mathcal{L}_X$  is indexed by elements of  $X$  and consists of the upper sets  $X_{\geq x} := \{y \in X : y \geq x\}$ .

**Fact 1.26.** ([DP, §5]) The assignments  $(X, \leq) \mapsto \mathcal{L}_X$  and  $\mathcal{L} \mapsto \mathcal{J}(\mathcal{L})$  define a pair of mutually inverse bijections between the set of finite posets and the set of finite distributive lattices.

However, in our applications, we are primarily interested in distributive lattices of vector subspaces. The set  $\mathcal{L}(U)$  of all vector subspaces of a given vector space  $U$  forms a poset under inclusion and a lattice under the operations of intersection and sum:

$$\forall V_1, V_2 \subset U \quad \begin{cases} V_1 \leq V_2 \stackrel{\text{def}}{\Leftrightarrow} V_1 \subset V_2; \\ V_1 \wedge V_2 := V_1 \cap V_2, \quad V_1 \vee V_2 := V_1 + V_2. \end{cases}$$

**Definition 1.27.** For a given collection of vector subspaces  $V_1, \dots, V_k \subset U$ , we define the lattice  $\mathcal{L}(V_1, \dots, V_k; U)$  as the smallest sublattice of  $\mathcal{L}(U)$  that contains all  $V_i$ .

Note that while the lattice  $\mathcal{L}(U)$  may be infinite, its sublattice  $\mathcal{L}(V_1, \dots, V_k; U)$  is always finite. The following proposition is a well-known fact from linear algebra (see, e.g., [PP, Chapter 1, Proposition 7.1] and references therein):

**Proposition 1.28.** *The lattice  $\mathcal{L}(V_1, \dots, V_k; U) \subset \mathcal{L}(U)$  is distributive if and only if there exists a basis  $\{e_1, \dots, e_n\}$  of the ambient space  $U$  such that for each  $i = 1, \dots, k$ , the space  $V_i$  has a basis given by  $V_i \cap \{e_1, \dots, e_n\}$ .*

In particular, under the assumptions of Proposition 1.28, the mapping:

$$\psi : V_i \mapsto V_i \cap \{e_1, \dots, e_n\} \subset \{e_1, \dots, e_n\}$$

defines an embedding of the distributive lattice  $\mathcal{L}(V_1, \dots, V_k; U)$  into the Boolean lattice of subsets of the basis  $\{e_1, \dots, e_n\}$ , where the set  $\psi(V_i)$  forms a basis of the space  $V_i$ .

From now on, we consider a distributive lattice  $\mathcal{L}$  of vector subspaces of a given vector space  $U$ , generated by its subset of  $\vee$ -irreducible vector subspaces:

$$\mathcal{J}(\mathcal{L}) := \{V_1, \dots, V_k\}.$$

We assume that  $(0)$  and  $U$  belong to  $\mathcal{J}(\mathcal{L})$  and that the induced poset  $\mathcal{J}(\mathcal{L})$  is bounded:

$$i \leq j \stackrel{\text{def}}{\iff} V_i \subset V_j.$$

Any element in the distributive lattice is a join of the  $\vee$ -irreducible elements, what means that any subspace  $V \in \mathcal{L}$  satisfies:

$$V = \vee_{V_i \subset V} V_i = \sum_{V_i \subset V} V_i.$$

The set  $\mathcal{J}(\mathcal{L})$  defines a standard filtration applied to both  $U$  and any subspace  $V$  in  $\mathcal{L}$ :

$$(1.29) \quad \mathcal{F}_{\mathcal{L}}^i V := V \wedge V_i = V \cap V_i, \quad i \leq j \Rightarrow \mathcal{F}_{\mathcal{L}}^i V \subset \mathcal{F}_{\mathcal{L}}^j V.$$

Let  $E := \{e_1, \dots, e_{\dim V}\}$  be a given common basis of  $\mathcal{L}$ . To each  $\vee$ -irreducible space  $V_i$ , we associate a subset  $E^i := \{e_1^i, \dots, e_{r_i}^i\} \subset E$ , consisting of basis elements that belong to  $V_i$  but not to any other proper subspace  $V_j \subsetneq V_i$ . From this definition, we obtain:

**Fact 1.30.**      • *Each space is spanned by the union of basis elements:  $V_i = \text{span}\langle \cup_{V_j \subset V_i} E^j \rangle$ .*

- *The elements from  $E^i$  form a basis for the space*

$$(1.31) \quad \mathsf{K}_i := V_i \Big/ \sum_{V_j \subsetneq V_i} V_j$$

*called the minimal subquotient in  $\mathcal{L}$ .*

- *The subquotient*

$$\mathcal{F}_{\mathcal{L}}^i V / \mathcal{F}_{\mathcal{L}}^{<i} V \simeq \begin{cases} \mathsf{K}_i, & \text{if } V_i \subset V, \\ 0, & \text{otherwise.} \end{cases}$$

- *The basis of the quotient  $\bar{V} = V'/V''$  for  $V', V'' \in \mathcal{L}$  consists of the (disjoint) union of the sets  $E^i$  for indices  $i$  satisfying  $(V_i \subset V')$  &  $(V_i \not\subset V'')$ .*

For any subset  $V_{i_1}, \dots, V_{i_r} \subset \mathcal{L}_V$  of  $\vee$ -irreducible subspaces, we define:

$$\mathcal{J}^{\leq i_1, \dots, i_r} := \{V_j : (j \leq i_1) \ \& \ \dots \ \& \ (j \leq i_r)\} \subset \mathcal{J}.$$

Then we have the equality:

$$V_{i_1} \wedge \dots \wedge V_{i_r} = \vee_{V \in \mathcal{J}^{\leq i_1, \dots, i_r}} V = \vee_{V \in \max(\mathcal{J}^{\leq i_1, \dots, i_r})} V.$$

The latter follows from the fact that all  $V_i$ 's are  $\vee$ -irreducible.

**Proposition 1.32.** *Each subquotient  $\bar{V} := V'/V''$  of two subspaces  $V', V''$  in a distributive lattice  $\mathcal{L}$  admits a resolution by a direct sum of  $\vee$ -irreducible elements.*

*Proof.* We prove by induction on  $i$  that the minimal quotient  $K_i$  (defined in (1.31)) has a resolution by  $\vee$ -irreducible elements  $V_j$  with  $j \leq i$ .

The base case is clear: when  $i$  is minimal, we have  $K_i = V_i$ .

Suppose the statement holds for all  $j < i$ . Then, by gluing these resolutions, we obtain a resolution for any quotient  $\bar{U} := U_1/U_2$  with  $U_1 \subsetneq V_i$ . Let  $V_{j_1}, \dots, V_{j_r}$  be the maximal  $\vee$ -irreducible subspaces satisfying  $V_j \leq V_i$ . Then, we have the short exact sequence:

$$0 \longrightarrow V_{j_1} + \dots + V_{j_r} \longrightarrow V_i \longrightarrow K_i \longrightarrow 0.$$

Applying inclusion-exclusion, we obtain the long exact sequence:

$$0 \rightarrow \bigcap_{s=1}^r V_{j_s} \rightarrow \dots \rightarrow \bigoplus_{\substack{J \subset [1,r] \\ |J|=l}} \bigcap_{s \in J} V_{j_s} \rightarrow \dots \rightarrow \bigoplus_{s=1}^r V_{j_s} \rightarrow \bigoplus_{s=1}^r V_{j_s} \rightarrow 0.$$

By induction, each intersection is resolved by  $\vee$ -irreducible elements, hence so is  $K_i$ .  $\square$

**Proposition 1.33.** *Let  $F : \mathcal{J}(\mathcal{L}) \rightarrow \mathcal{J}(\mathcal{L})$  be an increasing monotone idempotent on the poset  $\mathcal{J}(\mathcal{L})$  of  $\vee$ -irreducible elements in a given distributive lattice  $\mathcal{L}$ . Then the image  $F(\mathcal{J}(\mathcal{L}))$  forms a set of  $\vee$ -irreducible elements in the distributive sublattice  $\mathcal{L}_F \subset \mathcal{L}$  generated by  $F(\mathcal{J}(\mathcal{L}))$ .*

*Moreover, for every  $l \in F(\mathcal{J}(\mathcal{L}))$ , the minimal subquotients in the sublattice  $\mathcal{L}_F$  are given by:*

$$(1.34) \quad K_l^F := V_{F(i)} \bigg/ \sum_{V_{F(j)} \subset V_{F(i)}} V_{F(j)} \simeq \bigoplus_{l: F(l)=F(i)} V_l \bigg/ \sum_{V_j \subset V_l} V_j.$$

*Proof.* The proof follows from a direct application of Birkhoff's bijection between posets and distributive lattices, as discussed in Example 1.25.

First, note that any sublattice of a distributive lattice remains distributive. Second, any  $\vee$ -irreducible element of the original lattice that belongs to the sublattice remains  $\vee$ -irreducible within it. Finally, using Lemma 1.6, we conclude that there are no additional  $\vee$ -irreducible elements in  $\mathcal{L}_F$  beyond those in the set  $\{V_{F(i)} : V_i \in \mathcal{J}(\mathcal{L})\}$ . Isomorphism (1.34) follows from a direct examination of the elements of the basis  $E$ . Specifically, the union  $\bigcup_{l \in F^{-1}(F(i))} E^l$  constitutes a basis for both the left-hand and right-hand sides of (1.34), establishing the claim.  $\square$

## 2. BUBBLE-SORT ALGORITHM FOR ARBORESCENT POSETS WITH (ANTI)LINEARIZATION

This section is purely combinatorial and independent of the rest of the paper. We generalize the "bubble-sort" algorithm for arborescent posets  $\mathbf{S}$  with (anti)linearization and describe the Bruhat partial order and its properties on the set of monotone functions over these posets.

Later in the paper, these posets will appear as the set of staircase corners  $\mathbf{Sc}_{\overline{\pi}}$  (Proposition 4.11). The Bruhat poset of monotone functions described in this section will resurface in §4, playing a key role in defining the vertical and horizontal weights of  $DL$ -dense arrays (Definition 4.13), which form the main combinatorial foundation for the generalization of Howe duality for staircase matrices.

### 2.1. Arborescent Posets with Consistent (Anti)Linearization.

**Definition 2.1.** • A poset  $(\mathbf{S}, \succ)$ , together with an order-reversing monomorphism  $v : (\mathbf{S}, \succ) \hookrightarrow ([1m], <)$  that satisfies one of the two following equivalent conditions:

$$(2.2) \quad \begin{aligned} \forall s, t, r \in \mathbf{S}: [(v(s) < v(t) < v(r)) \& (s \succ r)] &\Rightarrow (t \succ r); \\ \Leftrightarrow \forall s \in \mathbf{S} \exists a_s \leq b_s \leq m : \mathbf{S}_{\succeq s} = v^{-1}([a_s, b_s]). \end{aligned}$$

is called an arborescent poset with consistent anti-linearization.

- A poset  $(\mathbf{S}, \prec)$ , together with an order-preserving embedding  $h : (\mathbf{S}, \prec) \hookrightarrow ([1n], <)$  that satisfies one of the two following equivalent conditions:

$$\begin{aligned} \forall s, t, r \in \mathbf{S}: [(h(s) < h(t) < h(r)) \& (s \prec r)] &\Rightarrow (t \prec r); \\ \Leftrightarrow \forall s \in \mathbf{S} \exists a_s \leq b_s \leq n : \mathbf{S}_{\succeq s} = h^{-1}([a_s, b_s]). \end{aligned}$$

is called an arborescent poset with consistent linearization.

Assumption (2.2) ensures that the Hasse diagram of  $(\mathbf{S}, \prec)$  forms a forest, where the set of minimal elements corresponds to the roots. Indeed, if this were not the case, there would exist three elements  $s, t, r \in \mathbf{S}$  such that  $s \succ t$ ,  $s \succ r$ , with  $t$  and  $r$  being incomparable, and  $v(s) < v(t) < v(r)$ . Consequently, the subset  $\mathbf{S}_{\succeq r}$  would contain  $s$  and, therefore, the preimage of the interval  $[v(s), v(r)]$ . However,  $t \notin \mathbf{S}_{\succeq r}$ , leading to a contradiction.

**Definition 2.3.** Given an arborescent poset  $\mathbf{S}$  with consistent linearization or anti-linearization  $v : \mathbf{S} \hookrightarrow [1m]$ , a composition  $\overline{d} := (d_1, \dots, d_m) \in \mathbb{Z}_{\geq 0}^m$  is called  $\mathbf{S}$ -dominant if the following conditions hold:

$$\begin{cases} l \notin v(\mathbf{S}) \Rightarrow d_l = 0; \\ s \succ t \in \mathbf{S} \Rightarrow d_{v(s)} \geq d_{v(t)}. \end{cases}$$

**Remark 2.4.** For an arborescent poset  $\mathbf{S}$ , the order-reversing embedding  $v : \mathbf{S} \hookrightarrow [1m]$  is a consistent anti-linearization if and only if its opposite order-preserving embedding  $\text{op} \circ v : \mathbf{S} \hookrightarrow [1m]$  is a consistent linearization. Here,

$$\text{op}(l) := m - l + 1$$

is the standard order-reversing involution of the interval  $[1m]$ .

Accordingly, a composition  $\overline{d} := (d_1, \dots, d_m) \in \mathbb{Z}_{\geq 0}^m$  is  $\mathbf{S}$ -dominant for the anti-linearization  $v$  if and only if the opposite composition  $\overline{d}^{\text{op}} := (d_m, \dots, d_1)$  is dominant for the linearization  $\text{op} \circ v$ .

From this point onward, we fix an arborescent poset  $\mathbf{S}$  with a consistent **anti**-linearization  $v : \mathbf{S} \hookrightarrow [1m]$ . Similar definitions and results hold for a consistent **linearization** by considering the opposite order on the linearly ordered set  $[1m]$ .

**Definition 2.5.** A composition  $\bar{d} := (d_1, \dots, d_m) \in \mathbb{Z}_{\geq 0}^m$  is called:

- $\mathbf{S}$ -admissible if the following inequality holds for all  $k \leq m$ :

$$\#\{j = 1, \dots, k : d_j > 0\} \leq \#\mathbf{S}_{\leq k},$$

where  $\mathbf{S}_{\leq k} := v^{-1}([1, k])$ .

- $\mathbf{S}[l]$ -dominant if the following implications hold:

$$\begin{cases} l \notin v(\mathbf{S}) \Rightarrow d_l = 0; \\ l \in v(\mathbf{S}) \Rightarrow \forall s \in \mathbf{S} : s \succ v^{-1}(l) \text{ we have } d_{v(s)} \geq d_l; \end{cases}$$

- $\mathbf{S}[\leq k]$ -dominant if  $\bar{d}$  is  $\mathbf{S}[l]$ -dominant for all  $l \leq k$ .

In particular, a composition  $\bar{d}$  is  $\mathbf{S}$ -dominant if and only if it is  $\mathbf{S}[\leq m]$ -dominant.

**Example 2.6.** Below is a pictorial example of an arborescent poset with consistent anti-linearization. The elements of the poset  $\mathbf{S}$  are represented by blue dots, edges in the Hasse diagram of  $\mathbf{S}$  are drawn as edges, and each cell corresponds to an element of the linearly ordered set  $[1m]$ :

$$\begin{array}{|c|c|c|c|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & & \bullet & \bullet & \bullet & \bullet & & \\ \hline \end{array} := \left\{ \begin{array}{l} \mathbf{S} := \{s_1, s_2, s_3, s_5, s_6, s_7, s_8\}, \\ s_1 \succ s_3, s_2 \succ s_3, s_5 \succ s_8, s_6 \succ s_7 \succ s_8, \\ v : \mathbf{S} \hookrightarrow [19], \text{ with } v(s_i) = i. \end{array} \right\}$$

We hope that the following pictorial illustration of various compositions for the aforementioned arborescent poset  $\mathbf{S}$  will help in understanding Definition 2.5:

$$\begin{array}{l} \begin{array}{|c|c|c|c|c|c|c|c|c|c|} \hline 3 & 4 & 1 & 0 & 4 & 6 & 3 & 2 & 0 & \\ \hline \end{array} \quad - \quad \text{is } \mathbf{S}\text{-dominant} \Leftrightarrow \mathbf{S}[\leq 9]\text{-dominant,} \\ \begin{array}{|c|c|c|c|c|c|c|c|c|c|} \hline 3 & 4 & 1 & 0 & 4 & 2 & 3 & 0 & 6 & \\ \hline \end{array} \quad - \quad \text{is } \mathbf{S}[\leq 6]\text{-dominant and } \mathbf{S}\text{-admissible,} \\ \begin{array}{|c|c|c|c|c|c|c|c|c|c|} \hline 3 & 4 & 1 & 8 & 4 & 6 & 3 & 2 & 0 & \\ \hline \end{array} \quad - \quad \text{is } \mathbf{S}[\leq 3]\text{-dominant, but not } \mathbf{S}\text{-admissible.} \end{array}$$

**Notation 2.7.** For each partition  $\lambda$ , we define the following subsets of  $\mathbb{S}_m \lambda$ , consisting of compositions whose multiset of nonzero elements equals  $\lambda$ :

$\mathbb{A}_{\mathbf{S}}^v(\lambda)$  – the set of  $\mathbf{S}$ -admissible compositions;

$\mathbb{D}_{\mathbf{S}}^v(\lambda)$  – the set of  $\mathbf{S}$ -dominant compositions;

$\mathbb{B}_{\mathbf{S}[\leq k]}^v(\lambda)$  – the set of  $\mathbf{S}[\leq k]$ -dominant and  $\mathbf{S}$ -admissible compositions.

We equip these sets with the poset structure induced from the Bruhat partial order on  $\mathbb{S}_m \lambda$ .

Note that every  $\mathbf{S}$ -dominant composition is also  $\mathbf{S}$ -admissible, leading to the following sequence of poset embeddings:

$$\mathbb{D}_{\mathbf{S}}^v(\lambda) = \mathbb{B}_{\mathbf{S}[\leq m]}^v(\lambda) \subset \mathbb{B}_{\mathbf{S}[\leq m-1]}^v(\lambda) \subset \dots \subset \mathbb{B}_{\mathbf{S}[\leq 0]}^v(\lambda) = \mathbb{A}_{\mathbf{S}}^v(\lambda).$$

For each  $k = 1, \dots, m$ , we denote by  $\tilde{\mathbf{S}}_k$  the poset  $\mathbf{S}_{\leq k} \sqcup \{k+1, \dots, m\}$ , where the partial order is inherited from  $\mathbf{S}_{\leq k}$ , while the remaining elements are incomparable with all

others. The poset  $\tilde{\mathcal{S}}_k$  is evidently arborescent, and the natural map  $\tilde{v} : \tilde{\mathcal{S}}_k \rightarrow [1m]$ , which coincides with  $v$  on  $\mathcal{S}_{\leq k}$  and acts as the identity on the complement, defines a consistent anti-linearization. From the definitions, we immediately obtain the following isomorphism of sets:

$$(2.8) \quad \mathbb{B}_{\mathcal{S}_{[\leq k]}}^v(\lambda) = \mathbb{D}_{\tilde{\mathcal{S}}_k}^{\tilde{v}}(\lambda) \cap \mathbb{A}_{\mathcal{S}}^v(\lambda).$$

The main goal of this section is to demonstrate that the posets  $\mathbb{D}_{\mathcal{S}}^v(\lambda)$  inherit many desirable properties of the Bruhat graph  $\mathbb{S}_m\lambda$ . One of the key features is the generalization of the classical bubble-sort algorithm, which defines a monotone idempotent map  $\mathbf{bs}_{\mathcal{S}} : \mathbb{A}_{\mathcal{S}}^v(\lambda) \rightarrow \mathbb{D}_{\mathcal{S}}^v(\lambda)$ . This map is constructed as a composition of monotone idempotents  $\mathbf{bs}_{\mathcal{S}}^k : \mathbb{B}_{\mathcal{S}_{[\leq k]}}^v(\lambda) \rightarrow \mathbb{B}_{\mathcal{S}_{[\leq k]}}^v(\lambda)$ . We have successfully proven that for a regular partition  $\lambda$ , the posets  $\mathbb{D}_{\mathcal{S}}^v(\lambda)$  are bounded, graded, subthin, and EL-shellable. In contrast, for  $\lambda$  with repeated parts, the poset may not be graded; however, we conjecture that it remains shellable. Let us now discuss the differences between the regular and non-regular cases.

The posets  $\mathbb{D}_{\mathcal{S}}^v(\lambda)$  and  $\mathbb{D}_{\mathcal{S}}^v(\lambda')$  are isomorphic whenever there is an isomorphism between the ordered multisets  $\lambda$  and  $\lambda'$ . In particular, if  $\lambda$  is "regular" (i.e., all parts of  $\lambda$  are distinct), then the set  $\mathbb{D}_{\mathcal{S}}^v(\lambda)$  of  $\mathcal{S}$ -dominant compositions does not depend on  $\lambda$  and will be denoted simply as  $\mathbb{D}_{\mathcal{S}}^v$ .

**Lemma 2.9.** *Suppose that  $\mathcal{S}$  is an arborescent poset with a consistent anti-linearization  $v : \mathcal{S} \hookrightarrow [1m]$ . Then the monotone projection  $\pi_{\lambda} : \mathbb{S}_m \twoheadrightarrow \mathbb{S}_m\lambda$  and its adjoint embedding  $\psi_{\lambda}^+ : \mathbb{S}_m\lambda \hookrightarrow \mathbb{S}_m$  (defined in (1.21)) restrict to a monotone projection and its adjoint monotone embedding on the set  $\mathbb{D}_{\mathcal{S}}^v$  of  $\mathcal{S}$ -dominant weights:*

$$(2.10) \quad \mathbb{D}_{\mathcal{S}}^v = \mathbb{D}_{\mathcal{S}}^v(\lambda + \delta) \begin{array}{c} \xleftarrow{\psi_{\lambda}^+} \\ \perp \\ \xrightarrow{\pi_{\lambda}} \end{array} \mathbb{D}_{\mathcal{S}}^v(\lambda).$$

*Proof.* It follows directly from Definition 2.5 that if  $\bar{a} \in \mathbb{S}_m(\lambda + \delta)$  is  $\mathcal{S}$ -dominant, then the corresponding composition  $\pi_{\lambda}(\bar{a})$  is also  $\mathcal{S}$ -dominant. Conversely, if  $\bar{b} \in \mathbb{S}_m\lambda$  is  $\mathcal{S}$ -dominant, then  $\psi_{\lambda}^+(\bar{b})$  is also  $\mathcal{S}$ -dominant.  $\square$

Consequently, the composition  $\psi_{\lambda}^+ \circ \pi_{\lambda}$ , also denoted by  $\psi_{\lambda}^+$ , is a non-increasing monotone idempotent on  $\mathbb{D}_{\mathcal{S}}$ .

**Example 2.11.** Suppose that  $\mathbb{T} = \mathbb{T}_1 \sqcup \dots \sqcup \mathbb{T}_k$  is a disjoint union of linearly ordered sets  $\mathbb{T}_q = (s_{m_{q-1}+1} \succ \dots \succ s_{m_q})$  and that  $v : \mathbb{T} \hookrightarrow [1m]$  is a consistent surjective anti-linearization given by:

$$v(s_i) = i.$$

Then, for each composition  $\bar{d} \in \mathbb{Z}_{\geq 0}^m$ , we define a composition  $\mathbf{bs}_{\mathbb{T}}(\bar{d})$  by sorting the elements within each connected component of  $\mathcal{S}$ . More precisely,  $\mathbf{bs}_{\mathbb{T}}(\bar{d})$  is determined by the following conditions, satisfied for all  $q = 1, \dots, k$ :

- The following multisets are equal:

$$\{d_{m_{q-1}+1}, \dots, d_{m_q}\} = \{\mathbf{bs}_{\mathbb{T}}(\bar{d})_{m_{q-1}+1}, \dots, \mathbf{bs}_{\mathbb{T}}(\bar{d})_{m_q}\}.$$

- The values in  $\mathbf{bs}_{\mathbb{T}}(\bar{d})$  are arranged in non-increasing order:

$$(2.12) \quad \mathbf{bs}_{\mathbb{T}}(\bar{d})_{m_{q-1}+1} \geq \dots \geq \mathbf{bs}_{\mathbb{T}}(\bar{d})_{m_q}.$$

By construction, we have  $\mathbf{bs}_\mathbb{T}(\bar{d}) = \mathbf{bs}_\mathbb{T}(\mathbf{bs}_\mathbb{T}(\bar{d}))$ , and for any  $\bar{a}, \bar{b} \in \mathbb{S}_m(\lambda)$  satisfying  $\bar{a} \prec_{\text{Br}} \bar{b}$ , we obtain:  $\mathbf{bs}_\mathbb{T}(\bar{a}) \preceq \mathbf{bs}_\mathbb{T}(\bar{b})$ . This confirms that  $\mathbf{bs}_\mathbb{T}$  is a monotone idempotent in the sense of Definition 1.5 (see, e.g., [BB, §2.5]).

Now, suppose that  $\lambda = (\lambda_1^{k_1}, \dots, \lambda_l^{k_l})$  with  $\lambda_1 > \dots > \lambda_l \geq 0$ . Then the corresponding set of  $\mathbb{T}$ -dominant compositions has the following structure:

$$\mathbb{D}_\mathbb{T}(\lambda) \simeq \mathbb{S}_{\bar{m}} \backslash \mathbb{S}_m \lambda \simeq \mathbb{S}_{\bar{m}} \backslash \mathbb{S}_m / \mathbb{S}_{\bar{\lambda}},$$

where

$$\mathbb{S}_{\bar{m}} := \mathbb{S}_{m_1} \times \dots \times \mathbb{S}_{m_r - m_{r-1}}, \quad \mathbb{S}_{\bar{\lambda}} := \mathbb{S}_{k_1} \times \dots \times \mathbb{S}_{k_l} = \text{Stabilizer}(\lambda).$$

In other words,  $\mathbb{D}_\mathbb{T}(\lambda)$  can be identified with a double coset of a Weyl group modulo two parabolic Weyl subgroups:

$$\begin{array}{ccc} \mathbb{S}_m = \mathbb{S}_m(\lambda + \delta) & \begin{array}{c} \xleftarrow{\psi_\lambda^+} \\ \xrightarrow[\perp]{\pi_\lambda} \end{array} & \mathbb{S}_m \lambda = \mathbb{S}_m / \mathbb{S}_{\bar{\lambda}} \\ \downarrow \mathbf{bs}_\mathbb{T} & & \downarrow \mathbf{bs}_\mathbb{T} \\ \mathbb{D}_\mathbb{T}^v = \mathbb{S}_{\bar{m}} \backslash \mathbb{S}_m & \begin{array}{c} \xleftarrow{\psi_\lambda^+} \\ \xrightarrow[\perp]{\pi_\lambda} \end{array} & \mathbb{D}_\mathbb{T}^v(\lambda) = \mathbb{S}_{\bar{m}} \backslash \mathbb{S}_m / \mathbb{S}_{\bar{\lambda}} \end{array}$$

Note that, as a poset, the double quotient  $\mathbb{S}_{\bar{m}} \backslash \mathbb{S}_m / \mathbb{S}_{\bar{\lambda}}$  may be non-graded, as illustrated in Example 2.42. However, all covering relations in the Hasse diagram of the induced Bruhat graph are given by appropriate transpositions. We refer to [Stm, Ko] for a discussion on double quotients in the context of Coxeter groups.

**2.2. Hasse diagram of  $\mathbf{S}$ -dominant compositions.** In this section, we assume that  $(\mathbf{S}, v)$  is an arborescent poset with **anti-linearization** and focus exclusively on  $\mathbf{S}$ -dominant compositions. From the definition, it follows that whenever  $j \notin v(\mathbf{S})$ , we have  $d_j = 0$  for any  $\mathbf{S}$ -dominant composition  $\bar{d}$ . Thus, the  $j$ 'th entry of any composition in  $\mathbb{D}_\mathbf{S}^v(\lambda)$  does not influence any conclusions in this section, as it is always equal to 0. Therefore, without loss of generality, we assume that  $v : \mathbf{S} \hookrightarrow [1m]$  is surjective, making it an isomorphism.

**Definition 2.13.** A pair  $(ij)$  of indices is called a minimal  $\mathbf{S}$ -disorder for an  $\mathbf{S}$ -dominant composition  $\bar{d} := (d_1, \dots, d_m)$  if the following conditions hold:

- $v^{-1}(i)$  and  $v^{-1}(j)$  are incomparable in  $\mathbf{S}$ .
- $(ij)$  forms a disorder, meaning that  $i < j$  and  $d_i > d_j$ .
- For all  $l$  in the interval  $(i, j) \subset [1m]$ , the following implications hold:

$$(2.14) \quad \left\{ \begin{array}{ll} v^{-1}(l) \prec v^{-1}(i) & \Rightarrow d_l \leq d_j, \\ v^{-1}(l) \succ v^{-1}(j) & \Rightarrow d_l \geq d_i, \\ v^{-1}(l) \text{ is incomparable with } v^{-1}(i) \text{ and } v^{-1}(j) & \Rightarrow d_l \notin [d_j, d_i]. \end{array} \right.$$

**Lemma 2.15.** If  $(ij)$  is a minimal  $\mathbf{S}$ -disorder for an  $\mathbf{S}$ -dominant composition  $\bar{d}$ , then the composition:

$$(ij) \cdot \bar{d} := (\dots, d_{i-1}, \mathbf{d}_j, d_{i+1}, \dots, d_{j-1}, \mathbf{d}_i, d_{j+1}, \dots)$$

is also  $\mathbf{S}$ -dominant, and  $\bar{d} \prec_{\text{Br}} (ij) \cdot \bar{d}$ .

*Proof.* Since each element  $d_l$  for  $l = i + 1, \dots, j - 1$  does not belong to the open interval  $(d_j, d_i)$ , it follows that  $\bar{d}$  is less than  $(ij) \cdot \bar{d}$  in the Bruhat partial order. The first two implications in (2.14) ensure that the  $\mathbf{S}$ -dominant inequalities remain valid for  $(ij) \cdot \bar{d}$ .  $\square$



It is worth mentioning that under the assumptions of Lemma 2.15, if all elements of the composition  $\bar{d}$  are pairwise distinct, then  $\bar{d} \prec (ij) \cdot \bar{d}$  is a covering relation in the Bruhat graph  $\mathbb{S}_m \lambda$ . Indeed, by the minimality condition, there is no  $l \in \{i+1, \dots, j-1\}$  such that  $d_i > d_l > d_j$ .

**Lemma 2.16.** *Suppose that  $\lambda$  is regular and  $\bar{a}, \bar{b} \in \mathbb{D}_S^v(\lambda)$  is a pair of Bruhat-comparable  $S$ -dominant compositions ( $\bar{a} \prec_{\text{Br}} \bar{b}$ ). Then there exists a minimal  $S$ -disorder  $(ij)$  for  $\bar{a}$  such that  $\bar{a} \prec_{\text{Br}} (ij)\bar{a} \preceq_{\text{Br}} \bar{b}$ .*

*Proof.* First, observe that if  $S$  is a disjoint union of linearly ordered sets, then the statement follows from Example 2.11.

We proceed by induction on the number  $m$  of elements in  $S$ . Recall that the Hasse diagram of  $S$  is a forest of rooted trees, where smaller elements are closer to the roots. If  $S$  consists of a single tree, then  $v^{-1}(m)$  is the unique minimal element of  $S$ . Thus,  $a_m = b_m = \lambda_m$ , where  $\lambda_m$  is the smallest part of  $\lambda$ . Then we can remove the root of  $S$  and apply induction on  $m$  with the truncated anti-linearization  $S_{< m} \rightarrow [1m-1]$  and the corresponding  $S_{< m}$ -dominant compositions  $(a_1, \dots, a_{m-1})$  and  $(b_1, \dots, b_{m-1})$ .

If the Hasse diagram of  $S$  contains multiple connected components, we decompose  $S$  according to  $v$ :

$$(2.17) \quad S = T_1 \sqcup T_2 \sqcup \dots \sqcup T_k, \quad |T_q| = m_q.$$

For  $q < q'$  and  $s \in T_q, s' \in T_{q'}$ , we have  $v(s) < v(s')$ , and we define a linear order on each tree  $T_q$  following  $v$ :

$$t \prec t' \stackrel{\text{def}}{\iff} \begin{cases} \exists q: t, t' \in T_q, \\ v(t) > v(t'). \end{cases}$$

The poset  $T = \sqcup_{q=1}^k T_q$  is a disjoint union of linearly ordered sets  $T_q$  and thus satisfies Example 2.11. Therefore, we have a monotone idempotent  $\text{bs}_T$  that maps  $\bar{d}$  to a  $T$ -dominant composition  $\text{bs}_T(\bar{d})$ , as described in (2.12).

If  $\text{bs}_T(\bar{a}) = \text{bs}_T(\bar{b})$ , then the multisets of values of  $\bar{a}$  and  $\bar{b}$  restricted to each  $T_q$  coincide, allowing us to proceed separately on each  $T_q$ .

Otherwise, there exists a transposition  $(ll')$  in the parabolic Bruhat graph such that:

$$\text{bs}_T(\bar{a}) \prec (ll')\text{bs}_T(\bar{a}) \preceq \text{bs}_T(\bar{b}).$$

Since all elements of  $\lambda$  are distinct, there exists a unique pair  $(i, j)$  such that:

$$a_i = (\text{bs}_T(\bar{a}))_i > a_j = (\text{bs}_T(\bar{a}))_j.$$

Note that  $(ll')$  is a minimal disorder for  $T$ -dominant composition  $\text{bs}_T(\bar{a})$  what follows that

$$\forall p = m_{q-1} + 1, \dots, m_{q'} \text{ we have } \text{bs}_T(a)_p \notin [(\text{bs}_T(\bar{a}))_{l'}, (\text{bs}_T(\bar{a}))_l] \iff a_p \notin [a_j, a_i].$$

Consequently  $(ij)$  is a minimal  $S$ -disorder for  $\bar{a}$  and, moreover,  $(ij)\bar{a} \preceq \bar{b}$ .  $\square$

**Theorem 2.18.** *The restriction of the Bruhat partial order to the subset  $\mathbb{D}_S^v(\lambda) \subset \mathbb{S}_m \lambda$  of  $S$ -dominant compositions is generated by the relations  $\bar{a} \prec (ij)\bar{a}$ , where  $(ij)$  is a minimal  $S$ -disorder for  $\bar{a}$ .*

*Proof.* First, suppose that  $\lambda$  is regular. Lemma 2.15 implies that  $\bar{a}$  is less than  $(ij)\bar{a}$  in the Bruhat partial order. Lemma 2.16 establishes that the comparison  $\bar{a} \prec (ij)\bar{a}$  defines an edge in the Hasse diagram. Indeed, assume that  $\bar{a} \prec_{\text{Br}} \bar{b} \in \mathbb{D}_{\mathbb{S}}^y(\lambda)$ . Then there exists a chain of  $\mathbb{S}$ -dominant compositions:

$$\bar{a} = \bar{c}_0 \prec \bar{c}_1 \prec \bar{c}_2 \prec \dots \preceq \bar{b}$$

such that each  $\bar{c}_i$  differs from  $\bar{c}_{i+1}$  by a transposition associated with a minimal  $\mathbb{S}$ -disorder and that  $\bar{c}_k \preceq \bar{b}$ . Since the Bruhat graph is finite, this chain cannot be infinite. Consequently, there exists an index  $l$  such that  $\bar{c}_l = \bar{b}$ .

If  $\lambda$  has repeating elements, we use the monotone projection  $\pi_\lambda$  and its adjoint  $\psi_\lambda^+$  from (2.10). Specifically, consider an edge  $\bar{a} \prec \bar{b}$  in the Hasse diagram of  $\mathbb{D}_{\mathbb{S}}^y(\lambda)$ . Then  $\psi_\lambda^+(\bar{a}) \prec \psi_\lambda^+(\bar{b})$  in  $\mathbb{D}_{\mathbb{S}}$ , and the latter must be connected by a chain of edges in the Hasse diagram of  $\mathbb{D}_{\mathbb{S}}$ :

$$\psi_\lambda^+(\bar{a}) = \bar{c}_0 \prec \dots \prec \bar{c}_{k-1} \prec \bar{c}_k \prec \dots \prec \bar{c}_l = \psi_\lambda^+(\bar{b}),$$

such that for all  $i \neq k$ , we have  $\pi_\lambda(\bar{c}_{i-1}) = \pi_\lambda(\bar{c}_i)$ . Consequently,  $\bar{a}$  differs from  $\bar{b}$  by a transposition in a disorder, and the corresponding disorder must be minimal since it arises from a minimal disorder in  $\mathbb{D}_{\mathbb{S}}$ .  $\square$

**2.3. Bubble-sort.** In this section, we continue working with an arborescent poset  $\mathbb{S}$  equipped with a consistent order-reversing anti-linearization  $v : \mathbb{S} \rightarrow [1m]$ . The **bubble-sort algorithm** described below is an inductive procedure for reordering the elements of an  $\mathbb{S}$ -admissible composition  $\bar{d}$  into an  $\mathbb{S}$ -dominant composition  $\text{bs}_{\mathbb{S}}(\bar{d})$ :

$$(2.19) \quad \text{bs}_{\mathbb{S}} := \text{bs}_{\mathbb{S}}^m \circ \dots \circ \text{bs}_{\mathbb{S}}^1.$$

In each step  $k$  (denoted by  $\text{bs}_{\mathbb{S}}^k$ ), only the first  $k$  elements of the composition are rearranged. Thus, we explain the procedure for compositions of length  $k$ . The  $k$ 'th step  $\text{bs}_{\mathbb{S}}^k$  follows this recursive procedure to ensure  $\mathbb{S}[k]$ -dominance:

**Algorithmic Definition 2.20.** *Suppose that the composition  $\bar{d} = (d_1, \dots, d_k)$  is  $\mathbb{S}[< k]$ -dominant and  $\mathbb{S}$ -admissible but not  $\mathbb{S}[k]$ -dominant. Then:*

- Find the index  $j$  as the maximal element of  $[1m]$  with respect to the linear order  $\leq$ , among the set of maximal elements of  $\mathbb{S}$  with respect to the partial order  $\prec$ , whose weight is less than  $d_k$ . That is,


$$(2.21) \quad j := \max_{\leq} \left\{ v \left( \max_{\preceq} \{ s \in \mathbb{S}_{<k} : d_{v(s)} < d_k \} \right) \right\}.$$

- Repeat this procedure for the composition obtained by swapping  $d_j$  and  $d_k$  in  $\bar{d}$ :

$$(jk) \cdot \bar{d} := (d_1, \dots, d_{j-1}, \mathbf{d}_k, d_{j+1}, \dots, d_{k-1}, \mathbf{d}_j)$$

until the composition becomes  $\mathbb{S}[k]$ -dominant.

**Example 2.22.** Let us show an example of bubble-sort process  $\text{bs}_{\mathbb{S}}$  applied to an  $\mathbb{S}$ -admissible

composition  for the arborescent poset  $\mathbb{S}$  with antilinearisation from Example 2.6. As suggested in 2.20 we apply  $\text{bs}_{\mathbb{S}}^k$  on the truncated compositions. We



**Proposition 2.26.** *The Hasse diagram of  $\mathbb{D}_S^v$  is a subgraph of the Bruhat graph  $\mathbb{S}_m$ . In particular,  $\mathbb{D}_S^v$  is graded, and the rank function in  $\mathbb{D}_S^v$  is given by:*

$$\text{rk}_{\mathbb{D}_S^v}(\bar{d}) := \#\{s, t \in S : (s \neq t) \ \& \ (v(s) < v(t)) \ \& \ (d_{v(s)} < d_{v(t)})\}.$$

*Proof.* If all  $\lambda_i$ 's are distinct, then the non-strict inequalities in (2.14) become strict. Consequently, a minimal  $S$ -disorder  $(ij)$  for  $\bar{a}$  defines an edge  $\bar{a} \rightarrow (ij)\bar{a}$  in the Bruhat graph  $\mathbb{S}_m$ . Thus, the rank function counts the number of (dis)orders among incomparable elements in  $S$ .  $\square$

**Corollary 2.27.** *The poset  $\mathbb{D}_S^v$  is subthin.*

*Proof.* Consider two compositions  $\bar{a}, \bar{b} \in \mathbb{D}_S^v(\lambda)$  such that the rank difference between  $\bar{a}$  and  $\bar{b}$  is exactly 2. Since the Hasse diagram of  $\mathbb{D}_S^v$  is a subgraph of  $\mathbb{S}_m$ , we know that the open interval  $(\bar{a}, \bar{b})^{\mathbb{S}_m}$  in  $\mathbb{S}_m$  consists of exactly two elements (since  $\mathbb{S}_m$  is thin). By assumption, at least one of these elements must belong to  $\mathbb{D}_S^v$ , which implies that  $\mathbb{D}_S^v$  is subthin.  $\square$

The following proposition generalizes a well-known property of symmetric groups to the case of  $S$ -dominant compositions. Note that this result holds only for regular  $\lambda$ , as demonstrated in Example 2.42.

**Proposition 2.28.** *Suppose that two incomparable  $S$ -dominant compositions,  $\bar{b} = (il)\bar{a}$  and  $\bar{c} = (jk)\bar{a}$ , are obtained from an  $S$ -dominant composition  $\bar{a}$  by applying transpositions in two different minimal  $S$ -disorders. Then, there exists an  $S$ -dominant composition  $\bar{d}$  that can be obtained from both  $\bar{b}$  and  $\bar{c}$  by applying transpositions corresponding to appropriate minimal disorders:*

$$(2.29) \quad \begin{array}{ccc} & \bar{a} & \\ (il) \swarrow & & \searrow (jk) \\ \bar{b} & & \bar{c} \\ \exists(pq) \searrow & & \swarrow \exists(iq) \\ & \exists \bar{d} & \end{array} \quad , \quad \begin{array}{ccc} & \exists \bar{d} & \\ \exists(pq) \swarrow & & \searrow \exists(iq) \\ \bar{b} & & \bar{c} \\ (il) \searrow & & \swarrow (jk) \\ & \bar{a} & \end{array} .$$

Here, the arrows indicate an increase in the Bruhat order within the set  $\mathbb{D}_S^v(\lambda)$ . Moreover, we assume that either  $(l < k)$  or  $(l = k)$  and  $(i < j)$ , and that the sets  $(\{i < l\} \cup \{j < k\})$  and  $(\{p < k\} \cup \{i < q\})$  are identical.

Conversely, if the composition  $\bar{a}$  is obtained from  $\bar{b}$  via the minimal disorder  $(il)$  and from  $\bar{c}$  via the minimal disorder  $(jk)$ , then there exists a composition  $\bar{d}$  such that both  $\bar{b}$  and  $\bar{c}$  can be obtained from  $\bar{d}$  by applying minimal disorders.

*Proof.* The values of  $\bar{a}$ ,  $\bar{b}$ , and  $\bar{c}$  coincide for all indices other than  $i, j, k, l$ . Therefore, it suffices to prove the statement of the lemma for a poset of cardinality 3 or 4, depending on the size of the set  $\{i, j, k, l\}$ .

If the cardinality of  $\{i, j, k, l\}$  is 4, then the transpositions  $(il)$  and  $(jk)$  commute. Moreover, since all elements of  $\lambda$  are distinct, the transposition  $(il)$  remains a minimal disorder for  $(jk)\bar{a}$ , and vice versa. On the other hand, there are exactly five different arborescent posets with an anti-linearization of cardinality 3 (see Example A.1), and in all cases, the set  $\mathbb{D}_S^v$  is isomorphic to the (parabolic) Bruhat graph. The only case where two distinct minimal disorders exist corresponds to a poset in which all elements are incomparable.  $\square$

**Proposition 2.30.** *The bubble-sort operation  $\text{bs}_{\mathbb{S}}^k : \mathbb{B}_{\mathbb{S}[\leq k]}^v \rightarrow \mathbb{B}_{\mathbb{S}[\leq k]}^v$  is a nonincreasing, monotone idempotent (in the sense of Definition 1.5).*

*Proof.* By Corollary 2.24, we already know that  $\text{bs}_{\mathbb{S}}^k$  is a nonincreasing idempotent. Now, we prove the remaining monotonicity property by induction on the Bruhat partial order.

According to Theorem 2.18, the edges in the Hasse diagram of  $\mathbb{S}_{<k}$ -dominant compositions correspond to minimal disorders for the poset  $\tilde{\mathbb{S}}_{k-1}$ , as discussed in (2.8). Suppose that  $(il)\bar{d} \succeq_{\text{Br}} \bar{d}$  represents an edge in the Bruhat subgraph of  $\mathbb{S}$ -admissible,  $\mathbb{S}_{<k}$ -dominant compositions. We now analyze the effect of the bubble-sort algorithm on these compositions. If  $(il) = (jk)$ , then  $\text{bs}_{\mathbb{S}}^k((il)\bar{d}) = \text{bs}_{\mathbb{S}}^k(\bar{d})$ . On the other hand, Proposition 2.28 states that if  $(il)$  and  $(jk)$  are distinct, then the compositions obtained after one step of the bubble-sort algorithm remain distinct and differ by a transposition corresponding to a minimal  $\mathbb{S}_{<k}$ -disorder. Consequently, they remain connected by an edge in the Hasse diagram.  $\square$

**Corollary 2.31.** *The bubble-sort process  $\text{bs}_{\mathbb{S}} : \mathbb{S}_m \rightarrow \mathbb{D}_{\mathbb{S}}^v$  is a nonincreasing, monotone idempotent on  $\mathbb{S}_m$ .*

*Proof.* By Definition (2.19),  $\text{bs}_{\mathbb{S}}$  is constructed as a sequential composition of nonincreasing, monotone idempotents  $\text{bs}_{\mathbb{S}}^k$  for  $k = 1, \dots, m$ , where each  $\text{bs}_{\mathbb{S}}^k$  is applied to the image of the previous step  $\text{bs}_{\mathbb{S}}^{k-1}$ .  $\square$

**Corollary 2.32.** *The poset  $\mathbb{D}_{\mathbb{S}}^v$  is bounded.*

*Proof.* Since the Bruhat graph  $\mathbb{S}_m$  is bounded, its image under a monotone idempotent must also be bounded.  $\square$

Suppose we are given an EL-labelling  $\mathcal{E}_{\mathbb{S}_m}$  of the Bruhat graph  $\mathbb{S}_m$ , as described in (1.20). Then, for each pair of comparable elements  $\bar{a} \prec_{\text{Br}} \bar{b}$ , we assign the path:

$$\text{path}_{\mathbb{S}_m}(\bar{a} \rightarrow \bar{b}) := (\bar{a} \prec \bar{y}_1 \prec \dots \prec \bar{y}_{k-1} \prec \bar{b}),$$

such that

$$\mathcal{E}_{\mathbb{S}_m}(\bar{a} \prec \bar{y}_1) < \mathcal{E}_{\mathbb{S}_m}(\bar{y}_1 \prec \bar{y}_2) < \dots < \mathcal{E}_{\mathbb{S}_m}(\bar{y}_{k-1} \prec \bar{b}).$$

This path is assumed to be unique and lexicographically minimal due to the EL-property.

**Lemma 2.33.** *Suppose that  $\bar{a} \prec \bar{b} \prec \bar{c}$  is a sequence of three adjacent elements in the Hasse diagram of  $\mathbb{D}_{\mathbb{S}}^v(\lambda)$ . Then the maximal chain  $\text{path}_{\mathbb{S}_m}(\bar{a} \rightarrow \bar{c})$  in the Bruhat graph  $\mathbb{S}_m\lambda$  with increasing  $\mathcal{E}$ -labellings contains an  $\mathbb{S}$ -dominant composition  $\bar{d}$  such that  $\bar{a} \prec \bar{d} \prec \bar{c}$ . Moreover, if  $\bar{d} \neq \bar{b}$ , then the induced  $\mathcal{E}_{\mathbb{S}_m}$ -labeling along the alternative path is decreasing:*

$$\mathcal{E}_{\mathbb{S}_m}(\bar{a} \prec \bar{b}) > \mathcal{E}_{\mathbb{S}_m}(\bar{b} \prec \bar{c}).$$

*Proof.* Since the edges in the Hasse diagram of  $\mathbb{D}_{\mathbb{S}}^v$  correspond to transpositions, we can restrict the problem to the cases where  $\#\mathbb{S} \leq 4$  and analyze the combinatorics of the  $\mathcal{E}_{\mathbb{S}_m}$ -labeling in  $\mathbb{S}_3$  and  $\mathbb{S}_4$ . Note that assumption (2.2) plays a crucial role in this argument.  $\square$

**Proposition 2.34.** *The induced  $\mathcal{E}_{\mathbb{S}_m}$ -labeling of the Hasse diagram of  $\mathbb{D}_{\mathbb{S}}^v$  is an EL-labeling.*

*Proof.* Let  $\bar{a} \prec \bar{b}$  be a pair of comparable  $\mathbb{S}$ -dominant compositions. Suppose that the length of the maximal chain between  $\bar{a}$  and  $\bar{b}$  in  $\mathbb{D}_{\mathbb{S}}^v$  is  $k$ . By Proposition 2.26, the distance between  $\bar{a}$  and  $\bar{b}$  in  $\mathbb{S}_m$  is also  $k$ . We prove by induction on  $k$  that the path  $\text{path}_{\mathbb{S}_m}(\bar{a} \rightarrow \bar{b})$  with the minimal  $\mathcal{E}_{\mathbb{S}_m}$ -labeling consists of  $\mathbb{S}$ -dominant compositions.

The base case ( $k = 1$ ) is trivial. For the induction step, assume that the claim holds for chains of length less than  $k$ . By Lemma 2.33, starting with any maximal chain from  $\bar{a}$  to  $\bar{b}$  that passes through  $\mathbf{S}$ -dominant elements, if the labeling is not increasing, then there exists a subsequence of three adjacent elements in the chain where the labeling is decreasing. Applying Lemma 2.33, we can replace this subsequence with another maximal chain in  $\mathbb{D}_{\mathbf{S}}$  from  $\bar{a}$  to  $\bar{b}$  with a lexicographically smaller labeling. This completes the induction step.  $\square$

Let us summarize in one theorem the key properties obtained for the set of  $\mathbf{S}$ -dominant compositions in the case of regular  $\lambda$ .

**Theorem 2.35.** *The following properties hold for the poset  $(\mathbb{D}_{\mathbf{S}}^v, \prec_{\text{Br}})$  of  $\mathbf{S}$ -dominant compositions:*

- *The covering relations are given by transpositions associated with minimal disorders.*
- *The poset  $\mathbb{D}_{\mathbf{S}}^v$  is bounded, graded, subthin, and EL-shellable.*
- *The bubble-sort map  $\text{bs}_{\mathbf{S}} : \mathbb{S}_m \rightarrow \mathbb{D}_{\mathbf{S}}^v$  is a monotone, nonincreasing idempotent.*

*Proof.* The covering relations were described in Theorem 2.18. Corollary 2.32 ensures that  $\mathbb{D}_{\mathbf{S}}$  is bounded, while Corollary 2.27 establishes the subthin property. The EL-shellability is proved in Proposition 2.34. Finally, from Corollary 2.31, we know that  $\text{bs}_{\mathbf{S}}$  is a nonincreasing, monotone idempotent.  $\square$

**Corollary 2.36.** *If  $\lambda$  is regular, then the following identity holds for the Möbius function on the poset  $\mathbb{D}_{\mathbf{S}} = \mathbb{D}_{\mathbf{S}}(\lambda)$ :*

$$\mu^{\mathbb{D}_{\mathbf{S}}}(\bar{a}, \bar{b}) = \begin{cases} (-1)^{\text{rk}(\bar{b}) - \text{rk}(\bar{a})}, & \text{if the interval } (\bar{a}, \bar{b})^{\mathbb{S}_m} = (\bar{a}, \bar{b})^{\mathbb{D}_{\mathbf{S}}^v}, \\ 0, & \text{otherwise.} \end{cases}$$

*In other words, the Möbius function is equal to  $\pm 1$  if every intermediate composition  $\bar{c} \in \mathbb{S}_m \lambda$  satisfying  $\bar{a} \prec_{\text{Br}} \bar{c} \prec_{\text{Br}} \bar{b}$  is  $\mathbf{S}$ -dominant, and it is zero otherwise.*

*Proof.* The poset  $\mathbb{D}_{\mathbf{S}}(\lambda)$  is graded, EL-shellable, and subthin, which implies that each interval in this poset is either a ball or a sphere (Fact 1.13). As mentioned in Corollary 1.14, computing the Möbius function requires determining when an interval is not thin. This happens precisely when no elements are lost when restricting the interval  $(\bar{a}, \bar{b})^{\mathbb{S}_m \lambda} \subset \mathbb{S}_m \lambda$  to the subposet  $\mathbb{D}_{\mathbf{S}}^v(\lambda)$  of  $\mathbf{S}$ -dominant weights.  $\square$

**2.4.2. Parabolic case.** From now on we suppose that partition  $\lambda = (\lambda_1^{k_1}, \dots, \lambda_l^{k_l})$  may have repeating elements and the anti-linearization  $v : \mathbf{S} \hookrightarrow [1m]$  may be nonsurjective.

**Lemma 2.37.** *The set  $\mathbb{A}_{\mathbf{S}}(\lambda)$  of  $\mathbf{S}$ -admissible compositions is the interval  $[\lambda_+^{\mathbf{S}}, \lambda_-]$  in the Bruhat poset  $\mathbb{S}_m \lambda$  if  $l(\lambda) \leq \#\mathbf{S}$ . On the other hand, if  $l(\lambda) > \#\mathbf{S}$ , then  $\mathbb{A}_{\mathbf{S}}(\lambda) = \emptyset$ .*

*Here,  $\lambda_+^{\mathbf{S}}$  denotes the unique  $\mathbf{S}$ -dominant composition whose  $l$ 'th nonzero element is equal to  $\lambda_l$ , while the remaining elements are zero.*

*Proof.* First, notice that the total number of nonzero elements in an  $\mathbf{S}$ -admissible composition cannot exceed the size of  $\mathbf{S}$ . Thus, any  $\mathbf{S}$ -admissible composition belongs to  $\mathbb{S}_m \lambda$  with  $l(\lambda) \leq k$ . Second, while decreasing the Bruhat order of an  $\mathbf{S}$ -admissible composition  $\nu \in \mathbb{S}_m \lambda$ , one can attempt to shift all zero elements to the right while preserving  $\mathbf{S}$ -admissibility. What remains is determining the position of the zeros. Ultimately, this reduces to analyzing the Bruhat graph  $\mathbb{S}_k \lambda$  where  $k = \#\mathbf{S}$ . The latter has a unique minimal element,  $\lambda_+$ .  $\square$

**Proposition 2.38.** *For any partition  $\lambda$  with repeating elements, the idempotent  $\text{bs}_S^k$  commutes with the parabolic projection  $\pi_\lambda$  and its adjoint embedding  $\psi_\lambda^+$ :*

$$(2.39) \quad \begin{array}{ccc} \mathbb{B}_{S[<k]} = \mathbb{B}_{S[<k]}(\lambda + \delta) & \xleftarrow[\pi_\lambda]{\psi_\lambda^+} & \mathbb{B}_{S[<k]}(\lambda) \\ \downarrow \text{bs}_S^k & & \downarrow \text{bs}_S^k \\ \mathbb{B}_{S[\leq k]} = \mathbb{B}_{S[\leq k]}(\lambda + \delta) & \xleftarrow[\pi_\lambda]{\psi_\lambda^+} & \mathbb{B}_{S[\leq k]}(\lambda) \end{array}$$

*Proof.* The proof follows from a straightforward comparison of these two idempotents based on the following simple observation:

$$(\lambda_i \geq \lambda_j) \Rightarrow (\lambda + \delta)_i > (\lambda + \delta)_j.$$

This ensures that while performing bubble-sort in  $\mathbb{B}_{S[<k]}$ , we first reorder the elements of the composition  $\bar{d}$  that become equal in  $\mathbb{B}_{S[<k]}(\lambda)$  and do other reorderings afterward.  $\square$

**Corollary 2.40.** *The bubble-sort idempotent  $\text{bs}_S^k$  is monotone for all  $\lambda$ .*


*Proof.* The idempotent  $\text{bs}_S^k : \mathbb{B}_{S[<k]}(\lambda) \rightarrow \mathbb{B}_{S[\leq k]}$  is the composition of three monotone maps  $\pi_\lambda \circ \text{bs}_S^k \circ \psi_\lambda^+$  and is consequently also monotone.  $\square$

**Corollary 2.41.** *The poset  $\mathbb{D}_S^v(\lambda)$  is bounded.*

*Proof.* The poset  $\mathbb{A}_S^v$  is bounded (since it is an interval in the Bruhat graph), which implies that its image under the monotone idempotent is also bounded.  $\square$

Unfortunately, not all properties of the Bruhat graph carry over to the parabolic case, as seen in the following example.

**Example 2.42.** The poset  $\mathbb{S}_2 \setminus \mathbb{S}_4 / \mathbb{S}_2$  is not graded:



The following conjectures have been verified in several specific cases. Notably, Conjecture 2.43 implies Conjecture 2.44, which is essential for determining coefficients in the Cauchy identity (5.18) for staircase matrices.

**Conjecture 2.43.** *For any partition  $\lambda$  with repeating elements, the poset  $\mathbb{D}_S(\lambda)$  is shellable.*

**Conjecture 2.44.** *The Möbius function  $\mu^{\mathbb{D}_S(\lambda)}(-, -)$  on the poset  $\mathbb{D}_S(\lambda)$  takes values in the set  $\{-1, 0, 1\}$ .*

**2.4.3. Arborescent Poset with Consistent Linearization.** We conclude this section by considering the opposite case of an arborescent poset  $\mathbb{T}$  enhanced with an consistent **linearization** (Definition 2.1).

**Theorem 2.45.** *Suppose that  $(\mathbb{T}, \prec)$  is an arborescent poset with an order-preserving consistent linearization  $h : \mathbb{T} \hookrightarrow [1n]$ . Then, if  $l(\lambda) \leq \#\mathbb{T}$ , the following properties hold for the set  $\mathbb{D}_{\mathbb{T}}^h(\lambda)$  of  $\mathbb{T}$ -dominant compositions:*

- *The set  $\mathbb{D}_{\mathbb{T}}^h(\lambda)$  is empty if  $l(\lambda) > \#\mathbb{T}$ .*
- *If  $l(\lambda) \leq \#\mathbb{T}$ , then:*
  - *The covering relations in the Bruhat partial order on the set  $\mathbb{D}_{\mathbb{T}}^h(\lambda)$  of  $\mathbb{T}$ -dominant compositions are given by transpositions in minimal disorders.*
  - *There exists a nondecreasing monotone projection  $\mathbf{bs}_{\mathbb{T}}^{\text{op}} : [\lambda_+, \lambda_-] \rightarrow \mathbb{D}_{\mathbb{T}}^h(\lambda)$  from the interval in the Bruhat graph  $\mathbb{S}_n \lambda$  to the set of  $\mathbb{T}$ -dominant compositions.*
  - *The poset  $\mathbb{D}_{\mathbb{T}}^h(\lambda)$  is bounded.*
- *If  $l(\lambda) = \#\mathbb{T}$ , then the poset  $\mathbb{D}_{\mathbb{T}}^h$  is graded, subthin, and EL-shellable.*

*Proof.* This follows from Remark 2.4 and the results already established for the consistent order-reversing anti-linearization of  $\mathbb{T}$ . □

### 3. DISTRIBUTIVE LATTICE OF DEMAZURE SUBMODULES

**3.1. Demazure Modules.** Let  $\mathfrak{g}$  be a semi-simple (or reductive) Lie algebra, and let  $\mathfrak{b} = \mathfrak{b}^+, \mathfrak{b}^-$  be its positive and negative Borel subalgebras, with  $\mathfrak{h} = \mathfrak{b}^+ \cap \mathfrak{b}^-$  being the Cartan subalgebra. Let  $\Phi = \Phi_+ \sqcup \Phi_-$  denote the corresponding root system, and let  $P$  be the weight lattice of  $\mathfrak{g}$  with  $P_+$  as the subset of dominant weights. The Weyl group  $W$  acts naturally on the weight lattice  $P$  and for any given weight  $\nu \in P$ , there exists a unique element  $\sigma_\nu \in W$  of minimal length such that  $\nu = \sigma_\nu(\lambda)$  for some  $\lambda \in P_+$ .

Let  $V_\lambda$  be the irreducible finite-dimensional  $\mathfrak{g}$ -module with highest weight  $\lambda$ . The  $\nu$ -weight subspace of  $V_\lambda$  is one-dimensional, and we denote by  $v_\nu$  a generator of this space (in particular,  $v_\nu \in V_\lambda$  is an extremal weight vector). M. Demazure introduced ([Dem1]) the submodule

$$D_\nu := U(\mathfrak{b}^+)v_\nu \subset V_\lambda,$$

defined as the submodule generated by the extremal vector  $v_\nu$ , and established the following key observation:

**Fact 3.1.** *For all  $\mu, \nu \in W\lambda_+$ , the following equivalence holds:*

$$\nu \prec_{\text{Br}} \mu \iff D_\nu \subset D_\mu.$$

These modules were later named Demazure modules and were described in terms of generators and relations:

**Theorem 3.2** ([J]). *For any  $\nu \in P$ , the defining relations for the Demazure  $\mathfrak{b}$ -module  $D_\nu$  are given by:*

$$(3.3) \quad e_\alpha^{\max\{-(\alpha^\vee, \lambda), 0\}+1} v_\nu = 0, \quad \forall \alpha \in \Phi_+,$$

where  $e_\alpha \in \mathfrak{b}_n$  is the Chevalley generator corresponding to the root  $\alpha$ .



We will also need the van der Kallen modules  $K_\lambda$  introduced by van der Kallen in [vdK], which are  $\mathfrak{b}$ -modules defined as:

$$K_\nu = D_\nu / \sum_{\mu \prec \nu} D_\mu = D_\nu / \sum_{\substack{\mu \in W\lambda \\ D_\mu \subsetneq D_\lambda}} D_\mu.$$

and shown to be cyclic  $\mathfrak{b}$ -modules defined by the following relations:

$$(3.4) \quad e_\alpha^{\max(1, \max\{-\langle \alpha^\vee, \nu \rangle, 0\})} v_\nu = 0, \quad \forall \alpha \in \Phi_+.$$

**Remark 3.5.** *The main example for us is  $\mathfrak{g} = \mathfrak{gl}_n$ , where  $\mathfrak{b}^+ = \mathfrak{b}_n^+$  and  $\mathfrak{b}^- = \mathfrak{b}_n^-$  are the Lie subalgebras of upper-triangular and lower-triangular matrices, respectively, and  $\mathfrak{h}_n = \mathfrak{b}_n^+ \cap \mathfrak{b}_n^-$  is the diagonal Cartan subalgebra. The space  $\mathfrak{h}_n$  is spanned by the diagonal matrix units, and its dual basis is denoted by  $\{\varepsilon_i : i = 1, \dots, n\}$ . In this case,  $P_+ \subset P$  consists of weights  $\lambda = \sum_{i=1}^n \lambda_i \varepsilon_i$  such that  $\lambda_i \geq \lambda_{i+1}$  for all  $i$ . Thus, weights can be identified with compositions, and dominant weights are in bijection with partitions of length at most  $n$ . The Weyl group  $W$  is isomorphic to the symmetric group  $\mathbb{S}_n$ .*

*The weight  $\lambda = \sum_{i=1}^n \lambda_i \varepsilon_i$  can be considered as a weight of  $\mathfrak{gl}_m$  for any  $m \geq n$ . In this setting, the corresponding composition is given by  $(\lambda_1, \dots, \lambda_n, 0, \dots, 0)$ .*

For an  $\mathfrak{h}_n$ -module  $M$  and  $\mu \in P$ , let  $M(\mu) \subset V$  be the weight  $\mu$  subspace. We define the character as follows:

$$(3.6) \quad \text{ch}_{\mathfrak{h}_n} M := \sum_{\mu=(\mu_1, \dots, \mu_n)} x_1^{\mu_1} \dots x_n^{\mu_n} \dim M(\mu).$$

In particular, for a partition  $\lambda_+$ , the character of the corresponding irreducible highest weight  $\mathfrak{gl}_n$  module  $V_{\lambda_+}$  is given by the Schur function  $s_{\lambda_+}(x_1, \dots, x_n)$ . The Demazure modules  $D_\nu$  and the van der Kallen modules  $K_\nu$  are labeled by compositions (i.e., arbitrary weights  $\nu \in P$ ). One has

$$\kappa_\nu(x_1, \dots, x_n) = \text{ch}_{\mathfrak{h}} D_\nu(x_1, \dots, x_n), \quad a_\nu(x_1, \dots, x_n) = \text{ch}_{\mathfrak{h}} K_\nu(x_1, \dots, x_n),$$

where  $a_\nu(x)$  and  $\kappa_\nu(x)$  are the Demazure atoms and key polynomials (see [Al, P] and references therein).

In this paper, we deal with left and right modules over the Borel subalgebra  $\mathfrak{b}_m \subset \mathfrak{gl}_m$ . To emphasize the distinction, we use the upper index **op** for right modules. Note that the right  $\mathfrak{b}_m$ -action decreases the weight, whereas the left action increases it. In particular, the right  $\mathfrak{gl}_m$ -module  $V_\lambda^{\text{op}}$ , is generated by a cyclic vector  $v_\lambda$  as a  $\mathfrak{b}_m$ -module.

**Notation 3.7.** *For each weight  $\nu$ , along with its corresponding dominant weight  $\lambda_+$  and antidominant weight  $\lambda_- = w_0 \lambda_+$ , we define the right Demazure  $\mathfrak{b}_m$ -module  $D_\nu^{\text{op}}$  as the submodule of the right  $\mathfrak{gl}_m$ -module  $V_{\lambda_+}^{\text{op}}$  generated by the extremal vector of right weight  $\nu$ .*

Note that we have an opposite inclusion for opposite (right) Demazure  $\mathfrak{b}_n^+$ -modules:

$$\nu \preceq_{\text{Br}} \mu \iff D_\nu^{\text{op}} \supset D_\mu^{\text{op}}.$$

In particular, the opposite van der Kallen modules are described as follows:

$$K_\nu^{\text{op}} = D_\nu^{\text{op}} / \sum_{\mu \succ_{\text{Br}} \nu} D_\mu^{\text{op}}.$$

Moreover, the Cartan subalgebra  $\mathfrak{h}_m \subset \mathfrak{b}_m$  acts on the opposite (right) Demazure and van der Kallen modules, yielding the opposite key polynomials and opposite Demazure atoms:

$$\kappa^\lambda(y_1, \dots, y_m) = \text{ch}_{\mathfrak{h}_m} D_\lambda^{\text{op}}, \quad a^\lambda(y_1, \dots, y_m) = \text{ch}_{\mathfrak{h}_m} K_\lambda^{\text{op}}.$$

Directly from the definition we have

$$(3.8) \quad \kappa^\lambda(y_1, \dots, y_m) = \kappa_{w_0\lambda}(y_m, \dots, y_1), \quad a^\lambda(y_1, \dots, y_m) = a_{w_0\lambda}(y_m, \dots, y_1).$$

Here,  $w_0$  is the longest element in the Weyl group. In the case of the symmetric group, it reverses the composition:

$$w_0(\lambda_1, \dots, \lambda_n) = (\lambda_n, \dots, \lambda_1).$$

Note that both  $\kappa^\lambda(y_1, \dots, y_m)$  and  $a^\lambda(y_1, \dots, y_m)$  contain the term  $y^\lambda$ , which serves as the leading term in the Bruhat order.

**Example 3.9.** Let  $\lambda = \lambda_+$  be a partition (i.e., a dominant composition), and let  $\lambda_- = w_0\lambda_+$  be the corresponding antidominant composition. Then we have the following isomorphisms:

$$D_{\lambda_-} \simeq V_{\lambda_+}, \quad \dim(D_{\lambda_+}) = \dim(D_{\lambda_-}^{\text{op}}) = 1, \quad D_{\lambda_+}^{\text{op}} \simeq V_{\lambda_+}.$$

Consequently, we obtain the following equalities for the corresponding polynomials:

$$\kappa_{\lambda_-}(x) = s_\lambda(x) = \kappa^{\lambda_+}(x), \quad \kappa_{\lambda_+}(x) = a_{\lambda_+}(x) = x^{\lambda_+}; \quad \kappa^{\lambda_-}(x) = a^{\lambda_-}(x) = x^{\lambda_-}.$$

**3.2. Demazure Modules Form a Distributive Lattice.** In this section, we provide a proof of Theorem 3.10 below. Although the statement of the theorem appears to be well known to experts, we were unable to find a precise reference. We are grateful to Michel Brion for explaining the key idea of the proof to us.

Let  $G$  be a simple Lie group, and let  $V_\lambda$  be its irreducible finite-dimensional highest weight representation. Demazure modules naturally arise in the geometry of the flag variety  $F = G/B$ . In particular, it is known that there exists a natural line bundle  $\mathcal{L}_\lambda$  associated with  $\lambda \in P_+$  such that

$$H^0(F; \mathcal{L}_\lambda) \simeq V_\lambda^*, \quad H^0(X_w; \mathcal{L}_\lambda) \simeq D_{w\lambda}^*.$$

Here,  $X_w \subset F$  is the Schubert variety corresponding to  $w$ . (See, e.g., [Ku] for an introduction to the geometry of flag varieties). Let  $D_{w\lambda}^\perp \subset H^0(F; \mathcal{L}_\lambda)$  denote the annihilator of  $D_{w\lambda}$ :

$$D_{w\lambda}^\perp := \ker(H^0(F, \mathcal{L}_\lambda) \rightarrow H^0(X_w, \mathcal{L}_\lambda)) \simeq \ker(V_\lambda^* \rightarrow D_{w\lambda}^*).$$

**Theorem 3.10.** *The subspaces  $\{D_{w\lambda} : w \in W\}$  of the integrable representation  $V_\lambda$  generate a distributive lattice, denoted  $\mathcal{L}_D(V_\lambda)$ . Equivalently, the dual lattice generated by  $\{D_{w\lambda}^\perp : w \in W\} \subset L(V_\lambda^*)$  is distributive.*

*Moreover, Demazure modules constitute a set of  $\vee$ -irreducible elements of this lattice  $\mathcal{L}_D(V_\lambda)$ .*

*Proof.* From the definition of the Bruhat order, we know that for any dominant weight  $\lambda$  and any pair of comparable elements in the Weyl group  $W$ , we have:

$$w_1 \prec_{\text{Br}} w_2 \in W \Rightarrow D_{w_1\lambda} \subset D_{w_2\lambda} \Leftrightarrow D_{w_1\lambda}^\perp \supset D_{w_2\lambda}^\perp.$$

Consequently, for any collection of pairwise incomparable elements  $w_1, \dots, w_k$  in the Weyl group  $W$ , we obtain the inclusion

$$(3.11) \quad D_{w_1\lambda}^\perp + \dots + D_{w_k\lambda}^\perp \subset \bigcap_{w: \forall i \ w \preceq w_i} D_{w\lambda}^\perp.$$

We now show that this inclusion is actually an equality. In other words, we demonstrate that if a section  $s \in H^0(F, \mathcal{L}_\lambda)$  vanishes when restricted to the intersection of Schubert varieties  $\cap_{i=1}^k X_{w_i}$ , then it can be written as a sum  $s = s_1 + \dots + s_k$  of  $k$  sections, where each  $s_i$  vanishes on  $X_{w_i}$ . Denote by  $I_Y$  the ideal sheaf of the structure sheaf of  $F$  consisting of functions vanishing on a subvariety  $Y \subset F$ . In particular, for Schubert varieties  $X_w$ , we have

$$D_{w\lambda}^\perp \simeq H^0(F; \mathcal{L}_\lambda \otimes I_{X_w}).$$

Since the schematic intersection of Schubert varieties is reduced, we obtain

$$\begin{aligned} I_{X_{w_1} \cap \dots \cap X_{w_k}} &= I_{X_{w_1}} + \dots + I_{X_{w_k}}; \\ I_{X_{w_1} \cup \dots \cup X_{w_k}} &= I_{X_{w_1}} \cap \dots \cap I_{X_{w_k}}. \end{aligned}$$

Thus, we have an exact sequence of ideal sheaves (originating from the inclusion-exclusion principle):

$$0 \longrightarrow I_{\cup_{i=1}^k X_{w_i}} \longrightarrow \bigoplus_{\substack{s=1 \\ i \neq s}}^k I_{\cup_{i=1}^k X_{w_i}} \longrightarrow \dots \longrightarrow \bigoplus_{s=1}^k I_{X_{w_s}} \twoheadrightarrow I_{\cap_{i=1}^k X_{w_i}} \longrightarrow 0.$$

Due to the Frobenius splitting of the flag variety  $F$ , we know that for any subset  $\{u_1, \dots, u_m\} \subset W$ , the higher cohomology vanishes (see [BrKu, Theorems 1.4.8 and 2.3.1]):

$$H^{>0}(F; \mathcal{L} \otimes I_{\cup_{i=1}^m X_{u_i}}) = 0.$$

This leads to an exact sequence of sections of the line bundle  $\mathcal{L}$ :

$$\begin{aligned} 0 \rightarrow H^0(F; \mathcal{L} \otimes I_{\cup_{i=1}^k X_{w_i}}) \rightarrow \bigoplus_{\substack{s=1 \\ i \neq s}}^k H^0(F; \mathcal{L} \otimes I_{\cup_{i=1}^k X_{w_i}}) \rightarrow \dots \rightarrow \\ \rightarrow \bigoplus_{s=1}^k H^0(F; \mathcal{L} \otimes I_{X_{w_s}}) \rightarrow H^0(F; \mathcal{L} \otimes I_{\cap_{i=1}^k X_{w_i}}) \rightarrow 0. \end{aligned}$$

Thus, we obtain the inclusion-exclusion formula for the intersection of Demazure modules and a surjective map

$$\bigoplus_{i=1}^k D_{w_i\lambda}^\perp \twoheadrightarrow H^0(F; \mathcal{L} \otimes I_{\cap_{i=1}^k X_{w_i}}) = \bigcap_{w: \forall iw \leq w_i} D_{w\lambda}^\perp.$$

From this, it follows that the inclusion (3.11) is actually an equality, establishing that the lattice of submodules  $D_{w\lambda}^\perp$  of the integrable module  $V(\lambda)^*$  is distributive.

It remains to show that Demazure submodules are  $\vee$ -irreducible elements of the lattice  $\mathcal{L}_D(V_\lambda)$ . This follows from the observation:

$$\forall U \in \mathcal{L}_D(V_\lambda), \quad D_{w\lambda} \subset U \Leftrightarrow v_{w\lambda} \in U.$$

□

It is worth mentioning that all subspaces of the lattice  $\mathcal{L}_D(V_\lambda)$  are  $\mathfrak{b}$ -submodules of  $V_\lambda$  and, in particular,  $\mathfrak{h}$ -modules. Consequently, each quotient  $U_1/U_2$  has a well-defined character  $\chi$  (see (3.6)). Let us outline the corollaries of §1.3, which apply naturally to our situation:

**Corollary 3.12.** *Let  $U \in \mathcal{L}_D(V_\lambda)$ , and let  $X_U \subset W\lambda$  be the subset of the parabolic Bruhat graph consisting of those  $w\lambda$  such that*

$$U \supset D_{w\lambda} \Leftrightarrow \text{the extremal vector } v_{w\lambda} \in U.$$

*Then we have:*

- $U \simeq \sum_{\mu \in X_U} D_\mu$ ;
- For the filtration  $\mathcal{F}_\mu U := U \cap D_\mu$ , the associated graded space has the following description:

$$\mathcal{F}_\mu U / \sum_{\nu \prec \mu} \mathcal{F}_\nu U \simeq \begin{cases} \mathbb{K}_\mu, & \text{if } \mu \in X_U; \\ 0, & \text{if } \mu \notin X_U. \end{cases}$$

where  $\mathbb{K}_\mu$  is the van der Kallen module;

- The character of  $U$  is equal to the sum of Demazure atoms:

$$\text{ch}(U) = \sum_{\mu \in X_U} a_\mu(x).$$

- The character of  $U$  is a linear combination of key polynomials  $\{\kappa_\nu : \nu \in W\lambda\}$  with integer coefficients.

*Proof.* The van der Kallen module  $K_{w\lambda}$  coincides with the one described in (1.31). Each van der Kallen module contains a unique nonzero image of an extremal vector from  $V_\lambda$ . Thus, the first three items of the corollary follow directly from the general theory outlined in Fact 1.30. The last item can be deduced either from Proposition 1.32 or from the Möbius inversion formula (Fact 1.8).  $\square$

**3.3. Sublattice Indexed by S-Dominant Weights.** Let us now apply the combinatorial results from §2 and the findings of the preceding section §3.2 to the specific case of  $\mathfrak{g} = \mathfrak{gl}_n$ ,  $W = \mathbb{S}_n$ , and an arborescent poset with (anti)linearization.

**Corollary 3.13.** *Suppose that  $(\mathbb{T}, h)$  is an arborescent poset with linearization (Definition 2.1) and that  $\lambda$  is a partition of length at most  $\#\mathbb{T}$ . Then the set*

$$\{D_{\bar{\alpha}} \subset V_\lambda : \bar{\alpha} \in \mathbb{D}_{\mathbb{T}}(\lambda)\}$$

*forms the collection of  $\vee$ -irreducible elements of the distributive sublattice  $\mathcal{L}_D^{\mathbb{T}}(\lambda)$  of  $\mathcal{L}_D(\lambda)$  generated by these elements.*

*Moreover, the minimal subquotients:*

$$(3.14) \quad \mathbb{K}_{\mathbb{T}, \bar{\alpha}} := D_{\bar{\alpha}} \Big/ \sum_{\bar{\tau} \in \mathbb{D}_{\mathbb{T}}(\lambda) : \bar{\tau} \prec \bar{\alpha}} D_{\bar{\tau}}$$

*admit a filtration induced by  $\mathcal{L}_D(\lambda)$ , indexed by the elements of the Bruhat graph  $W\lambda$ , such that the associated graded module satisfies:*

$$(3.15) \quad \text{gr}^{\mathcal{F}} \mathbb{K}_{\mathbb{T}, \bar{\alpha}} \simeq \bigoplus_{\bar{b} \in \mathbb{S}_n \lambda : \text{bs}_{\mathbb{T}}^{\text{op}}(\bar{b}) = \bar{\alpha}} \mathbb{K}_{\bar{b}}.$$

*The modules  $\mathbb{K}_{\mathbb{T}, \bar{\alpha}}$  are referred to as generalized van der Kallen modules.*

*Proof.* Recall that the opposite bubble-sort operation  $\text{bs}_{\mathbb{T}}^{\text{op}}$  defines an increasing monotone idempotent on the interval of Bruhat graph, whose image consists of  $\mathbb{T}$ -antidominant compositions (Theorem 2.45). Consequently, the assumptions of Proposition 1.33 are satisfied, and it remains to carefully apply these results in our setting.  $\square$

**Corollary 3.16.** *We have the following two equivalent descriptions of the  $\mathfrak{h}_n$ -character of the minimal subquotients:*

$$(3.17) \quad \text{ch}_{\mathfrak{h}_n}(\mathbb{K}_{\top, \bar{a}})(x) = \sum_{\substack{\bar{c} \in \mathbb{D}_{\top}(\lambda) \\ \bar{c} \leq_{\text{Br}} \bar{a}}} \mu^{\mathbb{D}_{\top}(\lambda)}(\bar{c}, \bar{a}) \kappa_{\bar{c}}(x) = \sum_{\bar{b}: \text{bs}_{\top}^{\text{op}}(\bar{b}) = \bar{a}} a_{\bar{b}}(x).$$

Here,  $\mu^{\mathbb{D}_{\top}(\lambda)}(\bar{c}, \bar{a})$  is the Möbius function on the poset  $\mathbb{D}_{\top}(\lambda)$  discussed in Subsection 1.1,  $\kappa_{\bar{c}}(x)$  is the key polynomial, and  $a_{\bar{a}}(x)$  is the Demazure atom associated with the composition  $\bar{a}$ .

*Proof.* The first equality follows from the description (3.14) of the generalized van der Kallen module, while the second follows from the associated graded structure given in (3.15).  $\square$

Let us also formulate a similar statement for the right Demazure modules:

**Corollary 3.18.** *Suppose that  $(\mathcal{S}, v)$  is an arborescent poset with a linearization and that  $\lambda$  is a partition of length at most  $\#\mathcal{S}$ . Then the right Demazure submodules*

$$\{D_{\bar{a}}^{\text{op}} \subset V_{\lambda}^{\text{op}} \simeq V_{-w_0\lambda} : \bar{a} \in \mathcal{S}_n\lambda\}$$

*form the set of  $\vee$ -irreducible elements of the distributive lattice  $\mathcal{L}_D^{\text{op}}(\lambda)$ . The subset of Demazure submodules indexed by  $\mathcal{S}$ -dominant compositions constitutes a subset of  $\vee$ -irreducible elements of the distributive sublattice generated by them:*

$$\mathcal{L}_D^{\mathcal{S}}(\lambda)^{\text{op}} := \langle D_{\bar{a}}^{\text{op}} : \bar{a} \in \mathbb{D}_{\mathcal{S}}(\lambda) \rangle \subset \langle D_{\bar{a}}^{\text{op}} : \bar{a} \in \mathcal{S}_n\lambda \rangle =: \mathcal{L}_D(\lambda)^{\text{op}}.$$

*The minimal subquotients:*

$$\mathbb{K}_{\mathcal{S}, \bar{a}}^{\text{op}} := D_{\bar{a}}^{\text{op}} \Big/ \sum_{\mu \in \mathbb{D}_{\mathcal{S}}(\lambda): \mu \succ \bar{a}} D_{\mu}^{\text{op}}$$

*have the following two descriptions of their characters:*

$$(3.19) \quad \text{ch}_{\mathfrak{h}_m}(\mathbb{K}_{\mathcal{S}, \bar{a}}^{\text{op}}) = \sum_{\substack{\bar{c} \in \mathbb{D}_{\mathcal{S}}(\lambda) \\ \bar{c} \geq_{\text{Br}} \bar{a}}} \mu^{\mathbb{D}_{\mathcal{S}}(\lambda)}(\bar{c}, \bar{a}) \kappa_{\bar{c}}(y) = \sum_{\bar{a}: \text{bs}_{\mathcal{S}}(\bar{a}) = \bar{a}} a_{\bar{a}}(y).$$

It is worth mentioning that, on one hand, any subspace  $D \in \mathcal{L}_D^{\top}(\lambda)$  admits a resolution by a direct sum of Demazure modules (Proposition 1.32). Consequently, whenever we are able to show that the Möbius function  $\mu^{\mathbb{D}_{\top}(\lambda)}(\bar{c}, \bar{a})$  takes values in  $\{0, 1, -1\}$  (Corollary 2.36 and Conjecture 2.44) we will be able to extract an explicit description of this resolution for the minimal subquotients  $\mathbb{K}_{\top, \bar{a}}$ , generalizing the classical BGG resolution.

#### 4. STAIRCASE SHAPES, CORNERS AND $DL$ -DENSE ARRAYS

In the first two subsections, §4.1–§4.2, we recall the main combinatorial objects and notations introduced in [FKhM3]. In §4.4, we develop new combinatorial structures, relate them to those introduced in §2, and apply them in §4.5.

**4.1. Young Diagrams in Staircase Shape.** The main focus of this note is the combinatorics surrounding staircase matrices, which arise as an output of the Gaussian elimination process. The shapes of these staircase matrices are indexed by Young diagrams, drawn in a reversed form.

Let us fix a positive integer  $m$  and a sequence of integers  $\bar{n} := (n_1, \dots, n_m)$  satisfying

$$0 < n_1 \leq n_2 \leq \dots \leq n_m.$$

The corresponding Young diagram (or staircase shape) is defined as

$$\mathbf{Y}_{\bar{n}} = \{(i, j) : 1 \leq j \leq m, 1 \leq i \leq n_j\} \subset \mathbb{Z}_{>0}^2.$$

This diagram is visualized as a collection of  $m$  columns of heights  $n_1, \dots, n_m$ , arranged from left to right. In particular, the first index  $i$  corresponds to the row, while the second index  $j$  denotes the column number. For example, the cells in the top row have the form  $(1, j)$  for  $1 \leq j \leq m$ .

**Example 4.1.** Here is an example of a Young diagram  $\mathbf{Y}_{\bar{n}}$  associated with  $\bar{n} = (1^3 3^3 4^2)$ :

$$\mathbf{Y}_{(1^3 3^3 4^2)} = \begin{array}{cccccccc} \square & \square & \square & \square & \square & \square & \square & \square \\ & & & \square & \square & \square & \square & \square \\ & & & & \square & \square & \square & \square \\ & & & & & \square & \square & \square \\ & & & & & & \square & \square \end{array}.$$

The reason for rotating the visualization of the Young diagram is that our primary focus is on staircase matrices, which we want to be acted upon by upper-triangular matrices from both the left and the right.

**Notation 4.2.** *The subspace of rectangular  $n_m \times m$  matrices consisting of matrices with zero entries outside the cells belonging to the Young diagram  $\mathbf{Y}_{\bar{n}}$  is called the set of staircase matrices of shape  $\mathbf{Y}_{\bar{n}}$  and is denoted by  $\text{Mat}_{\bar{n}}$ .*

**4.2. Poset of Staircase Corners.** In this subsection, we associate each partition  $\bar{n}$  with a subset of indices  $\mathfrak{S}_{\bar{n}} \subset \mathbf{Y}_{\bar{n}}$ , called the staircase corners. We define a partial order on this set and describe some fundamental properties of the resulting poset.

For a cell  $(i, j) \in \mathbf{Y}_{\bar{n}}$ , we define the partition  $\bar{n}^{ij} = (n_1^{ij} \leq \dots \leq n_{m-1}^{ij})$  obtained from  $\bar{n}$  by removing the  $i$ -th row and the  $j$ -th column:

$$(4.3) \quad n_k^{ij} := \begin{cases} n_k, & \text{if } k < j \text{ \& } n_k < i, \\ n_k - 1, & \text{if } k < j \text{ \& } n_k \geq i, \\ n_{k+1} - 1, & \text{if } k \geq j. \end{cases}$$

There is a natural bijection between the cells of  $\mathbf{Y}_{\bar{n}}$  with the  $i$ -th row and  $j$ -th column removed and the cells of  $\mathbf{Y}_{\bar{n}^{ij}}$ , denoted by  $\pi_{i,j}$ :

$$(4.4) \quad \pi_{i,j}(s, t) := \begin{cases} (s, t) & \text{if } s < i \text{ \& } j < t, \\ (s - 1, t) & \text{if } s > i \text{ \& } j < t, \\ (s, t - 1) & \text{if } s < i \text{ \& } j > t, \\ (s - 1, t - 1) & \text{if } s > i \text{ \& } j > t. \end{cases}$$

**Definition 4.5.** *The subset  $\mathfrak{S}_{\bar{n}} \subset \mathbf{Y}_{\bar{n}}$  of staircase corners is defined inductively by the following properties:*

- *In each row and each column, at most one cell (staircase corner) belongs to  $\mathfrak{S}_{\bar{n}}$ .*

- If  $n_j > n_{j-1}$ , then the corner cell  $(n_j, j)$  of the Young diagram  $\mathbf{Y}_{\bar{n}}$  belongs to  $\mathbf{Sc}_{\bar{n}}$ .
- For any  $(i, j) \in \mathbf{Sc}_{\bar{n}}$ , another cell  $(s, t)$  belongs to  $\mathbf{Sc}_{\bar{n}} \setminus \{(i, j)\}$  if and only if  $\pi_{ij}(s, t) \in \mathbf{Sc}_{\bar{n}_{ij}}$ . In other words, for any staircase corner  $(i, j) \in \mathbf{Sc}_{\bar{n}}$ , any other staircase corner is also a staircase corner of the diagram obtained by removing the  $i$ -th row and  $j$ -th column.

**Remark 4.6.** Suppose that the  $j$ 'th column of the Young diagram  $\mathbf{Y}_{\bar{n}}$  does not contain a staircase corner. Then by erasing the  $j$ -th column we get a bijection between  $\mathbf{Sc}_{\bar{n}}$  and  $\mathbf{Sc}_{\bar{n}_j'}$ , where  $\bar{n}_j' := (n_1 \leq \dots \leq \widehat{n}_j \leq \dots \leq n_m)$ . Similarly, if the  $i$ 'th row of the Young diagram  $\mathbf{Y}_{\bar{n}}$  does not contain a staircase corner and  $\bar{n}_i'$  is the partition then there is a bijection between  $\mathbf{Sc}_{\bar{n}}$  and  $\mathbf{Sc}_{\bar{n}_i'}$  where  $\bar{n}_i'$  is a partition with  $i$ 'th row omitted.


It was shown in [FKhM3, Lemma 1.24] that for any partition  $\bar{n}$ , the set  $\mathbf{Sc}_{\bar{n}}$  of staircase corners is well-defined. Moreover,  $\mathbf{Sc}_{\bar{n}}$  forms a "rook placement" in the Young diagram  $\mathbf{Y}_{\bar{n}}$ , meaning that for any cell  $(i, j) \in \mathbf{Sc}_{\bar{n}}$ , there are no other staircase corners in the same row or column. Evgeny Smirnov explained to us in private communication that the "rook placement" of staircase corners  $\mathbf{Sc}_{\bar{n}}$  is maximal under an appropriate order.

We equip the set  $\mathbf{Sc}_{\bar{n}}$  of staircase corners with the following partial order:

$$(4.7) \quad (i, j) \succeq (i', j') \stackrel{\text{def}}{\iff} \begin{cases} i \geq i', \\ j \leq j'. \end{cases}$$


In other words, an element is enlarged by moving in the down-left direction.

Below, we provide a few examples of the set  $\mathbf{Sc}_{\bar{n}}$  of staircase corners along with the Hasse diagram of the corresponding partial order. We represent the elements of  $\mathbf{Sc}_{\bar{n}}$  as blue dots and draw edges in the Hasse diagram oriented from smaller to larger elements. The first examples include rectangular shapes, triangular shapes, and a small mixture of both:



For a larger example  $\bar{n} = (3^4 5 9^2 13^{10} 16^2)$ , we attempt to visualize the algorithm for identifying the staircase corners. Specifically, we begin by coloring in lime the cells that belong to the rows and columns of the corner cells of the Young diagram. Next, we examine the remaining cells and mark in yellow the hooks associated with the corners of the Young diagram obtained after removing the lime cells. We continue this process, using a new color at each step, and mark the centers of the identified staircase corner cells with blue dots.

Finally, we overlay a Hasse diagram on top of the visualization.



Let us introduce special notations for the row index and the column number of a cell of the Young diagram  $\mathbf{Y}_{\bar{n}}$ :

$$\text{hor}((i, j)) := i, \quad \text{vrt}((i, j)) := j.$$

Since each row and each column contains at most one staircase corner, it follows that the row index and the column number define a linearization  $\text{hor} : \mathfrak{Sc}_{\bar{n}} \hookrightarrow [1, n_m]$  and an anti-linearization  $\text{vrt} : \mathfrak{Sc}_{\bar{n}} \hookrightarrow [1, m]$ :

$$(i, j) \prec (i', j') \in \mathfrak{Sc}_{\bar{n}} \Leftrightarrow \begin{cases} \text{hor}((i, j)) < \text{hor}((i', j')), \\ \text{vrt}((i, j)) > \text{vrt}((i', j')). \end{cases}$$

The following lemma (proven in [FKhM3, Lemma 1.24]) describes the fundamental properties of the poset  $(\mathfrak{Sc}_{\bar{n}}, \prec)$  of staircase corners.

**Lemma 4.9.** (i) *The Hasse diagram of the poset  $(\mathfrak{Sc}_{\bar{n}}, \preceq)$  forms a forest, with smaller elements positioned closer to the root of a tree. In other words,  $\mathfrak{Sc}_{\bar{n}}$  is an arborescent poset, where the map  $\text{hor} : \mathfrak{Sc}_{\bar{n}} \hookrightarrow [1, n_m]$  provides a consistent linearization, and the map  $\text{vrt} : \mathfrak{Sc}_{\bar{n}} \hookrightarrow [1, m]$  provides a consistent anti-linearization.*

(ii) *For any  $s \in \mathfrak{Sc}_{\bar{n}}$ , the subset  $\mathfrak{Sc}_{\bar{n}}\{\succeq s\}$  of elements greater than or equal to  $s$  forms an interval with respect to the linear order on  $\text{vrt}(\mathfrak{Sc}_{\bar{n}})$ :*

$$\forall s \in \mathfrak{Sc}_{\bar{n}} \quad \exists j_s \in [1, m] : \text{vrt}(\mathfrak{Sc}_{\bar{n}}\{\succeq s\}) = [j_s, \text{vrt}(s)].$$

Similarly, for the map  $\text{hor}$ :


$$\forall s \in \mathfrak{Sc}_{\bar{n}} \quad \exists i_s \in [1, n_m] : \text{hor}(\mathfrak{Sc}_{\bar{n}}\{\succeq s\}) = [\text{hor}(s), i_s].$$

(iii) *For any two incomparable elements  $s, t \in \mathfrak{Sc}_{\bar{n}}$ , the vertical and horizontal comparisons are equivalent:*

$$\text{vrt}(s) < \text{vrt}(t) \in [1, m] \Leftrightarrow \text{hor}(s) < \text{hor}(t) \in [1, n_m].$$



**Example 4.10.** In the following pictorial description, we consider the Young diagram  $\mathbf{Y}_{(2\ 3^3\ 5\ 7^4)}$ . The bullets denote the elements of the arborescent poset  $\mathbf{S}_{\bar{n}}$  and edges represent covering relations in this poset.



The following proposition describes the inverse procedure. Specifically, we seek a necessary and sufficient condition for a poset equipped with a consistent (anti)linearization to correspond to a staircase shape  $\mathbf{Y}_{\bar{n}}$ .

**Proposition 4.11.** *For any given arborescent poset  $(\mathbf{S}, \prec)$  with a consistent order-preserving linearization  $h : (\mathbf{S}, \prec) \rightarrow ([1, n_m], <)$  satisfying the following properties:*

$$\begin{cases} \forall s \in \mathbf{S}, \exists a(s), b(s) \text{ such that } h(\mathbf{S}_{\succeq s}) = [a(s), b(s)] \subset [1, n_m], \\ h^{-1}(n_m) \neq \emptyset, \end{cases}$$

*there exists a partition  $\bar{n} := (n_1 \leq \dots \leq n_m)$  and an order-preserving bijection  $\psi : \mathbf{S} \xrightarrow{\cong} \mathbf{S}_{\bar{n}}$ , such that  $h = \text{hor} \circ \psi$ .*

*Similarly, for any given arborescent poset  $(\mathbf{S}, \prec)$  with a consistent antilinear injection  $v : \mathbf{S} \rightarrow [1, n_m]$  that maps the subsets  $\mathbf{S}_{\succeq s}$  to subintervals of  $[1, m]$ , there exists a partition  $\bar{n} := (n_1 \leq \dots \leq n_m)$  and an order-preserving bijection  $\psi : \mathbf{S} \xrightarrow{\cong} \mathbf{S}_{\bar{n}}$ , such that  $v = \text{vrt} \circ \psi$  (provided that  $v^{-1}(1) \neq \emptyset$ ).*

*Proof.* It suffices to prove the second statement since the first follows by transposing the Young diagram  $\mathbf{Y}_{\bar{n}}$ .

The partition  $\bar{n}$  is constructed as follows:

$$\begin{cases} \text{If } \exists s \text{ such that } v(s) = k, \text{ then } n_k := \#\{t \in \mathbf{S} : v(t) < k\} + \#\{t \in \mathbf{S} : t \preceq s\}; \\ \text{If } v^{-1}(k) = \emptyset, \text{ then } n_k := \#\{t \in \mathbf{S} : v(t) < k\}. \end{cases}$$

□

Note that every arborescent poset  $\mathbf{S}$  with an isomorphic (anti)linearization admits a realization as a subset of staircase corners for an appropriate  $\bar{n}$ . However, if the (anti)linearization  $v : \mathbf{S} \hookrightarrow [1, m]$  is not surjective, it is necessary that each connected component of the Hasse graph of  $\mathbf{S}$  maps onto a closed interval of  $[1, m]$ .

**4.3. DL-dense arrays.** The *DL*-dense arrays defined below play a central role in this paper as well as in our previous work [FKhM3]. First, we recall the definition, introduce the Bruhat partial order on this set and establish several key properties of this partial order in Section 4.4.

**Definition 4.12.**

- A map  $A : \mathbf{Y}_{\bar{n}} \rightarrow \mathbb{Z}_{\geq 0}$  that assigns a nonnegative integer  $A_{i,j}$  to each cell  $(i, j) \in \mathbf{Y}_{\bar{n}}$  is called an array of shape  $\mathbf{Y}_{\bar{n}}$ .
- The total sum  $|A| := \sum_{(i,j) \in \mathbf{Y}_{\bar{n}}} A_{i,j}$  is called the degree of the array  $A$ .

- The sum of the entries in each row is called the horizontal weight of the array:

$$\text{hor}(A) := \left( \sum_{j=1}^m A_{1j}, \dots, \sum_{j=1}^m A_{n_m j} \right) \in \mathbb{Z}^{n_m}.$$

- The vertical weight of an array is defined as

$$\text{vrt}(A) := \left( \sum_{i=1}^{n_1} A_{i1}, \dots, \sum_{i=1}^{n_m} A_{im} \right) \in \mathbb{Z}^m,$$

i.e., it is the collection of column sums.

**Definition 4.13.** An array  $A$  of shape  $\mathbf{Y}_{\bar{n}}$  is called DL-dense if

$$\begin{cases} (i, j) \notin \mathbf{S}_{\bar{n}} \Rightarrow A_{i,j} = 0; \\ (i, j) \prec (i', j') \in \mathbf{S}_{\bar{n}} \Rightarrow A_{i,j} \leq A_{i',j'}. \end{cases}$$


**Remark 4.14.** The set of DL-dense arrays is in one-to-one correspondence with the set of order-preserving  $\mathbb{Z}_{\geq 0}$ -valued functions on the poset  $(\mathbf{S}_{\bar{n}}, \prec)$ .

**Definition 4.15.** For each partition  $\lambda = \{\lambda_1 \geq \dots \geq \lambda_k \geq 0\}$  (whose length is at most  $\#\mathbf{S}_{\bar{n}}$ ), we denote by  $\text{DL}_{\bar{n}}(\lambda)$  the set of DL-dense arrays  $A$  such that the multiset  $\{A_s : s \in \mathbf{S}_{\bar{n}}\}$  coincides with the multiset  $\{\lambda_i\}_{i=1}^k$ .

**Remark 4.16.** We have the following decomposition of the set of all DL-dense arrays of shape  $\mathbf{Y}_{\bar{n}}$  with respect to degree and partition:

$$\text{DL}_{\bar{n}} = \bigsqcup_{N \geq 0} \bigsqcup_{\substack{\lambda \vdash N \\ l(\lambda) \leq \#\mathbf{S}_{\bar{n}}}} \text{DL}_{\bar{n}}(\lambda).$$

**Example 4.17.** Here is an example of a DL-dense array of shape  $\mathbf{Y}_{(2,3,3,4)}$ :



**Remark 4.18.** The term array was introduced by Danilov and Koshevoy for rectangular shapes (see, e.g., [DK1, DK2]). They consider arrays as collections of balls placed in the cells of the diagram  $\mathbf{Y}_{\bar{n}}$ . One of the key features of their construction is the elegant bi-crystal structure, where balls can move up and down but only within consecutive columns or rows. The D-dense property means that moving a ball downward is not allowed, while the L-dense property prohibits moving any ball to the left.

The main difference between our approach and the one proposed in [DK1] is that the crystal structure alone is insufficient for our purposes. Instead, we must consider the action of all roots, not just the simple ones, as in the case of crystals.

Nevertheless, it is possible to define D-dense and L-dense arrays separately, describe the combinatorial densification maps, and relate this process to the bubble-sort map discussed in §2.3. We will explore these ideas in a separate paper.

**Proposition 4.19.** *The horizontal (vertical) weight of a  $DL$ -dense array is an  $\mathbf{Sc}_{\bar{n}}$ -dominant composition of length  $n_m$  (resp.  $m$ ), and the maps  $\text{hor}$  and  $\text{vrt}$  define bijections:*

$$\mathbb{S}_{n_m} \lambda \longleftarrow \mathbb{D}_{\mathbf{Sc}_{\bar{n}}}^{\text{hor}}(\lambda) \xleftarrow[\text{hor}]{\simeq} \text{DL}_{\bar{n}}(\lambda) \xrightarrow[\text{vrt}]{\simeq} \mathbb{D}_{\mathbf{Sc}_{\bar{n}}}^{\text{vrt}}(\lambda) \longrightarrow \mathbb{S}_m \lambda.$$

Here,  $\mathbb{D}_{\mathbf{Sc}_{\bar{n}}}^{\text{hor}}(\lambda)$  denotes the set of  $\mathbf{Sc}_{\bar{n}}$ -dominant compositions associated with the arborescent poset  $\mathbf{Sc}_{\bar{n}}$  equipped with the consistent linearization  $\text{hor} : \mathbf{Sc}_{\bar{n}} \hookrightarrow [1, n_m]$ . Similarly,  $\mathbb{D}_{\mathbf{Sc}_{\bar{n}}}^{\text{vrt}}(\lambda)$  consists of  $\mathbf{Sc}_{\bar{n}}$ -dominant compositions associated with the consistent antilinearization  $\text{vrt} : \mathbf{Sc}_{\bar{n}} \hookrightarrow [1, m]$ .

*Proof.* Since each row and each column of the Young diagram contains at most one staircase corner, and as noted in Remark 4.9, we know that  $\text{vrt}$  defines an order-reversing dominant antilinearization of  $\mathbf{Sc}_{\bar{n}}$ , while  $\text{hor}$  defines an order-preserving dominant linearization. On the other hand, each row and each column of the Young diagram contains at most one staircase corner, which implies that a  $DL$ -dense array  $A$  is uniquely determined by its horizontal (or vertical) weight. Furthermore, from the definition, we see that  $A$  is  $DL$ -dense if and only if  $\text{hor}(A)$  is  $\text{hor}(\mathbf{Sc}_{\bar{n}})$ -dominant (respectively,  $\text{vrt}(A)$  is  $\text{vrt}(\mathbf{Sc}_{\bar{n}})$ -dominant).  $\square$

**4.4. Bruhat order on  $\text{DL}_{\bar{n}}(\lambda)$ .** Let  $\mathbf{Y}_{\bar{n}}$  be a given staircase shape, let  $\mathbf{Sc}_{\bar{n}}$  be the corresponding poset of staircase corners (defined in §4.2), and let  $\lambda$  be a partition whose length does not exceed the size of  $\mathbf{Sc}_{\bar{n}}$ . The goal of this section is to define and describe the basic properties of the Bruhat partial order on the subset  $\text{DL}_{\bar{n}}(\lambda)$  of  $DL$ -dense arrays, where the multiset of nonzero elements coincides with  $\lambda$ .

First, recalling Remark 4.6, we can restrict our attention to the case where each row and each column of  $\mathbf{Y}_{\bar{n}}$  contains a staircase corner. This means that  $n_m = m = \#\mathbf{Sc}_{\bar{n}}$ , and we will work under this assumption for the remainder of Section 4.4.

**Definition 4.20.** *A pair of staircase corners  $(ij)$ - $(i'j')$   $\in \mathbf{Sc}_{\bar{n}} \times \mathbf{Sc}_{\bar{n}}$  is called a minimal  $DL$ -disorder for a  $DL$ -dense array  $A$  iff*

- staircase corners  $(ij)$  and  $(i'j')$  are incomparable in  $\mathbf{Sc}_{\bar{n}}$ :

$$(i < i') \text{ and } (j < j'),$$

- $(ij)$ - $(i'j')$  is a disorder:  $A_{ij} > A_{i'j'}$ ,
- the following implications hold for all staircase corners  $(kl) \in \mathbf{Sc}_{\bar{n}}$ :


$$(4.21) \quad \left[ \begin{array}{ll} (k > i) & \& (l < j) \\ (k > i') & \& (l < j') \end{array} \right] \Rightarrow A_{kl} \geq A_{ij},$$

$$(4.22) \quad \left[ \begin{array}{ll} (k < i) & \& (l > j) \\ (k < i') & \& (l > j') \end{array} \right] \Rightarrow A_{kl} \leq A_{i'j'}.$$

$$(4.23) \quad (i < k < i') \& (j < l < j') \Rightarrow A_{kl} \notin [A_{i'j'}, A_{ij}].$$

Let us give a pictorial description of the necessary inequalities for a minimal  $DL$ -disorder:

- A minimal disorder  $(ij)$ - $(i'j')$  is drawn in rose,
- Assumptions (4.21) are drawn in green,
- Assumptions (4.22) are blue,
- Assumption (4.23) is drawn in orange.



**Remark 4.24.** *The conditions that must be verified for  $(ij)$ - $(i'j')$  to be a minimal  $DL$ -disorder are only those represented in dark green ( $k < i$ )& $(j < l < j')$  and dark blue ( $k > i'$ )& $(j < l < j')$ , as well as the inequalities in the orange square ( $i < k < i'$ )& $(j < l < j')$ . The remaining blue and green inequalities follow directly from the  $DL$ -dense property of  $A$ . However, we suggest keeping all the assumptions to maintain the symmetry of both the conditions and the diagram.*

Here is a pair of examples of  $DL$ -disorders for a  $DL$ -dense array from  $DL_{(23^24)}(2^21^2)$ :



$(32)$ - $(44)$  is a minimal  $DL$ -disorder,  $(21)$ - $(44)$  is a non-minimal  $DL$ -disorder.

**Lemma 4.25.** *The following conditions are equivalent for any  $DL$ -dense array  $A$ :*

- $(ij)$ - $(i'j')$  is a minimal  $DL$ -disorder for  $A$ ;
- $(ii')$  is a minimal  $\text{hor}(\mathbf{S}_{\bar{\pi}})$ -disorder for  $\text{hor}(A)$ ;
- $(jj')$  is a minimal  $\text{vrt}(\mathbf{S}_{\bar{\pi}})$ -disorder for  $\text{vrt}(A)$ .

*Proof.* This follows from a direct comparison of inequalities (4.21)–(4.23) and Definition 2.13, which are visualized in the Figure above.  $\square$

**Corollary 4.26.** *For any two  $DL$ -dense arrays  $A$  and  $B$ , the following inequalities are equivalent:*

$$\text{hor}(A) \prec_{\text{Br}} \text{hor}(B) \Leftrightarrow \text{vrt}(A) \prec_{\text{Br}} \text{vrt}(B).$$

*In other words, the horizontal and vertical Bruhat orders on  $DL_{\bar{\pi}}(\lambda)$  coincide.*

*Proof.* Recall that  $\text{vrt}$  defines an isomorphism between  $DL_{\bar{\pi}}(\lambda)$  and  $\mathbb{D}_{\mathbf{S}_{\bar{\pi}}}^{\text{vrt}}(\lambda)$ . The covering relations in the latter poset are given by minimal  $\text{vrt}(\mathbf{S}_{\bar{\pi}})$ -disorders for  $\mathbf{S}_{\bar{\pi}}$ -dominant weights (Theorem 2.18). Similarly, the covering relations for the horizontal weights are also given by minimal  $\text{hor}(\mathbf{S}_{\bar{\pi}})$ -disorders, which are known to be the same covering relations thanks to Lemma 4.25.  $\square$

In particular, it follows that the set  $DL_{\bar{\pi}}(\lambda)$  admits a canonical Bruhat partial order, and the corresponding poset is bounded, graded, subthin, and EL-shellable if  $\lambda$  is regular.

**4.5. Double Demazure Distributive Lattice.** The combinatorial structures defined in the preceding sections allow us to construct a distributive lattice of Demazure submodules. This will later be used in the description of the main subject of this paper – Howe duality for staircase matrices.

**Theorem 4.27.** *For each partition  $\lambda$  of length at most  $\#\mathbf{S}_{\bar{\pi}}$ , the tensor product of  $\mathfrak{b}_n$ - $\mathfrak{b}_m$ -Demazure submodules*

$$\{D_{\text{hor}(A)} \otimes D_{\text{vrt}(B)}^{\text{op}} : A, B \in DL_{\bar{\pi}}(\lambda)\}$$

*in the product of two integrable representations  $V_{\lambda} \otimes V_{\lambda}^{\text{op}}$  forms the set of  $\vee$ -irreducible elements of a distributive lattice, which we denote by  $\mathcal{L}_{D \times D^{\text{op}}}^{\bar{\pi}}(\lambda)$ .*

*Proof.* Thanks to Proposition 4.19, we know that the poset  $DL_{\bar{\pi}}(\lambda)$  is isomorphic to the poset of  $\mathbf{S}_{\bar{\pi}}$ -dominant weights  $\mathbb{D}_{\mathbf{S}_{\bar{\pi}}}^{\text{hor}}(\lambda)$ . Consequently, by Corollary 3.13, the Demazure submodules  $\{D_{\text{hor}(A)} : A \in DL_{\bar{\pi}}(\lambda)\}$  form the set of  $\vee$ -irreducible elements of a distributive lattice  $\mathcal{L}_D^{\mathbf{S}_{\bar{\pi}}}(\lambda)$  of left  $\mathfrak{b}_n$ -submodules in the  $\mathfrak{gl}_n$ -module  $V_{\lambda}$ . Similarly, by Corollary 3.18, the opposite Demazure submodules  $\{D_{\text{vrt}(B)}^{\text{op}} : B \in DL_{\bar{\pi}}(\lambda)\}$  form the set of  $\vee$ -irreducible elements of the distributive lattice  $\mathcal{L}_{D^{\text{op}}}^{\bar{\pi}}(\lambda)$  of right  $\mathfrak{b}_m$ -submodules in the right  $\mathfrak{gl}_m$ -module  $V_{\lambda}^{\text{op}}$ .

A lattice of vector spaces is distributive if and only if it has a common basis. Therefore, if the distributive lattice  $\mathcal{L}$  has a common basis  $E$  and the distributive lattice  $\mathcal{L}'$  has a common basis  $E'$ , then the Cartesian product  $E \times E'$  defines a common basis for the tensor product of lattices  $\mathcal{L} \otimes \mathcal{L}'$ . This ensures that the tensor product remains distributive. Moreover, the tensor product of  $\vee$ -irreducible elements is also  $\vee$ -irreducible.  $\square$

Recall that for each  $\vee$ -irreducible subspace  $D$  in a distributive lattice  $\mathcal{L}$  of vector spaces, one assigns a minimal subquotient  $K_D$  (see (1.31)). By direct inspection, we observe that the minimal subquotients in the tensor product of distributive lattices correspond to the tensor product of minimal subquotients.

Consequently, the minimal subquotients in the lattice  $\mathcal{L}_{D \times D^{\text{op}}}^{\bar{\pi}}(\lambda)$  are indexed by pairs of  $DL$ -dense arrays and are isomorphic to the product of the generalized van der Kallen modules introduced in (3.14):

$$K_{A,B}^{\bar{\pi}} := K_{\mathbf{S}_{\bar{\pi}}, \text{hor}(A)} \otimes K_{\mathbf{S}_{\bar{\pi}}, \text{vrt}(B)}^{\text{op}}.$$

In particular, its  $\mathfrak{h}_n$ - $\mathfrak{h}_m$ -character has the following description based on the formulas (3.17) and (3.19):

$$\begin{aligned} (4.28) \quad \text{ch}_{\mathfrak{h}_n - \mathfrak{h}_m}(K_{A,B}^{\bar{\pi}}) &= \text{ch}_{\mathfrak{h}_n}(K_{\mathbf{S}_{\bar{\pi}}, \text{hor}(A)}) \cdot \text{ch}_{\mathfrak{h}_m}(K_{\mathbf{S}_{\bar{\pi}}, \text{vrt}(B)}^{\text{op}}) = \\ &= \left( \sum_{C \preceq A} \mu^{\text{DL}_{\bar{\pi}}(\lambda)}(C, A) \kappa_{\text{hor}(C)}(x) \right) \left( \sum_{C \succeq B} \mu^{\text{DL}_{\bar{\pi}}(\lambda)}(B, C) \kappa_{\text{vrt}(C)}(y) \right). \end{aligned}$$

Note that the summation in the first term runs over all  $DL$ -dense arrays less than or equal to  $A$ , while the summation in the second term ranges over  $DL$ -dense arrays greater than or equal to  $B$ .

## 5. HOWE DUALITY FOR STAIRCASE MATRICES

This paper is motivated by the combinatorics of various Cauchy-type identities for staircase matrices  $\mathbf{Mat}_{\bar{n}}$  that we introduced in [FKhM3]. All these identities are based on the description of the associated graded components of different filtrations of  $S^N(\mathbf{Mat}_{\bar{n}})$ . These filtrations originate from the highest weight category structures of left  $\mathfrak{b}_{n_m}$ -modules and right  $\mathfrak{b}_m$ -modules. In this paper, we propose considering weaker filtrations, which are still sufficient for computing the characters.

**5.1. The Bi-Module  $\mathbf{Mat}_{\bar{n}}$  of Staircase Matrices.** In this section, we fix a collection of integers  $\bar{n} := (n_1 \leq \dots \leq n_m)$  with  $n_1 > 0$ , and we denote the largest column  $n_m$  by  $n$ . In particular, the Young diagram  $\mathbf{Y}_{\bar{n}}$  is a subdiagram of a rectangular Young diagram  $\mathbf{Y}_{n^m}$ . The space of staircase matrices  $\mathbf{Mat}_{\bar{n}}$  consists of linear functions  $A : \mathbf{Y}_{\bar{n}} \rightarrow \mathbb{k}$ . In other words,  $\mathbf{Mat}_{\bar{n}}$  is the subspace of rectangular  $n \times m$  matrices whose entries vanish outside  $\mathbf{Y}_{\bar{n}}$ .

The space  $\mathbf{Mat}_{\bar{n}}$  is acted upon from the left by the Borel subalgebra  $\mathfrak{b}_n$  and from the right by the Borel subalgebra  $\mathfrak{b}_m$  of upper triangular matrices (via left and right multiplication). These actions commute, yielding a bi-module structure. In what follows, we are particularly interested in the bi-module  $S(\mathbf{Mat}_{\bar{n}})$ , the symmetric algebra of  $\mathbf{Mat}_{\bar{n}}$ .

For each  $(i, j) \in \mathbf{Mat}_{\bar{n}}$ , we denote by  $v_{ij}$  the matrix unit that sends the cell  $(i, j)$  to 1 and all other elements to zero. The set  $\{v_{ij} : (i, j) \in \mathbf{Y}_{\bar{n}}\}$  forms a basis of  $\mathbf{Mat}_{\bar{n}}$ . To each array  $A$  of shape  $\mathbf{Y}_{\bar{n}}$ , we associate the monomial

$$v^A := \prod_{i,j} v_{ij}^{A_{ij}}.$$

The set  $\{v^A\}$ , where  $A$  ranges over all arrays of degree  $N$ , forms a basis of the  $N$ 'th symmetric power  $S^N(\mathbf{Mat}_{\bar{n}})$ .

**Remark 5.1.** *The left action of the matrix unit  $E_{ij} \in \mathfrak{b}_n$  ( $i \leq j \leq n$ ) and the right action of the matrix unit  $E_{ij} \in \mathfrak{b}_m$  on matrix units  $\{v_{ij} : (ij) \in \mathbf{Y}_{\bar{n}}\}$  and arrays can be summarized as follows:*

$$\begin{aligned} E_{ij}v_{ab} &= \delta_{j,a}v_{ib}, & v_{ab}E_{ij} &= \delta_{b,i}v_{aj}; \\ E_{ii}v^A &= (\text{hor}(A))_i v^A, & v^A E_{ii} &= (\text{vrt}(A))_i v^A. \end{aligned}$$

**Definition 5.2.** *The left  $\mathfrak{h}_n$ -weight  $\nu$ -subspace  ${}_{\nu}S(\mathbf{Mat}_{\bar{n}})$  of  $S(\mathbf{Mat}_{\bar{n}})$  is spanned by the elements  $v^A$  such that  $\text{hor}(A) = \nu$ . Similarly, the right  $\mathfrak{h}_m$ -weight  $\bar{d}$ -subspace  $S(\mathbf{Mat}_{\bar{n}})_{\bar{d}}$  of  $S(\mathbf{Mat}_{\bar{n}})$  is spanned by the elements  $v^A$  such that  $\text{vrt}(A) = \bar{d}$ . The notation  ${}_{\nu}S(\mathbf{Mat}_{\bar{n}})_{\bar{d}}$  is used for the left-right weight  $(\nu, \bar{d})$  subspace of  $S(\mathbf{Mat}_{\bar{n}})$ .*

Note that the left  $\mathfrak{b}_n$ -action on the weight subspace  ${}_{\nu}S(\mathbf{Mat}_{\bar{n}})_{\bar{d}}$  increases the weight  $\nu$  while keeping  $\bar{d}$  unchanged, whereas the right  $\mathfrak{b}_m$ -action decreases the weight  $\bar{d}$  while keeping  $\nu$  unchanged.

The following proposition was proved in [FKhM3, Prp.4.3] based on a direct inspection of the  $\mathfrak{b}_n$ - $\mathfrak{b}_m$ -action:

**Proposition 5.3.** *The monomials  $v^A$  where  $A$  belongs to  $\bigcup_{\lambda \vdash N} \text{DL}_{\bar{n}}(\lambda)$ , the set of DL-dense arrays of total weight  $N$ , generate the  $\mathfrak{b}_{n_m}$ - $\mathfrak{b}_m$ -bimodule  $S^N(\mathbf{Mat}_{\bar{n}})$ .*

**Example 5.4.** For a rectangular Young diagram  $\mathbf{Y}_{n^m}$ , the set of staircase corners  $\mathfrak{S}_{n^m}$  forms a totally ordered set of size  $\min(m, n)$  (see Picture (4.8)). Consequently, the set  $\mathbf{DL}_{n^m}(\lambda)$  consists of a single element, which we denote by

$$A_\lambda := \begin{array}{|c|c|c|c|c|c|} \hline & & & & \lambda_5 & \\ \hline & & & \lambda_4 & & \\ \hline & & \lambda_3 & & & \\ \hline & \lambda_2 & & & & \\ \hline \lambda_1 & & & & & \\ \hline \end{array},$$

that has nonzero elements on the secondary diagonal starting from the bottom-left corner.

**5.2. Howe Duality and Cauchy-Type Identities.** The famous Howe duality introduced by R. Howe in [Ho] (see also [GW]) states the following isomorphism of  $\mathfrak{gl}_n$ - $\mathfrak{gl}_m$ -bimodules:

$$(5.5) \quad S^N(\mathbf{Mat}_{n \times m}) \simeq \bigoplus_{\substack{\lambda \vdash N \\ l(\lambda) \leq \min(n, m)}} V_\lambda^{\mathfrak{gl}_n} \otimes (V_\lambda^{\mathfrak{gl}_m})^{\text{op}}.$$

Here,  $V_\lambda^{\mathfrak{gl}_n}$  denotes the integrable  $\mathfrak{gl}_n$ -representation with highest weight  $\lambda$ . We are interested in the action of the Borel subalgebras  $\mathfrak{b}_n$ - $\mathfrak{b}_m$  of upper-triangular matrices rather than the full matrix Lie algebras. In particular, the maximal Demazure  $\mathfrak{b}$ -submodules of the Borel subalgebra coincide with the corresponding irreducible finite-dimensional  $\mathfrak{gl}_n$ -module:

$$V_\lambda^{\mathfrak{gl}_n} \simeq D_{\lambda_-}, \quad (V_\lambda^{\mathfrak{gl}_n})^{\text{op}} \simeq D_{\lambda_+}.$$

Moreover, the action of upper-triangular matrices defines a standard partial order on the set of dominant weights (partitions):

**Definition 5.6.** We say that partitions  $\lambda \geq \nu$  (with  $\lambda, \nu \vdash N$ ) if and only if

$$\forall r = 1, \dots, n \quad \lambda_1 + \dots + \lambda_r \geq \nu_1 + \dots + \nu_r.$$

Although the explicit formulas for the generators of the summands on the right-hand side of (5.5) are somewhat intricate, it is straightforward to identify representatives with respect to the following left ( ${}^\lambda \mathcal{F}$ ) and right ( $\mathcal{F}^\lambda$ ) filtrations indexed by partitions  $\lambda \vdash N$ :

$$(5.7) \quad \begin{aligned} \mathcal{F}^\lambda (S^N(\mathbf{Mat}_{n \times m})) &:= \left( \bigoplus_{\nu \geq \lambda} S^N(\mathbf{Mat}_{n \times m})_\nu \right) \mathbf{U}(\mathfrak{b}_m); \\ {}^\lambda \mathcal{F} (S^N(\mathbf{Mat}_{n \times m})) &:= \mathbf{U}(\mathfrak{b}_n) \left( \bigoplus_{\nu \geq \lambda} {}^\nu S^N(\mathbf{Mat}_{n \times m}) \right). \end{aligned}$$

In particular, isomorphism (5.5) can be weakened to the following isomorphisms:

$$\mathcal{F}^\lambda \left/ \sum_{\nu \geq \lambda} \mathcal{F}^\nu (S^N(\mathbf{Mat}_{n \times m})) \right. \simeq {}^\lambda \mathcal{F} \left/ \sum_{\nu \geq \lambda} {}^\nu \mathcal{F} (S^N(\mathbf{Mat}_{n \times m})) \right. \simeq D_{\lambda_-} \otimes D_{\lambda_+}^{\text{op}}.$$

Moreover, both quotients are cyclic  $\mathfrak{b}_n$ - $\mathfrak{b}_m$ -modules generated by the monomial

$$v^{A_\lambda} := v_{n,1}^{\lambda_1} \cdot v_{n-1,2}^{\lambda_2} \cdot v_{n-3,3}^{\lambda_3} \cdots$$

where  $A_\lambda$  is the unique  $DL$ -dense array of rectangular shape  $n^m$  whose multiset of elements is equal to  $\lambda$ . (See Example 5.4 for details).

The embedding of the space of staircase matrices into the space of rectangular matrices induces an embedding of the corresponding symmetric tensors, and the symmetric algebra inherits the left and right filtrations by weights:

$$\begin{aligned} \iota_{\bar{n}} : \mathbf{Mat}_{\bar{n}} &\hookrightarrow \mathbf{Mat}_{n \times m}; & \iota_{\bar{n}} : S^N(\mathbf{Mat}_{\bar{n}}) &\hookrightarrow S^N(\mathbf{Mat}_{n \times m}); \\ \mathcal{F}^\lambda S^N(\mathbf{Mat}_{\bar{n}}) &:= S^N(\mathbf{Mat}_{\bar{n}}) \cap \mathcal{F}^\lambda S^N(\mathbf{Mat}_{n \times m}). \end{aligned}$$

It follows that for each partition  $\lambda \vdash N$  with  $l(\lambda) \leq \min(n, m)$ , we have the following quotient map of  $\mathfrak{b}_n$ - $\mathfrak{b}_m$ -bimodules:

$$(5.8) \quad \bar{\iota}_{\bar{n}}^\lambda : \mathcal{F}^\lambda \left/ \sum_{\mu > \lambda} \mathcal{F}^\mu (S^N(\mathbf{Mat}_{\bar{n}})) \right. \hookrightarrow \mathcal{F}^\lambda \left/ \sum_{\mu > \lambda} \mathcal{F}^\mu (S^N(\mathbf{Mat}_{n \times m})) \right. \simeq V_\lambda^{\mathfrak{gl}_n} \otimes (V_\lambda^{\mathfrak{gl}_m})^{\text{op}}.$$

and the analogous quotient map for the left filtration.

**Proposition 5.9.** *For each  $DL$ -dense array  $A \in \mathbf{DL}_{\bar{n}}(\lambda)$ , the monomial  $v^A$  belongs to  $\mathcal{F}^\lambda S^N(\mathbf{Mat}_{n \times m})$  (respectively to  ${}^\lambda \mathcal{F}(S^N(\mathbf{Mat}_{n \times m}))$ ). Moreover, the  $\mathfrak{b}_n$ - $\mathfrak{b}_m$ -submodule of  $D_{\lambda_-} \otimes D_{\lambda_+}^{\text{op}}$  generated by the vector  $\bar{\iota}_{\bar{n}}^\lambda(v^A)$  is isomorphic to the Demazure submodule:*

$$(5.10) \quad D_{\text{hor}(A)} \otimes D_{\text{vrt}(A)}^{\text{op}}.$$

*Proof.* The module  $S^N(\mathbf{Mat}_{n \times m})$ , as well as its submodule  $V_\lambda \otimes V_\lambda^{\text{op}}$ , is a  $\mathfrak{gl}_n$ - $\mathfrak{gl}_m$ -module and consequently admits a linear action of the product of two Weyl groups, i.e., the symmetric groups  $\mathbb{S}_n \times \mathbb{S}_m$ . Moreover, this action maps the highest weight vector to an extremal vector in an irreducible representation.

Since  $A_\lambda \in \mathbf{DL}_{n \times m}(\lambda)$  is the unique  $DL$ -dense array of rectangular shape, the monomial  $v^{A_\lambda}$  represents the highest weight vector in  $\mathcal{F}^\lambda / \mathcal{F}^{> \lambda}(S^N(\mathbf{Mat}_{n \times m}))$  (see Example 5.4). The monomial  $v^B$  represents an extremal vector in this subquotient whenever the array  $B$  belongs to the  $\mathbb{S}_n \times \mathbb{S}_m$ -orbit of  $A_\lambda$ . Since  $A$  is  $DL$ -dense, it belongs to this orbit, implying that  $\iota_{\bar{n}}(v^A)$  is an extremal vector of  $V_\lambda \otimes V_\lambda^{\text{op}}$ . Consequently, the  $\mathfrak{b}_n$ - $\mathfrak{b}_m$ -submodule generated by it is the Demazure submodule  $D_{\text{hor}(A)} \otimes D_{\text{vrt}(A)}^{\text{op}}$ .  $\square$

Note that the Demazure submodules (5.10) often intersect inside  $V_\lambda \otimes V_\lambda^{\text{op}}$ . For example,

$$(5.11) \quad A \preceq_{\text{Br}} B \in \mathbf{DL}_{\bar{n}}(\lambda) \Leftrightarrow \begin{cases} D_{\text{hor}(A)} \subset D_{\text{hor}(B)}, \\ D_{\text{vrt}(A)}^{\text{op}} \supset D_{\text{vrt}(B)}^{\text{op}} \end{cases} \Rightarrow \\ \Rightarrow \left( D_{\text{hor}(A)} \otimes D_{\text{vrt}(A)}^{\text{op}} \right) \cap \left( D_{\text{hor}(B)} \otimes D_{\text{vrt}(B)}^{\text{op}} \right) = D_{\text{hor}(A)} \otimes D_{\text{vrt}(B)}^{\text{op}}.$$

However, thanks to Theorem 4.27, we know that  $D_{\text{hor}(A)} \otimes D_{\text{vrt}(B)}^{\text{op}}$  are  $\vee$ -irreducible elements of the distributive lattice  $\mathcal{L}_{D \times D^{\text{op}}}^{\bar{n}}(\lambda)$ .

**Theorem 5.12.** *For each partition  $\lambda \vdash N$  with  $l(\lambda) \leq \#\mathfrak{S}_{\bar{n}}$ , the associated graded component of the left (and right) filtrations (5.7) is isomorphic to the following sum of  $\mathfrak{b}_n$ - $\mathfrak{b}_m$ -Demazure (bi)-submodules:*

$$(5.13) \quad \text{gr} \mathcal{F}^\lambda := \mathcal{F}^\lambda \left/ \sum_{\nu > \lambda} \mathcal{F}^\nu (S^N(\mathbf{Mat}_{\bar{n}})) \right. \simeq \sum_{A \in \mathbf{DL}_{\bar{n}}(\lambda)} D_{\text{hor}(A)} \otimes D_{\text{vrt}(A)}^{\text{op}} \subset V_\lambda^{\mathfrak{gl}_n} \otimes (V_\lambda^{\mathfrak{gl}_m})^{\text{op}}.$$



Moreover, the corresponding  $\mathfrak{b}_n$ - $\mathfrak{b}_m$ -sub-bimodule is an element of the distributive lattice  $\mathcal{L}_{D \times D^{\text{op}}}^{\bar{\pi}}(\lambda)$  whose associated graded module with respect to the standard filtration  $\mathcal{F}_{\mathcal{L}}$ , defined in (1.29), is isomorphic to the following sum:

$$(5.14) \quad \bigoplus_{B \preceq C \in \text{DL}_{\bar{\pi}}(\lambda)} K_{\mathfrak{S}_{\bar{\pi}}, \text{hor}(B)} \otimes K_{\mathfrak{S}_{\bar{\pi}}, \text{vrt}(C)}^{\text{op}}.$$

*Proof.* Thanks to Proposition 5.9, we know that the elements  $v^A$  for  $A \in \text{DL}_{\bar{\pi}}(\lambda)$  belong to  $\mathcal{F}^\lambda$  and generate the summands on the right-hand side of (5.13). Consequently, the surjectivity of (5.13) follows. On the other hand, this also implies that  $v^A$  does not belong to  $\mathcal{F}^\nu$  for any  $\nu > \lambda$ . After Proposition 5.3, we know that monomials assigned to  $DL$ -dense arrays generate the symmetric algebra. It follows that  $\{v^A : A \in \text{DL}_{\bar{\pi}}(\lambda)\}$  generate the subquotient  $\mathcal{F}^\lambda / \mathcal{F}^{>\lambda}$ , proving the injectivity of (5.13).

To describe the set of minimal subquotients  $K_{A,B}$  appearing in the filtration  $\mathcal{F}_{\mathcal{L}}$ , it remains to identify the set of  $\vee$ -irreducible elements of the lattice  $\mathcal{L}_{D \times D^{\text{op}}}^{\bar{\pi}}(\lambda)$  that belong to the right-hand side of (5.13). As mentioned in (5.11), the  $\vee$ -irreducible subspace of  $D_{\bar{a}}$  is  $D_{\bar{b}}$  with  $b \preceq a$ , and on the other hand,  $D_{\bar{c}}^{\text{op}}$  is a subspace of  $D_{\bar{a}}^{\text{op}}$  if and only if  $\bar{c} \succeq \bar{a}$ . Thus, the  $\vee$ -irreducible subspaces of  $D_{\text{hor}(A)} \otimes D_{\text{vrt}(A)}^{\text{op}}$  consist of the products

$$D_{\text{hor}B} \otimes D_{\text{vrt}(C)}^{\text{op}} \text{ with } (\text{hor}(B) \preceq \text{hor}(A)) \ \& \ (\text{vrt}(C) \succeq \text{vrt}(A)) \Leftrightarrow B \preceq A \preceq C.$$

Since, in the right-hand side of (5.13), we sum over all possible  $A$ , we can omit the intermediate array  $A$  and conclude that products  $D_{\text{hor}B} \otimes D_{\text{vrt}(C)}^{\text{op}}$  form the complete set of  $\vee$ -irreducible elements of  $\text{gr}\mathcal{F}^\lambda$ . The indexing set of the set of minimal subquotients of  $\text{gr}\mathcal{F}^\lambda$  coincides with the set of  $\vee$ -irreducible submodules in it, what ensures the summation (5.14).  $\square$

We now compute the  $\mathfrak{h}_n$ - $\mathfrak{h}_m$ -character of  $\text{gr}\mathcal{F}^\lambda$  based on Theorem 5.12: Let us compute the  $\mathfrak{h}_n$ - $\mathfrak{h}_m$ -character of  $\text{gr}\mathcal{F}^\lambda$  based on Theorem 5.12:

$$(5.15) \quad \begin{aligned} \text{ch}_{\mathfrak{h}_n, \mathfrak{h}_m}(\text{gr}\mathcal{F}^\lambda) &\stackrel{(5.14)}{=} \sum_{B \preceq A \in \text{DL}_{\bar{\pi}}(\lambda)} \text{ch}_{\mathfrak{h}_n}(K_{\mathfrak{S}_{\bar{\pi}}, \text{hor}(B)}) \text{ch}_{\mathfrak{h}_m}(K_{\mathfrak{S}_{\bar{\pi}}, \text{vrt}(A)}^{\text{op}}) = \\ &= \sum_{A \in \text{DL}_{\bar{\pi}}(\lambda)} \left( \sum_{B \preceq A \in \text{DL}_{\bar{\pi}}(\lambda)} \text{ch}_{\mathfrak{h}_n}(K_{\mathfrak{S}_{\bar{\pi}}, \text{hor}(B)}) \right) \text{ch}_{\mathfrak{h}_m}(K_{\mathfrak{S}_{\bar{\pi}}, \text{vrt}(A)}^{\text{op}}) = \\ &= \sum_{A \in \text{DL}_{\bar{\pi}}(\lambda)} \text{ch}_{\mathfrak{h}_n}(D_{\text{hor}(A)}) \text{ch}_{\mathfrak{h}_m}(K_{\mathfrak{S}_{\bar{\pi}}, \text{vrt}(A)}^{\text{op}}) \stackrel{(3.19)}{=} \sum_{A \in \text{DL}_{\bar{\pi}}(\lambda)} \kappa_{\text{hor}(A)}(x) \left( \sum_{\substack{\bar{d} \in \mathbb{S}_n \lambda \\ \text{bs}_{\bar{\pi}}(\bar{d}) = \text{vrt}(A)}} a^{\bar{d}}(y) \right) = \\ &\stackrel{(3.19)}{=} \sum_{A \in \text{DL}_{\bar{\pi}}(\lambda)} \kappa_{\text{hor}(A)} \left( \sum_{B \succeq A} \mu^{\text{DL}_{\bar{\pi}}(\lambda)}(A, B) \kappa^{\text{vrt}(B)}(y) \right) = \sum_{A \preceq B \in \text{DL}_{\bar{\pi}}(\lambda)} \mu^{\text{DL}_{\bar{\pi}}(\lambda)}(A, B) \kappa_{\text{hor}(A)}(x) \kappa^{\text{vrt}(B)}(y). \end{aligned}$$

Finally, we reproduce the main identities from [FKhM3], known as "Cauchy identities for staircase matrices":

**Corollary 5.16.** *The following identities hold for any Young diagram  $\mathbf{Y}_{\bar{n}}$ :*

$$(5.17) \quad \prod_{(i,j) \in \mathbf{Y}_{\bar{n}}} \frac{1}{1 - x_i y_j} = \sum_{A \in \text{DL}_{\bar{n}}} \kappa_{\text{hor}(A)}(x) \cdot \left( \sum_{\bar{d}: \text{bs}_{\bar{n}}(\bar{d}) = \text{vrt}(A)} a^{\bar{d}}(y) \right),$$

$$(5.18) \quad \prod_{(i,j) \in \mathbf{Y}_{\bar{n}}} \frac{1}{1 - x_i y_j} = \sum_N \sum_{\substack{\lambda \vdash N \\ l(\lambda) \leq \#\mathbf{s}_{\bar{n}}}} \sum_{\substack{A \geq B \\ A, B \in \text{DL}_{\bar{n}}(\lambda)}} \mu^{\text{DL}_{\bar{n}}(\lambda)}(A, B) \kappa_{\text{hor}(A)}(x) \kappa^{\text{vrt}(B)}(y).$$

*Proof.* The left-hand side of these identities represents the  $\mathfrak{h}_n$ - $\mathfrak{h}_m$ -character of the symmetric algebra  $S^*(\text{Mat}_{\bar{n}})$ . In contrast, the right-hand side follows from the two different presentations of the character of the associated graded factors  $\text{gr}\mathcal{F}^\lambda$  described in (5.15).  $\square$

Note that in this paper, we use the bubble-sort operation  $\text{bs}_{\bar{n}}$  instead of its variant  $\text{hbs}$ , referred to as half-bubble-sort in our previous work [FKhM3], which is somewhat more cumbersome.

## REFERENCES

- [Al] P. Alexandersson, *The symmetric functions catalog*, <https://www.symmetricfunctions.com>.
- [AQ1] S. Assaf, D. Quijada, *A Pieri rule for key polynomials*, *Sém. Lothar. Combin.* 80B (2018), Article # 78, 12 pp.
- [AQ2] S. Assaf, D. Quijada, *A Pieri rule for Demazure characters of the general linear group*, <https://arxiv.org/abs/1908.08502>.
- [AE] O. Azenhas, A. Emami, *An analogue of the Robinson-Schensted-Knuth correspondence and non-symmetric Cauchy kernels for truncated staircases*, *European J. Combin.* 46 (2015), 16–44.
- [AGL] O. Azenhas, T. Gobet, C. Lecouvey, *Non-symmetric Cauchy kernel, crystals and last passage percolation*, *Tunisian J. Math.* 6(2) (2024), 249–297.
- [BW] D. Betea, M. Wheeler, *Refined Cauchy and Littlewood identities, plane partitions and symmetry classes of alternating sign matrices*, *J. Comb. Theory Ser. A*, 137 (2016), 126–165.
- [Bi] G. Birkhoff, *Rings of sets*, *Duke Math. J.* 3(3) (1937), 443–454.
- [B1] A. Björner, *Posets, regular CW complexes and Bruhat order*, *European J. Combin.* 5(1) (1984), 7–16.
- [BB] A. Björner, F. Brenti, *Combinatorics of Coxeter groups*, *Grad. Texts in Math.*, Vol. 231, Springer, New York, 2005.
- [BW1] A. Björner, M. Wachs, *Bruhat order of Coxeter groups and shellability*, *Adv. Math.* 43(1) (1982), 87–100.
- [BW2] A. Björner, M. Wachs, *Generalized quotients in Coxeter groups*, *Trans. Amer. Math. Soc.* 308(1) (1988), 1–37.
- [BC] A. Borodin, I. Corwin, *Macdonald processes*, *Probab. Theory Relat. Fields*, 158(1–2) (2014), 225–400.
- [BP] A. Borodin, L. Petrov, *Integrable probability: From representation theory to Macdonald processes*, *Probab. Surv.* 11 (2014), 1–58.
- [BrKu] M. Brion, S. Kumar, *Frobenius Splittings: Methods and Applications*, *Progress in Math.*, Vol. 231, Birkhäuser, Boston, 2005.
- [CK] S.-I. Choi, J.-H. Kwon, *Lakshmibai-Seshadri paths and non-symmetric Cauchy identity*, *Algebr. Represent. Theory* 21(6) (2018), 1381–1394.
- [DK1] V. Danilov, G. Koshevoy, *Bi-crystals and crystal  $(GL(V), GL(W))$ -duality*, Kyoto University, RIMS, 2004.
- [DK2] V. Danilov, G. Koshevoy, *Arrays and the combinatorics of Young tableaux*, *Russian Math. Surveys* 60(2) (2005), 269.
- [DK] G. Danaraj, V. Klee, *"Shellings of spheres and polytopes."* *Duke Math. J.* 41 (1974), 443–451
- [DP] B. Davey, H. Priestley, *Introduction to Lattices and Order*, 2nd ed., Cambridge Univ. Press, 2002.
- [Dem1] M. Demazure, *Désingularisation des variétés de Schubert généralisées*, *Ann. Sci. École Norm. Sup.* (4) 7 (1974), 53–88.

- [Dem2] M. Demazure, *Une nouvelle formule des caractères*, Bull. Sci. Math. 98(3) (1974), 163–172.
- [FKhM3] E. Feigin, A. Khoroshkin, Ie. Makedonskyi, *Cauchy identities for staircase matrices*, <https://arxiv.org/abs/2411.03117>.
- [FL] A. M. Fu, A. Lascoux, *Non-symmetric Cauchy kernels for the classical groups*, J. Comb. Theory Ser. A, 116 (2009), 903–917.
- [GW] R. Goodman, N. R. Wallach, *Representations and Invariants of the Classical Groups*, Cambridge University Press, 2000.
- [G] R. L. Graham, *Handbook of Combinatorics*, Elsevier, 1995.
- [Ho] R. Howe, *Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond*, The Schur Lectures (1992) (Tel Aviv), 1995, 1–82.
- [Hu] J. Humphreys, *Reflection Groups and Coxeter Groups*, No. 29, Cambridge University Press, 1992.
- [J] A. Joseph, *On the Demazure character formula*, Ann. Sci. École Norm. Sup. (4) 18(3) (1985), 389–419.
- [vdK] W. van der Kallen, *Longest weight vectors and excellent filtrations*, Math. Z. 201(1) (1989), 19–31.
- [Ka] M. Kashiwara, *The crystal base and Littelmann’s refined Demazure character formula*, Duke Math. J. 71(3) (1993), 839–858.
- [Ko] M. Kobayashi, *Construction of double coset system of a Coxeter group and its applications to Bruhat graphs*, <https://arxiv.org/abs/1907.11801>.
- [Ku] S. Kumar, *Kac-Moody Groups, Their Flag Varieties and Representation Theory*, Vol. 204, Springer Science & Business Media, 2012.
- [Las] A. Lascoux, *Double crystal graphs*, In Studies in Memory of Issai Schur, Birkhäuser Boston, 2003.
- [Mas] S. K. Mason, *An explicit construction of type A Demazure atoms*, J. Algebraic Combin. 29(3) (2009), 295–313.
- [Ok] A. Okounkov, *Infinite wedge and random partitions*, Selecta Math. (N.S.), 7(1) (2001), 57–81.
- [OR] A. Okounkov, N. Reshetikhin, *Correlation function of Schur process with application to local geometry of a random 3-dimensional Young diagram*, J. Amer. Math. Soc. 16(3) (2003), 581–603.
- [PP] A. Polishchuk, L. Positselski, *Quadratic Algebras*, Vol. 37, American Mathematical Society, 2005.
- [P] A. Pun, *On decomposition of the product of Demazure atoms and Demazure characters*, University of Pennsylvania, 2016.
- [Q] D. Quillen, *Higher algebraic K-theory: I*, In Higher K-Theories: Proceedings of the Conference held at the Seattle Research Center of the Battelle Memorial Institute, August 28 to September 8, 1972, Springer, Berlin, Heidelberg, 2006, 85–147.
- [St] R. P. Stanley, *Weyl groups, the hard Lefschetz theorem, and the Sperner property*, SIAM J. Algebraic Discrete Methods 1(2) (1980), 168–184.
- [St1] R. P. Stanley, *Enumerative Combinatorics: Volume 2*, Cambridge University Press, First edition, 2001.
- [Stm] J. R. Stembridge, *Tight quotients and double quotients in the Bruhat order*, Electron. J. Combin. (2004), Article R14.
- [W] M. Wachs, *Poset topology: tools and applications*, <https://arxiv.org/abs/0602226>.
- [Wl] J. W. Walker, *Homotopy type and Euler characteristic of partially ordered sets*, European J. Combin. 2(4) (1981), 373–384.

## APPENDIX A. PICTORIAL EXAMPLES

We present several illustrative examples of:

- Arborescent posets  $\mathbf{S}$  with a consistent (anti)linearization.
- The Hasse diagram of the poset  $\mathbb{D}_{\mathbf{S}}(\lambda)$  ordered by the Bruhat partial order.
- The Hasse diagram of the poset  $\text{DL}_{\overline{\pi}}(\lambda)$  of  $DL$ -dense arrays, along with its vertical and horizontal embeddings into the Bruhat graph.

Note that all our pictorial descriptions of the posets are slightly reversed compared to standard notation. Specifically, we draw the minimal element at the top and the maximal element at the bottom. However, the corresponding Bruhat (sub)graphs are presented in the standard orientation. This reversal occurs because we consider the dual order to the Bruhat order, based on comparisons of Demazure modules.

**Example A.1.** Here is a list of arborescent posets of cardinality 3 equipped with a surjective, consistent anti-linearization:

$$\mathbf{S}^1 = \boxed{\begin{array}{c} \curvearrowright \\ \bullet \\ \bullet \\ \bullet \end{array}}, \quad \mathbf{S}^2 = \boxed{\begin{array}{c} \curvearrowright \\ \bullet \\ \bullet \\ \bullet \end{array}}, \quad \mathbf{S}^3 = \boxed{\begin{array}{c} \bullet \\ \bullet \\ \bullet \end{array}}, \quad \mathbf{S}^4 = \boxed{\begin{array}{c} \bullet \\ \bullet \\ \bullet \end{array}}, \quad \mathbf{S}^5 = \boxed{\begin{array}{c} \bullet \\ \bullet \\ \bullet \end{array}},$$


$$\mathbb{D}_{\mathbf{S}^1} \simeq \mathbf{S}_2, \quad \mathbb{D}_{\mathbf{S}^2} \simeq \mathbf{S}_1, \quad \mathbb{D}_{\mathbf{S}^3} \simeq \mathbf{S}_2 \setminus \mathbf{S}_3, \quad \mathbb{D}_{\mathbf{S}^4} \simeq \mathbf{S}_2 \setminus \mathbf{S}_3, \quad \mathbb{D}_{\mathbf{S}^5} \simeq \mathbf{S}_3.$$

Note that if the Hasse diagram of an arborescent poset  $\mathbf{S} = \{s_1, \dots, s_k\}$  with a consistent antilinearization  $v : \mathbf{S} \xrightarrow{s_i \mapsto i} [1, k]$  is connected, then  $s_k$  is the unique minimal element of  $\mathbf{S}$ , and we obtain the following isomorphism of posets of dominant compositions:


$$\mathbb{D}_{\mathbf{S}}^v((\lambda_1 \geq \dots \geq \lambda_k)) \simeq \mathbb{D}_{\mathbf{S}'}^v((\lambda_1 \geq \dots \geq \lambda_{k-1})), \quad \text{where } \mathbf{S}' := \{s_1, \dots, s_{k-1}\}.$$

**Example A.2.** There exists a unique arborescent poset  $\mathcal{S}$  of cardinality 4 whose Hasse diagram has multiple connected components and is not a disjoint union of linearly ordered sets, which are covered by Example 2.11.


The corresponding poset is the set of staircase corners for the shapes  $\mathbf{Y}_{(23^24)}$  and  $\mathbf{Y}_{(13^24)}$ . These two shapes differ by a transposition, leading to two distinct antilinearizations that can be visualized as vertical map  $\mathbf{vrt}$  and opposite to the horizontal map  $\mathbf{hor}$ .




These two different anti-linearizations appear as the vertical map  $\mathbf{vrt}$  and the opposite to the horizontal map  $\mathbf{hor}$  of the same shape  $\mathbf{Y}_{(23^24)}$ . Below we draw the Bruhat graph for this set  $\mathbf{DL}_{(23^24)}$  together with the vertical embedding (the elements in the image are filled in orange).



The next picture shows the horizontal embedding of the same poset  $DL_{(2,3,3,4)}$ :




Here is an example of the Hasse diagram of the smaller poset  $DL_{\bar{\pi}}(\lambda)$  of the same shape  $\bar{\pi} := (2, 3, 3, 4)$  but the partition  $\lambda = (2, 2, 1, 1)$  has equal elements. So the poset is embedded in the parabolic Bruhat graph:



**Example A.4.** Let us now present the next nontrivial example of an arborescent poset with cardinality 5. We consider the set of staircase corners  $\mathbf{Sc}_{\bar{\pi}}$  corresponding to the partition  $\bar{\pi} := (23^25^2)$ .

Below, we illustrate the Hasse diagram of  $\mathbf{Sc}_{\bar{\pi}}$  along with the antilinearization  $vrt$ , as well as the Hasse diagram of the poset  $DL_{\bar{\pi}}$ .



ANTON KHOROSHKIN:  
DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, MOUNT CARMEL, 3498838, HAIFA, ISRAEL  
*Email address:* khoroshkin@gmail.com

IEVGEN MAKEDONSKYI:  
YANQI LAKE BEIJING INSTITUTE OF MATHEMATICAL SCIENCES AND APPLICATIONS (BIMSA), No. 544,  
HEFANGKOU VILLAGE, HUAIBEI TOWN, HUIROU DISTRICT, BEIJING 101408.  
*Email address:* makedonskii\_e@mail.ru