HALFSPACE REPRESENTATIONS OF PATH POLYTOPES OF TREES

AMER GOEL, AIDA MARAJ, AND ÁLVARO RIBOT

ABSTRACT. Given a tree T, its path polytope is the convex hull of the edge indicator vectors for the paths between any two distinct leaves in T. These polytopes arise naturally in polyhedral geometry and applications, such as phylogenetics, tropical geometry, and algebraic statistics. We provide a minimal halfspace representation of these polytopes. The construction is made inductively using toric fiber products.

1. INTRODUCTION

A polytope is given either by its vertex representation (\mathcal{V} -representation) or halfspace representation (\mathcal{H} -representation). The \mathcal{V} -representation describes a polytope as the convex hull of its vertices and the \mathcal{H} -representation defines a polytope as the intersection of halfspaces. Therefore, a \mathcal{V} -representation is a parametric description of the polytope, and an \mathcal{H} -representation is an implicit description of the polytope. We study *path polytopes* of trees, which are defined by their \mathcal{V} -representation as follows: given two distinct leaves i, j in a tree T with vertex set V and edge set E, let $i \leftrightarrow j$ be the set of edges in the path in T between i and j, and let $\mathbf{c}^{i\leftrightarrow j} \in \{0,1\}^{|E|}$ be the indicator vector for the edges used in $i \leftrightarrow j$. The path polytope of the tree T, denoted P_T , is the convex hull of the vectors $\mathbf{c}^{i\leftrightarrow j}$ for any two distinct leaves i, j in T. Our goal is to find its \mathcal{H} -representation, which provides a membership test for P_T . The Fourier-Motzkin elimination algorithm [10] outputs the \mathcal{H} -representation given the vertices of the polytope, but its time complexity grows exponentially with the dimension of the polytope. Therefore, computing the \mathcal{H} -representation of path polytopes with this algorithm is infeasible for large trees. In Theorem 1.1 we provide an explicit description of the \mathcal{H} -representation of any path polytope.

Path polytopes arrive naturally in graph theory, combinatorics, and applications. In tropical geometry, phylogenetic trees are parametrized by the path in [11], related to the tropical Grassmannian [14]. A key motivation of this work is the growing recognition that many statistical models defined on trees or graphs are parametrized by paths between their nodes. Examples include Brownian motion tree models, where the parametrization is given by paths among leaves in a phylogenetic tree [1]; staged tree models, parametrized by the paths from the root to a leaf on a rooted tree [6]; and (colored) Gaussian graphical models on block graphs [2, 13], parametrized by paths between any two nodes, among others.

An explicit description of the halfspaces for P_T in terms of the tree structure provides a better understanding of the polytope and is useful. For instance, polytopes arising from monomial parametrizations of log-linear models, such as our polytope induced by the path parametrization, have shown to be useful in Maximum Likelihood Estimation (MLE) problems [7, 8]; given a normalized vector of counts for a log-linear model, the MLE exists if and only if this vector belongs to the relative interior of the corresponding polytope. Here, the halfspace description of the polytope gives a membership test. In [9], the authors use halfspace representations to learn causal polytree structures from a combination of observational and interventional data. In fact, the path parametrization has already shown essential for all the progress related to the MLE of Brownian motion tree models [1, 3]. Therefore, we anticipate that the halfspace representation of our path polytope will be valuable for statistical applications in the models discussed earlier.

Before presenting our main result, let us introduce the necessary notation. A graph is a tuple G = (V, E) where V is a set of nodes and E is a set of unordered pairs of nodes, which are called edges. We assume $1 < |V| < \infty$. A path in a graph is a sequence of edges that connects two nodes. A tree is a graph in which any pair of nodes is connected by exactly one path. Given a vertex u in a tree T = (V, E), let $N(u) = \{v \in V \mid \{u, v\} \in E\}$ be the neighborhood of u. The degree of a vertex u in T is $\deg(u) = |N(u)|$. Leaves are nodes of degree one, so connected to exactly one other node. Let $Lv(T) \subset V$ denote the set of leaves of T, and $Int(T) = V \setminus Lv(T)$ denote the set of internal nodes in T. A star tree is a tree T with |Int(T)| = 1. See Figure 1.



FIGURE 1. Example of a graph, a tree, a path (dashed edges), and a star tree.

Let $E_{\text{leaf}}(T) = \{\{u, v\} \in E \mid u \in \text{Lv}(T) \text{ or } v \in \text{Lv}(T)\}$ be the set of edges that have a leaf as an endpoint. Let \mathbb{R}^E be the vector space with basis elements indexed by the set E, so $P_T \subset \mathbb{R}^E$. Our main result is the following.

Theorem 1.1. Given a tree T = (V, E) with |V| > 3 and no internal nodes of degree 2, the path polytope P_T has dimension |E| - 1 and a minimal \mathcal{H} -representation is given by

(1)
$$\begin{cases} x_{\{u,v\}} & \geq 0 \quad \text{for all } \{u,v\} \in E, \text{with } \deg(u), \deg(v) \neq 3 \\ -x_{\{u,v\}} + \sum_{w \in N(u) \setminus \{v\}} x_{\{u,w\}} & \geq 0 \quad \text{for all } u \in \operatorname{Int}(T) \text{ and all } v \in N(u) \\ \sum_{\{u,v\} \in E_{\operatorname{leaf}}(T)} x_{\{u,v\}} & = 2. \end{cases}$$

Remark 1.2. The condition that T has no internal nodes of degree 2 is without loss of generality: we show that one can collapse all nodes of degree 2 to and get a new tree such that their path polytopes are isomorphic. By minimal \mathcal{H} -representation we mean that no halfspace is redundant.

Remark 1.3. The internal nodes of degree 3 are special for the following reason. We show that the polytope P_T can be expressed as toric fiber products of pyramids over second hypersimplices. The second hypersimplex $\Delta_{n,2}$ has 2n facets for n > 3 while $\Delta_{3,2}$ has only 3 facets. As such, internal nodes of degree 3 (which correspond to $\Delta_{3,2}$) induce fewer facets of P_T , and therefore fewer halfspaces.

First, we check that every vertex in P_T satisfies the conditions implied by the \mathcal{H} -representation given in Theorem 1.1, so P_T is included in the polytope defined by that \mathcal{H} -representation. Then we prove that these halfspaces are sufficient to define P_T , and that none of these halfspaces is redundant. To do so, we express P_T as the toric fiber product of polytopes coming from subtrees of T, allowing us to compute its facets inductively. **Structure of the paper.** In Section 2 we review literature on polytopes and toric fiber products that is relevant for this paper. In Section 3 we show path polytopes are inductively constructed as toric fiber products of pyramids over path polytopes on star trees. Then, in Section 4, we describe the facets and their respective halfspaces of path polytopes of trees, concluding the proof of Theorem 1.1.

2. PRELIMINARIES

This section consists of three parts. First, we explain how a tree can be decomposed into a gluing of star trees, following the ideas in [1]. Second, we recall the basic notions related to polytopes and define path polytopes of trees. Third, we recall toric fiber products of polytopes.

2.1. Gluing of trees. Given two trees $T_1 = (V_1, E_1)$, $T_2 = (V_2, E_2)$, and two edges $e_1 \in E_{\text{leaf}}(T_1)$ and $e_2 \in E_{\text{leaf}}(T_2)$, the *gluing* of T_1 and T_2 along e_1, e_2 is the new tree $T = T_1 *_{e_1, e_2} T_2$ obtained as the union of T_1 and T_2 by identifying $e_1 \sim e_2$. That is, if $e_i = \{v_i, k_i\}$ with $k_i \in \text{Lv}(T_i)$ for i = 1, 2, then $V(T) = (V_1 \sqcup V_2) \setminus \{k_1, k_2\}$ and $E(T) = (E_1 \cup E_2 \cup \{v_1, v_2\}) \setminus \{e_1, e_2\}$. See Figure 2 for an example.



FIGURE 2. Gluing T_1 and T_2 along edges $\{1, 4\}$ and $\{8, 5\}$ to form T.

Lemma 2.1. Let T = (V, E) be a tree on k internal nodes, with respective degrees $n_1, \ldots, n_k > 2$. Then T can be expressed as a gluing of k star trees S_{n_1}, \ldots, S_{n_k} .

Proof. We argue by strong induction on k. If k = 1 the statement is trivial, since a tree with one internal node is a star tree. Let k > 1 and consider two internal nodes $u_1, u_2 \in V$ such that $\{u_1, u_2\} \in E$. Let T_1 be the subtree of T consisting of all nodes v in T such that the path between v and u_1 does not pass through u_2 , and define T_2 analogously. For i = 1, 2, let $T'_i = (V'_i, E'_i)$ where $V_i = V(T_i) \sqcup \{v_i\}$ and $E'_i = E(T_i) \sqcup \{\{u_i, v_i\}\}$, where v_i is a newly introduced leaf of T_i that connects to the internal node u_i . Then $T = T'_1 *_{\{u_1, v_1\}, \{u_2, v_2\}} T'_2$ and $Int(T'_1), Int(T_2) < k$, so the claim follows by induction.

2.2. Path polytopes on trees. We follow the notation from [16].

A set $P \subset \mathbb{R}^d$ is a *polytope* if there exist points $\mathbf{v}_1, \ldots, \mathbf{v}_k \in \mathbb{R}^d$ such that

$$P = \operatorname{conv}(\{\mathbf{v}_1, \dots, \mathbf{v}_k\}) = \left\{\sum_{i=1}^k \lambda_i \mathbf{v}_i \mid \lambda_i \ge 0, \sum_{i=1}^k \lambda_i = 1\right\}.$$

The set $\mathcal{V}(P) = {\mathbf{v}_1, \dots, \mathbf{v}_k}$ is called a \mathcal{V} - representation of P. Given a vector $\mathbf{a} \in \mathbb{R}^d$ and a scalar $b \in \mathbb{R}$, a linear inequality $\mathbf{a}^\top \mathbf{x} \leq b$ is *valid* for P if it is satisfied for all points $\mathbf{x} \in P$. A *face* of P is any set of the form

$$F = P \cap \{ \mathbf{x} \in \mathbb{R}^d \mid \mathbf{a}^\top \mathbf{x} = b \},$$

Page 3

where $\mathbf{a}^{\top}\mathbf{x} \leq b$ is a valid inequality for P. The *dimension* of a face F is the dimension of its affine hull $\operatorname{aff}(F)$, i.e., the smallest affine space containing F. We denote $\dim(F) = \dim(\operatorname{aff}(F))$. Similarly, the dimension of a polytope P is $\dim(\operatorname{aff}(P))$. A d-dimensional polytope is called a d-polytope. Given a face $F = P \cap \{x \in \mathbb{R}^d \mid \mathbf{a}^{\top}\mathbf{x} = b\}$, the equation $\mathbf{a}^{\top}\mathbf{x} = b$ is called the supporting affine space of F. A face F of dimension $\dim(P) - 1$ is called a facet of P, and a corresponding valid inequality $\mathbf{a}^{\top}\mathbf{x} \leq b$ is a halfspace of P. The set of affine spaces that contain P together with a set of halfspaces which describe the facets of P is called an \mathcal{H} -representation of P. Given two positive integers k, n, the (n, k)-hypersimplex is

$$\Delta_{n,k} = \left\{ (x_1, \dots, x_n) \in \mathbb{R}^n \mid \sum_{i=1}^n x_i = k, \ 0 \le x_1, \dots, x_n \le 1 \right\}.$$

When k = 1, $\Delta_{n,1}$ is the standard simplex Δ_n .

Consider a tree T = (V, E). Recall that for two vertices $u, v \in V$, $u \leftrightarrow v$ denotes the set of edges in the path that connects u and v, and the *edge indicator vector* $\mathbf{c}_T^{u \leftrightarrow v} = (c_e)_{e \in E} \in \mathbb{R}^E$ has $c_e = 1$ if $e \in u \leftrightarrow v$ and $c_e = 0$ otherwise. For simplicity, we will use $\mathbf{c}^{u \leftrightarrow v}$ instead of $\mathbf{c}_T^{u \leftrightarrow v}$ if there is no risk of ambiguity. The *path polytope* of the tree T is

$$P_T = \operatorname{conv}(\{\mathbf{c}_T^{i \leftrightarrow j} \mid i, j \in \operatorname{Lv}(T)\}) \subset \mathbb{R}^E.$$

The next important notion for us is *free join* of two polytopes, which refers to their union when they lie in *skew* affine spaces, i.e., affine spaces that are neither parallel or intersecting. More precisely, let P, Q be two polytopes such that $\dim(\operatorname{conv}(P \cup Q)) = \dim(P) + \dim(Q) + 1$. Then $\operatorname{conv}(P \cup Q)$ is called the *free join* of P and Q and is denoted $P \bigotimes Q$. We are interested in taking the convex hull of path polytopes of trees with the origin. The next result shows that it is a free join, see Figure 3.

Proposition 2.2. Given a tree T with at least two edges, the polytope P_T lives in the hyperplane defined by $\sum_{e \in E_{\text{leaf}}(T)} x_e = 2$. Consequently, $\operatorname{conv}(P_T \cup \{\mathbf{0}\}) = P_T \bigotimes \{\mathbf{0}\}$.

Proof. Every vertex $\mathbf{c}^{i \leftrightarrow j}$ of P_T satisfies the equation $\sum_{e \in E_{\text{leaf}}(T)} x_e = 2$. Hence, P_T lives in this hyperplane that does not contain the origin. This implies that $\text{conv}(P_T \cup \{\mathbf{0}\})$ is the free join $P_T \bigotimes \{\mathbf{0}\}$.



FIGURE 3. Path polytope of a star tree and its free join with the origin.

The following result describes the faces of the free join of two polytopes, which is needed to compute facets of path polytopes inductively.

Lemma 2.3 ([12, Proposition 2.1]). The faces of $P \bigotimes Q$ are precisely the sets of the form $F \bigotimes G$, where F is a face of P and G is a face of Q (including $F = \emptyset$ or P, and $G = \emptyset$ or Q).

2.3. Toric fiber products of polytopes. The notion of toric fiber product was introduced by Sullivant in [15] on toric ideals. We follow the notation from [9], which adapts this concept to polytopes.

An *integral* polytope is a polytope whose vertices all have integer coordinates. Given an integral polytope $P \subset \mathbb{R}^n$, a projection $\pi : P \to \mathbb{R}^m$ is called *integral* if $\pi(P)$ is an integral polytope. Given integral polytopes P_1 and P_2 and integral projections $\pi_1 : P_1 \to Q$ and $\pi_2 : P_2 \to Q$ for some polytope $Q = \pi_1(P_1) = \pi_2(P_2)$, the *toric fiber product* of P_1 and P_2 is

 $P_1 \times_Q P_2 = \operatorname{conv}(\{(\mathbf{x}, \mathbf{y}) \in \mathcal{V}(P_1) \times \mathcal{V}(P_2) \mid \pi_1(\mathbf{x}) = \pi_2(\mathbf{y})\}).$

This gives the toric fiber product polytope $P_1 \times_Q P_2$ by its \mathcal{V} -representation. We can retrieve the facets $P_1 \times_Q P_2$ when Q is a standard simplex using the following result.

Lemma 2.4 ([5, Lemma 3.2]). Let P_1 and P_2 be two polytopes. Let $\pi_i : P_i \to \mathbb{R}^n$, for i=1,2, be integral projections such that $\pi_1(P_1) = \pi_2(P_2) = \Delta_n$. Then all facets of the toric fiber product $P_1 \times_{\Delta_n} P_2$ are of the form $F_1 \times_{\Delta_n} P_2$ or $P_1 \times_{\Delta_n} F_2$, where F_i is a facet of P_i .

3. PATH POLYTOPES OF TREES VIA TORIC FIBER PRODUCTS

We use toric fiber products to construct the path polytope of a tree using path polytopes of its subtrees. More precisely, just as a T can be formed using gluing on stars S_{n_1}, \ldots, S_{n_k} , we show that the polytope P_T can be obtained by toric fiber products on $P_{S_{n_1}} \otimes \{0\}, \ldots, P_{S_{n_k}} \otimes \{0\}$. This is done by defining integral projections π_1 and π_2 to align corresponding indicator vectors. The resulting matched vectors describe connected paths in the new tree, forming a larger indicator vector for the glued structure. However, some paths between leaves of one subtree need not be glued with paths from different subtrees, requiring the inclusion of the origin in the toric fiber product.

Consider two trees T_1 and T_2 , and two edges $e_1 \in E_{\text{leaf}}(T_1)$ and $e_2 \in E_{\text{leaf}}(T_2)$. We call gluing integral projections for (T_1, T_2, e_1, e_2) to a pair of integral projections $\pi_i : P_{T_i} \bigotimes \{\mathbf{0}\} \to \Delta_3$ (i = 1, 2) such that $\pi_1(\mathbf{0}) = (0, 0, 1), \pi_2(\mathbf{0}) = (1, 0, 0),$

$$\pi_1(\mathbf{c}_{T_1}^{i \leftrightarrow j}) = \begin{cases} (1,0,0) & \text{if } e_1 \notin E(i \leftrightarrow j) \\ (0,1,0) & \text{if } e_1 \in E(i \leftrightarrow j), \end{cases} \quad \pi_2(\mathbf{c}_{T_2}^{i \leftrightarrow j}) = \begin{cases} (0,1,0) & \text{if } e_2 \in E(i \leftrightarrow j) \\ (0,0,1) & \text{if } e_2 \notin E(i \leftrightarrow j). \end{cases}$$

Example 3.1. Consider two trees T_1 and T_2 , and two edges $e_1 \in E_{\text{leaf}}(T_1)$ and $e_2 \in E_{\text{leaf}}(T_2)$. Let $\{\mathbf{b}_e^{(i)} \mid e \in E(T_i)\}$ denote the standard basis of $\mathbb{R}^{E(T_i)}$ for i = 1, 2. Let $\pi_i : \mathbb{R}^{E(T_i)} \to \mathbb{R}^3$ for i = 1, 2 be defined by

$$\pi_{1}(\mathbf{b}_{e}^{(1)}) = \begin{pmatrix} \frac{1}{2} \mathbb{1}_{E_{\text{leaf}}(T_{1})}(e) - \mathbb{1}_{\{e_{1}\}}(e) \\ \mathbb{1}_{\{e_{1}\}}(e) \\ -\frac{1}{2} \mathbb{1}_{E_{\text{leaf}}(T_{1})}(e) + 1 \end{pmatrix}, \quad \pi_{2}(\mathbf{b}_{e}^{(2)}) = \begin{pmatrix} -\frac{1}{2} \mathbb{1}_{E_{\text{leaf}}(T_{2})}(e) + 1 \\ \mathbb{1}_{\{e_{2}\}}(e) \\ \frac{1}{2} \mathbb{1}_{E_{\text{leaf}}(T_{2})}(e) - \mathbb{1}_{\{e_{2}\}}(e) \end{pmatrix},$$

where $\mathbb{1}_{S}(x)$ denotes the characteristic function for the set S. Then π_1, π_2 is a pair of gluing integral projections.

Example 3.2. Let $T_1 = S_3$ with center 1 and leaves 2, 3, 4, let $T_2 = S_3$ with center 5 and leaves 6, 7, 8, and let $T = T_1 *_{\{1,4\},\{8,5\}} T_2$, as in Figure 2. Let π_1, π_2 be gluing integral projections and let $Q = (P_{T_1} \bigotimes \{0\}) \times_{\Delta_3} (P_{T_2} \bigotimes \{0\})$. The tables below show the \mathcal{V} -representation for each of the polytopes involved, revealing that $P_T \cong Q$ (by mapping the edges $\{1,4\}, \{8,5\}$ to $\{1,5\}$ and leaving the rest fixed). We use colors to denote which vectors are matched together by the gluing integral projections, using the color code given by $\Delta_3 = \operatorname{conv}((1,0,0), (0,1,0), (0,0,1))$.

$\mathcal{V}(P_{T_1} \bigotimes \{0\})$	$\mathbf{c}^{2\leftrightarrow 3}$	$c^{2\leftrightarrow 4}$	$\mathbf{c}^{3\leftrightarrow 4}$	0	$\mathcal{V}(Q)$	$\mathbf{c}^{2\leftrightarrow 3}$	$\mathbf{c}^{6\leftrightarrow7}$	$\mathbf{c}^{2\leftrightarrow 6}$	$\mathbf{c}^{2\leftrightarrow7}$	$\mathbf{c}^{3\leftrightarrow 6}$	$\mathbf{c}^{3\leftrightarrow7}$
{1,2}	1	1	0	0	{1,2}	1	0	1	1	0	0
{1,3}	1	0	1	0	{1,3}	1	0	0	0	1	1
{1,4}	0	1	1	0	{1.4}	0	0	1	1	1	1
$\mathcal{V}(P_{T_2} \bigotimes \{0\})$	$\mathbf{c}^{6\leftrightarrow7}$	$\mathbf{c}^{8\leftrightarrow 6}$	$\mathbf{c}^{8\leftrightarrow7}$	0	{ 5 8}		0	1	1	1	1
{5,8}	0	1	1	0	(5,0)	0	0	1	1	1	1
{5,6}	1	1	0	0	{5,6}	0	I	I	0	I	0
{5,7}	1	0	1	0	{5,7}	0	1	0	1	0	1
	_	$\mathcal{V}(P_T)$	$\mathbf{c}^{2\leftrightarrow 3}$	$\mathbf{c}^{6\leftrightarrow7}$	$\mathbf{c}^{2\leftrightarrow 6}$	$\mathbf{c}^{2\leftrightarrow7}$	$\mathbf{c}^{3\leftrightarrow 6}$	$\mathbf{c}^{3\leftrightarrow7}$			
		{1,2}	1	0	1	1	0	0			
		{1,3}	1	0	0	0	1	1			
		{1,5}	0	0	1	1	1	1			
		{5,6}	0	1	1	0	1	0			
		{5,7}	0	1	0	1	0	1			

Theorem 3.3. Consider a gluing of two trees $T = T_1 *_{e_1,e_2} T_2$. Let $\pi_i : P_{T_i} \bigotimes \{\mathbf{0}\} \to \Delta_3$ (i = 1, 2) be a pair of gluing integral projections for (T_1, T_2, e_1, e_2) . Then,

 $P_T \cong (P_{T_1} \bigotimes \{\mathbf{0}\}) \times_{\Delta_3} (P_{T_2} \bigotimes \{\mathbf{0}\}).$

Proof. Let $Q = (P_{T_1} \bigotimes \{\mathbf{0}\}) \times_{\Delta_3} (P_{T_2} \bigotimes \{\mathbf{0}\})$. It suffices to construct an affine transformation ϕ that gives a bijection between the vertices of Q and the vertices of P_T . Let $e_1 = \{u_1, k_1\}, e_2 = \{u_2, k_2\}$ where $k_i \in \operatorname{Lv}(T_i)$ for i = 1, 2. Let $\{\mathbf{a}_e \mid e \in E(T_1) \cup E(T_2)\}$ be the standard basis of $\mathbb{R}^{E(T_1)} \times \mathbb{R}^{E(T_2)} \cong \mathbb{R}^{E(T_1) \cup E(T_2)}$, and let $\{\mathbf{b}_e \mid e \in E(T)\}$ be the standard basis of $\mathbb{R}^{E(T)}$. Consider the affine map given by $\phi(\mathbf{a}_{e_1}) = \phi(\mathbf{a}_{e_2}) = \frac{1}{2}\mathbf{b}_{\{u_1, u_2\}}$ and $\phi(\mathbf{a}_e) = \mathbf{b}_e$ if $e \neq e_1, e_2$.

The vertices of Q can be divided in three classes, one for each vertex of Δ_3 . First, given two distinct leaves $i, j \in \operatorname{Lv}(T_1) \setminus \{k_1\}$, the vertex $(\mathbf{c}_{T_1}^{i \leftrightarrow j}, \mathbf{0}) \in \mathcal{V}(Q)$ is mapped to $\mathbf{c}_T^{i \leftrightarrow j} \in \mathcal{V}(P_T)$ under ϕ . Second, given two distinct leaves $i, j \in \operatorname{Lv}(T_2) \setminus \{k_2\}$, the vertex $(\mathbf{0}, \mathbf{c}_{T_1}^{i \leftrightarrow j}) \in \mathcal{V}(Q)$ is mapped to $\mathbf{c}_T^{i \leftrightarrow j} \in \mathcal{V}(P_T)$ under ϕ . Finally, given $i \in \operatorname{Lv}(T_1) \setminus \{k_1\}$ and $j \in \operatorname{Lv}(T_2) \setminus \{k_2\}$ the vertex $(\mathbf{c}_{T_1}^{i \leftrightarrow k_1}, \mathbf{c}_{T_2}^{k_2 \leftrightarrow j}) \in \mathcal{V}(Q)$ is mapped to $\mathbf{c}_T^{i \leftrightarrow j} \in \mathcal{V}(P_T)$ under ϕ . We have considered all the vertices of both Q and P_T , so $P_T \cong Q$.

4. PROOF OF OUR MAIN THEOREM

In this section, we prove Theorem 1.1. We use the decomposition from Theorem 3.3 together with Lemma 2.3 and Lemma 2.4 to obtain explicit descriptions for the facets of P_T . Lemma 2.1 show that every tree can be decomposed into star trees. The following result shows that Theorem 1.1 is true for star trees, which will serve as a base case of our inductive argument.

Lemma 4.1. Let $S_n = (V, E)$ be the star tree on $n \ge 3$ leaves, and let u be the only internal node of S_n , that is, $N(u) = Lv(S_n)$. Then, $P_{S_n} = \Delta_{n,2}$. Hence, if $n \ge 4$ an \mathcal{H} -representation of P_{S_n} is given by

$$\begin{cases} x_{\{u,v\}} & \ge 0 \quad \text{for all } \{u,v\} \in E \\ -x_{\{u,v\}} + \sum_{w \in N(u) \setminus \{v\}} x_{\{u,w\}} & \ge 0 \quad \text{for all } v \in N(u) \\ \sum_{e \in E} x_e & = 2. \end{cases}$$

The corresponding set of facets is $\mathcal{F} = \{F^{\{u,v\}} \mid \{u,v\} \in E\} \cup \{G^{(u,v)} \mid v \in N(u)\}$ where

$$F^{\{u,v\}} = \operatorname{conv}\left(\{\mathbf{c}^{i\leftrightarrow j} \mid i, j \in \operatorname{Lv}(S_n), \{u,v\} \notin i \leftrightarrow j\}\right)$$
$$G^{(u,v)} = \operatorname{conv}\left(\{\mathbf{c}^{i\leftrightarrow v} \mid i \in \operatorname{Lv}(S_n) \setminus \{v\}\}\right).$$

When n = 3, the first class of halfspaces is redundant and the sets $F^{\{u,v\}}$ are not facets, but the rest of the statement holds.

Proof. By construction, we have

$$P_{S_n} = \operatorname{conv}\left(\left\{ (x_{\{u,v_1\}}, \dots, x_{\{u,v_n\}}) \in \{0,1\}^n \mid \sum_{e \in E} x_e = 2 \right\} \right) = \Delta_{n,2}.$$

The case n = 3 is shown in Figure 1. Assume that $n \ge 4$. For a star tree, $E_{\text{leaf}}(T) = E$, so $\sum_{e \in E} x_e = 2$, by Proposition 2.2. By definition of $\Delta_{n,2}$, the halfspaces of P_{S_n} are $0 \le x_e \le 1$ for all $e \in E$. Subtracting $2x_e \le 2$ from $\sum_{e \in E} x_e = 2$ we get the desired \mathcal{H} -representation. Finally, given any $v \in \text{Lv}(S_n)$, we have $F^{\{u,v\}} = P_{S_n} \cap \{x_e = 0\}, G^{(u,v)} = P_{S_n} \cap \{x_e = 1\}$, and $\dim(F^{\{u,v\}}) = \dim(G^{(u,v)}) = n - 2 = \dim(P_{S_n}) - 1$, so the statement follows. \Box

We are now ready to compute the dimension of path polytopes. Proposition 2.2 implies that P_T has codimension at least one. The next result implies that the codimension of P_T is one provided that T has no internal nodes of degree two.

Proposition 4.2. Given a tree T = (V, E) with |V| > 2, the dimension of the path polytope P_T is $|E| - |\{u \in V \mid \deg(u) = 2\}| - 1$, and P_T is contained in the linear space defined by

$$\begin{cases} x_{\{u,v\}} - x_{\{u,w\}} &= 0 \text{ for all } u \in \text{Int}(V) \text{ such that } N(u) = \{v,w\} \\ \sum_{e \in E_{\text{leaf}}(T)} x_e &= 2. \end{cases}$$

Proof. Let $u \in V$ such that $N(u) = \{v, w\}$. For any pair of distinct leaves $i, j \in Lv(T)$, we have $\{u, v\} \in i \leftrightarrow j$ if and only if $\{u, w\} \in i \leftrightarrow j$. Hence, every point in the polytope satisfies the first set of equations. The polytope also satisfies the last equation by Proposition 2.2. These equations are linearly independent, so the dimension of the polytope is at most

$$|E| - |\{v \in V \mid \deg(v) = 2\}| - 1.$$

Page 7

We show that it is equal to this quantity. Consider a gluing of two trees $T = T_1 *_{e_1,e_2} T_2$. Let $\pi_i : P_{T_i} \bigotimes \{\mathbf{0}\} \to \Delta_3$ (i = 1, 2) be a pair of gluing integral projections for (T_1, T_2, e_1, e_2) . Then

$$\dim(P_T) = \dim((P_{T_1} \bigotimes \{\mathbf{0}\}) \times_{\Delta_3} (P_{T_1} \bigotimes \{\mathbf{0}\})) =$$

= dim(P_{T_1} \bigotimes \{\mathbf{0}\}) + (P_{T_2} \bigotimes \{\mathbf{0}\}) - dim(\Delta_3) =
= (dim(P_{T_1}) + 1) + (dim(P_{T_2}) + 1) - 2 =
= dim(P_{T_1}) + dim(P_{T_2}).

Hence, if $T = S_{n_1} *_{e_1, e_2} \cdots *_{e_{r-1}, e_r} S_{n_r}$ where S_{n_i} is a star tree, then $\dim(P_T) = \sum_{i=1}^r \dim(P_{S_{n_i}})$ and $|E| = 1 + \sum_{i=1}^r (n_i - 1)$. Finally, $\dim(P_{S_2}) = 0$ and $\dim(P_{S_n}) = n - 1$ for n > 2 by Lemma 4.1, so the statement follows.

Theorem 1.1 omits the case when T has internal nodes of degree 2. This is because the polytope P_T is isomorphic $P_{T'}$ where T' is formed from T by merging edges $\{u, v\}$ and $\{u, w\}$ to the edge $\{v, w\}$ for any vertex u of degree 2. Both trees have the same set of leaves. This follows from the following result.

Proposition 4.3. If $T = T' *_{e_1,e_2} S_2$, then $P_T \cong P_{T'}$.

Proof. Let $f \in E(T)$ be the edge resulting from identifying e_1 with e_2 , and let $g \in E(T)$ be the edge in S_2 distinct from e_2 . Let $\{\mathbf{a}_e \mid e \in E(T')\}$ be the standard basis of $\mathbb{R}^{E(T')}$, and let $\{\mathbf{b}_e \mid e \in E(T)\}$ be the standard basis of $\mathbb{R}^{E(T)}$. Consider the affine map $\phi : \mathbb{R}^{E(T')} \to \mathbb{R}^{E(T)}$ given by $\phi(\mathbf{a}_e) = \mathbf{b}_e$ if $e \neq e_1$ and $\phi(\mathbf{a}_{e_1}) = \mathbf{b}_f + \mathbf{b}_g$. The map ϕ is injective and maps the vertices of $P_{T'}$ to the vertices of P_T , so $P_T \cong P_{T'}$.

Propositions 4.2 and 4.3 imply that we can assume without loss of generality that our trees do not have internal nodes of degree two, so that the corresponding path polytope has codimension one.

Theorem 4.4. Let T = (V, E) be a tree with $|E| \ge 3$ and with no internal nodes of degree 2. Then, the facets of P_T are

$$\{F_T^{\{u,v\}} \mid \{u,v\} \in E, \deg(u), \deg(v) \neq 3\} \cup \{G_T^{(u,v)} \mid u \in \operatorname{Int}(T), v \in N_T(u)\}\$$

where

$$F_{T}^{\{u,v\}} = \operatorname{conv}\left(\{\mathbf{c}^{i\leftrightarrow j} \mid i, j \in \operatorname{Lv}(T), \{u,v\} \notin i \leftrightarrow j\}\right) = P_{T} \cap \{x \in \mathbb{R}^{E(T)} \mid x_{\{u,v\}} = 0\}, \text{ and}$$

$$G_{T}^{(u,v)} = \operatorname{conv}\left(\{\mathbf{c}^{i\leftrightarrow j} \mid i, j \in \operatorname{Lv}(T), \{u,v\} \in i \leftrightarrow j \text{ or } \{u,w\} \notin i \leftrightarrow j \text{ for any } w \in N_{T}(u)\}\right) =$$

$$= P_{T} \cap \{x \in \mathbb{R}^{E(T)} \mid -x_{\{u,v\}} + \sum_{w \in N(u) \setminus \{v\}} x_{\{u,w\}} = 0\}.$$

Proof. We prove it by strong induction on r = |Int(T)|. The case r = 1 follows from Lemma 4.1. Suppose the claim holds for all trees with at most r - 1 internal nodes and let T have r internal nodes. Let $T = T_1 *_{e_1,e_2} T_2$, where T_1 and T_2 are trees with fewer than r internal nodes (see Lemma 2.1). Let $e_1 = \{u_1, k_1\}$ and $e_2 = \{u_2, k_2\}$ with $k_1 \in \text{Lv}(T_1)$ and $k_2 \in \text{Lv}(T_2)$. For simplicity of notation, let $\hat{P}_{T_i} = P_{T_i} \bigotimes \{\mathbf{0}\}$ for i = 1, 2. Hence, $P_T \cong \hat{P}_{T_1} \times_{\Delta_3} \hat{P}_{T_2}$ by Theorem 3.3. The facets of P_T are isomorphic to facets of the form $\{F_1 \times_{\Delta_3} \hat{P}_{T_2} \mid F_1 \text{ facet of } \hat{P}_{T_1}\}$ and $\{\hat{P}_{T_1} \times_{\Delta_3} \hat{P}_{T_2} \mid F_1$ facet of $\hat{P}_{T_1}\}$ and $\{\hat{P}_{T_1} \times_{\Delta_3} \hat{P}_{T_2} \mid F_1$ facet of $\hat{P}_{T_1}\}$ and $\{\hat{P}_{T_1} \times_{\Delta_3} \hat{P}_{T_2} \mid F_1$ facet of $\hat{P}_{T_1}\}$ and $\{\hat{P}_{T_1} \times_{\Delta_3} \hat{P}_{T_2} \mid F_1 \text{ facet of } \hat{P}_{T_1}\}$ and $\{\hat{P}_{T_1} \times_{\Delta_3} \hat{P}_{T_2} \mid F_1 \text{ facet of } \hat{P}_{T_1}\}$ and $\{\hat{P}_{T_1} \times_{\Delta_3} \hat{P}_{T_2} \mid F_1 \text{ facet of } \hat{P}_{T_1}\}$. $F_2 \mid F_2$ facet of \hat{P}_{T_2} by Lemma 2.4. By Lemma 2.3 and the induction hypothesis, the facets of P_{T_i} for i = 1, 2 are

- (1) $F_{T_i}^{\{u,v\}} \bigotimes \{\mathbf{0}\}$ for $\{u,v\} \in E(T_i)$ with $\deg(u), \deg(v) \neq 3$,
- (2) $G_{T_i}^{(u,v)} \bigotimes \{\mathbf{0}\}$ for $u \in \text{Int}(T_i)$ and v in the neighborhood $N_{T_i}(v)$ of v in T_i , and

(3)
$$P_{T_i}$$
.

We focus on the facets of the form $\{F_1 \times_{\Delta_3} \hat{P}_{T_2} \mid F_1 \text{ facet of } \hat{P}_{T_1}\}$. By symmetry, the same reasoning holds for the facets $\{\hat{P}_{T_1} \times_{\Delta_3} F_2 \mid F_2 \text{ facet of } \hat{P}_{T_2}\}.$

<u>Case 1:</u> Consider sets of the form $(F_{T_1}^{\{u,v\}} \bigotimes \{0\})$ for $\{u,v\} \in E(T_1)$. First, let $\{u,v\} = e_1 = e_1$ $\{u_1, k_1\} \in E(T_1)$. Then,

$$(F_{T_1}^{\{u_1,k_1\}} \bigotimes \{\mathbf{0}\}) \times_{\Delta_3} \hat{P}_{T_2} = \operatorname{conv} \left((F_{T_1}^{\{u_1,k_1\}} \times \{\mathbf{0}\}) \cup (\{\mathbf{0}\} \times F_{T_2}^{\{u_2,k_2\}}) \right).$$

which is isomorphic to $F_T^{\{u_1, u_2\}}$. Note that

$$F_{T_1}^{\{u_1,k_1\}} \times \{\mathbf{0}\} \subset L_1 \coloneqq \left\{ x \in \mathbb{R}^{E(T_1) \cup E(T_2)} \mid x_e = 0 \text{ for all } e \in E(T_2), \sum_{e \in E_{\text{leaf}}(T_1) \setminus \{e_1\}} x_e = 2 \right\},$$

$$\{\mathbf{0}\} \times F_{T_2}^{\{u_2,k_2\}} \subset L_2 \coloneqq \left\{ x \in \mathbb{R}^{E(T_1) \cup E(T_2)} \mid x_e = 0 \text{ for all } e \in E(T_1), \sum_{e \in E_{\text{leaf}}(T_2) \setminus \{e_2\}} x_e = 2 \right\}.$$

Since L_1 and L_2 are skew linear spaces, we have the free join $(F_{T_1}^{\{u_1,k_1\}} \times \{\mathbf{0}\}) \bigotimes (\{\mathbf{0}\} \times F_{T_2}^{\{u_2,k_2\}})$. By induction hypothesis, $F_{T_i}^{\{u_i,k_i\}}$ is a facet of P_{T_i} $(\dim(F_{T_i}^{\{u_i,k_i\}}) = \dim(P_{T_1}) - 1 = E(T_i) - 2)$ if and only if $deg(u_i) > 3$, for i = 1, 2. Hence,

$$\dim(F_T^{\{u_1,u_2\}}) = \dim(F_{T_1}^{\{u_1,k_1\}}) + \dim(F_{T_2}^{\{u_2,k_2\}}) + 1 \le (|E(T_1)| - 2) + (|E(T_2)| - 2) - 1 = |E(T)| - 2 = \dim(P_T) - 1$$

with equality if and only if $\deg(u_1), \deg(u_2) > 3$. So $F_T^{\{u_1, u_2\}} = P_T \cap \{x \in \mathbb{R}^{E(T)} \mid x_{\{u_1, u_2\}} = 0\}$ is a facet of P_T if and only if $\deg(u_1), \deg(u_2) > 3$.

Next, let $\{u,v\} \in E(T_1) \setminus \{e_1\}$ such that $\deg(u), \deg(v) \neq 3$. Assume that $u \in Int(T_1)$, so $\deg(u) \ge 4$. We have

$$(F_{T_1}^{\{u,v\}} \bigotimes \{\mathbf{0}\}) \times_{\Delta_3} \hat{P}_{T_2} = \operatorname{conv} \left(\{ (\mathbf{c}_{T_1}^{i \leftrightarrow k_1}, \mathbf{c}_{T_2}^{k_2 \leftrightarrow j}) \mid \mathbf{c}_{T_1}^{i \leftrightarrow k_1} \in F_{T_1}^{\{u,v\}}, \mathbf{c}_{T_2}^{k_2 \leftrightarrow j} \in P_{T_2} \} \\ \cup \{ (\mathbf{c}_{T_1}^{i \leftrightarrow j}, \mathbf{0}) \mid \mathbf{c}_{T_1}^{i \leftrightarrow j} \in F_{T_1}^{\{u,v\}}, i, j \neq k_1 \} \\ \cup \{ (\mathbf{0}, \mathbf{c}_{T_2}^{i \leftrightarrow j}) \mid \mathbf{c}_{T_2}^{i \leftrightarrow j} \in P_{T_2}, i, j \neq k_2 \} \right),$$

which is isomorphic to $F_T^{\{u,v\}}$. Fix two leaves $i, j \in \operatorname{Lv}(T)$ such that $\mathbf{c}_T^{i \leftrightarrow j} \notin F_T^{\{u,v\}}$, that is, $\{u,v\} \in i \leftrightarrow j$. Note that $F_T^{\{u,v\}} \subset \{x \in \mathbb{R}^{E(T)} \mid x_{\{u,v\}} = 0\} \not\ni \mathbf{c}_T^{i \leftrightarrow j}$. Hence, we have the free join $F_T^{\{u,v\}} \bigotimes \mathbf{c}_T^{i \leftrightarrow j}$. We show that $\operatorname{aff}(P_T) = \operatorname{aff}(F_T^{\{u,v\}} \bigotimes \mathbf{c}_T^{i \leftrightarrow j})$. Indeed, consider any two leaves i', j' such that $\mathbf{c}_T^{i \leftrightarrow j} \in P_T \setminus F_T^{\{u,v\}}$. Without loss of generality, suppose that $i \neq i', \{u,v\} \notin i \leftrightarrow i'$, and $\{u,v\} \in i \leftrightarrow v$ (permute the roles of i, j and i', j' if needed). Since $\deg(u) > 3$, there exists a leaf $k \in \operatorname{Lv}(T) \setminus \{i, i'\}$ such that $\mathbf{c}_T^{i \leftrightarrow k}, \mathbf{c}_T^{i' \leftrightarrow k} \in F_T^{\{u,v\}}$. We need to consider two cases. If Page 9

 $\deg(v) = 1, \text{ then } j = j' = v \text{ and } \mathbf{c}_T^{i' \leftrightarrow j} = \mathbf{c}_T^{i \leftrightarrow j} + \mathbf{c}_T^{i' \leftrightarrow k} - \mathbf{c}_T^{i \leftrightarrow k} \in \operatorname{aff}(F_T^{\{u,v\}} \bigotimes \mathbf{c}_T^{i \leftrightarrow j}). \text{ If } \deg(v) > 3,$ then there exists a leaf $l \in \operatorname{Lv}(T) \setminus \{j, j'\}$ such that $\mathbf{c}_T^{j \leftrightarrow l}, \mathbf{c}_T^{j' \leftrightarrow l} \in F_T^{\{u,v\}}$, so

$$\mathbf{c}_T^{i'\leftrightarrow j'} = \mathbf{c}_T^{i\leftrightarrow j} + \mathbf{c}_T^{i'\leftrightarrow k} - \mathbf{c}_T^{i\leftrightarrow k} + \mathbf{c}_T^{j'\leftrightarrow l} - \mathbf{c}_T^{j\leftrightarrow l} \in \operatorname{aff}(F_T^{\{u,v\}} \otimes \mathbf{c}_T^{i\leftrightarrow j}).$$

We have seen that every vertex of P_T lies in $\operatorname{aff}(F_T^{\{u,v\}} \otimes \mathbf{c}_T^{i \leftrightarrow j})$, so $\operatorname{aff}(P_T) = \operatorname{aff}(F_T^{\{u,v\}} \otimes \mathbf{c}_T^{i \leftrightarrow j})$. Therefore, $\dim(F_T^{\{u,v\}}) = \dim(P_T) - 1$, so $F_T^{\{u,v\}} = P_T \cap \{x \in \mathbb{R}^{E(T)} \mid x_{\{u,v\}} = 0\}$ is a facet of P_T .

<u>Case 2</u>: Consider sets of the form $(G_{T_1}^{(u,v)} \bigotimes \{\mathbf{0}\}) \times_{\Delta_3} \hat{P}_{T_2}$ where $u \in \text{Int}(T_1)$ and $v \in N_{T_1}(u)$. We have

$$(G_{T_1}^{(u,v)} \bigotimes \{\mathbf{0}\}) \times_{\Delta_3} \hat{P}_{T_2} = \operatorname{conv} \left(\{ (\mathbf{c}_{T_1}^{i \leftrightarrow k_1}, \mathbf{c}_{T_2}^{k_2 \leftrightarrow j}) \mid \mathbf{c}_{T_1}^{i \leftrightarrow k_1} \in G_{T_1}^{(u,v)}, \mathbf{c}_{T_2}^{k_2 \leftrightarrow j} \in P_{T_2} \} \\ \cup \{ (\mathbf{c}_{T_1}^{i \leftrightarrow j}, \mathbf{0}) \mid \mathbf{c}_{T_1}^{i \leftrightarrow j} \in G_{T_1}^{(u,v)}, i, j \neq k_1 \} \\ \cup \{ (\mathbf{0}, \mathbf{c}_{T_2}^{i \leftrightarrow j}) \mid \mathbf{c}_{T_2}^{i \leftrightarrow j} \in P_{T_2}, i, j \neq k_2 \} \right)$$

which is isomorphic to $G_T^{(u,v)}$ if $(u,v) \neq (u_1,k_1)$, and to $G_T^{(u_1,u_2)}$ if $(u,v) = (u_1,k_1)$. Suppose that $(u,v) \neq (u_1,k_1)$, the same reasoning applies to the other case. Fix two leaves $i, j \in Lv(T)$ such that $\mathbf{c}_T^{i \leftrightarrow j} \notin G_T^{(u,v)}$, so there exist $w_1, w_2 \in N_T(u) \setminus \{v\}$ such that $\{u, w_1\}, \{u, w_2\} \in i \leftrightarrow j$. Note that $G_T^{(u,v)} \subset \{x \in \mathbb{R}^{E(T)} \mid x_{\{u,v\}} = \sum_{w \in N(v) \setminus \{v\}} x_{\{u,w\}}\} \not\ni \mathbf{c}_T^{i \leftrightarrow j}$. Hence, we have the free join $G_T^{(u,v)} \bigotimes \{\mathbf{c}_T^{i \leftrightarrow j'}\}$. We show that $\operatorname{aff}(P_T) = \operatorname{aff}(G_T^{(u,v)} \bigotimes \mathbf{c}_T^{i \leftrightarrow j})$. Indeed, consider any two leaves i', j' such that $\mathbf{c}_T^{i' \leftrightarrow j'} \in P_T \setminus G_T^{(u,v)}$. Pick a leaf $k \in \operatorname{Lv}(T)$ such that $\{u, v\} \in i \leftrightarrow k$, so $\mathbf{c}_T^{i \leftrightarrow k} \in G_T^{(u,v)}$. For the same leaf k, we also have $\mathbf{c}_T^{j \leftrightarrow k}, \mathbf{c}_T^{j' \leftrightarrow k}, \mathbf{c}_T^{j' \leftrightarrow k} \in G_T^{(u,v)}$, so

$$\mathbf{c}_T^{i'\leftrightarrow j'} = \mathbf{c}_T^{i\leftrightarrow j} + \mathbf{c}_T^{i'\leftrightarrow k} - \mathbf{c}_T^{i\leftrightarrow k} + \mathbf{c}_T^{j'\leftrightarrow k} - \mathbf{c}_T^{j\leftrightarrow k} \in \operatorname{aff}(G_T^{(u,v)} \bigotimes \mathbf{c}_T^{i\leftrightarrow j}).$$

This implies that $\operatorname{aff}(P_T) = \operatorname{aff}(G_T^{(u,v)} \otimes \mathbf{c}_T^{i \leftrightarrow j})$. Therefore, $\dim(G_T^{(u,v)}) = \dim(P_T) - 1$, so $G_T^{(u,v)} = P_T \cap \{x \in \mathbb{R}^{E(T)} \mid x_{\{u,v\}} = \sum_{w \in N(v) \setminus \{v\}} x_{\{u,w\}}\}$ is a facet of P_T .

<u>Case 3:</u> Consider sets of the form $P_{T_1} \times_{\Delta_3} \hat{P}_{T_2}$. We have that

$$P_{T_1} \times_{\Delta_3} \hat{P}_{T_2} = \operatorname{conv} \left(\left\{ (\mathbf{c}^{T_1, i \leftrightarrow k_1}, \mathbf{c}^{T_2, k_2 \leftrightarrow j}) \mid i \in \operatorname{Lv}(T_1) \setminus \{k_1\}, j \in \operatorname{Lv}(T_2) \setminus \{k_2\} \right\} \\ \cup \left\{ (\mathbf{c}^{T_1, i \leftrightarrow j}, \mathbf{0}) \mid i, j \in \operatorname{Lv}(T_1) \setminus \{k_1\} \right\} \right)$$

is included in $\hat{P}_{T_1} \times_{\Delta_3} (G_{T_2}^{(u_2,k_2)} \bigotimes \{\mathbf{0}\}) \cong G_T^{(u_2,u_1)}$, so it is not a new facet of P_T .

We are now ready to prove Theorem 1.11 that is, (1) is a minimal \mathcal{H} -representation of the path polytope P_T .

Proof of Theorem 1.1. Consider a tree T with no internal nodes of degree 2. The last equality in (1) comes from Proposition 2.2. Every vertex $c^{i \leftrightarrow j}$ satisfies the inequalities and therefore every point in the polytope also satisfies the inequalities. By Theorem 4.4, every facet of P_T is the intersection of the polytope with a hyperplane corresponding to the boundary of a halfspace in the \mathcal{H} -representation given in (1). Hence, (1) is a minimal \mathcal{H} -representation. Lastly, $\dim(P_T) = |E| - 1$ by Proposition 4.2.

Corollary 4.5. Given a tree T = (V, E), let \mathcal{H} be the halfspaces given in Theorem 1.1. Then, P_T has dimension $|E| - \{u \in V \mid \deg(u) = 2\} - 1$ and a halfspace description

$$\mathcal{H} \cup \{x_{\{u,v\}} - x_{\{u,w\}} = 0 \mid u \in \operatorname{Int}(T), N(u) = \{v,w\}\}.$$

Remark 4.6. Note that the \mathcal{H} -representation from Corollary 4.5 is not minimal when internal nodes of degree 2 are present: some inequalities $x_{\{v,u\}} \ge 0$ are redundant. For example, let $N(v) = \{u, w\}$. Then $x_{\{v,u\}} - x_{\{v,w\}} = 0$ and $x_{\{v,u\}} \ge 0$ imply $x_{\{v,w\}} \ge 0$.

Remark 4.7. We expect the techniques presented in this paper to be useful to other polytopes defined by suitable "gluing" integral projections, provided that the polytopes are contained in a hyperplane that does not contain the origin (using Lemma 2.3). However, when the path polytope is constructed from all vectors $c^{i \leftrightarrow j}$ for any two nodes *i* and *j*, not only leaves (e.g., see [4]), the resulting polytope is full-dimensional. Hence, in this case one should seek alternative approaches.

ACKNOWLEDGMENTS

AG and AM were supported by the National Science Foundation under Grant DMS-2306672. AG was supported by an REU at the University of Michigan funded under this grant. AR was supported by fellowships from "la Caixa" Foundation (ID 100010434), with code LCF/BQ/EU23/12010097, and from RCCHU. Most of the work was done in Pierce Hall (SEAS) at Harvard University.

REFERENCES

- [1] Tobias Boege, Jane Ivy Coons, Christopher Eur, Aida Maraj, and Frank Roettger. Reciprocal maximum likelihood degrees of brownian motion tree models. *Le Matematiche*, 76(2):383–398, 2021.
- [2] Jane I Coons, Aida Maraj, Pratik Misra, and Miruna-Ştefana Sorea. Symmetrically colored gaussian graphical models with toric vanishing ideals. *SIAM Journal on Applied Algebra and Geometry*, 7(1):133–158, 2023.
- [3] Jane Ivy Coons, Shelby Cox, Aida Maraj, and Ikenna Nometa. Maximum likelihood degrees of brownian motion tree models: Star trees and root invariance. *arXiv preprint arXiv:2402.10322*, 2024.
- [4] Jane Ivy Coons, Joseph Cummings, Benjamin Hollering, and Aida Maraj. Generalized cut polytopes for binary hierarchical models. *Algebraic Statistics*, 14(1):17–36, 2023.
- [5] Rodica Dinu and Martin Vodička. Gorenstein property for phylogenetic trivalent trees. *Journal of Algebra*, 575:233–255, 2021.
- [6] Eliana Duarte and Christiane Görgen. Equations defining probability tree models. *Journal of Symbolic Computation*, 99:127–146, 2020.
- [7] Eliana Duarte, Benjamin Hollering, and Maximilian Wiesmann. Toric fiber products in geometric modeling. In *International Conference on Geometric Science of Information*, pages 494–503. Springer, 2023.
- [8] Stephen E Fienberg and Alessandro Rinaldo. Maximum likelihood estimation in log-linear models. *The Annals of Statistics*, pages 996–1023, 2012.
- [9] Benjamin Hollering, Joseph Johnson, and Liam Solus. Hyperplane representations of interventional characteristic imset polytopes, 2024.
- [10] Rui-Juan Jing, Marc Moreno Maza, and Delaram Talaashrafi. Complexity estimates for fourier-motzkin elimination. arXiv preprint arXiv:1811.01510, 2019.
- [11] Diane Maclagan and Bernd Sturmfels. *Introduction to tropical geometry*, volume 161. American Mathematical Society, 2021.
- [12] Peter McMullen. Constructions for projectively unique polytopes. Discrete Mathematics, 14:241–266, 1975.
- [13] Pratik Misra and Seth Sullivant. Gaussian graphical models with toric vanishing ideals. *Annals of the Institute of Statistical Mathematics*, 73:757–785, 2021.
- [14] David Speyer and Bernd Sturmfels. The tropical grassmannian. Adv. Geom, 4:389-411, 2004.
- [15] Seth Sullivant. Toric fiber products. arXiv preprint math/0602052, 2006.
- [16] Günter M Ziegler. Lectures on polytopes, volume 152. Springer Science & Business Media, 2012.

AMER GOEL, UNIVERSITY OF MICHIGAN

Email address: amergoel@umich.edu

AIDA MARAJ, MAX PLANCK INSTITUTE OF MOLECULAR CELL BIOLOGY AND GENETICS, AND CENTER FOR SYSTEMS BIOLOGY DRESDEN

Email address: maraj@mpi-cbg.de

ÁLVARO RIBOT, HARVARD UNIVERSITY

Email address: aribotbarrado@g.harvard.edu